

## PRELIMINARY RESULTS ON INTERMITTENT BEHAVIOUR OF MILLISECOND PULSAR SAX J1808.4-3658

M. T. Menna,<sup>1</sup> A. Papitto,<sup>1</sup> L. Burderi,<sup>1</sup> T. Di Salvo,<sup>2</sup> L. Stella,<sup>1</sup> and N. R. Robba<sup>3</sup>

**We analyzed RXTE data from the burst of the year 2000 of the X-ray millisecond binary pulsar SAX J1808.4-3658 with the intent of determining the new orbital parameters.**

We used the observations of SAX J1808.4-3658 performed by the Rossi X-Ray Timing Explorer (RXTE) when the source was again detected in outburst during the period January-March 2000 (Wijnands et al. 2001). In particular we examined the data from the Proportional Counter Array (PCA) (Jahoda et al. 1996). We first applied barycentric correction to the data using the optical coordinates of the source (Roche et al. 98).

This burst is much weaker than the previous 1998 burst. We extracted a series of light curves to determine intensity levels and have therefore focused our attention on the observation of the 2nd of Feb 2000, which has a count rate, background subtracted, consistent with a constant value of about 143 cts/s, one of the highest levels, where pulsations are most likely to be detected. Having verified the presence of pulsing behaviour calculating a power spectrum density which clearly shows the harmonic line at about 401 Hz, we then proceed to perform a series of epoch folding searches at the nominal spin period to follow the spin evolution along time and determine new orbital parameters.

The epoch folding searches are to be performed folding a number of spin cycles long enough to have enough photons to detect a signal, but short enough so that the spin period is not affected by the orbital motion. We fold over intervals of 100 s. We report in Fig. 1 an example of one of the subsets we found of Time (in seconds) *vs* Spin Period (in units of  $10^{-3}$  s). We then performed a global fit along all the time span of the observation (16 days) with a constant plus sinus function of the kind  $f(x) = c + A \sin(\nu t + \phi)$  and obtained the following values for the physical quantities: spin period




Fig. 1. Example plot of Time *vs* Spin Periods. The continuous curve represents the best fit.

$P_{spin} = 2.493918328 \cdot 10^{-3} s + / - 1 \cdot 10^{-9}$ , orbital period  $P_{orb} = 7248.95 s + / - 0.09 s$ , epoch of  $\pi/2$  mean longitude  $T_{nod} = 51576.1112 MJD + / - 2 \cdot 10^{-4}$ .

The puzzling behaviour that emerged is that, notwithstanding the constant flux level, on this time scale pulsations are not present in all the intervals. It can be seen in Fig. 1 by the lack of points along some zones of the curve which has been thoroughly explored. We are currently investigating in more detail the presence of this intermittent pulsating behaviour in the 2000 outburst data (Menna et al. 2004, in preparation).

### REFERENCES

- Roche, P. et al. 1998, IAUC, 6885  
 Jahoda, K. et al. 1996, Proc. SPIE, 2808, 59  
 Wijnands, R. et al. 2001, ApJ, 560, 892  
 Menna, M.T. et al. 2004, in preparation

<sup>1</sup>Osservatorio Astronomico di Roma, Monteporzio Catone (Roma), via di Frascati 33, 00040, Italy (menna@mporzio.astro.it).

<sup>2</sup>Astronomical Institute University of Amsterdam.

<sup>3</sup>Dipartimento di Astronomia, Università di Palermo.