

AD-A054 885

DELAWARE UNIV NEWARK DEPT OF STATISTICS AND COMPUTER--ETC F/G 12/1
ON THE MERITS OF AN 'APPROXIMATION' TO THE BUSY PERIOD OF THE G--ETC(U)
MAR 78 V RAMASWAMI, D M LUCANTONI AFOSR-77-3236

UNCLASSIFIED

TR-78/3

AFOSR-TR-78-0984

NL

| OF |

AD
A054885




END
DATE
FILMED
7-78

DDC

2

AD A 054885



AD No. _____
DDC FILE COPY

Department of
STATISTICS AND COMPUTER SCIENCE

DDC
RECEIVED
JUN 12 1978
B

UNIVERSITY OF DELAWARE
Newark, Delaware 19711

Approved for public release;
distribution unlimited.



AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)
NOTICE OF TRANSMITTAL TO DDC
this technical report has been reviewed and is
approved for public release IAW AFR 190-12 (7b).
Distribution is unlimited.
A. D. BLOSE
Technical Information Officer

2

6. ON THE MERITS OF AN APPROXIMATION TO THE BUSY PERIOD OF THE GI/G/1 QUEUE.

by

10 V. Ramaswami*
Purdue University & University of Delaware
and
David M. Lucantoni
University of Delaware

16 2304 17 A5 18 AFOSR 19 TR-78-0984

12 15p.

Department of Statistics
and
Computer Science

9 Technical Report, No. 78/3 14 TR-
11 March 1978

15 *Research sponsored by Air Force Office of Scientific Research Air Force Systems Command USAF, under Grant No. AFOSR-77-3236 at the Department of Statistics & Computer Science, University of Delaware, Newark, DE 19711. The U.S.G. is authorized to reproduce & distribute reprints for governmental purposes notwithstanding any copyright notation hereon.

DISTRIBUTION STATEMENT A
Approved for public release;
Distribution Unlimited

DDC
R
JUN 12 1978
REGISTRY

391 807

B SB

ABSTRACT

An approximation for computing the busy period characteristics in the GI/G/1 queue, suggested by D.Gross and C.M.Harris in their book "Fundamentals of Queueing Theory", is shown to be lacking in accuracy, and is therefore quite undesirable for practical use.

KEY WORDS

GI/G/1 queue, busy period, Phase Type distribution, Phase Type Renewal Process, computational probability.

ACCESSION	
NTIS	NTIS Section <input checked="" type="checkbox"/>
DDC	DDC Section <input type="checkbox"/>
UNCLASSIFIED	<input type="checkbox"/>
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	AVAIL. and/or SPECIAL
A	

1. Introduction:

In a recent book on queueing theory by Gross and Harris[1], an approximate method is suggested for obtaining the busy period characteristics of the GI/G/1 queue. Through a simple computational example, we demonstrate that this method can be severely inaccurate. It is possible to construct (simple) examples where the approximation is arbitrarily bad. Our discussion will show that the approximation could result in a gross under-estimation of the duration of the busy period and is therefore undesirable for use in practical situations.

We will draw our example from the class of GI/M/1 queues with inter-arrival times following a generalized Erlang distribution. These queues belong to the class of PH/M/1 queues, which in turn are a subclass of the N/G/1 queues, analyzed in detail by Ramaswami [5]. In Section 2, we provide a brief discussion of Phase Type distributions and Phase Type renewal processes and follow this in Section 3 with a discussion of the PH/M/1 queue with particular reference to its busy period. In Section 4 we discuss the approximate method suggested by Gross and Harris obtaining the special form of their approximate formula for the mean length of the busy period. Finally in Section 5, we present our computational example and make some further remarks on the approximation.

2. PH-Distributions and PH-Renewal Processes:

A phase type probability distribution (PH-distribution), introduced by M.F.Neuts[2], is obtained as the distribution of the time till

absorption in a finite Markov chain with an absorbing state into which absorption is certain. We shall briefly review such models.

Consider a continuous-time Markov chain with state space $\{1, \dots, n, n+1\}$ for which the states $1, \dots, n$ are transient and $(n+1)$ is absorbing. It is assumed that starting in any transient state the chain gets absorbed in $(n+1)$ with probability one. The infinitesimal generator of such a Markov chain then has the form

$$Q = \begin{bmatrix} T & \underline{T}^0 \\ \underline{0} & 0 \end{bmatrix}$$

where T is an $n \times n$ non-singular matrix with $T_{ii} < 0$, and $T_{ij} \geq 0$, $i \neq j$. Also $\underline{T}^0 \geq \underline{0}$ is an n -vector satisfying $T\underline{e} + \underline{T}^0 = \underline{0}$, where $\underline{e}' = (1, \dots, 1)$. We assume that we are also given an initial probability vector $(\underline{\alpha}, 0)$ for the Markov chain.

It is now easily seen that for the above Markov chain the time till absorption in the state $(n+1)$ has c.d.f.

$$F(x) = 1 - \underline{\alpha} \exp(Tx) \underline{e}, \quad x \geq 0.$$

Any probability distribution $F(\cdot)$ so obtained will be called a Phase Type Distribution, and the pair $(\underline{\alpha}, T)$ is called a representation. The mean of $F(\cdot)$ is given by $\lambda^{-1} = -\underline{\alpha} T^{-1} \underline{e}$.

In [3] Neuts considered the renewal process obtained by restarting the Markov chain Q instantaneously after each absorption (renewal) by performing a multinomial trial with probabilities $\underline{\alpha}$ and outcomes $1, \dots, n$

and called it a Phase Type Renewal Process (PH-Renewal Process). This results in another continuous-time Markov chain with state space $\{ 1, \dots, n \}$ and infinitesimal generator

$$Q^* = T + T^0 A^0,$$

where $A^0 = \text{diag}(a_1, \dots, a_n)$ and $T^0 = (T_1^0, \dots, T_n^0)$. This Markov chain describes the "phase" of the PH-renewal process. In [2] it has been shown that the representation (\underline{a}, T) of $F(\cdot)$ may be so chosen that the matrix Q^* is irreducible. We shall assume that this is indeed the case.

For the PH-renewal process described above, let $N(t)$ denote the number of renewals in $(0, t]$ and $J(t)$ denote the phase at time $t+$ and define

$$P_{ij}(v, t) = P\{N(t)=v, J(t)=j \mid J(0)=i\}, \quad v \geq 0, \quad 1 \leq i, j \leq n.$$

In [3] it is shown that the generating function of the matrices $P(v, t) = (P_{ij}(v, t))$ is given by

$$\tilde{P}(z, t) = \sum_{v=0}^{\infty} z^v P(v, t) = \exp\{(T+zT^0A^0)t\}, \quad |z| \leq 1, \quad t \geq 0.$$

Also by differentiating,

$$M(t) = \frac{\partial}{\partial z} \tilde{P}(z, t) \Big|_{z=1^-} = \sum_{k=1}^{\infty} \frac{t^k}{k!} \sum_{v=0}^{k-1} Q^{*v} T^0 A^0 Q^{*k-1-v} \quad (2.1)$$

where the (i, j) -th entry of $M(t)$ is

$$M_{ij}(t) = E\{N(t) I_{[J(t)=j]} \mid J(0)=i\}.$$

Note that

$$m(t) = \underline{a}M(t)\underline{e} = \underline{a} \sum_{k=1}^{\infty} \frac{t^k}{k!} Q^{*k-1} \underline{1}^0 \quad (2.2)$$

is the expected number of renewals in (0, t).

3. The PH/M/1 Queue:

We now consider a GI/M/1 queue in which arrivals occur according to a Phase Type renewal process. Such a model will be denoted by PH/M/1. We let μ denote the service rate.

Letting $X(t)$ and $J(t)$ denote respectively the queue length (i.e., the number of customers in the system) and the phase of the arrival process at time $t+$, it is easily seen that $\{(X(t), J(t)): t \geq 0\}$ is a Markov chain with state space $\{0, 1, \dots\} \times \{1, \dots, n\}$ and infinitesimal generator

$$\tilde{Q} = \begin{bmatrix} T & T^0 A^0 & 0 & 0 & 0 & \dots & \dots \\ \mu I & T - \mu I & T^0 A^0 & 0 & 0 & \dots & \dots \\ 0 & \mu I & T - \mu I & T^0 A^0 & 0 & \dots & \dots \\ 0 & 0 & \mu I & T - \mu I & T^0 A^0 & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \end{bmatrix}$$

The infinitesimal generator \tilde{Q} is of a form studied by V. Wallace [7] under the name quasi-birth-and-death processes. A queueing model with infinitesimal generator of the same form has also been studied by Neuts [4].

For the Markov chain \tilde{Q} above we let $\tilde{G}_{jj'}(x)$, $x \geq 0$, $1 \leq j, j' \leq n$, denote the probability that, starting in the state $(i+1, j)$, $i \geq 0$, the first visit to the set of states $\underline{i} = \{(1, k) : 1 \leq k \leq n\}$ occurs no later than x and that at the epoch of such a first passage the phase of the system is j' .
Let

$$G_{jj'}(s) = \int_0^\infty e^{-sx} d\tilde{G}_{jj'}(x), \quad \operatorname{Re} s \geq 0, \quad 1 \leq j, j' \leq n$$

and let $G(s)$ denote the $n \times n$ matrix of entries $G_{jj'}(s)$.

Theorem 3.1

i) The matrix $G(s)$ satisfies the equation

$$G(s) = [sI - (T - \mu I)]^{-1} \mu I + [sI - (T - \mu I)]^{-1} T^0 A^0 G^2(s) \quad (3.2)$$

for all $s \geq 0$.

ii) If the queue is stable, i.e., if $\rho = \lambda/\mu < 1$, then $G = G(0+)$ is stochastic.

iii) G is the minimal solution in the set of sub-stochastic matrices to the equation

$$G = (\mu I - T)^{-1} [\mu I + T^0 A^0 G^2] \quad (3.3)$$

iv) The matrix G can be computed using the recurrence formulas

$$\left. \begin{aligned} G_0 &= 0 \\ G_{k+1} &= (\mu I - T)^{-1} [\mu I + T^0 A^0 G_k^2], \quad k \geq 0. \end{aligned} \right\} \quad (3.4)$$

Proof:

Equation (3.2) follows from a standard first passage argument by considering the first time the queue length goes either down or up. That the queue

is stable iff $\rho = \lambda/\mu < 1$ is well-known. The proofs of the other claims in the theorem are similar to those in Neuts[4] and hence omitted.

Remark: In [5] it is shown that the matrix G is strictly positive and thus has a unique invariant probability vector \underline{g} satisfying $\underline{g}G = \underline{g}$, $\underline{g}\underline{e} = 1$.

Theorem 3.5:

Let
$$\underline{\mu}^* = - \left. \frac{d}{ds} G(s)\underline{e} \right|_{s=0+}.$$

Then

$$\underline{\mu}^* = -(I - G + \tilde{G})[(T + T^0 A^0) + (T^0 A^0 - \mu I)\tilde{G}]^{-1} \underline{e}, \quad (3.6)$$

where \tilde{G} is the $n \times n$ matrix each of whose rows is \underline{g} .

Proof:

Differentiating (3.2) with respect to s and letting $s \rightarrow 0$, we get

$$[(T - \mu I) + T^0 A^0 + T^0 A^0 G] \underline{\mu}^* = -\underline{e}. \quad (3.7)$$

Since $I - G + \tilde{G}$ is non-singular and since $\mu I + (T - \mu I)G + T^0 A^0 G^2 = 0$, we have that

$$[(T - \mu I) + T^0 A^0 + T^0 A^0 G](I - G + \tilde{G}) = (T + T^0 A^0) + (T - \mu I + 2T^0 A^0)\tilde{G}$$

Using this in (3.7) we get (3.6) by noting that

$$(T + T^0 A^0)\tilde{G} = (T + T^0 A^0)\underline{e}\underline{g} = \underline{0}\underline{g} = 0$$

Remark: Note that μ_i^* is the expected duration of a busy period which starts with 1 customer and in phase i .

Theorem 3.8:

$$\underline{g}\underline{\mu}^* = 1/[\mu(1-\rho)] \quad (3.9)$$

Proof:

This formula is established for the more general $N/G/1$ queue in [5].

Remark: Equation (3.9) provides a powerful computational check on the accuracy of numerical computations.

Algorithm to compute the mean length of the busy period:

Step 1: Compute G using (3.4)

Step 2: Compute the invariant probability vector \underline{g} of G

Step 3: Compute $\underline{\mu}^*$ using (3.6)

Step 4: Verify (3.9)

Step 5: Compute the mean length of the busy period $E(BP) = \underline{g}\underline{\mu}^*$

4. The Cross-Harris Approximation:

In their book [1], Gross and Harris obtain a "nice, neat" expression for the expected length of the busy period in a GI/G/1 queue. By treating the arrival process as though it were memoryless and mimicking the derivation of the mean length of the busy period in an M/G/1 queue, they "show" that the expected length of the busy period in a GI/G/1 queue is approximately equal to

$$E(BP) \approx E(S) / [1 - \int_0^{\infty} m(t) dB(t)] \quad (4.1)$$

where S denotes the service time and $B(\cdot)$ its c.d.f., and where $m(t)$ is the expected number of arrivals in $(0, t]$.

Note that for the PH/M/1 queue

$$\begin{aligned} \int_0^{\infty} m(t) dB(t) &= \underline{\alpha} \int_0^{\infty} \sum_{k=1}^{\infty} \frac{t^k}{k!} Q^{*k-1} \underline{1}^0 \mu e^{-\mu t} dt \\ & \text{by (2.2)} \\ &= \mu^{-1} \underline{\alpha} (\underline{I} - \mu^{-1} Q^*)^{-1} \underline{1}^0, \end{aligned}$$

whence (4.1) may be written as

$$E(BP) \approx 1 / [\mu - \underline{\alpha} (\underline{I} - \mu^{-1} Q^*)^{-1} \underline{1}^0] \quad (4.2)$$

The existence of the inverse in (4.2) follows from the fact that Q^* is a stable matrix, i.e., $\text{Re } \delta_i < 0$ for every eigenvalue δ_i of Q^* .

5. The Computational Example:

For our example we considered a queue with a generalized Erlang distribution for inter-arrival times which, as a PH-distribution, has the representation

$$\underline{\alpha} = (1, 0, 0, 0, 0)$$

$$T = \begin{bmatrix} -1 & 1 & 0 & 0 & 0 \\ 0 & -1 & 1 & 0 & 0 \\ 0 & 0 & -10 & 10 & 0 \\ 0 & 0 & 0 & -10 & 10 \\ 0 & 0 & 0 & 0 & -10 \end{bmatrix} \quad \text{and} \quad \underline{T}^0 = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 10 \end{bmatrix}$$

The service time distribution was assumed to be exponential with parameter $\mu = 0.483$. It may be verified that for this queue $\rho = 0.900$.

In implementing the algorithm at the end of Section 3 to compute the (exact) value of the expected length of the busy period, we terminated the iterations in (3.4) using the criterion

$$\max_{1 \leq i, j \leq n} |G_{k+1}(i, j) - G_k(i, j)| < 10^{-8}.$$

Then a linear extrapolation, based on the last two iterates, was used to make the computed matrix G stochastic. The invariant probability vector \underline{g} of G was computed using Wachter's method [6]. All these were programmed in APL, and a copy of the program may be obtained by writing to either of the authors. We summarize the results of our computations below:

$$\underline{\mu}^* = (14.55679, 20.58772, 29.53159, 30.85796, 32.24840)$$

$$\underline{g} = (0.27996, 0.54192, 0.05679, 0.05938, 0.06194)$$

$$\underline{g}\underline{\mu}^* = 20.73940$$

$$\underline{\mu}^{-1}(\underline{1}-\rho)^{-1} = 20.73940$$

$$\underline{\alpha} \underline{\mu}^* = 14.55679$$

The approximation to the expected length of the busy period was computed using (4.2). The computed value of the "approximation" for E(BP) turned out to be 5.95040 which, when compared to the true value 14.55679, is grossly inaccurate.

A close examination of the model here reveals why the "approximation" fares so poorly. The assumption of a memoryless property for the arrival process in the example amounts to resetting the phase of the PH-renewal process at 1 immediately after every departure. This, for our model, has the effect of "slowing down" the arrivals immediately after each departure and leads to under-estimating the duration of (and the number served during) a busy period. What one needs to recognize is that the resetting of the phase to 1 is repeated several times during a busy period and not just at the end of the first service completion alone. As ρ gets larger and closer to 1, one expects a busy period to consist of a larger number of service completions making the approximation progressively worse as ρ increases. Further by considering a general matrix T of the form

$$T = \begin{bmatrix} -a & a & & & \\ & -a & a & & \\ & & -b & b & \\ & & & -b & b \\ & & & & -b \end{bmatrix}$$

for defining the generalized Erlang distribution and by making 'a' smaller and 'b' larger, one can construct examples where the approximation is worse for queues with even the same value for ρ .

Acknowledgment: We express our gratitude to Professor M. F. Neuts for suggesting this problem and for his guidance.

REFERENCES

- [1] Gross, D. and Harris, C. M. (1974)
Fundamentals of Queueing Theory pp. 306-308
John Wiley & Sons, New York
- [2] Neuts, M. F. (1975)
Probability Distributions of Phase Type
in--Liber Amicorum Professor Emeritus H. Florin--
Department of Mathematics, University of Louvain, Belgium, 173-206.
- [3] Neuts, M. F. (1976)
Renewal Processes of Phase Type
Tech. Report #76/8, Dept. of Statistics & Computer Science, University
of Delaware, Newark, DE 19711
forthcoming in Nav. Res. Log. Quart.
- [4] Neuts, M. F. (1977)
The M/M/1 Queue with Randomly Varying Arrival and Service Rates
Tech. Report #77/11, Department of Statistics & Computer Science,
University of Delaware, Newark, DE 19711
- [5] Ramaswami, V. (1978)
The N/G/1 Queue and its Detailed Analysis
Tech. Report #78/1, Dept. of Statistics & Computer Science, University
of Delaware, Newark, DE 19711
- [6] Wachter, P. (1973)
Solving Certain Systems of Homogeneous Equations with Special Reference
to Markov Chains
M.Sc. Thesis, Department of Math, McGill University, Montreal, Canada,
77 pp with Appendices.
- [7] Wallace, V. (1969)
The Solution of Quasi Birth and Death Processes arising from Multiple
Access Computer Systems
Ph.D. Thesis, Systems Engineering Lab., Univ. of Michigan, Ann Arbor, MI,
Tech. Report No. 07742-6-T.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
1. REPORT NUMBER AFOSR-TR- 78-0984	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER	
4. TITLE (and Subtitle) ON THE MERITS OF AN "APPROXIMATION" TO THE BUSY PERIODS OF THE GI/G/I QUEUE		5. TYPE OF REPORT & PERIOD COVERED Interim	
		6. PERFORMING ORG REPORT NUMBER	
7. AUTHOR(s) V. Ramaswami and David M. Lucantoni		8. CONTRACT OR GRANT NUMBER(s) AFOSR-77-3236	
9. PERFORMING ORGANIZATION NAME AND ADDRESS University of Delaware Dept. of Statistics & Computer Science Newark, DE 19711		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61102F 2304/A5	
11. CONTROLLING OFFICE NAME AND ADDRESS Air Force Office of Scientific Research/NM Bolling AFB, Washington, DC 20332		12. REPORT DATE March 1978	
		13. NUMBER OF PAGES 13	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED	
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.			
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)			
18. SUPPLEMENTARY NOTES			
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) GI/G/I queue, busy period, Phase Type distribution Phase Type Renewal Process, computational probability.			
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) An approximation for computing the busy period characteristics in the GI/G/I queue, suggested by D. Gross and C.M. Harris in their book "Fundamentals of Queueing Theory", is shown to be lacking in accuracy, and is therefore quite undesirable for practical use.			

DD FORM 1473

1 JAN 73

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)