

Ecological site R010XY008ID South Slope Granitic 12-16 PZ PUTR2/PSSPS

Last updated: 9/23/2020
 Accessed: 04/19/2024

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Dave Franzen and Jacy Gibbs Intermountain Range Consultants 17700 Fargo Rd. Wilder, ID 83676
Contact for lead author	Brendan Brazee, State Rangeland Management Specialist USDA-NRCS 9173 W. Barnes Drive, Suite C, Boise, ID 83709
Date	03/25/2008
Approved by	Kendra Moseley
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:** can occur on this site. If rills are present they are likely to occur on the steeper slopes and immediately following wildfire. Coarse surface gravels limit rill development.

2. **Presence of water flow patterns:** can occur on this site. When they occur they are short and disrupted by cool season grasses and tall shrubs and are not extensive. Water infiltration is generally rapid for the site.

3. **Number and height of erosional pedestals or terracettes:** pedestals are rare on this site. Terracettes are common. Significant accumulation of coarse surface fragments develops on the uphill side of larger perennial grasses and shrubs. This accumulation is from concentrated flow and hoof/ foot traffic. Terracettes are a natural occurrence on the site.

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** is approximately 50-60%. This site is naturally unstable due to coarse surface fragments.

5. **Number of gullies and erosion associated with gullies:** do not occur on this site.

6. **Extent of wind scoured, blowouts and/or depositional areas:** usually not present due to coarse textured, gravelly soil surface.
-
7. **Amount of litter movement (describe size and distance expected to travel):** fine litter in the interspaces may move up to 3 feet following a significant run-off event. It generally moves onto terracettes. Coarse litter generally does not move except on the steeper slopes. Litter is also moved mechanically by hoof/ foot traffic.
-
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Values should range from 4 to 6 but needs to be tested.
-
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** structure ranges from weak fine granular to moderate medium granular. The A or A1 horizon is typically 2 to 10 inches thick. Soil organic matter (SOM) ranges from 1 to 5 percent.
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** bunchgrasses, especially deep-rooted perennials, slow run-off and increase infiltration. Tall shrubs accumulate snow in the interspaces.
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** not present.
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant: cool season deep rooted perennial bunchgrasses.
- Sub-dominant: tall shrubs > perennial forbs.
- Other:
- Additional:
-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** foothills big sagebrush and bitterbrush will become decadent in the absence of fire and ungulate grazing. Grass and forb mortality will occur as tall shrubs increase. Some grasses and forbs are susceptible to hoof/ foot traffic.
-
14. **Average percent litter cover (%) and depth (in):** additional litter cover data is needed but is expected to be 5-10 percent to a depth of <0.1 inches. Under mature shrubs litter is >0.5 inches deep and is 90-100 percent ground cover.
-
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** is 800 pounds per acre (888 kilograms per hectare) in a year with normal temperatures and precipitation.

Perennial grasses produce 55-65 percent of the total production, forbs 10-20 percent and shrubs 20-30 percent.

16. **Potential invasive (including noxious) species (native and non-native).** List species which **BOTH** characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is **NOT** expected in the reference state for the ecological site: include cheatgrass, leafy spurge, dalmatian toadflax, bulbous bluegrass, rush skeletonweed, musk and scotch thistle and diffuse, Russian and spotted knapweed.
-

17. **Perennial plant reproductive capability:** all functional groups have the potential to reproduce in most years.
-