



Some Nutritional and Medicinal Importance of Nigerian Walnut “*Tetracarpidium conophoram*”

Uchejeso Obeta^{1*}, Alexander Lawrence², Muhammad Akram³, Fahad Said Khan^{3,4}

¹Department of Medical Laboratory Management, Federal School of Medical Laboratory Science, Jos-Nigeria.

²Department of Haematology, Federal Medical Centre, Lokoja-Nigeria.

³Department of Eastern Medicine, Government College University Faisalabad-Pakistan.

⁴Department of Research and Development, Qarshi Foundation Lahore, Pakistan.

ABSTRACT

Nigerian walnuts (*Tetracarpidium conophoram*) are used as a diet and medicine is on the increase to the extent that it is almost scarce in various regions where they are marketed. This study took a brief review on the nature, classification, properties, phytochemical components, nutritional values, and medicinal properties. There is rapid growth towards research and developments involving Nigerian walnut for drug development. From this study, it is discovered that Nigerian walnut has shown anti-oxidant, anti-inflammatory, anti-chelating nature, and high blood pressure control activities. Also, it has shown anti-lipidemic, anti-diabetes, anti-malarial, antimicrobial /antibacterial, and anticancer nature. The walnut has given reproductive assistance and aided sperm production and sperm quality, wound healing, anti-ulcer, and stomach disorders treatment features. The Nigerian walnut has energizing and weight management and diseases prevention capacity. It aids in bone health, skin or other tissues repairs, and could go a long way to assist in other health challenges. There has also encouraged usage in the aspect of complementary and alternative medicine. There seems to be a close relationship between walnut and bitter kola in nature, diet and medicinal properties. Researchers and drugs producing companies should improve in their studies for more drug discoveries that could involve Nigerian walnut as it could assist in a Nobel discovery for treatment of difficult to treat diseases in the world.

Key Words: Walnut, Ukpa, Nutrition, Medicine, Diseases, Bitter kola

eIJPPR 2021; 11(5):34-40

HOW TO CITE THIS ARTICLE: Obeta U, Lawrence A, Akram M, Said Khan F. Some Nutritional and Medicinal Importance of Nigerian Walnut “*Tetracarpidium conophoram*”. Int J Pharm Phytopharmacol Res. 2021;11(5):34-40. <https://doi.org/10.51847/ZtQ0ccTGPe>

INTRODUCTION

Nutritional and Medicinal plants and vegetables have been advocated for the management of difficult to treat diseases [1, 2]. Herbal and complementary medicine has been advocated [3] and approved by WHO [4] in the cases of need. Nigerian walnut is discussed here for their nutritional and medicinal importance in Nigeria and across the world. This article shall stimulate more studies on walnut especially in the areas where research on them have not been studied.

The Nigerian Walnut with the Botanical Name (*Tetracarpidium conophoram*) contains an important oilseed crop that is grown in the tropical Sederhana regions of the world. You are a climbing shrub Dalam Spora plukenetia. The Bukan or Adala, as it was named so because the Walnut even has superficial similarity with the Walnut. Adalah A nature in Central West Tropical Africa

from Togo to Congo to Sierra Leone. Adalah is abundantly available in Nigeria, Cameroon, Republic of the Congo. It prefers the Hedges of the Dalam rainforest semi-shady places; secondary forest a low shrub; Iklan farms a kulat altitude 250-1400 (820-4590 strengthen appropriate). Although it is well registered in Sierra Leone, clearly Bukan originally did not come from Sierra Leone, as Bukan is registered in Liberia and Ghana, but the reproach Presence of Dalam Sierra Leone as a cause of the Return of Slaves as the reason why Bukan is known to come to Krio from Yoruba (Nigerian). Plukenetia conophora is the only plukenetia species native to Central West Africa. Others had to strengthen Platine originating from another part of Africa, of the Indian subcontinent, of South East Asia of America [5].

The scientific name of walnut is *Tetracarpidium conophoum* and classifies according to Arranz *et al.*, [6] as follows:

This is an **open access** journal, and articles are distributed under the terms of the Creative Commons Attribution-Non Commercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

Corresponding author: Uchejeso Obeta

Address: Department of Medical Laboratory Management, Federal School of Medical Laboratory Science, Jos-Nigeria.

E-mail: ✉ uchejesoobeta@gmail.com

Received: 28 June 2021; **Revised:** 21 September 2021; **Accepted:** 26 September 2021



Kingdom	Plantae
Order	Malpighiales
Family	Euphorbiaceae
Sub Family	Acalyphoideae
Genus	Plukenetia
Species	Tetracarpidium

MATERIALS AND METHODS

This study sort for information required to investigate the topic based on the nutritional and medicinal importance of Nigerian walnut. The required materials were obtained from scholarly sources such as peer-reviewed journals and books. The used articles mostly published by Nigerians were containing extensively the required information readers can easily understand.

Not only that, the used articles and books were also based on research experience conducted in and outside Nigeria using the Nigerian walnut. The content of used journals and books were generated through searching the keywords of the study in Google which includes Walnut, Nigerian walnut, Nutritional values of walnut, Medicinal uses of walnut, Phytochemicals of Nigerian walnut, and Studies on Nigerian walnut.

RESULTS AND DISCUSSION

The search and literature reviews revealed much useful information about the Nutritional and medicinal uses of Nigerian walnut ranging from the local names of walnut in Nigeria, phytochemical components, physical and chemical properties of walnut, and their pictorial nature as shown in **Figures 1-3**.

Names of walnut in Nigeria

The scientific name is *Tetracarpidium conophoum*. The common name is walnut otherwise called Nigerian walnut or African walnut. This is because it came from *Juglan nigra* (the family Juglandaceae because of African origin. Though the walnut is called Nigerian walnut, different nationalities in Nigeria have specific names given to the plant/seeds based on their locality or nationality. Such Nigerian names for walnut are as follows: Ukpa in Igbo nation, Ukwa in Igala and Idoma, Ekporo as called by Efik and Ibibios, Okhue or Okwe as it is called by Edo people, Gawadi bairi in Hausa, and Awusa or Asala in Yoruba nationality [5, 7, 8].

Description of the walnut

Tetracarpidium conophorum has a long history of cultivation of food as a plant species in West Africa. Guinea and West and Central Africa distribute and consume it. Capsule bear larvae are green when growing and greenish-yellow when growing. The leaves are about

53 inches wide and about 3 inches long by rotating. Only the wings are shown solely to give a sense of proportion. The taste of hard fruit and eucalyptus. The seeds take 4 to 6 months to mature and are found in the local market between June and September. It covers every pillar and especially the trees that surround it. It is a botanical made from cocoa beans and cola and used. Because it is based on the high consumption of endosperm oil used by several people in Nigeria, Sierra Leone, and the province of Borge in Congo. It grows well along the coast of Africa and is believed to come from the southwest. It is common for farmers to grow food in tropical Africa on farms and gardens, only for families and for the local market. *Tetracarpidium conophorum* is a climber that grows to about 6-18m in height when it is born on the stem. When a successful tree gets taller, the tallest tree grows from the ground onto other trees. They are well received in warm conditions. It has a stem that can reach 70 meters or more. It climbs taller trees elsewhere to achieve greater sunlight and maturity. In some cases, the wood can be broken and one stays in place until it is sold. You must be around 16cm round and dark gray, but your current green hair is still small. The roots are very beautiful. Leaves are (10 cm x 5 cm) short, oval, with serrated sides. They turned and pointed carefully to the head. The leaves have three nerve endings with leaf petioles up to 5 cm long [9].

Ndukwu and Ejirika [10] describe the physical characteristics of Nigerian beans including: water content - 28.2% by weight, mass (m) at 5.5.6m 6.5g, shell thickness - $0.135 \pm 0, 04$ cm, core mass - 4.22 ± 0.26 g, core water - 28.9% Wb, porosity - 45%, grain density - 0.815 g / cm^3 , grain density - 0.45 g / cm^3 , the percentage of corn - 76% in the grain area - 15.41 cm^2 . Characteristics can influence behavior. Physical increase in water content, arable area, and variation in grain size. The fruits on the tree are shown in **Figure 1**; Ripe fruit content is shown in **Figure 2** while dried fruit is shown in **Figure 3**.



Figure 1. Showing the African Walnut seed in the process of maturation.



Figure 2. Ripped African Walnut seed showing the internal structure

Fantastic wild berries are grown in traditional farming systems in tropical lowlands. It can tolerate any type of soil as long as it is well drained and can hold water.

It is a deciduous tree, new, with few branches, sometimes appearing as a candelabra. The seed is a 6-10 cm long capsule with an important capsule containing a subglobular seed of 1–1.5 cm long and a thin brown shell-like a small nut, hence the name in English [11].

Peanuts are the property of any plant of the genus *Juglans* (family Juglandaceae), in particular, the Persian or English walnut *Juglans regia*. Technically, nuts are hard-boiled or rooted fruit, so it is not a good vegetable. It is used in cooking as a side dish for green walnuts or as a well-ripened vegetable. *Juglans* oriental bean curd, *nigra* is the smallest meat on the market, as well as *Juglans cinerea* bean buttery. Walnuts are a protein that contains essential proteins and fatty acids.

African walnut Apart from nuts, research has disclosed that other parts of this plant are very crucial for medicinal, remedial, and medicinal purposes [12].



Figure 3. Showing matured Walnut seed in a shell.

Characteristics of Nigerian walnut

Beans are single-walled round fruits, usually eaten when ripe. After perfect cooking, the elimination of the husk affirm the wrinkled walnut shells that are usually found on both sides (3D marks can also begin). During adolescence, the chest breaks and stiffens and he knows it's time. The

cortex covers the core of tissue, which is usually divided into two parts by a divider. Cells are often like shells. Dip the walnuts in brown beans, which contain antioxidants. Antioxidants protect fruit oils from oxygen, thus preventing acidity [7].

The beans do not arrive during the growing season, no more than mid-spring. They store chemicals in the soil to inhibit weeds from contending to grow. For this reason, you shouldn't plant flowers or gardens near them. Oatmeal is made up of 14% carbohydrates, 15% protein, 4% water, and 65% fat comprising 7% fiber [8, 13].

Storage of walnut

Nuts, like other fruits, must be properly prepared and stored. Poor conservation makes walnuts vulnerable to attack by insects and fungal diseases; the latter, which produce aflatoxins, are potent carcinogens. The beans should be removed completely [5].

The ideal low-temperature range is -3 to 00°C (27 to 320°F) and low humidity for home and industrial storage. However, this cooling technology is not available in developing countries where walnuts are produced in large quantities. There, the nuts are best stored at temperatures up to 250°C (770°F). Temperatures above 300 degrees Celsius (860 degrees Fahrenheit) and humidity above 700% can cause rapid leaks and severe damage. With humidity above 75% , fungal strains that secrete harmful aflatoxins can develop [7, 14].

Phytochemistry of Nigerian walnut

Tetracarpidium conophorum has unprecedented health benefits, although it is still on the list of lesser-known foods. Phytochemical analysis of *T. conophorum* root leaves showed that it is involved in active compounds such as oxalates, phytates, tannins, alkaloids, flavonoids, and terpenoids. The presence of these phytochemicals indicates different aspects of the use of *T. conophorum* in herbal medicine [12].

Nowajkbe *et al.* [15] Saponins 8.37 and 5.03 mg / kg were found separately in boiled nuts / dried nut juice and peanut powder. Eggs and lentils contain protein (14.92%), oil (45.84%), crude fiber (1.14%), micronutrients (3.52%) and carbohydrates (15.38%) and others such as tannins (0.89 mg / 100). g, oxalate (1.28 mg / 100 g), phytic acid (3.105 mg / 100 g), trypsin inhibitor (1.84 mg / 100 g), saponins (985.0 mg / 100 g) and alkaloids (40.91 mg / 100 g).

Arenola and Adesina [16] reported that the proteins, fats, ashes, and intestinal contents of the Nigerian beans were reduced and cooked.

The study by Akpogheli *et al.* [17] identified 3.18% of Ash, 39.27% of water, 8.40% of fiber, 5.19% of fat, 20.74% of protein, and 23.22% of sugar, while minerals revealed 4029.14 mg of K per kg, 3480.00 mg of Na per kg, $3.014.28$ mg of Ca per kg, 726.11 mg of Mg per kg, 68.00

mg of Fe per kg, 24.01 mg of Zn per kg, 19.00 mg of Mn per kg, and 14.00 mg of Cu per kg. Chijoke *et al.* [18] also reported Nigerian beans containing alkaloids for 2.29 mg per 100g, glycosides -2.19 mg per 100 g, saponins - 8.07 mg per 100 g, flavonoids -0.02 mg per 100 g, tannins -0.89) per 100g, reducing sugars -4.10 mg per 100g and soluble carbohydrates for 1.06 mg per 100g. Likewise, the economy is exposed to a high content of water (31.40%), nutrients (6.01%), fiber (8.66%), proteins (28.85%), starch (21.30%), and high energy value (234.57 kcal) [19].

Oddee and others [20] recorded Phenols (7.44 mg / ml and 7.04 mg / ml), flavonoids (3.5 mg / ml and 1.66 mg / ml) and ascorbic acid (54.56 mg / kg and 44.00 mg / kg) for raw and ripe beans respectively.

Onawumi *et al.*, [21] studies have shown water (29%), fats (5.63%), fiber (14.92%), proteins (16.62%), ash (12.89%) and starch (20.94%) contain alkaloid (2670). mg / kg) and microtannins (0.56 mg / kg).

Nutritional and medical needs of Nigerian nuts

There are two types of nuts, in shells or shells. The body can be whole, divided, or partially due to the structure. Cooked walnuts are most commonly used, but they can be used in other foods. Pickled walnuts, which are all fruits, can be sweet or salty depending on the preservative solution. Peanut butter can be made at home or bought raw and cooked. All beans can be eaten on their own (raw, roasted, or pickled), as part of a mix like muesli or as a side dish. For example, peanut soup and peanut bread are enriched with peanuts [22].

Barber and Obinna-Echem [23] recommended Nigerian walnut flour as a better substitute to wheat flour based on the nutritional composition.

Popular with bean curd, ice cream toppings, and walnuts are utilized as a garnish in few dishes, Nicino is an alcoholic wine made from green, fresh fruit made with wine. alcohol is added to the syrup. Nut oil is available in the market and is utilized as a food, especially in salad dressings. It has a low smoke area, which averts its use for frying [5, 7].

At 100 grams of food, walnuts provide 2,> 40 kilojoules (654 calories) and a rich diet of minerals, mainly 163% AV manganese and vitamin B. Good functions of the human body include Thiamine, Riboflavin, Folate, Cyanocobalamin (Vitamin B1, B2, B3, B5, B6, B12, C, and E), Niacin, Manganese, Arginine, Pyridoxine, Selenium, Acrobatic Acid, Melatonin, Pantothenic. Acid, tocopherol, ellagic acid, polyphenols, omega-3 fatty acids, zinc oleic acid, calcium, phosphorous, potassium, sodium, H₂O, carbohydrates, and proteins [5, 8].

Various studies have shown a few healing properties of nignut walnut. Wound healing is a complex prepare in which the skin or other tissues repair themselves after an injury. It has been found that the use of *T. conophorum*

showed a rapid therapeutic effect of wound healing after 5% or 10% *T. conophorum* treatment for 8 days [24].

As a follow-up to the wound healing property of Nigerian walnut, an anti-ulcer feature has been shown. Ezealisiji *et al.* [25] and Anosike *et al.* [26] showed anti-ulcer, cytoprotective, and wound healing properties in their studies.

The utilize of Nigerian walnut in the therapy of gastrointestinal ailments and the control of high blood pressure was described by Ayoola *et al.* [27]. Nwachoko and Jack [28] also found that a hot aqueous extract of Nigerian walnuts protected rats from castor oil-induced diarrhea.

Antimicrobial /antibacterial studies have shown that Nigerian walnut has the potential. Akinwande [29] and Ajaiyeoba and Fadare [30] reported antimicrobial activities of Nigerian walnut.

Nigerian walnut has shown some anti-chelating nature in a study. An in-vitro chelating nature of Nigerian walnut showed a dose-dependent decrease in chelating nature [31].

Based on anti-inflammatory features of Nigerian walnut, Olaniyi *et al.* [32] and Oladokun *et al.* [33] showed that Nigerian walnut had a significant level of inflammation inhibition.

Amaze *et al.* [12] investigated the antioxidant properties of Nigerian walnut leaf extract and concluded that dried leaves have higher antioxidant activity than fresh leaves. The antiperoxidative activity of *T. conophorium* leaves was assessed by measuring their ability to lower malondialdehyde (MDA) levels in the reproductive organ and accessory glands of rats. The decrease of lipid peroxidation in conceptive and frill organs that extract Nigerian walnut has been reported by Akomolafe *et al.* [34], Kanu, and Okorie [8].

Walnut has anti accident properties that help the body to fight stress or oxidative stress which causes loss of red blood cells in the body, remove tree radicals help to support the immune system, help to protect against environmental and chemical toxins, help in sequencing agents of heavy metals and significantly help or protect quick aging [35]. Some studies around the reproductive assistance of Nigerian walnut, Akomolafe *et al.* [36] shows that Nigerian Walnut fights off pregnancy sicknesses and boost the development of the fetal brain.

Obianime *et al.* [37] and Obianime *et al.* [38] demonstrated that Nigerian walnut aided sperm production and quality in male guinea pigs. This was proven by Akomolafe *et al.* [39] and Ikpeme *et al.* [40] who showed that Nigerian Walnut improves sperm quality. Acumulavi *et al.* [34] Determination of the effect of Nigerian death adjustment. Akomolafe and Oboh [34] also showed a significant increase ($p < 0.05$) in lactate dehydrogenase (LDH), glucose-6-phosphate dehydrogenase (G-6PDH), glycogen

content, and 3- and 17-activity -hydroxysteroids (HSD) in testes and epididymis. Zn and Se content is present in the biochemical parameters of the testis, but also with a significant decrease in cholesterol. Studies showed a significant increase in serum testosterone, LH, and FSH levels, sperm count, motility, and viability, and a decrease in sperm abnormalities at $p < 0.05$, indicating the importance of spermatogenesis, parameters, and volume.

Acumulavi *et al.* [41] showed a significant increase ($p < 0.05$) in LDH activity, G-6PDH activity, glycogen content, and HSD activity 3 and 17 ° in testis, epididymis, zinc, and silicon; There is also a decrease in the number of spermatozoa in the testes, their viability and a significant increase in the number of sperm abnormalities, a set of criteria for sperm analysis and, consequently, a decrease in fertility hormone levels when drinking alcohol. Alcohol is probably not good for male fertility problems.

Akpogheli *et al.* [17] reported the diseases prevention capacity of walnut because of the presence of Omega-3 fatty acids (eicosapentaenoic acid called EPA and docosahexaenoic corrosive called DHA). Based on anticancer activities, some studies demonstrate that prostate and breast cancer can be reduced by consuming Nigerian walnut [42].

Nigerian walnut consumption is also important in anti-diabetes. Onwuli *et al.* [43]; Ogunyinka *et al.* [44] Ogbonna *et al.* [45] reported that Nigerian walnuts can reduce high glucose levels in the blood. It is also anti-lipidemic as reported by Ezealisiji *et al.* [46] and Analike *et al.* [47].

Kanu, *et al.* [48] demonstrate that Nigerian walnut aids in bone health. It seems good for bone growth and strength. Anti-malarial properties of Africa Walnut have been studied by Dada and Ogundolie [49]. In the cause of the study, Ogundolie *et al.* [50] used ethanol seed extract and discovered that it increased packed cell volume (PCV), red blood cells (RBC), hemoglobin (HGB), and platelet (PLT). This may provide another mechanism where malaria is well treated and hematological parameters may be increasing which is different from the medical approach of withdrawing haematinics during malaria treatment.

Nigerian walnut has shown energizing and weight management properties and these properties no doubt are traced to their physiochemical properties [15-20, 48]

Other important aspects of walnut use are still being researched. Walnut, according to Anosike *et al.* [26], can be used to treat the gastrointestinal tract and can help reduce premature deliveries in pregnant women because it has culinary uses, such as soup preparation; it can boost cognitive function due to the presence of vitamin B6 and omega-3 unsaturated fats, and it can also help as an anti-aging agent due to the presence of vitamin B6 and omega-3 unsaturated fats.

CONCLUSION

Nigerian walnut is an emerging plant/fruit that has great potential for nutrients and drugs. The Nigerian walnut could be related to bitter kola in nature, properties, and usage. The use of walnut should be encouraged among healthy and unhealthy individuals. Walnut has shown numerous medicinal properties that could solve one or more health-related challenges.

From this study, it is discovered that Nigerian walnut has shown anti-oxidant, anti-inflammatory, anti-chelating nature, and high blood pressure control activities. Also, it has shown anti-lipidemic, anti-diabetes, anti-malarial, antimicrobial /antibacterial, and anticancer nature. The walnut has given reproductive assistance and aided sperm production and sperm quality, wound healing, anti-ulcer, and stomach disorders treatment features. The Nigerian walnut has energizing and weight management and diseases prevention capacity. It aids in bone health, skin or other tissues repairs, and could go a long way to assist in other health challenges.

It is recommended that research should continue so that the use of walnut in complementary and alternative medicine could one day lead to modern medicine discovery.

Acknowledgments: All Nigerians are showing interest in herbal and complementary medicine research.

Conflict of interest: None

Financial support: None

Ethics statement: None

REFERENCES

- [1] Obeta MU, Ikeagwulonu RC, Ohanube AKG, Jwanse IR. Some Igbo Indigenous Plants with Anti-COVID-19 Properties; A Chapter 3 in Alternative Medicine – Update Eds. Mohammad Akram IntechOpen UK. 2021:33-56. doi:10.5772/intechopen.94244
- [2] Obeta MU, Ohanube GAK, Akram M. The truth about vitamin c enriched plants and vegetables used in Nigeria for the management of COVID-19 Edelweiss. Food Sci Tech. 2021;2:7-9.
- [3] Ozioma EO, Chinwe OA. Herbal medicines in African traditional medicine. Herb Med. 2019;10:191-214. doi:10.5772/intechopen.80348
- [4] WHO. World Health Organization. Traditional Medicine Strategy. 2014-2023. 2016.
- [5] Oke OS, Oyaniyi T, Adewumi OT, Bamigboye OT, Lawah MO, Jatto KA, et al. Economic, nutritional and medicinal values of African walnut (*Tetracarpidium*

- conophorum) in Nigeria (Hutch. & Dalziel): a review. *J Res For Wildl Environ.* 2020;12(2):80-90.
- [6] Arranz S, Pérez-Jiménez J, Saura-Calixto F. Antioxidant capacity of walnut (*Juglans regia* L.): contribution of oil and defatted matter. *Eur Food Res Technol.* 2008;227(2):425-31.
- [7] Ayodeji AE, Aliyu N. *Tetracarpidium conophorum* (African walnut) Hutch. & Dalziel: Ethnomedicinal uses and its therapeutic activities. *J Med Plants Econ Dev.* 2018;2(1):1-0. doi:10.4102/jomped.v2i1.47
- [8] Kanu AM, Kalu JE, Okorie AC. Nutritional and health values of African walnut (*Tetracarpidium conophorum*). *Int J Sci Technol Res.* 2015;4(9):215-20.
- [9] Sherrow V. *Encyclopedia of hair: a cultural history.* Greenwood Publishing Group; 2006. pp.267-355.
- [10] Ndukwu MC, Ejirika C. Physical properties of the African walnut (*Tetracarpidium conophorum*) from Nigeria. *Cogent Food Agric.* 2016;2(1):1232849. doi:10.1080/23311932.2016.1232849
- [11] Oyekale KO, Odutayo OI, Esan EB, Ogunwemimo KO, Denton OA, Bolaji DT. Comparative studies on phytochemical and proximate composition of four morphologically distinct segments of the conophor seedling (*Tetracarpidium conophorum* Hutch & Dalziel). *Braz J Biol Sci.* 2015;2(3):91-100.
- [12] Amaeze OU, Ayoola GA, Sofidiya MO, Adepoju BAA, Adegoke AO, Coke HAB. Evaluation of Antioxidant Activity of *Tetracarpidium conophorum* (Mull. Arg) hutch and dalziel Leaves. *Oxid Med Cell Longev.* 2011;12:9-11.
- [13] Edem CA, Dosunmu MI, Bassey FI, Francesca I. Determination of proximate composition, ascorbic acid and heavy metal content of African walnut (*Tetracarpidium conophorum*). *Pak J Nutr.* 2009;8(3):225-6.
- [14] DS V, Remedies BF. *A Comprehensive Study.* B. Jain Publishers; 2004. pp.3-9.
- [15] Nwaoguikpe RN, Ujowundu CO, Wesley B. Phytochemical and biochemical compositions of African walnut (*Tetracarpidium conophorum*). *J Pharm Biomed Sci.* 2012;20(9):1-5.
- [16] Arinola SO, Adesina K. Effect of thermal processing on the nutritional, antinutritional, and antioxidant properties of *Tetracarpidium conophorum* (African Walnut). *J Food Process.* 2014:1-4.
- [17] Akpogheli JO, Esemefade JU, Okoh R, Ugochukwu GC. The nutritional assessment of the seed of walnut (*Plukenetia conophora*) seed purchased in an open market in Warri, Delta State. *Nigeria J Chem Soc.* 2016;41(1):125-9.
- [18] Chijioke OC, Anosike C, Ani CC. Studies on the phytochemical and nutritional properties of *Tetracarpidium conophorum* (black walnut) seeds. *J Glob Biosci.* 2015;4(2):1366-72.
- [19] Chikezie UN. Phytochemical and proximate composition of *Tetracarpidium conophorum* (African Walnut) seeds. *Int J Res Stud Biosci.* 2017;5(10):25-31.
- [20] Udedi SC, Ani ON, Anajekwu BN, Ononamadu CJ, Igwilo IO, Ibeabuchi CG, et al. Nutritional composition and antioxidant activity of African walnut, (*Tetracarpidium conophorum*). *J Appl Biochem.* 2014;107:170-80.
- [21] Onawumi OO, Faboya OO, Ayoola PB. Chemical evaluation and nutritive values of African walnut leaf (*Plukenetia conophora* Mull. arg.). *Int J Herb Med.* 2013;1(3):122-6.
- [22] Chijioke OC, Anosike C, Ani CC. Studies on the phytochemical and nutritional properties of *Tetracarpidium conophorum* (black walnut) seeds. *J Glob Biosci.* 2015;4(2):1366-72.
- [23] Barber LI, Obinna-Echem PC. Nutrient composition, physical and sensory properties of wheat-African walnut cookies. *Sky J Food Sci.* 2016;5(4):24-30.
- [24] Ezealisiji KM, Omotosho AE, Udoh RU, Agbo MO. Wound healing activity of n-hexane and methanol extracts of *Tetracarpidium conophorum* (Mull. Arg.) Hutch (African Walnut) in Wistar rats. *Malays J Pharm Sci.* 2014;12(1):79-88.
- [25] Ezealisiji KM, Ijeomah SC, Agbo MO. Anti-ulcer activity of African walnut *Tetracarpidium conophorum* nuts against gastric ulcers in rats. *Asian Pac J Trop Dis.* 2014;4(1):670-3. doi:10.1016/S2222-1808(14)60772-6
- [26] Anosike CA, Abonyi O, Etaduovie SE. Effect of methanol extract of *Tetracarpidium conophorum* seed on indomethacin-induced ulcer in rats. *Glob Vet.* 2015;14(6):848-52.
- [27] Ayoola PB, Adeyeye A, Onawumi OO, Faboya OO. Phytochemical and nutrient evaluation of *Tetracarpidium conophorum* (Nigerian walnut) root. *Int J Res Rev Appl Sci.* 2011;7(2):197-202.
- [28] Nwachoko N, Jack IR. Phytochemical screening and anti-diarrhea activities of *Tetracarpidium conophorum* induced in albino rats. *Sky J Biochem Res.* 2015;4(4):21-4.
- [29] Akinwande OO. Phytochemical, antimicrobial, proximate and elemental studies of the leaves of *Tetracarpidium conophorum* (Mull. Arg.) Hutch and Dalziel', Ph.D. dissertation, Department of Chemistry Faculty of Science. Ahmadu Bello University, Zaria Nigeria, 2015:1-168.
- [30] Ajaiyeoba EO, Fadare DA. Antimicrobial potential of extracts and fractions of the African walnut–*Tetracarpidium conophorum*. *Afr J Biotechnol.* 2006;5(22):2322-5.

- [31] Olabinri BM, Eniyansoro OO, Okoronkwo CO, Olabinri PF, Olaleye MT. Evaluation of chelating ability of aqueous extract of *Tetracarpidium conophorum* (African walnut) in vitro. *Int J Appl Res Nat Prod.* 2010;3(3):13-8.
- [32] Olaniyi FE, Bambidele IO, Omokehinde AO, Ayodeji AA. Antiinflammatory activities of the chloroform extract of the fruit of *Tetracarpidium conophorum* (Mull. Arg.) (Nigeria Walnuts). *J Adv Med Pharm Sci.* 2016;6(1):1-7. doi:10.9734/JAMPS/2016/22898
- [33] Oladokun BO, Omisore ON, Osukoya OA, Kuku A. Anti-nociceptive and anti-inflammatory activities of *Tetracarpidium conophorum* seed lectin. *Sci Afr.* 2019;3:e00073. doi:10.1016/j.sciaf.2019.e00073
- [34] Akomolafe SF, Oboh G. Walnut leaf extract acts as a fertility agent in male Wistar albino rats—A search for herbal male fertility enhancer. *J Complement Integr Med.* 2018;15(2):1-14.
- [35] Ozyigit II, Uras ME, Yalcin IE, Severoglu Z, Demir G, Borkoev B, et al. Heavy metal levels and mineral nutrient status of natural walnut (*Juglans regia* L.) populations in Kyrgyzstan: nutritional values of kernels. *Biol Trace Elem Res.* 2019;189(1):277-90. doi:10.1007/s12011-018-1461-4
- [36] Akomolafe SF, Oboh G, Akindahunsi AA, Afolayan AJ. *Tetracarpidium conophorum* ameliorates oxidative reproductive toxicity induced by ethanol in male rats'. *BMC Complement Altern Med.* 2015;15(1):439-925.
- [37] Obianime AW, Uche FI. The comparative effects of aqueous extract of *Tetracarpidium conophorum* seeds and Proviron on the sperm parameters of male guinea pigs. *Asian Pac J Trop Med.* 2009;2(6):11-4.
- [38] Obianime AW, Uche FI. The effects of aqueous extracts of *Tetracarpidium conophorum* seeds and proviron on the sperm parameters of male guinea pigs. *Asian Pac J Trop Med.* 2010;3(1):21-4. doi:10.1016/S1995-7645(10)60024-7
- [39] Akomolafe SF, Oboh G, Akindahunsi AA, Afolayan AJ. Antiperoxidative activity of *Tetracarpidium conophorum* leaf extract in reproductive organs of male rats'. *Evid Based Complement Alternat Med.* 2015:1-8.
- [40] Ikpeme EV, Ekaluo UB, Udensi O, Ekerette EE, Ekpo PB, Asuquo BO. Sperm quality and hormone profile of male albino rats fed with seeds of African walnut (*Tetracarpidium conophorum*, Mull). *Annu Res Rev Biol.* 2014;4(9):1379-86.
- [41] Akomolafe S, Oboh G, Olasehinde T, Oyeleye S, Ogunsuyi O. Modulatory effects of aqueous extract from *Tetracarpidium conophorum* leaves on key enzymes linked to erectile dysfunction and oxidative stress-induced lipid peroxidation in penile and testicular tissues. *J Appl Pharm Sci.* 2017;7(1):51-6. doi:10.7324/JAPS.2017.70107
- [42] Nwaoguikpe RN, Ujowundu CO, Wesley B. Phytochemical and biochemical compositions of African walnut (*Tetracarpidium conophorum*). *J Pharm Biomed Sci.* 2012;20(9):1-5.
- [43] Onwuli DO, Bown H, Ozoani HA. Anti-hyperglycemic effect of *Tetracarpidium conophorum* in alloxan-induced diabetic female albino rats. *Int Sch Res Notices.* 2014;1(4). doi:10.1155/2014/124974
- [44] Ogunyinka BI, Oyinloye BE, Adenowo AF, Kappo AP. Potentials of some plant-derived foods in the management of diabetes and associated complications. *Afr J Tradit Complement Altern Med.* 2015;12(6):12-20. doi:10.4314/Ajtcam.V12i6.2
- [45] John OO, Monday UP, Nneka IW, Ogom OG, Uwaifiokun OC, Benedict NL. Effect of root and leaf extracts of *Tetracarpidium conophorum* on liver enzyme levels in alloxan induced diabetic rats. *J Pharm Res Int.* 2015;5(4):241-8. doi:10.9734/BJPR/2015/13407
- [46] Ezealisiji KM, Stanley CN, Ekanem ES. Evaluation of anti-cholesterol activity of ethyl acetate and n-hexane extracts of *Tetracarpidium conophorum* (Mull. Arg.) Hutch and Dalziel (African walnut) towards hypercholesterolemic rats. *Int J Pharmacogn Phytochem Res.* 2016;8(8):1372-6.
- [47] Analike R, Ahaneku JE, Njoku MC, Ahaneku GI, Ezeugwunne IP, Ogbodo EC. Effects of *Tetracarpidium conophorum*—“Nigerian Walnuts” on blood lipids, lipoproteins and glucose values in adult Nigerians. *Int J Innov Res Adv Stud.* 2017;4(7):67-71.
- [48] Kanu AM, Kalu JE, Okorie AC, Olabinri BM, Eniyansoro OO, Okoronkwo CO, et al. Evaluation of chelating ability of aqueous extract of *Tetracarpidium conophorum* (African walnut) in vitro. *Int J Appl Res Nat Prod.* 2015;3(3):1-13.
- [49] Dada EO, Ogundolie OO. In vivo antiplasmodial activity of raw ethanolic seed extract of *Tetracarpidium conophorum* in Swiss albino mice infected with *Plasmodium berghei*. *J Adv Med Pharm Sci.* 2016;9(2):1-8. doi:10.9734/JAMPS/2016/26789
- [50] Ogundolie OO, Dada EO, Osho IB, Oloruntola DA. Effects of raw ethanolic seed extract of *Tetracarpidium conophorum* on hematological and histopathological parameters in Swiss albino mice infected with *Plasmodium berghei*. *J Appl Life Sci Int.* 2017;12(2):1-14. doi:10.9734/JALSI/2017/33244