

AMERICAN DRUG INDEX

AMERICAN DRUG INDEX

1963

CHARLES O. WILSON, Ph.D.

*Professor of Pharmaceutical Chemistry, School of Pharmacy,
Oregon State University*

AND

TONY EVERETT JONES, Ph.D.

*Assistant Professor of Pharmaceutical Chemistry, School of Pharmacy,
University of Colorado; Director of
Pharmaceutical Research, Carbisulphoil Company*



J. B. LIPPINCOTT COMPANY

Philadelphia and Montreal

Copyright © 1963, by J. B. Lippincott Company


Copyright © 1958 1958 1959, 1960, 1961, 1962 by J. B. Lippincott

This book is fully protected by copyright and, with the exception of brief extracts for review, no part of it may be reproduced in any form without the written permission of the publishers

Distributed in Great Britain by
Pitman Medical Publishing Co., Limited, London

Library of Congress Catalog Card Number 55-6286

Printed in the United States of America



5.11
WIL-63

PREFACE

The American Drug Index has been prepared for the identification and correlation of the many pharmaceuticals available to the medical and allied professions. The need for this index has become acute as the number of drugs and drug products in all their tremendous variety has multiplied.

The organization of the index is fundamentally alphabetical with extensive cross-indexing. Names listed are generic (also called non-proprietary, public name or common name); brand (also called trade-mark, proprietary or specialty), chemical, U.S.P., N.F., N.N.D. and A.D.R. Synonyms that are in general use also are included. All names used for a pharmaceutical occur in alphabetical order with the pertinent data given under the brand name and the name by which it is made available. Data included are generic names, chemical names, manufacturer, pharmaceutical forms, size, dosage and use. Generic names occur in alphabetical order and here the recognition of the drug is indicated by U.S.P. (United States Pharmacopeia), N.F. (National Formulary), N.N.D. (New and Nonofficial Drugs) and A.D.R. (Accepted Dental Remedies). The information is in accord with the latest editions of these books: U.S.P. XVI, N.F. XI, N.N.D. 1962, A.D.R. 1962.

Because of the multiplicity of specialty names (brand names) used for the same therapeutic agent or the same combination of therapeutic agents, it was apparent that some correlation could be done. As an example of this, please turn to Aluminum Hydroxide Gel. Here under the generic name are given the brand names. Following are combinations of aluminum hydroxide gel organized in a manner to point out relationships among the many products. Reference then is made to the specialty name or names having the indicated composition. Under the specialty name are given manufacturer, composition, available forms, sizes, dosage and use.

The multiplicity of generic names for the same therapeutic agent has complicated the nomenclature of these agents. Examples of multiple generic names for the same chemical substance are (1) parabromdylamine, brompheniramine; (2) acetaminophen, p-hydroxy acetanilid, N-acetyl-p-aminophenol; (3) d-isoephedrine, d-pseudoephedrine; (4) methapyrilene HCl, thenylpyramine HCl; and (5) pyrilamine, pyranisamine, pyranilamine, pyraminyl and anisopyradamine.

Generic names recognized by the United States Pharmacopeia Committee, National Formulary Committee and the American Medical Association Council on Drugs are preceded by an asterisk.

The American Drug Index should be most useful to physicians, pharmacists, dentists, nurses, and to sales personnel, students and teachers in the fields incorporating pharmaceuticals. The cross-indexing feature permits the finding of drugs or drug combinations when only one major ingredient is known. For example, a combination of aluminum hydroxide gel and mag-

PREFACE

vi

nesium trisilicate is available. This combination can be found by looking under the name of either of the two ingredients and in each case the specialty names are given.

CHARLES O. WILSON

TONY E. JONES

FOREWORD

By Joseph B. Jerome, Ph.D.
Assistant Secretary Council on Drugs
American Medical Association

Identification of drugs, as of all common objects, benefits from the use of proper nomenclature. Physicians, educators, pharmacists and the associated health professions all have a need for an up-to-date and accurate publication which will enable these individuals to identify specific preparations despite the number of designations which normally come into use during the development of each commercial drug.

The authors of the *American Drug Index* are primarily concerned with recording trademarks, nonproprietary names and, in the case of mixtures, a statement of composition together with dosage form information. Since this book is in the nature of a compilation and is intended for research students as well as those engaged in the active practice in the health professions, it is necessary to include in the confines of such a work as this all nonproprietary names and as many trademarks as have come into use in the American medical press. The features that serve to distinguish the *American Drug Index* are its attempts to provide up-to-date and complete nomenclature information and the special recognition given to formally adopted nonproprietary names (names adopted by the U.S.P., the N.F., or the A.M.A.-U.S.P. Nomenclature Committee).

Under present world conditions, a new drug discovery in one country, should it prove significant, very readily crosses national borders and becomes available in other countries as well. The only readily accessible designation which crosses national borders along with the drug is its nonproprietary name.

Some confusion which has arisen in the past in drug nomenclature has resulted from the absence of one known, formally adopted nonproprietary name for each drug. Part of the problem lies in the failure of some manufacturers to cooperate in a centralized selection process using the existing mechanism (at the present time, the A.M.A.-U.S.P. Nomenclature Committee) to formally adopt an appropriate nonproprietary name for each drug.

The Council on Drugs of the American Medical Association provides a secretarial service to the A.M.A.-U.S.P. Nomenclature Committee and recommends the following specific procedure to all drug manufacturers marketing new drugs.

1. The Council encourages all manufacturers who have discovered or developed a new drug to submit a proposed nonproprietary name (or names) to the Secretary, Council on Drugs,* at such time as its preliminary

* 535 N. Dearborn St., Chicago 10, Ill

clinical evaluation indicates that more extensive clinical testing is warranted, or sooner if possible. These submissions will be referred to the joint Nomenclature Committee of the American Medical Association United States Pharmacopoeia for its consideration.

2. The initial submission of nonproprietary terminology should be accompanied by disclosure of the manufacturer's code number, the chemical name (Chemical Abstracts index names only are to be used) and the structural formula of the compound when these are known or by a description of the source and general method of preparation that accurately defines the product. The submission should also indicate the general pharmacological class or area of intended clinical use of the drug and, when available, the trade name which is to be applied.

3. The Nomenclature Committee and Council staff will review the proposed name(s) and, if necessary, negotiate with the manufacturer to ensure that a convenient and appropriate nonproprietary name is adopted for each new drug. The guiding principles for devising international nonproprietary names of the World Health Organization are regarded as helpful but not definitive in achieving this purpose.

4. Upon agreement between the manufacturer and the Nomenclature Committee on a suitable nonproprietary name, and with the express permission of the manufacturer (which can be given with the initial submission of a proposed nonproprietary name), the Council then transmits the proposed name to the World Health Organization, the British Pharmacopoeia Commission and the representatives of the French Codex, the Nordic Pharmacopoeia, the United States Pharmacopoeia and the National Formulary for their simultaneous consideration. In the case of serums or vaccines, the proposed terminology is also transmitted to the Division of Biologic Standards of the National Institutes of Health.

5. In the absence of serious objection from the foregoing agencies or known conflict with established nonproprietary or trade names, after a waiting period of 30 days, the proposed name is considered to be acceptable to all concerned. The adoption is recorded as final by the Council and the manufacturer is informed. While this adoption is for national use, the Council staff will exert its best efforts to effect international adoption of the same terminology through the mechanisms of the World Health Organization.

6. All nonproprietary names adopted in accordance with the provisions of this cooperative nomenclature program will be the designations used when and if the drugs are admitted to the United States Pharmacopoeia or the National Formulary.

The early formal adoption of an appropriate nonproprietary name coupled with the increasing publicity given such United States Adopted Names (USANs) by the A.M.A. U.S.P. Nomenclature Committee and the implementation accorded such names by the use of identifying notations in the American Drug Index represent significant advances toward clarifying existing drug nomenclature problems.

FOREWORD

By Lloyd C. Miller, Ph.D.

Director of Revision
United States Pharmacopoeia

The *American Drug Index* meets a profound and long-felt need in a unique fashion. Considering how sorely it was needed by the world of medicine and pharmacy, the wonder is that efforts to provide the Index first bore fruit no earlier than five years ago. Nothing like it existed before, and, in view of the immense labor involved in its preparation, it is likely to remain the only one of its kind—the most comprehensive current listing of drug nomenclature extant.

In simplest terms, the *American Drug Index* is a special kind of dictionary. To debate the need for almost any kind of dictionary has long been regarded as utterly pointless. Yet the Index goes far beyond merely defining terms. It distinguishes among the multiple names by which drugs are known, indicating in a helpful way the status of each name. It provides in condensed form an immense amount of information on the products to which the terms apply, including the quantitative composition of their various forms and the sources from which they are available. The conciseness with which all these facts are set forth is enough to make the compilation exceedingly useful in areas beyond those in which it is indispensable.

In an annual publication such as this, it is not inappropriate to take note of events having a bearing on drug nomenclature such as the proposals now before the Congress. If enacted, the latter will require a government agency to publish a list of the names by which drugs are known. Because of limitations imposed upon a government publication of this kind, such as attend the necessary use of trade names in a list that inevitably would assume an aura of legal status, no government list could approach the *American Drug Index* in comprehensiveness or utility. In any case, the duplication of function would be deplorable. It seems safe to say, in retrospect, that had the Index been available only a few years earlier, federal intervention in this area might not have been suggested.



FOREWORD

By Edward G. Feldmann, Ph.D.

*Director of Revision
National Formulary*

Many publications are presently available which are promoted on the basis of the information that they will provide the reader to identify and describe specific drug products on the American market. While undoubtedly most of them serve a useful function, this writer knows of none which provides as complete coverage and, therefore, is nearly as useful as the *American Drug Index*.

The editors have wisely restricted the types of information covered in their book to a number of well defined areas, and thus have been able to provide a unique service to the reader which is relatively exhaustive without becoming cumbersome.

The drug names given in the *National Formulary* and the *United States Pharmacopeia* must, of necessity, be limited to the nonproprietary or so called generic names, because of the official status accorded these two books under federal and state drug laws. Yet in order to make suitable use of the information provided in the official compendia it is necessary to correlate the information provided in the *NF* and *USP* monographs with pertinent aspects of the corresponding marketed forms of those drugs, namely their trade names, manufacturer's names, dosage forms available and concentrations in the dosage forms.

The American Pharmaceutical Association has been acutely aware of the particular need to correlate the nonproprietary or official names given in the *NF* and the *USP* with the brand or trade names under which many of these drugs are marketed, distributed and prescribed. In an effort to meet this need the A.P.A. prepared a booklet titled *Proprietary Names of Official Drugs*. This booklet was first issued when *USP XV* and *NF X* were official and has undergone three revisions since then. The *American Drug Index*, however, has the added advantage of supplying similar information for drugs not in the official compendia. It also supplies other information in addition to the drug names for all of the drugs including those which are *NF* or *USP* items.

Consequently it is a pleasure to accord a high recommendation to the 1963 edition of *ADI* which will surely prove to be as valuable and useful as the earlier volumes in the series.

- A.A.S. COMPOUND.** (Tilden) Aspirin 5 gr., nux vomica ext. 1/10 gr., hyoscyamus ext. 1/8 gr., galtemium tr. 3 min., cascara sagrada ext. 1 gr./Tab. Bot. 100s, 1000s.
Use: Analgesic.
- ABAC.** (Mossengill) Aspirin 5 gr., caffeine 1/4 gr., Pot. bromide 1/4 gr., Sod. bromide 1/4 gr./Tab. Bot. 100s, 1000s.
Use: (1-2 q. 3 h.) Analgesic sedative.
- ABAC FOR CHILDREN.** (Mossengill) Aspirin 1 gr., caffeine 1/10 gr., Pot. bromide 1/20 gr., Sod. bromide 1/20 gr., aromatics q.s./Tab. Bot. 1000s.
Use: (1 q. 4 h.) Analgesic.
- ABACTAL.** (Mossengill) Aspirin 5 gr., caffeine 1/4 gr., Pot. bromide 1/4 gr., Sod. bromide 1/4 gr., phenobarbital 1/2 gr./Tab. Bot. 100s, 1000s.
Use: (1-2 t.i.d.) Analgesic sedative.
- ABASIN.** (Winthrop Labs) Acetylcorbromol, Tab. (4 gr.) Bot. 100s.
Use: (1-2 t.i.d.) Sedative.
- ABOCILLIN 800 M., A.D.R.** (Abbott) Procaine penicillin G 600,000 u., buffered Pot. penicillin G 200,000 u. for aq. inj. Vial, 1 cc., Box 25s, 5 cc., 1s, 25s.
Use: I.M.; bacterial infections.
- ABOCILLIN-DC, A.D.R.** (Abbott) Procaine penicillin O in aq. susp. 600,000 u. 1 cc. Absoject disposable syr. w/needle, 1s, 25s.
Use: I.M.; penicillin therapy.
- ABO-VAC.** See: A.C.D. Solution. (Abbott).
- AB-CEDET.** (Beard). B-1 10 mg., B-2 10 mg., B-6 10 mg., B-12 6 mcg., A 12,500 U.S.P. u., D 1,250 U.S.P. u., calcium pantothenate 5 mg., C 100 mg./Tab. Bot. 100s, 1000s.
Use: Therapeutic vitamin preparation.
- ABDEC DROPS.** (Parke, Davis) Vit. A 5000 u., D 1000 u., B-1 1 mg., B-2 1.2 mg., B-6 1 mg., sod. pantothenate 5 mg., nicotinamide 10 mg., C 50 mg./D 6 cc. Bot. 15 cc., 50 cc.
Use: Vitamin deficiencies.
- ABDEC KAPSELS.** (Parke, Davis) Vit. A 10,000 u., D 1000 u., B-1 5 mg., B-2 3 mg., B-6 1.5 mg., B-12 2 mcg., di-pantothenol 10 mg., nicotinamide 25 mg., Vit. C 75 mg., mixed tocopherals 5 mg./Cap. Bot. 50s, 100s, 250s, 1000s.
Use: (1 daily) Multivitamin therapy.
- A-B-D-G.** (Blue Line) Vit. A 5000 u., D 1000 u., B-1 1 mg., B-2 2 mg./Cap. Bot. 100s.
Use: (1 daily) Dietary supplement.
- A-B-O-G VITAMIN.** (Hortan & Converse) Vit. A 5000 u., D 1000 u., B-1 333 u., riboflavin 2000 mcg./Tab. Bot. 100s, 250s, 500s, 1000s.
Use: (1 daily) Multivitamin therapy.
- A-B-D-G VITAMINS.** (Upjohn) Vit. A 2500 u., D 250 u., B-1 1.5 mg., B-2 1 mg., nicotinamide 10 mg./Cop. Bot. 50s, 100s, 250s, 1000s.
Use: (1 daily) Multivitamin therapy.
- ABOOL W/MINERALS.** (Parke, Davis) Vit. C 40 mg., B-2 2 mg., B-12 1.5 mcg., B-6 0.5 mg., B-1 1 mg., nicotinamide 12 mg., A 3000 u., D 1000 u., col. pantothenate 2.5 mg., folic acid 0.1 mg., w/iron, copper, zinc, magnesium, calcium, phosphorus, iodine, manganese, cobalt, and pot./Cap. Bot. 100s.
Use: Vitamin-mineral supplement.
- ABOOL W/VIT. C Cap.** (Parke, Davis) Vit. A 5000 u., D 1000 u., B-1 2.5 mg., B-2 2.5 mg., B-6 0.5 mg., B-12 1 mcg., nicotinamide 20 mg., Vit. C 50 mg., pantothenic acid (as Cal. salt) 5 mg./Cap. Bot. 25s, 100s, 250s, 500s.
Use: (1 daily) Vitamin deficiency.
- A-B-M-C OINTMENT.** See: Lembrose, Oint. (Wyeth).
- ABMINTHIC, N.N.D.** 1960. (Pfizer) Dithiazanine iodide, 3 3'-Diethylthiodicarbocyanine iodide. 200 mg./Tab. Bot. 100s.
Use: Anthelmintic.
- ABOOL.** See: Methiodal Sodium, (Winthrop Labs).
- A-B-S & C.** (Hortan & Converse) Atropine 1/120 gr., belladonna lvs ext. 1/8 gr., cascara sagrada 1/2 gr./Tab. Bot. 100s, 1000s.
Use: (1 h.s.) Laxative.
- ABSOIN.** (Lammon) N-acetyl-p-aminophenol, 150 mg.; salicylamide, 150 mg.; phenyltoloxamine dihydrogen citrate, 25 mg.; phenylephrine HCl, 2.5 mg./Tab. Bot. 100s
Use: Sinus headache.
- ABSOBBABLE CELLULOSE COTTON OR GAUZE, N.N.D.** 1959. See: Oxidized Cellulose. (Various Mfrs.).
- ABSOBBABLE DUSTING POWDER, U.S.P., N.N.D.** 1959. See: Bio-Sorb, Paw. (Ethicon)
- ABSOBBABLE GELATIN FILM, N.N.D.** 1961. Sterile nonantigenic, absorbable, water-insoluble, gelatin film. See: Gelfilm. (Upjohn)

***ABSORBABLE GELATIN SPONGE, USP**
NND 1962, ADR Gelatin sponge See
Geltaam, Preps (Upjohn)

ABSORBABLE SURGICAL DRESSINGS, See
Hema Pak (Johnson & Johnson)

***ABSORBABLE SURGICAL SUTURE, USP**
ADR (Various Mfr) Surgical Gut, Surgical
Catgut, Catgut suture

ABSOBENT BASE (Upsher Smith) Oint
Base Cetyl alcohol, stearyl alcohol, lauryl
sulfate, methyl p hydroxybenzoate & hydro
carbons Jar 1 5, 6, 25 lb
Use Ointment base

ARSORBENT OINTMENT BASE,
See: Plastibase Hydrophilic (Squibb)
Polysorb (Faugera)

ARTEN (Paul Maney) Rauwolfia serpentina
alkaloids ext Tab (1 mg & 2 mg) Bot
100s, 1000s
Use (2 daily) Hypotensive and tranquiliz
ing agent

ARUPHENINE CITRATE. (Chemo Pura) Di
ethylaminoethyl phenylethylacetate citrate
Pow for mfg
Use Antispasmodic

ARUPHENINE HYDROCHLORIDE (Chemo
Pura) Diethylaminoethyl phenylethylacetate
hydrochloride Pow for mfg
Use Antispasmodic.

A-C TROCHES (Abbott) Anesthesin ¼ gr,
colocid & licorice Rot 24s, 100s, 1000s
Use Throat & mouth infections

***ACACIA, USP, ADR** (Penick) Acacia
senegal, Rumarabic Mucilage, pow, ¼ lb,
1 lb, tears, ¼ 1 lb, syr; whole, ¼ 1 lb
Use: Demulcent, emulsifier

W/Mag oxide & veg mucinoid See: Aco
resil, Pow (Sherman)
W/Veg mucinoid See: Acarojel Pow
(Sherman)

ACA. COMPOUND. (Eowell) Aspirin 3½
gr, phenacetin 2½ gr, caffeine ½ gr/
Cap or Tab Bot 100s, 1000s Also with
codeine phos ¼, ½, 1 and 1 gr See:
AFC

Use (1 2 q 3 h) Analgesic & antipyretic

A-C-A-COMPOUND (Sutliff & Case) Acet
anilid 130 mg, Camphor 16 mg, Capsicum
16 mg, Alain 4 mg, Podophyllum 16 mg,
Acanite Root 0.65 mg, Atropine Sulfate 0.03
mg, Tr Gelsemium 0.2 cc /Tab Bot 1000s
Use: Analgesic & antipyretic.

ACAP, (Barre) Vit A 50,000 u /Cap Bot
100s
Use: Vitamin A deficiencies

ACAPNOID. (Chemica) Belladonna pow
1/10 gr, phenobarbital ½ gr, stramonium
* Generic name recognized by NF USP or A.M.A.

pow 1 gr, ephedrine sulfate ¼ gr /Tab
Bot 100s, 500s, 1000s

Use: Sedative & antispasmodic
ACARALUM (Sherman) Carbohydrate ura
nides (veg mucinoid) 50%, acacia 20%,
aluminum hydroxide gel 25%, Mag oxide
5% Pow Can R oz

Use (2 dr p r n) Gastric antacid
ACAROJEL (Sherman) Carbohydrate ura
nides (veg mucinoid) 50%, acacia 50%.
Pow Can R oz
Use Laxative

ACAROSIL (Sherman) Carbohydrate ura
nides (veg mucinoid) 50%, acacia 25%,
Mag oxide 25% Pow Can R oz
Use (1 dr p r n) Gastric antacid

ACCESSORONE (Upjohn) Vit A 22 600 u,
D 267 u red bone marrow ext 0.62 cc,
iron & ammonium citrate 0.259 Gm, molt
ext, yeast ext, citrum conc qs /fl oz
Rot 10 fl oz, 24 fl oz
Use Dietary supplement

ACCO. (Assoc Concentrates) Protein, 32%,
carbohydrate 54%, ash 5.75%, fat 0.75%,
crude fiber 1.61%, Vit B 1 2 mg, R 2 3
mg, nicotinic acid 15 mg Jar 15 oz
Use: Nutritional supplement

A-C-B SOLUTION, USP Sod citrate, citric
acid & dextrose in sterile pyrogen free
solution

Abbott—(Abbo Vac) 150, 250, 500 cc
(non Vac) 250 & 500 ml (Pliopob) w/Blood
collection set 500 cc

Raxter—600 cc bot with 70 120 or 300
cc of sol; 1000 cc with 500 cc of sol

Cutter—500 cc bot with 75 or 120 cc of
sol; 650 cc bot with 80 or 130 cc of sol
Use Anticoagulant for preparation of
plasma or whole blood

ACE-CAT-EDINE, ADR 1955 (Paul Maney)
Aspirin 3½ gr, acetophenetidin 2½ gr,
caffeine ½ gr/cap or Tab Cap, Bot
100s, 1000s; Tab, Na 1, Bot 100s 1000s;
also w/phenobarbital, Bot 1000s See
AFC

Use (1 2 q 3 h) Analgesic & antipyretic

ACECOLINE. (Angla French) Acetylcholine
chloride 20 50, 100 or 200 mg w/saligenin
40 mg/cc in propylene glycol Amp, 1 cc
Box 6s

Use SC, 1 M; parasympathetic agent

ACE-JEL (Ortho) Acid vaginal jelly buffered
to pH of 4.0, containing ricinoleic acid,
boric acid, oxyquinoline sulfate & propyl
ester parahydroxybenzoic acid in water
dispersible base Tube with or without ap
plicator

ACENOCOUMARIN. See: Sintrom, Tab
(Gargy)

- * **ACENOCOUMAROL**, N. N. D. 1962. 3-(α -acetonyl-p-nitrobenzyl)-4-hydroxycoumarin. See: Sintram, Tab. (Geigy)
- ACEPROMAZINE**. 10(3-dimethylaminopropyl)-3-acetylphenothiazine maleate. See: Atrovel (Ayerst), Plegicil (Ayerst)
Use: Tranquilizer.
- ACETABAR**, (Columbus) Phenobarbital $\frac{1}{4}$ gr., acetophenetidin $2\frac{1}{2}$ grs., acetylsalicylic acid $3\frac{1}{2}$ grs., caffeine $\frac{1}{4}$ gr./Tab. or wafer. Bots. 100s, 1000s.
Use: (1-2 q. 4 h.) Analgesic-sedative.
- ACETALAX**, (G. F. Harvey) Diacetyldihydroxyphenalisanin. Cap. 5 mg., Bot. 100s, 1000s.
Use: (1-2 h.s.) Laxative.
- p-**ACETAMIDO BENZOIC ACID SALT** of 2-dimethylaminoethanol, See: Deones, Tab. (Riker)
- * **ACETAMINOPHEN**, N.P. See: N-Acetyl-*p*-aminophenol.
- ACETAMINOPHENOL**. Acetaminophen. See: N-acetyl-*p*-aminophenol.
- ACETANILID**, A.D.R. (Various Mfrs.) (Acetylaminobenzene, acetylaniline, onifabesin) N-Phenylacetamide Cry. Bot. 1 oz., $\frac{1}{4}$ lb., 1 lb.
Use: Antipyretic, analgesic.
Powder Compound, N.P. (Various Mfrs.)
W/Acetophenetidin, salol & caffeine. See: Ferbrisol, liq. (Tilden).
W/Anesthesia, ephedrine alkaloid, bismuth subnitrate, petrolatum, menthol, eucalyptol. See: Nosalene, No. 1, liq. (Abbott).
W/Camphor, sopsicum, oilain, podophyllum, ozonite root, atropine sulfate, sr. gelsemium. See: A.C.A. Compound, Tabs. (Sutliff & Cose).
W/Quinine sulfate, podophyllin, oilain, aconite root, capsicum, atropine sulfate, and camphor. See: C.C.G., Tab. (Noyes).
- ACETARSOL**. See: Acetarsone, Tabs. (Abbott)
- * **ACETARSONE**, N.F., N.N.R. 1950s Tablets, N.F. (Abbott) 3-Acetamido-4-hydroxyphenylarsonic acid (Acetarsol, Acetphenarsone, Amarsol, Dynarsol, Ehrlich 594, Uinarsol, Orarsol, Osarsol, Foraxyl, Stavarsol.) Tab. (0.25 Gm.). Bot. 100s, 500s.
Use: Amebiasis & trichomoniasis.
W/Hydrolyzed carbohydrates. See: Devagan, Tabs. (Winthrop Labs.).
W/Polymyxin B, Neomycin. See: Neosparin vag. Supp. (Burroughs Wellcome).
- ACETARSONE SALT OF ARECOLINE**. See: Dracarbil, N.F. (Winthrop Labs.)
- * **ACETAZOLAMIDE SODIUM**, U.S.P., N. N. D. 1962. Sodium salt, U.S.P.
See: Diamax, Tabs. (Lederle).
Diomax Sequels, Cops. (Lederle)
- ACET-DIA-MER-SULFONAMIDE**, N.F., N.N.R.
* Generic name recognized by N.F., U.S.P. or A.M.A.
1935. Sulfacetamide, Sulfadiazine & Sulfamerazine.
See: Buffanamide, Susp. (Tulag)
Cefazine, Tab. (Bowman Bros.)
Dorsulfas, Susp. & Tabs. (Smith-Dorsey)
Incarposul, Susp. & Tabs. (Blue Line).
Famsulla, Susp., Tabs. (Merrell).
Sulfadiazine, sulfamerazine & sulfacetamide, Susp., N.F. Tab. N.F.
Sultrid (Taylor).
Tilamide, liq. (Tilden).
Tricambisol, Susp. & Tabs. (Schering)
Trizyl, Susp. & Tabs. (Central).
W/Gantrisin & Pot. penicillin G. See: Trio nacil, Tab. (Flint, Eaton)
W/Hyoscymus Ext. See: Suramate, Tab. (Patch)
W/Sulfamethoxine.
See: Delamets, Tab. (Durst).
Deltamide, Tab. or Susp. (Armour)
Katrosul, Tabs., Susp. (Chicago).
Quad-Suffo, Tab. (Zimmer).
W/Sulfamethazine, Pot. penicillin G. See: Deltamide w/Penicillin, Tab. or Supp. (Armour)
W/Urea, neocalamine, talc, glycerin, camphor, menthol. See: Neofanamide, liq. (McNeil)
- ACETEST**, (Ames). Sod. nitroprusside, disodium phosphate, aminoacetic acid, ioclose. Tab. Bot 100s, 250s.
Use: Test for acetone as diacetic acid in urine.
- ACETICYL**. See: Acetylsalicylic Acid.
- ACETIDINE**, A.D.R. (Merck Sharp & Dohme) Aspirin 6 pt., phenocellin 4 pt., caffeine 1 pt. Caps., pink, 5 gr., Bot. 100s, 1000s; white, 5 gr., Bot. 12s, 36s, 100s, 1000s; Tab., pink or white, 5 gr., Box, Bot. 100s, 1000s. See: A.P.C.
Use: (1-2 q. 3 h.) Analgesic, antipyretic.
W/Codaine phosphate $\frac{1}{4}$ gr., A.D.R. Tab. 100s, 500s.
Use: (1-2 q. 3 h.) Analgesic, antipyretic.
W/Codaine phosphate $\frac{1}{2}$ gr., A.D.R. Tab. 100s, 500s.
Use: (1-2 q. 3 h.) Analgesic, antipyretic.
- ACETILUM ACIDULATUM**. See: Acetylsalicylic Acid. (Various Mfrs.)
- ACTIBRO, PEDIATAB No. 1**. (Columbus Pharm.) Bromisovalum $\frac{1}{2}$ gr., ethyl salicylate carbonate 1 gr., acetophenetidin $\frac{3}{2}$ gr., caffeine $\frac{1}{16}$ gr./tab. Bot. 100s, 1000s.
Use: (1-2 q. 3 hr.) Sedative.
- ACETO-CORT**. (Mallard) Hydrocortisone 1.5 mg., aspirin 5 gr., aluminum hydroxide gel 100 mg./Tab. Bot. 100s, 1000s.
Use: Treatment of mild to moderate rheumatoid arthritis.

ACETOHEXAMIDE

***ACETOHEXAMIDE** (Eli Lilly) N [p acetyl phenylsulfonyl]-N cyclohexylurea
See: Dymelor
Use: Blood sugar lowering compound

ACETOL. See: Acetylsalicylic Acid (Various Mfr)

ACETOLACTIN (G P Harvey) Acetphenolsolin 5 mg /Cap Bot 100s 1000s
Use: Laxative or purgative

***ACETOMEBUCTOL**, NND 1962 See: Merbak, Prep (Schieffelin)

ACETONE or **DIACETIC ACID TEST**. See: Acetest, Tab (Ames)

ACETONYL (Upjohn) Granular effere salt Aspirin as sod salt 7½ gr w/Pol & Sod citrates & tartrates, calcium lactate, Sod phosphate & magnesium sulfate/2 oz Bot 4 oz
Use: (1 lps) Analgesic & antifebrile

ACETOPHEN, (Smith Dorsey) Phenobarbital ½ gr, aspirin 3 gr, acetophenetidin 2 gr, hyoscyamus ext ½ gr /Cap Bot 100s 1000s
Use: (1 lid) Sedative—analgesic

***ACETOPHENAZINE MALEATE**, 2 acetyl 10 {3 [4 (p hydroxyethyl)piperazinyl]propyl} phenothiazine dimaleate
See: Tindol (Scharing Corp)

***ACETOPHENETIDIN**, USP, NNR 1941, ADR; Tablets, USP (Various Mfr) (Phenacetin p Ethoxyacetanilide) Cry or Pow 1 oz, ¼ lb, 1 lb, 5 lb
Use: Antipyretic, analgesic
W/Aspirin, caffeine See: APC

ACETOSAL. See: Acetylsalicylic Acid (Various Mfr)

ACETOSALIC ACID. See: Acetylsalicylic Acid (Various Mfr)

ACETOSALIN. See: Acetylsalicylic Acid (Various Mfr)

ACETOULPONE SODIUM 4,4' diaminodiphenylsulfone 2 N-acetylsulfonamide See: Promacetin, Tab (Parke, Davis)

ACETO-VIB. (Mallard) Aspirin 5 gr, veratrum viride Pow 1 gr, rutin 20 mg, Vit C 30 mg /Tab Bot 100s, 500r 1000s
Use: Capillary dilator

ACETPHIN-ACETYL COMPOUND. (First Texas) Aspirin 3 gr, acetophenetidin 2 gr, caffeine ½ gr, gelurem tr 1 min /Cap Bot 100s, 500s, 1000r See APC w/gelurem

Use: (2-3 cap daily) Analgesic & antipyretic

ACETPHINABEINIL. See: Acetorsone, Tab (Abbott)

ACETPHENOLETIN (Chemo Pure) Pow Pig for mg

* Generic name recognized by N.F., USP or A.M.A.

See: Diacetylhydroxyphenylisatin, Prep (Various Mfr)

Use: Laxative
ACETPYBOGALL, NNR 1951 Pyrogallol triacetate

***ACETRIZOATE SODIUM**, NND 1962 Injection USP 3 Acetylaminio 2,4,6 Iriado benzoate sodium

See: Thixakon (Mallinckrodt)
Urokon Sodium, Amp (Mallinckrodt)
W/Polyvinylpyrrolidone See: Salpix, Viol (Ortho)

ACETRIZOIC ACID, USP XV NNR 1951 3 Acetylaminio 2,4,6 Iriodobenzoic acid
See: Acetrizoate Sod (Mallinckrodt)

ACETRYPTIN HYDROCHLORIDE, (Warner Lambert) 5 acetyltryptamine hydrochloride
Use: Antihypertensive agent

ACET-THIOGIN SODIUM. See: Theophylline Sod Acetate (Various Mfr)

ACETYCOL (Warner Chilcott) Catcheine salt cyclated 0.25 mg, p aminobenzoic acid 162 mg aspirin 325 mg, thiamine 5 mg, ascorbic acid 20 mg, niacin 15 mg /Tab Bot 100s 500s, 1000s
Use: (1 lid) Analgesic in arthritis, gout & neuritis

ACETYL ADALIN. See: Acetylcarbromal (Various Mfr)

p-ACETYLAMINOBENZALDEHYDE THIOSEMICARBAZONE. (Amithiozone, Antibacterium A Benzothiazon, Conleben, Myuzone, Neustob Tebathion, Thiomicid Thio

paramizone, Thiocetozone)
See: Panrone, Tab (Panray)
Teabozone, Tab (Consol Midland)
Tribione, Tab (Schenley)

Use: With streptomycin in the treatment of tuberculosis

ACETTLAMINO BENZENE. See: Acetanilid (Various Mfr)

2-ACETYLAMINO-3-NITROTHIAZOLE. See: Tritheton Tab (Ortho)

N-ACETYL-p-AMINOPHENOL. Acetaminophen

See: Amdil, Tab (Breon)
Anelix, Elix (Kirkman)
Apamide, Tab (Ames)
Iliadodyne, Liq (Lamman)
Fibro Gelic (First Texas)
Fibrolin Elix (Buffington)
Fendon Tab (APC)

Leitemp (Reid Labs)
Lyseca Syrup (Lloyd, D & W)
Metolid, Tab (Columbus)
Nacetyl (Allison)

Nabs, Tab (Narwich Pharmacal)
Pafuss (Sulfin & Case)
Rystal, Syr (Graft)

- Tempo, 5 gr. (Mead)
 Tussopop (Sulliff & Case)
 Tylenol, Elix & Tabs. (McNeil)
BUFFERED TABLETS.
 See Apamide-Ves, Tab. (Ames)
 W/Acetophenetidin, caffeine, Vit. B-1. See: Gesik, Tab. (Schilling)
 W/Azelophenelidin, phenylpropranolamine HCl, phenyltoloxamine dihydrogen citrate. See: Sinutab, Tab. (Warner-Chilcott)
 W/Acetylcarbromol. See: Apromol, Tab. (Ames).
 W/Adiphenin HCl, Methampyrone, phenyltoloxamine citrate. See: Anafac, Cap. (Pharmafac)
 W/A.P.C. See: Dalara Compound, Tab (Buffington)
 W/Ascorbic acid, salicylamide. See: Arthralgen, Tabs. (Whittier)
 W/Ascorbic acid, salicylamide, prednisone. See: Arthralgen PR, Tabs. (Whittier)
 W/Aspirin & caffeine. See: Acetylduo, Tab. (Chicago Pharm.). Trigesic, Tab. (Squibb).
 W/Aspirin, caffeine, aluminum hydroxide, calcium gluconate. See: Supac-B, Tab. (Misslon)
 W/Aspirin & caffeine with codeine. See: Trigesic w/Codeine, Tab. (Squibb).
 W/Aspirin, phenylephrine HCl, nospapine, caffeine, pyrilamine maleate, methapyrilene fumarate. See: Tussopop, Tab. (Sulliff & Case)
 W/Aspirin, trimethyl xanthine, chlorpheniramine maleate, phenylephrine HCl. See: Thorexlin Cold Capsules, (Gillelle Labs)
 W/Butaborbital, d-amphetamine sulfate, acetylsalicylic acid, salicylamide. See: Stuart Prolar, Tab. (Stuart)
 W/Butaborbital sod. See: Bulapap (Tollby-Naxon) Nacetyl, Tab. (Allison)
 W/Butaborbital Sod. & methamphetamine HCl. See: Algaxon, Tab. (McNeil)
 W/Chlorpheniramine maleate, Pot. alpha-phenoxethyl penicillin, phenylephrine HCl, phenylpropranolamine HCl, phenyltaxisine citrate. See: Syndecon, Tab., Sol. (Bristol)
 W/Chlorpheniramine maleate, dexamethorphan HBr, phenylephrine, ascorbic acid. See: Thorphan Cold Syr. (Bryant)
 W/Chlorpheniramine maleate, phenylpropranolamine HCl, nospapine, glyceryl guaiacolate, ammonium chloride. See: Mycotaxin, Syr. (Myers Carter)
 W/Chlorprophenpyridamine maleate, phenylephrine HCl. See: Novahistine Singlet, Tab. (Pitman-Moore)
 W/Chlorprophenpyridamine maleate, phenylephrine HCl, phenylpropranolamine HCl, ephedrine sulfate. See: Pyrels, Tab. (Bernhoff)
 W/Chlorzoxazone, codeine phosphate. See: Parafan w/Codeine, Tab. (McNeil)
 W/Catchicine, zoxazolamine. See: Triurate, Tab. (McNeil)
 W/Demerol. See: Demerol APAP (Brean)
 W/Dexamethorphan hydrobromide, chlorpheniramine maleate, pyrilamine maleate, phenylephrine HCl, caffeine, ascorbic acid. See: Tacol, Tabs. (Massengill)
 W/Diadol, mephesisin, homatropine methylbromide. See: Steri-Exe, Tab. (Buffington's)
 W/Diallylbarbituric acid, mephesisin, homatropine methylbromide. See: Stri-Exe, Tab (Buffington's)
 W/Dihydrocodeinone bitartrate, homatropine methylbromide, chlorpheniramine maleate, phenylephrine HCl. See: Hycamine Compound, Tabs (Endo)
 W/Dihydrocodeinone bitartrate, meperidine HCl. See: Demerol Compound, Tab. (Brean)
 W/Dihydrocodeinone bitartrate, phenylephrine HCl, pyrilamine maleate, pot. guaiacol sulfonate, sod. citrate, citric acid, chloroform. See: Cadigestic, Syr. (Central)
 W/Glyceryl guaiacolate. See: Ryslat, Syr (Grall)
 W/Guayanesin, phenylephrine HCl. See: Papanesin, Tab (Superior Pharm.)
 W/Hyoscin hydrobromide, butobarbital sodium. See: Pallacin, Tab. (Denob)
 W/Isometheptene mucate, dichlorophenazone. See: Midrin, Caps. (Carrick)
 W/d-Methorphan HBr, chlorpheniramine, phenylephrine. See: Romliar CF, Syr. (Hoffmann-La Roche)
 W/Methyliso-octenylamine mucate, salicylamide. See: Migrad Tab. (Crystal)
 W/Methyliso-octenylamine mucate, salicylamide, and caffeine. See: Nygram, Cap. (Dumas-Wilson)
 W/Neo-synephrine HCl, thenyldiamine HCl caffeine. See: Neo-Synephrine Compound Cold Tablets, (Winthrop Labs.)
 W/Neostigmine bromide, hyoscin hydrobromide. See: Darth, Tabs. (Denob)
 W/Noscapine alkaloid, phenylpropranolamine, glyceryl guaiacolate, chloroform. A.R. See: Coryx Suspension, liq (Marian)
 W/Phenindamine tartrate, sulfodimethazine caffeine. See: Madricidin, Cap. (Roche)
 W/Phenylephrine HCl, atropine sulfate, pheniramine maleate. See: TPC, Tab. (Tennessee Pharm.)
 W/Phenylephrine HCl, chlorpheniramine maleate, dexamethorphan HBr, ascorbic acid. See: Thorphan, Syr. (Bryant)

ACETYLANILINE

6

- W/Phenylephrine HCl, chlorpheniramine maleate nescapine, salicylamide Vit C See Nascaps Cap (Tablet Rack)
- W/Phenylephrine HCl chlorpheniramine maleate See Navahistine Siegfelt Tabs (Pitman-Moore)
- W/Phenylephrine HCl, isothipendyl HCl, dimethoxanate HCl, ammonium chloride sodium citrate chloroform See: Catheter Compound, Liq (Ayers)
- W/Phenylephrine, d methorphan HBr, pyrrolamine maleate, potassium guaicolosulfonate alcohol See Thormal Liq (Central)
- W/Phenylephrine HCl phenylpropanolamine hydrochloride dextromethorphan hydrobromide ascorbic acid chlorpheniramine maleate See Thorphan CF, Tabs (Bryant)
- W/Phenylpropanolamine ammonium chloride See: Zeston, Tab (Standex)
- W/Phenylpropanolamine glyceryl guaiacolate, nescapine See Coryz, Tab (Marion)
- W/Phenylpropanolamine HCl, phenyltoloxamine dihydrogen citrate, pot guaicolosulfonate See Conex Syrup (Lloyd, Dabney & Westerfield)
- W/Phenylpropanolamine HCl, pyrilamine maleate phenyltoloxamine citrate methopysilene HCl, phenylephrine HCl, chlorpheniramine maleate See Lychisil, Caps (Leyden Labs)
- W/Phenyltoloxamine dihydrogen citrate & phenylpropanolamine HCl See Conex Plus, Tab (Lloyd Dabney & Westerfield)
- W/Phenyltoloxamine dihydrogen citrate, salicylamide, di-amphetamine phosphate metoprine See: Acogestic, Cap (Strassenburgh)
- W/Phenyltoloxamine dihydrogen citrate, salicylamide Vit C, citrus bioflavonoids caffeine, sedemic amphetamine sulfate See: Euphenes, Tab (Lloyd Dabney & Westerfeld)
- W/Pipenzolate methylbromide, phenobarbital See: Pipal Pediatric Antipyretic (Stokeside)
- W/Salicylamide See: Pedagogic (Columbus)
- W/Salicylamide, acetylsalicylic acid, butabarbital d-amphetamine sulfate See: Stuart Prolar, Tab (Stuart)
- W/Salicylamide acetylsalicylic acid d-amphetamine sulfate, butobarbital, tyroxyamine sulfate, atropine sulfate hyoscyamine hydrobromide See: Stuart Prolar II Tab (Stuart)
- W/Salicylamide caffeine See: Matca Analgesic, Tabs (Matca)
- W/Salicylamide, caffeine, phenylephrine HCl, trisamine See: Confomal-CP, Caps (Physicians Products)
- W/Salicylamide chlorpheniramine maleate, phenylephrine HCl, phenylpropanolamine HCl caffeine ascorbic acid See: Trisamine Forte Cap (Barre)
- W/Salicylamide chlorpheniramine maleate, disopredrine hydrochloride See: Isogestic, Tabs (Arnar Stone)
- W/Salicylamide, codeine phosp, caffeine See Cylamide Tab (Chicago Pharmacol Co) Also with phenobarb
- W/Salicylamide, Dipyrone, Vit C See Dipralan Forte, Tab (Arnar Stone)
- W/Salicylamide, glyceryl guaiacolate, pyrilamine maleate phenylephrine HCl See: Guaiogestic 1sq (Columbus)
- W/Salicylamide, disopredrine HCl, chlorpheniramine maleate See Isogestic, Tab (Arnar Stone)
- W/Salicylamide, phenobarbital, caffeine See Cylamide w/Phenobarbital, Tab (Chicago Pharmacol Co)
- W/Salicylamide phenyltoloxamine dihydrogen citrate phenylephrine HCl See Absinal, Tab (Lemmon)
- W/Salicylamide, potassium guaicolosulfonate pyrilamine maleate ammonium chloride sodium citrate phenylephrine hydrochloride See Tussabar (Bruce)
- W/Sod salicylate phenylephrine glyceryl guaiacolate See Pediluss, Syr (Sutliff & Case)
- W/Terpin hydrate dextromethorphan, phenylephrine HCl, chlorpheniramine maleate, ascorbic acid See Syntussin Compound Cap (Fees Cameron)
- W/Tetracycline base phenyltoloxamine citrate See Tetrax Ap, Syr (Bristol)
- W/Thephoria madriban, caffeine See: Madradia Cap (Roche)
- W/Trimepazine tartrate, phenylpropanolamine HCl See Coplexen, Liq (Smith, Kline & French)

ACETYLANILINE, See Acetanilid (Various Mfr)

ACETYL-BETA-METHYLCHOLINE CHLORIDE, USP NNR 1947 See: Mecharyl Chloride Amp (Merck Sharp & Dohme)

ACETYL-BROMO-DIETHYLACETYL-CARBAMIDE, See: Acetylacetyl bromat (Various Mfr)

ACETYL-CAP-PHEN. (Smith Miller & Patch) Aspirin 3 gr., acetophenetidin 1 gr., caffeine 1/2 gr., Dovers Pow 3/16 gr / Cap Bot 100s, 1000s

Uses: (1 1 d) Analgesic.

ACETYL-CARB (Sherman) Aspirin & Sodium bicarb Tab 500s, 1000s

Uses: (1 2 q 3 h) Analgesic

* **ACETYLCARBROMAL**. (Chemo Puro, Meyer)
Acetyl adalin, acetyl-bromo-diethylacetyl-
carbamide. Pow. for mfg.
See: Abasin. (Winthrop Labs.)
Sedamyl. (Schenley).
Use: Sedative.

W/N-acetyl-p-aminophenol. See: Apromol,
Tab. (Ames)

W/Apomide. See: Apromol, Tab. (Ames).

W/Bromisovalum.
See: Dayted, Tab. (Toilby-Nason).
Sedayson, Tab. (Toilby-Nason).

W/d-Desoxyephedrine HCL.
See: Euphosed-S, Tab. (Schenley)

**W/Hyascyamine sulfate, atropine sulfate
& hyascine HBr**. See: Disperdon, Tab.
(Standard).

W/Mephenesin & resperline. See: Amril, Tab.
(Amfr-Grant)

ACETYLCOLINE CHLORIDE. (Various Mfr.).
Leader—Amp. (0.1 Gm., 0.2 Gm.) 1 cc-
12s.
Merck—Amp. (0.1 Gm.). Vial 10 Gm.
Use: S.C., I.M., I.V. Parasympathomimetic
agent & vasodilator. Paroxysmal lachry-
mation.

W/Saligenin. See: Acecoline, Amp. (Anglo-
French).

**ACETYLCOLINE-LIKE THERAPEUTIC
AGENTS**. See: Cholinergic agents.

* **ACETYLDIGITOXIN**, N.N.D. 1962. The
natural-occurring cardiac glycoside defined as
the alpha acetyl ester of digitoxin. See:
Acyfanid, Tab. (Sandoz)

ACETYLDUO. (Chicago Pharm) Acetyl-
paminophenol 2.5 gr., aspirin 3.5 gr., cof-
feine 0.5 gr./Tab. Bot. 100s, 1000s.
Use: (1 q. 3-4 h) Analgesic.

ACETYLIC COMPOUND, (Centrol) Salicyla-
mide 30 mg., acetophenetidin 30 mg.,
atropine sulfate 0.03 mg., aconite root 1.5
mg., camphor monobromated 8.0 mg., phen-
olphthalein 6.0 mg./Tab. Bot. 100s, 1000s
Use: Analgesic, antipyretic

ACETYLIN. See: Acetylsalicylic Acid. (Various
Mfr.).

ACETYLMETHIONINE. (Chemo Puro) Methio-
nine 78%. Pow. for mfg.
Use: Nutrient.

ACETYLMETHIONINE CALCIUM. (Chemo
Puro) Methionine 71%, calcium 9.5%. Pow.
for mfg.
Use: Nutrient.

ACETYLMIDICEL. See: Midicel Acetyl Susp.
(Parke, Davis)

ACETYLPHENYLISATIN. See: Diacetylhy-
droxyphenylisatin.

ACETYLRÉSÖRCINOL. See: Resorcinol Mono-
acetate. (Various Mfr.).

* Generic name recognized by N.F., U.S.P. or A.M.A.

ACETYLSAL. See: Acetylsalicylic Acid.

ACETYLSALICYLATE ALUMINUM. See: Alu-
minem Acetylsalicylate.

* **ACETYLSALICYLIC ACID**, U.S.P., A.D.R.,
N.N.R. 1947; Capsules, N.F.; Tablets, U.S.P.
(Various Mfr.) Acidum acetylsalicylicum, 2
acetoxybenzoic acid, Acetillum Acidulum,
Acetophen, Acetal, Acetalar, Acetalaric
Acid, Acetosalin, Aceticyl, Acetylin, Acetyl-
sol, Acylpyrin, Aspro, Helicon, Rhodine,
Solacelin, Solcalogen, Saletin.
See: Aspirin Preps. (Various Mfr.).
A.S.A., Tab. (Lilly)
Empirin (Burroughs Wellcome).
Fentagesic (Kremers-Urban)

Salts.

See: Aluminem Acetylsalicylate.

Calcium Acetylsalicylate.

Use: Analgesic & antipyretic.

W/Acetophenetidin & caffeine. See: A.P.C.
preparations. (Various Mfr.).

W/Mixtures. See: Aspirin preparations. (Vari-
ous Mfr.).

**ACETYLSALICYLIC ACID, ACETOPHENETI-
DIN AND CAFFEINE**, N.F. See: A.P.C.,
Prep. (Various Mfr.).

ACETYLSALICYLIC ACID ENTERIC COATED,
See: Asteric, Tab. (Brewer)

ACETYLSTROBELINE. See: Asmotone, Cap.
(National Research).

* **ACETYLSULFAMETHOXYPYRIDAZINE**,
N.N.D. 1962 3-(N-Acetylsulfanilamide)-6-
methoxypyridazine. See: Kynex Acetyl
Pediatric Suspension (Lederle)

N¹-ACETYLSULFANILAMIDE, (Albucid, p-
Aminobenzenesulfonacetamide; Sulfacet)
Sulfacetamide, N-Sulfanilylacetamide) See:
Sulamyd. (Schering).

Use: Sulfonamide therapy.

* **ACETYLSULFISOXAZOLE**, U.S.P., N.N.D.
1962. N¹-acetyl-3,4-dimethyl-5-sulfanilami-
doisoxazole.

See: Ganthrin Acetyl. (Hoffmann-La Roche)
Lip-Ganthrin. (Roche).

ACETYLTANNIC ACID, N.N.R. 1941.

ACHEASE. (Philadelphia Ampoule) Phenyl-
toloxamine dihydrogen citrate 44 mg,
acetophenetidin 150 mg., salicylamide 150
mg., caffeine 32 mg./Tab. Bot. 100s, 1000s.
Use: Tension headache.

ACHLORHYDRIA DETERMINATION. See:
Diagnex.

ACHLORHYDRIA THERAPY.

See: Acidal. (Various Mfr.)

Acidol-Pepsin, Cap. (Winthrop Labs.)

Acid-V, Cap. (Merck Sharp & Dohme)

Aclor, Cap. (Cole Chem. Co.)

Chlor-Ethamine, Tab. (Pitman-Moore)

Glyco-HCl, Tab. (Burnham)

- Glutamic Acid HCl
Muitipsin, Tab (Norgine)
Normacid Tab (Stuart)
- ACHROCIDIN** (Lederle) Tabs. Achromycin HCl 125 mg, phenacetin 120 mg, caffeine 30 mg, salicylamide 150 mg, chlorothen citrate 25 mg / Tab Bot 24s
- Byr. Achromycin HCl 125 mg, phenacetin 120 mg, salicylamide 150 mg, Vit C 25 mg, pyrilamine maleate 15 mg, methyl paraben 4 mg, propylparaben 1 mg / 5 cc Syr Bot 4 fl oz
- Use Antibiotic antihistamine, analgesic therapy
- * **ACHROMYCIN, USP** NND 1962 (Lederle) Tetracycline HCl
Cap. (50 mg & 100 mg) Bot 25s & 100s (250 mg) Bot 16s & 100s
- Ear Solution (50 mg vial and a 10 cc vial of 5% benzocaine in propylene glycol)
Intravenous (100, 250 & 500 mg) Vial
Intramuscular (100 & 250 mg) Vial
Oral Susp: (250 mg / 5 cc) Bot 15 Gm, dry pow 1 oz
Oint, (3%) Tube ½ & 1 oz
Ophth Oil Susp: 1%, Bot w/dropper 4 cc
Ophth, Oint (1%) Tube ½ oz
Ophth Sterilized Vial (25 mg) w/drop per
- Pediatric Drops, Liquid (100 mg / cc) Bot w/dropper 10 cc
- Pharyngets (15 mg) Pkg 10s
- Burgical Powders: Vial (200 mg / Gm) 5 Gm
- Syr. (125 mg / 5 cc) Bot 2 & 16 oz
Tab: (50 & 100 mg) Bot 25s & 100s; (250 mg) Bot 50s 250s & 1000s
Tab Soluble, (50 mg) Tube 40s, 100s
Troches (15 mg) Pkg 25s, 250s
- W/Hydrocortisone Oint (Lederle) Tetracycline HCl 3%, Hydrocortisone 2% Tube 5 Gm
- Use: Pyogenic Infections of the skin
- W/Hydrocortisone (Lederle) Ophth Oint Tetracycline HCl 1%, Hydrocortisone 1.5% Tube ½ oz
- Use: Eye Infections and skin Infections
- W/Phenacetin, caffeine, salicylamide, chlorothen citrate See: Achrocidin, Syr, Tab (Lederle)
- ACHROMYCIN NASAL SUSPENSION** (Lederle) Tetracycline HCl 375 mg, hydrocortisone acetate 2 mg, phenylephrine HCl 0.125% / cc Bot squeeze 15 cc
- Use: Upper respiratory infections
- ACHROMYCIN BYRUP.** (Lederle) Tetracycline HCl 125 mg, methylparaben 0.08%, propylparaben 0.02%, alcohol 1% / 5 cc Bot 2 oz, 1 pt
- Use Antibiotic therapy
- ACHROMYCIN-V.** (Lederle) Achromycin 250 mg buffered w/citric acid & Sod citrate / Cap Bot 16s, 100s Pink Caps; Achromycin 100 mg w/citric acid & Sod citrate / Cap Bot 25s, 100s Byr. Achromycin 125 mg buffered w/citric acid & Sod citrate / 5 cc Bot 2 oz 1 pt Pediatric Drops: Achromycin 100 mg w/citric acid & Sod citrate / cc Dropper Bot 10 cc
- Use: Treatment of infections caused by both gram negative and gram positive bacteria
- ACNROSTATIN-V.** (Lederle) Tetracycline citric acid & Sod citrate buffered 250 mg, nystatin 250M u / Cap Bot 16s, 100s Oral Susp tetracycline citric acid & Sod citrate but 125 mg & nystatin 125M u / 5 cc Bot 60 cc
- Use Blood spectrum antibiotic w/antifungal therapy
- ACIBAN.** (Lederle) Calcium caseinate 0.4 Gm calcium carb 0.1 Gm / Tab Bot 50s 250s 1000s
- Use: (1 2 1 2 d) Antacid
- ACICHEK.** (Valentine) Dihydroxy aluminum aminoacetate 250 mg, Sod carboxymethyl cellulose 150 mg, glycine 30 mg / Tab Bot 100s, 1000s
- Use Gastric hyperacidity in peptic ulcer
- ACID ACRIFLAVINE.** See: Acriflavine Hydrochloride (Various Mfr)
- ACID CARBOLIC, NUTGALL, AND ATROPINE COMPOUND.** (Upjohn) Phenol 6 gr, nutgall (equiv) 35 gr, atropine ¼ gr, benzocaine 10% / 10 gr / oz Oint Tube 1 oz Jar 1 lb
- Use: Rectally, local anesthetic, analgesic & antipruritic
- ACID CITRATE DEXTROSE SOLUTION.** See: ACD Sol
- ACID CLOAK CREME.** (Dome) pH 5.0 Creme Tube 1 oz, Jar 4 oz, 1 lb
- Use: Restore skin to normal pH
- ACI-DERMA** (Princer) An aqueous emulsion of glyceryl monostearate, cetyl alcohol, sodium lauryl sulfate, propyl and benzyl p hydroxy benzoates and propylene glycol buffered to pH 2.3 Bot 4 oz
- Use: Restore the acid mantle of the skin
- ACID HISTAMINE PHOSPHATE.** See: Histamine Phosphate (Various Mfr)
- ACIDIRON** (Walker Labs) Ferrus sulfate exsic 3 gr, hydrochloric acid diluted 125 mg / Tab Bot 100s
- Use: Anemias
- ACID MANTLE CREME.** (Dome) Aluminum acetate in specially prep water soluble
- * Generic name recognized by N.F., U.S.P. or A.M.A.

- hydrophilic cream of pH 4.2 Tube 1 oz. Jar 4 oz., 1 lb.
 Use: Skin prophylactic agent in restoring normal skin acidity.
- ACID MANTLE LOTION.** (Dane) Alaminem acetate lotion. pH 4.5. Bot. plastic, 1 oz. Bot. 4 oz. squeeze, 1 pt., ½ and 1 gal.
 Use: Dermatitis.
- ACIDOL.** See: Betaine Hydrochloride. (Various Mfr.).
- ACIDOLATE.** (White) (Hypoallergenic liquid skin cleanser.) Sulfated vegetable oils, liquid petrolatum & water. Bot. 8 oz., & gal.
 Use: Cleanses skin surfaces without irritation.
- ACIDOL-PEPSIN.** (Winthrop Labs.) Betaine HCl 4 gr., pepsin 3.5 gr./Cap. Bot. 50s.
 Use: (1-2 t.i.d.) Achlorhydria.
- ACIDORIDE.** (Abbott) d-Glutamic acid HCl Cap. (5 gr.), Bot. 100s, 500s, 1000s.
 Use: Achlorhydria.
- ACIDOTHYN.** (Flint-Eaton) Glutamic acid HCl. (Each cap. is equivalent to 10 min. of Dilute Hydrochloric Acid U.S.P.) Bot. 100s, 500s, 1000s.
 Use: (1-2) Achlorhydria.
- ACID TRYPAFLAVINE.** See: Acriflavine Hydrochloride. (Various Mfr.).
- ACIDULIN.** (Lilly) Glutamic acid HCl. (Equivalent to 0.62 cc. [10 min.] of diluted hydrochloric acid) Pulvules. Bot. 100s, 500s, 1000s.
 Use: (1-2) Achlorhydria.
- ACIDUSOL.** (Texas) Oxyquinoline sulfate, lactic acid, zinc sulfocarbolate, citric acid, dextrose, buffered solution. Bot. 8 oz.
 Use: Vaginal cleanser.
- ACIGLUMIN.** See: Glutamic Acid Hydrochloride. (Various Mfr.).
- ACI-JEL.** (Ortho) Acetic acid 0.92%, lincioleic acid 0.70%, oxyquinoline sulfate 0.025%, boric acid 3.0%, and glycerine 5% in a vegetable gum base. Jelly. Tube with applicator, 3 oz.
 Use: Vaginal antiseptic.
- * **ACINITRAZOLE.** 2-acetamide-5-nitrothiazole. Trichard See: Trithean (Ortho)
- ACITAMIN.** (Massengill) Glutamic acid HCl 0.324 Gm., thiamine HCl 1 mg., riboflavin 1 mg., nicotinamide 5 mg./cap. Bot. 100s, 500s, 1000s.
 Use: (1-2) Achlorhydria.
- A-C-K.** (G. F. Harvey) Aspirin 5 gr., menadione 1/200 gr., ascorbic acid ½ gr./Tab. Bot. 100s, 1000s
 Use: Salicylate therapy.
- A-C-K BUFFERED.** (G. F. Harvey) Aspirin 333 mg., ascorbic acid 33.3 mg., menadione
- 0.33 mg., calcium carbonate 60 mg./tab. Bot. 100s, 1000s
 Use: (1-2 q. 3 h.) Analgesic.
- ACKIE.** Superior Tab. Bot. 1000s
- ACKLETIN.** (Associated Concentrates) Iecithin, soya 8 gr. with vitamins/Cap. Bot. 100s, 250s.
 Use: (1 to 2 cap. daily) Fat metabolism.
- ACLOR.** (Cale) Glutamic acid HCl Cap. (5 gr.) Bot. 100s
 Use: (1-2) Achlorhydria.
- A.C.N.** (Person & Covey) Vitamin A palmitate 25M u., ascorbic acid 250 mg., niacinamide 25 mg., hesperidin 50 mg./Tab. Bot. 100s, 1000s.
 Use: Acne vulgaris.
- ACNAVEEN.** (Aveena Corp.) Colloidal oatmeal, kaolin, sulfur 3%, and hexachloraphen 0.5%. Lot, Paw, Pkg. 4 oz.
 Use: Topically, cleansing agent in the treatment of acne
- ACNE CORT DOME.** (Dane) Hydrocortisone ¼%, sulfur 4%, 2,2'-thiobis (4,6-dichlorophenol) 1%, resorcinol monoacetate 3%. Cream, Tube 1 oz. Lotion, Squeeze Bot. 1 oz.
 Use: Treatment of acne.
- ACNE-DERM.** (Lannett) 5% Dispersible sulfur, 3% zinc sulfate. Bot. 2 oz.
 Use: Skin conditions.
- ACNE-DOME.** (Dane) Regulars sulfur 4% (bithionol), 2,2'-thiobis (4,6-dichlorophenol) 1%, resorcinol monoacetate 3%. Double strength. sulfur 8%, 2,2'-thiobis (4,6-dichlorophenol) 1%, resorcinol monoacetate 3%. Creme 1 oz. Lotion Bot. 2 oz., 1 pt.
 Use: Treatment of acne.
- ACNE-DOME CLEANSER.** (Dane) Sulfur 2%, bithionol 1%, salicylic acid 2% in a colloidal Soy Bean complex. Medicated w/sponge 4 oz.
 Use: Medicated skin cleanser.
- ACNE MIXED "U.B.A."** (Lilly) Undenatured antigens of bacillus acne & staphylococci standardized to 0.05 mg. nitrogen/cc. Vial 20 cc.
 Use: Pustular acne.
- ACNE STROBACTERINE MIXED VACCINE.** (Merck Sharp & Dahme) Acne bacillus 800 million, Staphylococcus aureus 2000 million, Staphylococcus albus 2000 million/cc. Vial 5 cc.
 Use: Certain types of acne.
- ACNESTAT.** (Dane) Hydrocortisone free alcohol 35 mg., tyrothricin 3 mg., hexachlorophene 0.25%, colloidal sulfur 5%, resorcinol monoacetate 2%, and isopropyl 1D%/15 Gm. Tube 15 Gm., 1 oz.
 Use: Treatment of acne.

ACNE VACCINES

10

ACNE VACCINES. (Sherman) *Acno bacillus* 100 million *Colon bacillus* 300 million *Staphylococcus albus* 1000 million *Staphylococcus aureus* 1000 million/cc Vial 12.5 cc, 20 cc

Use SC with increasing dosage acute & chronic acne infections

ACNOMEL. (Smith, Kline & French) Resorcinol 2%, sulfur 8%, hexachlorophene 0.25% w/bentonite propylene glycol alcohol & titanium dioxide Cream Tube 1½ oz Cake. Sulfur 4% resorcinol 1% hexachlorophene 0.25% 1 oz containers

Use Acne therapy

ACNOTABS (Parke) Pancreatin bile salts pepsin Vit A C/Tab Bot 72s

Use Treatment of acne

ACOBRYONNA (Lemmon) Aconite pow 1.33 mg, byronia pow 1.33 mg, belladonna pow ext 0.16 mg /Tab Bot 100s

Use (2 q 3 h) Antipyretic & diaphoretic

ACOBRYONNA COMP. (Lemmon) Aconite pow 4.66 mg, belladonna pow ext 0.24 mg, byronia pow 1.33 mg, mercury iodide 1/100 gr, Sod salicylate 1 gr /Tab Bot 100s

Use (1 3 q 4 h) Follicular tonsillitis

ACOGESIC (Strosenburgh) Phenyltoloxamine dihydrogen citrate 25 mg, ocelyl p-aminophenol 300 mg, salicylamide 200 mg, di-
Amphetamine phosphate 2 mg, metoprine 0.5 mg /Cap Bot 100s

Use (1 2 q 4 h) Analgesic

ACOLAC BACTERIAL VACCINE. (U S Std Products) *Corynebacterium acnes* 1000 million, *Staphylococcus aureus* 500 million *Staphylococcus albus* 500 million/cc Vial 10 cc, 30 cc

Use: Acute or chronic infections caused by acne

ACON, NND 1962 (Endo) Water soluble vitamin A Cap (25 000 & 50 000 units/Cap) Bot 100s 1000s Soln (25 000 units/cc) Dropper Bot 30 cc

Use: Vitamin A therapy

ACONITE POWDER W/Belladonna pow ext, & byronia pow See: Acobryonna Tab.

(Lemmon)

W/Atropine sulf aminophylline, veratrum V See: Aconite Tab (Mallard)

ACR-ALLANTOMIDE (National Drug) Amino acidine HCl 1.500 sulfanilamide 10% allantoin 2%, in a water miscible base Tube 1 oz, Jar 1 lb

Use: Topically bacteriostatic

ACRIFLAVINE, NF X ADR NNR 1952 (Euflavine, Genacrine Neutroflavine) Acriflavine neutral, a mixture of 2,8-diamino-10-methylacridinium chloride & 2,8-diamino-

acrididine (Lilly) Tab 1½ gr Bot 100s, 1000s

Use External antiseptic

ACRIFLAVINE, BRILLIANT-GREEN, METHYLOSANILINE CHLORIDE MIXTURE, NNR 1953 See: Dymilal, Pow (McNeil)

ACRIFLAVINE HYDROCHLORIDE, NF X, ADR, NNR 1950 (Various Mfr) Hydrochloride form of acriflavine, NF Acid acriflavine, acid trypanflavine, flavine, trypanflavine

National Amine—Pow, Bot (1, 5, 10, 25 & 50 Gm) Tab (1½ gr) Bot 50s & 100s

Use: External bacteriostatic

ACRISAN EAR DROPS. (Recsei) Aminoacridine propionate 0.35% & Tris Histin 1% in propylene glycol Ear drops, ½ oz.

Use: Anti infective

ACRISAN THROAT SPRAY. (Recsei) Aminoacridine ascorbate 0.2% w/benzyl alcohol 0.5% Bot 4 oz

Use: Anti infective

ACRISORCIN 9-aminocridinylum 4-hexylresorcinolate See: Aksinol (Schering)

ACROSURGIC STERILE TOPICAL SPRAY POWDER. (American Cyanamid) Sterile tetracycline HCl 710 mg in powder form with chlorinated fluorinated hydrocarbons as propellant Aerosol 3 oz

Use Topical antibiotic in the prophylaxis and treatment of cutaneous infections

ACROYD (Parke Davis) Oxymetholone 10 mg /Tab Bot 30s

Use Anabolic

ACRYLIC CATIONIC RESIN. See Corba Resin (Lilly)

Vacid (Fester)

A C T. (Tilden) Pot bromide 11 mg, ammonium bromide 11 mg, Sod bromide 11 mg, aspirin 0.33 Gm, caffeine 33 mg /Tab Bot 100s, 1000s

Use (1 2 as needed) Analgesic w/sedative

ACTAMER. (Monsanto Chem) See: Bithional, USP

ACTASAL PEDIATRIC DROPS (Purdue Frederick) Choline salicylate 104.5 mg /O 6 ml Bot W/calibrated dropper—60 cc

Use Antipyretic, anti-inflammatory and analgesic for infants and children

ACTASE, NND 1962 (Ortho) Fibrinolytic (human) 30 000 u/vial with water 10 cc. Use (Dissolve in 10 cc of water, dilute to 250 cc for IV) Intravascular blood clots

ACTH, NND 1962 (Adrenocorticotrophic hormone Adrenocorticotrophin) See Corticotropin USP

National Drug—20 units/ml vial 2 ml, 10 ml

Parke Davis—25 units/vial 40 units/vial

- Sig Labs.—40 units/ml., vial 5 ml.; 80 units/ml., vial 5 ml.
- Testogon—40 units/ml., vial 5 ml.
- ACTH.** (National Drug) Corticot solution of ACTH. Inj. 20 U.S.P. units/cc.) 2 cc. & 10 cc. I.M. or I.V. drip.
- ACTN GEL, PURIFIED.**
- Hort Labs.—40 units/ml., vial 5 ml.; 80 units/ml., vial 5 ml.
- Intro Prods.—40 units/ml., dosette 5 ml.; 80 units/ml., dosette 5 ml.
- King Pharm.—40 units/ml., vial 1 ml.
- National Drug—40 units/ml., vial 1 ml., 5 ml.; 80 units/ml., vial 5 ml.
- Testogon—40 units/ml., vial 5 ml.; 80 units/ml., vial 5 ml.
- Vitarline—40 units/ml., vial 5 ml.
- Wesley Pharm.—40 units/ml., vial 5 ml.; 80 units/ml., vial 5 ml.
- ACTN (PITUITARY ADRENOCORTICOTROPIC HORMONE).** (Parke, Davis) Vial 25 units, 40 units (one unit is equivalent to 1 mg. International standard).
- ACTHAR, N.N.D. 1962.** (Armour) Corticotropin, U.S.P. Vial 10, 25, 40 mg. (lyophilized)
- ACTHAR GEL.** (Armour & National Drug) Corticotropin, U.S.P. Disp. Cart. (20 mg./cc.) 1 cc., Vial 5 cc. Disp. Cort. (40 mg./cc.) 1 cc., Vial 5 cc.
- ACTHAR GEL HIGHLY PURIFIED.** (Armour) H. P. acthar gel, a new, highly purified, repository ACTH preparation that may be given subcutaneously or intramuscularly. 20, 40 & 80 u. per cc., 5 cc. multiple dose vials. Supplied in a sterile 1 cc. B-d cartridge with B-2 disposable cartridge syringe, in potencies of 40 Armour u. per cc. See: Corticotropin, U.S.P.
- Use: Stimulant of adrenal cortex, anti-inflammatory, antiallergic & useful in collagen diseases.
- ACTINAR SOLUTION, N.N.D. 1962.** (Armour) Corticotropin, 10 u./cc. Amp. 1 cc., Vial 10 cc.
- Use: I.V., I.M., S.C., corticotropin therapy.
- ACTICORT INJ.** (Testogon) Repository corticotropin 40 U.S.P. u., Vit. B-12 1000 mcg. in 1% gelatin with 0.5% phenol as preservative/cc. Vial 5 cc.
- Use: Vitamin/hormone therapy.
- ACTIDIL, N.N.D. 1962.** (Burroughs Wellcome) Triprolidine HCl. Tab. (2.5 mg.) Bot. 100s. Syr. (1.25 mg./5 cc.) Bot. 1 pt.
- Use: Antihistamine.
- ACTIFED.** (Burroughs Wellcome) Triprolidine HCl 2.5 mg., pseudoephedrine HCl 60 mg./Tab. Bot. 100s, 1000s. Triprolidine HCl 1.25 mg., pseudoephedrine HCl 30 mg./5 cc. Syr. Bot. 1 pt.
- Use: Common cold.
- ACTIFED-C.** (Burroughs Wellcome) Codeine phosphate 10 mg., triprolidine HCl 2 mg., pseudoephedrine HCl 30 mg., glyceryl guaiacolate 100 mg./5 cc. Bot. pt.
- Use: Treatment of the common cold.
- ACTIFIELD.** (Burroughs Wellcome) Triprolidine HCl, 2.5 mg.; pseudoephedrine HCl, 60 mg./Tab. Bot. 100s. Triprolidine HCl, 1.25 mg.; pseudoephedrine HCl, 30 mg./5 cc. Syrup, Bot. 1 pt.
- Use: Cold, allergier.
- ACTILAMIDE SOLUTION.** (Broemmel) Sulfanilamide 0.40% and Chloraminol 0.13%. Nasal 1 oz.; Ophth., Drop, Bot. plastic 15 cc., 4 oz.; Gargle or Spray 1 pt.
- ACTINOMYCIN D.** Under study.
- Use: Treatment of certain malignant tumors.
- ACTINOSPECTACIN.** (Abbott, Upjohn) Antibiotic Under study.
- ACTIVATED CARBON-SILICON.** (Inorganic Bioelements Inc) Activated carbon 50%, active silicon 27.5%, oxides of aluminum 6%, mag. 3.5%, iron 1.5%. Pow. Bot. 3 oz.
- Use: Irritations of G.I. tract.
- ACTIVATED CHARCOAL CAPSULES.** (Requa) Activated vegetable carbon, N.F.; 4 gr./Cap. Bot. 30s
- Use: Digestive tract irritations & adjunct in the treatment of acne vulgaris.
- *ACTIVATED 7-DEHYDROCHOLESTEROL,** U.S.P., A.D.R. See: Vitamin D-3. (Various Mfr.)
- ACTIVATED ERGOSTEROL.** See: Calciferol, U.S.P.
- ACTIVOL** (Cerbisulphal) Liquid soap. Bot. 4 oz., 1 pt., 1 gal.
- Use: Surgical scrub-ups.
- ACTO-CORTIN GEL.** (Myers-Carter) Adrenal cortex ext. 40 & 80 mcg./cc. Vials 5 cc. Also w/Vial B-12 1000 mcg./cc. Vial 5 cc.
- ACTOL** (Massungill) Neomycin sulf. 65 mg., polymyxin B sulf. 5,000 u./5 cc. Sol. Bot. 1 pt.
- Use: Treatment of intestinal infections.
- ACTOPHIN.** (Biochemical) Vit. D 50,000 u., bile salts 32 mg./Cap. Bot. 50s, 100s, 500s.
- Use: Cholelithic with vitamin D supplement.
- ACTROPE.** (United Lab.) Corticotropin, U.S.P. Vials 25, 40 & 100 mg./cc.
- Use: Adrenocorticotrophic therapy.
- ACTYLATE TABLETS.** (Kinney) Ammonium salicylate 1½ gr., Pot. salicylate 1½ gr., strontium salicylate 1½ gr., p-aminobenzoic acid 4 gr., Vit. C ½ gr./Tab. Bot. 100s.
- Use: (2 q. 3 hr.) Rheumatic disorders.

* Generic name recognized by N.F., U.S.P. or A.M.A.

ACTYLATE PLUS E

12

ACTYLATE PLUS E (Kinney) Cortisone acetate 5 mg, amn salicylate $1\frac{1}{2}$ gr, pot salicylate $1\frac{1}{2}$ gr, strontium salicylate $1\frac{1}{2}$ gr, pot paminobenzoate 5 gr, vit C $\frac{1}{2}$ gr/tab Bot 50s

Use (1 2 1 1 d) Rheumatoid arthritis

ACUTEL (Ortho) Prothrombin time plasma Vial 0.5 ml

Use Diagnostic agent

ACUTRAN (Irwin Neisler) Amphetamine

Use Anorexic agent

ACUTUSS-AS (Rosane) Claphed anal HC

7.5 mg diphenylpyraline HCl 0.5 mg phenylephrine HCl 2.5 mg, glyceryl guaiacolate 25.0 mg acetaminophen 162.5 mg / Tab Bot 50s

Use Acute coughs

ACYLANID TABLETS (Sandaz) NND 1961

Acetyl digitoxin Tab (0.1 mg) pink, 100s & 1000s; (0.2 mg), 100s, 1000s

Use Digitalin therapy

ACYLYPRIN See Aspirin (Various Mfr)

A & D CAPSULES (Rowell) Vit A 5000 u

Vit D 300 u / Cap Bot 100s 1000s

Use (1 cap 1 d) Vitamins A & D therapy

A-D DROPS (Walker) Vit A 2000 u, Vit D

300 u / drop Bot w/dropper 15 cc, 60 cc

Use (m p) Vitamin A & D deficiencies

A & D OINTMENT (White) Vitamin A & D

In the ratios as found in cod liver oil Tube $1\frac{1}{2}$ oz, 4 oz, Jar 1 lb 5 lb

Use Promotes healing of tissue

A & D OINTMENT W/PREDNISOLONE

(White) Vitamins A and D plus 0.5% prednisolone Tube 10 gm, 25 gm

Use Thermal or chemical irritation, allergic dermatoses etc

A & D VITAMIN CONCENTRATE W/Vio-

lateral (Horton & Converse) Vit A 5000 u,

Vit D 1700 u / Tub Bot 100s, 250s, 500s, 1000s

Use Nutritional supplement

ADABEE & ADABEE-M (Robins) A 25,000

USP u, D 1000 USP u B 15 mg, B 2 10 mg, B-6 5 mg, nicotinamide 50 mg,

calcium pantothenate 10 mg, C 250 mg,

Adabee M Fe 15 mg, I 0.15 mg, Cu 1 mg,

Mn 1 mg, Mg 6 mg, Zn 15 mg, K 5 mg,

Ca 103 mg, P 80 mg added / Tab Bot 100s 500s

Use: Dietary supplement

ADAC (Adac Sales) liquid 3 oz

ADACAL (Salinas Pharm) Tabs 100s

ADALIN, NNR 1947 (Winthrop Labs)

Carbamol Tab (5 gr) Bot 25s

Use: (1 2 m p) Sedative & hypnotic

ADAMIN (Gold Leaf) Vit A 5000 u D 1000

u, B-1 1 mg, B-2 2 mg / Cap Bot 100s 1000s

Use: (1 daily) Dietary supplement

ADANON HYDROCHLORIDE, NND 1959.

(Winthrop Labs) Methadone HCl

Amp (5 mg/cc) 2 cc Box 10s, 25s, 100s

Elixir (5 mg / 5 cc) Bot 1 pt

Syr (1.65 mg / 5 cc) Bot 1 pt

Tab (2.5, 5.0, 7.5, 10 mg) Bot 100s, 500s

Vial (5 & 10 mg / cc) 20 cc. Box 1s.

Use Narcotic, analgesic

A D C. DROPS W/Vitamin B 6 (Parke, Davis)

Vit A 5000 u, O 1000 u, C 50 mg, B 6

1 mg / D 6 cc Dropper bot 15 cc, 50 cc

Use Vitamin deficiencies in infants and children

ADCAPS, (Blue Line) Vit A 5000 u, Vit D

500 u / Cap Bot 100s 1000s

Use (1 daily) Vitamin A & D

ADCETS (Walker) Vit A 5000 u, D 1000 u,

C 50 mg / Vitasphere Bot 50s, 250s

Use (1 daily) Dietary supplement

ADCOSULFAMINE SUSP. (Associated Drug)

Sulfadiazine 2.5 gr, sulfamerazine 2.5 gr,

sulfamethazine 2.5 gr, phenylephrine HCl

5 mg chlorpheniramine maleate 2 mg,

ammonium chloride 0.5 gr / 5 cc Bot pts

Use Cold Therapy

ADCO TRIAMINE LIQ. (Associated Drug)

Pyriminone maleate 12.5 mg, chlorphenira-

mine maleate 2.0 mg, phenylpropanol-

amine HCl 10.0 mg, caffeine 1/3 gr / 5 cc

Bot pts

Use Cold Therapy

ADEF, (Endocrine) Vit A 20,000 u, D 4000

u, E 10 u, F (Vitamin concentrate) 300

mg / 1 cc Bot w/dropper 30 cc, 4 oz

Use (1 or 2 cc daily) Fat & water metabo-

lism disorders

ADEF-C (Endocrine) Vit A 5000 u, D 1000

u, E 5 u, F (Vitamin concentrate) 100 mg,

C 50 mg / Cap Bot 100s

Use (1 to 2 daily) Mucous membrane A

skin allergies

ADEFLO (Upjohn) Vit A 5000 u, D 3 1000

u ascorbic acid 50 mg, pyridoxine hydro-

chloride 1 mg, fluoride (as sodium fluoride)

0.5 mg / O 6 cc.

Use Prevention of dental caries

ADEFLO PLUS, (Upjohn) Sodium fluoride

0.5 mg Vit A 12 mg, D 10 mCg, C 75

mg thiamine mononitrate 2 mg, Vit B-2

2 mg, nicotinamide 18 mg Vit B 6 1 mg,

calcium pantothenate 5 mg, Vit B 12 2

mCg / Tab Bot 100s

Use Prevention of dental caries

ADEFLO PRENATAL, (Upjohn) Vit A 6000

u, D 400 u, B 1 15 mg B 2 2.5 mg, C

100 mg, nicotinamide 20 mg, Vit B-6 10

mg, calcium pantothenate 10 mg, Vit B 12

2 mcg folic acid 0.4 mg, fluoride 1 mg,

- calcium 250 mg., iron 15 mg./Tab. Bot. 100s.
 Use: Dietary supplement.
- ADEMOL.** (Squibb) Flumethiazide, 6-Trifluoromethyl-7-sulfamyl-1,2,4-benzothiadiazine. Tabs. 500 mg. Bot. 60s
 Use: Oral diuretic.
- ADENINE SULFATE.** (Schwarz) Pow. Bot. 5 Gm.
 Use: Granulocytopenia.
- ADENO-B-12.** (Kirk) Adenosine 25 mg., vit. B-12 60 mcg./cc. Vial. 10 cc.
 Use: Treatment of tendinitis, bursitis, neuritis.
- ADENOLIN W/B-12 GEL.** (Lincoln) Adenosine-5-monophosphate 25 mg., Vit. B-12 150 mcg., & niacin 10 mg. in gelatin/cc. Vial 15 cc.
 Use: Neuritis, bursitis & related conditions.
- ADENOLIN FORTE.** (Lincoln) Adenosine 5-monophosphate 25 mg., vit. B-12 300 mcg., methionine 25 mg., niacin 10 mg., gelatin 17% /cc. Vial.
 Use: Treatment of arthritis.
- ADENOMINE.** (Morton) Vial, 10 cc.
- ADENOPLEX.** (U. S. Standard Products) Adenosine 5-monophosphate 25 mg., Vit. B-12 50 mcg., niacin 25 mg., B-1 10 mg./cc. Vial 10 cc.
 Use: I.M., treatment of collagen diseases.
- ADENOSINE W/Vit. B-12.** See: Adeno-B-12, Vial, (Kirk)
- ADENOSINE COMP. GEL.** (Vitamax) Adenosine-5-monophosphate sod. 25 mg. & sod. nicotinate 25 mg. & B-1 50 mg. w/gelatin/cc. Vial 10 cc.
 Use: I.M., bursitis, neuritis & others.
- ADENOSINE WITH B-12.** (Mallard) Adenosine-5-phosphoric acid 25 mg., Vit. B-12, crystalline 100 mcg., benzyl alcohol 2%/cc. Vial 10 cc.
- ADENOSINE DIPHOSPHATE.** (Schwarz) (ADP, Adenyldiphosphoric acid, Adenosine-diphosphoric acid) Pow. Bot. 2 Gm.
- ADENOSINEDIPHOSPHORIC ACID.** See: Adenosine Diphosphate. (Schwarz).
- ADENOSINE-5-MONOPHOSPHATE.** Adenylic acid. * Phosphate. See: Cardimane, Amp. (Endo)
 My B Den, Amp. (Blischoff)
 W/Nicotinic acid, Vit. B-12.
 See: Vastron, Amp. (Wampale Labs.)
 Adenalin-B-12 Gel (Lincoln)
 W/Vit. B-12.
 See: Adenosine w/B-12, Inj. (Mallard)
 Adenovite, Vial (Metropolitan Labs.)
 Cabacine, (New Mexico Pharm. Co)
 Cabaden, Amp. (Rand)
 Cyadesine, Inj (Superior Pharm)
 Cyanosine, Gel. (Vitamax)
- Denylex Gel (Lloyd, Dabney & Westerfield)
 Horvaden, Vial (G. F. Horvey)
 Mycaine, Vial (Myers-Carter)
 Thybadee, Vial. (Bell Pharmacol)
 TY-B, Amp. (Paul Maney)
 W/Vit. B-12, niacin, B-1
 See: Adenoplex, Vial (U.S. Standard Products)
- Adenosine Comp. Gel (Vitamax)
 B-Denose, Vial (B. F. Ascher)
 W/Vit. B-12, nicotinic acid, procaine HCl, and benzyl alcohol. See: Vibruken, Vial, (Philadelphia Ampoule Labs.)
 W/Vit. B-12, sod. nicotinate, procaine. See: Vastron, Amp. (Wampale)
- ADENOSINE-3-MONOPHOSPHATE AS THE SODIUM SALT.** See: Cardimane, Amp. (Endo).
- ADINOSINE TRIPHOSPHATE.** (Various Mfr.) (ATP, Adenytriphosphoric acid, Adenytriphosphoric acid) Pow. Bot. 1, 5, 100 Gm.
- ADENOVIRUS.** (Pitman-Moore) Adenoviruses propagated in monkey kidney tissue culture. Vial 5 cc.
 Use: Immunization against infections of adenoviruses.
- ADENOVIRUS VACCINE.** (Parke, Davis) Contains adenoviruses serotypes 3, 4, and 7 or adenoidal-pharyngeal-conjunctival viruses. Vials 5 cc.
 Use: Immunization (1 cc. I.M.) against colds.
- ADINOVITE.** (Metropolitan Labs.) Adenosine 5-monophosphate 25 mg., Vit. B-12 60 mcg./cc. Vial 10 cc., Box 10s.
 Use: Treatment of bursitis, pruritus, dermatitis, ulcerations accompanying varicose veins.
- ADENYLDIPHOSPHORIC ACID.** See: Adenosine Diphosphate. (Schwarz).
- ADENYLIC ACID.** See: Adenosine-5-monophosphate.
- ADENTIC ACID SALTS.** See: Ironyl, amp (Physiol. Chem.) Procaine adenyate.
- ADENYLIC ACID SOD. SALT.** See: Adenosine-5-monophosphate.
- ADENYLTROPHOSPHORIC ACID.** See: Adenosine Triphosphate.
- ADENYLTRIPHOSPHORIC ACID.** See: Adenosine Triphosphate.
- ADEPSINE OIL.** See: Petrolatum Liquid. (Various Mfr.).
- ADESTRIN OINTMENT.** (Brean) Fish liver oils rich in vitamins A & D. Oint. Tubes 1 oz., ½ lb. Jar 1 lb.
 Use: Emollient in the treatment of warts.
- ADETATE.** (Merck Sharp & Dohme) d-Adipetamine sulfate 30 mg., vit. B-1 30 mg., sod. glycerophosphate 240 mg., pot. glycerophosphate 60 mg./30 cc. Bot. 1 pt., 1 gal.
 Use: Additive stimulant.

ADEVAN (Walker) Vit A 10 000 u, D 1000 u, thiamin HCl 10 mg, riboflavin 10 mg pyridoxine HCl 2 mg, calcium pantothenate 10 mg, niacinamide 50 mg, ascorbic acid 100 mg alpha lipoic acid 10 mg /Cap Bot 100s 1000s

Use Multivitamin D deficiencies

ADEX (Squibb) Vit A 5000 u, D 800 u / Tab Bot 80s, 250s

Use: (1 daily) Vitamin A & D deficiencies

ADHES-ILLIN, ADR 1955 (Falaris) Penicillin G Sod 10 000 u carboxymethyl cellulose 23 1%/Gm of Oint Tube 1 5 Gm Use Typically penicillin therapy

ADIPEX, (Lemmon Pharmacal) Amobarbital 50 mg homotropine methylbromide 7 5 mg, methamphetamine HCl 10 mg /Tab Bot 100s 500s 1000s

Use: Treatment of depressive and anxiety states

ADIPHENINE (Chemo Puro) 2 Diethylamino ethyl diphenylacetate Pow 1 lb for mfg Use: See Transentine (Ciba)

ADIPHENINE AND PHENO (West ward) Adiphenine HCl 50 mg, phenobarbital 20 mg /Tab Bot 100s 1000s

Use: Treatment of peptic ulcer, gastritis, cholecystitis, biliary dyskinesia spastic colitis, gastro-intestinal spasm, certain gynecologic and urologic disorders

ADIPHENINE HYDROCHLORIDE (Biorganic) (2 Diethylaminoethyl diphenylacetate hydrochloride Pow 1 lb for mfg use

See: Senti, Tab (Vitorine)

Spasmi, Tab (Cole)
Transentine, Amp, Tab (Ciba)

Use: Antispasmodic.

W/Aluminum hydroxide kaolin & phenobarbital See: Africate Tab (Malford).

W/Dihydrochloric Acid, desoxycholic acid See Chalad Tab (Malford)

W/Hexamethylenamine Sod acid phosphate Myocymus est, trit cum est See: Hexased Tab (Malford)

W/Methampyrone n-acetyl p aminophenol, phenyltoloxamine citrate See: Antac, Caps (Pharmafac)

W/Phenob See: Spasmi, Tab (Cole)

W/Thiamin HCl pyridoxine HCl, phenyltoloxamine citrate See: Tapp, Caps (Pharmafac)

ADJUDITS (Fres-Cameron) Dextro-amphetamine phosphate 5 mg, Vit A 1500 u, B 1 0 34 mg, B 2 D 67 mg, B 6 0 33 mg, nicotinic acid 3 mg, ascorbic acid 20 mg, Va D-2 150 u /Trache Jar 36s

Use Anti-obesity

ADLEAF, [Gold Leaf] Vitamins A & D Amp (A 100,000 u, D 10,000 u/ec.) 6s, 25s,

* Generic name recognized by N F U S F or A.M.A.

100s Cap (A 5000 u, D 850 u) Bot. 100s, 1000s Tob (A 5000 u, D 500 u) Bot 100s, 1000s

Use Vitamin A & D therapy

AD-NIL (Medics) Levamphetamine Laeva 1-phenyl 3 amino propane sulfate 15 mg / Cap Bot 100s

Use Obesity

ADNOGEN-FL. (Pitman Moore) Adenovirus and influenza virus vaccines Vial 5 cc

Use Multiple antigen vaccine designed to immunize against both influenza and adenovirus infections

ADOLPH'S SALT SUBSTITUTE. (Adolph's) Mono potassium glutamate, glutamic acid, pot chloride, tricalcium phosphate Bot Shaker

ADONIDINE, (Delta) Bot 1 Gm

Use Cardiac stimulant

ADP See Adenosine Diphosphate (Schwarz)

A-D PERCOMORPN (Abbott) Vit A 5000 u, D 1870 u /Cap Bot 100s

Use (1 daily) Vitamin A & D therapy

A-D PERCOMORPN LIVER OIL WITH OTHER FISH LIVER OIL & VIOSTEROL. (Abbott) Vit A 60 000 u, D 8500 u /Gm Bot w/dropper 10 cc or 50 cc

Use Vitamin A & D therapy

ADRENAL CORTEX ACTIVE PRINCIPLE ON CNARCOAL. See: Cortisarbale, Tabs (Schieffelin)

ADRENAL CORTIX EXTRACTS, NND 1961 (Purified Extract of Adrenal Glands)

Influences the metabolism of proteins & carbohydrates Also promotes the formation of carbohydrates from proteins & fats Metabolism of electrolytes (K & Na ions) & water is affected Na International unit standard for the hormones of the adrenal cortex (0 1 mg 17 hydroxycorticosterone = 1 glycogen unit = 2 5 rat units = 50 dog units)

See: Acto Cortin Gel (Myers Carter)

Adrena-Cortin, Amp (Harrower)

ADRENAL CORTEX NORMONES

See Gluco Corticoids

Mineral Corticoids

* **ADRENAL CORTEX INJECTION, N F,**

NND 1958 (Aqueous) (Various Mfr)

Mixed natural adrenocortical hormones

See: Adrenalex, Vial (Darwin)

Cortile, Vial (Lincoln)

Eschalin, Vial (Parke Davis)

Recortex, Vial (Tastagar)

ADRENAL CORTEX IN OIL INJECTION.

USF

ADRENAL CORTEX PREPARATIONS.

See: Adrena Cortin, Aqueous, Amp, Vial (Harrower)

- Coritin**, vial. (Lincoln)
Corinalal, Cap. (Harrower).
Cortisarbate, Tab. (Schieffelin).
Escholin, Vial (Parke, Davis).
Glycortol, Pills. (Schieffelin).
Lipo-Adrenal Cortex, In oil, Vial. (Upjohn)
W/Anterior pituitary fresh gland, ovarian substance fresh gland, posterior pituitary. See: **Metrocrine**, Vial. (Metro).
W/Vitamin C. See: **Cortalex**, Tab. (Upjohn).
W/Vit. C, Vit. B 6 & Calcium glucoacetate. See: **Adrenalex-Oral**, Cap. (Darwin).
ADRENALIN, (Darwin) Cortisone acetate. Vial (200 mcg/cc.) 30 cc. Cap. Bot. 100s, 1000s.
 Use: I.V., I.M., adrenal insufficiency.
ADRENALIN-ORAL (Darwin) Adrenal cortex ext. 20 mcg, Vit. C 15 mg., Vit. B-6 5 mg., calcium gluconate 150 mg./Cap. Bot. 100s, 1000s.
 Use: (1-3 daily) Adrenal insufficiency.
ADRENALIN(E), A.D.R., N.N.R. 1949, See: **Epinephrine**. (Various Mfr.)
ADRENALIN, (Parke, Davis) Epinephrine. In Oil: Ampuls (1:500) 1 cc., 12s, 25s, 100s. Oint.: (1:1000) Tube ½ oz. Supp. (1:1000) Box 12s.
ADRENALIN BORATE, A.D.R. 1953.
ADRENALIN CHLORIDE SOLUTION, A.D.R. (Parke, Davis) Epinephrine HCl Principle of the medullary portion of suprarenal glands. Amp. (1:1000) 1 cc., 12s, 25s, 100s. Bot. (1:1000) 30 cc., (1:100) 5 cc. Steri-Vials and 1 cc. amp. boxes of 12, 25 & 100. Solution 1:100, in 5 cc. bottles, is a concentrated solution for arial inhalation only. Vial (1:1000) in 1 cc., 10s.
 Use: Sympathomimetic agent.
W/Cocaine HCl. See: **Codrenin**, liq. (Parke, Davis).
W/Ephedrine sulf. See: **Adrephine**. (Parke, Davis)
W/Procaine. See: **Adrocaine**, Sol. & Tab. (Parke, Davis).
ADRENALIN HYDROCHLORIDE. See: **Epinephrine Hydrochloride**. (Various Mfr.)
ADRENALONE. (No Mfr. currently lists) (Kephrene, Stryphnane). See: **Local anesthetic**.
ADRENAMINE. See: **Epinephrine**. (Various Mfr.)
ADRENATRATE. (Crookes Barnes) 1-epinephrine bitartrate 1% and 2%/Vials 7.5 cc. Use: Mydriosis
ADRENERGIC-BLOCKING AGENTS.
 See: **Aprisaline HCl**, Amp., Tab. (Ciba).
Arlidin (Arlington-Funk)
Benadaine, Amp. (Merck Sharp & Dahme)
Beexazoline HCl (Ciba).
Dibenzyline (Smith, Kline & French).
Ergot-Related Products.
Etamon Chloride, Vial (Parke, Davis).
Ganglionic Blocking Agents.
Hydergine, Amp., Tab. (Ciba).
Hydralazine HCl.
Ilidar (Eoche)
Octin (Knoff).
Phenoxylbenzamine HCl.
Phentolamine Methanesulfonate (Ciba).
Piperaxan. (Merck Sharp & Dahme)
Priscoline (Ciba).
Regitine, Amp., Tab. (Ciba).
Tolazoline HCl.
Yahimbine.
ADRENINE. See: **Epinephrine**. (Various Mfr.)
ADRENOCROME, N.N.R. See: **Carbozochrome Salicylate** N.N.D. 1961
ADRENOCROMAZONE. See: **Carbozochrome Salicylate**, N.N.D. 1961
ADRENOCROME MONOSEMICARBAZONE. See: **Carbozochrome Salicylate**.
ADRENOCORTICOTROPIC HORMONE, N.N.D. 1962. ACTH acts by stimulating the endogenous production of cortisone and the final result is similar whichever substance is administered. The nature of action of each is unknown. Both are necessary in the metabolism of carbohydrates, protein & fat & exert profound effects on neuromuscular metabolism. See: **Corticotropin**, U.S.P.
ADRENO-CORTIN. (Hawes) Purified ext. of adrenal glands, 2½ units/cc. Amp. 1 cc. Box 5s, 100s. Vial 10 cc. Box 1s, 10s, Vial 30 cc.
 Use: I.M., S.C., I.V., Addison's disease.
ADRENO-MIST. (Smith, Kline & French) Epinephrine HCl 1:100, chlorobutanol base in glycerin, Sod. bisulfite. Sol. Bot. ¼s, ½s, 1 oz.
 Use: Inhalation, paroxysms & symptoms of asthma.
ADRENOSEM SALICYLATE, N.N.D. 1962 (Massengill) Carbozochrome salicylate. Amp. (5 mg/cc.) 1 cc., 5s; Syr. (2.5 mg/cc) Rot. 4 oz.; Tab. (1 mg) sug. coated orange. Bot. 50s; Tab. (2.5 mg) yellow, Bot. 50s.
 Use: Orally, I.M., to increase capillary resistance and suppresses bleeding from a capillary bed.
ADREPHINE. (Parke, Davis) Adrenalin 1:10,000, ephedrine sulfate 2%, dist. water, q.s. Amp. 1 cc., 12s, 100s. Bot. 1 oz. Inhal. aet. Rot. w/dropper 1 oz., 16 oz.
 Use: Shrinking mucous membrane
ADRESTAT, N.N.D. 1962. (Organon) Carbozochrome salicylate 65 mg, hesperidin 50 mg., ascorbic acid 100 mg., sod. menadiol diphosphate 5 mg/Cap or Loz. Cap. Box 30s. Loz. Box 20s, Amp 1 cc.

Cortilin, vial. (Lincoln)
 Cortinorol, Cap. (Harrower).
 Cortisarbole, Tab. (Schieffelin).
 Eschafin, Vial (Parke, Davis).
 Glycortal, Pills. (Schieffelin).
 Lipo-Adrenol Cortex, in oil, Vial. (Upjohn)
 W/Anterior pituitary fresh gland, ovarian
 substance fresh gland, posterior pituitary.
 See: Metracrine, Vial. (Metro).
 W/Vitamin C. See: Cortalex, Tab. (Upjohn).
 W/Vit. C, Vit. B-6 & Calcium gluconate. See:
 Adrenalex-Oral, Cap. (Darwin).
ADRENALEX, (Darwin) Cortisone acetate.
 Vial (200 mcg./cc.) 30 cc. Cap. Bot. 100s,
 1000s.
 Use: I.V., I.M.; adrenal insufficiency.
ADRENALEX-ORAL, (Darwin) Adrenal cortex
 ext. 20 mcg., Vit. C 15 mg., Vit. B-6 5 mg.,
 calcium gluconate 150 mg./Cap. Bot. 100s,
 1000s.
 Use: (1-3 daily) Adrenal insufficiency.
ADRENALIN (E), A.D.R., N.N.R. 1949. See:
 Epinephrine. (Various Mfr.)
ADRENALIN, (Parke, Davis) Epinephrine.
 In Oil: Ampuls (1:500) 1 cc. 12s, 25s, 100s.
 Oint. (1:1000) Tube ½ oz. Supp. (1:1000)
 Box 12s.
ADRENALIN BORATE, A.D.R. 1953.
ADRENALIN CHLORIDE SOLUTION, A.D.R.
 (Parke, Davis) Epinephrine HCl. Principle of
 the medullary portion of suprarenal glands.
 Amp. (1:1000) 1 cc., 12s, 25s, 100s. Bot.
 (1:1000) 30 cc., (1:100) 5 cc. Steri-Vials and
 1 cc. amp. boxes of 12, 25 & 100. Solution
 1:100, in 5 cc. bottles, is a concentrated
 solution for oral inhalation only. Vial
 (1:1000) in 1 cc., 10s.
 Use: Sympathomimetic agent.
 W/Cocaine HCl. See: Codrenin, liq. (Parke,
 Davis).
 W/Ephedrine sulf. See: Adrephine. (Parke,
 Davis)
 W/Procaine. See: Adrocaine, Sol. & Tab.
 (Parke, Davis).
ADRENALIN HYDROCHLORIDE. See: Epinephrine
 Hydrochloride. (Various Mfr.)
ADRENALONE. (No Mfr. currently lists)
 (Kephrene, Stryphnone). See: Local anes-
 thetic.
ADRENAMINE. See: Epinephrine. (Various
 Mfr.)
ADRENATRATE. (Crookes Barnes) 1-epinephrine
 bitartrate 1% and 2%/Vials 7.5 cc.
 Use: Mydriasis
ADRENERGIC-BLOCKING AGENTS.
 See: Apresoline HCl, Amp. Tab. (Ciba).
 Arlidin (Arlington-Funk)
 Benodaine, Amp. (Merck Sharp & Dahme)
 Benzoxaline HCl (Ciba).
 Dibenzyline (Smith, Kline & French).

Ergot Related Products.

Etamon Chloride, Vial (Parke, Davis).
 Ganglionic Blocking Agents.
 Hydergine, Amp., Tab. (Ciba).
 Hydralazine HCl.
 Midar (Rache).
 Octin (Knoll).
 Phenoxymethamine HCl.
 Phenolamine Methanesulfonate (Ciba).
 Piperaxan. (Merck Sharp & Dahme)
 Prisolaline (Ciba).
 Regiline, Amp., Tab. (Ciba).
 Tolazoline HCl.
 Yohimbine.

ADRENINE. See: Epinephrine. (Various Mfr.).

ADRENOCROME, N.N.R. See: Carbazochrome Salicylate. N.N.D. 1961

ADRENOCROMAZONE. See: Carbazochrome Salicylate. N.N.D. 1961

ADRENOCROME MONOSEMICARBAZONE. See: Carbazochrome Salicylate.

ADRENOCORTICOTROPIC HORMONE, N.N.D. 1962. ACTH acts by stimulating the endogenous production of cortisone and the final result is similar whichever substance is administered. The nature of action of each is unknown. Both are necessary in the metabolism of carbohydrates, protein & fat & exert profound effects on neuromuscular metabolism. See: Corticotropin, U.S.P.

ADRENO-CORTIN, (Harrower) Purified ext. of adrenal glands, 2½ units/cc. Amp. 1 cc. Box 5s, 100s. Vial 10 cc. Box 1s, 10s; Vial 30 cc.

Use: I.M., S.C., I.V., Addison's disease.

ADRENO-MIST, (Smith, Kline & French) Epinephrine HCl 1:100, chlorobutanol base in glycerin, Sod. bisulfite. Sol. Bot. ½, ½, 1 oz.

Use: Inhalation, paroxysms & symptoms of asthma.

ADRENOSEM SALICYLATE, N.N.D. 1962 (Messingill) Carbazochrome salicylate. Amp. (5 mg./cc.) 1 cc., 5s; Syr. (2.5 mg./cc.) Bot. 4 oz.; Tab. (1 mg.) sug. coated orange, Bot. 50s; Tab. (2.5 mg.) yellow, Bot. 50s.

Use: Orally, I.M.; to increase capillary resistance and suppresses bleeding from a capillary bed.

ADREPHINE, (Parke, Davis) Adrenalin 1:10,000, ephedrine sulfate 2%, dist. water, q.s. Amp. 1 cc., 12s, 100s. Bot. 1 oz. Inhal. and Bot. w/dropper 1 oz., 16 oz.

Use: Shrinking mucous membrane

ADRESTAT, N.N.D. 1962. (Organon) Carbazochrome salicylate 65 mg., heparidin 50 mg., ascorbic acid 100 mg., sod. menadiol diphosphate 5 mg./Cap. or Loz. Cap. Box 30s, Loz. Box 20s, Amp. 1 cc.

- Ure Prevention and control of capillary bleed ng
- ADRESTAT F** (Organon) Adrenochrome semicarbazone 5 mg (as carboxochrome salicylate 130 mg)/cc Amp 1 cc w/disp syringe Box 5s Vial 5 cc
Ure IM, control at capillary bleeding
- ADRIN** (Merk Sharp & Dahme) Injection epinephrine HCl 1:1000 Bot 1 oz
Use SC, IM, IC Vasoconstrictor, antispasmodic
- ADRIZINE** (Key Corp) d Amphetamine sulfate 5 mg/Tab Bot 100s, 1000s
Ure CNS stimulant
- ADROCAINE SOLUTION & TABLETS** (Parke Davis) Adrenalin 1:100 000, procaine HCl 1½%, chlorbutane 0.5% Vial 50 cc Tab (hypo) one tab dissolved in 30 cc of water makes 0.1% procaine and adrenalin 1:100 000 solution
Ure Local anesthesia
- ADROYD** (Parke Davis) 2 methylidhydrotestosterone Oxymetholone 2.5 mg, 5 mg 10 mg/Tab Bot 30s
Use Restore or maintain proper metabolic balance in postsurgical patients
- ADSODIL** (Cawley) Tablets 75 1000s
- ADORMONE** (Herring) Complete set of reagents required
Use Pregnancy test
- *ADENOVIRUS VACCINE** (Parke Davis) Aqueous preparation of Adenoviruses types 3 4 and 7 in approximately equal proportions grown in monkey kidney tissue cultures and inactivated with formaldehyde See Adenovirus Vaccine (Types 3 4 and 7) (Parke, Davis)
- ADVICIN** (Schering) Diphenhydramethyl sulfate (Prantal) 2% undecylenic acid 5%, salicylic acid 3%/Cream Tube 50 Gms Powder Can 2 oz
Ure: Topical antilungal
- A-E** (Warner Chilcott) Methyltestosterone 2.5 mg, ethinylestradiol 0.01 mg/tab
Ure Anabolic agent
- AEA COMPOUND TABLETS** (Haack) Amnaphylline 2 gr, ephedrine HCl ½ gr, amobarbital ½ gr/tab Bot 100s 1000s
Ure: Sedative with diuretic
- AERODRIN** (Burroughs Wellcome) Polymyxin B sulfate 5000 u, neomycin sulf 5 mg, methaxamine HCl 5 mg/cc Sol Bot w/drapp 15 cc Spray Bot 15 cc
Ure: Relief & prevention of infection in allergic rhinitis
- AEROHALOR** (Abbott) Aerahalor Cartridges to introduce dry solid aerosols into the upper respiratory tract and lungs Individual boxes
* Generic name recognized by NF, USP or AMA
- AERONALOR CARTRIDGES PENICILLIN POTASSIUM** (Abbott) Penicillin G potassium 100,000 u/disposable sinter cartridge Vial 3 cartridges Box 4s
- AEROLONE COMPOUND** (Lilly) Clopate NCl 0.5 Gm, aludrine HCl 0.25 Gm, in propylene glycol/100 cc Sal Bot 1 fl oz
Use: Asthma, status asthmaticus & emphysema
- AEROPLAST, NND 1962** (Aeroplast Corp) Yeasote Ethyl acetate acetone solution of copolymers of hydroxyvinyl chloride-acetate and sebacic acid 9.3%, and modified maleic resin ester 3.1% Spray Pkg 3 oz, 6 oz, 12 oz
Use: A spray on* surgical dressing for wounds burns, and abrasions
- AEROSOL OY, ADR** See Dioctyl Sodium Sulfosuccinate, USP (Am Cyanamid)
- AEROSPORIN OTIC SOLUTION** (Burroughs Wellcome) Polymyxin B sulfate 0.1% (10,000 units/cc) in acidified propylene glycol Bot w/drapp 10 cc
Ure Topically (3-4 drops tid) Infections at the ear
W/Bacitracin See Polysporin and Bacitracin Oint (Burroughs Wellcome)
- AEROSPORIN SULFATE, NND 1962** (Burroughs Wellcome) Polymyxin B sulfate 50 mg (500,000 units/Tab) Tab, foil wrapped Box 12s; Vial (20 cc), sterile pow (500,000 u) See: Polymyxin B sulfate (Pfizer)
Use Orally for bacillary dysentery, parenteral form for hospital use only
W/Bacitracin See Polysporin
W/Bacitracin & neomycin sulfate See Neosporin, oint & Ophth Oint (Burroughs Wellcome)
W/Bacitracin neomycin, hydrocortisone See Cortisporin
- AET** (Abbott) 2 Aminoethylisothiaurium bromide HBr Under study
Use Treatment of cancer
- AETHYLIS CHLORIDUM** See: Ethyl Chloride (Various Mfr)
- AFAXIN, NND 1961** (Winthrop Labs) Oleovitamin A Cap 10,000 u, Box 25s, 100s Cap 25,000, 50,000 u, Box 100s & Bot 1000s In oil, 65,000 u per Gm, Bot 30 cc
Use: Vitamin A therapy
- AFENIL, NNR 1951** (Knoll) Calcium chloride (w/ea) Amp (10%)
Use: IV (10 cc) Calcium chloride therapy
- AFF** (Invenex) Ferrus Fumarate 195 mg ascorbic acid 100 mg/Tab Bot 100s
Use: Treatment of anemias
- A-FIL** (Texas Pharm Co) Menthyl anthranil

- late 5%, titanium dioxide in vanishing cream. Tubes 2 oz. and Sticks
 Use: Sun screen
- APKO BALM.** (American Pharmaceutical) Phenylmercuric acetate 0.012%, phenol 1%, and benzocaine. Tube 1 oz.
 Use: Antiseptic & analgesic dressing.
- APKO-CARB.** (American Pharmaceutical) (Granular efferv. salts.) Sod. bicarb., Pot. bicarb., citric acid, tartaric acid, magnesium sulfate, Sod. phosphate, calcium lactate & Sod. chloride. Bot. 4, 8, 1 lb.
 Use: (2 dr. m.p.) Antacid.
- APKO-HIST.** (American Pharmaceutical) Pyrilamine maleate 25 mg. Tob. Bot. 50s.
 Use: (1 t.l.d.) Antihistaminic.
- APKO-HIST W/A.P.C.** (American Pharmaceutical) Pyrilamine maleate 12½ mg., aspirin 3½ gr., acetophenetidin 2½ gr., caffeine ½ gr./Tob. Bot. 50s.
 Use: (1 t.l.d.) Antihistaminic with analgesic & antipyretic.
- APKOL.** (American Pharmaceutical) Mineral oil, as light & heavy. Emulsion, plain, 1 pt., 1 qt.
 Use: Laxative.
- AFKOPIL.** (American Pharmaceutical) Belladonna ext. 1/12 gr., atropin ¼ gr., phenolphthalein ½ gr., ipecac 1/15 gr./pill. Bot. 100s, 1000s.
 Use: (1-2 h.s.) Laxative.
- AFKO-SAL.** (American Pharmaceutical) Salicylamide. Tob. (5 gr.) Bot. 36s.
 Use: (1-2 q. 3 h.) Analgesic.
- APKO-THRICIN.** (American Pharmaceutical) Tyrothricin 2 mg., benzocaine 5 mg./oz. Bot. 12s. Cream (0.5 mg./Gm.) Tube 1 oz. Jar 1 lb.
 Use: Antibiotic therapy.
- APRODEX.** (Layan Labs) Methyltestosterone 5 mg., yohimbine HCl 5 mg., nux vomica 5 mg./Cap. Bot. 100s.
 Use: Aphrodisiac.
- * **AGAR, U.S.P., N.N.B. 1949.** (Various Mfr.) Agar-agar, Bengal Gelatin, Ceylan.
 Use: Bulk laxative.
 W/Mineral oil.
 See Agaral, Emulsion. (Warner). Cremagal. (Cremagal Lab). Petrogalar. (Wyeth).
- AGEAPLEX.** (Foy Labs) Vials, 10 ml.
- AGENA.** (Bio-Pharmaceuticals) Estragen 10,000 u., Vit. A 100,000 u./oz. Oint. Tube w/vaginal applicator or refill 1½ oz. Jar.
 Use: Hormone and vitamin therapy
- A-GIC.** (A-Gic Corp.) An ointment made from the juice of aloe vera leaves. Tube 2, 4, 8, 16 oz.
 Use: Topically, burn therapy.
- * Generic name recognized by N.F., U.S.P. or A.M.A.
- AGNIN.** See: Lanalin Anhydrous. (Various Mfr.).
- AGNOLIN.** See: Lanalin Anhydrous. (Various Mfr.).
- AGOFOLLIN.** See: Estradiol. (Various Mfr.).
- AGORAL.** (Warner Chilcott) Phenolphthalein & mineral oil in an emulsion containing agar-gel, tragacanth, egg-albumen, acacia, glycerin and water. Bot. 6, 10, 16 fl. oz.
 Use: Laxative.
- AGORAL, PLAIN.** (Warner Chilcott) Emulsion of mineral oil and an agar-gel. Bot. 6, 12, 16 oz.
 Use: Laxative.
- AGUAPD.** (Chicago Pharm.) Oxyquinoline sulfate, boric acid, Sod. chloride, magnesium sulfate. Pow. Bot. 4½ oz.
 Use: Acid douche solution.
- AGURIN, N.N.B. 1928.** See: Theobromine Sod. Acetate. (Various Mfr.).
- A-N TABLETS.** (Boyle) Pyrilamine maleate. Bot. 16s, 40s, 100s, 1000s.
 Use: Antihistaminic.
- AHR-639.** (Robins) 1-[3,3-diphenyl-2-pyrrolidone]. Under study.
 Use: Antidote for barbiturate poisoning
- AIDANT.** (Noyes) Acetophenetidin 2 gr., aspirin 3 gr., caffeine ¼ gr., gelsemium tr. 2 min., methyl salicylate syr., q.s./Tob. Bot. 100s, 1000s.
 Use: Analgesic.
- AKALON-T-5.** (Strosenburgh) Methscopolamine 5 mg., Tuozole 20 mg./Cap., Bot. 50s
 Also available in double strength.
 Use: Peptic ulcer, hyperacidity.
- AKINETON, N.N.D. 1961.** (Knoll) Biperiden HCl. 1-bicycloheptenyl-1-phenyl-3-piperidinopropanol-1 HCl 2 mg./Teb. Bot. 100s, 1000s.
 Use: Anticholinergic and myospasmolytic for Parkinson's disease.
- AKINETON HCl, N.N.D. 1962.** (Knoll) Biperiden lactate 5 mg. in aqueous 1.4% sodium lactate solution/cc. Boxes, six 1 cc. vials.
 Use: Relief of Parkinson-like dystonic crises
- AKRINOL** (Schering) Acrirarin
 Use: Topical use tinea versicolor
- A.L.A.** (Woods) Quinine sulfate, Iron chloride & magnesium sulfate & citric acid Sol. Bot. 4 oz.
 Use: (isp.) Malarial shills & fever.
- ALACTA.** (Mead-Johnson) Protein 4.1%, fat 1.5%, carbohydrates 3.8%, minerals 0.9%. Can 1 & 4 lb.
 Use: Infant food.
- ALASTID.** (Norgine Lab) Homatropine methylbromide 1/24 gr., dried aluminum hydroxide 7½ gr., phenobarbital ½ gr., mag-

- nesium Iritacide 2½ gr/Tab Bot 100s
Use (I m p) Antispasmodic with antacid
- ALATAL** (L I Drug) Aluminum aspirin 3½ gr acetophenetidin 2 gr caffeine ⅓ gr isobutylalylbarbituric acid ¾ gr/Tab 100s
Use Sedative and analgesic
- AL-AY.** (Bowman) Phenylephrine HCl 5 mg pyrilamine maleate 10 mg acetylsalicylic acid 150 mg caffeine 15 mg, aminoacetec acid 150 mg/Tab Bot 100s
Use Nasal decongestant
- * **ALAZANINE TRICHLORPHATE** (Parke Davis) 3 ethyl 2 [3 (3 ethyl 2 benzothiazol inylidene)propenyl] benzothiazolium 2 4 5 trichlorophenoxide salts with two formula weights of 2,4,5(trichlorophenol)
Use Anthelmintic
- ALSACIDE**, NND 1962 (Blue Line) Benzyl benzoate 50% Lotion Bot 1 pt 1 gal
Use Scabies
- ALBA-DOME Cream & Lotion** (Dome) Cream contains 20% and lot contains either 5 or 10% monobenzone (monoben zyl ether of hydroquinone) Cream 2 oz jars Lotion 2 oz bottles
Use Inflammatory skin conditions
- ALBAMYCIN SODIUM**, NND 1962 (Up john) Navobiacin sod 250 mg/Cap Bot 16s, 100s
Use: Wide range antibiotic therapy
W/Tetracycline See Panalba (Upjohn)
- ALBAMYCIN MIX-O-VIAL** (Upjohn) In upper compartment, N N Dimethyl aceta mide 10% v/v in sterile water for injection; lower compartment navobiacin sodium (Al bamyacin) 500 mg; nicotinamide 175 mg Bot 25s
- ALBAMYCIN SYR**, NND 1962 Navobia cin col 125 mg/5 cc Bot 2 oz, 1 pt
- ALBA-PENICILLIN.** (Upjohn) Albamyacin sod 250 mg, penicillin G pot 250,000 u/Cap Bot 16s 100s
Use: Effective against gram positive and several important gram negative organisms
- ALBOLENE** (McKesson & Robbins) Petrolatum liquid
- AL-BORO-PHEN** (Smith Barney) Boric acid 27 Gm, carbolic acid 0.324 Gm, am monium alum 0.324 Gm, eucalyptus oil 0.13 Gm, methyl salicylate 0.13 Gm, men thal 0.065 Gm, thymal D 0.32 Gm./oz. Pow Jar 7 oz, 1 lb, 5 lb
Use Vaginal antiseptic
- ALBUCIO.** See Sulfacetamide
- ALBUMIN (HUMAN) SALT-POOR** (Cutter) Normal Human Serum Albumin, USP, NNR 1954 Vial 20 cc (5 Gm of normal serum albumin) w/20 cc buffered diluent 50 cc (12.5 Gm of albumin) w/50 cc.
e Generic name recognized by NF, USP or A.M.A.
- buffered diluent Packed w/administration set
Use Shock, hypoproteinaemia, renal dis eases & cirrhosis
- * **ALBUMIN, NORMAL HUMAN SERUM,** USP NND 1962 See Albumin, Human (Salt Poor) 25% (Cutter)
- ALBUMIN TANNATE**, NNR 1925 (Na Mir currently lists) Albutannin, tannalbrn, tannin albuminate)
Use Diarrhea
- ALBUMIN TEST** (Northwest) Sulfosalicylic Acid
Use Diagnostic for urine albumin
- ALBUMIN TEST SOLUTION.** (Endo) Sulfo salicylic acid 10%, Aq menthol 50% Liq Bot 4 oz, 8 oz, 16 oz
- ALBUMISOL**, NND 1962 (Merck Sharp & Dohme) Normal serum albumin component of human blood 5%/cc Bot 250 cc, 500 cc w/intravenous set Also available as 25% concentration Vial 20 cc or 50 cc w/intravenous set
Use In cases of shock and burns
- ALBUSTIX** (Ames) Firm paper reagent strips impregnated with tetrabromphenol blue, citrate buffer and a protein adsorbing agent Bot 120s
Use Diagnostic, to indicate the presence of albumin in the urine
- ALBUTEST.** (Ames) Tab containing tetra bromphenol blue, citrate buffer, protein adsorbing agent and an inert carrier Bot 100s, 500s
Use Colorimetric urine protein estimations
- * **ALBUTOIN** (Baxter) 3 allyl 5 isobutyl 2 thi ohydantoin
Use: Anticonvulsant
- AL-CAROID** (American Ferment Co) Mag oxide 7 gr, mag carbonate 16 gr, bismuth subcarbonate 24 gr, caralid 24 gr, cal car bonate 182 gr, sod bicarbonate 183 gr/oz Pow Pkg 2 oz, 4 oz, 1 lb, Tab 1 tablet is equiv to ½ teaspoonful of the powder Bot 20s 50s, 100s, 500s, 1000s
- ALCAWAL.** (Wollau) Aluminum hydroxide 2 gr, calcium carb 1 gr, magnesium tri silicate 4 gr/Tab Bot 50s
Use (1 2 p c) Antacid
- ALCETIN** See: Dextrogesic, Tab (Pitman Moore)
- ALCID H M B** (Maltbie) Susp of hydrated alumina with homatrapine methylbromide 5 mg/8 oz Liq Bot pt gal
Use: Peptic ulcers
- ALCID LIQUID** (Maltbie) Susp of hydrated alumina, aluminum hydroxide gel Liq Bot pt, gal
Use: Antacid adsorbent

ALCITEX. (Elder) Phenobarbital $\frac{1}{8}$ gr., atropine sulfate 1/500 gr., cal. carbonate $3\frac{1}{2}$ gr., magnesium carbonate $2\frac{1}{2}$ gr., cerium oxalate $\frac{1}{2}$ gr./Tab. Bot. 100s, 500s, 1000s
Use: Sedative w/antispasmodic

ALCOHOL, 5% IN BECLYSYL, 5% IN WATER. (Abbott) Alcohol, dehyd. 50 cc., dextrose 50 Gm., thiamine HCl 10 mg., riboflavin 10 mg., nicotinamide 250 mg., pyridoxine HCl 5 mg., in water for inj./1000 cc. Bot. 1s, 48s.
Use: I.V., nutritional.

ALCONOL-DEXTROSE SOLUTIONS. (Don Baxter) 5% Alcohol, 5% dextrose in distilled water, 100 cc.; 5% alcohol, 5% dextrose in normal saline, 1000 cc.; 5% alcohol, 5% dextrose with vitamins in distilled water, 1000 cc.; 5% alcohol, 5% dextrose with vitamins in normal saline, 1000 cc.; 5% alcohol, 25% dextrose in distilled water, 1000 cc.

ALCONOL IN VITADIX-B. (Cutter) Alcohol 5%, dextrose 5% w/Vit. B-1 5 mg., Vit. B-2 7.5 mg., nicotinamide 50 mg., Vit. B-6 3 mg./liter, Salfiflask 1000 cc.
Use: I.V., 500 to 2000 c., Vit. B complex.

ALCON-EPRIN 12 and 50. (Alcon) Phenylephrine HCl 0.12% and 0.5% plus Benzalkonium chloride. Dropper Bot. 30 cc.

ALCON-EPRIN 25. (Alcon) Phenylephrine HCl 0.25% plus Benzalkonium chloride. Dropper Bot. 30 cc. Spray Pkg. 30 cc.

ALCO-PAN. (Bell) Panlothyenyl Alcohol 250 mg./cc. Viol 10 cc I.M. inj.
Use: Establishes normal peristalsis after abdominal surgery.

ALCOPARA. (Burroughs Wellcome) Bephenium Bromide 5 Gm./pk.

Use: Roundworm and hookworm infections

ALCOPHOBIN. (Consol. Midland Corp) Disulfiram 0.5 Gm./tab. Bot. 50s, 100s, 250s, 1000s.

Use: Alcoholicism.

ALCRISTA TABLETS OF IPECAC. (Lilly) Ipecac alkaloids 10 gr./Tab. Bot. 100s, 500s.

Use: (1-3 t.i.d.) Amebic dysentery.

ALDACTAZIDE. (Searle) Spironolactone 75 mg., hydrochlorothiazide 25 mg./Tab Bot. 40s, 100s.

Use: Treatment of edema as diuretic

ALDACTONE, N.N.D. 1962 (Searle) Spironolactone 100 mg./Tab. Bot 20s, 200s.

Use: Edema. Diuretic.

ALDACTONE-A. (Searle) Spironolactone 25 mg./Tab Bot. 40s, 100s.

Use: Edema.

ALDARSONE, N.N.D. 1959. (Abbott) (Phenar-

* Generic name recognized by N.F., U.S.P. or A.M.A.

sone Sulfoxylate) Sodium 3-amino-4-hydroxy-phenylarsinate-N-methanal sulfoxylate. Amp. (1 Gm.) 6s, 25s, 100s.

Use: I.V., Trypanosomiasis & neurosyphilitis

Vaginal Suppositories. (Abbott) Aldarson 0.5

Gm., Icolin 2.5 Gm./Vial. Box, single dose, Box 25s, 100s.

Use: Trichomonas vaginalis vaginitis.

* **ALDEFORM DISKS.** (Paul J. Mandaback) Paralormaldehyde 93% tabs. Bot. 100s

Use: Release of formaldehyde

AIDIAZOL-M. (Massengill) Susp.—Sulfadiazine and sulfamerazine 0.75 Gm. each with Sod. citrate 3 Gm./15 cc. Bot. pt., gal. Tab.—Sulfadiazine and sulfamerazine 0.12 Gm. each with Sod. citrate 0.25 Gm./Tab Bot. 100s, 1000s

Use: Di-sulfonamide therapy.

ALDIAZOL-S. (Massengill) Susp.—Sulfadiazine 0.75 Gm., sulfathiazole 0.75 Gm., Sod. citrate 22 gr., Sod lactate 27 gr./15 cc Bot. pt., gal. Tab.—Sulfadiazine 0.12 Gm., sulfathiazole 0.12 Gm., Sod. citrate 0.25 Gm./Tab. Bot. 1000s

Use: Dual sulfonamide therapy.

ALDIMATE. (Crestmed). Dihydroxy aluminum aminoacetate, 0.5 Gm./Tab. Bot. 100s.

Use: Nonsystemic antacid

ALDINAMIDE. 2-Carbomoyl pyrazine, See. Pyrazinamide, Tab. (Merck Sharp & Dahme)

ALDO TABLETS. (Warren-Teed) Strychnine sulf. 0.8 mg., belladonna ext. 5 mg., phenolphthalein 15 mg., alain 30 mg., Ipecac 4 mg./Tab. Bot. 1000s

ALDOCORTIN. See: Aldosterone

AEDOMET. (Merck and Co., Inc) Methyldopa
Use: Antihypertensive agent

ALDOSTERONE. 11-beta-21-Dihydroxy-3, 20-dioxo-4-pregnen-18-al. (Electrocortin, Aldocortin)

AIIMBARE. (Meyer) Tab Bot. 100s, 1000s

ALEPHED. (Massengill) Allylphenylbarbituric acid $\frac{1}{2}$ gr., ephedrine sulfate $\frac{1}{2}$ gr./Cap Bot. 100s, 1000s

Use: (1 q. 3-4 h) Antispasmodic.

ALEPSAL. (Fougera) Phenobarbital $1\frac{1}{2}$ gr., belladonna pow. $\frac{1}{2}$ gr., caffeine 2/5 gr./Tab. Bot. 20s.

Use: (1 t.i.d.) Sedative.

ALERCAP. (Ethical Pharm) Pyranisamine maleate 25 mg., Dover's pow. $\frac{1}{2}$ gr., aceto-phenetidin $3\frac{1}{2}$ gr., aspirin 2 gr., camphor $\frac{1}{2}$ gr., caffeine $\frac{1}{2}$ gr., atropine sulfate 1/500 gr./Cap. Bot. 100s & 1000s.

Use: Analgesic, antihistaminic, antipyretic.

- ALBDEX.** (Wyeth) Maltose 50%, dextrins 46% ash 0.5% Can 1 lb
Use Infant food
- ALERIN-TD** (Finley) Chlorpheniramine maleate 8 mg/Cap Bot 50s; 250s TD Caps 12 mg Bot 50s 250s
Use Treatment of allergic conditions
- ALESTONIC** (Merrell) Meratol HCl 2 mg 8 1 10 mg, B 2 5 mg B 6 1 mg niacinamide 50 mg choline 100 mg inositol 100 mg and 7 minerals/3 tbsp (45 cc) Bot 1 pt
Use Appetite stimulation
- ALESTROL** (Teragar) Safflower a 1715 mg Vit B 6 1.5 mg, E 0.03%/Cap Bot 100s
Use: Reduction of elevated blood cholesterol levels
- ALEUDIN** (National Drug) (Isopropylar terenol) Tab (10 mg) Bot 50s 1000s
Use Bronchodilator
- ALVAVIS** (Winthrop Labs) Triton WB 1339 0.125% Sod bicarb 2% glycerine 5% solution Bot 500 cc & 60 cc
Use Aerosolization for tuberculosis & other types of chronic pulmonary diseases
- ALFABETAMIN** (Breon) Vit A 5000 u, D 800 u, B 1 2 mg B 2 3 mg, C 75 mg, eiacin 20 mg, pantothenic acid 1 mg/Cop Bot 100s, 500s 1000s
Use: (1 daily) Multivitamin therapy
- ALFABETAMIN STBONGER** (Breon) Vit A 12,500 u, D 1000 u, C 150 mg, B 1 5 mg B 2 5 mg B 6 1 mg, eiacinamide 50 mg, pantothenic acid 10 mg/Cop Bot 100s 500s
Use Multivitamin therapy
- ALFACOL** (Ascher) dl Alpha tocopheryl acetate Cap (100 mg) Bot 100s 1000s; Vial (200 mg/cc in sesame oil) 10 cc
Use: I.M. orally vitamin E therapy
- ALFLOSONE ACETATE**, NND 1962 (Merck Sharp & Dohme) Fluorocortisone acetate Lot 0.1 & 0.25%, plastic squeeze Bot 15 cc Oint 0.1 & 0.25%, Tube 5 Gm, 15 Gm
Use: Many types of dermatitis and skin inflammations
- ALGIC.** (Spencer) Chlorpheniramine maleate 3 mg, phenyltoloxamine dihydrogen citrate 50 mg, rancephedrine HCl 25 mg/Tab Bot 100s
Use Antihistaminic therapy
- ALGIC-S A.** (Spencer) Chlorpheniramine maleate 6 mg, phenyltoloxamine dihydrogen citrate 100 mg, rancephedrine HCl 60 mg/Tab Bot 50s
Use Anti-allerg c
- ALGIN, ADR** See: Sodium Alginate, NF
- ALGINATES**, Cat & Sod W/Diactyl sod sol
- fosuccinate, aethioquinone derivative See: Combinate (Mead Johnson)
- ALGINEX.** (Pitman Moore) (Sulfathiazole) Sulfathiazole 10%, tyrothricin 0.05%, 2 chloro 4 phenyl phaeol 0.1%, methyl p hydroxy benzoate D 1% in a greaseless base, Tube 1 oz
Use Topically, antibacterial & fungicide
- ALGLYN, NND 1962** (Brayten) Dihydroxy aluminum omeoacetate Tab (7 1/2 gr) Bot 50s, 100s
Use Antacid
- ALGOSON** (McNeil) Acetyl p-aminophaeol 300 mg, buisol Sod 7.5 mg/Tab Bot 100s, 1000s
Use: (1 q 4 h) Analgesic and sedative
- ALNYDROX TETANUS TOXOID, NNB 1954** (Cutter) Tetanus Toxoid, Aluminum Hydroxide Adsorbed, USP. Vial 1 cc & 5 cc
- ALIDASE, NND 1962** (Searle) (Hyaluronidase) Amp, serum type 500 viscosity units, Pkg 6s, 25s, 100s 500 V units=150 TB units
Use S.C. more rapid dispersal and absorption of solutions
- ALINESE.** (Mackell) Aluminum hydroxide 7 gr, magnesium trisilicate 1 1/2 gr, reduced iron 1/4 gr/Tab Bot 50s, 250s, 500s Susp.—Aluminum hydroxide 6%, magnesium trisilicate 6% Bot 12 oz
Use: Antacid
- ALIPCO.** (Drug Products) Strychnine sulfate 0.5 mg belladonna lvs ext 54 mg, phenolphthalein 32 mg, aloin 13 mg, podophyllin 4 mg, ipecac 4 mg, capsicum oleoresin 2.6 mg/Cop Bot 100s 1000s
Use (1 or 2 hs p n) Adult laxative
- ALKABAB** (Brewer & Co) Atropine sulf 0.2 mg phenobarbital sod 15 mg, calcium carbonate 120 mg, aluminum hydroxide gel 200 mg/Tab Bot 100s
Use Spasm of gastrointestinal tract
- ALKA-CORD.** (Sutliff & Case) Xanthoxylum 8 gr Avena Sativa 8 gr, Ginger 4 gr, Gentian 4 gr, Sodium Bicarbonate 4 gr, Berberis Aquifolium 8 gr/fl oz Bot pt, gal
- ALKAGEL, NND 1961** (Lantien) Aluminum hydroxide gel dried Tab (0.32 Gm & 0.65 Gm) Bot 100s 1000s
Use Antacid
- ALKAGESTABS.** (Blue Line) Sod bicarb, papain, pancreatin, ginger pow & rhubarb Tab Bot 100s, 1000s
Use (2 3 q d) Antacid
- ALKALINE ABOMATIC** (Sutliff & Case) Sodium Bicarbonate 0.325 Gm, Sodium Borate 0.325 Gm, Sodium Chloride 0.325 Gm, Sodium Salicylate 20 mg, Sodium Benzoate 20 mg/Tab Bot 1000s

ALKALINIZING AGENTS are intended to restore the alkali reserve of the body (blood) and induce an alkaline urine.

Sen: Alka-Vess, Tab. (Ames).
Alka-Zann, Pow. (Warner).
Alknts, Tab. (Upjohn)
Alvesen, Tab. (Burroughs-Wellcome).
Anachloric, Tab. (Upjohn)
Antacids, Gastric.
Caknam, Pow. (Paxton)
Cal-Bis-Ma, Pow., Tab. (Warner-Chilcott)
Cito Lactate, liq. (Dantley-Evans)
Citroalka, liq. (Paske, Davis).
Citrocarbonate, Pow. & Syr. (Upjohn)
Citra-Soda, Pow. (Abbott).
Effervescent Granules. (Reed & Carnrick).
Enlacarb, Pow. (Reed & Carnrick).
Lactate-Ringer's Solution. (Varlaus Mfr.).
Nacitra, Tab. (McNeil).
Neutralizer (Upjohn)
Sodanco, Pow. (Erwin-Nisler) :
Tri-Calsale, Pow. (Paxton)

ALKALOL. (Alkalol Co) Thymol, eucalyptol, menthol, camphor, benzoin, Pot. alum, Pot. chlorate, Sod. bicarb., Sod. chloride, sweet birch oil, spearmint oil, pine & cassia oil. Bot. 8 oz., 1 pt.

Use: Eyes, nose, throat & all inflamed mucous membranes.

ALKAM. (First Texas) Dihydroxy aluminum aminoacetate. Tab. (7½ gr.) Bot. 100s, 500s, 1000s.

Use: Antacid.

ALKAM COMPOUND. (First Texas) Dihydroxy aluminum aminoacetate 7½ gr., homatropine methylbromide 1/100 gr., phenobarbital ¼ gr./Tab. Bot. 100s, 500s, 1000s.

Use: (1-2 p.c.) Antacids & antispasmodic.

ALKA-PHEN. (Rowell) Homatropine methylbromide 1/20 gr., phenobarbital ¼ gr., pepsin 3 gr., charcoal 5 gr./cap. Bot. 100s, 500s, 1000s.

Use: (1 t.i.d.) Antispasmodic.

ALKARHEIN. (Merritt) Rhubarb 1.3 Gm., glycerin of pancreatin 0.5 cc., combined in an mixer base with Sod. bicarb., cinnamon, peppermint and other aromatics, alcohol 17%/3D cc. Bot. pt., gal.

Use: Stomachic and gastric antacid.

ALKARINSE. (Massengill) Sod. bicarbonate 5 gr., sod. borate 5 gr., sod. chloride 5 gr., aromatics q.s./Cap. and Tab. Cap. Bot. 500s, Tab. Bot. 100s, 1000s.

Use: Nonirritating cleansing application for nasal passages.

ALKASOL. (Stadmd) Disodium hydrogen chloride

Use: Systemic alkaliizer

* Generic name recognized by N.F., U.S.P. or A.M.A.

ALKASPASMOL. (Walker) Belladonna alkaloid, equal to ext. ⅓ gr., aluminum hydroxide 250 mg., benzocaine 8 mg., magnesium trisilicate 250 mg., phenobarbital 8 mg./Tab. Bot. 100s, 1000s.

Use: (1 q. 3-4 h) Antacid & antispasmodic.

* **ALKAVERVIR**, N.N.D. 1962. Mixture of *Veratrum viride* alkaloids. See: Veriloid, Prep. (Riker)

W/Alsnroxylon.

Sen: Rau-Vartin, Tab. (National Drug)

Rauiloid-Veriloid. (Riker)

W/Mannitol hexanitrate & phenobarbital

See: Nitranital P.V. (Metrell).

Veratrabar (Smith-Dorsey).

ALKA-ZANE. (Warner-Chilcott) Sod. citrate 2.66 Gm., Sod. bicarb. 1.64 Gm., magnesium phosphate 0.246 Gm./5 Gm. of powder. Bot. 1½, 4, 8 oz.

Use: Alkalinizing agent

ALKERAN (Burroughs Wellcome) Phenylalanine mustard.

Use: Treatment of cancer.

ALKETS, TABLET. (Upjohn) Col. carb. 12 gr., Mag. carb. 2 gr., Mag. oxide 1 gr. Tab., Bot. 100s, 500s, 1000s.

Use: (1 or 2 t.i.d.) With Pamine to combat constipation.

ALKOFANONE. See: Clafanone, Tab. (Hoffman-LaRoche)

ALKRON. See: Parathion.

ALKYLDIMETHYLBENZYL AMMONIUM CHLORIDE. See: Baby Care, Cream. (Inveox)

W/Methylrosaniline chloride, polyoxyethylene-phenol, polyethylene glycol terdidethylthioether. See: Hyva, Tab. (Holland-Rantos)

ALKYLBENZYLDIMETHYLAMMONIUM CHLORIDE. See: Benzalkonium Chloride, Prep

ALLAMINE. (Denob) Chlorpheniramine maleate, 4 mg.; phenylpropanolamine HCl, 25 mg./Tab Bot 100s, 250s. also available w/Prednisone.

Use: Respiratory, nasal, ophthalmic and dermatologic allergies.

ALLANTOIN. (National Drug) (Glyoxylic diureid) Oint. (2%) Jar 1, 4 oz., 1 lb., 5 lb
Use: Wounds and resistant ulcers to stimulate the growth of healthy tissue.

ALLANTOIN. (Massengill) 2% in a water-washable base. Oint. 1 lb.

Use: Typically, aid in the promotion of granulation.

ALLANTOIN. See: Cutanal, (Summers).
W/9-Aminoacridine, sulfanilamide. See: AYC-Improved, Oint. (National Drug)

W/Acetic acid & aluminum hydroxide See Allucea Cap (Kremers Urban)
 W/Benzalkonium chloride di desoxyephedrine hydrochloride See Narakon Lq (Chesebrough)
 W/Coal tar extract See Alphosyl Lubricating Cream (Reed & Carnrick)
 W/Coal Tar extract Hexachlorophene See Sebical Cream (Reed & Carnrick)
 W/Coal tar hydrocarbons See Alphosyl HC Lot (Reed & Carnrick)
 W/Coal tar extract squalane See Skayfos Lotion (Lee Calbert)
 W/Crystal violet, urea See Urantain Violet, Oint (Irwin Neiler)
 W/Metaphen (Abbott) Allantoin 2%, metophen 1.500 Cream, Tube 1 oz, 4 1/4 oz, 1 lb
 Use Topically, tissue stimulant with antiseptic
 W/Methylbenzethonium chloride phenylmercuric acetate 9 aminoacridine HCl, succinic acid, lactose See Vogitab Vog Tab (Aicher)
 W/Neomycin sulfate dipiperdon HCl See Ro Derm (Lemmon)
 W/Neomycin sulfate vitamin A, D See YAD Cream (Walker)
 W/Phenylmercuric acetate methylbenzethonium chloride 9 aminoacridine HCl succinic acid, lactose See Vogitabs, Tab (Aicher)
 W/Sulfadiazine, topocaine See Sulfona mets Lot (National Drug)
 W/Sulfanilamide See Allantoin, Oint (National)
 W/Sulfanilamide & aminoacridine HCl See ACR Allantoin Oint (National)
 W/Sulfanilamide & sulfathiazole See Allantoin with sulfathiazole, Oint (National Drug)
 W/Sulfanilamide sulfathiazole & ephedrine sulfate See Allantoin Liquid with Ephedrine (National Drug)
 W/Tar solution, methamine See Altara, Gel (Domet)
 W/Tyralthricin & desoxyephedrine See Tyro Cadin, Sol (Recei)
 W/Trypsin zirconium carbonate, ethyl aminobenzoate, benzalkonium chloride Amersol 9 See Trypsigal Cream (Brucel)
 W/Vitamin A & D See Andoin, Cream (Uimer).

ALLANTOIN COMPOUND, OINTMENT
 (Upjohn) Allantoin 2%, cod liver oil concentrate 182%, secondary-amylicresols 0.5% in a eancreay base Tube 1 oz
 Use: Aid in the repair of tissue injuries & the promotion of epithelization
ALLANTOINIDE (National Drug) Allantoin

2%, sulfanilamide 10%, in a greasy water miscible base Oint. Tube 1 oz. Jar 4 oz, 1 lb, 5 lb

Use: Topically, to promote healing
ALLBIE, (Robins) Thiamine HCl 15 mg, pyridoxine HCl 5 mg, riboflavin 10 mg, nicotinamide 50 mg, calcium pantothenate 10 mg /Cap Bot 100s, 500s, 1000s
 Uses (1 daily) Vitamin B complex therapy
W/Vitamin C ascorbic acid 300 mg /Cap Bot 100s, 500s, 1000s
 Use (1 daily) Deficiencies of the water soluble vitamins

ALLBEX, (Approved Pharm Corp) Vit B1 5 mg B 2 1 mg, B 6 .25 mg, calcium pantothenate 3 mg, niacinamide 20 mg, ferrous sulfate 194.4 mg, inositol 10 mg, choline 10 mg, Vit B 12 (conc) 1 mcg /Cap Bot 50s, 100s, 1000s
 Use: Vitamin deficiencies

ALLERCORB, (Plesner) Ascorbic acid Tab 50 mg Bot 100s, 500s, 1000s
 Use: Vitamin C therapy.

ALLERCREME, (Texas Pharm) Ammoniated mercury 5% Lotion, Nursery, 6 oz; Lot on Skin, 6 oz; Oil, Detergent, 8 oz, 1 pt 1 gal; Oint 1 lb; Powder, Nursery 4 oz
 Use Topically, skin medications

ALLERCREME CLEANSING LOT, (Texas Pharm) White oil in water emulsion Bot 2 oz, 8 oz
 Use Skin allergic to regular cosmetics
ALLERCREME EYE SHADOW, (Texas Pharm) Available in 4 colors; free of irritating ingredients /Golden container
 Use Allergenic cosmetic

ALLERCREME SOAP SHAMPOO, (Texas Pharm) Pure, mild castile type shampoo used w/ either hard or soft water /Bot 8 oz
 Use Hypo allergenic shampoo

ALLERCREME SUNTAN LOTION, (Texas Pharm) Sendore Lot /Bot 4 oz
 Use Sun protective lotion

ALLERCREME ULTRA EMOLLIENT, (Texas Pharm) Refined derivatives of lanolin, sorbitol, cetyl alcohol petrolatum blend and purified water /Jar 2 oz
 Use Relieve and prevent dry skin

ALLERCUR, NND 1962 (Roerig) Clemizol HCl 1 p chlorobenzyl 2 pyrrolidylmethyl benzimidazole 40 mg & 20 mg /Tab Bot 100s
 Use Antihistamine

ALLERSEE, (Pharmacroll) Chlorpheniramine maleate 1 mg, methopropylene fumarate 5 mg phenylpropanolamine HCl 25 mg - V C 37.5 mg /Tab Bot 24s, 48s
 Use: Respiratory allergies

ALLERGAN, (Allergan) Boric acid 12% chlorobutanol 0.5%, antipyrine 0.1%, phenol 0.15%, methyl cellulose 0.12%, plus

- ethylenediamine tetraacetic acid. Bot. plastic 60 cc.
 Use: Ophthalmic lubricant and contact lens fluid.
- ALLERGESIC.** (Bryant) Phenyltoloxamine dihydrogen citrate 1%, neomycin sulf. 0.2% tytrathricin 0.02%, bithional 0.5%, zincum oxide 4%, dipiperon HCl 0.25%. Lotion, Plastic sqz, Bot. 4 oz.
 Use: Treatment of dermatitis caused by poison ivy, oak, and sumac.
- ALLERGEX.** (Hollister-Slier) Silicones, triethylene glycol, mineral oil with fungicide. Bot. ½ pt., 1 pt., 1 qt.
 Use: Control of house dust allergens.
- ALLERGIN.** (Myers-Carter) Methopropylene HCl 25 mg./cc. Vial 10 cc. Also Forte 100 mg./cc. Vials 10 cc.
- ALLERGI-MIST #1.** (Allergimist) Guanine, 0.001%; uric acid, 0.003%; thymine, 0.004%; indole, 0.005%; glycine, 0.066%; glutamic acid, 0.088%; alevaire, 0.222%; ethyl alcohol, 0.666%; organic dye, 0.014%; liquid phenol, 0.044%; sodium chloride, 0.015%; acetylcholine, 0.026%; amino acid derivatives, 0.026%/sol. 20 cc. Plastic nebulizers 20 cc.
 Use: Antiallergic.
- ALLERGI-MIST #2.** (Allergimist) Guanine, 0.001%; uric acid, 0.003%; thymine, 0.004%; indole, 0.005%; glycine, 0.066%; glutamic acid, 0.088%; alevaire, 0.222%; ethyl alcohol, 0.666%; organic dye, 0.014%; liquid phenol, 0.044%; sodium chloride, 0.015%; acetylcholine, 0.037%; amino acid derivatives, 0.037% with distilled water, q.s./20 cc. Plastic nebulizers: 20cc.
 Use: Antiallergic.
- ALLERGY PREPARATIONS.**
 See: Antihistamine Preparations.
 Cortisone Preparations.
 Hispril, Preps. (Smith, Kline & French)
 Vitamin A & D Preparations.
- ALLERMIST.** (Warran-Teed) Phenylephrine HCl ¼%, theinylpyromine HCl 0.2%, methylbenzethonium chloride 0.01% in a buffered solution. Bot. ¼ oz.
 Use: Aerosol nasal decongestant.
- ALLIMIN.** (Van Patten) Garlic concentrate 4% gr., parsley concentrate 2½ gr./Tab. Bot. 25s, 60s, 125s, 250s.
 Use: (f. i. d.) Peripheral vasodilator.
- ALLOBARRITAL.** See: Diallyl Barbituric Acid (Various Mfr.).
- ALLOBARRITONE.** See: Diallyl Barbituric Acid. (Various Mfr.).
- ALLOCAINE.** See: Procaine Hydrochloride. (Various Mfr.).

- ALLONAL.** (Roche Lab) Alurate 1 gr., ocelaphenedin 3 gr./Tab. Box 50s
 Use: (f. i. d.) Sedative.
- ALLUCEE.** (Kremers-Urbain) Allantoin 60 mg., ascorbic acid 50 mg., aluminum hydroxide 100 mg./Cap. Bot. 100s, 1000s.
 Use: (f. q. 3-4 h.) Peptic ulcer.
- ALLYDYNE.** (Irwin-Neisser) Mustard oil, menthol, thymol, camphor, methyl salicylate, turpentine & benzoic acid. Bot. pl., gal.
 Use: Topically, rubefacient & counter-irritant.
- ALLYL BARRITURIC ACID, N.F. X., N.N.R. 1952;** allyl-isobutylbarbituric acid Tablets, N.F. X. (Various Mfr.). See: Sandoptal, Tab (Sandoz)
 Use: Sedative.
- W/Aminopyrine, caffeine, See: Optalidon, Tab. (Sandoz)
 W/A.P.C. See: Fiorinal, Tab. (Sandoz)
 W/Methamphetamine sulfate, carbamide. See: Diabese, Tabs. (Philadelphia Lab).
 Diarobese, Tabs. (Riber).
- * **ALLYLESTRINOL.** 17 α -allyl-17 β -hydroxy-4-estrene. See: Gestonin (Organon Inc)
- ALLYLGUIACOL.** See: Eugenol. (Various Mfr.).
- ALLYL-ISOBUTYLBARBITURIC ACID.** See: Allyl Barbituric Acid, N.F.
- ALLYLISOPROPYLMALONYLUREA, N.N.R. 1946.**
 See: Alurate (Roche Lab)
 Aprabarbital, N.F.
- ALLYL ISOTHIOCYANATE, N.F. VIII.** (Oil of mustard) (Various Mfr.).
 Use: Counterirritant in neuralgia.
- W/Methyl salicylate, camphor, etc. See: Allydyne, Liq. (Irwin Neisser)
- N-ALLYLNORMORPHINE HCl, U.S.P. N.N.D. 1960.** *Nalorphine. See: Nalline HCl, Prep. (Merck Sharp & Dahme)
- 5-ALLYL-5-PHENYLBARBITURIC ACID.** (Alphenal, Alphenate) (Various Mfr.). See: Alphenem, Cap. (Massengill).
 Use: Sedative & hypnotic.
- W/Aminophylline. See: Alphyllin, Cap. (Massengill)
- W/Cyclopentyl-allyl-barbituric acid. See: Cyalph, Tab. (Massengill)
- W/Ephedrine sulfate. See: Alephed, Cap. (Massengill).
- * **ALLYLPRODINE.** 3-Allyl-1-methyl-4-phenyl-4-propionylpiperidine
- 5-ALLYL-5-SEC-BUTYLBARBITURIC ACID.** See: Iatuseal, Cap. (Winthrop Lab.)
- ALMADON.** (Benton) Aluminum hydroxide-magnesium carbonate 5 gr., hyoscyamine sulfate 0.019 mg., phenobarbital 8.1 mg.

* Generic name recognized by N.F., U.S.P. or A.M.A.

- atropine sulfate 0.0097 mg, hyaline hydrobromide 0.00325 mg /Tab Bot 100s
 Use: Antacid & antispasmodic
- ALMAG.** (Drug Products) Aluminum hydroxide 7½ gr, magnesium trisilicate 7½ gr / Tab Bot 100s, 1000s
 Use: Antacid
- ALMAGUCIN, NNR 1955** Antacid mix of gastric mucin, dried aluminum hydroxide gel USP & Mag trisilicate USP
 See: Mucolin, Liq, Tab (Harrower)
 Trimucolon Tab (Winthrop Stearns)
- ALMEBEX** (Meyer) Thiamine hydrochloride 10 mg, riboflavin 20 mg, niacinamide 50 mg, pyridoxine HCl 0.4 mg, pantothenic acid 0.6 mg, choline 330 mg, inositol 160 mg, biotin preserved with benzoic acid 0.1%/1 dr Bot pl, gal Also available w/B 12 5 mcg/dr
 Use: Vitamin deficiencies
- ALMEDERM CREAM, (Meyer)** (1% Hexachlorophene in a washable cream base) 1 oz, 1 lb
 Use: Antibacterial
- ALMEDIOL TABLETS, (Meyer)** β Estradiol 0.125 mg Bot 100s, 1000s
 Use: Buccal or sublingual administration
- ALMEDIOL VIAL, (Meyer)** β Estradiol 1 mg, procaine HCl 2% Sod bisulfite 0.1% benzyl alcohol 2%, In aq propylene glycol/cc Vial 10 cc
 Use: Estrogen therapy
- ALMEFRIN** (Meyer) Phenylephrine HCl 1% Bot pl
 Use: Intranasal vasoconstrictor
- ALMEFROL** (Meyer) d Alpha tocopherol 100 mg/Cap Bot 100s, 1000s
 Use: Vit E therapy
- ALMEGESIC** (Meyer) Pyrilamine maleate 25 mg, hyoscyamine HBr 0.04 mg, hyoscine HBr 0.008 mg, aspirin 2½ gr, acetophenol 2½ gr, caffeine ½ gr/Tab Bot 100s, 1000s
 Use: (1 q 1 d) Analgesic with antihistaminic
- ALMEHIST CAPS** (Meyer) Pyrilamine maleate 50 mg, hyoscyamine HBr 0.016 mg, hyoscyamine HBr 0.008 mg, d omphetamine sulfate 15 mg/Cap Bot 100s A 1000s
 Use: Allergic symptoms & motion sickness
- ALMEHIST SYRUP, (Meyer)** Pyrilamine maleate 25 mg, hyoscyamine HBr 0.008 mg, hyoscyamine HBr 0.040 mg/dr Bot pl, gal
 Use: Antihistaminic
- ALMELOSE, (Meyer)** Sod carboxymethyl cellulose D.5 Gm Bot 100s, 500s
 Use: Antacid
- ALMENIC, (Meyer)** Vit B 12 cryst 1 mcg, A 1250 u, D 250 u, ascorbic acid 25 mg, thiamine HCl 1.25 mg, riboflavin 1.25 mg, niacinamide 12.5 mg, pyridoxine HCl 0.25 mg, calcium pantothenate 1.25 mg, d alpha tocopherol 2.5 mg, folic acid 1 mg, ferrous sulfate, dried, 3 gr, Sod molybdate 0.2 mg, copper sulfate 1 H₂O 10 mg, cobalt sulfate 0.5 mg/Cap Bot 100s, 1000s
 Use: Vitamin deficiencies
- ALMERAZINE, (Meyer)** Syrup, pl, gal
- ALMERET** (Meyer) Vitamin B 12 Cop—Vitamin B 12 25 mcg/Cop Bot 100s Syr—Vit B 12 5 mcg/5 cc Bot 8 oz Vial—Vit B 12 30 mcg & 1000 mcg/cc 10 cc
 Use: Vit B 12 therapy
- ALMERET FORTE, (Meyer)** Vitamin B 12 100 mcg/cc Vial 10 cc, 20 cc
- ALMESCON, (Meyer)** Sod estrone sulfate 1.25 mg/Tab Bot 100s, 1000s
 Use: Estrogen therapy
- ALMESED** (Meyer) Pentobarbital 7½ mg, phenobarbital 7½ mg, hamotropine methyl bromide 2½ mg/5 cc of elixir Bot pl, gal
 Use: (1 dr 1 id) Sedative & antispasmodic
- ALMESTIGMINE, (Meyer)** Physostigmine salicylate 0.6 mg atropine sulfate 0.4 mg, Sod bisulfite 10 mg, dextrose 5%/cc Vial 20 cc
 Use: Parasympathetic & mydriatic agent
- ALMESTRONE** (Meyer) Estrone 10000 IU, progesterone 10 mg, β estradiol 0.5 mg, in sesame oil/cc Vial 10 cc
 Use: EM, estrogen and corpus luteum therapy
- ALMETNINE** (Meyer) Methscopolamine HBr 5 mg, mephobarbital 50 mg/cap Bot 30s, 100s
 Use: (2 daily) Anticholinergic in the treatment of peptic ulcers
- ALMETRIFON POWDER** (Meyer) Chiniolon 75%, boric acid USP 5% Sod lauryl sulfate 0.05%, lactose & dextrose qs Bot 2 oz
 Use: Amebicide
- ALMETRIFON TABLET, (Meyer)** Chiniolon 0.33 Gm Sod lauryl sulfate 0.04 Gm, corn starch, lactose qs/Vog Supp Bot 12s, 100s, 1000s
 Use: Amebicide
- ALMETROPIN** (Meyer) Vial 500 units/cc, 10 cc
- ALMEX SYRUP, (Meyer)** Vit B 12 1.0 mcg, B E 10 mg, B 2 ED mg, B 6 0.5 mg, niacinamide 50 mg, rice bran ext 1350 mg, ferrous gluconate 630 mg, copper sulfate ED mg, cobalt acetate 0.2 mg/cc Bot 1 oz, 4 oz 8 oz
 Use: Hemohalic.

ALMEZYME CAPSULES. (Meyer) Mylase-100 30 mg., Prolase-300 30 mg., dehydrocholic acid 100 mg., hyoscyamine HBr 0.1 mg., hyosine HBr 0.0065 mg., atropine sulfate 0.02 mg./Cap. Bot. 100s.

Use: Gastrointestinal indigestion.

ALMEZYME TONIC. (Meyer) Mylase-100 10 mg., Prolase-300 10 mg., Vit. B-12 25 mcg./5 cc. Bot. 4 oz.

Use: Improve digestion and absorption.

ALMGOH. (Maeg) Aluminum hydroxide gel and magnesium hydroxide. Bot. 100s Tab. Use: Antacid.

ALMINATE. (Bristol) Not listed. Dihydroxy aluminum aminoacetate Tab. (7½ gr) Bot 100s, 500s.

Use: Peptic ulcers and hyperacidity.

ALMOCARPINE. (Ayers) Pilocarpine HCl ½%, 1%, 2%, 3%, and 4%. Steri-Tainer drop bot. 15 cc. each.

Use: Reduction of intraocular tension.

ALMORA. (Durst) Magnesium gluconate 0.5 Gm/Tab. Pkg. 50s, Bot. 500s, 1000s.

Use: Sedative in dysmenorrhea.

ALOBILE. (Flint, Eaton) Bile salts ½ gr., phenolphthalein ½ gr., cascara sagrada ext. ½ gr., aloin ½ gr./Tab Bot. 100s, 500s, 1000s.

Use: (1-2 hrs) laxative.

ALOCAS #1. (Sutliff & Case) Cascara sagrada ext. 16 mg., aloin 16 mg., resin podophyllum 11 mg., belladonna ext. 8 mg., oleoresin ginger 4 mg./Tab Bot 1000s.

ALOCAS #2. (Sutliff & Case) Cascara sagrada ext. 16 mg., aloin 16 mg., resin podophyllum 11 mg., belladonna ext. 8 mg., strychnine 1.1 mg., oleoresin ginger 4 mg./Tab. Bot. 1000s.

ALOE EXTRACT W/Desoxycholic acid. See: Chalmadin, Tabs. (Ames).

ALOE VERA ACTIVE PRINCIPEE. See: Alva-gel, Oint. (Kenyon)

ALOE VERA JUICE OINTMENT.

See: A-Gic, Oint. (A-Gic Corp.)

Ray-Burn Oint. (Casmus Drug)

ALOIN, N.F. (Penick) A mixture of Cryst. pentosides from various aloes

Use: Laxative.

W/Amphetamine sulfate, phenobarbital, thyroid, & atropine sulfate See: Nika, Cap. (P. Maney)

W/dl-Amphetamine sulfate, thyroid, atropine sulfate & phenobarbital. See: Nika-Caps Ph 479, Cap. (Paul Maney).

W/Belladonna ext, Ipecac, & phenolphthalein.

See: Atkapi, pill. (American Pharmaceutical)

Alahen, pill. (Parke, Davis).

Apa, pills (Hance).

W/Belladonna lvs. ext., strychnine, & cascara sagrada. See: A-B-S & C, Tab. (Horton & Converse).

W/Belladonna lvs. ext., Strychnine Sulfate, Phenolphthalein, Ipecac, Podophyllin, Capsicum oleoresin. See: Alipca, Tab. (Drug Prods).

W/Bile Salts, Phenolphthalein, Cascara Sagrada ext. See: Alabile, Tab. (Eaton).

W/dl-Deoxyephedrine HCl, Thyroid, Atropine Sulfate.

See: Amphyrad, Tab. (Hance).

Myrabie, Tab. (Meyer).

W/Phenolphthalein, Belladonna lvs. ext., Ipecac Powder, Strychnine Sulfate. See: Aloth, Tab. (Mallard).

W/Phenolphthalein, belladonna ext., strychnine, ipecac paw. See: Pheben, Pill (Sutliff & Case)

W/Phenolphthalein, Ox Bile, Cascara Sagrada. See: Glox, Tab. (Cale).

W/Strychnine, Belladonna lvs. ext. & Cascara Sagrada. See: A-B-S & C, Tab. (Horton & Converse)

ALOIN, BELLADONNA, CASCARA, AND PODOPHYLLUM PILLS, N.F. (Merrell) "Hinkle's Pills," Cascara ext. 16 mg., aloin 16 mg., podophyllum resin 10 mg., belladonna ext. 8 mg., ginger oleoresin 4 mg./pill. Bot. 100s, 1000s.

Use: Laxative with antispasmodic.

ALOKRIEN. (Chicago Pharm) Aluminum hydroxide gel. Liquid, Bot. 1 pt., 1 gal. Use: Antacid.

ALOPECTOSE POWDER. (Strassenburgh) Osmo-Kaolin 16 Gm., aluminum hydroxide, dried, 3.5 Gm., pectin 1.7 Gm., beta-lactase 7.1 Gm./oz. of pow. Jar 1½ oz.

Use: Gastric antacid.

ALOPECTOSE TABLETS. (Strassenburgh) Osmo-Kaolin 7 gr., aluminum hydroxide 2 gr., pectin 1 gr., beta-lactase 1 gr./Tab. Bot. 100s, 500s, 1000s.

Use: Gastric antacid.

W/Metopine. (Strassenburgh) Same ingredients as Alopectose tablets with atropine methylnitrate 1/120 gr./Tab. Bot. 100s, 500s, 1000s.

Use: Gastric antacid with antispasmodic.

ALOPHEN PILLS. (Parke, Davis) Aloin ½ gr., belladonna ext 1/24 gr., ipecac 1/15 gr., phenolphthalein ½ gr./Pill or Cap. Bot. 30s, 100s, 1000s.

Use: Chronic constipation.

AEOSTORIN. (Farostan) Vit. D 50,000 u., iodocasein 73 mg., irradiated yeast 325 mg./Tab. Bot. 75s.

Use: Vitamin D therapy.

ALOTHAL. (Mallard) Phenolphthalein ½ gr., aloin ½ gr., belladonna lvs. ext. 1/12 gr.

ipocac paw 1/15 gr strychnine sulfate
1/80 gr/Tab Bot 1000s
Use laxative

ALOXED. (Chicago Pharm) Aluminum hydroxide 4% homotropine methylbromide 5 mg/oz Bot 6 oz, 1 pt, 1 gal
Use (1 2 dr q 2 4 hr) Antispasmodic in peptic ulcer therapy

***ALOXIDOHE** 3 Allyl 5 methylloxaxalidine 2 4 diane Malidone

ALLEGORA (Smith Orsey) Paregoric 1/2 fl oz bismuth subgallate 27 gr, aluminum hydroxide gel 27 gr/fl oz Susp Bot 2 fl oz 1 pt

Use Antispasmodic in simple diarrhea

ALPEN, NND 1962 (Schering) Powder, after reconstitution potassium phenethicillin 125 mg/tp Bot 60 cc Tabs, potassium phenethicillin 125 or 250 mg/Tab Bot 25s 100s

Use Infections due to penicillin sensitive bacteria

***ALPHA AMYLASE** A concentrated form of alpha amylase produced by a strain of non pathogenic bacteria
See Sucramase (Ryston Co)

ALPHA CHYMAR (Armour) A lyophilized form of crystallized alpha chymotrypsin Vial (750 u/10 cc (dry) vial Vial 5 cc diluent Reconstituted 150 u/cc
Use (150 u/cc) To dissolve the zonule fibers of the lens of the eye

ALPHA-CHYMOTRYPSIN See Alpha Chymar Vial (Armour)

ALPHACOL KAPS (Alabama Pharm Co) Each capsule contains Chlorpheniramine maleate 2 mg ext belladonna 1/4 gr phenylpropranolamine HCl 1/2 gr phenacetin 2 gr aspirin 2 gr, calcium gluconate 2 gr

ALPHACOL SYRUP. (Alabama Pharm Co) Each 5 cc contains Chlorpheniramine maleate 2 5 mg, phenylephrine HCl 2 5 mg, pheniramine maleate 6 25 mg chloroform 6 75 mg 1 Menthol 0 5 mg, alcohol 6% aromatic syrupy vehicle

ALPHA-DELTALIN (Lilly) Vit A 5000 u 0 500 u/No 1 Gelseal (Gelseal No 2 X 3) Gelseals No 1, Bot 100s 1000s Gel seals No 2 Bot 100s, 1000s
Use (1 2 daily) Vitamin A & D deficiencies

ALPHADIN (Alphaden) Calcium, phosphorus, iron, iodine, Vit D Tab (5 gr) Bot 100s, 500s 1000s
Use: (1 1 d) Supplement the diet

ALPHA-DOH (Alabama Pharm Co) Each tablet contains: Phenobarbital (1/2 gr) 16 2 mg, atropine sulfate 0 012 mg, hyoscyne

* Generic name recognized by N.F., U.S.P. or A.M.A.

hydrobromide 0 005 mg, hyoscyamine hydrobromide 0 180 mg

ALPHADROL. (The Upjohn Co) Fluprednisolone 0 75 mg or 1 5 mg/Tab Bot 25s 100s, 500s

Use Corticosteroid

ALPHA-ESTRADIOL, NHR 1954 (Nov known to be beta Estradiol) See: Estradio (Various Mfr)

W/Carboxymethyl Cellulose, Tween 80, Benzyl Alcohol See: Estro D (New Mexico Pharm Co)

ALPHA-ESTRADIOL BENZOATE, HHR 1955 See: Estradiol benzoate (Various Mfr)

ALPHA-HYPOPHAMINE, HNR 1949

See: Oxytacin Injection

ALPHA-KERI. (Westward) Kerohydric (a de waxed oil soluble fraction of lanolin), mineral oil, and a nonionic emulsifier Bot 8 fl oz

Use: Treatment of dry skin

ALPHA-PHEHOXYETHYL PENICILLIN. See Phenethicillin

ALPHALIN (Lilly) Vit A 10 000 u Gelseals Bot 100s, (25,000 u) Bot 100s, 1000s (50,000 u) Bot 100s, 1000s
Use: Vitamin A deficiency

ALPHAMOLS (V.C.A. Labs) Vit A 33,330 u, D 1000 u, B 1 12 mg, B-2 8 mg, B-6 1 mg, C 150 mg, E 10 mg, B 12 5 mg-nicotinamide 150 mg, calcium pantothenate 3 mg, folic acid 0 5 mg/Gelucap Bot 50s, 100s 250s, 1000s

Use (1 daily) Multivitamin therapy

***ALPHAPRODIHE HYDROCHLORIDE, NHD 1962** ADR di alpha 1,3 dimethyl 4 phenyl 4 propionoxy piperidine HCl See Nisental (Rache Labs)

ALPHA REDISOL. (Merck & Co, Inc) Cry hydroxocobalamin 1000 mcg/cc Vial 10 cc
Use Pernicious and megaloblastic anemia

ALPHASEM. (Massengill) Allylphenylbarbituric acid Cap (1 1/2 gr) Bot 100s, 500s
Use (1 q 3 h) Sedative

W/Aminaphylline See: Alphylin Cap (Massengill)

W/Cyclophen See Cycalph Cap (Massengill)

W/Ephedrine Sulfate See Alephed, Cap (Massengill)

ALPHASTEROL. (Meyer) Vit A, Syn Cap (25,000 u) Bot 100s 1000s, (50 000 u) Bot 100s, 1000s

Use Vitamin A deficiencies

ALPHASUL (Alabama Pharm Co) Each 5 cc contains D 5 Gm (7 7 gr) micro crystalline sulfanamides as follows Sulfadiazine 0 167 Gm, sulfamerazine 0 167 Gm sulfa methazine 0 167 Gm

- ALPHATINIC.** (Alabama Pharm. Co.) Each capsule contains: liver with stomach, nat U.S.P., 150 mg., Vit. B-12 with intrinsic factor concentrate $\frac{1}{2}$ U.S.P. oral unit, Vit. B-12 7.5 mcg., terrazo sulfite 300 mg., ascorbic acid 75 mg., lolic acid 1 mg.
- ALPHA-TOCOPHEROL.** See: Tocopherol, Alpha. (Various Mfr.).
- ALPHEAL.** See: 5-Allyl-5-phenylbarbituric acid. (Massengill).
- ALPHEATE.** See: 5-Allyl-5-phenylbarbituric acid. (Massengill).
- ALPHOSYL LOTION.** (Reed & Corrick) Allantoin 2%, selected coal tar extracts 5%. Bot. 8 fl. oz.
Use: Psoriasis.
- ALPHOSYL LUBRICATING CREAM.** (Reed & Corrick) Allantoin 2%, coal tar extract 5%/Tube 60 Gm.
Use: Dermatologic; psoriasis therapy.
- ALPHOSYL-HC.** (Reed & Corrick) Allantoin 2%, coal tar 5%, hydrocortisone $\frac{1}{4}$ % lot. Bot. 4 fl. oz., cream tube 1 oz.
Use: Acute, inflammatory psoriasis.
- ALPHYLLIN.** (Massengill) Allylphenylbarbituric acid $\frac{1}{4}$ gr., aminophylline $1\frac{1}{2}$ gr./Cap. Bot. 100s, 500s, 1000s
Use: (1-2) Diuretic myocardial stimulant, bronchial dilator.
- ALPRINE.** (Ulmer) Diethylbarbituric acid $\frac{3}{4}$ gr., acetophenetidin 2 gr., aspirin $3\frac{1}{2}$ gr./Tab. Bot. 100s, 500s, 1000s.
Use: (1 q 3 h.) Sedative and analgesic.
- ALPROCOLS.** (Drug Products) Belladonna ext 4.2 mg., stromonium ext. lvs. gr. 0.26, procaine HCl gr. $\frac{1}{4}$, aluminum subacetate, benzoic acid, ichthammol, ergotin, nutgall, aromatic oils and menthated cacao bullet/rectal Supp. Box 12s, 144s.
Use: Astringent & anesthetic.
- * **ALSEROXYLON, N.N.O.** 1962. A fat-soluble alkaloidal fraction extracted from the root of *Rauwolfia serpentina*; 1 mg. is equiv. to 0.2 mg. of reserpine.
See: *Rauwolfia Serpentina* Alkaloidal Ext. (Alseroxylon)
- ALSEROXYLON-ALKAVERVIR.** A mixture of partially purified extracts of *Rauwolfia serpentina* and *Veratrum viride*.
See: *Rauvera*, Tab. (Smith-Darsey)
Rauwolfia-Veratroid, Tab. (Riker)
Rau-Verlin, Tab. (National Drug).
- ALSICAL.** (Smith Darsey) Phenobarbital $\frac{1}{4}$ gr., belladonna ext. 1/6 gr., with benzocaine, calcium carb., magnesian oxide, magnesium trisilicate & dried aluminum hydroxide/trp. of Paw. or Tab. Paw., Cos 4 ox.; Tab., Bot. 100s.
Use: Antacid and adsorbent.
- * Generic name recognized by N.F., U.S.P. or A.M.A.
- ALSULFA.** (Dook) Sellur. silicon dioxide, aluminum hydroxide. Cream; 4 oz., 1 lb., 5 lb. Lotion; 4 oz., 1 pt., 1 gal.
Use: Topically, fungicidal.
- ALTA-FUR, H.N.D.** 1962. (Eaton) Withdrawn
Furaltadone 50 mg or 250 mg./Tab. Bot. 20s, 100s.
Use: Orally, treatment of systemic bacterial infections.
- ALTAFA GELL.** (Dane) Allantoin 2%, tar solution 2%, in Ringer's solution with a colloidal soybean complex base. Tube 1 oz. Jar 7 oz., 1 lb.
Use: Treatment of psoriasis.
- ALTAFA-CORT GELL.** (Dane) Hydrocortisone 2.5 mg./Gm. with allantoin, tar, and soybean complex gel. Tube 1 oz. Bot. 4 oz
Use: Treatment of psoriasis.
- ALTEPOSE.** (Merck Sharp & Dohme) Propadrine 50 mg., thyroid 40 mg., & Detinol 25 mg./tab. Bot. 100s, 1000s.
Use: Depress appetite.
- ALTERDERM.** (Massengill) Calcium sulfide 15 mg., arsenic trioxide 0.6 mg., powd. berberis 60 mg., ext. echinacea 30 mg., ext. butternet bark 15 mg., ext. nux vomica 3 mg./Tab. Bot. 1000s.
Use: Alterative.
- * **ALTHIAZIDE.** (Pfizer) 3-allylthiamethyl-3,4-dihydro-6-chloro-7-sulfamoyl-1,2,4-benzothiazine-1,1-dioxide.
Use: Hypotensive and diuretic.
- ALTHOSE.** (Wyeth) Methadone 10 mg./fl. oz. Syr. Bot. 1 pt., 1 gal.
Use: (1 dr. q. 3-4 h.) Sedative narcotic expectorant.
- ALTRAD.** (Ascher) β -estradiol 1.0 mg./cc. Vial 10 cc.
Use: I.M., estrogen therapy.
- ALTRICATE.** (Mallard) Aluminum hydroxide 4 gr., kaolin 2 gr., phenobarbital $\frac{1}{2}$ gr., adiphenine HCl 10 mg./Tab. Bot. 100s, 500s, 1000s.
Use: Antacid with antispasmodic & sedative.
- ALTRIGEL.** (Vitarine) Aluminum hydroxide Gel with magnesium trisilicate $7\frac{1}{2}$ gr./Tab. or tsp. Bot. 100s, 1000s and 1 pt. and 1 gal.
- ALUBELAP.** (Haack) Aluminum hydroxide gel 5 gr., phenobarbital $\frac{1}{2}$ gr., belladonna, pow. ext. 1/46 gr./Tab. Bot. 100r, 1000s.
Use: Antacid, sedative & antispasmodic.
- ALUCAINE.** (Jenkins Labs) Benzocaine 5%, carbolic acid 1.39%, ichthammol 1.39%, balsam peru 0.46%, alum excis. 3.7% w/oil code, oil eucalyptus in oint. base. Oint. Tube 1 oz.
Use: Relief of pain in burns, scalds, cuts, carbuncles, and hemorrhoids.

- ALUCEN.** (Central) Methscopolamine nitrate 1 mg aluminum hydroxide magnesium carbonate 380 mg /Tab Bot 100s, 500s
Use Management of peptic ulcer
- AL-U-CBEME,** NND 1961 (MocAllister)
Aluminum hydroxide equivalent to 4% aluminum oxide Susp Bot 1 pt, 1 gal
Use Antacid
- ALUDRINE HCl,** NND 1959 (Lilly) Isoproterenol HCl, USP
Use Sublingually sympathomimetic agent
W/Cyclopentamine (Clonidine HCl) atropine sulfate & procaine HCl See Aerolone Compound Sol (Lilly)
- W/Ephedrine Sulfate, Theophylline & Phenobarbital** See Nephthalin pediatric Tab (Leeming)
- ALUDBOX** (Wyeth) Aluminum hydroxide gel 1 tsp w/magnesium hydroxide ¼ tsp Susp, Bot 12 fl oz, Tab, Bot 60s.
Use Antacid
- ALUDBOX SA** (Wyeth) Ambutanium bromide 25 mg butobarbital 8 mg, W/aluminum hydroxide and magnesium hydroxide/Tab or 5 cc Tab Bot 100s, Susp 12 fl oz
Use Hyperacidity
- ALUGEL.** (Morton & Converse) Aluminum hydroxide gel dried gr 10/Tab Bot 100s
Use Gastric antacid
- ALUOGIC PALATABS** (Central) Aluminum hydroxide 150 mg, mag trisilicate 60 mg, homatropine methylbromide 1 mg, pectin 25 mg benzoic acid 12 mg, opium 12 mg & Camphorated Tr of opium 0.3 ml/Tab Bot 100s 1000s
Use Antacid and antispasmodic
- ALUKALIN** (Activated kaolin) See Lysin improved (Maltbie)
- ALUKOIDS.** (Gold Leaf) Aluminum hydroxide dried gel colloidal kaolin, pectin aromatics Tab, Bot 1000s
Use Antacid
- ALULOTION SULFATHIAZOLE.** (Wyeth)
Kaolin in alumina gel w/sulfathiazole (5%) Bot of 2 fl oz
Use Pyogenic infection Applied topically
- ALUMASABS** (Gold Leaf) Aluminum hydroxide, dried gel 2 gr, magnesium trisilicate 4 gr kaolin 2 gr, phenobarbital ½ gr, atropine sulfate 1/300 gr Tab, Bot 100s, 500s, 1000s
Uter Antacid sedative & antispasmodic
- ALUMACABS** (Gold Leaf) Aluminum hydroxide, dried gel ½ gr calcium carb. 3½ gr, magnesium carb 2 gr & aromatic Tab, Bot 100s, 500s, 1000s
Use Antacid
- ALUMAPEC.** (Lemman) Aluminum hydroxide
• Generic name recognized by N.F., U.S.P. or A.M.A.
- 5 gr, magnesium trisilicate 5 gr, pectin ½ gr/Tab Bot 100s
Use Antacid
- W/Phenobarbital Sod ½ gr, & Atropine Sulfate 1/500 gr (Lemman)** Bot 100s
Use (2 4 q 4 h) Antacid with antispasmodic
- ALUMI-CREAM** (Invenex) Aluminum powder 10% or 20%, in a base of cad liver oil with vit A and vit D zinc oxide, starch lanolin sesame oil, alkyl dimethylbenzyl ammonium chloride Bot 1½ oz
Use Protects skin in colostomy, ileostomy
- ALUMINOID** (Chatham) Colloidal aluminum hydroxide 5 gr /Cap Box 48s
Use Gastric antacid
- ALUMINOSTOMY,** (Richards Pharm) Aluminum pow 18%, with zinc oxide and zinc stearate in a bland water repellent ointment Jar: 1 & 5 oz and 1 lb
Use: An occlusive dressing for skin protection; colostomies and ileostomies
- ALUMINIUM** See: Aluminostomy (Richards Pharm) Aluminum Paste USP, Ladd's Paste
- ALUMINIUM ACETATE,** See: Buro-Sol, cream & powder (Dook)
W/Chamoille powder See Chamo, Creme & Pow (Dome)
W/Hydrophobic Cream pH 4.2 See: Acid Mantle Creme (Dome)
- * **ALUMINIUM ACETATE SOLUTION, USP** (Various Mfr) See Buro Sol Pow Cont (Dook) Domebain, Pow (Dome Chem)
- ALUMINIUM ACETOTARTRATE,** (Mallinckrodt) Powder, Bot ¼ lb, 1 lb
Use Antiseptic and astringent on mucous membranes
- * **ALUMINIUM ACETYSALICYLATE, NF** See Aspirin Outlet (Abbott)
- ALUMINUM AMINOACETATE, DIHYDROXY,** See Dihydroxy aluminum aminoacetate
- ALUMINIUM ASPIRIN**
W/Acetophenetidin, caffeine, isobutylalyl barbituric acid See: Alatal, Tab (L L Drug)
- W/Chlorpheniramine maleate, Phenylephrine HCl See SINCO, Tab (Leyden)
- W/Codaine phosphate, papaverine HCl phenylephrine HCl chlorpheniramine maleate Vit C See Calrex Compound Caps (Bowell)
- W/Dexamethasone Orphenadrine HCl See Defenor Tab (Schering)
- W/Dihydrocodeine bitartrate acetophenetidin caffeine See Dracogest No 3 Tab (Carstana)
- W/Methapyrine fumarate, scopolamine HBr See Drowz Tabs (Person & Convey)

- W/Orphenadrine HCl, dexamethasone. See: Delenar, Tab. (Schering)
- W/Phenylephrine HCl, chlorpheniramine maleate, phenindamine tartrate. See: Dalca, Tabs. (B. F. Ascher)
- W/Phenylramidol. See: Analexin AF, Tab. (Irwin, Neisler)
- ALUMINUM CAPRYLATE** W/Phenylmercuric Acetate, Urea, Citric Acid, Boric Acid, Lactose, Dextrose. See: Mono-Serts, Tab. (Wintersmith).
- ALUMINUM CARBONATE BASIC, NND 1961.** See: Basaljel, Susp. (Wyeth)
- ALUMINUM CHLORHYDROXIDE COMPLEX.** See: Chlorhydral (Reheis)
- ALUMINUM DIHYDROXYAMINOACETATE.** See: Dihydroxy Aluminum Aminoacetate. (Various Mfr.)
- ALUMINUM DIHYDROXY SODIUM CARBONATE.** See: Roloids, Tabs. (American Chicle Co)
- ALUMINUM GLYCINATE, BASIC.** See: Dihydroxy Aluminum Aminoacetate, N.F.
- * **ALUMINUM HYDROXIDE GEL, U.S.P., N.N.D. 1961.** Aq. susp. of aluminum hydroxide that is equivalent of 3%-42% aluminum oxide.
- See: Alcid, Susp. (Maltbie).
Alakreen, Susp. (Chicago).
Al-U-Creme, Susp., N.N.D. (MacAllister)
Amphojel, Susp., N.N.D. (Wyeth)
Co-Lu-Gel, Susp. (Ulmer)
Creamalin, Susp., N.N.D. (Winthrop Labs)
Flugel, Susp. (Breon)
Gelumina, Susp. (American Pharm.)
Hartgel, Susp. (Hart)
Hydraxal, Susp. (Blue Line).
Vanogel, Susp. (Van Pelt & Brown)
- W/Amylolytic enzyme, proteolytic enzyme, cellulolytic enzyme See: Halazyme, Tabs. (Halca)
- W/Dexamethasone, acetylsalicylic acid, ascorbic acid See: Ropad, Tabs. (Pitman-Moore)
- W/Guar cellupectinoid, magnesium oxide, magnesium trisilicate. See: Belvis, Tabs (Carrick)
- W/Homatropine Methylbromide. See: Alcid HMB, Susp. (Maltbie).
Alodox, Susp. (Chicago Pharm.)
Alutropin, Susp. (Campbell).
Hartgel HMB, Susp. (Hart)
- W/Homatropine Methylbromide & Magnesium Trisilicate. See: Trigelma H.M., Susp. (Buffington).
- W/HMB and phenobarb. See: Hartgel HMB, Tab. (Hart)
- W/HMB 1/4 gr., phenobarb. 1/6 gr., kaolin 3 gr. See: Kalpen, Tab. (Cole)
- * Generic name recognized by N.F., U.S.P. or A.M.A.
- W/Kaolin. See: Koomagma, Susp. (Wyeth)
- W/Kaolin Col., Pectin, Aromatics. See: Katpac, Liq. (Wyeth)
Pectalum, Liq. (Smith, Miller & Patch)
- W/Magnesium Hydroxide. See: Aludrox, Susp. (Wyeth)
Bidrox, Susp. (McNeil)
Creamlets, Tab. (Winthrop Lab.)
Maalox, Tab. (Rorer)
- W/Magnesium Trisilicate. See: Alinese, Susp. (Haskell).
A.M.T., Susp. (Wyeth)
Endo-Magsol, Susp. (Endo).
Gelusil, Susp. (Warner Chilcott)
Magsol, Susp. (Endo)
Malcagel, Susp. (Upjohn).
Milk of Trinesium, Susp. (Abbott).
Tsi-Creamolate, Susp. (Winthrop Labs)
Tridroxal, Susp. (Warren-Teed).
Trigelma, Susp. (Buffington)
Trilax, Susp. (Blue Line).
Trisogel, Susp. (Lilly).
- W/Magnesium trisilicate, kaolin. See: Kao Magnin, Syr. (Sulliff & Case)
Silmocat, Cap & Tab. (Amfre)
- W/Magnesium trisilicate, red iron. See: Alnese, Susp. (Haskell)
- W/Paregoric & Bismuth Subgallate. See: Alpegora, Susp. (Merck Sharp & Dohme)
- W/Pectin, Magnesium Carb. & Saccharin. See: Alupac, Susp. (Pitman-Moore).
- W/Sulfur, silicon dioxide. See: Alsulfa, Cr. 1 lot. (Doak)
- * **ALUMINUM HYDROXIDE GEL, DRIED U.S.P., N.N.D 1961.** Tabs. U.S.P.
See: Altagel, Tab., N.N.R. (Lonlean).
Alugel, Tab. (Horton & Converse).
Aluminad, Cap. (Chatham).
Aluside, Tab. (Harco).
Amphojel, Tab. (Wyeth)
Creamalin, Cap. & Tab. (Winthrop Lab)
Co-Lu-Gel, Cap. & Tab. (Ulmer)
Hartgel, Tab. (Hart)
Hycala, Tab. (Columbus).
- Aluminum Hydroxide Gel Dried Combinations on pp. 30-33.**
- ALUMINUM HYDROXIDE GLYCINE.** See Dihydroxy aluminum aminoacetate.
- ALUMINUM HYDROXY MAGNESIUM AMINOACETATE.** See: Glycomine AM (Hormelin)
- ALUMINUM HYDROXIDE MAGNESIUM CARBONATE.** See: Res O-Gel (Sanford)
- ALUMINUM MAGNESIUM HYDROXYCARBONATE.** Used with Pranyl (Crookes-Barnes)
- * **ALUMINUM NICOTINATE.** See: Nicotex (Walker Labs.)

Aluminum Hydroxide Gel Dried Combinations

| Aluminum Hydroxide | Others | Manufacturer | Brand |
|--------------------|---|----------------------|------------------------------|
| 4 gr. | Adiphenin HCl 10 mg, Phenobarbital ½ gr., kaolin 2 gr. | Mallard | Altricate, Tab |
| 2 gr | Aminophylline ½ gr | Massengill | Aminadrox, Tab |
| 2.5 gr | Aminophylline 5 gr, ethyl aminobenzoate 0.5 gr | Irwin, Neisler | Cardalin, Tab |
| 2.5 gr, | Aminophylline 5 gr., ethyl aminobenzoate 0.5 gr., phenobarbital ½ gr | Irwin, Neisler | Cardalin Phen, Tab |
| ½ gr | Aminophylline 1 gr., ephedrine HCl 1/12 gr., phenobarbital ½ gr., ethyl aminobenzoate ½ gr | Irwin, Neisler | Dainite Pediatric, Tab. |
| 100 mg. | Ascorbic acid 50 mg, allantoin 60 mg | Kremers Urban | Allucee Cap |
| 75 mg. | Aspirin 325 mg, ascorbic acid 20 mg, prednisone 0.75 mg | Schering | Sigmogen, Tab |
| 75 mg | Aspirin 500 mg, dexamethasone 0.25 mg | Merck Sharp & Dohme | Decagastic, Tab |
| 65 mg | Aspirin 300 mg, Rauwolfia serpentina whole root 25 mg, ascorbic acid 25 mg | Flint Eaton | Rauylate, Cap |
| 2 gr | Atropine methylnitrate 1/120 gr., kaolin 7 gr. pectin 1 gr. beta lactose ½ gr | Strazenburgh | Alopectose w/Metropine, Tab |
| 2 gr | Atropine sulfate 1/300 gr., kaolin 2 gr., Mag trisilicate 4 gr., phenobarbital ½ gr | Gold Leaf | Alumobarbs, Tab |
| 75 mg | Atropilgite activated 90 mg, peetobarbital sodium 7.5 mg, 2 diethylaminoethyl diphenylacetate HCl 250 mg. | Palmedica | Spasmasorb, Tab |
| 250 mg | Belladonna alkaloids ½ gr., Mag trisilicate 250 mg., benzocaine 8 mg., phenobarbital 8 mg | Walker | Alkasposmol, Tab |
| 5 gr | Belladonna ext 1/16 gr., Mag trisilicate 2½ gr. | Upjohn | Malcatabs w/Belladonna, Tab. |
| 0.15 Gm | Belladonna ext 5 mg., Mag trisilicate 0.3 Gm. | Rorer | Triolidin, Tab |
| 5 gr | Belladonna ext 1/46 gr., phenobarbital ½ gr. | Hoeck | Alubelop, Tab |
| 5 gr | Belladonna ext ½ gr phenobarbital ½ gr | Wyeth | Sebella, Tab. |
| 2½ gr | Belladonna ext 1/24 gr., Cal carbonate 6 gr., chlorophyllin 1% | Warren Teed | W T, Tab. |
| 250 mg | Butalbarital Sod 10 mg., methscopolamine bromide 2.5 mg., Mag trisilicate 250 mg | Staff Research Labs. | Eulctin, Tab |
| 100 mg | Butazolodin 100 mg., magnesium trisilicate 150 mg., homatropine methylbromide 1.25 mg | Geigy | Butazolodin Alko, Cop. |
| ½ gr | Cal carbonate 3½ gr., Mag carbonate 2 gr | Gold Leaf | Alumcarb, Tab |
| 2 gr | Cal carbonate 1 gr., Mag trisilicate 4 gr. | Wallow | Alcawal, Tab |
| 1½ gr | Carboxymethylcellulose Sod 3½ gr | Superior Pharmacal | Cal Luna, Tab. |
| 120 mg | Cortisone acetate 2.5 mg., salicylate 300 mg., Cal ascorbate 60 mg., Cal carb 60 mg | Massengill | Solcart, Tab. |

Aluminum Hydroxide Gel Dried Combinations

| Aluminum Hydroxide | Others | Manufacturer | Brand |
|--------------------|--|----------------------|---------------------------|
| 0.9 Gm. | Ergaine 0.45 Gm., resorcinol 0.1 Gm., Mag. trisilicate 0.15 Gm., Cal. carbonate 0.105 Gm., Mag. carbonate 0.06 Gm. | Organon | Trevidal, Tab. |
| 250 mg. | Gastric mucin 160 mg., magnesium glycinate 250 mg. | Barrows Chemical Co. | Mucogel, Tab. |
| 250 mg. | Gastric mucin 160 mg., mag. glycolate 250 mg., phenobarbital 6 mg., belladonna ext., 5 mg. | Barrows Chemical Co. | Mucogel-8, Tab. |
| 0.25 Gm. | Gastric mucin 0.16 Gm., Mag. trisilicate 0.45 Gm. | Harrow | Mucatin, Tab. & Liq. |
| 240 mg. | Gastric mucin 120 mg., magnesium trisilicate 360 mg., hama-irap'ne methylbromide 6.2 mg. | Maney | Gast-Rex, Tab. |
| 4 gr. | Gastric mucin 2½ gr. Mag. trisilicate 7 gr., homatropine methylbromide 1/85 gr. | Harrower | Muco-Seth, Tab. |
| 4 gr. | Gastric mucin 1 gr., Mag. trisilicate 2 gr., homatropine methylbromide 1.2 mg. phenobarbital ¼ gr. | Lemman | Antosed, Tab. |
| 2½ gr. | Homatropine methylbromide 1/48 gr., Mag. trisilicate 2½ gr. | Buffington | Trigelmo H M, Tab. & Liq. |
| 375 mg. | Homatropine methylbromide 2.5 mg., Mag. trisilicate 125 mg. | Crales | Trapasil, Tab. |
| 300 mg. | Homatropine methylbromide 2.5 mg., Mag. trisilicate 120 mg., pectin 50 mg. | Central | Alucon, Tab. |
| 2 gr. | Homatropine methylbromide 1/48 gr. Mag. trisilicate 3 gr., Mag. hydroxide 1 gr. & pectin ½ gr. | Walker | Gel-A-Nutrin, Tab. |
| 129.6 mg. | Homatropine methylbromide 2.5 mg., magnesium trisilicate 194.4 mg., phenobarbital 8 mg. | Blue Line | Neutrotabs, Tab. |
| 3 gr. | Homatropine methylbromide 1/150 gr., phenobarbital ½ gr. | Irwin, Nelster | Homatropal, Tab. |
| 7½ gr. | Homatropine methylbromide 1/24 gr., phenobarbital ¼ gr., Mag. trisilicate 2½ gr. | Rosmar | Alased, Tab. |
| 100 mg. | Homatropine methyl bromide 1.25 mg., prednisone 1.25 mg., phenylbutazone 50 mg., magnesium trisilicate 150 mg. | Geigy | Sterozolidin, Cap. |
| 100 mg. | Hydrocarisone 1.5 mg., aspirin 5 gr. | Mallard | Aceta-Cart, Tab. |
| 24 Gm. | Koolin 24 Gm., Cal. carbonate 76 Gm., bifs. subcarbonate 16 Gm., acerola, sod. chloride, dextrose | Lilly | Alamin, Pow. per 100 Gm. |
| 1 gr. | Kaolin 8 gr., pectin ½ gr. | Gald Leaf | Alukaids, Tab. |
| ½ gr. | Mag. carbonate 2 gr., Cal. carbonate 3½ gr. | Gold Leaf | Alumacarb, Tab. |
| — | Mag., carbonate, Col. carbonate, Taka Diastase | Parke, Davis | Antacid, Tab. |
| 10 gr. | Mag. carbonate 1¼ gr., pectin 2½ gr. | Pitman-Moore | Alupez, Tab. & Liq. |
| 1 tsp. | Mag. hydroxide ¼ tsp. | Wyeth | Aludrox, Susp. & Tab. |

Aluminum Hydroxide Gel Dried Combinations

| Aluminum Hydroxide | Others | Manufacturer | Brand |
|--------------------|--|-----------------|----------------------------|
| Present | W/magnesium carbonate 5 gr., hyoscyamine sulf 0.0319 mg, phenobarbital 31 mg., atropine sulf 0.0097 mg, hyoscyne hydrobromide 0.00325 mg | Benton | Almodon, Tab |
| 400 mg | Magnesium hydroxide 100 mg., scopolamine HBr 0.65 mg | Columbus | Hycadrox Tab |
| 200 mg | 1 hyoscyamine HBr 0.125 mg., butylscopolamine methylsulfate 1 mg. | Reed & Carnrick | Modulof, Tab |
| 25% | Magnesium hydroxide 200 mg., scopolamine methylsulfate 50% | Sherman | Acaralum, Pow |
| 260 mg | Mag. oxide 5%, acacia 20%, vegetable mucin 50% | Central Midland | Neutrolon CompoSite, Tab. |
| 324 mg | Magnesium trisulfate 488 mg | Winthrop Labs | Tri Creamale, Tab. |
| 162 mg | Mag trisulfate 162 mg | G F Harvey | Merdoxine, Tab. |
| 7½ gr. | Mag trisulfate 7½ gr | Durg Products | Almag, Tab |
| 10 gr. | Mag trisulfate 1½ gr | Endo | Melcolabs, Tab |
| 10 gr. | Mag trisulfate 5 gr | Upjohn | Trisogel, Tab & Liq. |
| 1½ gr | Mag trisulfate 4½ gr | Buffington | Trigelma, Tab & Liq. |
| 2½ gr | Mag trisulfate 2½ gr | Wyeth | A M T, Tab |
| 2.5 gr. | Mag trisulfate 3.85 gr | tonnell | Silmogel, Tab. |
| 3 gr | Mag trisulfate 6 gr., ascorbic acid 20 mg | Massengill | Cimodrox, Tab |
| 2 gr. | Mag trisulfate 4 gr., cal carbonate 1 gr. | Wallou | Aicarval, Tab |
| 4 gr. | Mag trisulfate 4 gr., chlorophyllin 25 mg | Walker | Chlorogel, Tab |
| 0.25 Gm | Mag trisulfate 0.45 Gm., gossyp mucin 0.16 Gm | Harrow | Mucolin, Tab |
| 375 mg. | Mag trisulfate 125 mg., homatropine methylbromide 2.5 mg | Crookes | Trapatul, Tab |
| 4 gr | Mag trisulfate 7 gr., gastric mucin 2 Gm. | Winthrop Labs. | Trimucolon, Tab |
| 2½ gr. | Mag trisulfate 2½ gr., homatropine methylbromide 1/48 gr. | Buffington | Trigelma H.M., Tab & Liq |
| 2 gr. | Mag trisulfate 5 gr., homatropine methylbromide 1/60 gr., kaolin 2 gr., methyl cellulose 1½ gr | Satliff & Case | Magnal, Tab. |
| 7 gr | Mag trisulfate 1½ gr., reduced iron ¼ gr. | Haskell | Alinese, Tab |
| 2 gr | Mag trisulfate 4 gr., kaolin 2 gr | Massengill | Kamodrox, Pow. & Tab. |
| 2 gr. | Mag trisulfate 4 gr., kaolin 2 gr | Elder | Magkaolin, Tab. |
| 25% | Mag trisulfate 50%, kaolin 25% | Amire | Silmocol, Cap., Tab & Pow. |
| 4 gr. | Mag trisulfate 7 gr., methylcellulose 1 gr | Hack | Neosorb, Tab & Susp. |
| 4 gr | Mag trisulfate 7½ gr., pancreatin troyoz ¼ gr. | Bobit | Gelozyme, Tab |

Aluminum Hydroxide Gel Dried Combinations

| Aluminum Hydroxide | Others | Manufacturer | Brand |
|--------------------|--|---------------------|-----------------------------------|
| 5 gr. | Mag. trisilicate 5 gr., pectin 1/2 gr. Mag. trisilicate, peppermint oil | Lehman Harvey | Alumopoc, Tab. Merdosine, Tab. |
| Present | W/Prednisolone 2.5 mg. or 5 mg. and magnesium trisilicate | Merck Sharp & Dohme | Co-Hydraltra, Tab. |
| Present | W/Prednisone 2.5 or 5 mg. and magnesium trisilicate. | Merck Sharp & Dohme | Co-Delltra, Tab. |
| 3 gr. | Mag. trisilicate 6 gr., Vit. C 20 mg. | Messengill | Cinadrox, Tab. |
| 2 1/2 gr. | Mag. trisilicate 5 gr., methoprine 1 mg., duodenum 1/4 gr. | Sirosenburgh | Metrocin, Tab. |
| 2 gr. | Mag. trisilicate 4 gr., phenobarbital 1/8 gr., atropine sulfate 1/600 gr., kaolin 2 gr. | Gold Leaf | Alumabarba, Tab. |
| 1 gr. | Mephensin 3 gr., Sod. sal. 5 gr., secobarbital 1/4 gr., phenobarbital 1/8 gr., Vit. C 5/6 gr. | Geriatric | Gerisal, Tab. |
| 200 mg. | Meprabamate 200 mg., prednisolone 1.0 or 2.0 mg. | Merck Sharp & Dohme | Mepralone, No. 1 or 2, Tab. |
| 2 gr. | Metiprin 1 mg., kaolin 1 1/2 gr., b-lactose 1 gr., pectin 1 gr. | Sirosenburgh | Metropectin, Tab. |
| 2 gr. | Pectin 1 gr., kaolin 7 gr., b-lactose 1 gr. | Sirosenburgh | Alopectase, Tab. |
| 150 mg. | Phenylbutazone 50 mg., prednisone 1.25 mg., magnesium trisilicate 150 mg., homotrope methylobromide 1.25 mg. | Gelgy | Sterozoline, Cap. |
| 200 mg. | Prednisolone 2.0 mg., aspirin 300 mg., sod. ascorbate 60 mg. | Merck Sharp & Dohme | Tempogen Forte, Tab. |
| 250 mg. | Procaine HCl 150 mg., eminoxystropine tropate HCl 1/260 gr., pentobarbital Sod. 1/6 gr., phenobarbital 1/6 gr., epoxy-aminoxystropine tropate HBr 1/260 gr., Vit. B-6 10 mg. | Narrower | Formalit, Tab. |
| 5 mg. | Propenpyridamine maleate 10 mg., scopolamine hydrobromide 0.15 mg., ethyl aminobenzoate 1 mg. | Davis & Sly | Doptren, Tab. |
| 120 mg | Salicylamide 250 mg., pyrilamine maleate 12 mg., phenylephrine HCl 5 mg., acetphenethdin 120 mg., caffeine 15 mg. | Central | Cenogestic, Tab. |
| 75 mg. | Salicylamide 325 mg., trimecetonone 0.5 mg., Vit. C 20 mg. | Lederle | Aristogestic, Cap. |
| 240 mg. | Scopolamine methylobromide 2 mg., mag. hydroxide 60 mg. | Elder | Amproxa, No. 293 Tab. |
| 125 mg. | Scopolamine methyl nitrate 1 mg., Mag. carb. 12.5 mg., Sod. dodecyl sulfate 12.5 mg. | Sirosenburgh | Skopalate P.A.M., Cap. |
| 300 mg. | Scopolamine methylobromide 2.5 mg., magnesium trisilicate 150 mg. | Buffington's | Trigelma w/Mescosime, Tab. |
| 1 Gm. | Sippylic 32.77 Gm., Mag. trisilicate 2 Gm., W/Vitamins | Purdue Frederick | Sippylics, Pow. / 8 tsp. |
| 2 1/4 gr. | Syntropan 3/40 gr., Col. carbonate 1.1 gr., Mag. peroxide 1 1/2 gr. | Rache | Syntrogel, Tab. |
| 1.5 Gm. | Vitamins 0.2 mg., minerals 1/2 gr. | Harrower | Colalain, Tab. |
| 5 gr. | Vitamins 0.2 mg., minerals 1/2 gr. | Hort | Hortgel H.M-3, Tab. |

- ALUMINUM OXYQUINOLATE** (Chemo Pura)
ALUMINUM PENICILLIN, NND 1962, ADR 1955 (Hynson Westcott & Dunning)
 Aluminum penicillin 50,000 units & Sod benzoate 0.3 Gm/Tab Pkg 12s
 Use Oral penicillin
- * **ALUMINUM PHOSPHATE GEL**, NF
 NND 1962 Aluminum phosphate 4.15%
 See Phosphaljel Susp (Wyeth)
- ALUMINUM PROTEINATE** W/Magnesium glycinate See Malumel (Rawell)
- ALUMINUM SALT OF THE AMINO ACID GLYCINE** See Dihydroxy aluminum amino acetate Prep (Various Mfr)
- ALUMINUM SILICATE SODIUM** See Ludoxon Tab (Schering)
- * **ALUMINUM SUBACETATE SOLUTION**, USP (Various Mfr) Aluminum acetate basic solution
 Use As an astringent wash
 W/Eucalyptol, Corbolic Acid Zinc Oxide, & Ichthyol See Azocol, Oint (Patch)
 W/Phenol zinc oxide ichthyol eucalyptol See Azocol Oint (Patch)
- ALUPEC LIQUID** (Pitman Moore) Aluminum hydroxide gel (Al₂O₃ gr 18) pectin gr 4 magnesium carb 4.6 gr saccharin sol 1/4 gr and preser/ft oz Bot 4 oz, 12 oz
 Use Gastric antacid
- ALUPEC TABLET**, (Pitman Moore) Aluminum hydroxide gel 10 gr, pectin 2 1/2 gr, magnesium carb 1 1/4 gr/Tab Bot 60s, 500s
 Use Gastric antacid
- ALUPINAL**, (Maltbie) Atropine sulfate 1/1000 gr aluxalin 7 1/2 gr phenobarbital 1/2 gr/Tab Bot 100s, 1000s, 5000s
 Use (1 2 tid) Antispasmodic & antacid
- ALURATE** (Roche Lab) Apobarbital NF Allylisopropylbarbituric acid
- ALURATE SODIUM**, NNR 1955 (Roche Lab)
- ALURATE VERDUM ELIXIR**, NNR 1955 (Roche Lab) Both elixir alurate verdum (green) and elixir alurate (red) provide the same amount of alurate (40 mg) of allylisopropylbarbituric acid per tsp Bot 6 oz 16 oz & 1 gal
 Use (1 dr q 3-4 h) Sedative, hypnotic
 W/Acetaphenetidin See Allanal Tab (Roche Lab)
- ALU SCOP**, (Lloyd, Dabney, & Westerfield) Methscopolamine bromide 1.25 mg, dicycloxyaluminum aminoacetate 450 mg, sed lauryl sulf 20 mg, magnesium hydroxide 37.5 mg/cap or 7.5 cc Cap Bot 100s, 500s
 * Generic name recognized by NF, USP or AMA
- Use: Peptic ulcer and gastro intestinal distress
- ALUTED** (Hart) External coal contains isoproterenol 5 mg; internal coal contains theophylline 2 gr, ephedrine 1/2 gr, secobarbital 1/2 gr/Tab Bot 24s, 100s Also available in half strength for pediatric use
 Use (1 2 qid) Treatment of asthmatic patients
- AUTROPIN** (Campbell) Homatropine methylbromide 1/100 gr/4 cc of aluminum hydroxide gel Bot 4 oz, 12 oz
 Use: (1 2 dr q 3 h) Antispasmodic & antacid
- AUTROPIN FORTE**, (Campbell) Homatropine methylbromide 5 mg in 4 cc aluminum hydroxide gel Bot 4 oz
 Use (1 tsp tid pc hs) Antispasmodic & antacid
- AUXIDE** (Harco) Aluminum hydroxide gel, dried Tab 10 gr Bot 100s, 1000s
 Use Antacid
- ALVAGEL** (Kenyon) 55% Aloe vera active principle in oint base Tube 1/2 oz, 1 oz Jar 4 oz 1 lb
 Use Topically, for sunburn eczema, dermatitis x ray & radiation burns
- ALVININE** (Wampole) Biphenamine HCl 1% aleyl alcohol 1% w/inethonolamine lauryl sulfate 1 2 propionediol, lauric acid diethanolamide Bot 4 oz, squeeze
 Use Dandruff and seborrhea
- ALVODINE** (Winthrop Labs) Piminoadine ethanesulfonate Ethyl 4 phenyl 1 [(3 phenylamino)propyl] piperidine 4 carboxylate or ethanesulfonate (20 mg/cc) Amp 1 cc, Box 25s, 100s Tabs 50 mg, Bot 100s, 500s
 Use Analgesic
- ALYSINE** (Merrell)
 Elixir—Natural Sod salicylate 2.3 Gm, Pot bicarb 3.3 Gm Sod citrate 1.1 Gm in an aromatized elixir with 10% alcohol/30 cc Bot 1 pt & 1 gal
 Powder—Natural Sod salicylate 2.46 Gm, natural magnesium salicylate 3.44 Gm, natural cal salicylate 3.44 Gm, Sod bicarb 6.15 Gm, calcium carb 6.15 Gm, magnesium carb 6.15 Gm (95 gr)/30 Gm Bot 1 oz, 1/4 lb & 1 lb
 Use: Analgesic & antirheumatic
- AEZINOX**, NND 1962 (Smith, Miller & Patch) Dihydroxy aluminum aminoacetate 77 gr/Tab Bot 100s 500s
 Use Hyperacidity and peptic ulcer
 W/Phenobarbital and Hamatropine Methyl Bromide (Smith, Miller & Patch) Dihydroxy aluminum aminoacetate 77 gr, phenobarbital 16.2 mg, hamatropine methyl bromide 0.65 mg/Tab Bot 100s 500s

ALZINOX MAGMA, N.N.D. 1962. (Smith, Miller & Patch) Dihydroxy aluminum aminoacetate 7.7 gr./5 cc. Bot. 8 fl. oz.

Use: Hyperacidity and peptic ulcer

W/Phenobarbital and Hamatropine Methyl Bromide. (Patch) Dihydroxy aluminum aminoacetate 7.7 gr., phenobarbital 8.1 mg., hamatropine methyl bromide 0.65 mg./5 cc. Bot. 8 fl. oz.

AMADIL. (Breon) N-Acetyl-p-aminophenol 300 mg./Tab. Bot. 100

Use: Analgesic and antipyretic.

* **AMANOZINE NCL**. 2-amino-4-anilino-5-triazine HCl.

AMARSAN. See. Acetarsone, Tab. (Abbott)
AMATEEN CONCENTRATE. (Dumas-Wilson) Protein 70%, total N 11%, amino N 5.6%, with Vit. B complex, ash 7.2%. Can 8 oz., 1 lb.

Use: Protein hydrolysate therapy.

AMAYITE, (United Specialties) Vit. A 7500 U.S.P. u., B-1 2.4 mg., B-2 2.7 mg., B-6 3 mg., B-12 10 mcg., B-15 5 mg., C 110 mg., D 400 U.S.P. u., niacin 10 mg., niacinamide 10 mg., iron 15 mg., calcium carbonate 200 mg., folic acid 0.3 mg./Tab. Bot. 60s, 250s, 500s

Use: Mixed vitamin deficiencies.

AMBAR. (Robins) Methamphetamine HCl 3.33 mg., phenobarbital 1/2 gr./tab. Bot. 100s, 300s.

Use: Tension-anxiety and fatigue states

AMBAR No. 1 EXTENTARS. (Robins) Methamphetamine HCl 10 mg., phenobarbital 1 gr./Extentab. 1/2 released for immediate action, 1/2 released gradually. Bot. 100s.

Use: Tension-anxiety and fatigue states.

AMBAR No. 2 EXTENTARS. (Robins) Methamphetamine 15 mg., phenobarbital 1 gr./Tab. Bot. 100s.

AMATEEN CONC. (Dumas-Wilson) Protein hydrolysate from yeast Pkg. 1#

* **AMAZONE**. 1,4-Benzoquinone amidinohydrazone thiosemicarbazone hydrate Iversol

* **AMBENONIUM CHLORIDE**, N.N.O. 1962. N,N'-bis-(2-diethylaminoethyl) oxamide bis-2-chlorobenzyl chloride See Mytelase, Tab. (Winthrop Labs)

AMBENYL EXPECTORANT. (Parke, Davis) Ambadryl HCl 24 mg., Benadryl HCl 56 mg., codeine sulf. 3 gr., ammonium chloride 8 gr., Pal. quatacal sulfonate 8 gr., menthol q.s., alcohol 5% /oz. Bot. 1 pt., 1 gal.

Use: (1-2 tsp. q. 3-4 h) Cough.

AMERLITS. (Rahm & Haas) Amberlite resins
Use: Gastric acidity.

* Generic name recognized by N.F., U.S.P. or A.M.A.

AMBITAL. (Lemmon) Aminophylline 1 1/2 gr., phenobarbital 1/4 gr., aluminum hydroxide 2 gr., cal. carbamate 3/4 gr./Tab. Bot. 100s
Use: (1-2 t.i.d.) Sedative vasodilator.

AMBODEX TIMED CAPSULE. (Testagar) d-Amphetamine HCl 15 mg., amobarbital 60 mg./Tab. yellow & brown. Bot. 100s, 1000s.

Use: (1 in morning) Restores emotional ability.

AMBODRYL HYDROCHLORIDE. (Parke, Davis) (Bromazine) Bromadiphenhydramine HCl Kapsels (25 mg.) Bot. 100s Elixir 10 mg./4 cc. Bot. 1 pt. Steri-Vials 5 mg./cc 10 cc. (N.V. or I.M.) Pow. Rot 1/2 oz.

Use: (1 q. 3-4 h.) Antihistaminic agent.

W/Ambadryl HCl, dihydrocodeinone, ammonium chloride & other cough agents. See: Ambenyl, Syr. (Parke, Davis).

AMBROL. (Smith, Miller & Patch) Secobarbital 50 mg., phenobarbital 50 mg., aspirin 195 mg., Vit. B-1 5 mg./Tab. Yellow. Bot. 100s.

Use: (1 h.s.) Sedative & hypnotic.

* **AMBOMYCIN**. (Pfizer) Antibiotic isolated from both filtrates of *Streptomyces ambosiensis*.

Use: Anti-neoplastic agent.

* **AMRUCETAMIDE**, 2-(di-n-butylamino)-2-(p-methoxyphenyl)acetamide

AMRUSAL. (Lemmon) Salicylamide 0.23 Gm., acetophenetidin 0.15 Gm., di-desoxyephedrine HCl 2 mg., ascorbic acid 15 mg./Tab. Bot. 100s

Use: Pain and depression.

* **AMRUTONIUM BROMIDE**. (Wyeth) Ethyldimethylammonium {3-Carbonyl-3,3-diphenyl propyl} ethyl dimethylammonium bromide. 10 mg./Tab. Bot. 100s.

Use: Ulcer therapy.

W/Rutobarbital, Aludrox. See: Aludrox SA (Wyeth)

* **AMBUTOXATE HYDROCHLORIDE**. 2-diethylaminoethyl-4-amino-2-butoxybenzoate HCl. See: Symlocaine (Winthrop Lab.)

AMCHLOR. (Brewer) Ammonium chloride Tab. E.C. (15 gr.). Bot. 100s, 500s, 1000s.

Use: Diuresis in cardiac edema

AMCORESE DAKAPS. (Amco Drug) Thyroid 90 mg., levo-amphetamine sulfate 15 mg., hamatropine methylbromide 0.72 mg., acetophenetidin 5 mg., mephobarbital 15 mg./Cap. Bot. 100s

AMDELATE. (Abbott) Ammonium mandelate Elixir—(8 Gm./fl. oz.) Bot. 1 pt., 1 gal. Syr.—(12 Gm./fl. oz.) Bot. 1 pt., 1 gal.

Use (1 dr. t.i.d.) Urinary antiseptic.

AM-DEX. (Superior) d-Amphetamine sulf. 5 mg./Tab. Bot. 100s and 1000s

AMDIL (Breon) N Acetyl p Aminophenol
300 mg / Tab Bot 100s
Use: Pain and reduction of fever

AMDROLIN INJ (Lincoln) Sal in oil 25 50
or 100 mg testosterone propionate per cc
in physiological saline Sal in Oil; 25 mg /
cc 50 mg / cc 100 mg / cc—10 cc Vial
Aqueous Susp 25 mg / cc 50 mg / cc,
100 mg / cc—10 cc Vial
Use: Androgen

AMEBACIDES

See Acetarsone NF

Anayodin Pills (Ams)
Balorsen Tab (Endo)
Camalorm Tab (Parke Davis)
Carbarsone Paw & Tab (Lilly)
Chinacal, Tab (Various Mfr)
Diadahydroxyquin, USP
Diadaquin, Tab (Searle)
Emeline HCl USP
Entero Violum, Tab (Ciba)
Fumid I (Abbott)
Glurubin, Tab (Messengill)
Milibis, Tab (Winthrop Labs)
Milibis Arolen, Tab (Winthrop Labs)
Moebiquin Tab (Consolidated Mid Co)
Quinoxyl, Tab (Winthrop Labs)
Thiocarbarsone (Lilly)
Treparsil Tab (Wollau)
Violum (Ciba)

AMEBAN See Carbarsone (Various Mfr)

AMECHOL See Methocholine Chloride

AMEDARB (Smith Miller & Polch) Amino
phylne 1 1/2 gr ephedrine sulfate 3/4 gr
phenobarbital 1/2 gr / Tab Bot 1000s
Use: (111d) Sedative ant spasmotic

AMEND S SOLUTION (Leeming) Organic
protein bound potassium free iodine equiv
alent to 14%, iodine Sol 2 oz
Use: Iodine therapy

AMERCID (Messengill) Codeine phosphate
1/12 gr, emmanium chloride antimony
Pot Iartrete Tolu balsam glycerin citric
acid Liq Bot 1 pl 1 gal
Use (1/2 dr q 2 h) Narcotic cough prepa-
ration

**AMERICAINE CLEAR OINTMENT, LIQUID
AND AEROSOL SPRAY.** (Arnor Stone)
Benzocaine 20% and benzethonium chlo-
ride 0.1% in bland water soluble vehicle
Oint 1 oz tube with rectal applicator 1/2
lb Jars 1 lb, 7 lb Liquid Bot 4 oz, Aero
3 5.3 & 11 oz
Use: Surface pain & itching

**AMERICAINE WITH CHLOROPHYLL OINT-
MENT.** (Arnor Stone) Same as "Clear" with
0.16% chlorophyll added Tube with rectal
applicator 1 oz Jar 1/2, 1 & 7 lb
Use: Surface pain & itching

AMERICAINE SUPPOSITORIES. (Arnor Stone)
Benzocaine 20% benzethonium chloride
0.1% / suppos Box 12s
Use: Topical anesthetic

**AMERICAINE INTRATRACHEAL LUBRI-
CANT.** (Amer Hosp Supply) Oint 20%
benzocaine Tube 1 oz

AMERICAINE-NEOMYCIN. (Arnor Stone)
Benzocaine 20%, neomycin sulfate 5 mg /
Gm Oint Tube 1/2 oz
Use: Topically, anesthetic and combat in-
fection

AMERTAN. (Lilly) (Tonic Acid Compound)
5% Tonic acid w/Merthiolate 1:5000
Jelly, Tube 1 oz, 12s, 5 oz, 12s, Jar
1 lb

Use: Antiseptic protective in burns
AMESEC. (Lilly) Aminophylline 2 gr, ephed-
rine HCl 3/4 gr, omytal 3/4 gr Enseals Bot
100s, 500s, 1000s Pulvules; Bot 100s, 500s
Use: Nocturnal asthma & allergic mani-
festations

AMETHOCAINE HYDROCHLORIDE, USP.
NND 1961, ADR See: Pontocaine HCl
(Winthrop Labs)

**AMETHONE HYDROCHLORIDE, NND
1962** (Abbott) (Amalonone HCl) 3 Beta
diethylaminoethyl 3-phenyl-2 benzofuran-
one hydrochloride
Concentrated—9%, Abba-Vial 20 cc
Use: Urologic anesthetic

AMETRIFON GERMICIDAL DETERGENT.
(Meyer) Benzethonium chloride 2.5% in a
solution containing a nonionic detergent
Liq Bot 8 oz, 1 gal
Use: Antiseptic cleansing agent

AMFADROL (Columbus) Methyl androstene-
diol 1 mg, diamphetamine sulf 2.5 mg /
Tab Bot 100s, 1000s
Use: Premature aging and senility

AMFEBARBS. (Gold Leaf) Amphetamine
sulfate, rac, 5 mg, phenobarbital 1/4 gr /
Tab Bot 100s, 1000s
Use: To restore emotional equilibrium

d-AMFETASUL, NND 1962 (Pitman Moore)
d-Amphetamine sulfate Tab (5 mg) Bot
100s, 1000s
Use: Central nervous system stimulant

AMFREXIN. (Amfre) Sod lauryl sulfate &
papain, acidulated to a pH of 4.0 Pkg
4 oz, 8 oz
Uses: Douche (1 tsp of paw / qt of water)

AMDESIC. (Kramers Urban) N acetyl p
aminophenol 2 1/2 gr, chlorphenopyrida-
mine maleate 1/30 gr, noscapine 1/30 gr,
1 hyoscyamine sulfate 1/1000 gr, aspirin
3 1/2 gr, caffeine 1/4 gr / Tab Bot 25s, 100s
Use: Symptomatic treatment of colds

AMGESIC NO 2. (Kramers Urban) Phenyl-
ephine HCl, 10 mg; N acetyl p amino-

phenol, 160 mg.; chlorprophenpyridamine maleate, 2 mg.; 1-hyascyamine sulfate, 0.0625 mg.; acetylsalicylic acid, 230 mg.; and caffeine, 15 mg./Tab. Bat. 25s, 100s. Use: Common cold.

AMIBARSON, See: Carbarsone. (Various Mfr.).

AMI-CAL (First Texas Pharm., Inc.) Vit. A—Palmitate 1000 U, Vit. D 400 U, R-12 Crystallin 2 Mcgm., calcium gluconate 200 mg., ferrous gluconate 20 mg., thiamin HCl 1 mg., riboflavin 0.5 mg., calcium ascorbate 50 mg., L-lysine 150 mg., niacinamide 25 mg./Cap. Bat. 1000s; apothecary jars 100s. Use: Nutritional supplement to pregnant women.

AMIDOFEBRIN, See: Aminopyrine. (Various Mfr.).

AMIDONE HYDROCHLORIDE, See: Methadone Hydrochloride. (Various Mfr.).

AMIDOPHEN, (Lilly) Aminopyrine 3½ gr., acetophenoidin 1 gr., caffeine ½ gr., hyascyamus ext. ¼ gr./pulvule. Bat. 100s, 500s.

Use: [1 q. 3-4 h.] Antipyretic analgesic.

* **AMIDOTRIZOATE, SODIUM**, See: Diatrizoate Sodium.

AMIDOPYRAZOLINE, See: Aminopyrine (Various Mfr.).

AMIDOPYRINE, N.N.R. 1949, A.D.R. See: Aminopyrine & Pyramidan. (Various Mfr.)

AMIDOXINE, (Mallard) Pyridoxine HCl 100 mg., thiamine HCl 50 mg. In saline solution/cc. Vial 10 cc., 5s, 10s, 25s.

Use: I.M., nausea & vomiting of pregnancy.

AMIDRYL, See: 2-(Benzhydryloxy)-N,N-dimethylethylamine hydrochloride.

AMID-SAL, (Glenwood) Salicylamide. Tab. (3½ gr.) Rot. 100s.

Use: (1-2 i.i.d.) Analgesic.

AMIFEDS, (Gold Leaf) Aminophylline 1½ gr., ephedrine sulfate ¾ gr., phenobarbital ½ gr./Tab. Bat. 100s, 1000s.

Use: [1 p. 3 h., p.r.n.] Sedative diuretic.

AMIGEN, N.N.D. 1962, (Mead Johnson) Protein hydrolysate enzymatic hydrolysate of casein. Amigen 5%, dextrose 5%, in 125, 500, and 1000 cc.; amigen 5%, dextrose 10% in 1000 cc.; amigen 5%, levupen 10% in 1000 cc.; amigen 3½%, dextrose 3½% in ½ lactated Ringer's solution in 250 cc. Bask; amigen 5%, dextrose 5% and alcohol 5% in 1000 cc.

Use: Provides protein nutrients.

AMINARSON, See: Carbarsone. (Various Mfr.).

AMINE OXIDASE INHIBITOR

See: Catran, Tab. (Lakeside)

Marplan, Tab. (Roche Labs)

* Generic name recognized by N.F., U.S.P. or A.J.A.

Marsilid, Tab. (Eoche Labs.)

Nardil, Tab. (Warner-Chilcott)

Niamid, Tab. (Pfizer)

AMINE RESIN, See: Polyamine Methylene Resin.

AMINET SUPPOSITORIES, (Ames) Aminophylline 0.5 Gm., pentobarbital Sod. 0.1 Gm., benzocaine 0.06 Gm. in the new Bischoff base. Also in quarter and half strength for children and prophylaxis, Box 12s.

Use: Bronchial asthma.

AMINICOTIN, See: Nicotinamide. (Various Mfr.).

AMINITE, (Mallard) Aminophylline ½ gr., veratrum viride ext. 1 gr., acanite pow. 1/5 gr., atropine sulfate 1/1000 gr./Tab. Rot. 100s, 500s, 1000s.

Use: Hypertension.

* **AMINITROZOLE**, 2-Acetylamino-5-nitrothiazole. See: Trithean, Tab. (Ortho)

* **AMINOACETIC ACID, N.F., N.N.E. 1941**, (Various Mfr.) (Glycine, glycoyl) available as elixir, pow., & tab.

See: Glycine Solv. (Abb).

GlycoElixir (Squibb).

Use: Myasthenia gravis.

W/Calcium carbonate, Aluminum hydroxide, magnesium carbonate. See: Glytabs, Tabs. (Deseret)

W/L-lysine. See: Lixocal Elix. (Lanett) Lycal Elix. (Konnelt)

W/Phenylephrine HCl pyrilamine maleate, acetylsalicylic acid, caffeine. See: Al-Ay, Tabs (Bowman)

W/Vit. B-1, B-2, niacin. See: Amothile, Elixir. (Harvey)

AMINOACETIC ACID & CALCIUM CARBONATE, N.N.R. 1955. See: Titrolac, Liq. & Tab. (Schenley).

AMINOACID & PROTEIN PREP.

See: Aminoacetic Acid, N.F.

Arginine.

Glutamic Acid

Histidine HCl

Iysine

Methionine,

Phenylalanine

Plasma Hydrolysate, N.N.R. 1954.

Protein Hydrolysate, N.N.R. 1955.

Thyroxine

AMINOACETIC ACID—HCl, See: Glyco-HCl (Burnham)

AMINOACETIC ACID HI, See: Glyco-HI (Burnham)

AMINO ACIDS, (Stuart) Protein hydrolysate 5 Gm / lbs. Pow. 6 oz.

Use: Protein depleted patients.

W/Dextrose. See: C.P.H., Soln. (Cutter)

W/Vit B 12 (Stuart) Amino acids 0.65 Gm
 Vit B 12 1 mcg /Tab Bol 100s, 500s
 Use Dietary supplement

AMINOACRIDINE-ASCORBATE W/Benzyl
 alcohol See: Acrisan Throat Spray (Reese)

AMINOACRIDINE CAPRYLATE W/Benzylal
 cahal in propylene glycol See: Acrisan
 Sol (Reese)

* **AMINOACRIDINE** 9-aminoacridine
 Use Bacteriostatic agent

9-AMINOACRIDINE HCL (Various Mfr)

W/Hydracortisone acetate tyrothricin phen
 ylmurcuric acetate, polysorbate 80, urea lac
 tase See Aquacort Vag Supp (Webster)

W/Phenylmercuric acetate methylbenze
 thanium chloride succinic acid allantoin
 lactose See Vag tabs, Tab (Ascher)

W/Phenylmercuric acetate methylbenze
 thanium chloride tyrothricin, urea succinate
 acid lactose See Trisert Vag nal, Tab,
 Paw (Ulmer)

W/Phenylmercuric acetate sulfacetamide
 benzalkonium chloride See Kenhex, Yagi
 nal Preps (Kendall)

W/Phenylmercuric acetate tyrothricin urea,
 lactose and sod lauryl sulf See: Triserts,
 Vag Supp (Webster)

W/Sulfonamide & allantoin
 See ACE—Allantamide Oint (National
 Drug)

A Y C Improved, Cream and Supp (Na
 tional Drug)

W/Tetracaine HCl, phenylephrine HCl, Vit
 A & Vit D See: Sema d, Supp (Mossengill)

9-AMINOACRIDINE PROPIONATE See:
 Trichoyal, Jelly (Reese)

9 AMINOACRIDINE UNDECYLENATE W/N
 myristyl 3 hydroxybutylamine HCl methyl
 benzethonium Cl succinic acid See: Cen
 asert Improved Vag Supp (Central)

**p-AMINO BENZENE-SULFONYLACETYL-
 AMIDE**. See Sulfacetamide

p-AMINO BENZOATE CALCIUM W/Salicyl
 amide Vit C, mephenesin See Causolia
 Tab (Grant)

p-AMINO BENZOATE, POTASSIUM See
 Potaba, Paw Tab (Glenwood)

W/Ammonium salicylate & ascorbic acid
 See Arnyl Sod Free Tab (Armour)

Pabalate—Sodium free Tab (Robies)

W/Cortisone acetate, ammonium salicylate &
 ascorbic acid See Neocylate w/Cortisone
 Tab (Central)

W/Hydracortisone free alcohol Pot salicyl
 ate & Vit C See: Arnyl F, Cap (Armour).

W/Salicylamide salicylate pot dicalcium
 glutamate and Vit C See: Glutalate Cap
 (Dumas Withan)

* Generic name recognized by NF USP or AMA

p-AMINO BENZOATE SODIUM.

See: PABA Sod, Tab (Various Mfr)

Nataba, Paw, Tab (Glenwood)

W/Hydracortisone, salicylamide, ascorbic
 acid See: Sylopar HC, Tab (Bunnington).

W/Phenobarbital calcichline salicylate, Vit
 B 1 & aspirin

See: Doforal Tab (Wally Pharm Co)

W/Sodium salicylate & ascorbic acid

See Pabasy, Tab (Ives Cameron)

Pabalate, Liq & Tab (Robins)

Salicin, Tab (Taylor)

W/Salicylamide sod salicylate, ascorbic
 acid bulabarbital sod See: Bisalate, Tab.
 (Allison Lab)

W/Salicylate Sod, Calcichline & Vitamins.

See: Sali Zem No 2, Tab (Zemmer)

W/Sod salicylate, dihydroxy aluminum
 aminoacetate & ascorbic acid See: Elpagan,
 Tab (Smith, Miller & Patch)

W/Salicylamide & ascorbic acid See: Syla
 par, Tab (Bunnington)

**p-AMINO BENZOIC ACID, NF; Tablets,
 NF (PABA)**

Available as cap, bol, paw, sal & tab
 Messengill—Sol (10% Bol) pt

Use: Dietary supplement & onliricketsial
 agent

W/Aspirin, ascorbic acid See: Pabodyne
 Tab (Blue line)

W/Aspirin, ascorbic acid, physostigmine
 salicylate, hamatropine methyl bromide
 See: Mabisyl, Tab (B F Ascher)

W/Aspirin calcichline salicylate, ascorbic
 acid, niacin, & thiamine See: Acetylcal, Tab

W/d Calcium pantlhenate See: Pabacol,
 Int (Ives Cameron)

W/Dipyron, salicylamide & Vit B 12 See:
 Cabasal, Tab (Rand Pharm)

W/Dipyron, salicylamide Vit C, B 12 See
 Dipal, Tab (US Standard Products)

W/Hydracortisone, aspirin, ascorbic acid
 See: Pabirin AC, Cap (Smith Dorsey)

W/Mephenesin & ammonium salicylate See:
 Sol Phenesis Tab (Marian)

W/Mephenesin, salicylamide, vit C, B-1
 stacinamide menadione See: Mybasol
 Tab (Lloyd, Dabney, & Westerfield)

W/Salicylamide, ascorbic acid & arganidin
 See Artamide Tab (Wampale)

W/Sod salicylate & ascorbic acid
 See Cempaba Tab (Sly)

Hysylate Tab (Warren Teed)

Nucarsal Tab (Kendall)

Purpasil C, Tab (Meyers Rosser)

W/Sod salicylate calcichline & ascorbic acid
 See PB Sol w/Calcichline Tab (Ulmer)

Salical Tab (Taylor)

p-AMINOBENZOIC ACID, SALTS.

See: p-Aminobenzoate potassium.

p-Aminobenzoate sodium.

AMINOBENZOYLGLYCINE. See: p-Aminohippuric Acid.***AMINOCAPROIC ACID.** 6-aminohexanoic acid. See: Coprocid (Merck & Co., Inc.)**AMINOCARDOL.** See: Aminophylline. (Various Mfrs.)

AMINO-CONCEMIN. (Merrell) Protein hydrolysate (45% aminoacids) 6.75 Gm., thiamine HCl 3.0 mg., riboflavin 2.0 mg., niacinamide 15.0 mg., pyridoxine HCl 1.0 mg., peptonized iron, N.F. 0.4 Gm., rice bran ext. 0.5 Gm. In a special base containing Liver Fraction II N.F./45 cc. Bot. 1 pt., 1 gal.

Use: Hypochromic microcytic anemia.

AMINODROX. (Massengill) Aminophylline 1½ gr., aluminum hydroxide gel, dried 2 gr./Tab. Bot. 100s, 1000s. Tab. w/phenobarbital ¼ gr., Bot. 100s, 1000s.

Use: (1-2 p.r.n) Diuretic w/antacid.

AMINODROX-FORTE. (Massengill) Aminophylline 3 gr., aluminum hydroxide gel 4 gr./Tab. Bot. 100s, 1000s. Also available with phenobarbital ¼ gr.

AMINO-ETHYL-PROPANOL. See: Aminoisobutanol

W/Bromotheophyllin

See: Pamobrom

AMINOFORM. See: Methenamine. (Various Mfrs.)

AMINOGEN. (Christina) Vit. B complex & folic acid. Amp. 2 cc. Vial 10 cc. Box 12s & 24s, 100s. Sol., Oral, Bot. 9 oz.

Use: Protein hydrolyzate therapy.

***AMINOGLUTETHIMIDE, N.N.D. 1962.** 2-(p-aminophenyl)-2-ethylglutarimide. See: Eliphen, Tab. (Ciba)

p-AMINOHIPPURATE SODIUM, N.F. N.N.D. 1962 (Merck, Sharp & Dohme) Amp. (10 Gm) 50 cc.

Use: 1 V., Diagnostic aid for renal plasma flow.

p-AMINOHIPPURIC ACID, N.F. (PAHA, 4-Aminobenzoylglycine, p-Aminobenzoylglycine).

AMINOISOBUTANOL. 2 Amino-2-methylpropanol 1.

See: Butophyllamine (Merrell)

Pamobrom for combinations

AMINOISOMETRADINE. See: Elicton, Tab. (Searle)

***AMINOMETRADINE, N.N.D. 1962.** 1-Allyl-6-amino-3-ethylpyrimidine-2,4-dione.

Mictine.

See: Mictine, Tab. (Searle)

Mincard Tab. (Searle)

* Generic name recognized by N.F., U.S.P. or A.M.A.

AMINOMYCIN. (Warner-Lambert Res. Inst) Withdrawn and replaced by nonproprietary name, *Perimycin*.

AMINONAT, N.N.D. 1959. Protein hydrolysates (oral)

***AMINOPENTAMIDE SULFATE, N.N.D. 1962.** α,α-Diphenyl, γ-dimethylamino valeramide. See: Centre, Tab. & Drops. (Bristol)

W/Butobarbital sod. See: Spasmodyl (Columbus)

AMINOPEPTODRATE, N.N.D. 1959. See: Caminoids, Gran. (Arlington)

AMINOPHEN. (Elder) Aminophylline 1½ gr., phenobarbital ¼ gr./Tab. Bot. 100s, 500s, 1000s

Use: Diuretic w/sedative

***AMINOPHTHLINE, U.S.P., N.N.R. 1954; Injection, U.S.P.; Suppositories, U.S.P.; Tablets, U.S.P.** (Aminocardol, Aminophyllin, Cordophyllin, Carena, Diaphyllin, Genophyllin, Inophyllin, Metaphyllin, Phyllindan, Theofamine, Theophylline)

See: Dura-Tab S.M. Aminophylline, Tab. (Wyne)

Reclad-Aminophylline. (Wampole)

Use: Diuretic & myocardial stimulant.

Aminophyllin Combinations on pp. 40-41. Aminophylline w/Phenobarbital on p. 41.

AMINOPHTHLINE w/BENADRYL. See: Hydrolyllin (Searle)

AMINOPHTHLINE INJECTION, U.S.P. Theophylline ethylenediamine. Amp. 3¼ gr., 7½ gr. (Various Mfrs.)

AMINOPHTHLINE SUPPOSITORIES, U.S.P. 7½ gr., 3½ gr. (Various Mfrs.) Pediatric 0.125 gr. Box 12s (Wyeth). See: Neutraphylline. (Merck Sharp & Dohme)

AMINOPHTHLINE TABLETS, U.S.P. Plain or enteric coated 1½ gr., 3 gr. (Various Mfrs.)

AMINOPHTHLINE-PHENOBARBITAL No. 1. (Suthff & Case) Aminophylline 1½ gr., Phenobarbital ¼ gr. Tab. Bot. 100s, 1000s.

AMINOPHTHLINE-PHENOBARBITAL No. 2. (Suthff & Case) Aminophylline 3 gr., Phenobarbital ¼ gr./Tablet. Bot. 100s, 1000s.

1-2-AMINOPROPANE SUCCINATE. See: Cydri, Tabs., Granucaps. (Tutag)

***AMINOPTERIN SODIUM, N.N.D. 1962** (Lederle) Sodium N-[p-(2,4-diaminopterid-6-ylmethylamino-benzoyl)-L-glutamate.

AMINOPTERIN TABLETS. (Lederle) 4-Aminopteroylglutamic acid. Tab. (0.5 mg) Bot. 100s.

Use: Acute leukemia in children.

4-AMINO-PTEROYLGLUTAMIC ACID. See: Aminopterin, Tab. (Lederle)

AMINOPTERINE, N.F. X., A.D.R., N.N.R. 1949;

Aminophylline Combinations

| Aminophylline | Others | Manufacturer | Brand |
|---------------|---|----------------------|--------------------------------------|
| ½ gr. | Aconite 1/5 gr., atropine sulfate 1/1000 gr., veratrum viride ext 1 gr | Mallard | Aminite, Tab |
| 1½ gr. | Allylphenylbarbital ¼ gr. | Massengill | Alphylin Tab |
| 3 gr. | Aluminum hydroxide 4 gr | Massengill | Aminadrox, Tab |
| 5 gr | Aluminum hydroxide 2.5 gr., benzocaine 0.5 gr | Irwin, Neisler | Cardolin Tab |
| 2 gr | Amytal ½ gr., ephedrine HCl ½ gr | Lilly | Amesec Pulvules |
| 195 mg | Butabarbital sod 16 mg., methamphetamine HCl 1 mg., glyceryl guaiacolate 100 mg., phenylephrine HCl 12 mg., aluminum hydroxide-magnesium carbonate 172 mg | Santford | Res O Phyllin Cap |
| 180 mg | W/Butabarbital 15 mg., phenylpropionolamine HCl 25 mg., chlorpheniramine maleate 2 mg., aluminum hydroxide 60 mg. magnesium trisilicate 60 mg | Voile | Asmacal, Tab |
| 100 mg | Butital Sod 10 mg., racemic desoxyephedrine HCl 2.5 mg | McNeil | Drenal, Tab |
| 100 mg | Delival 30 mg | Merck Sharp & Dahme | Delphyneid, Cap |
| 200 mg | Delival 30 mg | Merck Sharp & Dahme | Delphyneid Forte, Cap |
| 250 mg | W/Dextromethorphan HBr 20 mg., alcohol | Ferradale | Liximinal AT/5 cc |
| 100 mg. | Diphenhydramine 25 mg | Searle | Hydrylin, Tab |
| 25 mg | Diphenhydramine 6.25 mg., Pot iodide 15 mg., chloroform 8 mg | Searle | Hydrylin Comp., liq |
| 100 mg. | Diphenhydramine 25 mg., rac. ephedrine HCl 25 mg | Searle | Hydryline w/rac. ephedrine |
| 150 mg | Diphenhydramine 37.5 mg., rac. ephedrine sulfate 25 mg | Searle | Hydrine Tab |
| 3 gr. | Ephedrine HCl ¾ gr., amobarbital ¾ gr | Hoeck | A E.A., Tab |
| 125 mg | Mephobarbital 25 mg., ephedrine HCl 25 mg., In histin 25 mg | Reiser | Asmaton Tab |
| 125 mg | KI 250 mg., mephobarbital 30 mg., ephedrine HCl 20 mg., In histin 30 mg | Reiser | Neb-Asmaton, Cap |
| 100 mg | Pentaerythritol tetranitrate 15 mg | Kramers Urban | Amitrale, Tab |
| 100 mg | Pentaerythritol tetranitrate 10 mg., aluminum hydroxide 150 mg | Warner Chilcott | Peritrate W/Aminophylline, Tab |
| 0.5 Gm | Pentobarbital Sod 0.1 Gm | Wyeth | Wyeth, Supp., Tab. |
| 0.25 Gm., | Pentobarbital Sod 0.5 Gm | Smith Dorsey | Neulraphylline & Pentobarbital, Supp |
| 0.5 Gm | Pentobarbital Sod 0.1 Gm., benzocaine 0.06 Gm | Bischoff | Amineal, Supp |
| 3 gr. | Pentobarb Sod ¼ gr., ephedrine HCl ¼ gr., benzocaine ¼ gr., aluminum hydrox 2½ gr | Irwin, Neisler | Definite, Tab, |
| ½ gr., | Phenobarbital ¼ gr | Elder | Aminaphen, Tab |
| ½ gr. | Phenobarbital ¼ gr., ecamite pow 1/5 gr., veratrum viride ext 1 gr atropine sulf 1/1000 gr | Mallard | Veraphyllin, Tab |
| 1½ gr | Phenobarbital ¼ gr., ephedrine sulfate ¾ gr | Smith Müller & Patch | Amedarb, Tab |

Aminophylline Combinations

| Aminophylline | Others | Manufacturer | Brand |
|---------------|---|----------------|--------------------|
| 97 mg. | Phenobarbital 16 mg., rutin 30 mg. | Sulliff & Case | Ama-Ru, Tab. |
| 100 mg. | Quercetin 15 mg., ascorbic acid 25 mg., raterpine 50 mcg. | Schenley | Rufaminol-RQ, Tab. |
| 3 gr. | Quinine sulf., 4 gr. | Walker | Quinamm, Tab. |
| 100 mg. | Reserpine 0.1 mg., ephedrine sulf., 16 mg. | Harvey | Serphedrine Cap. |

Aminophylline with Phenobarbital

| Aminophylline | Phenobarbital | Others | Manufacturer | Brand |
|---------------|-----------------|---|-----------------------|---------------------------------|
| 1½ gr. | ¼ gr. | Aluminum hydroxide 2 gr. | Lemmon | Ambital, Tab. |
| 1½ gr. | ¼ gr. | Aluminum hydroxide 2.5 gr., bentocaine 0.5 Gm. | Messingill | Aminodrox w/Phenobarbital, Tab. |
| 5 gr. | ¼ gr. | Ephedrine ¾ gr. | Irwin, Neisler | Cardalfin-Phen, Tab. |
| 1½ gr. | ¼ gr. | Ephedrine HCl ¼ gr., Pot. iodide 3 gr. | Kremers-Urban | Enafen, Tab. |
| 2 gr. | ½ gr. | Ephedrine HCl 24 mg., pyritamine molecule 12.5 mg. | Paythress | Madrone, Tab. |
| 65 mg. | 8 mg. | Ephedrine sulfate ¾ gr. | Raymer | Barbahist, Tab. |
| 1½ gr. | ½ gr. | Ephedrine sulfate ¼ gr. | Gold Leaf | Amifed, Tab. |
| 2 gr. | ¼ gr. | Ephedrine sulfate ½ gr., ephorbia pilulifera ext. 1½ gr. | Harvey | Ampafed, Tab. |
| 1½ gr. | ½ gr. | Ephedrine sulfate ¾ gr. | Warren-Teed | Euphed-ital, Tab. |
| 1½ gr. | ¼ gr. | Ephedrine sulfate ¾ gr. | Smith, Miller & Patch | Amedarb |
| 1½ gr. | ½ gr. | Ephedrine sulfate ¾ gr., Raybanol 1 gr. | Raymer | Rayphyllin, Cap. |
| 1 gr. | ½ gr. | Ephedrine HCl 1/12 gr., ethyl aminobenzoate ¼ gr., aluminum hydroxide ½ gr. | Irwin, Neisler | Dalnite Pediatric, Tab. |
| 65 mg. | 8 mg. | Ephed. sulf., 24 mg., pyritamine molecule 12.5 mg./tab. | Raymer | Barbahist, Tab. |
| 0.5 Gm. | (Sod.) 0.05 Gm. | Pentobarbital 0.05 Gm., benzocaine 0.1 Gm. | Ferrows | Arbec, Supp. |
| 2½ gr. | ¼ gr. | Pot. iodide 2 gr. | Central | Amiophen, Tab. |
| 2½ gr. | ¼ gr. | Pot. iodide 2 gr. | Searle | Kiophyllin, Tab. |
| 1½ gr. | ¼ gr. | Pot. iodide 1 gr. | Kremers-Urban | Phylliphen, Tab. |
| 2½ gr. | ¼ gr. | Pot. iodide 2 gr. | Raymer | Rayminatol No. 2, Tab. |
| 1½ gr. | ½ gr. | Racophedrine ¾ gr. | Searle | Amadrine, Tab. |
| 1½ gr. | ¼ gr. | Rutin | Searle | Ruphyllin, Tab. |
| 45 mg. | ¼ gr. | Rutin 10 mg., mannitol hexanitrate 25 mg. | Calumbut | Hexo-Rutalin, Cap. |
| 100 mg. | ¼ gr. | Rutin 20 mg., veratrum viride 95 mg. | Adams | Verufamin, Tab. |
| 100 mg. | ¼ gr. | Rutin 60 mg., Vit. C 25 mg. | Schenley | Rufaminol, Tab. |

- Elixir, N F X; Tablets N F X (Various Mfr)
 (Amidofebrin Amidopyrazoline, Anaf
 brina Navamidon, Pyradane) Dimethyl
 amino-phenyl dimethylpyrazolane
 See: Pyramidon, Prep (Winthrop Labs)
 Use Antipyretic, analgesic
- W/Acetaphenetidine, caffeine & hyoscyamus
 ext
 See Amidaphen, Pulv (Lilly)
 Ancylol Tab (Massengill)
 Andafve, Cap (Massengill)
- W/Acetaphenetidin, caffeine & pheobar
 bitol See Nalgised Tab (Chicago Pharm)
- W/Acetaphenetidin caffeine phenobarbital
 & magnesium oxide See Campiric, Tab
 (Harco)
- W/Amytal See: Amytal w/Aminopyrine
 Cap (Lilly)
- W/Butehalol See Amina Neanal, Tab (Ab
 bott)
- W/Caffeine & ginger oleoresin See: Ami
 feine Cap (McNeil)
- W/Caffeine & raybanal See: Ray Pyrine
 Cap (Raymer)
- W/Dial
 See Barprine, Tab (Ulmer)
 Cibalgine Amp & Tab (Ciba)
- W/Hyoscyamus ext, phenacelin, caffeine
 See Amidaphen, Tab (Lilly)
- W/Isobutylalyl barbituric acid caffeine
 See Opialidan Tab (Sandoz)
- W/Phenobarbital
 See: Amylatene Compound, Cap (First
 Texas)
 Phenapyrine, Tab (Cole)
 Pyramidal, Tab (Winthrop Labs)
- W/Phenobarbital, hyoscyamus ext See: Ami
 tabs, Tab (G F Harvey)
 Nyased, Tab (Walker)
- W/Potassium bromide See: Gynalgas, Ellixr
 (Wyeth)
- AMINOQUIN NAPHTHOATE.** See: Pamo
 quine Naphthoate (Winthrop Labs)
- ***p-AMINOSALICYLATE CALCIUM, N F,**
 NND 1962 Tab NF (Dumas Wilson)
 7½ gr Bat 1000s
 See Parosal Calcium, Tab (Parroy)
- ***p-AMINOSALICYLATE, POT, NND 1962**
 See Parosal Pot Tab (Parroy)
- ***p-AMINOSALICYLATE SODIUM, USP**
 NND 1962 Tab, USP (Various Mfr)
 See: Pamisyl Sod Amp Tab (Parke Davis)
 Pasora Sod Tab (Smith Dorsey)
 Pasem Sod, Tab (Massengill)
 Pasmed Sod, Tab (Intermedico Corp)
- * Generic name recognized by N F, USP
- Parapas Sod Tab (Gold Leaf)
 Parasal Sod, Tab (Parroy)
- W/Isosozid See: Isopacaln, Tab (Consol
 Midlaed)
- W/Ameasaliclylic acid See: Neoposalate,
 Tab (Dumas Wilson)
- W/Succinic acid See: Tubertote, Tab (Heun)
- ***AMINOSALICYLIC ACID, U S P, NND**
 1962; Tablets U S P, PAS, p Ameasaliclylic
 acid
 See: Pamisyl, Tab (Parke, Davis).
 Para Pas Tab (Gold Leaf)
 Parasal Tab (Parroy)
 PAS Tab (Lilly)
 Propasa, Tab (Merck Sharp & Dohme)
 Rezipos, Paw (Squibb)
 Uses: Adjunct to streptomycin in tuberculosis
- W/p Aminosalicylate Sod See: Neoposalate,
 Tab (Dumas-Wilson)
- W/Isosozid
 See Dipasic (Parroy)
 Parasal INH (Parroy)
- W/Isosozid buffered See: Parasal INH 20
 Buffered (Parroy)
- p-AMINOSALICYLIC ACID SALTS.**
 See Aminosalicylate Calcium
 Aminosalicylate Potassium
 Aminosalicylate Sodium
- AMINOSERP TABLETS.** (Nysco) Aminophyl
 line 100 mg, reserpine 01 mg/Tab Bot
 1000s
 Use: Hypotensive in angina pectoris
- AMINOSOL, NND 1962** (Abbott) Modified
 fibrin hydrolyzate solids (amino acids &
 peptides) 50 Gm/1000 cc Solution (5%)
 Bot 1000 cc, 6s, 48s; Bot 500 cc, 6s, 48s
 Use: IV, to maintain a nitrogen balance
 5% W/Dextrose 5%. (Abbott) Sol, Bot
 1000 cc, 6s, 48s; Bot 500 cc, 6s, 48s; Bot
 250 cc, 6s, 48s
 Use: IV, nutritional
- 5% W/Dextrose 5% in Alcohol 5% (Abbott)
 Sol, Bot, 1000 cc, 6s, 48s.
 Use: IV, nutritional
- 5% W/Dextrose 5% & Sodium chloride
 D3% (Abbott) Sol, Bot 1000 cc, 6s, 48s.
 5% W/Fructose 10%/1000 cc Abbo-Liter
 container
 Use: IV, nutritional
- p-AMINOTOLUENE SULFONAMIDE.** See:
 Sulfabenzamine
- AMINOTRATE PHOSPHATE, NND 1962**
 Triethylate Phosphate See Triethanolamine
- Triethylate Biphosphate, Preps
- AMINOXYTROPINE TROPATE HCl** Atro
 pine N oxide HCl Genotropine HCl
- AMIODOXYL BENZOATE** (Frost) Vial (1
 Gm) Box 6s
 Use Rheumatoid arthritis

AMIOPHEN. (Central) Aminophylline 2½ gr., phenobarbital ¼ gr., Pot. Iodide 2 gr./Tab. Bot. 100s, 1000s.

Use: (I.I.d.) Myocardial stimulant, diuretic & vasodilator.

* **AMIPHENAZOLE NCl.** 2,4-Diamino-5-phenylthiazole HCl. See: Doptaxale.

* **AMISOMETRADINE, N.F., N.N.D.** 1962. 6-Amino-1-methyl-3-methylpyrimidine-2,4-dione. See: Ralictan, Tab. (Searle)

AMITABS. (G. F. Harvny) Aminopyrine 2½ gr., phenobarbital ¼ gr., hyalcyamus nxl. ½ gr./Tab. Bot. 1000s

Use: Antipyretic, analgesic and sedative

AMITIOZONI.

See: Panrone, Tab. (Panroy)

Teabazonn, Tab (Consol. Midland)

Tibionn, Tab. (Schenley)

AMITRATE, (Kremers-Urban) Aminophylline 100 mg., penicerythritol tetranitratn 15 mg., aluminum hydroxide 150 mg./Tab. Bot. 100s & 500s.

Use: (I.I.d.) Coronary vasodilator, diuretic.

AMITRENE, (Normond) Amphetamine sulfate, racemic. Tab. (5 mg. & 10 mg.) Bot. 100s. Vial (20 mg./cc.) 30 cc. (d-amphetamine sulfate). Tab. (5 mg) Bot. 100s.

Use: Central nervous system stimulant.

* **AMITRIPTYLINE HCl,** J-(3-Dimethylamino-propylidene)-1,2,4,5-dibenzocyclohepta-1,4-diene. Tryptizol hydrochloride. See: Etavil, Tabs. & Vials. (Merck Sharp & Dahme)

AMMENS MEDICATED POWDER. (Bristol-Myers) Oxyquinoline, zinc oxide, talc and starch. Pkg. 2 oz., 5.5 oz.

Use: Heat rash, prickly heat or diaper rash

AMMERZO. (Eldnr) Ammoniated mercury 10%, zinc oxide 7% in combination with aromatic oils. Oint., Jar 1 lb.

Use: Antiseptic skin preparation.

AMMIVIN. (National Drug) (Khallin, Crystalline) Tab. E.C. (10 mg) Bot. 100s. Tab. E.C. (20 mg) Bot. 40s, 100s. Vial (50 mg./cc.) 10 cc.

Use: Orally & I.M., angina pectoris, coronary insufficiency & bronchial asthma.

AMMOIDIN. See: Methaxsafen, N.N.D. 1958

AMMONERIC. (Sutliff & Case) Ammonium chloride 7½ gr./Tab. Bot. 1000s

Use: Diuretic

AMMONIATED MERCURY, U.S.P.; Oint., U.S.P.; Ophth. Oint. U.S.P. (Various Mfr.).

See: Aftercreme (Tnxal).

W/Zinc oxide. See: Ammerzo, Oint. (Eldnr)

AMMONIUM CHLORIDE, U.S.P.; Capsuls. U.S.P. XY; Tablets, N.F. X. Cap (5 gr. & 7½ gr.) Tab. plain or E.C. (5 gr. & 7½ gr) (Various Mfr.).

* Generic name recognized by N.F., U.S.P. or A.M.A.

See: Amchlor, Tab. (Brewer).

Ammaneric, Tab. (Sutliff & Case)

Use: Diuretic, expectorant.

W/Am. and sod. mandelatin. See: Mandchlor, El. (Vanpelt & Brown)

W/d-Amphetamine sulfate, homatropine methylbromide & Vit. B complex. See: Pre-Mens W/d-amphetamine, Tab. (Purdue Frederick).

W/Dihydrocodeinane bilartrate, phenylephrine NCl, chlorpheniramine maleate, potassium guaicol sulfonate, sodium citrofn. See: EFRICON EXPECTORANT (Lannett)

W/Dihydrocodeinone, pyrilamine maleate, dl-desoxyephedrine HCl, pot guaicol sulfonate, tartar emelic, chloroform. See: Bonodene, Syr. (Banar)

W/Homatropine methylbromide & Vit. B complex. See: Pre-Mens, Tab. (Purdun Frederick).

W/Mandelic acid as the monoethanolamine salt. See: Mandelic Acid & Ammonium Chloride, Enseals (Lilly).

W/Phenylephrine HCl, pyrilamine maleate, d-metharphon HBr, citric acid. See: Pylal dine Pediatric, Syr (Vanpelt & Brown)

W/Phenylpropanolamine, acetaminophen.

See: Zesta, Tab. (Slandex)

W/Quinine sulfate, belladonna R. ext., camphor, opiate fl. ext. See: Caryza, Tab. (Merck Sharp & Dahme)

W/Theophyllinn, phenobarb. See: Borphylin, Tab. (Mollard)

AMMONIUM CHLORIDE, ENSEALS. (Lilly)

Ammonium chloride. Tab. (7½ gr., 15 gr., lumed disintegrating) Bot. 100s, 500s, 1000s.

Use: (20 to 60 gr. t.i.d.) Acidification of the urine.

AMMONIUM GLUTAMATE. See: Gustamole, Pow. (Arlington)

AMMONIUM MANDELATE, U.S.P. XIV. Ammonium salt of mandelic acid.

Squibb—Tab. I.C. (5 gr.). Plain (7½ gr) Bot. 200s, 1000s.

Upsher-Smith—Syrup, N.N.R. 1947 (8 Gm./fl. oz.) Bot. pt., gal.

See: Amdelete, Elix., Syr. (Abbott).

Mandelic Acid, Elix. (Lilly).

Use: Orally, urinary antiseptic.

W/Sodium mandelatin & ammonium chloride See: Mandechlor, Elix. (Vanpelt & Brown).

AMMONIUM SALICYLATE, N.F. X. (Various Mfr.)

W/Pot. p-aminobenzoate & ascorbic acid. See: Arnyl Sodium Free, Tab. (Armour).

Also w/phenobarb.

W/Caffeine, terpin hydrate, ammonium gluconate, glycine, benzalkonium chloridn

- See Martusin Special, Liq (M R Thompson)
- W/Carbonyl acetate Pot paminobenzoate & ascorbic acid See Neacylate w/Carbonyl, Tab (Central).
- W/Mephenesin, paminobenzoic acid See: Sal Phenesin, Tab (Marion)
- W/Pot paminobenzoate, ascorbic acid, & methamphetamine HCl See Neacydex, Tab (Central)
- W/Pot guaracol sulfonate tr labels, eye talu balsam See Bramul Syr (Hoag)
- AMMONIUM SULFATE.** See: Dolamin, Amp (G F Harvey)
- AMMOPHYLLIN** See Aminophylline, USP (Various Mfr)
- AMMORID**, (Kinney) Benzethonium chloride & zinc oxide in a lanolin absorption base Tube 2 oz
Use Topically dermatologic ointment
- AMMORID DIAPER RINSE**, (Kinney) Methyl benzethonium chloride 5%, plus deodorizing & buffering agents in dry powder form ready soluble in water Bot 240 Gm
Use Protection against diaper rash
- AMNESTROGEN**, NND 1962 (Squibb) Conjugated estrogenic substances NNR 1954 Tab (0.3 mg) Bot 100s (0.625 mg & 1.25 mg) Bot 100s, 1000s (2.5 mg) Bot 25s 100s
Use Menopausal syndrome
- AMNIOTIN**, NNR 1949 (Squibb) Estrogenic substances in oil Vial (10 000 IU, 20 000 IU & 50 000 IU/cc) 10 cc
Use Intravaginal & IM; atrophic disorders of the genitalia
- AMNI VASNAGA PRODUCTS** See Khellin (Various Mfr)
- * **AMOBARBITAL**, USP NNR 1950, A DR; Elixir, NF Tablets USP (Various Mfr) 5 ethyl 5 isocarbonylbarbituric acid
See ANS Tab (Kenwood Labs)
Amytal (Lilly)
Use Hypnotic at intermediate duration
W/d Amphetamine sulfate belladonna ext See G I Tran Tab (Kendall)
- W/d Amphetamine sulf, and Vitamin complex with minerals
See: Daycap B Cap (Ethical Pharm)
Quadamine, Timed Cap (Tutag)
Timed Arda Cap (Testagar)
- W/Aspirin, phenacetin caffeine secobarbital sodium, buffered w/aluminum and magnesium hydroxides See Buffadyne W/Barbiturates Tabs (Lemman)
- W/Chloralpyrine See: A N S Tablets (Kenwood Labs)
- W/Dexedrine sulfate See: Dexamyf Spanuls (Smith, Kline & French).
- * Generic name recognized by NF, USP or AMA
- W/Dextro amphetamine HCl
See: Ambodex Timed Cap (Testagar)
- Amphetamine W/Amobarbital, Cap (West Ward)
- Am Phat Unadoy Ph 758, Cap (Pau Maney)
- Amyaldex Unicelles #1 & #2 (His Pharm)
- Dodexal, TD Cap (Central)
- Deeobarb, Tab TD (Nysco Labs)
- Dexamyf Spanuls (Smith, Kline & French)
- Deeibar B, Tab (Barre Drug Co)
- Evrader Plus, Tempules (Evrans)
- Sla Jaydex, Tab (Mid Atlantic)
- W/Deerlroomphetamine sulfate
See Domopal, Tab (Hoag)
- Lowebarb CTR, TR Cap (Lowe)
- Mildex Cap (Bonar)
- Tydex Plus TR Cap (Tyler)
- W/Dextro Amphetamine sulfate, acetyl salicylic acid, phenacetin See: Daptrial, Tab (Smith, Kline & French)
- W/Ephedrine HCl & aminophylline See: AEA, Tab (Heack)
- W/HMS See Sed Tens (Lemman)
- W/Nomatrapine methylbromide, methamphetamine HCl See: Adipex, Tab (Lemman)
- W/Mephenesin See Sonazol, Tab (Tulag)
- W/Mephenesin, nomatrapine methylbromide salicylamide physostigmine sol See: Myamphetane, Tab (Testagar)
- W/Methamphetamine, thyroid See: Obestetab (Lemman)
- W/Methicopalamine Bromide See: Scoline Amobarbital, Tab (Kendall)
- W/Phenobarbital, d amphetamine sulfate, 7 minerals 9 vitamins See Amvical X, Tab (Stuart)
- W/Salicylamide, phenacetin, d amphetamine sulfate, hyoscyamine sulfate See Sedamosal, Cap (Finley)
- W/Secobarbital, phenobarbital See Tritol Tabs (Testagar)
- W/Theophylline ephedrine, alcohol See Asmafield, Lrg (Canfield)
- W/Thyroid methamphetamine See: Obestetab (Lemman)
- * **AMOBARBITAL SODIUM**, USP, NNR 1950; Capsules, USP; Sterile, USP; Injection (Various Mfr)
Amp (1 1/2 gr) 6s, 12s, 25s, 100s (3 1/4 gr) 6s, 12s, 25s, 100s (7 1/2 gr) 6s, 12s, 25s 100s (Various Mfr)
Cap (3 gr) Bot 1000s (Amer Drug Prods)
Pow: Bot 1/2 lb, 1 lb Pkg 5 lb (Various Mfr)
Vials (15 1/2 gr) 10 cc (Various Mfr)
See: Amytal Sod (Lilly)

Use: Hypnotic of intermediate duration.

W/Secobarbital Sod. See: Dusotal, Cap. (Harvey)

AMOBESL. (Cohn) Dextro-amphetamine phosphate, dibasic, 5 mg., thiamine HCl 2 mg., carboxymethylcellulose sod. 165 mg./Tab. Bot. 100s, 500s, 1000s.

Use: Orally, reducing aid.

AMOCAL Jr. (Jenkins Labs) Glycyrrhiza compound 30 min., ipecac pow. 1/16 gr., amm. chloride 1/4 gr., Iodized Cal. 1/16 gr., w/licarice ext./Tab. Bot. 100s, 1000s.

Use: Expectorant and anodyne.

AMODEX, Timed. (Tastogor) d-Amphetamine HCl 1/4 gr., amobarbital 1 gr./Cap. Cap. Timed, Bot. 100s, 1000s.

AMODIN Timed Cap.: d-amphetamine HCl 7.5 mg., amobarbital 30 mg./Cap. Bot. 100s, 1000s

Use: (1 a.m.) Obesity.

* **AMODIAQUINE HYDROCHLORIDE**, U.S.P. N.N.D. 1962. Tab. U.S.P. 4 (3'-Diethylamino-methyl-4-hydroxyaniline)-7-chloroquinoline dihydro. (Parke, Davis) See: Camoquin Hydrochloride. Infatab (Parke, Davis)

W/Primaquine. See: Camoprim, Tab.

AMODRINE (Searle) Aminophylline 1 1/2 gr., racephedrine 3/8 gr., phenobarbital 1/8 gr./Pholn or E.C. Tab. Bot. 100s, 1000s.

Use: (1 i.i.d.) Antispasmodic.

AMOEIBICON. (Consol. Midland Corp.) Glycobiarsol 0.25 Gm./Tab. Bot. 50s, 250s, 1000s.

Use: (2 i.i.d. for one week) Intestinal amebiasis.

AMOL. Mono-n-onyl-hydroquinone ether. Sani B-E-I, Pow. (Marck Sharp & Dahme)

* **AMOLANONE HYDROCHLORIDE**, N.N.D. 1962. Sani Amethone Hydrochloride. (Abbott).

AMOLATE. (Mallord) Pyrilamine maleate 80 mg., phenylephrine HCl 20 mg., ammonium chloride 8 gr., Sod. citrate 16 gr., chloroform 2 gr., menthol, q.s./fl. oz. Bot. 3 oz., 1 gal.

Use: Antihistaminic cough therapy.

AMONDRIN. (Brean) Ephedrine HCl 1/2 gr., ammonium chloride 194 mg., calcium cresofate 16.2 mg., Pat. gualecol sulfonate 194 mg. & benzocaine 2 mg./Tab. Bot. 1000s.

Use: Cough preparation.

AMO-PELLCAPS No. 1. (Kirkman) Dextro-amphetamine sulf. 10 mg., amobarbital 60 mg./cap. Bot. 100s, 500s, 1000s.

Use: Mental & emotional distress.

AMO-PELLCAPS No. 2. (Kirkman) Dextro-amphetamine sulf. 10 mg., amobarbital

100 mg./cap. Bot. 100s, 500s, 1000s. Timed disintegration.

Use: Mental and emotional distress.

* **AMOPYROQUIN HCl**. 4-(7-chloro-4-quinolylamino)-a-1-pyrrolidyl-a-cresoldihydrochloride. See: Propoquin. (Parke, Davis)

AMO-RU. (Suliff & Cosn) Aminophyllin 97 mg., phenobarbital 16 mg., rulin 30 mg./Tab. Bot. 100s, 1000s

Use: Diuretic and myocardial stimulant w/ sedative

AMORYN. (Plessner) Estrogens conjugated, water-soluble mixture. Tab. (0.625 mg & 1.25 mg) Bot. 100s, 1000s.

Use: Conjugated estrogens therapy.

AMOSENE. (Ferndale) Bot. 100s.

Use: Fast-acting tranquilizer.

AMOTHILE. (G. F. Harvey) Vit. B-1 1 mg., B-2 0.5 mg., niacin 5 mg., aminoacetic acid 5 gr./5 cc. of mixt. Bot. 4, 8, 16 oz., 1 gal. Use: Vitamin B complex therapy.

* **AMOTRIPHINE HCl**. (Winthrop Labs) 3-Di-methylamino-1,1,2-tris (4-methoxyphenyl)-1-propene HCl.

Use: Cardiovascular agent.

AMOTROCIN. (McNeil) Tyrothricin 1 mg., cetyltrimethylbenzyl ammonium chloride 5 mg., orthosin (benzocaine) 3.5 mg./lozenge. Box 12s, Bot. 100s, 1000s.

Use: Minor throat irritations.

AMOXIN. (Farosin) Not listed 4-Toluene-sulfanylamino-1-acetylhydroxy-2-benzene carboxylic acid with iodoacetate & aspirin. Tab. (0.4 Gm), Vials 50s.

Use: (1 q. 3 h.) Analgesic.

AMPA. (Sherman) Aspirin 2 1/2 gr., acetophenetidin 2 1/2 gr., amphetamine sulfate, racemic, 1/26 gr./Tab. Bot. 100s, 1000s.

Use: (1-2 q. 4 h.) Analgesic with central nervous system stimulant.

AMPATED. (G. F. Harvey) Aminophyllin 2 gr., phenobarbital 1/4 gr., ephedrine sulfate 1/4 gr./Tab. Bot. 1000s, 5000s.

Use: Hypertension with sedative.

AMPAVE. (Marck Sharp & Dahme) Amphetamine phosphate, racemic, 5 mg., caffeine 30 mg./Tab. Bot. 100s, 1000s.

Use: Depress appetite.

AMPERONE. (Kramers-Urban) Dinastrol 0.5 mg., methyltestosterone 2.5 mg., pentobarbital sod. 300 mg., d-amphetamine sulf. 5.0 mg./Tab. Bot. 60s.

Use: Indicated for relief of menopausal symptoms accompanied by anxiety and tension.

AMPHAPLEX. (Palmedico) Methamphetamine saccharate, methamphetamine HCl, amphetamine sulfate, dextroamphetamine sulfate.

* Generic name recognized by N.F., U.S.P. or A.M.A.

5 mg of each in the 20 mg tab 25 mg of each in the 10 mg tab Bot 100s

Use Appetite control

AMPHARB (G F Harvey) Amphetamine sulfate 10 mg, phenobarbital ¼ gr/tab Bot 1000s 5000s

Use Diuretic with sedative

AMPHATE (Starck) Amphetamine phosphate 5 mg/Tab Bot 100s, 1000s

AMPHATIN (Morning) (Normand) Amphetamine HCl racemic 5 mg thyroid 60 mg atropine sulfate 0.18 mg, Vit A 2500 u 0.300 u ascorbic acid 50 mg, thiamine HCl 0.9 mg riboflavin 1.25 mg niacinamide 10 mg folic acid 0.75 mg/Tab Red Bot 30s 100s, 1000s

Use Anti obesity with vitamins

AMPHATIN (Afternoon) (Normand) For administration in afternoon Amphetamine HCl racemic 5 mg, thyroid 60 mg atropine sulfate 0.18 mg, alain 15 mg, Vit A 2500 u, D 300 u, ascorbic acid 50 mg thiamine HCl 0.9 mg, riboflavin 1.25 mg niacinamide 10 mg folic acid 0.75 mg/Tab Green Bot 30s 100s, 1000s

Use: Anti obesity with vitamins

AMPHCAPS 10 & 15 Mg caps (Manne Labs) Dextra amphetamine sulfate in a base with timed disintegration through a 6 to 10 hour period/10 mg and 15 mg/cap Bot Caps of 10 mg and 15 mg/100s and 1000s

Use Anorectic

***AMPHACLORAL**, α methyl N (2,2,2 trichloroethylidene)phenethylamine See: Acu Iron (Irwin Neisler)

AMPHEDASE (Parke Davis) d Amphetamine sulfate 25 mg, nicotinamide 25 mg thiamine HCl 5 mg, ascorbic acid 50 mg, laktal d ostate 300 mg/Kapsel Bot 100s, 500s Use (12 p.c.) Asthenia, depression, obesity or chronic alcoholism

w/Phenobarbital (Parke Davis) Amphetamine plus phenobarbital ¼ gr Tab, Bot 100s 1000s

Use: (1 or 2 p.c.) Depression, faulty nutrition or chronic alcoholism

AMPHEDRINE (Van Felt & Brown) Amphetamine sulfate racemic Tab 10 mg, Bot 100s 500s, 1000s; d Amphetamine sulfate Tab 5 mg, Bot 500s, 1000s

Use: Central nervous system stimulant

AMPHEDRINE-M (Van Felt & Brown) Levamphetamine sulfate 5 mg, Metylcellulose 350 mg/Tablet Bot 100s 1000s

Use: Appetite depressant

AMPHEDROXYNHYDROCHLORIDE, NND 1962 (Lilly) Methamphetamine HCl Elixir—

* Generic name recognized by NF USP or AMA

25 mg/4 cc Bot 1 pt, 1 gal Tab.—5 mg Bot 25s, 100s, 1000s

Use Central nervous system stimulant

***AMPNENIDONE**, 1 m aminophenyl 2 [1H] pyridane See: Darwal (Mallibie)

AMPNEREX, (Royce) d Amphetamine sulf. mg/Tab Bot 100s, 1000s

AM-PHET, (Paul Maney) Amobarbital 60 mg d amphetamine sulfate 15 mg/Cap Bot 100s 1000s

Use Treatment of obesity and depressly states

AMPNETAMINE W/AMOBARBITAL, (West Ward) No 1s; d Amphetamine sulfate 11 mg, amobarbital 60 mg/TD Cap Bot 100s 1000s

No 2s; d Amphetamine sulfate 15 mg amobarbital 100 mg/TD Cap Bot 100s 1000s

Use Sedative w/CNS stimulant

d-AMPHETAMINE CARBOXYMETHYLCELLULOSE SALT, See Bontril, TR Tab (O W Carrick)

AMPNETAMINE COMBINATIONS, (Dumas Wilson) d Amphetamine HCl 10 mg, thyroid 1 gr, alain ¼ gr, atropine sulfate 1/360 gr/Tab Bot 100s

Use Anti obesity therapy

Amphetamine Combinations on pp 47 to 51

AMPHETAMINE HYDROCHLORIDE, USP XIV, NNR 1949 Rac amphetamine HCl methylphenethylamine HCl dl 1 phenyl 2 aminopropane HCl dl methylphenethylamine HCl rac desoxyephedrine HCl Amp—20 mg/cc 1 cc (Various Mfr) Cap—12 mg (Luitgarten & Vitamix) Tab—5 mg 10 mg, 15 mg (Various Mfr)

Use Vasoconstrictor, central nervous system stimulant

d-AMPHETAMINE HYDROCHLORIDE (Various Mfr) Dextra amphetamine hydrochloride Cap.—5 mg, 10 mg, (Various Mfr) Tab—5 mg, 10 mg (Various Mfr).

Use Vasoconstrictor, central nervous system stimulant

***AMPHETAMINE**, Leva See Levamphetamine

***AMPHETAMINE PHOSPHATE**, Dibasic NF, NND 1962; Tab NF, Inj NF rac amphetamine phosphate Cap—5 mg 10 mg Tab—5 mg 10 mg (Various Mfr) See Amphetamine, Tab (Starck)

Bar Dex, Tab (Barre Drug) D etamine, Vial (Key Corp) Monaphos, Tab (G F Harvey) Profetamine, Amp & Tab (Clark & Clark) Racephen, Tab (Ives Cameron)

Amphetamine Combinations

| Amphetamine | Others | Manufacturer | Brand |
|---------------|--------------|---|-----------------------|
| d (sulf.) | 2.5 mg. | Acetophenetidin 2½ gr., pentobarbital sod. ¼ gr., aspirin 3½ gr. | Amnar-Stone |
| dl (phos.) | 2 mg. | Acetyl-p-aminophenol 300 mg., salicylamide 200 mg., metoprine 0.5 mg., phenyltoloxamine dihydrogen citrate 25 mg. | Sirasenburgh |
| d.HCl | ¼ gr. | Amobarbital 1 gr. | Testagor |
| d (sulf.) | 10 or 15 mg. | Amobarbital 60 or 100 mg. | West-Ward |
| d (sulf.) | 15 mg. | Amobarbital 60 mg. | Barre |
| d (sulf.) | 15 mg. | Amobarbital 1½ gr. | Paul Maney |
| d (sulf.) | 15 mg. | Amobarbital 60 or 100 mg. | Hiss Pharm., Central |
| d (sulf.) | 15 mg. | Amobarbital 60 mg. | Nystco Labs. |
| d (sulf.) | 10 or 15 mg. | Amobarbital 60 or 100 mg. | Smith, Kline & French |
| d (sulf.) | 3 mg. | Amobarbital ½ gr. | Barre |
| d (sulf.) | 10 mg. | Amobarbital 1 gr. | Smith, Kline & French |
| d (sulf.) | 15 mg. | Amobarbital 1 gr. | Hoag |
| d (sulf.) | 5 mg. | Amobarbital 1½ gr. | Evron |
| d (sulf.) | 10 mg. | Amobarbital 30 mg. | Evron |
| d (sulf.) | 15 mg. | Amobarbital 60 mg. | Evron |
| d (sulf.) | 15 mg. | Amobarbital 95 mg. | Tyler |
| d (sulf.) | 15 mg. | Amobarbital 100 mg. | Evron |
| d (sulf.) | 5 mg. | Amobarbital ½ gr., aspirin 2½ gr., phenacetin 2½ gr. | Smith, Kline & French |
| d (sulf.) | 5 mg. | Amobarbital 32 mg., belladonna ext. 16 mg. | Kendall |
| d (sulf.) | 15 mg. | Amobarbital 45 mg., plus vitamins & minerals | Testagor |
| d (sulf.) | 10 mg. | Amobarbital 45 mg., plus vitamins & minerals | Ethical Pharm. |
| d (sulf.) | 15 mg. | Amobarbital 64 mg., vitamin complex and minerals | Tutag |
| d (res. comp) | 16.25 mg. | Amobarbital 45 mg., w/vitamins and minerals | Sirasenburgh |
| d (sulf.) | 10.5 mg. | dl-Amphetamine resin complex 6.25 mg. dl-Amphetamine sulf. 10.5 mg. | Tutag |
| dl (phos.) | 15 mg. | d-Amphetamine sulf. 15 mg., butobarbital sod. 45 mg. | Lynn |
| d (sulf.) | 2.5 mg. | Aspirin 4 gr., acetophenetidin 2 gr. | G. F. Harvey |
| dl (phos.) | 5 mg. | Alarax, 5 mg., 9 vitamins 11 minerals/capsule. | Roerig |
| dl (phos.) | 1 mg. | Atrop. methyl nitrate 1 mg., carbonylmethylcellulose sod. 200 mg. | Sirasenburgh |
| dl (phos.) | 5 mg. | | |

Amphetamine Combinations

| Amphetamine | | Others | | Manufacturer | Brand |
|-------------|---------|---|--|---------------------|------------------------|
| d (phos.) | 5 mg | Atropine methyl nitrate, phenacetin 2½ gr., aspirin 3½ gr | | Lloyd | Espinal, Tab. |
| d (sulf.) | 15 mg | Butabarbital ¼ gr., acetophenolsolone 5 mg | | Direct Labs | Nilax, Cap |
| d (sulf.) | 5 mg. | Butabarbital ½ gr., acetylsalicylic acid 2.5 gr., salicyl amide 2.5 gr., N-acetyl p-aminophenol 2.5 gr., hyoscine sulfate 0.1214 mg., atropine sulfate 0.0258 mg., hyoscyne HBr 0.0088 mg | | Stuart | Stuart Prelar 8, Tab |
| d (sulf.) | 5 mg | Butabarbital 0.5 gr., acetylsalicylic acid 2.5 gr., salicyl amide 2.5 gr., acetaminophen 2.5 gr | | Stuart | Stuart Prelar, Tab |
| d (sulf.) | 5 mg | Butabarbital sod 10 mg, methylcellulose 350 mg | | Carrick | Bantri, Tab |
| d (phos.) | 2.5 mg. | Butabarbital sod 8 mg, homatropine methylbromide 0.65 mg., bullogen, end vitamins | | Columbus | Obesoids, Tab |
| d (phos.) | 5 mg. | Caffeine 150 mg | | Irwin Neusser | Stimalose, Tab |
| d (phos.) | 5 mg | Caffeine 50 mg | | Merck Sharp & Dohme | Ampane, Tab |
| d (phos.) | 5 mg | Carbamazepine methylcellulose sod, 8-1 | | Cole Chem Co | Amcabese, Tab. |
| d (non) | 10 mg | Chlorpheniramine tartrate 8 mg., pyrilamine tartrate 15 mg | | Irwin Neusser | Nalartan Tab |
| d (sulf.) | 5 mg | Dienestrol 0.5 mg, methyltestosterone 2.5 mg., peefobarbital sod 30 mg | | Kremers Urban | Amperone, Tab |
| l (sulf.) | 5 mg | Dramamine 50 mg | | Scarfe | Oromamine O, Tab |
| (sulf.) | 2.5 mg | Ethisterone 5 mg., amn chloride 5 gr., 1 hyoscyamine HBr 0.0625 mg | | Columbus | Gymens, Tab |
| (sulf.) | 2.5 mg | Ethinyl estradiol 0.01 mg., methyl testosterone 2.5 mg plus Gerrat formula | | Iederte | Gevrestin Cap |
| (sulf.) | 1.25 mg | Ethinyl estradiol 5 mg., methyltestosterone 1.25 mg., nicotinic acid 50 mg., Vit C 25 mg | | Lemmon | Stimogen, Tab |
| sulf) | 30 mg | Glycerophosphates sod & pot., vit B 1 | | Merck Sharp & Dohme | Adelate, Liq. |
| ulf) | 15 mg | Homatropine methylbromide 0.72 mg., acetophenolsolone 5 mg., mephobarbital 15 mg., thyroid 90 mg | | Amco Drug | Amcobese Dakaps |
| sulf) | 2.5 mg | Homatropine MB 0.5 mg., caffeine 0.5 mg., amn chlor., 5 gr., vit B complex | | Purdue Frederick | Pre-Mens W/Amphetamine |
| (sulf.) | 2.3 mg | Hyoscine HBr 0.024 mg., hyoscyamine HBr 0.12 mg., pyrilamine maleate 75 mg | | Meyer | Histalme Duracaps |
| (sulf.) | 2.5 mg | Mephenezin 4 gr | | P.S. -L | .. |

Amphetamine Combinations

| Amphetamine | Others | Manufacturer | Brand |
|-------------|--|------------------------|-------------------------|
| d (sulf.) | Mephensin 250 mg., butobarbital 8 mg. | Crytal | Cryodes, Tab. |
| d (sulf.) | Mephensin 250 mg., butobarbital sod. 8 mg. | Paul Maney | Evidox, Tab. |
| d (phas.) | Mephensin 300 mg., butobarbital sod. 10 mg. | Cole | Mutas, Tab. |
| d (sulf.) | Mephensin 200 mg., chloral hydrate 200 mg. | Perron & Carey | Sometal, Cap. |
| d (sulf.) | Mephensin 400 mg., phenobarbital 1/8 gr., kametropine 1/5 mg. | Adams | Setosamine, Tab. |
| d (phas.) | Mephobarbital 8 mg., w/Vitamins | Rowell | Vio-Dosase, Tabs. |
| d (sulf.) | Mepramate 400 mg. | Lederle | Bomodes, Tab. |
| d (sulf.) | Mepramate 15 mg. | Lederle | Bomades Sequels, Caps. |
| d (sulf.) | Methandiol 1 mg., vit. B complex | Columbus | Carovoid, Tab. |
| d (phas.) | Methapriline HCl 25 mg., phenacetin 2 1/2 gr., aspirin 3 1/2 gr. | Pilmon-Moore | Cohistine, Tab. |
| d (phas.) | Methapyrilene HCl 25 mg. | Irwin Neisler | Obeetz T.F., Tab. |
| d (sulf.) | Methylcellulose 150 mg. | Columbus | Amladol, Tab. |
| d (phas.) | Methylandrostenediol 1 mg. | Irwin Neisler | Obeall, Tab. |
| d (phas.) | Methylcellulose 150 mg. | Irwin-Neisler | Obeall, liq./os. |
| d (phas.) | Methylcellulose 9 gr. | Beed & Cornick | Cyclin, Tab. |
| d (phas.) | Methylcellulose 500 mg. | Irwin, Neisler | Obeall liq. per os. |
| d (sulf.) | Methylcellulose 9 gr. | Sutlin & Case | Celladrine, Tab. |
| d (sulf.) | Methylcellulose 500 mg., vit. B-1 | Ulcacin | Obealin, Tab. |
| d (phas.) | Methylcellulose 0.5 Gm., vit. B-1, C | Lincoln | Obeall Complex, Tab. |
| d (sulf.) | Methylcellulose 100 mg., B complex | Lederle | Zelca-4 |
| d (phas.) | Methylcellulose 200 mg., 10 vit. & 11 minerals | Lemon | Amphelcel, Tab. |
| d (phas.) | Methylcellulose 7 1/2 gr., dicol. phos. 3 gr. | Wallace | Appetrol, Tab. |
| d (sulf.) | Milawa 400 mg. | Whitner | Pambromat, Tab. |
| d (sulf.) | Pambrom 50 mg., carbromal 150 mg., salicylamide 150 mg. | Canlight | Hyssen-AC, Tab. |
| d (sulf.) | Pentobarbital sod. 65 mg. | U.S. Standard Products | Damilal, Tab. |
| d (sulf.) | Pentobarbital sod. 1/4 gr., acetophenetidin 2 1/2 gr., aspirin 3 1/2 gr. | Myco Labs. | Sedes, Tab. T.D. |
| d (sulf.) | Pentobarbital 60 mg., phenobarbital 30 mg. | Morton Pharm. Co. | Tymafast, Cap. |
| d (sulf.) | Pentobarbital sod. 20 mg., Vit. C 50 mg. | Smith, Miller & Patch | Temporid, 5 cc. or Tab. |
| d (sulf.) | Pentylenetetrazol 100 mg., w/coffaine | Grey Pharm. | Dexzyme, Tab. |
| d (sulf.) | Pentylenetetrazol 120 mg., nicotie 50 mg., vit. B-1, B-2, C | Finley | Sedamosal, Cap. |
| d (sulf.) | Phenacelin 120 mg., salicylamide 200 mg., amobarbital 15 mg., hyascyamine suff. 0.05 mg. | | |

| Amphetamine | | Others | | Manufacturer | Brand |
|-------------|---------|---|---|-----------------------|-------------------------------|
| d (sulf) | 15 mg | Phenobarbital | 34 mg | Wynn Pharmacal Co | Dexatal, Tab |
| d (sulf) | 10 mg | Phenobarbital | ¼ gr | G F Harvey | Amphorb, Tab |
| d HCl | | Phenobarbital | | Dumas Wilson | Celudex, Tab |
| d (sulf) | 5 mg | Phenobarbital | ¼ gr | Gold Leaf | Amfebarb, Tab |
| d (phas) | 5 mg | Phenobarbital | ¼ gr, & 1.5 mg | Smith Miller & Patch | Bluban, Tab |
| d (phas) | 3 mg | Phenobarbital | ¼ gr, aspirin 5 gr | Clark & Clark | Pen Phetamine, Tab |
| d (sulf) | 5 mg | Phenobarbital | 16 mg, methylcellulose 160 mg, choline, methionine, B-12 & minerals | Lakeside | Obelip, Cap |
| d (sulf) | 5 mg | Phenobarbital | ¼ gr, methylcellulose 200 mg, B vit & minerals | Stuart | Amvical Cap |
| d (sulf) | 5 mg | Phenobarbital | 16 mg, methylcellulose 200 mg, mag trisilic 200 mg, vit B complex | Smith Dorsey | Nexarin, Cap |
| d (phas) | 15 mg | Phenobarbital | ¼ gr, solycylamide 3½ gr | Irwin Neuster | Analdine, Tab |
| d HCl | 5 mg | Phenobarbital | ½ gr, thyroid ½ gr | Pharm Dist | Lakabie, Tab. |
| d (sulf) | 5 mg | Phenobarbital | ¼ gr, thyroid ½ gr | Barre | Barre Formula |
| d (sulf) | 10 mg | Phenobarbital | ¼ gr, thyroid ½ gr | Pharm Dist | Lakabie, Tab |
| d (sulf) | 15 mg | Phenobarbital | ¼ gr, thyroid ½ gr | Pharm Dist | Lakabie, Tab |
| d (phas) | 5 mg | Phenobarbital, thyroid | atropine three strengths | Clark & Clark | Clarkotabs, Tab |
| d (sulf) | 15 mg | Phenobarb. ¼ gr, thyroid 3 gr, atrop self | 1/180 gr, atropine | Paul Maney | Nika Caps Ph 749 |
| d HCl | 5 mg | Phenobarbital | ¼ gr, thyroid ¼ gr, melasse, atropine sulf | Dumas Wilson | Antibesity, Tab. |
| d HCl | 5 mg | Phenobarbital | ¼ gr, vitamins | Rowell | Vio Dex, Cap |
| d (sulf) | 1/26 gr | Phenacetin | 2½ gr, aspirin 2½ gr | Sherman | Ampo Tab |
| d (phas) | 1/25 gr | Phenacetin | 2½ gr, aspirin 2½ gr | Irwin, Neuster | Threaze, Tab. |
| d (sulf) | 2.5 mg | Phenacetin | 2½ gr, aspirin 3½ gr | Hiss | Amphabar, Tab. |
| d (sulf) | 2.5 mg | Phenacetin | 2½ gr, aspirin 3½ gr | Pitman Maare | Dexagesit, Tab |
| d (sulf) | 2.5 mg | Phenacetin | 2½ gr, aspirin 3½ gr | Hiss | Amphobar, Tab. |
| d (phas) | 1/26 gr | Phenacetin | 2 gr, aspirin 4 gr also with codeine | G F Harvey | Harvadyne No. 4, Tab. |
| d (sulf) | 15 mg | Prachlorperazine dimaleate | 7.5 mg | Smith, Kline & French | Extatrol, Spansule |
| d (sulf) | 2.5 mg | Pyrilamine maleate | 12.5 mg | KLP, Lab | Diapep, Tab. |
| d HCl | 2 mg | Pyrilamine maleate | 25 mg | Testagar | Pymadex, Tab, or vial per cc. |
| d HCl | 5 mg | Pyrilamine maleate | 25 mg | Elder | Minukol, Tab |
| d (sulf) | 1.5 mg | Pyrilamine maleate 50 mg, hyoscin HBr 0.016 mg, hyoscyamine HBr 0.08 mg | | Meyer | Almehist, Cap. |

Amphetamine Combinations

| Amphetamine | Others | Manufacturer | Brand |
|-------------|--|-----------------------|-------------------------------------|
| d (phas.) | 1 mg. Pyrilamine maleate 12 mg., methapyrilene fumarate, solicylamide | Irwin, Heisler | Amsellin, Cap. |
| d.HCl | 6 mg. Pyrilamine maleate 25 mg., methapyrilene HCl 25 mg. | Testagor | Pymodes, Timed, Cap. |
| d (sulf.) | 5 mg. Phenamine maleate 10 mg. | Riber | Rouwidrene, Tab. |
| d (sulf.) | 15 mg. Rousolfio terp. alk. est. 1 mg. | Nyco Labs. | Secodes, Tab. T.D. |
| d (sulf.) | 2.5 mg. W/Secobarbital 60 mg. | Parke, Davis | Amphedose, Cap. |
| d (sulf.) | 10 mg. Toladiastase, nicotinamide, vit. B-1, C | New Eng. Ph. | Dastheroid, Tab. |
| d (sulf.) | 5 mg. Also W/Phenobarb. 1/4 gr. | Dumas-Wilson | Nubete, Tab |
| d (sulf.) | Thyroid 1 gr. | Lowe | Lowebese CTR, T. R. Cap. |
| d (sulf.) | Thyroid 16 mg. | Harco | Destlebs, Tab. |
| d. | Thyroid 3 gr., olein 1/4 gr., atropine sulf, 1/180 mg., phenobarbital 1/4 gr. | Taylor | Obetrie, T. R. Tab. |
| d (sulf.) | 5 mg. Thyroid 3 gr., atropine sulf, 2/360 gr., olein 1/4 gr., phenobarbital 1/4 gr. | Gold Leaf | Ampholdr, Tab. No. 1 |
| d.HCl | 10 mg. Thyroid 1 gr., atropine sulf, 1/360 gr., olein 1/4 gr., phenobarbital 1/4 gr. | Dumas-Wilson | Amphetomies Combina- tions, Tab. |
| d (phas.) | Thyroid, atropine sulf, olein | Various Mfr. | Ampholds, Tab, No. 2 |
| d (phas.) | Thyroid, atropine sulf | Varlaus Mfr. | Amphofin, Tab. |
| d.HCl | 5 mg. Thyroid 1 gr., atropine sulf, 1/360 gr. | Gold Leaf | Ampholdr, Tab, No. 3 |
| d (sulf.) | 5 mg. Thyroid 1 gr., phenobarb. 1/4 gr. | Normand | Amroid, Tab. |
| d (HCl) | 10 mg. Thyroid 1 gr., phenobarbital 1/4 gr. | Dumas-Wilson | Obesity, Cap. |
| d (HCl) | 15 mg. Thyroid 50 mg., phenobarbital 25 mg. | Dumas-Wilson | Raphetabz, Tab. |
| d (sulf.) | 2.5 mg. Vit. B-1, B-2, nicotin | Smith, Kline & French | Adjulets, Trache |
| d (phas.) | 5 mg. Vit. B-1, B-2, C, niacinamide | Straenburgh | Obet T.C. |
| d (phas.) | 5 mg. Vit. A, D, B-1, B-2, B-6, C, niacinamide | Wyeth | |
| d (sulf.) | 15 mg. Vit. B-1, B-2, A, D, B-12, C, niacinamide thenate | Columbus | |
| d (sulf.) | 5 mg. Vit. B-1, B-2, B. 6, niacinamide & pantothenol/cc. | Vitamix | Amphetplex, Sol./cc. |
| d (sulf.) | 5 mg. Vitamins (8) and minerals (11) | Roeig | AMPlus, Cap. |
| d (sulf.) | 10 or 15 mg. Vitamins and minerals | Ethical Pharm. | Daycap, Cap. |
| d (sulf.) | 0.5 mg. Vitamins and hormones | Roeig | Neabee, liq./5 cc |

AMPHETAMINE PHOSPHATE

52

- Raphetamine Elixir NND 1958, Tab & Vial (Strassenburgh)
- * **AMPHETAMINE PHOSPHATE**, Dextro, NND 1962 NFs Tab NP Dextro amphetamine phosphate Tab (5 mg) (Various Mfr) See: Dextro Proletamine, Tab (Clark & Clark)
- * **AMPHETAMINE SULFATE**, USP, NNR 1957, Tablets NF (Racemic or dl form) Cap—(5 10 mg) (Various Mfr) Tab—(5 10 mg) (Various Mfr) Vials—(20 mg/cc) (Various Mfr)
- See Amirene Tab (Normand)
Amphetamine Tab (Van Pelt & Brown)
Amphids S Tab (Gold Leaf)
Benzedrine, Amp. Tab., NNR (Smith, Kline & French)
Linampheta, Tab (Lincoln)
- * **AMPHETAMINE SULFATE**, Dextro USP NND 1962 Dextro amphetamine sulfate Cap—(2 5 5 10 mg) (Various Mfr) Tab USP—(5, 10 mg) (Various Mfr) Vials USP—(20 mg/cc) 30 cc (Gatham)
- See: Adrixine, Tab (Key Corp)
Am Dex, Tab (Superior)
D Antelasul, Tab (Pilman Moore)
Amilrene, Tab (Normand)
Amphetamine Tab (Van Pelt & Brown)
Amphex, Tab (Royce)
Amphex Tab (Algra)
Amsustain, Tab (Key)
Cradex Cap (Craig)
Dadex, Cap (Central)
D Citramine, Tab (Prestan)
DAS, Tab (Stayer)
D Ale Ph 747, Cap (Paul Money)
Dexalone Tabs (Wyna)
Dexedrine Elix Granules Spanules (Smith, Kline & French)
Dex OB Tab (Tully Pharmaceuticals)
Dextrosule Cap (Starck)
Dex Sule Cap (Normal)
Diacurb Perle (Tutag)
Evradox, Tempules (Evrans)
Hetamine Tab (Dumas Wilson)
Lawedex CTR (Lowe)
Medex Tab (Medca)
Nilex Caps (Direct Labs)
Obesant, Tab (Lincoln)
Pellcaps Tab (Kirkman)
Tydex TR Cap (Tyler)
- d-**AMPHETAMINE TANNATE**
See: Synalan, Tab (Irwin Neisler)
W/Secobarbital See: Seca Synalan, Tab (Neisler)
- leva-**AMPHETAMINE**, Levamphetamine See Ad N1 Tab (Medica)
Amphetamine Tab (Vanpelt & Brown)
- * Generic name recognized by NF, USP or AMA
- Lavaba Tab (Superior)
Levonor, Tab (Nordmark)
- AMPHETOCEL** (Lemmon) d Amphetamine phosphate 2.5 mg, dicalcium phosphate 1 gr, methocel 7½ gr/Tab Bol 100s
Use: (2 fl d) Appetite depressant.
- AMPHETOID** (Horlan & Converse) No 1—Methamphetamine HCl 5 mg, thyroid 1 gr, atropine sulfate 1/360 gr, alain ¼ gr/Tab No 2—Methamphetamine HCl 5 mg, thyroid 1 gr, atropine sulfate 1/360 gr/Tab No 3—Methamphetamine HCl 5 mg, thyroid 1 gr, phenobarbital ¼ gr/Tab Bol 100s 500s
Use: Central nervous system stimulant
- AMPHETOPLEX**, (Vitamix) d Amphetamine sulfate 5 mg, thiamine 25 mg, riboflavin 5 mg, pyridoxine 5 mg, Vit B 12 30 mcg, niacinamide 50 mg, panthenol 5 mg/cc Soln
Use Appetite depressant
- AM-PHET UNODAY Ph. 758**, (Paul Money) Amobarbital 1½ gr, dextroamphetamine sulfate 15 mg/cap Bol 30s
Use: Depression and sedation
- AMPHEX**, (Algra) d Amphetamine sulfate Tab (5 mg) Bol 100s, 1000s
Use: Central nervous system stimulant
- AMPHOBAR**, (Hiss) Amphetamine sulfate 2.5 mg, phenacetin 2½ gr, aspirin 3½ gr/tab Bol 100s
Use: Analgesic with central nervous system stimulant
- AMPHOBESE**, (Starr) No 1—Methamphetamine HCl 10 mg, thyroid 1 gr, alain ¼ gr, atropine 1/360 gr/Tab No 2—Methamphetamine HCl 10 mg, thyroid 1 gr, atropine 1/360 gr/Tab No 3—Methamphetamine HCl 10 mg, thyroid 1 gr, phenobarbital ¼ gr/Tab Bol, Sets: 250s 750s 1000s, 3000s
Use: Anti obesity
- AMPHOCORTIN**, (Warner Chilcott) A water washable cream containing Amphetamine, neomycin and hydrocortisone
- AMPHOIDS** (Gold Leaf) No 1—Amphetamine sulfate racemic, 5 mg, thyroid 1 gr, atropine sulfate 1/360 gr, alain ¼ gr/Tab No 2—Amphetamine sulfate, racemic 5 mg, thyroid 1 gr, atropine sulfate 1/360 gr/Tab No 3—Amphetamine sulfate, racemic, 5 mg, phenobarbital ¼ gr, thyroid 1 gr/Tab 1000s 3000s
- AMPHOIDS-S** Gold Leaf Amphetamine sulfate 10 mg, each
Use: Central nervous system stimulant
- AMPHOJEL**, NND 1961 (Wyeth) Aq susp of aluminum hydroxide, equivalent to 4%

aluminum oxide. Bot. 12 fl. oz. Tab. (5 gr.)
Box 30s, Bot. 100s.
Uter: Gastric & duodenal ulcer or hyper-
acidity.
Without flavor. (Wyeth) Bot. 12 fl. oz.
W/Magnesium trisilicate. (Wyeth) Contains
magnesium trisilicate 12%. Bot. 12 fl. oz.
Uter: Gastric antacid.
W/Mineral oil. (Wyeth) Liquid petrolatum
10%. Bot. 12 fl. oz.
Tablet (Dried aluminum hydroxide gel)
(Wyeth) Tab. (5 gr., 10 gr.) Box 30s, Bot.
100s.
* **AMPHOMYCIN**. An antibiotic produced
by *Streptomyces canus*.
AMPHOPHEN. (Gold Leaf) Aspirin 3½ gr.,
acetophenetidin 2½ gr., caffeine ½ gr.,
phenobarbital ½ gr./Tab. Bot. 100s, 1000s.
Uter: Analgesic, sedative.
* **AMPHOTERICIN B**, U.S.P., N.N.D. 1962. A
polyene antibiotic substance obtained from
cultures of *Streptomyces nodosus*. Inj
U.S.P. See: Fungizone (Squibb)
W/Tetracycline phosphate potentiated. See:
Mystecin-F, Prep. (Squibb)
AMPHYL. (Lehn and Fink) A combination of
ortho-phenyl phenol and p-tert amyl phenol
w/pol. ricinoleate in propylene glycol &
alcohol. Bot. 1 pt., 1 gal.
Use: (¼ to 2%) General disinfecting pur-
pose
AMPHYROID. (Hance) di-Desoxyephedrine
HCl 1/12 gr., thyroid 1 gr., atropine sulfate
1/360 gr., alain ½ gr./Tab. Bot. 250s,
1000s.
Uter: Anti-obesity.
* **AMPICILLIN**, (Bristol) 6-[(1- α -aminophenyl-
acetamido) penicillanic acid. Penbritin.
Uter: Antibiotic.
AMPLICAPS. (Hart) Vit. A 5000 u., D 1000
u., B-1 1.5-mg., B-2 2.0 mg., B-6 0.1 mg.,
C 37.5 mg., niacinamide 20 mg., calcium
pantothenate 1 mg., Vit. E 2.27 mg./Cap.
Bot. 100s, 1000s.
Uter: (1 daily) Multivitamin therapy.
AM PLUS. (Roerig) Dextro-amphetamine sul-
fate 5 mg., Vit. A 5000 u., D 400 u., B-1
2 mg., B-2 2 mg., B-6 0.5 mg., niacinamide
20 mg., Vit. C 37.5 mg., Cal. pantothenate
3 mg., Col. 242 mg., cobalt 0.1 mg., copper
1 mg., iodine 0.15 mg., iron 3.33 mg.,
Mang. 0.33 mg., molybdenum 0.2 mg., Mag-
2 mg., Phos. 187 mg., Pot. 1.7 mg., zinc
0.4 mg./cap. Bot. 100s.
Uter: (1 daily a.m.) Anti-obesity.
AMPLUS IMPROVED. (Roerig) Vitamins ap-
proximately one-half of Amplus include
* Generic name recognized by N.F., U.S.P. or A.M.A.

hydroxyzine HCl 5 mg & d-Amphetamine
sulf. 5 mg./Cap. Bot. 100s.
Uter: Obesity control.
AMPOPHOS. (Burlington) Strychnine cacody-
late 1/120 gr., Sod. glycerophosphate 1½
gr., iron cacodylate 1/6 gr., procaine HCl
1%, water q.s./cc. 1 cc. ampuloid & 30 cc.
vial.
Uter: S.C., I.M., for secondary anemia
AMPROL. (Merck & Co., Inc.) Procaine peni-
cillin 4 Gm., streptomycin 30 Gm./Tan.
Uter: Coccidiostat.
AMPROTROPINE PHOSPHATE, N.N.R. 1950.
2-Oethylamino-2,2-dimethylpropyl tropate.
See: Syntropan, Prep. (Rache Lab).
AMPYRIMINE PHOSPHATE. (Smith, Kline
& French) 2,4,7-triamino-5-phenylpyrimido-
[4,5-d]-pyrimidine, phosphate hemihydrate.
Uter: Diuretic.
AMPYROX. (Elder)
Elixir: Scopolamine methylbromide 1 mg.,
butabarbital sod. 10 mg./5 cc. Bot. 1 pt.
Tab.—276 Scopolamine methylbromide 2
mg./tab. Bot. 100s.
Tab.—281 Scopolamine methylbromide 2
mg., butabarbital sod. 15 mg./tab. Bot.
100s.
Tab.—293 Scopolamine methylbromide 1
mg., aluminum hydroxide 250 mg., magne-
sium hydroxide 60 mg./tab. Bot. 100s.
Uter: Anticholinergic.
AMRIL. (Amfre-Grant, Inc.) Acetylcarbromal
200 mg., mephensin 150 mg., reserpine
0.05 mg./Tab. Box 50s, 100s.
Uter: Sedative.
AMROID. (Dumas-Wilson) Amphetamine HCl
10 mg., thyroid 1 gr., phenobarbital ¼ gr./
Tab. Bot. 100s, 1000r.
Uter: Control and management of obesity.
AMSALIN CAPSULES. (Irwin, Neizer) Pyr-
rilamine maleate 12 mg., phenylpyramine
tartrate 12 mg., salicylamide 150 mg., d-
amphetamine phosphate 1 mg./Cap. Bot.
100s, 500s, 1000s.
Uter: Functional dysmenorrhea.
AMSODYNE. (Elder) Phenobarbital ¼ gr.,
belladonna ext. ½ gr., acetylsalicylic acid
5 gr., caffeine ¼ gr./Tab. Bot. 100s, 500r,
1000s
Uter: Analgesic and sedative
AMSUSTAIN. (Key Corp.) Dextro-ampheta-
mine sulfate. Tab. (7½ & 15 mg) Bot. 30s,
500s
Uter: Control of appetite
A-M-T. (Wyeth) Aq. surp. of aluminum hy-
droxide gel with 12.5% of magnesium tri-
silicate. Bot. 12 fl. oz.
Uter: Gastric antacid.
Tablets. (Wyeth) Aluminum hydroxide 2.5 gr.

- w/magnesium trisilicate 3.85 gr Tab, box 30, bottle 100s
 Use Gastric antacid
- AMTENS** (Hoack) Reserpine 0.1 mg, homatropine methylbromide 0.5 mg, NH_4Cl 500 mg, caffeine 30 mg /Tab Bot 100s
 Use Tranquilizer
- AMUREX** (Reid) dl Methionine 200 mg /Cap Bot 500s 1000s
 Use Diaper rash
- AMVERO** (Mallord) Veratrum viride ext 32 mg, choline methionine tartrate 100 mg, mannitol hexanitrate 16 mg, rutin 20 mg, phenobarbital 16 mg /Tab Bot 100s, 500s, 1000s
 Use Functional hypertension
- AMVICEL** (Stuart) d Amphetamine sulfate 5 mg, Vit A 1700 u, C 170 u, B₁ 1 mg, B₂ 1 mg, B₆ 0.15 mg, niacin 10 mg, pantothenic acid 16 mg, methylcellulose 225 mg, Vit B₁₂ 1 mcg, phenobarbital $\frac{1}{4}$ gr, with calcium, iron, P₆₀, copper, iodine, cobalt, zinc & manganese /Cap Bot 100s
 Use Anti obesity
- AMVICEL-X10** (Stuart) Phenobarb 15 mg, amobarbital 40 mg, d amphetamine sulf 10 mg, 7 minerals 9 vit including B₁₂ /Tab Bot 100s
 Use: Obesity control
- AMVICEL-X15** (Stuart) Phenobarbital 20 mg, amobarbital 60 mg, d amphetamine sulfate 15 mg, 7 minerals 9 vitamins including B₁₂ /Tab Bot 100s
 Use: Obesity control
- AMVITDL** (Walker) Choline dihydrogen citrate 10 mg, dl methionine 15 mg, dl tryptophane 10 mg, l histidine HCl 20 mg, thiamine 10 mg, niacin 40 mg, riboflavin 2 mg, pyridoxine 5 mg/cc Amp 1 cc, Box 6s, 100s
 Use: IM (12 cc 3 times weekly) Lipotropic in liver disorders
- AMYALDEX UNICELLES #1** (Hiss Pharm) Dextroamphetamine sulfate 15 mg, amobarbital 60 mg /cap Bot 100s
 Use: (one a.m.) Delayed action, sedative w/CNS stimulant
- AMYALDEX UNICELLES #2** (Hiss Pharm) Dextroamphetamine sulfate 15 mg, amobarbital 100 mg /cap Bot 100s
 Use: (one a.m.) Delayed action, sedative w/CNS stimulant
- AMYDRICAIN HYDROCHLORIDE**, N N R 1947 (No Mfr currently lists) Alypen hydrochloride
 Use: Local anesthetic.
- AMYL-AMINOENZDATE** W/Ethyl para-amino-benzoate, cod liver oil, Sod phosphate
 * Generic name recognised by N.F., U.S.P. or A.M.A.
- panate See: Ultracain, Oint (Herzog)
- AMYLASE**. See: Fartizyme, Tabs (Breon)
 W/Protease lipase See: Trulase, Tabl (Smith, Miller & Patch)
- AMYLCAINE HYDROCHLORIDE**, N N E 1944 See Amyline hydrochloride (Novo coll)
- *AMYLENE NYDBATE**, U.S.P., N N R 1941 Tertiary amyl alcohol (Various Mfr)
 W/Tribromethanal See: Averlin (Winthrop Stearins)
- *AMYL NITRITE**, U.S.P. Isoamyl nitrite Burroughs Wellcome—Vaporate (0.18 cc (0.3 cc) Box 12s
 Lilly—Acpisals (0.18 cc, 0.3 cc) Box 12s
 Parke, Davis—Amp (0.31 cc) Box 12s 100c
 See: Vaporate, Amp (Burroughs Wellcome)
 Use Inhalation, coronary vasodilator in angina pectoris
- AMYLOCAINE HYDROCHLORIDE**, Benzoyl ester of methyl ethyl dimethyl aminomethyl carbinal See Slavaine (Wallau)
- AMYLOPINE**, (First Texas) Phenobarbital W/Atropine (First Texas) Phenobarbital $\frac{1}{4}$ gr, atropine sulfate 1/1000 gr /Tab Bot 100s, 500s, 1000s
 Use Sedative antispasmodic
 W/Belladonna (First Texas) Phenobarbital Sod 1 gr, belladonna pow, ext $\frac{1}{4}$ gr /Tab Bot 100s, 500s, 1000s
 Use Sedative-antispasmodic
 W/Ephedrine (First Texas) Phenobarbital $\frac{1}{4}$ gr, ephedrine sulfate $\frac{1}{4}$ gr /Cap Bot 100s 500s, 1000s
 Use Sedative vasoconstrictor
- AMYLOFENE COMPOUND**, (First Texas) Phenobarbital 1 $\frac{1}{2}$ gr, aminopyrine 3 $\frac{1}{2}$ gr /Cap Bot 100s, 500s, 1000s
 Use Sedative analgesic
- AMYLOFENE SODIUM**, (First Texas) Phenobarbital Sod Cap (1 $\frac{1}{2}$ or 3 gr) Bot 100s 500c 1000s
 Use Hypnotic
 W/Piperidolate HCl, cellulolytic enzyme proteolytic enzyme, pancreatin 3X, taurocholic acid See Doctilose, Tabl (Lake-side)
- AMYLOLYTIC ENZYME**
 W/Piperidolate hydrochloride cellulolytic enzyme, proteolytic enzyme, pancreatin 3X taurocholic acid See Doctilose, Tabl (Lake-side)
 W/Proteolytic enzyme d sorbitol See Kuzyme Cap (Kremers Urban)
 W/Proteolytic, cellulolytic See Trienzyme Tabs (Testagar)
 W/Proteolytic enzyme, cellulolytic enzyme aluminum hydroxide gel See: Halozym[®] Tabs (Holco)

W/Proteolytic enzyme, lipolytic enzyme. See: Formulose, Tabs. (Kremers-Urban)

W/Pyridosine HCl, pyrilamine maleate, proteolytic enzyme See: Nemisis, Tabs. (Beutlich)

W/Proteolytic enzyme (Papain), homatropine methylbromide, d-sorbitol. See: Converzime, Liq. (Ascher)

AMYLISINE-BENZOCAINE, A.D.R. (Novocoll)
Oint.—Amylsine base 4 Gm., benzocaine 20 Gm., clove oil 2 Gm./100 Gm. of lanolin-petrolatum base. Sol.—Amylsine base 4 Gm., benzocaine 18 Gm., clove oil 37 Gm., & wintergreen oil
Use: Topically, anesthetic.

AMYLISINE HYDROCHLORIDE, N.N.D. 1962, A.D.R. (Novocoll) (Amylcaine HCl) 2-Amyl-aminocethyl p-aminobenzoate HCl Pow. Vials (5 Gm.); Bot. (28.3 Gm.); Sol. (4%) Bot. 30 cc.
Use: 2 or 4% sol. for corneal anesthesia where mydriasis is not desired.

AMYLTRICRESOL, SEC., N.N.R. 1952, A.D.R. (Upjohn) Prepared by the reaction of cresol with secondary-amylic alcohol.
Use: Antiseptic.

AMYTAL, U.S.P., N.N.R. 1950. (Lilly) Amobarbital.
Elixirs (2 gr./oz.), Bot. 1 pt., 1 gal.; (4 gr./oz.), Bot. 1 pt., 1 gal.
Pow.: Bot. ½ oz., 4 oz.
Tab.: (8 mg.), Bot. 100s, 500s; (16 mg.), Bot. 100s, 500s; (32 mg.), Bot. 100s, 500s; (50 mg.), Bot. 100s, 500s; (D.1 Gm.), Bot. 100s, 500s.
Use: Hypnotic.

W/Acetylsalicylic acid. (Lilly) Amytal ¾ gr., acetylsalicylic acid 5 gr./pulfules, Bot 100s, 500s.
Use: Analgesic sedative.

W/Aminophylline. See: Aminophylline w/ Amytal, Pulv. (Lilly).

W/Aminophylline & ephedrine HCl. See: Amesec, Cop. (Lilly)

W/Aminopyrine. (Lilly) Amytal 1½ gr., aminopyrine 3½ gr./pulfule. Bot. 100s, 500s.
Use: (1 t.i.d., h.s.) Analgesic-sedative.

W/Etarin sulfate. See: Co-Etarin, Tab. (Lilly)

W/Ephedrine. See: Ephedrine w/amytal, Cop. (Lilly).

W/Ephedrine HCl & aminophylline. See: Amesec, Cop. (Lilly).

W/Foveril phosphate. See: Foveril & Amytal, Tab. (Lilly).

AMYTAL SODIUM, U.S.P., N.N.R. 1950. (Lilly) Amobarbital sodium.
Amp.s (1 gr.) Pkg. 6s, 100s; (1½ gr.) Pkg.

6s, 100s; (3¼ gr.)/pkg., (7½ gr.)/pkg., (15½ gr.)/pkg.—each pkg. contains amp. of water for injection.

Amp.: (3¼ gr.), Pkg. 6s, 25s, 100s; (7½ gr.), Pkg. 6s, 25s, 100s; (15½ gr.), Pkg. 6s, 25s, 100s, without amp. of water for injection.

Enseals: (3 gr.), Bot. 500s.

Pulfules: (1 gr.), Bot. 100s, 500s; (3 gr.), Bot. 100s, 500s.

Supp.: (3 gr.), Pkg. 12, 144

Use: Orally, i.m., i.v., rapid-acting sedative.

W/Secalon sodium. See: Tuinol, Cop. (Lilly).

ANABILE (Irwin, Nelsler) Conjugated cholic acid from ext. of ox bile, Tab. 1½ gr., Bot. 100s, 500s, 1000s

Use: (1 to 2 t.i.d.) Choleric.

ANABOLIC NORMONE. These agents stimulate constructive processes leading to retention of nitrogen and increasing the body protein.

See: Depo-Norlesterone. (Upjohn).

Fluoxymesterone.

Methyl Androstenediol. (Vorious Mfr.).

Stanolone, N.N.D.

ANACHLORIC. (Upjohn)

A (Bismuth subcarb. 10 gr., calcined magnesio 10 gr., Sod. bicarb. 10 gr./Tab.)

B (Bismuth subcarb. 10 gr., calcined magnesio 10 gr./Tab.)

C (Bismuth subcarb. 10 gr., Sod. bicarb 20 gr./Tab.)

D (Calcined magnesio 10 gr., Sod. bicarb. 10 gr./Tab.)

E (Col. carb. 10 gr./Tab.)

G (Bismuth subcarb. 10 gr., Col. carb. 10 gr., Sod. bicarb. 10 gr./Tab.) Bot. 100s, 1000s.

Use: Antacid.

ANACIN. (Whitehall) Aspirin, acetophenetidin 3 gr., caffeine, Tabs., Pkg. 4s, 72s, 30s, 50s, 100s

Use: Analgesic

ANADREX. (Anahist) 1-Desoxyephedrine, methyl salicylate, menthol, eucalyptus and spearmint oils. Inhaler.

Use: Treatment of nasal and sinus congestion.

ANADROL. (Syntex) Oxymetholone 2.5 mg. (2-hydroxy-methylene-17 α -methyl-17 β -hydroxy-3 androstanone) Tab. Bot. 50s
Use: Anabolic steroid therapy.

ANADYN. (Paul Maney) Salicylamide 1¼ gr., acetophenetidin 1¼ gr., caffeine alkaloid ½ gr./Tab. Bot. 1000s.

Use: (1-2 q. 3 h) Analgesic-antipyretic.

W/Phenobarbital. (Barlow-Maney) Phenobarbital ½ gr., acetophenetidin 2½ gr., salicylamide 3½ gr., caffeine alkaloid ½

gr/Tab Also half strength for children
Bot 1000.

Use (1 2 q 3 h) Analgesic sedative

ANAFAC. (Pharmalac) Methampyrone 300 mg, *n*-acetyl paminophenol 100 mg, ad phenine HCl 10 mg, phenyltoloxamine citrate 20 mg/Cap Bot 100s

Use: Analgesic

ANAFEBRINA. See: Aminopyrine (Various Mfr)

ANAFUNG. (Deritin Chemical) Sodium undecylenate sodium propionate, 3,4,4-tri chloroacarbon lide, salicylic acid, menthol camphor, cream Tube 1 oz Shaker cap 2 oz Aerosol can 6 oz

Use Fungal infections

ANAGINE (Mastengill) Aspirin 5 gr, acetophenetidin 3½ gr, caffeine ½ gr, atropine sulfate 1/300 gr/Cop Bot 500s, 1000s

Use: (1 Cop q 3-4 h) Analgesic

ANAGILL. (Mastengill) Aspirin 6½ gr, caffeine ½ gr, gelicemium lr 1½ min, propylene glycol/3 cc Bot 1 pt, 1 gal

Use (1 2 dr q 3 h) Analgesic

ANAHIL (Irwin, Neisler) Cholic acids 1½ gr as the Sod salts/Tab Bot 100s, 500s, 1000s

Use: Chronic gallbladder disease

ANALIST, NND 1962 (Anahist Co) Thonzylamine hydrochloride Tab Bot 15s, 40s, 100s

Use: Antihistaminic

ANALIST, SUPER (Anahist Co) Thonzylamine HCl 25 mg, ascorbic acid 20 mg, aspirin 3½ gr phenacelin 1½ gr, caffeine ½ gr/Tab or 5 cc syr

Use Colds

ANALBALM. (Central) Methyl salicylate, menthol, camphor liq 1 pt, 1 gal

Use Topically counterirritant to reduce inflammation

ANALDINE. (Irwin Neisler) d Amphetamine phosphate 15 mg, salicylamide 3½ gr, phenobarbital ¼ gr/Tab Bot 100s 500s, 1000s

Use: (1 q 3-4 h) Analgesic, relaxant

ANALDYNE. (Mastengill) Aspirin 2½ gr, acetophenetidin 1½ gr, phenobarbital ¼ gr, hyoscyamus ext 1/6 gr/Cop Bot 100s, 500s, 1000s

Use: (1 q 3-4 h) Analgesic, relaxant

ANALEPTICS. Usually a term applied to agents with stimulant action, particularly on the central nervous system See also central nervous system stimulants

See Amphetamine Salts (Various Mfr)
Bluban, Tab (Patch)
Camphor in Oil (No Mfr currently lists)
Cardiozal (Bilhuber Knoll)

Coramine, Amp, liq (Bilhuber Knoll)

Dextro Amphetamine Salts. (Various Mfr)

Ephedrine Salts (Various Mfr)

Labelline, Amp. (Sandoz)

Methamphetamine Salts (Various Mfr.)

Metrazol, Amp (Bilhuber Knoll).

Nikethamide (Various Mfr).

Pentylene tetrazol.

Picrotoxin, Amp (Various Mfr).

Sodium Sulfinate (Various Mfr).

Saduxin (Brewer).

Strychnine Salts (Various Mfr)

ANALEPTONE. (Reed & Carnrick) Pentylene tetrazol 200 mg, nicotinic acid 1½ gr/4 cc Elixir, Bot 8 oz, 1 pt Tab, Pentylene tetrazol 100 mg, nicotinic acid 50 mg-pepsin/Tab Bot 100s

Use: Vasomotor and cortical centers stimulant

ANALEPTONE-ANABOLIC. (Reed & Carnrick) Pentylene tetrazol 50 mg, methyltestosterone 0.83 mg, ethinyl estradiol 0.0015 mg, strontium salicylate 450 mg/Tablet Bot 100s

Use: Degenerative processes of geriatrics

ANALEXIN. (Irwin, Neisler) Phenylramidol NCI 200 mg/Tab Bot 100s, 500s Caps 400 mg Bot 50s

Use: Relief of pain and associated muscle tension

ANALEXIN-AP. (Irwin, Neisler) Phenylamidol 100 mg, aluminum aspirin 300 mg/Tab Bot 100s

Use: Relief of pain and associated muscle tension

ANALEXIN SYRUP. (Irwin, Neisler) Phenylamidol salicylate 100 mg/5 cc Bot 1 pt

Use: Relief of pain

ANALEXIN-400. (Irwin, Neisler) Phenylamidol HCl 400 mg/Cap Bot 50s

Use: Analgamyloxants; analgesic; skeletal muscle relaxant

ANALGECAIN (Metro) Procaine (base) 0.5%, benzocaine 3.0%, butyl aminobenzoate 1.0%, benzyl alcohol 5.0% in sesame oil Vial 10 cc.

Use: Injection (2.5-3 cc) Anal fissures, pruritus ani & ulcers

ANALGEMUL (Tilden) Camphor, menthol & methyl salicylate in a vegetable oil base Bot 4 oz, 1 pt, 1 gal

Use: Topically, rubefacient, counterirritant

ANALGEMUL OINTMENT. (Tilden) Methyl salicylate 1%, histamine dihydrochloride 0.1%, methyl salicylate 10%/Tube 1½ oz

Use: Analgesic; hyperemic counterirritant

ANALGESIC BALM. Menthol w/methyl salicylate in a suitable base

Lilly—½ oz, 1 oz, 1 lb, 5 lb, 10 lb

Parke Davis—½ oz, 1 lb, 5 lb

Merck Sharp & Dohme—1 oz., 1 lb.
Squibb—1 oz.
Use: Topically, analgesic-counterirritant.

ANALGESIC BALM. (Massengill) No. 2 methyl salicylate, menthol, camphor, eugenol, oil of eucalyptus, in a nongreasy base. No. 3 chloral hydrate 10 gr./oz., methyl salicylate, camphor, menthol and essential oil, in a nongreasy base. No. 2 Tebe, 1 oz., for 1 lb. No. 3 for 1 oz., 1 lb.
Use: Anodyne, analgesic and counterirritant.

ANALGESIC CALAMINE OINTMENT. See Caligesc Oint. (Merck Sharp & Dohme)

ANALGESICS, A.D.R. 1953. See preparations at: Acetanilid-type, Antipyrine-type, Aspirin, Cinchophen, Narcotic, Salicylamide.

ANALGESINE. See: Antipyrine. (Various Mfr.)

ANALGESTINE. (Mallord) Dover's powder $\frac{1}{4}$ gr., acetophenetidin $1\frac{1}{2}$ gr., camphor monobromated $\frac{1}{4}$ gr., aspirin 2 grs., caffeine anhyd. $\frac{1}{8}$ gr., atropine sulfate 1/500 gr./Cap. Bot. 100s, 500s, 1000s.
Use: Analgesic.

ANALGIA A.P.C. (Merrell) Aspirin 0.23 Gm., acetophenetidin 0.15 Gm., caffeine 15 mg./Tab. Bot. White 100s, White or Orange 1000s.
Use: Analgesic-antipyretic.

ANAMEBA. (Chicago) Iodochlorhydroxyquin, 125 mg., bacitracin methylene disalicylate equiv. to 5,000 U.S.P. u. of bacitracin activity./Tab. Bot. 24c.
Use: Oral omeblide.

ANANASE. (Rare) Proteolytic enzymes 50,000 u./Tab. Bot. 100c.
Use: Inflammation.

ANASMA. (Amre-Grant) Ephedrine sulfate 25 mg., theophylline calcium salicylate 100 mg., methapyrilene HCl 20 mg., potassium iodide 200 mg./Tab. Bot. 100c, 1000s.
Use: Antispasmodic-Sedative.

ANASORB. (Marion Labs.) Ferrus fumarate 300 mg., D-carbitol 300 mg., Vit. B-12 5 mcg., Intrinsic factor $\frac{1}{4}$ u., folic acid D.5 mg., ascorbic acid 75 mg./Tab. Bot. 60s, 500s.
Use: Hemolitic.

ANATOID SUPPOSITORIES. (Miller Lab.) Bismuth salicylate 4.41%, benzocaine 3.67%, menthol 1%, resorcin D.59%, E-hydroxyquinoline 0.37%, & q.c. with base to 100%. Box 9s.
Use: Rectally, anesthetic.

ANATOLA. (Parke, Davis) Vit. A palmitate syn. Amp.—(50,000 units) 1 cc., 10s. Cap.—(25,000 units), boxes 100s, 500s, (50,000 units). Boxes—30s, 100s, 500s.
Use: Vitamin A deficiency.

ANATOX. (Anotax) Tab. Bot. 20s.

ANATOXIN. (Roman). See: Diphtheria Toxoid, U.S.P.

ANATUSS TABS. (Maynard) Glyceril guaiacolate 50 mg., N acetyl-P-aminophenol 200 mg., d-methorphan HBr 10 mg., phenyl-ephedrine NCl 5 mg., phenylpropanolamine NCl 25 mg., chlorpheniramine maleate 2 mg., Vit. C 100 mg./Tab. Bot. 100s.
Use: Relief of coughing.

ANAYODIN, N N R. 1950. (Bischoff) Chiniofon pills E.C. 0.25 Gm. Bot. 50s, 100s, 1000s. Pow., Bot. 25s, 100s.
Use: (1-2 q.l.d.) Acute and chronic amebiasis.

ANCILLIN. (Smith, Kline & French) Sodium Diphenicillin.
Use: Antibiotic.

ANCYTAL. (Massengill) Aminopyrine $3\frac{1}{2}$ gr., acetophenetidin 1 gr., caffeine $\frac{1}{2}$ gr., hyoscyamus ext. $\frac{1}{4}$ gr./Tab. Bot. 100s, 500s
Use: (1 tablet p.r.n.) Analgesic.

AND-EST INJ. (Ulmer) Estrone 2 mg., testosterone 25 mg./cc. aq. susp. Vial 10 cc.
Use: (1 cc. I.M.) Menopausal syndrome, androgen-estrogen therapy.

ANDESTERONE. (Lincoln) Estrogens conjugated 1.25 mg., methyl testosterone 10 mg./Tab. (Also, 0.325 mg., 0.65 mg./Tab) Bot. 100c, 1000s. Aq. Sulp. 15 cc.
Use: Androgen and estrogen therapy.

ANDIN. (Estr) Vit. A 50,000 u., Vit. D 10,000 u./cc. Amp. 2 cc., Vial 10 cc.
Use: Vitamin A & D therapy.

ANDIRA. (Paul Money) Citrus pectin $2\frac{1}{2}$ gr., calcium gluconate $1\frac{1}{2}$ gr./tab. Bot. 100s, 1000s. Pow. 4 oz., 1 lb.
Use: (2 q. 4 hrs.) Diarrhea.

ANDIFIVE. (Massengill) Aminopyrine $3\frac{1}{2}$ gr., acetophenetidin 1 gr., caffeine $\frac{1}{2}$ gr., hyoscyamus ext. $\frac{1}{4}$ gr./Cap. Bot. 500s
Use: Analgesic

ANDOIN. (Ulmer) Allantoin 1% w/vitamin A & D. Cream 1 oz., 1 lb.
Use: Topically, stimulates granulation.

ANDOSIX. (Massengill) Acetylsalicylic acid 5 gr., phenobarbital $\frac{1}{2}$ gr., hyoscyamus ext. 1/6 gr./Cap. Bot. 100s, 1000s.
Use: Analgesic and sedative

ANDOTROPE. (Rawell) Pituitary gonadotropins. Vial 5 cc.
Use: Pituitary hypofunction.

ANDRESTRAQ. (Central) Estrogenic substance natural 20,000 units, testosterone 10 mg., procaine HCl 1.0% & chlorbutanol/ cc. Vial 10 cc.
Use: I.M. (1 or 2 cc. weekly) Menopause symptoms.

ANDRESTRAQ-M

58

ANDRESTRAD-M (Central) Same formula as Andretraq with testosterone increased to 25 mg Vial 10 cc

ANDREX INJ. (Ries) Testosterone 20 mg, estrane (0.5 mg) 5,000 IU, Vit C 100 mg, B 1 25 mg, B 2 5 mg, B 6 5 mg, niacinamide 25 mg, pantothenol 10 mg/cc Vial (multiple dose) 20 cc

Use Androgenic therapy

ANDROBINE. (G P Harvey) Methyl testosterone 10 mg, yohimbine HCl 4 mg / Tab Bot 100s

Use Androgenic agent

ANDRODIOL (Carrick) Diolastene Methyl androstenediol Hypoglossals (10 mg) Bot 30s, 100s (25 mg) Bot 30s, 100s

Use: SL buccally, orally, anabolic nitrogen therapy

ANDROGENS Substances which possess masculinizing activities

See Andretraq, Vial (Central)
Andrusal, Vial (C D Smith)
Cavaterone, Amp (Cavendish)
Malesirone, Vial, Tab (Kirk)
Masenate Amp, Vial (Schieffelin)
Metandren Ling, Tab (Ciba)
Methyl Testosterone (Various Mfr.)
Neo Hambreol Tab (Organon)
Neo Hambreol (M) Tab (Organon)
Orchical Vial (Cale)
Oretan, Tab (Schering)
Oretan P, Tab (Schering)
Oretan M, Tab (Schering)
Perandren, Amp, Vials (Ciba)
Testandrene, Tab (Carrick).
Testosterone (Various Mfr.)
Testrane, Tab, Vial (Miller)
Testryl Vial (Squibb)
Virasterone, Amp (Endo).

ANDROGEN-ESTROGEN THERAPY.

See: Dieneutral with Methyltestosterone
Estradiol Esters with Methyltestosterone
Estradiol Esters with Testosterone
Estrogenic Substance, Conjugated with Methyltestosterone
Estrogenic Substance Mixed with Methyltestosterone
Estrogenic Substance Mixed with Testosterone
Estrane with Testosterone

ANDRO-HART. (Hart Drug Corp) Testosterone propionate in oil Vial (25 mg/cc) 10 cc.

Use: I.M., androgen therapy.

ANDROID (Brown Pharmaceutical) Methyl testosterone 2.5 mg, glutamic acid 50 mg, thyroid extract 10 mg, thiamine hydrochloride 10 mg / Tab Bot 100s, 500s

* Generic name recognized by N.F., U.S.P. or A.M.A.

Use: Hypogonadism, male impotence in male frigidity

ANDROID-H P. (Brown) Thyroid extract X mg, methyl testosterone 5 mg, glutamic acid 50 mg, thiamine HCl 10 mg / Tab Bot 100s, 500s

Use Impotency and hypogonadism in male

ANDROLAN AQUEOUS. (Lannett) Testosterone 25 mg/cc, susp Vials 10 cc. Also as 50 mg/cc

Use: I.M., androgen therapy

ANDROLAN. (Lannett) Testosterone propionate 25 mg/cc in oil Vials 10 cc

Use: I.M. androgen therapy

ANDROLIN, NND 1962 (Lincoln) (Testosterone Aq Susp) Testosterone 25 mg, 50 mg/cc, 100 mg/cc Vial 10 cc

Use: I.M., androgen therapy

ANDROLIN IN OIL, NND 1958 (Lincoln) Testosterone propionate 25 mg, 50 mg/cc; 100 mg/cc Vial 10 cc

Use I.M., androgen therapy

ANDRO-MEDICONE (Medicone) Thyroid ext 1/12 gr, strychnine sulfate 1/60 gr, yohimbine HCl 1/12 gr / Tab Bot 60s, 600s 1000s

Use: Tonic & stimulant

ANDRONAQ, NND 1962 (Central) Testosterone Vials (25 mg/cc) 10 cc (50 mg/cc) 10 cc Vials (100 mg/cc) 10 cc

Use: I.M., androgen therapy

ANDRONATE, NND 1962 (Central) Testosterone Propionate (25 mg 1 cc) Vials 10 cc

Use I.M. Androgen Therapy

ANDROPHEROL. (Central) Testosterone propionate 25 mg & dl-alpha tacopheral acetate 100 mg/cc Vials 10 cc

Use I.M., androgen and vitamin E therapy

ANDROSOL SUSP., NNR (C. D. Smith) Testosterone Vials (25 mg) 10 cc, 30 cc (100 mg/cc) 10 cc

Use: I.M. (25 to 75 mg weekly), androgen therapy

* **ANDROSTANAZOLE.** 17b hydroxy 17a methylandrostan [32c] pyrazole (Winthrop Labs)

ANDROSTANOLONE. See Stanolone
ANDROSTANE-17-(beta)-ol-3-one. See Stanolone

ANDROTEST P. See: Testosterone Propionate

ANDROTOCOPHEROL (New Mexico Pharm Co) Alpha Tacopheral Acetate 100 mg, Testosterone Propionate 25 mg, Chlorbutanol 0.5% / cc Vials 10cc

ANDROVARIN (Warren Teed) Testosterone 50 mg, estrone 5 mg, procaine HCl 0.5%

ocacia 1%, thimerosal 0.01% in an isotonic aqueous solution/cc. Vial 10 cc.

Use: I.M. (1 cc. weekly), androgen and estrogen therapy.

ANDRO-WOLFIA (Ortega) Methyltestosterone 5 mg., rauwolfia serpentina 50 mg./Tab. Bot. 100s, 500s.

Use: Treatment of gonadal insufficiency.

ANDRUSOL SUSPENSION, N.N.D. 1962. (Smith, Miller & Patch) Testosterone suspended in isotonic sod. chloride solution. 25 mg./cc. Vial 10 cc., 30 cc.s 100 mg./cc. Vial 10 cc.

ANDRUSOL-P, N.N.D. 1962. (Smith, Miller & Patch) Testosterone propionate in peanut oil. Amp. (25 mg./cc.) 15 cc., (50 mg./cc.) 10 cc., (100 mg./cc.) 10 cc.

Use: I.M. (25 to 75 mg. weekly), androgen therapy.

AN-DU-HIST TT, (Phillips Roxane) Chlorpheniramine maleate 8 mg., phenyltoloxamine dihydrogen citrate 25 mg., d-amphetamine sulfate 5 mg./Tab. Bal. 500s.

Use: Treatment of allergies.

ANECTINE CHLORIDE, N.N.D. 1962. (Burrroughs Wellcome) Succinylcholine chloride, diacetylcholine chloride. Vials, multiple dose (20 mg./cc.) 10 cc. Sterile Sol. (50 mg. & 100 mg./cc.) Amp. 10 cc.

Flo-Pak, Plastic inj. unit (500 or 1000 mg.) Box 12s.

Use: I.V., short-acting skeletal muscle relaxant

ANELIX, (Kirkman) N-Acetyl-p-aminophenol. 120 mg./5 cc. Elx. Bot. 1 pl.

Use: Analgesic.

ANELMID, See: Delvez, (Lilly)

ANERGEX, (Mulford Colloid Labs.) A specially prepared ext. of Toxicodendron quercitulum. Vial 8 ml.

Use: I.M., 1 ml. daily for 6 to 8 days, treatment of allergic diseases

ANERTAN. See: Testosterone Propionate, Amps. & Vials, 17-Methyltestosterone, usually tablets.

ANESACAINE, (New Mexico Pharm Co) Nupercaine 0.5 mg., Quinine HCl 6.5 mg., Benzyl Alcohol 2%./cc. Vials. 100 cc.

ANESTHESIN, N.N.R. 1948, A.O.R. Ethyl-p-aminobenzoate N.F. See Benzocaine.

Use: Local anesthetic.

ANESTHETICS, BASAL

See: Avertin. (Winthrop Labs.)

Cyclopropene. (Various Mfr.).

Ethyl Chloride. (Various Mfr.).

Evipal Sodium. (Winthrop Labs.)

Neraval, Vial (Schering)

Pentothal Sodium. (Abbott).

Suritol. (Parke, Davis).

Viadril, Vial (Pfizer)

Vinethene. (Merck Sharp & Dahme)

ANESTHETICS, INJECTABLE.

See: Amylocaine HCl. (Wallace).

Amylsine. (Novaco).

Apothesine HCl. (Parke, Davis).

Blockain (Breon)

Butethamine. (Novaco).

Chlorprocaine HCl.

Cyclaine. (Merck Sharp & Dahme)

Dibacaine HCl. (Ciba).

Eucaine HCl. (No Mfr. currently lists).

Hexylcaine HCl. (Merck Sharp & Dahme)

Intracaine HCl. (Squibb).

Isocaine HCl. (Various Mfr.).

Larocaine HCl. (Rache)

Lidocaine HCl. (Astra).

Lucaine HCl. (Maltbie)

Metycaine HCl. (Lilly).

Monocaine HCl. (Novaco).

Nesacaine HCl. (Maltbie)

Nupercaine HCl. (Ciba).

Piridocaine HCl.

Pentacaine. (Winthrop Labs.)

Primocaine HCl. (Novaco)

Procaine Salts. (Various Mfr.).

Propoxycaïne HCl.

Quatane HCl. (Smith, Kline & French).

Tetracaine HCl. (Winthrop Labs.)

Unocaine HCl. (Novaco).

Xylacaine HCl. (Astra)

ANESTHETICS, OPHTHALMIC.

See: Amylsine HCl. (Novaco).

Butacaine Sulfate. (Abbott).

Butyn Sulfate. (Abbott).

Cocaine HCl.

Dorsacaine HCl. (Smith-Dorsey)

Holocaine HCl. (Winthrop Lab.)

Metycaine HCl. (Lilly).

Neopaine HCl. (Novaco).

Ophaine HCl. (Squibb).

Pentacaine. (Winthrop Labs.)

Procaine Salts. (Various Mfr.).

Quatracaine HCl. (Abbott).

ANESTHETICS, SPINAL.

See: Cyclaine. (Merck Sharp & Dahme)

Diothane HCl. (Merrell)

Diperodon HCl. (Merrell).

Hexylcaine HCl. (No Mfr. currently lists).

Halocaine HCl. (Upjohn).

Lucaine HCl. (Maltbie).

Metycaine HCl. (Lilly).

Phenacaine HCl. (Upjohn).

Pentacaine. (Winthrop Labs.)

Spinocaine. (Winthrop Labs.)

Tetracaine HCl. (Winthrop Labs.)

Unocaine HCl. (Novaco).

ANESTHETICS, TOPICAL.

See: Amethane HCl. (Abbott)

Biphenamine HCl. (Sayhyun).

- Bulacaine Sulfate (Various Mfr)
 Bulamben Picrate (Abbott)
 Butyl Aminobenzoate (Abbott)
 Cromolone (Geigy)
 Cycloine (Merck Sharp & Dohme)
 Daphine (Sayhyun)
 Dibucaine (Ciba)
 Diathone HCl (Merrell)
 Dyclone Creme Sol (Pitman Moore)
 Ethyl Aminobenzoate (Various Mfr)
 Eurox (Geigy)
 Hexylcaine HCl (No Mfr currently lists)
 Intracaine HCl (Squibb)
 Lorocaine HCl (Rache Labs)
 Lidocaine HCl
 Metylcaine HCl (Lilly)
 Manocaine HCl (Navacol)
 Orihalarm (Winthrop Labs)
 Penthesine Methanesulfonate (Sandoz)
 Pontocaine HCl (Winthrop Labs)
 Pramoxine HCl (Abbott)
 Proposein (No Mfr currently lists)
 Quatracaine HCl (Abbott)
 Quatone (Smith Kline & French)
 Stavaine HCl
 Surfaca ne HCl (Lilly)
 Tetracaine HCl (Winthrop Labs)
 Tranthane HCl (Abbott)
 Xylacaine HCl (Astra)
- ANESTHONE CREAM, IMPROVED** (Parke Davis)
 Adrenalin 1:5000 ephedrine HCl 2%, ethyl aminobenzoate 10% Tube ½ oz
 Use: Relief of inflammatory conditions of accessible mucous membrane
- ANESTRO**, (Mallard) Testosterone 25 mg estrone 2 mg/cc Vial 10 cc, 5s, 10s 25s
 Use: 1 M, androgen and estrogen therapy
- ANETHAINE** See Pontocaine Hydrochloride (Winthrop Stearns)
- ANETIC**, (Chicago Pharm) Aesculus hamamelis collinsiana stramonium, ichthammol oil of tar & phenol No 2 contains benzocaine Oint Tube 1 oz Jar 1 lb
 Use: Hemorrhoids
- ANEURINE HYDROCHLORIDE**, NND 1952
 ADR See: Thiamine HCl USP Prep (Various Mfr)
- ANEXSIA W/CADERNE** (Massengill) A.P.C w/Cadene Cap (¼ gr, ½ gr) Bot 100s 500s Tob (¼ gr, ½ gr 1 gr) Bot 100s 500s
 Use: (1 2 q 3 h) Analgesic antipyretic
- ANEXSIA-D** (Massengill) D hydrocodone bitartrate 7 mg, acetophenetidin 0.15 Gm aspirin 0.23 Gm, caffeine 30 mg/Tab Bot 100s, 500s 1000s
 Use: Analgesic
 * Generic name recognized by N.F., U.S.P. or A.M.A.
- ANGIN**, (Eaton) Estrone 2 mg, testosterone 25 mg/cc Aq susp Vial 10 cc
 Use: S.C., estrogen and androgen therapy
- ANGISAL** (Smith Dorsey) Pentobarbital ¼ gr, caffeine ¼ gr, salicylamide 3½ gr, phenacetin 2½ gr/Tab. Bot 100s, 500s, 1000s
 Use: (1 q 3 h) Sedative, analgesic.
- ANGICAP**, (Pro-Acet) Pentaerythritol tetranitrate 30 mg/Trmed Cap Bot 100s, 250s
 Use: (1 bid) Angina pectoris
- ANGIO-CONRAY**, (Mallinckrodt) (Sodium Iothalamate 20%) Iothalamic acid
 Use: Radiographic contrast medium
- * ANGIOTENSIN AMIDE**, (Crba) L-Aspartoyl 3 L valyl angiotensin octapeptide See Hypertension CIBA
- ANGITET**, (G F Harvey) Pentaerythritol tetranitrate 10 mg/Tab Bot 100s, 1000s
 Use: Angina pectoris
- ANHYDROHYDROXYPROGESTERONE**, USP XIII Ethisterone, USP
 Use: Corpus luteum hormone deficiency threatened abortion
- ANHYDRON**, (Eli Lilly) Cyclothiazide
- ANICOR** (Ensley Scott) Syrup Bot 4 oz
- * ANILAMATE**, (Upjohn) Methylcarbomale of salicylanilide
 Use: Anti-inflammatory compound
- * ANILERIDINE HCl and Phos**, NND 1962 Ethyl 1 (4 aminophenethyl)-4 phenylisopropionate
 See: Lestrine HCl (Merck Sharp & Dohme) Tertine Phos (Merck Sharp & Dohme)
- ANION EXCHANGE RESINS**,
 See: Polyamine Methylene Resin, NND 1959
- * ANISINODIONE**, 2 p anisyl indandione 1, 3 (C₁₈H₁₂O₃) See Miradon (Schering)
- ANISOPYRADAMINE**, See: Pyrramine Maleate
- 2-p-ANISYLINDANEDIONE-1, 3** See: Miradon Tabs (Schering)
- ANNIDALIN**, See: Thyamol Iodide (Various Mfr)
- ANNOPHAL**, (Therachemicals) Phenobarbital ¼ gr, atropine 0.0214 mg, with hyoscyamine sulfate & hyoscyamine HBr/Cap or Tab Cap, Bot 100s, 500s; Tab, Bot 500s
 Use: (1 2 q 3 h) Antispasmodic, sedative
- ANOBALM, STRONGER**, (Sullivan & Case) Chloral hydrate 50%, methyl salicylate 20%, menthol 10%, eugenol 15%, lanolin 250%, petrolatum wax base Oint 1 lb
- ANOZEN**, (Bentex Pharm) Acetophenetidin 3 gr aspirin 3 gr, hyoscyamus ext ¼ gr phenobarbital ¼ gr/cap Bot 100s, 1000s
 Use: Analgesic w/sedative

- AN-O-DYN.** (Paul Money) Phenylpropanolamine HCl 12.5 mg, pyrilamine maleate 12.5 mg, prophepyridamine maleate 12.5 mg, salicylamide 5 gr., Vit. C 20 mg., aluminum magnesium hydroxy carbonate 2 gr./Cap. Bot. 100s, 500s.
Use: Treatment of the common cold.
- ANODYNE Jr.** (Jenkins Labs.) Acetophenelidin ½ gr., caffeine 1/10 gr., sod. bicarbonate 1/5 gr., amm. carbonate 1/32 gr., licorice ext./tab. Bot. 100s, 1000s.
Use: Analgesic.
- ANODYNINE.** See: Antipyrine.
- ANODYNON.** See: Ethyl Chloride.
- ANODYNOS-DHC.** (Breon) Dihydrocodonine bitartrate 5 mg, aspirin 227 mg, N-Acetyl-p-Aminophenol 150 mg, caffeine 32.4 mg./Tab. (Also available as A.P.C. plain.) Bot. 100s, 500s.
Use: Narcotic, analgesic.
- ANOITOL.** (Suliff & Case) Phenol 1.00%, aluminum acetate 3.75%, camphor 1.00%, eucalyptol, thymol, menthol, methyl salicylate, zinc oxide, emollient base q.s. Oint. 1 lb., 5 lb.
Use: Protective & antiseptic
- ANOROID.** (Bryant) Tyrothricin 0.01%, hexachlorophen 0.5%, benzocaine, bismuth subgallate, bismuth subiodide, balsam peru, zinc oxide. Oint. Tube 1½ oz.
Use: Rectally, treatment of hemorrhoids
- ANOTRIX.** (Elder) Elix. 400 Scopolamine methylbromide 1 mg, butobarbital sod. 10 mg., and alcohol 10%/5 cc. Bot. 1 pt., 1 gal.
- ANOVLAR.** Norethindrone plus ethinylestradiol.
Use: Oral contraceptive
- ANOVO.** (Chicago Pharmacol) Strychnine phosphate 1/150 gr., suprarenal ½ gr., iron phosphate 1/100 gr., Pot. phosphate 1/100 gr./Pills, E.C. Bot. 100s, 1000s.
Use: Hypotension, nervousness & melancholia.
- ANPITRONE.** (Gold Leaf) Chariotic ganadotropin. Vial (5000 u. & 10,000 u.) 10 cc.
Use: I.M., cryptorchidism.
- ANPLEX.** (Chicago) liver, crude, 2 vs Vit. B-12 10 mcg., folic acid 5 mg./cc. Amp. 1 cc. Box 6s, 12s, 25s, 100s. Amp. 10 cc. Box 1s.
Use: I.M., antianemia therapy.
- ANPLEXALT VIAL 10A.** (Chicago Pharm.) Liver containing Vit. B-12 equivalent to 20 mcg. with Vit. B-12 25 mcg. & folic acid 5 mg. added/cc. Vial 10 cc., 1s, 6s, 30s.
Use: I.M., anemias & sprue.
- ANPLEXFORT.** (Chicago) Vit. B-12 30 mcg., folic acid 10 mg, liver inj. 10 mcg./cc. Vial 10 cc.
Use: I.M., antianemias.
- ANPLEXIX.** (Chicago) Vit. B-12 60 mcg., folic acid 10 mg., liver inj. 10 mcg./cc. Vial 10 cc.
Use: I.M. antianemias.
- ANPLEXUL.** (Chicago Pharm.) Same as Anplex.
- ANPLEXUL VIAL 110A.** (Chicago Pharm.) Liver, crude, with Vit. B-12 activity equivalent to 2 mcg. of cyanocobalamin, Vit. B-12 cryst. 10 mcg, folic acid 5 mg./cc. Vial 10 cc., 1s, 6s, 100s; Amp. 1 cc., 1s.
Use: I.M., anemias of all types.
- ANQUIL.** (Dumas Wilson) Reserpine. Tab. (0.10, 0.25, 0.50, 1, 4 & 5 mg.) Bot. 100s, 1000s.
Use: Tranquillizing agent.
- A. N. S. TABLETS.** (Kenwood Labs.) Chloralpyrine (dichloralantipyrine) 5 gr, omobarbital 1 gr./Tab. Bot. 30s.
Use: Insomnia
- ANSAC.** (Massengill) Digitalin 1/50 gr., oxydendron arboreum ext. ½ gr., sambucus canadensis ext. ½ gr., squillin ¼ gr., sparteine sulfate ¼ gr./Tab. Bot. 100s, 1000s
Use: Cardiac stimulant
- ANSADO OINTMENT, N.N.D. 1962.** (Rorer) Salicylanilide 4.5%, Jar 4 oz., 1 lb.
Use: Topically, fungicidal ointment.
- ANSEMCO.** (Massengill) No. 1 A.P.C. No. 2 A.P.C. w/tr. gelsemium, Cap. No. 1 & 2, Bot. 500s, 1000s. No. 2 1000s only.
Use: Orally (1-2 q. 4 h) Analgesic & antipyretic.
- ANSIV** (Abbot) 1-[2-methoxyphenyl]-4-[3-methoxypropyl]-piperazine phosphate.
Use: Lowers blood pressure through peripheral vasodilation.
- ANSODYNE.** (Elder) Phenobarbital ¼ gr., belladonna ext. ⅓ gr., aspirin 5 gr., caffeine ¼ gr./Tab. Bot. 100s, 1000s.
Use: Analgesic, sedative.
- ANSOLYSEN, N.N.D. 1962.** (Wyeth) Pentamethyl-pyrrolidinium tartrate. Pentamethyl-1,5-bis(1'-methyl-pyrrolidinium bitartrate). Tab. (20, 40, 100 & 200 mg) Bot. 100s. Vial (10 mg/cc) 10 cc.
Uses Orally, S.C., I.M., ganglionic blocking agent, hypertension.
- ANSPAS.** (Dumas-Wilson) Hyoscyne HBr 0.0065 mg., atropine sulf. 0.0194 mg., hyoscyamine sulf. 0.1037 mg., phenobarb. 16 mg./Tab or 5 cc. Bot 100s and Elixir 1 pt., 1 gal.
Use: Antispasmodic.
- ANTABUSE, N.N.D. 1961.** (Ayerst) Disulfiram,

- tetraethylthiuram disulfide Tab (0.5 Gm)
Bot 50s 1000s
Use: Orally, in alcoholism
- ANTACID.** (Sutliff & Case) Bismuth subgallate 0.2 Gm, Magnesium Carbonate 0.2 Gm Sodium Bicarbonate 1.5 Gm Calcium Carbonate 1.9 Gm / 3.9 Gm Powder Jars 2½ oz, 4½ oz, 1b 5 lb
- ANTACID.** (Smith, Miller & Patch) Calcium carbonate 3½ gr, Magnesium carbonate 2 gr, Bismuth subnitrate ½ gr/Tab Bot 100s 1000s
- ANTACID TABLETS** (Parke Davis) Aluminum hydroxide gel, magnesium carb, calcium carb Taka diastase Box 36s
Use: Antacid
- ANTACID NO 3** (National Drug) Phenobarbital ½ gr, atropine sulfate 1/300 gr magnesium hydroxide 5 gr cal carbonate 5 gr/Tab Bot 100s
Use: Antacid, antispasmodic and mild sedative
- ANTACIDEN** (Tilden) Sodium bicarbonate calcium carbonate oil of peppermint, magnesium carbonate Bot 4 oz
Use Gastric antacid
- ANTACIDS, GASTRIC.** A substance that counteracts or neutralizes acidity
See: Alkalinizing Agents
Aluminum Carbonate Basic
Aluminum Hydroxide
Aluminum Phosphate
Anion Exchange Resins
Calcium Carbonate
Dihydroxy Aluminum Aminoacetate
Dihydroxy Aluminum Sod Carb
Magnesium Carbonate
Magnesium Oxide
Magnesium Trisilicate
Mucin
Mucatin
Polyamine Methylene Resins NND
Tritalac
- ANTAGEL** (Aftison) Aluminum magnesium hydroxy carbonate 10 gr/Tab Bot 50s
Use: Antacid
- ANTAGONISTS OF CURARIFORM DRUGS**
See: Tension Chloride (Rache Labs)
Neostigmine Methylsulfate (Rache Labs)
- ANTAMINE.** (Grove) Pyrilamine maleate Tab Bot 12s, 36s
Use: Antihistaminic.
- ANTAR.** (Ulmer) Thenylpyramine 2%, active coal tar principles 1% Oint Jar 1 oz, 1 lb
Use: Topically, dermatoses
- ANTASED.** (Lemmon) Aluminum hydroxide 4 gr, homatropine methylbromide 12 mg, gastric mucin 1 gr., magnesium trisilicate
- ANTAZOLINE HYDROCHLORIDE, U.S.P. XV** NND 1961, Tablets, U.S.P. XV (Antolan) 2-(N-Benzylaminomethyl) 2-imidazoline HCl
See: Antistine Hydrochloride, Sol & Tab (Ciba)
W/Pyribenzamine See: Dibistine, Tab. (Ciba)
- *ANTAZOLINE PHOSPHATE, N.F., NND 1962** Ophth Saln NF Histolab See Antistine Phosphate Oph Sal (Ciba)
W/Boric acid sodium carbonate anhydrous sodium chloride, phenylmercuric acetate See Azalone Ophthalmic (Smith, Miller & Patch)
W/Naphazoline, boric acid, phenylmercuric acetate, sod chloride, sod carbonate anhydrous See Vasocin A Ophthalmic, Sol (Smith, Miller & Patch)
- ANTICIBUS.** (Rand) Vit B 1 5 mg, Vit B-2 3 mg, niacinamide 10 mg, calcium pantothenate 0.5 mg, pyridoxine HCl 0.25 mg strychnine sulfate 1.47 mg/fl oz Bot 1 pt, 1 gal
Use: Vitamin supplement
- ANTEPAR, NND 1962** (Burroughs Wellcome) Piperazine citrate Syr (equiv. to 100 mg of piperazine hexahydrate/cc) Bot 4 fl oz, 1 pt, 1 gal Tab (500 mg) Bot 100s 1000s Wafers (500 mg) Box 28s
Use: Orally, anthelmintic, to eradicate pinworms and roundworms
- ANTERIOR PITUITARY, NND 1959** See Pituitary, Anterior
- ANTEX** (Bentex) Estrone 3,333 IU/oz salicylic acid methyl salicylate, sodium lauryl sulfate in a greaseless base Tube 1½ oz
Use: Acne vulgaris
- ANTHALAZINE.** (Bowman) Piperazine hexahydrate 100 mg/cc Syr Bot 4 oz, 1 pt, 1 gal Oz Bot 1000s
Use: Treatment of pinworms and roundworms
- ANTHALLAN.** (Medica Chemica) 3 Di(N-butyl)aminomethyl tri hydroxy benzoturanone 85 mg /Cap Bot 100s, 500s, 1000s
Use: Dermatomycosis
- ANTHAPHYLLINE.** (Medica Chemical Corp) Mixture of antihistal and amnaphylline
Use: Asthma
- ANTHELCOLE.** (Cole) Piperazine citrate 275 mg/Tab Bot 100s 500s, 1000s Ltg Bot 1 pt
Use: Orally, eradication of pinworm and round worm infestation
- ^a Generic name recognized by N.F., U.S.P. or A.M.A.

Hygromycin (Lilly)
 Kenamycin
 Kontrex (Bristol)
 Magnomycin (Pfizer)
 Mycostalin (Squibb)
 Neomycin (Various Mfr.)
 Novobiocin (Various Mfr.)
 Nystalin (Squibb)
 Oleandomycin
 Oleandomycin, Acetyl
 Oxytetracycline
 Penicillin (Various Mfr.)
 Polymyxin B (Various Mfr.)
 N (Pyridinamethyl) Tetracycline
 Ristocelin (Abbott)
 Seromycin (Lilly)
 Spiromycin (Paulenc, Montreal)
 Sponin (Abbott)
 Sireptomycin (Various Mfr.)
 Synleirin (Bristol Lab)
 Terramycin (Pfizer)
 Tetracycline (Various Mfr.)
 Tylasin (Lilly)
 Tyrothricin (Various Mfr.)
 Vancemycin
 Viomycin (Various Mfr.)

ANTICEPHALGICS. Agents used to prevent or cure a headache usually of the recurring type over the region of the external carotid artery.

See Cafergal Tab (Sandaz)
 D hydroergotamine [DHE] (Sandaz)
 Gynergen, Amp, Tab (Sandaz)
 Wigaine Tab (Organon)

ANTI-CHLOR (Klug) Chlorpheniridine maleate USP 4 mg / Tab Bot 100s
 Use May fever urticaria, allergic eczema drug sensitivity serum sickness

ANTICHOLINERGIC AGENTS.

See Ambulanium [Wyeth]
 Aminopentamide
 Anirenyl Syr, Tab (Ciba)
 Banthine, Amp, Tab (Searle)
 Benztropine Methanesulfonate
 Cont I, Tab (Lakeside)
 Cenline, Soft, Tab (Bristol)
 Cogentin Methanesulfonate Tab (Merck Sharp & Dahme)
 Darbid (Smith, Kline & French)
 Darisone, Tab (Merck Sharp & Dohme)
 Diphenmethanil Methylsulfate
 Elaine, Pulv (Lilly)
 Hexacyclium Methylsulfate
 Isopropamide Iodide
 Kemadin Tab (Burroughs Wellcome)
 Mepiperphenadol Bromide
 Methantheline Bromide
 Methscopolamine Bromide
 Manodial, Cap (Winthrop Labs)
 Murel, Tab (Ayerst)

Oxyphenonium Bromide
 Pamine, Tab (Upjohn)
 Pothilon, Tab (Lederle)
 Penthenate Bromide
 Pipal, Tab (Lakeside)
 Prantal, Cr, Tab, Vial (Schering)
 Probanthine, Tab (Searle)
 Procydiline HCl
 Propantheline Bromide
 Stopalate Nitrate, Tab (Sirasenburgh)
 Tral, Tab (Abbott)
 Tricalad, Tab (Burroughs Wellcome)
 Traclyamal Chloride
 Tridihexethyl
 Valethamate Bromide

ANTICOAGULANT ACID CITRATE DEXTROSE SOLUTION, USP. See A.C.D. Solution (Various Mfr.)

ANTICOAGULANTS.

See: Acenocoumarin
 Bishydroxycoumarin (Various Mfr.)
 Coumadin, Amp (Endo)
 Cumopryon (Abbott)
 Cyclocoumarol
 Donitone, Tab (Schiefelin)
 Dicumaryl (Various Mfr.)
 Dipaxin, Tab (Upjohn)
 Diphenadione
 Eridione, Tab (Eric, Kirk & Gary)
 Ethyl Biscoumacetate, Tab (Geigy)
 Hedulin, Tab (Walker)
 Hemostatics
 Heparin (Various Mfr.)
 Indon Tab (Parke, Davis)
 Koagomin, Vial (Chatham)
 Liqueomin Sad, Vial (Organon)
 Phenindione, Tab (Schiefelin)
 Prothromodin, Tab (G F Harvey)
 Sintrom, Tab (Geigy)
 Tramezan, Tab (Geigy)
 Warfarin Sad

ANTICOIDS (Gold Leaf) Magnesium carb. calcium carb. bismuth subnitrate Tab Bot 100s

Use (prn) Anacid

ANTICONVULSANTS. Agents that inhibit muscular spasms originating in the central nervous system

See Rencchlorpropamide
 Celonin, Capsules (Parke Davis)
 Dilantin Sodium, Cap (Parke, Davis)
 Diphenylhydantoin Sodium (Various Mfr.)
 Ethotoin
 Eklipen Tab (Ciba)
 Gemand, Tabs (Abbott)
 Glutamic Acid (Various Mfr.)
 Hydantal, Tab (Sandaz)
 Mebaral, Tab (Winthrop Labs)
 Mebaroin, Tab (Winthrop Labs)
 Mephobarbital

Chlor Trimefan (Schering)
 Clistin Maleate (McNeil)
 Cyclizine HCl
 Decapryn (Meirell)
 Dafen (Schenley)
 Dimenhydrinate
 Dimetane (Robins)
 Di Paralene (Abbott)
 Diphenhydramine HCl (Parke, Davis)
 Diphenylpyraline HCl
 Doxylamine Succinate
 Dramamine (Searle)
 Histadyl HCl (Lilly)
 Isath pendyl
 Marezine HCl (Burroughs Wellcome)
 Methaphenilene HCl (Various Mfrs)
 Methapyrilene HCl (Various Mfr)
 Neohelramine (Nepero)
 Perazil (Burroughs Wellcome)
 Parabromdylamine Maleate
 Paracathinaxamine Maleate (Burroughs Wellcome)
 Phenindamine Tartrate
 Phenylalaxamine Dihydrogen Citrate
 Phenergan (Wyeth)
 Pheniramine Maleate
 Promethazine HCl
 Prophepyndamine (Schering)
 Pyralizine HCl
 Pyribenzamine (Ciba)
 Pyrilamine Maleate (Various Mfr)
 Pyrobutamine
 Pyron I (Lilly)
 Pyrolate (Upjohn)
 Tecaryl HCl Tab
 Tegathen (Lederle)
 Thelad I HCl (Winthrop Labs)
 Thenyldamine HCl (Winthrop Labs)
 Thenyline HCl (Abbott)
 Thepharin (Rache Lab)
 Theruh sin (Ayerst)
 Thanylam ne HCl
 Trimefan (Schering)
 Tr pelennam ne
 Tarant I (Winthrop Labs)
 Vibor ne HCl (Pfizer)
 Zalamine (White)

ANTI HUMAN PRECIPITIN SERUM. (Parke Davis, Nyland) Bot wide mouth 1 cc.
 Use: Diagnostic aid for the identifying human blood in blood stains.

ANTILEUKEMIA. See Leukemia.

ANTIMALARIAL AGENTS

See Amodiaquin HCl
 Arelen D phosphate Tab (Winthrop Labs)
 Atebr ne D hydrochloride, Tab. (Winthrop Labs)
 Cemaquine Hydrochloride, Tab. (Parke Davis)

* Generic name recognized by N.F., U.S.P. or A.M.A.

Chloroquinide Hydrochloride, Tab (Lilly)
 Chloroquine Phosphate (Winthrop Labs.)
 Caco Quinine, Susp (Lilly)
 Daraprim, Tab (Burroughs Wellcome)
 Equamine (Various Mfrs)
 Guanafat Hydrochloride, Tab (Lilly)
 Hydroxychloroquine Sulfate
 Paludrine HCl (Ayerst)
 Pamaquine Naphthoate, Tab (Winthrop Labs)
 Penlaquine Phosphate, Tab (Abbott)
 Ploaquent (Winthrop Labs)
 Plasmochin Naphthoate, Tab (Winthrop Labs)
 Primaquine Disphosphate, Tab (Winthrop Labs)
 Pyrimethamine
 Quinacrine Hydrochloride, Tab
 Quinine Salts (Various Mfr)
 Toloquine, Pow (Mallinckrodt)

ANTI-M A ANTI-N SERUM. (Lederle) Anti A colored with a yellow dye Anti N colored with a green dye Vial 25 tests

Use Diagnostic aid for the detection of M & N substances in human red blood cells

ANTI-M LECTIN. (Hyland) Extract from seed of *Iberis amara*
 Use Blood typing reagent for blood factor M

ANTI-N LECTIN (Hyland) Extract from seed of *Vicia graminea*
 Use Blood typing reagent for blood factor N

ANTIMONYL POTASSIUM TARTRATE See Antimony Potassium Tartrate, U.S.P.

***ANTIMONY POTASSIUM TARTRATE** U.S.P. Tartar emetic (Various Mfr)
 Use: Schistosomiasis leishmaniasis, expectorant & emetic

W/Dilaudid HCl, Sod citrate, chloroform See Dilaudid, Cough Prep (Tabletack)

W/Methapyrilene HCl, Sod citrate ammonium chloride, ephedrine HCl See Semcal Prep (Mallinckrodt)

W/Pot guaiaaculsulfonate & chloroform See Kalitran Syr (National).

W/Thenylpyramine HCl, ammonium chloride Sod citrate menthol, aromatics See Histo comp Syrup (Approved Pharm.)

ANTIMONY PREPARATIONS. See Antimony Potassium Tartrate (Various Mfr)

Antimony Sodium Thioglycollate (Various Mfr).

Antimony Thioglycollamide (H W & Dering)
 Fuadin, Amp (Winthrop Labs)
 Neostam, Vial (Burroughs Wellcome)
 Stibamine (Burroughs Wellcome)

- Sibaphen (Winthrop Labs)
Tartar Emetic (Various Mfr.).
- ANTIMONY SODIUM THIOGLYCOLLATE**, N.F., N.N.R. 1949; Injection, N.F. (Various Mfr.)
Use: Schistosomiasis, leishmaniasis & filariasis.
- ANTIMONY THIOGLYCOLLAMIDE**, N.N.R. 1951. (Hynson, Westcott & Dunning) Amp (0.4% solution) 10 cc.
Use: I.M., I.V., prolozoacide.
- ANTIMONY TRIIODIDE**. W/iodoform, mental, camphar. See: Camiral, Amp. (C. D. Smith)
- ANTIMUMPS SERUM (Human)**. (Hyland Labs.) Gamma globulin fraction of blood from healthy human donors. Each 2.5 cc. is equivalent in antibody content to no less than 25 cc. of original hyperimmune serum Vial 2.5 cc.
Use: I.M., treatment of mumps.
- ANTI-NAUSEANTS**.
See: Apolamine, Tab. (Winthrop Labs)
Bonamine. (Pfizer)
Compazine, Tab. (Smith, Kline & French)
Chlorelone, Prep. (Parke, Davis)
Dromamine, Tab. (Searle)
Emetrol, Liq. (Klinney).
Genescopolamine
Marezine. (Burroughs Wellcome)
Mornidine, Amp. & Tab. (Searle).
Mollon sleekness.
Pipamoxine
Probutylin (Rorer)
Pyridoxine HCl Preparoliant (Various Mfr.).
Thorazine, Tab. (Smith, Kline & French)
Tigan, Cap. (Roche Labs.)
Trelafon, Tab. (Schering)
- ANTI-NAUSEA SUPPRETTES**. (Webster) Pyridamine maleate 25 mg., pentobarbital sod. $\frac{1}{2}$ gr./Supp. Box 12s
Use: Nausea
- ANTI-NEOPLASTIC**
See: Leukemia
- ANTI 5-NITRO-2-PURALDOXIME**. See: Micalur (Eaton)
- ANTINOGAN**. (Bristol) Polysaccharide from an actinomycete.
Use: Cancer therapy
- ANTI-OBESITY AGENTS**.
See: Thyroid
Amphetamine Preparations (Various Mfr.).
Aracen, Tab. (Central)
Dextro-amphetamine Preparations (Various Mfr.).
Diethylpropion.
Ionomine, Tab. (Shassenburg)
Levanar, Tab. (Nordion)
Leva-Amphetamine

- Methamphetamine Preparations (Various Mfr.)
Preludin, Tab. (Geigy)
Tenuate (Merrell)
Tepanil, Tab. (National Drug)
Vitamins with Antiobesity Agents.
- ANTI-PAC COMPOUND**. (Lawitt Labs)
Phenylephrine HCl 5 mg.; salicylamide 0.23 gm., acetophenetidin 0.15 gm.; caffeine 0.03 gm.; ascorbic acid 50 mg.; hesperidin complex 50 mg.; chlorpraphenpyridimine maleate 2 mg./tab. Bot. 30s, 100s
Use: Cold therapy
- ANTI-PARASYMPATHOMIMETICS**. See: Parasympatholytic agents
- ANTI-PARATYPHOID AGGLUTINATING SERUM**. (Parke, Davis) Paratyphoid bacillus "A" (Salmonella paratyphi) Vial 5 cc.
Use: Diagnostic aid in the test for B paratyphosis "A" & "B."
- ANTI-PELLAGRA VITAMIN**. See: Nicotinic acid.
- ANTIPEOL**. (Myers) Bacterial antigen from Staph. albus, aureus & citreus; Strep. hemolyticus and viridans, and B. pyocyanus. Liq., Bot. 60 cc., 100 cc.; Oint. 4 oz.
Use: Topically, onlvirus therapy.
- ANTI-PERNICIOUS ANEMIA PRINCIPLE**.
See: Vitamin B 12.
- ANTI-PERTUSSIS SERUM (Human)**. N.N.D. 1939. (Hyland Labs) Gamma globulin fraction of blood from healthy human donors. Each 2.5 cc. is equivalent to antibody content to no less than 25 cc. of original hyperimmune serum. Vial 2.5 cc
Use: I.M., treatment of whooping cough.
- ANTI-PERTUSSIS SERUM (Rabbit)**, N.N.R. 1952. (Wyeth) Refined and concentrated Vial (20,000 units) 4 cc.
Use: Immunization or treatment of whooping cough.
- ANTI-PROTOZOAN AGENTS**.
See: Anayodin, Pills (Bischoff)
Antimony Preparations.
Arsenic Preparations.
Bismuth Preparations.
Chinifon (Various Mfr.).
Dihydroxyquinoline.
Diadoquin, Tab. (Searle).
Emetine Hydrochloride (Various Mfr.).
Iodoxyquinoline Sulfonate Sodium (Bischoff).
Iodoxyquinoline
Naphthyl Sodium (Winthrop Labs.)
Ouinaxyl, Tab. (Burroughs Wellcome)
Suramin Sodium (Winthrop Labs)
Viofarm (Ciba).
Yodoxin, (Yodoxin, Inc)
- ANTIPYRETIC AGENTS**.
See: Acetanilid Preparations

- Chlor Trimetan (Schering)
- Chistin Maleate (McNeil)
- Cyclizine HCl
- Decapryn (Merrell)
- Diaten (Schenley)
- Dimenhydrinate
- Dimetane (Robins)
- Di Paralene (Abbott)
- D phenhydramine HCl (Parke, Davis)
- D phenylpyraline HCl
- Doxylamine Succinate
- Dramamine (Searle)
- Mistadyl HCl (Lilly)
- Isath pendyl
- Marezine HCl (Burroughs Wellcome)
- Methaphenitene HCl (Various Mfr)
- Methapyrilene HCl (Various Mfr)
- Neohetramine (Nepera)
- Perazi (Burroughs Wellcome)
- Parabromdylamine Maleate
- Paracarbinoxamine Maleate (Burroughs Wellcome)
- Phenindamine Tartrate
- Phenyltoloxamine Dihydrogen Citrate
- Phenergan (Wyeth)
- Pheniramine Maleate
- Promethazine HCl
- Propenpyridamine (Schering)
- Pyralazine HCl
- Pyribenzamine (Ciba)
- Pyrilamine Maleate (Various Mfr)
- Pyributamine
- Pyranil (Lilly)
- Pyralozate (Upjohn)
- Tacaryl HCl, Tab
- Togathen (Lederle)
- Thenfed I HCl (Winthrop Labs)
- Thenylid amine HCl (Winthrop Labs)
- Thenylene HCl (Abbott)
- Ikepharin (Roche Lab)
- Therush st n (Ayerst)
- Thansylam ne HCl
- Trimetan (Schering)
- Tripeleennam ne
- Torant I (Winthrop Labs)
- V bar ne HCl (Pfizer)
- Zelamine (White)

ANTI HUMAN PRECIPITIN SERUM (Parke, Davis Hyland) Bot wide mouth 1 cc. Use: Diagnostic aid for the identifying human blood in blood stains

ANTILEUKEMIA. See: Leukemia

ANTIMALARIAL AGENTS

- See Amodiaquin HCl
- Arolex D phosphate, Tab (Winthrop Labs)
- Atobrine D hydrochloride, Tab (Winthrop Labs)
- Camaquin Hydrochloride, Tab (Parke, Davis)

* Generic name recognized by N.F., U.S.P. or A.M.A.

- Chloroguanide Hydrochloride, Tab (Lilly)
- Chloroquine Phosphate (Winthrop Labs)
- Coco Quinine, Susp (Lilly)
- Daraprim, Tab (Burroughs Wellcome)
- Equinine (Various Mfr)
- Guanafol Hydrochloride, Tab (Lilly)
- Hydroxychloroquine Sulfate
- Paludrine HCl (Ayerst)
- Pamaquine Naphthoate, Tab (Winthrop Labs)
- Pentaquine Phosphate, Tab (Abbott)
- Plaquenil (Winthrop Labs)
- Plasmochin Naphthoate, Tab (Winthrop Labs)
- Primaquine Disphosphate, Tab (Winthrop Labs)
- Pyrimethamine
- Quinacrine Hydrochloride, Tab
- Quinine Salts (Various Mfr)
- Totoquine, Pow (Mollinckstadt)
- ANTI-M & ANTI-N SERUM**, (Lederle) Anti M colored with a yellow dye, Anti N colored with a green dye Vial 25 tests Use Diagnostic aid for the detection of M & N substances in human red blood cells
- ANTI-M LECTIN** (Hyland) Extract from seed of *Iberis amara*. Use Blood typing reagent for blood factor M
- ANTI-N LECTIN** (Hyland) Extract from seed of *Vicia graminea* Use Blood typing reagent for blood factor N
- ANTIMONY POTASSIUM TARTRATE**, See Antimony Potassium Tartrate USP
- * **ANTIMONY POTASSIUM TARTRATE**, USP Tartar emetic (Various Mfr) Use Schistosomiasis leishmaniasis expectorant & emetic
- W/Dilaudid HCl, Sod citrate, chloralform See Dilocol Cough Prep (Tablerock)
- W/Methapyrilene HCl Sod citrate ammonium chloride ephedrine HCl See: Sental Prep (Massengill)
- W/Pot guanacolsulfonate & chloralform See Kaitron Syr (National)
- W/Thenylpyramine HCl, ammonium chloride Sod citrate, menthol, aromatics See: Hiscamp Syrup (Approved Pharm)
- ANTIMONY PREPARATIONS.** See: Antimony Potassium Tartrate (Various Mfr)
- Antimony Sodium Thioglycollate (Various Mfr)
- Antimony Thioglycollamide (H W & Duning)
- Fuadin, Amp (Winthrop Labs)
- Neostam, Vial (Burroughs Wellcome)
- Stibamine (Burroughs Wellcome)

- Stibophen (Winthrop Labs.)
Tortor Emetic (Various Mfr.)
- ANTIMONY SODIUM THIOGLYCOLLATE**, N.P., N.N.R. 1949; Injection, N.P. (Various Mfr.)
Use: Schistosomiasis, leishmaniasis & filari-
osis.
- ANTIMONY THIOGLYCOLLAMIDE**, N.N.R. 1951. (Hynson, Westcott & Dunning) Amp. (0.4% solution) 10 cc
Use: I.M., I.V., protozoicide.
- ANTIMONY TRIIODIDE**, W/iodoform, menthol, camphor. See: Camitol, Amp. (C. D. Smith)
- ANTIMUMPS SERUM (Human)**. (Hyland Labs) Gamma globulin fraction of blood from healthy human donors. Each 2.5 cc. is equivalent in antibody content to no less than 25 cc. of original hyperimmune serum
Vial 2.5 cc.
Use: I.M., treatment of mumps
- ANTINAUSEANTS**,
Sina Apolamine, Tab. (Winthrop Labs)
Bonamine (Pfizer)
Compazine, Tab. (Smith, Kline & French)
Chloratone, Prep. (Parke, Davis)
Dramamine, Tab. (Searle)
Emetrol, Liq. (Kinney).
Cenescopolamine
Moresine. (Burroughs Wellcome)
Moronidine, Amp. & Tab. (Searle).
Motion sickness.
Pipamazine
Probutylin (Rorer)
Pyridazine HCl Preparations (Various Mfr.)
Thorazine, Tab. (Smith, Kline & French)
Tigon, Cap. (Roche Labs.)
Trelafon, Tab (Schering)
- ANTINAUSEA SUPPLETTS**. (Webster) Pyrilamine maleate 25 mg., pentobarbital sod. 1/2 gr./Supp. Box 12s
Use: Nausea
- ANTI-NEOPLASTIC**
See: Leukemia
- ANTI 5-NITRO-2-PURALDOXIME**. See: Micalofur (Eaton)
- ANTINOGAN**. (Bristol) Polysaccharide from an actinomyces.
Use: Cancer therapy.
- ANTI-OBESITY AGENTS**.
See: Thyroid
Amphetamine Preparations (Various Mfr.)
Arocen, Tab. (Central)
Dextro-amphetamine Preparations (Various Mfr.)
Diethylpropion.
Ionomine, Tab (Strassenburg)
Levonor, Tab (Nardson)
Levo-Amphetamine.
- Methamphetamine Preparations (Various Mfr.)
Prelodin, Tab. (Geigy)
Tenuate (Merrell)
Tepanil, Tab (National Drug)
Vitamins with Antiobesity Agents.
- ANTI-PAC COMPOUND**. (Lawitt Labs)
Phenylephrine HCl 5 mg; salicylamide 0.23 gm; acetaphenetidin 0.15 gm; caffeine 0.03 gm.; ascorbic acid 50 mg; hesperidin complex 50 mg; chlorphenpyridamine maleate 2 mg /tab. Bot 30s, 100s
Use Cold therapy
- ANTIPARASYMPATHOMIMETICS**. See: Parasympatholytic agents.
- ANTI-PARATYPHOID AGGLUTINATING SERUM**. (Parke, Davis) Paratyphoid bacillus "A" (Salmonella paratyphi) Vial 5 cc.
Use: Diagnostic aid in the test for B paratyphus "A" & "B."
- ANTI-PELLAGRA VITAMIN**. See: Nicotinic acid.
- ANTIPEOL** (Myers) Bacterial antigen from Staph. albus, aureus & citreus; Strep. hemolyticus and viridans, and B. pyocyaneus. Liq. Bot. 60 cc., 100 cc.; Oint. 4 oz.
Use: Topically, antiviral therapy.
- ANTI-PERNICIOUS ANEMIA PRINCIPLE**. See: Vitamin B-12.
- ANTIPTUSSIS SIRUM (Human)**, N.N.D 1959. (Hyland Labs.) Gamma globulin fraction of blood from healthy human donors
Each 2.5 cc. is equivalent to antibody content to no less than 25 cc. of original hyper-immune serum. Vial 2.5 cc.
Use: I.M., treatment of whooping cough.
- ANTIPTUSSIS SIRUM (Rabbit)**, N.N.R 1952. (Wyeth) Refined and concentrated Vial (20,000 units) 4 cc.
Use: Immunization or treatment of whooping cough.
- ANTIPROTOZOAN AGENTS**.
See: Anayodin, Pills (Bischoff).
Antimony Preparations.
Arsenic Preparations.
Bismuth Preparations.
Chinifolan (Various Mfr.).
Diadohydroxyquinoline.
Diadoquin, Tab. (Searle).
Emetine Hydrochloride (Various Mfr.).
Iodoxyhydroxyquinoline Sulfanate Sodium (Bischoff).
Iodoxyhydroxyquinoline
Naphthide Sodium (Winthrop Labs)
Quinoxyl, Tab. (Burroughs Wellcome)
Suramin Sodium (Winthrop Labs)
Valofan (Ciba).
Yodoxin, (Yodoxin, Inc.)
- ANTIPYRETIC AGENTS**.
See: Acetanilid Preparations

- N Acetyl p Aminophenol
 Antipyrine Preparations
 A P C Preparations
 Aspirin Preparations
 Salicylamide
- * **ANTIPYBINE**, N P N N R 1940 (Various Mfr) Analgesine anodynine, dimethylazoquinazine, azydimethylquinizine, parodyne phenylone pyrazoline, sedatine
 Use Analgesic antipyretic
- W/Anesthesin glycerin See E B O, liq (First Texas)
- W/Benzocaine & glycerin, ear drops
 See Auregan, liq (Doha)
- Lanaurine, liq (Lanett)
- W/Benzocaine, sed mercuric iodide See Auracaine Sal (Ethical Pharm)
- W/Rorate See Callyrum, Soln (Wyeth)
- W/Caffeine citrate, isodopyrine & labeline sulfate See Felsol Pow (Amer Felsol)
- W/Chloral hydrate See Sominet, Tab (National Drug)
- W/Glycerin See: Roy Aurelone, liq (Roy mer)
- W/Hydrocortisone alcohol d/d desoxyephedrine hydrochloride pyrimidine maleate See Binalgan HC Sol (Doha)
- W/Hydrocortisone polymyzin B sulfate, neomycin sulfate d busaine HCl See Otocort liq (Lemman)
- W/Phenylephrine HCl, benzocaine See Tymogest W 7, liq (Warren Teed).
- W/Salicylic acid See: Solipyrine Pow Tab (Redel).
- W/Sod bromide, Sed salicylate & caffeine See: Kremasa Sal (Tilden)
- W/Sulfathiazole Sed, propyl p amine benzoate See: Sulf Aurelone liq (Roy mer)
- W/Urea benzocaine & glycerine ear drops See: Aurazine, Sol (Smith Dorsey)
- ANTIPYRINE TYPE ANALGESICS**
 See: Amidopyrine (Various Mfr).
 Aminopyrine (Various Mfr).
 Antipyrine
 D pyrene (Chemo Pura)
 Isodopyrine (No Mfr currently lists)
 Loridon (Roche Lab)
 Meluber, N N B 1940
 Methapyrene (Winthrop)
 Nevald n, Amp, Tab (Winthrop Labs)
 Navalg n (Winthrop Labs)
 Phenazone N N R 1940 (Various Mfr)
 Pyramiden, Prop, N N R 1940 (Winthrop Labs)
- ANTIRHEUMATIC SERUM**, (Lederle) A serum that has been refined and concentrated from hyperimmunized horses Vial (1000 u).
 Use: Rab es prophylaxis
- * Generic name recognized by N P, U S P or A M A

ANTIRHEUMATIC PREPARATIONS.

- See p Aminobenzoic Acid and salts (col pot, sad)
- Ammonium Salicylate
 Anti arthritic Hormones
 Colchicine preps
 Genhsate Sodium
 Glucuranolactone
 Royals with Succinate, Tab (Roymer)
 Salicylamide preps
 Sodium Salicylate
- ANTI-RH TYPING SERUM** (Numon), (Lederle) Anti Rh (Anti D) 85%—Tubes 10 tests Vial, with pipette, 5 cc Anti Rh, (Anti C plus D) 17%—Tube, 10 tests; Vial with pipette 5 c
 Use Diagnostic aid in the determination of Rh₊ (D) and Rh₊ (C plus D) factors in red blood cells
- ANTIBICKETSIAL AGENTS**, N N R, 1949
 See p Aminobenzoic Acid (Various Mfr)
 p Aminobenzoate, Sod (Various Mfr)
 Aureomycin (Lederle)
 Chloramycetin (Parke Davis)
 Rocky Mountain Spotted Fever Serum (Bobbitt) (Lederle, Merck Sharp & Dahme)
 Terramycin (Pfizer)
- ANTI-RUST TABLETS** (Winthrop Labs) Sod Carb, Monohyd 3 16 Gm, Sod nitrite 10 50 Gm /lab Bot 50s, 500s
 Use Rust preventive
- ANTI-RUST**, (Merrell) Sod nitrite Tab, Bot 50s
- ANTISCOBBUTIC VITAMIN** See Ascorbic Acid
- ANTISEPTIC, CHLORINE, ACTIVE.**
 See Antiseptic N Chloro Compounds
 Hypochlorite Preps
- ANTISEPTIC, N-CHLORO COMPOUNDS**
 See: Azochloramide (Wallace & Tiernan)
 Chloramine T (Various Mfr)
 Chlorazene (Frost)
 Chlorozadin (Wallace & Tiernan)
 Dichloramine T (Various Mfr)
 Nalazone (Abbott)
- ANTISEPTIC, DYES.**
 See: Acriflavine (Various Mfr)
 Acriflavine Base (Various Mfr)
 Acriflavine, HCl (Various Mfr)
 Acriflavine Neutral (Various Mfr)
 Aminoacridine, HCl
 Bismuth Violet Prep (Tablerock)
 Brilliant Green
 Crystal Violet
 Dimazon (Heilbratt)
 Dymizal (McNeil)
 Rhazozene (Squibb)
 Euclein
 Gentian Violet (Various Mfr)

Mallophene (Mallinckrodt)
 Methylrosaniline Chloride (Various Mfr.)
 Methyl Violet.
 Pyridium. (Warner-Chilcott)
 Sereium (Squibb).

ANTISEPTIC, IODINE, ACTIVE.

See: Bursaline (Burnham Saluble Iodine)
 Globaline, Tab. (Wallace and Tiernon)
 Heliogen, Tab. (Penick & Co)

ANTISEPTIC, MERCURIALS.

See: Acelomeractol
 Merbak (Schieffelin).
 Mercarbamide (Upjohn).
 Mercosols (Various Mfr.).
 Mercresin (Upjohn)
 Mercirachrome (Hynsan Wescott & Dunning)
 Mercurophen. (Merck Sharp & Dohme)
 Meradicein (Hynsan Wescott & Dunning)
 Merihialate (Lilly).
 Metaphen (Abbott).
 Nitramersal (Abbott)
 Phenylmercuric Acetate (Various Mfr.)
 Phenylmercuric Borate (Various Mfr.)
 Phenylmercuric Chloride (Various Mfr.)
 Phenylmercuric Nitrate (Various Mfr.)
 Phenylmercuric Picrate (Various Mfr.)
 Thimerosal

ANTISEPTIC, PHENOLS.

See: Anthralin (Various Mfr.).
 Betanophthal (Mallinckrodt).
 Bithional. (Winthrop Labs.)
 p Chlorophenol. (No Mfr. currently lists)
 Chlorothymol (Merck)
 Cool Tar Products (Various Mfr.).
 Creosote (Various Mfr.).
 Cresols (Various Mfr.).
 Dichlorophene.
 Dihydroxy Anthranol
 Eugenol (Various Mfr.)
 Guaiacol (Various Mfr.)
 Hexachlorophene (Various Mfr.)
 Hexylresorcinol. (Merck Sharp & Dohme)
 Ienigallol (Knoll)
 Methylparaben (Various Mfr.).
 o-Phenylphenol (Various Mfr.)
 Oxyquinoline Salts (Various Mfr.)
 Phenol (Various Mfr.).
 Picric Acid (Various Mfr.).
 Propylparaben (Various Mfr.)
 Pyrogallol (Various Mfr.)
 Pyrogallol Monoacetate (Various Mfr.)
 Pyrogallol Triacetate (Various Mfr.)
 Resorcinol (Various Mfr.)
 Resorcinol Monoacetate (Various Mfr.)
 Thymol (Various Mfr.).
 Trinitrophenol (Various Mfr.).

ANTISEPTIC, SURFACE-ACTIVE AGENTS.

See Bactine (Miles)
 Benzalkonium Chloride. (Winthrop Labs)

Benzethonium Chloride (Parke, Davis)
 Ceepryn Chloride (Merrell).
 Cepacof (Merrell).
 Cetyl Pyridinium Chloride (Merrell)
 Diaporene Chloride (Homemaker's)
 Methylbenzethonium Chloride (Various Mfr.)

Phemeral Chloride (Parke, Davis)
 Roccal. (Winthrop Labs)
 Sulstide (Merck Sharp & Dohme)
 Thonzonium Bromide (Nepero)
 T-Cap Cream (Wallace & Tiernon)
 Zephiran Chloride. (Winthrop Labs)

ANTISEPTICS.

See: Dyes.

Furacin (Eaton).
 Iodine Products
 Mercurials.
 N Chloro Compounds.
 Phenols.

Surface Active Agents.

ANTISPASMIN CITRATE. (Roymer) Sestrin.

N-ethyl-2,2'-diphenyl-dipropylamine citrate
 Tab. 120 mg. Bot. 100s, 500s, 1000s

See: Profenid. (C. D. Smith)

Use. Relaxes smooth muscle.

W/Phenobarbital, (Roymer) Antispasmin citrate 120 mg phenobarbital 15 mg / Tab Bot. 100s 1000s

ANTISPASMODICS. Usually refers to agents that combat the muscarinic effect of liberated acetylcholine. The relief of smooth muscle spasms. Parasympatholytic agents. See: Abuphenine Salts (Chemo Pura)

Adiphenine.
 AntichoEnergie Agents
 Antispasmin, Tab. (Roymer)
 Artane (Jedlerie).
 Atropine Methylbromide.
 Atropine Methylnitrate.
 Atropine Salts.
 Belladonna Products.
 Bellotoline (Sondaz)
 Benzyl HCl (Merrell)
 Cycloplegia
 Dibukine. (Merck Sharp & Dohme)
 Droxyline Phosphate (Lilly).
 Disipol, Tab. (Riker)
 Ethaverine HCl (Various Mfr.)
 Genascopolamine.
 Homatropine Methylbromide (Various Mfr.)
 Hyoscyamine Salts (Various Mfr.)
 Marplan, Prep (Rache)
 Methscopolamine
 Murel, Tab (Ayerst)
 Neotropine HCl (Warner)
 Octin (Knoll)
 Papaverine Salts (Various Mfr.)
 Parkinsonism
 Pavafrane (Searle).

Poveril (Lilly)
 Profenil (C D Smith)
 Scopodex (Lobica Debrulle)
 Scopolamine Amineoxide
 Scopolamine Salts
 Sesiron (C D Smith)
 Suovil Tab (Merck Sharp & Dohme)
 Synliron (Roche)
 Trasenine (Ciba)
 Trocaine HCl (Paythress)
 Valerhamate Bromide

ANTISTERILITY VITAMIN See Vitamin E

ANTISTINE HCl, NND 1958 (Ciba) Antazoline hydrochloride

Use Antihistamine
 W/Pyribenzamine See Dibistine (Ciba)

ANTISTINE PHOSPHATE, NND 1962 (Ciba) Antazoline phosphate Sol Oph (0.5%) Bot w/dropper 15 cc

Use: Antihistamine
ANTISTINE-PRIVINE (Ciba) Antistine HCl 0.5% & privine HCl 0.025% Bot 1 oz w/dropper; 16 fl oz

Use Nasal decongestant with antihistamine
ANTITHROMBIN W/Protrombin Thrombokinase See Neo Hemoplastin Vial (Parke Davis)

ANTITHYROID AGENTS

See: Iothaluroc I
 Itrum I Tab (Ciba)
 Methiocal, Tab (Schwartz)
 Methimazole
 Methylthiouracil (Various Mfr)
 Muraol, Tab (Organon)
 Propylthiouracil (Various Mfr)
 Tapazole (Lilly)
 Thiomecil (Phys Drug)
 Thiouracil Tab (Parke Davis)

ANTITOXINS

See Botulism Antitoxin (Lederle),
 Diphtheria Antitoxin USP
 Gas Gangrene Antitoxin NF
 Tetanus Antitoxin USP

ANTITUBERCULOSIS AGENTS

See: p Aminosalicylates (No Co, K) (Varr
 ous Mfr)
 Benzopas Tab (Dorsey)
 Diasone Sod (Abball)
 Dihydrostreptomycin (Various Mfr)
 Glucosulfone Sodium (Parke Davis)
 Isoniazid (Various Mfr)
 Promacetin (Parke Davis)
 Promin (Parke Davis)
 Promizole Tab (Parke, Davis)
 Pyrazinamide Tab (Merck Sharp & Dohme)
 Rez pas (Squibb)
 Seromycin (Lilly)
 Streptomycin (Various Mfr)
 Teebazone Tab (Consal Midland)

* Generic name recognized by NF, USP or A.M.A.

Thiazolsulfone (Parke, Davis)
 Thiouanatinamide (Hoffmann La Roche)
 Tubercate Tab (Heun)
 Viomycin (Parke, Davis; Pfizer)

ANTI-TULARENSE AGGLUTINATING

SERUM (Lederle) Vial 1 cc
 Use Diagnostic agent in the identification of Pasteurella tularensis

ANTI-TUSSIVE, (Canright) Dexametharphon HBr 10 mg potassium guaiacol sulfonate 125 mg terpin hydrate 100 mg, phenyl propanolamine HCl 125 mg, pyrilamine maleate 125 mg /Tab Bot 60s
 Use Antitussive

***ANTIVENIN, BLACK WIDOW SPIDER, USP NND 1962** (Merck Sharp & Dohme) (Latrodectus Mactans) Antitoxic serum (horse) Lyovac Vial 2.5 cc
 Use Treatment of black widow spider symptoms

ANTIVENIN, SNAKE, POLYVALENT, NND 1962 (Wyeth) Crocaline antivenin polyvalent USP (Neotric Crocalidae) Rattlesnake copperhead & moccasin antitoxic serum Pkg, Combination 15 cc
 Use Bites by crocalid snakes of the United States

ANTIVERT (Roerg) Meclizine 12.5 mg, nicotinic acid 50 mg /Tab Bot 100s Syr (meclizine HCl 6.25 mg, nicotinic acid 25 mg /5 cc Bot 1 pt)
 Use Vertigo

ANTIVY, (Ciba) Pyribenzamine hydrochloride 2%, zirconium oxide 4%, Lat Bot 80 ml
 Use: Control of dermatitis due to poisonous oak and sumac

ANTIXEROPHTHALMIC VITAMIN, See: Vitamin A

ANTOPIC (Wyeth Dorsey) Pyrilamine maleate 5%, calamine, zinc oxide & benzocaine in a water soluble base Tube (1 1/2 oz)
 Use: Anesthetics w/local anesthetic

ANTOSEN (Squibb) Propenpyridamine 25 mg, codeine phosphate 65 mg alcohol 5% /R oz Bot 1 pt, 1 gal
 Use (1 dr q 3 h) in common cough
 Sedative effect

ANTORA-H TD (Maynard) Phenobarbital 1/4 gr hyoscyamine sulf 0.3858 mg atropine sulf 0.0750 mg, hyscine HBr D0222 mg, pentaerythritol tetranitrate 30 mg /imed disintegration cap Bot 60s 250s
 Use Prevention of attacks in angina pectoris

ANTRENYL BROMIDE, NND 1962 (Ciba) (Oxyphenantrum Bromide) Inj Sals Vial (1 & 2 mg /cc) 10 cc Syr (5 mg /4 cc)

- Bot. 1 pt. Tab. (5 mg) Bot. 100s, 500s, 1000s.
 Use: Anticholinergic activity.
- ANTRENYL BROMIDE PEDIATRIC DROPS.** (Ciba) Oxyphenonium bromide 1 mg/drop. Bot. 5 cc. w/dropper.
 Use: Anticholinergic.
- ANTRENYL BROMIDE PHENOBARBITAL.** (Ciba) Oxyphenonium bromide 5 mg. Phenobarbital 15 mg./Tab. Bot. 100s.
 Use: Anticholinergic with sedative.
- ANTROCOL.** (Paythress) Phenobarbital gr. ¼, colloidal sulfur gr. ½, atropine sulfate gr. 1/200/Tab. Bot. 100s, 500s.
 Use: (q. 3 h. p.r.n.) Sedative, antispasmodic
- ANTRYPOL.** See: Suramin Sod. (Winthrop Labs.)
- ANTUITRIN-GROWTH.** (Parke, Davis) Anterior pituitary extract containing growth factors. Vials (8 rat growth units/cc) 10 cc.
 Use: I.M., stimulating growth.
- ANTUITRIN-S.** (Parke, Davis) Chorionic Gonadotropin. Vial (5000 I.U./cc) when 10 cc. diluent is added.
 Use: I.M., cryptorchidism and functional uterine bleeding.
- ANTURANE, N N D, 1962.** (Geigy) Sulfapyrazone 100 mg/Tab. Bot. 100s
 Use: Treatment of chronic gout.
- ANUCAINE.** (Calvin Chem.) Procaine, butyl-p-aminobenzoate, benzyl alcohol in sweet almond oil. Amp. 5 cc., Box 6s, 24s, 35s, 100s.
 Use: I.M., anesthetic for rectal treatment.
- ANUSCAINE.** See: Analgescaine, Vial (Metra)
- ANUSOL.** (Warner-Chilcatt) Bismuth subgallate (2.25%), bismuth resorcin compound (1.75%), Nicaraguan balsam (medicinal) (3.00%), zinc oxide (11.00%), boric acid (5.0%), cocoa butter ointment base q.t. ad 100.00%. Boxes 6s, 12s, 24s
 Use: Rectal suppository.
- ANUSOL-NC.** (Warner-Chilcatt) Same as Anusol plus hydrocortisone 10 mg/Supp. Box 12s.
- ANUSOL UNGUENT.** (Warner-Chilcatt) Bismuth subgallate, bismuth resorcin compound, Nicaraguan balsam, zinc oxide & boric acid. Tube 1 oz.
 Use: Topically, irritative & inflammatory disorders of the skin.
- ANZOL SOLUTION, A D R, 1955.** (Veltex Co.) Benzethonium chloride ¼% w/Sol. carb & Sod. nitrite. Bot. 1 gal.
 Use: Antireptic
- AOLAN.** (Duke) Lactalbumin suspension Amp. 2 cc., box 6s, 100s; 5 cc., box 6s, 5Dr; 10 cc., box 6s, 50s.
 Use: Non-specific protein therapy.
- AORAL.** (Marsengill) Vit. A Cap (25,000 u) Bot. 100s, 500s; (50,000 u.) Bot. 100s.
 Use: Vitamin A therapy.
- APAC.** (Chicago Pharm) Arpirin 3½ gr., acetophenetidin 2½ gr., caffeine ¼ gr., phenobarbital ¼ gr./Tab. Bot. 100s, 1000s.
 Use: Analgesic, redative.
- APAMIDE.** (Ames) N-acetyl p aminophenol. Tab. (0.3 Gm). Bot. 100r.
 Use: Analgesic, antipyretic.
- W/Acetylcaramol.** See: Apromol, Tab. (Ames).
- APAMIDE-VES.** (Ames) N-acetyl-p-aminophenol 0.3 Gm., in a citrate carbonate base/Tab. Pkg. 50r.
 Use: (1-2 q. 4 h) Analgesic & antipyretic.
- A-P-A-P.** (Kendall) N-acetyl p-aminophenol. 2½ gr. Tab. Bot. 100s, 500s.
 Use: Relief of pain.
- APATATE DROPS.** (Kenwood Labs) Thiamine HCl 15 mg, Vit. B-12 25 mcg/cc. Bot. w/dropper 15 cc., 30 cc.
 Use: Orally; nutritional supplement.
- A.P.C., N.F.** (Vanauis Mfr.) Aspirin, phenacetin & caffeine.
 See: A.C.A. Compound, Cap., Tab. (Eowell). Ace Caf-Edine, Tab. (Paul Maney) Acetidine, Cap., Tab. (Merck Sharp & Dahme)
 Anacin, Tab. (Whitehall)
 Analgia No. 1, Tab. (Merrell)
 Ansemco No. 1, Tab. (Massengill)
 Aphodyne, Tab. (Gold Leaf)
 A.S.A. Compound, Pulv. (Lilly)
 Ascadin, Pulvoids (Drug Prods.)
 At-Co-Phen, Cap. or Tab. (Ulmer)
 Aspodyne, Tab. (Lemman).
 Aspirafine Comp. No. 3 (McNeil).
 Arpirotabs, Tab. (Brewer).
 Aspodyne, Tab. (Blue Line).
 Empirin Comp. Tab. (Burroughs Wellcome).
 P.A.C. Compound, Cap. or Tab. (Upjohn)
 Phen-acetyl, Tab. (Smith-Dorsey)
 Toc-55, Tab. (Norwich)
- W/Codeine phosphate.** (Lilly) Tab. ¼, ½ & 1 gr.
 See: Acetidine w/Codeine, Tab. (Merck Sharp & Dahme)
 Anxasia w/Codeine, Tab. (Massengill).
 Cadcol, Tab. (Horton & Converse).
 Cogesic-16, Tab. (Puget Sound)
 Empirin compound w/Codeine phosphate, Tab. (Burroughs Wellcome).
- W/Demerol.** (Winthrop Labs.) Arpirin 3 gr., phenacetin 2½ gr., caffeine ¼ gr., demerol HCl ½ gr./tab. Bot. 100r.
 Use: Analgesic.
- W/Gelsemium.** See Table, p. 72.
- W/Phenobarbital**
 See: Acetobar, Tab. & Water (Calumbar).
 Amphaphen, Tab. (Gold Leaf)

- APAC, Tab. (Chicago).
 Aphaphen, Tab. (Gold Leaf).
 Asborphen, Tab. (Lemman).
- A-P-C NO. 2.** (Sutliff & Case) Aspirin 260 mg., Acetophenetidin 130 mg., Caffeine 16 mg./Tab. Bot. 1000s.
- A-P-C PEDIATRIC.** (Sutliff & Case) Aspirin 2 gr., Acetophenetidin 1 gr., Caffeine $\frac{1}{2}$ gr./Tab. Bot. 1000s.
- A-P-CILLIN.** (White) Procline penicillin G 100,000 u., pheeytoloxamine dihydrogen citrate 25 mg., acetylsalicylic acid 150 mg., phenacetin 120 mg., caffeine 30 mg./Tab. Bot. 50s, 500s.
 Use: Secondary bacterial infection.
- A-P-CILLIN-200.** (White Labs.) Procline penicillin G 200,000 u., acetylsalicylic acid 150 mg., phenacetin 150 mg., caffeine 45 mg., dipheylpyrrolidine HCl 2 mg./tab. Bot. 24s, 100s, 500s.
 Use: 1 q.l.d. respiratory infections.
- APCO.** (Hence) Aloin $\frac{1}{2}$ gr., belladonna ext. 1/12 gr., Ipecac 1/15 gr., phenolphthalein $\frac{1}{2}$ gr./Pill. Bot. 100s, 1000s.
 Use: Laxative.
- APCOPHIS.** (Donley-Evans) Phenylephrine HCl 10 mg., chlorpropenpyridamine maleate 2 mg., pyrilamine maleate 6 mg., methoprene HCl 3 mg., aspirin 3 $\frac{1}{2}$ gr., phenacetin 2 $\frac{1}{2}$ gr., caffeine $\frac{1}{2}$ gr./Tab. Bot. 100s, 500s.
 Use: Common cold.
- APERENS.** (Broyten) Cassia ougustifolia (senna) 275 mg./Tab. Bot. 50s.
 Use: Laxative.
- APERGEL.** (Northwest Labs.) Carboxymethylcellulose Sod. Sol. 5%. Bot. 1 pt. Tab. (0.5 Gm.) Bot. 100s.
 Use: Antacid—laxative.
- APESTRIN.** (Harrower) Gonadotropic substance of chorionic origin, obtained from human pregnancy urine. Vial (1000 I.U. or 5000 I.U.) 10 cc. size with solvent.
 Use: I.M., cryptorchidism & functional uterine bleeding.
- APEXOL.** (Roerig) Vit. A nat. 50,000 u. Cap. Box 50s.
 Use: Vitamin A therapy.
- APG-HORM.** (Canfield) Equine pituitary gonadotropic hormone 125 u./5 cc. Vial.
 Use: Treatment of amenorrhea.
- APHCO.** (American Pharm.) Benzocaine, bismuth subgallate, bismuth oxyiodide, zinc oxide, bismuth resorcin sulfonate, peruvian balsam, & boric acid Oint. 1 $\frac{1}{2}$ oz. Supp. Box 12s.
 Use: Hemorrhoidal therapy.
- APHODYNE.** (Gold Leaf) Aspirin 3 $\frac{1}{2}$ gr., acetophenetidin 2 $\frac{1}{2}$ gr., caffeine $\frac{1}{2}$ gr./Tab. Bot. 100s, 1000s. Juniors (approx. $\frac{1}{4}$ the above amounts.) Bot. 100s, 1000s.
 Use: Analgesic, antipyretic.
- APHOPHEN.** (Gold Leaf) Aspirin 3 $\frac{1}{2}$ gr., acetophenetidin 2 $\frac{1}{2}$ gr., caffeine $\frac{1}{2}$ gr., phenobarbital $\frac{1}{8}$ gr./Tab. Bot. 100s, 1000s.
 Use: Analgesic—sedative.
- APNRODISIAC.** See: Pego Palo Faridom (Quimica Dominicana)
- APITOL.** (Premier Pharm. Corp) Liquid 4 oz.
- APITOR.** (Estra) Chorionic gonadotropin. Vial (5000 u.) 10 cc. w/diluent.
 Use: I.M. (100 to 1000 units 2 or 3 times weekly) for cryptorchidism.
- APISED.** (First Texas Chem.) d-Amphetamine sulfate 2.5 mg., phenylpropanolamine HCl 25 mg., theobromine 150 mg./Tab. Bot. 100s, 500s.
 Use: Appetite depressant.
- "A.P.L."** (Ayerst) Chorionic gonadotropin. Solution (500 I.U., 1000 I.U./cc.) Vial 10 cc. Dry form, package provides one "Secule" containing 20,000 I.U. and one 10 cc. vial sterile diluent.
 Use: Chorionic gonadotropin therapy.
- APOLAMINE.** (Winthrop Labs) Leminal 15 mg., atropine sulfate 0.1 mg., scopolamine HBr 0.2 mg., benzocaine 100 mg., riboflavin 4 mg., pyridoxine 2 mg., nicotinamide 25 mg./tab. Bot. 100s.
 Use: Antinausea.
- APOLARTHROX.** (Roerig) Vit. A 30,000 u., D 25,000 u./Cap. Box 60s.
 Use: Vitamin A & D therapy.
- *APOMORPHINE HYDROCHLORIDE, U.S.P.** Tablets, N.F.; Pow., Bot. 5 gr., 15 gr., $\frac{1}{2}$ oz., 1 oz. (Various Mfr) Lilly, H.T. (1/10 gr.) Tube 20s, 100s, 500s, 1000s.
 Use: Emetic.
- APOTHELINE HYDROCHLORIDE, N.R.** 1946. (Parke, Davis) Diethylaminoethyl cianamate hydrochloride.
- APPELLA.** (Winthrop Labs.) Apple Pow. Bot. 7 & 18 oz.
 Use: Diarrhea.
- APPETROL.** (Wallace) d-Amphetamine sulfate 5 mg., Miltown 400 mg./Tab. Bot. 50s.
 Use: Anorexiant.
- APPETUM.** (Philo. Ampoule) Vitamin B-1 10 mg., B-12 25 mcg./tab. Bot. 28s.
 Use: (Chew 1 daily) Improve appetite.
- APPLE JUICE CONCENTRATE.** See: Infase, Liq. (Borden).
- APPLE POWDER.** See: Appella, Pow. (Abbott)
- APPLIDERM No. 1.** (White) Menthol 0.2%, hexachlorophene 0.1% in an aq. soln. of

* Generic name recognized by N.F., U.S.P. or A.M.A.

- Epinephrine and isopropyl alcohol. Spray pack
size 3 oz.
- Use Careful
- APPLIDERM No. 2. (W314) Menthol 0.2%,
benzocaine 0.1%, in an emulsion
base. Tube 1 1/2 oz.
- Use Careful
- APPLIDERM No. 3. (W315) Water in oil
emulsion consisting of petrolatum with
various emulsifiers as the emulsifying
agent. Tube 1 1/2 oz.
- Use - It is protective
- APPLIDERM No. 4. (W316) 0.5% Benzoin
and 2% ppt. Vaseline is a stabilized
preparation, emulsion base. Tube 1 1/2 oz.
- Use - Substrate & clean area
- APPLIDERM No. 5. (W317) Salicylic acid
2% w/w 20%, in an anhydrous, water
insoluble base. Tube 1 1/2 oz.
- Use Careful
- APPLIDERM No. 6. (W318) Crude coal tar
2% in a water-in-oil emulsion base. Tube
1 1/2 oz.
- Use Careful
- APPLIDERM No. 7. (W319) Undecylenic acid
1% benzocaine 0.1%, in a hydro-
philic emulsion base. Tube 1 1/2 oz.
- Use Careful
- APPROLONE HYDROCHLORIDE, NND
1917 K. Bar preparation hydrochloride
Sup 17 mg 1 cc. 34 Tab. 10 mg, 20
mg, 30 mg & 100 mg. Bot 100, 300,
500.
- Use - Effective in therapy of hypertension's
disease.
- See Supplement Prod. Files
- APRILONIN EXCISE K. Bar. Approval
11 mg. 100 mg 100 mg. Bot 100.
- Use - Suspension
- ARPROBATEL, M.P. 163.8 1912 (24%
suspension) and 25% suspension
suspension. Bot. 100 mg. 100 mg. Bot 100.
See Supplement Prod. Files
- See Long Acting Approval
- See also Suspension 100 mg. Bot 100.
- ARPROBATEL EXCISE, M.P. 163.8 1912
100 mg. 100 mg. Bot 100. 100 mg. Bot 100.
- See Long Acting Approval
- See also Suspension 100 mg. Bot 100.
- ARPROBATEL EXCISE (25% suspension) 100 mg. Bot 100.
- See Long Acting Approval
- See also Suspension 100 mg. Bot 100.
- ARPROBATEL EXCISE (100 mg. Bot 100).
- See Long Acting Approval
- See also Suspension 100 mg. Bot 100.
- ARPROBATEL EXCISE (100 mg. Bot 100).
- See Long Acting Approval
- See also Suspension 100 mg. Bot 100.

- Use: (1 q 3-4 h) Sedative, analgesic.
- APROTIVE. (Approved Pharm) Dicalcium
Phos 900 mg, ferrous sulfate 30 mg, Va
B 1 2 mg, B-2 2 mg, D 400 u./Cap. Bot.
100s, 1000s, 5000s
- Use Prenatal therapy
- A.P.S. Aspirin, phenacetin and salicylamide
- APYRON. See: Magesium acetylsalicylate
(Vimmary).
- AQUASAB. (Webster) Supp. 1/2 gr
12s, 1 gr 12s, 1 1/2 gr, 12s, 3 gr, 12s
- AQUA-B-ASCORBIC ACID. (Upjohn) Vit B 1
12.5 mg, B 2 3 mg, B-6 1.5 mg, d panto-
thyl alcohol 15 mg, nicotinamide 50 mg,
ascorbic acid 75 mg/cc. Vial 10 cc.
- Use IV or I.M., nutritional supplement
- AQUACHLORAL. (Webster) Chloral hydrate
Caps 3 1/2 gr, 7 1/2 gr. Bot 100s. Supp. 1/2
gr, 10 gr, 15 gr. 12s. Syr. (7 1/2 gr / 5 cc)
1 pt
- Use: (Rectal) sedative
- AQUACILLIN. (Schenley) Procaine penicillin
G crystalline 100,000 units/cc. Vial 1 cc.
5 cc, 10 cc.
- Use: I.M., respiratory penicillin therapy
- AQUACILLIN-A.S. (Schenley) Aq. susp. pro-
caine penicillin G, 300,000 units/cc. Vial
1 cc, with disp. syr. Vial 5 cc, 10 cc.
- Use: I.M., penicillin therapy
- AQUACILLIN-A.S./600. (Schenley) Aq.
susp. procaine penicillin G, 600,000 units
cc. Vial 1 cc, with disp. syr. Also available
to 1,000,000 units.
- Use: (Double strength) penicillin therapy
- AQUACILLIN A.S./One Million. (Schenley)
Penicillin G procaine 1,000,000 u./dose
Vial, Aq. 2 vial
- Use: Penicillin therapy
- AQUACORE SUPPRETTES. (Wm A. Webster
Co.) Hydrocortisone acetate, 10 mg; phen-
thol, USP, 2.5 mg; phenylmercuric
borate, 3 mg; 8-aminocapric acid, 3 mg;
polyorbate 90, USP, 10 mg; urea 20 mg
borate, 50 mg / Supprett Jars of 14
- Use: Vaginal
- AQUACRINE. (Endocrine) Estrone 20,000
IU/cc. Vial 10 cc.
- Use: Menopausal syndrome, post-menopausal
- AQUADOL. (Parsons Drug) Estrad-ol in oil
susp. injections 100 mg, 0.66 mg, or
1.33 mg/cc. 10 cc or 25 cc.
- Use: I.M., estrogen replacement therapy
- ARPROBATEL EXCISE (Parsons Drug) (Schenley)
0.66 mg, progesterone 25 mg / cc. Vial 10
cc
- Use: Progesterone (Parsons Drug) (Schenley) 0.66
mg, progesterone 25 mg / cc. Vial 10 cc
- Use: Progesterone (Parsons Drug) (Schenley) 0.66
mg, progesterone 25 mg / cc. Vial 10 cc
- Use: Progesterone (Parsons Drug) (Schenley) 0.66
mg, progesterone 25 mg / cc. Vial 10 cc

© Copyright 1962 by Parke-Davis & Co., Detroit, Mich. All rights reserved.

AQUAGEN. (Lemman) Beta-estradiol aq. susp. Vial (0.2 mg., 0.5 mg./cc.) 10 cc., 30 cc.

Use: I.M. (0.2 mg.-0.5 mg. daily) for estrogenic hormone deficiencies.

AQUAGESTIN. (Harvey Lab.) Natural estrogenic substances 25,000 I.U. & progesterin 25 mg./cc. Vial 10 cc.

Use: I.M., estrogen and androgen therapy.

AQUA IVY. (Strong Cabb) Poison Ivy ext., alum precipitated. Vial 3 cc. & 7.5 cc. w/diluent.

Use: I.M., treatment of & prophylaxis of poison ivy dermatitis.

AQUA IVY AP. N.N.D. 1962. (Syntex Chem. Co) Ext. of poison ivy and oak leaves. Bot. 5 cc. (1:50 dil.), 10 cc. 1:5 dil.) Orol Tabs. (1.2 mg. of alum ppt. pyridine ivy). Bot. 50s, 100s.

Use: Immunization against poison ivy.

AQUAKAY. See: Menadione, U.S.P.

AQUAMEPNTON INJECTION. (Merck, Sharp & Dahme) Phytanadiol (Vit. K-1) 10 mg./Amp. 1 cc./Box 6s. 1 mg./D5 cc. Amps. Box 25s.

Use: Control of hemorrhage.

AQUAMIC QLEUM A. (Biopharmaceuticals) Water-miscible vit. A 25M and 50M u./Cap. Bot. 100s, 1000s. Qint. 3000 M.u./oz.

AQUANEPHTON. (Merck Sharp & Dahme) See: *Phytanadiol Sodium Diphosphate.

AQUAPHOR. (Duke) Water-miscible ointment base. Tube 2 oz. Jar 1 lb., 5 lb., 50 lb.

Use: Water-miscible ointments.

AQUAPLEX. (New Mexico Pharm. Co) Vit. A 10,000 I.U. D 1,000 I.U., E 2 mg., C 75 mg., Nicotinamide 25 mg., S-1 10 mg., B-2 2 mg., B-6 3 mg., Panthelol 2 mg., Senzyl Alcohol 2%, Gentisic Acid Ethanolamide 1.5%/2cc. 10cc. Vials.

AQUAPRIOL. (Estra) Beta-Estradiol 1.0 mg. & progesterone 12.5 mg./cc. Vial, aq. susp. 10 cc.

Use: I.M. (1 cc. daily), corpus luteum & estrogenic therapy.

AQUAPROPEN. (Massengill) Penicillin G procaine 300,000 units, penicillin G Pot. 100,000 units/dose in 5-dose vial. Vial 20 cc./diluent.

Use: I.M., penicillin fortified therapy.

AQUARIUS. (Frasst) Hydrachlarathiazide 25 mg., 50 mg./Tab.

Use: Diuretic.

AQUASIL. (Lentheric) Silicone lat.

Use: Skin protective.

AQUASOL A. N.N.D. 1962. (U. S. Vitamin) Vit. A, nat., water soluble. Amp. 2 cc. (50,000 u.), 50,000 u./cc. Vial 10 cc. Cap.

(25,000, 50,000 or 100,000 u./Cap) Bot. 100s, 500s, 1000s. Sol. (50,000 u./cc.) Bot. 15 cc. or 30 cc. with dropper.

Use: Orally, I.M., vitamin A therapy.

AQUASOL A W/PANTHODERM. (U. S. Vitamin) Vit. A 200,000 u., panthelol 2% in a water-miscible base. Oint. Tube 1 oz. Use: Topically, emollient & healing aid.

AQUASOL A & D. (U. S. Vitamin) Aq. sol. of fish liver oil containing Vit. A 3,636 u., & Vit. D 727 u./0.1 cc. Bot. w/dropper 15 cc., Bot. 240 cc.

Use: Orally, vitamin A & D therapy.

AQUASOL A-C-D. (U. S. Vitamin) Vit. A 5000 u., D 1000 u., C 75 mg./0.6 cc. Bot. w/dropper 15 cc., 30 cc.

Use: Orally, multivitamin therapy.

AQUASOL E. (Funk) dextra Alpha-tocopherol. Cap. (30 & 100 mg) Bot. 100s, 500s, 1000s. Eq. (50 mg./cc.) Bot. 15 cc., Amp. 2 cc., Box 6s, 10 cc.

Use: Vitamin E therapy.

AQUASPENIQN ESTROGENIC SUBSTANCE. (Pitman-Moore) Estrogenic substance obtained from pregnancy urine. Vial 20,000 I.U. (2 mg./cc.) 10 cc.

Use: I.M., estrogenic replacement therapy.

AQUASPERSE ACID DRQPS. (White) Vit. A 5000 u., D 1000 u., C 50 mg./0.6 cc. Sol. w/dropper 15 cc. & 50 cc.

Use: Multi-vitamin therapy.

AQUA-STRONE. (Kirk) Estrogenic substance. Vial 10 & 30 cc. (20,000 I.U./cc.)

Use: I.M., estrogenic replacement therapy.

AQUASTRONE. (Neon) Estrogenic substances nat. Vial (20,000 u./cc. aq. susp.) 10 cc., 30 cc.

Use: I.M. (0.5 cc. weekly); menopausal symptoms.

AQUASUSPENSION G. (Pitman-Moore) Penicillin G procaine 300,000 u./cc. Vial, Aq. susp., 10 cc.

Use: Penicillin therapy.

AQUASYNTH. (U. S. Vitamin) Vit. A palmitate. Cap. (25,000 u.) Bot. 100s, 500s, 1000s (50,000 u) Bot. 100s, 500s, 1000s. Use: Vitamin A therapy.

AQUA-TESTERONE. (Endocrine Co) Testosterone crystalline aq. susp. Vial (25 mg./cc.) 10 cc.

Use: I.M. (1 cc. weekly); androgen therapy.

AQUATYL. (Irwin, Neisler) Dioctyl sod. sulfosuccinate 50 mg., cholic acid 100 mg./Tab. Bot. 60s

Use: Fecal softener

AQUAVITIN. (Kramers-Urban) Vit. B-1 10 mg., riboflavin 0.15 mg., niacinamide 15 mg., Vit. B-6 0.5 mg., panthelol 2.5 mg., Vit. C 30 mg./cc. Vial 10 cc., Box 25s, 100s.

* Generic name recognized by N.F., U.S.P. or A.M.A.

Use I.V., vitamin K complex with vita
min C.

AQUISTRONE (Afas) Natural estrogen
20,000 LU./cc. Vial 10 cc., 30 cc.

Use I.M., menopausal disturbances.

AQUINONE See Menadione USP

AQUINOL (Estr) beta Estradiol aq susp
Vial (10 mg./cc.) 10 cc.

Use I.M. (1 mg 1 to 2 times weekly)
menopausal symptoms.

ARALIN PHOSPHATE, N.M.O. 1962 (Win
throp Labs) Chloroquinol phosphate Tab
125 & 250 mg., Bot 100s 1000s.

Use: Crabs; suppression & treatment of
malaria. Also for extra-intestinal amebiasis
(amoebic hepatitis). Also rheumatoid arthritis

W/Anelina MCL, Plaqueol sulfate See: Tri
quin Tab. (Winthrop Labs.)

W/M/3a. See Milbils w/Aralen Diphos
phate Tab. (Winthrop Labs.)

ARAMINE BITARTRATE, N.M.O. 1962 (Merck
& Dahme) Meteron not bitartrate
1/2% Sol. Bot 1 oz. w/M/1 atomizer Bot
1 oz. w/dropper 1/2% 1% Amp 1 cc.,
See 12s, Vial 10 cc.

Use Decongestant

ARANTHOL (Kant) 2 Methylene-6-hy
droxy-4-methylheptane Amp (100 mg./cc.)
1 cc., Tab. (122 mg) Bot 25; 100s Sterile
sola (100 mg./cc.) Vial 10 cc.

Use Analgesic & anesthetic

ARIC SUPPOSITORIES (Fellows) Penta
barbital Sod CCS Gm., phenobarbital
Sod CCS Gm., quinalbarbital 0.50 Gm.,
ethyl amobarbital 0.10 Gm./Supp Bot
12s

Use Somnolytic & sedative

ARCOLASE (Sharp) Trizyme 2E mg., Epso
11 mg./Tab Bot 50s

Use Digestive disorders

ARBUTIN, N.M.R. 1937 (Abbott, Merck
& Sharp & Dohme) [Fellows] Hydroquinone Beta
D-glucopyranoside

Use Diuretic

ARCOCOL (Sharp) Sharp & Dohme) Not
solub. but act. 2 cc. calcium hypophos
phate 24 mg., and hypophosphate 0.1 Gm.,
pot hypophosphate 0.1 Gm., quinine hypo
phosphate 1 mg., and low act. CCS Gm./
3" in Bot per

Use Stimulant to the metabolic functions

ARCOCAL (Lanar) Lanarol 68 Gm.,
quinalbarbital 25 mg., Vx 1 11.5 mg./
11 cc. Sol., Bot 12 cc

Use: Convuls 12 cc daily treatment of
epilepsy

ARCOLASE (Sharp) Trizyme 11 mg (Sharp
& Dohme) Same composition by N.F., USP or A.S.A.

lytic 30 mg., proteolytic 6 mg., celluloly
2 mg./lipase 25 mg./Tab Bot 50s

Use: Digestive disorders

ARCOLASE PLUS, (Arco) Trizyme omyolytic
30 mg., proteolytic sulfate 0.02 mg., hy
cyanine sulfate 0.10 mg./Tab Bot 50s

Use: Treatment of gastrointestinal
disturbances

ARCOSTERONE ENANTINATE INJ. (Arco)
Testosterone enanthate 100 mg./cc. 10 cc
multiple dose vials

Use: Androgen

ARCYLATE (Bentex Pharm) Sod thiosally
ole Amp (100 mg) 2 cc Vial 50 cc

Use I.M., arthritis, rheumatism and related
conditions

ARECOLINE ACETARSONE SALT, See: Dr
carb 1 NF

ARECOLINE HYDROBROMIDE, NF (Mc
Inckrodt) Tablets NF Hydrobromide of a
alkaloid obtained from the dried ripe seed
of areca catechu Tab (91 to 109%
arecoline hydrobromide)

ARFONAO, N.N.D. 1962 (Roche Lab) (1,
methaphan camphorsulfonate) d 3,4(1,
dibenzyl 2'-keto-imidazolido) 1,2 trimethy
lene thiophanum d camphor sulfonate
Amp (50 mg./cc) 10 cc., Box 6s & 25s
Use: Short-acting vasodilator for control
hypertension

ARGININE HCL, N.N.D. 1962 See: A
rginine (Gray Pharm)

ARGININE GLUTAMATE, See: Modusol
(Abbott)

ARGIVENE, N.N.D. 1962 (Gray Pharm)
Arginine HCl 20 Gm Vial 50 cc

Use: I.V. Encephalopathies associated w/
disturbances of the liver

AROYN, N.M.R. 1949 (Abbott) See: M/
Silver Protein (Various Mfr)

AROYPUVIB (Barnes) Argypulv 20% loaf
40%, lactose 40% Pow Pig Cop (2 Gm
12s Pig Bot (7 Gm) 3

Use: Vaginal anti-infective

AROTOL, N.M.R. 1920 (Barnes) Mild h
over protein. Crystals Bot 1 oz., 4 oz Sol
Fon (5%, 10%, 15%, 20%, 25%) Pig 1
oz., 1/2 oz., 1 oz., 2 oz., 4 oz; Tab 2
Gm 1 Bot 37s Pig 3 Bot Stabilized w/
Bot 1/2 oz., 1 oz

Use: Topical nasal and ophthalmic in
fections

W/Keolis & lactose See: Argypulv 20
pow (Barnes)

ARISTOCORT, N.N.D. 1962 (Federal) Triac
etalone Tab. 1 mg Bot 50s 500s 2 ml
4 mg Bot 20s 100s 500s 1000s 87
2 mg/3 cc. Bot 4 cc

Use Anti-inflammatory steroid

W/Chlorpheniramine maleate, ascorbic acid.
See: Aristamin, Tab. (Lederle)
Use: Indicated in rheumatoid arthritis, hay fever, bronchial asthma.

ARISTOCORT CREAM, N.N.D. 1961. (Lederle) Triamcinolone acetonide 0.1%. Tube 5 & 15 Gm., 0.5% Tube 5 Gm., 15 Gm.

ARISTOCORT CREAM 0.5%, (Lederle Labs) Triamcinolone acetonide 0.5%, methylparaben U.S.P. 0.16%, propylparaben U.S.P. 0.04% in water base w/glyceryl monostearate, squalone, polyorbate 80 U.S.P., spermaceti U.S.P., stearyl alcohol U.S.P., and sorbitol solution N.F./Tube 5 Gm., 15 Gm.
Use: Treatment of dermatoses.

ARISTOCORT DIACETATE FORTE. (Lederle) Triamcinolone diacetate 40 mg./cc. Vials 1 & 5 cc.
Use: Parenteral steroid.

ARISTOCORT DIACETATE INTRALESIONAL. (Lederle) Triamcinolone diacetate 25 mg./cc. Vial 5 ml.

ARISTOCORT DIACETATE PARENTERAL. (Lederle) Triamcinolone diacetate 25 mg./cc. Vial 5 cc.
Use: I.S., I.A., I.M. Rheumatoid arthritis.

ARISTOCORT DIACETATE SYRUP. (Lederle) Triamcinolone diacetate (2 mg/5 cc.) Bot. oz.
Use: Corticosteroid hormone.

ARISTOCORT OINT. (Lederle) Triamcinolone acetonide 0.1%. Tube 5 & 15 Gm.

ARISTODERM FOAM, N.N.D. 1962. (Lederle) Triamcinolone acetonide 0.1%. Aerosol./Press. Bot. 7½ cc. & 15 cc.
Use: Dermatoses.

ARISTOL, N.N.R. 1949. (Merck; Winthrop abr.) See: Thymal iodide.

ARISTOGESIC. (Lederle) Triamcinolone 0.5 mg., salicylamide 325 mg., aluminum hydroxide 75 mg., ascorbic acid 20 mg./Cap. 100s.
Use: Rheumatoid arthritis and other moderate rheumatic conditions.

ARISTOMIN. (Lederle) Aristocort 1 mg., chlorpheniramine maleate 2 mg., ascorbic acid 75 mg./Cap. Bot. 30s, 100s.
Use: Allergic ailments

ARCAPS. (Arlington-Funk) Ephedrine HCl 6 mg., ascorbic acid 100 mg., aspirin 130 mg., phenobarbital 26 mg./Cap. Bot. 50s, 100s
Use: Relief from asthma, common cold & hay fever.

ARLIDIN, N.N.D. 1962. (Arlington-Funk) Nydrin HCl. Phenyl-1-butyl-norsupifen. 1-(p-hydroxyphenyl)-2-(1-methyl-3-phenyl-propyl-amino)propanal HCl.

Am.p. (5 mg./cc.) T cc., Box 6r, 25r, 100s; Tab. (6 mg.) Bot. 50s, 100s.
Use: Orally; vasodilator, increases peripheral circulation and cardiac output.

ARMAZIDE, N.N.D. 1960. (Armour) Isoniazid. Tab. (50 mg.) Bot. 100s.
Use: Streptomycin-resistant tuberculosis.

ARMIDE. (Warren) Salicylamine 320 mg., salicylsalicylic acid 160 mg./Tab. Bot. 100r. No. 2—Salicylamide 320 mg., salicylsalicylic acid 160 mg., butobarbital 16 mg./Tab. Bot. 100s.
Use: Aesthetic in treatment of dental pain

ARNE TIMESULES. (Hoag) Ferrus fumarate 330 mg., thiamine 5 mg./Cap. Bot. 30s.
Use: Aemia.

ARNEL-PR SUSP. (Arnell) Sodium phenobarbital 2 gr., atropine sulfate 1/40 gr., in colloidal susp. of aluminum and magnesium hydroxides/oz. Bot. 12 oz.
Use: Gastric hyperacidity.

AROBON. (Nertle) (Pitman-Moore) A powder made from specially prepared carob flour, high in pectin, lignin and hemicellulose. Jar 5 oz. Alra wafers Bot. 12s.
Use: Orally, symptomatic relief in all types of diarrhea

ARRESTIN. (Johnson & Johnson) d-Meltharphon.
Use: Antitussive.

ARSAMBIDE. See: Carbarsone (Various Mfr.).

ARSECLOR. See: Dichlorophenarsine hydrochloride (Parke, Davil).

ARSENEIN. (Eliel) Strychnine arsenate 1/64 gr., quinine arsenate 1/32 gr., iron arsenate 1/32 gr., nucleic acid 1/50 gr./Tab. Bot. 100r, 500s, 1000s
Use: Appetite stimulant
Arseamide Sod.

ARSENIC COMPOUNDS.
See: Acetarsan.
Aldarsone (Abbott).
Arsphenamine.
Arstinal (Endo).
Bolarzen (Endo).
Eisorphen (Endo).
Carbarsone (Various Mfr.).
Copper Cocodylate (Kramers-Urbac).
Dichlorophenarsine (Parke, Davil).
Ferric Cocodylate (Farnsworth).
Glycobarsol. (Winthrop Labs.)
Mapharsen (Abbott).
Mercaptoarsenat (Endo).
Milibis. (Winthrop Labs.)
Neoarsphenamine (Abbott).
Oxophenarsine HCl (Abbott).
Phenarsone (Abbott).
Sodium Cocodylate (Various Mfr.).
Thiocarbarsone (Lilly).
Tryparsamide (Endo).

W/Black pepper See: Asiatic, Tab (Hortae and Convesse)

ARSENO 39. See Osophenarsine hydrochloride (Parke, Davis)

ARSENOBENZENE. See Arsphenamine (Abbott)

ARSENOXIDE See: Osophenarsine hydrochloride (Parke, Davis)

ARSENPHENOLAMINE. See: Arsphenamine (Abbott)

ARSPHENAMINE, USP XIII, NNR 1949 (Abbott; Mesck) (Arsenobenzene, Arsenobenzol, Arsphenolamine, Ehrlich 606, Salvorsan) 3,3'-Diamino-4,4'-dihydroxyarsenobenzene dihydrochloride

Use: Antisyphilitic & in other spirochetal infections

***ARSTHINOL, NND 1962 2-(3'-acetamido 4-hydroxyphenyl) 1,3-dithia 2-oxacyclopentane 4-methanol mercaptoselenol** See: Selenarsen Tab (Endo)

ARTAMIDE (Wampole) Salicylamide 0.25 Gm, p-aminobenzoic acid 0.25 Gm, ascorbic acid 20 mg, organidin 20 mg / Tab Bot 100s, 500s

Use: (2 1 1 d) Rheumatic arthritis, rheumatic fever, osteoarthritis & gout

W/Calcichine (Wampole) Salicylamide 250 mg, p-aminobenzoic acid 250 mg, Vit C 20 mg, organidin 20 mg, calcichine 0.25 mg / Tab Bot 100s

Use: 1 2 q 3 hr; gout and gouty arthritis

ARTAMIDE-HC (Wampole) Salicylamide 0.25 Gm, p-aminobenzoic acid 0.25 Gm, Vit C 20 mg, hydrocarthosol 25 mg / Cap Bot 100s

Use: Treatment of rheumatic arthritis

ARTANE HYDROCHLORIDE, NND 1962 (Lederle) Trihexyphenidyl HCl Elixir (2 mg / 5 cc), Bot 1 pt, 1 gal; Tab (2 mg & 5 mg), Bot 100s, 1000s

Use: (1 1 1 d) Parkinsonism

ARTANE SEQUELS (Lederle) Trihexyphenidyl HCl 5 mg / Tab Bot 30s 500s

Use: Parkinsonism

ARTEMINYL (Cale Chem) Phenobarbital sod 1/2 gr, ephedrine sulf 1/2 gr, theophylline 2 gr, isopropylartereol HCl 1/6 gr / Tab Bot 100s

Use: Asthmatic paroxysms

ARTESENOL, ADR See: Levophed, Amp (Winthrop Labs)

ARTESINE. (Canfield) Pentylentetrazol 100 mg, nicotinic acid 50 mg / Tab or 5 cc of Elixir Tab Bot 100s; Liq Bot 1 pt

Use: Treatment of senile psychosis

ASTERINE (Canfield) Vit B 12, 50 mcg; B 1, 20 mg; B 2, 2 mg; B 6, 5 mg; ino

*Generic name recognized by NF, USP or A.M.A.

cinamide, 75 mg; choline chloride, 20 mg; pantheol, 5 mg; refined liver, 2 mcg; peptylenetetrazole, 10 mg / cc. 10 cc. multiple dose vials

Use: Geriatric syndrome

ARTERO-CYN. (Filet, Eaton) Pot. thiocyanate. Elixir 2 gr / 5 cc Bot 1 pt, 1 gal

Use: (1 dr 1 id) Hypertension therapy

ARTERODYL TT. (Columbus) Tolazoline HCl 75 mg / TT Tab Bot 100s

Use: Arteriosclerosis Peripheral vasodilator

ARTHO-B-12. (Columbus) Physostigmine salicylate 0.6 mg, hamatropia methylbromide 1.4 mg, Vit B 12 100 mcg / cc. Vial 10 cc

Use: Parasympathetic stimulant

ARTNO-E. (Columbus) Physostigmine salicylate 0.6 mg, hamatropia methylbromide 1.4 mg, alpha-tocopherol acetate 75 mg / cc Vial 10 cc

Use: 1 M (1 cc, daily), rheumatoid arthritis

ARTNO-STERONE. (Columbus) Desoxycorticosterone acetate 5 mg, Sod ascorbate 500 mg / 2 cc Amp 2 cc, Box 12s, 25s, 100s

Use: 1 M (2 cc daily), Addison's disease

ARTHOTABS. (Columbus) Acetylsalicylic acid 0.325 Gm, succinic acid 0.195 Gm, calcium succinate 0.130 Gm, ascorbic acid 30 mg / Tab Bot 72s

Use: (1 2 1 1 d) Arthritis

ARTHOTABS No. 2. (Columbus) Sod salicylate 0.215 Gm, succinic acid 0.108 Gm, calcium succinate 0.108 Gm, ascorbic acid 20 mg / Tab Bot 126s

Use: (1 2 1 1 d) Arthritis

ARTHRALGEN. (Whittier) Methochlorine chloride 1/4%, thymol 1%, menthol 10% methyl salicylate 15%, in a super-absorbable washable oint base Tube 1 oz, Jar 8 oz

Use: Topically, to increase blood flow and relieve pain.

ARTHRALGEN & ARTHRALGEN PR (Whittier) Calcylamide 250 mg, N-acetyl-p-aminophenol 250 mg, Vit C 25 mg / Tab PR has prednisone 1 mg added Bot 100s

Use: Antiarthritic

ARTHRITIS VACCINE. (Hallister Steir) Killed streptococci, staphylococci & B. xerosis Pk 9 5 cc

Use: 5 C (50 000 organisms weekly) or thrills

ARTHRITIS VACCINE. (Sherman) Microformans of Crowe 500 million, Streptococcus hemolytic, nonhemolytic, visidans 500 million, Pseudodiphtheroid bacillus 200 million / cc Vial 12.5 cc, 20 cc, 50 cc

Use: 5 C (10 000 organisms weekly), rheumatic conditions

ARTHRITIS VACCINES,

See: Arthritis Vaccine (Sherman) (Pitman-Moore) (Hollister-Stain)

Bacterial Intravenous Proteins (Cutter).

S.I.P. (Cutter)

Rheumatoid Arthritis Vaccines (Ledette).

Rheumatism Phylacogen (Parke, Davis).

Strepto Combined Vaccine (National).

Streptococcus Immunogen Arthritis (Parke, Davis).

Streptococcus Vaccine Arthratropic (Lilly).

Streptococcus Vaccine (Lilly).

ARTHROPAN. (Purdue Frederick) Choline salicylate 870 mg./tsp. Sol. S, 16 oz.

Use: Analgesic or antipyretic, in the treatment of rheumatoid arthritis.

ARTICON. (Walker) Hydrocarbons acetate 5 mg., salicylamide 250 mg., mephenesin 200 mg./Tab. Sol. 24s, 100s, 1000s.

Use Antiarthritic.

ARTISONE ACETATE, See: 21-Acetoxyprogesterone

ARZOL. (Arzol Chem. Co.) Silver nitrate applicators 75% pks, 100s

Use: Caustic

A-S PERCOMORPH LIVER OIL w/Other fish liver oils & vitamin (Abbott) Vit. A 60,000 u., D 8,500 u./Gm. Sol. w/dropper 10 cc. & 50 cc.

Use: (0.6 cc. daily) Vitamin A & D therapy.

A.S.A. w/Calcium carbonate, methylprednisolone. See: Medapria, Tab. (Upjohn)

A.S.A. (Lilly) (Aspirin) Acetylsalicylic acid. Essalts (5 & 10 gr.) Sol. 100s, 1000s; Pulvules (5 & 10 gr.) Sol. 100s, 1000s; Supp. (5 & 10 gr.) Pkg. 6s; Tab. (1 gr. pink & 5 gr.) Sol. 100s, 1000s.

A.S.A. COMPOUND. (Lilly) Acetylsalicylic acid

Use: Analgesic, antipyretic.

acid 3½ gr., acetophenetidin 2½ gr., caffeine ½ gr./Pulv. or Tab. Sol. 100s, 1000s of each.

Use: Analgesic, antipyretic.

A.S.A. COMPOUND W/Codaine. (Lilly) No. 1 w/codaine phosphate ½ gr.; No. 2 w/codaine phosphate ¼ gr.; No. 3 w/codaine phosphate ½ gr.; No. 4 w/codaine phosphate 1 gr.

Use: Analgesic, antipyretic.

ASAL. (Lemman) Ascorbic acid ¼ gr., salicylamide 5 gr./Tab. Sol. 100s.

Use: (1-2 i.i.d.) Analgesic.

ASALCINE. (Lemman) Ascorbic acid 15 mg., calchicine 1/250 gr., salicylamide 5 gr./Tab. Sol. 100s.

Use: (1-2 q. 4 h.) Analgesic.

ASARUM COMPOUND. (First Texas) Quinine sulfate 8 gr./fl. oz. Sol. 1 pt., 1 gal.

Use: (1-2 dr. q.i.d.) Auricular fibrillation.

* Generic name recognized by N.F., U.S.P. or A.M.A.

A.S.B. (Newman) Calcium carb., magnesium carb., bismuth subcarb., Sod. bicarb. & kaolin. Pow., Cap 3 oz.

Use: Antacid.

ASBAPHEN. (Lemman) Aspirin 3½ gr., acetophenetidin 2½ gr., caffeine ¼ gr., phenobarbital ¼ gr./Tab. Sol. 100s, 1000s.

Use: (1 q. 3 h.) Analgesic & sedative.

ASCADIN. (Drug Products) Aspirin 3½ gr., acetophenetidin 2½ gr., caffeine ½ gr./Pulvaid Sol. 1000s.

Use: (1 p.c.n.) Analgesic & antipyretic.

AS-CA-PHEN. (Ulmer) Aspirin 3½ gr., phenacetin 2½ gr., caffeine ½ gr./tab. or cap. Sol. 100s, 1000s.

Use: Analgesic

ASCO. (Sutcliffe & Case) Aspirin 3 gr., Daver's Powder ¼ gr., Acetophenetidin 1½ gr., Camphor Monobromated ¼ gr., Caffeine ½ gr., Atropine Sulfate 1/500 gr./Tab. or cap. Sol. 100s, 1000s.

ASCORAMIDE. (Warren-Teed) Nicotinamide 50 mg., ascorbic acid 50 mg., lactose q.s./Tab. Sol. 100s, 1000s.

Use: Vincent's infection, gingivitis & bleeding gums.

ASCORB. (Harper) Vit. C 100 mg./Tab. Sol. 100s.

Use: Vitamin C deficiency.

ASCORBACINE. (Tastagar) Procaine HCl 250 mg., ascorbic acid 150 mg./cap. Sol. 100s.

Use: (1 q. 3-4 h.) pruritus.

ASCORBAJEN Jr. (Jenkins Labs.) Ascorbic acid 30 mg./tab. Sol. 100s, 1000s.

Use: Dietary adjunct.

* **ASCORBIC ACID,** U.S.P., N.N.R. 1954, A.D.R.; Injection, U.S.P.; Tablets, U.S.P.; 3-Oxo-L-gulofuranolactone (enal form) (Anti-scorbic vitamin—vitamin C.) Injections (Various Mfr.). Amp. (100 mg./cc.) 1 cc., 2 cc., 5 cc., (200 mg./cc.) 5 cc., (500 mg.) 2 cc., 5 cc., (1000 mg.) 5 cc., 10 cc. Vial—(100 mg./cc.) 10 cc., 30 cc., (250 mg./cc.) 10 cc.

Tab. (25 mg., 50 mg., 100 mg., 250 mg., 500 mg.) (Various Mfr.)

See: Allercarb, Tab. (Plessner).

Ascorbajen, Jr., Tab. (Jenkins Labs)

Contaxin, Tab. (Winthrop Labs.)

Cebiasse Dulcet, Tab. (Abbott)

Cecon (Abbott)

Ce-Mi-Lin, Lozenge (E. S. Miller)

Cevalin, Tab. (Lilly)

Cevex (Walker Tab 1)

C-Quin, Tab. (E. S. Miller)

Lemasorb, Tab. (Ascher).

Vicia, Tab. (Brewer).

Cap (Various Mfr) 25 mg, 100 mg, 250 mg 500 mg

Liq See Cecan (Abbott)

Cevalin Drops (Lilly)

Cevex (Walker)

Ce Vi Sol (Mead Johnson)

Liqui Cee Liq (U S Standard)

Pow. See: Cebiane (Merck Sharp & Dohme)

ASCORBIC ACID INJ, U S P

See Ascorbin amp (Lakeside)

Cantoxin, amp (Winthrop Lab)

Genalate, amp (Abbott)

Cevalin amp (Lilly)

Natracarb Injectable (U S Vit)

Scorbate Injection (Kirk)

Vicin, amp (Brewer)

ASCORBIC ACID SALTS

See: Bismuth Ascorbate

Calcium Ascorbate

Ferrous Ascorbate

Sodium Ascorbate N N O

ASCORBIN, (Lakeside) Sod ascorbate Amp 2 cc (500 mg) Box 25s

Use IM, SC, IV vitamin C therapy

ASCORBIN R (Lakeside) Vit B 1 50 mg B 2 10 mg, B 6 50 mg, C 1000 mg, niacin 250 mg, pantothenic acid 50 mg / 10 cc Vial 10 cc Box 25s, 100s

Use Vit B complex with Vit C

ASCORBO THIAMINE COMPOUND (Meyer) Vit B 1 10 mg, B 2 5 mg, B 6 5 mg niacinamide 100 mg cal pantothenate 10 mg, thaline chloride 5 mg, para amino benzoic acid 5 mg, vit C 150 mg, E 3 I U, cal glycerophosphate 120 mg / Cap Bot 100s

Use: Metabolic disturbances

ASCOR-RUTIN (American Pharm) Two strengths: Ascorbic acid 100 mg, rutin 20 mg / Tab; ascorbic acid 300 mg, rutin 60 mg / Tab Bot 100s, 500s

Use: Vitamin C and bioflavonoid therapy

ASCORUTIN (Lemman) Ascorbic acid 50 mg, rutin 50 mg / Tab Bot 100s, 1000s

ASCORUTIN, (U S 3rd Prads) Rutin & ascorbic acid Tab (20 mg, 100 mg) Bot 100s, 1000s; (60 mg, 300 mg) Bot 100s, 500s, 1000s

Use: Increase capillary fragility

ASCORUTIN-BIO (Arnar Stone) Ascorbic acid 100 mg, rutin 25 mg, heiperidie 100 mg / Tab Bot 100s

Use: Abnormal capillary fragility and permeability

ASCRIPTIN (Rorer) Aspirin 300 mg, mag aluminum hydroxide 150 mg / tab Bot 100s, 500s, also available w/ Cadeine 1/2 gr, Bot 50s

Use: Orally (1 2 q 3-4 h) Analgesic

ASDRIN (Cale) Isopropylarterenal HCl 10

mg / Tab Bot 100s, 1000s

Use: Bronchodilator

ASEPTICROME. See: Merbromin (Various Mfr)

ASEPTOGILL (Mastengill) Thymal iodide bismuth oxyiodide, boric acid, zinc oxide & dermalol Pow 1 oz

Use: Antiseptic & astringent

ASEPTUROL, (Wallau) Methenamine, lithium benzoate Sod benzoate & piperazine Bot 4 oz

Use Orally (2 dr q 3 h), urinary antiseptic

A S F (Anti Stress Formula) (Roerig) Vit B 1 10 mg B 2 10 mg, niacinamide 100 mg, calcium pantothenate 20 mg, Vit B 6 2 mg, folic acid 1.5 mg, Vit B 12 4 mcg C 300 mg, K 2 mg / Cap Bot 30s, 100s

Use (1 2 daily) Multivitamin supplement

ASIAN FLU VACCINE, (Merck Sharp & Dohme) Each cc of vaccine contains 200 CCA v Vial 10 cc

Use Prophylaxis against influenza

ASTATIC TABLETS, (Horton & Converse) Arsenic Itraxide 1/9 gr, black pepper 1 gr / Tab Bot 100s

Use Tonic

ASTUM (Drug Products) Carbolic acid 1% aluminum acetate, ichthammal, zinc oxide aromatic oils in a petrolatum stearin base Tube 1 1/2 oz Jar 1 lb 5 lb

Use Astringent & soothing dressing

ASMACOL (Vale) Butobarbital 15 mg, am inophylline 180 mg, phenylpropanolamine HCl 25 mg chlorpheniramine maleate 2 mg, aluminum hydroxide 60 mg, mag nesium trisilicate 60 mg / Tab Bot 100s

Use Bronchospasm due to asthma

ASMAFIELD ELIXIR, (Canfield) Theophylline 130 mg, amobarbital 10 mg, ephedrine HCl 20 mg / 15 cc Bot Pts, Gals

Use Bronchial asthma and hay fever

ASMASAN, (Recces) Aminophyllin 125 mg Mephobarbital 25 mg, Ephedrine HCl 25 mg, tri histin 25 mg / Tab Bot 100s

Use: Allergies

ASMATONE, (National Research) Acetyl strobaline base Cap (5 1/2 gr) Bot 15s 30s, 100s, 500s, 1000s

Use: (1 q 3 h) Antihistamine

ASMI-NIST, (Cale) Theobromine 150 mg, ephedrine sulf 25 mg, phenylpyramine HCl 10 mg / Tab Bot 100s, 500s, 1000s

Use: Bronchial asthma

ASMINYL, (Cale) Ephedrine sulfate 1/2 gr, pheoobarbital Sod 1/2 gr & theophylline 2 gr / Tab Bot 100s liq Bot 4 oz

Use: (1 q 4 h) Vasconstrictor with sedative

ASOPHYLLIN, (Crystal) Outer coat: Isoproterenol 10 mg; Inner coat: racephedrin

HCl 16 mg., phenobarbital 8 mg., neohyline 130 mg./Tab. Bot. 100s.

Use: Treatment of asthma.

ASPACAM. (Smith-Dorsey) Aspirin 3½ gr., acetophenetidin 2½ gr., w/camphor monobromate, gelsemium, strychnine sulfate, cloin, podophyllin & atropine sulfate/Tab. (Also without laxative.) Bot. 100s, 1000s.

Use: Antipyretic & analgesic.

ASPARTIC ACID SALTS. See: Spartase, Tab. (Wyeth)

ASPARTOCIN. (Lederle) An antibiotic from *Strept. griseus* var. *spiralis*.

Use: Antibiotic.

ASPERGILLUS ORYZAE ENZYME. Diastase See: Taka-Diastase (Parke, Davis).

***ASPERKINASE.** (Smith Kline and French) Proteolytic enzyme mixture derived from *aspergillus oryzae*.

ASPHAC-B. (Central) Aspirin 4 gr., acetophenetidin 2 gr., dried aluminum hydroxide gel 2 gr., caffeine anhydrous ¼ gr., gelsemium tincture 3 min./Tab. Bot. 100s, 500s, 1000s.

Use: Analgesic

ASPHAC-G. (Central) Aspirin 4 gr., acetophenetidin 2 gr., caffeine ¼ gr., gelsemium fr. 3 min./Tab. Also w/caffeine ¼ & ½ gr. Bot. 100s, 500s, also available in Half Strength, & Infant Palatabl

User [1 q. 4 h.] Analgesic.

ASPHAMAL. (Central) Salicylamide 4 gr., acetophenetidin 2 gr., racephedrine HCl 1/12 gr., pyranisamine maleate 1/5 gr./Tab. Bot. 100s, 500s, 1000s

User Camman cold.

ASPHAMAL-D. (Central) Salicylamide 4 gr., acetophenetidin 2 gr., racephedrine HCl 5 mg., pyrilamine maleate 12 mg., dover's powder 24 mg./Tab. Bot. 100s, 500s, 1000s

Use: Analgesic with antihistaminic agent.

ASPHEDER. (Central) d-Desoxyephedrine HCl 1/24 gr., butobarbital ¼ gr., salicylamide 4 gr. & acetophenetidin 2 gr./Tab. Bot. 100s, 500s, 1000s

Use: Analgesic & antidepressant.

ASPIBARB. (Lemman) Aspirin 5 gr., phenobarbital ¼ gr./Tab. Also in half strength for children. Bot. 100s, 1000s.

Use: [1 q. 4 h.] Analgesic-sedative.

ASPIDRINE. (Notick) Aspidosperma fr. 240 min., strantium iodide 25 gr., hydriodic acid 10% 16 min., ephedrine HCl ½ gr., helianthus syr. 240 min./tab. Bot. 1000s.

Use: Respiratory stimulant.

ASPIDYNE. (Lemman) Aspirin 3½ gr., acetophenetidin 2½ gr., caffeine ½ gr./Tab. Also for children: Aspirin 1 gr., acetophenetidin ½ gr., caffeine ¼ gr./Tab. (W/

* Generic name recognized by N.F., U.S.P. or A.M.A.

Caffeine phosphate ¼ gr./Tab.) Bot. 100s, 1000s.

Use: [I.I.d.] Analgesic.

ASPIDYNE COMP. (Lemman) Dover's powder ½ gr., acetylsalicylic acid 2 gr., acetophenetidin 1½ gr., caffeine ½ gr., atropine sulfate 1/500 gr., camphor monobromate ½ gr./Tab. or Cap. Bot. 1000s.

Use: Analgesic-antipyretic.

ASPIFEDRINE. (Lemman) Phenobarbital ½ gr., Hyoscyamus ext. ½ gr., salicylamide 3 gr., acetophenetidin 2 gr., di-desoxyephedrine HCl 1/20 gr./Tab. Bot. 1000s.

Use: Analgesic-antipyretic.

ASPIRFENS. (Gold Leaf) Aspirin 5 gr., phenobarbital ¼ gr./Tab. Bot. 100s, 1000s.

Use: Analgesic-sedative.

ASPIRIN, U.S.P., N.N.R. 1947, A.D.R. (Various Mfr.) (A.S.A., Acetaphen, Acetol, Acetosal, Acetosafin, Acetylcy, Acetylin, Acetylsal, Emprin, Saletin.) Acetylsalicylic acid

See: Aspidrops, Flavored. (C. D. Smith)

Aspirin E.C.

Aspirin Jr., Tab. (Jenkins Labs.)

Crystal, gran. (Armour)

Pirsel. (Testagar)

Spiro-Dine, liq. (Warren-Teed)

Aspirin-Barbiturate Combinations on pp. 82 and 83.

ASPIRIN DULCET. (Abbott) Aluminum acetylsalicylate 1½ gr./dulcet. Bot. 50s, 100s.

Use: Analgesic.

ASPIRIN ENTERIC COATED.

See: A.S.A. (Tilly)

Asteric, Tab. (Brewer)

Ecotrin, Tab. (Smith, Kline & French)

ASPIRIN SACCHARINATED W/Benzoesulfimide (Saccharin). See: Theryl, Tab. (Church)

ASPIRIN SALTS.

See: Aluminum Acetylsalicylate

Calcium Acetylsalicylate.

ASPIRIN-SECOCARBITAL, Suppratee. (Webster).

No. 1 (Aspirin 1 gr., secobarbital Sod ½ gr.)

No. 2 (Aspirin 3 gr., secobarbital Sod ¾ gr.)

No. 3 (Aspirin 5 gr., secobarbital Sod 1 gr.)

No. 4 (Aspirin 10 gr., secobarbital 1½ gr.) Jar 12s.

Use: Orally; analgesic, antipyretic.

ASPIRIN Jr. (Jenkins Labs.) Aspirin 2½ gr./tab. Bot. 100s, 1000s.

Use: Analgesic.

ASPIROCAL. (McNeil) Butisol ½ gr., aspirin 4 gr., calcium gluconate 2 gr./Tab. Bot. 100s, 1000s.

Use: [1 q. 3 h.] Analgesic-sedative.

| Aspirin | | Others | Manufacturer | Brand |
|---------|--|--------|----------------|----------------------------|
| 5 gr | Amytal ¾ gr | | Lilly | Amytal w/Acetylsol Cap |
| 3½ gr. | Barbital 1½ gr., acetophenetidin 2 gr | | Warner | Peralga Tab |
| 4 gr | Butobarbital ¼ gr., acetophenetidin 2 gr, d-desoxyephedrine HCl 1/24 gr | | Central | Asphedex Tab |
| 2.5 gr | Butobarbital 0.5 gr., salicylamide 2.5 gr., acetaminophen 2.5 gr., d-amphetamine sulf 5 mg | | Stuart | Stuart Pralar, Tab |
| 2.5 gr | Butobarbital 0.5 gr., salicylamide 2.5 gr., acetaminophen 2.5 gr., d-amphetamine sulf 5 mg., hyoscyamine sulf 0.1384 mg atropine sulf 0.0258 mg., hyoscyne HBr 0.0066 mg | | Stuart | Stuart Probar B Tab |
| 4 gr | Bulisol ¼ gr. Cal gluconate 2 gr | | McNeil | Asp rocol, Tab |
| 5 gr | Cyclopal ¾ gr | | Upjohn | Cyclopal & Aspirin, Tab |
| 3½ gr | Diallylbarbital ¾ gr., acetophenetidin 2 gr | | Ulmer | Alpine Tab |
| 3½ gr | Diallylbarbital ¼ gr., acetophenetidin 2½ gr., caffeine ¼ gr gelatinum tr 1 min | | Buffington | Dorodol Tab |
| 224 mg | Hexobarbital 100 mg., dihydrohydroxycodeinone HCl 4.5 mg dihydrohydroxycodeinone terephthalate 0.38 mg, homatropine terephthalate 0.38 mg., phenocetin 160 mg caffeine 32 mg | | Endo | Perobarb, Cop |
| 5 gr | Ipral 1 gr | | Squibb | Ipral Aspirin Tab |
| 5 gr | Mephobarbital 16.2 mg d amphetamine HCl 2.5 mg., phenocetin 162 mg | | Rowell | Mapho Oex Tab |
| 300 mg | Nembutal 30 mg | | Abbott | Nembu Gestic, Cop |
| 5 gr | Pentobarbital ½ gr | | Pyrrhress | Syrinin, Tab |
| 5 gr | Pentobarbital ½ gr | | Abbott | Nembutal & Aspirin, Cop |
| 5 gr | Pentobarbital ½ gr | | Massengill | Nipirin, Cop |
| 5 gr | Pentobarbital Sod ¾ gr & 1½ gr | | Novacol | Pembrin, Cop |
| 3½ gr | Pentobarbital Sod ¼ gr., d-amphetamine sulfate 2.5 mg., acetophenetidin 2½ gr | | Arnor Stone | Polygestic, Tab. |
| 5 gr | Phenobarbital ¼ gr | | Lemman | Aspiorb, Tab |
| 5 gr | Phenobarbital ¼ gr | | Gold Leaf | Aspirfens, Tab |
| 5 gr | Phenobarbital ¼ gr | | Winthrop | Lumalgin Tab |
| 2 gr | Phenobarbital ½ gr. | | Jenkins Labs. | Solibar Jr., Tab |
| 5 gr | Phenobarbital ½ gr | | Sutliff & Case | Aspirin Phenobarbital, Tab |
| 4 gr | Phenobarbital ¼ gr acetophenetidin 2 gr | | Warner-Tand | Aspiral Tab |

Aspirin-Barbiturate Combinations

| Aspirin | Others | Manufacturer | Brand |
|---------|---|--------------------|--|
| 4 gr. | Phenobarbital ¼ gr., acetophenetidin 2 gr. | Harvey | Harvadol, Tab. |
| 3½ gr. | Phenobarbital ¼ gr., acetophenetidin 2½ gr. | Burroughs Wellcome | Empiral, Tab. |
| 3½ gr. | Phenobarbital ¼ gr., acetophenetidin 2½ gr., caffeine ¼ gr. | Lemman | Atbarphen, Tab. |
| 5 gr. | Phenobarbital ¼ gr., di-amphetamine phosphate 3 mg. | Clark | Pen-Phelamine, Tab. |
| 325 mg. | Phenobarbital 8 mg., p-aminobenzoate Sod. 15 mg., calcichine sulfate 0.1 mg., Vit. B-1 25 mg. | Wally Pharm. Co. | Dalarol, Tab. |
| 130 mg. | Phenobarbital 26 mg. - ascorbic acid 100 mg., ephedrine HCl 26 mg. | Arlington | Artcapt, Cap. |
| 180 mg. | Phenobarbital 15 mg., belladonna alkaloids 0.025 mg., acetophenetidin 120 mg. | Rorer | Paodon, Cap. |
| 5 gr. | Phenobarbital ¼ gr., belladonna ext. ¼ gr., caffeine ¼ gr. | Elder | Anodyne, Tab. |
| 5 gr. | Phenobarbital ½ gr.,gelsemium ext. 3/40 gr., caffeine ¼ gr. | Cole | Phac, Tab. |
| 5 gr. | Phenobarbital ½ gr., hyoscyamus ext. 1/6 gr. | Mastengill | Andasix, Cap. |
| 0.5 gr. | Phenobarbital ¼ gr., hyoscyamus ext. ¼ gr. | Rorer | Lemaspirin |
| 162 mg. | Phenobarbital 16.2 mg., acetophenetidin 194 mg., hyoscyamine sulf. 0.031 mg., prophepyridamine maleate 12.5 mg., phenylephrine HCl 10 mg. | Robins | w/Hyoscyamus, Tab. Phenaphen Plus, Tab. |
| 3 gr. | Phenobarbital ¼ gr., hyoscyamus ext. ½ gr., acetophenetidin 2 gr. | Frasst | Acetaphen, Cap. |
| 2½ gr. | Phenobarbital ¼ gr., hyoscyamus ext. 1/6 gr., acetophenetidin 1½ gr. | Mastengill | Analdyne, Cap. |
| 2½ gr. | Phenobarbital ¼ gr., hyoscyamus ext. ½ gr., acetophenetidin 2½ gr. | Mastell | Haiamal, Tab. |
| 2½ gr. | Phenobarbital ¼ gr., hyoscyamus alkaloids 0.025 mg., acetophenetidin 2½ gr. | Vanpelt & Brown | Hycetin, Tab. |
| 5 gr. | Phenobarbital ½ gr., Sod. & Pot. bromides each ¼ gr., caffeine ¼ gr. | Massengill | Abacol, Tab. |
| 3 gr. | Sandapal ¼ gr., caffeine ½ gr., phenacetin 2 gr. | Sandaz | Florinal, Tab. |
| 1 gr. | Secobarbital | Webster | Aspirin-Secobarbital |
| 3 gr. | Secobarbital ¾ gr. | | Supprethcs, No. 1 |
| 5 gr. | Secobarbital ¾ gr. | | No. 2 |
| 10 gr. | Secobarbital ¾ gr. | | No. 3 |
| 195 mg. | Secobarbital 50 mg., phenobarbital 50 mg., Vit. B-1 5 mg. | Patch | No. 4 Ambral, Tab. |

ASPIROFEINE (McNeil) Aspirin 3½ gr., caffeine ½ gr., acetophenetidin 2½ gr./Cap Bot 100s, 1000s Also w/Daversa Pow ¼ gr./Tab

Use Analgesic-antipyretic

ASPIROTABS (Brewer) Aspirin 3½ gr., phenacetin 2½ gr., caffeine ½ gr./Tab Bot 100s, 1000s. Also w/cocaine ¼ & ½ gr

Use Analgesic.

ASPIROPS (Smith Miller & Patch) Aspirin 3 gr. in a homogenous mixture of aromatic flavors Bot 100s, 500s

Use: (1-2 p.m.) Analgesic.

ASPIVITE (Marck Sharp & Dohme) Aspirin 5 gr., menadione 0.25 mg., ascorbic acid 35 mg./Cap Bot 100s, 1000s

Use: (1 q 3-4 h) Analgesic.

ASPODYNE (Blue Line) Aspirin 3½ gr., acetophenetidin 2½ gr., caffeine ½ gr./Tab Bot 100s, 1000s.

W/Cocaine phosphate ¼ gr Bot 100s, 1000s.

Use (1 p.m.) Analgesic-antipyretic.

ASPOGIN, NND 1939 Dihydroxyalumi-
num acetate

ASPRID-C, (Maney) Prednisone 0.75 mg., salicylamide 5 gr., ascorbic acid 50 mg./Tab Bot 100s, 1000s.

Use: Treatment of mild rheumatic disorders

ASPRO See: Acetylsalicylic acid

A-B-R, (Morton & Converse) Vit C 300 mg., pyridox 40 mg., Vit B-1 10 mg., riboflavin 10 mg., niacinamide 100 mg., calcium pantothenate 40 mg., folic acid 2.5 mg., Vit B-12 4 mg./2 Cap Bot 60s, 250s, 1000s.

Use Therapeutic vitamin therapy

ASTERIC TABLETS, (Brewer) Acetylsalicylic acid 5 or 10 gr./E.C. Tab Bot 100s, 1000s, 5000s, 10,000s

Use Analgesic

ASTERIC COMPOUND, (Brewer) Aspirin 300 mg., ascorbic acid 15 mg., calcium pantothenate 1 mg./Tab. Bot. 100s

Use Treatment of rheumatoid arthritis

ASTEROL, NND 1962 (Roche Lab) 2-D-methylamino-5-(beta-ethylaminoethoxy) benzothiazole dihydrochloride Di-amthazole dihydrochloride Oint 5% asterol in carbomer base) 3 oz., tubes and 3 lb. jars.

Pow 15% asterol in fine, dry powder) 1½ oz. shaker top container. Tinct 15% asterol in 70% isopropyl alcohol) 2 oz & 16 oz. bottles

Use Antirheumatoid drug

ASTHMACAINE (Parke-Davis Labs) Products from hydrolysis of 1099 mg cocaine (¼

gr) cocaine—0.002 mg., atropine sulfate 1.79 mg., sod nitrite 38.41 mg/cc Also available in 1½ strength Bot 3D & 60 cc

Use Relief from asthmatic distress

ASTRAFER, NND 1962 (Astra) Dextriferron (2%) 5 cc amp Box 1Ds 2D mg/cc Fe+3

Use IV Iron deficiency, late pregnancy

ASTRINGENT COMPOUND, (Upjohn) Zinc sulfate, zinc chloride, Pot, alum & boric acid Pow, Pkg 3 oz

Use: Vaginal cleanser

AT-7 Neechloraphene in Dial soap (Aurum)

A.T. 10, See Dihydratachysterol

ATABEE 10 (Finley) Vit C, 500 mg; thio mine mononitrate, 15 mg; riboflavin, 10 mg; pyridoxine HCl, 2 mg; nicotinamide 50 mg; calcium pantothenate, 10 mg/Cap Bot 100s

Use: Water soluble B complex and ascorbic acid vitamin deficiencies

ATARGINE DIHYDROCHLORIDE, NND 1961 (Winthrop Labs) Quinacrine hydrochloride Tab (0.1 Gm) 100s, 500s

Use: Malaria of all types

W/Asalen phosphate, Plaquenil sulfate See Triquin, Tab (Winthrop)

ATARAXIC AGENTS,

See: Acepromazine

Alisoylone

Atarax (Roerig)

Atrovet (Ayerst)

Azocyclanot HCl

Benactyzine HCl

Bucylzine

Clopidiamine

Caisoprodol

Chlorpromazine

Camazine (Smith, Kline & French)

Dartal (Searle)

Depral (Wallace)

Deserpidine

Dimethylone (National Drug)

Dornwal, Tab (Maltbie)

Ectylurea

Ethchlorvynol

Equanil (Wyeth)

Franquel (Merrell)

Fluphenazine

Harmony (Abbott)

Hydroxyzine

Isopropylid Phos

Lebrum, Caps (Roche)

Levanil, Tab (Squibb)

Isomeprobamate

Marsilid Phos (Roche Lab)

Mellonal, Tab (Sandoz)

Mepazine

Meprobamate

Meprotab (Wallace)

- Meprospon (Wallace)
 Meraltran (Merrell)
 Methylphenidate.
 Milltown (Wallace)
 Moderil (Pfizer)
 Nostyn (Ames)
 Ozonamide.
 Pacotal (Warner-Chilcott)
 Permlit, Tab. (White)
 Perphenazine
 Phenoglycodol
 Phenergon (Wyeth)
 Phenyltoloxamine
 Phobex (Lloyd, Dabney)
 Pipradrol.
 Placidyl (Abbott)
 PRN (Bristol)
 Prochlorperazine
 Protizin, Tab. (Squibb)
 Promazine
 Promethazine.
 Promazolone
 Qulacin, Tab. (Merrell)
 Rauwolfia serpentina
 Rescinnamine (Roemer)
 Reserpine
 Ritalin (Ciba)
 Soft-Tran-Tab. (Stuart)
 Sparine (Wyeth)
 Stelazine, Tab. (Smith, Kline & French)
 Suvilil (Merck Sharp & Dahme)
 Suvren (Ayerst)
 Sycotrol, Tab. (Reed & Cornrick)
 Tentone, Tab. (Lederle)
 Thiopropazate
 Thioridazine
 Thorazine (Smith, Kline & French)
 Trancln, Tab. (Schering)
 Trancopol, Cap. (Winthrop Labs.)
 Trifluorpromazine.
 Trifluoperazine
 Trilatan (Schering)
 Uttron (Lilly)
 Vesprin
 Vistoril (Pfizer)
- ATARAX, N N D, 1962.** (Roerig) Hydroxyzine HCl. 1-p-Chlorobenzhydryl-4,2-(2-hydroxyethoxy)ethylthiethylamine HCl. Tab. (10 mg, 25 mg, 50 mg, 100 mg) Sol. 100s. Syr. (10 mg./5 cc.) Sol. 1 pt. Amp. (50 mg./cc) 2 cc., 5s. Vial (25 mg./cc) 10 cc
 Use: Treatment of tension.
 W/Penta-erythrityltetranitrate. See: Cortraz, Tab. (Roerig)
- W/d-Amphetamine, 9 vitamins; 1) mineralz/capsule. See: Amplus Improved. (Roerig)
- ATARAXOID 2.5.** (Pfizer) Prednisolone 2.5 mg. hydroxyzine hydrochloride 10 mg./Tab. Sol. 100s.
- S.O.** (Pfizer) Prednisolone 5 mg., hydroxyzine hydrochloride 10 mg./Tab. Bot. 30s, 100s
 Use: To control anxiety in arthritis, asthma and allergic dermatoses
- ATARVET.** (Ayerst) Acepromazine.
- ATAV.** (Cole) Vit. A 50,000 u./Cap. Bot 100s
 Use: Vitamin A deficiencies.
- ATHA-SPRAY** (Youngs Rubber Co) N-Tri-chloromethylmercapto-4-cyclohexene-1,2-dicarboximide and benzalkonium chloride. 3 oz. aerosol con.
 Use: Fungicidal medication.
- ATHEMOL.** (Meyer) Theobromine magnesium oleate 200 mg./Tab. Bot. 100s, 1000s.
 Use: Arterio sclerosis.
- ATHEROL.** (U. S. Standard Products) Safflower Oil 1.15 Gm., Vit. B-6 3 mg./Cap. Bot. 100s.
 Use: Reduction of blood cholesterol.
- ATHEROXIN.** (Gray Pharm) Refined corn oil (Maydol) 15 Gm. with pyridoxine HCl 3.33 mg./fl. oz. Emulsion. Bot. 24 oz.
 Use: Orally, to decrease cholesterol blood-level
- ATOPNAN.** (Warner-Chilcott) (Cinchophen) Phenytycinchoninic acid. Pow., Bot. 30 Gm.; Tab. (gr. 5) Box 20s, (gr. 7.5) Box 20s; Bot. 50s, 100s, 500s.
 Use: (5 gr. to 1 Gm.) Rheumatism, gout & neuralgia.
- ATOPHIN.** (Blochemical) Vit. D 50,000 u., bile salts 32 mg./Cap. Bot. 100s, 1000s.
 Use: Vitamin D therapy.
- ATP.** See: Adenosine Triphosphate (Various Mfr.)
- ATRALOSE.** (Warner Chilcott) Homatropine methylbromide 1 mg., phenobarbital 8 mg & cellohyf (methylcellulose) 500 mg./Tab. Bot. 100s, 500s
 Use: (1-2 t.l.d.) Antispasmodic with bulk laxative.
- ATRATAN.** (Neisler) Atropine Iannate 1 mg./Tab. Bot. 50s.
 Use: Relief from ureteral colic symptoms.
- ATRAVET.** (Ayerst) Acepromazine.
 Use: Tranquillizer.
- ATRINAL.** (Rache Lab) Atropine sulfate. W/Papaverine HCl, pantopon. See: Spasmalgin, Tab., liq. (Rache)
- ATROCAL.** (Drug Products) Atropine sulfate 1/500 gr., calcium carb. ppt. 10 gr. & aromatics/Pulroid. Bot. 100s, 1000s.
 Use: (1-4 daily) Antispasmodic-antacid.
- ATROCHOLIN.** (Blue Line) Dehydrocholic acid 2 gr., homatropine methylbromide 1/100 gr./tab. Bot. 100s, 1000s.
 Use: (1 t.l.d. p.c.) Antispasmodic & hydrocholeretic.
- ATRODYNE.** (Lemman) Atropine sulfate 0.012

mg, acetylsalicylic acid $\frac{3}{4}$ gr acetophenetidin $\frac{1}{2}$ gr, caffeine $\frac{1}{15}$ gr, menthol $\frac{1}{60}$ gr/Tab Bot 1000s

Use Analgesic antipyretic

ATROLOSE. (Dumas Wilson) Atropine sulfate $\frac{1}{150}$ gr, phenobarbital $\frac{1}{4}$ gr, yeast concentrate 1 gr, Sod carboxymethylcellulose 4 gr/Cop Bot 100s, 1000s

Use: (1 2 t i d., m p) Antispasmodic sedative

ATROMAG (Lemman) Phenobarbital Sod $\frac{1}{8}$ gr, magnesium trisilicate 7 gr, Sod bicarb $2\frac{1}{2}$ gr, hyoscyamus ext $\frac{1}{16}$ gr, atropine sulfate $\frac{1}{1000}$ gr/Tab Bot 1000s

Use Antacid-antispasmodic

ATROPAL. (Irwin, Neisler) Atropine sulf $\frac{1}{500}$ gr phenobarbital $\frac{1}{8}$ gr, mag trisilicate $2\frac{1}{2}$ gr, aluminum hydroxide $2\frac{1}{2}$ gr./Tab Bot 100s, 300s, 1000s

Use: Hyperacidity, nervous dyspepsia

ATROPHYSINE. (Lonnert) Physostigmine sulfate 0.6 mg, atropine sulf. 0.6 mg/cc Vials 10 & 30 cc

Use: I.M., antispasmodic for skeletal muscle

***ATROPINE, N F** (dl Hyoscyamine) dl Tropyl Itrate

W/Phenol Nutgall & benzocaine See: Carbolic Acid, Nutgall & Atropine Compound (Upjohn).

Atropine-Hyoscyamine-Hyoscyamine Combinations (See also Billedonna Products) on pp 87 to 88.

ATROPINE METHYLNITRATE. (Various Mfr) dl Hyoscyamine methylnate

See: Ekamine, Tab (Lloyd)
Eumydrin Tab (Wintrop Labs)
Horvatrate, Tab (G F Harvey)
Metante Tab (Drug Frods)
Metropne Elis & Tab (Straßenburgh).
Tidum Tab (Amco Drug)

W/Aluminum aminoacetate See: Koloyd Tab. (Lloyd)

W/Aluminum hydroxide, pectin osmakolin, beta lactose
See: Alopectate w/metropine Tab (Straßenburgh)

Metropect n, Tab (Straßenburgh)

W/Amphetamine phosphate racemic & amphetamine phosphate & Sod carboxymethylcellulose See: R phetacet, Tab (Straßenburgh)

W/dl Amphetamine phosphate epine & acetophenetide See: Espal Tab (Lloyd).

W/A.P.C. See Metrocaps Cop (Straßenburgh)

W/Barbital methocapamine nitrate See Icolate Tab (Centra)
W/Duodenum pow, aluminum hydroxide, a Generic name recognized by N.F., U.S.P. or A.M.A.

mag trisilicate See Metrocin, Tab (Straßenburgh)

W/Mephobarbital, hametropine methylbromide, & hyoscyne HBr

See Spascaps, T D Tab (Nysca Labs)

Trascine Unicettes (Hiss Pharm)

W/Oxidized bile acids, ox and hog bile ext, phenobarbital See G B S, Tab (G F Harvey)

W/Phenobarbital

See Ekamine w/Phenobarbital, Tab (Lloyd).
Horvatrate w/Phenobarbital, Tab (G F Harvey)

Metante w/Phenobarbital, Tab (Drug Frods)

W/Phenobarbital & dihydroxy aluminum aminoacetate See Horvatrate A, Tab (G F Harvey)

W/Phenylazo diamine pyridine HCl, sulface amide See Tridium, Tab (Amco)

W/Racemic amphetamine phosphate, dextro-amphetamine & Sod carboxymethylcellulose See Biphetacet, Tab (Straßenburgh).

W/Sulfacetamide, phenylazo diamine pyridine HCl See Cytizem, Tab (Zemmer)

ATROPINE-N-OXIDE HCl. See Atropine Oxide HCl

***ATROPINE OXIDE HYDROCHLORIDE** (Xitrium) atropine N oxide hydrochloride

See Xro genatropine HCl (Xitrium)

***ATROPINE SULFATE, USP, ADR Inj**

USP Tablets USP (Various Mfr) Tab—Hypodermic, $\frac{1}{200}$, $\frac{1}{500}$, $\frac{1}{100}$, $\frac{1}{50}$ gr Orally, $\frac{1}{500}$ $\frac{1}{250}$, $\frac{1}{200}$, $\frac{1}{150}$ $\frac{1}{100}$ gr

See Atropisal (Iso Sol)

Parasympatholytic & antispasmodic agent

Use: Mydriasis

Atropine Sulfate Combinations on pp 90 and 91.

Atropine Sulfate with Phenobarbital on p 92.

***ATROPINE TANNATE.** See: Atrotan, Tab. (Nasler)

ATROPINE TARTRATE. (Squibb) Tin tube (2 mg) solution

Use: I.V., I.M., combats the paralyzing action of nerve gas

ATROPISOL. (Crookes Barnes) Atropine sulf $\frac{1}{4}$ to 4% buffered soln drop bot & disp-perlets

Use: Mydriasis

ATRO-SALCO. (Pitman Moore) Dover's powder $\frac{1}{8}$ gr, atropine sulfate $\frac{1}{1000}$ gr, acetophenetidin $2\frac{1}{2}$ gr, aspirin $3\frac{1}{2}$ gr, gallesium tr 2 min, camphor monobromated $\frac{1}{4}$ gr/Cop Bot 100s, 1000s

Use: (1 t i d.) Analgesic and reduces respiratory secretions

Atropine-Hyoscine-Hyoscyamine Combinations (See also Belladonna Products)

| Atropine sulf. | Hyoscine HBr. | Hyoscyamine HBr. | Others | Manufacturer | Brand |
|-------------------------------------|---------------------------------------|--|---|--|---|
| 0.06 mg. 0.0258 mg. | 0.035 mg. 0.0086 mg. | 0.305 mg. sulf. 0.1384 mg. | Acetaminophen 2.5 gr., aspirin 2.5 gr., sulfi- cylamide 2.5 gr., bu- tobarbital 0.5 gr., di- amphetamine sulf. 5 mg. | Smith, Kline & French Stuart | Prydon, Spansule Stuart Prelor-B, Tab. |
| 0.02 mg. 0.0194 mg. | 0.006 mg. 0.0065 mg. | sulf. 0.1 mg. sulf. 0.1037 mg. | Acetylcarbamol Butobarbital Sod. 16.2 mg./dr | Standard Pharm. Zemmer | Disperdon, Tab. Zemarine Elix. |
| 0.02 mg. 0.02 mg. 0.0194 mg. | 0.006 mg. 0.007 mg. 0.0065 mg. | 0.1 mg. 0.1 mg. sulf. 0.1037 mg. | Chloral hydrate 250 mg. Chlorbutanol 250 mg. Dehydracetic acid 250 mg., phenobarbital 16.2 mg. | Fellows Wampale Whittier | Felo-Ral, Cap. or 5 cc. Butadonna, Cap. Bellabil, Tab. |
| 0.02 mg. | 0.0065 mg. | 0.1 mg. | Dehydrocholic acid 100 mg., prolose-300 30 mg., mylase-100 30 mg. | Meyer | Almesyme, Cap. |
| 0.019 mg. 1/1000 gr. | 0.006 mg. | 0.225 mg. sulf. 1/2000 gr. | Delvalnal Aethenamine 2 gr., phenyl salicylate 1/2 gr., methylene blue 1/10 gr., benzoic acid 1/2 gr. | Merck Sharp & Dohme Blaine | Delkadon, Tab. Uriseplin, Tab. |
| 0.0382 mg. | 0.0195 mg. | sulf. 0.3111 mg. | Pentobarbital sod. 43 6 mg | Columbus | Belokoids TT, Tab. |
| 0.024 mg. 0.06 mg. 0.0194 mg. | 0.0072 mg. 0.035 mg. 0.0065 mg. | 0.123 mg. 0.305 mg. sulf. 0.1037 mg. | Phenobarbital 0.016 mg. Phenobarb. 1 gr. Phenobarbital 16 mg. | Narmond Smith, Kline & French Dumas-Wilson | Narital, Tab. Prydonal, Spansule Anspas. Tab or Elix / 5 cc. |
| 0.0194 mg. | 0.0065 mg. | sulf. 0.1037 mg. | Phenob. 1/2 gr./1/2 gr. | Robins | Donnatal also NO2 |

| Atropine salt | Hyostine HBr | Hyoscyamine HBr | Others | Manufacturer | Brand |
|---------------|--------------|-----------------|---|-------------------|------------------------------|
| 0.0194 mg | 0.0065 mg | 0.1037 mg | Phenobarbital 1/4 gr she coat, oct valoid 1 gr | Hoag | Barcole 8 Tabs |
| 0.073 mg | 0.0272 mg | 0.2858 mg | Phenobarbital 48.6 mg | Vanpelt & Brown | Barb donna CR Tabs |
| 0.06 mg | 0.0195 mg | sulf 0.321 mg | Phenobarbital 48.6 mg | Roberts | Melinal Tab |
| 0.0194 mg | 0.0065 mg | sulf 0.1037 mg | Phenobarbital 1/4 gr | Bonar | Carenal Tab or El x /5 cc |
| 0.024 mg | 0.0072 mg | 0.123 mg | Phenobarbital 16 mg | Narmand | Narlat Tab |
| 0.015 mg | 0.0048 mg | 0.18 mg | Phenobarbital 1/4 gr | P. Iman Moore | Hytrana Tab El /5 cc. |
| 0.022 mg | 0.06 mg | sulf 0.04 mg | Phenobarbital 1/4 gr | Ch. coga Pharm | Hypnaldyne Tab |
| 0.0194 mg | 0.0065 mg | sulf 0.1037 mg | Phenobarbital 1/4 gr | Nycto | Spastolate Cap |
| 0.0194 mg | 0.0065 mg | 0.1037 mg | 16.2 mg | Carlone | Baritone Cap |
| 0.0194 mg | 0.0065 mg | 0.104 mg | Phenobarbital 1/4 gr | G. F. Harvey | Spasids Tab |
| 0.0385 mg | 0.021 mg | sulf 0.36 mg | Phenobarbital 1/4 gr | Infermed ca | Sustated Tab |
| 0.052 mg | 0.0195 mg | sulf 0.311 mg | Phenobarbital 50 mg | Eth cal Pharm | Bellatonic T D Cap |
| 0.025 mg | 0.0074 mg | sulf 0.1286 mg | Phenobarbital 16 mg | Vanpelt & Brown | Barb donna D A Tab |
| 0.02 mg | 0.007 mg | 0.1 mg | Phenobarbital 16 mg | Warren Teed | Bar Dan Tab El /5 cc. |
| 0.0214 mg | percent | percent | Phenobarbital 1/4 gr | Therachem cois | Annaphal Tab |
| 0.0171 mg | 0.0068 mg | sulf 0.0856 mg | Phenobarbital 1/4 gr | Sulf H & Case | Bar Cy Am ne Amp /cc. |
| 0.0171 mg | 0.0068 mg | sulf 0.0856 mg | Phenobarbital 1/4 gr | Sulf H & Case | Bar Cy A Tab Tab |
| 0.0195 mg | 0.0065 mg | 0.1040 mg | Phenobarbital 1/4 gr | Metro | Sedospom Tab |
| 0.0194 mg | 0.0065 mg | sulf 0.137 mg | Phenobarbital 1/4 gr | Eth cal | Bellatronic Cap |
| 0.02 mg | 0.006 mg | 0.103 mg | Phenobarbital 1/4 gr | Harton & Canversa | Sedala d |
| 0.0194 mg | 0.0065 mg | sulf 0.1037 mg | Phenobarbital 1/4 gr | Bentex Phr | Benedanna Tab |
| 0.02 mg | 0.003 mg | sulf 0.1 mg | Phenobarbital 8 mg calcium carbonate 225 mg | Crestmed | Doncarb Tab |
| 0.0194 mg | 0.0065 mg | 0.1037 mg | Phenobarbital 1/4 gr pec in 2 gr cool n 90 gr | Rob ns | Donnagel products |
| 0.0097 mg | 0.0033 mg | 0.0519 | Phenobarbital 1/4 gr ds hydroxyalum num aminooacetate 0.5 Gm | Rob ns | Dannalate |
| 0.0194 mg | 0.0065 mg | sulf 0.1037 mg | Phenobarbital 1/4 gr d hydroxy aluminum aminooacetate 0.5 Gm. 100 to 200 | Rob ns | Donnagel Susp /az |

Atropine-Hyoscine-Hyoscyamine Combinations (See also Belladonna Products)

| Atropine sulf. | Hyoscine HBr. | Hyoscyamine HBr. | Others | Manufacturer | Brand |
|----------------|---------------|-------------------|---|--------------------|----------------------|
| 0.0097 mg. | 0.00325 mg. | sulf. 0.0519 mg. | Phenobarbital 8.1 mg., aluminum hydroxide-magnesium carbonates 5 gr. | Benton | Almadon, Tab. |
| 0.0194 mg. | 0.0065 mg. | sulf. 0.1037 mg. | Phenobarbital 16.2 mg., glycine 150 mg., cal. carbonate 350 mg. | Lemor | DancoI, Tab. |
| 0.0097 mg. | 0.0033 mg. | sulf. 0.0518 mg. | Phenobarbital 8.1 mg., pepsin 150 mg., pantoic acid 300 mg., bile salts 150 mg. | Robins | Dannazyme, Tab. |
| 0.02 mg. | 0.0065 mg. | 0.2 mg. | Phenorb. 8 mg., sod. phenorb. 8 mg. | United Labs. | BorbeIoid, Tab. |
| 0.1 mg. | 0.1 mg. | 0.007 mg. | Phenorb. 16 mg., Tokadiostole 2 1/2 gr. | Parke, Davis | Bordose, Liq./4 cc. |
| 0.0242 mg. | 0.0081 mg. | sulf. 0.1296 mg. | Phenorb. 1/4 gr., vi. 8.1 12.5 mg. | First Texas | Hysial, Tab. |
| 0.08 mg. | 0.003 mg. | 0.12 mg. | Phenylsodiominopyridine HBr 100 mg. | Edwards | Urogesic, Tab. |
| 0.0195 mg. | 0.0065 mg. | sulf 0.1040 mg. | Phenylazodiominopyridine HCl 50 mg., sulf. fencloamide 500 mg. | Vitomix | Pylosetamide, Tab. |
| 0.0195 mg. | 0.0065 mg. | sulf. 0.104 mg. | Phenylazodiominopyridine 50 mg., sulfacetamide 500 mg. | Richard & Williams | Uricin, Tab. |
| 0.0140 mg. | 0.0042 mg. | sulf. 0.0702 mg. | Pyrilamine maleate 12.5 mg., tolyclamide 3 1/2 gr., acetophenetidin 2 1/2 gr., caffeine 1/2 gr. | Van Pelt & Brown | Pyr-Aspro, Tab. |
| 0.02 mg. | 0.006 mg. | sulf. 0.1 mg. | Reserpine 0.1 mg. | Lemmon | Redana/Tab. or 5 cc. |
| 0.025 mg. | 0.0074 mg. | sulf. 0.1286 mg. | Reserpine 0.1 mg. | Yanpelt & Brown | Raucanna, Tab. |
| 0.02 mg. | 0.007 mg. | 0.105 mg. | Reserpine 0.1 mg. | Walker | Serpedan, Tab. |
| 0.0097 mg. | 0.00325 mg. | sulf. 0.05185 mg. | Sulfacetamide 25 mg., phenylazo-diaminopyridine HCl 50 mg. | Columbus | Sulfid B-A, Tab. |

Atropine Sulfate Combinations

| Atropine Sulfate | Others | Manufacturer | Brand |
|----------------------|--|---------------------|--|
| 1/2000 gr | Acetophenetidin 2 gr, quinine sulfate 1/4 gr, camphor 1/4 gr, piperazine 1/40 gr, oil n 1/16 gr | Upjohn | Special Formula No 2, Tab or Cap |
| 0.03 mg | Atropinetidin 8 mg, strontium salicylate 32.5 mg, phenolphthalein 6.5 mg, camphor monobromated 32.5 mg, acanin 1.6 mg | Suit H & Case | Phenac Tab |
| 1/1000 gr | Aminophyllin 1/2 gr, veratrum viridum nat 1 gr, acenit powder 1/5 gr | Mallard | Amin in Tab |
| 1/360 gr | Amphetamine HCl 10 mg, thyro d 1 gr, aloin 1/2 gr | Dumas Wilson | Amphetamine Combina ions Tab |
| 1/360 gr, 0.18 mg | Amphetamine sulf 5 mg, thyro d 1 gr, aloin 1/2 gr di Amphetamine HCl 5 mg, thyro d 50 mg, aloin 50 mg vt B complex | Marco | Dastabs |
| 1/300 gr | Aspirin 3 gr, am chlor 3 gr, caffeine 1/2 gr | Narmand | Amphalin Tab |
| 1/1000 gr | Aspirin 2 gr, acetophenetidin 1 gr, caffeine 1/2 gr, phenylephrine HCl 5 mg, salicylamide 1 gr | Patch Clapp | Pradolain Tab Emagr n Forte Tab |
| 1/500 gr | Aspirin 3 gr, Dovers powder 1/4 gr, acetophenetidin 1 1/2 gr, camphor monobromated 1/2 gr, caffeine 1/2 gr | Suit H & Case | Asca Tab or Cap |
| 1/1000 gr | Aspirin 1 1/2 gr, phenacetin 1 1/2 gr, caffeine 1/2 gr, phenolphthalein 1/5 gr & camphor | Upjohn | Special Formula No 1, Tab |
| 0.012 mg | Aspirin 1/2 gr, phenacetin 1/2 gr, caffeine 1/15 gr, menthol 1/60 gr | Lemmon | Alrodyne Tab |
| 1/600 gr | Caffeine 1/10 gr, camphor 1/10 gr | Waldor Corp | Cafrese Tab |
| 1/500 gr | Cal carb 10 gr | Drug Prod | Alreacal Cap |
| 0.125 mg | Chloral hydrate 250 mg, cal bromide 0.5 Gm | Fellows | Fello Sed/4 cc |
| 1/250 gr | Chlorpheniramine maleate 12 mg, ephedrine HCl 1/2 gr | Marsh Emory Labs | Histolon Tabs |
| 1/300 gr | Demerol 50 mg /cc | Winthrop Labs | Demerol Atropine Amp |
| 0.12 mg | Desoxyephedrine HCl 7 1/2 mg | Eastern | Delatropin Loz |
| 1/360 gr | di Desoxyephedrine HCl 1/12 gr, thyro d 1 gr, aloin 1/2 gr | Hance | Amphyroid, Tab |
| 1/360 gr | di Desoxyephedrine HCl 5 mg, thyro d 1 gr, aloin 1/4 gr | Meyer | Myrobie Tab |
| 1/350 gr | Also without Aloin Alto w/Phenobarb 1/2 gr di Desoxyephedrine 1/12 gr, thyro d 1 gr | Meyer | Myrobie No 2, Tab |
| 1/300 gr | Also without Thyro d Alto w/Phenobarb 1/4 gr Di allylbarbituric acid 1/2 gr, mag carb 2 1/2 gr, cal carb 3 1/2 gr, | Buffington | Dia Tropine, Tab |
| 0.25 mg | bis, subcarb 1 gr | Endo | Hymarphon Atropine, Tab, or Amp /cc |
| 0.025 mg | Di hydromorphanone HCl 2 mg | Endo | |
| | Diphenoxylate HCl 2.5 mg | | |

Atropine Sulfate with Phenobarbital

| Atropine Sulfate | Phenobarbital | Others | Manufacturer | Brand |
|------------------|---------------|----------------------------|---------------------|---------------------------|
| 0.04 mg | 8 mg | | Central | Phentropine, Tab |
| 1/500 gr. | 1/16 gr. | | Massengill | Barbitone, No 1 Tab |
| 1/1000 gr. | 1/8 gr. | | First Texas | Amylatene & Atropine, Tab |
| 1/200 gr. | 1/4 gr. | Alulet (Keolin) 2 1/2 gr | Testagar | Bar Trapin, Tab |
| 1/1000 gr. | 1/8 gr. | d Amphetamine sulf 5 mg. | Malibie | Alupinal, Tab |
| 1/300 gr. | 1/16 gr. | d Thyroid 1 gr. | Beil Pharmacol | Celudex, Tab |
| 1/120 gr. | 1/4 gr. | d Thyroid 3 gr. | Low | Lowebese CTR, Cap |
| 1/120 gr. | 1/4 gr. | d Atropine sulf 15 mg. | Paul Maney | Niko Caps, Ph 749 |
| 1/300 gr. | 1/8 gr. | d Atropine HCl 5 mg. | Dumas | Antibese Tab |
| 1/500 gr. | 1/8 gr. | d Thyroid 1/4 gr. | Warren Teed | Asopine, Tab |
| 1/1000 gr. | 1/8 gr. | Aspirin 4 gr. | Zimmer | Biazophen, Tab |
| 1/500 gr. | 1/8 gr. | Blomith (Vibrit) 2 gr. | Sutliff & Case | C P A Tab |
| 1/500 gr. | 1/8 gr. | Calc carb 10 gr | Elder | Alcifax, Tab |
| 1/500 gr. | 1/8 gr. | Calc carb 3 1/2 gr. | | |
| 0.03 mg | 18.2 mg | Hyoscine HBr 0.13 mg | Haack | Cymtal Tab |
| 0.02 mg | 8 mg | Hyoscyamine HBr 0.2 mg. | United Labs | Barbaleid Tab |
| | | barbitol Sod 8 mg | | |
| 0.01 mg | 1/4 gr. | Hyoscyamine HBr 0.05 mg. | Haackell | Belnessum, Tab |
| 1/5000 gr. | 1/8 gr. | Mag trisulfate 7 gr | Vonpell & Brown | Silalaid Tab |
| 1/5000 gr. | 1/8 gr. | Hyoscyamine HBr 1/250 gr. | Vonpell & Brown | Silalaid, Tab |
| 1/1000 gr. | 1/8 gr. | Mag trisulfate 7 gr | Temman | Altromag, Tab |
| 1/1000 gr. | 1/8 gr. | Hyoscyamine sulf 1/250 gr. | | |
| 1/1000 gr. | 1/8 gr. | Hyoscyamus est 1/16 gr. | | |
| 1/300 gr. | 1/8 gr. | Mag trisulfate 7 gr | C D Smith | Matrobarb, Tab |
| 1/1000 gr. | 1/10 gr. | Mag trisulfate 5 gr | Warren Teed | Triotal Tab |
| 0.16 mg | 32 mg | Scopolamine HBr 0.22 mg | Ives Cameron | Compomal, Tab |
| 1/2000 gr. | 1/4 gr. | Scopolamine HBr 1/500 gr | Haack | Cymtal, Tab |
| 0.1 mg | 15 mg | Scopolamine HBr 0.2 mg. | Winthrop Labs | Apolomine, Tab. |
| | | pyridosane HCl 2 mg. | | |
| | | nicotina mide 25 mg. | | |
| 1/200 gr. | 1/4 gr. | Sulfur 1/2 gr | Peaythress | Anfracol, Tab |
| 2/360 gr. | 1/4 gr. | Thyroid 3 gr. | Taylor | Obetiro, T R Tab |
| 1/200 gr. | 1/4 gr. | d Amphetamine sulf 15 mg. | Rond | Cynidol, Elix & Tab |
| 1/200 gr. | 1/4 gr. | olein 1/4 gr. | Physicians Products | Tropinol, Liq. |
| 1/200 gr. | 1/4 gr. | Vitamins | Dumas W'lian | Atrolase, Cap |
| 1/150 gr. | 1/4 gr. | Vitamins | | |
| 1/150 gr. | 1/4 gr. | Yeast concentrate 1 gr. | | |
| | | Sed CMC 4 gr | | |

See also atropine hyoscyamine combinations

ATROSTIGMINE. (Harvey) Physostigmine salicylate 1 mg., atropine sulfate 0.65 mg./cc. Vial, 10 cc.

Use: Anterior poliomyelitis, rheumatoid arthritis, bursitis, myasthenia gravis.

ATTAPULGITE, ACTIVATED W/Dihydrostreptomycin, polymyxin B sulf., pectin. See: Polymagma, Susp. [Wyeth]

W/Pectin. See: Spensin, susp. (Ives-Cameron)

W/Pentobarbital sodium, 2-diethylaminoethyl diphenylacetate hydrochloride, aluminum hydroxide. See: Spasmatarb, Tab. (Palmedica)

W/Polymyxin B sulfate, dihydrostreptomycin base (as sulfate), pectin, hydrated alumina powder. See: Polymagma, Tab. [Wyeth]

W/Zinc oxide, colloidal sulfur, resorcinol, isopropanol. See: Sebasorb Lotion (Summec Labs)

ATTAPULGITE W/Dihydrostreptomycin salt, polymyxin B sulf., pectin, alumina gel. See: Spensin-PS, Susp. (Ives-Cameron)

AUBROMID. (McNeil) Gold tribromide 6/20 gr./30 cc. of elixir. Bot. 1 pt., 1 gal. Use: (1 dr. q. 4 h.) Control of cough.

AURACORT. (Columbus) Hydrocortisone 2.5 mg., polymyxin B sulfate 2000 u., neomycin 3.5 mg., procaine hydrochloride 10 mg., propylene glycol, glycerin/cc. Flat drop bot. 7.5 cc.

Use: Inflammation of ear.

AURALCAINE. (Ethical Pharm.) Antipyrine 5%, benzocaine 1%, red mercuric iodide 1.500, acetone 10%, alcohol 5%, glycerol base. Bot. ½ oz. w/dropper.

Use: Inflammation in the middle ear.

AURALGAN. (Doha) Antipyrine 0.81 Gm., benzocaine 0.21 Gm., in dehyd glycerine 17.98 Gm. fl. oz. Bot. with/dropper ½ fl. oz. Use: Antiphotogenic & analgesic ear drops.

AURASOL, Improved. (Rorer) Procaine butyrate 4%, magnesium sulfate 20.4%, with aromatics & glycerin. liq., Bot. w/dropper 15 cc.

Use: (2-4 gtt. in the ear) Relief of ear-ache in otitis media.

AURATHION. (Bruce) Gold sod. thiosulfate 10 mg., sod. thiosulfate 50 mg., sod. bisulfite 34 mg. w/benzyl alcohol/cc. Vial 10 cc.

Use: Rheumatoid arthritis.

AURAZINE. (Smith-Dorsey) Urea 10%, antipyrine 5%, benzocaine 0.5% & glycerine q.s./fl. oz. Bot. w/dropper ¼ fl. oz., ½ fl. oz., Bot. 1 pt.

Use: Analgesic & anesthetic ear drops.

AURCLOID, N.N.D. 1962. See: Radio-Gold (Au¹⁹⁹) Colloid.

* Generic name recognized by N.F., U.S.P. or A.M.A.

***AUREOMYCIN PREPARATIONS, N.N.D. 1962.** Capsules, U.S.P.; Injection, U.S.P.; Ophthalm., U.S.P., A.D.R. Ch'ortetracycline NCI U.S.P. (Lederle)

Calcium Oral Drops, Susp. (100 mg./cc.) Vial 10, 20 cc.

Calcium Syr. (125 mg./5 cc.) Bot. 4 oz. & 16 oz.

Cap. U.S.P. (50 mg.) Bot. 25s, 100s; (100 mg.) Bot. 25s; (250 mg.) Bot. 16s, 100s.

Ear Soln. (50 mg. pow.) (diluent propylene glycol 10 cc.) dropper bottle

Intravenous, U.S.P. Vial (100 mg., 250 mg. or 500 mg./vial)

Oint. Topically (3%) Tube ½ oz., 1 oz.

Oint. Ophth. (1%) Tube ¼ oz.

otic, Crystalline Vial (50 mg.) W/vial of propylene glycol w/benzocaine 5%.

Pharyngeal (15 mg.) Box 10s.

Pow. Spersal (Jar (75 Gm.—25 doses) (37.5 Gm.—12 doses)

Pow., Surgical Vial 5 Gm.

Sol., Ophth., U.S.P. Vial (25 mg.) w/dropper.

Tab., Soluble, Crystalline (50 mg./tab.) Bot. 40s, 100s.

Trachea (15 mg./trache) Bot. 25s, 250s.

W/Triple Sulfas. (Lederle) Aureomycin 125 mg., sulla meth-dia-mer, 167 mg/Tab Bot. 12s, 100s, 1000s.

Use: Orally, broad spectrum antibiotic with triple sulfonamide therapy.

***AUREOTOPE, N.N.D. 1962 (Squibb) Radio-Gold au 198. 20-50 mc per cc. Vials of 20 to 800 cc**

Use: Cancer therapy.

AURISTIL. (Blue Line) Benzocaine 9%, hexyl-resorcinol 0.1%, menthol 4%, camphor 2%. Dropper, rubber-top droppers, 12i. Use: Analgesic for ear-ache.

AURI-TUSSIN. (Zemmer) Gold tribromide 1/30 gr./10 min. Bot. ½ oz.

Use: Control of cough.

AUROBROM. (Professional Pharm. Co) Auric bromide 1/20 gr./10 min. Bot. dropper 15 cc., 30 cc., & oz., 1 pt.

Use: (10 to 15 gtt-drops i.i.d.) Whooping cough

AUROCIIN. (Christina) Gold naphthyl sulf-hydryl derivative. Amp (5% ac 12.5%) 2 cc.

Use: Antirheumatic gold therapy.

AUROLIN. See: Gold sodium thiosulfate (Various Mfg.)

AUROL-SULFIDE. (Hille) Gold sulfide, colloid. Bot. (1%) ½, 1 & 4 oz. for oral use only; Vial (2%) 15, 30 & 60 cc.

Use: I.M., arthritis.

AUOMERCAPTOACETANILID AUROTHIO-

- GLYCANIDE, NND 1961** See *Laurae*, Vial (Endo)
- AUBOPIN.** See Gold sodium Thiosulfate (Various Mfr)
- AUROSAN** See Gold sodium Thiosulfate (Various Mfr)
- *AUROTHIOGLUCOSE, NF, NND 1962;** Injection NF Gold thioglucose See *Solganal*, Amp (Schering)
- *AUROTHIOGLYCANIDE, NND 1962** Aurathio glycanilide, alpha auromercapto acetonilid See *Lauran Vial* (Endo)
- AUROTHIOGLYCOLANILID** See: Aurathio glycanide NND 1959
- AUROTHIOMALATE, SODIUM.** See Gold Sodium Thiomolate, NF
- AUTRINIC** (Lederle) Intrinsic factor concentrate See *Falvin with Autrinic* (Lederle)
Use: Increase absorption of Vit B 12
- AVAX, (Walker)** Crystalline vit A 25,000 u Bot 100s 1000s 50,000 u Bot 100s, 1000s 100,000 u Bot 1000s
- AVAZYME** (Wampole) Crystalline chymotrypsin 50,000 Wampole units/Tab Bot 48s 250s.
Use: Anti-inflammatory agent; enzyme preparation; chymotrypsin preparation
- AVC IMPROVED.** (National Drug) Cream containing 9 aminoacridine 0.2%, sulfacetamide 15% & allantoin 2% with lactose in a water-miscible base Tube w/ or without applicator 4 oz
Use: Trichomoniasis, moniliasis & mixed vaginal infections
- AVC SUPPOSITORIES** (National Drug) 9 Aminoacridine HCl 14 mg, sulfanilamide 1.05 Gm, allantoin 140 mg W/lactose buffered to acid pH Box 12s
Use: Vaginitis
- AVEENO** (Fougera) Gum fraction of oat meal, Pkg 18 oz, 4 lb
Use: Colloid bath therapy
W/Aluminum acet, col acet See: *Bur Veer*, *Pow* (Aveeno)
- AVEENO BAR** (Aveeno Corp) Colloidal oat meal (75%), hexachlorophene, hypochlorite derivative adjusted to pH 5.7 Cete 3% oz
Use: A soap free lathering bar for those who will avoid soap
- AVEENO "Oiled"** (Aveeno) Aveeno colloid oatmeal impregnated with a high percentage (75%) of liquid petrolatum and refined olive oil Hermetically sealed cans 15 oz
Use: Dry skin conditions
- AVEENO OINTMENT.** (Aveeno Corp) Aveeno Colloidal Oatmeal, stearic acids, and hexachlorophene 0.5% Tube 1 oz, 4 oz
*Generic name recognized by NF, USP or AMA
- Use For Itching or irritated skin conditions
- AVEENO POWDER** (Aveeno) Baby Powder Colloidal oatmeal, zinc oxide, hexachlorophen 1/4%, p-chloro meta xylenol 1/4% Pkg 3 3/4 oz 8 oz unscented **Foot Powder** Colloidal oatmeal, zinc oxide, boric acid p-chloro meta xylenol 1% Pkg 3 3/4 oz
Use: Topically, dermatologic
- AVEENO SOAP SUBSTITUTE.** (Aveeno Corp) Aveeno Colloidal Oatmeal in an emollient petrolatum base Tube 3 3/4 oz
Use: To cleanse sensitive skin
- AVENTYL.** (Lilly) Natriptyline
- AVERTIN.** NNR 1957 (Winthrop Labs) Tribromoethanol
W/Amylene hydrate, NNB 1955 (Winthrop Labs) Tribromoethanol solution, USP Bot 25 cc, 100 cc
Use: Rectal, general anesthesia
- A-V-G** (First Texas) Resorcin 2 1/2%, beta naphthol 5%, balsam peru 7 1/2%, sulfur ppt 20% Oint Tube 1 oz Jar 1 lb, 5 lb
Use: Antiseptic
- AVICAP.** (Burroughs Wellcome) Vit A 5000 u D 500 u, C 75 mg, B 1 2 mg, B 2 3 mg & nicotinamide 20 mg /Cap Bot 100s 1000s
Use: (1 daily) Multivitamin therapy
- AVINAR.** (Armour) Uredopo
Use: Anti neoplastic agent
- A-VISOL.** (Hance Bros and White) Water miscible vit A 25 M and 50 M u /Cap Bot 100s, 1000s
- A-VITA.** (Brewer) Vitamin A Capsules 25,000 u Bot 25s, 100s, 1000s Amp (25,000 u /cc) 1 cc Box 6s, 12s, 100s
Use: Vitamin A deficiency
- A-VITA-O Na 2.** (Pitman Moore) Vit A 5000 u, O 500 u /Cap Bot 100s
Use: (1 daily) Vitamins A & D deficiencies
- "A" VITAMIN IN OIL.** (Squibb) Vit A 100,000 u /cc Amp 1 cc Box 5s, 25s, 100s
Use: I.M., vitamin A deficiency
- "A" VITAMIN SYNTHETIC.** (Squibb) Vit A 50,000 & 25,000 u /Cap Bot 30s 100s 1000s
Use: Vitamin A deficiency
- AVITUM** (Ives Cameron) Vit A palmitate syn Cap Vit A 50,000 & 25,000 u Bot 30s, 100s
Use: Vitamin A deficiencies
- AVLOSULFON.** (Ayerst) Diaminodiphenylsulfone (Dapsone) Tab 100 mg Bot 100s
Use: Treatment of all forms of leprosy
- A.V.M** (Luka) Dextro-amphetamine sulfate 15 mg, vit A, 6600 USP units C, 400 USP units B 1, 160 mg; B 2, 250 mg; nicotinamide, 1550 mg; C, 3000 mg; ferrous sulfate 2000 mg; cobalt sulfate 0.49 mg

copper sulfate, 2.80 mg; sodium molybdate, 0.45 mg; zinc sulfate, 3.90 mg; potassium iodide, 0.13 mg / Cap. Bot. 100s.

Use: Obesity.

AVOSYL, N.N.D. 1959. (Schenley) Mephensin. Tab. (0.25 Gm) Bot. 100s, 1000s.

Use: Muscle relaxant

AXEFEN. (Warren-Teed) APC/Tabs. 1000s

AXENE. (Warren-Teed) Phenacelin 0.12 Gm and aspirin 0.25 Gm./Tab. Bot. 100s, 1000s

AXEROPHTNOL. See: Vitamin A.

AXIOUEL. (McNeil) Valnoctamide.

Use: Tranquilizing drug.

AXOFOR. (Warren-Teed) Phenobarbital ¼ gr., acetophenetidin 2 gr., acetylsalicylic acid 3 gr., dipyron 1 gr./cap. Bot. 100s.

Use: (1 l.i.d.) Non-narcotic, sedative, analgesic.

AXOPINE. (Warren-Teed) Atropine sulfate 1/500 gr., phenobarbital ¼ gr., acetylsalicylic acid 4 gr., acetophenetidin 2 gr./Tab. Bot. 100s, 1000s.

Use: (1 l.i.d.) Antispasmodic—analgesic.

AXOTAL. (Warren-Teed) Phenobarbital ¼ gr., acetophenetidin 2 gr., acetylsalicylic acid 4 gr./Tab. Bot. 100s, 1000s.

Use: (1 q. 3 h.) Analgesic—sedative.

* **AZACYCLONOL HCl**, N.N.O. 1962. Alpha, alpha-diphenyl-4-piperidinemethanol HCl. See: Franquel, Tab. (Merrell)

AZALONE OPHTHALMIC. (Smith, Miller & Patch) Antozoline phosphate 0.5%, boric acid 1.2%, w/sod. carbonate anhydrous, sod. chloride, and phenylmercuric acetate 0.002%. Bot. plastic squeeze w/dropper-lip, 15 cc.

Use: Antihistamine ophthalmic therapy.

* **AZAMETHONIUM BROMIDE**, 2-Methyl-3-azapentamethylenebis(ethyl-dimethyl-ammonium) dibromide. Peniomide.

* **AZAPETINE PHOSPHATE**, N.N.D. 1962. 6 Ally-6,7-dihydro 5H-dibenz (c,e) azepine, azepine. See: Hidar (Roche Lab)

* **AZATNIOPRINE**. (Burroughs Wellcome) 6-(1-methyl-4-nitroimidazol-5-ylthio) pterine. Use: Anti-leukemic compound

AZEPHINE (Azapetine). See: Hidar. (Roche Lab)

* **AZETEPA**. (Lederle Labs) P,P-bis (1-azidiryl) N-ethyl-N(1,3,4-thiadiazol-2-yl) phosphinic amide.

Use: Anti-neoplastic agent.

AZINOL. (Scheffelin) Zinc oxide, oleum & phenol. Oint. Tube 1 oz.

Use: Astringent & antiseptic.

AZOCHLORAMID, N.N.R. 1955. (Wallace & Tiernon) Chloroazodin, N.F., N,N'-Dichloroazodicarbonamidine, alpha, alpha, azo-bis(chloroformamidine). Soln. in Triacetin

* Generic name recognized by N.F., U.S.P. or A.M.A.

(1:500). Bot. 2, 8 oz., 1 pt., 1 gal. Saline Mixture, Bot. to prepare one gallon of 1:3300 soln. Tab. (to prep. 2 oz. of 1:3300 soln.) Jar 100s, 500s. Oint. (1:1000) Tebe 1 oz. Jar 1 lb.

Use: Topically in wound infections.

AZOCOL. (Patch) Aluminum subacetate, eucalyptol, carbolic acid, zinc oxide & schthynate. Tube 1½ oz. Jar 1 lb.

Use: Antiseptic—astringent.

AZODETTES. (Plessner) Sulfacetamide 125 mg, selfadiazine 125 mg., phenylazodiamino-pyridine 50 mg./Tab. Bot. 100s, 500s

Use: Sulfonamide therapy

AZODRINE SET. (Prema) Racemic epinephrine 2.25%, glycerite of epinephrine. Set—10 cc. with oral nebulizer, refill vial 10 cc.

Use: Bronchial asthma.

AZODYNE HCl. See: Phenylazodiaminopyridine HCl, Prep. (Various Mfrs.)

AZO GANTRISIN. (Roche Lab.) Gantrisin 500 mg., phenylazo-diamino-pyridine HCl 50 mg./Tab. Bot. 100s, 500s.

Use: (2 q i d) Urinary tract infections.

AZO GANTANOL. (Roche Lab) Sulfamethoxazole 500 mg., phenazopyridine HCl 100 mg./Tab. Bot. 100s

Use: Urinary anti-infective.

AZO KYNEX. (Lederle) Sulfamethoxy-pyridazine HCl 125 mg., phenylazodiaminopyridine HCl 150 mg./Tab. Bot. 100s.

Use: Sulfo therapy.

AZOLATE. (Amfre-Gran) Phenylazo-diamino pyridine HCl 25 mg., methenamine mandelate 250 mg., dl-methionine 50 mg./Tab. Bot. 100s.

Use: Urinary tract infections.

AZO-MANDELAMINE. (Warner-Chilcott) Phenylazo-diamino-pyridine HCl (Pyridium) 50 mg., methenamine maleate 500 mg./Tab Bot. 100s.

Use: Urinary infections.

AZOPYRINE. See: Azulfidine, Tab. (Pharmacia)

AZO-SUL. (Cambridge) Sulfadiazine, 1.67 gr., sulfamethazine 1.67 gr., sulfacetamide 1.67 gr., phenylazo-diamino-pyridine 50 mg., atropine sulfate 0.0194 mg., hyascyamine 0.1307 mg., hyoscine 0.0065 mg., phenobarbital 16 mg./Tab. Bot 100s

Use: Anti-infective, analgesic—cystitis, pyelitis.

AZO-SULFA B/A. (Tutag) Phenylazodiamine pyridine HCl 50 mg., hyoscine HBr D0065 mg., atropine sulf. 0.0194 mg., hyascyamine sulf. 0.1307 mg. in core, and self-acetamide 125 mg in outer coating/Tab Bot. 100s, 1000s.

Use Analgesic and antispasmodic w/zul fanamide therapy in urinary tract infections

AZOSULFAMIDE See Neoprontosil, Amp. Tab (Winthrop Labs)

***AZOTOMYCIN**. (Pfizer) Antibiotic isolated from broth filtrates of *Sireptomyces ombafaciens*

Use Anti neoplastic agent

AZOTREX. (Bristol) Tetracycline phosphate equiv to 125 mg of tetracycline HCl, sulfamethazole 250 mg phenylazadiamidesopyridine HCl 50 mg /Cap Bot 24s Syr Tetra cycline 125 mg Sulfamethazole 250 mg / 5 ml Bot 4 oz

Use: Treatment of urinary tract infections

AZO-UROQUID-A (Beach) Methesamine mandelate 0.5 Gm, O methanine 200 mg, phenylsodiopyridine HCl 75 mg, homatropine methylbromide 5 mg /E C Tab Bot 100s

Use Treatment of urinary infections

***AZOVAN BLUE**. Tetrasodium salt of 4,4'-di [7 (1 amine 8 hydroxy 2,4 disulpho-naphthylazo)] 3,3 bitolyl See Evans Blue Dye

AZTEC (Arnor Stone) Theophylline 125 mg, d isoephedrine HCl 25 mg, phenyltoloxamine 25 mg /Tab Bot 100s

Use Bronchial asthma and hay fever

AZULFIDINE, NND 1962 (Pharmacia) Sulfazoxazole sulfapyridine, ozopyrine Tab (0.5 Gm) Bot 100s

Use Ulcerative colitis

AZURE A CARBACRYLIC RESIN DIAGNOSTIC TEST, NND 1961 See Diognex Blue (Squibb)

***AZURESIN**, USP NND 1962 Azure A Carbacrylic resin 3 amine-7 dimethylamino phenozolium chloride See Diognex Blue (Squibb)

BABE-E-VITA DROPS. (Lincoln) Vit A 5000 u, D3 1000 u, C 60 mg, B1 1 mg, B2 0.6 mg, calcium pantothenate 3 mg, nicotinamide 10 mg, Vit B12 5 mcg, B6 D5 mg /0.6 cc Bot 30 cc.

Use Orally, multiple vitamin therapy

BABY-GARD (Lavenex) Cod liver oil, zinc oxide, starch, lact tanolin alkyl dimethyl benzyl ammonium chloride/squeeze bot 1 oz

Use Dermatologic cream

BACID. (U S Vitamin) A specially cultured strain of human *Lactobacillus acidophilus* sodium carboxymethylcellulose 100 mg /Cap Bot 50s

Use Functional diarrhea

BACIDRIN (Upjohn) Bacitracin 200 u.

* Generic name recognized by NF, USP or A.M.A.

ephedrine HCl 5 mg, myristyl-gamma picolinium chloride 1:10,000/cc. Bot. w/ dropper 15 cc

Use: Upper respiratory infections

BACIGUENT OINTMENT, NND 1962 (Upjohn) Bacitracin 500 u /Gm. Oint Tube ½ oz, 1 4 oz Ophth Oint Tube 1 dr

Use: Topically, pyogenic skin infections.

BACI-JEL. (Research Prods) Oint (bacitracin 500 u /Gm) Tube ½ oz Ophth. oint (bacitracin 500 u /Gm) Tube ½ oz.

Use: Pyogenic infections

BACILLET. (Abbott) Penicillin G Pot. 20,000 u, bacitracin 50 u /troche Bot. 10s, 100s 1000s

Use: Orally (1 q 2 h), infections of the oral cavity

BACILYRIUM. (Commercial Solvents) Bacitracin 500 u /cc w/D 0.5% cyclomethycaine Bot. 10 cc

Use: Topically, eye infections

BACIMYCIN OINTMENT. (Walker) Bacitracin 500 u, neomycin sulfate 5 mg /Gm Tube ½ oz, 100 Gm

Use: Topically, wide spectrum antibiotic in simple burns & abrasions

BACIMYCIN OPHTHALMIC OINT, (Walker) Bacitracin 500 u, neomycin sulfate 5 mg /Gm Oint, Tube ½ oz

Use Superficial infections of the eye

BACIMYCIN TABLETS. (Walker) Bacitracin 2500 u, neomycin sulfate 25 mg /Tab Bot 30s, 100s

Use Orally, Gram negative and Gram positive organisms

BACIMYCIN W/HYDROCORTISONE.

(Walker) Hydrocortisone 10 mg, bacitracin 500 u, neomycin sulf 5 mg /Gm. Oint Tube ½ oz

Use Inflammatory skin infections

***BACITRACIN**, USP, NND 1962, ADR Oint USP An antibiotic produced by a strain of *Bacillus subtilis*

Ointment, USP; Ophth Oint

Available forms (Various Mfr)

Diagnostic Tabs

Ointment

Ophthalmic Oint

Soluble Tabs

Systemic Use Vial

Topical Use, Vial

Troches

Vaginal Tabs

See Baciguent, Oint (Upjohn).

Baci Jel, Oint, Ophth oint (Research Prods)

Ginebatin, Vag Supp (Commercial Solvents)

Parentracin, Vial (Commercial Solvents)

- Topiracin, Oint., Ophth. Oint., Vial (Commercial Solvents).
- W/Benzocaine.
See: Baci-Trache (Upjohn).
- W/Caritrons acetate. See: Caritrons-Bacitracin Ophth. Oint. (Merck Sharp & Dahme)
- W/Cyclamethycaine. See: Bacillyrium, Sol. (Commercial Solvents).
- W/Desoxyephedrine HCl (Commercial Solvents) Bacitracin 250 u. & desoxyephedrine HCl $\frac{1}{4}\%$ /cc. Bot. 15 cc. (Bacitracin nasal).
- W/Ephedrine HCl. See: Bacidrin, Sol (Upjohn).
- W/Homarylamine HCl, tyrothricin, neomycin sulf., benzocaine. See: Pentozets, Trache. (Merck Sharp & Dahme)
- W/Hydrocortisone, neomycin sulfate. See: Bacimycin W/Hydrocortisone, Oint. (Walker)
- W/Hydrocortisone, tyrothricin, neomycin sulf., benzocaine. See: Hydrozets, trache. (Merck Sharp & Dahme)
- W/Iodochlorhydroxyquin, Methylene disalicylate. See: Anamaba, Tabs (Chicago)
- W/Neomycin base, hydrocortisone. See: Hydroderm, Oint. (Merck Sharp & Dahme)
- W/Neomycin sulfate.
See: Bacimycin, Oint., Ophth. Oint. & Tab. (Walker).
- Bacitracin-Neomycin, Ophth. Oint. (Paul Money) & (Abbott)
- Myctracin, Oint. (Upjohn).
- Neobacin, Tab. (Commercial Solvents)
- W/Neomycin sulfate, & propyl p-aminobenzoate. See: Du-Biotic, Traches (White)
- W/Neomycin, polymyxin B sulfate.
See: Neo-Polycin Oint. & Ophth. Oint. (Pitman-Moore).
- Neo-Polycin Ophth. Oint. (Pitman-Moore)
- Neoparin, Oint. & Ophth. Oint. (Burroughs Wellcome)
- Tribact, Oint. (Dermatologicals)
- Spectra-Biotic, Oint (American Pharmaceutical Co.)
- W/Neomycin sulf., polymyxin B sulf., hydrocortisone acetate. See: Neopolycin-HC, Oint., Ophth. Oint. (Pitman-Moore)
- W/Neomycin & polymyxin. See: Wybiotic Traches (Wyeth).
- W/Neomycin sulfate, polymyxin B sulfate
See: Dermospor, Oint. (Lowe)
- W/Neomycin sulfate, tyrothricin & benzocaine. See: Tetrazets, Traches. (Merck Sharp & Dahme)
- W/Penicillin G potassium.
See: Bacilllets, Traches (Abbott).
- Candicillin, Trache (Preme)
- Chailletr, Traches (Chase Chemical)
- Cillorels, Traches (Bristol)
- W/Phenacaine HCl. (Commercial Solvents)
- Bacitracin 1000 u. & phenacaine HCl $\frac{1}{4}\%$ /Gm. Ophth. Oint. Tube $\frac{1}{2}$ oz.
- W/Polymyxin B sulfate.
See: Candeltes, Traches (Pfizer).
- Polycin, Oint. & Ophth. Oint. (Pitman-Moore).
- Polysporin, Oint. & Ophth. Oint. (Burroughs Wellcome).
- W/Polymyxin B sulfate and Isobutyl p-aminobenzoate. See: Poly-Tracin, Traches. (Hance)
- W/Polymyxin B sulfate and neomycin sulfate
See: Trimixtin, Oint. (Hance)
- W/Polymyxin B sulfate, neomycin sulfate & hydrocortisone free alcohol.
See: Cortisporin, Ophth. Oint. (Burroughs Wellcome).
- W/Polymyxin B & streptomycin calcium chloride complex. See: Multibiotic Gauze Pads (Bio-Rama).
- W/Streptomycin sulfate & polymyxin B sulfate. See: Bio-Band Adhesive Strips (Multibiotic Corp. of America)
- W/Tyrothricin. See: Tyrotrac, Oint. (Merck Sharp & Dahme)
- W/Tyrothricin, neomycin sulfate & benzocaine. See: Tetrazets, Traches. (Merck Sharp & Dahme)
- W/Tyrothricin & benzocaine. See: Tracnets, Traches. (Merck Sharp & Dahme)
- *BACITRACIN METHYLENE DISALICYLATE.**
See: Anamaba (Chicago)
- &BACITRACIN-NEOMYCIN.** (Paul Money)
Neomycin sulfate equiv. to 3.5 mg. of base, bacitracin 500 u./Gm. Topical Oint. Tube $\frac{1}{2}$ oz., Ophth. Oint. $\frac{1}{2}$ gr.
Use: Treatment of infections due to gram-positive or gram-negative bacteria
- BACITRACIN ZINC W/Neomycin sulf., polymyxin B, benzalkonium chloride.** See: Biotres, Oint. (Central)
- BACI-TBOCHE.** (Upjohn) Bacitracin 1000 u., benzocaine 3 mg./trache. Bot. 20s
Use: Orally, antibiotic B local anesthetic trache
- BACTERIAL INTRAVENOUS PROTEIN.** (Cutter) Staph. albus & aureus 300,000, streptococci 300,000, pneumococci, H. influenzae, N. catarrhalis & B. Friedländer 100,000/cc. Vial 5 B 20 cc.
Use: I.V., arthritis vaccine.
- BACTINE.** (Miles) Methylbenzethonium chloride w/polyethylene glycol, mono-*is*-octyl phenyl ether, chlorothymol & alcohol. Bot 1 $\frac{1}{2}$ oz., 6 oz., 1 pt., 1 gal.
Use: Cationic detergent with germicidal & bactericidal activity.
- BACTO-SENSITIVITY DISKS.** (Lilly) Vial 10s
Use: To determine the sensitivity & resistance of micro-organisms to erythramycin

BACRATYCYN

98

BACRATYCYN (Wallace) Tyrothricin 1000 mg/Gm Oint Pkg 55 Gm

Use Topically, skin infections

BACULIN (Amfre Grant) Diiodohydroxyquinoline 100 mg, phenylmercuric acetate 3 mg sodium lauryl sulfate 3 mg, lactose dextrose, potassium alum and papain/vag Tab Bot 50s, 100s, 1000s Also available as powder

Use: (2 intervaginally, bid) for vaginal infections

BAKER'S MODIFIED MILK. (Baker) Modified cows milk with animal & vegetable fats lactose, dextrose, Vit A 2500 u, D 800 u B 1 0.6 mg, B 2 1.2 mg, niacin 5 mg, & iron ammonium citrate/lb, or qf liq 14 1/2 oz Pow 1 lb

Use Infant food

BALANCEL-FORTE. (Meyer) A vitamin mineral combination w/B 12 Caps Bot 100s

Use Supplement during pregnancy and lactation

BALANTROL. (Pritchard) Tablets, TD 15 mg Bot 30s 250s

BAL IN OIL, USP, NND 1962 ADR (Hynson, Westcott & Dunning) Dimercaprol, British anti Lewisite, 2,3-dimercaptopropanol Amp (10% in peanut oil) 3 cc Box 10s

Use IM, arsenic antidote

BALARSEN, NND 1962 (Endo) (Arsithinal, Metacyparsenal) It is the interaction of the arsenoxide form of acetarsone & dimercaprol (BAL) Tab (100 mg) Bot 25s 100s 1000s Sol 1% Bot 60 cc Vag supp (50 mg) Box 6s

Use Orally (10 mg/kg of body weight/day for five days) amebicidal agent

BALCORT. (Endo) Arsithinal and hydrocortisone

Use Treatment of Trichomonas vaginitis

BALMEX MEDICATED LOTION. (Mocsis) Fraction of lanolin, hexachlorophene, oil lanolin, balsam Peru, silicone Bot 4 fl oz

Use: Diaper rash

BALNETAR LIQUID. (Westwood) 2.5% crude coal tar, USP, lanolin derivative mineral oil and a nonionic emulsifier /tsp Bot 8 oz

Use: Dermatologic

BALTRINSIC SCT-7. (Simpson) Vit B 12, 25 mcg; stomach substance, 50 mg /Tab Bot 100s, 500s 1000s, 5000s

Use: Anemia

BALVIS (Carrick) Guar cellupectinoid 275 mg, magnesium oxide 70 mg, aluminum hydroxide gel 80 mg, magnesium trisilicate 160 mg /Tab Bot 40s, 100s

Use: Peptic ulcer and hyperacidity

BAMADEX. (Lederle) Meprobamate 400 mg

* Generic name recognized by NF, USP or A.M.A.

d amphetamine sulfate 5 mg /Tab Bot 100s

Use Appetite depressant for the overweight

BAMADEX SEQUELS. (Lederle) Meprobamate 300 mg, d amphetamine sulf. 15 mg /sust release cap Bot 30s

Use Obesity

***BAMETHAN** 2 n Butylamino 1 p hydroxy phenylethanol Vasculit sulphate

B-AMIN. (Buffington) Vit B 1, Tab (1 mg & 5 mg) Bot 100s, 1000s

Use Orally, thiamine therapy

BAMITE. (First Texas Chem) Pentobarbital sod 15 mg, d amphetamine sulf 10 mg, Vit B 12 10 mcg, ferrous gluconate 60 mg, methyl cellulose 200 mg /Tab Bot 100s

Use Anorexic agent in obesity

BANALG (Cofe) Menthol, camphor, methyl salicylate & eucalyptus in a greaseless base Bot 2 oz, 1 pt,

Use Topically, a counterirritant

BANARIN. (Pharmaceutical) Capsules, 10s, 20s

BANASIL. (Ulmer) Reserpine 0.25 mg /tab Bot 100s, 1000s (1 mg) Bot 100s Vial (2.5 mg/cc) 10 cc

Use Tranquilizing agent

BANAUSEA. (Amfre Grant) Scopolamine HBr 0.5 mg, benzocaine 100 mg, methoppyriene NCl 10 mg, mag hydroxyaminacetate 100 mg, B 6 5 mg /Tab Bot 60s, 120s, 1000s

Use Treatment and control of nausea and vomiting in pregnancy

BANCARIS. (Burroughs Wellcome) Thienum clarylate

Use Canine hookworm

BANDAMYCIN Spray Bandage. (New Med Techniques) Neomycin 3.5 mg per Gm of solid concentrate, freon as propellant; film former is plasticizing citric acid derivative

Can 2 oz

Use: Bandage for minor cuts, burns

BANESIN FORTE. (Kendall) Salicylamide 200 mg, mephenesin 200 mg, acetaminophen 200 mg, butabarbital Sod 10 mg, ascorbic acid 20 mg /Tab Bot 100s

Use Orally; muscular relaxant, analgesic

BANOCIDE. See Diethylcarbamazine USP

BANTHINE, NND 1962 (Searle) Methanethine bromide (beta-diethylaminoethyl xanthene-9-carboxylate methobromide) Amp, serum type, 50 mg Box 6s, 25s, & 100s

Tab 50 mg Bot 100s

Use: Orally (50-100 mg, q 6 h) anti cholinergic preparation

W/Phenobarbital. (Seorie) Banthine 50 mg., phenobarbital 15 mg./Tab. Bot. 100s.

Use: (1-2 q. 6 h.) Anticholinergic with sedative.

BANULAX. (Merrell) Dicyclamine HCl 30 mg., sod. carboxymethylcellulose 6 Gm., sod. lauryl sulfate 60 mg., casonthronal 180 mg./oz.

Use: Laxative.

BARACEL. (Barre) Zinc phenosulfonate $1\frac{1}{2}$ gr., cerium oxalate 2 gr., koolin 90 gr., pectin 2 gr., methyl cellulose 5%/R. oz. Bot. 6 oz., 12 oz., gal.

BARBADEX. (Kirkmon) Dextro-Amphetamine sulf. 5 mg., omobarbital $\frac{1}{2}$ gr./tab. Bot. 100s, 1000s, 5000s.

Use: Mental and emotional distress.

BARBA-NIACIN. (Cole) Phenobarbital $\frac{1}{2}$ gr., niacin $\frac{1}{4}$ gr./Tab. Bot. 100s, 1000s.

Use: Sedative.

BARBA-NIACIN FORTE. (Cole) Phenobarbital $1\frac{1}{2}$ gr., niacin $\frac{3}{4}$ gr. Bot. 100s.

Use: Sedative/hypnotic.

BARBATIN. (Fuller) Aspirin $2\frac{1}{2}$ gr., phenacolin $2\frac{1}{2}$ gr., phenobarbital $\frac{1}{4}$ gr./Tab. Bot. 100s, 500s.

Use: Analgesic-sedative.

BARBATRO, NO. 1. (Massengill) Phenobarbital $\frac{1}{4}$ gr., atropine sulfate 1/500 gr./Tab. Bot. 500s, 1000s

BARBATRO, NO. 2. (Massengill) Phenobarbital $\frac{1}{4}$ gr., atropine sulfate 1/250 gr./Tab. Bot. 500s, 1000s

BARBATRO, NO. 3. (Massengill) Phenobarbital $\frac{1}{4}$ gr., atropine sulfate 1/500 gr./Tab. Bot. 1000s

BARBELINE. (Cole) Phenobarbital $\frac{1}{2}$ gr., ephedrine HCl $\frac{1}{4}$ gr., ext. iobella $\frac{1}{4}$ gr./Tab. Bot. 100s, 500s, 1000s.

Use: Orally (1-2 q. 3 h), for paroxysms of bronchial asthma.

BARBELOID. (United Labs.) Hyoscyamine HBr 0.2 mg., atropine 0.02 mg., hyoscyne HBr 0.0065 mg., phenobarbital 8 mg., & Sod. phenobarbital 8 mg./Tab. Bot. 100s, 1000s.

Use: Orally (1-2 q. 3-4 h), antispasmodic & sedative.

BARBENYL. See, Phenobarbital.

BARB-EPHED. (Sutliff & Cose) Phenobarbital 16 mg., ephedrine HCl 24 mg./Tab. Bot. 100s, 1000s

BARBEPHINE. (Chicago Pharm.) Theophylline 2 gr., ephedrine $\frac{1}{2}$ gr., phenobarbital $\frac{1}{2}$ gr./Tab. Bot. 100s, 500s, 1000s.

Use: Orally (1-2), for relief of bronchial asthma.

BARBICAINE. (Cutter) Procaine HCl 50 mg.

* Generic name recognized by N.F., U.S.P. or A.M.A.

pentobarbital 4 mg., phenobarbital 4 mg./cc. Bot. 15 cc.

Use: Treatment of colic and cardiospasm
BARBIOLIN. (Taylor Labs.) Secobarbital Sod. 30 mg., pentobarbital Sod. 30 mg., phenobarbital 30 mg./Tab. Bot. 100s.

Use: Sedative & hypnotic.

BARBIDEX. (Flint-Eaton) di-Desoxyephedrine HCl 5 mg., phenobarbital 16 mg./Tab. Bot. 100s, 500s, 1000s.

Use: Orally (1-2 m.p.), improvement of mood.

BARBIDONNA. (Vanpelt & Brown) Phenobarbital $\frac{1}{4}$ gr., belladonna alkaloids 0.13 mg./Tab. or fl. dram. Elix., Bot. 1 pt., 1 gal. Tab., Bot. 100s, 500s, 1000s.

Use: Orally (1 Tab. or tsp.), antispasmodic & sedative.

BARBIDONNA-CR. (Vanpelt & Brown) Phenobarbital 48.6 mg., hyoscyamine SO_4 0.3858 mg., atropine SO_4 0.0750 mg., hyoscyne HBr 0.0222 mg./Tab. Bot. 50s.

Use: Antispasmodic.

BARBIDONNA D.A. (Vanpelt & Brown) (outer-coat) Hyoscyamine sulfate 0.1286 mg., atropine sulfate 0.025 mg., hyoscyne HBr 0.0074 mg., (core) phenobarbital 16 mg./Tab. Bot. 100s, 1000s.

Use: Before-meal spasmolysis and between-meal sedation.

BARBIDONNA No. 2. (Vanpelt & Brown) Phenobarbital $\frac{1}{2}$ gr., belladonna alkaloids 0.13 mg./Tab. Bot. 100s, 500s, 1000s.

Use: Antispasmodic & sedative.

BARBIFLORA COMPOUND. (First Texas) Phenobarbital Sod. 6 gr., pasiflora incarnata 32 gr./fl. oz. Elix. Bot. 1 pt., 1 gal.

BARBIPHENYL. See, Phenobarbital.

BARBIPI. (Chicago Pharm.) Phenobarbital 0.5 gr./pill. Bot. 100s, 1000s.

Use: Orally (1-2 q. 3-4 h.), induce sleep & hypnosis.

BARBISODITE. (Chicago Pharm.) Barbital Sod. 1 gr., Sod. nitrite 2 gr./pills. E.C. Bot. 500s, 1000s, 6000s.

Use: Orally (1), treatment of hypertension & angina pectoris.

BARBITA. (Chicago Pharm.) Phenobarbital $\frac{1}{4}$ gr./pill. Bot. 100s, 1000s, 6000s, 13,000s.

Use: Orally, sedative.

BARBITAB PLUS. (Barre) d-Amphetamine phosph. 5 mg., phenobarb. 16.2 mg., methylcellulose 162.5 mg., Vit. B-1 1.5 mg., B-2 1 mg., B-6 0.1 mg., B-12 0.25 mcg., niacinamide 10 mg., cal. pantothenate 0.5 mg., folic acid 0.1 mg., diac. phos. 200 mg./Tab., Bot. 100s, 1000s

* **BARRITAL, N.F.** N.N.R. 1949, A.D.R.; Tablets, N.F. (Barbitone, Deba, Dormonal,

Hypnogene, Malonal, Sedoval, Uronal
Veronal, Vesperial) Diethylbarbituric acid,
diethylmalonylurea See: Nauronidol, Eliz
(Schieffelin)
Use Orally, sedative, hypnotic
W/Aspirin & acetophenetidin See: Peralga,
Tab (Warner Chilcott)
W/Dihydroergotamine, phenobarbital Sod
scopolamine HBr, sandoptal See: Plexonal,
Tab (Sandoz)
W/Hyascyamus, chamomile & passiflora See:
Insomniat Tab (Burlington)
W/Hyascyamus ext, passion flower ext, &
valerian See: Neuradol, Tab (Tidann)
***BARBITAL SODIUM, N.F., N.N.B. 1949**
A.D.R.; Tablets, N.F. (Barbitone Sodium,
Diethylbarbiturate Monosodium, Dimethyl
malonylurea Sodium, Embinal, Medinal,
Veronal Sodium) See: Medinal, Eliz & Tab
(Warner).
Use: Hypnotic, sedative
W/Chloroform, ext valerian See: Neurinose,
liq & Teh (Faugerol).
W/Pentaerythritol tetranitrate See: PET B,
Caps (Barre Drug)
W/Phenobarbital Sod, sandoptal Sod,
scopolamine HBr, & Dihydroergotamine
methanesulfonate (D.H.E. 45) See: Plexonal,
Teh (Sandoz).
W/Sodium nitrite See: Barbitadite, Pills
(Chicago Pharm.)
BARBITONE, N.H.R. 1949, A.D.R. See: Bar-
bitol
BARBITONE SODIUM, A.D.R. See: Barbitol
Sodium
BARBITURATE-ASPIRIN COMBINATIONS.
See: Aspirin Tehle
BARBITURATES, INTERMEDIATE DURA-
TION.
See: Aprebarbital (Chemo Pure).
Butobarbital (Various Mfr.).
Butethal (Various Mfr.).
Delvinal (Merck Sharp & Dohme)
Diallylbarbituric Acid (Various Mfr.)
Lorvate (Winthrop Lab)
Prebarbital (Squibb).
Vinbarbital (Merck Sharp & Dohme)
BARBITURATES, LONG DURATION.
See: Barbital (Various Mfr.)
Mebaral (Winthrop Labs)
Mephobarbital (Winthrop Labs)
Phenobarbital (Various Mfr.)
BARBITURATES, SHORT DURATION.
See: Aethylbarbituric Acid (Sandoz)
(Sandoz).
Amobarbital (Various Mfr.)
Brevital (Various Mfr.)
Cyclobarbital (Winthrop Lab.)
Cyclopent (Upjohn)
*Generic name recognized by H.F., U.S.P. or A.M.A.

Cyclophen (Massengill)
Hexethal Sol (Parke, Davis)
Medomax (Gargy)
Ortal (Parke, Davis)
Pentobarbital Salts (Various Mfr.)
Phenadorn (Winthrop Labs)
Secobarbital (Various Mfr.)
BARBITURATES, TRIPLE.
See: Barbiclin, Tab (Taylor Labs)
Butseco, S.C.T. (Bowman)
Ethobral, Cap (Wyeth)
BARBITURATES, ULTRASHORT DURATION
See: Evral (Winthrop Labs)
Hexobarbital (Winthrop Labs)
Neraval (Schering)
Pentathal Sod (Abbott)
Surital Sod (Parke, Davis)
Thramylol Sod (Parke, Davis)
Thiopental Sod (Abbott)
BARBITWIN, (Paul Maney) Pentobarbital
sodium 1 gr butobarbital 3/4 gr/Tab Sol
100s 1000s
Use: Sedative
BARBIVIS (Chicago Pharm) Phenobarbital
15 gr/pill Sol 100s, 1000s
Use: Orally (1); sedative & hypnotic
BARBODEX (P. J. Koylto Co) No 1, 100s
No 2, 100s
BARBONATE (Bristol) Dihydroxy aluminum
aminoacetate 0.5 Gm, phenobarbital 8 mg,
belladonna alkaloids 0.16 mg, oil pepper
mint qs/Tab Sol 100s, 500s
Use: Orally (1 2 q 4 h), management of
gastric & duodenal ulcers; hyperacidity &
spasmodic pain
BARBOSEC, N.M.O. 1961, (Rawell) Seco-
barbital Sod 3/4 gr, 1 1/2 gr/Cap Sol 100s,
500s, 1000s
Use: (1 q 3 4 h) Sedative, hypnotic.
BARBRADINE, (Paul Maney) Butobarbital
1/6 gr, neophylline 2 1/2 gr, racephedrine
HCl 3/4 gr/Tab Sol 100s, 1000s
Use: Hay fever and asthma
BARBTHEO, (Smith, Millar & Patch) Theobromine
Sod acetate 5 gr, dicalcium phos-
phate 2 1/2 gr, phenobarbital 1/2 gr/Cap
or Tab Cap, Sol 100s, 1000s, Tab, Sol
100s, 1000s
Use: Orally, as a coronary vasodilator,
sedative & diuretic.
BARCOLE, (Haag) Phenobarbital 1/2 gr, acti-
vated charcoal 1 gr/Tab Sol 100s
Use: long-acting sedative
BARCOLE-B, (Haag) Phenobarbital 1/2 gr +
hyoscyamine 1037 mg, atropine sulfate
.0194, hyoscyamine HBr 006S, activated char-
coal 1 gr./Tab Sol 100s
Use: Antispasmodic
BAR-CY-AMINE, (Sullivan & Case) Pheno-

barbital 16.0 mg., hyoscyamine sulfate 0.0856 mg., atropine sulfate 0.0171 mg., hyoscyne hydrobromide 0.0068 mg., alcohol 9.0%/5 cc. Bot. 1 pl., 1 gal.

Use: Orally (1 dr. q. 3-4 h), antiparasympathetic redative.

BAR-CY-A-TAR. (Sullivan & Case) Phenobarbital $\frac{1}{4}$ gr., hyoscyamine sulfate 0.0856 mg., atropine sulfate 0.0171 mg., hyoscyne hydrobromide 0.0068 mg./Tab. Rot. 1000s

Use: Antiparasympathetic sedative

RARDASE. (Parke, Davis) Belladonna ext. $\frac{1}{6}$ gr., phenobarbital $\frac{1}{4}$ gr., taka-diastore $2\frac{1}{2}$ gr./tab. Rot. 100s, 1000s, liq. Phenobarbital $\frac{1}{4}$ gr., takadistore $2\frac{1}{2}$ gr., hyoscyamine sulfate 0.1 mg., hyoscyne 0.007 mg., atropine 0.02 mg./4 cc. Bot. 1 pt.

Use: (1-2 i.i.d.) Anticholinergic, sedative & starch-digestant.

BAR-DEX. (Barre) d-Amphetamine phosp. 5 mg./Tab. Bot. 100s, 1000s

BAR-DEX TD. (Barre) d-Amphetamine sulf. 15 mg. lime dirint./Cap. Bot. 25r, 300s

RARDIF. (Paul Money) Methocel 400 cps. 22 gr., aluminum hydroxide 2.0 gr., sod. lauryl sulfate 0.22 gr./45 gr. Paw. Bot. 4 oz., 1 lb.

Use: X-ray investigation of the gastro-intestinal tract

BARDIOL. (Barry) Estradiol. Aq. surp. or in Oil Vial (0.1 mg./cc.) 10 cc., 30 cc.; Vial (0.2 mg./cc.) 10 cc., 30 cc. Aq. susp. only: Vial (0.5 mg./cc.) 10 cc., 30 cc.

Use: I.M., estrogen therapy.

BAR-DON. (Warren-Teed) Hyoscyamine HBr 0.100 mg., hyoscyne HBr 0.007 mg., atropine sulfate 0.02 mg., phenobarbital 16.670 mg./5 cc. Tab. contain phenobarbital sodium. Elix., Sol. 1 pt., 1 gal. Tab., Rot. 100r, 500r, 1000s.

Use: Orally, redative & antispasmodic.

RAR-DOX. (Barry-Martin) Pyridoxine KCl 10 mg./Tab. Bot. 100s and 1000r. Vial (25 or 50 or 100 mg./cc.) 10 cc.

RARD-PARKER FORMALDEHYDE. (Bord-Parker) Isopropyl alc. 65.26%, methanol 27.5%, formaldehyde 8%, hexachlorophene 0.5% & water 23.39%. Bot. 1 gal.

Use: Bactericide

RARESTROGEN. (Barre) Conjugated estrogenic substances (water soluble). Tab. 1.25 mg. Bot. 100s, 1000s.

RARI-STRESS. (Barre) Vit. R-1 10 mg., B-2 10 mg., niacinamide 100 mg., Vit. C 300 mg., B-6 2 mg., B-12 4 mcg., folic acid 1.5 mg., cal. pantothenate 20 mg./Cap. Bot. 100s.

Use: Stress vitamin therapy.

RARITAR PLUS. (Barre) d-Amphetamine

* Generic name recognized by N.F., U.S.P. or

phosphate 5 mg., phenobarbital 16.2 mg., methylcellulose 162.5 mg., Vit. B-1 1.5 mg., R-2 1 mg., B-6 0.1 mg., niacinamide 10 mg., Cal. pantothenate 0.5 mg., folic acid 0.10 mg., Vit. B-12 0.25 mcg., dicalcium phosphate 200 mg./Tab. Bot. 100r, 1000s.

Use: Anorectic in obesity.

* **BARIUM SULFATE, U.S.P., N.N.R. 1949.** A.D.R. (Mallinckrodt) Barium sulfate U.S.P. for X-ray diagnosis. Carister 1 lb., 5 lb., Drum 10 lb., 25 lb., 100 lb., 250 lb.

Use: Orally or rectally, X-ray diagnosis.

BARIUM SULFATE PREPARATION, N.N.R. 1949.

See: Baropaque Surp. (National Synthetic).

Ray-Tec Lap Packs (Johnson & Johnson)

Rugar (McKesson & Robbins)

Skraybart (Merck Sharp & Dohme)

Ultrapaque-A, R and C (Bell-Craig)

BARIUM SULFATE SURGICAL SPONGES.

See: Ray-X Spanger (Parke, Davis).

BARIZOL. (Barre) Dipyrone 50 mg., Mephensin 125 mg., solicylamide 125 mg., and Vit. C 50 mg./Cap. (yellow) Bot. 100s, 1000s

Use: Analgesic-relaxant

BARMAL. (Paul Money) Dehydrocholic acid 3 gr., hyoscyamine HBr $\frac{1}{300}$ gr., magnesium trisilicate $1\frac{1}{2}$ gr., dried aluminum hydroxide gel $1\frac{1}{2}$ gr./Tab. Bot. 100s, 1000s.

Use: Gastro-intestinal distress.

BAROLOID. (Bell-Craig) Microbarium sulfate. Consists 10 lbs., 50, 100, 250 lbs.

Use: Contrast medium for radiography of the alimentary tract.

BARONE. (Barre) Dipyrone. Vial 500 mg./cc. 30 cc.

Use: Antirheumatic and analgesic.

BAROPAQUE, N.N.R. 1949. (Bell-Craig) 95% Barium sulfate, 4% flavoring & 1% suspension medium. Pkg. $4\frac{1}{2}$ oz. doses.

Use: Orally or enema, alimentary tract radiography.

RAROSMIN. See: Diarmin. W/Methenamine & resarcinol. See: Buchform, Tab. (Strassenburgh).

RARPHYLLIN. (Mallard) Ammonium chloride 5 gr., theophylline $1\frac{1}{2}$ gr., phenobarbital $\frac{1}{4}$ gr./Tab. Rot. 100s, 500s, 1000r.

Use: Orally (1-2 i.i.d.), hypertension & sedative.

BARPLEX. (Masrengill) Phenobarbital 20 mg., Vit. R-1 5 mg., Vit. B-2 2 mg., nicotinamide 25 mg./5 cc. or Tab. Bot. 1 pt., 1 gal. Tab. Rot. 100s, 1000r

Use: Sedative w/Vit. R Complex therapy

BARPRINE. (Ulmer) Diallylbarbituric acid

A.M.A.

- $\frac{1}{2}$ gr. aminopyrine $3\frac{1}{2}$ gr/Tab Bot 100s
1000s
Use Orally (1 2 1 1 d), sedative & analgesic
- BARRE FORMULA.** (Barre) d Amphetamine sulf 5 mg. phenobarb 1/6 gr., thyroid $\frac{1}{2}$ gr/Tab Bot 100s, 1000s
Use Central nervous system stimulant
- BARTDNE** (Cartone) Phenobarbital 16.2 mg., scopolamine HBr 0.0065 mg., atropine sulfate 0.0194 mg., hyoscyamine sulfate 0.1037 mg./Cap Bot 100s
Use Antispasmodic
- BARTRATE** (Davis & Sly) 5 Ethyl 5 isopentyl barbituric acid 50 mg. atropine methyl sulfate 0.5 mg./Tab Bot 100s 500s
Use: Treatment of peptic ulcer
- BAR-TRDPIN** (Testagor) Atropine sulfate 1/200 gr., phenobarbital $\frac{1}{4}$ gr/Tab Bot 100s 100s 1000s
Use Orally (1 2 q 3-4 h), antispasmodic & sedative
- BARTRDPIN TIMED**, Capsules (Testagor) Atropine sulfate 1/66 gr., phenobarbital $\frac{1}{4}$ gr./cap Bot 100s
Use Timed & integrating antispasmodic
- BARIBATE** (Plant Eaton) Sod pentobarbital $\frac{1}{4}$ gr., phenobarbital $\frac{1}{4}$ gr./Cap Bot 100s 500s 1000s
Use Orally (1 2 1 1 d), sedative & hypnotic
- BARISU.A.C.** (Barre) Dihydrocodeinone bitartrate 0.01 Gm., phenyltolazamine dry hydrochloride 0.1 Gm., pot guaicol sulfate 0.525 Gm., emm chlor 0.224 Gm., iperac F est 0.12 cc., labelia 0.2935 Gm., sod am citrate 12 Gm., sanguinaria 0.231 Gm., menthol and talc q.s./30 ml
- BARTWD** (Umeis) Secobarbital sod $1\frac{1}{2}$ gr., dry barbituric acid $\frac{1}{4}$ gr./tab Bot 100s 500s
Use (1 h) sedative
- BASALIL**, NND 1962 (Wyeth) Basic aluminum carbonate gel Bot 12 fl oz
Use Orally prophylaxis of renal stone
- BASIBGIN** (Sandoz) Ergonovine tartrate water soluble 0.25 mg./tab or 0.2 mg./Amp Bot 10s 100s, 500s, 1000s. Amp 1 or Bar 6s 12s 50s 100s
Use Orally, IV., SC. or IM., erythracis
- BASIX**, (Kalamberg) Polymonoeratin Cap 0.75 Gm 1 Bot 100s, 1000s
Use (2 q 2 h) Management of gastric hyperacidity
- BASIC ALUMINUM AMINOACETATE**, NND 1957 See Dihydroxy Aluminum Aminoacetate
- BASIC ALUMINUM CARBONATE**, NND 1942 See Basalil Sulf (Wyeth)
- BASIC ALUMINUM GLYCINATE**, NND 1957 See Dihydroxy Aluminum Aminoacetate
- 1959 See: Dihydroxyaluminum aminoacetate
- BASIC BISMUTH CARBONATE**. See: Bismuth Sebcarbonate, U.S.P
- BASIC BISMUTH GALLATE**. See: Bismuth Subgallate (Various Mfr)
- BASIC BISMUTH NITRATE**, NNR 1943 See: Bismuth Subnitrate (Various Mfr)
- BASIC BISMUTH SALICYLATE**. See: Bismuth Subsalicylate, U.S.P
- BASIC FORMULA VITAMIN**, ADR 1954 (Squirbb) Vit B 1 10 mg., B 2 5 mg., niacinamide 150 mg., Vit C 150 mg./Tab Bot 30s, 100s, 250s, 1000s
Use: Vitamin deficiencies
- BASIC FORMULA VITAMIN**, (West Ward) Vit B 1 10 mg., B 2 5 mg., niacinamide 150 mg., Vit C 150 mg./Tab Bot 100s, 1000s See: Vitelin, Tab (Squirbb)
- BASIC FUCHSIN**, NF See: Carbol Fuchsin Solution NF
- BASIGETS ULTIMA** (C B Kendall) Methyl androstenediol 2.5 mg., ethinyl estradiol 5 mcg., vit A 4000 u, vit D 400 u, vit B 12 3 mcg vit B 1 1 mg., vit B 2 2 mg., niacinamide 10 mg., pyridoxine hydrochloride 0.5 mg., pantothenyl alcohol 1 mg., monosodium glutamate 50 mg., vit C 30 mg., cobalt chloride 5 mg., iron sulfate 75 mg., DL methionine 5 mg., Llysine 50 mg., hesperidin complex 25 mg., yeast prote hydrolysate 50 mg./Tab Bot 100s, 500s
Use Geriatric anabolic and multivitamin supplement
- BASSORAN PLAIN AND BASSORAN WITH CASCARA** (Merrell) Plain Sterculia gum 87%, magnesium trisilicate 8.7%, with cascara sterculia gum 75%, magnesium trisilicate 7.5%, aromatic cascara 72 minims/oz Bot 7 oz, 24 oz
Use Orally, chronic constipation
- BASTRAN** (Wampole) Vit B 12 75 mcg. Sod metacinate 20 mg., adenosine 5 monophosphoric acid 25 mg., procaine HCl 1%/cc Amp 5 cc.
Use IM., treatment of bursitis and osteoarthritis
- BAX**, (McKesson & Robbins) Vit A 5000 u, D 500 u, B-1 333 u, B 2 2 mg., C 30 mg., niacinamide 10 mg., calcium pantothenate 1 mg., Vit B-6 D2 mg./Cap Bot 100s 250s
Use Orally, multivitamin therapy
- BAYRR 205**, NND 1954 See Naphuride Sod. Amp (Winthrop Lab.) Serum Sodium, U.S.P XIV (No Mfr currently lists)
- B-B-B** (Paternal Drug) Benzyl benzoate

- 25% in an aromalized base. Bot. 4 fl. oz., 1 pt., 1 gal.
Use: Externally, scabicide.
- BCG VACCINE, N.N.D. 1962.** See: Tuberculosis Vaccine.
- B.C.N. IMPROVED.** (Hoock) Vit. B-1 10 mg., ascorbic acid 150 mg., niacinamide 150 mg., Vit. B-6 5 mg./Tab. Bot. 100s, 1000s.
Use: Orally, vitamin therapy.
- B-COMPLEX.** (Mallard) Vit. B-1 100 mg., B-2 0.9 mg., B-6 1.0 mg., nicotinamide 100 mg./5 cc. or Tab. Amp. 5 cc., 30 cc., Tab. Bot. 100s, 500s, 1000s. See: Vitamin B complex tables.
Use: Orally, vitamin B complex therapy.
- W/Vit. C.** (Smith-Dorsey) Vit. B-1 0.25 mg., B-2 0.5 mg., B-6 0.125 mg., ascorbic acid 7.5 mg., calcium pantothenate 0.125 mg., niacinamide 2.5 mg./wafer. Bot. 100s, 250s.
Use: Vitamin B complex with vitamin C.
- B-COMPLEX ELIXIR.** (Wyeth) Vit. B-1 1.3 mg., B-2 1.3 mg., B-6 0.5 mg., pantothenic acid 0.8 mg., niacin 6.3 mg. & rice bran/5 cc. Elix. Bot. 4 oz., 32 oz.
Use: Orally, vitamin B complex therapy.
- B-COMPLEX FORTIFIED.** (Walker) Yeast extract & liver 300 mg., thiamine HCl 10.0 mg., riboflavin 3.0 mg., pyridoxine HCl 0.2 mg., calcium pantothenate 30 mg., niacinamide 20 mg., Vit. B-12 50 mcg., folic acid 0.67 mg., biotin 0.1 mg., p-aminobenzoic acid 10.0 mg., inositol 20.0 mg., choline dihydrogen citrate 20.0 mg./Cap. Bot. 100s, 1000s.
Use: Multiple vitamin B deficiencies.
- B-COMPLEX INJECTABLE.** See: Hypobeta, Vial (Merck Sharp & Dohme)
- B-COMPLEX LIQUID.** (Smith, Miller & Patch) Vit. B-1 12 mg., B-2 8 mg., niacinamide 40 mg., panthelol B mg., Vit. C 60 mg W/ rice bran/fl. oz. Bot. 4 oz., 1 pt., 1 gal.
Use: Vit. B-complex therapy
- B-COMPLEX PABENTERAL.** (Parke. Davis) Vit. B-1 10 mg., B-2 2 mg., B-6 5 mg., pantothenic acid 10 mg., niacin 50 mg./cc. Vial 10 cc.
Use: Injectable vitamin B complex therapy.
- B-COMPLEX SYRUP.** (Abbott) Vit. B-1 2 mg., B-2 1.5 mg., niacin 20 mg. & yeast extract/5 cc. Bot. 12 oz., 1 gal.
Use: Orally, vitamin B complex therapy.
- B-COMPLEX TABLET.** (C. D. Smith) Vit. B-1 2 mg., B-2 1 mg., niacinamide 10 mg., panthelol 1 mg., Vit. C 15 mg., B-12 1 mcg., pyridoxine HCl present/Tab. Bot. 1000s
Use: Vit. B-complex therapy
- B-COMPLEX WITH C.** (Sulliff & Case) Vit. B-1 3 mg., B-2 3 mg., C 30 mg., B-6 1.5 mg., pantothenic acid 2 mg., niacin 15 mg., yeast 5 mg./Tab. Bot. 100s, 1000s.
Use: Orally, vitamin B complex w/Vit. C.
- B COMPLEX AND IRON.** (Smith, Miller & Patch) Ferrous sulfate 200 mg., Vit. B-1 12 mg., B-2 0.4 mg., Niacinamide 4 mg., Calcium pantothenate 0.24 mg., Vit. B-12 1 mcg./Tab. Bot. 100s, 1000s.
- B-COMPLEX W/LIVER EXTRACT.** (Squibb) Vit. B-1 10 mg., B-2 3 mg., niacinamide 100 mg., crude liver ext./1 cc. Vial 10 cc., 1e. 10s.
Use: I.M. (1 cc. daily) in B complex deficiencies and severe nutritive failure.
- B-COMPLEX VITAMIN.** (Squibb) Thiamine mononitrate 2.0 mg., riboflavin 2.0 mg., niacinamide 15.0 mg., pyridoxine HCl 0.125 mg., pantothenic acid 2.0 mg., yeast 100.0 mg., Vit. B-12 2.0 mcg., folic acid 0.134 mg./Tab. Bot. 30s, 100s.
Use: Orally, dietary supplement.
- B-COMPLEX W/VIT. C.** (Abbott) Vit. B-1 3 mg., B-2 3 mg., nicotinamide 20 mg., Vit. B-6 0.1 mg., pantothenic acid 3 mg., ascorbic acid 30 mg., liver fraction #2 0.33 Gm./Cap. Bot. 100s, 250s, 500s, 1000s.
Use: Orally, vitamin B complex with vitamin C therapy.
- B-DENOSE.** (B. J. Ascher) Adenine-5-monophosphoric acid 25 mg., Vit. B-12 50 mcg., B-1 10 mg., nicotinic acid 25 mg./cc. Vial 10 cc.
Use: Treatment of varicose vein
- BDEP.** Same as Benzathine Penicillin G.
- B-DOX.** (Lannett) Vit. B-1 100 mg., B-6 100 mg./cc. Vials 10 cc.
Use: I.M. or I.V., nausea and vomiting
- BEADOX, N.N.D. 1960** (Preme) Pyridoxine HCl. Tab 10 mg., 25 mg. Bot. 100s.
- BEASY.** (Blaine) Rescinnamine 0.1 mg., B-1 10 mg., B-12 25 mcg./Tab Bot 100s, 1000s.
Use: Antihypertensive.
- BEATINE.** (Preme) Thiamine hydrochloride. U.S.P. Elix. (2000 u./fl. oz.) Bot. 1 gal. Vial (50 mg/cc) 10 cc., 30 cc. Vial (100 mg/cc) 10 cc., Tab. (5 mg., 10 mg., 25 mg., 50 mg., 100 mg., 250 mg) Bot. 100s, 1000s.
- BEBALY.** (Baine) Vit. B-1 10 mg., B-12 25 mcg./5 cc. Liq Bot pt., gal.
Use: Vitamin supplement.
- BE-BE CREAM.** (Crookes-Barnes) Silicone, zinc oxide, catamine, starch, hexylresorcinol Tubes 1 & 2 3/4 oz. Jar 1 lb
Use: Skin protector and diaper rash.
- BECAPLETS.** (Breon) Vit. B-1 3 mg., B-2 2 mg., B-6 0.1 mg., Cal. pantothenate 1 mg., niacinamide 10 mg./Cap. Bot. 100s, 1000s.
Use: Orally, vitamin B complex therapy.
W/Vit. C. (Breon) Same as Becaplets with

vitamin C 50 mg added Bot 100s, 1000s
Use: Orally, vitamin B complex with Vit C therapy

BECEMOR, (Chicago Pharmacol) Vit C 1500 mg, B 1 200 mg, B 2 100 mg, B 6 50 mg B 12, concentrate, containing B 12 activity of 0.5 mcg, folic acid 0.5 mg nicotinamide 1000 mg, calcium pantothenate 50 mg /Tab Bot 40s 100s 1000s
Use: Orally dietary supplement

BECEMAX (Winthrop Labs) Vitamins B 1 5 mg, B 2 3 mg, B 6 3 mg C 50 mg niacin 40 mg, pantothenic acid 3 mg /10 cc Vial 10 cc Formerly Becejex

BECEVITE, (Carol) B 1 15 mg B 2 10 mg, B 6 5 mg, C 250 mg /calcium pantothenate 10 mg, nicotinamide 50 mg hesperidin purified 20 mg /Tab Bot 100s

Use: Pre and post surgical cases; hasten wound healing in peptic ulcers

BECLAMIDE N Benzyl b chlorpropionate (Nydrane)

BECLYSYL W/ALCOHOL. See Alcohol in Beclysyl (Abbott)

BECLYSYL 5% IN SALINE, (Abbott) Dextrose 50 Gm, Sodium chloride 9 Gm, thiamine HCl 10 mg, riboflavin 10 mg, nicotinamide 250 mg, pyridoxine HCl 5 mg, in water for injection/1000 cc; 1000 cc, Bot 1s, 6s, 48s; 500 cc, Bot 1s 6s 48s
Use: IV, dietary supplement

BECLYSYL 5% IN WATER, (Abbott) Dextrose 50 Gm, thiamine HCl 10 mg, riboflavin 10 mg, nicotinamide 250 mg, pyridoxine HCl 5 mg, in water for injection/1000 cc; 1000 cc, Bot 1s, 6s, 48s; 500 cc, Bot 1s, 6s, 48s
Use: IV, dietary supplement

BECLYSYL 10% IN SALINE, (Abbott) Dextrose 100 Gm, Sodium chloride 9 Gm, thiamine HCl 10 mg, riboflavin 10 mg, nicotinamide 250 mg, pyridoxine HCl 5 mg, in water for injection/1000 cc; 1000 cc, Bot 1s 6s, 48s; 500 cc, Bot 1s, 6s, 48s
Use: IV, dietary supplement

BECLYSYL 10% IN WATER, (Abbott) Dextrose 100 Gm, thiamine HCl 10 mg, riboflavin 10 mg, nicotinamide 250 mg, pyridoxine HCl 5 mg, in water for injection/1000 cc; 1000 cc, Bot 1s, 6s, 48s; 500 cc, Bot 1s, 6s, 48s
Use: IV, dietary supplement

BICOJECTION, (Endocrine) Vit B-1 50 mg, B-2 1 mg, B-6 2 mg, nicotinamide 100 mg, pantothenol 10 mg, Vit B-12 15 cc

10 mcg), Vit B 2 0.3 mg, niacinamide 50 mg, Vit. B 6 0.3 mg /2 cc Vial 30 cc
Use: IM, anemias

BECOMCO ELIXIR, (Brannon) Vit B-1 3 mg B 2 2 mg, B 6 1.5 mg, B-12 1 mcg, nicotinamide 10 mg, calcium pantothenate 0.85 mg, ferric ammonium citrate 180 mg, liver fraction # 1, 200 mg /5 cc Bot 1 of, 1 gal

Use: Orally, conditions of malnutrition

BECOMCO ELIXIR PLUS, (Breon) Soluble ferric pyrophosphate 265 mg, B 1 3 mg B 2 2 mg B 6 1.5 mg, niacinamide 10 mg, Cal pantothenate 0.85 mg B 12 1 mcg /5 cc Bot 1 pt, 1 gal

Use: Iron and multivitamin tonic

BECOPERLES (Endocrine) Vit A 5000 U 1000 U, B 1 3 mg, B 2 3 mg, nicotinamide 20 mg calcium pantothenate 5 mg, Vit B 6 0.25 mg, E 5 IU/perle Bot 100s

Use: Orally vitamin therapy

BECOPLEX, (Endocrine) Vit B-1 5 mg, B 2 3 mg, B 6 0.5 mg, B 12 1 mcg niacinamide 20 mg, panthenol 3 mg, choline 20 mg, liver NF 100 mg /Tab Bot 100s

BECOPLEX CONC DROPS, (Endocrine) Vit B 12 10 mcg, B 1 5 mg, B 2 1 mg, B 6 1 mg, niacinamide 20 mg, panthenol 10 mg, choline 50 mg, inositol 20 mg, stebran conc 300 mg /cc Bot 30 cc

BECOPLEX FORTIFIED (Blue Line) Vit B 1 1 mg, B 2 2 mg, B 12 1 mcg, exc larrous sulfate 136 mg, dex liver 200 mg /Cap Bot 100s

Use: Orally, vitamin with liver & iron therapy

BE-COPLEX W/C (Cowley Pharm) Cap 100s

Use: Dietary supplement

BICOPEX W/IRON, (Blue Line) Ferrrous sulfate dried 1 gr, yeast 7 1/2 gr /Tab. Same as Becoplex with above ingredients added Bot 100s

Use: Orally (2 cc), dietary supplement

BECOTIN (illy) Thiamine chloride 10 mg riboflavin 10 mg, pyridoxine HCl 5 mg, nicotinamide 50 mg, pantothenic acid 25 mg, Vit B 12 1 mcg, liver preparation & stomach tissue material, des, 0.35 Gm /pulsule Bot 100s, 500s, 1000s

Use: Orally, vitamin B complex deficiencies
W/Vit C (illy) Thiamine chloride 10 mg, riboflavin 10 mg, pyridoxine HCl 5 mg, nicotinamide 50 mg, pantothenic acid 25 mg, Vit B 12 1 mcg, ascorbic acid 150 mg, liver preparation & stomach tissue material, des, D 35 Gm /pulsule Bot 100s 500s, 1000s

Use: Orally, vitamin B complex and vitamin C deficiencies.

BECOTIN-T. (Lilly). Vit. B-1, 15 mg.; B-2, 10 mg.; B-6, 5 mg.; nicotinamide, 100 mg.; pantothenic acid, 20 mg.; B-12, 4 mcg.; C, 300 mg.; liver preparation and stomach-tissue material, des., 125 mg./Tab. Bot. 100s, 1000s, 5000s.

Use: Vit. B-complex and vit. C deficiencies.

BEDADOL. (Ives-Cameron) Phenobarbital $\frac{1}{4}$ gr. with vitamin B complex/Tsp. or Tab. 100s. Elixir 1 pt.

BEDOCE. (Lincoln) Cryst. Vit. B-12 30 mcg., B-6 mg., 200 mcg., 1000 mcg./cc. Vial 5 cc. or 10 cc.

Use: I.M., vitamin B-12 therapy.

BEDOCE-GEL. (Lincoln) Vit. B-12 200 mcg. or 1000 mcg./cc. in 17% gelatin solution. Vial 10 cc.

BEDOCEPLIX. (Lincoln) Vit. B-1 20 mg., niacinamide 80 mg., Vit. B-2 2 mg., calcium pantothenate 2 mg., 4 mg., Vit. B-12 10 mcg./2 cc. Vial 30 cc.

Use: I.M., vitamin B complex therapy.

BE-DODEC. (Schleffelin) Vit. B-12 cryst, 15 mcg./1 cc. Amp. 1 cc. Box 6s.

Use: I.M., hemolapoletic.

BEDOXINE. (Massengill) Pyridoxine HCl. Amp. (50 mg.) 10 cc., 6s, 25s, 100s. Vial (100 mg.) 10 cc., Tab. (25 or 50 mg.), Bot. 100s.

Use: I.M., vitamin B-6 therapy.

BEECHWOOD CREOSOTE. See: Creosote, N.F.

BEECHWOOD CREOSOTE W/Guaiacol, aminoacids and polypeptides. See: Peptonooids-Creosote, Cough Prep. (Arlington)

BEE-GEL. (Myers-Carter) Vit. B-1 50 mg., B-2 1 mg., B-6 5 mg., niacinamide 50 mg., pantothenic acid 5 mg./cc. Vial 10 cc.

W/B-12 15 mcg./cc. Vial 10 cc.

W/Vit. C 25 mg./cc. Vial 10 cc.

BEE-12-GEL. (Myers-Carter) Vit. B-12 200 mcg. & 1000 mcg./cc. Vial 10 cc.

BE-EM-COTES. (Foul Maney) Sod. salicylate 6 gr., ascorbic acid 20 mg., calcium p-aminobenzoate 3 gr., menadiene 0.5 mg./Tab. Bot. 100s, 1000s.

Use: Orally (1-2 q. 3 h.), analgesic & antipyretic.

BESIX. (Testagar) Pyridoxine HCl 25 mg./Tab. Bot. 100s, 1000s

BEE STING THEBAPY. See Formo-Quieocaine (Fronswarth Lab.)

BEE VENOM SOLUTION, LYOVAC. (Merck Sharp & Dohme) Vial 1 cc. with 1 cc. of diluent equal to 10 stings.

Use: Desensitization to bee sting and arthritis.

BEFEREX. (Central) Iron-ammanium citrate

3.0 Gm., yeast-malt-rice bron ext. 1.6 ml., Vit. B-1 2 mg., Vit. B-2 4 mg., niacinamide 15 mg., Vit. B-6 1.5 mg., col. pantothenate 60 mg./fl. oz. Sol., 1 pt., 1 gal.

Use: Dietary supplement.

BEFOLEX. (Central) Folic acid 1.7 mg., ascorbic acid 30.0 mg., niacinamide 10 mg., ferrous sulfate 19.4 mg., Vit. B-1 1 mg., B-2 2 mg., B-6 0.5 mg., calcium pantothenate 1 mg., yeast ext. 150 mg./Tab.

Bot. 100s, 500s, 1000s.

Use: Orally, treatment of macrocytic anemias.

BEHEPAN. See: Vitamin B-12.

BEHEPARON HEMATINIC. (Kieney) Vit. B-1 $\frac{1}{2}$ mg., B-2 0.5 mg., pantothenic acid 0.2 mg., niacin 5 mg., Vit. B-6 0.2 mg., whole liver $8\frac{1}{2}$ gr., ferrous sulfate exsic. 3 gr./Cap. Bot. 100s, 1000s.

Use: Orally, vitamin with liver & iron therapy.

BEJECTAL. (Abbott) 4 cc. of solution No. 2 mixed with the amount of solution No. 1 in the large vial yields 10.6 cc. of a solution in which each 10 cc. contains: thiamine HCl 100 mg., riboflavin 20 mg., nicotinamide 750 mg., pyridoxine HCl 50 mg., calcium pantothenate 50 mg., benzyl alcohol, as a preservative 0.9%, water for injection, to make 10 cc. 10 cc. size (2 bottles consisting of solution No. 1 and solution No. 2), Box 6s.

Use: I.M., vitamin B complex therapy.

W/Vitamin C. (Abbott) Vit. B-1 20 mg., B-2 3 mg., B-12 2 mcg., niacinamide 75 mg., Vit. B-6 5 mg., C 100 mg., Sod. pantothenate 5 mg./cc. Vial 10 cc. Box 5s.

Use: Vitamin B complex deficiencies.

W/Liver. (Abbott) Thiamine HCl 10 mg., riboflavin 0.5 mg., niacinamide 10 mg., pantothenic acid 5 mg., choline 12 mg., in crude liver ext., & Vit. B-12 2 mcg., phenol 0.5% as a preservative/1 cc. Vial 10 cc. Box 6s.

Use: I.M., vitamin B complex therapy.

BEJECTAL IMPROVED W/VIT. C. (Abbott) Vit. B-1 200 mg., B-2 30 mg., B-12 20 mcg., niacinamide 750 mg., Vit. B-6 50 mg., C 1000 mg., Sod. pantothenate 50 mg./10 cc. Vial 10 cc.

Use: (1 cc. daily) Severe vitamin deficiencies.

BEJEX. (Abbott) Vit. B-1 10 mg., B-2 10 mg., B-12 25 mcg., nicotinamide 250 mg., B-6 5 mg., C 500 mg., sod. pantothenate 50 mg./5 mL. Uservial container 5 ml.

Use: I.M., I.V., Vitamin B complex supplement.

BELAKOIDS TT (Columbus) Phenobarbital sod. 486 mg., hyoscyamine sulfate 0.3111 mg., atropine sulfate 0.0582 mg., hyoscyne

hydrabamide 0.0195 mg /T Y Tab Bot
100s 500s

Use Antispasmodic

BELAP. (Hoeck) #0—Phenobarbital gr $\frac{1}{4}$
belladonna ext gr $\frac{1}{4}$ /Tab; #1—pheno-
barbital gr $\frac{1}{4}$, belladonna ext gr $\frac{1}{4}$ /
Tab; #2—phenobarbital gr $\frac{1}{2}$ belladonna
ext gr $\frac{1}{4}$ /Tab Bot 100s, 1000s

Use Orally sedative with antispasmodic
action

BELBABB (Haskell) Phenobarbital gr $\frac{1}{4}$
atropine hyoscyamine & scopolamine
equivalent to belladonna Ir 0.5 cc/Tab
Cap or fl dr of Elixir Tablets #2 con-
tain the same alkaloid content but with
phenobarbital gr $\frac{1}{2}$ Cap—Bot 100s
500s 1000s Elix—Bot 6 fl oz, 1 pl,
1 gal Tab—Tube 20s Bot 100s 500s
1000s Tab #2, Bot 100s 500s 1000s

Use Orally sedative & antispasmodic

BELBABB-B (Haskell) Phenobarbital $\frac{1}{4}$ gr,
atropine sulf 0.0240 mg, hyoscyamine HBr
0.1280 mg, hyoscyamine HBr 0.0072 mg, vit
B 1.5 mg, B 2.2 mg cal pantothenate 2.5
mg B 6 0.5 mg, niacinamide 10 mg, B 12
2 mg/tab Bot 100s 500s

Use (2-4 daily) Antispasmodic w/nutri-
tional supplement

BELBABB TRISULETS (Haskell) Phenobarbital
& belladonna alkaloids Bot 30s 100s

Use Antispasmodic sedative

BELBARMINE (Haskell) d Oetoxyspedrine
HCl 5 mg, phenobarbital 32 mg, hyoscy-
mine HBr 0.0072 mg, atropine sulf 0.024 mg
hyoscyamine HBr 0.128 mg/Tab Bot 100s
1000s 5000s

Use Treatment and control of obesity

BELBOTONE (Kendall) Tr belladonna 48
min, Sod bromide 60 gr, pepsin, pan-
creatin, dextrose lactic acid hydrochloric
acid, carminative aromatics & distilled
water q s /fl oz Bot 8 oz

Use Orally, mild sedative & antispasmodic

BELDASE (Finley) Pepsin NF, 200 mg s
charcoal (activated), 300 mg, hyoscyamine
sulfate 0.10 mg, atropine sulfate, 0.02 mg,
hyoscyamine HBr, 0.006 mg /Cap Bot 100s

Use Digestive disturbances an antispas-
modic

BELXON. (American Pharmaceutical) Vit
B-1 1.5 mg, B 2 1 mg, pantothenic acid
0.3 mg, Vit B 6 0.25 mg, niacin 5 mg
with yeast & liver/Cap Bot 100s, 500s,
1000s Vit B 1 3 mg, B 2 12 mg, ni-
acin acid 12 mg, & fresh liver/fl oz
Elixir Bot 1 pl, 1 gal

Use Orally dietary supplement

BELFER (Durst) B 12 5 mcg, B-6 1 mg, C 25

mg folic acid 0.2 mg, iron gluconate 50
mg, sorbitol 100 mg/Tab Bot 100s
Use: Anemia

BELGLYN, TAB. (Brayten) Dihydroxy alumi-
num aminoacetate 7.7 gr, belladonna alk
1/400 gr/Tab Bot 100s

Use: (1 or 2 p.c.h.s.) Antispasmodic, antacid

BELINTRON. (Amer Pharm Co) Liver trac-
tion #2 3 gr B 12 w/intrinsic factor conc.
 $\frac{1}{2}$ u, B 12 5 mcg, folic acid 1 mg, ferrous
sulfate 300 mg, B 1 2 mg, B 2 2 mg, C 75
mg, niacinamide 10 mg, copper proteinates
1 mg, betaine HCl 2 gr/cap Bot 60s

Use (2 daily) Anemias

BELIVIRON. (Horton and Converse) Vit B-1
10 mg, B 2 0.5 mg, B 6 1 mg, niacinamide
10 mg, pantothenyl alcohol 1 mg, ferrous
factors 20 mg, liver extract eq to 5 mcg,
B 12/cc Vial 10 cc

Use IM, B complex therapy, hematinic

BELLABIL. (Whittier) Dehydrocholic acid 250.0
mg, hyoscyamine sulfate 0.1037 mg, atro-
pine sulfate 0.0194 mg, hyoscyamine HBr
0.0065 mg, phenobarbital 16.2 mg/Tab
Bot 100s

Use Antispasmodic, sedative

BELLABULGARA. (Lederle) Ext Bulgarian
belladonna root standardized to yield to
0.4 mg of total alkaloids/Tab Bot 100s
1000s

Use Orally, antispasmodic

BELLADENAL. (Sandoz) Bellatoline 1/250
gr, phenobarbital $\frac{3}{4}$ gr/tab, or spars
tab Bot 100s

Elixir: (Bellafoline 1/1000 gr, phenobarbi-
tal 3/16 gr/4 cc) Bot 1 pl, 1 gal
Supp: (Bellafoline 1/120 gr, phenobarbi-
tal 1 1/2 gr) Box 6s, 30s

Use Orally, antispasmodic & sedative

BELLADONNA ALKALOIDS.

See: Bellat, Tab (VanPelt & Brown)
Frydon, Spansule (Smith, Kline & French)

Bellation, Tab (Merck Sharp & Dohme)

W/A.P.C., & pyranisamine. See: Pyr Aspro

Tab (VanPelt & Brown)

W/Aspirin, acetaphenolindin, & caffeine

See: Bellaspro Tab (VanPelt & Brown)

W/Chloral hydrate See: Felo Bal, Cap

(Fellows)

W/Chlorbutanal See: Bufadonna Cap

(Wampole)

W/Dihydroxy aluminum aminoacetate See

Belglyn Tab (Brayten)

W/Dihydroxy aluminum aminoacetate, phe-

nobarbital

See: Barbonate, Tab (Bristol)

Malglyn Camp, Tab (Bristol)

W/Dehydrocholic acid See: Decholin w/

Belladonna, Tab (Ames)

w/c

- bismuth oxyiodide, bismuth subcarbonate & balsam Peru wyanoidr, Supp. (Wyeth).
- W/Ergotamine tartrate, acetophenetidin & caffeine. See: Wigraine, Tab. (Orgonon)
- W/Ergotamine tartrate, phenobarb. See: Bellergal, Tab. (Sandoz)
- W/Phenobarb. See: Belladenal, Elix., Tab. & Supp., (Sandoz)
- W/Phenobarbital and vitamin B complex. See: Belbarb-B, Tab. (Haskell)
- ***BELLADONNA EXTRACT**, N.F., A.D.R.; Ointment, N.F.
- Belladonna Extract Combinations on p. 108.
- ***BELLADONNA LEAF**, U.S.P., Tr. U.S.P., A.D.R. 1955.
- W/Phenobarbital & caffeine. See: Alepsol, Tab. (Fougera)
- W/Phenobarbital & benzocaine. See: Gastrolic, Tab. (Mallord).
- W/Phenobarbital, stramonium & ephedrine sulfate. See: Acopnoid, Tab. (Chemico)
- W/Phenobarbital, theophylline, ephedrine HCl, pyrilamine maleate, potassium iodide, magnesium trisilicate. See: Theafed Pedalric, Tab. (Sutliff & Case)
- ***BELLADONNA LEAF FLUID EXTRACT**, N.F.
- W/Camphor & alcohol. See: Comphella. Modified (Pitman-Moore). Also w/ephedrine.
- BELLADONNA PLASTER**, N.F.
- W/Phenobarbital, kolin colloidal. See: Kophebel, Tab. (Elder).
- Belladonna Products and Phenobarbital Combinations on pp. 109 and 110.
- ***BELLADONNA TINCTURE**, U.S.P., A.D.R. 1954. See: Toladonna, Tab. (Rexall).
- W/Mucilage, licorice. See: Vanul, Liq. (Vanguard Pharmaceuticals)
- W/Sod. bromide, pepsin, pancreatin, dextrase, lactic acid, hydrochloric acid. See: Belbratone, Tr. (Kendall)
- BELLADONNA, TRISILICATE & CHARCOAL**. (Raymet) Belladonna ext. 1/12 gr., magnesium trisilicate 7 1/2 gr., activated charcoal 1 gr./Tab. Bot. 100s, 500s, 1000s.
- Use: Orally, antispasmodic & gastric antacid.
- BELLAFOLINE**. (Sandoz) Levo-hyoscyamine maleate. Amp.—(0.5 mg.) 1 cc. injection, Tab.—(0.25 mg.) Oral use, Bot. 100s, 1000s
- Use: Antispasmodic.
- W/Pentobarbital cod. & catergol. See: Cotergal-PB, Supp. (Sandoz)
- W/Phenobarbital. See: Belladenal, Elix., Supp., Tab. (Sandoz)
- BELLAL**. (Vanpelt & Brown) Belladonna alkaloids 0.13 mg (equivalent to belladonna tr. min. 7)/Tab. Bot. 100s, 500s, 1000s.
- Use: Orally, antiriparmonic.
- BELLASPRO**. (Vanpelt & Brown) Acetylcholinic acid 3 1/2 gr., acetophenetidin 2 1/2 gr., caffeine 1/2 gr., belladonna alkaloids (equivalent to Tr. belladonna min. 3.8)/Tab. Bot. 100s, 500s, 1000c.
- Use: Orally (1-2 t.i.d.), antiriparmonic with analgesic & antipyretic action.
- W/Codeine. (Vanpelt & Brown) Tablets # 1 contain in addition to Bellaspro tablets, codeine phosphate 1/4 gr./Tab. Tablets # 2 contain in addition to Bellaspro tablets, codeine phosphate gr. 1/2/Tab. Bot. 100s.
- Use: See Bellaspro
- BELLATURIC**. (Ethical Pharm) Hyoscyne HBr 0.0065 mg, hyoscyamine sulfate 0.1037 mg, atropine sulfate 0.0194 mg, phenobarbital 16.2 mg./Cap., Tab. or 5 cc. Cap. & Tab., Bot. 100c; elixir, Bot. 1 pl.
- Use: Orally, antispasmodic.
- BELLATURIC T-D**. (Ethical Pharm) Phenobarbital 50 mg., atropine sulf. 0.0582 mg., hyoscyamine sulf. 0.311 mg., hyoscyne HBr 0.0195 mg./T.D. Cap. Bot. 30c
- Use: Sedative and antispasmodic
- BELLERGERAL**. (Sandoz) Bellafoline 0.1 mg., ergotamine tartrate 0.3 mg., phenobarbital 200 mg./Tab. Bot. 100c, 500c, 1000c, Space-Tabs., Bot. 100c
- Use: Orally (1), sedative w/antispasmodic.
- BELLIRON**. (Harvey Lab) Liver Ext. 2 mcg., ferrous lactate 20 mg., Vit. B-1 10 mg./cc. Vial 30 cc.
- BELLKAGEL**. (Ferndale) Liq. Bot. 8 oz.
- Use: Treatment of bacillary dysentery.
- BELNESIUM**. (Haskell) Phenobarbital 1/2 gr., atropine sulfate 0.01 mg., hyoscyamine hydrobromide 0.05 mg., magnesium trisilicate 7 gr./Tab. Bot. 50s, 250s, 500c, 1000c.
- Use: Orally (1-2 t.i.d.), antiriparmonic & gastric antacid.
- BELPHEDRIBARB**. (Haskell) Ephedrine sulfate 3/4 gr., phenobarbital 1/2 gr., belladonna alk. equivalent to belladonna tr. 0.5 cc./Tab. Bot. 50s, 500s.
- Use: Orally (1-2), sedative & antiriparmonic action.
- BELSAL**. (Setliff & Case) Benzoic acid 10%, salicylic acid 5%, benzocaine 1%, thymol 2%, lanolin, petrolatum, perfume, color q.s. Oint. 1 lb.
- Use: Fungicidal
- BELVIS**. (Carnick) Guar cellupectinoid 275 mg., magnesium oxide 70 mg., aluminum hydroxide gel 80 mg., magnesium trisilicate 160 mg./Tab. Bot. 100s.
- Use: Treatment of peptic ulcer.
- * Generic name recognized by N.F., U.S.P. or A.M.A.

Belladonna Extract Combinations

| Belladonna Extract | Others | Manufacturers | Brand |
|--------------------|--|---------------------|----------------------------|
| 0.16 mg | Acetate 1.33 mg, hyoscyne 1.33 mg | Leamon | Acabryanna, Tab |
| 1/20 gr | Acetate 1/10 gr., ammonium chloride 1/2 gr., camphor 1/2 gr., quinine sulfate 1/2 gr. | Merck Sharp & Dohme | Caryza, Tab. |
| 1/12 gr | 1/2 gr. quinine sulfate 1/2 gr. | Hance | Apco Pill |
| 1/12 gr | Alain 1/2 gr., ipacat 1/15 gr., phenolphthalein 1/2 gr. | Amer Pharm | Alkoni Pill |
| 1/12 gr | Alain 1/2 gr., ipacat 1/15 gr., phenolphthalein 1/2 gr., strychnine sulfate 1/80 gr. | Mallard | Alotkol, Tab |
| 1/12 gr | Alain 1/2 gr., catara 1/2 gr., strychnine 1/120 gr. | Horton & Converse | A. S. & C., Tab |
| 1/2 gr | Alain 1/2 gr., catara 1/2 gr., phenolphthalein 1/2 gr. podophyllin 1/2 gr. | BuFington | Catecaphen, Tab |
| 1/2 gr | Alain 1/2 gr., pectic 4 mg., phenolphthalein 32 mg., strychnine sulfate 0.5 mg., podophyllin 4 mg., capsicum oleoresin 2.6 mg. | Drug Prod | Alipco Tab |
| 3.4 mg | Anarbarital 32 mg., d. amphetamine sulf 5 mg. | Kendall | G. I. Iron Tab |
| 1.6 mg | 3. methyl subcarb 14%, Sod bicarb 25%, toatin 58% | Zerec | Kaacorb Pow |
| 1/2 gr | Bulbarbital sod 30 mg. | Denab | Insabel TD, Tab |
| 30 mg | Bulbarbital 16 mg., Methylpulsioxone 15 mg. | Flough | Sibel Tab |
| 1.6 mg | Bulbarbital 16 mg. | McNeil | Bulbarb, Tab, Elix |
| 1.5 mg | Bulbarbital 16 mg. | Merck Sharp & Dohme | Coscara Comp., Tab |
| 3 gr | Castora seg. oil 15 mg., oila 15 mg., podophyllin 10 mg., ginger oleoresin 4 mg. | Jenkins Labs | Donnafed Jr., Tab |
| 1/20 gr | Ephedrine HCl 1/16 gr., salicylamide 1/2 gr. | Upjohn | Malcatab w/Belladonna, Tab |
| 1/16 gr | Mag trisulfate 2.5 gr., aluminum hydroxide 5 gr. | Forest | Trialdin, Tab |
| 5 mg | Mag trisulfate 0.3 Gm., aluminum hydroxide 0.15 Gm. | Forest | Trisdonna, Tab |
| 1/12 gr | Mag trisulfate 7 1/2 gr., charcol 1 gr. | Taitby Naton | Lairdell, Liq & Tab |
| 5 mg | Mephensin 200 mg., nicotinic acid 25 mg. | McNeil | Careni, Tab |
| 7.5 mg | Methapyrilene HCl 25 mg., rac desoxyephedrine HCl 1.25 mg. | Abbott | Nembudonna, Elix |
| 10 mg /isp | Nembutal Sod 15 mg /isp | Abbott | Nembudonna, Cap # 1 |
| 1/6 gr. | Nembutal Sod 1/4 gr. | Abbott | Nembudonna, Cap # 2 |
| 1/6 gr | Nembutal Sod 1/4 gr. | Elder | Donobarb, Tab |
| 1/6 gr | Phenobarbital 1/2 gr., aluminum hydroxide 5 gr. | Hoack | Alubelap, Tab |
| 1/46 gr | Phenobarbital 1/2 gr., ethoverene HCl 20 mg. | Armour | Tri Syrac, Tab |
| 4.1 mg | Phenyltoloxamine 20 mg., ethoverene HCl 20 mg. | Superior Pharm. | Coryza, Tab |
| 1/20 gr | Quinine sulf 1/2 gr., amm chlor 1/2 gr., camphor 1/2 gr., acetate | Superior Pharm. | Cephe's Tab |
| 1/50 gr | Sod salicylate 2 gr., phenacetin 2 gr., camphor 1/10 gr. col 1/2 gr., capsicum 1/10 gr. | | |

Belladonna Products and Phenobarbital Combinations

| Belladonna | Pheno- barbital | Others | Manufacturer | Brand |
|----------------------|---------------------|---|--------------------|------------------------------|
| Alkaloïds 0.13 mg. | ¼ gr. | Aluminum hydroxide 250 mg., mag. trisilicate 250 mg., benzocaine 8 mg. | Vonpelt & Brown | Barbidonna, Elixir & Tab. |
| Alkaloïds | ¼ gr. | Aspirin 180 mg., acetophenetidin 120 mg. | Robins | Dannetel, Cap., Elixir, Tab. |
| Alkaloïds | ¼ gr. | Aspirin 5 gr., caffeine ¼ gr. | Merk Sharp & Dohme | Tranidon, Elixir |
| Tr. 0.5 cc. | ¼ gr. | Benzocaine 2 gr. | Haskell | Belbarb, Cap., Elixir & Tab. |
| Ext. ½ gr. | ¼ gr., ¼ gr., ½ gr. | Benzocaine, aluminum hydroxide, Cal. carb., Mag. oxide, Mag. trisilicate | Heack | Belap, Tab. |
| Ext. 1/6 gr. | Sod. ¼ gr. | Cal. carb. 6 gr., Mag. trisilicate 3½ gr., chloro- phyll 1% | Meyers-Rasser | Pheno-Donna No. 2, Tab. |
| Tr. ½ gr. | ½ gr. | Benzocaine 1/16 gr., aluminum hydroxide 2½ gr., Cal. carb. 6 gr., Mag. trisilicate 3½ gr., chloro- phyll 1% | Rozall | Teladonna, Tab. |
| Ext. ¼ gr. | 8 mg. | Benzocaine 8 mg., Mag. trisilicate 250 mg., alumi- num hydroxide 250 mg. | Walker | Alkospasmol, Tab. |
| Alkaloïds 0.0265 mg. | 15 mg. | Caffeine 2/5 gr. | Rarer | Foodon, Tab. |
| Ext. ½ gr. | ¼ gr. | Chorcoal 1 gr. | Elder | Ansodyne, Tab. |
| Lvs. ½ gr. | ¼ gr. | Dihydroxy aluminum aminoacetate 0.5 Gm. | Mallard | Castrolic, Tab. |
| Ext. 1/6 gr. | ¼ gr. | Ephedrine sulfate ¼ gr., stramonium 1 gr. | Smith-Dorsey | Al-Si-Cat, Pow. & Tab. |
| Ext. 1/24 gr. | 1/16 gr. | Ergotamine tartrate 0.3 mg. | Warren-Teed | W.T. Tab. |
| Ext. 1/6 gr. | ¼ gr. | Gastric mucin 160 mg., mag. glycinate 250 mg., aluminum hydroxide gel 250 mg. | Smith-Dorsey | Simaphen, Pow. |
| Alkaloïds ¼ gr. | 8 mg. | Hyoscyamus ext. ¼ gr. | Walker | Alkospasmol, Tab. |
| Pow. ½ gr. | 1½ gr. | Koalin 7½ gr. | Faugera | Aleptol, Tab. |
| Ext. 1/12 gr. | ½ gr. | Ox bile salts 3½ gr. | Rorer | Chardonne, Tab. |
| Alkaloïds 0.162 mg. | ¼ gr. | Pentobarbital Sod. 3/16 gr. | Broyten | Malglyn Camp., Tab. |
| Tr. 0.5 cc. | ½ gr. | | Hastell | Belphedribarb, Tab. |
| Pow. 1/10 gr. | ½ gr. | | Chemica | Acapnoid, Tab. |
| Alkaloïds 0.1 mg. | 20 mg. | | Sandaz | Bellergal, Tab. |
| Ext. 5 mg. | 6 mg. | | Barrows Chem. Co. | Mucagai-8, Tab. |
| Ext. ¼ gr. | ¼ gr. | | Meltra | Sposticen, Tab. |
| Root, ¼ gr. | ¼ gr. | | Elder | Kaphebel, Tab. |
| Ext. 1/24 gr. | ¼ gr. | | Smith-Dorsey | Panazolin, Tab. |
| Ext. ¼ gr. | 3/16 gr. | | Kirkman | Duadon, Tab. |

Belladonna Products and Phenobarbital Combinations

| Belladonna | Pheno- barbital | Others | Manufacturer | Brand |
|---------------------|--------------------|-----------------------|--------------|-------------------------------|
| Alkaloids 1/600 gr | 1/4 gr | Sulfur 1/8 gr | Hart Drug | Taludan, Tab |
| Ext 1/6 gr | 1/4 gr | Toto Diosase 2 1/2 gr | Parke Davis | Bardase, Tab |
| Alkaloids 2 mg. | 1/4 gr. | Vitamins | Wyeth | Beplete w/Bellad., Cop & Liq |
| Alkaloids 0.25 gr | 5 mg | Vit B complex | Ives Comaran | Donosedo, Elixir & Tab |
| Bellatrine 1/200 gr | 1/4 gr | | Sandoz | Belladenol, Elixir Supp & Tab |

* Generic name recognized by N.F., U.S.P. or A.M.A.

* BEMEGRIDE, N N O 1962 beta Ethyl beta methylglutarimide

See: Megimide (Abbott)

Mikedimide (Parroy)

BEMEX. (Malford) Vit B 1 100 mg, B 2 2 mg, B 6 4 mg, niacinamide 100 mg panthelol 4 mg/cc Vial 30 cc

Use IM vitamin B therapy

BEMINAL FORTE (Ayerst) Vit B 1 300 mg B 2 30 mg B 6 50 mg pantothenic acid 50 mg niacin 700 mg/10 cc Vial 10 cc

Use Injectable vitamin B complex

W/Vitamin C (Ayerst) Thiamine HCl 250 mg riboflavin 12.5 mg, nicotinamide 50 mg, pyridoxine HCl 3 mg, calcium pantothenate 100 mg, Vit C 250 mg, Vit B 12 w/intrinsic factor conc 1/9 u/cap Bot 100s 1000s

Use Orally, vitamins B & C

BEMINAL WITH IRON ELIXIR, (Ayerst) Vit B 12 10 mcg, thiamine 25 mg, riboflavin 12.5 mg niacinamide 50 mg, pyridoxine 1 mg D panthelol 10 mg, iron pyrophosphate 400 mg D sorbitol 7.87 Gm, alcohol 15%/20 cc

Use Dietary supplement

W/Iron & Liver (Ayerst) Vit B 1 1 mg, B 2 2 mg, B 12 2 mcg, B 6 0.5 mg, nicotinamide 10 mg, calcium pantothenate 1 mg yeast 100 mg, liver fraction II 100 mg ferrous carbonate 200 mg/Cop Bot 100s 1000s

Use Orally, in certain anemias

BEMINAL TABLETS, (Ayerst) Thiamine HCl 50 mg, riboflavin 30 mg, nicotinamide 100 mg, pyridoxine HCl 0.2 mg, calcium pantothenate 0.5 mg, lactic acid 0.25 mg brewers' yeast 1650 mg/Tab Bot 50s 100s 1000s

Use Vitamin B complex deficiencies

W/Phenobarbital (Ayerst) Thiamine HCl 5 mg, riboflavin 2 mg, nicotinamide 10 mg Vit B 12 1 mcg, pyridoxine 0.5 mg, D panthelol 2 mg, ascorbic acid 25 mg phenobarbital 16 mg/Tab Bot 100s, 500s

Use (1 tab 11d) Sedative w/vitamin B complex therapy

BEMOCIN, (Ayerst) Thiamine HCl 250 mg riboflavin 12.5 mg nicotinamide 750 mg Vit C 2000 mg, B 12 30 mcg, folic acid 0.67 mg, pyridoxine HCl 10 mg, calcium pantothenate 100 mg, dl methionine 400 mg, inositol 400 mg, choline bitartrate 600 mg/cap Bot 100s, 1000s

Use: Orally, dietary adjunct in alcoholism

BEMOTINIC CAPSULES. (Ayerst) Ferrous sulfate extic. 300 mg., vit. B-12 w/intrinsic factor concentrate $\frac{1}{2}$ u., folic acid 1.0 mg., B-1 10 D mg., vit. C 100 mg./cap. Bot. 100s, 1000s.

Use: Orally, deficiency anemia.

BEMOTINIC LIQUID. (Ayerst) Ferric ammonium citrate 200 mg, Vit. B-12 4 mcg., gastric tissue ext. 450 mg, folic acid D.33 mg., thiamine HCl 1.5 mg., riboflavin 1.0 mg., pyridoxine HCl D.2 mg./5 cc. Bot. 1 pl., 1 gal.

Use: (1-2 tsp) Nutritional anemias.

BEMULTRA. (Merck Sharp & Dohme) Vit. A 25,000 u., Vit. D 2000 u., thiamine HCl 15 mg., riboflavin 10 mg., pyridoxine HCl 1 mg., Vit. B-12 10 mcg., calcium pantothenate 10 mg., niacinamide 150 mg., ascorbic acid 150 mg., mixed tocopherols 10 mg./Cap. Bot. 100s

Use: Orally, high potency, therapeutic multivitamin.

BENA-13 ELIXIR. (Crystal) Vit B-12 25 mcg, Vit. B-1 10 mg./5 cc. Liq. Bot. pl.

Use: Appetite stimulant.

BENACINE. (Parke, Davis) Benadryl HCl 25 mg., hyoscine HBr 0.325 mg./Tab. Bot. 100s.

Use: (1 p.r.n.) Motion sickness.

BENACOL. (Seydel) Not listed Benzocaine benzoate 5%, benzyl alcohol 5%. Ampuls, 3 & 5 cc.

Use: Local anesthetic.

* **BENACTYZINE HYDROCHLORIDE, NND.** 1962. 2-Diethylaminoethyl benzilate (Nutrilin) Cevonal, Luclid, Nutinal.

See: Phobex, Tab. (Lloyd, Dabney & Westerville)

Suavifil, Tab. (Merck Sharp & Dohme)

W/Meprabamate See: Deprol. (Wallace)

BENADEX. (Fuller) Benzocaine 2%, aluminum acetate 2%, homamolis ext. 2%, phenol 1%. Tube 1 oz. Jar 1 lb.

Use: Hemorrhoids.

BENADRYL W/Aminophyllin. See: Hydryllin Preps. (Searle)

W/Chlorotheophyllinate. See: Dramamine. (Searle)

BENADRYL COUGH PREP. See: Benylin Expectorant (Parke, Davis).

BENADRYL HYDROCHLORIDE, NND 1962. (Parke, Davis) Diphenhydramine hydrochloride. 2-(Benzohydroxy)-N,N-dimethyl-ethylamine hydrochloride.

* Generic name recognized by N.F., U.S.P. or A.M.A.

Amp. (50 mg./cc) 1 cc. Box 10s.

Cap.: (25 mg.) Bot. 100s, 1000s.

Creom: (2%) Tube 1 & 2 oz.

Elixir: (10 mg./4 cc.) Bot. 1 pl., 1 gal.

Emplets: (50 mg) Bot. 100s, 1000s.

Kap seals: (50 mg) Bot. 100s, 1000s.

Pow.: Vial 14.17 Gm.

Sol.: (10 mg./cc) 10 cc., 30 cc.

Use: Antihistaminic.

W/Ambadryl HCl, dihydrocaineane, ammanium chloride & other caught agents
See: Ambenyl, Syr. (Parke, Davis).

W/Aminophylline See: Hydryllin (Searle)

W/Ammonium chloride, sod citrate, chloroform, menthol, chloroform. See: Benylin Expectorant. (Parke, Davis)

W/8-Chlorotheophylline. See: Dramamine (Searle)

W/Ephedrine sulfate. (Parke, Davis) Benadryl hydrochloride 50 mg., ephedrine sulfate 10 or 25 mg./Cap. Bot. 100s, 1000s.

Use: Provides the complementary effects of the original antihistaminic & the sympathomimetic.

W/Ephedrine hydrochloride & chloretane
See: Benadryl, Sol. (Parke, Davis)

W/Hyoscine HBr. See: Benacine, Tab. (Parke, Davis)

W/Zirconium carbonate hydrated. See: Zira-dryl, Cream (Parke, Davis).

BENADRYL HCl LOTION OR CREAM. See: Caladryl (Parke Davis)

BENA-FEDRIN. (Parke, Davis) Benadryl HCl D 1%, ephedrine HCl 1%, chloretane 0.5%. Bot. w/dropper, 1 oz.

Use: Nasal congestion.

* **BENANEEBIN HCl.** 3-(2-aminoethyl)-1-benzyl-5-methoxy-2-methylindole HCl.

BENAPEN. See: Benethamine

BENASEPT. (Blue Line) Benzalkonium chloride 1.1000 Sol., stainless. Bot. 1 pl., 1 gal. Tincture 1.1000 (alcohol 50% & acetone 10%). Bot 1 pt., 1 gal. Vaginal Jelly W/ applicator, Tube $1\frac{1}{2}$ oz.

Use: Topically, cationic germicidal.

BENAT W/B-12 TABLETS. (National Drug) Whole liver conc. 100 mg., Vit. B-12 conc. 10 mcg., B-1 5 mg., B-2 2 mg., nicotinamide 50 mg., Cal. pantothenate 5 mg., Vit. B-6 1 mg./Tab. Bot. 24s, 100s.

Use: Orally or I.M., vitamin therapy.

W/B-12 injectable. (National Drug) Vit. B-1 10 mg., Vit. B-2 2 mg., nicotinamide 50 mg., pantothenyl alcohol 1 mg., Vit. B-6

BEN-CAINE B.B.

112

- 1 mg, Vit B 12 30 mcg, liver inj 0.5 ml / cc Vial 10 cc
 Use I.M. vitamin B complex
- BEN CAINE B B** (Ulmer) Benzocaine 9%, n butylaminobenzoate 9% Tube w/applca for 1 oz Jar 1 lb
 Use Topically local anesthetic
- BEN CO FED** (Ode Peacock) Codeine phosphate 1 gr, ephedrine sulfate $\frac{1}{4}$ gr benzyl alcohol 7.47 min, sorbital 180 gr nannitol 45.3 gr / fl oz Bot 8 oz
 Use Orally (1 dr q 3-4 h) relief of coughs
- BENEDICTIN** (Merrell) Bently HCl 10 mg decopryn succinate 10 mg, pyridoxine HCl 10 mg / Tab Bot 100s
 Use Nausea and vomiting of pregnancy morning sickness
- * **BENDOBFLUMETHIAZOLE, N N O 1962 3** Benzyl 3:4 d hydro 7 sulphamoyl 6 trifluoro methyl benzo 1 2 4 thio d oxine 1 1 d oxide
 See Naturel n (Squ bb)
- W/Potassium Chloride See Naturel n W/K (Squ bb)
- W/Zauwolfia serpent na See Roultra X N (Squ bb)
- BENECAP**, (Bentex Pharm) Vit A 5000 u, D 1000 u B 1 1.5 mg, B-2 2 mg, B-6 0.1 mg, C 37.5 mg, Cal pantothenate 1 mg, nicotinamide 20 mg / cap Bot 100s 1000s
 Use: Multiple vitamin therapy
- BENEDONNA** (Bentex Pharm) Hyoscyamine sulfate 0.1037 mg, atropine sulfate 0.0194 mg, hyoscyne HBr 0.0065 mg, phenobarbital $\frac{1}{4}$ gr./tab Bot 100s 1000s
 Use: Antispasmodic with sedative
- BENEDRIN** (Bene) Isopropylarterenal HCl 10 mg, ethoverline HCl 30 mg, theophylline 125 mg, butobarbital 10 mg / Tab Bot 100s 500s
 Use: Asthma and hay fever
- BENELIXIR**, (Bentex Pharm) Vit A 4500 u, B-1 1 mg, B-2 0.75 mg, C 60 mg, D 1000 u, niacinamide 10 mg / 6 cc. Bot 8 oz
 Use: Dietary supplement
- BENEMID, NND 1962** (Merck Sharp & Dohme) probenecid 0.5 Gm / Tab Bot 100s 1000s
 Use: Orally (1 daily/wk) increase excretion of uric acid
- V/Calch cne See: Calbenemid Tab (Merck Sharp & Dohme).
- V/Pot penicillin G See: Bemonden, Prep (Merck Sharp & Dohme)
- * Generic name recognized by N.F., U.S.P. or A.M.A.
- BENENATAL** (Bentex Pharm) Vit A 1333 u D 133 u B 1 0.33 mg, B 2 0.667 mg C 16.6 mg, niacinamide 3.3 mg, dicalcium phosphate anhydrous 210 mg, ferrous sulfate dried 12 mg / cap Bot 100s, 1000s
 Use Vitamin and mineral supplement
- BENENATAL THERAPEUTIC**, (Bentex Pharm) Vit A 2000 u, D 333 u, B 1 1.5 mg, B-1 mg, B 12 1 mcg, folic acid 0.5 mg, Vit C 35 mg, niacinamide 7 mg, ferrous sulfate dried 12 mg, dicalcium phosphate anhydrous 261.9 mg, Cal gluconate 15 mg / cap Bot 100s, 1000s
 Use Dietary supplement
- BENIRON** (Bentex Pharm) Liver desic. 200 mg ferrous sulfate dried 136.1 mg, Vit B 1 1 mg, B 2 2 mg, B 12 1 mcg / Cap Bot 100s 1000s
 Use: Dietary supplement
- * **BENITHAMINE** N Benzyl 2 phenylethyl amine salt of benzylpenicillin Benopen
- BENETRYCIN** (Supreme) Tyrothricin 2 mg, benzocaine 5 mg / Troche Bot 15s Cold Caps Bot 24s Syr 4 oz, 1 gal
 Use Mouth & throat infections
- BENEVITE**, (Bentex Pharm) Vit A 5000 u D 1000 u B 1 2.5 mg, B-2 2.5 mg, B 6 0.5 mg C 40 mg, niacinamide 20 mg, Cal pantothenate 5 mg, folic acid 0.5 mg, Vit E 2 mg / cap Bot 100s, 1000s
 Use Multiple vitamin therapy
- BENGAL GELATIN** See: Agar
- BENHY**, (Sulfin & Case) Benzocaine 65 mg hydrastis 65 mg, tannic acid 65 mg, boric acid 65 mg, thymol iodide 16 mg, bismuth subiodide 32.5 mg, eucalyptol q.s. / SUPP Box 12s
- BENIZOL**, (Various Mfr) Pentylene tetrazol 100 mg, nicotinic acid 50 mg / Tab or 5 cc. Bot 100s 1 pt
 Use: 1 2 tabs or 1 tsp 1 d; in senile cases to restore sense of responsibility
- BENIO**, (Strassenburgh) Benzyl benzoate 25%, benzyl alcohol 2%, emulsion Bot 6 oz, 1 pt, 1 gal
 Use: Topically, scabicide
- BENODAMINE HYDROCHLORIDE, NND 1959** (Merck Sharp & Dohme) Piperaxan HCl 2% sol in saline for IV inj AMP 10 cc
 Use: Diagnostic agent for epinephrine producing tumors
- BIOQUIN, NND 1962** (Elder) Monobeneyl ether of hydroquinone (Monobenzone)

Oint. 20%, Tube 1/4 oz., Jar 1 lb. Lotion 5%, Bot. 4 oz.

Use: Topically, hyperpigmentation.

***BENOXINATE HYDROCHLORIDE**, U.S.P. N.N.D. 1962. Ophth. Soln. U.S.P. 2-diethylaminoethyl-4-amino-3-n-butoxy benzoate HCl. See Dorsacaine HCl Prep. (Smith-Dorsey)

BENSULFOID, (Paythress) 33% elemental sulfur by weight. Pow. 1 oz. Tab. Bot. 100s, 500s.

Use: Acne, seborrhea, etc.

BENSULFOID LOTION, (Paythress) Colloidal sulfur 2%, resorcinol 2%, thymol 0.5%, hexachlorophene 0.5%, zinc oxide 6%, alcohol (by vol.) 12%, perfume, cosmetic colors, emulsifiers./Lotion Bot. 2 oz.; Powder Bot. 1 oz.

Use: Topical treatment of acne.

BENTHAMINE, (Bentex Pharm.) Thiamine 1 mg., phenobarbital 1/2 gr./tab. Bot. 100s, 1000s.

Use: Sedative

BENTYL HCl, N.N.D. 1962, (Merrell) Dicyclamine HCl. (Diethylamine carbethoxybicyclohexyl HCl). Cap. 1/16 gr. Bot. 100s, 500s, 1000s. Syr. 1/16 gr./5 cc. Bot. 1 pt., 1 gal. Inj. 10 mg./cc. Amp. 2 cc. Vial. 10 cc.

Use: Antispasmodic.

W/Dacryl succinate, pyridoxine HCl. See Benedictin, Tab. (Merrell)

W/Mag. oxide, aluminum hydroxide & methylcellulose. See Kalantyl Gel, Tab. (Merrell).

W/Phenobarbital (Merrell). Cap. Bentlyl HCl 1/6 gr. or 1/2 gr., phenobarbital 1/4 gr. Bot. 100s, 500s. Syr. Bentlyl HCl 1/6 gr., phenobarbital 1/4 gr./5 cc. Bot. 1 pt., 1 gal.

Use: Antispasmodic & sedative.

W/Phenobarbital repeat action. (Merrell) Bentlyl HCl 10 mg., phenobarbital 15 mg. & bentlyl HCl 10 mg. In an enteric case. Tab., Bot. 100s, 500s

BENTYSON, (A.P.C.) Tyrothricin 1 mg., neomycin sulf. 5 mg., benzocaine 5 mg./Loz. Vial 12s.

Use: Minor sore throats.

BENULAX, (Merrell). Dicyclamine HCl, 15 mg; sodium lauryl sulfate, 30 mg; modified polyglucose gum, 27 Gm; casanthranol, 90 mg / 15 cc. Bot. 6 oz.

Use: Constipation, colitis.

BENULONE, (Merrell). Dicyclamine HCl, 15 mg; sodium lauryl sulfate, 30 mg; modified polyglucose gum, 27 Gm; casanthranol, 90 mg / 15 cc. Bot. 6 oz.

* Generic name recognized by N.F., U.S.P. or A.M.A.

Red polyglucose gum, 2.7 Gm./15 cc. Bot. 12 oz.

Use: Chronic constipation.

BENYLATE, N.N.D. 1962. (Braun) Benzylbenzoate 25%, emulsified. Bot. 4 fl. oz., 1 pt., 1 gal.

Use: Topically, scabicides & pediculocides.

BENYLIN EXPECTORANT, (Parke, Davis) Benadryl HCl 80 mg., ammonium chloride 12 gr., Sod. citrate 5 gr., chloroform 2 gr., menthol 1/10 gr., alcohol 5%/fl. oz. Bot. 1 pt., 1 gal.

Use: Orally, coughs.

***BENZALKONIUM CHLORIDE**, U.S.P., N.N.D. 1962. Sol., U.S.P., A.D.R. (Winthrop Labs.) Alkyl. dimethyl-benzylammonium chloride, zeohrol. Sol. 1:5000. (Alcan) Droptainer, Ophth. 15 cc.

See Benzsept, Sol. & Tr. (Blue Line).

Germicin (Consolidated Mid.)

Hyamina 3500 (Rohm & Haas)

Hyva Yog., Tab. (Holland-Rantos)

Pheneene Germicidal Solution Tincture.

(Ulmer)

Reccal (Winthrop Labs.)

Zephiren Chloride. (Winthrop Labs.)

Use: Surface antiseptic

W/Allantoin, di-desoxyephedrine HCl. See: Norekon, liq. (Chesebrough).

W/Bacitracin zinc, polymyxin B, neomycin sulf. See: Biotres, Oint. (Central)

W/Cetyl dimethyl ethyl ammonium bromide, isopropyl alc., sod. nitrite. See: Cetylclide, Soln. (Cetylite Inc)

W/Glycerin. See: Larakon, Sol. (Baybank).

W/Glycine, ammonium gluconate, ammonium salicylate, caffeine, terpin hydrate. See: Martuslin Special, liq. (M. R. Thompson).

W/Phenylmercuric acetate, sulfacetamide, 9 aminoacridine HCl. See: Kenhex, Vaginal Prep. (Kendall)

W/Polymyxin B, neomycin sulfate, zinc bacitracin. See: Biotres, Oint. (Central Pharmaceutical)

W/Pyrilamine maleate, phenylephrine HCl methylparaben, propylparaben.

See: Detergel Aromatic Spray (Bruce)

W/Pyrilamine maleate, zincum oxide & p-chloro-m-xyleneol. See: Rhucen, Oint. (Central Pharm.)

W/Tyrothricin & Isobutyl-p-an-nob... See: Cyclo-Thrycin, Loz. (Pace)

BENZAPAS, (Smith-Dorsey, & Bent... Calcium benzoyl-p-aminonitric...

Bot 4 Gm Pk & 1 lb Tabs 0.5 Gm Bot 100s

Use (10 to 15 Gm daily) Tuberculosis

BENZARSOL (Pittman Moore) Polybenzarsol Poly(methylene 4 hydroxybenzenearsonic acid)

Use: Treatment of amoebiasis

***BENZATHINE PENICILLIN G, USP**
NND 1962 ADR NND benzylethylene diamine di(benzylpenicillin) Benzethacil Dibencil Penidural See Penicillin G Benzathine USP

BENZAZOLINE HYDROCHLORIDE. Toluazoline HCl USP NND 1962 See Prisoline Prep (Ciba)

BENZCARBIMINE See Duolar (Armour)

***BENZCHLORPROPAMIDE** See Hibicon (Lederle)

BENZEDRER INHALER, NND 1962 (Smith Kline & French) Propylhexedrine 250 mg & aromatics Single plastic tubes

Use Relieves congestion of head colds allergic rhinitis & sinusitis

BENZEDRINE, NNR 1953 (Smith, Kline & French) Racemic amphetamine Sol 1% Bot 1 R oz 12 R oz

Use: Instilled for nasal decongestion & vasoconstrictor

W/Cocaine sulfate aspirin phenacetin See Edrisal with Codeine Tab (Smith, Kline & French)

BENZEDRINE SULFATE, NND 1958 (Smith Kline & French) Racemic amphetamine sulfate Sol Amp (20 mg) 1 cc Sponules 15 mg Bot 30s Tab 5 mg, 10 mg Bot 100s, 1000s

Use: Effective in symptomatic treatment of many mild psychogenic depressive states

***BENZENE HEXACHLORIDE, GAMMA, USP**, NND 1962 Pow, Sol, Oint, USP Gamma benzene hexachloride hexachlorocyclohexane lindane

Fidelity Lab—Pow 50%, Pkg 1 lb, 5 lb Imperial—Pow 50%, Pkg 1 lb, 4 lb 12%, Pkg 1 lb, 4 lb

See Gamiso Cream (Texas Pharmacal) Gexane Oint & liq, NND (Strasbourg)

Kwell, Cream, Oint & Shampoo, NND (Commercial Solvents)

***BENZESTROL**, NF NND 1962 (Schiefelahn) Elixir, Tabs NF Synthetic estrogen (2,4 di(p hydroxyphenyl) 3 ethyl hexane) (Octofollin)

* Generic name recognized by NF, USP or AMA

Amp: (5 mg/cc) 1 cc, Box 12s

Elixir: (2 mg/dl) Bot 1 pt

Sol in oil: (5 mg/cc) Vial 10 cc

Tab: (0.5, 1, 2, 5 & 25 mg) Bot 100s 1000s

Vaginal Tab: (0.5 mg) Bot 100s

Vial (5 mg/cc) 10 cc.

Use: Synthetic estrogen

W/Phenobarbital (Schieffelin) Benzestrol 1 mg phenobarbital 1/4 gr/Tab Bot 100s.

BENZETNACIL. Dibenzylethylenediamine dipenicillin G (DBED) See: Penicillin G Benzathine, USP

BENZETHIDINE. See 1 (2 benzylasethyl)-4 phenylpiperidine 4 carboxylic acid ethyl ester

***BENZETHONIUM CHLORIDE, USP**; Sol USP NND 1962 ADR p Tertiary acyl phenoxy ethoxy ethyl ammonium chloride Phemeride

See Ammoid Diaper Rinse (Kinney)

Anzol Sol (Veltex)

Hyacide liq (Fawcett)

Nyamine 1622 (Rohm & Noas)

Phemerol Chloride, Sol & Tr (Parke Davis)

Phemithyn (Flint, Eaton)

W/Non-ionic detergents See Ametrifan

Germicidal Detergent, liq (Meyer)

W/Sod carb and sod nitrite See Anzol Soln (Veltex)

W/Tyrothricin See Tyrothricin

W/Zinc oxide See Ammoid, Oint & Rinse (Kinney) See Trihexyphenidyl HCl

BENZETHONIUM CHLORIDE MONOHYDRATE. See Phemithyn, Sol (Flint, Eaton)

BENZHEXOL. *Trihexyphenidyl

alpha-BENZHYDROL NCl, Diphenylhydroxy

corbinol See: Franquel, Inj (Merrell)

BENZILATE. See: Diphenylacetate

***BENZILONIUM BROMIDE**. 1 ethyl 3 pyrrolidinyl benzilate ethylbromide See Parlyn (Parke Davis)

***BENZINDOPYRINE HYDROCHLORIDE**

(Irwin Neisler) 1 benzyl 3 [2 (4 pyridyl) ethyl] indole hydrochloride

Use Psychotherapeutic agent

***BENZMALECENE** a N [2 methyl 2,3 di p & Co) 1 benzyl 3 [2 (4 pyridyl) ethyl] indole HCl

Use Psychotherapeutic agent

***BENZMALECENE**, a N [2 methyl 2,3 di p chlorophenylpropyl) maleamic acid

BENZIOL (Bentex) Pentylene tetrazol 200 mg, nicotinic acid 100 mg/tab or 10 cc

Use: Senile patients

BENZODENT, (Peter, Strong) Benzocaine, eugenol, hydroxyquinoline sulphate in an adhesive ointment base. Tube ¼ oz., 1 oz. Use: Analgesic in adjustment to dentures.

BENZOCAINE, N.N.R. 1949, A.D.B., Ethyl aminobenzoate N.F., anesthetic, orthesin, porothesin

Use: Local anesthetic.

See: Americaine, preps. (Arnar-Stone).

Benzolin, Oint. (Brean).

* **BENZOCHLOROPHENE SODIUM**. The sodium salt of ortho-benzyl-para-chlorophenol

BENZOCOMP OINTMENT. (Maltbie) Benzocaine 1%, phenol 1%, zinc oxide 5%. Oint., Tube 1 oz., Jar 1 lb.

Use: Topically, local anesthetic & antiseptic.

BENZOCONES, (Fuller) Supp. Benzocaine 2 gr., hamamelis ext. 1 gr., bismuth subcarb. 1 gr., cocoa butter/Supp. Oint. Benzocaine 2%, aluminum acetate 2%, hamamelis ext. 2%, phenol 1%. Box 12s with Oint.

Use: Recl. Hemorrhoids.

* **BENZODIPA**, (Armour) benzyl[bis(aziridinyl) phosphinyl] carbamate. See: Dualor.

BENZODIOXANE PIPEROXAN. See: Benzodaine (Merck Sharp & Dohme)

BENZO-NEXYL, (Rowell) Hezylresorcinol 2.4 mg., benzoic acid 1/12 gr., benzocaine 1/5 gr./Loz. Box 24s.

Use: Throat anesthetic & antiseptic.

* **BENZOIC ACID**, U.S.P. (Various Mfr.) Pkg ¼ lb., 1 lb.

Use: Fungistatic & fungicidal.

W/Salicylic acid, benzocaine & chlorinated thymol. See: Ben-Sal-Caine, Oint. (Pitman-Moore).

W/Salicylic acid & chloromercuri corvazol. See: Monarden, Oint. (Smith-Dorsey).

W/Salicylic acid. See: Whitfield's Oint. (Merck Sharp & Dohme)

* **BENZOIC AND SALICYLIC ACID OINTMENT**, U.S.P. See: Whitfield's Oint. (Merck Sharp & Dohme)

BENZON W/Quilloja Ext., cool tar ext., nitromersol, & isopropyl alcohol. See: Kori cide, Sol. (Kormel Co.)

BENZOL. Usually refers to benzene.

BENZOLEX OINT. (Klug) Benzocaine 0.75%, cod liver oil 4%, lanolin 11%, zinc oxide 8%, balsam peru 1.5%, bismuth subgallate 1.5%, special oint. base, q.s. Tube 1 oz.

Use: Hemorrhoids.

* Generic name recognized by N.F., U.S.P. or A.M.A.

BENZOLIN OINTMENT. (Brean) Dibucaine NCI 0.5%. Tube 1 oz., Jar 1 lb., 5 lb.

Use: Topically, local anesthetic.

* **BENZONATATE**, N.N.D. 1962. *o*-Methoxy-poly(ethyleneoxy)ethyl *p*-butylaminobenzoate. See: Tessalon, Cap. (Ciba)

BENZOPYRROLATE. See: Benzopyrrolonium.

* **BENZOPYRROLONIUM BBOMIDE**. (Robins) 1-methyl-3-pyrrolidyl benzilate methobromide.

Use: Anticholinergic.

* **BENZOOQUINONIUM**. See: Mytolon, Vial (Winthrop)

BENZO-SALICYLIC COMPOUND OINTMENT. (Upjohn) Acid benzoic 12%, acid salicylic 6%, chlorothymol ¼%. Jar ¼ oz., 2 oz.

Use: Topically, keratolytic & fungicidal.

BENZOSOL. See: Guaiacol Benzoate

BENZOSULFIMIDE, U.S.P., A.D.B. See: Saccharin

W/Aspirin saccharinated. See: Theryl, Tab. (Church).

* **BENZOYLPAAS**. Calcium benzoyl-*p*-aminosalicylate.

See: Benzopas (Dorsey Labs)

Therapas (Barnes-Hind)

N'-BENZOYLSULFANILAMIDE. See: Sulfo-benzamide.

W/Sulacetamide, sulfathiazole, urea. See: Sulzin, Tab. (Ortho)

* **BENZPHETAMINE HCl**. *N*-Benzyl-*N*- α -dimethylphenethylamine HCl, dextra. See: Didrex, Tab. (Upjohn)

* **BENZPIPERTLON**, (Sandoz) 3-Benzyl-1-(1-methyl-4-piperidyl)-4-phenyl-3-pyrazolin-5-one.

Use: Connective tissue disorders.

* **BENZPYRINIUM BROMIDE**, N.F., N.N.D. 1962. 1-Benzyl-3-(dimethylcarbamoyloxy)pyridinium bromide. See: Stigmonene Bromide, Amp. (Warner-Chilcott)

* **BENZOQUINAMIDE**, (Roerig) *N,N*-diethyl-1,3,4,6,7,11b hexahydro-2-hydroxy-9,10-dimethoxy-2*H*-benzo[*a*]quinolizina-3-carboxamide acetate. See: Quontril.

BENZSULFOID LOTION. (Paythress) Col-lodal sulfur 2%, resorcinol 2%, thymol 0.5%, hexachlorophene 0.5%, zinc oxide 0.5%, alcohol 12%. Sol. 1 oz.

Use: Treatment of acne.

* **BENZTHIANIDE**. 3-Benzylthiomethyl-6-chloro-7-sulfamyl-2*H*-1,2,4-benzothiadiazine-1,1-dioxide. See: Urese (J. B. Roerig)

- ***BENZTHIAZIDE**. 3 benzylthiamethyl 6 chloro 7 sulfamyl 1,2,4 benzothiadiazine 1,1 dioxide See: Naclex, Tab (Robies)
- ***BENZTROPINE METHANESULFONATE**, NF NND 1962 Tabs NF 3 Dipheyl methoxytiapone methanesulfonate Tropine benzohydril ether methanesulfonate See: Cagentin Methanesulfonate, Tab (Merck Sharp & Dahme)
- BENZYDROFLUMETHIAZIDE**. 3 benzyl 3,4 dihydra 6 (trifluoromethyl) 1,2,4 benzthia diazine 7 sulfanamide, 1,1 dioxide See Benzdrufumethiazide
- ***BENZYL ALCOHOL**, NF, NNB 1950 Phenylcarbinol
Use Antiseptic, local anesthetic
- BENZYL p-AMINOSALICYLIC ACID** See: Benzapas, Tab (Smith Kline)
- ***BENZYL BENZOATE**, NF; Lat, NF NND 1962 Saponated NF
See Atbaide Lat (Blue Line)
BBS Lat (National Drug)
Benlatex Lat (Texas)
Benzlate Lotion, NND 1958 (Brean)
Benzylats, Cap (Merck Sharp & Dahme)
Calebent Lotion (Cale)
Scabenzole (Hart)
Vanazole (VanPelt & Brown)
Vanxanate, Lat (Veltex)
Zylate (Upjohn)
Use 10 to 30% emulsion in scabies
W/Benzyl alcohol See Benzol Emulsion (Strasbourg)
- W/DDT (Chlorophenothane) & benzocaine, NNR 1954
See Benzamol D, Emulsion (Irwin, Neisler)
Enbin Emulsion NNR (W Cooper)
Paracin Emulsion (Flint, Eaton)
Pubicide, Emulsion (Warren Teed).
- W/DDT, benzocaine, polyoxyalkylene derivatives of saibitan mono oleate See Topotide, Lat (Lilly)
- W/Isopropyl alcohol See Zylate, Sol (Upjohn)
- W/Tyrosine & benzocaine See Tyrosabe, Sol (Merck Sharp & Dahme)
- BENZYL BENZOATE CAPSULES**. See Benzylats (Merck Sharp & Dahme)
- BENZYL BENZOATE MISCIBLE**. (Hysac Westcott and Dunning) A 20% alcoholic soln Bot 2 oz
Use Orally, antispasmodic
- BENZYL BENZOATE, SAPONATED**, NF Triethanolamine 20 Gm, oleic acid 80 Gm, benzyl benzoate qs 1000 cc
- BENZYL CARBINOL**. See: Phenylethyl Alcohol, NF
- * Genetic name recognized by NF, USP or A.M.A.
- BENZYLEPHEDRINE**. See: Bephedin (Winthrop Labs)
- BENZYLETS**, (Merck Sharp & Dahme) Benzyl benzoate 0.3 cc/Cap. Gelatin Globules Bot 24s
Use: Antispasmodic
- BENZYLOG** See: Sulfabenzamine
- BENZYL PENICILLIN 2-DIETHYLAMINO-ETHYL ESTER HI**. (Panathamate HI), Neopenil, Estapee (None listed)
- BENZYL PENICILLIN G, POTASSIUM**. See Penicillin G Potassium, USP, NND 1960
- BENZYL PENICILLIN G, SODIUM**. See Penicillin G, Sodium, USP; NND 1960
- BENZYL SALICYLATE W/Camphor**. See: Camphro Salyt, Amp (Faugera)
- BENZYL SUCCINATE**. (Hynson, Westcott & Dunning) Dibenzyl succinate Tab. 4 gr Vial 40s
Use: Orally (1 fld), antispasmodic
W/Mephenesin, salicylamide See: Exposmus Tab (M. H. Smith)
- BENZYL SUCCINATE, MONO**. See: Tyrbonal (Raymer)
- BEPADIN CAPSULES**. (Ives Cameron) Vit B 1 3 mg, B 2 4 mg, niacinamide 20 mg
Cal pantothenate 2 mg, Vit B-6 0.3 mg-yeast & liver conc/Cap Bot 100s, 250s 1000s
Use Orally, vitamin supplement
- BEPADIN W/Vit C** (Ives Cameron) Vit B-1 2 mg, B 2 3 mg, niacinamide 20 mg, Cal pantothenate 0.25 mg, Vit B 6 0.33 mg, ascorbic acid 75 mg, yeast & liver conc/Cap Bot 100s, 250s, 1000s
Use: Orally, vitamin supplement
- BEPADIN W/Vit C & Vit B 12** (Ives Cameron) Vit B 1 2 mg, B 2 3 mg, B 6 0.33 mg, B 12 2 mcg, C 75 mg, niacinamide 20 mg, Cal pantothenate 0.25 mg-yeast 45 mg, liver 45 mg/Cap Bot 100s
Use Vitamin B complex
- BEPADOL ELIXIR** (Ives Cameron) Phenobarbital 1/4 gr, Vit B 1 1 mg, B 2 0.5 mg B 6 0.15 mg, niacinamide 5 mg, cal pantothenate 0.3 mg/Tsp Bot 1 pt
- BEPADOL TABLETS** (Ives Cameron) Phenobarbital 1/4 gr, Vit B 1 3 mg, B 2 2 mg B 6 0.5 mg, niacinamide 10 mg, cal pantothenate 1 mg, liver fraction 2 NF 65 mg, yeast 175 mg/Tab Bot 100s
- BEPENTA**. (Strasbourg) Tab Vit B 1 0.5 mg, B 2 1 mg, B 6 0.075 mg Cal pantothenate 0.075 mg, niacin 5 mg/Tab Bot 100s 1000s Syr Vit B 1 2 mg, B 2 4 mg

B-6 0.33 mg., pantothenic acid 0.33 mg., niacin 20 mg./fl. oz. of Syr. Sol. 1 pt., 1 gal.

Use: Orally, vitamin supplement.

BEPHAN. (Sandoz) Belladonna alkaloids as the moleate salts 0.5 mg., aluminum hydroxide-glycine 450.0 mg., mag. oxide 60.0 mg./Spacetab. Bot. 100s. [See pHan, Tabs.]
Use: Sustained anticholinergic action.

BEPHEDIN. (Winthrop Labs.) Benzyl ephedrine

W/Isoproterenol, phenorb, theophylline. See: Isuprel-Franol (Winthrop Labs.)

W/Luminal & theophylline. See: Franol, Tab. (Winthrop Labs.)

***BEPHENIUM BROMIDE.** N-benzyl-N,N-dimethyl-N-(2-phenoxyethyl)ammonium bromide. See: Alcopara (Burroughs Wellcome)

BEPNOID. (Smith, Miller & Patch) Methamphetamine 5 mg., thyrald ½ gr., Vit. B-1 1 mg., B-2 2 mg., B-6 0.1 mg., pantothenic acid 1 mg., niacin 10 mg./Tab. Bot. 100s, 1000s.

Use: Orally (1 l.l.d., o.c.), anti-obesity therapy.

BEPLETE. (Wyeth) Phenobarbital ¼ gr., Vit. B-1 1.5 mg., B-2 1 mg., B-6 0.33 mg., B-12 1.66 mcg., niacinamide 10 mg., d-Cal. pantothenate 0.2/Tab. or 4 cc. Bot. 100s, 1000s. Elixir, Bot. 1 pt., 1 gal.

Use: Orally, sedative with vitamin B complex therapy.

W/Belladonna. (Wyeth) Phenobarbital 0.25 gr., Vit. B-1 1.5 mg., B-2 1.0 mg., B-6 0.33 mg., B-12 1.66 mcg., niacinamide 10 mg., pantothenate 0.2 mg., total belladonna alkaloids 0.2 mg./4 cc. or cap. Bot. 1 pt. or 100s.

Use: Orally, antispasmodic & sedative with vitamin B complex.

BEPRON CAPS. (Wyeth) liver conc. 4.6 gr., Vit. B-12 12 mcg., dried ferrous sulfate 12 gr./Cap. Bot. 100s, 500s.

Use: Orally, treatment of anemias.

BEPRON FORTIFIED. (Wyeth) liver conc. 2.33 Gm., iron saccharate 3 gr., Vit. B-1 2 mg., B-2 2 mg., niacin 15 mg., Vit. B-12 15 mcg./oz. Liq., Bot. 1 pt. Also available as tablets. Bot. 100s, 1000s.

Use: Antianemias.

BEPRON LIQUID. (Wyeth) liver conc. 36 gr./fl. oz. equiv. of crystalline Vit. B-12 15 mcg., iron 4 gr. as saccharated ferrous iron/fl. oz. Bot. 1 pt., 1 gal.

Use: Orally, treatment of anemias.

BEB-EX. (Ives-Cameron) Cal. succinate 2.8 gr., acetylsalicylic acid 37 gr./Tab. Bot. 100s, 1000s.

Use: Orally (1-2 q. 3 h.), for treatment of arthritic & rheumatic disorders.

BERITAB. (Chicago) Vit. B-12 w/intrinsic factor ½ oral u./Tab., cream color. Bot. 30s.
Use: (1 daily) Vitamin, hemotonic.

BIRLA-CF. (Berkeley) N-acetyl-p-aminophenol 2½ gr., caffeine 10 mg., ascorbic acid 20 mg., phenylephrine HCl 5 mg., pheniramine moleate 10 mg./Tab. Bot. 100s.
Use: Cold Therapy.

BIBOCCA-C. (Roche Lab.) Thiamine HCl 10 mg., riboflavin 10 mg., niacinamide 80 mg., pyridoxine HCl 20 mg., pantothenol 20 mg., ascorbic acid 100 mg. & d-Biotin 0.2 mg./2 cc. Amp. 2 cc., Box 6s, 25s, 100s. Vial. 20 cc.

Use: I.V., I.M., water-soluble vitamin therapy.

BIBOCCA-C 500. (Roche Lab.) In addition to Berocca formula, this product contains 500 mg. Vit. C. Amp. 2 cc., Box 6s, 50s

BIROID. (Walker Vitamins) Thyrald ½ gr., thiamine HCl 3 mg./Cap. Bot. 50s, 500s, 1000s.

Use: Orally, in hypothyroidism.

BEBUBIGEN. N.N.D. 1962. (Upjohn) Vit. B-12. Cap. 25 mcg., Bot. 50s. Vial I.M. (10 mcg./cc.) 10 cc., I.M. or I.V. Cryst. (30 mcg./cc.) 10 cc., (30 mcg./cc.) 10 cc., 1000 mcg./cc. 5 cc. or 10 cc.

Use: Hematopoietic.

BESERTAL. (Central) Phenobarbital 15 mg., reserpine 0.0625 mg., Vit. B-1 5 mg./Tab. Bot. 100s, 500s, 1000s

Use: Hypertension

BESTA TABS. (Palmedica) Vit. B-1 15 mg., B-2 10 mg., niacinamide 100 mg., calcium pantothenate 20 mg., B-12 4 mcg., B-6 5 mg., Vit. C 300 mg./Tab. Bot. 100s.

Use: Vitamin supplement.

BETABOR. (Sly) Sulfathiazole 2%, beta lactose 78%, boric acid 20%. Vaginal Cap or Tab., Bot. 24s.

Use: (1 q. 2 or 3 days) For treatment of leukorrheal conditions.

BETACAINE HYDROCHLORIDE. See:

Eucaine Hydrochloride.

BETA-CEVALIN. (Lilly) Vit. B-1 5 mg., B-2 5 mg., B-6 2.5 mg., B-12 1 mcg., C 75 mg., Cal. pantothenate 12.5 mg., niacinamide 25 mg., liver prep. & stomach-tissue desic. 0.455 Gm./Pulv. Bot. 100s, 500s.

Use: Orally (1-3 daily), vitamin B complex & vitamin C deficiencies.

BETA CILIN. (Wirth) Inj 300,000 u./cc Vial 10 cc.

Oral Susp. 300,000 u./5 cc. Bot. 60 cc.

Tab. 200,000 u. Bot. 100s.

BETA-CONCEMIN ELIXIR. (Merrell) Thio-

mine HCl 32 mg, riboflavin 16 mg, niacinamide 80 mg, pyridoxine HCl 8 mg, liver fraction #2 2.6 Gm/30 cc Bot 12 oz, 1 gal

Use Orally, hematinic with nutritional supplement

BETA-CONCEMIN FERRATED (Merrill) Thio mine HCl 10 mg, riboflavin 2 mg, niacinamide 10 mg, pyridoxine HCl 1 mg, ferrous gluconate 5 gr, Vit B 12 activity 33 mcg / Cap or dr Cap, Bot 100s, 1000s, Elixir, Bot 12 oz, 1 gal

Use Orally (12 dr or Cap 1/d) therapeutic hematinic & nutrient tonic

BETA-CONCEMIN TABLETS (Merrill) Thio mine HCl 4 mg, riboflavin 2 mg, niacinamide 10 mg, pyridoxine HCl 1 mg, choline dihydrogen citrate 12.5 mg / Tab Bot 100s, 1000s

Use Orally, hematinic with nutritional supplement

RETACURON. (Lakeside) Vit B 1 9 mg, B 2 3 mg, B 6 12 mg, niacinamide 60 mg, pantothenic acid salt 12 mg, ferrous pepsinate 274 mg, copper gluconate 16 mg & rice bran/8 oz Bot 1 pt

Use Orally, hematinic

BETADINE, NND 1962 (Tailby-Nason) Poly vinylpyrrolidone iodine complex (1% available iodine) Bot 1/2 oz Shampoo Plastic

Aerosol Spray 3 oz

Liq Bot 8 oz, 1 pt

Oint 1 oz Tube

Shampoo Bot 4 oz

Surgical Scrub 1 pt

Swab Aids 100s

Yog Douche Bot 8 oz

Yog Gel Tube W/App'l. 3 oz

Use: Topical antiseptic

BETA-ESTRADIOL. See: Estradiol USP

BETA-EUCAININE HYDROCHLORIDE. See: Eucaine Hydrochloride

BETA FOPELEX CAPSULES. (M. R. Thompson)

Vit B-1 1 mg., B-2 2 mg., R 12 1 mcg, ferrous glycinate 25 mg, folic acid 1 mg, niacinamide 10 mg, liver substance 130 mg., yeast 130 mg / Cap Bot, 42s, 500s

Use: Orally (1 1/d) hematinic & vitamin supplement

BETA FOPELEX DROPS. (M. R. Thompson)

Vit B-1 1 mg, B 2 2 mg, ferrous glycinate 10 mg, aminoacetic acid 130 mg, folic acid 0.5 mg, niacinamide 5 mg/cc. Bot w/dropper 30 cc.

Use: Orally, hematinic & dietary supplement

BETA FOPELEX ELIXIR. (M. R. Thompson)

Vit B-1 1 mg, B-2 1 mg, B-12 1 mcg, niacinamide 10 mg, folic acid 1 mg, amino-

acetic acid 0.325 Gm, ferrous glycinate 25 mg, rice bran ext, yeast & liver fraction #2 250 mg / 5 cc Bot, 8 oz, 1 gal

Use Orally, hematinic

***BETA-GLUCURONIDASE**. See: Ketodase 1uq (Warner Chilcott)

***BETAHISTINE HYDROCHLORIDE** (Spencer) 2 (2 methylaminoethyl)pyridine hydrochloride See Serc

BETA-HYPOPONAMINE See Vasopressin

BETAINE. W/Glycocyamine See Betalys amine (Internal Minerals & Chem)

RETAIN-CHOLINE W/Inositol, dehydrocholic acid, ox bile & liver R. See: Hepoxyn, Tab (Fadacine)

BETAINE HYDROCHLORIDE, NNR 1940. (Various Mfr) Acidal HCl, lysine HCl

W/Dehydrocholic acid, choline bitartrate & methionine See Chalitrapic, Tab (Buffington)

W/Homatropine methylbromide, desoxycholic acid, ginger, pancreatin See: Convertin H, Tab (B F Ascher)

W/Mephenesin, ethaverine HCl See: Mepherin, Tab (G F Harvey)

W/Oleoresin ginger, pancreatin, desoxycholic acid & chlorophyllin See Convertin, Tab (Ascher)

W/Pancreatin, ketocholic acids, bile salts See: Pradose, Tab (Miller Labs)

W/Pepsin See: Acidal Pepsin, Cap (Winthrop Labs)

W/Pepsin & methylcellulose See: Normacid Tab

W/Thyroid, steapsin See: Chylipase, Tab (Columbus)

BETAJEL. (Floralose) Fibrous, gel forming portions of psyllium seed w/dextrose & beta lactase Rox 12 oz

Use Orally, bulk stimulation, lubrication & laxative

RETAKAR. (Endo) Karaya gum w/Vit B complex Can 4 oz, 1/2 lb, 1 lb

Use Orally, laxative

RETAL (Central) Phenobarbital 15 mg, Vit B 1 5 mg / Tab Bot 100s, 1000s

Use: Sedative

BETA-LACTOSE. (Borden) Pkg 1 lb, 5 lb

Use: Infant feeding

BETALIN-12. (Lilly) Vit B 12 cryst Amp Vial (30, 100/cc) 10 cc, 1000 mcg / cc 5 cc

Use: IM, pernicious anemias

BETALIN COMPLEX, AMPULS. (Lilly) Thiamine HCl 10 mg, riboflavin 4 mg, niacinamide 150 mg, pantothenic acid 5 mg, pyridoxine HCl 10 mg, B 12 5 mcg / 2 cc Amp 2 cc, Pkg 6s, 100s, 10 cc, Pkg 25s, 50s, 100s

Generic name recognized by N.F. USP or A.M.A.

Use: I.M., I.V., vitamin B complex deficiencies.

BETALIN COMPLEX, ELIXIR. (Lilly) Thiamine chloride 2 mg., riboflavin 1 mg., pyridoxine HCl 0.5 mg., Vit. B-12 1 mcg., pantothenic acid 2 mg., nicotinamide 5 mg., w/liver fraction 400 mg./fl. dr. Bot. 1 pt., 1 gal. Use: Vitamin B complex deficiencies.

BETALIN COMPLEX, F.C. (Lilly) Thiamine chloride 12.5 mg., riboflavin 3 mg., nicotinamide 50 mg., pantothenic acid 2.5 mg., pyridoxine HCl 5 mg., ascorbic acid 75 mg., benzyl alcohol 2%/cc. Amp. 2 cc., Box 6s, 100s, Vial 10 cc., Box 25s, 100s.

Use: I.M. or I.V., vitamin B complex & vitamin C.

BETALIN COMPLEX, SYRUP. (Lilly) Thiamine HCl 1.5 mg., riboflavin 1 mg., pyridoxine 0.6 mg., pantothenic acid 1 mg., nicotinamide 7.5 mg./5 cc. Bot. 1 pt., 1 gal.

Use: (1-2 tsp. daily) Vitamin B complex deficiencies.

BETALIN COMPOUND. (Lilly) Thiamin chloride 1 mg., riboflavin 2 mg., pyridoxine HCl 0.5 mg., pantothenic acid 3.333 mg., nicotinamide 10 mg., Vit. B-12 1 mcg., liver prep. & stomach-tissue material, desic. 0.33 Gm./Pulvule. Bot. 100s, 500s.

Use: Orally, vitamin B complex deficiencies.

BETALIN 3, N.N.D. (Lilly) Thiamine chloride, Amp. (100 mg./cc.) 1, 5, 10 & 30 cc. Elixir (6.66 mg./fl.oz.) Bot. 1 pt., 1 gal. Tab. (5, 10, 15, 25, 50 & 100 mg.) Bot. 100s, 500s.

Use: I.M., I.V., orally; Vit. B-1 deficiency.

BETA-L TEST. (Hyland) Test reagent is a specific antiserum obtained by hyperimmunization of animals w/human low-density lipoprotein./60 test kit containing antiserum, capillary pipettes, hematocrit tubes, magnifying glass, millimeter scales and card slides.

Use: Diagnostic reagent.

* **BETAMETHASONE.** 9 α -Fluoro-11 β -17 α -21-trihydroxy-16 β -methylpregna-1,4-diene-3,20-dione. 9 α -Fluro-16 β -methylprednisolone. See: Celestone (Schering Corp)

BETA-METHISCHOL SYRUP. (U. S. Vitamin) Betaine monohydrate 3.5 Gm., litcholine citrate 1.4 Gm., dl-methionine acetylated 0.24 Gm., inositol 0.5 Gm., Vit. B-12 24 mcg., liver conc. 0.8 Gm./oz. Bot. 1 pt., 1 gal.

Use: (2-3 dr. t.i.d.) lipotropic.

BETANAPHTHOL BENZOATE, N.N.B. 1940

Mallinckrodt—Pow. 1 oz., 1/4 lb., 1 lb

Mesck—Pow. 1 oz., 1/4 lb., 1 lb

Use: Antiseptic & parasiticide

* Generic name recognized by N.F., U.S.P. or A.M.A.

BETAPAN. (Brewer) Vit. B-1 0.18 mg., B-2 100 mcg., B-6, nicotinic acid, folic factor & other components of B complex as supplied by liver/Cap. Bot. 100s, 500s.

Use: Orally (1-2), in the treatment of anemias.

BETAPHOS. (Alphaden) Sod. acid phosphate w/effervescent alkaline buffer salt base Pow., Bot. 1/2 lb., 1 lb.

Use: Orally; chologogue & laxative.

BETAPLEXIN. (Winthrop Labs.) Vit. B-1 1.5 mg., B-2 0.5 mg., B-6 0.1 mg., nicotinamide 5 mg., pantothenic acid 0.275 mg./Cap. or Tab. Bot. 100s, 500s.

Use: Orally, vitamin B complex therapy.

BETAPLEXIN ELIXIR. (Winthrop Labs.) Vit. B-1 3 mg., B-2 1 mg., B-6 0.3 mg., nicotinamide 10 mg., pantothenic acid 0.75 mg./5 cc. Bot. 1 pt., 1 gal.

Use: Orally, vitamin B complex therapy.

BETAPLEXIN SYRUP. (Winthrop Labs.) Vit. B-1 3 mg., B-2 1 mg., B-6 0.675 mg., nicotinamide 10 mg., pantothenic acid 1.8 mg./5 cc. Bot. 1 gal.

Use: Orally, vitamin B complex therapy.

BETAPRONE. (Testogor) Purified grade (99% minimum purity) beta-propiolactone. Amp 3 cc., 5 cc., 10 cc.

Use: Sterilization of vaccines, tissue grafts and plasma.

BETA-PYRIDYL-CABBINOL. Alcohol corresponding to nicotinic acid. See: Roncolol, Elixir & Tab. (Roche Lab)

BETAXON. (Warren-Teed) Thiamine HCl 3 mg., ferrous sulfate 1 Gm./fl. oz. Syrn. Bot. 1 pt., 1 gal.

Use: Orally, hematinic with vitamin B-1 supplement.

BETASCORBATE. (Upjohn) Vit. B-1 0.36 mg., B-2 0.1 mg., nicotinic acid 5 mg., Cal. pantothenate 0.2 mg., Vit. B-6 0.08 mg., ascorbic acid 15 mg./Tab. Bot. 50s, 100s, 500s.

Use: Orally, dietary supplement.

BETASYAMINE. (Internat. Minerals & Chem.) Betaine & glycocyanine in various combined dosage forms.

Use: Polyomyelitis, arthritis & degenerative heart disease.

BETASYNPLEX ELIXIR. (Winthrop Labs.) Vit. B-1 0.66 mg., B-2 0.66 mg., B-6 0.33 mg., Cal. pantothenate 0.33 mg., nicotinamide 5 mg./dr. Bot. 1 pt.

Use: Vitamin B complex.

BETASYNPLEX TABLETS. (Winthrop Labs.) Vit. B-1 0.666 mg., B-2 0.666 mg., B-6 0.333 mg., Cal. pantothenate 0.333 mg., nicotinamide 5 mg./tab. Bot. 100s.

Use: Orally, vitamin B complex therapy.

BETASYNPLEX VIAL (Winthrop Labs) Vit B 1 30 mg, B 2 10 mg, B 6 5 mg, nicotinamide 100 mg, cal pantothenate 5 mg / dose Amp single dose Vial 10 dose Also available w/Vit B 12 5 mcg/dose Also available w/Vit C 500 mg/dose

Use I.M., vitamin B complex therapy

BETHIOLIN (Lincoln) Cysteine 10 mg, thiamine 30 mg, riboflavin 2 mg, niacin 100 mg, pantothenic acid 4 mg, Vit B-6 2 mg, foreign protein 100 mg / 2 cc Amp 2 cc, Vial 30 cc, 60 cc

Use I.M., as nonspecific antirheumatic therapy

ETATRIN CAPSULES (Gateway Labs) Vit B 1 2 mg; B 2 3 mg; B 6 1 mg, cyanocobalamin 5 mcg; Vit B 12 with IFC 1/12 USP u, niacinamide 10 mg; ascorbic acid 50 mg; ferrous fumarate 10 mg / Cap Bot 100s 1000s

Use Vitamin Combination

ETAXIN, NND 1958 (Winthrop Labs) Thiamine HCl Elixir, 6 mg / fl oz Bot B fl oz, 32 fl oz 1 gal Tab 5 mg 10 mg, 25 mg Bot 100s 500s 50 mg Bot 25s 100s 500s, 1000s; 100 mg Bot 100s 500s 1000s Vial (250 mg) 10 cc; (500 mg & 1000 mg) 10 cc; (3000 mg) 30 cc

Use I.M. for treatment of beriberi & alcohol & other types of neuritis

BETAZOLE, HCl, USP, NND 1962 Inj USP 3-(b-aminoethyl) pyrazole d HCl See Histalog Amp (Lilly)

ITAZYME (Barrows) Vit B 1 25 mg, B 1 (Cocisoboxylase from yeast) 2.5 mg, B 2 10 mg, niacinamide 50 mg, calcium pantothenate 25 mg, B-6 25 mg, menadione (2 methyl 1-4 naphtho quinone) 2 mg, choline HCl 10 mg, high potency yeast concentrate 200 mg, Brewers type yeast 200 mg, yeast protein hydrolysate 1 Gm, sodium nucleate 100 mg, glutamic acid 50 mg, glycine 50 mg, cysteine HCl 25 mg, Iron, manganese magnesium, zinc / 3 Tabs Bot 100s 1000s

BETEX, IMPROVED (Horton & Converse) Vit B 1 100 mg, B-2 2 mg, B 6 1 mg, niacinamide 20 mg, pantothenyl alcohol 5 mg / cc, Vial 10 cc, 20 cc

Use I.M. + vitamin B complex therapy

BETHANECHOL CHLORIDE, U.S.P., NND 1962, Inj USP; Tab USP Carbonyl methylcholine chloride See Urecholine Chloride, Amp & Tab (Merck Sharp & Dohme)

Use: (Orally 5 to 30 mg; 2.5 to 5 mg subcut) Parasympathomimetic

BETHANIDINE HYDRIODIDE (Berroughs) * Generic name recognized by N.F., USP or A.M.A.

Wellcome 1 Benzyl 2,3 dimethylguanidine N1

Use: Hypotensive

BETNIAMIN, (Massengill) Thiamine HCl Vial (100 mg/cc) 10 cc Cap. 3 1/2 mg, 10 mg 15 mg, Bot 100s, 1000s Elixir 6 mg / fl oz Bot 1 pt, 1 gal

Use Orally, vitamin B-1 deficiencies

BETNILE, (G F Harvey) B 6 10 mg, B 2 10 mg, niacinamide 100 mg, B 6 2 mg, Vit B 12 activity 4 mcg, 1 lysine monohydrochloride 800 mg, aromatic vehicle qs / 1 oz Elix Bot 1 pt

Use: Nutritional supplement

BETHROID, (Massengill) Thyroid 30 mg Vit B 1 3 mg, nicotinamide 10 mg, Vit B 2 3 mg / Tab Bot 100s, 1000s

Use: Thyroid medication

BETOCAINE, (Warsen Teed) Vitamins B 1 100 mg, B 2 5 mg, B-6 5 mg, niacinamide 40 mg, pantothenol 5 mg & procaine HCl 10 mg / cc Vial 10 cc

BETOLAKE W/Vit C (Lakeside) Vit B-1 50 mg, B 2 10 mg, B-6 50 mg, C 250 mg; niacinamide 250 mg, Sed pantothenate 50 mg / 5 cc Vial 10 cc Box 25s, 100s

Use I.M., vitamin B complex

BETONE (J E Dahl) Vit A 25,000 u D 1000 u, Thiamine HCl 10 mg, riboflavin 5 mg, ascorbic acid 150 mg, niacinamide 150 mg / Cap Bot 100s

Use Multivitamin therapy

BETONE (Lemman) Vit B 12 3 mcg, liver desc. NF 0.1 Gm, ferrous gluconate 0.1 Gm, Vit B 1 1 mg, B 2 0.5 mg, Cal. pantothenate 0.5 mg, niacinamide 10 mg-Mang, copper, cobalt / Tab Bot 100s

Use Orally, hematinic

BEVATINE, NND 1962 (Smith Dorsey) Vit B-12 cryst Vial (100 mcg 1000 mcg) 10 cc

Use I.M., vitamin B 12 therapy

BEVATINE-C, (Smith Dorsey) Vit B 1 20 mg, B 2 2.5 mg, B-6 2.5 mg, C 100 mg, niacin 100 mg, pantothenic acid 5 mg / cc Vial 5 cc

Use: Vitamin B complex w/vitamin C therapy

BEVEX, (Metropoliton) Lives Inj, Vit B-12 activity 2 mcg, B-12 10 mcg, niacinamide 10 mg, Vit B-6 0.5 mg, B 1 10 mg, B 2 0.5 mg, Cal pantothenate 3 mg / cc Amp 1 cc, Box 100s; Vial 10 cc, 30 cc

Use I.M., hematinic & treatment of anemia

BEVIDORAL, Filmtabs, NND 1960 Not listed (Abbott) Vit B 12 & duodenal ext concentrate Tab (equiv to 1/2 USP u) Bot 25s 100s

Use: Orally; pernicious anemia

BEVIDOX CONCENTRATE, N.N.D. 1962. (Abbott) Cyanoecobalamin, Vit. B-12. Cap. 25 mcg. Bot. 100s. Tab., Dulcet 10 mcg. Bot. 100s. Vial (30 mcg./cc.) 10 cc., (150 mcg./cc.) 5 cc., (1000 mcg./cc.) 5 cc., 10 cc.
Use: Treatment of macrocytic & pernicious anemia.

BEVIMIN. (Lemmo) Vit. B-12 8 mcg., B-1 8 mcg., B-2 2 mcg., niacinamide 20 mcg., cal. gluconate 250 mg., ferric ammonium citrate 250 mg., Mang. glycerophosphate 60 mg./fl. oz.
Use: Orally, treatment of anemias & other B complex deficiencies.

BEVIMINE, (Maney) Thiamine 10 mg., riboflavin 5 mg., pyridoxine hydrochloride 1 mg., niacinamide 50 mg., calcium D-pantothenate 5 mg., Vit. B-12 3 mcg., desiccated liver 100 mg., dried yeast 50 mg., choline bitartrate 20 mg., inositol 10 mg., ascorbic acid/Tab. Bot. 100s.
Use: Treatment of deficiencies of vitamin B complex and vitamin C.

BEVIPLEX, SUPER. (Lincoln) Thiamine HCl 25 mg., niacinamide 25 mg., riboflavin 2½ mg., pyridoxine 5 mg., Sod. pantothenate 5 mg., benzyl alcohol 1%, distilled water q.s./cc. Vial 20 cc.
Use: I.M., vitamin B complex therapy.

BEVIPLEX, SUPER-FORTIFIED. (Lincoln) Thiamine HCl 50 mg., niacinamide 100 mg., riboflavin 2 mg., pyridoxine 2 mg., pantothenic acid 2 mg./cc. Vial 30 cc.
Use: Vitamin B complex therapy.

W/Vit. C. (Lincoln) Thiamine HCl 25 mg., niacinamide 25 mg., riboflavin 2½ mg., pyridoxine 5 mg., Sod. pantothenate 5 mg., ascorbic acid 50 mg./cc. Vial 20 cc.
Use: Vitamin B complex therapy.

BEVITAL (Key Corp.) Pentobarbital Sod. 1½ gr./Tab. Bot. 100s

BEVITAM. (Merrell) Vit. B-12 as a zinc tartrate B-12 complex combined with sufficient free crystalline B-12. 500 mcg./cc. Vial 5 cc. w/diluent. Repasitory Inj. Vit. B-12 2500 mc./5 cc. Vial 5 cc.
Use: i.M., Vit. B-12 therapy.

BEVITE. (Lemmo) Vit. B-12 2 mcg., liver desic. N.F. 0.1 Gm., thiamine mononitrate 1 mg., riboflavin 0.5 mg., Cal. pantothenate 0.5 mg., niacinamide 10 mg./Tab. Bot. 50s.
Use: Vitamin therapy in treatment of anemias.

BEVITE. (Key Corp.) Vitamin B-12, Crystalline. Vial (30 mcg./cc.) 10 cc., 30 cc. (100 mcg./cc.) 10 cc. (1000 mcg./cc.) 10 cc.

BEYBONS. (Vitamin Corp.) Vit. B-1 6 mg., B-2 3 mg., B-6 0.5 mg., C 50 mg., niacinamide 10 mg., Cal. pantothenate 3 mg., folic acid 0.1 mg., Vit. B-12 1 mcg., liver &

yeast q.s./Cap. Bot. 50s, 100s, 250s, 1000s.
Use: Vitamin therapy & dietary supplement

BEWON, ELIXIR. (Wyeth) Thiamine chloride 1.5 mg./fl. oz. Bot. 1 pt., 1 gal.
Use: Promoting appetite.

BEXEL. (McKesson & Robbins) Vit. B-1 500 u., B-2 2 mg., Cal. pantothenate 0.5 mg., niacin 10 mg., Vit. B-6 0.25 mg./Cap. Bot. 40s, 100s, 250s.
Use: Vitamin therapy.

BEXELETES. (McKesson) Vit. A 5000 u., D 1000 u., B-1 2 mg., B-2 2 mg., C 50 mg., niacinamide 25 mg., B-6 1 mg., folic acid 0.1 mg., cyanocobalamin 2 mcg., calcium pantothenate 5 mg., D-alpha tocopheryl acetate 2 IU/lessel. Bot. 120s, 300s.
Use: Vitamin deficiencies.

BEXELETES-M. (McKesson) Same formula as Bexeletes plus: iron (ferrous sulfate) 25 mg., iodine (potassium iodide) 0.15 mg., cobalt (sulfate) 0.1 mg., copper sulfate 1 mg., manganese sulfate 0.2 mg., molybdenum (sodium molybdate) 0.2 mg., magnesium sulfate 1 mg., zinc sulfate 0.5 mg., potassium sulfate 5 mg./lessel. Bot. 100s, 250s.
Use: Vitamin deficiencies.

BEXII, N.N.D. 1962. (Chicago Pharm) Vit. B-12 cryst. Amp. (15 mcg./cc.) 1 cc., Box 12s, 25s, 100s. Vial (30 mcg./cc.) 10 cc.
Use: I.M., hematinic in anemias.

BEXII-M. (Chicago Pharm.) Vit. B-12 cryst. Vial (1000 mcg./cc.) 10 cc., Pkg. 1c, 6s, 30s.
Use: I.M., anemias.

BEXIFORT, (Chicago Pharm.) Vit. B-12 cryst Vial (100 mcg./cc.) 10 cc.
Use: Anemias.

BEXINE. (Ascher) Thiamine HCl 25 mg., riboflavin 2 mg., pyridoxine HCl 5 mg., nicotinamide 100 mg., pantothenic acid 5 mg./cc. Vial 10 cc.
Use: I.M., vitamin therapy.

W/Vit. B-12 fortified. (Ascher) Thiamine HCl 25 mg., riboflavin 2 mg., pyridoxine HCl 5 mg., niacinamide 100 mg., pantothenic acid 5 mg., Vit. B-12 cryst. 30 mg./cc. Vial 10 cc.
Use: I.M., vitamin therapy.

BEXITAR, N.N.D. 1960. (Chicago Pharm) Vit. B-12 with intrinsic factor ½ u./Tab. Bot. 30s, 100s.
Use: Pernicious anemia.

BEXIVER SOLUTION. (Breon) Crude liver 2 u./cc. W/Vit. B complex. Vial 10 cc.
Use: I.M., injectable liver.

BEXO-C. (Mallard) Vit. B-1 25 mg., B-2 5 mg., B-6 5 mg., C 50 mg., niacinamide 50 mg., pantothenal 5 mg./cc. Vial 10 cc.
Use: Treatment of multiple Vitamin B complex deficiencies.

BEXOID. (C. D. Smith) Thyroid 30 mg, Vit B-1 2 mg, B-2 2 mg, niacinamide 10 mg, Cal pantothenate 1 mg, Vit B-6 0.1 mg / Tab Bot 100s, 1000s

Use: Orally, thyroid therapy w/vitamin supplement

BEXOSAL. (Ascher) Not listed Salicylamide 250 mg, di desoxyephedrine HCl 1 mg, thiamine HCl 10 mg, riboflavin 5 mg, niacinamide 20 mg, ascorbic acid 50 mg / Tab Bot 100c, 1000s

Use: Orally (1 2 1 1 d), analgesic, CNS stimulant & vitamin supplement

BEXTRAL. (Central Pharm) High potency B complex factors Vial 10 cc, 30 cc

BEXTRAL B-12. (Central) Vit B-1 25 mg, B-2 2 mg, B-6 5 mg, nicotinamide 100 mg, pantothenic acid 5 mg, Vit B-12 30 mcg / cc. Vial 10 cc, 30 cc

Use: I.M., vitamin B complex therapy

BEZON. (Whittier) Vit B-1 5 mg, B-2 3 mg, B-6 0.5 mg, B-12 1 mcg, C 50 mg, niacinamide 20 mg, Cal, pantothenate 3 mg / Cap Bot 30s, 100s Available as Forte Tab

Use: Orally, vitamin B complex w/vitamin C therapy

BEZON FORTE. (Whittier) Thiamine mononitrate 25 mg, riboflavin 12.5 mg, niacinamide 50 mg, calcium pantothenate 10 mg, pyridoxine HCl 5 mg, ascorbic acid 150 mg / cap Bot 30s, 100s

BEZYMIN. (Crocker) Not listed Pancreatic substances equiv to pancreatin USP 450 mg, malt dextrose 50 mg, ox bile ext 100 mg, glutamic acid HCl 100 mg, Vit B-1 5 mg, B-2 1 mg, niacinamide 10 mg, Vit C 25 mg / Tab Bot 50s, 100s, 500s

Use: Orally, faulty digestion

B-FERRATED. (Upjohn) Ferrous sulfate 1.3 Gm, Vit B-1 12 mg, B-2 2 mg, nicotinamide 50 mg / R or Elisir, Bot 1 pt, 1 gal

Use: Orally, hematinic w/vitamin B complex

B-F-I POWDER. (Merck Sharp & Dohme) Biz-muth formic-iodide, zinc phenolsulfonate, biz-muth subgallate, emel, Per alum, boric acid, menthol, eucalyptol, thymol & inert diluents Can 1/4 cc, 1 1/2 oz & 1/2 lb

Use: Antiseptic Brit-acid dressing & surgical dressing pow

B-FOL-C. (Pitman Moore) Vit B-1 1 mg, B-2 1.5 mg, B-6 0.1 mg, C 50 mg, niacinamide 5 mg, Cal pantothenate 0.5 mg, yeast 65 mg / Tab, Bot 100s, 500s

Use: Orally, vitamin B complex.

B-FOLIDOX. (Abbott) Vit B-12 25 mcg, folic acid 1.7 mg / Cap Bot 100s, 1000s

Use: Orally, macrocytic anemias.

* Generic name recognized by N.F., U.S.P. or A.M.A.

B-FORTEX. (Plechner) Vit B-1 100 mg, B-2 1 mg, B-6 10 mg, niacinamide 100 mg, Cal pantothenate 10 mg / cc Vial 10 cc.

Use: I.M., vitamin B complex therapy

B-G-PHOS' ELIXIR. (Merck Sharp & Dohme) Thiamine HCl 4 mg, riboflavin 2 mg, Cal glycerophosphate 0.12 Gm, Sod glycerophosphate 0.25 Gm, Pot glycerophosphate 25 mg, Mang glycerophosphate 15 mg, Vit B-12 25 mg / 30 cc Bot 1 pt, 1 gal

Use: Orally, stimulate appetite.

B-2 OR G VITAMIN. See: Riboflavin

BIALAMICOL. 3,3'-Diallyl 5,5'-bisdiethylaminomethyl 4,4'-dihydroxydiphenyl, d hydrochloride See Camofarm (Parke, Davis)

BIALLYLAMICOL. See Camofarm (Parke, Davis)

BIATRON ELIXIR. (National Drug) Vit B-1 8 mg, B-2 2 mg, nicotinamide 5 mg, ferric ammonium citrate 4 gr, Cal gluconate 4 gr, Mang glycerophosphate 1 1/2 gr / 8 oz Bot 8 fl. oz, 1 gal

Use: Orally (2 dr f i d), deficiency of B vitamins & minerals

BIATRYN-C. (National Drug) Vit B-1 1 mg, B-2 1 mg, C 15 mg, nicotinamide 10 mg, ferrous sulfate 3 gr, liver fraction #2 3 gr / Tab Bot 100s, 1000s

Use: Orally (2 1 1 d), anemias.

BICAREX. (Massengill) Ox bile ext 1 gr, phenolphthalein 1/2 gr, capsicum oleoresin 1/40 gr, cascara ext, 1/2 gr, papain 2 gr, nux vomica ext 1/16 gr / Tab Bot 100s, 500s, 1000s.

Use: Orally (1 3 daily) choleric & chologogue

BICNOLAN. (Klug) Ox bile extract NF, 1.5 gr, cascara ext 5 gr, phenolphthalein 5 gr, oleoresin capsicum 0.05 gr / Tab Bot 100s

Use: Choleric laxative

BICNOLATE. (Blue Line) Desic. ox gall 2 1/2 gr, Sod salicylate 1/2 gr, phenolphthalein 1/2 gr, cascara 1/2 gr / Tab Bot, 100s.

Use: Orally (1 1 1 d. o c); choleric & chologogue

BICILLIMYCIN ALL-PURPOSE. (Wyeth) Bicillin 600M u, procaine penicillin G 300M u, Pot. penicillin G 300M u, streptomycin 250 mg, dihydrostreptomycin 250 mg / dose. Vial, single dose.

Use: Penicillin and streptomycin therapy

BICILLIN, NND 1962, ADR. (Wyeth) Benzathine penicillin G, USP

Drops (150M u / cc) Bot w/dropper 10 cc

Susp: (150,000 u / 5 cc) (300M u / 5 cc) Bot 2 fl. oz.

Tab: (100M u) Bot 100c (200,000 u) Vial 36s

UBER (600 M u./cc.) 1 cc.
VIAL (300M u./cc.) 10 cc.
 Use: long-acting penicillin therapy.

W/Procaine penicillin G, Pot. penicillin G, streptomycin sulfate & dihydrostreptomycin sulfate. See: Bicillinmycin All Purpose (Wyeth).

BICILLIN, Inj., long acting. (Wyeth) Tubex (600,000 u.) 10s. Vial (300,000 u./cc.) 10 cc. Disp. Syr. (900,000 u.) 1.5 cc., (1,200,000 u.) 2 cc., (2,400,000 u.) 4 cc.

BICILLIN ALL-PURPOSE. (Wyeth) Dibenzyl-ethylenediamine dipenicillin G 600M u. W/procaine penicillin G 300M u. & Pot. penicillin G 300M u./dose. 2 cc. Vial 1 & 5 dose.
 Use: I.M., high blood levels w/prolonged effect.

BICILLIN C-R 600. (Wyeth) Bicillin 300,000 u., procaine penicillin G 300,000 u./1 cc. Tube. Vial 1 dose. 2 doses.

BICILLIN C-R 300. (Wyeth) Bicillin 150M u., procaine penicillin G 150M u./cc. Vial 10 cc.

BICILLIN L-A. (Wyeth) Bicillin 200M u./tab. long acting. Vial 36s.
 Use: long acting penicillin therapy.

BICILLIN-SULFAS. (Wyeth) Bicillin 150,000 u. w/sulfadiazine, sulfamerazine & sulfathiazole 0.167 Gm. of each/3 cc. or tab. Bot. 2 oz., Tab., Bot. 36s.
 Use: Orally, triple sulfonamide w/penicillin therapy.

BICILLIN-VEE. (Wyeth) Benzathine penicillin O 100M u., penicillin V 100M u./tab. Bot. 26s.
 Use: Orally, penicillin therapy.

BICBEAL. (G. F. Harvey) Nux vomica P. ext. 1/8 gr., bile salts compound 2 gr., pancreatin 1 gr., sod. bicarbonate 1/2 gr., secretin 1/4 gr./Tab. Bot. 100s, 1000s
 Use: Cholagogue

BIDOLAB. (Forrest Company) Diactyl sod. sulfosuccinate 3/4 gr., ox bile 1 gr./Tab. Bot. 30s, 100s
 Use: Treatment of constipation

BIDROLIN. (Armaur) Dehydrocholic acid 1.9 gr., choline bitartrate 2 1/2 gr./Tab. Bot. 100s, 1000s.
 Use: Orally (2 p.c., l.i.d.), hydrocholeric & lipotropic.

BIDSOX. (McNeil) Aluminum hydroxide 400 mg., Mag. hydroxide 100 mg./3 cc. Susp. Bot. 12 fl. oz. Also available as tablets Bot. 100s, 1000s
 Use: Orally, gastric antacid & adsorbent.

BIDUPAN. (Cavendish) Ox bile ext., conc. pancreatin, duodenal substance & charcoal. Tab. Bot. 50s, 100s, 500s, 1000s.

Use: Orally, choleric & cholagogue.

BIFACTON, N.N.D. 1962. (Organon) Vit. B-12 w/intrinsic factor conc., 1/4 & 1/2 anti-anemia u. of Vit. B-12 w/intrinsic factor conc./Tab. Bot. 30s.
 Use: Pernicious anemia.

BI-FER-LIN. (Barlow-Money) Vit. B-1 1 mg., B-2 2 mg., B-6 0.01 mg., ferrous gluconate 5 mg., liver fraction #2 5 mg., pantothenic acid 2 mg., nicotinamide 5 mg., folic acid 0.75 mg./Cap. or Tab. Bot. 100s, 1000s.
 Also, above formula available w/Vit. B-12 3 mcg./dr. or Tab. Elixir, Bot. 1 pt., 1 gal./Tab. Bot. 100s, 1000s.
 Use: Orally (2-4 t.i.d.), secondary anemias.

BIFIBON. (Massengill) Ferrous sulfate 0.2 Gm., Vit. B-1 2 mg., Vit. B-2 2 mg., niacinamide 15 mg./Tab. Bot. 100s
 Use: Iron deficiency anemias

BIFRAN. (Maltbie) Methamphetamine HCl 5 mg., dehydrocholic acid 200 mg., pentobarbital 15 mg./Tab. Bot. 100s, 500s, 1000s.
 Use: Orally, antiobesity.

BI-GER-MONE. (Kirk) Testosterone propionate 25 mg., estradiol benzoate 1 mg. in sesame oil/cc. Vial 10 cc. Aq. Susp. contains estrone 2 mg., testosterone 25 mg./cc. Vial 10 cc.
 Use: I.M., menopause & functional uterine bleeding.

BINISTOL. (Sullin & Cote) Pyrilamine maleate 25 mg., methapyrine lumate 25 mg./Tab. Bot. 100s, 1000s
 Use: Antihistaminic

BINISTIN. (Ingram) Pyrilamine maleate 6 mg., chlorpheniramine maleate 1 mg., salicylamide 200 mg., phenylephrine HCl 5 mg., Vit. C 100 mg./Cap. Box 18s. Bot. 100s.
 Use: Antihistaminic and analgesic.

BI-KETOLAN. (Merrell) Oxidized or keto-form bile acids 250 mg. Tab. Bats. 100s & 1000s.
 Use: Hepatic dysfunction.

BLAMIDE. (Rossar Labs.) Dehydrocholic acid 2 gr., ox bile 2 gr., homatropine methylbromide 1/48 gr., phenobarbital 1/4 gr./Tab. Bot. 50s.
 Use: (1 t.i.d.), antispasmodic w/cholagogue

BLAZE. (Central) Bile salts compound 0.15 Gm., papain 0.12 Gm., duodenal substance 8 mg., sod. bicarbonate 30 mg./Tab. Bot. 100s, 1000s
 Use: Biliary stimulant and laxative

BLICAIN. (Cofe) Purified ox bile 3 gr., aloin 1/2 gr., cascara ext. 1/2 gr./Tab. Bot. 100s.
 Use: Orally, choleric & cholagogue.

BILCO CREAM. (Tablas) lactic acid, oleic acid, citric acid, triethanolamine, methyl &

propyl p hydroxybenzoate in glyceryl mono-
 stearate Cream 3 oz
 Use Contraceptive cream
BILCO JELLY, (Tablax) Lactic acid, boric
 acid, methyl & propyl p hydroxybenzoate
 in veg gum Tube 3 1/2 oz
 Use Contraceptive
BILE ACIDS, OXIDIZED. Note also dehydro-
 cholic acid
 See Bru Col, Tab (United Lab)
 Bi Ketalan, Tab (Merrell)
 Bru Col Tab (United Lab)
 W/Atropine methyl nitrate, ox & hog bile
 ext., phenobarbital See G B S, Tab
 (G F Harvey)
 W/Ox bile, steapsie, phenobarbital & homo-
 tropina methylbromide See Oxochalin,
 Tab (Columbus)
BILE EXTRACT (Various Mlr) Pow 1/4 lb.,
 1 lb See: Bilisalin, Tab (Harrower)
 W/Dehydrocholic acid See: Bilicholan, Tab
 (Harrower)
 W/Diactyl Sod Sulfosuccinate See Bidrolor,
 Tab (Armour)
BILE EXTRACT, OX, NF, NNR 1949 Pur-
 fied oxgall
 Armour—Tab 3 gr, 5 gr, Bot 100s, 500s,
 1000s
 Burroughs Wellcome—Tablet 4 gr. Bot
 100s
 Lilly—Enseal 5 gr, Bot 100s, 500s, 1000s
 C. D Smith—Tab 5 gr, Bot 1000s
 Stoddard—Tab 3 gr, Bot 100s, 500s,
 1000s
 See: Bicol, Tab (McNeil)
 Bilein (Abbott)
 Bilisalin, Tab (Harrower)
 Bilsa, Emule (Superior Pharm)
 Chalacole, Cap (Cole Chem)
 Glycotaura, Cap & Tab (Pfyson)
 W/Betaine, choline, Inositol, dehydrocholic
 acid & liver B See: Hepozyn, Tab (Endo-
 crine).
 W/Cascara, aloin See Bilcain, Tab (Cole)
 W/Cellulose, pepsin glutamic acid HCl, pan-
 creatin See: Kanulose Tab (Smith Dorsey)
 W/Calcyntn comp ext., cascara sagrada ext.,
 padophylin & hyoscyamus ext See: Bileo
 Secrn Compound Tablets (First Texas)
 W/Dehydrocholic acid See: Bilicholan, Tab
 (Harrower)
 W/Dehydrocholic acid, homotropine methyl
 bromide, phenobarbital See: Bilomide, Tab
 (Nargine Labs)
 W/Dehydrocholic acid & pancreatic sub-
 stance See: Pancald, Tab (Strassenburgh)
 W/Dehydrocholic acid, pepsin, pancreatin
 See: Neabile, Tab (Haug Drug Co)
 W/Desoxycholic acid, oxidized bile acids,
 pancreatin See: Bilogen, Tab (Organon)

W/Diactyl sod sulfosuccinate See: Bidrolor,
 Tab (Armour)
 W/Iron lactate, columba, chamomile &
 rhubarb See: Cholea, Cap (C. D. Smith).
 W/Ketocholanic acids See: Cholanor, Tab.
 (Maltbie)
 W/Nux vomica ext., Sod bicarb & Cal-
 phosphate See: Dunokol, Tab. (Cole Chem
 Co)
 W/Oxidized bile acids, steapsin, pheno-
 barbital & homotropine methylbromids
 See: Oxochalin, Tab (Columbus).
 W/Pancreatin, duodenal sub., charcoal. See:
 Bidupan, Tab. (Covendish)
 W/Pancreatin, duodenal subst & papain
 See: Miller Zyme, Tab. (Miller)
 W/Pancreatin, papain & duodenal substances
 defatted & desic. See: Korozyme, Tab
 (Karmel Co)
 W/Pancreatin, pepsin & dehydrocholic acid
 See: Ro-Bile, Tab (Rowell)
 W/Phenolphthalein, cascara sagrada See:
 Bilsnots, Tab (Endo)
 W/Phenolphthalein, Sod. salicylate, cascara
 & menthol. See: Chalaxal, Tab (C. D
 Smith)
 W/Phenolphthalein, cascara ext., cap oleo-
 resin, nux vomica ext., papain See: Bicarex,
 Tab (Messengill)
 W/Sod pentobarbital, homotropine methyl
 bromide, dehydrocholic acid See: Homo-
 chol, Tab (Lemmon)
BILE EXT., OX & HOG, W/Atropine methyl
 nitrate, oxidized bile acids, phenobarbital
 See G B S, Tab. (G F. Harvey)
BILE PRODUCTS.
 See Bile Salts
 Dehydrocholic Acid
 Desoxycholic Acid
 Ketocholanic Acid
BILE-LIKE PRODUCTS.
 See: Gallagen, Tab (Messengill)
 Zanchol, Tab (Searle)
BILE SALTS, N N B 1949 Sod glycocholate
 & taurocholate Note also Bile Ext., Ox,
 NNR, and oxidized bile acids
 See: Bilein, Cap & Tab (Abbott)
 Chobile, Tab (Irwin, Nessler)
 Glycotaura, Tab (Hynson).
 Ox Bile Ext., N N B 1949
 Oxidized Bile Acids
 Plebsin, Tab (Plesner)
 W/Belladonna & nux vomica comp (Merrell)
 Bile salts 60 mg, belladonna leaf ext 5
 mg, nux vomica ext 2 mg, phenolphtha-
 lein 30 mg, Sod salicylate 15 mg, aloin
 15 mg/Tab. Bot 1000s
 Uses: Orally, laxative w/antispasmodic
 W/Cascara sagrada ext See: Calapa, Tab
 (Carnrick)

W/Cascara sagrada, capsicum oleoresin, peppermint oil. See: Neo-Tarocal, Tab. (Plessner).

W/Cascara & phenolphthalein. See: Glycolauron w/cascara & phenolphthalein, Tab. (Hynsan).

W/Cascara, phenolphthalein, aloin. See: Alabile, Tab. (Flint, Eaton)

W/Cascara sagrada, phenolphthalein, capsicum oleoresin & peppermint oil. See: Tarocal, Tab. (Plessner).

W/Cascara ext., Sod. salicylate, capsicum oleoresin & peppermint oil. See: Duachal, Tab. (Plessner).

W/Dehydrocholic acid & pancreatic substance. See: Depancol, Tab. (Chilcott).

W/Iron. See: Biron, Cap. (Lilly)

W/Ketocholanic acids, pancreatin, betaine HCl. See: Pradase, Tab. (Miller Labs)

W/2-Methyl naphthoquinone. See: Chola-K, Tab. (Buffington)

W/Pancreas. See: Chobile-Pan, Tab. (Irwin, Nelsler)

W/Pancreatin, sod salicylate, sod. bicarb. See: Cholagestin, liq. (Strang Cobb)

W/Papain, duodenal substance, Sod. bicarb. & phenolphthalein. See: Cholagic, Tab. (Central).

W/Papain, cascara sagrada ext., phenolphthalein & capsicum oleoresin. See: Tarocal Compound, Tab. (Plessner).

W/Phenolphthalein, papain & ginger. See: Cholatal Compound, Tab (Upjohn).

W/Phenolphthalein, capsicum oleoresin & cascara ext. See: Verocalate, Tab. (Standard).

W/Richoleic acids. See: Byleric, Cap. (Merrell)

W/Sod. tauracholate & Sod. glycocholate. See: Cholatal, Cap. (Upjohn)

W/Veratrum viride, Sod. nitrite, ocanite & cascara. Also available w/phenobarbital & rulin. See: Palansars, Tab. (Tailby-Nason).

W/Vit. D.

See: Actaphen, Cap. (Biochemical).

Ataphan, Tab. (Warner-Chilcott)

W/Vit. K. See: Chola K, Tab. (Buffington)

BILE, WHOLE, DESICCATED W/Ketocholanic acid See: Supligril, Tab. (Amer. Ferment).

W/Pancreatin, mycozyme diastase, pepsin, ext. nux vomica. See: Enxobite, Tab. (Mallard)

BILEIN, N.N.R. 1949 (Abbott) Bile salts obtained from ox bile. Cap. 5 gr. Bot. 100s. Enterabs 3 gr. Bot. 100s. Tab. 3 gr. Bot. 100s.

Use: Orally, choleric & chologogue.

W/Dehydrocholic acid See: Hydro-Bilein, Tab. (Abbott).

* Generic name recognized by N.F., U.S.P. or A.M.A.

W/Sorlate & dehydrocholic acid. See: Sorlate-Bilein, Tab. (Abbott).

BILEO-SECRETIN COMPOUND. (First Texas) Ox bile ext. 1 gr., calcynth comp. ext. 1½ gr., cascara sagrada ext. ½ gr., podophyllin 1/12 gr., hyascyamus ext. ¼ gr./Cap. or Tab. Bot. 100s, 500s, 1000s.

Use: Orally, chologogue & laxative.

BILEZYME. (Geriatric) Amylolytic enzyme 30 mg., proteolytic enzyme 6 mg., dehydrocholic acid 200 mg., desoxycholic acid 50 mg./Tab. Bot. 42s, 100s, 500s.

Use: Digestive aid.

BILEZYME PLUS. (Geriatric) Phenobarbital 8 mg., hamotropine methylbromide 2.5 mg., gerinase 30 mg., geriprotase 6 mg., dehydrocholic acid 200 mg., desoxycholic acid 50 mg./Tab. Bot. 42s, 100s.

Use: Constipation

BILIARY COMPLEX. See: Veribile, Tab. (Sharp & Dohme)

BILICHOLAN. (Harrower) Dehydrocholic acid 1½ gr., bile ext. 3½ gr./Tab. Bot. 100s, 1000s.

Use: Orally, hydrocholeric & choleric.

BILISALIN. (Harrower) Bile ext. 0.23 Gm. Bot. 100s.

Use: Orally, chologogue.

BILITAH. (U. S. Standard Prods) Tab. 1000s.

BIOGIN. (Organon) Ox bile ext 120 mg., oxidized mixed ox bile acids 75 mg (equiv. to 1½ gr. dehydrocholic acid), desoxycholic acid 30 mg., pancreatin 250 mg./Tab. Bot. 100s, 1000s.

Use: Orally, choleric & digestant.

BILOPAQUE. (Winthrop) Sodium Tyropanoate.

BILORIC. (Arcum) Pepsin 90 mg.; oxbile 560 mg.; dehydrocholic acid 159 mg.; hamotropine methylbromide 20 mg./Cap. Bot. 100s.

Use: Antispasmodic.

BILSON. (Lilly) Iron bile salts. Pulvules 2½ gr., 5 gr. Bot. 100s, 500s.

Use: Orally, liver dysfunction, gallbladder disease.

BILSA. (Superior Pharm.) Oxgall. Empute 5 gr. Bot. 1000s.

Use: Orally, chologogue.

* **BIMETHOXYCAINE LACTATE** for Bis-[*o*-methoxyphenyl]isopropylamine lactat. See: Isocaine Lactate (Upjohn)

BINASOL. (Rorer) Sod. bismuth tartrate 50 mg/cc Vial 30 cc. Box 12s, 25s, 100s.

Use: LM (1-2 ec. q. 5 days for 12 weeks), syphilis.

BINESTRO. (Ascher) Pat. Estrone Sulf. In

Bot. $\frac{1}{2}$ fl. oz. in plastic atomizer to deliver fine mist spray or drapper bot.

Use: An anti-allergic, antibacterial & decongestant nasal solution.

BIOMYDRIN TABLETS. (Warner-Chilcott) N-acetylamino phenol 150 mg., acetophenetidide 150 mg., phenylpropanolamine HCl 12 mg., phenyltoloxamine dihyd. cit. 6 mg., thonzylamine HCl 6 mg., chlorpheniramine maleate 0.5 mg./Tab. Bot. 24s.

Use: Common cold.

BIOMYDRIN F W/HYDROCORTISONE. (Warner-Chilcott) Hydrocortisone D02%, thonzonium bromide 0.05%, neomycin sulfate 0.1%, gramicidin 0.005%, thonzylamine HCl 1%, phenylephrine HCl 0.25%. Squeeze Bot. 15 cc.

Use: Treatment of hay fever, allergic rhinitis & sinusitis.

BIOMYDRIN OPHTHALMIC. (Warner-Chilcott) Neomycin sulf. 0.1%, gramicidin 0.005%, thonzylamine HCl 0.125%, boric acid 1.2%, phenylephrine HCl 0.125%, thimerosal 0.002%. Sol. drapper Bot. 10 cc.

Use: Ocular infection.

BIOMYDRIN OTIC. (Warner-Chilcott) Thonzonium bromide 0.05%, neomycin sulfate 0.1%, gramicidin 0.005%, thonzylamine HCl 1.0%. Plastic Bot. $\frac{1}{2}$ oz.

Use: Otitis media & otitis externa.

BIOPAN. (Bell Pharmacol) Cobalt chloride 20 mg., copper sulf 4 mg., ferrous sulf. 300 mg., ascorbic acid 20 mg./tab. Bot. 100s.

Use: Anemias.

BIOPAR, N.N.O. 1962. (Armour) Cryst. Vit. B-12 6 mcg., Intrinsic factor 30 mg./Tab. Bot. 30s.

Use: Orally, treatment of anemias.

BIOPAR, FORTE. (Armour) Vit. B-12 w/intrinsic factor concentrate $\frac{1}{2}$ u., Yrt. B-12 25 mcg./Tab. Bot. 30s.

BIOS I. See: Inositol, N.F.

BIOCORB. (Coral) Citrus bioflavonoids 200 mg., Vit. C 200 mg./Cap. Bot. 100s, 1000s.

Use: Antihemorrhagic.

BIOSOL-M. (Upjohn) Neomycin sulfate 200 mg, methscopolamine bromide 2 mg./cc.

Use: Treatment of bacterial diarrheas.

BIO-SORB, N.N.D. 1962. (Ethicon) Absorbable Dusting Powder, U.S.P. Storch-derivative dusting pow., N.N.R. 1954. Canister 5 lb.

Use: Topically, protective & absorptive.

BIOSULFA 125 M. (Upjohn) Pot. penicillin G 125,000 u., sulfadiazine 0.167 Gm., sulfamerazine 0.167 Gm., sulfamethazine 0.167 Gm./tab. Bot. 50s, 500s, 1000s.

Use: Orally (2-4 tabs., then 1-2 q. 4-6 h.). sulfonamide & penicillin therapy.

BIOSULFA 250 M. (Upjohn) Pot. penicillin G 250,000 u. & sulfadiazine, sulfamerazine, sulfamethazine, each 0.167 Gm./Tab. Bot. 50s, 500s, 1000s.

Use: Orally (2-4 stat. then 1-2 q. 4-6 h.), sulfonamide therapy w/penicillin.

BIOSULFA 500 M. (Upjohn) Pot. penicillin G 500,000 u., and sulfadiazine, sulfamerazine, sulfamethazine each 0.167 Gm./tab Bot. 50s.

Use: Sulfonamide therapy w/penicillin.

BIOSYNEPHRINE. (Winthrop Labs.) Phenylephrine HCl 0.5%, thenzylidamine HCl 0.05%, hydrocorfisono 0.02%, Neomycin sulf. 1 mg./cc. polymyxin B sulf. 3000 u./cc. Squeeze Bot. 15 cc.

Use: Nasal spray.

BIOTAL. (Cole) Phenobarbital $\frac{1}{2}$ gr., atropine sulf. 1/1000 gr., bis. subnitrate 2 gr., cerium oxalate 2 gr./Tab. Bot. 100s, 500s, 1000s.

Use: Sedative with antispasmodic.

BIOTANE BOZENGES. (Phillips Roxane) Neomycin 3.5 mg, polymyxin B sulfate 1000 U., benzocaine 5 mg, cetyl pyridinium Cl 2 mg./Lozenge. Bot. 100s.

Use: Irritations of mouth and throat.

BIOTEXIN. See: Nevoblocin.

BIOTIN, N.N.R. 1950, A.O.R.

Breon—Amp. (0.1 mg) 1 cc. Box 12s, 25s, 100s.

Roche Lab—Pow., Vial 100 mg. & 1 Gm. Tab. 1 mg Bot. 30s.

Use: Vitamin therapy.

See: Bioepiderm, Amp. (Breon).

Dermaject (Columbus Pharm.).

BIO-TOSMOSAN HC. (Oaha) Anhydrous glycerol, neomycin (from sulfate) 5.7 mg, gramicidin 7.6 mg, hydrocortisone alcohol 19 mg./15 cc.

Use: Otitis, external otitis and allergic otitis.

BIOTRES. (Central) Bacitracin zinc 200 u., neomycin sulfate 3 mg., polymyxin B 4,000 u., benzalkonium chloride 5 mg./Gm. Tube $\frac{1}{2}$ oz., Box 12s.

Use: Topically, treatment of primary or secondary pyoderms.

BIOTROPIN. (Biophysical) Anterior pituitary gonadotropins. Vial 300 m.u./cc. 10 cc.

Use: I.M. (1-2 cc. over a period of 2-3 months), gonadotropic deficiency.

BIOXATPHEN. (Zemmer) Phenobarbital $\frac{1}{2}$ gr., atropine sulfate 1/1000 gr., bismuth subnitrate 2 gr., cerium oxalate 2 gr./Tab. Bot. 100s, 500s, 1000s.

Use: Orally (1-2 t.i.d.); antispasmodic & sedative.

B.I.P. (Cutter) A vaccine of 6 organisms

containing one million killed bacteria/cc
Vials 5 and 20 cc

Use (0.1 cc 1 M. to start) Weekly

BI-PEN (Reed & Carnrick) Procaine penicillin
G 300,000 u, w/cryst Pot penicillin G
100,000 u/cc Vial 1, 5 or 10 dose

Use: 1 M. (400M u), combined penicillin
therapy

* **BIPERIDEN HCl**, NND 1962 Alpha-[Bicy-
clo[2,2,1]hept-5-en-2-yl] alpha phenyl-1
piperidine propanol HCl (1 bicycloheptenyl
1 phenyl 3 piperidino propanol 1) HCl

See Akineton HCl (Knoll)

* **BIPHENAMINE HYDROCHLORIDE**, β di-
ethylaminoethyl 3 phenylsalicylate hydro-
chloride

See Alvinine, Shampoo (Wampole)

Melsaphine (Sahyun)

BIPHETACEL, (Strosenburgh) Racemic am-
phetamine phosphate monobasic, 5 mg,
dextro amphetamine phosphate 5 mg,
metoprine (methyl tropine nitrate) 1 mg
Sod carboxymethylcellulose 200 mg/Tab
Bot 100s, 1000s

Use Orally (12 1/d, oc), control of
appetite in treatment of obesity

BIPHETAMINE, (Resin) (Strosenburgh) d Am-
phetamine resin and dl amphetamine resin
equal parts Three strengths 7½ mg, 12½
mg, & 20 mg/Cap Bot 100s

Use Appetite control (long acting)

BIPHETAMINE-T, (Strosenburgh) d amphet-
amine 625 mg, dl amphetamine 625 mg,
Tuazole (2 methyl 3 orthotolyl 4 quinazoline
as cation exchange complexes of sulfon-
ated polystyrene) 40 mg/Cap (black and
green 12½) d amphetamine 10 mg, di-
amphetamine 10 mg tuazole 40 mg/Cap
(black and red 20) Bot 50s

Use: Weight control

BIPBOSEP, (Massengill) Neomycin sulf 5
mg, polymyxin B sulf 5000 u, phenyl-
ephine HCl 5 mg, tween 20, thimerosal
1,100,000, as a preservative/cc Plus sqz
bot 15 cc, Bot 8 fl oz

Use: Upper respiratory tract involvements

BIBHUTOX, (Central) Ext of ivy & oak
Amp 1 mg/cc w/d solvent

Use: I.M. (1 cc & repeated in 1 or 2 weeks),
treatment of poison ivy & poison oak der-
matitis

BIROCAP TABLET, (Massengill) Ferrrous lac-
tate 4 gr., cupric gluconate ¼ gr., thia-
mine HCl 4 mg, riboflavin 4 mg, nicotina-
mide 20 mg/Tab Bot 100s, 500s, 1000s

Use: Orally hematitic reconstructive tonic

BISOLVIN IMPROVED, (Irwin, Neisler)
liver subs 7.5 gr, ferrous sulf 1.5 gr, Vit
B 1 1 mg, B 2 1 mg, B-6 0.1 mg, cal

* Generic name recognized by NF, USP or A.M.A.

paniathenolate D 5 mg, niacinamide 5 mg/
Cap Bot 100s, 500s, 1000s

Use: Secondary anemias

B-IRON, (Irwin, Neisler) Ferrous sulf 195
mg, Vit B 1 3.6 mg, B 2 5.4 mg, niacina-
mide 18 mg, B 6 D 1 mg, yeast ext and
liver 70 mg, wheat germ 140 mg/Tab
Bot 100s, 500s, 1000s

Use: Secondary anemias of pregnancy
anorexia

BIBONEX (Robins) Ferrous sulfate azide
3 gr, thiamine HCl 1 mg/Tab Ferrous
sulfate 16 gr, thiamine HCl 6 mg/fl oz
Tab Bot 100s, 1000s Syr Bot 1 pt, 1 gal

Use Orally, secondary anemias

BIS (ACETOXYPHENYL)OXINDOL. See
Diacetylhydroxyphenylisatin

* **BISACODYL**, NND 1962 Di-(p-acetoxy-
phenyl) 2 pyridylmethane See Dulcolax
Tab (Geigy)

BISALATE (Allison Lab) Sod salicylate 5
gr, salicylamide 2½ gr, Sod paramine
benzoate 5 gr, ascorbic acid 50 mg, buta-
barbital Sod ⅓ gr/Tab Bot 100s, 1000s

Use Rheumatic disorders

BISATIN, See Diacetylhydroxyphenylisatin

* **BISHYDROXYCOUMARIN**, USP, NND
1962 Cap, NF Tab USP See Dicumarol
(Various Mfr)

BISILAD, (Central Pharm) Bismuth subgallate
5 gr, colloidal kaolin 165 gr, Lugol's solu-
tion ½ min/fl oz Bot 6 oz, pts, gals

Use: Gastro intestinal protective and ad-
sorbent

BIS-LAN-OL, (U S Standard) Bismuth sub-
salicylate 2 gr, olive oil q s/cc Bot 30 cc,
60 cc, 500 cc

Use Syphilis

BISMAPHEN, (Smith Dorsey) Col carb 25
gr, Mag carb 8 gr, Sod bicarb 8 gr,
kaolin 15 gr, bismuth subgallate 3 gr,
benzocaine ¼ gr, phenobarbital ¼ gr,
belladonna ext 1/6 gr/tsp Paw Can 4 oz

Use: Orally, gastric hyperacidity

BISMABSEN, (Abbott) Bismuth arspenamine
sulfonate (Sulfarsphenamine bismuth) Amp
0.1 Gm Box 10s, 100s

Use: I.M. only, syphilis

BISMOL, (National Drug) Bismuth subsalicy-
late 2 gr, benzocaine 1%, guaiacol ½%,
Sod vanilate ¼%, cottonseed oil q s/
1 cc Injections 25 cc, 100 cc

Use: IV (12 cc every 5-6 days for 5 to
15 doses), syphilis

BISMO-BHODANOL, (Hille) Colloidal bis-
muth Vial ½ gr bismuth/cc 30 cc, 100 cc

Use: I.M. (2 cc weekly for 15 doses),
syphilis

BISOSALICATE COMPOUND (Blue Line)

- Cal. carb., Mag. trisilicate, bismuth subcarb., cerium oxalate, zinc phenolsulphonate. Bot. 1 pt., 1 gal.
Use: Orally, antacid.
- BISMU-C.** (Physiological Chemicals) Bismuth ascorbate 36.5 mg., glycerol 10%, chlorobutanol 0.5%, benzyl alcohol 2%/cc. Amp. 1 cc.
Use: I.M., bismuth in antiluetic therapy.
- BISMUCIN.** (Cole) Iodochlorhydroxyquin 130 mg., bismuth subcarb. 260 mg./tab. Bot. 100s, 500s, 1000s.
Use: Orally, intestinal amebiasis.
- BISMUCOBB.** (Cole Chem.) Bismuth ascorbate 30 mg., chlorobutanol 0.5%, benzyl alcohol 2%, glycerin 10%/cc. Amp. 1 cc. Box 10s, 25s.
Use: I.G. [1 cc. q. 5-6 days for 20 doses], syphilis.
- BISMU-C.** (Physiol. Chem.) Bismuth ascorbate 26.5 mg./cc. Amp. 1 cc.
- BISMU-KINO.** (Wampole) Bismuth oxycarb. 10 gr., eucalyptus gum 6 gr., chloroform 2 min. & phenyl salicylate, camphor, menthol & carminative oils of nutmeg & clove in soothing, demulcent base, w/alcohol 2%/fl. cc. Bot. 4 oz., 1 pt., 1 gal.
Use: Protective, carminative & mild astringent.
- BISMU-RNEA.** (Cole) Oplum pow. 1/40 gr., Krameria ext. 1 gr., bismuth subgallate 2 gr., pectin 1/2 gr., kaolin 5 gr./Tab. Bot. 100s
Use: Diarrhea.
- BISMUTH ARSPHENAMINE SULFONATS.**
See: Bismarsen, Amp. (Abbott).
- BISMUTH ASCORBATE.**
See: Bismu-C, Amp. (Physiological Chemicals),
Bismucarb, Amp. (Cole Chem.)
Bismuth Cevitamate, Amp. (Smith, Miller & Patch)
- BISMUTH BETANAPHTHOL.** (Mallinckrodt) Bi₂O₃ 75%, betanaphthol 15%.
Use: Intestinal astringent.
- BISMUTH-CESIUM OXALATE.** (Sutcliffe & Case) Bismuth Subnitrate 2 gr., Cerium Oxalate 2 gr., Procaine Hydrochloride 1/12 gr./Tablet. Bot. 1000s.
- BISMUTH CEVITAMATE.** (Smith, Miller & Patch) Sod. bismuth cevitamate 36.5 mg./cc., glycerin 10%, benzyl alcohol 2%. Amp 1 cc. Box 5s, 25s, 100s; Vials 10 cc.
Use: I.G. [1 cc. q. 5 days for 15 to 20 doses], syphilis.
- BISMUTH, COLLOIDAL METALLIC.** See: Bismo-Rhadanal, Vial. (Hille)
- * Generic name recognized by N.F., U.S.F. or A.M.A.
- * **BISMUTH GLYCOLYLABSANILATE.** Bis. methyl N-glycolylarsanilate.
See: Glycolarsol, N.F.
MITIBis (Winthrop Labs)
- BISMUTH NYDRORIDE.** See: Bismuth Magma, N.F.
- BISMUTH, INSOLUBLE PRODUCTS.**
See: Bismol (National Drug).
Bismo-Rhadanal (Hille)
Bismuth-Formic Iodide. (Merck Sharp & Dahme)
Bismuth B-Naphthal (Mallinckrodt).
Bismuth Oxyiodide (Mallinckrodt).
Bismuth Subgallate (Various Mfr.).
Bismuth Subsalicylate (Various Mfr.).
Bismuth Tribromophenate (N. Y. Quinine).
- * **BISMUTH MAGMA, N.F., N.N.R. 1949.**
See: Crama-Bismuth (Merck Sharp & Dahme)
Loc-Bismo (Hart).
- W/Kaolin, Pectin. See: Kaolin-Bismuth Pectin Cpd. (Pilman-Moore)
- BISMUTH METAL, AQUEOUS SUSPENSION.**
See: Bismo-Rhadanal, Vial. (Hille)
- BISMUTH OXYCARBONATE.** See: Bismuth Subcarbonate.
- BISMUTH OXYIODIDE.** (Mallinckrodt) Bismuth subiodide. Pow. 1/4 lb., 1 lb.
Use: Local antiseptic.
W/Bis. subgallate, thiogenol, Syntropan, Lanocaine. See: Airof, Supp. (Roche Lab)
- * **BISMUTH POTASSIUM TARTRATE, N.F. X, N.N.B. 1952; Inj., N.F. X, Basic bismuth Pot. bismuthotartrate.**
Brower—25 mg./cc. Amp. 2 cc.
Miller—Amp. (0.016 Gm./cc) 2 cc., Box 12s, 100s; Sol. 30 cc., 60 cc.
Baymer—(2 1/2%) Amp. 2 cc., Box 12s, 25s, 100s.
Use: I.M.; syphilis.
See: Bismuth Soluble, Amp. (Endo).
- BISMUTH RESORCIN SULFONATE.** W/Bis. subgallate, bis. oxyiodide, benzocaine, zinc oxide, peruvian balsam, boric acid. See: Aphca hemorrhoidal aint. (American Pharm.)
- BISMUTH SODIUM ARSANILATE.** See: Bismamar, Amp. & Vial (Kremer-Urban).
- BISMUTH SODIUM TARTRATE, N.N.R. 1952.**
Kirk—Amp. (20 mg) 2 cc., Box 12s, 100s; Vial 2 fl. oz.
Searle—1.5%, 3%. Vial 60 cc.
Use: I.M.; syphilis.
See: Binasol, Vial. (Rorer)
BST Solution. (Canfield)
- W/Pectin. See: Bismol, Liq. (Merrell)
- BISMUTH SODIUM THIOGLYCOLLATE,**
N.N.R. 1952. See: Thio-Bismol, Amp. (Parke, Davis).
- * **BISMUTH SODIUM TRIGLYCOLLAMATE,**

- USP NHD 1962 Tabs USP See Bis
 Irimate Tab (Smith Miller & Patch)
 Use Suppressant for lupus erythematosus
- BISMUTH SUBBENZOATE, (N Y Quinine)**
 Use Dusting powder for wounds
- ***BISMUTH SUBCARBONATE, USP (Various Mfr)** Tab NF
 Use Gastroenteritis, diarrhea
 See Crema-Carbonates (Merck Sharp & Dohme)
- W/Charcoal and ginger See: Harvocarb,
 Cap (G F Harvey)
- W/Iodochlorhydroxyquin See Bismucin, Tab
 (Cole)
- W/Phenyl salicylate, chloroform, eucalyptus
 gum, camphor See: Bismo Kino, Liq (Wam
 pole)
- ***BISMUTH SUBGALLATE, H P (Various Mfr)**
 Dermatol
 Use Various skin conditions
- W/Bis-oxylidide bis resorcin sulfonate
 benzocaine, zinc oxide peruvian balsam,
 boric acid See Aphco hemorroidal oint
 (American Pharm)
- W/Kaolin and benzoic acid See Koo
 Gallate, Liq (Pitman Moore)
- W/Kaolin colloidal opium tr, ginger See
 R. O. Kal Liq (Ulmer)
- W/Opium paw Krameria ext pectin koo
 lin See Rismu Rheo Tab (Cole)
- W/Zinc oxide, bis resorcin comp, nicora
 guan bal, boric acid See: Anusal, Supp
 (Warner)
- BISMUTH SUBIODIDE, See:** Bismuth oxy
 iodide
- ***BISMUTH SUBNITRATE, NF; Tab, NF**
 Use Gastroenteritis, omebic dysentery &
 locally for wounds
 See: Crema-Bismuth Sesp (Merck Sharp &
 Dohme)
- Loc Bismo, Susp (Hort)
- W/Atropine sulfate phenobarbital & cerium
 oxalate See: Bioxatphen Tab (Zemmer)
- W/Magnesium carb, calcem carb See:
 Anticoids, Tab (Gold Leaf)
- ***BISMUTH SUBSALICYLATE, USP; Sesp**
 USP, Basic bismuth salicylate
 Endo—Amp (D) 1 Gm bismuth in peanut
 oil/cc] 2 cc, Box 12s, 25s 100s, Bot 20 cc,
 60 cc, 100 cc.
 Mallinckrodt—Bot 1 oz, Carton ¼ lb,
 1 lb
 Parke, Davis—Amp (D) 13 Gm. in peanut
 oil/cc] 1 cc, Vial 30 cc., 60 cc.; Bot 500 cc.
 User, I.M. (D) 1 Gm weekly for 10 weeks)
 syphilis
- See: Ris Lon-al (U S Standard)
 B smol, Amp (National Drug)
 Bi-Sub-Sal, Susp (Dome Chemical)
- * Generic name recognized by N.F., U.S.P. or A.M.A.
- Stabilol, Amp (Squibb)
- W/Benzocaine, guaiacol & Sod ricinoleate
 in cottonseed oil See: Bismol, Vial (Na
 tional)
- W/Paregoric, pepsin, zinc sullocarbonate,
 solal & methyl salicylate See Parepeptol
 Liq (Chicago Pharm)
- BISMUTH TANNATE, (Various Mfr)** Tanbis-
 muth
 Use Astringent & protective in GI dis
 orders
- BISMUTH TRIBROMOPHENATE, (N Y Qui
 nino)**
 Use Intestinal antiseptic
- BISMUTH VIOLET, (Tabletack)** Bismuth with
 Hexomethyl pararosaniline chloride Oint
 1% Jar 1 oz, 1 lb Pili ½ gr, Bot 100s
 500s Sol ½%, Bot ½ oz, 6 oz, pl Tr
 ½%, Bot 6 oz, 1 pl, also w/benzoic acid
 salicylic acid
 Use Bactericidal & fungicidal
- BISMUTH, WATER-SOLUBLE PRODUCTS,**
 See: Bismuth Arspenamine Sulfonate (Bis
 mersen) (Abbott)
- Bismuth Cevitamate, Amp (C D Smith)
- Bismuth Potassium Tartrate (Various Mfr.)
- Bismuth Sodium Thioglycollate (Parke
 Davis)
- Bismuth Sodium riglycollamate (C D
 Smith)
- Bismuth Potassium Tartrate (Various Mfr)
- Bistrimate Tab (C D Smith)
- Thiabismol, Amp (Parke, Davis)
- Thiopentaton, Amp (Christina)
- BISNAMAR, (Kremer Urban)** Bismuth sod
 arseniate 0.09 Gm/2 cc Amp 2 cc, Box
 25s Vial (multiple dose) 30 cc, 60 cc
 Use I.M. (2 cc m p), syphilis
- BISTRIMATE, NND 1962 (Smith, Miller &
 Patch)** Bismuth sodium triglycollamate, 410
 mg [equivalent to 75 mg elemental bis
 muth]/Tab Bot 100s, 1000s
 Use Chronic sore throat
- BISTRIM BROMIDE, NND 1958 (Squibb)**
 Hexamethanum bromide 25 mg/cc & 100
 mg/cc Vial, 1D cc of each Tab (125 &
 250 mg) Bot 100s, 500s
 User: S.C., I.M., I.V. ganglionic blocking
 agent in peripheral vascular diseases
- RISTRIM CHLORIDE, NND 1958 (Squibb)**
 Hexamethanum chloride Vial 100 mg/cc
 1D cc
 Use: Ganglionic blocking agent
- ***BIS-TROPAMIDE, Tropicamide**
 See Mydracyl, Sol (Alcon)
- BI-SUB-SAL, (Dome Chemical)** Bismuth sub-
 salicylate 130 mg/cc in sesame oil Vial
 50 cc
 User: I.M., treatment of syphilis.

BI-SULFAZINE, N.N.R. 1955. (Warren-Teed)

Dia-mer-sulfanamide, sulfadiazole & sulfamerazine, 50 mg. each/cc. of susp. Bot. 1 pt., 1 gal.

Use: Orally, sulfanamide therapy.

BISULFON, N.N.R. 1955. (Veltex) Dia-mer-

sulfanamides, sulfadiazole & sulfamerazine, 50 mg. each/cc. w/Sad. lactate. Susp., Bot. 1 pt.

Use: Orally, sulfanamide therapy.

BITAVITA. (Chicago) Vit. A, 5000 U.S.P.

units; D, 500 U.S.P. units; C, 50 mg.; B-1, 2 mg.; riboflavin, 2 mg.; niacinamide, 20 mg.; calcium pantothenate, 5 mg.; folic acid, 0.25 mg.; B-6, 1 mg.; E, 0.68 I.U.; B-12, 5 mcq./Tab. Bot. 100s, 1000s, 6000s.

Use: Vitamins supplement.

***BITHIONOL, U.S.P.** Bis(2-hydroxy-3,5-di-

chlorophenyl)sulfide, XI-7.

See: Actamer. (Mansonia Chem.)

Bacteriostat CS-1 (Catalin)

Isothidol liq. (Hillco-Davis Chem.)

Use: Local anti-infective.

W/Allantoin salicylic acid. See: Domerine,

Shampoo (Dome)

W/Resorcinol monoacetate, sulfur. See: Acne-

Dome Creme (Dome).

W/Tyrosinase, Neomycin sulf., phenyltoloxamine dihydrogen citrate, zirconium oxide, dipiperdon HCl. See: Allergesic, lot. (Bryant)

***BITHIONOLATE SODIUM.** Disodium 2,2'-thiobis(4,6-dichlorophenoxy). See: Voecide BN (Vanderbilt Co)

BITRATE. (Arca) Nitroglycerine 1/200 gr. pentaerythritol tetranitrate 20 mg./Tab. Bot. 100s.

Use: Aegle pectoris.

BITBINSIC. (Elder) Ferrous glucoate 40 mg., duodenum whole desic. 260 mg., Vit. B-12 10 mg., cobalt chloride 15 mg./Cap. Bot. 100s, 500s, 1000s

Use: Anti-anemias.

BITUMINOUS ASPHALTITE.

See: Bitupal-HC (Faugera)

Bitupal, oint. (Faugera)

BITUPAL CREAM. (Aveeno or Faugera) Distilled bituminous asphaltite 15%, zinc oxide 8%, water washable base. Jars 2 1/2 oz., 1 lb.

Use: Antipruritic, dermatitis therapy.

BITUPAL-NC. (Aveeno or Faugera) Bituminous asphaltite, 15%; hydrocortisone, 1%; zinc oxide, 8% cream. Tubes 7 & 20 Gm.

Use: Treatment of various dermatoses.

BITUPAL OINTMENT. (Aveeno or Faugera) Distilled bituminous asphaltite. Tube 1 oz. & jar 16 oz.

Use: Treatment of neurodermatitis, atopic

* Generic name recognized by N.F., U.S.P. or A.M.A.

dermatitis, nummular eczema & tar-sensitive dermatoses.

BITUPAL-NC. (Aveeno) Bituminous asphaltite W/Hydrocortisone 1%. Cream Tube 7 Gm. Bituminous asphaltite plain in cream or ointment. Tube 1 oz., Jar 2 1/2 oz. Also available without Hydrocortisone.

Use: Treatment of dermatitis and related skin conditions.

BIVALENT GAS OXIGENE ANTITOXIN, N.N.R. 1952.

BIVAM. (U. S. Vitamin) Citrus bioflavonoid compound 100 mg., Vit. C 100 mg., A 6000 u., D 600 u., B-1 3 mg., B-2 3 mg., B-6 3 mg., B-12 3 mcg., K 1 mg., E 1 I.U., cal. lactate 1 Gm., ferrous glucoate 100 mg., niacinamide 25 mg., cal. pantothenate 5 mg., folic acid 0.5 mg., w/micerals/3 tabs. Bot. 100r, 300s, 1000s

Use: Nutritional supplement

B-JEN. (Fellows) Thiamine HCl 5 mg., fl. ext. gention 0.04 cc., fl. ext. taraxacum 0.06 cc., Phos. acid 0.02 cc., Isocalholic/4 cc.

Elixir, Bot. 8 fl. oz.

Use: Orally, tonic.

B-K-P MIXTURE. (Sutliff & Cass) Kaolin 40 gr., bismuth subcarbonate 20 gr., pectin 10 gr./fl. oz. Bot. 1 pt., 1 gal.

BLACK PEPPER W/Arsenic. See: Asiatic, Tab. (Horton & Converse)

BLACK WIDOW SPIDER ANTIVENIN,

U.S.P., N.N.D. 1962. Antivenin (Latrodetus Mactans). Antitoxic serum (horse). Lyovac, Vial 2.5 cc. (Merck Sharp & Dahme)

Use: I.M., 2.5 cc.

***BLASTOMYCIN.** (Parke, Davis) Filtrate of a culture of blastomyces dermatitidis grown on liq. sythelic media. Box, 2 vials of 0.01 cc. Blastomycin B 1 cc. vial of diluent. Use: Diagnosis of N. Amer. Blastomycosis (Gilchrist's disease).

BLAURONS. (G. F. Harvey) Ferrous carbonate 1 gr., cupric sulfate 0.06 gr., Vit. B-1 1 mg./Pill Bot. 100s, 1000s

Use: Secondary anemias

BLIMERASE. (Tutag) Sulfur 8%, salicylic acid 2.25% w/surface active agent, eucalyptine. Lotion, Bot. 4 oz., 1 pt.

Use: Treatment of acne

BLEPH. (Allergan) Bleph-10 contains sodium sulfacetamide 10%, Bleph-30 contains 30%. Plastic dropper bot. 15 cc.

Use: Ophthalmic.

BLEPHAMIOE. (Allergan) Sod sulfacetamide 10%, prednisolone acetate 0.2%, phenylephrine HCl 0.12% in a plastic polymer and wetting agent. Dropper Bot. 5 cc.

Use: Treatment of blepharitis.

BLESSAMIN. (M. R. Thompson) Vit. A acetate

7500 u., D 2 750 u., E acetate 3 g., K 2 mg, B-1 15 mg, B-2 6 mg, B-6 5 mg, B-12 2 mcg, folic acid 1.5 mg, C 150 mg, niacinamide 45 mg, calcium pantothenate 6 mg, 1 lysine 7.5 mg, protein enzymate 360 mg, 7 minerals and elements /3 cap Jar 90s, 240s
 Use: Prenatal vitamin and mineral supplement

BLEX. (Sutlin & Case) Vit. B-1 1.5 mg, B-2 2 mg, B-6 0.1 mg, niacinamide 10 mg, pantothenate 1 mg, with other 8 complex factors from yeast and liver/Cap Bot 100s, 1000s
 Use: B-complex deficiency

BLEXICON IMPROVED. (Sutlin & Case) Vit B-1 2 mg, B-2 2 mg, B-6 0.5 mg, cal. pantothenate 2 mg, niacinamide 20 mg, Vit B-12 2 mcg./4 cc. Syr 1 pt., 1 gal.
 Use: B-complex deficiency

BUNX LIQUID. (Barnes Hind) Sterile isotonic, alkaline borate buffer containing boric acid and sodium borate Bot 4 oz
 Use: Ophthalmic irrigation

BLISTONS. (Purepac) Chlorbutanol, menthol, besochlorophene, Iantrol Aerosol 2 oz.
 Use: Burn therapy

BLOCKAIN (Breon) Propoxycaïne HCl Sol 1/2%. Bot 30 cc.
 Use: Local anesthetic.

BLOOD, ANTICOAGULANTS. See: Anticoagulants.

BLOOD COAGULATION. See: Hemostatics
BLOOD FRACTIONS. (Cutter).
 See: Albumin Salt Poor, Human Fibrin Foam & Thrombin.

Normal Human Plasma Thrombin, Human
BLOOD GROUPING SERUM, U.S.P., N.N.R. 1955

Hyland—Anti A & Anti B Vial 2 cc., 5 cc
 Anti-A, Anti-B (Group O), Vial 5 cc.

Iederle—Anti A w/methylene blue, for slide test, 10 test Anti A & Anti B, Pkg 2 cc., 5 cc.

Ortho—Anti A absorbed, Vial 2 cc. Anti A or Anti B, Vial 5 cc. Anti A & Anti B, Vial 5 cc.

Marck Sharp & Dohme, N.N.R.—Type A, Vial 1 cc., 5 cc; Type B, Vial 1 cc., 5 cc.
 Use: Diagnostic aids

BLOOD GROUP SPECIFIC SERUM, N.N.R. 1955 (Marck Sharp & Dohme) Isoagglutinins A & B Vial 10 cc

Use: For cross testing of Group O blood for transfusion purposes.

BLOOD PLASMA. See: Plasma.

BLOOD PLASMA SUBSTITUTES

See Dextran-45 (Abbott)
 Expander (Commercial Solvents)
 Gamma (Bartol)

Plavolox. (Wyeth)

PVP Macraze (Schenley)

Vinisol (Abbott)

BLOOD VOLUME DETERMINATION. See: Evans Blue, N.N.R. 1954 (Warner Chilcott)

BLUBAN. (E. L. Patch) d Amphetamine phosphate 5 mg, phenobarbital 1/4 gr., thiamine 5 mg/Tab Bot 100s, 1000s
 Use: Orally (12 11 d), sedative & CNS stimulant

BLUKOR. (Brown Pharm Co) Chorionic gonadotropin 200 IU, thiamine chloride 25 mg, L glutamic acid 52.5 ppm, chlorbutanal 0.5%, procaine HCl 1%/cc. Vials 10 and 25 cc

Use: I.M. Impotence, coronary thrombosis, male climacteric & senility

BLU-PHEN. (Lemmon) Phnobarbital 1/4 gr./Tab Bot 100s

BLUTENE CHLORIDE, N.N.D. 1962 (Abbott) (Tolonium chloride), Toluidine blue O chloride, 3 amine 7 dimethyl amine 2 methyl phenoxethonium salt Tab 100 mg Bot 25s, 100s

Use: Orally (13 daily), functional uterine bleeding (menometrorrhagia)

B-N. (Eric, Kirk & Gary) Boetirocin, 500 units; neomycin sulfate, 5 mg/Oint 1/2 oz
 Use: Dermatitis, Impetigo, burns

B-NUTRON SYRUP. (Nion) Thiamine HCl 2 mg, riboflavin 0.5 mg, pyridoxine 0.2 mg, niacinamide 10 mg, ferrous gluconate 1 gr Mang Sod citrate 1/4 gr/5 cc. Bot 4 oz, 8 oz, 1 pt., 1 gal

Use: Orally, nutritional deficiencies.

B-NUTRON TABLETS. (Nion) Thiamine HCl 2 mg, riboflavin 3 mg, Cal pantothenate 0.2 mg, niacinamide 20 mg, ferrous gluconate 1 gr, Mang Sod citrate 1/4 gr/Tab Bot 100s, 500s

Use: Orally, nutritional deficiency states.

BO-CAR-AL. (Merck Sharp & Dohme) Boric acid, pot alum, phenol, eucalyptus oil methyl salicylate, thymol & menthol Bot 4 oz, 1 lb

Use: Douche

BOEPIDERM. (Breon) d Biotin 0.25 mg/cc Amp 1 cc.
 Use: Treatment of dermatitis & pruritus

BOLAX. (Overseas Pharm) Senna leaves 15%, senna pods, 25%, root of elder, 5% plus herbs of pansy, yarrow, fruits of caraway, root of licorice Pwd Pkg 3.4 oz.

In units of 12
 Use: Laxative

BOLDOOT TOWELETTES. (CMC) Eau de Cologne and a special cosmetic base Towel B, 20s, 50s.
 Use: Deodorize, sterilize and clean, cool and refresh the skin.

BONACAL PLUS. (Kenwood) Vitamins & minerals. Tab. Bot. 100s.

Use: (1 tab. t.i.d.) Multiple vitamins w/ minerals.

BONADENE. (Bonar) Dihydrocodeineone 1/6 gr., pyrilamine maleate 50 mg., di-desoxyephedrine hydrochloride 2 mg., potassium guaicol sulfonate 8 gr., ammonium chloride 8 gr., tarter emetic 1/12 gr., chloroform 2 minims/oz. Syr.

Use: Antitussive.

BONADETTES. N.N.D. 1962. (Pfizer) Meclizine HCl, Tab. 25 mg., Box 6s, 10s.

Use: Treatment of motion sickness.

BONADOXIN. (Roerig) Pyridoxine HCl 50 mg., bonamine 25 mg./Tab., pink & blue. Bot. 25s, 100s.

Drops—Meclizine dihydrochloride 8.33 mg., pyridoxine HCl 16.67 mg./cc. Bot. 30 cc.

Use: Antihousea.

BONADOXIN INTRAMUSCULAR. (Roerig) Meclizine HCl 25 mg., pyridoxine HCl 50 mg., plus lidocaine, benzyl alcohol and citric acid/cc. Amp. 1 cc./5s.

Use: Nausea and vomiting.

BONAMINE, N.N.D. 1960. (Pfizer) New name Bonine. Meclizine, 1-p-chlorobenzhydryl-4-m-methylbenzyl piperazine di HCl long-acting Tab. (25 mg.), Box 8s & 8ol. 100s, 500s; Chewing Tab. (25 mg.). Cortan 8s. Ellx. (12.5 mg./5 cc) 8ol. 1 pl.

Use: Orally (1-2 Tab.), motion sickness.

W/Pyridoxine HCl. See: Bonadoxin, Tab (Pfizer).

BONA-TUSS COUGH SYRUP. (Bonar) Pyrilamine maleate 5 mg., dihydrocodeineone bitartrate 1.666 mg., calcium iodide 1/10 gr., pot. guaicol sulfonate 1.334 gr., menthol 0.208334 gr., chloroform 1/16 minim/5 cc. Bot. 16 oz.

Use: Cough due to common cold.

BONAVITES. (Lloyd, Dabney & Westerfield) Vit. A (acetate) 6000 u., Vit. D 800 u., B-1 1.5 mg., B-2 1.8 mg., niacinamide 15 mg., B-6 1 mg., pantothenic acid 2.5 mg., cabalamin 3mCg., C 40 mg./chewable Tab. Bot. 100s.

Use: Vitamin supplement.

BONAZIN. (Crookes-Barnes) Zinc sulf. 1/4% w/boric acid & methylcellulose, Vial 15 cc.

BONBEC-TD. (Finley) Vit. C 150 mg., B-1 7.5 mg., B-2 5.0 mg., calcium pantothenate 50 mg., B-6 1.0 mg., nicotinamide 250 mg./Cap. Bot. 30s, 100s.

Use: Therapeutic multiple vitamin

BON-DU. (Warner-Teeb) Oxyquinoline benzoate, phenol, menthol, boric acid, thymol, menthol, eucalyptol, methyl salicylate &

alum. Jelly with or without applicator 4 1/4 oz. Pow., Bot. 3 1/2 oz., 16 oz.

Use: Vaginal cleanser.

BONE PHOSPHATE. (Upjohn) Dicalcium phosphate & purified bone phosphate 7 1/2 gr., each/loz. or Gm. of pow. loz., Bot. 1/2 lb. Pow., Bot. 4 oz., 1 lb.

Use: Mineral supplement.

BONE PHOSPHATE. (Levitt) Purified bone phosphate, Tab. 7 1/2 gr. Bot. 100s, 1000s

Use: Mineral supplement.

BONPANT. (C. D. Smith) Col. carb. 300 mg., ferrous fumarate 150 mg., col. ascorbate 60 mg., Vit. D 300 u., B-1 1 mg., B-6 1.5 mg., lemon bioflavonoid complex 50 mg./Tab. Bot. 100s, 1000s.

Use: Vitamin-mineral supplement in pregnancy and lactation.

BONINE, N.N.D. 1962. (Pfizer) Meclizine HCl 25 mg./Tab. or chewable Tab. Bot. 8s, 100s. Elixir (12.5 mg./5 cc.) 8ol. 1 pl.

Use: Motion sickness.

BONMIOTIN. (Crookes-Barnes) Physostigmine salicylate 1/4 and 1/2% w/Philacarpine HCl 2%. Vial 7.5 cc.

Use: Synergistic emetic

BONTRIL. (G. W. Carrick Co.) #1—Carboxyphen chemically equivalent to 10 mg. of dextroamphetamine sulfate, 27 mg. of bulbarbital, 20 mg./Triple-release Tab.

#2—Carboxyphen (chemically equivalent to 15 mg. of dextroamphetamine sulfate), 40 mg. of bulbarbital, 30 mg./Triple-release Tab. Bot. 50s.

Use: Obesity.

BORAX, U.S.P., A.D.R. See: Sodium Borate

BORIC ACID, U.S.P., A.D.R. Oint., N.F. Sol., N.F.; Cryst. or Pow.

Use: Mild antiseptic.

W/Alum, phenol, etc.

See: Al-Bora Phen. (Smith-Dorsey)

Ba-Car-Al. (Merck Sharp & Dahme)

W/Chloretane. See: Boro-Chloretane, Pow. (Parke, Davis)

W/Glyceric. See: Boroglycerin Glycerite, N.F.

BORIC ACID OINTMENT, N.F., A.D.R. (Various Mfr.) 5% to 10%. See: Borofax, Oint. (Burroughs Wellcome)

BORNATE, N.N.D. 1961. (Wyeth) Isobornyl thiocyanacetate. Lot. (5%) 2 fl. oz.

Use: Topically, head, crab & body lice.

BOROCAINE. (Halperin) Procaine borate. Oint. Tube 1 oz., 1 lb.

Use: Local anesthetic.

BOROFAX OINTMENT. (Burroughs Wellcome) Boric acid 5%. Tube 1/4, 1 1/2. Container 1 lb., Ophth oint., Tube 1/6 oz. Use: Topically, for chafed and chapped skin.

*BOROGLYCERIN GLYCERITE, NF A OB
1952 Boric acid 31 parts & glycerine 96
parts.

Use: Externally, in dermatitis

BOBOSAL (C. B. Kendall & Co) Powder
4 oz.

BOBOTANNIC COMPLEX, W/salicylic acid
ethyl alcohol See: Oncha-Phytex (Wynlit)

BOE-20-PHEN OINTMENT, (Ulmer) Boric
acid, zinc oxide, phenol, camphor 1 lb jar

Use: Bacteriostatic dressing ointment

BOVIBON, (Borchardt) Vit A 10 000 u, D
1000 u, thiamine HCl 6 mg, riboflavin 6

mg pyridoxine HCl 08 mg, Cal panto-
thenate 2 mg, niacinamide 30 mg, iron
(as iron & ammonium citrate) 30 mg, Mang

15 mg, mixed tocopherols (as antioxidants)
2 mg, lecithin (as stabilizer) 300 mg / 8 oz
Bot 8 oz, 1 gal

Use: Multivitamin iron supplement

BOE-ZO-PHEN, (Ulmer) Boric acid, zinc
oxide, phenol 0.5%, camphor Ointment
Jar, 1 lb

Use: Antiseptic astringent, and protective

BOTULISM ANTITOXIN, NNR 1949 (Red
cells) 10,000 u each Type A & Type B in
vial

Use: Prophylaxis & treatment of the toxias
of *C. botulinum*, Types A or B

BOUBONAL See: Ethyl Vanillin, USP

BOVILL (Irwin, Neisler) Methylcellulose 450
mg, thiamine mononitrate 2.5 mg / Tab
Bot 100s 500s 1000s

Use: Orally, laxative

BOVINE ALBUMIN 30% SOLUTION (Ar-
mour) Bot w/dropper 5 cc, Vial 50 cc.

Use: As diluent in Bk typing

BOYB CITRA CAPSULE, (Boyle) Hesperid n,
purifed 100 mg, vit C 50 mg, phenyl-
ephrine HCl 5 mg, prophepyridamine

maleate 6.25 mg, methapyrilene HCl 833
mg, pyrilamine maleate 833 mg, soli-
cylamide 200 mg, acetophenetidin 120 mg,
caffeine 30 mg / cap Bot 100s 1000s

Use: Treatment of the common cold

BOYLE CITRA COUGH SYRUP, (Boyle) Di-
hydrocodeine h tartrate 1.66 mg, hes-
perid n 833 mg, phenylephrine HCl 2.5

mg, vit C 30 mg, prophepyridamine
maleate 2.5 mg, pyrilamine maleate 333
mg, pot citrate 150 mg / 5 cc Bot 1 pt,
1 gal

Use: Treatment of the common cold

BOYLE HEMATINIC, (Boyle) Ferrous glu-
conate 27.4 gr, iron 200 mg, Vit B 1 12

mg, B 1 6 mg, niacinamide 108 mg, Vit
C 108 mg, liver desic 1.17 Gm, Vit B 12

6 General name recognized by N.F., USP & A.M.A.

w/intrinsic factor 1/5 u / 6 Tab Bot 100s
1000s

Use: Anemias

BOYLE NEMATINIC LIQUID (Boyle) Ferrous
gluconate 20 gr, Vit B 12 10 mcg, amino

acetic acid 1.5 Gm, Vit B 1 10 mg, B 2
10 mg, B 6 3 mg, niacinamide 60 mg

pyelhenol 65 mg, liver fraction #15 gr,
cobalt 0.05 mg, Mang 5 mg / 8 oz Bot
1 pt, 1 gal

Use: Treatment of iron deficiency anemias

BOYLE HEMATINIC W/VIT. B-12, (Boyle)
Ferrous gluconate 27.6 gr, iron 200 mg

Vit B 1 12 mg, B 2 6 mg, niacinamide
108 mg, Vit C 108 mg, liver desic 1.2

Gm Vit B 12 60 mcg, Vit. B 12 w/in-
trinsic factor 2/5 u / 6 Tab Bot 100s, 1000s

Use: Anemias

BOYLE PBE-NATAL, (Boyle) Dicalcium phos-
phate 0.75 Gm, ferrous gluconate 45 mg,
Vit A 6000 u, D 600 u, B 1 4 mg, B 2

2.5 mg, niacinamide 50 mg, Vit C 100
mg, Pot Iodide 0.15 mg, copper sulfate

10 mg, Mag sulfate 15 mg, w/cobalt,
Mang, zinc, molybdenum & nickel / 3 Tab
Bot 100s, 1000s.

Use: Pregnancy & lactation.

BOYLE THERAPEUTIC NEMATINIC, (Boyle)
Vit B 12 w/intrinsic factor 1/5 oral u, iron

200 mg, Vit B 12 30 mcg, folic acid 1.5
mg, Vit B 1 6 mg, B 2 3 mg, niacinamide

54 mg, Vit C 150 mg, copper sulfate 1
mg, cobalt sulfate 1 mg, liver desic 75

mg / Tab Bot 30s, 500s

Use: (1-3 daily) Anemias

BOZELLE, (Dumas Wilson) Tube 1 1/2 oz
Use: Emollient Creme

B-PLEX ELIXIR, (Wyeth) Vit B 1 24 mg
B 2 1 mg, brewer's yeast 15.6 Gm, nicot-
inic acid 8 mg / 4 dr Bot 8 fl oz, 1 gal

Use: Orally, vitamin B therapy

B PLEX 25, (Philadelphia Ampoule Labs)
Vit B 1 25 mg, B 2 1 mg, B 6 1 mg,
niacinamide 75 mg, cal pantothenate 1

mg / cc Vial 30 cc

Use: Vitamin B complex therapy

B PLEX-100, (Philadelphia Ampoule Labs)
Vit B 1 100 mg, B 2 2 mg, B 6 2 mg,
niacinamide 125 mg, cal pantothenate 10

mg / cc Vial 30 cc

Use: Vitamin B complex therapy

B PLEX FORTIFIED, (Philadelphia Ampoule
Labs) Vit B 1 100 mg, B 2 2 mg, B 6 5 mg,
niacinamide 125 mg, cal pantothenate 1

mg, B 12 10 mcg / cc Vial 30 cc

Use: Vitamin B complex therapy

B PLEX, SPECIAL FORMULA, (Philadelphia
Ampoule Labs) Vit B 1 10 mg, B 6 5 mg

mg, B 12 10 mcg / cc Vial 30 cc

Use: Vitamin B complex therapy

B-2 2 mg, niacinamide 50 mg, cal. pantothenate 10 mg./cc. Vial 10 cc.
Use: Dietary supplement.

B-P-5 MIXTURE. (Sulfin & Case) Bismuth salicylate 520 mg., solal 104 mg., hecure opium camphorated 375 cc., zinc phenol-sulfonate 52 mg., pepsin 260 mg., aromatics q.s./fl. oz. Bot. 1 pt., 1 gal.
Use: Diarrhea

BRACODIN. (McNeil) Ephedrine HCl 1/16 gr., quebracha lr. & Pot. iodide/S cc. Bot. 1 pt., 1 gal.
Use: Orally, non-narcotic cough therapy.

BRADOSOL BROMIDE. (Ciba) Dodecyl-dimethyl (2-phenoxyethyl) ammonium bromide 1.5 mg. and beezeacaine 2.5 mg./lozenge. Pkg. 24s.
Use: Oral infections.

BRAL. (Cole) Acetylsalicylic acid 5 gr., Sod. bromide 1 1/2 gr., Pot. bromide 1 1/2 gr., ammonium bromide 1 1/2 gr., caffeine alkaloid 1/4 gr./Tab. Bot. 100s.
Use: Orally (1-2 i.i.d.), sedative & analgesic.

BRAMOSO. (Tilden) Sod. bromide 14 gr., antipyrine 24 gr., Sod. salicylate 28 gr., caffeine 4 gr./oz. Bot. 1 pt., 1 gal.
Use: Orally (1 dr. q. 3-4 h. p.r.n.), sedative & analgesic.

BRASIVOL. (Stiefel) Aluminum oxide, hexachlorophene 1%, in a hypoallergenic base of neutral soap and surfactant cleansers. Fine—Jars 5 1/2 oz. Medium—Jars 6 1/2 oz. Rough—Jars 7 oz.

BRASIVOL BASE. (Stiefel Labs.) Cleansing paste containing a synergistic mixture of soaps and detergents, with hexachlorophene 1%. Jar 4 1/2 oz
Use: Degreasing cleanser; treatment for acne.

BREDATIVE-DHC. (Breco) Pentobarbital Sod. 1/4 gr., phenobarbital 1/4 gr., dihydrocodeine 5 mg. Tab., Bot. 100s, 500s.
Use: Analgesic, sedative.

BREMIL. (Borden) Fat 27.5%, protein 11.7%, carbohydrate 54.6%, minerals 3.9%, water 2.3%, Col. 5.1 Gm., Vit. A 25.7 u., D 6.43 u., thiamine 0.003 mg., riboflavin 0.008 mg., niacin 0.048 mg., Vit. C 0.40 mg., iron 0.064 mg./Gm. Cans 1 lb.
Use: Infant feeding.

BREMIL LIQUID. (Borden) Vit. A 2500 u., D 800 u., C 50 mg., B-1 0.4 mg., B-2 1 mg., elacin 6 mg., B-6 0.4 mg W/protein, fat, carbohydrate, calcium, phosphorus/qt. Cae 13 oz.
Use: Infant feeding.

BREOM. (Breon) Hydroxyquinoline beezeate 0.2% in aromatic quince & trogeacanth ve-

hide, alcohol 12%, methylparaben (0.1%). Plastic squeeze Bot. 2 oz.
Use: Skin antiseptic.

BREONEX-AO. (Breon) Vit. B-1 100 mg, B-2 5 mg., niacinamide 100 mg., B-6 5 mg., pantothenol 5 mg./cc. Vial 10 cc.
Use: I.M., 1-2 cc., vitamin B complex therapy.

BREONEX-C. (Breco) Vit. B-1 5 mg, B-2 3 mg, B-6 3 mg, C 50 mg, niacinamide 40 mg., pantothenol 3 mg./cc. Vial. 10 cc.
Use: Vitamin B deficiencies.

BREONEX-L. (Breon) Thiamine HCl 150 mg., riboflavin 50 mg., pyridoxine HCl 25 mg., Cal. pantothenate 25 mg., niacinamide 500 mg., B-12 10 mcg./Vial 10 cc. Pkg. 1 e., 6 w., 25 u.
Uses I.M., I.V., high potency vitamin B complex therapy.

BREONEX-L w/ Vit. C. (Breon) Breonex-L w/ Vit. C 50 mg./cc. Pkg. of 1, 6, 25 single doses.
Use: High potency B complex.

BREONEX-M. (Breon) Contains twice the amount of Breonex-L plus Vit. B-12 50 mcg. Vial w/diluent 30 cc.
Use: Multiple vitamin deficiencies.

BREONEX STRONGER. (Breon) Thiamine HCl 10 mg., riboflavin 0.5 mg., pyridoxine HCl 1 mg., niacinamide 25 mg./cc. Amp. 2 cc., Box 25s.
Use: I.M., I.V., vitamin B complex therapy.

BRETYLIUM TOSYLATE. N- α -bromobenzyl N-ethyl-N,N-dimethyl ammonium-p-toluene sulfonate. See: Dorenthln, Tab. (Burroughs Wellcome)

BREVITAL. N.N.D. 1962. (Lilly) Methohexital. Sodium α -di-1-methyl-5-[1-methyl-2-pentylyl]-5-allyl-barbiturate. Cryst. pow. form. In 500 mg rubber-stoppered amp. and 2.5 Gm. glass-sealed amp. Pkg. of 1 and 25 Amp. 5 Mg. Amp. and 25s. 2.5 Gm. Amp. and 25s. Dry ampoules. 1% sat—250 cc. of diluent to 2.5 Gm. ampoule; 500 cc. to 5 Gm. ampoule.
Use: (1%) Ultrashort-acting intravenous anesthetic.

BREVOZOL. (Lemmon) Tyrothricin 0.1%, beezeacaine 2.5%, 9-aminoacridine HCl 0.2%/Gm. Diet. tube 1/2 oz., 1 cc.
Use: Topically; antihistaminic w/local anesthetic & antiseptic.

BREWER'S YEAST. N.N.R. 1952 (Various Mfr) Pow., Tab. 6 gr. & 7 1/2 gr.
W/Diacetyldihydroxyphenylisotln & sod. carboxymethylcellulose. See: Zylax. Tab (Glidden)
W/Sod. carboxymethylcellulose. See: Zymelox. Tab. (Glidden)

BRIDASE. (Vio Bio Corp) Powder, 10 Gm.

BRILLIANT GREEN W/Benzocaine and Mag Sulf See: Osmopak (Irwin Neisler)

W/Acriflavine & methylrosaniline chloride Mix, N.N.R. 1953 See Dymixal, Faw (Mc Neil)

BRISTALIN. (Bristol) Bristamin 125 mg. Ipecac R ext, ammonium chloride, Sod citrate & menthol/S cc Syr Rat 4 oz
Use: Orally (1 dr q 2-3 h) antihistamine cough preparation

BRISTAMIN. (Bristol) Phenyltoloxamine dihydrogen citrate Lot 1%, Rat 6 R oz Tab 25 mg, 50 mg Tab Bat 100s
See: PRN, Tab (Bristol)

Use: Antihistamine therapy
W/Amm chlor, Ipecac R ext, sod cit menthol See: Bristalin, Syr (Bristol)

W/APC (Bristol) Bristamin 25 mg, aspirin 3 1/2 gr, acetophenetidin 2 1/2 gr, caffeine 1/2 gr/Tab Bat 100s

Use: (111d) Colds & allergic symptoms

W/APC & Procaine Pan O
See: A P Cillin, Tab (White)
Sristaphen Tab (Bristol)

W/Dihydrostreptomycin sulfate See: Dihydrostreptomycin Otic W/Sristamin, liq (Bristol)

BRISTAMIN LOTION W/ZIRCONIUM. (Bristol) Phenyltoloxamine dihydrogen citrate 1% with zirconium oxide 4% Lotion, Bot 6 oz See: Zirnox Let (Bristol)
Use: Treatment of Rheu dermatitis

BRISTAPIN (Bristol) Procaine penicillin G 100,000 u, aspirin 2 1/2 gr, phenacetin 2 gr, caffeine 1/2 gr, bristamin dihydrogen citrate 25 mg/Tab Sol 24s 100s
Use: Orally, penicillin w/antihistamine & analgesic action

BRISTAPEN-200 (Bristol) Procaine penicillin G 200M u, aspirin 2 1/2 gr, phenacetin 2 gr, caffeine 1/2 gr, bristamin dihydrogen citrate 25 mg/Tab Bat 24s 100s
Use: Orally, penicillin w/antihistamine and analgesic action

BRITISH ANTI-LEWISITE. Dimercapof See: RAL

BRO-R INJECTION (Broemmel) B-1 100 mg, R 2 2 mg, B-4 5 mg, panthenol 10 mg, niacinamide 100 mg, Vit B 12 30 mcg/ml Vial 10 ml
Use: I.M., S.C. treatment of vitamin deficiencies

BROCLORMUS. (Marrang M) Chloral hydrate 15 R, potassium bromide 15 gr, hyoscymus tr 7 min/S cc Elia Bot 1 pt, 1 gal
Use: Antiparalmed c, sedative and hypnotic

BRODATI (Wende) Ephedrine sulfate 25 mg, acetylcarbromol, 260 mg/Tab Bat 100s 1000s
Use: Bronchial asthma

BRODITAL. (U S Standard) Phenobarbital 1/4 gr, Pot sodida 2 1/2 gr, theobromine 2 1/2 gr/Tab Bat 100s, 1000s
Use: Orally (1 2 q 3 h) analgesic & sedative

BRO-LAC (Broemmel) Plastic drop Bat 15 cc
Use: Artificial tears sterile solution

BROMACRYLIDE. (Lederle Lab) N-(acrylamidomethyl) 3 bromopropionamide

BROMALEATE. A mixture of 2 amino-2 methyl 1 propanol and 8 bromotheophylline 50 mg (Pamobrom) with pyristamine moleate 30 mg

See: Neo Bromth, Tab (Broyten)
Neo Mensalem, Tab (Walker)
Neo Parbrom, Tab (Central)
Neo Pyrapor, Tab (Ulmer)

* **BROMAZINE** See Ambodryl (Forke, Davis)

BROMALATE. (Endo) Col gluconate 0.5 Gm, strontium bromide 0.5 Gm dextrose 1 Gm / Amp 10 cc Rox 6s, 25s, 100s
Use: I.V., antipruritic & sedative

* **BROMCLORENONE**, 6 bromo 5 chloro 2 benzoxalolone See: Vinyzene (Scientific Chemicals)

* **BROMELAINS**. (Dole, H, Pineapple Co.) Concentrate of proteolytic enzymes derived from pineapple plant Ananas Sativus See: Anonase (Rorer)

BROMETHOL. See: Avertin, Sol (Winthrop Labs)

BROMIC COMPOUND. (Pitman Moore) Sod bromide 14 gr, Cal bromide 1/4 gr, ammonium bromide 1/2 gr, lithium bromide 1/2 gr, Pot bromide 1/2 gr, cypridium 3 gr/5 cc Bat 1 pt, 1 gal
Use: Orally (1 2 dr q 3 h), sedative & hypnotic

BROMIDES NO. 1. (Merck Sharp & Dohme) Pot bromide 26 Gm, sod bromide 26 Gm, ammonium bromide 18 Gm, cal bromide D75 Gm, alcohol 14%/30 cc Bat 1 pt, 1 gal
Use: Orally (1 2 dr, prn) sedative & hypnotic

BROMIDES COMPOUND SPECIAL (Sullif & Case) Sod bromide 374 Gm, pot bromide 375 Gm, cal bromide 65 mg, lithium bromide 65 mg, ammonium bromide 130 mg, strawberry flavor/R oz Bat 1 pt, 1 gal
Use: Sedative

BROMIDES.
See: Bromazo (Tilden)
Bromo Flora (Warren Teed)

* Generic name recognized by N.F., U.S.P. or A.M.A.

- Elix. Liq. (Chicago Pharm.)
 Lonobrom, El. (Lonnell)
 Nervie, Liq. (Honce)
 Nervasine, Liq. (Dios)
 Neurabromyds, Liq. (Rohrer)
 Peacocks Bromide (Peacock)
 Pentabramides, Liq. (Merrill)
 Solomides, Tab. (Merck Sharp & Dohme)
- BROMIDES W/Alkalinizing agents.**
 See: Bromianyl, (Upjohn)
- BROMIDES W/Aspirin & Caffeine.**
 See: Abac, Cap. (Massengill)
 ACT, Tab. (Tildee)
 Bral, Tab. (Cole).
- BROMIDES W/Borbiturates.**
 See: Neo-Sedophen, Liq. (C. D. Smith)
 Nurodal, Liq. (Buffington)
 Sedalco, Liq. (Mallard)
- W/Cypripedium** See: Bramic Comp., Liq. (Pitman-Moore)
- W/Niacinamide HBr.** See: Broniacin, Liq. & Tab. (Massengill)
- W/Phenobarbital sod.**
 See: Abactal, Tab. (Massengill)
 Bromiphen, Tab. (McNeil)
 Bromobarb, El. (Hart)
 Sedophen, Liq. (C. D. Smith)
- BROMIDE SALTS.**
 See: Ammonium
 Calcium
 Ferrus
 Lithium
 Potassium
 Sodium
 Strontium
- BROMIDIA, (Bottle)** Chloral hydrate 1½ gr., Pot. bromide ½ gr., hyoscyamus ext. ¼ gr./5 cc. Bot. 4 oz., 1 pt.
 Use: Orally (1 dr. p. r. n.), sedative.
- * **BROMINDIONE, (U.S., Vitamin & Pharmaceutical Co.)** 2-(p bromophenyl)-1,3 indandione.
 See: Halinone.
 Use: Anticoagulant.
- BROMINYL, (Upjohn)** Sod. bromide 9.75 gr., col. bromide 3.75 gr., pot. bromide 2½ gr., sod. citrate, pot. tartrate, sod. lactate & mag. sulfate/isp. Bot. 4 oz.
 Use: Orally, sedative.
- W/Barbital, (Upjohn)** Same as Brominyl, w/ barbital 5 gr. added/isp. Bot. 2 oz.
 Use: Orally, sedative.
- BROMIPHEN, (McNeil)** Phenobarbital sod. ¼ gr., sod. bromide 3¾ gr., pot. bromide 3¾ gr./Tab. Bot. 100s, 1000s
 Use: Orally (1-2 q. 3 h.), sedative & hypnotic.
- * **BROMISOVALUM, N.F., N.N.R. 1951.**
 A.D.R. Tab. N.F. α-Bromoisovaleryl urea.
 * Generic name recognized by N.F., U.S.P. or A.M.A.
- Premo-Pow.** Bot. 1 oz., ¼ & 1 lb. Tab. [5 gr.] Bot. 100s, 1000s.
 See: Bramaral, N.N.R. (Knoll)
 Isoval, Tab. (Mission Pharm.)
 Kortryl (Direx Lab)
- W/Ethyl salicylate** carbonate, acetophenelidid & caffeine. See: Acetabra, Pediatob No. 1, (Columbus Pharm.)
 Use: Sedative & hypnotic.
W/Acetylcarbamol.
 See: Sedasyon, Tab. (Tailby-Nason).
 Daysed, Tab. (Tailby-Nosoe).
- W/Carbamol.** See: Tabarea, Tab. (Table-rock).
- W/Homatropine methylbromide, hyoscyamine HBr & scopolamine HBr.** See: Tadona, Tab. (Tanlerack Labs.).
W/Octie. See: Voletin, Tab. (Keoll)
- BROMORARR, (Hart)** Pheoobarbital 2 grs., sod. bromide 60 gr. Elix. Bot. 1 pt.
 Use: Sedative, hypnotic, antispasmodic and colmative.
- BROMOCHOLINE BROMIDE, (Anglo-French).**
 Not marketed. Used in Hypoton Tab.
- W/Chloral hydrate & methylacetylcholine.**
 See Hypoton, Tab. (Anglo-French).
- BROMODIETHYLACETYUREA.** See: Carbromal, N.F.
- * **BROMODIPHENHYDRAMINE HCl.** 2-(4-Bromodiphenylmethoxy)ethylidimethylamine. 4-Bromo-α-phenylbenzyl 2-dimethylaminoethyl ether. See: Ambodryl. (Parke, Davis)
- BROMO-FLORA, (Warren-Teed)** Strontium bromide 10.37 Gm., passiflora 5.18 Gm., alcohol 5% & aromatics/n. oz. Rd. 1 pt., 1 gal.
 Use: Orally (1-2 dr. q. 3-4 h.), sedative.
- BROMISOVALERYL UREA.** Alpha, bromoisovaleryl urea. See: Bromisovalum, N.N.R.
- BROMOPHIN.** See: Apomorphine HCl (Merck).
- BROMOSALICYL ALCOHOL.** See: Bromsalizol, Amp. (Hynson, Wescott & Dunning)
- BROMOSALIGENIN.** Bromosalicyl alcohol. See: Bromsalizol, Prep. (Hynson, Wescott & Dunesig)
- BROMOSPAZ, (Chicago Pharm.)** Homatropine methylbromide 5 mg., phenobarbital 16.2 mg./tab. Bot. 100s.
 Use: Antispasmodic with sedative.
- B-R BROMOTHEOPHYLLINE** W/2-amino-2-methyl-1-propanol. See: Pamobrom
- B-R BROMOTHEOPHYLLINE** W/Ethisterone, chlorpropenpyridamine maleate, salicylamide. See: C, Compound (Schering)
- BROMOTHEOPHYLLINATE AMINOISORUTANOL** See: Pamobrom
- BROMOTHEOPHYLLINATE PYRILAMINE.**
 See: Pyrabrom

BROMOTHEOPHYLLINATE PYRANISAMINE. See Pyrobrom

BROMPHENIRAMINE MALEATE, NND
1962 (3 p bromophenyl 3 pyrid 2' ypropyl
dimethylamine Parabromdylamine maleate
See Dimetane (Robins)

W/Phenylephrine HCl, phenylpropranolamine
HCl See Dimetapp Extentabs, Tabs,
Elixir (Robins)

BROMSALIZOL. (Hynson, Westcott & Dunning) Monobromosalicyl alcohol (Bromosaligenin) Sol 4% in oil Amp 10 cc; Tab 5 gr Bot 40s

Use: Orally ec I.M. (5 10 gr tid), anti
spasmodic.

BROMSULPHALEIN, NNR 1952 (Hynson,
Westcott & Dunning) Sulfobromophtholein
Sod USP, Phenottetramphtholein diso-
dium Amp (5% sol) 3 cc, 7.5 cc, Box
10s, 50s, 10 cc, Box 10s, 50s

Use: Diagnostic aid
BROMTH. (Broyten) Pyrobrom Tab 50 mg
Bot 100s

Use: Orally (1 2 pr n), antinausea & ma-
tion sickness

BROMUAL (Haug) Potassium guaiaecol sul-
fonate 2 gr, ammonium salicylate 4 gr,
lobelia tincture 20 minims, Tolu balsam
syrup 180 minims/oz Bot 3 oz

Use: Expectorant
BROMURAL (Kealf) NNR 1951, ADR
1955 (Bromisuralum, NF) Pow, Bot 1 &
8 oz Tab (5 gr) Tube 10s, Bot 100s, 500s,
1000s

Use: Orally (5 10 gr tid & hs); quick
acting sedative & hypnotic.

W/Octie See: Valactin, Tab (Kealf)

BRONCHONIST. (Sull ff & Case) Pot guaia-
ecol sulfonate 500 mg, pyrilamine 75 mg,
d desoxyephedrine HCl 3 mg/R oz Bot
1 pt, 1 gal.

Use: Orally (1 2 dr mp), see narcotic
ant histamine expectorant.

BRONCHONIST-NEO (Sull ff & Case) Glyc-
eryl guaiaecolate 250 mg, phenylephrine
HCl 25 mg, pyrilamine maleate 30 mg,
methoxyphenylene isonate 30 mg/B oz Bot
4 cc, pt, gal

BRONCHO-RECTAL SUPP. (5 mpson) Cre-
osote carbonate, 10 min; guaiaecol, 1/4 gr/
Supp Box 14s

Use: Decongestant

BRONCHO-TUSSIN NO. 1. (First Texas)
Morphine acetate 1/4 gr, terpin hydrate
4 gr, ammonium hypophosphite 8 gr, Pot
guaiaecol sulfonate 8 gr, alcohol 40%/R
oz Bot 1 pt, 1 gal

Use: Orally (1 dr q 3 h), cough sedative
& expectorant

* Generic name recognized by N.F.-U.S.P. or A.M.A.

BRONCHO-TUSSIN NO. 2. (First Texas)

Codaine phosphate 1 gr, terpin hydrate
4 gr ammonium hypophosphite 8 gr, Pot
guaiaecol sulfonate 8 gr, alcohol 40%/R
oz Bot 1 pt, 1 gal

Use: Orally (1 2 dr q 3 4 h), cough sedo-
tive & expectorant

BRONDECON. (Warner Chilcott) Oxtriphyt
line 200 mg, glyceryl guaiaecolate 100 mg/
Tab ar 2 tsp of elixir. Bot 100s, 1 pt

Use: Expectorant—bronchodilator.

BRONIACIN. (Massengill) Elixir: Ammonium
bromide 0.5 Gm, Pot bromide 0.5 Gm
eiacinamide HBr 20 mg/dr. Bot 1 pt, 1
gal Tab, Ammonium bromide 0.25 Gm
Pot bromide 0.25 Gm, niacinamide HBr
10 mg/Tab Bot 100s, 1000s

Use: Sedative

BRONKEPHRINE. (Breon) Ethylnorepinephrine
HCl 2 mg/cc isotonic sol Vial 10 cc

Use: S.C., I.M., I.V.; bronchodilatory in
bronchial asthma

BRONKODYL. (Columbus) Isopropylarterenal
HCl 10 mg (outer core), phenylpropranol-
amine HCl 25 mg, glyceryl guaiaecolate
100 mg (inner core)/Tab Bot 100s, 500s

Use: Asthma attacks

BRONKODYL PEDIATAB 17. (Columbus)
Isopropylarterenal HCl 5 mg, phenylpro-
pranolamine HCl 15 mg, glyceryl guaiaecol-
ate 60 mg/Tab Bot #17/100s

Use: Acute or chronic asthma

BRONKOMETER. (Breon) Isoethorine meth-
anesulfonate 0.6%, phenylephrine HCl
0.125%, thenyldiamine HCl 0.05%, free
propellants

Use: Bronchial asthma

BRONKOSPRAY. (Breon) Isoethorine HCl
1.0%, phenylephrine HCl 0.25%, thenyldi-
amine 1.0%, in buffered saline solution

Use: Bronchial asthma

BRONKOTABS (Breon) Ephedrine sulf 24
mg, theophylline 100 mg, thenyldiamine
HCl 10 mg, glyceryl guaiaecolate 100 mg;
phenobarb 8 mg/Tab Bots 100s

Use: Asthma

BRONKOTAS ELIXIR. (Breon) Glyceryl guaia-
ecolate, 50 mg, ephedrine sulfate 12 mg,
theophylline 15 mg, phenobarbital 4 mg,
chlorpheniramine maleate 1 mg/5 cc. Bot
pts

Use: Bronchial asthma

BRO-PARIN. (Broemmel Pharm) Heparin so-
dium 2000 u, polymyxin B sulfate 100 000
u, neomycin 20 mg, hydrocortisone 5 mg/
cc. in a polyethylene glycol 400 base
Dropper vial 5 cc

Use: External otitis, chronic weeping or
zema eczematoid dermatitis

BBO-PARIN-N. (Broemmel) Polymyxin B sulfate 50,000 u., neomycin 10 mg., heparin sod. 1000 u., hydrocortisone 5 mg., chlorbutanol 0.10%, in polyethylene glycol 400 base/cc., Drop Vial 5 cc.
Use: External otitis.

BRO-TUSSA. (Brown) Chlorpheniramine maleate 1 mg., phenylephrine 10 mg., caffeine 45 mg., pyrilamine maleate 12.5 mg., methatrylene HCl 12.5 mg., dextromethorphan HBr 10 mg., ascorbic acid 200 mg./Tab. Bot. 30s, 100s.
Use: Decongestant.

BROWN COMPOUND W/AMMONIUM CHLORIDE LOZENGE. (Parke, Davis) Fid. ext. glycyrrhiza 12 min., ammonium chloride 3 gr., antimony and potassium tartrate 1/40 gr., benzoic acid 1/20 gr., camphor q.s., oil anise q.s./Loz. Bot. 30s, 1 lb.
Use: Stimulate secretion in acute bronchitis.

BROWN COMPOUND W/AMMONIUM CHLORIDE WITHOUT OPIUM. (Merrell) Benzoic acid 1.8 mg., tartar emetic 0.9 mg., licorice ext. D.11 Gm., ammonium chloride 60 mg., oil anise q.s., camphor q.s./Tab. Bot. 1000s.
Use: Orally, cough preparation.

BROWN MIXTURE. (Merrell) Opium tr., camphorated, 12 cc., ethyl nitrite spirit 3 cc., antimony & Pot. tartrate 0.024 Gm., flax. glycyrrhiza 12 cc., alcohol 10%/100 cc./Bot. 1 gal.
Use: Orally, cough preparation.

BROWN MIXTURE. (Merck Sharp & Dohme) Mixture opium & glycyrrhiza, Comp. N.F. Bot. 1 pl., 1 gal.
Use: Orally, cough preparation.

BROXIL. (Beechem Research Lab) Pot. 6-(alpha phenoxypropionamide) penicillanic acid.
Use: Synthetic penicillin.

BROXOLIN. (Breon) Oxyquinoline sulfate, boric acid, Sod. chloride & Mag sulfate. Pow. Bot. 6 oz.
Use: Vaginal douche cleanser.

BROXOLIN. (Breon) Glycobiarsol 250 mg./3 cc. Cream, Tube 110 cc. w/applicator
Use: Trichomonos vaginalis, monilia albicans

BROZANTHINE. (Blue Line) Corbramol 1 gr., theobromine 5 gr., Cal.-urea-iodide 2 1/3 gr./Tab. Bot. 100s, 1000s.
Use: (1-2 t.i.d) Sedative & myocardial stimulant.

BRUCCELLA ABORTUS. (Parke, Davis) Vial 5 cc.
Use: Diagnosis of undulant fever in man.

BRUCCELLA ABORTUS ANTIGEN (FEBBILE). (Ledele) Vial 5 cc.

Use: Diagnostic reagent for macroscopic rapid slide agglutination method.

BRUCCELLA ANTIGEN RAPID. (Pitman-Moore) Vial 5 cc.

Use: For brucellosis in acute or chronic form.

BRUCCELLA VACCINE, N.N.R. 1953. Undulant fever vaccine.

National Drug—2500 million killed Br. abortus & Br. suis/cc. Vial 30 cc.

Pitman-Moore—1000 million killed Br. abortus & Br. suis/cc. Vial 20 cc.

Use: I.M., S.C., undulant fever.

BRUCCELLERGEN. (Merck Sharp & Dohme) Brucella protein nucleate. Vial 1 cc.

Use: Diagnosis of brucellosis (undulant fever) by the intradermal test.

BRUCILLIN ANTIGEN. (Merck Sharp & Dohme) Prepared from Br. abortus, Br. melitensis & Br. suis. Vial 2 cc.

Use: Active phase of brucellosis.

BRU-COL. (United Labs) Mixed oxidized bile acids. Tab. 3 1/2 gr. Bot. 100s, 1000s.

Use: Cholecystitis, cholangitis & hepatic dysfunction.

B-SOBBIC. (Deseret) Vit. C 500 mg., B-1 25 mg., B-2 10 mg., B-12 2 mcg., calcium pantothenate 10 mg., nicotinamide 75 mg., hesperidin 50 mg./Tab Bot. 100s.

Use: Vitamin supplement.

B-SOBBIC. (Werner-Chilcatt) Thiamine HCl 1 mg., ascorbic acid 30 mg./tab. Box 50s, 250s.

Use: Orally, vitamin B-1 w/vitamin C deficiencies.

BSP. See: Bromsulpholein.

BSP LIQUID. (Glidden) Ringer's sol. 85%, colamine 63%, isopropyl alcohol 4%, methyl cellulose 1.3%. Sol. Bot. 4 oz.

Use: Applied to buttocks & back for prevention of bedsores.

BST SOLUTION. (Confield) Bismuth sodium tartrate 1.5% or 3%. Vial 10 cc.

Use: Treatment at all stages of syphilis.

B-SYN-CO. (Pitman-Moore) Thiamine HCl 3 mg., riboflavin 0.33 mg., pyridoxine HCl 1 mg., niacinamide 10 mg./cc. or Tab Vial 25 cc. Bot. 500s.

Use: Orally, vitamin B deficiencies

BZ #3. (Sullivan & Case) Bismuth salicylate 520 mg., salol 104 mg., zinc phenolsulfonate 104 mg., pepsin 260 mg., aromatics, color q.s./fl oz. Bot. 1 pl., 1 gal.

Use: Diarrhea.

B-TELVE. (Paul Money) Vit. B-12 25 mcg., B-1 10 mg./tab. Bot 100s.

Use: Dietary supplement.

B-TROPIC. (Vale) Tricholine citrate (47% thia-choline base) 6 Gm., inositol 2 Gm., thiamine HCl 3 mg., riboflavin 2 mg., nicotinic

- acid 20 mg, in flavored sugar free vehicle
Bot 1 oz, 1 pt, 1 gal Each Cap contains,
Choline dihydrogen citrate 375 mg, inositol
125 mg thiamine HCl 1 mg, riboflavin 0.5
mg, nicotinic acid 5 mg Cap, Bot 100s
1000s
Use: Orally (1 dr or 2 cap q 1 d) impaired
I₂ & oxygen metabolism
- B-TWELV.** (Sherman) Vit B 12 cryst 30
mcg or 50 mcg 100 mcg/cc Vial 10 cc
Use 1 M (1 mcg daily) hematopoietic
- B-TWELVORA.** (Sherman) Vit B 12 25 mcg/
Cap Bot 100s
Use Orally (1 or more pr n) hematopoi-
etic
- BUBARTAL SODIUM.** (Columbus) Butobarbi-
tal sod 125 mg w/polyethylene glycol
benzyl alcohol water/cc Vial 10 cc
Use 1 M, sedative
- BUBARTAL TT** (Columbus) Butobarbital so-
dum 60 mg/Tab Bot 100s, 500s
Use Sedative or hypnotic
- BUCAT.** (Sulliff & Case) Buchu ext 65 mg,
sod barate 260 mg, trifolium ext 65 mg,
carna silk ext 325 mg, hydrangea ext
325 mg, atropine sulfate 0.13 mg/Tab
Bot 100s 1000s
Use: Diuretic
- BUCHINATE** (O F Harvey) Buchu p ext
¼ gr, squill ½ gr, asparagus seed ext
¼ gr apocynin ¼ gr, podophyllin 1/20
gr, potassium nitrate 1 gr, oil juniper ¼
min/Pill Bot 1000s
Use Diuretic and mild laxative
- BUCHUFORM** (Strassenburgh) Methenamine
2 gr resarcinol 1 gr, barasmin ¼/Tab
Bot 100s 500s 1000s
Use Orally (1 2 1 d) urinary antiseptic
BUCHU See Barasmin
- BUCHU, JUNIPER & POTASSIUM ACETATE.**
NF (Merck Sharp & Dohme) Buchu 15 Gm,
juniper 7.5 Gm pot acetate 5 Gm alcohol
37%/100 cc Bot 1 pt, 1 gal
Use: Treatment of cystitis
- BUCHU, JUNIPER AND POTASSIUM ACE-
TATE NO 1 ELIXIR** (Parke Davis) Buchu
45 gr, juniper 12 gr potassium acetate
16 gr, alcohol 30%/fl oz Bot 1 pt, 1 gal
- BUCLADIN.** See: Stuart Buccladin, Tabs
(Stuart)
- *BUCLAMASE.** (Ryton) Amylase alpha 10
mg w/1250 Byston units of amylolytic
activity/mg Tabs, Bot 48s
Use Inflammation
- *BUCLIZINE HYDROCHLORIDE.** 1-(p shlar
benzhydyl) 4 (p tert butylbenzyl) disthyl
enediamine dihydrochloride (longilene)
See Sofron (Stuart)
Yiboxine Hydrochloride (Pfizer)
- * Generic name recognized by N.F., U.S.P. or A.M.A.
- W/Oxytetracycline,** glucosamine, pheno-
cetin salicylamide, caffeine See Car-
Tetracyclin Caps (Pfizer)
- W/Salicylamide,** caffeine, phenacetin, TAO
See: TAO AC, Cap (Roerig)
- W/Scopolamine,** atropine sulf, hyoscyamine
sulf, Vit B 6 See Stuart Buccladin, Tab
(Stuart)
- W/Tetracycline HCl,** phenacetin, coffee-
ine salicylamide See: Tetracyclin, Tab (Pfizer)
- BUFAZO.** (Sulliff & Case) Sulfacetamide 125
mg, sulfamethoxazole 125 mg, phenazopyri-
dine HCl 65 mg, citric acid buffered
base 150 mg / Cap Bot 100s, 1000s
Use: Urinary antibacterial and analgesic
- BUFFADYNE W/BARBITURATES** (Lemmon)
Aspirin 300 mg, phenacetin 150 mg,
caffeine 30 mg, secobarbital sodium 8 mg,
amobarbital 8 mg, buffered w/aluminum
and magnesium hydroxides/Tab Bot 100s
Use: Analgesic
- BUFFAGESIC.** (Miss) Tablets 100s, 1000s
- BUFFERIN, A O R** (Bristol Myers Co) Aspirin
5 gr, aluminum glycinate, mag carb (Dial-
minate) Bot 12s, 100s
Use: Mild analgesic
- BUFFONAMIDE, N N R 1955** (Tulag) Sulf
acetamide, sulfadiazine, sulfamerazine each
33 mg and Sod citrate 100 mg/cc Bot
2 oz, 1 pt
Use Triple sulfonamide therapy
- BUFFABS.** (Burlington) Col carbonate 3 gr,
mag carbonate 2 gr, bis subcarbonate 1
gr, glycine ¼ gr, aluminum hydroxide gel
½ gr/Tab Bot 100s, 1000s
Use: Antacid
- BUFOPTO CARBACEL.** (Professional Pharm)
Carbachol 0.75% or 1.5% w/methylcellu-
lose and benzalkonium chloride Bot 15 cc
Use: Glaucoma
- BUFOPTO-EFRICEL.** (Professional Pharm)
Phenylephrine HCl 10%, methylcellulose
½%, benzalkonium chloride 1:25 000 solu-
tion Plastic dropper Bot 5 cc
Use: Vasoconstriction and pupil dilation
- BUFOPTO METHULOSE.** (Professional Pharm)
Methylcellulose ¼% with zephiran chloride
as preservative Bot 15 cc, 30 cc
Use: Vehicle for ophthalmic medicaments
and contact lens fluid
- BUFOPTO MYTRATE.** (Prof Pharm) Epineph-
rine bitartrate 2%, benzalkonium 1:25 000,
sterile stable Methylulose soln Bot 15 cc
Use: Glaucoma
- BUFOPTO WITH PILOCEL.** (Professional
Pharm) Methylcellulose ophthalmic buffered
sol w/pilocarpine HCl as ¼%, ½%, 1%
2%, 3% & 4% Vial 10 cc
Use: Myosis

- BUFOPTO SULFACEL-15.** (Professional Pharm.) Sulfacetamide Sod. 15%, methylcellulose 1/2% In buffered solution. Plastic Bot. 15 cc.
Use: Ophthalmic bacterial infections.
- BUFOPTO SULFAPRED.** (Professional Pharmacal) Prednisolone acetate 0.25%, sodium sulfacetamide 10%, phenylephrine HCl 0.12%, propylene glycol, sodium phosphate monobasic, sodium phosphate dibasic, sodium thiosulfate in a methylcellulose base, preserved with methyl paraben 0.04, propyl paraben 0.02%. Clear plastic containers 5 cc.
Use: Allergic blepharitis.
- BUFOPTO-VERNACEL.** (Prof. Pharmacal) Prephenpyridamine maleate 0.5%, phenylephrine HCl, 1/2%. Plastic bot, 15 cc.
Use: Ophthalmic soln.
- BUFOPTO ZINC SULFATE.** (Professional Pharm.) Plastic Bot. 15 cc, Glass 30 cc.
Use: Conjunctivitis.
- BUFOSAL.** (Tablerack) Sod. salicylate 15 gr./dr. w/Col. carb. & Sod. bicarb. as granulated effler. Bot. 4 oz.
Use: Orally (1 tsp. i.i.d.), analgesic & antipyretic.
- BULATIN.** (Ulmer) Carboxymethylcellulose Sod. 6 gr., diacetylhydroxyphenyltatin 1 mg., dehydrated prune concentrate 2 gr./Tab. Bot. 100s, 500s.
Use: (2-3 with glass of water b.i.d.) Bulk laxative.
- BU-LAX.** (Ulmer) Sod carboxymethylcellulose. Tab. 0.50 Gm. Bot. 100s, 500s, 1000s.
Use: Orally (1-2 b.i.d.), laxative.
- BULKALOID.** (U.S. Standard) Methylcellulose 7 1/2 gr./Tab. Bot. 100s, 1000s.
Use: Orally, persistent constipation.
- BULKOGEN.** A mucin extracted from the seeds of *Cyanopsis tetragonaloba*
W/Hyascyamine HBr, scopolamine HBr, aluminum hydroxide, magnesium hydroxide. See: Hycodrox, Tab. (Columbus)
- BULKOSOL.** (Columbus) Bulkogen 500 mg. diacetyl sod. sulfosuccinate 40 mg./Cap. Bot 100s, 1000s.
Use: Constipation.
- BU-MA-LAC.** (Pitman-Moore) Milk protein, defatted w/Mag. butyrate 1%, procaine HCl 1%. Vial 20 cc.
Use: I.M., nonspecific protein therapy.
- BUMINTEST.** (Ames) Sulfasalicylic acid in effler. tab. Tab., Bot. 32s, 100s, 500s.
Use: Test for albumin in urine.
- * **BUNAMIODYL SODIUM, N.N.D. 1962**
See Orabitez, Cap. (Faugero)
- BUNESIA.** (McNeil) Hamotropine methylbromide 2.5 mg. (1/24 gr), butisal Sod. 10 mg. (1/6 gr.), Mag hydroxide 0.3 Gm. (5 gr.) w/aromatics/Tab. Bot. 100s, 500s, 1000s.
Use: (1 or 2 i.i.d., a.c. & h.s.), antispasmodic, sedative & antacid
- BUPENNA, Tab.** (Burrough Bros.) Penicillin G. Tab. 50 (50,000 u.); 100 (100,000 u.); 200 (200,000 u.); 250 (250,000 u.). Bot. 100s, 1000s. Available w/Triple Sulfo Bot. 24s, 100s.
- * **BUPNENINE.** 1-p-Hydroxyphenyl-2-(1-methyl-3-phenylpropyl-amino)propanol. Peridialol hydrochloride.
- BUBBO-CAINE.** (Rawell) Burbot liver oil 50%, benzocaine 5% & petrolatum. Oint., Tube 1 oz.
Use: Topically, vitamin therapy.
- BUBBO LIVER OIL, N.N.R. 1955.** (Rawell) liq. Vit. A 4480 u., D 640 u./Gm. Bot. w/dropper 60 cc, Bot. 8 fl. oz. Cap. Vit. A 2215 u., D 315 u. Bot. 50s, 100s, 250s. Oint. 50%, Tube 1 oz., 4 oz; Jar, 1 lb.
Use: Orally, dietary supplement, topically as orally, stimulant in healing.
- W/Benzocaine, boric acid, zinc oxide, phenol & camphor. See: Burnolatum, Oint. (Rawell).
- W/Benzocaine & petrolatum. See: Burbo-Caine, Oint. (Rawell).
- BUBN DRESSINGS,**
See: Foille, Preps. (Carbisulphol)
Zinox, Dressings (Hynson, Wercott & Ouning)
Zincosote Burn Dressing, N.N.D. 1958.
- BURNHAM SOLUBLE IODINE.** (Burnham Soluble) Iodine 1.1 mg./drop. Bot. 1/2, 1, 2, 4 oz.
Use: 10 to 30 gtt. t.i.d. in water 15 min. o.c.
- BURNOLATUM.** (Rawell) Benzocaine 2%, burbot liver oil, boric acid, zinc oxide, phenol & camphor. Oint. Tube 1 oz.
Use: Topically, relief of mild skin irritations caused by sunburn & insect bites.
- BURN-QUEL.** (Halperin) 1 oz. dispenser.
Use: Burn therapy.
- BURNTANIK.** (Hartan & Converse) When diluted with 4 parts water the sol. contains: Tannic acid 5%, p tertiary-amyl phenol 1:5000, benzyl alcohol 4%, ethyl alcohol 5% w/Sod, Cal. & Pot. chlorides to make isotonic. Pkg 4 oz., 16 oz.
Use: Burn therapy.
- BURO-SOL.** (Doak) Aluminum acetate. Cream. 4 oz., 1 lb. Paw. 4, 8, 16 oz.
Use: Topically, astringent & antiseptic.
- BURO-SOL ANTISEPTIC POWDER.** (Doak) Contents make a dilute Burow's Sol.—

- aluminum acetate sol—plus benzethonium chloride Pkg 12s
Bot Pow 4 oz, 1 & 5 lb
- BURO-SOL CONCENTRATE.** (Daak) Pow to make Buraw's Sol Pkg 8 oz 4s; 32 oz 16s
Use Dermatological
- BUROW'S SOLUTION, U.S.P.**
See Aluminum Acetate Solution
Buro Sol Concentrate (Daak)
- W/Cool tar sol & starch See Bur Tar Gen Oint (Texas)
- W/Starch See Bur Tar Gen (Texas Ford Co)
- W/Aveeno See Bur Vein (Faugera)
- ***BURBAMIODYL SODIUM, NND 1962**
Sodium 3 (3 butyrylamino 2,4,6 trisido phenyl) 2 ethylacrylate See: Orabiles (Faugera)
- BURBOLINE, NND 1959** (Burnham Soluble) Di glycolic hydriodide 2 mofes, combined with 2 atomic weights of iodine Tab 30.5 to 32% of active iodine Bot 50s
Use Soluble iodine for disinfection of drinking water (1 tab to qt)
- BUR-TAR-GEN** (Texas Pharm) Buraw's sol 5% coal tar sol starch in nongreasy base Oint 2 oz 1 lb
Use Topically antiseptic
- BUR-VEEN** (Faugera) Aveeno colloidal oat meal aluminum sulfate Cal acetate Pow Pkg, Box 6s 100s
Use: 1 Pkg dissolved in 3 pint water ss equiv to 120 Buraw's Solution with the added protection of Aveeno
- ***BUSULFAN, U.S.P., HND 1962** Tetra methylene dimethanesulfonate Tabs U.S.P., See Myleran Tab (Burroughs Wellcome)
- BUSULPHAN.** 1:4 Dimethanesulphonyloxy butane See Myleran
- ***BUTABARBITAL, (Various Mfr)** 5 sec butyl-5 ethylbarbituric acid
See Butobarb Unicelles (Hiss Pharm)
Butisal, Preps (McNeil)
Nocetyl (Allison)
Nilax Caps (Direct Labs)
- W/Acetaminophen phenacetin caffeine See: Butgetic Tabs (McNeil)
- W/Ambutonium bromide aluminum hydroxide mag hydroxide See Aludrox SA Susp, Tab (Wyeth)
- W/Aminaphylline phenylpropanolamine HCl chlorpheniramine maleate, aluminum hydroxide magnesium trisilicate See Asma cal Tabs (Vale)
- W/d Amphetamine sulf See Butacex Tab (Paul Maney)
- W/d Amphetamine sulf, acetphenolisatin See Nilax, Cap (Direct Labs)
- W/d Amphetamine sulf, acetylsalicylic acid
- * Generic name recognized by N.F., U.S.P. or A.M.A.
- salsylamide, acetaminophen See: Stuart Prolar, Tab (Stuart)
- W/d Amphetamine sulf, acetylsalicylic acid salicylomide, N Acetyl p Aminophenol, hyoscyamine sulf, atropine sulf, hyoscyine HBr See Stuart Prolar B, Tab (Stuart)
- W/Carboxyphen See: Bonril Timed No 2 Tab (G W Carnrick)
- W/Chlorpheniramine maleate, hyoscyine hydrobromide See: Pedro Sol (Warren Pharmacal)
- W/d Desoxyephedrine HCl, aspirin & acetphenetidin See: Asphedex, Tab (Central)
- W/Hydrochlorothiazide See: Butizide Pres Tab (McNeil)
- W/Hydrochlorothiazide, reserpine See: Sulterpazide Prestab (McNeil)
- W/1 Hyoscyamine See: Levamine, Tabs/Caps (Meyer)
- W/Mephenesin See: Halabar, Tab (Carnrick) Tolbusem, Tab (Zemmer)
- W/Mephenesin, d amphetamine sulfate See: Crystodex, Tab (Crystal)
- W/Methscopolamine nitrate, atropine methyl nitrate See: Scolote, Tabs (Central)
- W/Pentaerythritol tetranitrate
See Pentril B, Tempules (Evron)
Corolet T.D., Caps (Crystal)
- W/Pentaerythritol tetranitrate, nitroglycerine See Buto Penite, Tabs, Caps (Carnrick)
- W/Pentobarbital sodium. See: Barbitwin Tabs (Paul Maney)
- W/Pentobarbital Sod See: Twin Barb Improved, Tab (Ascher)
- W/Pentobarbital & phenobarbital See: Quess, Tab (Durst)
- W/Phenobarbital, secobarbital See: Duradorm, Tab (Lincoln)
- W/Secobarbital See Butasec Unicelles (Hiss Pharm)
- W/Theophylline, Ephedrine HCl See: Numa Dura Tabs (Wynn)
- W/Theophylline, Ethaverine HCl, Isopropyl arterenal See Benedrin, Tab (Bene)
- W/Triethanolamine trinitrate biphosphate See Metamine w/Butabarbital, Tab (Leeming)
- ***BUTABARBITAL SODIUM, NND 1962**
Sod 5 sec butyl 5 ethylbarbiturate Caps, Elixir Tabs, N.F.
See: Subartal TT Tabs (Columbus)
Butabarpal, Vial (Philadelphia Ampoule)
Butak, Tab (Hoack)
Butaxem, Tab and Elix (Zemmer)
Butisal Sodium (McNeil, N.N.R.)
Insalat, Tab (Denab)
Mebutal Tab (Medco)

Approved Pharm.—Elixir D.2 Gm./30 cc. Bot. 1 pt., 1 gal.

Bowman—Tab. (¼ gr., ½ gr.) Bot. 100s, 1000s, 5000s. Elix. (3 gr./oz.) Bot. 1 pt., 1 gal.

Use: Sedative & hypnotic.

W/Acetyl-p-aminophenol.

See: Butapap, Syrup (Physicians Products Co.)

Nacetyl, Tab. (Allison)

W/Acetyl-p-aminophenol & methamphetamine HCl. See: Algoston, Tab. (McNeil)

W/Aminopentamide hydragen sulfate. See: Spasmolyd, Tab. (Columbus)

W/Aminaphylline, phenylephrine HCl, methamphetamine HCl, glyceryl guaiacolsulfonate, & aluminum hydroxide-magnesium carbonate. See: Bes-O-Phyllin, Cap. (Santard)

W/d-Amphetamine sulf., d-amphetamine sulf. See: Robudex-G.R., Cap. (Lynn)

W/d-Amphetamine phosphate, homatropine methylbromide, bulkogen, Vit. A, Vit. D, Vit. B-1, B-2, niacinamide. See: Qbetolds, Tab. (Columbus)

W/d-Amphetamine sulf., mephenesin.

See: Butadex, Tab. (Paul Maney)

Muslox, Tab. (Cole)

W/d-Amphetamine sulfate & methylcellulose. See: Bontril, Tab. (Cornick)

W/Ascorbic acid, Sod. p-aminobenzoate, salicylamide, Sod. salicylate. See: Bisolate, Tab. (Allison Lab.)

W/Belladonna ext.

See: Insebel TD, Tab. (Denab)

W/Dehydrocholic acid, belladonna extract. See: Decholm-BB, Tabs. (Ames)

W/d-Desoxyephedrine HCl, aspirin, phenacetin. See: Asphedex, Tab. (Central Pharm.)

W/Homatropine methylbromide. See: Butathyl, Tab. and Elix. (Zemmer)

W/Homatropine methylbromide, profenil citrate. See: Fentropine W/Butabarbitol, Tab. (C. D. Smith)

W/Hyascine hydrobromide, n-acetyl-p-aminophenol.

See: Pallacin, Tab. (Denab)

W/Hyascine HBr, hyoscyamine sulfate, atropine sulfate. See: Zemarine Elixir. (Zemmer)

W/Mephenesin & d-amphetamine sulfate. See: Butadex, Tab. (Paul Maney)

W/Mephenesin & salicylamide. See: Banesin, Tab. (Kendall)

W/Methscopolamine bromide, aluminum hydroxide gel dried & Mag. trisilicate. See: Eutcin, Tab. (Staff Research)

W/Methylphysaloxane, belladonna extract. See: Sibel, Tabs. (Plough)

* Generic name recognized by N.F., U.S.P. or A.M.A.

W/Pentaerythritol tetranitrate. See: Perbutzem, Tab. (Zemmer)

W/Pentaerythritol tetranitrate, reserpine. See: Pentrasine, Tab. (McNeil)

W/Pentobarbital sodium. See: SED, Caps. (Layden Labs)

W/Pentobarbital Sod. & d-desoxyephedrine. See: Somnadex, Tab. (Central)

W/Pentobarbital sod., phenobarbital. See: Pyramid, Liq. (Barrel)

W/Phenobarbital, pentobarbital sod., secobarbital sod., belladonna paw. ext., phenyltoloxamine, ethaverine HCl. See: Tri-Synar Plus, Tab. (Armour Labs.)

W/Phenobarbital Sod., pentobarbital Sod. & secobarbital Sod. See: Quadra-Sed, Sol. (C. D. Smith)

W/Paldine methylsulfate. See: Nactival, Tabs. (McNeil)

W/Reserpine. See: Reserbutal, Tab. (Vitamix)

W/Scopolamine hydrobromide. See: Ampyrox, Tab. # 281 & Elix. (Elder)

W/Secobarbital Sod., pentobarbital Sod. & phenobarbital. See: Nidar, Tab. (Armour)

W/Secobarbital Sod. & phenobarbital. See: Butsero, S.C.T., Tab. (Bowman)

Ethabrol, Cap. (Wyeth)

BUTABAB UNICELES. (Miss) Butobarbital 45 mg./long acting tab. Bot. 30s.

Use: Long acting sedative.

BUTABABPAL. (Philadelphia Ampoule) Butobarbital ead. 125 mg. in polyethylene glycol/cc. Vial 10 cc.

Use: I.M., sedative.

BUTACAINE SULFATE, N.F. X, N.H.R. 1950, A.D.X. (Varrone Mfr.) Butelline, 3,4-Dibutylaminopropyl-p-aminobenzoate sulfate.

See: Butyn Sulfate (Abbott)

Butyl Metaphen, Dent. Oint. (Abbott)

Use: Surface anesthetic.

BUTACEX. (Paul Maney) d-Amphetamine sulfate 5 mg., butobarbital ½ gr./tab. Bot. 100s, 1000s.

Use: Mental & emotional distress, treatment of obesity.

BUTA-DEX SULES. (Normal Pharmacy Co) Capsules T.D. 10 mg., 15 mg., 100s, 500s, 1000s.

* **BUTADIAZAMID.** N-(5-butyl-1,3,4-thiadiazol-2-yl)-p-chlorobenzenesulfonamide (Lederle Lab)

BUTADONNA. (Wampole) Chlorbutanol 250 mg., hyoscyamine sulfate 0.1 mg., atropine sulfate 0.02 mg., hyoscyne HBr 0.007 mg./Cap. Bot. 100s.

Use: (1 t.i.d. o.c.) Spasmolytic, sedative

BUTA-DDN SULES. (Normal Pharmacy Co) Capsules T.D. 100s, 500s, 1000s

BUTADOR CAPS. (Physicians Products) Buta-

- barbital sodium 15 mg; acetaminophen 250 mg; salicylamide 200 mg; atropine sulfate 0012 mg; scopolamine hydrobromide 0048 mg; and hyoscyamine sulfate 024 mg/Caps Bot 100s 1000s
Use Analgesic
- BUTADOR 2, 3, and 4** (Physicians Products)
Butabarbital sodium 15 mg, acetaminophen 250 mg, salicylamide 200 mg, atropine sulfate 0012 mg, scopolamine hydrobromide 0048 mg and hyoscyamine sulfate 024 mg #2 with 15 mg codeine phosphate for moderate to severe pain #3 30 mg and #4 60 mg, codeine phosphate Bot 100s, 1000s 5000s
Use Analgesic
- BUTAGESIC** (Lowe) Butabarbital sodium 1/4 gr, N-acetyl p-aminophenol 3 1/2 gr, salicylamide 2 1/2 gr, aluminum hydroxide gel 1 gr, magnesium carbonate 1 gr/Tab Bot 100s 1000s
Use Sedative Analgesic
- BUTAK** (Hoack) Butabarbital sod 15, 30, & 100 mg/Tab Bot 100s & 1000s
Use Sedative
- BUTALBITAL**, Talbutal 5 allyl 5 sec butyl barbituric acid
See Latusate (Winthrop Labs)
Demeral W/Latusate (Winthrop Labs)
- BUTALGIN** Bee di Methodane Hydrochloride Prep
- ***BUTALLYLOHAL**, NF X NNB 1951; Tab NF X 5-(2 Bromoallyl) 5 sec butylbarbituric acid (Pernactan)
Use Hypnotic
- BUTAMBEN PICRATE**, NNR 1951 See: Butesin Picrate Prep
- BUTAMIN**, See Tufacaine Hydrochloride (Winthrop Labs)
- ***BUTAMOXANE HCl**, (Eli Lilly) 2 (n butyl-aminomethyl) 1,4 benzodioxane HCl
Use Tranquillizing agent
- ***BUTANISAMIDE**, 1 butyl 1 (a methoxy phenyl)urea
- BUTAPAP**, (Tailby Nason) Butabarbital sod 15 mg acetyl p-aminophenol 120 mg/5 cc Bot 1 pl
Use Relief of pain, restlessness
- BUTA-PENTILE**, (Carrick) Tab pentaerythritol tetranitrate 10 mg nitroglycerine 03 mg, butabarbital 16 mg; Cap pentaerythritol tetranitrate 30 mg, nitroglycerine 09 mg, butabarbital 48 mg Tabs Bot 100s; Caps Bot 50s
Use: Angina pectoris
- ***BUTAPEBAZINE** (Riker) 1 {10 [3-(4 methyl 1 piperazinyl) phenothiazin 2 yl]} 1 butanone
Use Psychotropic agent
- * Generic name recognized by NF, USP or AMA
- ***BUTAPHYLLAMINE**, (Merrill) (Not Listed)
Theophylline aminoisobutanol
W/Dicapryn, nethamine See: Nethoprin Cap & Syr (Merrell)
W/Phenobarbital & nethamine HCl See: Nethaphyl, Cap (Merrell)
- BUTASEC UNICELLES**, (Hiss Pharm) Secobarbital 1 1/2 gr, butabarbital 1/2 gr/cap Bot 100s, 500s
Use: (1 hr) Delayed action, sedative
- BUTATHYL**, (Zemmer) Butabarbital sodium 10 mg, hamatropine methylbromide 2.5 mg/Tab, dr Tab Bot 100s, Ellixir, Pl
Use: Antispasmodic and sedative
- BUTAZEM**, (Zemmer) Butabarbital sodium 15 mg, 30 mg, 01 Gm/Tab (#1, #2, #3) 02 Gm/ox Tab #1, #2, #3, Bot 100s Ellixir, Bot Pl
Use: Sedative
- BUTAZOLIDIN**, NND 1962 (Geigy) (Phenylbutolazone) 3,5-dioxo-1,2 diphenyl-4-butylpyrazolidine Tab 100 mg Bot 100s 250s, 1000s
Use: (300 to 600 mg daily) Analgesic & anti-inflammatory action
- BUTAZOLIDIN ALKA**, (Geigy) Butazolidin (phenylbutolazone) 100 mg, aluminum hydroxide 100 mg, mag trisilicate 150 mg, hamatropine methylbromide 1.25 mg/Cap Bot 100s
Use Anti-inflammatory, analgesic and antipyretic
- BUTELLINE**, See Butocaine Sulfate (Varlot Mfr)
- BUTESIN**, NNR 1950 (Abbott) n Butyl p-aminobenzoate Pow, Bot 5 Gm, 1 oz
Use Topically, local anesthetic
W/Procaine See: Zylcaine, Amp (Abbott)
- BUTESIN COMPOUND**, (Abbott) Butesin 2 1/2 gr, hydrostiz ext 1/2 gr, Hamamelis ext 1 gr, in cocoa base/Supp Box 12s
Use Rectally, local anesthetic suppository
- BUTESIN PICRATE**, Butamben picrate, NNR 1951
- BUTESIN PICRATE 1% W/Butesin 1% Ophth Oint**
Abbott—Contains 1% butein picrate (Butamben Picrate) micronized & butesin (Butyl Aminobenzoate) 1% in soft petrolatum base Tube 1/2 oz
Use Topically, local anesthetic & antipruritic
- BUTESIN PICRATE OINTMENT W/Metaphen** NNB 1951 (Abbott) Contains 1% butein picrate (Butamben Picrate, Abbott) & metaphen (Nikromersol, Abbott) 1 5000 Tube 1 oz 2 oz Jar 1 lb
Use Minor burnz of various kinds—electric steam, hot metal & scalds

- BUTETHAL**, N.F. X, N.N.B. 1951, A.D.R.; Tab. N.F. X (Various Mfr.) 5-Ethyl-5-butylbarbituric acid. See: Neonal, Tab. (Abbott)
Use: Sedative & hypnotic.
- W/Mephensin, d-amphetamine sulf.** See: Butodex, Tab. (Paul Maney)
- BUTETHAMINE FORMATE**, N.N.D. 1962. 2 Isobutyl aminoethyl-p-aminobenzoate formate. See: Monocaine Formate, Amp. (Novoca)
- BUTETHAMINE HYDROCHLORIDE**, N.F., N.N.D. 1962, A.D.B.; [n], N.F. See: Monocaine Hydrochloride (Novoca)
- BUTETHANOL**. See: Pantocaine Hydrochloride (Winthrop Labs.)
- BU-THICIN**, Troches. (Burrough Bros.) Tyrothricin 2 mg., benzocaine 5 mg./Troche. Bot. 16s, 500s.
Use: Mouth & throat infections.
- BUTIBEL Tab. and Elixir**, (McNeil Labs) Butisal Sod. 15 mg., belladonna ext. 15 mg./tab. or dr. Tab. Bot. 100s, 1000s; Elixir Bot. 1 pt., 1 gal.
Use: Antispasmodic with sedative.
- BUTIBEL R-A**, (McNeil) Butisal Sod. 20 mg., ext. belladonna 30 mg./Tab. Bot. 50s, 500s.
Use: Antispasmodic, sedative with prolonged action.
- BUTICAPS**, (Stanley Drugs) Vit. F from linoleic & linolenic acids, amino acids & lecithin. Cap., Bot. 100s.
Use: Orally, nutritional skin treatment.
- BUTIGETIC**, (McNeil) Butobarbital Sod. 15 mg., acetaminophen 200 mg., phenacetin 150 mg., caffeine 30 mg./Tab. Bot. 100s
Use: Analgesic W/sedative, for treatment of tension headache.
- BUTISEPAZIDE PRESTAB**, (McNeil) Butobarbital 30 mg., hydrochlorothiazide 25 mg., reserpine 0.1 mg./Green 25 Prestab. Butobarbital 30 mg., hydrochlorothiazide 50 mg., reserpine 0.1 mg./Orange 50 Tab. Bot. 50s.
Use: Hypertension.
- BUTISERPINE**, (McNeil) Butisal Sod. 15 mg., reserpine 0.1 mg./tab. or 5 cc. Bot. 100s, 1000s. Elix., Bot. 12 fl. oz.
Use: Hypertension.
- BUTISERPINE B-A**, (McNeil) Butisal sod. 30 mg., reserpine 0.2 mg./Tab. Bot. 50s.
Use: Prolonged action antihypertensive and sedative.
- BUTISOL**, Butobarbital.
W/Aspirin & Cal. gluconate. See: Aspraal, Tab. (McNeil).
W/Chloratam, belladonna ext., Ipecac syr. & Pot. chloride. See: Lectusal, Cough Prep. (McNeil)
- BUTISOL-HYOSCYAMUS**, (McNeil) Butisal
* Generic name recognized by N.F., U.S.P. or A.M.A.
- $\frac{1}{2}$ gr., hyoscyamus ext. $\frac{1}{2}$ gr./Tab. Bot. 100s, 1000s, 5000s.
Use: Orally, sedative & hypnotic.
- BUTISOL R. A.** (McNeil) Butisal Sod. 30 mg. or 60 mg./delayed action tab. Bot. 50s, 500s.
Use: Prolonged sedative.
- BUTISOL SODIUM**, N.N.D. 1961. (McNeil) (Butobarbital Sod.) Sod. 5-sec-butyl-5-ethylbarbiturate. Cap. (1 $\frac{1}{2}$ gr.) Bot. 100s, 500s, 1000s. Elixir (50 mg./5 cc) Bot. 1 pt., 1 gal. Tab. (1 $\frac{1}{2}$ gr., $\frac{1}{2}$ gr., $\frac{1}{4}$ gr., 1 $\frac{1}{2}$ gr.) Bot. 100s, 500s, 1000s.
Use: Intermediate sedative & hypnotic
W/Acetyl-p-aminophenol & methamphetamine HCl. See: Algoson, Tab. (McNeil).
W/Aminophylline & desoxyephedrine HCl racemic. See: Drenal, Tab. (McNeil)
W/Belladonna ext. See: Butibel, Tab. and Elixir, (McNeil Labs.).
W/Beserpine. See: Butiserpine, Tab. & Elix. (McNeil)
- BUTIZIDE PRESTAB**, (McNeil) Butobarbital 30 mg., hydrochlorothiazide 25 mg./Blue 25 Prestab. Butobarbital 30 mg., hydrochlorothiazide 50 mg./Gold 50 Tab. Bot. 50s.
Use: Hypertension.
- BUTOBEN**, (Merck Sharp & Dohme) Butyl p-hydrobenzoate. Bot. 1 oz., $\frac{1}{2}$ lb., 1 lb.
Use: Preservative against yeast.
- BUTODEX**, (Paul Maney) Mephensin 250 mg., d-amphetamine sulfate 2.5 mg., butobarbital Sod. 8 mg./Tab. Bot. 100s.
Use: (1-2 t.i.d.) Muscle relaxant.
- BUTOLAN**, Benzylphenyl carbamate. See: Diphenan. (Burroughs Wellcome).
- BUTAPTRONOXYL**, U.S.P. XV (Indoione) Butyl mesityl oxide.
Use: Insect repellent, used in Camp. Dimethyl Phthalate Solution, U.S.P.
- BUTRITANNIC**, (Rare) Procaïne butyrate 5%, ephedrine 0.5%, tannic acid 1 $\frac{1}{4}$ %, w/phenol & ammonium alum. Oint. Tube 1 oz., Jar 4 oz.
Use: Topically, local anesthetic w/antispasitics.
- BUTSECO**, S.C.T. (Bawman) Butobarbital Sod. $\frac{1}{4}$ gr., zecobarbital Sod. $\frac{1}{4}$ gr., phenobarbital $\frac{1}{4}$ gr./tab. Bot. 100s, 1000s.
Use: Sedative.
- BUT-TAB-GEN**, (Texas Pharm. Co) Buraw's soln., March 10 nongreasy base. Bot. 2 & 16 oz.
Use: Dermatitis.
- BUTYLATED HYDROXYANISOLE**, (Registered in Great Britain) A mixture of 2-tert. butyl-4-methoxyphenol and 3-tert. butyl-4-methoxyphenol.

BUTYLATED HYDROXYTOLUENE. (Registered in Great Britain) 2,6-Di-tert-butyl-p-cresol

BUTYL CHLORIDE, N.F.

Use: Anthelmintic (veterinary)

BUTYL AMINO BENZOATE, N.F., N.N.R. 1950 Scuratarme See: Butesin (Abbott)

Use: Surface anesthetic.
W/Benzocaine See: Benzocaine B.B., Oint (Uimer).

W/Benzyl alcohol, phenylmercuric borate & benzocaine See: Dermathyn, Oint (Fliet Ector).

W/Procaine base benzocaine & benzyl alcohol in sesame oil See: Analgesic, Vial (Metro).

W/Procaine, benzyl alcohol, in sweet almond oil See: Anacaine, Amp (Calvin Chem)

W/Sulfadiazine & tyrothricin See: Suthagen Troches. (Thompson).

BUTYL p-HYDROXYBENZOATE. See: Butabac (Merck Sharp & Dohme)

BUTYLPHENAMIDE. N-n butyl 2 phenyl salicylamide See: By-NA MID (Cutler Lab)

BUTYLAMINE HCl. (Eli Lilly) N-tert butyl N,1,1 trimethyl 2 propynylamine HCl

Use: Hypotensive

BUTYN METAPHIN DENTAL OINTMENT. (Abbott) Butyn 4%, metaphen 1:1500 concentration Ointment applicator syringe 4.5 Gm appl syr /66.

Use: Dental anesthetic and antiseptic.

BUTYN SULFATE, N.N.R. 1950, A.D.R. (Abbott) Butacaine sulfate Fow., Bot 5 Gm., 1 oz.

Use: Local anesthetic, effective topically

W/Metaphen (Abbott) Butyn sulfate 2%, metaphen (Nitromersol) 1,000, aphth. oint base q.s. Tube 1/2 oz., Box 1 doz

Use: Antiseptic ointment rapidly producing local anesthesia

BUTYN SULFATE, 2% SOLUTION (Abbott) Bot w/dropper, 1 fl. oz.

Use: Topical application for surface anesthesia.

B-VITIN. (Burrington) Vit B1 2 mg., B-2 3 mg., B-6 1 mg., Cal. pantothenate 1 mg., nicotinamide 20 mg./Pillm Bot 100s, 1000s

Use: Orally, vitamin therapy

B-VITIN-C. (Burrington) Thiamin 5 mg., riboflavin 5 mg., pyridoxine 5 mg., Cal. pantothenate 5 mg., nicotinamide 50 mg., ascorbic acid 150 mg., B-12 5 mg./Tab. Bot 100s

Use: Restricted diet

BYU See: Bromocriptin, U.S.P.

BYLINIC (CAPSULE). (Marrow) E-2e salts 60 mg., thiothamyl acid 0.66 Gm./Cap Bot 100s

Use: Cholegogic.

* Generic name recognized by N.F., U.S.P. or A.M.A.

BY-NA-MID LOY., OINT., PWD, TINCT (Cutler) Butylphenylsalicylamide (butylphenamide) 5%, all forms Lot 2 oz bot, Oint 1/2 oz tube, Pwd 1 oz, Tinct 1 oz Bot
Use: Athlete's foot and fungal infections
BZ55. See: Carbutamide

C, COMPOUND (Schering) Ethisterone 10 mg, chlorphenpyridamine maleate 2 mg, 8 bromotheophylline 25 mg, salicylamide 300 mg, caffeine 30 mg/Tab
Use: Treatment of premenstrual tension

CABBAGE JUICE. See: U Vita (Pitcher Pharmacal)

CACODYLATE W/Cinchophee, methenamine See: Cin Ca Mine "E," Vial (Kirk)

CACODYLIC ACID SALTS. See: Calcium Salt, Copper Salt, Ferric Salt, Iron Salt, Sodium Salt

CACODYNE IODINE (Res Medications Inc) Colloid iodine 0.2% in gelatin & Sod cacodylate 7 gr Isotonic Amp 10 cc 10s 100s

Use: Arterial diseases

CADE OIL. See: Juniper Tar, U.S.P. Tar Bath (Ar Ex)

CADMIUM SULFIDE, N.N.D. 1962 See: Capsebon (Pillman Moore)

Cyclospasmol (Ives Cameron)

CADROSON. (Merck Sharp & Dohme) Cal 3.5 Gm., phos DJS Gm., fluorine D.5 mg-iron 15 mg., Vit D 1000 u./6 Tabs Bot 100s

Use: (1 2 11 d) Dietary supplement

CAFARYL (Elder) Salicylamide 4 gr, pyrilamine maleate 25 mg, caffeine 1/2 gr./Cap-Bot 100s, 500s, 1000s

Use: Analgesic (1 2 q 3-4 h) Antihistamine & treatment of premenstrual tension

CAFED NOSE DROPS (Elder) Pot pepticin G 100,000 u., methamphetamine HCl 1/4 %/bot, Bot w/dropper 1/4 oz

Use: Nasal congestion & sinusitis

CAFERAN. (Vitarles) Ergotamine tartrate 1 mg, caffeine 100 mg./Tab Bot. 100s, 500s 1000s

Use: Migrates and tension headaches

CAFERDOT. N.N.R. 1955 (Sandoz) Ergotamine with caffeine, ergotamine tartrate 1 mg, caffeine 100 mg/Tab Bot 100s 500s, 1000s; Supp Ergotamine tartrate 2 mg, caffeine 100 mg Box 12s

Use: Orally & rectally for treatment of migraine & tension headaches.

CAFERDOT-PR. (Sandoz) Ergotamine tartrate

2 mg., caffeine 100 mg., pentobarbital sod. 60 mg., belfolofine 0.25 mg./supp. Pkg. 12s. Tob.: Ergolamine tartrate 1 mg., caffeine 100 mg., pentobarbital sod. 30 mg., bellafoline 0.125 mg. Bot. 100s.

Use: Treatment of migraine.

CAPTRATE. (Vitomix) Ergolamine tartrate 1 mg., caffeine 100 mg./Tob. Bot. 100s, 1000s

Use: Migraine

* **CAFFEINE**, U.S.P., A.D.R. (Various Mfr.) Guaranine; Methyltheobromine; Thein.

Use: Cardiac, respiratory & psychic stimulant.

CAFFEINE CITRATED, N.P. Tob., N.P., A.D.R. Bot. 1 oz., Pkg. ¼ lb., 1 lb. Tob. N.P. (Various Mfr.) See: A.P.C.

Use: Central nervous system stimulant.

CAFFEINE SODIO-BENZOATE, See: Caffeine and sod. benzoate, U.S.P.

* **CAFFEINE SODIUM BENZOATE**, U.S.P., N.N.R. 1947, A.D.R., In., U.S.P.; Tob. N.P. Approximately equal parts of caffeine and sod. benzoate. (Various Mfr.) Amp. (3¼ & 7½ gr.) 2 cc., Box 12s, 100s. Nypo Tob. (1 gr.) Tube 20s, 100s & Pow.

Use: Orally & i.M. (0.25-0.50 Gm.) Central nervous system stimulant.

CAFFEINE SODIUM SALICYLATE, N.F. (Various Mfr.) Bot. 1 oz., Pkg. ¼ lb., 1 lb.

Use: See Caffeine.

CAFFEINE-THEOPHYLLINE COMPOUND, w/Nux Vomica Ext. See: Xanthinus, Tob. (Coie).

CAPOTAN, (Premo) Acetophenetidol 3 gr., aspirin, caffeine and quinine sulfate. Tab., Tin 12s; Bot. 100s.

Use: (1-2 i.i.d.) Analgesic & antipyretic.

CAFROSE, (Wolter Corp) Atropine sulfate 1/600 gr., camphor 1/10 gr., caffeine anhydrous 1/10 gr./Tab. Bot. 100s.

Use: (1-2 q. 4 h.) Antispasmodic.

CAGOL, (Horvey) Gomenol 0.12 Gm., guaicol 0.1 Gm., eucalyptol 0.08 Gm., iodoform 0.2 Gm., camphor 0.05 Gm./2 cc. olive oil Amp. 2 cc.

Use: i.M., expectorant

CAJUPUT OIL IN PEANUT OIL. See: Cajundol, Liq. (Squibb).

CAKNAM, (Paxton) Sod. mag. & pot. citrate, with Cal. lactate, sod. phosphate, sod. bicarb.; total citrate 70% gran. eff. pow., Bot. 3½ oz.

Use: Orally to increase the alkali reserve.

CALA-COMP. (Sulliff & Cose) Calamine 20%, phenol 1%, benzocaine 3%, washable type base q.s. Oint. 1 lb., 5 lb.

Use: Antiseptic, local anesthetic, emollient

CAL-20 OINTMENT, (Ulmer) Calamine, zinc

* Generic name recognized by N.F., U.S.P. or A.M.A.

oxide, phenol, camphor, lanolin. 1 oz. jar, 1½ lb. jar.

Use: All-purpose dressing ointment.

CALABRYL, (Farke, Davis) 1% Benadryl HCl in specially prepared colomine base. Cream; tube, 1½ oz.; lotion, bot. 6 oz. Plastic Squeeze Bot. 80 cc.

Use: Antihistaminic & antipruritic.

CALAMATUM, (Toitby-Nasan) Colomine, zinc oxide, camphor, phenol. Tube 2 oz.

W/Benzocaine. Tube 2 oz.

W/Dazrin 2% (Methopyrilene HCl). Tube 1½ oz.

Use: Antipruritic & protective therapy.

CALAMER, (Schieffelin) Not listed. Calamine, menthol, salicylic acid, merbok. Bot. 8 oz., 16 oz., 1 gal.

Use: Topically, antiseptic skin therapy.

* **CALAMINE**, U.S.P. (Various Mfr.) Prepared. Pkg. ¼ lb., 1 lb. Lotion U.S.P., Lotion Phenolated U.S.P., Ointment, N.F.

Use: Astringent and mild antiseptic

See: Calatex, Lot. (Bayle)

CALAMCOLIN, (Upjohn) Colomine 5%, zinc oxide 5%, camphor-phenol 1%. Oint. 1½ oz., Jar 1 lb.

Use: Topically, protective & antiseptic.

CALATEX, (Bayle) Colomine lot. Plain, Bot. 4 oz., 8 oz., 1 pt. Menthalated & phenolated, Bot. 4 oz., 8 oz., 1 pt.

Use: Protective & antiseptic

CAL-A-TNESIN, (Walker Corp) Phenol 3/50 gr., cal. iodized 1/10 gr., benzocaine 1/12 gr., w/cottsoot, cubeb & ext. licorice/Tob. Bot. 100s.

Use: Relief of the discomfort of sore throats.

CAEAXIDE, (Elder) Calamine 40 gr., zinc oxide 20 gr., phenol 1¼/4 oz. Liq., Bot. 4 oz., 1 pt., 1 gal.

Use: Topically, protective & antiseptic.

CALBEDOCE, (Buffington) Purified bone phosphate 3¼ gr., dicalcium phosphate 3¼ gr., ferruz sulf. exsic. 1½ gr., vit. A 1000 u., D 500 u., B-1 1.5 mg., B-2 1.5 mg., niacinamide 5 mg., C 10 mg., B-12 2 mcg., folic acid 0.15 mg./tab. Bot. 100s, 1000s.

Use: Dietary supplementation in pregnancy and other periods of stress.

CALBEDOCE-LACTATE, (Buffington) Calcium lactate 7½ gr., ferruz sulfate exsic. 1½ gr., vit. A 1000 u., D 500 u., B-1 1.5 mg., B-2 1.5 mg., niacinamide 5 mg., C 10 mg., B-12 2 mcg., folic acid 0.15 mg./tab. Bot. 100s, 1000s.

Use: Phosphorus-free, dietary supplement in pregnancy.

CAL-BIS-MA, (Warner-Chilcott) Sod. bicarb. 9 gr., cal. carb. 5 gr., mag. carb. 3 gr., mag-

trisilicate 2 1/4 gr., bismuth subcarb 1/4 gr /
Tsp or Tab Paw, Pkg 3 oz, 12 oz; Tab
Bot 10s

Use: Hyperchlorhydria

CAL-BO-TAB. (Burlington) Dicalcium phos-
phate 3 3/4 gr., purified bone phosphate 3 3/4
gr., ext ferrous sulfate 1 1/2 gr., Vit D
500 u., niacinamide 10 mg., also trace ele-
ment cap, zinc, mang., mag., fluorine and
pot /Tab Bot 100s, 1000s

Use: Mineral & vitamin supplement

CALBRODIN (Upjohn) Codeine acetate 1/2
gr., chloroform 1 gr., Cal bromide, fl ext
belladonna root 1/4 min., benzoic acid 1/2
gr /A oz Bots 1 pt, 1 gal

Use: (1 tsp q 3 h) Sedative, expectorant

CALCETS. (Walker) Vit A 500 u., D 250 mg
cal phosphate 500 mg., w/trace elements/
Sphere Bot 60s

Use: Dietary supplement

CALCIBRONAT. (Sandoz) Double salt of cal
bromide galactogluconate Amp (1.25 Gm)
10 cc Box 5s, 20s, 100s; Gran, Trn, 100
Gm, 500 Gm 1 Tab effer, Tin 10s 50s

Use: Sedative

CALCI-C. (Physiological Chemicals) Cal as
corbate Tabs 3 gr Bots 100s, 300s 1000s

Use: Vitamin C therapy

CALCIPAS PHOS-FREE. (Nian) Dicalcium
carbonate 140 mg., cal gluconate 330 mg.,
Vit D 375 u /Tab Bot 100s, 500s

Use: (1 2 tld) Calcium & vitamin therapy

W/Iron Dicalcium phosphate 290 mg., Cal
gluconate 190 mg., ferrous gluconate 64
mg., Vit D 345 u /Tab Bot 100s 500s

(Nan).

Use: (1 2 tld) Calcium vitamin & iron
therapy

CALCICARS. (Hynson Westcott & Dunning)
Cal carb 10 gr., cinnamon oil 1/40 m/Tab
Bos, 3 tubes of 12s; Bot 100s, 500s, 1000s

Use: Orally (1 2 q 3 h) Gastric antacid

CALCI-D. (Irwin, Neisler) Dicalcium phos-
phate 7 1/2 gr., Vit D 400 u /Tab Bot 100s,
500s, 1000s

Use: Calcium therapy

CALCI-D-FERROUS. (Irwin, Neisler) Dical-
cium phosphate 7 1/2 gr., Vit D 400 u.,
ferrous sulfate, ext 35 mg /Tab Bot 100s,
500s, 1000s.

Use: Orally (1 2 tld) Dietary supplement

CALCIDIN. (Abbott) Available iodine com-
bination with lime and starch plus r
iodate

Pow. Bot 1 ex., 3 lb.

Tab. 1/3 gr (20 mg)

1 gr (60 mg) Bot

* Generic name recognif

(0.15 Gm), Bot 100s, 1000s; 5 gr (0.3
Gm), Bot 100s, 1000s

Use: Iodine therapy

CALCIDIN TROCHES. (Abbott) Calcidin 1/2
gr., anesthesin 1/4 gr., licorice ext., anise
oil Box 24s; Bot 100s

Use: Throat & mouth infections

W/Codeine phosphate, Troches (Abbott) Cal
iodide, ephedrine HCl, pentobarbital &
codeine phosphate Bot 25s

Use: Antitussive

CALCIDRINE, SYRUP. (Abbott) Codeine
phosphate 64.8 mg., nembulal sod 25 mg
ephedrine HCl 25 mg., cal iodide anh 910
mg /fl oz Bot 4 oz, 1 pt, 1 gal

Use: Treatment of cough due to colds

* **CALCIFEROL, USP, NND 1961, ADR**
Caps, Sain USP Vitamin D 2, viosterol
synthetic vitamin D, irradiated ergosterol
oxidized ergosterol

See:

(Ampuls & Vials)

Dee Osterol, Viol 50 000 u/cc (Brean)

Dee Ron, Viol 500,000 u/cc (Warren
Teed)

Derotal, Amps 50 000 u/cc (Brewer)

Ertron, Amps 500 000 u/cc (Whittier)

Hr Derotal Amps 200 000, 400,000, 600
000 u/cc (Brewer)

Metadec Amps 600,000 u/cc (Merck
Sharp & Dohme)

Sterogyl, Amps 600,000 u / 1.5 cc (Roussel)

(Cap & Tab.)

Condopals, Cap 50 000 u (Nian)

D Arthun, Cap 50 000 u (Horvey)

Davitin, Cap 50 000 u (Ives Cameron)

Decaps Cap 50 000 u (Cole)

Dee Osterol, Cap 50 000 u (Brean)

Dee Ron, Tab 50,000 u (Warren Teed)

Deltolm, Gelseal, 50 000 u (Lilly)

Derotal Cap 50,000 u (Brewer)

Diactol, Cap 50 000 u (Plessner)

Darel, Cap 50 000 u (Mastengill)

Drisdol, Cap 50 000 u (Winthrop Labs)

D Yohne Cap 50,000 u (Dorsey)

Ertron, Cap 50 000 u (Whittier)

Ds Derotal, Cap 200,000 u (Brewer)

Intron, Cap 100,000 u (Whittier)

Sterogyl (Roussel)

Vio-D Cap 50,000 u (Rowell)

(Solution)

Candel 50 M, Tsp (Nian)

Drisdol, Sol 250 u/drop (Winthrop Labs)

W/Bale calts See: Actophen, Cap (Bio-
chemical)

* whole & Vit E See: Viderol, Cap
, Davis

c acid & Vit E See: Evanol

- W/Iodo-Caseln & yeast. See: Alastorin, Tab. (Farastan)
- W/Phenol & benzocaine. See: Col-Vi-Nol, Oint. (Uimer)
- W/Sod salicylate. See: Ergasteron, w/Sodium salicylate, Cap. (Aldrich)
- W/Sod. salicylate & Vit. C. See: Deslyate, Cap. (Walker)
- W/Vit. A. See: Vitamin A w/Vit. D, Preps. (Various Mfr.)
- W/Vit. A, sol. lactate, ferrous gluconate, aluminum hydroxide, benzocaine. See: Calciloc, Tab. (Irwin, Neisler)
- W/Vit. A, sarotene & dibasic cal. phosphate. See: Supplomat, Cap. (Ayerst)
- W/Vit. A & sulfathiazole. See: Sulfavitin, Oint. (Chicago)
- W/Vit. A & Vit. C. See: Vit. A w/Vit. C, & Vit. D, Preps. (Various Mfr.)
- W/Vit. C, calcium, phos., fluorine. See: Calcigen, Cap. (Boyle)
- CALCIFORTIS-D.** (United Labs) Vit. D 150 u., Ca. 137 mg., phos. 137 mg./Tab. Bot. 100s, 1000s.
Use: Calcium therapy.
- CALCIDEN** W/Vit. C. (Boyle) Cal. 0.112 Gm., phos. 0.046 Gm., fluorine 0.076 mg., Vit. C 10 mg., Vit. D 125 u./Cap. Bot. 100s.
Use: Dental caries & calcium deficiencies.
- CALCIGRANULES.** (Walker Vitamins) Dibasic cal. phosphate 3 Gm., calciferol (Vit. D-2) 400 u./Tsp. Bot. 250 Gm.
Use: Calcium & Vitamin-D deficiencies.
- CALCILAC.** (Irwin, Neisler) Vit. A 1500 u., D 150 u., Cal. lactate 350 mg., ferrous gluconate 50 mg., aluminum hydroxide 150 mg., benzocaine 35 mg./Tab. Bot. 100s, 300s.
Use: Nutritional supplement in pregnancy
- CALCINATA IMPROVED.** (Niant) Iron 15 mg., Vit. A 5000 u., D 400 u., B-1 2 mg., B-2 2 mg., B-6 2 mg., B-12 2 mcgm., cal. pentothenate 5 mg., niacinamide 10 mg., C 50 mg., folic acid 0.3 mg., K 0.5 mg., iodine 0.15 mg., calcium 215 mg./Tab. Bot. 90s
Use: 1-2 tid; prenatal vitamin and mineral supplement.
- CALCIPHOS.** (Knoll) Cal. & iron salt of inositol-hexaphosphate. Pow. Bot. 3 oz & 6 gr. Tab. Bot. 100s, 500s.
Use: Calcium supplement during periods of increased demands on mineral reserves.
- CALCI-PLUS.** (Zemmer) Six vitamins plus calcium lactate and ferrous sulfate/Tab. Bot. 100s.
Use: Dietary supplement in pregnancy, convalescence.
- CALCISALIN.** (Warner-Chilcott) Vit. A 6000 u., D 400 u., C 100 mg., B-1 1.5 mg., R 2 2.5 mg., B-6 0.5 mg., niacinamide 15 mg., sol. pantothenate 5 mg., Vit. 5-12 6 mcg., cal. lactate 2 Gm., ferrous sulfate 50 mg., folic acid 0.25 mg., aluminum hydroxide gel 1.5 Gm./6 Tabs. Bot. 100s, 300s
Use: Dietary supplement & prenatal therapy.
- CALCIUM ACETYSALICYLATE.** Kalmopyrin, Isalsetal, soluble aspirin, tycolale.
See: Crystar, Pow. (Armour)
Sumorin. (Staley Drug Prods.)
Use: As aspirin.
- *CALCIUM ACETYSALICYLATE-CARBAMIDE.** See: Caluric, Tab. (Smith-Dorsey)
- W/Triaminic. See: Ursiev, Tab. (Dorsey)
- CALCIUM ALGINATE.**
See: Celginace (Mead Johnson)
Combinate (Mead Johnson)
W/Sodium alginate complex. See: Kalpin, Pow. (Wallace)
- *CALCIUM AMINOSALICYLATE, N.F., N.N.D. 1962.** Tab., N.F. (Dumas-Wilsae) 7½ gr. Bot. 1000s.
See: Parasol Calcium (Parroy)
Pasara, Cap., Tab. (Smith-Dorsey)
Use: (2-4 Gm. daily) Tuberculosis therapy.
- CALCIUM ASCORBATE.** See: Calc-C, Tab. (Physiological Chemicals).
- W/Benzyl alcohol and dextrose injection. See: Calcicorbate, Amp., Tabs. (Cole)
- W/Cal. gluconate and benzyl alcohol. See: Cal-Uote, Amp. (Paul Mooney)
- W/Cortisone acetate, aluminum hydroxide gel dried, Sod. salicylate & Cal. carb. See: Salfort, Tab. (Mastengill)
- *CALCIUM 4-BENZAMIDOSALICYLATE.** Calcium 4-benzamido-2-hydroxybenzoate. Aminocyl 8-PAS. See: Therapas, Pow. (Barnes-Hind)
- CALCIUM BENZOYL-p-AMINOSALICYLATE.** See: Cal. 4-Benzamidosalicylate.
- CALCIUM BIS-DIOCTYL SULFOSUCCINATE.** See: Dioctyl Calcium Sulfosuccinate.
- *CALCIUM BROMIDE, N.F. (Various Mfr.)** Pow. Pkg. ¼ lb., 1 lb.
Use: Sedative & anti-epileptic.
- W/Galactogluconate, double salt. See: Calcibronat, Amp. (Sandaz)
- CALCIUM CACODYLATE.** (Ingram) Vial 2 gr./cc. Box 12s, 25s, 100s.
- *CALCIUM CARBONATE, PRECIPITATED, U.S.P., A.D.B. (Various Mfr.)** Precipitated chalk, Tab., N.F.
Use: Antacid.
- W/Amino asetic acid, aluminum hydroxide, magnesium carbonate. See: Glytals (Desret)
- W/ASA, methylprednisaloe. See: Medaprin, Tab. (Upjohn)
- W/Aspirin, ascorbic acid, & menadione. See: A-C-K Buffered, Tab. (G. F. Harvey)

* Generic name recognized by N.F., U.S.P. or A.M.A.

actyl Sod sulfasuccinate See Celginace Tab & Gran (Mead Johnson)

W/Diactyl Sod succinate anthraquinone derivatives (Penstim) See: Combinate, Tab or gran (Mead Johnson)

CALCIUM SUCCINATE W/Aspirin

See Ber Ex Tab (Pan Pharm)

Dalcin, Tab (Dalcin)

W/Aspirin succinic acid, calcium succinate See Arthatabs Tabs (Calumbus)

W/Sod salicylate, succinic acid vit C See Arthatabs No 2 (Calumbus)

CALCIUM UNDECYLENATE, See: Cal Desene Pow (Maltbie)

W/Hydrocortisone, neomycin See: Caldecort (Maltbie Lab)

CALCIUM UBEA IODIDE W/Carbromal, theabromine See: Brozanthine, Tab (Blue Line)

CALCI-VITA, (Endocrine) Bone meal 0.3 Gm, Vit A 200 u, Vit D 400 u, Vit C 30 mg, Vit B 1.1 mg/Tab Bot 100s

Use (1-2 daily) Tissue development in pregnancy & lactation, prevention & treatment of dental caries & dental neurogia

CALCIWAFERS (Neon) Dicalcium phosphate 580 mg, Cal gluconate 380 mg, Vit D 750 u/Wafer Box 50s 250s

Use: (1-3 daily) Tissue development in pregnancy & lactation

CALCREOSE, NNR 1957 (Maltbie) Cal compounds of cresol guaiacol and other phenols present in creosote Tab plain or EC, 4 gr Bot 100s 500s 1000s Also available with calcium iodized Syr Bot 4 oz, 1 pt 1 gal

Use (1-3 daily) Expectoant

CAL-C-TOSE, (Roche Lab) Vit A 2000 u, Vit D 1000 u, Vit B 1.2 mg, Vit B 2 1 mg, Vit C 75 mg, niacinamide 5 mg, cal 200 mg, phos 160 mg, iron 10 mg/2 tsp Pkg 12 oz and 5 lb

Use: Dietary supplement

CALDEBROM (G F Harvey) Cal gluconate 0.5 Gm, strontium bromide 0.5 Gm, dextrose 0.5 Gm/10 cc Amp, Box 10s 100s, 250s

Use Epilepsy & allied conditions

CALDECORT, (Maltbie Lab) Hydrocortisone acetate 10 mg, cal undecylenate 3D mg, neomycin sulf 5 mg/Gm oint tube 7 Gm & 28 Gm

Use Dermatitis

CAL-DESENE, (Maltbie) Calcium undecylenate 15% in talc Pow shaker pkg 2 oz Use Fungus infection Diaper rash

CALDEX, (Endo) Cal gluconate 570 mg, cal D saccharate 50 mg, dextrose 1.4 Gm/10 cc Amp, Box 6s 25s 100s

Use: Calcium therapy

CALETTES #1 & #2, (Crystal) Bone meal 420 mg, ferrous sulfate 200 mg, Vit A 1000 IU, D 500 IU, C 25 mg, B 1.1 mg, B 2.1 mg, niacinamide 3 mg, K iodide 0.065 mg, Ca-sulfate 10 mg, Cu sulfate 10 mg, Mg-oxide 110 mg, Mn sulfate 140 mg, K sulfate 110 mg, Zn-oxide 125 mg, Na molybdate D4D mg #2 same but is phosphorus free/Tab Bot 100s

Use: Dietary supplement for pregnancy, lactation, anemias

CALFERBEE, (Smith, Miller & Patch) Cal phosphate tribasic (anhy) 400 mg, ferrous sulf 100 mg, Vit D 400 IU, ascorbic acid 25 mg, thiamine HCl 1 mg, riboflavin 1 mg/tab Bot 100s, 1000s; Calferbee NB 250 tab in EvenFlo Nursing Bottle

Use: Mineral Vitamin therapy

CALFERBEE LACTATE, (Smith, Miller & Patch) Calcium lactate 333 mg, ferrous sulf 100 mg, Vit D 400 u, Vit C 25 mg, B 1.1 mg, B 2.1 mg, aluminum hydroxide 125 mg/Tab Bot 100s, 1000s, and Calferbee Lactate NB 250s

Use: Dietary supplement

CALGAGE-S, (Squibb) Test Kit Bot 60 ml with test tube cap, & 2 droppers

Use: Test for urine calcium

CAL-GLU-DIN, (Paul Money) Ferrous gluconate 2½ gr, Vit D 400 u, B 2.1 mg, B 1.1 mg, Cal lactate 7½ gr/Tab Bot 100s

Use: Dietary supplement

CALGLYPNOS-D, (Mellibie) Cal gluconate 3 gr, dicalcium phosphate B gr, Vit D 450 u/Tab Bot 100s, 500s, 1000s

Use (1-2 daily) Calcium & Vitamin D therapy

CALGLUQUINE, (Sandoz) Not listed Cal gluconogalactogluconate 137.5 mg, quinine gluconate 60 mg/cc Vials 5 & 10 cc

Use: Uterotonic

CAL-G-SOL, (Harvey) Calcium gluconate 0.5 Gm, cal levulinate D.5 Gm/10 cc Vial 10 cc

CALHISTINE TABS. & ELIXIR, (Caldwell & Blair) Phenylephrine HCl 5 mg, methapyrilene fumarate 125 mg/Tab and Tip Elixir also contains chloroform 1 min, menthol qs, wild raspberry base qs Bot Tab 100s 1000s Elixir pt, gal

Use: Nasal congestion, bronchial coughs of allergic origin

CALIGESIC OINTMENT, (Merck Sharp & Dohme) Calamine prepared B Gm, benzoic acid 3 Gm, hexylated m creal DD.5 Gm/100 Gm Tube, 1½ oz, 4 oz

Use: Local dermatitis & inflammatory conditions

CAL I-M. (Myers-Carter) Calcium levulinate 1.5%, cal. lactate 1%. Vials 10 cc.

CALINESE. (Haskell) Cal. carb. 500 mg., mag carb. 60 mg., mag. trisilicate 200 mg / Tab. Bot. 100s, 500s.
Use: Antacid

CALIOBEN. See: Calcium Iodobehenate

CALIPHOS. (New Mexico Pharm. Co.) Calcium glycerophosphate 1%, calcium levulinate 1½%, chlorobutanol 0.55%/cc. Liq. Vials 100 cc.

CALIRAD SUPES. (Winthrop Labs) Vit. A 1500 u., vit. D 300 u., vit. B-1 1 mg., vit. B-2 1 mg., vit. C 30 mg., nicotinamide 8 mg., pot. iodide 0.1 mg., dibasic cal. phosphate 0.58 Gm., ferrous gluconate 50 mg., menadiolone 0.15 mg./cop. Bot. 50s, 250s.
Use: Vitamin & mineral supplement in pregnancy & lactation.

CALMALOID. (G. F. Hervey) Colomine 8½%, phenol 2%, zinc oxide 4½%, rose water. Oint. Tubes 1 oz.; Jars, 1 lb.
Use: Topically, antiseptic.

CALMIRIN. (Kremers-Urban) Aspirin 4 gr., cal. glutamate 2 gr./tab. Bot. 100s, 1000s.
Use: {1-2 q. 2-4 h} Analgesic & antipyretic.

CALMITOL. (Leeming) Camphorated chloral, hyoscyamine elcote, camphor, menthol with chloroform, ether and alcohol. Liq. Bot. 2 oz.
Use: Antiseptic solution.

CALMITOL OINTMENT. (Leeming) Same as calmitol liquid with zinc oxide added to lanolin-petrolatum base. Tube 1½ oz.; Jar 1 lb.
Use: Topically, antiseptic ointment.

CALMOZOL LOTION. (Nationol) Colomine, zinc oxide, phenol, isopropyl alcohol, in a glycerinated vehicle. Lotion 4 oz., 1 pt., 1 gal.
Use: Topically, protective & antipruritic.

CALMULSION. (Upjohn) Dicalcium phosphate 100 gr., Vit. D 2700 u./fl. oz. of emulsion Bot. 12 fl. oz., 1 gal.
Use: (1-2 tsp. t.i.d.) Dietary supplement for calcium & vitamin-D therapy.

CAL-O-B. (Warren-Teed) Cal. glycerophosphate 500 mg., ferrous gluconate 200 mg., vit. D 1,000 u., B-6 1 mg., C 50 mg., w/trace minerals/tab. Bot. 100s
Use: Pregnancy & lactation.

CALOCYLATE. (Kremers-Urban) Aspirin 5 gr., ascorbic acid 15 mg., cal. glutamate 2 gr./Tab. Bot. 100s, 1000s.
Use: (1-2 q. 3 h) Analgesic & antipyretic.

CAL-O-NOL. (Paul Money) Colomine, zinc oxide, camphor, & phenol. Also available with benzocaine Oint 1 oz., 1 lb
Use: Topically, protective & anesthetic with antiseptic.

W/Antihistamine. (Paul Money) Sulfur resorcinol, colomine, zinc oxide, bentonite magma, cal. hydroxide, and pyrilamine maleate 1%. Oint. 1 oz., 1 lb.
Use: Topically, dermatitis.

W/Resorcinol. (Paul Money) Sulfur, resorcinol, colomine, zinc oxide, bentonite magma, calcium hydroxide. Oint. 1 oz., 1 lb.
Use: Topically, dermatitis

CALOPNYLL ELIXIR. (Caldwell & Bloor) Theophylline 80 mg./15 cc. Bot. pts., gals.
Use: Angina and asthmatic attacks.

CALOBIGEN 1000. (Baxter) Sucrose 175 Gm., casein hydrolysate 60 Gm., abs. alcohol 19 cc., sod. chloride 3.1 Gm., pot. chloride 2.2 Gm., cal. glycerophosphate 1.4 Gm., mag. chloride, hexahydrate 1.1 Gm./1000 cc. Bot. 1 L.
Use: For tubal nutrition.

CALOTEX. (Texas Pharm) Colomine Lotion Bot. 1 pt. & 1 gal.

CALOXIDE. (Blue Line) Colomine, zinc oxide, phenol, glycerin. Lotion, 1 pt., 1 gal.; cream w/Benzocaine 1½ oz.; oint. 1½ oz.
Use: Topically, protective & antiseptic.

CALOXIDINE (IODIZED CALCIUM).
See: Calcium Iodized.

Sulphadine-Strassenburgh.

CALPANATE. (Consumers Vit.) d Calcium pantothenate 10 mg./Cop. Bot. 100s.

CALPHOSAN. (Carlton Corp.) 1% cal. lactate & cal. glycerophosphate. Amp 10 cc., Box 10s, 100s; Vials 60 cc., available w/Vit. B-12, Vit. 60 cc.
Use: I.M., I.V.; Calcium therapy.

CALPURATE. (Maltbie) Theobromine cal. gluconate 7½ gr./Tab. Bot. 100s, 500s, 1000s, Pow. Bot. 1 oz.
Use: (1-2 daily) Diuretic.

W/Phenobarbital. (Maltbie) Theobromine cal. gluconate 7½ gr., phenobarbital ¼ gr./Tab. Bot. 100s, 500s, 1000s.
Use: (1-2 daily) Diuretic with sedative

CAL-BON OS. (Rowell) Vit. A 500 u., D 500 u., C 60 mg., citrus bioflavonoids 60 mg., Vit. B-6 5 mg., B-12 1 mcg., K 1 mg., ferrous fumarate 200 mg., powdered base containing 8 minerals and trace elements 500 mg /Tab. Bot. 60s, 250s, 1000s.
Use: Nutritional supplementation.

CALSAMATE. (F. Traut) Aspirin 5 gr., cal. glutamate 2 gr., ascorbic acid 15 mg /Tab. Bot. 100s, 1000s.
Use: (1-2 q. 4 h) Analgesic.

CALSANS. (PodiatRex Ltd) Calcium, Vit. C, D and glutamic acid HCL. Phosphorus free. Tabs. 100s.

Use: Treatment of low calcium tetany.

CALSARBAIN. (Sutliff & Case) Cal. carbo-

note 195 mg, phenobarbital 16 mg, sod bicarbonate 195 mg, papain 65 mg, pan creatin 65 mg, benzocaine 16 mg, flavoring agents q s /Tab 100s, 1000s
Use: Antacid w/sedative

CALSCORBATE, (Cole) Cal ascorbate 500 mg, benzyl alcohol 1½%, dextrose 5%
Amp 5 cc & 10 cc, Box 10s, 25s

Use: (5 cc I G or I V) Calcium & vitamin C deficiencies

CALSCORBATE ORAL, (Cole) Cal ascorbate 30 mg, dicalcium phosphate 108 Gm, Vit D 2 800 u./Tab Bot 100s

Use: (1 2 T i d) Calcium & vitamin therapy
CALSECOF, (Massengill) Ethylmorphine HCl 1/24 gr, chloralform ½ gr, labelka 1¼ gr, tartar emetic 1/48 gr/5 cc Syr Bot 1 pt, 1 gal

Use: Anodyne, expectorant and bronchial sedative

CALSI-MAG, (Smith Dorsey) Cal carb 25 gr, mag carb 8 gr, sod bicarb 9 gr, koolin 15 gr, bismuth subnitrate 3 gr /Tsp Pow Cans 4 oz

Use: (1 2 Tsp q 3 h) Gastric hyperacidity

CALSISALIN (Harrower) Vit A 6000 u, D 400 u, C 100 mg, 8 l 1.5 mg, 8-2 25 mg, 8-6 0.50 mg, niacinamide 15 mg, cal panthothenate 3 mg Vit B 12 6 mcg, folic acid D 25 mg, cal lactate 2 Gm, ferrous sulfate 50 mg, aluminum hydroxide dried gel 1.5 Gm/6 Tab Bot 100s

Use: (2 3 T i d, p c) Dietary supplement

* **CALSULPHYDRYL**. See: Hydrosulphosol Preps (Lients).

CALTALE, (Parke Davis) Cal carb 10 gr, with pepsin & taka-d estase/Tab Bot 100s; Tin 1000s

Use: (1 2 pr n) Digestive & gastric antacid

CALTHY, (Sull N & Case) Eucalyptol 058% alcohol 19.6%, benzoic acid 28%, menthol .026%, boric acid 3.0%, methyl salicylate .07%, resorcinol .32%, thymol .066%, glycerin 1.427% liq Bot pt, gal

CALUATE, (Paul Maney) Cal ascorbate 500 mg, cal gluconate 500 mg, benzyl alcohol 1.5%/10 cc. Amp Box 6s, 25s, 100s

Use: (5-10 cc. weekly) Calcium & vitamin therapy

CALUBIN (Smith Dorsey) Soluble calcium acetylascrylate-carbamide eq to 300 mg aspirin/tab Bots 100s, 500s

Use: Analgesic.

CALVATIN-C, (Smith Dorsey) Bone phos phosphate 43 mg, irradiated yeast pow [Vit D 200 u] 35 mg, thiamine HCl 1 mg, riboflavin 1 mg, ferrous lactate 100 mg. C 25 mg /Tab Bot 100s, 1000s.

Use: Dietary supplement

* Generic name recognized by N.F., U.S.P. or A.M.A.

CALVEROL, (Elder) Cal citrate 4 gr, ferrous gluconate 1 gr, Vit D 500 u /Tab Bot 100s, 500s, 1000s

Uses Calcium and iron therapy

CAL-VI-NOL OINTMENT, (Ulmer) Cod liver oil w/vit A & D 1 oz lube, 1 lb jar

Use: Stimulate granulation in lesions

CALZINOCAINE, (Pitman Moore) Creme tube 1 oz, lb jar

CAL-ZO, (Ulmer) Calamine, zinc oxide, phenol D 5%, camphor, lanolin oint base

Cont jar 1 oz, 1¼ lb

Use: Antiseptic dressing

CAMATUSS EXPECTORANT, (Combridge) Codeine sulfate 10 mg, sodium citrate 300 mg, ammonium chloride 150 mg, tartar emetic D 5 mg, potassium guaiacollanate 300 mg, pyrimamine maleate 10 mg, alcohol 6%/Tsp Bot Pts

Use: Coughs and colds

CAMINOIDS, N N D 1962 (Arlington Funk) Protein Hydrolysates Oral Aminopeptidols

N N D 1959 Amino acids and polypeptides derived by enzymatic digestion Dry, granu

lor pow Bot B oz; Cans 1 lb, 5 lb, 10 lb

Use: Protein supplement

CAMIROL, (Smith, Muller & Patch) Camphor menthol, sodoform, antimony triiodide, in

peanut oil Amp 1 cc, Box 5s 25s, 100s; 2 cc, 5s, 25s, 100s; Vial 10 cc

Use: I.M. Respiratory infections

CAMOFORM (Parke, Davis) Bismolical Hy drochloride Diethylamylal HCl Diethyl di

ethylaminoethyl phenol dihydrochloride 1ab (250 mg) Bot 20s

Use: Amebitoid agent

CAMOLLIENS LOTION, (Upjohn) Calamine, zinc oxide, benzocaine, myristyl γ picolinium chloride lotion, 4 oz, 1 pt, 1 gal

Use: Topically, protective & antiseptic

CAMOLLIENT CREAM, (Upjohn) Calamine zinc oxide, camphas, phenol, Cream, Bot 1½ oz.

Use: Topically, protective & antiseptic.

CAMOPRIM, (Parke, Davis) Amodiaquin HCl (Camoquin) 350 mg, primaquine phosphate 35 mg /Tab Bot 100s Infotabi Camoquin

75 mg, primaquine 15 mg /Tab

Use: Against parasites of malaria

CAMOQUIN NYDROCHLOBIDE, N N D 1962 (Parke, Davis) Amodiaquin HCl, 4(7

Diethylaminomethyl-4 hydroxyquinoline)-7 chloroquinoline, dihydrochloride, dihydrate

Tab Q 2 Gm Folder 3s; Pkg 36s; Bot 100s

Use: (2 5 first dose) Antimalarial

CAMPHELLA, MODIFIED, (Pitman-Moore) Belladonna flex 1 min, camphor 2 gr, alcohol 14%/fl oz Bot 1 pt, 1 gal.

Use: (1-2 tsp. q 3 h) Check secretions of the upper respiratory tract.

CAMPHELLA W/Ephedrine. (Pitman-Moore) Same as Camphella, modified with addition of ephedrine sulfate 1 gr./fl. oz. Bot. 1 pt., 1 gal.

Use: (1-2 tsp q 3 h) Treatment of bronchial asthma, urticaria & pertussis.

CAMPNODAL-SPARTEINE. (Torigion) Sparfeine sulfate $\frac{1}{2}$ gr./2 cc. Amp. 2 cc., Box 12s, 25s, 100s.

Use: Cardiac, respiratory & central nervous system stimulant.

***CAMPHORATED PARACHLOROPHENOL,** N.F., A.D.R. 1955.

CAMPHORIC ACID ESTER of p-Tolyl-methylcarbinol as Diethanolamine Salt. See: Gallotens, Tab. (Massengill).

CAMPHRO-SALYL. (Fougera) Benzyl salicylate $\frac{1}{2}$ gr., camphor 1/10 gr., in olive oil/5 cc, Amp. 5 cc., Box 6s.

Use: 5 C., I.M., Antirheumatic.

CAMPOLON. (Winthrop Labs) liver Inj., crude, Amp. (2 U.S.P. inj. u./cc) 2 cc, Box 5s, 100s; Amp 5 cc., Box 3s; Vial 10 cc., Box 1s, 50s.

Use: I.M., hemotopoeitic.

CAMPOLON FORTE. (Winthrop Labs) liver Inj., 15 U.S.P. inj. u./cc, Amp. 1 cc., Box 3s; Vial, 5 cc. & 10 cc; Box 1s.

Use: I.M., hemotopoeitic.

CAMSO LINIMENT. (Sulliff & Case) Camphor, gum 1.25%, ammonium chloride 3%, ammonia water, stronger 1.60%, oil origonum 0.75%, oil turpentine 14%, soap 3.50%. Liniment 1 pt., 1 gal.

Use: Topically, counterirritant

CAMUSOL. (Central) Camphor 60 mg, eucalyptal 80 mg, guaiacol 100 mg, Iodoform 15 mg, oil niaouli 10 mg, in sesame oil/2 ml. Vial 30 cc.

CANDAVITE. (Jackson-Mitchell) Multivitamin preparation in candy. Bot 50s.

Use: Dietary supplement.

CANDETTES. (Pfizer) Bacitracin 50 u., polymyxin B 1000 u./Traches. Pkg. 10s.

Use: Throat & mouth infections.

CANDETTES COUGH-JEL. (Pfizer) Carbetapentone citrate 7.25 mg./tsp. raspberry jel.

Use: Cough.

CANDETTES LOZENGES. (Pfizer) Carbetapentone tannate 7.25 mg. Box 10s

Use: Suppress cough.

CANDETTES SYRUP. (Pfizer) Carbetapentone citrate 7.25 mg., pyrilamine maleate 12.5 mg., phenylephrine HCl 5 mg., amm. chlor. 50 mg./tsp. Bot. 2 oz.

Use: Cough.

CANDETTES TABLETS. (Pfizer) Phenylephrine HCl 5 mg., salicylamide 97 mg., phenacetin

*Generic name recognized by N.F., U.S.P. or A.M.A.

97 mg., caffeine 20 mg., pheniramine maleate 10 mg., Vit. C 20 mg./Tab. Bot. 24.

Use: Colds

CANDICILLIN. (Premo) Pot. penicillin G 25,000 u., bacitracin 50 u./Troche. Pkg. 10s.

Use: Vincent's Infection.

CANDYCON. (Allison) Chlorprophepyridamine maleate 2 mg., phenylephrine HCl 5 mg./Tab. Bot. 50s.

Use: Antihistamine-Decongestant.

CANICHRONE. (Nian) Vit. B-1 2 mg., Vit. B-2 2 mg., niacin 10 mg./Tab. Bot. 90s, 500s.

Use: Vitamin-B complex therapy.

CANTAXIN. (Winthrop Labs) Ascorbic acid Tab. (100 mg. & 500 mg.) Bot. 100s; Amp. (100 mg.) 2 cc.; Box 100s; Amp. (500 mg.) 5 cc., Box 5s, 20s. Solution in propylene glycol (100 mg./cc). Box 10 cc.

CANTHARADIN. See: Cantharone (Ingram)

CANTHARONE 1lg. (Ingram). Cantharidin 0.7% in vehicle with cellulose ethyl ether.

Bot. 15 cc

Use: Treatment for removal of warts

CANTIL, N.N.D. 1962. (Lakeside) Mepenzolate methylbromide, N-Methyl-3-piperidyl-diphenylglycolate methobromide. Tab. 25 mg., Bot. 100s

Use: Postganglionic parasympathetic inhibitor, in the treatment of varied colonic disorders.

CANTIL W/PHENDBARBITAL. (Lakeside) Cantil 25 mg., phenobarbital 16 mg./Tab Bot. 100s.

Use: Sedative with a postganglionic parasympathetic inhibitor.

CANTILYN. (Lakeside) Mepenzolate bromide 30 mg., kaolin 3 Gm, pectin 150 mg/15 cc. Cantilyn W/Neomycin in addition neomycin sulfate 300 mg Bot. 6 fl. oz

Use: Treatment of diarrhea.

CANTILYN W/NEOMYCIN 110. (Lakeside) Mepenzolate bromide 30 mg; kaolin 3 Gm; pectin 150 mg Neomycin 300 mg /tsp Bot. 6 oz.

Use: Antispasmodic Combination

CAPATHYN. (Pfizer-Eaton) Pyrethrin HCl 20 mg., acetylsalicylic acid 0.23 Gm., acetophenetidin 0.15 Gm., caffeine 0.03 Gm /Cap Bot 100s, 500s, 1000s.

Use: (1-2 q. 3-4 h) Antihistaminic, antipyretic & analgesic.

CAPILOID. (Pearson & Covey) Hesperidin 100 or 200 mg. ascorbic acid 100 mg./tab. Bot. 50s, 100s, 1000s

Use: (1-2 t.Ld.) Capillary fragility.

CAPILON. (Plessner) Citrus bioflavonoid complex 100 mg., rutin 100 mg., ascorbic acid 100 mg./Tab. Bot. 100s, 500s, 1000s.

Use: (1-2 t.Ld.) Capillary fragility.

CAPLA. (Wallace Labs) Mebutamate 2 Methyl-2 sec butyl 1,3 propanediol 300 mg /Tab Bot 100s

Use: Tranquilizer in treatment of hypertension

CAPRE. (Marian) Vit A 5000 u, D 400 u B 1 20 mg B 2 20 mg, B 6 50 mg, B 12 20 mcg, C 100 mg, calcium pantothenate 10 mg folic acid 0.3 mg, niacinamide 100 mg, iron 500 mg from 150 mg ferrous fumarate calcium 2000 mg, d sorbitol 100 mg /Tab, Bot 100s

Use: Vitamin and mineral deficiencies

* **CAPROCHLORONE.** Levo-gamma (alpha chloro benzyl) delta alpha gamma phenylcaproic acid

CAPROCID. (Merck) Aminocaproic acid

Use: Antifibrinolytic agent

CAPROKOL CAPSULES, N N R 1934 (Merck Sharp & Dohme) 0.15 Gm Hexylretoracinal in oil/Cap Bot 100s

Use: (1:1:1) Urinary antiseptic

CAPROKOL PILLS See Crystoids antihelminthic (Merck Sharp & Dohme)

* **CAPRIOMYCIN.** (Eli Lilly) An antibiotic derived from streptomyces capreolus

Use: Antibiotic

CAPROHEX OINTMENT, (Horton & Converse) Copper caprylate 0.5%, zed caprylate 10% caprylic acid 3%, water soluble base q.s Tube 1 oz 16 oz

Use: Topically, fungicidal

CAPRYLATE COPPER, W/Sad caprylate & caprylic acid See Capronex Oint (Horton & Converse)

CAPRYLATE-PROPIONATE MIXTURES, N N R 1954 See Sapranol, Preps (Wyeth)

CAPRYLATE, SALTS

See Aluminum Caprylate

Copper Caprylate

Kaprylex, Cap (Strassenburgh)

Sodium Caprylate

Zinc Caprylate

CAPRYLATE SODIUM, INJ, N N R (Chemo

Pural) Paw Bulk

Ingram—Amp 33%, 1 cc Pkg 12s 25s, 100s

Strassenburgh—Sol 0.2 Gm/cc Bot 60

cc, 480 cc 3 B 3 1

Use: Fungus infections of skin

See Sodium Caprylate Preps

W/Copper caprylate & caprylic acid See:

Caprinex Oint (Horton & Converse)

W/Zinc caprylate

See: Caprylic Comp, N N R 1954

Naprylate, N N R 1954 (Strassenburgh)

CAPRYLIC ACID W/Copper caprylate &

sad caprylate See Capronex Oint (Horton & Converse)

* Generic name recognized by N F, U S P or A M A

W/Copper caprylate, salicylanilide, p chloro xyleneol See: Priculex, Oint (Elder)

CAPRYLIC COMPOUND, N N R 1955 See: Naprylate, Preps (Strassenburgh)

CAPRYLIUM DOUCHE CREME POWDER

& TABLETS. (Strassenburgh) Pkg 3 Gm; Box

12; Vaginal Tab replaces the Naprylate

Vaginal Suppositories, Pkg 24s; Vaginal

Crema, new name for Naprylate Vaginal

Crema, Tube 90 Gm

CAPSIRON, H H D 1962 (Pilman Moore)

Cadmium sulfide susp 1% in a shampoo

base Plast Sqz Bot 4 oz

Use: Treatment of seborrheic dermatitis

CAPSICUM OLEORESIN, H F. (Various Mfr)

W/Camphor, corn oil, cajuput oil and tur

pentine oil See: Capsolin, Oint (Parke

Davis)

CAPSOLIN. (Parke, Davis) Oleoresin capsi

cum camphor, croton oil, cajuput oil &

turpentine oil Tube 1 1/4 oz; Jars 1 lb,

5 lb

Use: Topically, counterirritant

CAPSULUGOL'S. (Frost) Lugol's sol 0.3 cc/

Cap Bot 21s, 50s, 100s

Use: Orally, hyperthyroidism

CAPTIVITE-12. (Merck Sharp & Dohme) Vit

A 3000 u, vit D 1000 u, thiamine HCl 1.5

mg, riboflavin 1.2 mg, niacinamide 10

mg, Vit B 12 crystalline 5 mcg, ascorbic

acid 40 mg/dr Bot 4 oz

Use: (1 tsp daily) Dietary supplement

* **CAPTODIAMINE HCl,** 4 Butylthio e

phenylbenzyl 2 dimethylaminoethyl sulph

ide Covatin hydrochloride See: Sufran

Tab (Ayeris)

* **CARAMIPHEN ETHANEDISULFOHATE**

W/Phenylephrine hydrochloride, pheninda

mine tartrate See: Dandril, Tab (Whitehall)

W/Chlorpheniramine maleate, phenylpro

panolamine HCl, Isopropamide See: Tusa

Ornade, Caps & Liq (Smith, Kline and

French)

* **CARAMIPHEN HYDROCHLORIDE,** N N D

1962 1 Phenylcyclopentanecarboxylic acid

2 diethylaminoethyl ester hydrochloride

See: Candettes (Pfizer)

Ponparnit, Preps (Geigy)

Use: Proposed for muscular spasm & tremor

in Parkinson's disease

CARBACEL Ophth Sol (Professional) Sterile

buffered aqueous methacellulose solution

containing carbachol USP in 3/4% and

1 1/2% strengths with benzalkonium chlo

ride 15 cc plastic direct application bats

Use: Chronic non congestive glaucoma

* **CARBACHOL,** U S P. Carbamylcholine chlo

ride Choline chloride carbamate, Lentin

Carbolin

See: Bulepto Carbocel, Sol. (Professional Pharm.)

Carbamiotin (Crookes-Barnes)

Carbafin (Merck Sharp & Dohme)

Daryl, Amp., Tab. (Merck Sharp & Dohme)

Use: Parasympathomimetic agent.

W/Methylcellulose. See: Isopto Carbachol, Soln. (Alcon)

* **CARBACRYLAMINE RESINS**, N.N.D. 1962.

See: Carbo-Resin, Pow. (Lilly)

CARBACRYLIC RESIN, QUININE, N.N.D.

1959. See: Diognex (Squibb)

CARBAK. (Hoack Labs.) Mephenesin 4 gr., carbromal 4 gr./tab. Bot. 100s, 1000s.

Use: (1 q.i.d.) Sedative for tension and stress.

CARRAMIDE, N.P., N.N.R. 1950, A.D.R.

1953. See: Urea (Various Mfr.)

CARRAMIDE COMPOUNDS,

See: Abasin, Tab. (Winthrop Labs.)

Acetylcarbromal (Various Mfr.)

Adalin, Tab. (Winthrop Labs.)

Apramol, Tab. (Ames)

Bromisovalum (Various Mfr.)

Bromural, Tab. Knoll

Corbrital, Elix. & Kap. (Parke, Davis)

Carbromal (Various Mfr.)

Carbropent, Tab. (Blue Line)

Car-Sed-Inc, Tab. (Schenley)

Isoval, Tab. (Mission)

Sedomyl, Tab. (Schenley)

Sedormid, Tab. (Roche)

Ureophil, Pow. (Abbott)

CARBAMIDE PEROXIDE. See: Urea-Peroxide.

CARBAMINE. (Key Corp.) Synthetic urea, citric acid, sodium bicarbonate/6 Gm. packet, Box 12s.

Use: Treatment of peptic ulcers.

CARBAMINO PNEYL-ARSONIC ACID.

See: Carbarsone (Various Mfr.)

CARBAMIOTIN. (Crookes-Barnes) Carbachol

3%, 1½% & 3% solutions. Vial 15 cc

Use: Miotic.

CARBAMYLCHOLINE CHLORIDE, U.S.P. See:

Carbachol, Prep. (Merck Sharp & Dohme)

CARRAMYLMETNYLCHOLINE CHLORIDE,

U.S.P., N.N.D. 1962.

See: Bethanechol chloride, U.S.P.

Urecholine chloride (Merck Sharp & Dohme)

CARBANESIN. (F. Traut) Mephenesin 4 gr.,

carbromal 2½ gr./Tab. Bot. 100s, 1000s.

Use: (1-2 t.i.d.) Sedative & relaxant.

* **CARRARSONE**, U.S.P., N.N.R. 1954; Cop-

N.F., Supp., U.S.P., Tab., U.S.P. (Lilly).

N-carbamylarsonic acid. Amobarb, ameban,

amiblorson, arsambide, fenarsone,

* Generic name recognized by N.F., U.S.P. or A.M.A.

leucarsonone, aminarsonone, amebarsone p-Car-

baminophenylarsonic acid.

Pow.: Vial (2 Gm.); Box 6s; Bot. 1 oz.

Cap. U.S.P.: (0.25 Gm.) Bot. 20s, 500s.

Supp. U.S.P.: (3 24 Gm.) Bot. 20s, 500s.

Tab. U.S.P.: (0.05 Gm.) Box 40s, 500s.

Tab. U.S.P.: (0.25 Gm.) Bot. 20s, 500s.

Use: Acute & chronic amebiasis & tricho-

moniasis.

CARRATIAZOLE OINTMENT. (Rare) Sulfathiazole 5%, cool tar 1%, aint. base & alcohol 3%. Jars 25 Gm., 4 oz., 1 lb.

Use: Topically, for sulfathiazole susceptible skin infections.

* **CARRAZOCROME SALICYLATE**, N.N.D.

1962. Adrenochrome Monosemicarbazone

Sod. Salicylate Complex.

See: Adrenasem Salicylate (Massengill)

Adretal, Preps. (Organon)

* **CARRENZIDE**. (Warner-Lambert Res. Inst.)

Ethyl 3-(α -methylbenzyl)carboxylate.

Use: Anti-depressant, monoamine oxidase inhibitor.

* **CARRETAPENTANE CITRATE**, N.F., N.N.D.

1962. 2-(Diethylaminoethoxy) ethyl 1-phenyl-

cyclopentyl-1-carboxylate citrate

See: Leucarbonate, Tabs. (Loulsons)

Toclase, Syr. Tab. (Pfizer)

W/Salicylamide, phenacellin, caffeine. See: Leucarbonate W/S.P.S., Caps. (Loulsons)

W/Terpin hydrate & chloroform. See: Toclaste Expectorant Compound (Pfizer)

CARRETAPENTANE TANNATE W/Chlorpheniramine tannate, ephedrine tannate, phenylephrine tannate. See: Rynatuss

Tablets (Irwin, Neisler)

W/Ephedrine tannate, phenylephrine tannate, chlorpheniramine tannate, pyrilamine tannate. See: Rynafon Expectorant, Liq (Irwin, Neisler)

CARREINOXYSTRINDOYL METHYLRESERPATE. See: Singaserp, Tab. (Ciba).

CARRITINIL SALICYLATE. See: Sal-Ethyl Carbonate, Pow., Tab. (Parke, Davis)

* **CARBIMAZOLE**. 1-Ethoxycarbonyl-2,3-dihydro-3-methyl-2-thiazimidazole. Neo-Mercazole.

CARBINOL. See: Methonal. (Various Mfr.)

CARBINOXAMINE, DEXTRO FORM. See: Twiston, Tab (McNeil)

* **CARBINOXAMINE MALEATE**, N.F., N.N.D. 1962. Tabs., N.F. Paracarbinoxamine 2-(p-

chloro- α -[2-dimethylaminoethoxy]benzyl) pyridine maleate

See: Citrine Maleate, Elix., Tab. (McNeil)

CARROCAINE HCl. (Winthrop) For infiltration and nerve block—mepivacaine HCl,

1% or 2% sterile saline soln. For caudal and peridural block—mepivacaine, 1%, in

sterile modified Ringer's soln 1%/50 cc /
infil n b; 2%/50 cc /infil n b; 1%/30cc /
caudal

Use Nerve block

CARBO-CORT. (Dome) Creme: Hydrocorti-
sone alc 5 mg /Gm, coal tar soln 3%
in acid mantle vehicle Jar 4 oz Lellans
Hydrocortisone alc. 2.5 mg /Gm, coal tar
soln 3% in acid mantle vehicle Rot 4 oz
Use Dermatoses

CARBO-CORT FORTE (Dome) Hydrocorti-
sone alcohol 1/2% coal tar solution 10%
Creme 1/2 oz

Use Dermatoses

CARBO DOME. (Dome Chemicals) Liquor
carbonis detergent 5% 10% or 15% in
acid mantle vehicle Cream Tube 1 oz, 4
oz 1 lb Lat on 2 oz, 4 oz, 1 pt
Use Recalcitrant dermatoses

CARBO-HIST CREAM (Poul Maney) Liquor
carbonis detergens 5%, pyrilamine male-
ate 2%, benzocaine 1%, menthol 0.5%
Tube 2 oz; Jar 1 lb

Use Eczema, seborrheic dermatitis & ec-
zemataid dermatitis

*** CARBOL-FUCHSIN SOLUTION, NF, NNR**
1955 (Veltex) Original fuchsin formula
known as Castellani's Paint Fuchsin 0.3%,
baric acid 1%, phenol 45%, resarcinol
10% acetone 5%, alcohol 10%, water q s
Bat 30 cc, 60 cc, 120 cc, 480 cc

See Carfusin, Sol (Rorer)

Castadern, Lat (Lannett)

Castellani's Paint (Horton & Converse)

Use Local anti-infective

*** CARBOMYCIN, NND 1960 ADR** An-
antibiotic from *Streptomyces halstedii* See
Magnamycin Tab (Pfizer)

CARBON DIOXIDE, NNR 1947 See Ener-
vac Phormalax Supp (Pharmacia Labs)

CARBONIC ANHYDRASE INHIBITOR

See Ademal (Squibb)

Cardrose (Upjohn)

Daranide (Merck Sharp & Dohme)

Diamox (Lederle)

Diuril (Merck Sharp & Dohme)

Esidrix (Ciba)

Hydra Diuril (Merck Sharp & Dohme)

Naturetin (Squibb)

Neptazone (Lederle)

Oretic (Abbott)

CARBONIS DETERGENS, LIQUOR. See Coal
Tar Solution

*** CARBON TETRACHLORIDE, NF, NNR**
1949, NF Benznatarm (Mahnckrodt)

Use Hookworm

CARBO-RESIN, NND 1962 (Lilly) Paw B
Gm packets Pkg 24; Bat 1 lb

Use: Sodium removing resins

* Generic name recognized by NF, USP or A.M.A.

CARBOSE D. See: Carboxymethylcellulose
sodium, Prep (Various Mfr)

CARBOWAX 400, USP; 1500, 1540,
NF; and 4000, USP, NNR 1955 Poly-
ethylene glycol 1500, 1540 & 4000

CARBOXYMETHYLCELLULOSE SALT OF
DEXTROAMPHETAMINE. (Carbasynphen)
See Bantrel Timed Tab (G W Carrick)

*** CARBOXYMETHYLCELLULOSE SODIUM,**
USP NND 1962 ADR 1956, Carbose
D, CMC (Merck-Paw Co)

See Almelase Tab (Meyer Lab)

Aperget, Sol & Tab (Northwest)

Bu Lax, Tab (Ulmer)

Carmethose, Liq NND 1958 (Ciba)

Cel U Sol, Tab (Wintersmith)

Cethyase, Tab. (Ascher)

CMC, Tab (Amer Pharm Co)

Natulose, Tab (National Drug)

Thylase Sod, Tab NND 1958 (Jackson
Mitchell)

W/Acetphenalisatin, prune conc See
Bulatin, Tab (Ulmer)

W/Aluminum hydroxide See: Cel Luma
Tabs (Superior Pharm)

W/rac Amphetamine phosphate & d Amphet-
amine phosphate See: Biphetacel, Tab
(Strassenburgh)

W/Atropine sulfate, phenobarbital & yeast
See: Alrolase, Cap (Dumas Wilson)

W/Brewers yeast See: Zymelose, Tab
(Glidden)

W/Brewers yeast & diacetylhydroxyphenyl-
isatin See Zylax, Tab (Glidden)

W/Diacetyldioxyphenylisatin

See Cethytin, Tab (Ascher)

Effergel, Paw (Stuart Co)

W/Digitoxin See: Foxalin, Tab (Orant)

W/Dihydroxy aluminum & aminoacetate
See Acichek, Tab (Valentine)

W/Dioctyl cad sulfasuccinate See: Stuart
Diatase, Cap (Stuart)

W/Dioctyl sodium sulfasuccinate, diacetyl-
hydroxyphenylisatin See: Dialose Plus
Cap (Stuart Co)

W/Kaolin & Cal carb See: Calostat, Liq
(Schenley)

W/Magnesium hydroxide See: Turicum
(Whittier)

W/Magnesium oxide See: Carmethose, Tab
(Ciba)

W/Magnesium oxide & trisentine See:
Carmethose Trisentine, Tab (Ciba)

W/Methylcellulose See: Ex Coloric waters
Water (Eastern Research)

W/Prune concentrate & diacetylhydroxy-
phenylisatin See: Prulose complex (Har-
rower)

W/Vit B1 See: Celbe, Tab (Cole)

W/Vit. B-1 & dextro-amphetamine phosphate. See: Amobese, Tab. (Cole).

3,3-CARBOXYMETHYLENE Bis-[4-Hydroxycoumarin]ethyl ester. See: Tromexan, Tab. (Geigy).

CARBOXYPHEN W/Butobarbital.

See: Bontril Timed No. 2, Tab. (G. W. Carrick)

CARBTRITAL ELIXIR. (Parke, Davis) Carbromol 6 gr., pentobarbital sod. 2 gr./fl oz. Bot. 1 pt.

Use: (1-2 tsp. q. 3-4 h.) Sedative.

CARBTRITAL KAPSEALS. (Parke, Davis) Carbromol 4 gr., & pentobarbital sodium 1½ gr. Half-strength, carbromol, 2 gr., & pentobarbital ¾ gr. Bot. 100s, 1000s.

Use: (1-2 p.m.) Hypnotic & sedative.

***CARBROMAL**, N.F., N.N.R. 1949. (Various Mfr.) Bramadiethylacetylurea, bramadal, nyclol, plonadalin, urodol.

See: Adalin, Tab. (Winthrop Labs.)

Kartryl. (Direct Lab.)

Use: Sedative & mild hypnotic.

W/Bromisovalum (Bromural). See: Tabarea, Tab. (Tablerack).

W/Dextroamphetamine sulfate, pambrom, & salicylamide. See: Pambromal, Tab. (Whittier)

W/Ephedrine hydrochloride. See: Ephdramol, Cap. (Blue Line).

W/Mephenezin.

See: Corbak, Tab. (Haack)

Carbanesin, Tab. (F. Traut)

W/Pentaerythritol tetranitrate See: Perased, Tab. (Mallard)

W/Pentobarbital Sod.

See: Carbrapent, Tab. (Blue Line).

Carbtrital, Kapseals. (Parke, Davis).

W/Scopolamine HBr. See: Car-Sed-Inc, Tab. (Raymer)

W/Scopolamine HBr, & Ray-Succinate See: Suxsed, Tab. (Raymer).

W/Theobromine, Cal-urea-iodide. See: Branxanthine, Tab. (Blue Line)

CARBROPENT. (Blue Line) Pentobarbital sod 1½ gr., carbromol 4 gr./Tab. Bot. 100s.

Use: (1-2 m.p.) Sedative & hypnotic.

CARB SEED HEMECCELLULOSE. See: Yacuas. (Ayerst).

CARBUNOL OINTMENT. (Blue Line) Chloral hydrate 20 gr., cade oil 20 gr., phenol 10 gr., salicylic acid 10 gr./oz. Jar. 2 oz., 1 lb

Use: Topically, local anesthetic with antiseptic.

***CARBUTAMIDE.** N-Butyl-N'-sulphanitylurea Bucral; BZ 55; Ivenot; Nadison.

Use: Oral hypoglycemic agent

CARCHOLIN. (Merck Sharp & Dohme) (Car-

* Generic name recognized by N.F., U.S.P. or A.M.A.

bachol U.S.P.) Corbamychochine chloride Powder, Bot. 0.5 Gm.

Use: Reduction of intraocular tension in glaucoma simplex.

CARDALIN. (Irwin, Neisler) Aminophylline 5 gr., aluminum hydroxide 2.5 gr., ethyl aminobenzoate 0.5 gr./Tab. Bot. 100s, 500s, 1000s.

Use: Congestive heart failure for the control of paroxysmal dyspnea & pulmonary edema.

CARDALIN-PHEN. (Irwin, Neisler) Aminophylline 5 gr., aluminum hydroxide 2.5 gr., ethyl aminobenzoate 0.5 gr., phenobarbital ¼ gr./Tab. Bot. 100s, 500s, 1000s.

Use: (1-2 t.i.d.) Congestive heart failure.

CARDELMYCIN. (Pfizer). See: Novobiocin

CARDENZ. (Boyer) Magnesium ascorbate 25 mg, magnesium oxide 15 mg, Vit. E 5 mg, Inositol 30 mg, magnesium p-aminobenzoate 10 mg, A 2000 i.U., B-6 1.5 mg., B-12 1 mcg, D 100 u., folic acid 0.3 mg, nicotinamide 20 mg, magnesium gluconate 200 mg, hesperidin complex 100 mg, iodine DDS 5 mg, potassium gluconate 50 mg /Tab Bot 100s.

Use: Atherosclerosis.

CARDIAMID. See: Nikethamide, (Varlau Mfr.).

CARDIAZOL. (Knott) Pentamethylenetetrazol (Metrozol) Amp. (0.1 Gm) 1 cc., Box 6s

Use: I.M., I.V., Central nervous system stimulant.

CARDID. (Cofe) Phenobarbital sod. ¼ gr., aminophylline 1½ gr., pot. iodide 1 gr./Tab. Bot. 100s, 500s, 1000s

Use: Treatment of bronchial and cardiac asthma.

CARDILATE. (Burroughs Wellcome) Erythryl tetranitrate 5, 10, or 15 mg /Sublingual Tab Bot. 100s.

Use: Prophylaxis and long-term treatment of patients with recurrent anginal pain.

CARDIOGRAFIN. N.N.D. 1962. (Squibb) Diazlozote methylglucamine 85% in a sterile, aqueous solution. Vial 50 cc

Use: Diagnostic agent.

CARDOL. (Smith, Miller & Patch) Crotaegus tr. 2 min., Digitalis tr. 2 min., strophanthus tr. 1 min., strychnine 1/100 gr./Tab Bot 100s, 1000s.

CARDIOMONE. (Enda) Adenosine-5 monophosphate 20 mg./cc. as the sodium salt Amp. 1 cc. Boxes 12s, 25s, 100s. Gel. (20 mg./cc., 100 mg./cc) Vials 10 cc.

Use: Vasodilator for angiospastic conditions as in angina pectoris.

CARDIOQUIN. (Purdue Frederick) Quinidine polygalacturonate 275 mg equivalent to

- quinidine sulfate 200 mg / Tab Bot 50s
Use Cardiac arrhythmias
- CARDOPHYLLIN** See Aminophylline (Various Mfr)
- CARDRASE**, NND 1962 (Upjohn) Ethox zolamide 125 mg / Tab Bot 25s, 100s 1000s Tab 62.5 mg Bot 100s
Use: Diuretic, carbonic anhydrase inhibitor
- CARENA**, See Aminophylline (Various Mfr)
- CARINOL** (Bonar) Phenobarbital $\frac{1}{4}$ gr, atropine sulfate 0.0194 mg, hyoscyne hydrobromide 0.0065 mg, hyoscyamine sulfate 0.1037 mg / tab or 5 cc Tab Bot 100s, 1000s Elix Bot 1 pt
Use Sedative and antispasmodic
- CARPUSIN**, NNR 1955 (Rorer) Carbolfuchsin paint, NNR 1955 Original fuchsin formula known as Castellini's Paint Fuchsin 0.3%, boric acid 1%, acetone 5%, alcohol 10%, water q.s Bot 10 cs, 120 cc
Use: Local anti infective Athlete's feet
- CARGENTOS**, NNR 1945 See Silver Protein Mild
- CARGET**, (Hoack Labs) Charcoal activated 200 mg, pepsin 32 mg pentobarbital sod 16 mg / tab Bot 100s, 1000s
Use: Digestant w/sedative
- CARICA PAPAYA ENZYME**, See Caroid (Amer Ferment)
Papain (Various Mfr)
W/Diastase See Caripeptic, liq (Upjohn)
- CARIEPTIC** (Upjohn) Carica papaya enzyme (papain) with diastase 14 gr / fl oz Bot 1 pt, 1 gal
Use: (2-3 tsp) Digestant
- * **CARISOPRODOL**, NND 1962 N isopropyl meprobamate isomeprobamate Nisopropyl 2 methyl 2 propyl 1,3 propanediol di-sorbamate
See Relo, Tab (Schering)
Same Tab (Wallace)
W/Acetophenetidin, caffeine See: Same Compound, Tabs (Wallace)
W/Acetophenetidin caffeine, sodium phosphate See: Same Compound W/Caffeine Tabs, (Wallace)
W/Prednisolone See: Sameart, Tab (Wallace)
- CARITOL** (Wyeth) Carotene, pro-vit A. Cap 25,000 U.S.P. μ with tacopherols 15 mg Bot 100s
Use: Vitamin A therapy
W/Vitamin D (Wyeth) Vit. A 15,000 μ , Vit D 3000 e / Cap Bot 100s
Use: Dietary supplement for vitamins A and D
- CARMITHOSE**, NND 1962 (Ciba) Tab., Sed carboxymethylcellulose 225 mg, mag oxide 75 mg Bot 100s; Liq 15%, Bot 12 fl oz, 1 gal
Use (1-2 tsd) Gastric antacid
- CARMITHOSE-TRASENTINE**, (Ciba) Sod carboxymethylcellulose 225 mg, mag oxide 75 mg, trosentine 25 mg / tab Bot 100s
Use: (1-2 q 3-4 h) Gastric antacid with antispasmodic
- CARNACTION**, (Covendish) Diaphragm muscle extract Bot oral sol 30 & 50 cc; Amp. for inj (1 cc), 12s, 50s, (2 cc) 12s, 50s, & 2 oz bot
Use Vasodilator
- CARNALAC**, (Carnation) Carnation evaporated milk, maltose dextrin syrup, vitamin D 12s
Use Prepared infant feeding formula
- CARNOSAL**, (Squibb) Elixir Bot 8 oz
- CAROB FLOUR**, See Araban, Paw (Nestle)
- CAROID**, A proteolytic enzyme obtained from Carica Papaya (Papain)
W/Bile salts & phenolphthalein (Americon Ferment) Bile salt compound $1\frac{1}{2}$ gr, nut vomica ex 1/16 gr, phenolphthalein $\frac{1}{2}$ gr, pulv saroid $\frac{1}{4}$ gr, cascara $\frac{1}{2}$ gr capsicum 1/80 gr / Tab Bot 20s, 50s, 100s 500s, 1000s
Use (1 tsd & 2 hs) Choleric & slow gogue
W/Charcoal (Amer Ferment) Tab, Bot 45s 150s, 500s, 1000s
W/Diastase (Amer Ferment) Tab, Bot 45s 150s, 500s, 1000s
W/Soda (Amer Ferment) Tab, Bot 45s 150s 500s, 1000s
- CAROTENE**, NNR 1954 (Wyeth) Pro-vit A Cap (5000 u/Gm. in oil) 50 cc. with dropper Bot; Sol (7500 u/Gm) Bot 50 cc. See Caritol, Cap (Wyeth)
Use Vitamin A therapy
W/Vit C, choline See: Ceetomin Tab. (Anglo-French)
W/Vit D See: Caritol W/D (Wyeth)
- * **CARPERIDINE**, Ethyl 1 (2 carbamylethyl)-4 phenylpiperidine 4 sarboxylate (Winthrop Lab)
Use: Anti tussive
- * **CARPHENAZINE MALEATE**, 1 (10-1) [4 (2 hydroxyethyl) 1 piperaziny] propylphenothiazin 2 yl) 1 propanone bis(hydrogen maleate See Prokeltazine Maleate (Wyeth)
- CARPHENOL** See: p Benzylphenyl Carbo-mate
- CARRBUTABARB** (Carrtane) Butobarbital sodium $\frac{1}{2}$ gr / Tab or 5 cc Tab Bot 100s pint, gal
Use: Sedative and hypnotic

CARRHIST ELIXIR. (Carrtane) Pyrilamine maleate 8 mg., phenindamine tartrate 5 mg., chlorpheniramine maleate 15 mg., phenylephrine HCl 5 mg / 5 cc. Bot. pts.
Use: Hay fever, allergic type of colds.

CARRITINIC-C. (Corrtone) Ascorbic acid 75 mg., Ferrous fumarate 200 mg / Tab. Bot 100s.

Use: Anemias.

CARTUSSIN-NN. (Corrtone) d-methorphan HBr 7.5 mg., glyceryl guaiacolate 43.2 mg., ammonium Cl 43.2 mg., pyrilamine maleate 4.12 mg., phenylephrine HCl 2.5 mg., sodium citrate 160.4 mg / 5 cc. Bot. 4 oz., pt.
Use: Coughs due to colds.

CABSALAM. Name approved by B.P.C. Benz-1,3-oxazine-2,4-dione O-Corbamoylsalicylic acid lactam.

CAB-SED-INE. (Raymer) Carbramol 250 mg., scopolamine hydrobromide 0.1 mg. / Tab. Bot. 100s, 500s, 1000s.

Use: (1-2 q. 3 h.) Mild sedative & spasmolytic actions.

CARTOSE. (Winthrop Labs.) Dextrins 41%, dextrose 17%, maltose 16%. Bot. 1 pt.
Use: Infant food.

CARTRAX. (Roerig) Atoxol 10 mg. & pentythryllyletranitrate 10 mg / Tab. yellow, 20 mg. / Tab. pink. Bot. 100s, 500s.
Use: Hypertension

CARYOLYSINE. See: Mustorgen (Merck Sharp & Dahme)

CARYOPHYLLIC ACID. See: Eugenol. (Various Mfr.)

CARYOPHYLLUS. See: Clove. (Various Mfr.)

CASACOMB. (Sulliff & Case) Sod. gentisate 150 mg., salicylamide 150 mg., d-Desoxyephedrine 0.5 mg., ascorbic acid 10 mg. / Tab. Bot. 100s, 1000s.

Use: (1-2 q. 3 h) Analgesic or antipyretic.

CASA-HEX. (Sulliff & Case) Mannitol hexanitrate 1/2 gr., phenobarbital 1/4 gr / Cap. Bot. 100s, 1000s.

Use: (1 t.i.d.) Vasodilator with sedative.

CASAKOL. (Upjohn) Palaxalkol, 250 mg & casanthranol, 30 mg / capsule or 5 cc. Bot. 16s, 100s; syrup, 4 oz., pt.

Use: Treatment chronic constipation.

CASALOSE. (Dumas-Wilson) Methylcellulose 1 gr., cascara sagrada ext. 1/2 gr. / Tab. Bot. 100s, 1000s.

Use: (1-2 t.i.d. & h.s.) Laxative.

* **CASANTHRANOL.** Cathartic anthranol glycosides of cascara sagrada. See: Peristim (Mead Johnson)

W/Diethyl sodium sulfosuccinate. See: Dicon, Tabs (Chicago Pharm)

W/Palaxalkol. See: Casakol, Syr. (Upjohn)

* Generic name recognized by N.F., U.S.P. or A.M.A.

CASATE SODIUM. (Sulliff & Case) Sod gentisate 0.5 Gm. / Tab. Bot. 100s.

Use: Treatment of arthritis.

CASCAPHEN. (Buffington) Belladonna lvs. ext. 1/2 gr., podophyllin 1/4 gr., aloin 1/4 gr., cascara sagrada ext. 1/4 gr., phenolphthalein 1/2 gr. / Tab. Bot. 100s, 1000s.

Use: (1-2 h.s.) Cathartic.

CASCARA GLYCOSIDES.

See: Casanthranol Casakol (Upjohn)

Peristaltine, Tab. (Ciba)

Peristim (Mead-Johnson)

* **CASCARA SAGRADA, U.S.P.** flex, N.F. (Various Mfr.)

W/Maltine. See: Maltine w/Cascara sagrada, liq. (Chickoff).

W/Methylcellulose & compound senna powder. See: Cellulose, Tab. (Babst).

CASCARA SAGRADA AROMATIC. (Merck Sharp & Dahme) Mullard formula, alcohol 18%. Not U.S.P., Bot. gal.

CASCARA SAGRADA EXTRACT, N.F. Tab., N.F.

W/Aspirin, nux vomica ext., Hyoscyamus ext. & Getsemium tr. See: A.A.S. Compound, Tab. (Tilden).

W/Belladonna lvs. ext., podophyllin, aloin & phenolphthalein. See: Cascophen, Tab. (Buffington).

W/Diethyl sodium succinate. See: Diethyl-Zem, Tab. (Zemmer)

W/Ox bile ext., phenolphthalein, sod. salicylate, menthol. See: Roystol, Tab. (Roemer).

W/Bile salts. See: Colopo, Tab. (Cornick)

W/Bile salts, papain, phenolphthalein, capsicum oleoresin. See: Toracol Compound, Tab. (Plessner).

W/Bile salts, phenolphthalein & capsicum oleoresin. See: Yeracolate, Tab. (Standard).

W/Bile salts, capsicum oleoresin & peppermint oil. See: Neo-Toracol, Tab. (Plessner).

W/Bile salts, phenolphthalein, capsicum oleoresin & peppermint oil. See: Toracol, Tab. (Plessner).

W/Dehydrocholic acid. See: Nublax, Tab. (Numotoxine).

W/Methylcellulose. See: Casalose. (Dumas-Wilson).

W/Phenolphthalein, aloin & ox bile. See: Glox, Tab. (Cole).

CASCARA SAGRADA FLUIDEXTRACT, N.F. See: Cas-Evac, liq (Parke, Davis).

Kasagra, liq. (Winthrop Labs)

CASCARA SAGRADA FLUIDEXTRACT AROMATIC, U.S.P.

W/Psyllium husk pow. & prune pow. See: Coxyllium, Pow. (Upjohn).

CASCAROMATIC W/Potassium thiocyanate. See: Thio-Cara, liq. (Sulliff & Case).

CASEC. (Mead Johnson) Cal caseinate Contains 88% protein & 16% cal Tins 3½ oz, 30 oz

Use: Concentrated protein diet

CASEIN PURIFIED. See Kirkavin, Vial (Kirk)

CASEMAL (Sutcliffe & Case) Sod Glyco taur cholate 1 gr, Sod salicylate 1 gr, phenol phthalein 1½ gr, Ext cascara sagrada 1½ gr, menthol 1/20 gr /Tab Bot 100s, 1000s

CAS-EVAC (Paiké Davis) Liq prep of cascara sagrada Bot 2 oz w/dropper, ¼ pt, 1 pt, 1 gal

Use Constipation

CASIDE (Endo) Sod iodide 20 mg, casein 10 mg, benzyl alcohol/cc Vial 30 cc, 60 cc

Use (1-2 cc q 3-4 days) Nonspecific protein therapy

CASNU. (Sutcliffe & Case) Asafetida 1 gr Fow Ext nux vomica 1¼ gr Fow Ext cascara sagrada 1 gr, ginger ¾ gr dios lose 1/20 gr, capsicum ½ gr /Tab Bot 1000s

CASSIA ACUTIFOLIA PADS W/Dehydrocholic acid See: Senabile, Tab (Purdue Frederick)

CASSIA ANGUSTIFOLIA SENNA

See Aperens, Tab (Brayten)

Roenten, Pow (Brayten)

CASTADIRM. (Lannell) A lotion of boric acid, resorcinol acetone & salin of Carbol Fuchsin Bot 1, 4 & 16 oz

Use: Eczematoid conditions of epidermatophytic origin

CASTELLANI'S PAINT. (Horion & Converse) Sol Bot 1 oz, 4 oz, 1 pt

See: Carbol Fuchsin Point (Yeltex)

Carfusin, Liq (Rorer)

CASTELLANI'S PAINT. (Vonpelt & Brown) Phenol 45%, resorcinol 10%, boric acid 1%, basic fuchsin 0.3%, acetone 5% Bot 4 oz, pt

Use Germicide

CASTOR OIL, USP, NNR 1941, Aromatic, Cap, NF See Neolaid (Lederle)

Use: Cathartic.

CASYLIUM. (Upjohn) Cascara flex, aromatic 29 min, psyllium husk pow 37½ gr, prune pow 12 gr /isp Fig 4 oz, 8 oz

Use: Laxative

CATAGEN. (Sherman) Estrogenic substance 2500 IU w/progesterone 25 mg/cc. Vial 5 cc.

Use: I.M. Estrogenic & corpus luteum therapy

CATARRHALIS COMBINED VACCINE. (Vario Mfr)

Use: Treatment of the common cold

* Generic name recognized by N.F., U.S.F. or A.M.A.

CATARRHALIS 'IMMUNOGEN'. (Paiké Davis)

Use Treatment of the common cold

CATHOCILLIN. (Merck Sharp & Dahme) Sod Novobiocin equiv to 125 mg of novobiocin, pot penicillin G 125,000 u/Cap Bot 16s, 100s

Use Antibiotic and penicillin therapy

CATHOCILLIN PORTE. (Merck Sharp & Dahme) Sod novobiocin 250 mg, pot penicillin G 250,000 u/Cap Bot 16s, 100s

CATHOMYCIN CALCIUM, NND 1962 (Merck Sharp & Dahme) Cal novobiocin NND 1958 125 mg /5 cc Syr Bot 2 oz, 1 pt

CATHOMYCIN SODIUM, NND 1962 (Merck Sharp & Dahme) Novobiocin Sod Cap 250 mg, Bot 16s, 100s Vial Lyovac (500 mg) 20 cc

Use (1 Gm initially, followed by 250 mg q s d) Staphylococcal infections

CATHOMYCIN SODIUM LYOVAC, NND 1962 (Merck Sharp & Dahme) Novobiocin sodium 500 mg /20 cc vial Vials 500 mg /20 cc

Use Staphylococci resistant to other antibiotics

CATNOZOLE. (Merck Sharp & Dahme) Cathomycin 125 mg and sulfa methylthiadiazole 375 mg /Tab Bot 24s, 100s

Use: Urinary tract infections

CATIONIC RESINS. See Resins, Sod & Pot Removing

CATOLAC BACTERIAL VACCINE. (U S Standard) M catarrhalis 400 million, pneumococcus, types I, II and III 300 million streptococcus hemolyticus 150 million streptococcus viridans 150 million, staphylococcus aureus 500 million, staphylococcus albus 500 million, Friedländer's bacillus 300 million in casein menstruum/cc Vial 10 cc

Use: Treatment of common & chronic colds

CATRON (Lakeside) Phenylisopropylhydrazine 3 mg and 6 mg /Tabs Bot 100s

Use Antidepressant Inhibit monoamine oxidase

CAUSALIN (Amfre Grant) Salicylamide 300 mg, calcium paminobenzoate 50 mg Vit C 50 mg, mephenesin 150 mg /Tab Bot 50s, 100s, 500s, 1000s

Use: Orally treatment of rheumatoid arthritis

CAV-ECOL. (Cavendish) Wheat germ oil 3 mg /Perle Bot 75s, 200s

Use: Vitamin E therapy

CAVED-5 TABS. (Hallandale) Liqueur bis meth subnitrate, aluminum hydroxide gel magnesium carbonate, sodium bicarbonate

- calamus, frangulin glycoside, starch and lalcum/Tab. Bot. 60s, 240s, 600s.
- CAVESTROGEN.** (Cavendish) Estrogenic substances, natural in oil. Amp. (2000 I.U., 5000 I.U., 10,000 I.U./cc.). Box 12s, 25s, 100s.
Use: I.M., Estragee therapy.
- CAVOLYSIN.** (Cavendish) Thyroid 65 mg., cal. iodobehenate 0.27 mg., phenolphthalein 16 mg., ascorbic acid 100 mg./Tab. Bot. 100s, 250s, 500s, 1000s.
Use: (1-2 I.I.d.) Anti-obesity.
- CAVON.** (Cole) Vit. A 1333 u., Vit. B-1 0.33 mg., Vit. B-2 0.67 mg., Vit. C 16.67 mg., Vit. D 1333 u., niacinamide 3.33 mg., dicalcium phosphate, dried, 0.21 Gm., ferrous sulfate, dried, 12 mg./Cap. Bot. 100s.
Use: Dietary supplement.
- CAVOSEPT.** (Cavendish) Dichlorisulfamide benzoate, Vaginal Tab. Tube 12s.
Use: Trichomonade.
- CAVOTESTRONE.** (Cavendish) Ext. of animal testis, 2 capan u./Amp. Amp., Box 4s, 12s.
Use: I.M., Androgen therapy for males.
- CAYTINE.** N.N.O. 1982. (Lakeside) *Protobylol HCl. α -(α -methyl-3,4-methylenedioxyphenethylamino)-methyl]-protocatechyl alcohol hydrochloride. Tab. 2 mg Bot. 50s. W/Pentaborbital 32 mg. Bot. 50s, Inhal 1:100, Bot. w/dropper 10 cc. Inf. 0.5 mg./cc. 1 cc. amp. Box 4.
Use: Bronchodilator effective orally, parenterally, and by inhalation.
- CAZOMETH.** (Blue Line) Methapyrilene HCl 1%, colamine, and zinc oxide. Lot. Bot. 4 fl. oz.
Use: Topically, antihistaminic and protective
- C-B TIME.** (Arca) Ascorbic acid 150 mg., Vit. B-1 10 mg., B-2 10 mg., B-6 5 mg., B-12 10 mcg., niacinamide 50 mg., calcium pantothenate 10 mg./Cap. Ascorbic acid 300 mg., thiamine HCl 15 mg., riboflavin 10 mg., pyridoxine HCl 5 mg., cabolamin concentrate 5 mcg., niacinamide 100 mg., d-panthelol 20 mg./5 cc. Caps., Bot. 30s, 100s. Liq. Bot. 4 oz.
Use: Vitamin supplements.
- C-B-U OINTMENT.** (Warren-Teed) Chlorophyll water soluble 1%, benzocaine 10% and urea 33%. Jar 1 1/2 oz., 16 oz., 1 lb.
Use: Topically, deodorant with local anesthetic.
- C-B VONE.** (U. S. Vitam) Vit. C 150 mg., citrus bioflavonoid compound 50 mg., B-1 25 mg., B-2 10 mg., B-6 1.5 mg., B-12 5 mcg., niacinamide 75 mg., panthelol 7.5 mg., choline 25 mg., inositol 15 mg., desiccated liver 100 mg., yeast 25 mg./Cap. Bot. 15s, 30s, 50s, 100s, 500s
Use: Nutritional supplement.
- C.C.G.** (Noyes) Acetaellid 2 gr., quinine sulfate 1/2 gr., podophyllin 1/40 gr., aloin 1/16 gr., acelite root 1/20 gr., capsicum 1/4 gr., atropine sulfate 1/2000 gr., camphor/Tab. Bot. 100s, 1000s.
Use: (1 I.I.d.) Analgesic.
- C.C.G. No. 2.** (Noyes) Dover's pow. 2/7 gr., ipecac 1/35 gr., atropine sulfate 1/2000 gr., acelite ext. 1/20 gr., capsicum 1/4 gr., podophyllin 1/40 gr., acetophenetidin 2 gr., camphor/Tab. Bot. 100s.
Use: (1 I.I.d.) Analgesic.
- CEANOTHUS AMERICANUS, EXTRACT.** See: Ceanothyn, Liq. (Flint, Eaton).
- CEANOTHYN.** (Flint-Eaton) Ceanothus americanus ext. Liq., Bot. 1 pt., 1 gal.
Use: (4 tsp. & may be repeated in 30 minutes) Increase the coagulability of the blood when orally administered.
- CEBIFOSTIS.** (Upjohn) Thiamine HCl 5.0 mg., riboflavin 5 mg., pyridoxine HCl 1.5 mg., cal. pantothenate 25 mg., nicotinic acid amide 50.0 mg., ascorbic acid 150.0 mg., folic acid 0.5 mg., Vit. B-12 activity 2 mcg./Tab. Bot. 100s, 500s.
Use: (1-3 daily) Vitamin-B complex therapy.
- CEBENASE.** (Upjohn) Intrinsic 1/6 anti-anemia oral v., Sod. ascorbate 150 mg., folic acid 3 mg., thiamine mononitrate 10 mg., riboflavin 10 mg., pyridoxine hydrochloride 2.5 mg., Cal. pantothenate 10 mg., nicotinamide 75 mg./Tab. Bot. 50s, 500s.
Use: Hematinic.
- CEBETINIC.** (Upjohn) Ferrous gluconate 5 gr., Vit. B-12 activity 5 mcg., folic acid 0.67 mg., thiamine HCl 0.5 mg., nicotinamide 10 mg., ascorbic acid 25 mg./Tab. Bot. 60s, 500s.
Use: Hematinic.
- CEBINOLDS.** (Hartan & Converse) Hesperidin complex 100 mg., ascorbic acid 100 mg./Cap. Bot. 25s, 100s, 1000s
Use: Capillary abnormalities
- CEBIONE.** N.N.R. 1954. (Merck Sharp & Dohme) Ascorbic acid, crystalline Vit. C, cevitamic acid. Cryst. Bot. 1 Gm.
Use: Antiscorbutic.
- CECOBEE.** (Lloyd, Dabney & Westerfield) Vit. B-12 100 mcg., B-1 50 mg., C 50 mg., niacinamide 75 mg., B-6 5 mg., B-2 5 mg., panthelol 6 mg./cc. Vial 20 cc.
Use: I.M., I.V., deficiencies in B-complex vitamins & Vit. C.
- CECON DULCET TABLETS.** (Abbott) Ascorbic acid 50 mg. Bot. 100s.
Use: Vitamin-C therapy.
- CECON SOLUTION.** (Abbott) Ascorbic acid

10% in propylene glycol Each drop from enclosed dropper supplies 2.5 mg ascorbic acid; each cc. contains 100 mg Bot w/dropper 10 cc., 50 cc

Use: Vitamin C therapy

CE-DE-FLOR (CDF). (Hoack) Vit C 30 mg Vit D 100 u., calcium fluoride 3 mg /Tab Bot 100s, 1000s

Use: Orally In prevention of dental decay
CEDESRON (Smith, Miller & Patch) Vitamin B 12 w/intrinsic factor conc., 1 u., B 12 10 mcg., ferrous fumarate 400 mg., ascorbic acid 200 mg thiamine mononitrate 10 mg pyridoxine HCl 3 mg., folic acid 2 mg /Tab Bot 30s, 100s

Use: Anemias

CEDILANID (Sandoz) Crystalline lanatoside C, Tab (0.5 mg) Bot 100s, 500s, 1000s
Use: Orally, d gitalization

CEDILANID-D. (Sandoz) Deslanoside, U.S.P.; desacetylanatoside C, Amp (0.4 mg /2 cc & 0.8 mg /4 cc) Box 6s, 30s, 100s
Use: Same as lanatoside C.

CEEBIC. (Parson & Convey) Vits B 1 25 mg B 12 10 mcg., B 6 5 mg, C 300 mg., niacinamide 100 mg., cal pantothenate 10 mg., hesperidin 30 mg /Tab Bot 100s

Use: Treatment of B complex and Vit C deficiencies

CEEPYRIN CHLORIDE, NND 1962 (Merrell)
Cetyl pyridinium chloride

Aq Sol 1:1000 (soton c) Bot 1 pt., 1 gal

Celef Sol (Hydroalcohol & acetone sol)

for 1 ml, 1r & aq sol, Bot 30 cc

Conc. Sol (10%) Bot 160 cc, 1 gal

Vaginal Suppl (1:1000) Box 12s

See: Cepical Laseoges, Sol (Merrell)

Use: Antiseptic

W/Zeac peroxide & Col Peroxide See:

Peroguent, Oval (Merrell).

CITAMIN (Aegla-Franck) Ascorbic acid

choline & carotene Tab Bot 100s.

Use: Treatment of capillary fragility

CHIST. (Rocky Mtn) Phenylephrine HCl 100

mg., thiopropoxypridamine maleate, 2

mg., methoxypraline HCl 550 mg., pyril

amine maleate 850 mg., salicylamide 227.0

mg., acetophenone d 120 mg., caffeine

a total 300 mg., Vit C 200 mg /Cap Bot

100s.

Use: Cold allerges, maintain capillary

integrity

CHISTEA. (Organon) Thiopropoxypridamine

maleate 10 mg., Vit C 100 mg., ephrin

220 mg., in an enteric base of sod

bitarb 1570 mg., Cal hydroxide & stirc

acid Tab. Vit 100s also in lots of 1000s.

Use: 112 a.i.d. Antipyretic with analgesic

properties.

CELBE. (Cale) Sod carboxymethylcellulose 0.5 Gm., thiamine HCl 2 mg /Tab Bot 100s, 500s, 1000s

Use: Chronic constipation

CELBENIN (Bristol & Beecham Res Lab Eng) 6 (2,6 dimethoxybenzamide) penicillin monohydrate

See Staphicillin (Bristol)

Use: Penicillin resistant staphylococci

CELESTONE. (Schering) Betomethasone 0.6 mg /Tab Bot 30s, 100s

Use: Anti-inflammatory steroid

CELGINACE (Mead Johnson) Calcium and sodium alginates 750 mg diacetyl sodium sulfasuccinate 50 mg /Tab or 1sp Tab Bot 30s 60s Granules Bot 147 Gm & 294 Gm

Use: Constipation in low residue diets

CELLASE 1000 (Baxter and Wallerstein) Cellulose Cellulolytic enzyme See: Cellulase

CELLODEX, TABS (Hiss Pharm) Methylcellulose, phenobarb, d-amphetamine HCl Bot 1000s

Use: Antiobesity

CELLODRINE (Sutliff & Case) Methylcellulose 500 mg., d Amphetamine sulfate 2.5 mg., thiamine mononitrate 3 mg /Tab Bot 100s 1000s

Use: (1 a.c.) Anti-obesity therapy

CELLOORAN. (Tilden) Methylcellulose, sugar cerlose (glucose) & vanillin Bot 4 oz

Use: (1-3 tsp q.i.d.) Laxative

CELLOSE (Dumas Wilson) Methylcellulose 6 gr., yeast concentrate 2 gr /Tab Bot 100s 1000s

Use: (1-3 t.i.d.) Laxative

CELLOSIZ W/ PHS. (Carbide & Carbon) Hydroxy Ethyl Cellulose

CELLOTHYL, NND 1962 (Warner Chilcott) Methylcellulose gran Bot 25 Gm 100

Gm; 1 Tab (0.5 gr) Bot 100s, 500s, 5000s.

Use: (1-3 t.i.d. & h.s.) Chronic constipation.

CELLUDIX. (Bell Pharmacal) d Amphetamine sulf 5 mg., thyroid 1 gr., methylcellulose 5 gr., atropine sulf 1/360 gr., phenobarbital 1/4 gr /Tab Bot 100s, 1000s

Use: Antiobesity

CELLULASE. A true cellulase derived from *Aspergillus niger* See: Cellase 1000 (Baxter)

W/Mylose, ptilase calcium carbonate magnesium glycinate See: Zylase Tabs (Bryant)

W/Pepsin glutamic acid pancreatin as bile salt See: Kanulase Tab (Smith Dorsey)

CELLULOSE. W/Hexachlorophene See: ZeaTab (Srlife)

CELLULOS METHYL ETHER. See: Methyl Cellulose, Prop (Various Mfr).

© Contains name recognized by N.F., U.S.P. or A.M.A.

- * CELLULOSE, OXIDIZED, U.S.P., N.N.D.** 1962, A.D.R. Cellulosic acid.
See: Hemo-Pak Oxytel (Parke, Davis)
Oxytel (Poike, Davis)
- CELLULOSIC ACID, N.N.D. 1962, A.D.R.** See: Oxidized Cellulose
- CELLULOXYLIC ENZYME.** See: Cellulose
- W/Amylolytic enzyme, proteolytic enzyme, aluminum hydroxide gel** See: Holazyme, Tabs. (Holco)
- CELLULOXYLIC, W/Amylolytic, proteolytic.** See: Trienzyme, Tabs. (Testagar)
W/Enzyme, proteolytic enzyme. See: Gustazyme, Tabs. (Geriatric Pharmaceutical)
- W/Piperidolate HCl, amylolytic enzyme, proteolytic enzyme, pancreatin, taurocholic acid.** See: Doctilase, Tabs. (Lakeside)
- CEL-LUMA.** (Superior Pharmacol) Ploin sod carboxymethylcellulose 3½ gr., aluminum hydroxide gel susp. 1½ gr./Tab. Bot. 100s.
Use: (1-2 q. 3-4 h.) Peptic ulcer & gastric hyperacidity.
- CEL-LUMA W/Phenobarbital & atropine.** (Superior Pharmacol) Phenobarbital 1/16 gr., atropine sulfate 1/2000 gr., sod. carboxymethylcellulose 3½ gr., aluminum hydroxide gel 1½ gr./Tab. Bot. 100s.
Use: (1-2 q. 3-4 h.) Peptic ulcer.
- CELLUMETH.** (Chicago Pharm.) Methylcellulose 0.5 Gm./Tab. Bot. 100s, 1000s.
Use: (1-3 t.i.d. w/water) Bulk laxative.
- CELLUSENA.** (Bobs) Methylcellulose ½ Gm., cascara pow. 0.04 Gm., comp. senna pow. 0.162 Gm./Tab. Bot. 100s.
Use: (1-2 t.i.d. & h.s.) Laxative.
- CELMOL.** (Smith, Miller & Patch) Codeine phosphate ½ gr., sod. citrate 7½ gr., citric acid 7½ gr., aq. ext. of cocillona 4 min., Ext. of lobelia 1 min./fl. oz. Bot. 1 pt., 1 gal.
Use: (1 tsp. q. 3-4 h.) Sedative expectorant.
- CELONTIN, N.N.D. 1962** (Parke, Davis) Methsuximide. Kapsals 300 mg., Bot 100s.
Use: (1 to 4 daily) Petit mal and psychomotor attacks.
- CELRON.** (Moffet) Caps. Bot. 100s, 1000s.
Use: Hypochromic (iron deficiency) anemia
- CEL-U-SOL.** (Wintersmith Chemical) Sod. carboxymethylcellulose. Tab. (7½ gr.) Bot. 100s.
Use: (1-3 daily w/large glass of water) Functional constipation.
- CELTROPIN.** (Campbell) Hamatropine methylbromide 5 mg, methylcellulose 800 mg/dr. Bot. 4 oz.
Use: (1-2 dr. t.i.d.) Antispasmodic.
- CE-MI-LIN.** (Smith, Miller & Patch) Ascorbic acid 100 mg/Losenge Bot 100s, 500s, 1000s
- CENAC.** (Central) Hexachlorophene 0.25%, colloidal sulfur 8%, resorcinol 2%, isopropyl alcohol 30%. Bot. 2 oz.
Use: Acne.
- CENAGESIC.** (Central) Salicylamide 0.25 Gm., pyrilamine maleate 12 mg, phenylsiphine HCl 5 mg, acetophenetidin 0.12 Gm., dried aluminum hydroxide gel 0.12 Gm., caffeine anhydrous 15 mg./Tab. Bot. 100s, 500s, 1000s.
Use: Common colds.
- CENAGESIC IMPROVED.** (Central) Phenylsiphine HCl 5 mg, pyrilamine maleate 12 mg, salicylamide 250 mg, acetophenetidin 120 mg, ascorbic acid 30 mg., caffeine 15 mg./Tab. Bot. 100s, 500s, 1000s.
Use: Common cold symptoms.
- CENALENE.** (Central) Pentylene tetrazol 100 mg., thiamine HCl 1.67 mg, niacinamide 7.5 mg, cyanocobalamin 2.3 mcg., alcohol 15%/5 ml. Bot. pt. Also available in Tabs.
Use: Geriatrics.
- CINAMAL.** (Central) Salicylamide 250 mg., acetophenetidin 120 mg, rocephedrine HCl 5 mg, pyrilamine maleate 12 mg., ascorbic acid 50 mg, Dover's powder 24 mg./cap. Bot. 18s, 100s, 500s, 1000s.
Use: (1-2 q. 4 h.) Common cold
- CENASERT IMPROVED VAG. TABS.** (Central) 9-aminoacridine undecylenate 1 mg., N-methyl-3-hydroxybutylamine HCl 1 mg., methylbenzethonium chloride 18 mg., succinic acid 12.5 mg./Tab. Bot. 30s, 100s, w/inserts.
Use: Vaginitis
- CENDIX.** New name for Dadox (Central).
- CENDIXAL.** New name for Dadoxal (Central)
- CENIRON PLUS.** (Central) Ferrous sulfate dried 200 mg., vit. B-12 w/intrinsic factor conc. 0.5 u., folic acid 2 mg, ascorbic acid 125 mg, mang. sulfate monohydrate 12 mg, copper sulfate 3 mg, cobalt chloride 0.4 mg/tab. Bot. 100s, 500s, 1000s.
Use: Hematinic.
- CENOLATE.** (Abbott) Ascorbic acid. Amp. (100 mg./cc.) 1 cc., Box 7s, 25s; Amp. (500 mg./cc.) 1 cc., Box 6s, 25s; Amp. (1000 mg./cc.) 2 cc., Box 6s, 25s.
Use: Vitamin-C therapy.
- CENALEX.** (Central) Pentylene tetrazol 100 mg, B-1 1.67 mg, niacinamide 7.5 mg. Vit. B-12 2.5 mcg. alcohol 15%/5 cc. Bot. pt., gals.
Use: Treatment of senile patient with cerebral arteriosclerosis.
- CENTALONE.** (Paschall) Placental extract. Oral/injectable. Bot 100 caps. 2.5 mcg. Inj. 10 cc. multiple dose vials, 10 mcg. per cc

* Generic name recognized by N.F., U.S.P. or A.M.A.

Use Oral 2 to 4 caps daily Injection $\frac{1}{2}$ to 1 cc two or three times per week
Rheumatoid arthritis and sclerod collagen diseases dermatology and gynecology

CENTRAL NERVOUS SYSTEM DEPRESSANT,

N.R. See Sedatives

CENTRAL NERVOUS SYSTEM STIMULANTS,

See Amphetamine (Various Mfr)

d Amphetamine (Various Mfr)

Caffeine (Various Mfr)

Camphor

Cafron (Lakeside)

Coramine, Amp (Knoll)

Deoner (Riker)

Desoxyephedrine (Various Mfr)

Labeline Amp (Sandoz)

Lorfan (Roche)

Morsilid (Roche)

Meg mide (Abball)

Merotran (Merrell)

Methamphetamine (Various Mfr)

Metrozal Amp, Pow Tab (Knoll)

Nalline (Merck, Sharp & Dahme)

Nardil, Tab (Warner Lambert)

Nikethamide (Various Mfr)

Pentylentetrazol (Various Mfr)

Pimasin (Ciba)

Ritolin, Tab (Ciba)

Sodium Succinate (Various Mfr)

Seduxin (Brewer)

Talranil (Geigy)

CENTRINE, NND 1962 (Bristol) Amino

penlamide (alpha alpha diphenyl, gamma

dimethyl aminovaleramide sulfate) Tab

(0.5 mg) Bot 100s, Bot w/drapper (0.5

mg/10 drops/fl oz)

Use: Anticholinergic, antispasmodic agent

W/phenobarbital, reserpine See: Neuro

Centrine, Tab (Bristol)

CENTROBAR (Hiss) Butobarbital $\frac{1}{4}$ gr, as

pirin $3\frac{1}{2}$ gr, acetophenetidin $2\frac{1}{2}$ gr,

caffeine $\frac{1}{4}$ gr, buffered with calcium car

bonate/Tab Bot 100s, 1000s

Use: Sedative and analgesic

CEPACOL (Merrell) Cepyrin chloride (brand

of Cetylpyridinium Chloride) 14000 & phos

phate buffers in a pleasantly flavored solu

tion safe for oral use Bot 5 oz, 14 oz,

1 gal

Use: Antiseptic solution for oral hygiene

CEPACOL ANTIBIOTIC TROCHES (Merrell)

Cetylpyridinium 1:1500 neomycin sulfate

2.5 mg, tyrothricin 1 mg, methyparaben

4 mg, propylparaben 1 mg. Benzocaine

5 mg in a candy base/lozenge 16s foil

wrapped in 4s

Use: Minor throat and mouth irritations

CEPACOL THROAT LOZENGES (Merrell)

Cepyrin chloride (brand of Cetylpyridinium

* Generic name recognized by NF, U.S.P. or A.M.A.

Chloride) 1:1500, benzyl alcohol 0.3%

Box 24s

Use Antibacterial lozenges for sore throat

*CEPARSOLATE SODIUM (Abball) See

Sodium Thiacetphenorsamide

CEPNAELINE HYDROCHLORIDE W/Sod

benzoate, sanguinaria, squill syr, aromatics

See Sedotussin, Liq (Lilly)

CEPNAL (Herron) Tabs Bot 100s

*CEPNALOTININ SODIUM (Lilly) 7 (2 Thienyl

acetamide) cephalosporanic acid sod salt

CEPNEB (Superior Pharm) Acetaphenetidin

2 gr belladonna ext 1/50 gr, sod salt

cytate 2 gr, gum camphor $\frac{1}{2}$ gr, capsicum

1/10 gr, iodized Col $\frac{1}{2}$ gr/Tab Bot 100s

Also available w/Daver's Powder

Use (1 q 3 h) Analgesic with antispas-

modic

CERAPON, Triethanolamine Polypeptide

Oleate Condensate (Purdue Frederick)

See: Cerumenex (Purdue Frederick)

Kerrid (Blair Lab)

CERAZOLE (SUSPENSION), (Massengill)

Microcrystalline sulfathiazole 12 Gm in a

cherry flavored suspension/15 cc Bot 1 pl,

1 gal

Use Sulfo therapy

CERBARTROL (Hermelin) Red bone marrow

ext Amp 1 cc, 5 cc

Use: I.M., treatment of rheumatoid arthritis

CERELEXIN (Upjohn) Comp Vit B 1 0.36

mg, riboflavin 0.1 mg, nicotinic acid 5 mg,

Cal pantothenate 0.2 mg, Vit B 6 0.08

mg, iron 10 mg & yeast/Tab; Syrup Vit

B 1 0.9 mg, riboflavin 0.2 mg, yeast liver

& malt/lsp Bot 8 fl oz

Use Vitamin B complex therapy

CERELOSE, See Glucose (Various Mfr)

CERETOL A (Barre) Mephenesin 0.5 Gm

salicylamide 0.2 Gm/Tab Bot 100s, 1000s.

Use Analgesic muscular relaxant

CERETOL B (Barre) Mephobarbital 21.6 mg

mephenesin 0.5 Gm/Tab Bot 100s, 1000s

Use: Sedative—muscular relaxant

CER-O-CILLIN, NND 1962 (Upjohn) Hypo-

allergenic Penicillin O Sodium Vial (200

000 u) 1s (500 000 u) 1s

Use: Orally, I.M., penicillin therapy

CER-O-CILLIN-CHLOROPROCAINE, NNE

1955 See: Depa Cer O Cillin (Upjohn)

CER-O-CILLIN, DEPO, (Upjohn) Hypoaller

genic penicillin O Chloroprocaine peni

cillin O 300 000 u and 1,500,000 u per

vial

Use: I.M. Penicillin therapy to allergi

c patients

CEROFORT (White Labs)

Drops—L lysine HCl 450 mg, B 12 25

mcg., B-1 10 mg., B-6 5 mg./1.5 cc. Bot. 24 cc.

Elixir—L-lysine HCl 790 mg., vit. B-1 10 mg., B-12 25 mcg., B-2 10 mg., B-6 2 mg., niacinamide 100 mg., panthenol 20 mg./1/2 pt. Bot. 8 fl. oz., 1 gal.

Tab.—L-lysine HCl 790 mg., vit. A 25,000 u., D 1,000 u., B-1 10 mg., B-2 10 mg., B-6 2 mg., B-12 4 mcg., folic acid 1.5 mg., cal. pantothenate 20 mg., niacinamide 100 mg., ascorbic acid 300 mg./Tab. Bot. 60s 500s

Use: Dietary supplement.

CEROL (Wyeth) Whole wheat flour, yellow-corn flour, wheat germ, maltose syrup, dehydrated whole papaya fruit, nontart dry solids, primary dried yeast, tricalcium phosphate, iron pyrophosphate, thiamine, riboflavin and niacinamide. Bot. 8 oz.

Use: Nutritional supplement.

CERO-SAL (Kahlenberg) Salicylic acid, cerium salicylate, n-butyl parahydroxybenzoate, thymol & chlorobutanol. Oint. Jars 1/2 oz.

Use: Topically, fungicidal & anesthetic.

CEROSI, (Ives-Cameron) Codeine phosphate 1 gr., phenindamine tartrate 60 mg., phenylephrine HCl 30 mg., chloroform 1 min., ipecac flax., glycerin, pol. guaiacolsulfonate, sod. citrate, citric acid/oz. Bot. 1 pt.

Use: Orally, antihistaminic cough therapy.

CEROSE, PEDIATRIC, (Ives-Cameron) Oestramethorphan HBr 7.5 mg., phenindamine tartrate 5 mg., phenylephrine HCl 5 mg., chloroform, 0.25 minims, fl. oz. Ipecac, glycerin, pol. guaiacolsulfonate 86 mg., sod. citrate 3 gr., citric acid 1 gr. & alcohol/5 cc. Bot. 1 pt.

Use: Respiratory infections and cough due to colds.

CER-O-STREPS, (Upjohn) Dihydrastreptomycin 0.5 Gm., & streptomycin 0.5 Gm. present as the sulfates, chlorapracaïne penicillin O 300,000 u., penicillin O sod. 100,000 u./vial Also available in one-half strength.

Use: I.M., penicillin & streptomycin therapy.

CER-O-STREP-ONE, (Upjohn) Streptomycin 1 gm., 2 chlorapracaïne penicillin O 300,000 u., penicillin O sod. 100,000 u. per vial

Use: I.M. Antibiotic therapy.

CER-O-STREP-ONE-HALF, (Upjohn) Streptomycin 0.5 gm., 2-chlorapracaïne penicillin O 300,000 u., penicillin O sod 100,000 u. per vial.

Use: I.M. Antibiotic therapy.

CERTANE DOUCHE POWDER, (Vogarell)

Oxyquinoline sulfate, boric acid, sod. chloro-

ride & sod sulfo-diactyl succinate. Bot 4 oz., 12 oz.

Use: Vaginal cleanser.

CERTANE JELLY, (Vogarell) Phenylmercuric acetate 0.02%, w/oxyquinoline sulfate, boric acid, sod. sulfo-diactyl succinate Tube 2 1/2 oz.

Use: Contraceptive jelly.

CERUMINEX SOLUTION, (Purdue Frederick) Cerapan 10% in propylene glycol with chlorobutanol 0.5%. Dropper bot. 8 cc., 15 cc.

Use: Removal of cerumen (ear wax).

CERVICAL GLUCOSE INDICATOR, a plastic applicator stick tipped with cotton impregnated with two enzymes, glucose oxidase (30 units) and peroxidase (22.74 units), plus a suitable yellow dye (FD & C Yellow No 5, 56.5 mcg. and orthotolidine (1 mg.). See: Ovulest (Eli Lilly & Co.)

CERVILAXIN, N.D. 1961. (National Drugs) Purified Relaxin, Vials 2 cc 8x 1s, 6s.

Use: Short labor, easier delivery.

CEVIRATE, (Physicians) Ferrous fumarate 300 mg., Vit. C 150 mg./Tab. Ferrous fumarate 100 mg., Vit. C 50 mg./Pedi. Tab Bot. 100s, 1000s.

Use: Iron deficiency anemia

CERVO, (Sutliff & Case) Chloroform 2 min., pot. guaiacol tulfonate 8 gr., onilmyon & pot. tartrate 1/12 gr., ammonium chloride 8 gr., diluted hydrocyanic acid 3 min./fl. oz. Syr. 1 pt., 1 gal.

Use: Expectorant for simple coughs

CERVO W/DIHYDROCODEINONE BITARTRATE, (Sutliff & Case) Same formula as Cervo except for the addition of Dihydrocodeinone bitartrate 10 mg./fl. oz. Bot. 1 pt., 1 gal.

CERVO W/MORPHINE, (Sutliff & Case) Same formula as Cervo except for the addition of morphine sulfate 15 mg/fl. oz. Bot. 1 pt., 1 gal.

CETALKONIUM CHLORIDE

W/Neomycin sulfate, hydrocortisone, pyrilamine maleate, dipiperdan HCl. See: Fent-O Augine Ear Drops, Liq (Canfield)

W/Neomycin sulf., hydrocortisone, pyrilamine maleate, dipiperdan HCl, urea. See: Otolin, Drops (Louisiani)

W/Pyritamine maleate, hydrocortisone, neomycin sulfate, dipiperdan HCl, urea. See: Tego-Otic, Soln. (Ortega)

CETAPNOL, (Texas Pharm.) Cetyl alcohol, stearyl alcohol, propylene glycol, sod. lauryl sulfate ointment base. Jar 1 lb., 5 lb. Bot. 1 pt.

Use: Washable ointment base.

CETAPRED, (Alcon) Sulfacetamide sodium

* Generic name recognized by N.F., U.S.P. or A.M.A.

- 10%, prednisolone 0.25% in a methyl cellulose vehicle Bot 5 cc, 15 cc
 Use Inflammatory conditions of the eye
- CETAVLON.** See: Cetyltrimethyl Ammonium Bromide
- CETAZINE, TABLET, NNR 1955** (Bowman)
 Acet Dia Mer Sulfonamides 83 mg each sulfa/Tab Bot 100s
- CETHYLOSE.** (Ascher) Sod carboxymethyl cellulose Tab (0.5 Gm) Bot 100s, 1000s
 Use (1 2 1 d followed with water) Laxative
- CETHYTIN.** (Ascher) Sod carboxymethyl cellulose 0.5 Gm with diacetylhydroxyphenyl isofin 1 1/2 mg /tab Bot 100s, 1000s
 Use (1 2 1 d) Laxative
- CETRIL** (Menlo Park) Cetrimide cetyltrimethylammonium bromide ethoxylated lanolin alcohols lactic acid aerosol propellant Aerasol bot 6 oz
 Use Antiseptic, skin cleanser
- CETRIMIDE, W/Cetyltrimethylammonium bromide** See Cetril (Menlo Park)
- CETRO-CIROSE.** (Ives Cameron) Codeine phosphate 1/12 gr, with pot guaiaacal sulfonate, chloroform, ipecac flexil sod citrate, citric acid & alcohol/5 cc Bot 1 pt, 1 gal
 Use (1 tsp q 3 h) Narcotic cough preparation
- CETROFEN.** (Warren Teed) Morphine sulfate 1/24 gr, w/pot guaiaacal sulfonate, tartar emetic, ammonium chloride, citric acid, sod citrate and with or without sugar/5 cc Bot 3 oz, 1 pt, 1 gal
 Use: (1 tsp q 3 h m p) Narcotic cough preparation
- CETYBEN.** (Sutiff & Case) Tyrothricin 20 mg, benzocaine 50 mg, cetylalkaneam chloride 30 mg/Trache Bot 100s, 1000s
 Use: Antibiotic
- CETYLCIDE.** (Cetylite Ind) Cetyl dimethyl ethyl ammonium bromide 65%, Benzalkonium chloride 65%, isopropyl alcohol 13%, Sod nitrite 11%, inert ingredients 63% Bot 1 gal
 Use: Disinfectant for dental instruments
- CETYLDIMETHYL BENZYL AMMONIUM CHLORIDE W/Choline Salicylate** See Pharmacy Concentrate (Furdue Frederick)
- CETYLDIMETHYLETHYL AMMONIUM BROMIDE, A.D.R.** See: Radical Germicidal Solution (Novacol)
- W/Benzalkonium chloride, Isopropyl alc, sod nitrite** See: Cetylcide, Salm (Cetylite Inc)
- W/Isopropyl alcohol & inert ingredients** See: Sulficide Sol (Sutiff & Case)
- *CETYLDIMETHYL PYRIDINIUM CHLORIDE, USP,**
 * Generic name recognized by N.F., USP or A.M.A.
- INN 1962 Soln, U.S.P.** See: Ceopryl Chloride (Marrell)
- CETYL-STEARIC BASE.** (Upsher Smith) Glycerol monostearate, cetyl alcohol, stearic acid, triethanolamine & mineral oil Jar 14 oz, 5 lb.
 Use: Water soluble ointment base
- CETYLTRIMETHYL AMMONIUM BROMIDE.** (Bio Labs) Cetrimide B.P., Cetavlon, CTAB
 See: Sulficide (Sutiff & Case)
 Use: Antiseptic Surface active agent
- CEVALIN.** (Lilly) Ascorbic acid
 Amp: (500 mg) 1 cc, Pkg 6s, 25s, 100s; (500 mg) 5 cc, Pkg 6s, 25s, 100s; (1 Gm) 5 & 10 cc, Pkg 6s, 25s, 100s
 Drops: (30 mg/0.3 cc) Bot 15 & 50 cc
 Tab (25 50, 100, 250 & 500 mg) Bot 100s, 500s, 1000s
 Use: IM, IV, S.C. & orally, vitamin-C deficiency
- CEVANOID.** (Kirkman) Ascorbic acid 100 mg, hesperidin 100 mg/cap Bot 100s, 1000s
 Use: Capillary fragility
- CEVEX, NNR 1954** (Walker) Ascorbic acid 5 mg/drop Bot, w/dropper 15 cc Tabs 25 mg, 50 mg, 100 mg, 250 mg Bot 100 1000s 500 mg Bot 100s
 Use Vitamin C deficiencies
- CEVICETYL.** (Chicago Pharm) Aspirin 5 gr Vit C 125 mg/Tab Bot 100s
 Use: Analgesic
- CEVIPABA.** (Sly) Sod salicylate 250 mg, paminobenzoic acid 250 mg, Vit C 20 mg/Tab Bot 100s, 500s, 1000s
 Use Analgesic
- CEVIPAK** (Nion) Chlorprophenpyridamine maleate 2 mg, citrus bioflavonoids 25 mg Vit C 25 mg, salicylamide 100 mg, aceto phenethidin 60 mg, caffeine 15 mg/Tab Bot 24s
 Use Treatment of colds
- CEVIRATE PEDIATRIC.** (Physicians Prod) Ferrous fumarate 100 mg, ascorbic acid 50 mg/Tab Bot 100s
 Use Treatment of anemia
- CE-VI-SOL.** (Mead Johnson) Ascorbic acid 100 mg/cc Bot w/dropper 15 cc, 50 cc
 Use: Vitamin C deficiencies
- CEVITAMIC ACID, NNR 1945** See: Ascorbic acid
- CEVITAN.** See: Ascorbic acid
- CEYLON GELATIN** See: Agar
- CHAMO-CREME & POWDER.** (Dome) Cream Chamomile flower & aluminum acetate pH 5.5 Tube 1 oz, 2 oz, 4 oz & 1 lb Pow Chamomile flower Pkg 12s, 100s
 Use: Antiphlogistic & antipruritic
- CHARDONNA.** (Rorer) Phenobarbital 1/2 gr,

- belladonna ext. 1/12 gr., activ. charcoal 1 gr./Tab. Bot. 100s.
 Use: (I.L.D.) Antispasmodic & sedative
- CNASILETS.** (Chase Chemical) Pot. penicillin G 20,000 u., bacitracin 50 u./Trache. Pkg. 20s.
 Use: Vincent's Infection.
- CNEE-O-GENF** (Keene) See: Ethinyl Estradiol 0.02, 0.05 and 0.5 mg/Tab. Bot. 100s, 1000s.
- CNELAFRIN.** See: Epinephrine.
- CNELEN.** See: Ethyl Chloride.
- CHEL-IRON, H.N.D. 1962.** (Kinney) (Ferrocitrate N.N.D.) Iron choline citrate complex 0.33 Gm./Tab. Bot. 100s.
 Use: Iron deficiency anemias.
- CHEL-IRON PEDIATRIC DBOPS.** (Kiewey) (Ferrocitrate N.N.D.) Iron choline citrate complex 0.133 Gm., equiv. to 16 mg of elemental iron and 48 mg. of choline base/cc. Bot. 30 cc.
 Use: Iron deficiency anemias.
- CNEL-IRON PLUS,** (Kinney) Iron choline citrate (Ferrocitrate N.N.D.) 200 mg, vit B-12 with intrinsic factor concentrate 1/2 u, ascorbic acid 50 mg, folic acid 0.5 mg, thiamine mononitrate 2 mg, riboflavin 1 mg, pyridoxine HCl 2 mg., niacin 25 mg., calcium pantothenate 5 mg/Tab. Bot. 100s.
 Use: Iron deficiency anemias.
- CHEL IRON—112.** (Kinney) Chel Iron 417 mg., Vit. B-1 10 mcg, B-12 25 mcg/5 ml. Bot. 4 oz.
 Use: Hemolitic
- CHEMIPEN, H.N.O. 1962.** (Squibb) Pot Phenethicillin 125 or 250 mg./Tab. Bot 24s, 100s Syrup after reconstitution 125 mg./5 cc. Bot. 12 dose.
 Use: Infections amenable to therapy with oral penicillin.
- CNENATAL.** (Miller) Dicalcium phosphate 2,000 mg, magnesium ascorbate 100 mg, magnesium glycinate 200 mg, hesperidic complex 60 mg., A 5,000 I.U., D 400 I.U., B-1 3 mg., B-2 3 mg., B-6 5 mg., B-12 9 mcg., elaciamide 30 mg, pantothenate 5 mg., magnesium P-aminobenzate 10 mg., Cu 1 mg., I.O. 12 mg., Mn 2 mg., Zn 20 mg., Fe 30 mg., K 20 mg., lysine 25 mg, tacopherols (mixed) 10 mg, folic acid 0.3 mg./3 Tabs. Bot. 100s.
 Use Prenatal vitamin-mineral combination
- CNERACOL** (Upjohn) Codeine phosphate 1/6 gr., pot. guaicol sulfonate, chloroform, ammonium chloride, antimony & pot. tartrate, white pine & wild cherry/5 cc. Bot. 2 oz., 4 oz., 1 pt., 1 gal.
 Use: (1 tsp. q. 3 h.) Narcotic cough preparation.
- CNERALIN SYRUP,** (Lanett) Pot. guaicol sulfonate 8 gr., chloroform 2 min., ammonium chloride 8 gr., antimony and pot. tartrate 1/12 gr. In white pine & wild cherry/fl. oz. Bot. 1 pt. & 1 gal.; also with codeine 1 gr./oz.
 Use: (1 tsp. q. 3 hr.) Cough preparation.
- CNERA SULFA SUSPENSION.** (Sutliff & Case) Sulfadiazine, sulfamerazine 0.26 Gm. each/5 cc. Bot. 1 pt., 1 gal.
 Use: Orally, sulfonamide therapy.
- CNERA SULTABS.** (Sutliff & Case) Sulfadiazine 0.26 Gm., sulfamerazine 0.26 Gm./Tab., Bot. 100s, 1000s.
 Use: Orally, sulfonamide susceptible organisms.
- CNER-EOMINE,** (Pitman-Moore) Ethylenediamine dihydroiodide 4 gr., ammonium chloride 16 gr., squill fr, 20 min, artificial syr. of cherry q.s./fl. oz, Bot. 1 pt., 1 gal. Also available w/codeine or morphine.
 Use: (1 tsp. q. 3-4 h., p.r.n.) Expectorant.
- CNERITUSSAB,** (Bruce) Concentrate, 1 part is diluted with 4 parts of syrup or honey, it contains per 30 cc. phenylephrine HCl 30 mg., ammonium chloride 500 mg, potassium guaicol sulfonate 500 mg, sodium citrate 500 mg, saccharin 60 mg, chloroform 0.1 ml. Bot. pts., gals.
 Use: Expectorant.
- CNERBIN.** (Ascher) Vit. A 4000 u., D 800 u, B-1 1.5 mg., B-2 1 mg., B-6 0.5 mg, niacinamide 12 mg., ascorbic acid 40 mg, cal. pantothenate 2.5 mg, B-12 1 mcg/4 Gm. Granules Bot. 4 oz., 10 oz.
 Use: Orally, dietary supplement.
- CNEBROSOTE,** (Knox Drug) Alcohol 2%, chloroform 2 min., Col. creosote sulfonate 4 grs., Col. guaicol sulfonate 4 grs., wild cherry bark 30 grs., white pine bark 30 grs., spikenard root 4 grs., balm of gilead buds 4 gr., bloodroot 4 grs., tartor emetic 1/12 gr., benzyl alcohol 4 min., saffron bark 2 grs., ammonium chloride 8 grs./fl oz.
 Use: Cough due to colds.
- CNEST RUB.** (Jahesoe & Johnson) Camphor, menthol, eucalyptol/stick.
 Use: Coughs & colds.
- CNESTAMINE B & 12.** (Leeds) Chloropheniramine maleate 8 or 12 mg./Cap. Bot. 50s.
 Use: Allergic states responsive to anti-histamines.
- CNESTER-VIT.** (Leeds) Vit. A 5000 U.S.P. units, D 1000 U.S.P. units, B-1 1 mg., B-2 2 mg., C 50 mg., niacinamide 15 mg, calcium pantothenate 3 mg., B-6 2 mg., B-12 5 mcg, L-lysine monohydrochloride 15 mg/Cap. Bot. 30s.

- ***CHINIOFON**, NF, NNB, 1950 Tab, NF
Use Multivitamin for pediatric use
- CHEW-A-MINS** (Stayer) Vit C 60 mg
Vit B 12 2 mg, folic acid 0.25 mg Vit
B 1 2.5 mg, Vit B 2 2.5 mg, Vit B 6 1 mg,
niacinamide 20 mg, calcium pantothenate
5 mg, Vit A 5000 u, Vit D 1000 u, Vit E
1 International u / Tab Bot 72s
Use Vitamin supplement
- CHILDREN'S COUGH SYRUP**, (G F Harvey)
Ipecac flexl 2 min, wild cherry flexl 2
min, squil flexl 8 min, senega flexl 8
min, menthol 1/16 gr, citric acid 1 1/2 gr /
8 oz. Bot 1 pt, 1 gal.
- CHINESE GELATIN**. See Agar
- CHINESE ISINGLASS** See Agar
- 7 Iodo 8 hydroxyquinoline 5 sulfonic acid
sodium salt Yalen
Abbot—Tab (0.25 Gm) Bot 50s, 500s
Endo—Tab (0.25 Gm) Bot 50s, 100s
500s
Winthrop—Tab (4 gr) Bot 100s
See Anayodin, Pill & Pow (Sitchoff)
Quinoxyl, Tab (Burroughs Wellcome)
Use: Amebic dysentery
W/Soric acid, sod, lauryl sulfate lactose &
dextrose See Almetrifon, Pow & Tab
(Meyer)
- CHINOSOL**, (Jewell) 8 Hydroxyquinoline sul-
fate Tab 7 1/2 gr Vial 6s Tri Tab
(3/5 gr) Bot 50s, 110s
Use: (Dissolve 1 Tab in 1 pint of water)
Antiseptic sol
- ***CHLOPHEDIAHOL HCl** α -(2 dimethylamino
ethyl)- α chlorobenzhydrol HCl See: ULO,
Syr (Riker)
- W/Diphenylpyraline HCl phenylephrine HCl
glyceryl guaiacolate, alcohol See Ulaminic
(Riker)
- W/Diphenylpyraline HCl, phenylephrine HCl
glyceryl guaiacolate, acetaminophen See:
Ulogesic, Tabs (Riker)
- CHLORADONNA** (Ulmer) Chloral hydrate
250 mg, hyoscyamine 0.1 mg, atropine
0.02 mg, hyoscine 0.006 mg/cap Bot
100s, 500s
Use: (1 2 p r n) Sedative & antispasmodic.
- CHLORAGEL**, (Walker) Aluminum hydroxide
4 gr, mag trisilicate 4 gr chlorophyll
fractions 25 mg / Tab Bot 100s, 1000s
Use: (1 3 daily) Antacid
- CHLOBAHIST TABS & SPAHCAPS** (Domed)
Chlorpheniramine maleate 4 mg / Tab
Chlorpheniramine maleate 8 or 10 mg /
Cap Bot 100s
Use: Symptomatic relief of allergic mani-
festations
- * Generic same recognized by NF, U S P or A.M.A.
- ***CHLORAL BETAINE**, (Mead Johnson)
Chloral hydrate betaine adduct
Use: Hypnotic, sedative
- CHLORAL, CAMPHORATED**, W/Hyoscy-
amine oleate, camphor, menthol, etc See:
Calmital, Liq & Oint (Leeming)
- W/Thenylpyramine HCl, menthol hyoscy-
amine oleate See: Calthenamine Cream
(Leeming)
- ***CHLORAL HYDRATE**, U S P; Caps, U S P
Fellows—Cap (3 3/4 gr) Bot 24s, 100s
(7 1/2 gr) Bot 50s
G F Harvey—Cap (3 3/4 gr) Bot 100s;
(7 1/2 gr) Bot 100s
Massengill—Cap (7 3/4 gr) Bot 100s, 500s
West Ward—Cap (3 3/4 gr, 7 1/2 gr) Bot
100s
See: Aquachloral, Supp (Webster)
Fellows, Cap (Fellows)
Hydral, Cap. (Person & Covey).
Lorinal, Cap & Drops (Arnar Stone)
Lycoral, Liq (Fellows)
Noctec, Cap & Sol (Squibb)
Reclules, Supp (Fellows)
Somnos, Cap & Elixir, (Merck Sharp &
Dohme)
W/d Amphetamine sulfate & mephene-
sin See Somatol, Cap (Person & Covey).
W/Antipyrine See: Sominal, Tab (National
Drug)
W/Code oil, phenol & salicylic acid See
Corbunal, Oint (Blue Line)
W/Col bromide & atropine sulfate See
Fello-Sed, Liq (Fellows)
W/Ergotamine tartrate caffeine, scapola-
mine aminoaxide HBr See Reclafol M
grasne Dis Applicators (Wampole Labs)
W/Hyoscine HBr See Chloralixine, Elixir
(Chicago Pharm)
W/Hyoscyamine, atropine & hyoscine
See: Chloradonna, Cap (Ulmer)
Feforal, Cap & Liq (Fellows)
W/Hyoscyamus ext & pot bromide
See: Bromidia (Battle)
Bromide & Chloral Comp (Merck Sharp &
Dohme)
Hypno Bromic, Sol (Wampole)
W/Menthol & methyl salicylate See: Chloro-
Salicylate, Oint (Kremers Urban)
W/Mephene- sin & scopolamine HBr See
Somersed, Cap (Columbus)
W/Methylalicylcholine & bromocholine
bromide See: Hypotal, Tab (Anglo-French)
W/Methyl salicylate & menthol
See: Chloral Methylal, Oint (Ulmer).
Chloro-Salicylate Oint (Kremers Urban).
Car Al Gesic, Oint (Ulmer)
W/Morphine HCl See: Papine, Liq (Battle)

W/Phenol, camphor, olive oil. See: Otagine, Liq. (G. F. Harvey)

W/Phenyltoloxamine citrate (Kremers-Urban) See: Loryl

W/Pot. bromide & hyoscyamus exl. See: Brocloreum, Syr. (Mossingill)

Bromidia, Liq. (Balfie)

W/Pot. bromide, hyoscyamine HBr, horse-nettle, wild lettuce & celery seed. See: Norkogan, Liq. (Tilden).

CHLORAL, PENTAERYTHRITOL. See: Perichlor, Cap. (Ives-Cameron)

CHLORALIXINE. (Chicago Pharm.) Chloral hydrate 40 gr., hyoscyamine HBr 1/19 gr., alcohol 14% /fl. oz. Bol. 1 pl., 1 gal. Use: (1-2 dr. h.s. in water) Sedative.

CHLORAL-METHYLOL. (Ulmer) Chloral hydrate, methyl salicylate and menthol Tebe 1 oz.; Jor 1 lb. Use: Topically, anesthetic & analgesic.

CHLORALPYRINE. See: Dichloralpyrine.

CHLORALPYRINE DICHLORALPYRINE. See: Dichloralantipyrine.

W/Amobarb.

See: A.N.S., Tab. (Kenwood Lab.)

Saminal, Tab. (National Drug)

CHLORAMATE. (Hiss Pharm) Chlorpropen-pyridomine moleate, 12 mg. Unicelles Tab. Bot. 30s & 100s. Vial 10 mg. 30 cc. 100 mg./cc. 5 cc. Use: Symptomatic relief of allergic conditions

* **CHLORAMBUCIL.** N.N.D. 1962. Tabs., U.S.P. gamma-p-Di-(2-chloroethyl)amino-phenylbutyric acid. See: Leukeran, Tab (Burroughs Wellcome)

* **CHLORAMINE-T.** N.F. X, N.N.R. 1949. A.D.R. Sod. paratoluene-sulfan chloramide, chloramine & chlorazone. Lilly—Tab. (0.3 Gm), Bol. 100s, 1000s. Mallinckrodt—Pow., 1 lb. Use: Antiseptic, deodorant

See: Chlorazene, Prep. (Frost)

W/Pot. iodide. See: Heliogen (Penick)

CHLORAMIPHENE. (Meirell) Under study. Use: Investigation, orally control of conception.

* **CHLORAMPNICOL.** U.S.P., N.N.D. 1962. A.D.R.; Cap., Inj., Ophth. Oint., Sterile Susp., U.S.P. See: Chloramycetin, Prep (Parke, Davis) Chloramphenicol Palmitate, U.S.P.

W/Paramomycin, hydrocortisone acetate, in a liquid petrolatum base. See: Numecort Ointment (Parke, Davis)

* **CHLORAMPHENICOL PALMITATE.** U.S.P., N.N.D. 1962. See: Chloramycetin Palmitate, Pediatric (Parke, Davis)

* **CHLORAMPHENICOL PANTOTHENATE**

COMPLEX. Pluriquimica, LDA (Portugal) A complex consisting of 4 parts of chloramphenicol to one part of calcium pantothenate See: Pantofenical.

* **CHLORAMPHENICOL SODIUM SUCCINATE.** N.N.D. 1962, U.S.P.; Inj., U.S.P. See: Chloramycetin Succinate, Inj. (Parke, Davis)

CHLORANIL. 2,3,5,6-Tetrachloro-1,4-benzoquinone. See: Kerolac. (Salem Pharm.)

CHLOR-ANODYNE. (Parke, Davis) Morphine HCl 2½ gr., diluted hydrocyanic acid 9 min., alcohol 67%, chloroform 46 gr., peppermint oil 1½ min., capsicum tr. ¾ min./fl. oz. Bol. 4 oz., 1 pt., 1 gal. Use: To relieve colic, neuralgia & spasmodic pains.

* **CHLORAZANIL HCl.** 2-Amino-4-(p-chloroanilino)-s-triazine HCl. See: Daquin. (Riker)

CHLORAZINE. N.N.R. 1950; A.D.R. 1953. (Frost) Chloramine-T, N.F., sod. p-toluene-sulfanchloramide. Pow., Pkg. No. 1 & No. 2; Bol 1 lb., 5 lb. Aromatic Pow. (5%), Bol. 2 oz., 4 oz., 1 lb., 5 lb. Tab. (0.3 Gm), Bol. 20s, 100s, 300s, 1000s

Use: Topically; antiseptic, deodorant.

CHLORAZODIN. See: Azachloramid

CHLORAZONE. See: Chloramine-T.

* **CHLORBETAMIDE.** Dichloro-N-(2,4-dichlorobenzyl)-N(2-hydroxyethyl)-acetamide. Menthonide, Pantolin

CHLORBUTANOL (Chlorbutal). See: Chlorbutanol, U.S.P.

W/Triethanolamine polypeptide plate condensate. See: Cerulav Draps, Otic. (Harvey)

CHLORBUTOL. N.N.R. 1946, A.D.R. See: Chlorobutanol

CHLORCYCLIZINE-CALAMINE CREAM. See: Di-Paralene Calamine Cream. (Abbott)

* **CHLORCYCLIZINE HYDROCHLORIDE.** U.S.P., N.N.D. 1962; Tab U.S.P., 1-(p-Chlorobenzhydryl)-4-methylpiperazine hydrochloride. Histamin.

See: Diparalene, Prep. (Abbott)

Perosil, Prep. (Burroughs Wellcome)

Use: Relief in allergic disorders.

W/Pseudoephedrine HCl. See: Fedrozil, Tab (Burroughs Wellcome)

W/Pseudoephedrine HCl, acetophenetidin, aspirin, caffeine. See: Emprozil, Tab. (Burroughs Wellcome)

W/Tronothane HCl, calamine, zinc oxide, menthol. See: Tronalen, Lot. (Abbott)

CHLORDANE. See: Wydone, Liq. (Wyeth)

* **CHLORDANTOIN.** 5-(1-ethylamyl)-3-trichloromethylthiazidantoin. See: Sporostacin Cr. (Ortho)

* **CHLORDIMORINE HCl.** γ-morpholino-propyl 2-chloro-4-phenylphenyl ether HCl

* **CHLORDIAZEPOXIDE.** 7-chloro-2-methyl-

* Generic name recognized by N.F., U.S.P. or A.M.A.

amino 5 phenyl 3H k 4 benzodiazepine 4
oxide HCl Methaminodiazepine HCl See
Librium (Roche Lab)

W/Clidinium bromide See Librax (Roche
Lab)

CHLORESIUM MUCINOID, TABLETS.

(Rystan) Water soluble chlorophyll a plus
aluminum hydroxide & mag trisilicate in
dehydrated powdered akra Tab Bot 50s
200s Box 25 powders

Use Peptic ulcer & peptic hyperacidity

CHLORESIUM NASAL SOLUTION. (Rystan)

Chlorophyll Bot ½ oz, 2 oz 8 oz

Use Relief & deodorization in acute A
chronic respiratory infections

CHLORESIUM OINTMENT, NND 1962 (Ry
stan) Water soluble chlorophyll a (0.5%)

in specially prepared water soluble base
Tube 1 oz, 4 oz

Use: Promote tissue growth

CHLORESIUM SOLUTION, NND 1962

(Rystan) Water soluble chlorophyll a
(0.2%) in isotonic saline solution Bot 2
oz 8 oz

Use Cleansing of purulent wounds

CHLORESIUM AEROSOL. (Rystan) Water

soluble chlorophyll Sol drop Bot ½ oz

Use Deodorant

CHLOR-ETHAMINE. (Pitman Moore) Ethyl

enediamine dihydrochloride Tab (5 gr)

Bot 100s, 500s, 1000s

Use: Urinary acidifier

CHLORETHYL. See: Ethyl Chloride

CHLORETONE, NHR 1950 (Parke, Davis)

Chlorobutanol liq 1 oz

Use: Orally, SC sedative, antiseptic &
local anesthetic.

CHLORETONE INHALANT. (Parke, Davis)

Chlorelone 1 Gm, camphor 25 Gm, men
thal 18 Gm, cinnamon oil D04 Gm/100

Gm. in liq petrolatum Bot. 2 fl oz, 1 pt

Use: Antiseptic nose spray

CHLOR-EX SPECIAL. (Sutliff & Case) Chloro-

form 3 min, lobelia herb 7½ gr, tartar

emetic ½ gr, syr talu/fl oz Syr 1 pt,
1 gal

Use: Expectorant

CHLOR-EX SPECIAL W/MORPHINE. (Sutliff

& Case) Chloroform 3 min, lobelia herb

7½ gr, tartar emetic ½ gr, morphine sul

tate 16 mg, syr talu, color and aromatics/
fl oz Syr 1 pt, 1 gal

Use: Expectorant

CHLORQUANIDE HYDROCHLORIDE See:

Chloroquanide HCl. (Verlous Mir)

CHLORHEXADOL. 2 methyl-4 (2',2',2'-

trichlor 1 hydroxyethoxy)-2 pentanol See:

Lors (Wallace Labs)

CHLORHERIDINE OIHYDROCHLORIDE

* Generic name recognized by N.F., USP or A.M.A.

(Chlorhexide) W/Benzocaine See: Hibitant
Lozenges (Ayers)

CHLORNYDROL. (Rehais) Aluminum chlor

hydroxide complex, Al₂(OH)₅Cl

Use: An astringent that is mildly acid in
deodorant and antiperspirant preps

CHLORHYDROXYQUINOLIN See: Quinalor
Comp (Squibb)

**CHLORINATED AND IODIZED PEANUT
OIL.**

See Chloriodized Oil

Iodochloral, Sol (Searle)

CHLORINE COMPOUND, ANTISEPTIC. See

Antiseptics, Chlorine

***CHLORIODIZED OIL, USP, NND, 1962**

Chlorinated and iodized peanut oil See
Iodochloral Sol (Searle)

***CHLORISONDAMINE CHLORIDE, NND**

1962 4,5,6,7 Tetrachloro 2 (2 dimethylam-

monethyl) scandoline dimethylchloride See
Ecolad, Tab (Ciba)

***CHLORMADINONE ACETATE.** (Eli Lilly)

d⁶ 6 chloro 17 α -acetoxyprogesterone

Use: Progestational agent

***CHLORMERODRIH, NND 1962** 3 Chloro

mercurs 2 methoxypropylurea (Mercloran)

See: Neohydrin, Tab (Lakeside)

***CHLORMETNAZANONE.** 2 (4 Chloro-

phenyl)-3 methyl-4-metathiazanone 1 diax

ide See: Tranccopal, Tab (Winthrop Labs)

***CHLORMEZANONE, Chlormelthiazanone**

See: Tranccopal, Tab (Winthrop Labs)

2 (4 chlorophenyl) 3 methyl-4 metathiazan

one 1 dioxide

W/Acetylsalicylic acid

See: Tranccaprin, Tab (Winthrop Labs)

W/Aspirin See: Tranccaprin, Tabs (Winthrop

Labs)

***CHLOROAZODIN, NF X, NNR 1955;**

ADR; Sol, NF X ADR Alpha alpha, Aso

bis (chloroformamidino) See: Azachloramid

Prep (Wallace & Tiernan)

***CHLOROBUTANOL, USP, NNR 1950**

AOR Sedatam (Various Mir)

See: Chloralene, Prep (Parke, Davis)

Clortran, Cap (Wampole)

Use: Anesthetic, antiseptic & hypnotic

W/Myoscyamine sulfate, atropine sulfate &

hyoscyamine HBr See: Butadonna, Cap (Wam

pole)

W/Myristyl gamma picolinium chloride See

Wet Tone B, Saln (Braemmel)

W/Pyrimin B sulfate, neomycin, heparin

sodium, hydrocarisane, in polyethylene

glycol 400 base See: Bra Parin H, Vial

(Braemmel)

W/Pyrimine maleate, ephedrine hydro-

chloride See: Salahist, Nasal Spray (Vita

mix Corp)

- CHLORODYNE.** (Merck Sharp & Dohme) Morphine HCl 0.19 Gm., chloroform 2.6 cc., diluted hydrocyanic acid 0.55 cc., capsicum tr. 0.1 cc., peppermint oil D.T. cc., alcohol 52% / 30 cc. Sol. 1 pl.
Use: Anodyne.
- CHLOROETHANE.** See: Ethyl Chloride. *
- CHLOROFORM, U.S.P., A.D.R. 1955, H.F.** Lin., Spirit, Water. Sol. 2 oz., ¼ lb., 1 lb.
Use: Inhalation anesthetic.
- CHLOROFORM AND MORPHINE ANODYNE.**
See: Chlorodyne, Liq. (Merck Sharp & Dohme)
Chlor-Anodyne. (Parke, Davis)
- CHLOROGIENE.** (Purdue Frederick) Water-soluble chlorophyll. Duchettes (3 Gm.) Box 12s
Use: Acid douche solution for deodorization & cleanser.
- CHLOROGIENE COLPOYARS.** (Purdue Frederick) Concentrated water-soluble chlorophyll "a," lactose & excipients, buffered to an acid pH. Vaginal tab., Box 12s.
Use: Vaginal cleanser.
- CHLOROGUANIDE HYDROCHLORIDE, U.S.P., N.N.B.** (Various Mfr) (Proguanil HCl) 1-(p-chlorophenyl)-5-isopropylbiguanidine HCl.
Guanalol HCl. (Lilly) N.N.R.
Poludrine HCl. (Ayerst).
Use: Antimalarial.
- CHLOROHYDRATES.** Same as Hydrochloride.
- CHLORO-IODOHYDROXYQUINOLINE.**
See: Iodochlorhydroxyquin, U.S.P.
- 3-CHLORO-7-iodo-8-QUINOLINOL, U.S.P.** See: Iodochlorhydroxyquin, U.S.P.
- CHLORO-O-LAVE.** (Raymer) Chlorophyll, boric acid, zinc sulfate, zinc phenosulfonate, alum, salicylic acid, oxyquinoline sulfate, thymol, menthol. Pow., 3 oz., Pkg. 12s.
Use: Vaginal douche.
- CHLOROMETHAPYRILENE CITRATE, U.S.P., N.N.D. 1962.** See: Chlorothen, Citrate, U.S.P.
- p-CHLOROMETAXYLEHOL.**
See: Aveno Baby Powder (Faugera)
Dett, Liq. (Atlantic Sales Corp.)
- CHLOROMYCETIN, N.N.D. 1962** (Parke, Davis) D-(—)-theo-2-Dichloroacetamidato-1-p-nitrophenyl-1,3-propanediol. Chloramphenicol, U.S.P.
Cap.s (50 & 100 mg) Bot. 25s, 100s.
Cream: (1%) Tube 1 oz.
I.M.s (1 Gm.) Vial.
I.V.: (0.5 Gm.) w/amp. 2 cc. of 50% sol. of N,N-dimethylacetamide, Pkg 10s.
Kapsals: (250 mg) Bot. 16s, 100s.
Ophth. Oint.s (1%) Tube ½ oz.
- Otic Drops:** (0.5%) Sol. 15 cc. (Propylene glycol)
Ophth. Sol.s (25 mg) Bot. w/dropper 15 cc. (dry)
Painfuls, Pediatric (125 mg / 4 cc.) Vial 60 cc.
Sensitivity Disks (10, 30 & 60 mcg) Bot 25s.
Sols (0.5 Gm.) Amp. (2 cc.) Box 6s.
W/Dihydrostreptomycin sulfate. See: Chlorostrep, Kapsol. (Parke, Davis)
W/Hydrocortisone acetate, polymyxin-B sulfate. See: Ophthocort Ophthalmic. (Parke, Davis)
- CHLOROMYCETIN-HYDROCORTISONE OPHTHALMIC SUSP.** (Parke, Davis) Chloromycetin 12.5 mg., hydrocortisone acetate 25 mg., borate buffer, phenol. Vial 5 cc. Susp. (dry) stable at room temperature for 10 days
Use: (First 2 days 2 gtt. q. 1-3 h., then 2 gtt. q. 3 or 4 h.) Inflammatory infections.
- CHLOROMYCETIN SUCCINATE, N.N.D. 1962.** (Parke, Davis) Chloramphenicol sodium succinate equal to 1 Gm Chloromycetin. Steril-vial 1 Gm.
* **CHLOROPHENOTHALE, U.S.P., N.N.R. 1957.** (Geiarol, Neacid, Dicophone) 1,1,1-Trichloro-2,2-bis(p-chlorophenyl) ethane.
See: DDT.
W/Benzocaine & benzyl benzoate.
See: Bensamol-D, Susp. (Irwin-Nelsler)
Enbin, Emul. (Wm. Cooper)
Paracin, Liq. & Oint. (Eaton)
Pucicide, Liq. (Warren-Teed)
Topocida, Susp. (Lilly)
W/Ethylaminobenzoate & benzyl benzoate, N.N.R. 1955.
See: Enbin, Susp. (Cooper & Nephew)
Paracin, Susp. (Eaton).
- CHLOROPHYLL "a" OINTMENT.** See: Chloresium Oint. (Rystan).
- CHLOROPHYLL "a" SOLUTION (Chlorophyllin).** See: Chloresium Sol. (Rystan).
W/Aluminum hydroxide & mag. trisilicate. See: Chloragel, Tab. (Walker).
W/Aluminum hydroxide, mag. trisilicate & okra pow. See: Chloresium Mucinoïd (Rystan).
W/Benzocaine & urea.
See: C-B-U, Oint. (Warren-Teed).
Ureacaine, Oint. (Chicago Pharm.).
W/Kaolin. See: Kaophyl, Cap. (Warren-Teed).
W/Oxyquinoline benzoate. See: Chlorothena, Oint. (Amar-Stone).
W/Phenobarbital, belladonna ext., benzoic acid, Cal. carb., mag. trisilicate & aluminum hydroxide. See: W-T Tab. (Warren-Teed).

* Generic name recognized by N.F., U.S.P. or A.M.A.

W/Propionate sod See: Praphyllin, Oint (Rystan)

W/Triple sulfa vaginal suppositories See: Chloro Sul, Supp (Columbus)

CHLOROPHYLL, OIL-SOLUBLE, W/Sulfa
See: Chloro Sul Vag Supp (Columbus Pharm)

CHLOROPHYLL OINTMENT (Warren Teed)
Chlorophyll ext water soluble 1% Tube 1 oz; Jar 1 lb

Use Topically, deodorize & promote healing

CHLOROPHYLL TABLETS

See Deadophyll, Tab (Wallace)

Nulla, Tab (Depre)

Olodex, Tab (Walker)

***CHLOROPHYLL, WATER-SOLUBLE, NND 1962** (Various Mfr) Chlorophyllin See See Chloresium Prep (Rystan)
Chloresium (Rystan)

CHLOROPHYLLIN,

Use Healing agent & deodorizer

Chlorogene, Duchettes (Purdue Frederick)

Chlorophyll

W/Tyrosine, carbamide & benzocaine See Tyrosyl Troches (U S Standard Prods)

***CHLOROPREDNISONE ACETATE, (Lilly)**

60 chloroprednisone 21 acetate

Use Topical corticoid compound

***CHLOROPROCAINE HCl, NND 1962**
beta, Diethylaminoethyl 2 chloro-4 amino benzoate HCl See Nesocaine Vial (Malt bis)

W/Epinephrine See Versocaine, Sol (Lederle)

***CHLOROPROCAINE PENICILLIN O, NND 1962** See Depo Cer O Cillin, Vial (Upjohn)

W/Penicillin O Sod, dihydrostreptomycin sulfate streptomycin sulfate See Cer O Streps, Vial (Upjohn)

CHLOROQUINE W/Milribis See: Milribis

W/Aralan Diphosphate, Tab (Winthrop Labs)

W/Quinacrine hydroxychloroquine See Triquin, Tab (Winthrop Labs)

***CHLOROQUINE PHOSPHATE, USP NND 1962**, Tab USP (7 Chloro 4(Di ethylamino-1 methylbutylamino) quinoline d phosphate Nivaquine See Aralan Diphosphate, Tab (Winthrop Labs)

Use: Febrile attacks of malaria

W/Aralan Diphosphate Tab (Winthrop Labs)

W/Glycobiarsol See: Mit bis

W/Salicylamide acetic acid See: Diphadyna Tabs (H 31)

5-CHLORO-SALICYLANILIDE W/Salicylanilide, 3,3' and 5,4 Dichlorosalicylanilide, undecylenic acid & zinc undecylenate See:

* Generic name recognized by NF, USP or A.M.A.

Salundek New, Oint. (Wallace & Tiernon)

CHLORO-SALICYLATE OINTMENT. (Kremers Urban) Chloral hydrate, menthal, methyl salicylate Tube 1/2 oz, 1 lb; Jars 1 lb, 5 lb

Use Topically, analgesic & counterirritant

CHLOROSTREP. (Parke, Davis) Chloromycetin 125 mg, dihydrostreptomycin sulfate 125 mg / 4 cc or cap Bot 12s Susp Bot 60 cc
Use Diarrhea and related intestinal infections

CHLORO-SUL. (Columbus) Oil soluble chlorophyll 15.6 mg, sulfadiazine 130 mg, sulfa methazine 130 mg, sulfa acetamide 130 mg, lactic acid 6.5 mg, methyl parahydroxybenzoate 6.5/Vaginal supp Box 12s
Use: Non specific infections of vagina & cervix

CHLOROTHELIA. (Arnor Stone) Water soluble chlorophyll w/oxiquinoline benzoate Sol 1 oz, 2 oz, 8 oz; Oint Tube 1 oz; Jar 1 lb; Bot 2 oz

Use Infections & slow healing wounds

CHLOROTHELIUM (Arnor Stone) Oxiquinoline benzoate & chlorophyll Oint Jar 1 lb
Use: To promote healing & antiseptic

***CHLOROTHEN 2** [(5 Chloro 2 thenyl)(2 di methylaminoethylamino) pyridine hydrochloride & citrate salts (Chlorothienylpyromine chloromethopyriline Pyrithen)
Use: Symptomatic relief in allergic disorders

***CHLOROTHEN CITRATE, NF, NND 1962** Tab (Whittier) NN Dimethyl N-(2 pyridyl) N (5 chloro 2 thenyl) ethylene diamine citrate Chloromethopyriline Chlorothienylpyromine pyrithen, Bot 100s
See: Togalhen, Tab (Lederle)
Use: (Oral 25 to 50 mg) Antihistaminic
In relief of allergic disorders

W/Acetophenetidin & caffeine
See: Fenoxodyne Compound, Tab (Breon)

W/Achromycin HCl, salicylamide, phenacetin, & caffeine See: Achromycin, Tab (Lederle)

W/Pyrimamine, thenylpyromine See: Dermo Fox, Liq (Recsei)

CHLOROTHENYL PYRAMINE. See: Chlorothen, Prep (Whittier)

CHLOROTHEOPHYLLINATE W/BENADRYL.
See: Dramamina, Tab (Searle)

***CHLOROTHIAZIDE, USP, NND 1962**
See: Diuril, Tab (Merck Sharp & Dahme)
W/Deserpidine See Oreticyl and Oreticyl Forte Tab (Abbott)

***CHLOROTHIAZIDE SODIUM, NND 1962**
See: Diuril, Lyovac (Merck Sharp & Dahme)
Dupres (Merck Sharp & Dahme)

- * **CHLOROTRIAMISENE**, H.F., N.H.D. 1962.
See: Tace, Cop. (Merrell)
W/Ergonovine maleate. See: Tace w/Ergonovine, Cop. (Merrell)
- * **CHLOROXYLEHOL**. See: Sepliderm. Preps (Aveeno)
W/Propyl-p-aminobenzoate, urea & benzyl alcohol.
See: Cenothetin, Oint. (Central)
Dunoguent, Oint. (Ascher)
- CHLORPACTIN**. (Guardian Chem.) Straight chain hydrocarbon with hypochlorous acid
See: Chlorpactin-WCS 90.
- * **CHLORPHEAMINE** w/phenylephrine HCl, neomycin, polymyxin B sulfate, benzocaine.
See: Rhioic, Loz. (Lemmon Pharm.)
W/Phenylephrine HCl, phenylpropanolamine HCl, phenyltoloxamine citrate. See: Holdcon. Tab. & Syr. (Bristol).
W/Phenylephrine, analgesics, and Vitamin C. See: Corlco. (Hoog).
- * **CHLORPHEHESIN CARRAMATE**. (Upjohn)
3-p-chlorophenoxy-2-hydroxypropyl carbamate.
- * **CHLORPHENIRAMINE MALEATE**, U.S.P., H.H.D. 1962. [n.], Syr., Tabs U.S.P. 2 [p-Chloro- α -pho(2-dimethylaminoethyl)benzyl] pyridine maleate.
See: Alerin-Td, Cops (Finley)
Chetamine, Tab (Leads)
Chloramate, Tab. (Hiss Pharm.)
Chlor-Trimeton Maleate, Prep. (Schering)
Histachlor, Tab. (Vitaminix Corp.)
Histirin, Tab. (Vitarine)
H-Sladur, Tabs. (Wynn)
Nascosed (Chicago)
Phenetron Injectable. (Lonnelt)
Pirlton, Tab. (4 mg) (West-Ward)
Poloramine, Tab. (Schering)
Teldrin Sponules. (Smith, Kline & French)
- d-CHLORPHEHIRAMINE MALEATE**. See: Poloramine (Schering)
- * **CHLORPHEHIRAMINE TANNATE**. W/Carbetapentane tannate, ephedrine tannate, phenylephrine tannate. See: Rynatess Tablets. (Irwin, Neisler)
W/Pyrimolamine tannate, d-amphetamine tannate. See: Nalerton, Tab. (Neisler)
W/Phenylephrine tannate, carbetapentane tannate, ephedrine tannate. See: Rynatess, Tabule (Irwin, Neisler)
W/Phenylephrine tannate, ephedrine tannate, carbetapentane tannate, pyrimolamine tannate. See: Rynatan Expectorant, Liq. (Irwin, Neisler)
- * **CHLORPHENOXAMINE HCl**, H.H.D. 1962.
beto-Dimethylaminoethyl [p-chloro- α -methyl-benzhydryl] ether HCl. See: Phenoxane (Pitman-Moore)
- * **CHLORPHEHTERMINE HCl**. (Worner-Lambert Res. Inst.) p-Chloro- α - α -dimethylphenethylamine HCl. See: Pre-Sole.
Use: Anorexic.
- CHLORPHTHALIDONE**. 1- α -3-(3'-sulfonyl-4'-chlorophenyl)-3-hydroxy-isaindoline. See: Hygraton (Geigy)
- * **CHLORPROMAZINE HCl**, U.S.P., H.H.D. 1962. Inj., Syr., Tab. U.S.P. 2-Chloro-10-(3-dimethylamino-propyl) phenothiazine. See: Thorazine, Amp., Supp., Syr., Tab. (Smith, Kline & French)
- * **CHLORPROPAMIDE**, N.H.D. 1962 (Pfizer)
N-Propyl-N'-(p-chlorobenzenesulfanyl) urea
See: Diabinese, Tab. (Pfizer)
Use: Hypoglycemic agent.
- CHLORPROPHEPYRIDAMINE MALEATE**, See: Chlorpheniramine Maleate, U.S.P.
- * **CHLORPROTHIXENE HCl**. (Hoffmann-La Roche) 2-chloro-9-(3-dimethylamino-propylidene)-thioxanthene HCl. See: Toractan.
- * **CHLOROQUINALDOL**, U.S.P., H.H.D. 1962. Oint., U.S.P. 5,7-Dichloro-8-hydroxyquinoline. See: Starasol, Oint. (Geigy)
- * **CHLORQUINOL**. A mixture of the chlorinated products of 8-hydroxyquinoline containing about 65 per cent of 5,7-dichloro-8-hydroxyquinoline. Ouxalin.
- * **CHLORTETRACYCLINE HCl**, N.F., H.H.D. 1962, A.D.R. Inj., Ophth., Cops., H.F. See: Aureomycin, Prep. (Lederle)
- * **CHLORTETRACYCLINE CALCIUM**, H.H.D. 1962. See: Aureomycin, Drops, Syr. (Lederle)
- * **CHLORTHALIDONT**. 2-chloro-5-[1-hydroxy-3- α -1-isindolyl] benzenesulfonamide
See: Hygraton, Tab (Geigy)
- CHLOR-TRIMETON COMPOUND**. (Schering)
[See: Demazin] Chlorprophepyridamine maleate 1.25 mg., phenylephrine HCl 2.5 mg/5 cc Syr. 1 pt. Repeatals Demazin Chlor-Trimeton maleate 4 mg, phenylephrine 20 mg Rot. 100s.
Use: Antihistaminic.
- CHLOR-TRIMETON CREAM**. (Schering) Tubes (1%) 50 Gm.
Use: Topically, antihistaminic to relieve allergic disorders.
- CHLOR-TRIMETON GLUCONATE** W/Metihortelone acetate See: Melretan, Nasal Spray (Schering)
- W/Phenylephrine HCl, gramicidin. See: Coricidin Nasal Mist, Saln. (Schering)
- CHLOR-TRIMETON MALEATE**, N.H.D. 1962. (Schering) Chlorprophepyridamine maleate. (Chlorpheniramine maleate) Amp. 10 mg./cc. 1 cc. 100 mg./cc. 2 cc) Box 6n Syr. (2 mg/4 cc.) Bar. 1 pt., 1 gal; Tab.

* Generic name recognized by N.F., U.S.P. or A.M.A.

(4 mg) Bot 100s, 1000s, Tab, repeat action (8, 12 mg) Bot 100s, 1000s

Use: {Oral, I.M. 2 to 4 mg} Antihistamine therapy
See Yeldrin, Spansules (Smith, Kline & French)

W/Aspirin, phenacetin, caffeine, phenyl ephrine See Coricidin D' Decongestant Tablets (Schering)

W/Cortisone acetate See Corticloran, Susp (Schering)

W/Aspirin & acetophenetidin See Coricidin Pediatric Medlets (Schering)

W/APC See Coricidin, Tab (Schering)

W/APC & penicillin G procaine See Coricidin w/Penicillin, Tab (Schering)

W/Dihydrocodeinane bitartrate, sod salt cyclate, sod citrate, caffeine, glyceryl guaiacolate See Coricidin, Syr (Schering)

W/Melicorin, ascorbic acid See Melieton Tab (Schering)

W/Salicylamide, phenacetin, caffeine, ascorbic acid, & methamphetamine HCl See Coricidin Forte, Cap (Schering)

CHLOR-TRIMETON RA, (Schering) Chlor Trimeton 8 mg, sod pentobarbital 50 mg / repeat action tab Bot 100s

Use: (1 q 8 h) antihistamine with sedative
CHLORYLIN (Schering) Trichlorethylene USP Bot 1 fl oz

Use: Analgesic Inhalant
CHLOR-X, (Mallard) Aluminum hydroxide 2 gr, magnesium trisilicate 4 gr, magnesium oxide 3 gr / Tab Bot 100s, 1000s
Pow 3 1/2 oz
Use: Antacid

***CHLORZOXAZONE**, NND 1962 See Paroflex, Tab (McNeil)

W/Acetaminophen, codeine phosphate See Paroflex w/Codeine Tab (McNeil)

CHOBELINE, (Reed & Cornrick) Choline chloride 2 Gm, thiamine HCl 167 mg, riboflavin 6.67 mg, niacinamide 333 mg / 30 cc. Bot 1 pt, 1 gal
Use: (1-4 tsp daily) lipotropic & anti-hepatotoxic

CHOBILE, (Irwin, Neisler) Cholic acid (conjugated as sod glycocholate & sod. taurocholate) 1 1/2 gr, ketocholonic acids 1 1/2 gr / Tab Bot 100s, 500s, 1000s
Use: (1-3 daily) Stimulates bile secretion

CHOBILE-PAH, (Irwin, Neisler) Sod salts of ox bile acids equivalent to 1 1/2 gr of cholic acid with ketocholonic acids 1 1/2 gr & pancreas deffatted, dried 1 1/2 gr / Tab Bot 100s, 500s, 1000s
Use: (1-3 daily) Stimulation of bile secretion

CHOCINATE (Sull & Case) Sod glycocholate 1 1/2 gr, sod salicylate 1 1/2 gr,
* Generic name recognized by N.F., U.S.P. or A.M.A.

sod succinate 1 1/2 gr / Tab Bot 100s, 1000s

CHOLACOLE, (Cole) Ox bile ext 5 gr / Cap Bot 100s

Use: (2 tsd w/water) Chalogogue & choleric

***CHOLACRYLAMINE RESIN**, (Merck) An anion exchange resin consisting of a water soluble polymer having a molecular weight equivalent between 350 and 360 in which aliphatic quaternary amine groups are attached to an acrylic backbone by ester linkages

CHOLAD, (Mallard) Dehydrocholic acid 1 1/2 gr, desoxycholic acid 1 1/2 gr, adiphenn HCl 10 mg / Tab Bot 100s, 1000s
Use To stimulate bile flow

CHOLADONNA, (Lemmon) Dehydrocholic acid 60 mg, ox bile ext 0.12 Gm, belladonna powd ext 4 mg, alain 15 mg / Cap Bot 100s
Use: Impairment of bile secretion

CHOLA-K, (Buffington) Bile salts 5 gr, menadiene 1 mg / Tab Bot 100s, 1000s
Use (1-3 daily) Bile salt therapy plus vita msn K

CHOLAN-DH, (Maltbie) Dehydrocholic acid 250 mg / Tab Bot 100s, 500s, 1000s
Use: (1 tsd, p.c) Stimulate bile flow

CHOLAN-HMB, (Maltbie) Phenobarbital 8 mg, dehydrocholic acid 250 mg, homotropine methylbromide 2.5 mg / Tab Bot 100s, 500s, 1000s
Uses (1-2 tid, p.c) Nonobstructive biliary disease

CHOLAN V, (Maltbie) Dehydrocholic acid 250 mg, homotropine methylbromide 5 mg / Tab Bot 100s, 500s, 1000s
Use: Choleric antispasmodic

CHOLANIC ACID, Dehydrodesoxycholic acid

CHOLANOR, (Maltbie) Ketocholonic acid 1 1/2 gr, ox bile, dehy 1 1/2 gr / Tab Bot 100s, 500s, 1000s
Use: (1-2 tid, p.c) Increases bile flow

CHOLARACE, (Warner Chilcott) In the coating Rocphedrine HCl 20 mg, pentobarbital 27.5 mg in the core, Choline theophyllinate 200 mg / Tab Bot 100s, 500s
Use: (1 tab q 3-4 hrs) Symptomatic relief of acute bronchospasm

CHOLASE, (Cutter) Cholinesterase
Use: I.V., dermatologic conditions

CHOLIATHOL, (Nysco) Choline bitartrate 200 mg, inasitol 100 mg, dl methionine 100 mg / Cap. Bot 100s
Use: Lipotropic agent

CHOLATOL, (Upjohn) Bile salts, with sod

toerocholate & sod. glycocholate. Cop., Bot. 40s, 500s, 1000s.

Use: (1-2 t.i.d., p.c.) Stimulate bile flow.

CHOLATOL COMPOUND. (Upjohn) Bile salts 2 gr., phenolphthalein $\frac{1}{2}$ gr., papain 1 gr., ginger 1/16 gr./Tab. Bot. 40s, 500s, 1000s.

Use: (1-2 t.i.d., p.c.) Stimulate bile flow.

CHOLATROPIN. (Lakeside) Dehydrocholic acid 4 gr., homotropine methylbromide 2 $\frac{1}{2}$ mg./Tab. Bot. 100s, 500s.

Use: (3-6 daily) Increase bile flow.

CHOLAVITE. (Ba Labs.) Dehydrocholic acid 100 mg., homotropine methyl bromide 1.5 mg., thiamine mononitrate 2.5 mg., riboflavin 2.5 mg., pyridoxine HCl 2.5 mg., niacinamide 10 mg., calcium pantothenate 2 mg., Vit. B-12 Intrinsic factor $\frac{1}{4}$ unit./Tab. Bot. 100s, 1000s.

Use: Diuretic with vitamin-mineral supplementation.

CHOLAXOL. (Smith, Miller & Patch) Phenolphthalein $\frac{1}{2}$ gr., sod. salicylate 1 gr., ox bile est. 1 gr., cascara ext. $\frac{1}{2}$ gr., menthol 1/20 gr./Tab. Bot. 1000s.

Use: (1-2 t.i.d.) Laxative with sedative.

CHOLA-ZEM. (Zemmer) Atropine sulfate 1/500 gr., sod. glycocholate 1 gr., sod. taurocholate 1 gr., nitroglycerin 1/500 gr., meg. sulfate 5 gr./Tab. Bot. 100s, 1000s.

Use: (1 t.i.d.) Cholagogue & chologogue with antispasmodic.

CHOLECYSTOGRAPHY AGENTS.

See: Chloridized Oil, (Searle).

Iodized Oil. (Various Mfrs.)

Iodoalphonic Acid. (Schering).

Iodobrassid. (Ciba).

Iodochlorol. (Searle).

Iodopanoic Acid. (Winthrop Labs.)

Iopahendylate Inj. (La Payette).

Lipiodine. (Ciba).

Lipiodal. (Fougero).

Lipiodal Radiologique Ascendant. (Fougero).

Monaphen. (National Synthetics).

Pontopaque. (La Payette).

Priodox. (Schering).

Telopaque. (Winthrop Labs.)

CHOLEDYL. N.N.D. 1962. (Worner-Chilton)

Choline theophyllinate. Tab. (100 mg. & 200 mg) Bot. 100s, 500s.

Use: (Children, 100 mg. t.i.d.; adults 200 mg. q.d.) Theophylline therapy.

CHOLELITH. (Parke, Davis) Sod. salicylate $\frac{1}{2}$ gr., sod. oleate $\frac{1}{2}$ gr., phenolphthalein $\frac{1}{2}$ gr./Pili. Bot. 100s, 500s, 1000s.

Use: (1-2 t.i.d.) Analgesic & antipyretic with laxative.

CHOLIOI LIQUID. (Smith, Miller & Patch) Ferric chloride 6 gr., ox bile ext. 3 gr., columbo 6 gr., chamomile flowers 13 gr.,

* Generic name recognized by N.F., U.S.P. or A.M.A.

rhubarb 13 gr., aromatics q.s., alcohol 25%./fl. oz. Bot. 4 oz., 1 pt., 1 gal.

CHOLETHALATES. (Elder) Phenolphthalein $\frac{1}{2}$ gr., hog bile ext. 1 gr., sod. salicylate 1 gr., eod. oleate 1 gr./Tab. Bot. 100s, 500s, 1000s.

Use: Laxative

* **CHOLERA VACCINE, U.S.P., N.N.B. 1952.**

Lederle—India Strolins, Vials 1.5 cc., 20 cc. Lilly—Vials 1.5 cc., 20 cc.

Wyeth—Vials 20 cc.

CHOLERETIC.

See: Bile Preps. and Forms.

Dehydrocholic Acid.

Desoxycholic Acid.

CHOLESTERIN. See: Cholesterol, U.S.P.

CHOLESTERIMIZED ANTIGEN. (Parke, Davis). 5 cc. Vial.

Use: Test in diagnosis of suspected syphilitic sera.

* **CHOLESTEROL, U.S.P.** (Various Mfr.) 5,6-Cholesten-3-ol (Cholesterol).

* **CHOLESTYRAMINE RESIN.** (Merck) A strongly basic anion exchange resin containing quaternary ammonium functional groups which are attached to a styrene-divinylbenzene copolymer (about 2% divinylbenzene)

CHOLEX SYRUP LIQUID. (Smith, Miller & Patch) Choline dihydrogen citrate 1 Gm./5 cc. Bot. 1 pt., 1 gal.

Use: Liver disorders

CHOLEX. (Bell-Craig) Meal of egg yolk, lecithin and glycerine for emptying of the gallbladder. Bot. 30 cc.

Use: Routine cholecystography

CHOLIC ACID. 3,7,12-Trihydroxycholanolic acid. (Cholic acid).

W/Dioctyl sod. seifosuccinate. See: Aquatyl, Tab. (Irwin, Neisler)

W/Bivanol lactate, cinnamyl ephedrine, pepper mint oil. See: Chovanol, Tab. (Winthrop Lab)

See: Cholic Acid, Oxidized

Dehydrocholic Acid

CHOLIC ACID, SALTS. See: Anobite, Tab. (Irwin, Neisler).

W/Ketoholanic Acids. See: Chobile, Tab. (Irwin, Neisler).

W/Methanamine. See: Pelamine, Tab. (Sandax).

W/Methylcellulose. See: Chaducell, Tab. (Mallinck).

CHOLIMETH, AROSOL. (Central) Trisholline citrate 1380 mg., Inositol 1002 mg., methionine 252 mg. Vit. B-12 15 mcg./15 cc. Bot. 1 pt., 1 gal.

Use: Lipotropic agent.

CHOLIMIN. (Vitarine) Choline dihydrogen

trate 1 Gm, dl methionine 0.33 Gm, inositol 0.25 Gm, Vit B 1 3.33 mg, B 2 3.33 mg B 12 3.33 mcg, w/liver stomach conc equiv to approx 12 Gm of fresh liver/3 Cap Bot 20s 50s 100s

Use: (1) (1 d) Lipotropic agent

CHOLINE BASE W/Inositol

See: Incholip, Liq (Flint Eaton)

Wychal, Cap & Liq (Wyeth)

W/Inositol & pyridoxine HCL See Vasculol

Liq (Schenley)

W/Betaine, Inositol, dehydrocholic acid ox bile, liver See: Hepozyn Tab (Endocrine)

* **CHOLINE BITARTRATE, N.F., A.D.R. 1955**

Cap, N.F., Tab

W/d Amphetamine, phenobarbital dl methionine & vitamins See: Obalip, Cap (Lakeside)

W/Ascorbic acid hesperidin complex dl methionine inositol pancreatin See: Nes per C Agotropic Cap (National)

W/Dehydrocholic acid See: Lipocholon, Tab (Moltbie)

W/Dehydrocholic acid methionine & betaine See: Cholotropic, Tab (Buffington)

W/Estratole, Vit B-12 & folic acid See: Menotrop, Tab (Lakeside)

W/Inositol See: Wychal Cap (Wyeth)

W/Inositol & dimethionine See: Cholathol Cap (Nysco)

W/Inositol, methionine & liver ext See: Lipophilate Cap (Brewer)

W/Inositol & multiple vitamins See: Cholines B Cap (Adams)

W/dl Methionine

See: Cholimeth, Tab (Central)

Methcolate, Tabs (Ascher)

W/Methionine, Inositol, folic acid, vit B 12 See: Delphicol, Cap (Lederle)

W/Inositol & dl methionine

See: Cholathol, Cap (Nysco)

Cholimeth Fortified (Central)

Lipocaps, Cap (Lakeside)

W/Inositol dl methionine ox bile See: Choline Compound Cap (Rowell)

W/d Pantothonyl alcohol See: Hopan

Choline Tab (Warren Teed)

W/Reserpine alkaloid, theophylline only drows See: Theracor, Tab (C. D. Smith)

W/Vit E and liver See: Geratose, Cap (Patch)

* **CHOLINE CHLORIDE, N.N.D. 1962 (Various Mfr)**

Abbott—Sol., Bot 1 pl, 1 gal

Upjohn—Syr 4 Gm./fl oz Bot 1 pl

Horton & Converse—Jelly (1%) Tube

1/2 oz

Tablerock—Elixir, Bot 1 pl—1 gal

Use: Liver disorders.

* Generic name recognized by N.F., U.S.P. or A.M.A.

W/Heparin Sod, folic acid nicotinamide benzyl alcohol See: Hepovite, Inj (Mallard)

W/Heparin Sod Vit B 12, folic acid & nicotinamide See: Hep Nine B, Vial (Columbus)

W/Methionine, inositol, B vitamins See: Chol Nine B Inj (Columbus)

CHOLINE CITRATE, TRICHOLOLINE CITRATE

W/Inositol methionine, B 12 See: Cholimeth aerosol (Central Pharm)

CHOLINE COMPOUND, (Rowell) Choline bitartrate 400 mg dl methionine 100 mg inositol 100 mg, ox bile ext 15 mg/Cap Bot 100s 500s 1000s

Use: Fat metabolism

* **CHOLINE DINYOROGEN CITRATE, N.F.**

N.N.D. 1962 Cap, Tab, N.F. 2 Hydroxy ethyl trimethylammonium citrate—U S Vit

0.5 Gm Bot 100s, 500s

Walker—Vit 250 mg, Bot 100s Tab, N.F.

Lily—10 gr Bot 100s 1000s

Parke Davis—10 gr Bot 100s 1000s

Walker Vit—500 mg Bot 100s 1000s

Use: Liver disorders

See: Cholix Syr, Liq (C. D. Smith)

Chothyn Tab, Syr, N.N.R. (Flint, Eaton)

Citrocholine Tab (United Lab)

W/Inositol & Methionine

See: Lipocaps Cap (Lakeside)

Lipotropic, Cap (Lakeside)

W/Inositol, polysorbate 80 See: Menichol

Liq (Iver Cameron)

W/Methionine See: Chamene Cap (Dumas Wilson)

W/dl Methionine inositol & vitamins See: Cholimin, Cap (Vitarine)

Chol Nine B Cap (Columbus Pharm)

C.M.I., Liq (Haskell)

W/Thyroid substance thiamine HCl & ribo-

flavin See: Methyroid Tab (Lakeside)

W/Vit B 12 & inositol See: Lipoliquid Liq

(Lakeside)

CHOLINE GLUCONATE SOLUTION, N.N.D.

1959 (Reed & Cornrick) Bot 1 pl

W/Inositol See: Sironositol, Liq

CHOLINE METHIONINE TARTRATE W/Ver-

trum viride, mannitol hexanitrate, ruf n &

phenobarbital See: Amvero Tab (Mallard)

* **CHOLINE SALICYLATE**

See: Arthropan Liq (Purdue Frederick)

Actasal Ped Drops (Purdue Frederick)

W/Cetyl dimethyl benzyl ammonium chlor-

ide See: Phoryclid n Concentrate (Purdue Frederick)

* **CHOLINE SUCCINATE, N.N.D. 1962**

* **CHOLINE THEOPHYLLINATE, N.N.D. 1962**

See: Cholelyt, Tab (Neperal)

W/Pentobarbital, rocephedrine NCl. See: Chlorace, Tab. (Nepero)

CHOLINERGIC AGENTS, (Parasympathomimetic Agents)
 See: Arecoline HBr,
 Bethanechol Chloride,
 Benzpyrinium Bromide,
 Carbachol,
 Carcholol. (Merck Sharp & Dohme)
 Deaner. (Riker)
 Doryl. (Merck Sharp & Dohme)
 Floropyl. (Merck Sharp & Dohme)
 Furmethide Iodide. (Smith, Kline & French).
 Furthrethonium Iodide.
 Guanidine HCl, Tab. (Davies, Rose).
 Humorsol (Merck Sharp & Dohme)
 Mechalyt Chloride & Bromide. (Merck Sharp & Dohme)
 Mestinan, Tab. (H. La Roche)
 Methacholine Chloride.
 Mytelosa (Winthrop Lab.)
 Neostigmine Bromide.
 Neostigmine Methylsulfate.
 Physostigmine Salicylate.
 Phospholine Iodide (Campbell Ph. Co.)
 Pilocarpine Nitrate.
 Prostigmine Bromide. (Roche).
 Prostigmine Methylsulfate. (Roche).
 Stigmonene Bromide. (Worner).
 Urecholine Chloride. (Merck Sharp & Dohme)

CHOLINERGIC BLOCKING AGENTS. See:
 Parasympatholytic agents.

CHOLINESTERASE, (Cutter Lab., Winthrop Labs.) See: Choline. (Cutter Lab.)

CHOLIPITENE, (Myers-Carter) Choline chlor. 200 mg., inositol 100 mg., dl-methionine 50 mg., Vit. B-12 6 mcg./2 cc. Vials 30 cc.

CHOLITROPIC, (Buffington) Dehydrocholic acid 175 mg., choline bitartrate 100 mg., methionine 50 mg., betaine 50 mg./tab. Bot. 72s, 500s.
 Use: Lipotropic agent.

CHOLI-VASCULARS, (Vit. Corp. of Amer.)
 Vit. E 30 mg, Inositol 150 mg., Vit. B-6 3 mg., Vit. B-1 10 mg., choline methionine 200 mg., Vit. B-12 5 mcg., Vit. C 30 mg./Cap. Bot. 100s, 1000s.
 Use: Lipotropic & multivitamin therapy.

CHOLMODIN, (Ames) Desoxycholic acid 1½ gr., aloes ext. ¼ gr./Tab. Bot. 50s, 500s.
 Use: (1-3 t.i.d.) Choleric & chologogue.

CHOL-NINE B, (Columbus) Choline dihydrogen citrate 320 mg., methionine 125 mg., inositol 50 mg., Vit. B-12 (cryst) 2 mcg., thiamin chloride 2.5 mg., riboflavin 2.5 mg., niacinamide 25 mg./Cap. Bot. 100s, 500s, 1000s.
 Use: Vitamin deficiencies.

CHOL-NINE B INJECTION, (Columbus)
 Choline chloride 75 mg., methionine 25 mg., inositol 15 mg., thiamin chloride 2 mg., riboflavin 2 mg., Vit. B-12 (cryst.) 5 mcg./cc. Amp. (1 cc.) Vial 10 cc.
 Use: B complex therapy.

CHOLOGESTIN, (Strong Cobb) Bile salts 2 gr., sod. sulfocylate 2½ gr., pancreatin 5 gr., sod. bicarb. 5 gr./15 cc. Bot. 12 fl. oz. See: Tablettestin. 3 tabs./15 cc. Bot. 100s. (Strong Cobb)
 Use: Choleric and chologogue.

CHOLEGIC, (Central) Bile salts 2 gr., papain ½ gr., duodenal substance ½ gr., sod. bicarbonate ½ gr., phenolphthalein ½ gr./tab. Bot. 100s, 500s, 1000s. Also available without phenolphthalein
 Use: (1-2 h.s.) Laxative.

CHOLOGRAFIM SODIUM, NND, 1962, (Squibb) Disodium salt of N,N'-diethyl bis-(3-amino-2,4,6-triadobenzoic acid). Sod. Iodipamide. Amp. (20%) 20 cc; 1 cc. for sensitivity testing.
 Use: Diagnostic agent.

CHOLOGRAFIM METHYLGLUCAMINE, NND, 1962, (Squibb) Iodipamide methylglucamine. Amp. 52% 20 ml. w/amp. 1 ml. for sensitivity testing
 Use: I.V., Cholangiography and cholecystography.

CHOLEREBIC, (Mallord) Desoxycholic acid 1½ gr./Tab. Bot. 100s, 1000s.
 Use: Choleric.

CHOLEXIN, (Flint Eaton Div., Baxter Labs., Inc.) Sodium Dextrothyroline
 Use: Cholesteropenic and lipopenic agent.

CHROMELIN COMPLEXION BLENDER, (Summers) Dihydroxyacetone 2%, Isopropanol 50%. Bot. 1 oz.
 Use: Cosmetic darkening of whitened skin areas.

CHOMENE, (Dumas-Wilson) Caps. Bot. 100s, 1000s.

***CHONDODENDRON TOMENTOSUM, NND, 1962.** See: Intocostin, Vial (Squibb)

CHONDRO-STANN, (Christina) Lin salt. 1 cc. Amp 12s, 24s, 100s
 Use: Tin therapy.

CHONDREUS, N.F., Extract, N.F., Mucilage, N.F. (Irish Moss)
 W/Petrolatum, liq. See: Kondremul (Patch)

CHORAND, (Smith, Miller & Patch) Gonadotropins, chorionic. Vial (10M u) 10 cc.
 Use: Cryptorchidism.

CHORON, (Massengill) Chorionic gonadotropin. Vial (5000 I.U.) 10 cc.
 Use: I.M., Cryptorchidism.

CHORIONIC GONADOTROPIN, NND, 1960. Injection. See: Gonadotropin, Chorionic, U.S.P.

CHOTHALOIN, (Elder) Phenolphthalein ½

* Generic name recognized by N.F., U.S.P. or A.M.A.

gr. cascara sagrada ext 1/2 gr, alum 1/2 gr, ex bile ext 1/4 gr / Tab Bot 100s, 500s, 1000s

Use: Laxative

CHOTHYN, N N D 1962 (Flint, Eaton) Choline dihydrogen citrate (syr 25% w/v) Cap (0.5 Gm) Bot 100s, 500s, 1000s, Syr 1 pt, 1 gal

Use: Lipotropic agent

CHOVANOL (Winthrop Labs) Chalc acid 1/2 gr, w/rivonal lactate, cinnamyl ephedrine & peppermint oil / tab Bot 100s

Use (1-2 HS) Choleretic & chologogue

CHROMARGYRE. See: Merbromin, Cryst (Prema)

CHROMELIN. (Summers Labs) Dihydroxy acetone 2%, rasopropanol 50% Applr Bot 1 oz

Use Dermatologic

CHRYSAZIN See: Danthron, N F

CHYLAMIN (Elder) Solicylomide 8 gr, colchicine 1/300 gr / Tab Sol 100s, 500s

Use: Analgesic

CHYLIPASE. (Columbus) Steapsin 600 mg, betaine 1200 mg, thyroid 30 mg / 6 Tab Bot 100s, 500s, 1000s

Use (6 daily) Correction of abnormal blood lipids; lipotropic agent

CHYMAR. (Armour) Chymotrypsin in oil, or Sod chloride solution 5000 u/cc Vial 5 cc

Use: i.M., Treatment of inflammations

CHYMAR BUCCAL (Armour) Chymotrypsin (proteolytic activity) 10 M. U. / Buccal Pkg 24s

Use: Treatment & management of inflammation

CHYMAR OINT. (Armour) Neomycin palmitate 3.5 mg & hydrocortisone HCl 1.25 mg & Proteolytic enzymes 10,000 u/Gm Tube 5 & 15 Gm

Use: Dermatological emul with proteolytic activity

CHYMASE. (Columbus) Vial 5 cc.

Use: Treatment of inflammation

CHYMOLASE. (Warren Teed) Chymotrypsin 5000 u with gelatin, 50 mg / cc Amp. 1 cc., Vial 5 cc

Use: Anti-inflammatory agent

CHYMOLASE ORAL (Warren Teed) Trypsin and chymotrypsin ratio of three to one 50,000 Warren Teed Units / Tab Sol 4Es

Use: Traumatic injury

CHYMORAL. (Armour) Trypsin and chymotrypsin, 50,000 Armour units (activity in ratio of approximately 6 to 1) / Tab Bot 4Es

Use: Oral anti-inflammatory

CHYMOTEST INJ. (Testagar) Proteolytic activity from chymotrypsin with gelatin, 5000

testagar units, 50 mg in water for injection / cc Vial 5 cc multiple dose

Use: Soft tissue inflammations

CNYMOTRYPSIN. An enzyme, a chymotrypsin, obtained in crystalline form from mammalian pancreas by aqueous acid extraction of its proenzyme, chymotrypsinogen, and subsequent conversion with trypsin to chymotrypsin

See Alpha Chymar Ophth (Armour)

Avazyme (Wampole)

Chymar, Vial (Armour)

Chymotest (Testagar)

Enxeon (Breon)

W/Gelatin

See Chymolase (Warren Teed)

Chylryp, Vial (B F Ascher)

W/Trypsin

Chymoral, Tab (Armour)

Ozenzyme, Tab (Natronol Drug)

CNYMOTRYPSIN INJECTION. (Philo Amoute Labs) Chymotrypsin 5000 u, gelatin 5%, with preservatives, in aqueous solution / cc Vials 5 cc

Use: Inflammation

CHYTRYP INJ. (Ascher) 5000 Ascher chymotrypsin units of proteolytic activity / ml Vial 5 ml multiple dose

Use: Acute inflammatory conditions

CIACIN. (Endo)

Amp I M, 1 cc: Copper acetate 0.63 mg + Iron cocodylate 13 mg, Sod. cocodylate 40 mg / cc Box 12s, 100s

Amp I V, 5 cc: Copper acetate 0.45 mg + Iron cocodylate 40 mg, Sod cocodylate 100 mg

Amp I V, 10 cc: Copper acetate 0.9 mg + Iron cocodylate 40 mg, Sod cocodylate 200 mg / 10 cc Box 6s, 25s, 100s.

Use: I M, I V, nutritional anemias

CISALGINE. (Ciba) Diallyl borbituric acid 1/2 gr, aminopyrine 3 1/2 gr / tab or 1 cc

Amp 2 cc, 5s; tab. 15s, 100s

Use: Sedative & analgesic.

CICOLINE. (Paul Maney) Chalc dihydrogen citrate 50%, Inositol 3 1/2%. Caps Bot 100s

Use: Cirrhosis of the liver.

CIDICOL (Upjohn) Ethylmorphine HCl 1/24 gr, w/Pat citrate, citric acid, pot gualacolate sulfonate / 5 cc Syr. Bot 1 pt, 1 gal

Use: (1-2 tsp. q 3 h) Narcotic cough preparation

CIGNOLIN, N N R 1947 See: Anthraquinone Oint (Abbott)

CLIFOMIDE. (Winthrop Labs) Pot penicillin G 300,000 u, sulfadiazine, sulfamerazine, sulfamethazine each 0.17 Gm / 5 cc Sulp-Bot 2 oz

Use: Orally, treatment of penicillin & sulphonamide susceptible organisms.

CILGRAM. (Columbus) Pot. penicillin G 100,000 u./5 cc. when dissolved as directed. Bot. 1 oz.

Use: Orally; penicillin therapy.

CILMICYN. (Wyeth) Penicillin fortified 400,000 u. w/dihydrostreptomycin 1 Gm./dose. Viol. 1 dose.

CILMICYN IMPROVED. (Wyeth) Procaine penicillin G 300,000 u., pot. penicillin G 100,000 u., dihydrostreptomycin base as sulfate 0.25 Gm., streptomycin base as sulfate 0.25 Gm./dose. Viol. 1 & 5 dose.

Use: Mixed or multiple infections.

CILLORAL. (Bristol)

Tab., 50,000, 100,000, 250,000 & 500,000 u. Pot. penicillin G/Tab. Bot. 12s, 100s.

Tab. Soluble; Pot. penicillin G 50,000 & 100,000 u., Bot. 24s, 100s.

Trachea; Procaine penicillin G 5000 u./Trache. Bot. 20s.

CILLORAL BUFFERED TABLETS. (Bristol) 100,000 u., 250,000 u., 500,000 u. Crystalline potassium penicillin G buffered w/calcium carbonate/Tab. Bot. 100s, 100,000 u. also Bot. 1000s.

Use: Parenteral penicillin therapy.

CIMADROX. (Messengill) Aluminum hydroxide gel, dried, 3 gr., mag. trisilicate 6 gr., ascorbic acid 20 mg., aromatics, q.s./Tab. Bot. 100s, 500s, 1000s. Comp. Pow. Bot. 8 oz.

Use: Gastric & duodenal ulcer.

CIMCIPUGA W/Sed. salicylate, gelsemium, pot. iodide, methyl salicylate. See: Realdine No. 3, Liq. (Messengill).

CINAPHYL, N.N.D. 1959. (Ascher) Theophylline sod. Glycinate. Tab. 0.16 Gm. & 0.33 Gm. Bot. 100s, 1000s.

Use: Diuretic.

CINRISAL (McNeil) Colchicine 1/250 gr., sod. salicylate 5 gr., ascorbic acid 15 mg./Tab. Bot. 100s, 1000s.

Use: (3 daily) Gouty arthritis.

CIN-CA-MINE "E." (Kirk) Cinchophen, cacodylate, methenamine. Viol 15 cc., Rox 6s, 25s, 100s.

Use: Urinary antiseptic.

CINCHONA BARK. (Various Mfr.)

Use: Antimalarial & tonic.

W/Anhydrous quinine, cinchonidine, cinchonine, quinidine & quinine. See: Totaquine. Pow. (Various Mfr.)

CINCHONINE. (Mallinckrodt) Alkaloid, Cryst. Pkg. 1 oz. Sulfate, Bot. 1 oz.

Use: As quinine dihydrochloride.

CINCHONINE SALTS. (Various Mfr.)

Use: As quinine dihydrochloride.

* Generic name recognized by N.F., U.S.P. or A.M.A.

CINCINOPHEN, N.F. X (Various Mfr.) Tab., N.F. X. Phenylcinchoninic acid, 2-Phenylquinoline-4-carboxylic acid

Use: As analgesic, antipyretic & in gout.

See: Atophan, Tab & Pow. (Warner-Chilcott).

W/Cacodylate, methenamine. See: Cin-Ca-Mine "E." (Kirk)

W/Mono-Iodo-cinchophen. See: Foroston, Cap. (Faroston).

CINCINOPHEN HYDROIODIDE. See: Oxy-Iodide, Tab. (Lilly).

CINCODYNE. (Centrol) Cinchophen 0.2 Gm., acetylsalicylic acid 0.2 Gm./Tab. Bot. 100s, 1000s

Use: Diuretic, analgesic, antipyretic

CINNAMON OIL, A D R. 1955; Water, (Various Mfr.)

W/Cat. carb. See: Calcicarb, Tab. (Hynson, Wescott & Dunning)

* **CINNARIZINE**, 1-Diphenylmethyl-4-trans-cinnamylpiperazine. See: Mitronal (G. D. Searle)

CINTRISUL 100 & 300. (Leeds) Penicillin G cryst. 100,000 or 300,000 units, sulfamerazine 0.167 mg., sulfadoxine 0.167 mg., sulfamethiazine 0.167 mg./Tab. Bot. 50s.

Use: Penicillin-triple sulfa therapy.

CIPAD. (Dumas-Wilson) Vit. D 400 u., B-1 2 mg., B-2 2 mg., ferrous sulfate ½ gr., dicalcium phosphate 7½ gr./Cap. Bot. 100s, 1000s.

Use: Vitamin with calcium & iron therapy.

CIRCULIN. (Fleming) Pentaoerythritol tetranitrate 40 mg./Tab. Bot. 100s, 1000s.

Use: Vasodilator.

CIRCULIN-R5. (Fleming) Pentaoerythritol tetracitrate 40 mg., butobarbital sodium 40 mg./Tab. Bot. 100s, 1000s.

Use: Vasodilator-sedative.

CIRIN. (Zemmer) Aspirin 5 gr., ascorbic acid 50 mg./Tab. Bot. 100s, 500s, 1000s.

Use: Analgesic.

* **CISMADINONE ACETATE.** (Eli Lilly) d'-6-chloro-17 α -acetoxyprogesterone acetate.

Use: Progestational compound.

CITHAL (Tablerock) Theobromine 4 gr., phenobarbital ½ gr., watermelon seed ext. 2 gr./Cap. Bot. 100s, 500s.

Use: (1-2 T.I.D.) Diuretic & sedative.

CITRA-DEL (Boyle) Pyrilamine 8.33 mg., methoprylone 8.33 mg., pheniramine 6.25 mg., hesperidin 100 mg., phenylephrine 5 mg., Vit. C 50 mg., salicylamide 200 mg., acetophenetidin 120 mg., caffeine 30 mg./Cap. Bot. 50s.

Use: Treatment of the common cold.

CITRA FORTE. (Boyle) Dihydrocodeinone B-tartrate 5 mg., hesperidin methyl chalcone

833 mg Vit C 30 mg, pheniramine maleate 2.5 mg, pyrrolamine maleate 333 mg
 pot citrate 150 mg / 5 cc Bot 1 pt Caps
 Bot 50s

Uses: Antitussive expectorant

CITRA H.P. (Boyle) Phenylephrine HCl 10 mg, prophenpyridamine maleate 625 mg, pyrrolamine maleate 833 mg, methapyrilene HCl 833 mg / Cap Bot 100s, 1000s

Uses: Common cold

CITRA LACTATE (Danley Evans) Sod lactate 40%, pot citrate 7½ gr / dr Bot 8 oz, 1 gal

Uses: Orally to increase the alkali reserve

CITRALKA. (Parke Davis) Sol of disodium hydrogen citrate 120 gr / fl oz Bot ½ pt, 1 gal

Uses: Urinary alkalization

CITRAMIN See: Helmitol, Tab (Winthrop Labs)

d-CITRAMINE (Preston) d Amphetamine sulf 5 mg / Tab Bot 100s, 1000s

CITRASULFAS (Upjohn) Sulfadiazine 15 gr, sulfamerazine 15 gr, sulfamethoxazole 15 gr, sod citrate 23¼ gr, alcohol 0.45% / fl oz Bot 1 pt 1 gal

Use: Triple sulfonamide therapy

CITRATED CAFFEINE, NF See: Caffeine Citrate

CITRATED NORMAL HUMAN PLASMA, NHR, USP. See: Plasma, Normal Human USP

CITRATE SALTS W/Calcium lactate sod phosphate & sod bicarb See: Calnom Effer Pow (Paxton)

CINTRIAMIDE (Lederle Labs) 3,4,5 trimethozycinnamide

Uses: Tranquilizer

CITRICOL. (Emerson) Pot guaiaccol sulfonate 8 gr, citric acid 7 gr, menthol 1/10 gr, sod citrate 20 gr, chloroform 2 min, wild cherry 5 min, wine ipecac 8 min / fl oz Bot 1 pt, 1 gal

Uses: (12 tsp q 3 h) Expectorant & sedative

CITRIN See: Vitamin P

CITRISAN (Chicago Pharmacol) Salicylamide 5 gr, ascorbic acid 85 mg, bioflavonoid complex 50 mg / tab Bot 50s, 100s, 1000s

Uses: (2 qid) Common cold

CITROBIC. (Blue Line) Hesperidin 100 mg, ascorbic acid 100 mg / Tab Bot 100s, 1000s

Use: Fragile capillaries

CITROCARBONATE (Upjohn) Sod & pot as citrates & tartrates, Cal lactate, sod phosphate, mag sulfate sod chlorides free sod bicarb 12½ gr / dr when dissolved Bot 2 oz, 4 oz, & 8 oz Tabs Bot 42s

Uses: Increase alkali reserve

* Generic name recognized by NF USP or A.M.A.

CITROCARBONATE, SYRUP. (Upjohn) Cit rates of sod and pot 7½ gr, cal lactate mag sulf, sod phosphate / Tsp Bot 1 pt, 1 gal

Uses: (2 to 4 tsp) Antacid

CITROCHOLINE. (United Lab) Choline di hydrogen citrate Tab 10 gr Bot 100s, 1000s

Uses: (1 2 Gm daily) Aid in fat metabolism

CITRO-CODEA. (First Texas) Codeine phosphate 1 gr, pot guaiaccol sulfonate 4 grs, sod citrate 30 gr, citric acid 6 grs / fl oz Bot 1 pt, 1 gal

Uses: (1 tsp q 3 h) Cough sedative & expectorant

CITRO-FLAV. (Direct) Bioflavonoid complex 25 mg, hesperidin complex 75 mg, Vit C 100 mg / Tab Bot 100s, 500s, 1000s

Uses: Vitamin supplement for abnormal capillary fragility

W/Salicylamide (Direct) Bioflavonoid complex 25 mg, hesperidin complex 75 mg, Vit C 50 mg, salicylamide 162 mg / Tab Bot 100s, 500s, 1000s

Uses: Muscular pains due to cold

CITROID COMPOUND. (Clayton) Citrus Bioflavonoid 100 mg, Vit C 50 mg, aspirin 4½ gr / Cop Bot 16s, 32s

Uses: Treatment of the common cold

CITRONIN (Parke, Davis) Ethylmorphine HCl 1/24 gr, w/citric acid, sod citrate pot guaiaccol sulfonate cos evoc, ipecac flexl & alcohol / fl dr Bot 1 pt, 1 gal

Uses: (1 tsp q 3 h) Narcotic cough preparation

CITRO-THIOL. (Rache Lab) Codeine phosphate 1/16 gr with pot guaiaccol sulfonate, citric acid, sod citrate & chloroform / 5 cc Bot 1 oz

Uses: (1 tsp q 3 h) Narcotic cough preparation

CITRO-TNYME. (Pitman Moore) Thyme 60 gr, ipecac syr 30 min, sod citrate 40 gr, benzoic acid 0.25% / fl oz Syr, Bot 1 pt 1 gal

Uses: (1 2 tsp p r n) Expectorant

CITROVORIUM FACTOR, NND 1962 See: Leucovorin Calcium

CITRUS BIOFLAVONOID COMPOUND

See: Bioflavonoid Cpds (Various Mfr)

C.V.P., Syr (U.S. Vitamin)

Koracitrin, Tab (Kormel)

Vitamin P

W/Ascorbic acid See: C.V.P., Cap (U.S. V.I. Corp)

W/Ascorbic acid, and aspirin See: Citroid Compound Cap (Clayton)

W/Ascorbic acid, and menadione.

See: Carperidin, Cap. (Dumas-Wilson)

Vitava, Cap. (Calumbus)

W/Ascorbic acid, hesperidin methyl chalcone, mucopolysaccharides. See: Hesperidone-C, Cap. (Cy Labs)

W/Ascorbic acid, phenyltoloxamine dihydrogen citrate, salicylamide, acetyl p-aminophenol, caffeine, racemic amphetamine sulf. See: Euphenex, Tab. (Lloyd, Dabney & Westerfeld)

W/Ascorbic acid, rutin. See: Capilon, Tab. (Plessner)

W/Ascorbic acid, rutin, Vit. B-6. See: Nion-dax, Tab. (Nion)

CLAPANONE. (Roche Lab.) Alkafanone. 3-Phenyl-3-sulfanyl-propiophenone. Tab. 0.5 Gm., Bot. 100s, 500s; Susp. (0.5 Gm./5 cc.), Bot. 16 oz.

Use: Nonspecific or simple diarrhea.

CLARETIN-12. See: Vitamin B-12.

CLABI-PORT. (Bio-Roma) Vit. A 25,000 u., D 1000 u., B-1 10 mg., B-2 5 mg., C 150 mg., B-12 2.5 mg., folic acid 1 mg., niacin 150 mg./Tab. Bot. 100s, 1000s.

Use: Multivitamin therapy.

CLARILAC. (Pitman-Moore) Milk protein hydrolyzate. Vial, 20 cc.

Use: I.M., nonspecific protein therapy.

CLARIN. N.N.D. 1962. (Thos. Leeming) Heparin pot. 1500 u./sublingual tab. Bot. 50s.

Use: (Under tongue). Clears lipemic serum. Management of atherosclerosis, hyperlipemia.

CLARKOTABS. (Clark & Clark) Racemic amphetamine phosphate 5 mg. w/thyroid, atropine & phenobarbital/Tab. Sets of 3 colors, regular set & high-potency set.

Use: (1 a.c., 1 i.d.) Anti-obesity therapy.

CLASOL. (Vitmix Corp) Naphazoline hydrochloride 0.05%, phenylmercuric acetate 1:5000. Bot. 1 oz.

Use: Nasal spray.

CLAYONE. (Maltbe) Rutin 50 mg., ascorbic acid 50 mg./Tab. Bot. 100s, 500s, 1000s.

Use: Orally, capillary permeability.

CLEAB-BUROW PWD. & PWD. PACKETS.

(Greco) 1:20 Burow's Sol. (Aluminum acetate, aluminum tartrate acetate, and sodium diacetate)/Packet or Tsp Packets 12s, 100s. Pwd. 4 oz cans, 1 lb.

Use: Wet dressing for allergic dermatitis, acutely inflamed skin conditions.

CLEFAMIDE. Name Approved by B.P.C. as: Dichloro-N-(2-hydroxyethyl)-N-(4-p-nitrophenoxybenzyl)acetamide Mebinol.

* Generic name recognized by N.F., U.S.P. or A.M.A.

* **CLEMIZOLE HCl.** N.N.D. 1962. 1-p-Chlorobenzyl-2-pyrrolidyl-methyl benzimidazole. See: Allercur, Tab. (Roerig)

Beotrol (Tailby-Nason Co., Inc.)

CLEREX. (Hygiene Research Inc) Modified carbohydrate gum base. Oint., Pkg. 3 oz., 4.5 oz.

Use: Skin protective against organic solvents.

* **CLIDINIUM BB OXIDE.** 1-Methyl-3-benziloyloxyquinuclidinium bromide. See: Quanzan (Roche)

W/Chloridiazepoxide. See: Librax (Hoffmann-La Roche)

CLIMACTIN. (Myers-Carter) Ant pit. gonad. hormone 5 Ru, B-1 25 mg., glutamic acid 52.5 ppm., procaine HCl 1%/cc. Vial 25 cc, Caps. Bot. 100s.

CLINICNBOME. (Wyeth) Vaginal smears, stain. Vial 1.025 Gm.

Use: Diagnostic aid, microscopic.

CLINILAB KIT. (Ames) Clinitest reagent tablets, Uristix reagent strips, Acetest reagent tab., lactates reagent tab., Hematest reagent tab./Kit. Complete kit.

CLINISTIX. (Ames) Glucose oxidase, peroxidase and orthotolidine. Diagnostic test for glucose in urine. Carton 12s, pkg. of 30 & 60 strips

Use: Diagnostic agent.

CLINISTIX IMPROVED. (Ames) Reagent strips—Bot. 30s, 60s, Dip-and-read urine test.

Use: Detection of glucose.

CLINITEST. (Ames) Tab. containing copper sulfate, sod. hydroxide & heat-producing agents. Patient's plastic set; Tab. refills, Bot. 36s; Box 24s; sealed in foil.

Use: For urine-sugar determination.

CLINOCAPINE HCl. See: Procaine HCl. (Taylor)

CLISTANAL. (McNeil) Clistin maleate 2 mg., aspirin 3½ gr., phenacetin 2½ gr., caffeine ½ gr./Tab. Bot. 100s, 1000s.

Use: Orally (1-2 q. 4 h); analgesic & antihistaminic.

CLISTIN EXPECTORANT. (McNeil) Carbinoxamine maleate 2 mg., ammonium chloride 2 gr., pot. guaiacal sulfonate 1 gr., sod. citrate 2 gr., w/chloroform & benzyl alcohol/oz. liq. Bot. 1 pt., 1 gal.

Use: (1 dr. q. 3 h) Antitussive.

CLISTIN MALEATE. N.N.D. 1962. (McNeil) Paracarbinoxamine maleate, 2-[p-chloro-alpha-(2-dimethylaminoethoxy)-benzyl] pyridine maleate, Tab. 4 mg. Bot. 100s, 1000s.

Rep action tab. Clistin R.A. 8 mg orange color 50s, 500s. Elixir (24 mg/oz) Bot 1 pt., 1 gal. Expectorant also available.

Use: Antihistamine.

W/A PC See: Clistinal, Tob (McNeil)
CLISTIN R-A (McNeil) Clistin molecule 6 mg in outer coating & 6 mg in inner core/Tab Bot 50s 500s Tab 8 mg, available
 Use Delayed antihistaminic action

***CLOBENZTROPINE** (Endo) 3-(p-chloro-*o*-phenylbenzoyloxy) tropone See Tepran

CLOCREAM (Upjohn) Vit A & D In vanesah Ing base Tube 1 oz; Jar 1 lb, 5 lb
 Use Topically, vitamin therapy

***CLOPENAMIC ACID** (Parke Davis) N [2,3-dichlorophenyl]anthranilic acid
 Use anti-inflammatory agent

***CLOMIPHENE CITRATE**, (Wm S Merrell) 1 [p- β (diethylaminoethoxy)phenyl] 1,2-diphenylchloroethylene dihydrogen citrate

***CLONITRATE** 3-chloro-1,2-propanediol di-nitrate See Dylate (Spencer Lab)

CLOPANE HCl, NND 1962 (Lilly) Cyclopentamine HCl Amp. 25 mg, 1 cc Pkg 6s, 100s Nasal sol 0.5% Bot 1 pt, 1 gal; 1% Bot 1 fl oz, 1 pt Sol 0.5% plastic squeeze Bot 1/2 oz
 Use: Relief of nasal congestion, vasoconstrictor & vasopressor

W/Pot penicillin G (Lilly) Cyclopentamine HCl 0.5% and 1 soluble tab pot penicillin G cryst 15,000 u Each cc. of 0.5% solution clopane HCl will contain 10,000 u of penicillin O Combination pkg 15 cc. of 0.5% solution clopane HCl, 1 soluble tab pot penicillin G cryst 150,000 u & drop per assembly
 Use: Acute & chronic sinusitis

W/Aludrine HCl See Aerolone Compound (Lilly)

W/Histolodyl See Histo Clopane, Cap (Lilly)

W/Histolodyl & pyronil See Co Pyronil Cap (Lilly)

CLOPHEDIANOL HCl (Roxane) See Acufuse AS

***CLOPHENOXATE HCl** (Winthrop Labs) 2 Dimethylaminoethyl p-chlorophenoxyacetate HCl
 Use: Cerebral stimulant

CLORACAINE, (Drug Frads) Aluminum subacetate, benzocaine, zinc oxide, thymal & chlorinated cottonseed oil Oint, Tube 1 1/2 oz, Jar 1 oz, 1 lb
 Use: Local anesthetic & antiseptic

CLORASEN, NNR 1949 See Dichloro Maphorsen, Amp (Parke, Davis)

***CLORDROLONE**, (Smith Kline & French) 4-chloro-17 β hydroxy 17 α methylster 4 en 3-one
 Use Anabolic agent

CLORPACTIN WCS-90, (Guardian Chem Corp) Monoxychlorosene Organic deriva
 * Generic name recognized by N.P., U.S.P. or A.M.A.

five of hypochlorous acid having active chlorine 2 Gm Bot. pkg 5 Bot
 Use: 2 Gm In qt water to treat infected area Bactericide, fungicide and sporicide

CLORPACTIN XCB, (Guardian Chem) Oxychlorosene Monoxychlorosene Four times activity of Clorpactin Wes 90 5 Gm Bot. pkg 4s
 Use: Topical anti-infective

CLORTRAN, (Wampole) Chlorobutanol Cap 0.25 Gm & 0.5 Gm Bot 100s
 Use: (0.25 Gm, q 4-6 h) Sedative & hypnotic

CLOVE OIL, W/Methyl sal, cinn oil, chlorobutanol See: Dentalane, liq (Parke, Davis)

CLUSIVITS, (Ayerst) Vit A 10,000 u, D 1000 u, C 50 mg, thiamine mononitrate 3 mg, B 2.5 mg, B-6 1 mg, d panthenol 5 mg, nicotinamide 25 mg, B 12 1/6 u, with intrinsic factor conc, plus eleven micrograms/Tab Bot 100s
 Use: Dietary supplement.

CLUSIVOL, (Ayerst) Vit A 25,000 u, D 1000 u C 150 mg, thiamine HCl 10 mg, riboflavin 5 mg, pyridoxine HCl 0.5 mg, panthenol, equiv to 10 mg, of Cal pantothenate, Vit B-12 w/intrinsic factor conc 1/6 unit, folic acid 0.5 mg, nicotinamide 50 mg, Vit E 0.5 IU, dl methionine 12.5 mg, lysine HCl 25 mg & trace minerals/1 Cap Bot 100s, 1000s
 Use: Nutritional reinforcement

CLUSIVOL GERIATRIC, (Ayerst) Premarin 0.125 mg, methyltestosterone 1.25 mg, Vit A 12,500 u, D 500 u, C 75 mg, B 1.5 mg, B 2.5 mg, B 6 0.5 mg, panthenol 1 mg, Vit B 12 w/intrinsic factor conc 1/6 u, folic acid 0.5 mg, nicotinamide 50 mg, biotin 0.05 mg, Vit E 0.5 IU, iron 25 mg, cal 100 mg, phosphorus 77 mg, copper 0.5 mg, cobalt 0.05 mg, manganese 0.5 mg, iodine 0.075 mg, potassium 2.5 mg, zinc 0.6 mg, magnesium 3 mg, lysine HCl 25 mg, dl methionine 12.5 mg/Cap Bot 100s, 1000s
 Use: Geriatrics

CLUSIVOL CHEW TABS, (Ayerst) Vit A 5000 USP u, O 1000 USP u, C 40 mg, B 1.2 mg, B 2.2 mg, B 6 1 mg, B 12 3 mcg, nicotinamide 10 mg, Ca-pantothenate 5 mg, Fe lum 3 mg, Mn 0.5 mg, Zn 0.5 mg, Mg 3 mg, Ca 40 mg, P 30 mg/Tab 50s
 Use: Nutritional reinforcement

CLUSIVOL SYRUP, (Ayerst) Vit A 2500 u 0.500 u C 20 mg, B 12 2 mcg, B 1 1 mg, B 2 1 mg, nicotinamide 5 mg, d panthenol 3 mg, B 6 0.6 mg iron, calcium phosphorus, manganese zinc, magnesium 1

- lysine HCl, Cysteine HCl, Inositol & choline/5 cc. Bot. 8. 20, 32 fl. oz.
Use: Nutritional supplement.
- CLU-VI-MINS.** (New Mexico Pharm. Co)
Vit. A 12,500 u., D 1000 u., B-1 5 mg., B 2 2.5 mg., niacinamide 40 mg., B-6 D.5 mg., calcium pantothenate 4 mg., lactic acid D.5 mg., B-12 1 mcg., C 75 mg., E 2 I.U., choline bitartrate 31.4 mg., dl-methionine 10 mg., inositol 15 mg., calcium 75 mg., phosphorus 58 mg., iron 30 mg., cobalt D.04 mg., copper D.45 mg., iodine D.075 mg., magnesium 3 mg., manganese D.5 mg., molybdenum 0.1 mg., potassium 2 mg., zinc D.5 mg./cap. Bot. 100s, 1000s.
Use: Vitamin & mineral supplement.
- CLYSEROL SOLUTION.** (Clyserol Labs.) Disodium phosphate 1.25 Gm., monosodium phosphate 5.83 Gm./100 cc. Plastic container 100 cc.
Use: Enema solution.
- CLYSODRAST,** 4-4'-(diacetoxydiphenyl)-(pyridyl-2) methane/1.5 mgs. lactic acid 2.5 gm./pack. Box of packs, 25s, 50s.
Use: For colonic evacuation (Barnes-Hind Barium Co)
- C M** (Massengill) Pepsin, bismuth subsalicylate, salol, zinc sulfocarbonate and ethyl alcohol in aromatic base. Bot. 4 oz.
Use: Simple diarrheas, nausea and heartburn.
- C.M.C. CAPSULES.** (Stuart) Sod. carboxymethylcellulose. Cap. 0.65 Gm. Bot. 100s.
Use: (3 daily w/water) laxative.
- CMC CELLULOSE GUM, NND, 1962.** See: Carboxymethylcellulose Sodium, Preps.
- CMC.** (Novacol Chem. Co) Eutectic mixture camphor and p-chlorophenol. Bot. 1 oz.
Use: Sterilizing root canals
- C.M.I.** (Hoskett) Choline dihydrogen citrate 3 Gm., methanine 0.6 Gm., inositol D.6 Gm., Vit. B-12 30 mcg./fl. oz. Bot. 1 pt., 1 gal.
Use: Lipotropic agent.
- C.M.S. COMPOUND.** (Jewell) Cal. carb., Mag. carb., Sod. bicarb., bismuth subcarb., kaolin, diastase, popain & peppermint oil. Pow., Con 4 oz.
Use: Antacid.
- COACTYN.** (Kinney) Hamatrapine methylbromide D.5 mg., phenobarbital 8 mg./5 cc. Bot. 3 oz., 1 pt.
Use: Antispasmodic with sedative.
- COAL TAR, U.S.P. Prep.**
See: Alphasyt (Reed & Carrick)
Appliderm No 6, Oint. (White)
Daxolan (Dome)
L.C.D. Camp., Oint. (Almay)
Nu-Kal-Tar, Cream & Lot. (Benef)
Ortho-Derma (G. F. Harvey)
Palytar Bath, Cap. (Stiefel Lab)
Supertah, Oint. (Talby-Nason)
Tarbonis, Cream (Tarbonis)
Tar Dermament, Lot. (Parke, Davis)
Tardook, Oint. (Doak)
W/Allantoin, hydrocortisone. See: Alphasyt-HC, Lot. (Reed & Carrick)
W/Hydrocortisone. See: Tarcartin, Cream. (Reed & Carrick)
W/Mercury oleate, salicylic acid, phenol & p-nitrophenol.
See: Prasal, Emulsion (Torch)
Estercol, Emulsion (less p-nitrophenol) (Torch)
W/Nitromersol, quillaja ext., benzoin, & Isopropyl alcohol. See: Kericide, Sol. (Kor-mel Co)
W/Quinolor. See: Tarquinor, Cream (Squibb)
W/Pyriminole maleate. See: Histar, Oint. (Tarbonis).
W/Salicylic acid, sulfur and menthol. See: Sol-Su-Tar, Oint. (Ulmer)
W/Sulfathiazole.
See: Corbathiazole, Oint. (Rorer).
Sol-Tarbonis, Oint. (Tarbonis).
W/Sulfur, camphor, chaulmoogra oil & zinc oxide paste. See: Pasta Durel, Oint. (Doak).
W/Sulfur & salicylic acid. See: Supertah w/sulfur & salicylic acid, Oint. (Talby-Nason).
W/Zinc oxide.
See: Pix-Gel, Oint. (Pitman-Moore)
Tarpaste (Doak)
W/Zinc oxide & starch.
See: Co-Go-Tar, Liq & Oint. (Texas Pharm)
Co-Tar-Ox, Oint. (Ulmer)
Daxolan, Oint. (Dome)
Kalsix (Dome)
- COAL TAR, DISTILLATE.**
See: Distarol, Cream & Oint. (Manhattan)
Lavorol, Liq (Doak)
Tarolba, Oint. (Upjohn)
Teroxide, Oint (Abbott)
W/Colamine & salicylic acid. See: Distarol w/colamine, Oint. (Manhattan).
W/Resorcinol & salicylic acid. See: Tasmel, Oint. (Smith, Kline & French).
W/Sulfur, menthol & starch. See: Dencolor, Liq & Oint. (Denver Chemical).
W/Sulfur & salicylic acid. See: Pragmatol, Oint. (Smith, Kline & French).
W/Zinc oxide. See: Ultraine, Oint (Bischoff).
W/Zinc oxide & starch. See: Kalsix A & D, Oint (Dome).
- COAL TAR EXTRACT** W/Allantoin. See: Alphasyt, Lot. (Reed & Carrick)
W/Allantoin, squalene. See: Skaylas Lotion (Lee-Colbert).
W/Allantoin, hexachlorophene. See: Sebical Cream (Reed & Carrick)

W/Nitramersal, quillaia See Koricide, Liq (Karmel)

COAL TAR PASTE. (No Mfr currently lists)

W/Zinc paste See Tarpaste, Paste (Doak)

COAL TAR SOLUTION, USP Liquor Carbonis Detergens LCD

See Carbo Dome (Dome)

Crealin Pearson (Merck Sharp & Dohme)

LCD Oint (Almay)

Ortho Derma (G F Harvey)

Wright's Saln (Faugera)

W/Burrows solution & aint base See Bur Tar Gen Oint (Texas Pharm)

W/Hydrocortisona alcohol micronized allon toin See Altara Cart Gell (Dome)

W/Ointment base See Torcreme, Oint (Strassenburgh)

W/Pyrimamine maleate benzocaine, menthol See Carbo Hist Cream (Paul Money)

W/Salicylic acid See Epidol (Epidol Co)

W/Sulfur, colloidal salicylic acid & zinc oxide See Teparor, Cream (Merck Sharp & Dohme)

COAL TAR SYNTHETIC OINT, See Torolin (Smith Dorsey)

CO-AMINO (Physiological Chemicals) Protein hydrolyzates of beef, yeast, & B complex factors w/added vitamins for 1 lb (454 Gm)

Use: Aminoacids therapy

COAX, (Cambridge) Aspirin 3½ gr, phenacetin 2½ gr, caffeine ¼ gr phenobarbital ¼ gr, pvd ext belladonna 162 mg phenylephrine 5 mg, chlorpheniramine 4 mg/Tab Bot 100s

Use Coughs and colds

COBACINE (New Mexico Pharm Co) Adenine 5 monophosphoric acid 25 mg Vit B 12 60 mcg, benzyl alcohol 1.5%/cc Liq Vials 10 cc, 30 cc

COBADEN (Rand) 25 mg of Adenine 5 monophosphoric acid & 60 mcg of Vit B 12/cc Vial 10 cc.

Use Stimulates erythropoiesis & specific for the treatment of Addisonian or pernicious anemia

COBALAMED. (Medics) Sterile sol of cyanocobalamin USP containing 200 mcg per cc in base of 4% gelatin USP and 0.9% sodium chloride USP Vial 10 cc

Use: Aqueous injections of B 12

***COBALAMIN CONCENTRATE,** NP See Vitamin B 12 Concentrate (Various Mfr)

COBALT W/iron, Vit B 12 & B 1 See: Hemo-Vital Liquid (Smith)

W/iron col, B 6, D See Glucodax, Cap (Kirkman)

* Generic name recognized by NF, USP or AMA

COBALT CHLORIDE W/Ferrous sulfate

See: Ferrabalt, Tabs (Testagar)

Rancavite, Prep (Lloyd)

W/Ferrous sulf, copper sulf, mang, sulf, vit C See: Ceniron, Tab (Central Pharm)

W/Ferrous sulf, mang sulf See: Colron Tab (Houg)

W/Ferrous sulf & vit B complex See: Hema-lavals, oval (Uimer)

COBALT THERAPY,

See Ceniron, Tab (Central)

Cabedox, Cap (Lincoln)

Cacuron Tab (Kendall)

Cafran, Tab (Houg)

Cumaliron, Syr (Borchardt)

Ferrabalt, Tabs (Testagar)

Glucodax, Cap (Kirkman)

Hemo-Vital, Liq (Smith)

Rancavite, Prep (Lloyd)

COBASAL. (Rand Pharm) Dipyrone 125 mg; p aminobenzoic acid 8 gr, salicylamide 6 gr, Vit B 12 5 mcg /2 tab Bot 100s 1000s. Use: 2 tab t i d; antirheumatic

COBASAL WITH COLCHICINE. (Rand) Colchicum seed ext 0.625 gr, dipyrone 0.125 Gm, para amino benzoic acid 8 gr, salicylamide 6 gr, Vit B 12 5 mcg /Tab Bot 100s, 1000s

Use Gout

COBAVITE. (Metro) Vit B 12 Vial 1000 mcg/cc, 10 cc; 100 mcg/cc, 10 cc; 30 mcg/cc, 10 cc; 30 mcg/cc, 30 cc; Amp 30 mcg/cc, Box 12s, 100s, 15 mcg/cc; Box 10s 100s

Use Anemias

COBODOCE. (Lincoln) Caball gluconate 25 mg, ferrous gluconate 200 mg, Vit B-12 5 mcg, dexte stomach subs 150 mg/Cap Bot 100s 500s

Use (i p c, t i d) Anti anemia

COBEPRIN. (Winthrop Labs) 1,3,4 Dihydroxynorephedrine Norhamoepinephrine Nordefrin

Use Vasconstrictor

W/Novocain See Novocain Cabetrin, Amp (Winthrop Labs)

COBEGEL (G F Harvey) Cyanocobalamin 300 mcg/cc in sesame oil gelled with 3% aluminum monostearate Vial 5 cc

Use Anemia

COBENZIL. (Abbott) Cadeine phosphate 1 gr, sod citrate 25 gr, ammonium chloride 5 gr, ipecac syr in 30 m, menthol & aromatics qs in benzinated syr/fl oz (Abbott x Benzinated Cough Syrup) Bot 1 pt, 1 gal

Use (1 tsp q 3 h) Expectorant syrup for coughs

COBETABON. (Warren Teed) Vit B 12, cryst

- 10 mcg., stomach pow. 100 mg., folic acid 1 mg., ascorbic acid 50 mg., ferrous sulfate, dried 194.4 mg., desic. liver 200 mg./Cop. Bot. 50s, 500s.
 Use: (1-3 daily) Anemias.
- COBEX.** (Metro) Vit. B-12 10 mcg., B-1 10 mg., B-6 10 mg., panthelol, 10 mg., niacinamide 25 mg., riboflavin 2 mg./cc. Vial 10 cc.
 Use: Vitamin B complex therapy.
- COBIC-FORTE.** (Confield) Cobalt gluconate 9 mg., peptonized iron 59 mg., folic acid 5 mg., liver injection equiv. to B-12 activity 5 mcg./2 cc. Vials 10 cc.
 Use: Anemias.
- COBIONE, N.N.D. 1959.** See: Vit B-12.
- * **COBRAVENOM.** See: Cobroxin, Amp. (Hynson, Westcott & Dunning)
 W/Formic acid & silicic acid. See: Nyloxin, Amp. (Hynson, Westcott & Dunning)
- COBRA VENOM SOLUTION.** (Hynson, Westcott & Dunning) Cobra venom, containing 10M u./cc. Amp. 1 cc., Box 10s.
- COBROXIN.** (Hynson, Westcott & Dunning) Cobra venom sol., the neurotoxic principle Amp. 1 cc. (50 mouse u.) Box 10s, 25s.
 Uses (S.C. & I.M.) Analgesic for intractable pain.
- * **COCAINE, N.F.** Alkaloid, Pow. (Mallinckrodt). Hydrochloride, U.S.P., Pow. (Mallinckrodt) Tob., N.F., 1.14 gr. & 2.28 gr. Bot. 100s. (Sharp & Dahme).
- * **COCAINE HYDROCHLORIDE, U.S.P., A.D.R. 1955; Tob., N.F.**
 W/Epinephrine. See: Codrenin, Sol. (Parke, Davis).
- W/Homatropine HCl. See: Homatropine HCl & cocaine HCl. Tablets, Ophthalmic. (Burroughs Wellcome)
- COCCIDIOIDES IMMITIS, MIXED STRAINS.** See: Coccidioidin, Vial. (Cutter).
- COCCIDIOIDIN.** (Cutter) Mixed strains of coccidioides immitis. Vial. 1:100 dilution 1 cc.; 1:10 dilution 1 cc.
 Use: Diagnostic aid of Coccidiomycosis.
- COCCULIN, N.N.D. 1961.** See: Picrotoxin. (Various Mfr.)
- CO-CILLIN TROCHES.** (Coastal Pharm.) Pot. penicillin G 10,000 u./Troche. Pkg. 20s.
 Use: Vincent's infection.
- COCIN.** (Dumas-Wilson) Cyanocobalamin 25 mcg., B-1 10 mg., niacinamide 5 mg./tsp. Bot. pl.
 Use: Appetite stimulant, nutritional supplement.
- COCI-TOL.** (Bullington) Ethylmorphine HCl ¼ gr., alcohol 6%, eucalypta tr. 24 min., euphorbia tr. 120 min., wild lettuce syr.
 * Generic name recognized by N.F., U.S.P. or A.M.A.
- 120 min., cascara sagrada ext. 8 gr., menthol 2/25 gr./fl. oz. Bot. 1 pt., 1 gal.
 Use: (1 tsp. q. 3 h.) Sedative for coughs.
- COCOA BUTTER, U.S.P.** See: Theobroma Oil.
- COCO-OIAZINE.** (Lilly) Sulfadiazine 0.5 Gm./dr. Bot. 1 pt.
 Use: Sulfadiazine therapy.
- COCONUT OIL W/Sucrose.** See: Ediol., Liq. (Schaeley).
- COCO-QUININE, N.N.R. 1949.** (Lilly) Quinine sulfate 10 gr./fl. oz. Bot. 3 fl. oz., 1 pt., 1 gal.
 Use: Malaria.
- COCO-SULFONAMIDES TRIPLEX.** (Lilly) Chocolate-flavored susp. containing in each 100 cc., sulfathiazole 3.33 Gm., sulfadiazine 3.33 Gm. & sulfamerazine 3.33 Gm. Bot. 1 pt.
 Use: Sulfonamide therapy.
- COCO-THIAZOLE.** (Lilly) Sulfathiazole 40 gr./fl. oz. Bot. 1 pt.
 Use: Sulfathiazole therapy.
- COCURON.** (Kendall) Cobalt, copper & iron. Tob., Bot. 100s, 500s. Drops 15 cc.
 Use: Orally (1 t.i.d., p.c.) erythrogenic.
- CODADYNE.** (G. F. Harvey) Codeine sulfate 1 gr., aspirin 3½ gr., acetophenetidin 2 gr., caffeine ½ gr./cap. Bot. 100s.
 Use: Analgesic with sedative.
- CODANIST SYRUP.** (Burroughs Bros) Codeine phosphate 8.7 mg., pyrilamine maleate 5 mg., chloroform, pot. guaicol sulfonate, ammonium chloride, menthol & raspberry/4 cc. Bot. 1 pt., 1 gal.
 Use: (1 tsp. q. 3-4 h.) Cough due to colds.
- CODALEX.** (Mottard) Dihydrocodanone bitartrate 1/6 gr., pot. guaicol sulfonate 8 gr., ephedrine HCl 0.5 gr., sod. citrate 20 gr., citric acid 5 gr./fl. oz. Bot. 3 oz., 1 pt., 1 gal.
 Use: (1 tsp. q. 3 h., m.p.) Cough due to colds.
- CODANOL (A.P.C.) VII, A, D, and C w/ Vitamin B complex.** Liq. Bot. 4 oz. Oint. Tube 1 oz.
 Use: Dietary supplement.
- CODANOL MALT.** (A.P.C.) Vit. A 5000 u., D 1200 u., B-1 1 mg., B-2 0.5 mg., niacinamide 10 mg., iron & ammonium citrates, green 3 gr./3 dr. Bot. 1 lb., 3½ lb.
 Use: Nematocin.
- CODASPRO No. 1 & No. 2.** (Vanpelt & Brown) New name for Bellaspro. Tob. W/Codeine.
- * **CODEINE, N.F., N.N.D., 1962, A.D.E.** (Various Mfr.) Cryst. or Pow. Bot. ¼ oz., ½ oz., ½ oz. & 1 oz.
- CODEINE ACETATE W/Chloroform.** Col.

bromide belladonna root flexi See Cal
braden, Liq (Upjohn)

Codeine combinations on pp. 189 to
191.

CODEINE METHYLBROMIDE. (Eucodin)

Use Narcotic cough depressant

***CODEINE PHOSPHATE, USP, ADR**; Inj
USP Tablets, USP, ADR 1955 (Various
Mfr) Pow Bol 1/4 oz, 1/2 oz, 1/2 oz &
1 oz

CODEINE PHOSPHATE TABLETS H.T. (Merck
Sharp & Dohme) 1/4 gr, 1/2 gr & 1 gr Bol
100s

***CODEINE SULFATE, NF, ADR** (Various
Mfr) Pow Bol 1/4 oz, 1/4 oz, 1/2 oz &
1 oz Tab, NF, 1/4 gr, 1/2 gr & 1 gr

CODELORTONE, See Prednisolone

CO-DELTRA (Merck Sharp & Dohme) Pred
nisone 2.5 mg or 5 mg, w/magnesium tris
silicate 50 mg & aluminum hydroxide gel
300 mg /Tob Bol 30s, 100s, 500s

Use: (1 q d) Hormone therapy

CODEMPERAL (Burroughs Wellcome) Na. 2;

Phenobarbital 1/4 gr, aspirin 3 1/2 gr, aceto
phenolidin 2 1/2 gr, codeine phosphate 1/4
gr /Tob Bol 100s; Na. 3, Codeine phos
phate 1/2 gr w/emperal/Tob Bol 100s

Use: Analgesic & sedative

CODESAL (Burlington) Na. 1; Codeine sul
fate 1/4 gr, acetophenolidin 2 1/2 gr, acetyl
salicylic acid 3 1/2 gr, caffeine 1/4 gr,
gelsemium lr 1 min /Tab; Na. 2, Codeine
sulfate 1/2 gr /Tob Bol 100s, 500s, 1000s

Use: (1 m p) Analgesic & antipyretic

CODESSIN. (Warren Teed) Codeine sulfate
1/6 gr w/Sod salicylate ammonium chlor
ide & tartar emetic/5 cc Syr Bol 2 oz,
1 pt, 1 gal

Use: (1 tsp q 3 h, m p) Narcotic cough
preparation

CODIAZINE (SUSPENSION). (Massengill)

Microcrist sulfadiazine 13 Gm (20 gr), sn
cocoa suspension/15 cc Bol 1 pt, 1 gal

Use: Sulfa therapy

CODIGESIC. (Central) Dihydrocodeinone bi
tartrate 10 mg, phenylephrine HCl 30 mg,
pyrilamine maleate 50 mg, acetyl p Amino
phenol 0.5 Gm, pot guaicolalsulfonate D 5
Gm, sod citrate 13 Gm, citric acid D 3
Gm, chloroform D 1 ml /oz Bol 2, 3, 4 oz,
pints, gal

Use: Respiratory disorders

CODIMAL DH SYRUP (Central) Pyrilamine

maleate 50 mg, dihydrocodeinone bitartrate
10 mg, Pot guaicolalsulfonate 500 mg,
Sod citrate, citric acid chloroform/30 cc
Bol 3 oz, 1 pt, 1 gal

Use: Ant histaminic expectorant

CODIMAL PH SYRUP. (Central) Codeine

* Generic name recognized by NF, USP or A.M.A.

phosphate 60 mg, phenylephrine hydro
chloride 30 mg, pyrilamine maleate 50
mg, potassium guaicol sulfonate 0.5 Gm,
sod citrate 13 Gm, citric acid 0.3 Gm,
chloroform 0.1 ml /fl oz Syr Bol 3 oz,
1 pt, 1 gal

Use: Sedative expectorant for coughs due
to colds

COOIS (Hoack) Salicylamide 7 1/2 gr, col
leine anhy 1/2 gr, dihydrocodeinone bitar
trate 1/4 gr /Tob Bol 100s, 1000s

Use: (1 q 3-4 h, m p) Cough depressant

CODISTAN. (Chicago Pharmacol) Dihydro
codeinone bitartrate 10 mg, pyrilamine
maleate 200 mg, glyceryl guaicolate 100
mg, menthol 6.5 mg, chloroform 0.75 mg /
oz Syr Bol 1 pt

Use: Treatment of coughs and colds

CODITRATE SYRUP. (Central) Dihydro
codeinone bitartrate 1/6 gr, pot guaicol
sulfonate 8 gr, sod citrate 20 gr, chloro
form 1.5 min, citric acid 5 gr /fl oz Syr
Bol 2, 3, 4 oz, 1 pt, 1 gal

Use: (1 tsp q 3-4 h, m p) Cough de
pressant

CODITRATE TABLETS. (Central) Dihydro
codeinone bitartrate 2.4 mg, pot guaicol
sulfonate 2 gr, sod citrate 5 gr /Tob Bol
100s, 500s

Use: (1 q 3-4 h, m p) Treatment of coughs

W/Parmol (Central) Dihydrocodeinone bi
tartrate 1/6 gr, pot guaicol sulfonate
8 gr, sod citrate 20 gr, citric acid 5 gr,
chloroform 1.5 min, pyrilamine male
ate 1/4 gr /fl oz Bol 1 pt, 1 gal

Use: (1 tsp q 3-4 h, m p) Cough de
pressant

COO LIVER OIL, ADR Emulsion, NF,
ADR 1955 Emulsion w/malt, NF, ADR
1954

Use: Vitamin A & D therapy

W/Malt ext (Burroughs Wellcome) VII A
645D e, Vit O 645 e Bol 10 fl oz, 20
fl oz

W/Metaphen (Abbott) Cod liver oil 70%
metaphen 1:2500 Tubes 1 oz, 3 3/4 oz;
Jar 1 lb, 5 lb

W/Viosterol, ADR 1954

Abbott—Vit A 2800 u, Vit D 255 u/
Gm Bol 12 fl oz

Squibb—Vit A 2000 u, Vit D 440 u/
Gm Bol 3 fl oz, 1 pt

W/Amyl, p aminobenzoate, benzocaine See
Ultrasin, Oint (Herzog)

W/Anesthesin, zinc oxide, hydroxyquinoline
See: Medicine Dressing (Medicine)

W/Benzocaine See: Marusan, Oint (Mas
sengill)

W/Phenol See: Gadoment, Oint (Patch)

| Cadeine | Others | Manufacturer | Brand |
|----------------------|--|----------------------|----------------------------------|
| (Phos.) 1 gr | Amm chlor 8 gr., pot guaiacal sulfonate 8 gr., tartar emetic 1/16 gr., chloroform 2 min., menthol 1/16 gr., white pine & wild cherry | Cole | Cren, Syr /A oz |
| (Phos.) 1/6 gr | Amm chlor, pot guaiacal sulfonate, tartar emetic, white pine, wild cherry | Upjohn | Charcoal/5 cc. |
| (Sulf.) 0.5 gr | Amm chlor 8 gr., senega, sod cit., grandelho, menthol | Chicago Pharm | Respi Sed/A oz |
| (Phos.) 1 gr | Amm chlor 8 gr., sod benzoate 4 gr | Blue Line | Syr codate/A oz |
| (Phos.) 1 gr | Amm chlor 5 gr., sod citrate, menthol, aromatics | Abbott | Cobensil/A oz |
| (Phos.) 1/4 & 1/2 gr | A P C belladonna oil (tr 3-8 min) | Vanpeit & Brown | Bellospro W/Cod, Tob No 1 & No 2 |
| (Sulf.) 1 gr | A P C, gelsemium tr 1 min | Burffington | Codestol No 2, Tob |
| (Sulf.) 1/4 gr | A P C, gelsemium tr 1 min | Burffington | Codestol, No 1, Tob |
| (Phos.) 1/2 gr | Aspirin 3 1/2 gr., phenacetin 2 1/2 gr., phenobarbital 1/4 gr | Burroughs Wellcome | Codemperial Tob No 3 |
| (Phos.) 1/4 gr | Aspirin 2 1/2 gr., phenacetin 2 1/2 gr., phenobarbital 1/4 gr hy atcyamus ext 1/4 gr | Hentzell | Hotacade Tob |
| (Sulf.) 1/40 gr | Atropine sulf 1/500 gr, ipccac, A P C. | Messengill | Dasin CS/Cop |
| (Acet) 1/2 gr | Bellodonna root fldxt 1/4 min., chloroform 1 gr., col bromide, benzoc acid | Upjohn | Colbrodein Liq/A oz |
| (Sulf.) 1/2 gr | Benzedrine sulf 2.5 mg., aspirin 2.5 mg., phenacetin 2.5 gr | Smith Kline & French | Edrisol, Tab |
| (Sulf.) 1/6 gr | Cal iodide, chloroform, wild cherry, tolu squill, yerba santa, citric acid, menthol | Winthrop Labs | Deka/5 cc |
| (Phos.) 1 gr | Chloroform, white pine poplar bud, glycerin alcohol | Parke, Davis | Cosodein/A oz |
| (Sulf.) 1 gr | Ephedra 30 gr., col guaiacal sulfonate 5 gr | Stratensburgh | Mor Phedoc/A oz |
| (Sulf.) 1/6 gr | Ephedra pot guaiacal sulfonate, tolu, squill, menthol | Lilly | Ephedrol/5 cc |
| (Phos.) 1 gr | Ephedrine sulf 1/4 gr., benzyl olc 7.47 min., sorbital 180 gr., mannitol 45.3 gr | Od Peacock | Ben Co Fed/A oz |
| (Phos.) 1/12 gr | Ipccac pot guaiacal sulfonate, chloroform, sod citrate, citric acid | Wyeth | Cetro-Cirose/5 cc |
| (Phos.) 1/6 gr, | Ipccac fldxt, squill, chloroform | Searle | Codofarm/5 cc |
| (Sulf.) 1/9 gr | Ipccac ext, squill ext sanguinaria, cherry, chloroform | Norwich | Teek/5 cc |
| (Sulf.) 1/2 gr | Ipccac squill, senega sanguinaria, menthol | Squirb | Senodin syr/A oz |
| (Sulf.) 1/6 gr | Ma huang, pot guaiacal sulfonate, tolu squill, menthol | Lilly | Ephedrol/5 cc |
| (Phos.) 1 gr | Phenidamine tartrate 60 mg., phenylephrine HCl 30 mg., chloroform 1 min, ipccac fldxt glycerin pot guaiacal sulfonate sod citrate, citric acid | Ives Cameron | Cerose 1 q/oz |

W/Vit A & D See: Col Vi Nol Ointment (Ulmer)
 W/Yeast phenyl mercuric nitrate See: Bio Dyne Oint (Whitehall)
 W/Zinc borosalicylate See: Vitaquent, Oint (Tailby-Nason)
 W/Zinc oxide & talcum See: Desitin, Oint (Desitin)
 W/Zinc oxide mag carb & lime water See: Desitin, Emulsion (Desitin)
COD LIVER OIL CONCENTRATE, N N R
 1952 ADP 1955 Concentrate of non saponifiable fraction of cod liver oil
 Clinodal—Bot 60 cc
 White—Cap, Liq & Tab
COD LIVER OIL EMULSION, N F W/Mall, N F
COD LIVER OIL, HIGH POTENCY
 Parke Davis—Vit A 2000 u, Vit D 250 u/Gm Bot 4 fl oz, 12 fl oz 1 pl; Cap 0.062 cc, 12 cc each 0.062 cc Box 12s 100s
 Patch—Vit A 2000 Vit D 200/Gm Bot 4 fl oz, 12 fl oz, 1 1/2 pl
COD LIVER OIL OINTMENT See: Burbo-Caine Oint (Rowell)
 Maruquent Oint (Massengill)
CODOFORM, (Searle) Codeine phosphate 1/6 gr Ipecac flex chloroform & squill/ 5 cc Bot 1 pl, 1 gal
 Use: (1 tsp q 3 h, m p) Narcotic cough preparation
CODONE (Lemmon) D hydrocodeinone bitartrate Tab 5 mg Bot 100s 1000s
 Use: (1 q 3-4 h) Treatment of coughs
CODRENIN (Parke, Davis) Cocaine NCl 9 1/3 gr, adrenol n 1/33 gr (1:15 000)/ fl oz Bot 1 oz
 Use: Local anesthetic
CODSAL (Horton & Converse) No 1: Codeine phosphate 1/4 gr, aspirin 3 1/2 gr, acetophenetidin 2 1/2 gr, caffeine 1/2 gr / Tab Bot 100s, 500s 1000s; No 2: Codeine phosphate 1/2 gr, aspirin 3 1/2 gr, acetophenetidin 2 1/2 gr, caffeine 1/2 gr./Tab Bot 100s 500s 1000s
 Use: Analgesic
CO-ELORINE, (Lilly) Elixir sulfate, tricyclanil sulfate 25 mg, amylal 8 mg / Pulvule Bot 100s 1000s Or Elorine chlor de 100 mg, amabarbitol 16 mg./Pulvule Bot 100s 1000s
 Use: (1 2) Spasmolysis without sedation
COENZYME-B (Barrons) Diphosphothiamin (Cocarbonylose) 25 mg / Cap Bot 50s (50 mg / 2 cc). Amp Box 100s
 Use: Coenzyme for Vitamin B 1
COESTRONE (Barry) Estrogens 20000 IU / cc Susp Vial 10 cc
 Use: I.M. estrogen therapy

COFERIC, (Crestmed) Ferrous sulfate 200 mg cobalt chloride 15 mg, ascorbic acid 100 mg, B-12 2 mcg / Tab Bot 100s.
 Use: Anemias
CO-FERRIN (Ulmer) Iron, peptonized, 30 mg; cobalt gluconate, 9 mg; folic acid 5 mg; Liver Injection USP, 5 mcg; procaine NCl, 1% / cc Parenteral; multi purpose anti anemic; 10 cc, 1s, 24s; 1 cc, 6s.
 Use: Resistant anemias
COFRON CAPSULES, (Abbott) Ferric ammonium citrate 0.6 gr (elemental iron 7 mg [1/10 gr]) copper sulfate 1/60 gr (elemental copper 0.27 mg [1/250 gr]) liver concentrate, N.F. & liver fraction 2 NF 0.5 Gm / Cap Bot 100s, 500s, 1000s.
 Use: Secondary anemias
COFRON ELIXIR WITH VITAMIN B-12 (Abbott) Ferric ammonium citrate 1.15 Gm representing approx 200 mg elemental iron copper sulfate 10.8 mg representing 2.66 mg elemental copper, Vit B-12 as concentrate, 6 mcg liver fraction No 1, NF 2 Gm in wine flavored base/fl oz Bot 12 fl oz, 1 gal
 Use: Nematotic
COFRON TABLETS, (Abbott) Soluble ferric pyrophosphate, 0.22 Gm (Fe 25 mg), copper sulfate 4 mg (Cu 1 mg) / Tab Bot 100s, 1000s
 Use: Hematinic.
CO-GA-TAR (Texas Pharm) Coal tar in 5% & 10% concentration, w/zinc oxide & corn starch liq 100 Gm; Oint 5% 1 lb.; Oint 10% 1 lb
 Use: Topically; antiseptic
CO-GEL (Arco) Amylolytic enzyme 3 mg; proteolytic enzyme 0.6 mg, cellulolytic enzyme 0.2 mg, w/calcium carbonate 300 mg, glycine 85 mg / Tab Bot 100s
 Use: Hyperacidity
COGENAT, (National Drug) Conjugated estrogens Tab (0.625 mg & 1.25 mg) Bot 100s, 1000s
 Use: Estrogen therapy
COGENTIN, NND 1962 (Merck Sharp & Dohme) Benzotropine Methanesulfonate Tropine Benzohydril Ether Methanesulfonate Tab (2 mg scored) Bot 100s, 1000s (1 mg / cc) Amp 2 cc, Box 6s
 Use: (0.5 2 mg daily) Parkinsonism
COGESIC, (Mead Johnson) Proclidine Hydrochloride
 Use: Analgesic
COGESIC-16 (Pacific Sales) Codeine phosph 1/2 gr, aspirin 3 1/2 gr, phenacetin 2 1/2 gr-caffeine 1/2 gr w/citric acid and sod bi carbonate/2 tabs Bot 16s
 Use: Analgesic
CONEMA, (Sull ff & Case) Ferrous sulfate 0.2

- Gm., manganese 5 mg., copper 0.2 mg., molybdenum 0.2 mg., cobalt 0.5 mg., zinc 1 mg., Vit. B-1 1 mg., niacinamide 10 mg., folic acid 0.3 mg., Vit. B-12 1 mcg./Tab. Bot. 100s, 1000s
 Use: Nutritional supplement
- COHISTINE**, (Pitman-Moore) Thénylpyramine HCl 25 mg., d-amphetamine phosphate 1 mg., aspirin 3½ gr., acetophenetidin 2½ gr./Tab. Bot. 100s, 1000s.
 Use: (1-3 daily) Antihistamine therapy.
- CO-HYDELTRA**, (Merck Sharp & Dahme) Prednisolone 2.5 mg. or 5 mg., w/magnesium trisilicate 50 mg. & aluminum hydroxide gel 300 mg./tab. Bot. 30s, 100s, 1000s
 Use: 1 (q.i.d.) Hormone therapy.
- CO-IRON**, (Haug) Ferrus sulfate dried 100 mg., cobalt chloride 10 mg., Mang. sulfate 15 mg./Tab. Bot. 100s.
 Use: Hematonic.
- COLACE**, N N O. 1962. (Mead Johnson) Diacetyl sodium sulfosuccinate. Cap. (30 mg. or 100 mg.) Bot. 30s. Liq. 1%, Bot. 30 cc. Syr. (20 mg / 5 cc.) Bot. 8 oz.
 Use: Orally, fecal softener.
 W/Cascara glycosides. See: Peri-Coloce.
- COLACTIN**, (Hillard Labs.) Scopolamine methylbromide 0.93 mg., guar gum 400 mg., diacetyl sod. sulfosuccinate 20 mg./Cap. Bot 30s.
 Use: Functional constipation.
- COLAGYN**, (Smith Lab. Inc.) Zinc sulfocarbonate, pot. oxyquinoline sulfate, lactic acid and boric acid. Jelly. Tube w/applicator & refill 6 oz.
 Use: Contraceptive.
- COLASCORE**, (O. F. Harvey) Colchicine 1/250 gr., ascorbic acid 15 mg. Sod salicylate 5 gr./tab. Bot. 1000s, 5000s.
 Use: Analgesic.
- COLAZINE SPENSOID**, (Drug Specialists) Sulfamethoxine, sulfadiazine, sulfamerazine, sulfacetamide, 0.125 Gm./5 cc. Bot. Ps
 Use: Infections responsive to sulfa therapy
- COLBAN**, (Mentholatum) Vit. C w/prophenpyridamine, phenacetin, caffeine and salicylamide. Bot. 35s.
 Use: Common cold.
- COLBENEMID**, (Merck Sharp & Dahme) Benemid 0.5 Gm., colchicine 0.5 mg /Tab Bot. 100s.
 Use: Chronic gout and gouty arthritis.
- COLBRON**, (Ciba) Ferrus sulfate, dried 2 gr., thiamine HCl 0.3 mg., liver concentrate 2½ gr./Cap. Bot. 100s.
 Use: Anemias.
- COLCAP**, (Arden) Pyrilamine maleate 12.5 mg., chlorpheniramine maleate 2 mg., phenylpropanol hydrochloride 5 mg., phenylephrine hydrochloride 2.5 mg., glyceryl guaiacolate 100 mg., salicylamide 150 mg., Vit. C 150 mg./Cap. Bot. 100s.
 Use: Relief of common cold.
- * **COLCHICINE**, (U.S.P.) Tab., U.S.P. (Various Mfr.)
 Abbott—Tab. 1/100 gr. Bot. 100s, 1000s.
 Lilly—Tab. 1/100 gr. Bot. 100s, 1000s.
 Merck Sharp & Dahme—Tab. 1/100 gr. Bot. 100s, 1000s.
 Use: Acute gout.
 W/Acetaminophen, Zoxazolamine See: Triurate, Tab. (McNeil)
 W/Aspirin, cimicitugin & phthalocin. See: R.M. Cole, Cap. (Cole)
 W/Aspizin & neocinchophen. See: Neo-Salicyphen, Cap. (Abbott).
 W/Benemid. See: Calbenemid (Merck Sharp & Dahme)
 W/Methyl salicylate. (Lilly) Colchicine 1/250 gr., methyl salicylate 3 min./Glabule. Bot. 100s, 1000s.
 Use: Orally, gout therapy.
 W/Methyl salicylate. (Parke, Davis) Colchicine 1/250 gr., methyl salicylate 3 min./Cap. Bot. 100s, 500s, 1000s.
 Use: Orally, gout therapy.
 W/Methyl salicylate. See: Colchi-Sol, Cap. (Huxley).
 W/Salicylamide.
 See: Chylamin, Tab. (Elder)
 Solamide w/Colchicine, Tab. (Columbus)
 W/Salicylamide, p-aminobenzoic acid, Vit. C & organidin. See: Aramide W/Colchicine, Tab. (Wampole Labs).
 W/Salicylamide & ascorbic acid. See: Asal-cine, Tab. (Lemmon)
 W/Sod. salicylate & ascorbic acid.
 See: Cinbisol, Tab. (McNeil).
 Colosorb, Tab. (G. F. Harvey)
 Mincalot, Tabs. (Testagar)
 Salcarbina, Tab. (Chicago Pharm.).
 W/Sod. salicylate, p-aminobenzoic acid & ascorbic acid. See: P-B-Sal C w/Colchicine, Tab. (Ulmer)
 W/Sod. salicylate, Sod. p-aminobenzoate & vitamins. See: Sol-Zem No. 2, Tab. (Zemmer)
 See: P-B-Sal w/Colchicine, Tab. (Ulmer)
 Solcal, Tab. (Taylor).
 W/Sod. salicylate & pot. iodide.
 See: Colsolide, Tab. (Chicago Pharm.).
 Solitum, Tab. (C. D. Smith).
 W/Sod. salicylate, Sod. iodide & Sod. cacodylate See: Sol-Sa-Col, Tab. (Breen).
 W/Strontium salicylate & methyl salicylate.
 See: Stronicaps, Cap. (Elder).

* Generic name recognized by N.P., U.S.P. or A.M.A.

W/Zoxazolamine acetaminophen
See Triurate Tab (McNeil)

COLONICINE SALICYLATE
W/Aspirin See Formosal Tab (First Texas)
W/Paba, aspirin, thiamin, ascorbic acid & niacin See Acetylcol Tab (Warner Chilcott)
W/Phenobarbital Sod p aminobenzoate
Vit B 1 B aspirin See Dalarol, Tab (Wolff Pharm Co)

COLONICINE SALICYLATED.
W/Aspirin, acetophenetidin & caffeine See Far Dyne Cap or Tab (First Texas)
W/Salicylamide, thenylpyrimine HCl, atropine sulfate & acetophenetidin See FCAH, Cap (First Texas)

COLCHI-SAL. (Huxley) Colchicine 1/250 gr w/methyl salicylate/Cap Bot 25s, 100s 500s
Use: Orally, analgesic & antirheumatic in gout therapy

COLDEX NO 2 Elix (Tyler) Phenylephrine HCl, 5 mg; prophepyridamine malate 12.5 mg; chloroform 13.5 mg; 1 menthol, 1 mg/Trp Bot 4 oz, Pls Gals
Use: Respiratory congestion

COLD VACCINES, INJECTABLE. Mollister Ster, Vial mixed 5 cc & 20 cc Sherman, Vial 12.5, 20 & 50 cc
See: Immunovac Cold Vaccine (Parke Davis) Pneumo-Strepto Vaccine (National Drug) Respiratory Vaccine (National Drug)

COLD VACCINES, ORAL.
Sherman—Oral Tabs
See: Entoral Pulvules (Lilly) Immunovac Tab (Parke Davis) M.V.R.I., Vial (Cutter) Neovaccine (Merck Sharp & Dahme) Oravax Tab (Merrill) Shervac Tab (Sherman) Vaccogen Tab (Merck Sharp & Dahme)

COLEBINZ. (Cole) Benzyl benzoate lotion Bot 2 oz, 4 oz, 1 pt
Use: Topically scabicide & treatment of fungus infections

COLINE. (Warren Teed) Phenolphthalein 1/2 gr, as bile ext 1 gr, cascara ext 1/2 gr, chionanthus 1/2 gr & Sod biphosphate 4 gr./Tab Bot 100s
Use: (2 hr) Choleric & chologogue

COLESITIN. (Victor Lemon) See: Lactishe Complex Plus

COLESITIN, FORTIFIED. (Victor Lemae) Pow Fig 9 oz, 3R oz

COLEY'S MIXTURE. (Parke, Davis) Toxic of strains of *Erysipelas streptococcus* B B prodigious Vials 1 & 20 cc.
Use: (I.M.) 0.03 cc followed by 1 cc intra-venously IV or I.M.

COLIBETA (Columbi) Vit B 12 w/intrinsic

factor concentrate 1/6 u, ferrous gluconate 165 mg, liver desic 100 mg, folic acid 0.5 mg, thiamine mononitrate 1 mg, riboflavin 1 mg, niacinamide 7.5 mg, Vit C 20 mg, w/cobalt, copper mong, zinc molybd/Cap Bot 100s, 500s 1000s
Use (1-3 daily) Nematonic

COLICELL. (Sutliff & Case) Whole bile ext 227 mg, ketocholanic acids 97 mg, methyl cellulose 130 mg, homatropine methyl bromide 2.5 mg/Tab Bot 100s, 1000s
Use Laxative w/antispasmodic

COLIMYCIN SODIUM METHANESULFONATE. (Warner Chilcott) See Colistimethate Sod

COLIMYCIN SULFATE. (Warner Chilcott) See: Coly Mycin

***COLISTIMETNATE SODIUM,** Sod Colistin Methansulfonate See Coly mycin injectable (Warner Chilcott)

COLISTIN BASE, W/Neomycin base, hydrocortisone acetate, thonzonium bromide polysorbate 80, acetic acid, sodium acetate See Coly Mycin Otic W/Neomycin and Hydrocortisone, (Warner Chilcott)

COLISTIN SULFATE. The sulfate salt of an antibiotic substance elaborated by *Aero bacillus colistinus* See Colimycin Sulfate (Wasees Chilcott)

COLITONE TABLETS. (Bowman) Dried yeast 200 mg, Vit B 12 w/intrinsic factor 1/20 v, pectin 250 mg/Tab Bot 100s, 500s 1000s, 5000s
Use: Digestion aid

COLLODAURUM. (Kohlenberg) Aurum colloid 0.13 mg/0.62 cc Bot 1 oz; Amp 5 cc & 2 cc Box 12s
Use (IV, topically) Palliative in malignancy

COLLOIDAL ALUMINUM HYDROXIDE, USP See: Aluminum Hydroxide Gel, USP
COLLOIDAL IRON COMPOUND. (Hille) Iron, copper, mang, zinc, iodine, Col pol-phos, mag, Sol Bot 12 fl oz
Use: Dietary mineral supplement

COLLOIDAL SILVER CHLORIDE, NF VIII NNR 1950 See: Lunasol (Hille)

COLLOIDAL SILVER IODIDE, NF, VIII, NNR 1950 See: Neo Silvol (Parke, Davis)

COLLO-SUL CREAM. (Crookes Barnes) Colloid sulfur 2% in greaseless emulsion base Jars 2 oz, 1 lb
Use: Acne and seborrhea

COLLO-SUL-HC. (Crookes) Sulfur 2%, Hydrocortisone acetate 1% Cream, Jar 1/2 oz
Use: Seborrheic dermatitis

COLLYRIUM. (Wyeth) Isotonic neutral borate

* Generic name recognized by N.F., U.S.P. or A.M.A.

sol. w/antipyrine 0.4%. Bot. 7 fl. oz. w/eye cup, 4 fl. oz. w/eye cup.
Use: Eye lotion.

COLLYRIUM, WITH EPHEDRINE. (Wyeth) Neutral isotonic sol. of boric acid & borax, containing 0.4% antipyrine & 0.1% ephedrine. Bot. 4 fl. oz. w/dropper.
Use: Eye lotion.

COLO-BAR. (Faugera) Concentrate of egg yolk in stable blend of vegetable fats with chocolate flavor. Each bar provides 140 calories, protein 6.7%, fat 67.8%, carbohydrate 24.5%./Bar. Box 12s.
Use: Gallbladder Evacuant.

COLOGEL. (Lilly) Methylcellulose 9 Gm./100 cc. Sol. Bot. 1 pt.
Use: (2-6 tsp. daily) Constipation.

COLOMANG. (Crookes) Mang 0.25% as colloidal hydroxide. Amp. 1 cc. & 2 cc. Box 6s, 25s, 100s. Vials 30 cc.
Use: Acne & furuncles.

COLON VACCINE. (Sherman) E. Coli 200 million, pneumococci 200 million, streptococci mixed 200 million, Staph. albus 300 million, Staph. aureus 300 million/cc. Vial 12.5, 20 cc. & 50 cc.
Use: Mixed bacterial infections.

COLOPO. (Cornrick) Bile salts 1 gr., cocca soda ext. 2 gr./Tab. Bot. 40s, 1000s.
Use: (1-2 h.s.) Choleraic & chelagogue

COLOVAC. (McNell) Yellow phenolphthalein 0.1 Gm., mag. hydroxide 0.3 Gm./Tab Bot. 100s, 1000s.
Use: Laxative.

COLPOTABS. (Purdue Frederick) Tyrothricin 5 mg, phenylmercuric acetate 3 mg, chlorophyll "a" 10 mg, Sod lauryl sulfate 7.5 mg, & beta-lactose Veg. Tab. Box 12s.
Use: Vaginal infections.

COLPROSTERONE. (Ayerst) Progesterone Vaginal Tab, 25 mg. silver foil and 50 mg. gold foil. Box 30s. "NS" Boxes of 15 w/applicator.

COLREX. (Rawell) Chlorpheniramine maleate 2 mg., phenylephrine HCl 5 mg., Vit C 100 mg., hesperidin 100 mg., aspirin 300 mg./Cap. Bot. 16s, 250s, 1000s.
Use: Common cold.

COLREX COMPOUND. (Rawell) Caffeine phosphate 16 mg, papaverine HCl 16 mg, aluminum aspirin 300 mg., phenylephrine HCl 5 mg, chlorpheniramine maleate 2 mg., vit. C 100 mg./Cap. Bot 100s.
Use: Relief of symptoms from common cold

COLSALIDE. (Chicago Pharmacol) Sod salicylate 5 gr., pat. lalide 1 gr., colchicine 1/150 gr./Tab. Tab. enteric coated. Bot. 100s, 1000s, 6000s.
Use: Gout & gouty arthritis & neuritis.

COLSARGEN. (Craikes-Barnes) Aq colloidal

isotonic susp. of metallic silver 0.05% Susp Bot. 4 fl. oz.
Use: Silver antiseptic.

CO-LU-GEL. (Ulmer) Aluminum hydroxide gel. Liquid, Bot. 1 pt., 1 gal.; Tab. 10 gr., Bot. 100s, 500s, 1000s.
Use: Antacid.

CO-LU-GEL M.T. LIQUID. (Ulmer) Co-Lu-Gel liquid with magnesium trisilicate 7.3%. Bot. 1 pt., 1 gal.
Use: (1-2 tsp.) Antacid.

COL-VI-NOL. (Ulmer) Vit. D 3600 u., w/phenol & benzocaine/ox. Oint. Jar or Tube 1 oz. Jar 1 lb.
Use: Topically, local anesthetic ointment.

COLY-MYCIN INJECTABLE. (Warner-Chilcott) Colistimethate sodium 150 mg., dibucaine HCl 8 mg, sodium citrate 50 mg, citric acid 10 mg/Vial. Vials 150 mg.
Use: Urinary tract infections.

COLY-MYCIN OTIC W/NEOMYCIN AND HYDROCORTISONE. (Warner-Chilcott) Colistin base as the sulfate 3 mg., neomycin base as the sulfate 3.3 mg, hydrocortisone acetate 0.5 mg., thonzonium bromide, polysorbate 80, acetic acid, sodium acetate/cc 5 cc. w/dropper.
Use: Acute and chronic external otitis.

COLY-MYCIN PEDIATRIC SUSPENSION. (Warner-Chilcott) Colistin base as sulfate 300 mg./70 ml. Sol. 70 ml.
Use: Acute bacterial diarrhea.

COMATENE. (New Mexico Pharm Co.) Choline Bitartrate 200 mg., Inositol 100 mg, dl-methionine 100 mg./Cap. Bot. 100s, 1000s.

COMBADINE. (Ives-Cameron) Sulfathiazole 0.166 Gm., sulfadiazine 0.166 Gm., sulfamerazine 0.166 Gm./Tab. Bot. 100s, 1000s.
Use: Sulfonamide susceptible infections.

COMBEVITA CAPSULES. (Merck Sharp & Dohme) Thiamine HCl 1 mg., liver B4 mg., riboflavin 1 mg., pyridoxine HCl 0.1 mg., nicotinamide 10 mg., cal. pantothenate 0.1 mg./cap. Bot. 100s, 500s.
Use: (3 caps. daily) Dietary supplement.

COMBEX W/Vitamin C Capsules. (Parke, Davis) Vit. B-1 10 mg., B-2 10 mg., B-12 1 mcg, nicotinamide 10 mg., pantothenic acid 3 mg, Vit. C 50 mg. liver concentrate D17 Gm, liver fraction No. 2 0.17 Gm./Cap. Bot. 100s, 500s, 1000s.
Use: Vitamin deficiency.

COMBEX INJECTABLE. (Parke, Davis) Vit. B-1 10 mg., Vit B-2 0.33 mg, nicotinamide 10 mg., Sod pantothenate 4 mg, Vit. B-12 2 mcg./cc. Vial 10 cc.
Use: Vitamin B complex.

COMBEX PARENTERAL STERI VIALS. (Parke, Davis) Vit. B-1 10 mg., B-2 0.22 mg, nico-

Inamide 10 mg, pantothenic acid 4 mg, B 12 as B mcg B 12, crystallin and liver extract for parenteral use 10 mcg/cc 10 cc vial

Use: Vitamin B complex deficiency

COMBEX KAPSEALS. (Parke, Davis) Vit B 1 10 mg, Vit B 2 0.33 mg, nicotinamide 10 mg, pantothenic acid (Sod salt) 4 mg, liver ext for parenteral use equiv Ia Vit B 12 2 mcg/cc Vial 10 cc Box 12s

Use: Vitamin B complex deficiency

COMBEX THERAPEUTIC KAPSEALS (Parke, Davis) Thiamine HCl 25 mg, riboflavin 15 mg, nicotinamide 100 mg, folic acid 2.5 mg, pyridoxine HCl 1 mg, pantothenic acid (as Sod salt) 10 mg, ascorbic acid 150 mg/Kapsel Bot 100s, 1000s

Use: Vitamin B complex therapy

COMBICHOLE. (F Traut) Dehydrocholic acid 2 gr, desoxycholic acid 1 gr/Tab Bot 100s, 1000s

Use: (12 hr) Stimulate bile flow

COMBIO. (Smith Kline & French) Darbid (isopropamide) as the iodide 5 mg, Compazine (prochlorperazine) as the dimaleate 10 mg/Sponsule Bot 30s 250s

Use: Gastrointestinal disorders

COMBIGEL (Diamond Labs) Succinylsulfathiazole 45 gr neomycin sulfate 142 mg mepiperphenidol 10 mg, kaolin 80 gr pectin 4 gr, Neojel 5 gr/Bot 6 fl oz

Use: Treatment of diarrhea

COMBIMYCIN. (CMC) Dihydrostreptomycin base 0.5 Gm, streptomycin base 0.5 Gm as the sulfates Vial, 1 Gm,

COMBINACE (Mead Johnson) Calcium & Sod alginates 750 mg, diactyl Sod succinate 50 mg, anthraquinone derivatives (Perishin) 30 mg/Tab or gran Tab Bot 30s, 60s Gran Bot 147 Gm, 294 Gm

Use: Constipation due to hard dry stool, inadequate bulk or motility

COMBINED BACTERIAL VACCINE (Parke Davis) E Cali 200 million pneumococcus 200 million streptococcus mixed 100 million, staph albus 500 million staph aureus 500 million/cc Vial 5 cc

Use: Mixed bacterial infections

COMBINED CARBONATES COMPOUND. (Rorer) Bismuth subcarb 28.5%, mag carb 62.5% & cerium oxalate 9% Pow, dispensing box

Use: Gastric hyperacidity

COMBINED DIPHTHERIA TETANUS. Taxovax N.N.R. 1954 (Parke, Davis, Lilly) Vial 1.5 cc, 1 Immunization; 7.5 cc, 5 Immunizations

COMBIOTIC. (Pfizer) Cryst procaine penicillin G 300,000 u, Sod penicillin G 100,

000 u, dihydrostreptomycin sulfate 0.5 Gm/dose Vial 5 doses

Use: Fortified penicillin with streptomycin therapy

COMBIOTIC P/S. (Pfizer) Procaine penicillin G 300,000 u w/penicillin G Sod 100,000 u w/dihydrostreptomycin 1 Gm as sulfate/1 cc (dry) Vial 1 cc & 5 cc. Also available w/dihydrostreptomycin 0.5 Gm/1 cc.

Use: Penicillin-Streptomycin therapy

COMBISTIX. (Ames) Protein test tetrabrom phenol blue, citrate buffer, protein-adsorbing agent; glucose test area glucose oxidase, orthotolidin and a catalyst; pH test area methyl red and bromthymol blue Box, strips, 125s

Use: Diagnostic agent

COMBISTREP. (Pfizer) 1/2 Streptomycin and 1/2 dihydrostreptomycin Vial 1 Gm & 5 Gm

COMBISUL. (Schering) Sulfadiazine 0.166 Gm, sulfathiazole 0.166 Gm, sulfamerazine 0.166 Gm/1 cc or Tab, Bot 100s, 1000s Susp Bot, 4 oz, 1 pt

Use: Triple sulfanamide therapy

COMBITAL. (Rond) Secobarbital 3/4 gr, butalbarbital 1/2 gr, phenobarbital 1/4 gr/Tab Bot 100s

COMBOGENIC. (Smith, Miller & Patch) beta Estradiol 0.20 mg, methyltestosterone 10 mg, thiomine HCl 10 mg, niacinamide 100 mg, riboflavin 10 mg, pyridoxine HCl 5 mg, Vit B 12 5 mcg, pantothenol 5 mg/Tab Bot 30s, 100s

Use: Menopause & dysmenorrhea

COMEDAB STICK. (Dome) Sulfur 2%, resorcinol 1%, hexachlorophene 1%, in a lubricating base/Stick Pkg 5.5 Gm

Use: Acne

COMEDOME STICK. (Dome) Colloidal sulfur 1%, hexachlorophene 1%, in a detergent base/Gm

Use: Acne

COMFORTINE. (Rorer) Calamine, zinc oxide boric acid Oint Tube 1 1/2 oz Jar 25 Gm, 100 Gm

Use: Antiseptic & protective ointment

COMIN. (Cole) Vit B 1 1.5 mg, B 2 2 mg, B 6 0.1 mg, niacinamide 10 mg, Cal pantothenate 1 mg, liver, yeast/Cap Bot 100s

Use: Vitamin B complex

COMPANOL. (Winthrop Labs) Scopolamine HBr 1/300 gr, atropine sulfate 1/400 gr laminal 1/2 gr/tab Box 12s Bot 50s, 500s

Use: Motion sickness

COMPAZINE. N.N.D. 1962 (Smith, Kline & French) Prochlorperazine Tab 5 mg, 10 mg & 25 mg Bot 50s 500s Amp. (5 mg/ cc) 2 cc Box 6s, 100s

- Concentrate**, 10 mg./cc., 4 oz. 12s, 3ds. Spacetate 10 mg., 15 mg. & 30 mg. Bot. 30s, 250s. 75 mg. Bot. 30s. Supp. (2.5 mg., 5 mg., 25 mg.) Box 6s. Syr. (5 mg./5 cc.) Bot. 4 oz. Vial. (30 mg.) 10 cc. Box 20s, 100s.
- W/d-amphetamine. See: Eskatrol, Cap. (Smith Kline & French)
- W/Darbid. See: Cambid, Spanzule. (Smith Kline & French)
- Use: Treatment of mental & emotional disturbances; also nausea & vomiting.
- COMPIRIN**. (Marco) Aminopyrine 3 gr., acetophenetide 1½ gr., caffeine ½ gr., phenobarbital 1/6 gr., mag. oxide & aromatics/Tab. Bot. 100s, 1000s.
- Use: (1-2 a. 3-4 h.) Analgesic with sedative.
- COMPLEXIN**. (Estra) Thiamine HCl 10 mg., riboflavin 0.1 mg., pyridoxine HCl 1 mg., niacinamide 10 mg./cc. Amp. 1 cc. Vial 10 cc. & 30 cc.
- Use: Vitamin B deficiencies.
- COMPLEXO-B**. (Mallord) Vit. B-1 25 mg., B-2 1 mg., B-6 1 mg., nicotinamide 50 mg., panthenol 1 mg./cc. Vial 30 cc.
- Use: Vitamin B therapy.
- COMPLIGEN**. (Pitman Moore) Aluminum phosphate adsorbed diphtheria toxoid, tetanus toxoid, poliomyelitis vaccine and pertussis vaccine/Vial 9 cc.
- Use: Primary immunisation of infants and children.
- COMPOCILLIN, N.N.B. 1957**. (Abbott) Nydrabamine penicillin O.
- COMPOCILLIN-V, Oral Susp., N.N.D. 1962**. (Abbott) Nydrabamine Penicillin V 300M u. (180 mg)/5 cc. Bot. 40 cc. & 80 cc. Granules.
- COMPOCILLIN-YK, FILMTABS, N.N.D. 1962**. (Abbott) Penicillin V Pat. 200M u./Tab. Bot. 50s, 100s, 400M u./Tab. Bot. 25s, 100s.
- COMPOCILLIN-YK W/SULFAS**. (Abbott) Pot. penicillin V 125 mg., sulfadiazine 167 mg., sulfamerazine 167 mg., sulfamethazine 167 mg./Filmtab. Bot. 50s.
- Use: Mixed infections.
- COMPOCILLIN-YK GRANULES**. (Abbott) Pot. penicillin-V 200,000u. (125 mg./tsp. Bot. 40 & 80 cc.
- Use: Oral penicillin therapy.
- COMPOCILLIN-YK-SULFAS GRANULES**. (Abbott) Potassium penicillin V 125 mg., sulfa-meth-dia-mer each 167 mg./5 ml. Bot. 60 ml.
- Use: Mixed infections.
- COMPOMAL**. (Ives-Cameron) Scopolamine HBr 0.22 mg., atropine sulfate 0.16 mg., phenobarbital 32 mg. Tab. Box 12s. Bot. 50s, 500s.
- Use: Motion sickness.
- COMPOUND 42**. See: Warfarin. (Various Mfr.)
- COMPOUND B**. See: Corticosterone. (Various Mfr.)
- COMPOUND E** (Kendall E) See: Cortisone Acetate. U.S.P.
- COMPOUND F**. See: Hydrocortisone. (Various Mfr.)
- COMPREN**. (Lilly) Vit. A 6,000 u., D 400 u., B-1 1.5 mg., B-2 3 mg., B-12 7.5 mcg., intrinsic factor conc. ½ oral u., C 100 mg., nicotinamide 15 mg., lactic acid 0.4 mg., iron 15 mg., B-6 6 mg., cal. pantothenate 4 mg., calcium 750 mg./3 pulv. Bot. 100s, 300s, 1000s.
- Use: Prenatal supplement.
- COMTEX**. (Estra) Thiamine HCl 5 mg., riboflavin 1 mg., pyridoxine HCl 1 mg., niacinamide 1 mg., Cal. pantothenate 10 mg., benzyl alcohol 2%/cc. Amp. 1 cc. Vial 10 cc. & 30 cc.
- Use: Vitamin B therapy
- COMYCIN**, (Upjohn) Tetracycline phosphate equiv. to tetracycline HCl 250 mg., nystatin 250,000 u./Cap. Bot. 16s, 100s. Half strength (125 mg.) Bot. 25s, 100s.
- Use: Tetracycline therapy and where non-fatal infections might be encountered
- COMYCIN HALF-STRENGTH**. (Upjohn) One-half formula of comycin. Bot. 25s, 100s.
- CONALSYN**. (H. E. Cenci) Chlorpheniramine maleate 8 mg., phenylephrine HCl 20 mg., methscopolamine nitrate 2.5 mg./Cap. Bot. 30s, 250s
- Use: Hay fever and similar allergies.
- CONAR**. (Massengill) Narcotine 10 mg., pheniramine maleate 10 mg., phenylephrine HCl 5 mg., w/chloroform, menthol/5 cc. Bot. 2 oz., 1 pt.
- Use: Aesthetic in the common cold
- CONAR EXPECTORANT**. (Massengill) Nascopine 10 mg., pheniramine maleate 10 mg., phenylephrine HCl 5 mg., glyceryl guaiacolate 100 mg., chloroform 12 mg., menthol 0.2 mg./5 cc. Bot. pt.
- Use: Relief of coughs.
- CONAR TROCHE**. (Massengill) Nascopine 5 mg., pheniramine maleate 5 mg., phenylephrine HCl 2.5 mg., menthol 0.1 mg., aromatics/Troche. Vial 12s, Bot. 100s.
- Use: Coughs & nasal congestion.
- CONCEMIN B**. (Merrill) Thiamin HCl 2 mg., riboflavin 1 mg., niacinamide 5 mg., ferrous sulfate 0.24 Gm., pyridoxine HCl 0.5 mg., choline dihydrogen citrate 6.25 mg./Cap. Bot. 100s, 1000s.
- CONCENTRATED OLEOVITAMIN A & D, N.N.B. 1955**. See: Oleovitamin A & D, Concentrated. (Various Mfr.)

CONCENTRATED TUBERCULIN

198

- CONCENTRATED TUBERCULIN.** See: Tuberculin, Old, U.S.P.
- CONDOCAPS.** (Nian) Vit D 50 000 u In oil/cap Bot 100s, 500s
Use: Orally, vitamin D deficiencies
- CONDO.** (Nian) Calciferol 50M u / 5/2 cc. Bot 8 oz
Use: Vitamin D therapy
- CONESTRON, NND 1962** (Wyeth) Estrogenic substances, water soluble Tab 0.625 & 1.25 mg Bot 100s, 1000s
Use: Estrogen therapy
- CONEX-DA.** (Lloyd, Dobney & Westerfield) Phenylpropanolamine HCl 50 mg, phenyltoloxamine dihydrogen citrate 50 mg / Dual action Tab Bot 100s
Use: Orally, relief of nasal congestion
- CONEX PLUS.** (Lloyd, Dobney & Westfield) Phenylpropanolamine HCl 25 mg, phenyltoloxamine dihydrogen citrate 25 mg, acetyl p aminophenol 250 mg / Tab Bot 100s
Use: Orally, analgesic and relief of nasal congestion
- CONEX SYBUP.** (Lloyd, Dobney & Westfield) Phenylpropanolamine HCl 25 mg, phenyltoloxamine dihydrogen citrate 25 mg, acetyl p aminophenol 250 mg, pot guaiaacolsulfonate 110 mg / 10 cc Bot 1 pt
Use: Treatment of cough due to colds
- CONFETS** (Schenley) Cryst penicillin G pot buffered 5000 u / Tab Tubes 12s
Use: Penicillin sensitive organisms
- CONGENS** (Blaine) Water soluble conjugated estrogens from pregnant mares urine 1.25 mg / Tab Bot 100s, 1000s
Use: Control of menopausal symptoms and senile vaginitis
- CONGO BED,** Injection
Use: Hemostatic in hemorrhagic disorders
- CONJUGATED ESTROGENS, NNR** See: Estrogens, Conjugated (Various Mfr)
- CONOVID** (Searle) The trademark in Britain for Enovid
- CONRAY.** (Mollinckradt) (meglumine sothalamate 60%) 0.6 Gm / cc Iothalamate Acid
Use: Radiographic contrast medium
- CONSDBIN HCl** (DuMont) Phenylephrine HCl 1/4 & 1% Bot 1 pt
- CONSOL.** (Merrill) Narcotine (Noscapine) 5 mg terpin hydrate 50 mg Doxylamine succinate 3 mg, menthol / 5 cc Bot 4 oz, 1 pt
Use: Treatment of cough due to colds
- CONSOLETS.** (Merrill) One troche equals 5 cc. of Consol Pk 24s
- CONSTONATE.** (Nysco Labs) Dicotyl sod sulfosuccinate 20 mg / cap Bot 100s 1000s
Use: Fecal softener
- CONSOLID.** (Ciba Pharm Prod, Inc.) See: Sulfachlorpyridazine
- CONTACT.** (Menley & James) Belladonna alkaloids 0.25 mg; phenylpropanolamine HCl 50 mg; chlorpheniramine maleate 40 mg / spansule Bot 12s, 20s
Use: Cold
- CONTEBIN.** See: p-acetylamino benzaldehyde thiosemicarbazone
- CONTNEX.** (Ingram) Ferrus gluconate 75 mg, folic acid 10 mg, thiamin HCl 15 mg, riboflavin 1 mg, niacinamide 15 mg / cc. Dropper Bot 30 cc
Use: Treatment of anemias
- CONTBA CREME, NNR 1955** (Cantrol) Phenylmercuric acetate 0.06%, cream base
Use: Contraceptive
- CONTRAMAL-CP.** (Physicians Prods) Acetyl p aminophenol 325 mg, salicylamide 225 mg, caffeine 30 mg, phenylephrine HCl 5 mg, trisamine 20 mg / Cop. Bot 100s.
Use: Analgesic.
- CONVALETS FILMTABS.** (Abbott) Vit B 1 10 mg, B 2 10 mg, B-6 2 mg, C 300 mg, B 12 4 mcg nicotinamide 100 mg, Cal pantothenate 20 mg, folic acid 1.5 mg / Tab Bot. 25s, 100s, 500s
Use: Dietary supplement
- CONVANAMINE EXPECTOBANT.** (Yanpelt & Brown) Phenylephrine HCl 375 mg, phenylpropanolamine HCl 625 mg, chlor pheniramine maleate 1 mg, pyrilamine maleate 625 mg, glyceryl guaiaacolate 100 mg / Tab Bot 100s
Use: Colds
- CONVERTIN.** (Ascher) Betaine HCl 130 mg; & ginger oleoresin 1/600 gr, surrounding enteric coated core of pancreatin, chlorophyllin 100 mg, desoxycholic acid 50 mg / tab Bot. 84s, 500s
Use: Promote apital digestion
- CONVEBTIN-H.** (B. F. Ascher) Hamotropine methylbromide 25 mg, betaine HCl 130 mg, ginger oleoresin 1/600 gr, pancreatin 625 mg, desoxycholic acid 50 mg / Tab Bot 84s, 500s
Use: Bladder dysfunction, liver disease postcholecystectomy syndrome
- CONVEBZYME** (Ascher) Proteolytic enzyme (papain) 10 mg, amylolytic enzyme 80 mg, homotropine methylbromide 25 mg, d sorbitol 4.4 Gm / 5 cc Bot pts
Use: Digestive disorders
- CONVEBZYME TABS.** (Ascher) Cellulolytic enzyme 5 mg, proteolytic enzyme 10 mg, amylolytic enzyme 30 mg, lipolytic enzyme 800 Ascher Units / Tab Bot 100s, 500s
Use: Improper digestion of food

COOK COLD TABLETS. (E. K. Cook) Bot. 25s, 100s.

Use: Treatment of common cold.

COOMASSIE BLUE. (Ayerst) Sodium Azobenzene.

Use: Agent for dye dilution studies of cardiac output.

COOPER CREME, N.N.R. 1953. (Whittaker) Trioxymethylene 0.04%, Sod. oleate 0.67%, Irihydroxyethylolene 7.91%, diethyl Sod. sulfasuccinate 0.5%, stearic acid 21.04%, aluminum silicate 2.34% & aromatics. Tube 75 Gm.

Use: Contraceptive.

* **COPARAFFINATE, N.N.D. 1962.** See: Iso-Par, Oint. (Medical Chem.)

COPAVIN. (Lilly) Codeine sulfate $\frac{1}{4}$ gr., papaverine HCl $\frac{1}{4}$ gr./Cap. Pulvules Bot. 100s, 500s, 1000s, Tab. 40s, 500s.

Use: Head colds with cough.

COPAVIN COMPOUND ELIXIR. (Lilly) Codeine sulfate $\frac{1}{2}$ gr., papaverine HCl $\frac{1}{2}$ gr., atoin $\frac{1}{4}$ gr., Sod. salicylate 20 gr./fl. oz. Bot. 1 pt., 1 gal.

Use: Head colds with cough.

COPIHE MIXIR. (G. F. Horvey) Codeine sulfate 1 gr., teipin hydrate 8 gr./fl. oz. Bot. 4 oz., 1 pt., 1 gal.

Use: Cough preparation.

COPLEXEN LIQ. (Smith Kline & French) Trimepazine 2 mg., acetaminophen 120 mg., phenylpropranolamine HCl 10 mg./5 cc. Bot. 12 oz.

Use: Cold complex in children.

COPPER ACITATE W/Iron cocodylate, sod. cocodylate. See: Clogan, Amp. (Endo)

COPPER CACODYLATE W/Iron cocodylate & Sod. cocodylate. See: Ferro-Cuprin, Amp. (Kramers-Urban).

COPPER CARBONATE. See: Cupran (Ayerst).

COPPER CAPRYLATE W/Caprylic acid salicylanide, p-chlor-m-xylanol. See: Pricutex, Oint. (Elder)

W/Sod. caprylate & caprylic acid. See: Capronex Oint. (Hartan & Conversa)

COPPER NEAD BITE THERAPY. See: Anivrenin, Snake Polyvalent.

COPPERIN. (Jewell) A—(Iron as Iron ammonium citrate 32 mg., copper as sulfate 1 mg./Cap.) B—(Iron as Iron ammonium citrate 32 mg., copper $\frac{1}{4}$ mg./Cap.) Bot. 100s, 500s.

Use: Orally (1 t.i.d., p.c.) secondary anemias.

COPPER SULFATE. See: Mancidene (Xtee)

COPPER UNDECYLENATE.

W/Sod. propionate, Sod. caprylate, propionic acid, undecylenic acid, salicylic acid & Sod.

* Generic name recognized by N.F., U.S.P. or A.M.A.

CORALLIN W/PHENOBARBITAL

diethylsulfosuccinate. See: Verdefam, Sal. (Texas).

W/Undecylenic acid. See: Kudelate, Oint. (Ulmer)

W/Zinc undecylenate, undecylenic acid & phenylmercuric acetate. See: Di-Spar, Oint. (Wintersmith).

COPRAL. (Brewer) Iron & ammonium citrate 0.75 Gm., liver ext. (1:20) 0.5 Gm., Vit. R-1 6 mg., B-2 4 mg., B-12 4 mcg., niacinamide 20 mg., red bone marrow 15 m./fl. oz. Bot. 8 oz., 1 pt., 1 gal.

Use: Anemias.

COPRANE. (Fuller) Salicylic acid 1%, camphor 1%, alcohol Bot. 8 oz.

Use: Antipruritic.

COPSAMINE, N.N.D. 1961. (Durst) Pyrilamine maleate. Tab. (25 mg.), Bot. 100s, 500s, 1000s.

Use: Orally, antihistamine.

CO-PYRONIL. (Lilly) Pyrabutamine 15 mg., Histodyl (phenylpyramine) 25 mg., Clapone HCl (cyclopentamine) 12.5 mg./Pulvula. Bot. 100s, 1000s.

Use: Allergic symptoms.

CO-PYRONIL PEDIATRIC. (Lilly) Pyronil 7 $\frac{1}{2}$ mg., Histodyl 12 $\frac{1}{2}$ Gm., Clapone HCl 6.25 mg./pulvula. Bot. 100s, 1000s.

Use: (1-2 t.i.d.) Allergic symptoms.

CO-PYRONIL SUSPENSION. (Lilly) Pyrabutamine 7.5 mg., Histodyl 12.5 mg., clapone HCl 6.25 mg./5 cc. Susp. Bot. 1 pt.

Use: Orally, antihistaminic.

CORALCAP-ASTHMA. (Coral). Aminophylline, 3 gr., ephedrine sulfate, $\frac{1}{2}$ gr., chlorpropenpyridamine maleate, 4 mg., phenobarbital, $\frac{1}{4}$ gr./Cap. Bot. 100s, 1000s.

Use: Bronchial asthma and hay fever.

CORAL COLD CAP. (Coral Tabs) No. 1, No. 2. Capsules. Bot. 100s, 1000s.

CORALET-T.D. (Crystal) Pentaerythritol tetranitrate 30 mg., butabarbital 50 mg./Cap. Bot. 100s.

Use: Hypertension.

COR-AL-GESIC. (Ulmer) Chloral hydrate 2%, methyl salicylate, menthol, Oint. Jar 1 lb.

Use: Analgesic and counterirritant.

CORALIXIN. (Crystal) Dihydroxypropyl theophylline 80 mg., glycerine 10%/5 cc. Bot. pt., gal.

Use: Treatment of cardiovascular disease

CORALLIN. (Crystal) Glyceryl trinitrate 0.4 mg., pentaerythritol tetranitrate 15 mg., neothylline 100 mg./Tab. Bot. 100s.

Use: Coronary dilator.

CORALLIN W/PHENOBARBITAL. (Crystal) Glyceryl trinitrate 0.4 mg., pentaerythritol

- tetranitrate 15 mg, neophylline 100 mg
phenobarbital 15 mg /Tab Bot 100s
Use Angina p and coronary insuf
- CORAMINE, NND 1962** (Ciba) Niketha
mide liq 25% Bot 1 oz, 3 oz, 1 pt Amp
25% 1½ cc & 5 cc Vial 20 cc
Use: Respiratory stimulant
- CORAVAR**, (Crystal) Ethaversion HCl 30 mg,
d hydroxypropyl theophylline 100 mg /Tab
Bot 100s
Use: Coronary vasodilator
- CORDAN CREAM** (Lilly) Flurandrenolone
0.5 mg; in water washable base/Gm Tube
7.5 Gm
Use: Dermatologic steroid
- CORDAN OINTMENT** (Lilly) Gm Flu
randrenolone 0.5 mg remulsified petrolatum
base/Gm Tube 15 Gm
Use: Dermatologic steroid
- CORDAN-N CREAM**, (Lilly) Flurandreno
lane (Cordran) 0.5 mg; neomycin sulfate
5 mg; in water washable base/Gm Tube
7.5 Gm, 15 Gm
Use: Dermatologic steroid
- CORDAN-N OINTMENT**, (Lilly) Flurandren
olone 0.5 mg; neomycin sulfate 5 mg;
emulsified petrolatum base/Gm Tube 7.5
& 15 Gm
Use: Dermatologic steroid
- CORDEX**, (Upjohn) Prednisolone 0.5 mg,
aspirin 5 gr/tab Bot 100s 300s Also
available as buffered Tabs (Cal carb 200
mg/Tab)
Use: Rheumatoid arthritis
- CORDEX FORTE**, (Upjohn) Prednisolone 1.5
mg, aspirin 5 gr/tab Bot 100s 300s Also
available as buffered Tabs (Cal carb 200
mg/Tab)
Use: Arthritis
- CORDAN** (Lilly) Flurandrenolone 0.5 mg/
Cream & Oint Tube 7.5 Gm and 15 Gm
Use: Topical corticoid compound
- CORDAN-N**, (Lilly) Cordran 0.5 mg, neo
mycin sulfate 5 mg./Cream & Oint Tube
7.5 Gm and 15 Gm
Use: Topical corticosteroids
- COR-D-TAR** (Der lin) Hydrocortisone alcoh
ol 1%, d odehydroxyquinoline 2%, coal
tar soap 2%, in a water-miscible base
Tube ½ oz, 1 oz
Use: Dermatitis
- CO-RE-OA POWDER**, ADR 1855 (Corege)
Karye gum 98186 Gm, Mag carb 0274
Gm, methyl salicylate, peppermint oil
menthol Rum resin/100 Gm Pkg ½, 1½
& 3½ oz
Use: Denture adhesive powder
- CORENOL** (McNeil) Chloral hydrate 2 mg,
barbitone est 8 mg, racemic dosage
- ephedrine HCl 125 mg /Tab Bot 100r
1000s
Use (12, 3-4 h) Antispasmodic with ant
histamine
- CORICIDIN**, (Schering) Chlor trimeton male
ate 2 mg, acetylsalicylic acid 3½ gr,
acetophenetidin 2½ gr, caffeine ½ gr/
Tab Tube 12s Bot 100s, 1000s
Use (14 daily) Common cold
W/Cocaine phosphate (Schering) Corcidin
w/cocaine phosphate ¼ or ½ gr/Tab.
Bot 100s
- CORICIDIN "D" DECONGESTANT**, (Scher
ing) Chlor trimeton maleate 2 mg, aspirin
0.23 Gm, phenacetin 0.16 Gm, caffeine
30 mg phenylephrine 10 mg/Tab Boxes
12s Cartons of 6 boxes Bot 100s
Use Nasal decongestant & treatment of
common cold
- CORICIDIN NASAL MIST**, (Schering) Chlor
Trimeton gluconate 3 mg, phenylephrine
HCl 5 mg, gramicidin 0.05 mg/cc Plastic
squeeze bot
Use Nasal decongestant
- CORICIDIN FORTE**, (Schering) (Corfaris)
Chlor trimeton maleate 4 mg, salicylamide
190 mg phenacetin 130 mg, caffeine 30
mg, ascorbic acid 50 mg, methamphetamine
HCl 125 mg/cap Bot 100s, 1000s
Use Orally, analgesic
- CORICIDIN PEDIATRIC MEDILETS** (Scher
ing) Chlor trimeton maleate 0.75 mg, as
pirin 80 mg, acetophenetidin 16 mg/
Medilet Bot 25s & 100s
Use: Antihistaminic with analgesic
- CORICIDIN-PENICILLIN SUSPENSION**,
(Schering) Pot penicillin G 250 m.u., chlor
trimeton 2 mg, Sod salicylate 112.5 mg/
5 cc. Buffered with Sod citrate Bot 60 cc.
at dry paw Ia which 42 cc of water is
added.
- CORICIDIN WITH PROCAINE PENICILLIN**
G. (Schering) Chlor trimeton maleate 2.0
mg, procaine penicillin G 150 000 u., as
pirin 0.15 Gm, acetophenetidin 0.12 Gm,
caffeine 0.02 Gm /Tab Bot 24s, 100s
Use: Penicillin with analgesic & antihis
taminic
- CORICIDIN, SYRUP**, (Schering) d Methorpan
NBr, chlortrimeton maleate 2 mg, Sod
salicylate 225 mg, Sod citrate 120 mg,
caffeine 30 mg, glyceryl guaiacolate 30
mg/5 cc. Bot 4 oz, 16 oz
Use: (1 dr q 3-4 hr) Antihistaminic cough
therapy
- CORICO**, (Hoag) Phenylephrine, chlorphe
namine, analgesics and Vitamin C. Coprules
Bot 100s
Use: Common cold
- CORIFORTE** See: Corcidin Forte

- CORILIN.** (Schering) Chlorphenylpyridomine maleate 0.75 mg., sod. salicylate 80 mg and glycine 25 mg./cc. Dropper bot. 30 cc. Use: Nasal congestion in infants.
- CORIZAHIST** (Mason Pharmaceuticals) d-Metharphan hydrabamide 60 mg., chlorpheniramine maleate 7.5 mg., ammonium chloride 500 mg., phenylephrine hydrochloride 30 mg./syr., phenylephrine hydrochloride 17 mg., pheniramine maleate 8 mg., pyrilamine maleate 8 mg., Vit C 50 Gm./Tab. Bot. 100s.
Use: Treatment of common cold
- CORLUTONE, N.N.D. 1962.** (Gold Leaf) Progesterone 10 mg./cc. in sesame oil. Vial 10 cc.
- CORMED.** See: Nikethamide. (Various Mfrs.)
- CORN OIL.** See: Lipomul-Oral (Upjohn).
- CORODENE** (Graff) Antihistamine, antitussive, two decongestants w/Acetyl p-aminophenol and a stimulant. Cap. Rot. 100s.
Use: Cold and influenza symptoms.
- CORON INJECTABLE.** (Savage) Peptonized Iron 100 mg., cobalt gluconate 9 mg./2 cc. Vials 100 cc.
Use: Treatment of iron deficiency anemia.
- COROVAS TYMCAPS.** (Amire-Grant) Penterythritol tetranitrate 30 mg., secobarbitalic acid 50 mg./cap. Bot. 60s, 120s, 500s, 1000s.
Use: 1 q. 12 h.; prolonged action coronary vasodilator.
- CORPA-GEN.** (Barry) Beta-estradiol 0.5 mg. & progesterone 5 mg./cc. Vial 10 cc.
Use: Threatened abortions.
- CORPARID.** (Sullin & Coe) Methamphetamine HCl 4.0 mg., d-amphetamine sulf. 2.0 mg., pentobarbital sod. 300 mg., Vit. C 50.0 mg., B-1 1.0 mg., B-2 1.0 mg., B-6 0.1 mg., B-12 0.2 mcg., cal. pantothenate 1.0 mg., niacinamide 3.0 mg./Cap. Bot. 100s, 1000s
Use: Vitamin supplement
- CORPERIDIN.** (Dumas-Wilson) Hesperidin methyl chalcone 100 mg., ascorbic acid 100 mg., menadiolone 0.75 mg./Cap. Bot. 100s, 1000s
Use: An aid in the maintenance of normal capillary function
Use: Control of eye inflammation.
- CORPLEX.** (Chicago Pharmacol) Bioflavonoid complex 100 mg., ascorbic acid 100 mg./Tab. Bot. 100s, 1000s
Use: To increase capillary resistance and decrease fragility
- CORPOMONE, N.N.D. 1961.** (Preme) Cryst. progesterone in oil. Vial (10 mg./cc.) 10 cc., (25 mg./cc.) 10 cc. Tabs. (10 mg.) Bot. 100s, 1000s, (25 mg.) Bot. 50s, 500s S.I. Tabs. (10 mg.) Bot. 100s, 1000s.
Use: Functional uterine bleeding.
- CORPORA LUTEA, N.N.R. 1931.** (Soluble extract) (Parke, Davis) Amp. 1 cc. Box 6s, 100s. Each cc. contains soluble ext. derived from 7 gr. fresh corpus luteum tissue.
Use: Corpus luteum preparation.
- CORPORA LUTEA, DESICCATED.** (Parke, Davis) Emplats 5 gr. Bot. 100s, Tablets 5 gr. Bot. 50s & 500s.
Use: Corpus luteum therapy.
- CORPUS LUTEUM, N.N.D. 1961.**
See: Corpora Lutea, Amp. & Tab. (Parke, Davis)
Hydroxyprogesterone Caproate Lutrexin, Tab. (Hynson, Wescott & Dunning)
- CORPUS LUTEUM EXTRACT, N.N.R. 1931.**
See: Lutein, Ext. Amp. (Hynson, Wescott & Dunning)
- CORPUS LUTEUM, EXTRACT (WATER SOLUBLE).**
National Drug—Ext. from 1.17 Gm. of corpus luteum tissue, Amp. 1 cc., Box 12s, 25s, 100s; Injectosal 25 cc.
Parke, Davis—Ext. from 0.454 Gm. of corpus luteum tissue, Box 6s, 100s.
- CORRECTIVE Jr.,** (Jenkins Labs) Ipecac pow. 1/32 gr., calomel 1/16 gr., bismuth subnitrate 2 gr., sod. bicarbonate 1 gr., pancreatin 9/40 gr., pepsin 1/20 gr., papain 9/40 gr./tab. Bot. 100s, 1000s.
Use: Laxative, emetic & digestive
- CORRECTOL.** (Pharmaco) Diethyl sod. sullosuccinate 100 mg., phenolphthalein 1 gr./Tab. Bot 30s
Use: Fecal softener and laxative
- CORTACET.** (Hart) Desoxycorticosterone acetate 5 mg./cc. Vial 10 cc.
- CORT-ACNE CREAM.** (Dermik) Hydrocortisone alcohol D 25%; N-Sulfanilylacетamide 8.5%; salicylic acid 1%; sulfur colloid 3% and 2,2'-methylenebis 0.5% in a greasy vegetable oil vehicle./Tube. Tube 1 oz.
- CORTADYNE.** (Miss) Cortisone acetate 2.5 mg., aspirin 300 mg., ascorbic acid 50 mg., aluminum hydroxide 120 mg., mag. hydroxide 40 mg./Tab. Bot. 100s
Use: Treatment of arthritis and rheumatism
- CORTAFAC.** (Pharmaco) Hydrocortisone alcohol D 25%; in a bland, white ionized base buffered to pH 4.5. Lot. 4 oz.
Use: Dermatologic.
- CORTALEX.** (Upjohn) Ascorbic acid 15 mg./tab., hydrocortisone 0.1 mg./tab. Bot. 40s, 100s, 1000s.
Use: Dermatologic
- CORTARIN CREAM.** (C & M Pharmacol)

- Hydrocortisone ¼% Iodochlorhydroxyquin 3%, LCD 5% special base Tube 2 oz
Use Dermatoses
- COR-TAR-QUIN** (Dome Chemicals) Micro sized hydrocortisone 0.25%, 0.5% or 1% w/liquid carbonis detergens 2%, diiodo hydroxyquinoline in acid mantle vehicle 1% Latan Sqz Bot ½ oz, 1 oz, 4 oz Cream Tube 5 Gm, ½, 1, 2 and 4 oz 1 lb
Use Dermatoses
- COR-TAR-QUIN FORTE.** (Dome) Hydrocortisone alcohol ½%, coal tar solution 10%, diiodohydroxyquinoline 3% Creme ½ oz
- CORTATE, NND 1961** (Schering) Desoxy corticosterone acetate in oil for I.M. inj: 1 cc Amp containing 5 mg, Box 6s, 50s; 10 cc Pellet 75 mg, Vial 3s Vials containing 5 mg/cc, Box 1s, 6s, Buccal Tabs 2 mg, Bot 30s, 100s
Use Addison's disease
- CORT-DOME.** (Dome) Hydrocortisone free alcohol pH 4.6 Cream (¼, ½, 1 & 2%) Tube 5 Gm, ½, 1, 2, 4 oz Jars (¼%) 4 oz, 1 lb Lotion (½, 1 & 2%) Bot ½, 1 oz, 2 oz, & 4 oz, ¼% 4 oz Supp. (25 mg) Box 12s
Use Topically: dermatitis
- CORTEF, NND 1962** (Upjohn) Hydrocortisone Tab 5 mg, Bot 50s; 10 mg & 20 mg, Bot 25s, 100s, 500s
- CORTEF ACETATE OINT., NND 1962** (Upjohn) Hydrocortisone acetate oint 10 and 25 mg/Gm Tube 5 Gm and 20 Gm
- CORTEF ACETATE SOLUTION, NND 1962** (Upjohn) Hydrocortisone acetate Aq susp, sterile 50 mg/cc Vial 5 cc
Use Intra-articular & Intrabursal Injection
- CORTEF-F OINTMENT, NND 1959** (Upjohn) Fluorohydrocortisone Oint (0.1 or 0.2%) Tube 5 Gm & 20 Gm
Use: Topically; skin diseases.
- CORTEF FLUID, NND 1962** (Upjohn) Hydrocortisone cyclopentylpropionate 10 mg/cc Oral susp Bot 4 oz
Use: Orally, hydrocortisone therapy
- CORTEF INTRAMUSCULAR.** (Upjohn) Hydrocortisone 50 mg, polysorbate 4 mg, carbonylmethylcellulose 5 mg, in sod chloride sol/cc Vial 5 cc
Use: I.M., adrenal failure & infections
- CORTEF ORAL SUSPENSION.** (Upjohn) Hydrocortisone cyclopentylpropionate equiv to 10 mg of hydrocortisone/5 cc Bot 4 oz
Use Orally, adrenal cort hormone
- CORTEF STERILE SOLUTION.** (Upjohn) Hydrocortisone Amp (100 mg) 20 cc Sax Te, 5s, 25s, 50s, 100s
Use: 20 cc added to 500 or 1000 cc IV sol; adrenal failure & infections
- * Generic name recognized by N.F., U.S.P. or A.M.A.
- CORTELL CREAM.** (Dermatologicals, Inc) Hydrocortisone ocelate, either ¼, ½, or 1%, in water washable base ¼, 4 oz Jars; ½ and 1%, Tubes ½ oz, 1 oz
Uses: Topical antipruritic and anti-inflammatory agent
- CORTENIL.** See: Desoxycorticosterone Acetate (Various Mfr)
- CORTICAL HORMONE PRODUCTS.**
See Adrenal Cortex Ext (Various Mfr.)
- Aristocort (Lederle)
Cortace (Upjohn).
Corticotropin
Cortisone Acetate (Upjohn).
Desoxycorticosterone Acetate (Various Mfr.).
Fludrocortisone
Hydrocortisone
Kenacort (Squibb)
Lipo Adrenal Cortex (Upjohn)
Magnacort (Pfizer)
Medrol (Upjohn)
Methylprednisolone
Prednisolone
Prednisone
Triamcinolone
- CORTICLORON, SUSPENSION.** (Schering) Cortisone ocelate 5 mg, Chlor trimeton molecule 2.5 mg/cc. Susp. Bot w/dropper 15 cc
Use: Nasal and ocular symptoms
- CORTICLORON, NASAL.** (Schering) Cortisone acetate 5 mg, chlortrimeton 3 mg/cc Squeeze bot 15 cc
Use: Nasal therapy
- CORTICOSTERONE, U.S.P.** (Kendalls Compound B) (Various Mfr)
- * **CORTICOTROPIN INJ, U.S.P., NND 1962;** ACTH, Adrenocorticotrophic or adrenocorticotrop(h)in hormone or corticotrophin
Armour—Vial 40 u.
Upjohn—Vial 5 cc 2 dosage forms 20 u/cc & 40 u/cc
Wilson—Vial 5 cc 2 dosage forms 20 u/cc & 40 u/cc
In dosage form 1 mg = 1 USP u, 1 IU & 1 Armour u.
See: Acthar, Vial (Armour)
Actrope, Vial (United)
Cortrophin, Vial (Organon)
Acthatrope (Spanner)
Solacthyl, Vial (Squibb)
- W/Zinc See Cortrophin Zinc, Vial (Organon)
- CORTICOTROPIN (ACTH) LYOPHILIZED, NND 1962** (G F Harvey) Vial (40 u) 5 cc
See Acthar, Lyophilized (Armour)
ACTH (Various Mfr)
- CORTICOTROPIN (ACTH), STERILE, U.S.P.**

N.N.R. 1962. (Upjohn) 25 U.S.P. u./Vial. 4D U.S.P. u./Vial 20 cc.

***CORTICOTROPIN-GEL PURIFIED, U.S.P., N.N.D.** 1962. (Wilson) Corticotropin in gelatin menstruum, Vial (40 u./cc. & 80 u./cc.) 5 cc.

Use: I.M., S.C., corticotropin therapy.

CORTICOTROPIN-GEL PURIFIED, N.N.D. 1962 (National Drug) Corticotropin in respiratory menstruum, Vial (40 u./cc. & 80 u./cc.) 5 cc.

CORTICOTROPIN HIGHLY PURIFIED, N.N.D. 1962. See: H. P. Acthar Gel, Disposable Cartridge & Vial. (Armour).

***CORTICOTROPIN INJECTION, REPOSITORY, U.S.P., N.N.D.** 1962.

See: Acthar Gel. (Armour)

ACTH Gel, Purified. (National Drug & Testagar)

Corticotropin Gel. (Organon)

Cortrophin-Zinc (Organon)

Cortrophin Gel (Organon)

Dnpo-ACTH. (Upjohn)

El-Acorte Gel. (Brnon)

H.P. Acthar Gel. (Armour)

Purified Corticotropin Gel. (Wilson)

Use: (I.M. 10 to 100 u. daily) Anterior pituitary hormone therapy.

CORTICOTROPIN SOLUTION, N.N.R. 1955. (Wilson) Adrenocorticotrophic activity derived from anterior pituitary. A sterile solution in 5 cc. multiple dose vial.

***CORTICOTROPIN-ZINC HYDROXIDE SUSPENSION, STERILE, U.S.P., N.N.D.** absorbed on zinc hydroxide See: Cortrophin-Zinc, Vial (Organon)

CORTIFAN CREAM 1%, N.N.D. 1962. (Schering) Hydrocortisone as free alcohol 10 mg./Gm. in water-washable base. Tube 10 Gm.

Use: Allergic dermatoses.

CORTILIN. (Lincoln) Adrenal cortical ext. Aq. euph. (7 mg. desoxycorticosterone act./cc.) Vial 10 cc.

CORTINAO. (Central) Aq. susp. of fine cryst. of 11-desoxycorticosterone acetate, each cc containing 5 mg. desoxycorticosterone acetate. Vials 10 cc.

Use: Cortical insufficiency.

CORTINAZIN, N.N.R. 1955. (Pfizer) Isoniazid. Isonicotinic hydrazide. Tab. 50 mg. Bot. 100s.

Use: Orally, antituberculosis therapy.

CORTINORAL. (Harrower) Adrenal cortex ext. Cap. ½ rat u. Bot. 50s.

Use: Orally, adrenal cortex therapy.

CORTIPHATE. (Travenol Labs) Hydrocortisone 21-phosphate, disodium salt. 50 mg./ml. Vial 2 ml.

* Generic name recognized by N.F., U.S.P. or A.M.A.

Use: IV., I.M. or S.C. adrenocortical insufficiency.

CORTIQUIN OINT. (Klug) Hydrocortisone, micronized, free alcohol 0.5%, liquor carbonis detergens 1.5%, diiodohydroxyquinoline 1.0%. Tubes 15 Gm.

Use: Dermatoses of fungus natura.

CORTISAL (Lincoln) Cortisone acetate 2.8 mg., aspirin 350 mg., magnesium trisilicate 100 mg./Tab. Bot. 100s

Use: Treatment of rheumatoid arthritis

***CORTISONE ACETATE, U.S.P., N.N.D.** 1962. Injection, Ophth. Susp., Tab. (Knoll's Comp. E.) (Upjohn).

See: Cortagon Acetate, N.N.R. (Schering).

Cortone Acetate, N.N.R. (Merck Sharp & Dohme)

Cortivate, Cap. (United Lab.).

Isopta-Cortisone, Sesp. Ophth. (Alcon Labs.).

W/Actylate. See: Actylate Plus E, Tab. (Ksney)

W/Ascorbic acid. See: Cortivita, Cap. (United Lab.)

W/Aspirin, ascorbic acid, aluminum hydroxide, mag. hydroxide. See: Cortodynn, Tab. (Hess)

W/Ammonium salicylate, Pot. p-aminobenzoate & Vit. C. See: Neocylain w/Cortisone, Tab. (Central).

W/Aspirin, magnesium trisilicate. See: Cortisal, Tab. (Lincoln)

W/Bacitracin. See: Cortann-Bacitracin, Ophth. Oint. (Merck Sharp & Dohme)

W/Chlor-Trimetan malate. See: Corticlaran, Sesp. (Schering).

W/Neomycin sulfate. See: Neocortan, Ophth. Oint. & Susp. (Upjohn).

W/Sod. salicylate, aluminum hydroxide gel dried, Cal. ascorbate & Cal. carb. See: Solcort, Tab. (Massengill).

W/Vit. C. See: Cortivate, Cap. (United Lab.).

***CORTISONE ACETATE, U.S.P., N.N.D.** 1962. (Upjohn) 11-Dehydro-17-hydroxycorticosterone. Ophth. Oint. 15 mg./Gm. Tabs. 5 mg., Bot. 50s; 10 mg., Bot. 24s, 100s; 25 mg., Bot. 20s, 100s, 500s.

CORTISORBATE. (Schiffelin) Adrenal cortex active principle absorbed on charcoal Tab. ½ rat u. 20s, 100s; 1 rat u. 20s, 100s. Use: Orally, adrenal insufficiency.

CORTISPORIN. (Burroughs Wellcome) Polymyxin B sulfate 5000 u., bacitracin 400 u., neomycin sulfate 5 mg., hydrocortisone free alcohol 10 mg./Gm. Oint. Ophth. Tube ½ oz. & Topically ½ oz.

Use: Anti-inflammatory & antibacterial.

CORTISPORIN LOTION. (Burroughs Wellcome) Polymyxin B sulfate 10,000 u., neo-

- mycin sulfate 5 mg, hydrocortisone 10 mg/cc Plus squeeze bot 10 cc
Use Anti-inflammatory, antipruritic
- CORTISPORIN OPHTHALMIC SUSPENSION.** (Burroughs Wellcome) Polymyxin B sulfate 10 000 u, neomycin sulfate 5 mg, hydrocortisone (1%) 10 mg, in a sterile, saline isotonic/ml Drop bot 5 cc
Use: Ophic, inflammation
- CORTISPORIN OTIC DROPS** (Burroughs Wellcome) Polymyxin B sulf 10 000 u, neomycin sulf 5 mg, hydrocortisone free alcohol 10 mg/cc Bot w/dropper 5 cc
Use Ear infections
- CORTISPRAY** (Walker) Hydrocortisone D 5% soln Squeeze bottle 15 cc
Use Skin itching & inflammation
- CORTIVITE.** (United Lab) Cortisone acetate 10 mg, ascorbic acid 100 mg/Cap Bot 40s, 100s, 1000s
Use: Stimulaton of the adrenal cortex
- CORTOGEN ACETATE, NND 1962** (Schering) Cortisone acetate (Kendall comp E1 Aq. susp 25 mg/cc multiple dose Vial 10 cc Ophth Susp 5 or 25 mg/cc Bot, w/dropper 5 cc 0.5% Bot w/dropper 5 cc Tab 25 mg Bot 20s 30s 100s
- CORTOLINE CREAM.** (Dermatologicals Inc) Hydrocortisone 1/2%, iodochlorhydroxyquinoline 3% Tube 1/2 oz 1 oz
Use: Topical antipruritic anti-inflammatory agent
- CORTOMIXIN.** (Braemmel) Polymyxine B sulfate 10,000 u, hydrocortisone 0.5% in a propylene glycol base/cc Vial 5 cc
Use: Local infections of the ear
- CORTOMYCIN.** (Bryant) Hydrocortisone acetate 10 mg, or 25 mg, neomycin 5 mg/Gm Oint Vial 5 Gm, 20 Gm
Use: Topically, skin infections
- CORTONE ACETATE, NND 1962** (Merck Sharp & Dahme) Cortisone acetate Ophth Susp. 2.5% & 0.5% Vial 5 cc Oint 1.5% 1/2 oz tube Orol Tab 25 mg, Bot 20s 40s, Tab 5 mg, Bot 50s Parenteral Susp 20 cc, 25 mg/cc, Vials 10 cc 50 mg/cc W/Bacitracin (Merck) Cortisone acetate 15 mg, bacitracin 1000 u/Gm Ophth Oint Tube 3.5 Gm
Use: Anti-inflammatory with antimicrobial activity
- CORTONE-BACITRACIN OPHTHAEMIC OINTMENT.** (Merck Sharp & Dahme) Cortisone acetate 15 mg, bacitracin 1000 u/Gm Tube 1 dr
Use: Treatment of infections of the eye
- CORTOPIX CREAM.** (Dermatologicals, Inc) Hydrocortisone 1/2%, liquor carbonis detergens 3% Tube 1/2 oz, 1 oz
Use: Subacute and chronic dermatoses
- CORT-QUIN CREME & LOTION.** (Dome Chemicals) Hydrocortisone 1/4%, 0.5% or 1% and diiodohydroxyquin 1% in Acid Mantle base Tube 0.5 oz, 1 oz Lotion 1/4% Bot 4 oz
Use Subacute and chronic dermatoses
- CORTRIL, NND 1962** (Pfizer) Hydrocortisone Tab. 10 mg, Bot 25s, 100s 20 mg Bot 20s, 100s Vaginal Tab: Hydrocortisone free alcohol 10 mg/vag tab Pkg 10s Oint. 1% or 2 1/2% Tube 1/6 oz
- CORTRIL ACETATE, NND 1962** (Pfizer) Ophth. Oint. 1/2 oz tube, hydrocortisone 0.5% & 2.5% Topical Oint. 1/6 oz tube hydrocortisone 1% & 2.5% Aq. Susp. 5 cc Vial, hydrocortisone 25 mg/cc
Use Anti-inflammatory
- W/Terramycin, (Pfizer) Hydrocortisone acetate 15 mg, terramycin HCl 5 mg/cc Susp Bot w/dropper 5 cc
Use (2 gtt tid) Broad spectrum antibiotic with anti-inflammatory therapy
- W/Terramycin, topical Oint (Pfizer) Hydrocortisone 10 mg, terramycin 30 mg/Gm Tube 1/2 oz See Terra Cortril (Pfizer)
Use Skin infections & inflammatory conditions
- CORTROPHIN GEL, NND 1962** (Organon) Corticotropin Gel Vial 25 USP & 40 USP u Pkg w/5 cc vial of solvent
- CORTROPHIN-ZINC, NND 1962** (Organon) Corticotropin 40 u, zinc 2 mg/cc Vial 5 cc Disp Syr (20 or 40 u/cc) 1 cc
Use IM & S.C., long acting ACTH therapy
- CORTUNON.** (Cartunon) Vasodilator principle from liver Amp (7 1/2%) 2 cc, Box 6s; Vial 20 cc; Cap 75 mg Bot 50s
Use Orally & injectable vasodilator
- CORTYZANOL** (Pfizer) Prednisolone 0.02 Gm tetrahydrozoline 0.1 Gm/100 ml or Prednisolone 0.02 Gm, tetrahydrozoline 0.05 Gm/100 ml
Uses Anti-inflammatory and vasoconstrictor
- CORYBAN** (Roerig) Salicylamide 220 mg, acetophenetidine 120 mg, caffeine 30 mg, propenpyridamine maleate 10 mg, ascorbic acid 20 mg, hesperidin purified 20 mg/cap Bot 12s
Use (1 2 q 4 h) Provides relief for the symptoms of the common cold and improves resistance to colds
- CORYBAN-D.** (Roerig) Phenylpropanolamine 25 mg, chlorpheniramine maleate 2 mg, phenacetin 160 mg, ascorbic acid 25 mg, salicylamide 220 mg, caffeine 30 mg/Tab Bot 24s
Use: Common cold
- CORYBIO.** (US Standard Products) Hesperidin 25 mg, ascorbic acid 12.5 mg, phenyltoloxamine 12.5 mg aspirin 113.5 mg.

- acetophenelidid 81 mg., caffeine 16 mg./Tab. Bot. 100s, 1000s.
Use: Common cold.
- CORYPNEDRINE.** (Wallou) Ephedrine NCI $\frac{1}{4}$ gr., aspirin $7\frac{1}{2}$ gr./Tab. Bot. 20s.
Use: Orally, analgesic (premenstrual tension).
- CORYZ.** (Marion) N-acetyl-p-aminophenol 325 mg., phenylpropanolamine 50 mg., glyceryl guaiacolate 100 mg., noscapine 20 mg./Tab. Bot. 50s, 250s, 1000s.
Use: Symptoms of the common cold.
- CORYZ SUSPENSION.** (Marion) Noscapine alkaloid 10 mg., phenylpropanolamine NCI 25 mg., glyceryl guaiacolate 50 mg., acetaminophen 160 mg., chloroform 4 mg./5 ml. Bot. T pt., gal.
Use: Treatment of common cold
- CORYZ A.T.** (Marion) Noscapine 30 mg., glyceryl guaiacolate 50 mg., alcohol 4%/5 cc. Bot. 8 oz.
Use: Antilussive.
- CORYZA.** (Superior Pharm.) Quinine sulfate $\frac{1}{2}$ gr., ammonium chloride $\frac{1}{2}$ gr., camphor $\frac{1}{2}$ gr., belladonna pow. ext. 1/20 gr., acanite pow. ext. 1/10 gr./Tab. Bot. 1000s.
Use: Rhinitis.
- CORYZA RX A (RICHARDS) C.C.T.** (Merck Sharp & Dohme) Belladonna ext. 3 mg., acanite root 24 mg., quinine sulfate 30 mg., ammonium chloride 30 mg., camphor 30 mg./Tab. Bot. 1000s.
Use: Rhinitis
- CORYZOL.** (Tilden) Ephedrine, benzocaine, camphor, menthol, pine oil, in petrolatum. Oint. Tube $\frac{1}{4}$ oz.
Use: Relief of irritation, inflammation and congestion of the nasal mucosa
- COSADEIN.** (Parke, Davis) Codeine phosphate 1 gr., chloroform 2 gr., white pine 32 gr., paper bud 4 gr. & glycerin 120 min., w/alcohol 20%/fl. oz. Bot. T pt., 1 gal.
Use: (1 tsp. q. 3-4 h. p.r.n.) Sedative, expectorant.
- CO-SALT.** (Arlington-Funk) Choline, potassium chloride, ammonium chloride, tri-calcium phosphate. Bot. 2 oz., 8 oz.
Use: Low-sodium diets.
- COSANTIN.** See: Diparcol. (May & Baker)
- COSANYL.** (Parke, Davis) Codeine sulf. T gr., euphorbia pilulifera fr. 120 min., squill comp. syr. 24 min., wild lettuce zyr. (bitterless) 8 gr., & menthol 2/25 gr. & alcohol 6%/fl. oz. Bot. 2 oz., 4 oz., T pt., 1 gal.
Use: (1 tsp. q. 3-4 h. p.r.n.) Sedative expectorant.
- COSA-SIGNEMYCIN.** (Pfizer) New Signemycin. Oleandomycin as triacetyloleandomycin 41.7 mg., glucosamine-potentiated tetracycline (Cosa-Tetracycl) 83.3. Cap. 125 mg. Bot. 25s, 100s; Cap. 250 mg. Bot. 16s, 100s.
Use: Various infections.
- COSA-SIGNEMYCIN PEDIATRIC DROPS.** New Signemycin Ped. Drops. (Pfizer) Cosa-Signemycin 100 mg./cc. Bot. 100 mg./cc/10 cc.
Use: Antibiotic.
- COSA-SIGNEMYCIN ORAL SUSPENSION.** New Signemycin Syr. (Pfizer) Cosa-Signemycin 125 mg/5 cc. Bot. 125 mg/5 cc/2 oz.
Use: Antibiotic.
- COSA-TERRABON SUSP.** (Pfizer) New Terramycin Syrup. Calcium di-oxytetracycline equivalent to 125 mg oxytetracycline with 125 mg. glucosamine or n-acetyl glucosamine/5 cc. Bot. 2 oz., 1 pt.
Use: Infections caused by Gram positive and Gram negative bacteria; rickettsiae, spirchetes and large viruses.
- COSA-TERRABON PED. DROPS.** New Terramycin Ped. Drops. (Pfizer) Calcium di-oxytetracycline with glucosamine 100 mg/cc. or 5 mg/drop. 10 cc. dropper bottle with calibrated plastic dropper.
Use: Common infections.
- COSA-TERRACYDIN CAPSULES** now named Terracydin Capsules.
- COSA-TERRAMYCIN.** (Pfizer) New Terramycin. Caps. 125 mg Bot. 25s, 100s; Caps 250 mg. Bot. 16s, 100s. Oral susp. 125 mg/5 cc. Bot. 2 oz. Pediatric 5 mg/Drop. Bot. 10 cc.
Use: Antibiotic.
- COSA-TERRASTATIN.** (Pfizer) New Terrastatin. Oxytetracycline 250 mg, glucosamine 250 mg., nystatin 250,000 u./Cap. Bot. 50s. Susp $\frac{1}{2}$ above 15 cc. Bot. 2 oz.
Use: Antibiotic.
- COSA-TETRABON ORAL SUSPENSION** now named Tetracycl Syrup.
- COSA-TETRABON PEDIATRIC DROPS** now named Tetracycl Pediatric Drops.
- COSA-TETRACYDIN.** (Pfizer) New Tetracyclin. Oxytetracycline w/glucosamine 125 mg, butizine HCl 10 mg, phenacetyl 120 mg, salicylamide 150 mg., caffeine 30 mg/Cap. Bot. 50s.
Use: Relief of the common cold.
- COSA-TETRACYCLIN.** (Pfizer) New Tetracycl. Caps. Tetracycline w/glucosamine. 125 mg., Bot. 25s, 100s; 250 mg. Bot. 16s, 100s. Oral Susp. (125 mg tetracycline w/glucosamine/5 cc) Bot. 2 oz. Pediatric Drops (5 mg/Drop). 100 mg/cc. Bot. 10 cc.
- COSA-TETRASTATIN.** (Pfizer) New Terrastatin. Glucosamine-potentiated tetracycline HCl 250 mg., nystatin 250,000 u./Cap. Bot.

- 16s 100s Susp ½ above/5 cc Bot 2 oz
COSIL, (Reed & Carnrick) (Dimethicose) Di methyl polysiloxane D1 Gm, magnesium hydroxide 1.25 Gm /15 cc
 Use Management of Constipation
- COSILLEX** (Warner Chilcott) Codeine phosphate 1/6 gr ammonium chloride, potassium citrate squil comp. syr., menthol, cocillana citric acid euphorbia & wild cherry/5 cc Bot 4 oz, 1 pt
 Use (1 tsp q 3-4 h, prn) Narcotic cough preparation
- COSMEDICAKE**, (Kay) Benzalkonium chloride, salicylic acid, hexachlorophene in a greaseless base Cakes 4 shades
 Use Management of acne vulgaris
- COSMOLINE** See Petrolatum
- COSMO-REY**, (Jeffery) Vit B 12 1000 mcg /cc Vial 10 cc
 Use Anemias
- COTARNINE BIPHENALATE**, See Styptol
- COTARNINE CHLORIDE**, NF VIII Cotaresee hydrochloride See Stypticin (Merck Sharp & Dohme)
- COTARNINE HYDROCHLORIDE**, (Merck Sharp & Dohme) See Cotarnine Chloride
 Use Narcotic and hemostatic
- COS-TAR-OK**, (Ulmer) Crude coal tar 5%, zinc oxide in aromatic oint base 1 oz Jar 1 lb
 Use Antipruritic
- COTAZYM** (Organon Inc) Lipase, 2,000 units having digestive power for 17 Gm; trypsin having digestive power for 34 Gm dietary protein; amylase having digestive power for 40 Gm dietary starch plus other enzymes from whole pancreas/Cap Bot 100s
 Use Pancreatic Enzyme Deficiency
- COTAZYM-B** (Organon) Pancreatic digestive enzymes (liponcreatin)—lipase, trypsin amylase, plus bile salts and cellulose/Tab Bot 24s
 Use Indigestion
- COTEL**, (Jeffery) Vit B 12 25 mcg /Tab Bot 100s
 Use Anemias
- COTETRIN** (Haskell) Pyramisamine maleate 20 mg, ephedrine sulfate 10 mg, ephedrine 2½ gr, acetophenetidin 2½ gr/Tab Bot 100s
 Use (1-2 q 4 h) Analgesic with antihistamine
- COTNAMID** (Lloyd, Dabney & Westerfield) Phthalylsulfacetamide 1 Gm, bismuth subcarbonate 1 Gm, kaolin 25 Gm, pectin 250 mg, sodium carboxymethylcellulose 250 mg, homatropine methylbromide D 625 mg, methscopolamine bromide 0.625 mg, hyoscyamine HBr 0.625 mg /oz. Susp Bot 1 pt
 Use Orally, bacteriostatic & antispasmodic
- COTNERA COMPOUND**, (Ayerst) Dimethazone HCl 25 mg, Isalhipendyl HCl (Therohistin) 2 mg, 1-phenylephrine HCl 5 mg, acetaminophen 100 mg, ammonium chloride 100 mg, Sodium citrate 50 mg, chloroform D 25%, alcohol 10%/5 cc Bot 1 pt, 1 gal
 Use Cold & cough syrup
- COTNERA SYRUP**, NND 1962 (Ayerst) Dimethazone HCl 25 mg /5 cc Bot pt, 1 gal
 Use Expectorant
- COTINAZIN**, NND 1960 (Pfizer) Isoniazid 50 mg /Tab Bot 100s, 1000s
 Use Treatment of tuberculosis
- COTLONE**, (Truxton) Prednisolone 5 mg /Tab Bot 100s, 500s, 1000s, 5000s
 Use Arthritis, bronchial asthma, allergic dermatoses
- COTONE**, (Truxton) Prednisone 5 mg /Tab Bot 100s, 500s, 1000s, 5000s
 Use Arthritis, bronchial asthma allergic dermatoses
- COTOPHEROL**, (Endo) d Tocopherol acetate equiv to 200 mg, dl alpha tocopheryl acetate/cc Vial 10 cc
 Use Vitamin therapy
- ***COTTONSEED OIL EMULSION**, NND 1962 See Lipomul IV (Upjohn)
- COTUSSIS**, (National Drug) Codeine phosphate 1/6 gr, terpin hydrate ½ gr w/chloroform, menthol & glycerin/5 cc Bot 3 oz, 1 pt, 1 gal
 Use (1 tsp q 3-4 h, prn) Narcotic cough preparation
- COUMADIN**, NF X, NND 1962 (Endo) Warfarin Sodium, 3 (alpha acetylbiphenyl) 4 hydroxycoumarin Sodium Amp (75 mg /10 cc) Tab 5, 7.5, 10, 25 mg, Bot 25s, 100s
 Use Orally & IV, anticoagulant
- COVANAMINE**, (Vanpelt & Brown) Phenylephrine HCl 15 mg, Phenylpropanolamine HCl 25 mg, Chlorpheniramine maleate 4 mg, pyrilamine maleate 25 mg /Tab Bot 50s, 250s; Lq: Phenylephrine HCl 3.75 mg, phenylpropanolamine HCl 6.25 mg, chlorpheniramine maleate 1 mg, pyrilamine maleate 6.25 mg in a sorbital base/5 cc Lq Bot pint
 Uses Oral decongestant and colds
- COVANAMINE TABLETS**, (Vanpelt & Brown) Phenylephrine HCl 3.75 mg, phenylpropanolamine HCl 6.25 mg, chlorpheniramine maleate 1 mg, pyrilamine maleate 6.25

mg., glyceryl gualacolate 100 mg./Tab. Bot. 100s, 500s.

Use: Treatment of symptoms due to cold.

COVANGESIC. (Voepel & Brown) Phenylephrine HCl 7.5 mg., phenylpropanolamine HCl 12.5 mg., chlorpheniramine maleate 2 mg., pyrilamine maleate 12.5 mg., aspirin 324 mg./Tab. Bot. 100s

Use: Colds.

COVANGESIC LIQUID. (VanPelt & Brown) Phenylephrine HCl 3.75 mg., phenylpropanolamine HCl 6.25 mg., chlorpheniramine maleate 1.0 mg., pyrilamine maleate 6.25, acetaminophen 120 mg./5 cc. Bot. 1 pt, 1 gal.

Use: Symptoms associated with common colds.

COVATIN. See: Surven.

COVICONE CREAM. (Abbott) Silicone, nitrocellulose, castor oil suspended in vasishing cream base. Tube 1 oz., 4 oz., Jar 1 lb.

Use: Typically for protecting skin from irritation.

CO-VI-FERRA. (Mallard) Cobalt chloride 15 mg., ferrous sulf. 5 gr., Vit. B-12 2 mcg., B-1 1.5 mg., B-2 0.5 mg., nicotinamide 5 mg./Cap. Bot. 100s, 1000s

Use: Anemias due to iron deficiency

COVINAL. (Myers-Carter) Vit. A, D, E, B-1, B-2, B-6, C, niacinamide, panthenol. Vials 30 cc.

COVISTEN. (Organon) Stenedial 5 mg. [methylandrostenedial], ascorbic acid 50 mg., thiamine HCl 2 mg., riboflavin 3 mg., pyridoxine HCl 1 mg., Vit. B-12 2.5 mcg., niacinamide 20 mg., folic acid 0.5 mg., Vit. A 5000 u., Vit. D 3 500 u., ferrous gluconate 75 mg., dibasic Cal phosphate 200 mg., mang hypophosphite 0.5 mg., dibasic mag. phosphate 20 mg./Tab. Bot. 100s.

Use: Tissue builder & dietary supplement for waning gonadal function.

COVITRAL. (Flint, Eaton) Vit. B-12 w/Intrinsic factor conc. 1/6 u., thalium bitartrate 166 mg., folic acid 0.33 mg., ascorbic acid 16.66 mg., pyridoxine HCl 0.83 mg./Cap. Bot. 100s, 500s, 1000s

Use: Vitamin, hematonic.

COVON (Deteret) #1 Tablets: Acetylsalicylic acid 3½ gr., phenacetin 2½ gr., caffeine ½ gr., dihydrocodeine bitartrate 7.5 mg./Tab. #2 Tablets: Acetylsalicylic acid 3½ gr., phenacetin 2½ gr., caffeine ½ gr., dihydrocodeine bitartrate 15 mg./Tab.

Use: Analgesic.

COZYME, NND 1962. (Trovant Lab) Dexpantenol, 250 mg./cc./Solin Vials 10 cc.

Use: Correct and prevent intestinal atony.

C.P.A. (Sutliff & Case) Phenobarbital ½ gr.

atropine sulfate 1/500 gr., Cal. carb. 10 gr./Tab. Bot. 100s, 1000s

Use: Antispasmodic & sedative.

C.P.H. (Cotter) Amino acids 50 Gm. & dextrose 50 Gm. Bot. 1000 cc.

Use: Protein hydrolysate.

C.P.T. (C.P.T. Lab) Chemical pregnancy test. Pkg. complete reagent set.

Use: Early diagnosis of pregnancy.

C-QUIN. (Smith, Miller & Polch) Ascorbic acid Tab. 500 mg. Bot. 100s, 500s, 1000s

CRACILIN TABS. (Craig) Potassium penicillin G 400,000 u./Tab. Bot. 100s.

Use: Infections.

CRADIX CAPS. (Craig) d-Amphetamine sulfate 15 mg./timed disintegration cap, Bot 100s.

Use: Psychogenic & appetite depression.

CRADIX B No. 1 and No. 2 Caps. (Craig) #1 d Amphetamine sulfate 10 mg., amobarbital 60 mg./timed dis. cap. #2 d-Amphetamine sulfate 15 mg., amobarbital 60 mg./timed dis. cap. Bot. 100s.

Use: Obesity control.

CRADOL. (Breon) Emulsion of methylbenzethonium chloride 1:1500 in an oxyhesterin base of liquid petrolatum and lanolin. Bot. 4 oz.

Use: Seborethia capitis (Cradle cap)

CRATAGANITE. (Smith, Miller & Polch) Veratrum viride 1 gr., opocyum tr. 1 min, Sod nitrite 1 gr., nitroglycerine 1/100 gr., crotaegus tr. 1 min., pot. nitrate 1 gr./Tab. Bot. 100s, 1000s.

Use: Vasodilator.

C-REACTIVE, PROTEIN ANTISERUM. See: CRP-Antiserum, Vial. (Schiffelin).

CREAMALIN, NND. 1961. (Winthrop Labs) Aluminum hydroxide gel 320 mg., mag. hydrox. 75 mg./Tab. Tab. Tin 12s, Bot 50s, 100s, 200s, 1000s. Cap. Box 24s Bot 100s, 1000s. Liq., flavored, Bot. 8 fl. oz., 12 fl. oz., 1 pt, 1 gal Unflavored 1 pt.

Use: Hyperacidity.

CREAMALIN H.M. (Winthrop) Aluminum hydroxide gel, dried, 0.23 Gm., kamatropise methylbromide 1 mg./Tab.

Use: Treatment of peptic ulcer.

CREAMLETS. (Winthrop) Dried aluminum hydroxide gel 160 mg., magnesium hydroxide 37.5 mg., in a mint flavored gum base. Tab Box. 12s.

Use: Relief of heartburn, indigestion and gastric hyperacidity.

CREMACAL. (Mabart) Calamine, benzocaine, phenol & glycerin in water miscible base. Oint. Tube 1 oz

Use: Analgesic, antipruritic.

CREMAGOL (Cremagol Lab) Emulsion of

- liq petrolatum & agar agar Bot 14 oz
 Also w/cascara also w/phenolphthalein
 Use Laxative
- CREMO-BISMUTH**, NNR 1949 (Merck Sharp & Dohme) (Milk of bismuth NF) Bismuth subnitrate 233 Gm (36 gr)/30 cc Bot 1 pt, 1 gal
 Use: Antacid & intestinal astringent
- CREMO-CARBONATES**, (Merck Sharp & Dohme) Chloroform 1 gr, bismuth subcarb 20 gr, mag carb 20 gr, col carb 10 gr/30 cc Bot 12 oz
 Use Antacid
- CREMODIAZINE** (Merck Sharp & Dohme) Sulfadiazine 3 Gm/30 cc Bot 1 pt
 Use: Sulfonamide therapy
- CREMULSION** (Creomulsion Co) Beechwood creosote cascara specac, menthol, white pine wild cherry Bot 4 8 fl oz
 Use Coughs and bronchial irritations due to colds
- CREMULSION FOR CHILDREN** (Creomulsion Co) Beechwood creosote cascara ipsecac, menthol, white pine, wild cherry Bot 4 fl oz
 Use Coughs and bronchial irritations due to colds, for children 1-14 years
- CREMOMYCIN SUSPENSION** (Merck Sharp & Dohme) Sulfisoxazine 3 Gm, kaolin colloid 3 Gm, pectin 0.3 Gm, neomycin sulfate 300 mg/30 cc Bot 8 oz
 Use Antidarrheal
- CREMOSUXIDINE**, (Merck Sharp & Dohme) Sulfisuxidazine 3 Gm/30 cc (Pectin 1% kaolin 10%) Bot 1 pt
 Use: Intestinal sulfa therapy
- CREMGTHALIDINE** (Merck Sharp & Dohme) Sulfathalidine 6 Gm/30 cc Bot 8 oz
 Use: Inflammatory conditions of colon
- CREMOTRES** (Merck Sharp & Dohme) Sulfamerazine 2 Gm, sulfadiazine 4 Gm, sulfamethazine 4 Gm/100 cc Bot 1 pt
 Use: Triple sulfa therapy
- CREMOTRES-A**, (Merck Sharp & Dohme) Sulfamerazine 2 Gm, sulfathiazine 4 Gm, sulfamethazine 4 Gm, sod lactate 33.3 Gm/100 cc Bot 1 pt
- CREN**, (Cole) Codeine phosphate 1 gr, chloroform 2 min, Pot guaiacal sulfonate 8 gr, ammonium chloride 8 gr, antimony & Pot tartrate 1/16 gr, menthol 1/16 gr, white pine comp syr 60 min, wild cherry 22 gr/fl oz Bot 1 pt, 1 gal
 Use: (1 tsp q 3 h) Sedative expectorant
- CREOLIN—Pearson** (Merck Sharp & Dohme) Cool tar disinfectant & germicide
 Use: Disinfectant
- * **CREOSOTE**, NF, ADR 1955 Wood creosote
 * Generic name recognized by NF USP or A.M.A.
- sole creosote, beechwood creosote Saponated Soln NF
- W/Calcium** as salts. See: Colcrease, Tab (Maltbie)
- W/Guaiacal, aminoacids, polypeptides** See: Peptoloids creosote, Liq (Arlington)
- W/Terpin hydrate** See: Creo Terpin, Liq (Wampole)
- W/Terpin hydrate comp** See: Ethocrea, Liq (Merck Sharp & Dohme)
- CREO TERPIN** (Wampole) Terpin hydrate w/creosote Bot, 3 oz, 6 oz, 1 gal
 Use (1 tsp q 3-4 h) Cough preparation
- CREOZETS** (Creomulsion Co) Beechwood creosote cascara, ipsecac, menthol, white pine wild cherry Box w/16 lozenges
 Use: Coughs, bronchial and throat irritation
- CREPHEX** (Schenley) Ephedrine sulfate .1 mg, pyramine moleate 12.5 mg, Cal cresol sulfonate 150 mg & chlorofan 5 mg/5 cc Bot 1 pt, 1 gal
 Use: (1 tsp q 3-4 h, m p) Cough syrup w/antihistamine
- CRESATIN**, NNR 1951 (Merck Sharp & Dohme) Metacresylacetate Bot 1 oz
 Use Infections of nose, ear, mouth & thn (fungus).
- * **CRESEL**, USP, ADR 1955 (Various Mfr)
 Mixture of 3 isomeric cresols
 Use Antiseptic & disinfectant
- CRESEL-CALCIUM**,
 See Colcrease (Maltbie)
 Kres Lumin (Winthrop Labs)
- CRESEL PREPARATIONS**,
 See Cresotin, Liq (Merck Sharp & Dohme)
 Creolin—Pearson (Merck Sharp & Dohme)
 Cresol (Various Mfr)
 Cresylane, Liq (Parke, Davis)
 Hydrosol, Liq (Blue Line)
 Phenolur, Liq (Squibb)
 Saponated Cresol Sol, NF
- CRESRADINE**, (Raymer) Iodized Cal creosote 2 gr, terpin hydrate 1 gr, hyoscyamus ext 1/8 gr, strychnine sulfate 1/100 gr, Phos 1/200 gr/Cap or Tab; Iodized Cal creosote 8 gr, camphorated opium tr 80 m, ipsecac syr 20 m, senega syr 10 min, sanguinaria syr 10 min, wild cherry syr 80 min/fl oz Bot Cap 100s, 500s 1000s; Syr. Bot 3 oz, 1 pt, 1 gal
 Use: Cough preparation
- CRESTABOLIC INJ.** (Nutrition Control P) Methylandrostenedial 25 mg/cc; Sustalactin/cc Vial 10 cc
 Use Anabolic Steroid
- CRESYLIC ACID**, Same as Cresol USP
- CRESYLIC ACID, SAPONIFIED**, See: Cresolane, Liq (Parke, Davis)

CREYLONE. (Parke, Davis) Cresylic acid, saponified. Bot. 1 gal.

Use: Antiseptic solution.

CRETA-TABS. (SuthR & Case) Calcium Carbonate 10 gr./Tab. Bot. 100s, 1000s.

CRINOGEN. (Merrell) Estrogenic substances (Equiv. to 1 mg. {10,000 I.U.} of estrone, 10 mg. progesterone/cc. Vials 5 cc.

Use: Estrogen & androgen therapy.

CRIODIN (Crookes) Iodine w/colloidal protein. Vial 30 cc.

Use: Iodine therapy.

CRISTALLOVAR. See: Estrone.

CRISTIVITE. (Hiss) Vit. B-12 25 mcg., R-1 10 mg./5 cc. Bot. 4 oz. Vial: Vit. B-12 100 mcg., R-1 100 mg./cc. Vial 10 cc.

Use: Anemias.

CROLIUM SUSPENSOID. (Kent).

W/Colamel 10%. Bot. 5 oz., 1 pt., 1 gal.

W/Copper sulfate 2%. Bot. 5 oz., 1 pt., 1 gal.

W/Sulfathiazole 1/2%. Bot. 5 oz., 1 pt., 1 gal.

W/Sulfur 8%. Bot. 5 oz., 1 pt., 1 gal.

Vehicle, plain, Bot. 5 oz., 1 pt., 1 gal.

Use: Topically, in dermatoses.

C-BON. (Rawell) Ascorbic acid 100 mg., ferrous gluconate 300 mg./Tab. Bot. 100s, 1000s.

Use: Anemias.

C-RON PEDIATRIC. (Zawell) Vitamin C 50 mg., ferrous fumarate 100 mg./chewable Tab. Bot. 100s, 1000s.

Use: Iron deficiency in children.

C-RON PRENATAL. (Rawell) Ferrous fumarate 150 mg., Vit. C 100 mg., A 3000 u., D 300 u., purified powdered bone 750 mg./Tab. Bot. 100s, 1000s.

Use: Prenatal supplement.

CRONETAL. See: Disulfiram.

***CROTALINE ANTIVENIN POLYVALENT,** U.S.P. N.D. 1962. (Wyeth) North and South American Antisnakebite serum. Pow. Inj. yields 10 cc. of restored serum.

Use: I.M.

***CROTAMITON,** N.D. 1962. See: Eurax, Cream & Lot. (Geigy)

C.R.P. ANTI-SERUM. (Schleffelin) C-reactive protein. Vial 1 cc.

Use: Diagnostic agent.

C.R.P. POSITEST. (Schleffelin) C-reactive Protein sufficient to produce a precipitate of 3-plus magnitude when mixed with C-reactive Protein Antiserum. Vial (0.5 ml.) 40 tests.

Use: Diagnostic agent.

CR-TEST. (Hyland) 2 glass slides, diluent,

* Generic name recognized by N.F., U.S.P. or A.M.A.

and sufficient reagent and capillary pipettes for 60 screening tests. Test kit.

Use: Detection of C-reactive protein.

CRU-R7. (M. R. Thompson) Vit. B-12 20 mcg., folic acid 5 mg., liver B-12 2 mcg./10 cc. Vial 10 cc.

CRUDR TUBERCULIN. See: Tuberculin, Old, U.S.P.

CRUPIC. (Jenkins Labs.) Paregoric 5 min., hyoscyamus ext. 1/20 gr., lpecac syr. 5 ml., licorice ext. 1 gr./tab. Bot. 100s, 1000s.

Use: Spasmodic croup and bronchitis.

CRYORETA. (Upjohn) Thiamine HCl 2 mg., riboflavin 2 mg., pyridoxine HCl 1 mg., Caf. parathenate 10 mg., nicotinamide 50 mg./5 cc. Bot. 8 oz.

Use: Vitamin R therapy.

***CRYPTENAMINE ACETATE & TANNATE,** N.D. 1962. Esters of alkaloids of *Veratrum viride*

See: Unitensin Acetate, Tab. (Irwin, Neisler)

Unitensin Tannate, Tab. (Irwin, Neisler)

CRYPTENAMINE, W/Methylclothiazide. See:

Diutensen, Tabs. (Irwin, Neisler).

CRYPTENAMINE, /Methylclothiazide, reserpine. See: Diutensen E, Tabs. (Irwin, Neisler)

W/Eserspine. See: Unitensin R, Tab. (Irwin, Neisler).

CRYSEALLINE TRYPSIN, N.N.R. 1941. Highly purified prep. of enzyme as derived from mammalian pancreas glands.

See: Porezyme, Vial. (National Drug)

Tryptar. (Armour)

CRYSTAL VIOLET. See: Methylrosaniline Chloride, U.S.P.

CRYSTAR. (Armour) Pediatric aspirin. Pkg.

1 gr. Box 24 packets.

Use: Orally, soluble aspirin for pediatric analgesic.

CRYSTICILLIN, N.N.D. 1962. (Squibb) Procaine penicillin G Vial 300,000 u. & 3,000,000 u.

Use: Infections caused by gram positive organisms.

CRYSTICILLIN UNIMATIC 300 A.S. (Squibb)

CRYSTICILLIN UNIMATIC 600 A.S. Procaine penicillin G in aq. susp. Unimatic syringe, 300,000 or 600,000 u. Vial 300,000 u., 1,500,000 u. and 3,000,000 u.

CRYSTIFOR 400. (Squibb) 400 Procaine penicillin G 300,000 u. w/Pot. penicillin G 100,000 u., Vial 1 dose or 5 dose.

Use: I.M., Penicillin-susceptible infections

CRYSTIMIN-GEL. (Hiss) Vit. B-12 200 mcg in a special gelatinous vehicle/cc. Vial 10 cc. 1000 mcg./cc. 10 cc.

Use: I.M., Vit. B-12 therapy

CRYSTIVITE SORBETTES. (Hiss) Thiamine
HCl 5 mg, B 12 5 mcg, lysine 5 mg,
d sorbitol 500 mg / Tab Bot 50s
Use: Tonic

CRYSTIVITE-FE SORBETTES (Hiss) Thiamine
HCl 5 mg, B 12 5 mcg, lysine 5 mg, Fe
fumarate 35 mg, d sorbitol 500 mg / Tab
Bot 50s
Use: Tonic

CRYSTODEX (Crystal) Mephesisin 250 mg,
d-amphetamine sulfate 2.5 mg, butabarb-
itol 8 mg / Tab Bot 100s 1000s
Use: Analgesic

CRYSTODIGIN, NNR 1950 (Lilly) Cryst
d gitoxin Tab 0.05 mg (Yellow) 0.1 mg
(White) and D 15 mg (Yellow) (scored) 0.2
mg Pkg 100s, 500s
Use: Congestive heart failure

CRYSTODIGIN SOLUTION (Lilly) Cryst
digitoxin Amp (0.2 mg/cc) 1 cc, 10 cc
Use: Congestive heart failure (Slow IV,
may be administered IM).

CRYSTOGEN, (Metro) Estrone Viot (2 mg /
cc) 10 cc; (5 mg / cc) 10 cc
Use: Menopausal syndrome

CRYSTOIDS ANTHELMINTIC. (Merck Sharp
& Dohme) Hexylresorcinol Large 0.2 Gm,
Vial 5 pills; small 0.1 Gm Vial 6 pills

CRYSTWEL, NND 1962 (U S Standard)
Cyanocobalamin USP, Vit B 12 Vials
(30 mcgm / cc) (60 mcgm / cc) (100
mcgm / cc) (1000 mcgm / cc) 10 cc

C.S.A. (A.P.C.) Aspirin 3 1/2 gr., Col succi-
nate 2.8 gr / EC Tab Bot 100s
Use: (1 2 q 3 h) Analgesic & antipyretic

C.S.R. UNIT. Unit represents the amount of
veratrum viride extract per Kg of body
weight which will just abolish the pressor
response to the carotid sinus reflex result-
ing from occlusion of the carotid arteries
in dogs

CTAB. See: Cetyl Trimethyl Ammonium Bro-
mide

CLFERTIL, (Lakeside) Vit B-12 3 mcg, fer-
rous sulfate basic, 200 mg, chlorophyllin,
Sod, Pot & copper salt 25 mg / Tab Bot
100s

Use (1 1 d) Hypashramic enemas

CUMALTRON (Borshardt) Iron & ammonium
citrate 8 gr., copper sulphate 1/20 gr.,
cobalt sulfate 1/60 gr., Mang chloride 1
gr., red bone marrow 10 gr., Vit B 1 1/60
gr./R oz Syr Bot 1 pt, 1 gal

Use: Secondary anemias

CUMERTILIN, NND 1962 (Endo) Mercuma-
tin (each cc contains 39 mg of mercury
& 50 mg of theophylline) Uncoated Tab-
lets are 67.7 mg of sumatlin equiv to 20
mg each of mercury & theophylline Amp

Generic name recognized by N.P. U.S.P. or A.M.A.

1 cc, 2 cc Boxes T2, 25 & 100s, 10 cc
Boxes 1, 10 & 100s, Tab Bot 100s 1000s
Use: Mercurial diuretic

CUMERTILIN SOOUM, NND 1962 (Endo)
Mercumatin Sod Mercury 39 mg, the-
ophylline 50 mg / cc Amp 1 cc, 2 cc,
Box 6s 12s 100s Tabs (677 mg) Bot
100s

Use: Mercurial theophylline diuretic

CUMETHAROL. Name Approved by B.P.C.
4,4-Dihydroxy 3,3'-(2-methoxyethylidene)
dicoumarol Dicoumoxone

CUMOPYRAN, NND 1958 (Abbott) Cyclo-
cumarol Tab 25 mg / Tab Bot 100s, 1000s
Use: Anticoagulant

CUM-VATINE, (Smith Dorsey) Vit A 25,000
u, D 1000 u, B 1 10 mg, B 2 5 mg, nic-
otinamide 150 mg & ascorbic acid 150
mg / Cap Bot 100s
Use: Oretory supplement

CUPERTIN CREAM, (Cooper, Tinsley) Di-
phenylpyridine HCl, 2% in hydrophylic oint
Tube 15 Gm

Use: Fungicidal Antipruritic

CUPREX, (Merck Sharp & Dohme) Copper
oleate w/tetrolin, acetone & liquid paraffin
Bot 3 oz, 1 pt

Use: Topically, pediculicide

***CUPRIC SULFATE, NF** W/Zinc sulfate &
camphor See Dolibour, Pow (Doak)

CUPRON, (Ayers) Ferrous carb 0.4 Gm-
cuprous carb 0.0026 Gm / Cop Bot 100s
Use: Orally, iron & copper therapy

CUPROUS CARBONATE W/Ferrous car-
bonate See: Cupron, Cop (Ayers)

CURARE, NND 1962, ADR

See: Chandodendron Tomentosum Est
NND 1959

Intocostin (Squibb)
d Tubocurarine (Various Mfr)

Use: Muscle relaxant.

CURARE ANTAOONIST.

See: Neostigmine Methylsulfate (Various
Mfr)

Temilon, Amp (Hoffmann La Roche)

CURRAL See: Diethyl Barbituric Acid (Vari-
ous Mfr)

CUTE-ASEPTO OINTMENT. (Hart) Salicylic
acid 4%, chlorobutanol 5%, oil of cade
0.3%, liquid menthol 3%, zinc oxide 4%
in a lanolin base Tubes 1 oz, jars 1 lb
Use: Fungus Infections

CUTE-ASEPTO POWDER (Hart) Sodium thio-
sulfate 10%, zinc oxide 10%, boric acid
14%, carbonic acid (phenol) 1.2% w/cam-
phor, oil of cloves liquid menthol in a
base of aluminum hydroxide, kaolin, mag

nesium carbonate and rice starch. Shaker top cans 2 oz.

Use: Fungus reinfection.

CUTEMOL EMOLLIENT BATH OIL (Summers Labs) Hypoallergenic lanolin fractions with additional emollient agents and nonionic dispersant /Bot. Bot. 6 oz.

Use: Bath additive.

CUTEMOL EMOLLIENT CREAM (Summers Labs) Allantoin 0.2%, hypoallergenic lanolin fractions, with additional emollient agents and a nonionic emulsifier in a high-aqueous content water-in-oil cream base. Tube 2 oz.

Use: Dermatologic.

CUTICURA MEDICATED CREAM (Patter D & C) Bithional, tyrosine, para-chlorometaxylenol, allantoin, menthol, cetyl alcohol. Cream. Tube. ½ oz.

Use: Dermatologic.

CUTICURA SOAP (Cuticura) 3,5,4-Tribromo salicylonilide 80%, brominated salicylonilides 20%. Cakes 3 ½ oz., 6 oz.

Use: Skin care.

CUTONE (Sutliff & Cose) Phenol 2%, zinc sulf. 1.1%, zinc oxide 4%, starch 1%, glycerin 6.8%. Tot. 1 pt., 1 gal.

Use: Antiseptic & protective

C.V.P. (U. S. Vit. Corp) Citrus bio-flavonoid comp. 100 mg, ascorbic acid 100 mg / Cap. Bot. 100s, 500s, 1000s. Also double strength as Dua-C.V.P.

Use: To increase capillary resistance & decrease fragility.

C.V.P. SYRUP (U. S. Vitamin) Citrus bio-flavonoid comp. 100 mg, ascorbic acid 100 mg./5 cc. Bot. 4 & 16 oz., 1 gal.

Use: (3-6 tsp. daily) Capillary tone & resistance.

C.V.P. w/VITAMIN K (U. S. Vitamin) Citrus bioflavonoid compound 100 mg, ascorbic acid 100 mg., menadione 0.66 mg./tab. or 5 cc. Syr., Bot. 4 oz., 1 pt., 1 gal. Tab., Bot. 100s, 1000s.

Use: To increase capillary resistance and maintenance of normal prothrombin levels.

C VITAMIN. See: Ascorbic Acid.

CYADESINE (Superior Pharm) Adenosine-5-monophosphate 25 mg, Vit. B-12 crystalline 60 mcg./cc. Vial 10 cc.

Use: Regulating carbohydrate metabolism.

CYALPH (Massengill) allylphenylbarbituric acid ¾ gr., cyclophen [Cyclopentenylallylbarbituric acid] ¾ gr./cap. Bot. 100s, 500s, 1000s

Use: (1-2 q. 3 h) Sedative hypnotic.

CYAMOPSIS TETRAGONOLOBA GUM. See: Regonal (Organon).

CYANAL (Lloyd Bros) Thiamine mononitrate

* Generic name recognized by N.F., U.S.P. or A.M.A.

10 mg, Vit. B-12 25 mcg, pyridazine HCl 5 mg /Tab. Bot 50s.

Use: To stimulate appetite.

CYANOBEX (Metropolitan Labs) Vitamin B-12 20 mcg, B-1 HCl 10 mg, B 6 HCl 3 mg, B-2 10 mg., pantothenol 4 mg., niacinamide 25 mg., liver inj. 20 mcg per cc. D.O.S cc./cc. Vials 10 cc.

Use: I.M., vitamin B complex therapy.

***CYANOCOBALAMIN, U.S.P., N.N.D. 1962.** A.D.2. Inj. U.S.P., N.N.R. See: Vitamin B-12, Progs.

CYANO-GIL (Mourry Biological) Cyanocobalamin 200 mCg or 1000 mCg, gelatin menstrum 15%, preservative/cc. Vial 10 cc. Use: Pernicious anemia.

CYANOPLEX (Myers-Carter) Liver Inj. 100 mg, ferrous gluconate 0.66 gr., B-2 0.3 mg, niacinamide 100 mg., dl-methionine 10 mg., L-cysteine HCl 1 mg./2 cc. Vial 30 cc.

CYANOSINE GEL (Vitamix) Adenosine-5-monophosphate cad. 25 mg. & B-12 100 mcg., with gelatine/cc. Vial 10 cc. Also avail. as HP with 100 mg/cc.

Use: I.M., beriberi, neuritis and other neuromuscular conditions

CYCLAMINE HYDROCHLORIDE, N.N.D. 1962 (Merck Sharp & Dohme) Hezylcaline HCl. Amp. (2.5% w/dextrose 10%) 2 cc; Amp. (2%) 20 cc; Vial (1%) 30 cc; Bot. (5%) 60 cc; Sol. Bot. (5%) 60 cc; Jelly (5%) Tube 1 oz.

Use: Local anesthetic.

CYCLAMATE CALCIUM & SODIUM, N.N.D. 1962, A.D.R Cyclohexylsulfamate dihydrate salt. See: Sucaryl Salts. (Abbott)

CYCLAMYCIN, N.N.D. 1962. (Wyeth) Triacetylfalendamyacin Vial 125 mg, 250 mg, 365 Oral Susp. (125 mg /5 cc) Bot 2 oz.

Use: Infectious diseases.

***CYCLANDELATE, N.N.D. 1962.** 3, 5, 5-Tri-methylcyclohexyl mandelate. See: Cyclospasmal (Ives-Cameron)

CYCLEX (Merck Sharp & Dohme). Hydrochlorothiazide, 25 mg; meprobamate, 200 mg /Tab Bot. 100s, 1000s

Use: Premenstrual tension with edema

***CYCLIZINE, U.S.P., N.N.D 1962.** Tab U.S.P. 1-Diphenylmethyl-4-methylpiperazine N-Benzhydryl-N methyl piperazine hydrochloride or lactate.

See: Marezine Hydrochloride. (Burroughs Wellcome)

W/Ergolamate tartrate, caffeine. See: Migrol. Tab. (Burroughs Wellcome)

CYCLOBAR. (Schieffelin) Cyclopentolate

- HCl, 1% neomycin citrate 0.55% / 3.54
Gm collapsible tube
Use Mydriatic with antibiotic
- ***CYCLOBARBITAL CALCIUM, NF X**,
N.N.R. 1951, Tab NF X Cyclobarbitone
namuron, palinum, cyclobarbitol Cal 5 (1
Cyclohexenyl) 5 Ethylbarbituric Acid See
Phanador (Winthrop Labs)
Use: Hypnotic & sedative w/intermediate
action
- CYCLOBARBITONE**. See: Cyclobarbitol
- CYCO-BELL** (Moore Kirk Labs) Hyoscina
HBr 0.0065 mg Hyoscyamine HBr 0.1037
mg, Atropine sulfata 0.0194 mg, Triple
Barb (33 1/3 each of phenobarbital sodium
butobarbital sodium pentobarbital sodium)
16.2 mg / Tab ar 5 cc Elixir 4 oz, 1 pt
Tablets Bot 100s
Use: Antispasmodic and sedative
- ***CYCLOBENZAPRINE** NCI 5 (3 Dimethyl
aminopropylidene) dibenzo(a,e)cyclohepta
triene HCl
- ***CYCLOCUMAROL**, N.N.D. 1959 See
Cumapyran, Tab Not listed (Abbott)
- CYLOGESTERIN**, (Upjohn)
Aq, Sus; Estrogenic substance 10 000 u &
Progesterone 10 mg / cc Vial 1 cc, Box 3s
25s 50s 100s, 5 cc, 1s, 5s, 25s, 50s, 100s
Estrogenic substance 20 000 u & progeste-
rone 25 mg / cc, Vial 5 cc, Box 5s, 25s,
50s, 100s
Tab; Estrogenic substance 10 000 u & pro-
gesterone 30 mg / Tab, Bot 15s
Use: Estrogenic & corpus luteum therapy
- CYLOGYL**, N.N.D. 1962 (Schiefelin) Cyclo-
pentolate HCl Sol 1/2 Bot 1/2 oz for
Caucasian races; Sol 1% Bot 1/2 oz for
dark skinned races 2% Bot 2 cc, 7.5 cc
Gel 1% Tube 3.54 Gm
Use: Cyclaplegic & mydriatic.
- CYCLOHEXALAMINE** (Parke Davis) 1 phen-
ylcyclohexylamine
- CYCLOHEXYSULFAMATE SODIUM**, N.N.D.
1962 See: Sularyl Sodium, Prep (Abbott)
- ***CYCLOMETHYCAINE**, N.N.D. 1962 See
Surfacaine Salts (Lilly)
- CYCLOMETHYCAINE AND THENYLPYRA-
MINE**. See: Histadyl and Surfacaine
(Lilly).
- CYCLOMYDRIL**, (Schiefelin) Phenylephrine
HCl 1%, cyclopentolate HCl 0.2% in a
sterile ophthalmic solution Bot 2 ml, &
7.5 ml
Use: Mydriatic
- CYCLONAL SODIUM**, (May & Baker) Nexa-
barbital sod, USP See Hexabarbital Solu-
ble (Various Mfr)
Use: Sedative
- CYCLOPAL**, (Upjohn) 5 Allyl 5-(2 cyclopen-
ten-1-yl) barbituric acid Cyclopentenyl allyl-
barbituric acid Cap 3/4 gr. & 2 1/4 gr Bot
100s, 500s, 1000s.
Use: Sedative
- W/Aspirin. (Upjohn) Cyclopal 3/4 gr, aspirin
5 gr./Tab Bot 100s, 500s.
Use: (1/2 tid, prn) Sedative with anal-
gesic
- W/Ephedrine See: Ephedrate, Cap (Up-
john)
- W/Ephedrine sulfate & theophylline See
Zantrate, Tab (Upjohn)
- CYCLOPAL W/P-A-C** See: P A C W/Cylo-
pal, Tab (Upjohn)
- CYCLOPEN**, (Massengill) Cyclopentenylallyl
barbituric acid 1 1/2 gr / Cap Bot 500s,
100s
Use: Sedative, hypnotic
- W/Alphaxem See: Cycalph, Cap (Massengill)
- ***CYCLOPENTAMINE HYDROCHLORIDE**,
NF, N.N.D. 1962 Inj, Soln NF See:
Clonane hydrochloride (Lilly)
- W/Aludrine, atropine sulfate & procaine HCl
See: Aerolone Compound, Sol (Lilly)
- W/Thenylpyramine. See: Histo Clonane
Cap, Liq (Lilly)
- CYCLOPENTENYL - ALLYL - BARBITURIC
ACID**.
See: Cyclopal, Cap (Upjohn)
Cyclopen, Cap (Massengill)
- W/Allylphenyl barbituric acid See: Cycalph
Cap (Massengill)
- ***CYCLOPENTHIAZIDE**, (Ciba) 6 chloro 3
(Cyclopentylmethyl) 3,4-dihydro 7 sulfamyl
2H, 1,2,4 benzothiadiazine 1,1 dioxide See
Navidrix
- ***CYCLOPENTOLATE HYDROCHLORIDE**,
USP, N.N.D. 1962 Ophth Soln, USP
beta dimethylaminoethyl (1 hydroxycyclopentyl)
phenylacetate HCl See: Cyclogyl, Sol
(Schiefelin)
- W/Phenylephrine HCl See: Cyclamylril, Sol
(Schiefelin)
- W/Neomycin citrate See: Cyclabar, Ophth.
(Schiefelin)
- CYCLOPENTYLPROPIONATE** See: Depo-
Testosterone, Vial (Upjohn)
- ***CYCLOPHOSPHAMIDE**, N.N.D. 1962 N,N-
bis(beta chloroethyl) N', arthro propylene
phosphoric acid ester diamide monohy-
drate See: Cycloxan, Tab (Mead Johnson)
- ***CYCLOPROPANE**, USP, N.N.D. 1962
A D R Trimethylene
Mallinckrodt—Cylinder 40 & 100 gal
Squibb—Cylinder 40, 100 & 200, 230 gal
Use: Inhalation anesthetic

CYCLOPYRROLATE. See: Cyclopyrronium Bromide.

* **CYCLOPYRRONIUM BROMIDE.** (Robins) 1-Ethyl-3-pyrrolidyl α -phenylcyclopentaneacetate methobromide.

Use: Anticholinergic.

* **CYCLOSERINE, N.N.D. 1962.** Is an antibiotic produced by Streptomycin orchidococcus.

See: Oxomycin, (Merck Sharp & Dohme) Seromycin, (Lilly)

CYCLOSPASMOL, N.N.D. 1962. (Ives-Cameron) Cyclandelate. 3,5,5-trimethylcyclohexyl mandelate 100 mg. and 200 mg./Tab. Bot. 100s.

Use: Vasospastic agent.

* **CYCLOTERENOL HCl.** (Mead Johnson) 3-(p-chlorophenyl)-5-methyl-3-pyrrolidinol HCl Use: Neurotropic bronchodilator.

* **CYCLOTHIAZIDE.** (Lilly) 6-chloro-3-(2-norbornenyl)-5,7-sulldimyl-3,4-dihydro-1,2,4-benzothiadiazine-1,1-dioxide. See: Anhydron (Lilly)

Use: Diuretic.

CYCLO-THRYCIN, (Hance) Tyrothricin 2 mg., Isobutyl p-aminobenzoate 5 mg., benzalkonium chloride 0.1%/Loz. Viol 16s. Bot. 1000s.

Use: Infections of throat & mouth.

CYCLOTROL SUSPENSION, (Hyrex) Sterile suspension of crystalline natural estrogen, 25,000 I.U./ml. equivalency of estrone, with progesterone U.S.P. 25 mg/ml Viol 10 cc Use: Estrogen.

CYCOTIN, (Reed & Carrick) Methylcellulose 300 mg. & d-amphetamine phosphate 2.5 mg./Tab. Bot. 100s, 500s, 1000s, 5000s.

Use: Appetite depressant.

* **CYCRIMINE HYDROCHLORIDE, N.F., N.N.D. 1962 Tabs., N.F. 1-phenyl-1-cyclopentyl-3-piperidino-1-propanol hydrochloride. See: Pagitane Hydrochloride (Lilly)**

CYDRIL, (S. J. Tutog & Co) Levamphetamine Succinate (1 isomer) alpha-2-phenylamino-propane succinate 7 mg/Tab 21 mg/Cap. Bot. Tabs. 100s, 500s, 1000s. Caps 100s, 1000s.

Use: Antiobesity agent for hunger control.

CYLAMIDE, (Chicago Pharmacol Co) N Acetyl-p-aminophenol 2½ gr., salicylamide 3½ gr., codeine phosph. ¼ gr., caffeine ½ gr./tab. Bot. 100s, 1000s.

Use: (1-2 t.i.d.) Analgesic.

W/Codeine or phenobarbital. (Chicago) N acetyl-p-aminophenol 2½ gr., salicylamide 3½ gr., codeine phosphate ¼ gr., caffeine ½ gr./Tab. Phenobarbital ¼ gr. in place of codeine. Bot. 100s.

* Generic name recognized by N.F., U.S.P. or A.M.A.

Use: Headaches, colds, grippe, muscular aches and pains.

W/Phenobarbital. (Chicago Pharmacol Co.) N-Acetyl-p-aminophenol 2½ gr., salicylamide 3½ gr., phenobarbital ¼ gr., caffeine ½ gr./tab. Bot. 100s, 1000s.

Uses (1-2 q. 4 h) Analgesic w/sedative.

CYLANDELATE, N.N.D. 1959. See: Cyclospasmol (Ives-Cameron)

CYLERT, (Nordson Pharm) *Magnesium Pemetine.

Use: Central nervous system stimulant.

CYMITAL, (Haack) Scopolamine HBr 0.13 mg, atropine sulfate 0.03 mg, phenobarbital 16.2 mg./Tab.

Use: Emotional stress.

CYNAL, (Lloyd Brothers) R₁₂ 25 mcg, with absorption-enhancing complex, B₁ 10 mg, and B₆ 5 mg./each "Cherrow-Chew" tab. Bot. 50s.

Use: Stimulate appetite.

CYNIDAL ELIXIR, (Rand) Atropine sulfate 1/200 gr., phenobarbital ¼ gr., Vit. B-1 1 mg, Vit. B-2 0.50 mg., niacinamide 5 mg, Cal. pantothenate 0.30 mg., pyridoxine HCl 0.15 mg., alcohol 16% by volume, other factors of the Vit. R-complex/cc, Bot. 1 pt., 1 gal.

CYNIDAL TABS, (Rand) Atropine sulfate 1/200 gr., phenobarbital ¼ gr., Vit. R-1 1 mg., B-2 0.50 mg., niacinamide 5 mg., Cal. pantothenate 0.30 mg., pyridoxine HCl 0.15 mg./Tab. Bot. 100s, 1000s.

* **CYPROHIPTADINE HYDROCHLORIDE, 1-methyl-4-[5-dibenzo[e,e]cyclohexetrenylidene]piperidine HCl. See: Perioctin HCl (Merck Sharp & Dohme)**

W/Dexamethasone. See: Dronactin, Tabs. (Merck Sharp & Dohme)

CYREN A. See: Diethylstilbestrol. (Various Mfrs.)

CYSTAMIN. See: Methenamine. (Various Mfrs.)

CYSTINE, (Various Mfrs) L-Cystine.

Delta—Bot. 100 Gm.

Mann—Bot. 25 Gm.

Use: An amino acid.

CYTITARS, (Walker Corp) Atropine sulfate 1/1000 gr., hydrangea 1¼ gr., uva ursi 1½ gr., triticum ¾ gr., corn silk 1 gr., in an eff. base/Tab.

Use: Cystitis.

CYSTIZEM, (Zemmer) Phenylazo-diaminopyridine HCl 50 mg, sulfacetamide 250 mg., atropine methylnitrate 1 mg./Tab. Bot. 100s.

Use: G-U antiseptic.

CYSTODYNE, (Tyree) Methenamine, buchu.

- uva ursi, hops, pareira brava, Hyoscyamus & Pot acetate Pow Bot 4 1/2 oz, 10 oz
Use Orally, urinary antiseptic
- CYSTORRHOL.** (Buffington) Hyoscyamus ext 1/16 gr, atropine sulfate 1/500 gr, bexo methenamine 2 1/2 gr, salol 1 gr, acid benzoic 1/10 gr/Tab Bot 100s, 1000s
Use (3 daily) Antispasmodic & urinary antiseptic
- CYTAL** (Cutter) A concentrated solution When diluted with 9 parts of water containing sorbitol 27 Gm, mannitol 0.54 Gm, methylben 0.5 mg propylben 0.1 mg & butylben 0.1 mg Liter bot
Use Urologic irrigating fluid Not used IV or subcu
- CYTELIN, NND 1962** (Lilly) Beta & dihydrobrato sitosterols Susp 20%, Bot 1 pl
Use Orally, lipotropic
- CYTINIC.** (Standard Pharmacol) Ferrous gluconate 200 mg, Vit B 12 10 mcg, duode-num subs 150 mg, liver subs des 100 mg, folic acid 0.67 mg, Vit C 50 mg/Cap Bot 50s 100s 1000s
Use (2 11 d) Anti anemia therapy
- CYTOBEX.** (Parke, Davis) Vit B 12 15 mcg/cc Amp 1 cc Box 25s
Use Pernicious anemias
- CYTOCYM.** See Thromboplastin (Various Mfr).
- CYTOFERIN** (Ayerst) Ferrous fumarate 200 mg, ascorbic acid 150 mg/Tab or 2 dr Bot 100s 1000s Liq Bot 8 fl oz
Use: (1 2 11 d) Anemias
- CYTOLAY.** (Armour) Chymatrysin, lyophilized 7 mg/Cap Bot 25s
Use: Diagnostic agent
- CYTOMEL, HND 1962** (Smith Kline & French) Sod lithium 1 Triiodothyronine sod Tab 5 mcg, 25 mcg, Bot 100s
Use: Orally, 25 to 50 mcg daily metabolic insufficiency
- CYTORA.** (Organon) Folic acid 0.75 mg ferrous sulfate essic, 100 mg, liver concentrate 150 mg, Vit C 25 mg, B 11 mg, B 2 1 mg, B-6 0.5 mg, niacinamide 5 mg, Cal pantothenate 1 mg, Vit B 12 w/ Intrinsic factor 0.05 mcg/Tab Bot 100s, 250s, 1000s
Use: Hematinic
- CYTOXAN, HND 1962** (Mead Johnson) Cyclophosphamide 2H 1,3,2-oxazophosphorine 2 [bis(2 chloroethyl)amino]tetahydro-2,2-oxide Inj Vial 100 mg, 200 mg Tab, 50 mg Bot 100s
Use: Certain forms of cancer
- CYTRIN** (Upjohn) Medroxyprogesterone acetate (Provera) 25 mg, ethasolamide (Cardose carbonic anhydrase inhibitor) 35 mg ectylurea (Levanil) 300 mg/Tab Bot 20s 400s
Use Premenstrual tension syndrome
- CZO W/ANTIHISTAMINE.** (Massengill) Calamine, prep 30 gr, zinc oxide 30 gr, phenylazylamine dihydrogen citrate 1%, glycerin & aromatics qs/fl az Bot 3 oz 1 pt, 1 gal
Use Dermatitis.
- CZO LOTION W/PHENOL.** (Massengill) Calamine, prep 30 gr, zinc oxide 30 gr, carbolic acid, liq 5 min, glycerin & aromatics qs/fl az Bot 3 oz, 1 pt, 1 gal
Use Emollient, astringent, antipruritic.
- W/Benzocaine** (Massengill) Calamine, prep gr 40, zinc oxide gr 20, benzocaine 2% aromatics/fl oz Lat Bot 3 oz, 1 pt, 1 gal
Use: Antipruritic & analgesic lotion
- DAA.** See Dihydroxy Aluminum Aminoacolate
- DACRIOSE** (Crookes Barnes) Isotonic solution of boric acid potassium chloride, sodium carbonate/Bal 120 cc
Use Ophthalmic irrigating solution
- DACTIL, NND 1962** (Lakeside) Pipedolate N ethyl 3 piperidyl diphenylacetate HCl Cap 50 mg Bot 50s
Use: (1 qid) Postganglionic parasympathetic, relieving smooth muscle spasm
- W/Phenobarbital** 16 mg Cap Bot 50s
Use Eutonic agent produces normal tons of viscous, antispasmodic
- W/Pipilol** See Tridal, Tab (Lakeside)
- DACTIL-OB.** (Lakeside) Piperidolate HCl 100 mg, ascorbic acid 50 mg, hesperidin complex 50 mg/Tab Bot 100s
Use Prevention of premature delivery
- DACTILASE.** (Lakeside) Piperidolate HCl 50 mg, standardized cellulolytic enzyme 2 mg amylolytic enzyme 15 mg, proteolytic enzyme 10 mg pancreatin 3X 100 mg, tauracholic acid 15 mg/Tab Bot 60s
Use Antispasmodic w/digestive enzymes
- DACTRON** (G F Harvey) Digitalis flow 1/4 gr, Strophanthus p ext 1.40 gr, cactus grand p ext 1/10 gr, strychnine sulfats 1/60 gr, nitroglycerin 1/200 gr/Pill Bot 100s, 1000s
Use: Cardiac stimulant and tonic
- DADEX** (Central) Dextroamphetamine sulf 15 mg/TD Cap Bot 100s, 500s
Use CNS stimulant
- DADEXAL.** (Central Pharmacol) Dextroamphetamine sulf 15 mg, amobarbital 60 mg/TD Cap Bot 100s, 500s
Use: CNS stimulant w/sedative
- DAILYVITE.** (Coral) Vit A 4000 u, Vit D 400 u, Vit B 1 1 mg, Vit B 2 12 mg, nicotinamide 10 mg, pyridoxine hydrochloride

ride 0.1 mg., calcium pantothenate 1 mg., folic acid 0.1 mg., Vit. B-12 1 mg., Vit. C 30 mg., Iron 5 mg., cabot 0.1 mg., copper 0.5 mg., iodine 0.05 mg., magnesium 1 mg., manganese 0.1 mg., zinc 0.1 mg. /Cap. Bot. 100s, 1000s.

Use: Vitamin-mineral supplement.

DAINITE TABLETS. (Irwin, Neisler) Day Tab. Sod. pentobarbital $\frac{1}{2}$ gr., aminophylline 3 gr., ephedrine HCl $\frac{1}{2}$ gr., ethyl aminobenzoate $\frac{1}{2}$ gr., aluminum hydroxide $2\frac{1}{2}$ gr. / Tab. Night Tab. Phenobarbital $\frac{3}{4}$ gr., Sod. pentobarbital $\frac{1}{2}$ gr., aminophylline 4 gr., ethyl aminobenzoate $\frac{1}{2}$ gr., aluminum hydroxide $2\frac{1}{2}$ gr. / Tab. Unit contains 50 day (pink) tab. & 20 night (black) tab.

Use: Therapy in treatment of bronchial asthma.

DAINITE-KL. (Irwin, Neisler) Potassium iodide 5 gr. & Ingrid. of Day tab. Cap. Bot. 50s, 500s, 1000s

Use: Treatment of asthma

DAINITE PEDIATRIC. (Irwin, Neisler) Aminophylline 1 gr., ephedrine HCl 1/12 gr., phenobarbital $\frac{1}{4}$ gr., ethyl aminobenzoate $\frac{1}{2}$ gr., aluminum hydroxide $\frac{1}{2}$ gr. / Tab. Bot. 100s.

Use: Treatment of asthma.

DAKIN'S SOLUTION. See: Sodium Hypochlorite Solution, Diluted, N.F.

DALACIN. (Upjohn) See: Streptovaricin.

DALACTUM. (Mead Johnson) Low fat milk & dextrin-maltose w/added Vit. D. Con 13 fl. oz.

Use: Feeding of premature & full term infants with low fat tolerance.

DALCA. (Ascher) Aluminum aspirin 300 mg., phenylephrine HCl 10 mg., chlorpheniramine maleate 2 mg., phenindamine tartrate 5 mg. / Tab. Bot. 100s, 500s

Use: Cold therapy.

DALDRIN. (Merck Sharp & Dahme) Codeine phosphate 60 mg., chloralform 60 mg., propadrine HCl 60 mg., menthol 12 mg., Ipecac flex. 0.12 cc., sod. citrate 1.2 Gm. / 30 cc. Bot. 1 pt., 1 gal.

Use: (1 tsp. 2-3 h) For relief of coughs.

DALE GOAT MILK. (Cutter) Fat 27%, protein 25%, lactose 38.5%, cal./fl. oz. 142, ash 6.5%, water 2%, total solids 98%. Paw. 8 oz., 1 lb.

Use: Dietary & digestive disturbances.

DALIBOUR POWDER. (Daak) Cupric sulfate 8%, zinc sulfate 28%, camphor w/buffer. Pkg. 4 oz.

Use: Wet dressing in treatment of infections of skin.

DALICIN. (Upjohn) Streptovaricin. Ueder study

Use: Treatment of tuberculosis

DALIDOME. (Dome Chemicals) Copper sulfate, zinc sulfate, camphor, specially buffered to pH 4 & Pow. Pkg. 2.2 Gm., 4 oz. & 1 lb. for making Dalibour sol.

Use: Bacteriostatic, astringent wet dressing for skin.

DA-11-KAPS. (Ulmer) Vit. A 5000 u., D 1000 u., thiamine HCl 3 mg., riboflavin 3 mg., pyridoxine 0.1 mg., ascorbic acid 37.5 mg., Cal. pantothenate 1 mg., nicotinamide 20 mg., mixed tocopherols 2 mg. / Cap. Bot. 100s, 500s, 1000s.

Use: Vitamin therapy.

DALIVIM FORTE. (Dalin) Iron ammonium citrate 18 gr., liver fraction 3 gr., Vit. B-12 30 mCg., A 30,000 u., D 3000 u., B-1 12 mg., B-2 4 mg., niacinamide 20 mg., Vit. B-6 2 mg., calcium pantothenate 12 mg., mixed tocopherols 30 mg., calcium glycerophosphate 65 mg., manganese glycerophosphate 17.5 mg. / oz. Bot. 4 oz., 12 oz.

Use: Hematitic and dietary supplement.

DALSOL. (U.S. Vitamin) Vit. D 50,000 u., A 5000 u., ascorbic acid 75 mg., thiamine 3 mg., riboflavin 3 mg., pyridoxine 0.33 mg., niacin 15 mg., niacinamide 15 mg., Cal. pantothenate 3 mg., alpha tocopherol 7.5 mg. / Cap. Bot. 100s, 250s, 500s, 1000s.

Use: Vitamin therapy.

DALYDT. (Hynson, Wescott & Dunning) 3,5-Dibromosalicylaldehyde. Dusting pow. 2%. Cantoleers 1 oz. Oint. 2%, 2 oz. Sol. 0.5%, Bot. 2 fl. oz.

Use: Topically, antiseptic against gram-negative bacteria & fungicidal.

DALZINATE. Not listed (Smith, Miller & Fitch) Phenobarbital $\frac{1}{2}$ gr., hamatropine methylbromide 150 gr., aluminum dihydroxy aminoacetate $7\frac{1}{2}$ gr., Vit. C 25 mg. / Tab. Bot. 100s, 500s.

Use: (1-2 fl. oz., p.c. & h.s.) Antispasmodic & antacid.

DAMASON. (Mason Pharmaceuticals) Codeine phosphate $\frac{1}{2}$ gr., acetylsalicylic acid $3\frac{1}{2}$ gr., phenacetin $2\frac{1}{2}$ gr., caffeine $\frac{1}{2}$ gr. / Cap. Bot. 100s

Use: Analgesic and antipyretic.

DAMASON-P. (Mason Pharmaceuticals) Dihydrocodeinone bitartrate 5 mg., hamatropine hydrobromide 0.38 mg., acetylsalicylic acid 224 mg., phenacetin 160 mg., caffeine 32 mg. / Tab. Bot. 100s

Use: Analgesic.

DAMBOSE. See: Inaitol, U.S.P.

D-AMFETASUL. (Pitman Moore) Dextroamphetamine sulfate. Tab. 5 mg. Bot. 100s.

Use: Central nervous system stimulant.

DAMITAL. (U.S. Standard Products) d-Amphetamine sulfate 2.5 mg., pentobarbital

- sad 1/4 gr, acetaphenelidin 2 1/2 gr, aspirin 3 1/2 gr / Tab Bot 100s, 1000s
Use (1 2 1 d) Treatment of pain, anxiety, & depression
- DANILONE**, NND 1962 (Schieffelin) Phenadione Tab 50 mg Bot 100s
Use Orally blood anticoagulant
- DANISH BENZYL LOTION** (Tilden) Benzyl benzoate 27% by weight, stearic acid lanolin, triethanolamine, water Bot 4 oz, pt gal
Use Scabicide
- DANISN OINTMENT** (Tilden) Sulfur 10%, potassium 5%, lanolin, petrolatum, water Jar 2 oz, lb
Use Scabicide
- DAN-OCTYL**, (Mid Atlantic) Danthran 50 mg, diethyl sodium sulfosuccinate 100 mg / cap Bot 30s
Use Treatment of constipation
- ***DANTHRON**, NF 18 Dihydroxy anthraquinone, dioxyanthraquinone
See: Dianone, Tab (Klug)
Darbonate, Prep (Schenley)
Ishzin Prep (Winthrop)
W/Calcium bis (diethyl sulfosuccinate) See Daxidan Cop (Lloyd Bros)
W/d Calcium pantothenate See: Modane Tab (Warren Teed)
V/Diacetylhydroxyphenylisatin See: Vio-lax, Tab (Rowell)
V/Diethyl sod sulfosuccinate
See: Diacetyl W/Danthran, Tab (West Ward)
Dorbantyl, Cap (Schenley)
Doxan, Cap (Lloyd Bros)
Daxinate Danthran, Cap (Lloyd Bros)
Doxitul, Tab (Crestmed)
V/Mg(OH)₂, prune concentrate powder See Danthro-Lax, Tab (Mallard)
- DANTHRO-LAX**, (Mallard) Danthran 100 mg, Mg(OH)₂ 3 gr, prune concentrate powder 3 gr / Tab Bot 100s, 1000s
Use: Functional constipation
- DAPANONE NBr**, (Merck Sharp & Dohme) See: Phenisonone HBr
- DAPRISAL**, (Smith Kline & French) d Amphetamine sulfate 5 mg, amobarbital 1/2 gr, acetylsalicylic acid 2 1/2 gr, phenacetin 2 1/2 gr / Tab Bot 50s, 500s
Use: (1 2 q 3 h) Analgesic w/central nervous system stimulant
- DAPSONE** (DIAMINODIPHENYLSULFONE), USP Tab USP See: Avlosulfon, Tab (Ayerst)
- DAPTA**, (Wyeth) Vit A 10,000 u, D 2000 u thiamine HCl 3 mg, riboflavin DS mg, nicotinamide 15 mg, B 6 2 mg, ascorbic acid 100 mg/cc. Bot w/dropper 15 cc., 30 cc.
* Generic name recognized by NF, USP or A.M.A.
- Uses Orally, vitamin therapy & prophylaxis for infants & children
- DAPTAZOLE**, Amiphenazole 2 4 Diamino-5-phenyl thiazole (JAMA, 163, 1341, 1957)
Use: Barbiturate and morphine antagonist
- DAPTEN** (Davis & Sly) Scopolamine HBr 0.15 mg, Prophepyridamine maleate 10 mg, Emecaine 6 gr (ethyl aminobenzoate and aluminum hydroxide 1 S) / Tab Bot 100s, 1000s
Use Nausea, morning sickness, radiation sickness
- DAQIN**, (Riker) Chlorazoxoni HCl 150 mg / Tab Bot 100s
Use Diuretic; Carbonic anhydrase inhibitor
- DARANIDE**, NND 1962 (Merck Sharp & Dohme) Dichlorphenamide Tab 50 mg Bot 100s
Use: Carbonic anhydrase inhibitor
- DARAPRIM**, NND 1962 (Burroughs Wellcome) (Pyrimethamine) 2,4 Diamino-5 p-chlorophenyl 6 ethylprimidine Tab 25 mg Box 30s Bot 100s
Use Orally, antimalarial,
- DARATHON** (Grant) Glyceryl guaiacolate 100 mg, Phenylpropanolamine HCl 12 mg, Pheniramine maleate 7 mg / 5 cc Bot 1 pt
Use Temporary relief of coughs due to colds
- DARSACIN**, (Crestmed) Phenylephrine HCl 10 mg; prophepyridamine maleate 6 mg; methoprylene HCl, 8 mg, pyrimidine maleate, 8 mg, acetylsalicylic acid, 240 mg; phenacetin, 120 mg; caffeine alkaloid, 30 mg; Vit C, 50 mg; hesperidin purified, 50 mg / Cap Bot 50s
Use Cold therapy
- DARBD**, NND 1962 (Smith, Kline & French) Isopropamide iodide, (3 Carbamoyl 3,3 d phenylpropyl) diisopropylmethyl ammonium iodide 5 mg / Tab Bot 50s
Use: Anticholinergic
W/Campazine See: Cambid Sponsule, Cap (Smith, Kline & French)
- DARCIL**, NND 1962 (Wyeth) Phenethicillin potassium 125 mg L 250 mg / Tab Bot 36s
Use: Infections due to penicillin susceptible organisms
- DAKENTININ**, (Burroughs Wellcome) Brevyl sum Tosylate 200 mg / Tab Bot 100s
Use: Antidrenergic, antihypertensive agent
- DARFEKIN**, (Dartell) Iron 100 mg, Vit B-12 12 mcg, folic acid 500 mcg, B-6 1 mg C 100 mg, liver stomach concentrate containing intrinsic factor activity 250 mg-glycine 112 mg / Tab Bot 100s & 500s
Use: Nematonic
- DARICON**, NND 1962 (Pfizer) Oxypentidine HCl 10 mg / Tab Bot 60s, 500s
Use: G-I & G-U disorders

- DARICON-PS.** (Pfizer) *Oxyphenylimine* HCl 5 mg., phenobarbital 15 mg./Tab. Bot. 100s.
Use: Anticholinergic with sedative.
- DARILAN.** (Davies, Base) Besorolol, peruvian balsam, bismuth subcarb., zinc oxide, carbonic acid, juniper tar. Oint., Jar 1 oz., 4 oz.
Use: Dermatitis.
- DARO.** (Testagar) See: Tridex, Tabs.
- DAR-PLEX.** (Darwin) Thiamine HCl 500 mg., riboflavin 50 mg., pyridoxine HCl 100 mg., ascorbic acid 100 mg., niacinamide 500 mg., Sod. pantothenate 100 mg., gentisic acid 2%, Vit. B-12 cryst. 50 mcg./Pkg. Bot. 100s. Thiamine HCl 25 mg., riboflavin 2.5 mg., pyridoxine HCl 5 mg., Vit. B-12 cryst. 2.5 mcg., ascorbic acid 50 mg., niacinamide 25 mg., Sod. pantothenate 100 mg., gentisic acid 2%/cc. Vial 20 cc.
Use: Multi-vitamin therapy.
- DARROWS-POTASSIC SALINE SOLUTION.** (Don Baxter, Inc.) See: Potassic Saline Injection, Lactated, U.S.P.
- DARROW'S SOLUTION, U.S.P.** (Don Baxter) Pot. chloride 260 mg., Sod. chloride 400 mg., Sod. r-lactate \$90 mg./100 cc. Bot. 150 cc., 500 cc.
Use: Supplement for loss of body fluids.
- DARSTINE BROMIDE.** Not listed (Merck Sharp & Dohme) Mepiperphenidol 1-(3-hydroxy-5-methyl-4-phenylhexyl)-1-methylperidinium bromide.
Use: Anticholinergic agent in peptic ulcer.
- DARTAL, N.N.D. 1962.** (Searle) Thiopropazate dihydrochloride Tab. 2, 5, 10 mg Bot. 50's, 100c.
Use: Anesthetic agent
W/Pro-Banthine 15 mg., Dartal 5 mg./Tab. Bot. 100s.
- DARTN** (Denab) N-acetyl-p-amino-phenol 500 mg., neostigmine bromide 1.5 mg., hyoscine hydrobromide D.125 mg./Tab. Bot. 100c
Use: Analgesic.
- DARTN W/COLCHICINE.** (Denab) N-acetyl-p-amino-phenol, 500 mg., neostigmine bromide 1.5 mg., hyoscine hydrobromide 0.125 mg., calcichine 0.25 mg./Tab. Bot. 100s, 500s
Use: Analgesic.
- D-ARTHIN.** (G. F. Harvey) Vit. D 50,000 u./Cap. Bot. 100s, 1000s
Use: Vitamin D deficiencies.
- DARTHSOINOL.** (Boeing) Vit. D 50,000 u., A 5000 u., ascorbic acid 75 mg., thiamine HCl 3 mg., riboflavin 2 mg., pyridoxine HCl 0.3 mg., niacinamide 15 mg., Cal. pantothenate 1 mg., mixed tocopherols (type 4) 4 mg./Cap. Bot. 100s.
Use: (1 p.c., 1 d.) Antiarthritic, nutritional.
- DARTRAROID.** (New Eng. Phr. Co.) d-Amphetamine sulf. 10 mg, thyroid 1 gr./Tab. Bot. 50s
- D. A. S.** (Stayner) d-Amphetamine sulf. 5 mg./Tab. Bot. 100s, 1000c
- DARVON, N.N.D. 1962.** (Lilly) Dextro Propoxyphene HCl 32 mg, 65 mg./Pulv. Bot. 100s.
Use: Analgesic
W/A.B.A., ULTRAN. See: Darvo-Tran. Pulvule (Lilly)
- DARVON COMP.** (Lilly) Dextro Propoxyphene 32 mg, aspirin 227 mg, acetophenetic 162 mg., caffeine 32.4 mg./Pulv. Bot. 100s
Use: Antipyretic and analgesic
- DARVON COMPOUND-65.** (Lilly) Dextro propoxyphene HCl 65 mg, acetylsalicylic acid 227 mg, acetophenetidin 162 mg., caffeine 32.4 mg./Pulvule. Bot. 100c
Use: Acute, chronic or recurrent pain.
- DARVO-TRAN.** (Lilly) Dextro propoxyphene HCl 32 mg., acetylsalicylic acid 325 mg, phenaglycodel 150 mg./pulvule. Bot. 100s.
Use: Pain accompanied by tension and anxiety.
- DASTROL, N.N.D. 1962.** (Evon) Meperhesin. Tab. 0.25 Gm, 0.5 Gm. Bot. 100s.
Use: Muscle relaxant.
- DASIKON.** (Massengill) Semikan HCl (methapyrilone HCl) 3/4 gr., caffeine 1/2 gr., acetylsalicylic acid 3 gr., acetophenetidin 2 gr., atropine sulfate 1/1000 gr./Cap. Bot. 100s, 500s, 1000s.
Use: Colds.
- DASIN.** (Massengill) Dover's pow. 1/2 gr., acetylsalicylic acid 2 gr., acetophenetidin 1 1/2 gr., camphor 1/4 gr., caffeine 1/4 gr., atropine sulfate 1/500 gr., Sod. sulfate, dried, 7 1/2 gr./Cap. Bot. 100s, 500c, 1000c.
Use: (1 l.d., p.r.n.) Analgesic, antipyretic, diaphoretic & sedative.
- DASIN FOR CHILDREN.** (Massengill) Dover's pow 1/4 gr., acetylsalicylic acid 1/2 gr., acetophenetidin 1/2 gr., camphor, mono-bromated, 1/16 gr., caffeine 1/32 gr., atropine sulfate 1/2000 gr./Tab. Bot. 100s, 500s, 1000s.
Use: (p.r.n.) Analgesic, antipyretic, diaphoretic & sedative
- DASIN-CS.** (Massengill) Codeine sulfate 1/40 gr., Ipecac 1/20 gr., acetophenetidin 1 1/2 gr., atropine sulfate 1/500 gr., acetylsalicylic acid 2 gr., camphor 1/4 gr., caffeine 1/4 gr./cap. Bot. 100s, 500s, 1000s.
Use: (1 l.d., p.r.n.) Analgesic, antipyretic, diaphoretic & sedative.
- DATA.** (Klug) Aminophylline 150 mg, di-desoxyephedrine 8.5 mg, atain 1/20 gr.,

- thyroid $\frac{3}{4}$ gr, phenobarbital $\frac{3}{8}$ gr /Tab Bot 100s
Use Obesity
- DATA-C.** (New Mexico Pharm) D alpha Tocopheryl acetate equivalent to 100 IU of Vit E /Cap Bot 100s
- D-ATE UNODA Ph 747.** (Paul Maney) d Amphetamine sulfate 15 mg /delayed action cap Bot 100s
Use 1 daily; day long control of appetite in weight reduction
- DATURINE NBr.** See Hyoscyamine Salts, NF
- DAVITAMIN FORTIFIED.** (Irwin, Nessler) Vit A 5000 u, D 1000 u, B 1 3 mg, B 2 25 mg, B 6 075 mg, B 12 15 mcg, niacinamide 20 mg, cal pantothenate 5 mg, folic acid 0.34 mg, w/Cal phos, iron, iodine mang, cobalt, molybd, copper, zinc, mag, pot, & d alpha tocopherol acetate 5 mg /Cap Bot 100s, 500s, 1000s
Use Vitamin and mineral supplement
- DAVITAMIN IMPROVED.** (Irwin, Nessler) Vit A 5000 u, D 1000 u, B 1 15 mg, B 2 2 mg, B 6 01 mg C 37.5 mg, E 2 IU, cal pantothenate 1 mg, niacinamide 20 mg /Cap Bot 100s, 500s, 1000s
Use: Treatment for multiple vitamin deficiencies
- DAVITAMIN THERAPEUTIC.** (Irwin, Nessler) Vit A 25,000 u, D 1000 u, B 1 10 mg, B 2 5 mg, C 150 mg, niacinamide 150 mg /Cap Bot 100s, 500s, 1000s
Use Multiple vitamin deficiencies
- DAVITAMON K.** See Menadione (Various Mfr)
- DAVITAPNOS.** (Smith Osorsey) Caps Dibasic Cal phosphate $7\frac{1}{2}$ gr, ergosterol, equiv to 400 u Vit D Bot 100s, 1000s Wafers Dibasic Cal phosphate 15 gr, ergosterol, equiv to 400 u Vit D Bot 100s, 250s, 1000s
Use: Calcium & vitamin D therapy for tissue building
- DAVITIN.** (Ives Cameron) Vit D, synthetic Cap 5000 u Bot 100s, 1000s
Use: Orally, vitamin D therapy
- DAXALAN** (Dane) Coat tar, crude, 2.88% w/zinc oxide & starch Oint $2\frac{3}{4}$ oz, 5 oz, 1 $\frac{1}{2}$ lb, 6 $\frac{1}{2}$ lb Pediatric (1.44%) $2\frac{1}{4}$ oz, 4 oz, 1 lb, 5 $\frac{1}{4}$ lb
Use: Topically, antiseptic skin ointment
- DAVID.** (Roerig) Xanthal HCl
Use: Psychotropic agent
- DAYALETS.** (Abbott) Vit A 10,000 u, D 1000 u, thiamine mononitrate 5 mg, riboflavin 5 mg, nicotinamide 25 mg, pyridoxine HCl 2 mg, Vit B 12 2 mcg, folic acid 0.25 mg, Cal pantothenate 5 mg, ascorbic acid 100 mg /Filmtab Bot 50s 100s, 250s, 1000s
Use One tablet daily
- DAYALETS-M.** (Abbott) Same vit as Dayalets plus 9 minerals /Filmtab Bot 100s 250s
- DAYAMIN CAPSULES.** (Abbott) Vit A 10,000 u, D 1000 u, thiamine HCl 5 mg, riboflavin 5 mg, nicotinamide 25 mg, pyridoxine HCl 15 mg, pantothenic acid (as Cal pantothenate) 5 mg, ascorbic acid 100 mg /Cap Bot 30s, 100s, 250s
Use: Multiple vitamin therapy
- DAYAMINERAL, FILMTABS.** (Abbott) New name for Maxilets tabs
- DAYCAP.** 10 mg or 15 mg (Ethical Pharm) d Amphetamine sulf 10 mg or 15 mg with Vit A 5000 u, D 1000 u, B 1 2 mg, B 2 2 mg, B 12 1 mcg, C 20 mg, B 6 01 mg, niacinamide 20 mg, cal pantothenate 1 mg, folic acid 0.1 mg with minerals /TD tab Bot 30s
Use: CNS stimulant with vitamins and minerals
- DAYCAP "B."** (Ethical Pharm) d Amphetamine sulf 10 mg, amobarbital 64 mg Vit A 5000 u, D 1000 u, B 1 2 mg, B 2 2 mg, B 12 1 mcg, C 20 mg, B 6 01 mg, niacinamide 20 mg, cal pantothenate 1 mg, folic acid 0.1 mg, Cu 1 mg, mang 1 mg, molybd 0.2 mg, pot 5 mg, Zn 12 mg /T O Cap Bot 150s
Use: Vitamin supplement with CNS stimulant and sedative
- DAYSED.** See Sedasyon (Tailby-Nason)
- DAYSED.** (New Eng Phr Co) Phenobarbital 1 gr /Cap time disint Bot 30s
- DAY-TABS.** (Fremo) Vit B 1 3 mg, B 2 2 mg, B 6 05 mg, Cal pantothenate 1 mg, niacinamide 5 mg, yeast 6 gr /Tab Bot 100s, 1000s
Use: Dietary supplement.
- DAYTEENS FILMTAB.** (Abbott) Vit A 5000 u, D 1000 u, B 1 2 mg, B 2 2 mg, niacinamide 20 mg, B 6 05 mg, B 12 2 mcg, calcium pantothenate 5 mg, C 50 mg, with Fe 10 mg, Cu 0.15 mg, I 0.1 mg, Mn 0.05 mg, Mg 0.15 mg, Ca 250 mg, P 193 mg /Filmtab Bot 100s, 250s
Use: Vitamin mineral formula for teenagers
- DBED** Dibenzylethylenediamine dipenicillin G See: Benzathine penicillin G
- DBI.** NND 1961 (U S Vitamin) Phenformin N β -Phenethylharmamidinylaminourea Tab 25 mg Bot 100s TD Caps 50 mg Bot 100s
Use: Orally, treatment of diabetes
- DCA** See Desoxyaceticosterone acetate (Various Mfr.)
- D.C.P. 34D.** (Parke, Davis) Each 10 gr rEP

- resents approximately 2½ gr. Cal. & 1 4/5 gr. Phos. Pow., 4 ox., 1 lb.; Tab. 7½ gr. Bot. 100s, 1000s. Wafer, Bot. 36s, 100s, 250s.
 Use: Correction of calcium and phosphorus deficiencies.
- DDT.** See Chlorophenothane, U.S.P.
- DEADLY NIGHTSHADE LEAF, U.S.P.** See Belladonna Leaf, U.S.P.
- DEANER, N.N.D. 1962.** (Riker Lab.) Deanol (2-Dimethylaminoethanol as the p-acetamidobenzoic acid salt) 25 mg./Tab. Bot. 100s, 500s. (100 mg./Tab.) Bot. 50s.
 Use: Neurasthenia and chronic fatigue states, mild depressions and chronic headache.
- DEANOL ACETAMIDOBEZOATE, N.N.D. 1962.** (2-Dimethylaminoethanol as the p-acetamidobenzoic acid salt). See: Deaner, Tab. (Riker Lab.)
- DEBA.** See: Barbitol (Various Mfr.).
- DEBROX (IPC),** New name for G.H.P. Bot. 15 cc.
- DECABAMATE.** (Merck Sharp & Dohme) Decadron 0.25 mg., meprobamate 200 mg./Tab. Bot. 100s.
 Use: Arthritic and rheumatic disorders, allergic, inflammatory skin and eye disorders.
- DECADRON, N.N.D. 1962.** (Merck Sharp & Dohme) Dexamethasone. 16- α -methyl-9- α -fluoro-prednisolone. Tabs. 0.75 and 0.5 mg. Bot. 100s, Elixir (0.5 mg./5 cc) Bot. w/ dropper 100 ml.
 W/Isoproterenol sulf. See: Prodecadion (Merck Sharp & Dohme)
 W/Meprbamate. See: Decobamate, Tab. (Merck Sharp & Dohme)
 Use: Corticosteroid therapy.
- W/Xylacaine.** (Merck Sharp & Dohme) Dexamethasone 4 mg., lidocaine HCl 10 mg./cc. Vial 5 cc.
 Use: Anti-inflammatory.
- DECADRON INJECTION.** (Merck Sharp & Dohme) 2 cc. disposable syringe of dexamethasone phosphate 4 mg. per cc. & needle.
- DECADRON PHOSPHATE, N.N.D. 1962.** (Merck Sharp & Dohme) Dexamethasone 21-phosphate sod. In. (4 mg./cc.) Vial 5 cc. Ophth. Solu. 0.1% Dropper Bot. 5 cc. Ophth. Oint. 0.05% Tube 3.5 Gm. Topical Cream 0.1% Tube 5 Gm., or 15 Gm.
- DECADRON AEROSOL SPRAY.** (Merck Sharp & Dohme) Dexamethasone 10 mg./Container (Pressurized) 90 Gm.
 Use: Dermatological conditions.
- DECADRON, RESPHALER.** (Merck Sharp & Dohme) Dexamethasone 21 phosphate 18
- mg. in total weight of 12.6 gms. Aerosol—170 sprays.
 Use: Bronchial asthma.
- DECA-DURABOLIN.** (Organon) Nandrolone decanoate 50 mg./cc. sesame oil 1 cc, amps. Box 4s.
 Use: I.M. anabolic maintenance.
- DECAGESIC.** (Merck Sharp & Dohme) Dexamethasone 0.25 mg., acetylsalicylic acid 500 mg., aluminum hydroxide 75 mg. in the form of a dried gel/Tab. Bot. 100s.
 Use: Rheumatoid arthritis.
- DECALATE.** (Normand) Dicalcium phosphate 5 gr., cal. gluconate 2½ gr., Vit. D 333 u./Tab. Bot. 100s, 500s, 1000s
 Use: Calcium therapy
- DECALBEX.** (Hart) Dicalcium phosphate 0.58 Gm., iron pyrophosphate 12.50 mg., Vit. D 330 u., thiamine HCl 3 mg., riboflavin 2 mg., niacinamide 1.25 mg., brewer's yeast pow. 125 mg./Cap. Bot. 100s, 500s, 1000s.
 Use: Dietary supplement.
- DECALBION.** (Mossengill) Dicalcium phosphate 10 gr., Vit. D 444 u., ferrous sulfate, dried, 1½ gr., thiamine HCl 3 mg., riboflavin 2 mg./Cap. Bot. 100s, 500s, 1000s.
 Use: Provide certain essential vitamins & minerals.
- DECALBION-C.** (Mossengill) Dicalcium phosphate 15 gr., Vit. D 444 u., ferrous sulfate, dried, 1 gr., thiamine HCl 3 mg., riboflavin 3 mg., niacinamide 15 mg., Cal. ascorbate 25 mg., Vit. B-12 1 mcg./Cap. Bot. 100s, 500s, 1000s.
 Use: Hematologic therapy.
- DECALBION FORTE.** (Mossengill) Dicalcium phosphate 15 gr., Vit. A 2500 u., D 250 u., ferrous sulfate dried 1 gr., thiamine HCl 3 mg., riboflavin 3 mg., niacinamide 15 mg., Cal. ascorbate 25 mg., Vit. B-12 1 mcg./Cap. Bot. 100s, 500s, 1000s.
 Use: Therapeutic doses of vitamin A in combination w/minerals & vitamins.
- *DECAMETHONIUM BROMIDE, N.N.D. 1962.** See: Syncurline, Vial. (Burrroughs Wellcome)
- DECAMETHYLENE bis-** (m-dimethylamino-phenyl-N-methylcarbamate) dimethabromide. See: Humorsol. Soln. (Merck Sharp & Dohme)
- DECAPRYN SUCCINATE, N.N.D. 1962.** (Merrell) (Doxylamine succinate) 2[alpha(2-Dimethylaminoethoxy)-alpha-methylbenzyl]pyridine succinate Syr. (5.25/5 cc). Bot. 1 pt., 1 gal.; Tab. (12.5 mg./Tab.), Bot. 100s, 1000s, (25 mg./Tab.) Bot. 100s, 1000s.
 Use: Antihistamine.
 W/Benlyl HCl, pyridoxine HCl. See: Bendectin, Tab. (Merrell)

* Generic name recognized by N.F., U.S.P., or A.M.A.

W/Butaphyllamine & nethamine HCl See: Nethaprin, Cap & Syr (Merrell)

W/APC Tablets (Merrell) Decapryn succinate 6 mg, aspirin 0.23 Gm, acetophenetidin 0.15 Gm, caffeine 30 mg/tab Bot 100s, 500s

Use: Antihistaminic w/analgesic

DECAPS. (Cale) Vit D, activ ergosterol Cap 50 000 u Bot 100s

Use Orally, vitamin D therapy

DECASPRAY AERASOL. (Merck Sharp & Dohme) Dexamethasone 10 mg (0.075 mg/second of spray) and neomycin sulfate 50 mg (0.375 mg/second of spray)/90 Gm Aerosol Can 90 Gm

Use Dermatitis with moist oozing lesions

DECATAB (Thompson) Tab Bot 100s 250s
DECA-VI-CAPS. (Mead Johnson) Vitamin A 3000 u, D 1000 u, B 1 1 mg, B 2 1.5 mg, niacinamide 10 mg, C 50 mg B 6 1 mg, pantothenic acid 3 mg, B 12 3 mcg, biotin 30 mcg/cap Bot 30s

Use Multiple vitamin therapy

DECA-VI-SOL TEASPOON VITAMINS. (Mead Johnson) Vit A 5000 u D 1000 u, ascorbic acid 60 mg, B 1 1.5 mg, B 2 1.5 mg, niacinamide 15 mg, B 6 1.2 mg, B 12 3 mcg, pantothenol 5 mg, biotin 40 mcg/5 cc, Bot 4 B, 12 oz

Use Dietary supplement

DECA-VI-SOL with IRON (Mead Johnson) Vit A 4000 u C 75 mg, D 400 u, B 12 3 mcg, thiamine 12 mg, riboflavin 1.5 mg, pyridoxine 12 mg, niacinamide 15 mg, panthenol 5 mg, biotin 40 mcg, iron 10 mg/Tab Bot 50s Chewable Tabs Bot 250s

Use Dietary supplement

DECAVITAMIN CAPSULES AND TABLETS. USP, Vit A 5000 u, D 400 u, C 75 mg, calcium pantothenate 5 mg, B 12 2 mcg, folic acid 0.25 mg, nicotinamide 20 mg B 6 2 mg, B 2 3 mg, B 12 2 mg/cap or tab Merrell—Bot 100s, 1000s Pfizer—Tabs 100s, 1000s

West Ward—Cap Bot 100s, 1000s

Use: 1 daily, vitamin therapy

DE-CHOLE. (United) Pyridoxine HCl 10 mg, Vit B 12 10 mcg, niacin 25 mg, niacinamide 25 mg, pangamic acid 10 mg, betaine HCl 150 mg, choline bitartrate 150 mg, inositol 100 mg/Cap Bot 60s, 250s

Use: Treatment of disturbed fat metabolism

DECLORISON ACETATE. 9-alpha, 11 beta, dichloro 1,4 pregnadiene 17-alpha 21 diol 3 20 diene 21 acetate See: Diaderm (Schering)

DECOLIN SODIUM, NNR 1953 (Ames) 20% aq. sol of 5ad dehydrocholate for IV inj; Amp 3 cc, 5 cc, 10 cc, Box 5s, 20s, 100s.

Use: Medical & surgical management of biliary tract disorders

DECNOLIN TABLETS, NNR 1953 (Ames) Dehydrocholic acid 3 1/2 gr/Tab Bot 100s 500s, 1000s, 5000s

Use: To promote drainage of biliary duct

W/Belladonna Tab (Ames) Dehydrocholic acid 3 1/2 gr & belladonna ext 1/6 gr/Tab Bot 100s, 500s

Use: Medical & surgical management of biliary tract disorders, w/antispasmodic

DECNOLIN-BB (Ames) Sodium butabarbital 15 mg, dehydrocholic acid 250 mg, belladonna ext 10 mg, (0.125 mg alkaloid)/Tab Bot 100s

Use Hydrachaleric; antispasmodic

DECHOTYL. (Ames) Dehydrocholic acid 200 mg, desoxycholic acid 50 mg, dioctyl sulfosuccinate sodium 50 mg Vit B 1 2 mg/Tab Bot 100s

Use Chronic constipation

DECICAIN, Tetracaine HCl

DECLOMYCIN HCl, NNB 1962 (Lederle) 7 chloro 6 demethyltetracycline HCl De

methylchlorotetracycline HCl 150 mg/cap Bot 16s & 100s Caps 75 mg Bot 25s

Pediatric Drops (60 mg/cc) Bot 10 cc w/dropper Susp (75 mg/5cc) Bot 2 oz

Syr. (75 mg 5 cc) Bot 2 oz

Use Broad spectrum antibiotic

DECLOSTATIN. (Lederle) Demethylchlorotetracycline (Declomycin) 150 mg, nystatin 250 000 u/Cap Bot 16s, 100s

Use Tetracycline sensitive infections

DECLOSTATIN SUSP. (Lederle) Demethylchlorotetracycline 75 mg as HCl nystatin 125 000 u/lip Bot 2 oz

DECO-DROPS. (Donley Evans) Vit A 5000 u D 1000 u, C 50 mg, B 1 1 mg, B 2 0.5 mg, B 6 2 mg, folic acid 0.1 mg, niacinamide 10 mg, panthenol 3 mg/D 6 cc Bot 30 cc

Use Multi vitamin supplement

DECOMINIC. (Flint, Egan) Chlorpheniramine 6 mg, pyriminone maleate 37.5 mg, phenyl

ephine HCl 15 mg, ascorbic acid 75 mg/Tab Bot, 50s, 250s

Use: Common cold, hayfever, etc

DECOPAL DROPS. (Philadelphia Labs) Sod salicylate 2 1/2 gr, phenylephrine HCl 2.5 mg, chlorpheniramine maleate 1.25 mg/0.4 cc Bot 30 cc

Use: Pediatric analgesic in the treatment of symptoms due to cold

DECORTIN, NND 1962 (Schiefelin) De

soxycorticosterone acetate Amp (5 mg/cc. sn oil) 1 cc, 10 cc

Use: 1 M., 5 C., adrenal cortex replacement therapy.

DECOSAL. (Donley Evans) Salicylamide 4 gr-

- ascorbic acid 40 mg., succinic acid 4 gr./
Tab. or 10 cc. Ellixr, Bot. 1 pt. Tab., Bot.
100s, 500s.
Use: Orally, analgesic & antipyretic.
- DECSORB Plain & w/Neomycin.** (Donley-
Evans). Activated attapulgites, 330 Gm.;
pectin, .60 Gm.; sodium & Potassium lac-
tates, 3.0 Gm.; glucose, 6.6 Gm.; sucrose,
9.0 Gm. w/Neomycin has neomycin sulfate,
300 mg.; added/30 cc Bot. 16 oz.
Use: Oral treatment of diarrhea.
- DECO-SPAS.** (Donley-Evans). 2-diethylamino-
ethyl diphenylacetate HCl, .025 Gm.; pen-
taborbital sodium, .0075 Gm.; activated
attapulgites, .090 Gm.; aluminum hydroxide,
.075 Gm./Tab. Bot. 100s, 500s.
Use: Antacid-Antispasmodic.
- DECOSTERONE, N N D 1962.** (Boray) De-
soxycorticosterone acetate, Aq. susp., Vial
(5 mg./cc.) 10 cc., in sesame oil, Vial (5
mg./cc.) 10 cc.
Use: I.M., adrenal cortex replacement ther-
apy.
- DECOSTRATE.** (E. S. Miller) Desoxycorticoste-
rone acetate 5 mg./cc. in cottonseed oil.
Vial 10 cc.
Use: Adrenal cortex & metabolic therapy.
- DECOTONIC.** (Donley Evans) Vit. B-12 25
mcg., B-6 5 mg., B-1 10 mg., folic acid 0.1
mg./Tab. Bot. 30s, 100s.
Use: Appetite stimulant.
- DECUPRYL.** (Crookes-Barnes) Copper undecy-
lenate, undecylenic acid, aerosol in fat sol-
vent, volatile-liquid base containing tetra-
chloroethylene & isopropanol. Cream, Jar
1 oz., 1 lb. liq., 1 & 4 oz. Pow., Can
2 oz.
Use: Ringworm of scalp, athlete's foot &
other skin fungus infections.
- DECUPRYL LIQUID.** (Crookes Barnes) Copper
undecylenate, undecylenic acid, dioctyl so-
dium sulfosuccinate in a volatile-liquid
base containing tetrachloroethylene and iso-
propanol/Bot. 1 oz.
Use: Ringworm of the scalp, athlete's foot
and other superficial fungus infections.
- DECYL.** (Paul Maney) Undecylenic acid 20%,
zinc undecylenate 0.5%, w/menthol &
benzocaine 2%. Pow., 1 oz. Cream, Tube
1 oz.; Jar 1 lb.
Use: Topically, fungus infections of skin.
- DEEGLANS.** (C. B. Kendall) Thyroid U.S.P.
1/2 gr., d-desoxyephedrine HCl 4 mg.,
thiamine HCl 3 mg., riboflavin 2 mg., nico-
linamide 10 mg., ascorbic acid 20 mg./
Tab. Bot. 100s, 500s.
Use: (1-3 daily a c.) Management of obesity.
- DEE-OSTEROL.** (Greon) Vit. D-2, actv. ergos-
terol. Cap. 50,000 u. Bot 100r. 500r.
- * Generic name recognized by N.F., U.S.P. or A.M.A.
- 1000s. Sol. 50,000 u./cc. in oil. Vial, rubber
capped, 30 cc.; Box 1s, 6s.
Use: Vitamin D therapy.
- DEE-RON.** (Warren-Teed) Vit. D 2, Tab.
50,000 u. Bot. 100s, 1000s. Vial 500,000
u./cc. 15 cc.
Use: Orally, I.M., vitamin D therapy.
- DEE-RONAL.** (Warren-Teed) Vit. D 50,000
u., ascorbic acid 30 mg., thiamine HCl 3
mg., riboflavin 1 mg., pyridoxine HCl 0.1
mg., Cal. pantothenate 0.5 mg., nicotina-
mide 10 mg., natural mixed tocopherols
3 mg./Tab. Bot. 100s, 1000s.
Use: (1-2 daily) Vitamin therapy.
- DEGALOL.** (Ames) (Desoxycholic acid) 3,12-
Dihydroxycholic acid. Tab. 1 1/2 gr. Bot.
100s
Use: Orally, a choleric.
- DEGESF.** (Barnes-Hind) Phenylephrine HCl
0.2%, benzalkonium chloride 0.01%, sod
bisulfite sterile soln. Bot. 15 cc
Use: Eye Irritations.
- DENYCOL.** (Rexall) Dshydrocholic acid. Tab
3 1/2 gr. Bot. 100s.
Use: Stimulate bile flow.
- N-N'-BIS-DEHYDROABITYLETHYLENEDIA-
MINE SALT OF PENICILLIN G.** See: Com-
pocillin, Susp. (Abbott)
- DEHYDROCHOLATE, SODIUM, N.F., N.N.R.**
1953.
See: Decholin Sodium, Amp. (Ames).
Dilabil, Sodium, Amp. (Breon).
- * **7-DEHYDROCHOLESTEROL, ACTIVATED,**
U.S.P. (Various Mfr.) Vitamin D-3.
- * **DEHYDROCHOLIC ACID, U.S.P., N.N.R.**
1953; Tab. U.S.P. (Various Mfr.) 3,7,12
Triketocholic acid Ketocholic acid acti-
vated cholic acid. Note bile acids oxidized
Use: Orally, hydrocholeric & choleric.
See: Cholan-DH, Tab. (Maltbie).
Dechalin, Tab. (Ames)
Dehychol, Tab. (Rexall)
Dilabil, Tab., N.N.R. 1953 (Breon).
Hykalex, Tab. (Chicago Pharm.)
Ketocholic acid.
Oxycholin, Tab. (Blue Line).
Fracholan, Tab. (Squibb)
- W/Adiphenine HCl, desoxycholic acid. See:
Cholad, Tab. (Mallard)
- W/Belladonna ext. See: Decholin W/Bella-
donna, Tab. (Ames).
- W/Belladonna and phenobarb. See: Nubific,
Tab. (Hobart Lab.)
- W/Betaine Choline, Inositol, ox bile & liver-
B. See: Nepazyn, Tab. (Endocrine).
- W/Bile, desic. See: Supligol, Tab. (American
Ferment)
- W/Bile, dehy. See: Cholanar, Tab. (Maltbie)

- W/Bile, homatropine methylbromide See B laric Caps (Arcum)
- W/Bile, homatropine methylbromide, pheno barbital See B lamide, Tab (Nargine Labs)
- W/Bile ext See Bilichalan Cap (Harrower)
- W/Bile ext, methyl cellulose, homatropine methylbromide See Calicell, Tab (Suliff & Case)
- W/Bile ext, pepsin, pancreaticin See Nea bile, Tab (Haug Drug Co)
- W/Bile salts and pancreatic substance See Depancof, Tab (Warner Chilcott)
- W/Bile & pancreatic substance See Pan croid, Tab (Strosenburgh)
- W/Bile salts pancreaticin betaine HCl See Pradase, Tab (Miller Labs)
- W/Bile salts See Hydro Bilein Tab (Abbott)
- W/Bile salts & Sorlate See Sorla Bilein Cap (Abbott)
- W/Bulbarbital sod bellodonna extract See Decholin BB Tabs (Ames)
- W/Cascara sagrada See Nubilax, Tab (Nu malazine)
- W/Cassio oculifolia pods active principles See Senabile Tab (Purdue Frederick)
- W/Cholic acid See Chabile, Tab (Irwin Neisser)
- W/Choline bitartrate
See Lipacholan, Tab (Maltbie)
Bidrolin Tab (Armour)
- W/Choline bit, methionine betaine See Chal tropic, Tab (Buffington)
- W/Desoxycholic acid
See Combichole, Tab (F Trout)
Doxychol K, Tab (Brean)
Fexchol Tab (Hart)
Ketox Tab (Ascher)
- W/Desoxycholic acid choline methionine tartrate & odiphenin HCl See Desacholine Tab (Mallard)
- W/Desoxycholic acid diactyl sulfasuccinate sodium, thiamine HCl See Dechoryl, Tab (Ames)
- W/Diactyl sodium sulfasuccinate See Neo lax, Tab (Central)
- W/Homatropine methylbromide
See: Atrochalin, Tab (Blue Line)
Cholan V, Tab (Maltbie)
Chalotropin Tab (Lakeside)
- W/Homatropine methylbromide & pheno barbital See Hytrophin, Tab (Chicago Pharmacal)
- W/Hyascyamine HBr, Mag trisilicate & aluminum hydroxide gel, dried See: Bar mat, Tab (Paul Maney)
- W/Hyascyamine sulfate, atropine sulfate, hyoscine HBr, phenobarbital See: BELLA Bit Tab (Whitmer)
- W/Methamphetamine HCl & pentobarbital See B Iron, Tab (Maltbie)
- W/Methamphetamine HCl & vitamin B-1 See Ganone, Tab (Endo)
- W/Pancreatin, pepsin & ox bile See Ra Bile Tab (Rawell)
- W/Phenobarbital & homatropine methylbromide See: Chalan HMB, Tab (Maltbie)
- W/Phenobarbital, homatropine methylbromide, geralase, gariprolase, desoxycholic acid See: Bilezyme Plus, Tabs (Genetric)
- W/Sod pentobarbital, homatropine methylbromide ox bile ext See Homochol, Tab. (Lemmon)
- W/Prolase 300, Mylase 100, hyascyamine HBr, hyoscine HBr, atropine sulfate See Almezyme Cap (Meyer)
- W/Teralasa, pancreatin, polysorbate 80 See Digalose, Cap (Boyle)
- DEHYDROCHOLIN** See: Dehydrocholic acid
- 11-DIHYDRO-17-HYDROXYCORTICOSTERONE**, See Cortisone
- DEHYDRODESOXYCHOLIC ACID** See: Cholic acid
- DEHYDROISOANDROSTERONE**, See Dan drone, Tab (Organon)
- DEHYDROLAX**, (Mallard) Tabs 100s 1000s
- DEIMAL**, (Boyle) Vit A 5000 u, D 500 u, thiamine 33 mg, riboflavin 2 mg, niacinamide 30 mg Vit B 12 10 mcg, folic acid D 17 mg, Vit B 6 0.2 mg, E 1 IU, C 90 mg pantothenol 1.67 mg choline bitartrate 25 mg, Cot Phos iron Mang, nickel iodine zinc copper cobalt, molybdenum Mag /Tab Bot 100s, 1000s
Use Vitamin & mineral therapy
- DEKA** (Winthrop Labs) Codeine sulfate 1/6 gr, w/cal iodide, chloroform, wild cherry lola, squill, yerba santa citric acid menthol, glycerin & alcohol/5 cc Bal 4 oz 1 pt, 1 gal
Use (1 tsp q 3-4 h, p r n) Narcotic cough preparation
- DELACURARINE**, (Squibb) d Tubocurarine chloride 27 mg, myricin 48 mg /cc. In o.l. Vial 5 cc
Use: Muscle relaxant
- DELADUMONE**, (Squibb) Testosterone enanthate 90 mg, estradiol valerate 4 mg/ml Vial 1 & 5 ml
Use IM (1-2 ml q 2 to 4 weeks) Androgen and estrogen therapy
- DELADUMONE 2X**, (Squibb) Testosterone enanthate 180 mg, estradiol valerate 8 mg/cc Vial 2 cc
Use Hormone preparation
- DELALUTIN**, NND 1962 (Squibb) 17-alpha hydroxyprogesterone caproate Vial (125 mg, 250 mg /cc) 2 cc, 10 cc
Uses IM, long-acting progestational agent
- DELATESTRYL SOLUTION**, NND 1962

- (Squibb) Testosterone enanthate 200 mg / cc. in oil. Vial 1 & 5 cc.
 Use: I.M., androgen therapy.
- DELATROPIN.** (Eastern) Methamphetamine HCl 7½ mg., atropine sulfate 0.12 mg / Laz. Bot. 60s.
 Use: (1 o.c., I.I.d.) Anti-obesity management.
- DELBENOL.** (Brown Pharm.) Safflower oil 1150 mg./Cap. Bot. 89s emulsion- linoleic acid 4500 mg., Vit. B 6 4.5 mg., mixed Vit. E 0.004%, glycerin, polysorbate 60, sorbitan trioleate, cal. cyclamate, sod benzoate, propyl gallate/15 cc. Bot. 1 pt.
 Use: Cholesterol control & nutritional supplement.
- DELCO GRANULES.** (Merck Sharp & Dohme) Each 100 Gm. contains protein 50 Gm. & carbohydrate 30 Gm. Bot. 1 lb., 5 lb.
 Use: Orally, dietary supplement.
- DELECTAVITES.** (White Lab) Vit. A 5000 u., D 1000 u., C 75 mg., E 2 u., B-1 2.5 mg., B-2 2.5 mg., B-6 1 mg., B-12 3 mcg., pantothenal 5 mg., nicotinamide 20 mg., folic acid 0.1 mg., biotin 30 mcg., rutin 12 mg., plus 9 minerals/nuggett, Bot. 30s, 90s.
 Use: Dietary supplement.
- DELENAR.** (Schering) Dexamethasone D 35 mg., orphenadrine HCl 35 mg., aluminum aspirin 375 mg./Tab. Bot. 100s.
 Use: Rheumatism and mild arthritis.
- DELESTROGEN, N.N.D. 1962** (Squibb) Estradiol valerate in sesame oil. Vial (10 mg./cc.) 1 cc., 5 cc.
 Use: I.M.; amenorrhea, menopausal syndromes.
- DELESTROGEN INJ.** (Squibb) Estradiol valerate in castor oil./Vial (40 mg/cc) 5 cc.
 Use: Menopausal syndrome, prostatic carcinoma.
- DELFIN.** (Ortho) Nonylphenoxypolyethoxyethanol 5% in an oil-in-water emulsion at pH 4.5. Vaginal cream, tube with or without applicator, 70 Gm.
 Use: Conception control.
- DELFIN-ETTES.** (Ortho) Delfin Vaginal Cream. Pre-filled single-use applicators. Pack/4s.
 Use: Contraceptive.
- DELINAL.** (National Drug) Propenzolate Hydrochloride
- DELKADON TABLETS.** (Merck Sharp & Dohme) Hyoscyamine HBr 0.225 mg., atropine sulfate 0.019 mg., scopolamine HBr 0.006 mg., delvinal (vinbarbital) 30 mg / tab. Bot. 100s, 1000s.
 Use: (1-2 I.I.d.) Antispasmodic agent & sedative.
- DELMOR.** (Merck Sharp & Dohme) Protein 50 Gm., cal. 600 mg., phos. 440 mg., iron 15 mg., thiamine HCl 10 mg., riboflavin 10 mg., pyridoxine HCl 1 mg., Cal. pantothenate 5 mg., niacinamide 100 mg., Vit. B-12 cryst. 5 mcg., ascorbic acid 100 mg., Vit. A 4000 u., D 400 u., whole liver substance of Vit. B-complex 1 Gm./100 Gm. Pow., Bot. 1 lb., 5 lb.
 Use: Provides supplementary amounts of essential minerals & vitamins.
- DELOMETS.** (Durst) Sulfacetamide 200 mg., sulfamethazine 150 mg., sulfamerazine 100 mg., sulfadiazine 50 mg./Tab. Bot. 36s.
 Use: Quodriac sulfonamide therapy.
- DELPHCOL.** (Lederle) Choline bitartrate 350 mg., dl-methionine 190 mg., Inositol 38 mg., folic acid 0.2 mg., Vit. B-12 2 mcg./Cap. Bot. 100s, 1000s.
 Use: Folate change in liver.
- W/Methionine sol.** (Lederle) Tricholine citrate 1.4 Gm., acetyl dl-methionine 1.22 Gm., inositol 0.3 Gm., folic acid 0.2 mg., Vit. B-12 12 mcg./15 cc. Bot. 1 pt.
- DELPNYNOID.** (Merck Sharp & Dohme) Aminophylline 100 mg., delvinal 30 mg. Forte, Aminophylline 200 mg., delvinal 30 mg./cap. Bot. 100s.
 Use: (1 I.I.d.) Diuretic w/sedative.
- DELTA-CORTEF.** N N D. 1962. (Upjohn) Delta-1-hydrocortisone (Prednisolone) Tab. 1 mg., Bot. 100s, 5 mg., Bot. 30s, 100s.
 Use: Orally, rheumatoid arthritis & intractable bronchial asthma.
- DELTA-CORTRIL.** See. Prednisolone
- DELTA-1-CORTISONE.** See. Deltasone, Tab. (Upjohn)
- DELTACORTONE.** See: Prednisone
- DELTA-OOME.** (Dome) Prednisone 5 mg./Tab. Bot 100s, 1000s.
 Use: Anti-inflammatory steroid.
- DELTA-5-HEMISUCCINOXYPRGENONE-LONE.** See. Panazolone (Dook)
- DELTA-1-HYDROCORTISONE.** See. Prednisolone.
- DELTA¹, 6-METHYL-HYDROCORTISONE.** See: Medrol, Tab. (Upjohn)
- DELTALIN.** (Lilly) Vit. D 2. Gulsuals 50,000 u. Bot. 100s, 500s.
 Use: Vitamin D deficiency.
- DELTALIN ALPHA NO. 1.** (Lilly) Vit. A 5000 u., D 500 u. Cap., Bot. 100s, 1000s.
 Use: Orally, vitamin A & D therapy.
- DELTALIN ALPHA NO. 2.** (Lilly) Vit. A 15,000 u., D 1500 u. Cap., Bot. 100s, 1000s.
 Use: Orally, vitamin A & D therapy.
- DELTALIN ALPHA BETA.** (Lilly) Vit. A 5000 u., D 500 u., B-1 1 mg., B-2 2 mg /Cap. Bot. 100s, 1000s.
 Use: Orally, multivitamin therapy.
- DELTAMIDE.** (Armour) Sulfadiazine 0.167 Gm., sulfamerazine D 167 Gm., sulfamethazine D 056 gr., sulfacetamide 0.111 Gm./

- W/Bile, homatropine methylbromide See Bilaric Caps (Arcum)
- W/Bile, homatropine methylbromide, phenobarbital See B lamide Tab (Norgine Labs)
- W/Bile ext See B lichenol, Cap (Harrower)
- W/Bile ext methyl cellulose homatropine methylbromide See Colicell, Tab (Suthin & Case)
- W/Bile ext pepsin, pancreatin See Neo-bile Tab (Haug Drug Co)
- W/Bile salts and pancreatic substance See Depancol, Tab (Warner Chilcott)
- & pancreatic substance See Pan Tab (Strasensburgh)
- alts pancreatin, betaine HCl See: } Tab (Miller Labs)
- alts See Hydro Rilein, Tab (Abbott)
- salts & Sorlate See: Sorla Rilein, Cap (Abbott)
- W/Butbarbital sod belladonna extract See Decholin BB Tabs (Ames)
- W/Coscoro sagrada See Nubilox, Tab (Nuzolizine)
- W/Cozzio oculifolia pads, active principles See Senabile, Tab (Purdue Frederick)
- W/Cholic acid See Chobile, Tab (Irwin Neisler)
- W/Choline bitartrate See Lipacholan, Tab (Maltbie)
- Bidrolin, Tab (Armour)
- W/Choline bi, methionine, boroine See Cholitropic, Tab (Buffington)
- W/Desoxycholic acid See Combichole Tab (F Trout)
- Doxychol K, Tab (Recon)
- Fkochol, Tab (Hart)
- Kelosox, Tab (Ascher)
- W/Desoxycholic acid, choline methionine tartrate & adiphenin HCl See Desocholine Tab (Mallord)
- W/Desoxycholic acid, dioctyl sulfosuccinate sodium, thiamine HCl See Decholyt, Tab (Ames)
- W/Dioctyl sodium sulfosuccinate See Neolox Tab (Central)
- W/Homatropine methylbromide See Atracholin, Tab (Blue Line)
- Cholan V, Tab (Maltbie)
- Cholatropin Tab (Lakeside)
- W/Homatropine methylbromide & phenobarbital See Hytrophen Tab (Chicago Pharmacal)
- W/Hyoscyamine HBr, Mag Irrisiclate & aluminum hydroxide gel, dried See Barbol, Tab (Paul Maney)
- W/Hyoscyamine sulfate, atropine sulfate, hyosine HBr, phenobarbital See BELLA BIL, Tab (Whitier)
- W/Methamphetamine HCl & pentobarbital See R Iran, Tab (Maltbie)
- W/Methamphetamine HCl & vitamin B₁. See: Ganone, Tab (Endo)
- W/Pancreatin, pepsin & ox bile See: Ro-Bile, Tab (Rowell)
- W/Phenobarbital & homatropine methylbromide See Cholion HMB, Tab (Maltbie)
- W/Phenobarbital homatropine methylbromide, gerilose, geriprotase, desoxycholic acid See: Bilezyme Plus, Tabs (Geriatric)
- W/Sod pentobarbital, homatropine methylbromide ox bile ext See: Homochol, Tab (Iraman)
- W/Prolase 300, Mylose 100, hyoscyamine HBr, hyosine HBr, atropine sulfate See: Almezyme, Cap (Meyer)
- W/Teralase, pancreatin, polysorbate 80 See: Digolose, Cap (Boyle)
- DEHYDROCHOLIN** See: Dehydrocholic acid
- 11-DEHYDRO-17-HYDROXYCORTICOSTERONE**, See Cortisone
- DEHYDRODESIOXYCHOLIC ACID**, See Cholanolic acid
- DEHYDROISOANDROSTERONE**, See Dian-drone, Tab (Organon)
- DEHYDROLAX**, (Mallord) Tabs 100's, 1000's
- DEIMAL**, (Boyle) Vit A 5000 u, D 500 u, thiamine 33 mg, riboflavin 2 mg, niacin amide 30 mg, Vit B 12 1.0 mcg, folic acid 0.17 mg, Vit B 6 0.2 mg, E 11 U, C 50 mg, panthenol 1.67 mg choline bitartrate 25 mg, Cal Phos, iron, Mang, nickel, t-dine zinc, copper, cobalt, molybdenum, Mag / Tab Bal 100's, 1000's
- Use Vitamin & mineral therapy
- DEKA**, (Winthrop Labs) Codeine sulfate 1/6 gr, w/col iodide, chloroform, wild cherry, tolu, squil, yerba santa, citric acid, menthol, glycerin & alcohol/5 cc Bal 4 oz, 1 pt, 1 gal
- Use (1 tsp q 3-4 h, p r n) Narcotic cough preparation
- DELACURARINE**, (Squibb) d Tubocurarine chloride 27 mg, myricin 48 mg/cc in at Vial 5 cc
- Use Muscle relaxant
- DELADUMONE**, (Squibb) Testosterone enanthate 90 mg, estradiol valerate 4 mg/ml Vial 1 & 5 ml
- Use I.M. (1.2 ml q 2 to 4 weeks) Androgen and estrogen therapy
- DELADUMONE 2X** (Squibb) Testosterone enanthate 180 mg, estradiol valerate 8 mg/cc Vial 2 cc
- Use Hormone preparation
- DELALUTIN**, NND 1962 (Squibb) 17 alpha-hydroxyprogesterone caproate Vial (125 mg, 250 mg/cc) 2 cc, 10 cc
- Use I.M., long-acting gestational agent
- DELAESTRYL SOLUTION**, NND 1962

- (Squibb) Testosterone enanthate 200 mg./cc. in oil. Vial 1 & 5 cc.
 Use: I.M., androgen therapy.
- DELATROPIN.** (Eastern) Methamphetamine HCl 7½ mg., atropine sulfate 0.12 mg./loz. Bot. 60s.
 Use: (1 a.c., t.i.d.) Anti-obesity management.
- DELBENOL.** (Brown Pharm.) Safflower oil 1150 mg./Cap. Bot. 87s emulsion linoleic acid 4500 mg., Vit. B-6 4.5 mg., mixed Vit. E 0.004%, glycerin, polysorbate 60, sorbitan trioleate, col. cyclamate, sod. benzoate, propyl gallate/15 cc. Bot. 1 pt.
 Use: Cholesterol control & nutritional supplement
- DELCO GRANULES.** (Merck Sharp & Dohme) Each 100 Gm. contains protein 50 Gm. & carbohydrate 30 Gm. Bot. 1 lb., 5 lb.
 Use: Orally, dietary supplement.
- DELECTAVITES.** (White Lab) Vit. A 3000 u., D 1000 u., C 75 mg., E 2 u., B-1 2.5 mg., B-2 2.5 mg., B-6 1 mg., B-12 3 mcg., pantothenol 5 mg., nicotinamide 20 mg., folic acid 0.1 mg., biotin 30 mcg., rutin 12 mg., plus 9 minerals/suggetl. Bot. 30s, 90s.
 Use: Dietary supplement.
- DELENAR.** (Schering) Dexamethasone 0.25 mg., orphenadrine HCl 15 mg., aluminum aspirin 375 mg./Tab. Bot. 100s.
 Use: Rheumatism and mild arthritis.
- DELESTROGEN, N M D. 1962** (Squibb) Estradiol valerate in sesame oil. Vial (10 mg./cc.) 1 cc., 5 cc.
 Use: I.M., amenorrhea, menopausal syndrome.
- DELESTROGEN INJ.** (Squibb) Estradiol valerate in castor oil./Vial (40 mg./cc) 5 cc.
 Use: Menopausal syndrome, prostatic carcinoma.
- DELLEN.** (Ortho) Nonylphenoxypolyoxyethanol 5% in an oil-in-water emulsion at pH 4.5. Vaginal cream, tube with or without applicator, 70 Gm.
 Use: Conception control.
- DELLEN-ETTES.** (Ortho) Dellen Vaginal Cream. Pre-filled single-use applicators. Pack/4s.
 Use: Contraceptive.
- DELINAL.** (National Drug) Propenzolate Hydrochloride.
- DELKADON TABLETS.** (Merck Sharp & Dohme) Hyascyamine HBr 0.225 mg., atropine sulfate 0.019 mg., scopolamine HBr 0.006 mg., delvinal (vinbarbital) 30 mg./tab. Bot. 100s, 1000s.
 Use: (1-2 t.i.d.) Antispasmodic agent & sedative.
- DELMOR.** (Merck Sharp & Dohme) Protein 50 Gm., cal. 600 mg., phos. 440 mg., iron 15 mg., thiamine HCl 10 mg., riboflavin 10 mg., pyridoxine HCl 1 mg., Col. pantothenate 5 mg., niacinamide 100 mg., Vit. B-12 cryst. 5 mcg., ascorbic acid 100 mg., Vit. A 4000 u., D 400 u., whole liver substance of Vit. E-complex 1 Gm./100 Gm. Pow., Bot. 1 lb., 5 lb.
 Use: Provides supplementary amounts of essential minerals & vitamins.
- DELOMETS.** (Durst) Sulfacetamide 200 mg., selfamethazine 150 mg., sulfamerazine 100 mg., sulfadiazine 50 mg./Tab. Bot. 36s.
 Use: Quadruple sulfonamide therapy.
- DELPNICOL.** (Lederle) Choline bitartrate 350 mg., di-methionine 190 mg., inositol 3E mg., folic acid 0.2 mg., Vit. B-12 2 mcg./Cap. Bot. 100s, 1000s.
 Use: Fructose change in liver.
- W/Methionine sat.** (Lederle) Tricholine citrate 1.4 Gm., acetyl di-methionine 1.22 Gm., inositol 0.3 Gm., folic acid 0.2 mg., Vit. B 12 12 mcg./15 cc. Bot. 1 pt.
- DELPNYNOID.** (Merck Sharp & Dohme) Aminophylline 100 mg., delvinal 30 mg. Forte, Aminophylline 200 mg., delvinal 30 mg./rap. Bot. 100s.
 Use: (1 t.i.d.) Diuretic w/sedative.
- DELTA-CORTEF, N M D. 1962.** (Upjohn) Delta-1-hydrocortisone. (Prednisolone) Tab. 1 mg., Bot. 100s, 5 mg., Bot. 30s, 100s.
 Use: Orally, rheumatoid arthritis & intractable bronchial asthma.
- DELTA-CORTRIL.** See: Prednisolone
- DELTA-1-CORTISONE.** See: Deltasone, Tab. (Upjohn)
- DELTA-CORTONE.** See: Prednisone
- DELTA-DOME.** (Dome) Prednisone 5 mg./Tab. Bot. 100s, 1000s.
 Use: Anti-inflammatory steroid.
- DELTA-5-NEMISUCCINOXYPREGNE-
 LONE.** See: Panzalone (Doak)
- DELTA-1-HYDROCORTISONE.** See: Prednisolone.
- DELTA-1,6-METHYL-HYDROCORTISONE.** See: Medrol, Tab. (Upjohn)
- DELTALIN.** (Lilly) Vit. D 2 Gelseals 50,000 u. Bot. 100s, 500s.
 Use: Vitamin D deficiency.
- DELTALIN ALPHA NO. 1.** (Lilly) Vit. A 5000 u., D 500 u. Cap., Bot. 100s, 1000s.
 Use: Orally, vitamin A & D therapy.
- DELTALIN ALPHA NO. 2.** (Lilly) Vit. A 15,000 u., D 1500 u. Cap., Bot. 100s, 1000s.
 Use: Orally, vitamin A & D therapy.
- DELTALIN ALPHA BETA.** (Lilly) Vit. A 3000 u., D 500 u., B-1 1 mg., B-2 2 mg./Cap. Bot. 100s, 1000s.
 Use: Orally, multivitamin therapy.
- DELTAMIDE.** (Armour) Sulfadiazine 0.167 Gm., sulfamerazine 0.167 Gm., selfamethazine 0.056 gr., sulfacetamide 0.311 Gm./

Tab or dr Bot 100s; Bot 4 oz 1 pt
 Use: Sulfanamide therapy
W/Penicillin (Armour) Sulfadiazine D 167
 Gm sulfamerazine 0 167 Gm, sulfamethiazine 0 056 Gm sulfacetamide 0 111 Gm
 Pot penicillin G 250 000 u/Tab or dr Bot 4 oz 1 pt
 Use: Sulfanamide w/penicillin therapy
DELTASMYL (Roussel) Prednisone 15 mg
 theophylline 120 mg ephedrine HCl 15 mg, phenobarbital 8 mg/Tab 8at 50s
 Use Asthma
DELTASONE, NND 1962 (Upjohn) Prednisone Tab 5 mg Bot 30s 100s
 Use: Orally rheumatoid arthritis & intractable bronchial asthma
DELTRA, NND 1962 (Merck Sharp & Dahme) Prednisone Tab 1 mg Bot 100s 1000s; 2.5 mg 5 mg Bot 30s 100s 500s 1000s
 Use: 20 30 mg daily; treatment of rheumatoid arthritis
DELTRA-STAB See: Prednisolone
DELTYL EXTRA (Givaudan Delawanna Inc)
 Mainly Isopropyl myristate
 Use: Anthelmintic Pinworms roundworms & whipworm
DELUTEVAL (Squibb) Estradiol valerate 2.5 mg
 hydroxy progesterone caproate 125 mg/cc Ampul 2 cc Also available in double strength Vial 5 cc
 Use Hormone preparation
DELVEX, NND 1962 (Lilly) Dithiazonine iodide Polymeth ne blue dye (3,3 diethyl thiodicarbonyl iodide) 50 100 200 mg/Tab Bot 50s 100 mg 200 mg/Tab Bot 100s
 Use: Anthelmintic Pinworm roundworm & whipworm
DELVINAL, NND 1962 (Merck Sharp & Dahme) Vinbarb tal Sodium
W/Atropine sulf hyascyamine HBr hyoscine HBr See: Delkadan Tab (Merck Sharp & Dahme)
DELVINAL-ASPIRIN CAPSULES (Merck Sharp & Dahme) Delvinal 30 mg, aspirin 0.3 Gm /cap Bot 100s, 1000s
 Use: Sedative & analgesic
DELVINAL SODIUM, NND 1962 (Merck Sharp & Dahme) Vinbarb tal Sod
 Amps (1 gr/cc) 5 cc sterile sol
 Caps Brown, 1/2 gr, Bot 100s, 500s 1000s; Orange 1 1/2 gr, Bot 100s, 500s 1000s; Orange & Brown, 3 gr, Bot 100s, 500s
 Elxirs (4 gr /fl oz) Bot 1 pt, 1 gal
 Pnw: Bot 1/2 oz
 Vials (1 gr/cc) 20 cc
 Use: General sedative
W/Aminoacetyl See: Delphynoid Cap (Merck Sharp & Dahme)
 *Generic name recognized by NF, USP or AMA

W/Aspirin See: Delvinal Aspirin, Cap (Merck Sharp & Dahme)
W/Belladonna alkaloids See: Delkadan, Tab (Merck Sharp & Dahme)
W/Diethylstilbestrol See: Delvinal diethyl stilbestrol, Tab (Merck Sharp & Dahme)
DELVINAL-DIETHYLSTILBESTROL (Merck Sharp & Dahme) Delvinal, sad vinbarbital 30 mg, diethylstilbestrol 0 1 mg/Tab Bot 100s, 1000s
 Use: Orally, sedative w/estrogen therapy
DEMAMIN (Schering) (Formerly Chlor Tri metan Compound) Syr Chlorpropenpyridamine molecula 1 25 mg, phenylephrine 2.5 mg/5 cc Bot 1 pt Repetabs Outer and inner layers each contain chlortrimeton molecula 2 mg, phenylephrine 10 mg Bot 100s
 Use Antihistaminic
***DEMECARIUM BROMIDE, NND 1962** De comethylene bis (m dimethylamino phenyl N methylcarbamate) dimethyl bromide See Numorsol Sol (Merck Sharp & Dahme)
DEMEROL APAP, (Breon) Demerol 50 mg & N acetyl p aminophenol 300 mg/Tab Bot 100s
 Use: Sedative analgesic agent
DEMEROL-ATROPINE, (Winthrop Labs)
 Demerol 50 mg, atropine sulf 1/300 gr/cc Amp 2 cc, Box 25s
 Use: I.M., S.C., analgesic
DEMEROL COMPOUND (Breon) Each central white layer, Demerol HCl 25 mg; outer green layer dihydrocodeinone bitartrate, 5 mg; outer pink layer, acetyl p aminophenol, 150 gm /Stratified Tab Bot 100s
 Use Analgesic
DEMEROL HYDROCHLORIDE, NND 1962
 Meperidine HCl
 Breon:
 Inj, Amp (50 mg/cc) 2 cc Vial 30 cc
 Amp (100 mg/cc) Vial 20 cc Tab (50 mg) Bot 100s
 Winthrop Labs
 Amps Sterile disposable u for instant Inj (100 mg) 2 cc, Box 5s 10s, 100s
 Amp - (50 & 100 mg) 2 cc, Box 5s 25s 100s
 Amps (50 75 mg) 1 ml Disposable syringe Box 10s
 Elxlr (50 mg /isp) Bot 16 fl oz
 Pow Vial 15 Gm
 Tabs (50 mg) Bot 25s 100s, 250s 1000s; (100 mg) Bot 25s 100s, 500s
 Vial (50 mg/cc) 30 cc, Box 1s
 Use: Narcotic analgesic
W/A.P.C. See: A.P.C. W/Demerol Tab (Winthrop Labs)
W/Lotusate (Winthrop Labs) Demerol HCl

100 mg., Lolusate 100 mg./Tab. Bot. 100s.
Use: Relief of pain and of the insomnia
that may accompany pain.

DEMEROL COMPOUND. (Breon) Meperidine HCl 25 mg., dihydrocodeinone bitartrate 5 mg., N-acetyl-p-aminophenol 150 mg./Tab. Bot. 100s.
Use: Analgesic.

DEMEROL-SCOPOLAMINE. (Winthrop Labs) Demerol 50 mg., scopolamine HBr 1/300 gr./cc. Amp. 2 cc. & Vial 30 cc.
Use: (1 or 2 cc. I.M.) Obstetric analgesic, amnesia and preoperative medication.

* **DEMETHYLZINE PHOSPHATE.** See: Anisiv (Abbott Lab.).

* **DEMETHYLCHLORTETRACYCLINE HCl,** N.N.D. 1962. 7-chlor-4-dimethylamino-1,4,4a,5,5a,6,11,12a-octahydro-3,6,10,12,12a-pentahydroxy-1,11-dioxo-2-naphthacecarboxamide. See: Declomycin, Cap. (Lederte).

W/Nystatin. See: Declostatin (Lederte).

DENAMONE. (Vio-Bin) 1.8 conc. wheat germ oil. Cap. 3 min. Bot. 100s, 500s, 1000s.
Use: Vitamin therapy.

DENCO ACETONE TEST. (Denver) Sod. nitroprusside, sod. carbonate, am. sulf.
Use: Diagnostic for urine acetone.

DENCO SUGAR TEST. (Denver) Bismuth salt.
Use: Diagnostic for urine sugar.

DENCOTAR. (Denver Chemical) Coal tar, specially processed 1½%, sulfur 1%, w/menthol & starch. Oint., Tube 1½ oz. Liq. shampoo, Bot. 4 oz.
Use: Topically, antiseptic in skin disorders.

DENTACEVIN. (Walker Labs.) Niacinamide 100 mg., ascorbic acid 100 mg./Tab. Bot. 50s, 500s, 1000s.
Use: Orally, vitamin C w/niacinamide

DENTALONE, A.O.B. 1935, (Parke, Davis) Cleve oil 63.4 cc., methyl salicylate 2.5 cc., cinamon oil 5 cc., chlorobutanol 38.3 cc., ctenamic aldehyde 5 cc./100 cc. Bot. 1 oz.
Use: Local anesthetic to relieve toothache.

DENTOCILLIN. (Andrew Jergens Co.) Pot. penicillin G 500 u./Gm. Paw., Cos 3 oz.
Use: Dental caries.

DENYL SODIUM. (Premo) See: Diphenylhydantoin Sodium (Various Mfr.).

DENYLEX OEL. (Lloyd, Dabney & Westfield) Adenosine-5-monophosphate 25 mg., vitamin B-12 150 mcg., niacin 10 mg., in 17% gelatin solution. Vial 10 cc.
Use: Neuromuscular disorders

DEO. (Davis) Ointment. Tube ½ oz Jar 1 oz.

DEODO. (Merck Sharp & Dohme) Can 2½ oz.
Use: Deodorant.

DEODOPHYLL. (Wallace) Chlorophyll. Tab. 100 mg. Bot. 75i.

* Generic name recognized by N.F., U.S.P. or A.M.A.

Use: Orally, neutralize breath & perspiration odors.

DEODORANT POWDER. See: Deodo, Paw. (Merck Sharp & Dohme)

DEOFED. (Drug Products) d-Desoxyephedrine HCl 5 mg./Cap. Bot. 100s, 1000s.

Use: Central nervous system stimulant.

DEORYCNOLIC ACID. See: Desoxycholic Acid (Various Mfr.).

DEORYCORTICOSTERONE ACETATE, N.N.O. 1961. See: Desoxycorticosterone acetate. U.S.P.

DEORYCORTONE ACETATE. (Metropolitan Labs) Desoxycorticosterone acetate. U.S.P. Vial (5 mg./cc) 10 cc. in sesame oil. Vial (5 mg./cc) 10 cc. in susp.
Use: Adrenal insufficiency.

DEPANCOL. (Warner-Chilcott) Dehydrocholic acid ½ gr., bile salts 1½ gr., pancreatic substance 4½ gr./Tab. Bot. E.C. Bot. 50s, 200s.
Use: Orally, choleric.

DEPINAR. (Armour) Vit. B-12 zinc tannate complex eq. to 500 mcg./cc. Vial 5 cc.
Use: Long-acting Vit. B-12 therapy.

DEPO-ACTH, N.N.D. 1962. (Upjohn) Adrenocorticotropin 40 u. or 80 u., gelatin 160 mg./cc. Vial 5 cc. Mix-O-Vial 40 u./cc 1 cc. & 5 cc.; 80 u./cc. 5 cc.
Use: I.M., S.C.; repository ACTH therapy.

DEPO-CER-O-CILLIN, N.N.D. 1962. (Upjohn) Chloroprocaine penicillin O Vial, 300,000 u. & 1,500,000 u. 1 dose & 5 dose.
Use: Hypo-allergenic penicillin.

DEPO-ESTRADIOL, N.N.O. 1962. (Upjohn) Depo-estradiol cyclopentylpropionate. Vial 1 mg./cc. cottonseed oil, 10 cc. Box 1s, 5s, 25s; 5 mg./cc. cottonseed oil, 5 cc. Box 1s, 5s, 25s.
Use: 1 inj. I.M./month; menopause symptoms.

DEPO-HEPARIN SODIUM. (Upjohn) Heparin Sod (20,000 u.) 200 mg., gelatin 180 mg., dextrose 80 mg./cc. Cartridge w/disposable syringe 1 cc., Box 1s, 5s, 25s.
Use: S.C., I.M., anticoagulant.

W/Vasocostrictors. (Upjohn) Heparin Sod. (20,000 u.) 200 mg., epinephrine HCl 1 mg., ephedrine HCl 10 mg., gelatin 180 mg., dextrose 80 mg./cc. Cartridge w/disposable syringe 1 cc., Box 1s, 5s, 25s.
Use: S.C., I.M., anticoagulant w/vasocostrictors.

DEPO-MEDROL, N.N.D. 1962. (Upjohn) Methylprednisolone acetate 40 mg./cc. suspended in sodium chloride injection. Vial 1 cc., 5 cc. (20 mg./cc) Vial 5 cc.

See: Medrol.

Uses: Anti-inflammatory.

DEPO-NORTESTONATE (Upjohn) 19 Nor-testosterone 17 cyclopentylpropionate Vial 100 mg/cc

Use: I.M., anabolic steroid therapy

DEPO-PENICILLIN, NND 1962 (Upjohn) Procaine penicillin G cryst 300,000 u, w/2% aluminum monostearate in peanut oil/cc Cartridge w/disposable syringe 1 cc Vial, multiple dose, 10 cc

Use: I.M., long-acting penicillin therapy

DEPO-PENICILLIN FORTIFIED. (Upjohn) Procaine penicillin G 300,000 u, Pot penicillin G 100,000 u, w/2% aluminum monostearate in peanut oil/cc Cartridge w/disposable syringe 1 cc Vial 10 cc

Use: I.M., long acting penicillin therapy

DEPO-PROVERA. (Upjohn) Medroxyprogesterone acetate 50 mg in aq susp /cc Vials 1 and 5 cc

Use: Long acting (16 da) progestational agent

DEPO-TESTADIOL. (Upjohn) Sterile sol testosterone cyclopentylpropionate 50 mg, estradiol, 17 cyclopentylpropionate 2 mg, chlorthalidone 5 mg, cottonseed oil/cc Vial 1 cc, 10 cc

Use: {1 cc I.M. q 38 weeks} Androgen estrogen therapy

DEPO-TESTOSTERONE, NND 1962 (Upjohn) Testosterone Cyclopentylpropionate Vial {25 mg, 50 mg, 100 mg/cc} 10 cc Box 5s 25s

Use: I.M., long acting androgen therapy

DEPROL (Wallace) Meprobamate 400 mg benactyzine HCl 1 mg/Tab Bot 50s

Use: Antidepressant

DEPROPANEX. (Merck Sharp & Dahme) Pancreatic ext, deproteinated Vial, 10 & 30 cc rubber capped

Use: I.M., vasodilator & antispasmodic

DERATOL. (Brewer) Vit D, actv ergosterol Amp (50,000 u) 1 cc, Bot 6s, 12s, 100s Cap (50,000 u) Bot 30s, 100s, 1000s

Use: Orally or I.M., vitamin D therapy

DERFULE. (Cate) Pow. of lpecac & opium (containing opium 1/40 gr) ¼ gr, atropine sulfate 1/500 gr, acetylphenelidin 1½ gr, aspirin 2 gr, camphor monobromated ½ gr, caffeine ½ gr/Cap Bot 100s, 500s, 1000s Also available in half strength

Use: (1 2) For headache & head colds

DERIFIL. (Rylan) Water soluble derivatives of chlorophyll 100 mg/Tab Bot 30s, 100s

Use: Designed to control color/fat or fleshy odors

DERIZENE. (Hallister Siler) Phenylephrine HCl Nasal drops, ¼ % sol, Bot 1 fl oz Ophth sol, ½ %, Bot ½ fl oz

Use: Vasoconstrictor & decongestant

W/Zinc sulfate ophth (Hallister Siler) Phen-

ylephrine HCl ½ %, zinc sulfate ¼ % sol Bot ½ fl oz

Use Vasoconstrictor w/antiseptic

DERMABASE. (Marcelle) Hypo allergenic oint base of fatty acid esters & alcohol Jar 16 oz, 5 lb

Use: Ointment base

DERMACOL. (Drug Products) Mercuric chloride, salicylic acid, boric acid, Sod benzoate, thymol, eucalyptol, hydrocyanic acid glycerol, alcohol Bot 1 pt, 1 gal

Use Antiseptic lotion

DERMADERM INJECTION. (Hyrex) Riboflavin 5 phosphate sodium ester 50 mg/ml representing riboflavin USP 35 mg/ml Vials 10 cc

Use Vitamin therapy

DERMAGEL CREME. (Bruce) Zirconium carbonate 4%, sod thiosulfate 2%, glycerin 3%, Isopropyl alcohol 8%, benzocaine 3% in a colloidal magnesium aluminum silicate base Plastic Bot 2 oz, 1 pt, 1 gal

Use Treatment of poison ivy

DERMAGILL. (Massengill) Carbolic acid, cresol, bismuth subgallate, zinc oxide, sulfur code oil Oint, Tube 2 oz, Jar, 1 lb, 5 lb

Use Antipruritic & parasiticide

DERMA-GUARD (Greer) Protective adhesive pow Can w/sifter top, 4 oz

Use: Protective powder for skin

DERMAJECT (Columbus) Riboflavin 5 monophosphate Sod 25 mg, B 6 5 mg, biotin 25 mg, B 12 500 mcg /cc Vial 10 cc

Use I.M., skin disorders

DERMALGEN. (National Drug) Alum, bismuth subgallate, acetanilid, thymol iodide boric acid, zinc stearate, phenol Pow, Bot w/sprinkler top, 1½ oz

Use Astringent, bacteriostatic dusting powder

DERMA-MEDICONE. (Medicone) Anesthesia 5%, oxyquinoline sulfate 1%, ephedrine HCl D 1%, oleo cadinumichthyal compound 1% Oint, Tube 1 oz, Jar 1 lb

Use: Antipruritic & vasoconstrictor

DERMA-MEDICONE-HC. (Medicone) Hydrocortisone acetate 10 mg, benzocaine 20 mg, oxyquinoline sulfate 10.5 mg, ephedrine HCl 11 mg, menthol 4.8 mg, Ichthammol 10 mg, zinc oxide 137 mg/Gm Tube 7 Gm & 20 Gm

Use: Dermatitis

DERM-ANODYNE. (Upjohn) Aconite tr ¾ oz, atropine 1¼ gr, menthol, camphor, capsicum oleoresin/oz Oint, Tube ½ oz, 1 oz

Use: Analgesic anodyne

DERMA-PAX. (Recsei) Trihlistin 0.5%, chloro-phenylpyramine, pyrrolamine & phenylpyro-

- mine w/benzyl alcohol, chlorobutanol & isopropyl alcohol, liq. 4 oz.
 Use: Topically, antipruritic w/antihistamine.
- DERMAR.** (A. J. Parker) Sulfur colloidal, alcohol, glycerin in a homogenized base. Bot. 4 oz.
 Use: Sulfur therapy in minor skin irritations
- DERMASPOR Ointment.** (Lowe) Neomycin sulfate, 5 mg; polymixin B sulfate, 5000 units, bacitracin, 400 units/Gm. Tube 15 Gm.
 Use: Primary skin and mucous membrane infections
- DERMASULF.** (Smith, Miller & Polch) Aq. sol. of polythionates essentially in form of tetrathionate & pentathionate. Sol. 2½%, Bot. 4 oz., 5%, Bot. 2 oz. Cream Squeeze Bot. 55 Gm.
 Use: Topical sulfur treatment.
- DERMATHYH.** (Flint, Eaton) Benzyl alcohol 3%, benzocaine 3.5%, butyl-p-aminobenzoate 1% & phenylmercuric borate. Tube 1 oz.
 Use: Topical local anesthetic.
- DERMATOL.** See: Bismuth subgallate. (Various Mfr.).
- DERMATONE.** (Chicago Pharmacol) Arsenic trioxide 1/12%, ammoniated mercury 4%, boric acid 6.66%, zinc oxide 4 1/6%. Tube 1 oz., Jar 8 oz., 1 lb., 5 lb.
 Use: Antiseptic ointment for skin.
- DERMATROL.** (Gordon) Cod liver oil, zinc oxide, talc, and balsam of peru in a petrolatum base.
 Use: Granulation stimulant in treatment of wounds.
- DERMA-VEE.** (Sherman) Streptococci, staphylococci, pneumococci & E. coli cultures in cream base. Oint., Tube ¼ oz.
 Use: Antivirus therapy.
- DERMEZE.** (Promo) Thienylpyramine HCl 2%, benzocaine 2%, tyrothricin D.25 mg./Gm. Cream, Tube 1 & 2 oz.
 Use: Topically; antihistaminic, antibiotic & anesthetic.
- DERMOLATE.** (White) High molecular weight-saturated sulfate octadecanoic acid, stearic acid as the Sod. salt. Cake 4 oz.
 Use: Routine skin cleansing.
- DERMO-PARIN.** (Broemmel) Polymyxin B sulf. 100,000 u., neomycin sulf. 20 mg., heparin sod. 2,000 u., hydrocortisone 5 mg./Gm. Oint. Tube 5 Gm.
 Use: Skin infections.
- DERMOPLAST.** (Daha) Benzocaine, menthol, benzethonium chloride. Aerosol Spray 6 oz., 12 oz. Bot. 3 oz.
- Use: Typically, anesthetic, antipruritic, astringent
- DERMOTAN OINTMENT.** (Horlan & Converse) Zinc oxide 10%, phenol 0.44% w/ aromatic oils in a bland base. 1 oz., 1 lb., 5 lb.
 Use: Skin irritations, burns
- DERMOVAN.** (Tosco Pharm. Co) Glycerol monostearate, spermaceti, mineral oil, glycerin & water. Vanishing-type base, 1 oz., 1 lb.
 Use: Ointment base.
- DERMYCIN.** (Chal-Yon) p-Nitrophenol Sod. Iodate. liq. 1 oz., 8 oz., 1 pt.
 Use: Topically, fungus of skin.
- DERONIL, H H D.** 1961. (Schering) Dexamethasone D.75 mg./Tab. Bot. 50s. 500s.
 Use: Hormone-corticosteroid.
- DES.** (Grant) Diethylstilbestrol. liq. 25 mg./tsp. Bot. 6 oz., 1 pt. Supp. 5 mg./25 mg. Box 17s. Tab. 0.1 mg., 0.5 mg., 1 mg., 2 mg., 5 mg., 10 mg., 25 mg., 50 mg., 100 mg. Bot. 100s; Pkg. 500s, 1000s.
 Use: Orally, estrogen therapy.
- DESA-B.** (Pitman-Moore) Desic. crystall. Vit. B complex factors. Vial 30 cc.
- DESACCHROMIN.** A nonprotein bacterial colloidal dispersion of polysaccharide.
- DESACETYL-LANATOSIDE C.** See: Deslanoside, U.S.P.
- DESACHOLINE.** (Moffard) Dehydrocholic acid 100 mg., desoxycholic acid 100 mg., choline methionine tartrate 100 mg., adphenin HCl 10 mg./Tab. Bot. 100s, 500s, 1000s.
 Use: Choleric.
- DESA-HIST.** (Deseret) Chlorpheniramine maleate 12 mg./Tab. Bot. 100s.
 Use: Antihistaminic
- DESA-HIST PF ELIXIR.** (Deseret). Chlorpheniramine maleate, 2 mg; 1-phenylephrine HCl, 3 mg; alcohol, 7.5% in a syrup base/5 cc. Bot. pts.
 Use: Systemic nasal decongestant.
- DESAMINE TABLETS.** (Starr) Methamphetamine HCl. Tab. 5 mg., White or Yellow, Bot. 1000s; 10 mg., White, Bot. 1000s.
 Use: Orally, central nervous system stimulant.
- DESBUTAL.** (Abbott) Desoxy HCl (d-desoxyephedrine HCl) 5 mg., nembul Sod. [pentobarbital Sod.] 30 mg./Cap. Bot. 100s, 1000s.
 Use: Depressive & anxiety states. Management of obesity.
- DESBUTAL GRADUMET.** (Abbott) Methamphetamine HCl 10 mg., pentobarbital sodium

60 mg / Tab or Methamphetamine HCl 15 mg pentobarbital sodium 90 mg / Tab Rot 100s 500s

Use: Treatment of obesity

DESCORNAQ. (New Mexico Pharm) Desoxy corticosterone Acetate 5 mg Chlorbutanol 0.3%, Polyoxyethylene Sorbitan Mono Ole ate 0.06% / cc liq Vials 10 cc

DESCORTERONE. (Kirk) Desoxycorticosterone acetate Vial 5 mg / cc in oil 10 cc

Use: I.M., adrenal cortex replacement therapy

DESCOTONE (Merck Sharp & Dahme) Desoxycorticosterone acetate Disk 2 mg Rot 100s Sol 5 mg / cc Vial 10 cc

Use: Adrenal insufficiency

DESENE (Maltbie) New name Cal Desene Calcium undecylenate 15% in a talc base Powder shaker pkg 2 oz

Use: Diaper rash

DESEX, NNR 1951 (Wallace & Tiernan)

Oint contains 5% undecylenic acid buffered at pH 6.5 by addition of triethanol amine & 20% zinc undecylenate Pow contains 2% undecylenic acid & 20% zinc undecylenate Sol contains undecylenic acid 10% in 40% prapanol, propylene glycol, triethanolamine & water, buffered at pH 6.5 Oint, Tube 1 oz; Jar 1 lb

Pow, Canister 1 1/2 oz; Carton 1 lb Sol., Rot 2 fl oz 1 pt

Use: Fungus infections

***DESERPINE, NND 1962** An alkaloid derived from Rauwolfia canescens 11 Desmethoxyreserpine (Rounormine) See Harmony, Tab (Abbott)

W/Chlorothiazide See: Orechyl and Orechyl Forte, Tab (Abbott)

W/Methylothiazide See: Enduranyl Tabs (Abbott)

W/Pentobarbital calc See: Harmony N, FilmTab (Abbott)

DESERPINE (Deseret) Reserpine 0.1 mg, D 25 mg, 1 mg, / Tab

Use: Treatment of hypertension

DESICOL KAPSEALS. (Parke, Davis) No 1 cap. containing 0.325 Gm of desic whole bile, equiv to approx 2.5 cc of fresh whole bile Rot 100s, 1000s

Use: To relieve bile deficiency.

DESIGUENT-BACITRACIN. (Upjohn) 500 w Bacitracin/Gm Oint, Jar 1 oz

***DESIPRAMINE HYDROCHLORIDE** (Lake side) 10 11 dihydro 5-(3 methylaminopropyl) 5H d benz [bf] azepine hydrochloride See: Narpramin

DESITABS. (Marco) No 1: Amphetamine sulfate 5 mg, thyroid

* Generic name recognized by N.F., U.S.P. or A.M.A.

1 gr., atropine sulfate 1/360 gr., aloin 1/4 gr / Tab (Also double strength)

No 2: Amphetamine sulfate 5 mg, thyroid 1 gr., atropine sulfate 1/360 gr / Tab (Also double strength)

No 3, Amphetamine sulfate 5 mg, thyroid 1 gr., phenobarbital 1/4 gr / Tab (Also double strength)

Use: Thyroid therapy

DESITIN. (Desitin) Cod liver oil, zinc oxide Mag carb, lime water Emulsion, Bot 4 oz

Use: External skin therapy

DESITIN COR-D-TAR-CREAM (Desitin) Hydrocortisone alcohol 1% diadalydroxy quinoline 2%, sal of coal tar 3% in white stainless, water miscible base / Tubes 1/2 oz, 1 oz

Use: Dermatologic

DESITIN HC. (Desitin) Hydrocortisone acetate 10 mg, Norwegian cod liver oil, lanolin, zinc oxide, bismuth subgallate, balsam peru, cocoa butter base / supp Sax 12s

DESITIN-HC OINTMENT. (Desitin) Desitin ointment formula plus hydrocortisone alcohol 1% Tubes 1/2 oz, 1 oz 1/4% Tube 1 1/2 oz

Use: Dermatitis pruritus

DESITIN LOTION (Desitin) Cod liver oil, zinc oxide, Mag carb, rose water & lime water Sol 4 oz

DESITIN OINTMENT. (Desitin) Cod liver oil zinc oxide & talcum Oint, Tube 1 oz, 2 oz, 4 oz Jar 1 lb

DESITIN POWDER (Desitin) Cod liver oil zinc oxide talcum & Mag oxide Can 2 oz, 2 1/2 oz

Use: Topically, antiseptic

DESITIN RECTAL. (Desitin) Cod liver oil zinc oxide, lanolin, talc, sod lauryl sulfate, petrolatum Oint Tube 1 1/2 oz

Use: Treatment of hemorrhoids

DESITIN SOAP. (Desitin) Cake, Pkg 3s

DESITIN SUPPOSITORIES (Desitin) Cod liver oil, zinc oxide, bismuth subgallate lanolin, peru balsam & cocoa butter Rox 12s

Use: Rectally, antiseptic & healing

DESIVER. (U S Vitamin) Liver, whole desic. 9 Gm / Tab Bot. 100s, 500s, 1000s

Use: Orally, antianemic therapy

***DESLANOSIDE, USP; Inj., USP** Des acetylanoside See: Cedilanid D, Amp (Sandaz)

ET-DESMETHOXY RESERPINE. Deserplano See: Harmony (Abbott)

Rounormine (Penick)

DESOCCHROMIN. See: Piramen, Vial (Travel)

DESODS (Calumbus) Thyroid U.S.P. 1/2 gr.

dl-desoxyephedrine HCl 3/40 gr., thiamine HCl 3/40 gr./Tab. Bot. 100s, 1000s.

Use: Obesity.

DESOMIDE, (Mallord Inc.) Dipyrone 1½ gr., salicylamide 1½ gr., dl-desoxyephedrine HCl 1.5 mg./tab. Bot. 100s, 1000s.

Use: Analgesic.

DESOXEDRINE W/Thyroid & atropine sulfate. See: Table Tab. (Testagar).

DESOXETS. (Abbott) Desoxyn HCl (Methamphetamine HCl) 5 mg., Vit. A 5000 u., D 500 u., thiamine mononitrate 3 mg., riboflavin 2.5 mg., Vit. B-12 activity 1 mcg., pantothenic acid (as Cal. pantothenate) 5 mg., ascorbic acid 50 mg./Tab. Bot. 100s, 1000s.

Use: (1 t.i.d., o.c.) Anti-obesity therapy w/ vitamin supplement.

DESOXO-5, (Sutliff & Case) d-Desoxyephedrine HCl 5 mg./Tab. Bot. 100s, 500s, 1000s.

Use: Adjunct in obesity.

DESOXYCHOLIC ACID, (Various Mfr.) Desoxycholic acid, 3,12-Dihydroxycholonic acid. Tab. 1½ gr.

Use: Choleric.

See: Cholorebic, Tab. (Mallord)

Deqalol, Tab. (Ames)

Drexolan, Tab. (Breon).

W/Aloes ext. See: Chalmodin, Tab. (Ames).

W/Betaine HCl, eleoressin ginger & pancreatin. See: Convertin, Tab. (Ascher).

W/Dehydrocholic acid.

See: Cambichole, Tab. (Trout)

Doxychol-K, Tab. (Breon).

Exachol, Tab. (Hort)

Keloxol, Tab. (Ascher).

W/Dehydrocholic acid, bile ext., sorbitan monooleate, oleic acid. See: Oxsorbil, Cap. (Ives-Cameron)

W/Dehydrocholic acid, choline methionine tartrate, & adiphenin HCl. See: Desocholine, Tab. (Mallord).

W/Dxhydrocholic acid, diactyl sulfosuccinate sodium. See: Dechoty, Tab. (Amxs)

W/Phenobarbital, hamalirapin methylbromide, gerilase, geriprotase, d-hydrocholic acid. See: Bilzyme Plus, Tab. (Geriatric)

DESOXY-COMP. No. 1. (Lemmon) Phenobarbital 15 mg., thyroid 15 mg., dl-desoxyephedrine HCl 5 mg., atropine sulfate 0.2 mg., strychnine sulfate 1 mg., elain 5 mg./Tab. Bot. 100s

Use: Treatment of obesity

DESOXY-COMP. No. 2. (Lemmon) Phenobarbital 15 mg., thyroid 60 mg., dl-desoxyephedrine HCl 5 mg., atropine sulfate 0.2 mg., strychnine sulfate 1 mg., elain 5 mg./Tab. Bot. 100s

Use: Treatment of obesity

* Generic name recognized by N.F., U.S.P. or A.M.A.

***DESOXYCORTICOSTERONE ACETATE**, U.S.P., N.N.D. 1962. Oil Inj., Buccal Tab., U.S.P., Pellets, N.F. (Various Mfr.)

[DCA, Deoxycosterone acetate, Cortate, Cortiron, Cortenil, Cortigen, Syncort, Syncortyl.] Aq. susp., 5 mg./cc. Vial 10 cc.; in sesame oil, 5 mg./cc. Vial 10 cc.

Uses I.M. or S.C. (2-5 mg.) adrenal cortical insufficiency.

See: Arcort, Vial (Moore & Co.).

Cortarmour, Vial & Sublingual. (Armour)

Cortacet, Vial. (Hart)

Cortate, Amp., Vial & Tab. (Schering).

Cortinaq, Vial. (Central Pharm.)

Decartin, Amp. (Schieffelin).

Decosterone, Vial (Barry).

Decostrate, Vial (Smith, Miller & Patch)

Deoxycortone Acetate, Vial (Metropolitan).

Dexcorterone, Vial (Kirk).

Decotone, Vial. (Marck Sharp & Dahme)

Deco Acetate, Sol., N.N.R. (Organon).

Dorcestrin, Vial (Smith-Dorsey).

Doxotone, Vial (Testagar)

Doxo, Vial (Mallord).

Percorten, Amp., Lingual, Pellet & Vial (Cibo).

Steroaq, Aq. Susp. (Ascher).

W/Chlorabutanol, Polyoxyethylene Sorbitan Monooleate. See: Descornaq. (New Mexico Pharm.)

W/Sodium ascorbate (Vit. C). See: Artho-Sterone, Amp. (Columbus).

***DESOXYCORTICOSTERONE TRIMETHYLACETATE**, U.S.P. See: Percorten Trimethylacetate, Vial (Cibo).

DESOXYCORTONE ACETATE, See: Desoxycorticosterone Acetate, Prep. (Various Mfr.)
Desoxyephedrine Hydrochloride Combinations on pp. 232 to 232.

DESOXYEPHEDRINE HYDROCHLORIDE, N.F., N.N.D. 1962. (Various Mfr.) Methamphetamine HCl, N.F.; dextro-1-phenyl-2-methylaminopropane, d-phenylisopropylmethylamine, d-N-methylamphetamine. Cap. 5 to 1D mg., Tab. 5 or 1D mg. Elixir 20 mg./fl. oz.

See: Amphetrazyn, Elixir & Tab., N.N.R. (Lilly).

Apamine. (Stillico)

Dzafed, Cap. (Drug Products)

Desamine, Tab. (Starr)

Dextoredrine. (Testagar)

Desoxo-5, Tab. (Sutliff & Case).

Desoxyephedrine, Tab. (Upjohn)

Desoxyn, Amp., Elixir, & Tab. (Abbott).

Dexstim, Aerial, Tab. & Vial. (Central)

Desyphed, Tab. (Winthrop Labs.)

Dexoval, Tab. (Valz)

Dexstim (Central Pharm.)

Desoxyephedrine Hydrochloride Combinations

| Desoxyephedrine HCl | Others | Manufacturer | Brand |
|---------------------|--|----------------|---------------------------|
| d, 1.25 mg. | Methapyrilene HCl 25 mg. | Walter Abbott | Hist-Oxa-Mine, Tab. |
| d, 2.5 mg. | Methapyrilene HCl 50 mg. | | Thenylene & Desosyn, Tab. |
| d, 2.5 mg. | Methapyrilene HCl 50 mg. | Massengill | Hisdria, Tab. |
| dl, 125 mg. | Methapyrilene HCl 25 mg., belladonna ext. 7½ mg. | McNeil | Corenil, Tab. |
| d, 6 mg. | Methylcellulose 300 mg. | Sly | Obetamine, Tab. |
| d, 5 mg. | Methylcellulose 2½ gr., thyroid ¼ gr., phenobarbital ½ gr. | Zemmer | Hyrobel, Tab. |
| 1 mg. | Methyltestosterone 2.5 mg., estrogenic subs. conj. 0.25 mg., vitamins, minerals | Paul Maney | Vitrimin, Tablets |
| d, 5 mg. | Pentobarbital Sod., 60 mg., butobarbital Sod., 30 mg. | Central | Somnalex, Tab. |
| d, 2.5 mg. | Pentobarbital Sod., ½ gr., homatropine MB 1/40 gr., methyl cellulose 5 gr. | Central | Redaxcel, Tab. |
| d, 5 mg. | Pentobarbital ¼ gr., mephobarbital ¾ gr. | Walker | Mepha-d, Tab. |
| dl, 5 mg. | Pentobarbital Sod. 20 mg., vitamins B-1, B-2, B-6, C | Massengill | Obedrin, Tab. |
| d, 3.33 mg. | Phenobarbital ¼ gr. | Rabins | Ambar, Tab. |
| dl, 5 mg. | Phenobarbital 16 mg. | Filtz-Eaton | Barbidex, Tab. |
| 2.5 mg. | Phenobarbital 30 mg., Dilantin 100 mg. | Parke, Davis | Phelanin, Capsules |
| dl, 1/20 mg. | Phenobarbital ¼ gr., hyoscyamus ext. ½ gr., salicylamide 3 gr., acetophenetidin 2 gr. | Lemmon | Aspifedrine, Tab. |
| dl, 5 mg. | Phenobarbital 16 mg., Mag. gluconate 7½ gr. | Kendall | Magilla, Tab. |
| dl, 5 mg. | Phenobarbital ¼ gr., thyroid 1 gr. | Meyer | Myrabile #3 |
| d, 5 mg. | Phenobarbital 15 mg., secobarbital Sod. 30 mg. | Massengill | Hypnarm, Pill |
| d, 2.5 mg. | Phenobarbital ¼ gr., dl.alpha.tocopheryl acetate 5 mg. | Brean | Transibarb, Cap. |
| d, 1 mg. | Premarin 0.25 mg., methyltestosterone 2.5 mg., vitamins | Ayerst | Mediotec, Cap. & Liq. |
| d, 0.5% | Pyrimamine maleate 2% | Lemmon | Pyraphed, Sol. |
| dl, 2 mg. | Pyrimamine maleate 50 mg., dihydrocodeine bitartrate 1/6 gr., w/omm. chloride, pot. guaifacal sulf., chloroform, tartar emetic | Taylor Labs. | Tussitor, Syr. |
| d, 3 mg./oz. | Pyrimamine 75 mg./oz., Pot. guaifacal sulf. 500 mg /oz. | Suffitt & Case | Bronchastil, Liq. |
| dl, 2.5 mg. | Reserpine 0.1 mg. | Wampole | Secolrest, Tab. |
| dl, 10 mg. | Reserpine 0.25 mg. | Ascher | Du-Orio, Tab. |
| dl, 1/30 gr. | Salicylamide 3½ gr., acetophenetidin 2½ gr. | Wampole | Tryado, Tab. |
| dl, 1 mg. | Salicylamide 250 mg., vitamins | Ascher | Bexasol, Tab. |
| d, 0.5 mg. | Salicylamide 150 mg., gentisate Sod. 150 mg., Vit. C 10 mg. | Suffitt & Case | Casacomb, Tab. |
| dl, 1.5 mg. | Salicylamide 1½ gr., dipyrone 1½ gr. | Mallord Inc. | Dasomide, Tab. |
| dl, 1/20 gr. | Salicylamide 3 gr., hyoscyamus ¼ gr., phenobarbital ½ gr., acetophenetidin 2 gr. | Lemmon | Aspifedrine, Tab. |

D-O-E, Elixir B Tab. (Breon)
 Doxyfed, Elixir B Tab. (Raymer)
 Drinalfa, Tab. B Viol. (Squab)
 Efraxine, Elixir B Tab. (Maltbie)
 Methamphetamine HCl (Various Mfr)
 Methedrine HCl, Amp. B Tab. (Rusroughs Wellcome)
 Methaxyn. (Kenny Pharm)
 Miller-Drine (Miller)
 Narodin, Tab. (Endo)
 Oxydrin (Grant)
 Phedrisax, Tab. (Ascher)
 Semoxydrine, Tab. & Elixir (Massengill)
 Stimdex, Tab. (Ulmer)
 Syedrax, Elixir & Tab. (McNell)
 W/Vitamins. See: Vitamin Tablets.
dl-DESOXYEPHEDRINE HYDROCHLORIDE.
 dl-Methamphetamine HCl.
 See: Delrex, Tab. (Mallord)
 Lanazine, Tab. (Lannett)
 Methomphin, Tab. (Rorer).
 Miller-Drine, Tab. (Miller).
 Oxyfed, Tab. (Cale).
 Oxydess, Elixir B Tab. (Chimedic).
 Premadrin, Tab. (Premo).
 Nermadrine Solution (Yanpelt & Brown).
 W/Terramycin hydrochloride. See: Terramycin Nasal W/Desoxyephedrine Hydrochloride, Sol. (Pfizer).
leva-DESOXYEPHEDRINE. See: Anodrex (Anahist).
DESOXYETS. (Abbott) Desoxyn 5 mg., Vit. B-12 1 mcg., A 5000 u., D 500 u., thiamin mononitrate 3 mg., riboflavin 2.5 mg., ascorbic acid 50 mg./Tab. Bot. 100s, 1000s.
 Use: (1 f.l.d., a.c.) Appetite depressant w/ vitamin supplement.
DESOXYN HYDROCHLORIDE, N.N.D. 1962. (Abbott) Methamphetamine HCl, Amp. (20 mg/cc), 1 cc., Box 6s, 25s. Elixir, 20 mg/B. oz. Bot. 1 pt., 1 gal. Tab. 2.5 mg., Bot. 25s, 100s, 1000s; 5 mg., Bot. 100s, 1000s. Gradumete 5, 10 B 15 mg Bot. 50s, 500s.
 Use: Central nervous system stimulant.
 W/Nembutol. See: Desbutal, Cap. (Abbott).
 W/Vitamins. See: Desoxets, Tab. (Abbott)
DESOXYN GRADUMET. (Abbott) d-Desoxyephedrine 5 mg., 10 mg., or 15 mg/Tab. Bot. 50s, 500s.
 Use: Appetite depressant; aid age menopause.
DESOXY NOREPHEDRINE. See: Amphetamine
DESOXY-PHEN. (Lemmon) Phenobarbital 30 mg., dl-desoxyephedrine HCl 10 mg./Tab. Bot. 100s
 Use: Treatment of narcolepsy
DESOXYRIBONUCLEASE. W/Fibrinolysin
 See: Elase, prep. (Parke Davis)
 * Generic name recognized by N.F., U.S.P. or A.M.A.

DESPLEX. (Amfre-Grant) Diethylstilbestrol 25 mg., Hesperidin complex 100 mg., vit. B-1 2 mg., B-2 2 mg., B-6 1 mg., C 50 mg., calcium pantothenate 10 mg., niacinamide 50 mg., folic acid 0.25 mg., B-12 0.5 mg./Tab. Bot. 30s, 100s, 500s.
 Also available with diethylstilbestrol potencies of 0.02, 0.05, 0.1, 0.25, 0.5, 1, 5, 10, 50 and 100 mg.
 Use: Treatment of habitual abortion.
DESTRIOL. See: Estriol.
DESYLATE. (Walker) Vit. D 5000 u., Sod. salicylate 5 gr., ascorbic acid 50 mg./Cap. Bot. 100s, 1000s.
 Use: (1-3 daily) Rheumatism & arthritis.
BESYPHED HYDROCHLORIDE, N.N.D. 1962. Methamphetamine HCl (Winthrop Labs.) d-Desoxyephedrine HCl Tab. 2.5 mg. Bot. 100s.
 Use: Central nervous system stimulant.
DETERDEL AROMATIC SPRAY. (Bruce) Pyrilamine maleate 0.25%, benzalkonium chloride 0.05%, phenylephrine HCl 0.25%, methylparaben 0.02%, propylparaben 0.02%. Bot. pts, gals.
 Use: Colds.
DETERDEL EMULSION. (Bruce) Neomydin 1:5000, pyrilamine maleate 0.25%, benzalkonium chloride 0.05%, phenylephrine HCl 0.25%, methylparaben 0.02%, propylparaben 0.02%, camphor 0.025%, menthol 0.25%, methyl salicylate 1:5000. Bot. pts, gals.
 Use: Intranasal decongestant.
DETERGENTS, Surface-Active.
 See: Acidolate. (White)
 Dermolate. (White)
 p-Hisoderm. (Winthrop Labs.)
 phisohek. (Winthrop Labs.)
 Terjolate. (White)
DETERGOL. (Dook) Not listed. Nonalkaline, super fatted detergent. Liq. Bot. 4 oz, 1 pt, 1 gal
 Use: Topically on dry, inflamed skin.
DETERJEL AROMATIC SPRAY. (Bruce) Tyralthrin 0.02%, pyrilamine maleate 0.25%, benzalkonium chloride 0.05%, phenylephrine HCl 0.25%, methylparaben 0.02%, propylparaben 0.02%. Bot. 1 pt.
 Use: Antibiotic decongestant.
DETHYLANDIAMINE. Thenyldiamine. See: Thendadil.
DETIGON HYDROCHLORIDE. See: *Chlaphedianol
DETONAL. (M. R. Thompson) Not listed. Secobarbital Sod. 3/4 gr in outer coating, phenobarbital in inner core 1/8 gr/Tab. Bot. 24s, 100s, 1000s.
 Use: (1-2) Prolonged sedative

- DETREX.** (Mallard) dl Desoxyephedrine HCl
10 mg / Tab Bot 100s, 500s, 1000s
Use Central nervous system stimulant
- ***DETROTHYRONINE, D 4** [3 iodo 4 hydroxyphenoxy] 3,5 diiodophenylalanine
(Smith Kline & French)
- DETT.** (Atlantis Sales Corp) F Chlorometa xylenol Bot 4 & 16 oz & 1 gal
Use Antiseptic
- DEVEGAN.** (Winthrop Labs) Combination of acetarsone (acetylaminohydroxyphenyl arsonic acid) w/hydrolyzed carbohydrates & small amount of boric acid Each tab contains 0.25 Gm of arsenical constituent
Pow., Bot. 1 oz., 8 oz.; Vial 10 Gm Tab, Box 25s, 250s
Use: Trichomonas vaginalis vaginitis
- DEVICAL.** (Premo) Dicalcium phosphate 6.5 Gm, Vit D 2000 u / 8 oz Bot 1 pt
Use: Dietary supplement
- DEVICAPS.** (Hart) Vit A 12,500 u, D 1000 u, Thiamine HCl 5 mg, riboflavin 5 mg, ascorbic acid 150 mg, niacinamide 50 mg, pyridoxine HCl 1 mg, Caf pantothenate 10 mg / Cap Bot 30s, 100s, 300s
Use: Multiple vitamin deficiencies
- DEXABAR.** (Medco) #1 Dextroamphetamine sulfate 10 mg, amobarbital 60 mg / Med sule #2 Dextro amphetamine sulfate 15 mg, amobarbital, 90 mg / Medsule Bot 30s, 100s
Use: Obesity control
- DEXABARS** (Nyco Labs) No 1—d Amphetamine sulf 10 mg, amobarbital 60 mg / tab TD
No 2—d Amphetamine sulf 15 mg, amobarbital 160 mg / tab
No 3—d Amphetamine sulf 15 mg, amobarbital 100 mg / tab
Bot 100s, 1000s
Use: Anesthesia
- DEXABESE.** (Danley Evans) Dextro-amphetamine, 5 mg; amobarbital, 16 mg; methyl cellulose, 125 mg / Tab Bot 100s, 500s
Use: Obesity control
- DEXAGESIC.** (Pitman Moore) Acetaphenetidin 2½ gr, aspirin 3½ gr, d amphetamine sulfate 2.5 mg / Tab Bot 100s, 1000s
Use: Orally, analgesic & central nervous system stimulant
- DEXALME.** (Meyer Chemical) Dextro amphetamine sulfate Tab (5 mg) Bot 100s, 1000s
Use: Central nervous system stimulant
- DEXALME L.A.** (Meyer Chemical) Dextro amphetamine sulfate Cap (10 & 15 mg) Bot 100s, 1000s Also w/sobarb 60 mg / Cap Bot 100s
Use: Long acting central nervous system stimulant (sympathomimetic).
- * Genetic name recognized by N.F. USP or A.M.A.
- DEXALONE** (Wynn) d amphetamine sulfate 10 and 15 mg / Dura Tab Bot 30s, 100s, 250s
Use Anesthesia
- ***DEXAMETHASONE, NND 1962** Hexadecardol 9 alpha fluoro 16 alpha methyl prednisolone
See Decadran (Merck Sharp and Dahme)
Deronil, Tab (Schering)
Gammacorten, Tab (Ciba)
Hexadral Tab (Organon)
Marsdex (African Lab)
Use Same as for prednisone or prednisolone
Dose Individualized according to disease
0.75 mg is equivalent to 4 mg triamcinolone 5 mg prednisone/prednisolone, 20 mg hydrocortisone or 25 mg cortisone
W/d Chlorpheniramine, Vit, C See Palanit Tab (Schering)
W/Aspirin aluminum hydroxide as the dried gel See Decagesic, Tab (Merck Sharp & Dahme)
W/Aspirin aluminum hydroxide ascorbic acid See Ropad, Tabs (Pitman Moore)
W/Cyproheptadine HCl See Dronadin Tabs (Merck Sharp & Dahme)
W/Dechlorpheniramine maleate, ascorbic acid See Palanit, Tab (Schering)
W/Mepramate See: Decabamate, Tab (Merck Sharp & Dahme)
W/Neomycin Sulf See Decaspray Aerosol (Merck Sharp & Dahme)
W/Orphenadrine HCl, aluminum aspirin See Delenor, Tab (Schering)
W/Phenyltolaxamine dihydrogen citrate See Taldex, Tabs (Pitman Moore)
W/Zylacaine See Decadran w/Zylacaine (Merck Sharp & Dahme)
- ***DEXAMETHASONE PHOSPHATE 50-DIUM, NND 1962**
See Decadran Phosphate (Merck Sharp & Dahme)
Rexphaler Decadran (Merck Sharp & Dahme)
W/Neomycin sulf See Neo Decadran Phosphate (Merck Sharp & Dahme)
- DEXAMINE.** (Myers Carter) d Amphetamine sulf 2 mg, pyrilamine maleate 2.5 mg / cc Vial 10 cc
- DEXAMPHETAMINE.** See: d Amphetamine
- DEXAMYL.** (Smith, Kline & French) Each tab & each 5 cc of elixir contains dexedrine sulfate 5 mg, amobarbital, ½ gr Tab; Bot 100s, 1000s Elixir 12 fl oz
Use: Mental & emotional distress
- DEXAMYL SPANSULES.** No 1: Dexedrine sulfate 10 mg, amobarbital 1 gr / Spansule

No. 2; Dexedrine sulfate 15 mg., amobarbital 1½ gr./Spansole. Bot. 30s, 250s.

Use: Mental & emotional distress.

DIXAPHET Tabs & Spanseps. (Dome) Dextroamphetamine sulfate, 5 mg./Tab. Dextroamphetamine sulfate, 10 or 15 mg./Cap Bot. 100s.

Use: Obesity control.

DIXASLIM Tabs, Spanscaps & Spanscap Forte. (Dome) Dextroamphetamine sulfate, 5 mg.; amobarbital, 30 mg./Tab Dextroamphetamine sulfate, 10 mg.; amobarbital, 60 mg./Cap Dextroamphetamine sulfate, 15 mg.; amobarbital, 100 mg./Spanscap Forte. Bot. 100s.

Use: Weight reduction when mental and emotional distress is cause of overeating.

DIXATABS. (Walker) Sod. chloride 7 gr., dextrose 3 gr., thiamine 1 mg., ascorbic acid 10 mg./Tab. Bot. 100s, 1000s.

Use: Orally, dehydration & heat cramps.

DEXATAL. (Wynn Pharmacol Co) No 1—d-Amphetamine sulfate 15 mg., phenobarbital ¼ gr./Dura-Tab. S.M.

No 2—d-Amphetamine sulfate 10 mg., phenobarbital ½ gr./Dura-Tab. S.M. Bot. 30s, 50s, 250s.

Use: Sustained medication, antidepressant.

***DEXOMPHENIRAMINE MALEATE.** N.N.D. 1962. dextro form of brompheniramine Dextro-parabromdylamine 1-(p-bromophenyl)-1-(2-pyridyl) 3-dimethylamino propane maleate See: Disamer, Tab, 5yr (White)

W/d-Isoprenaline. See: Disophrin, Tab. (White)

***DEXCHLORPHENIRAMINE MALEATE.** N.N.D. 1962. d-3-[p-chloro-alpha(2-dimethylamino-ethyl)benzyl]pyridine maleate. See: Dextro-Chlorpheniramine.

W/Desamethaxane, ascorbic acid. See: Pol-anil, Tab. (Schering)

DEXEDRINE SULFATE. N.N.D. 1962 (Smith, Kline & French) Dextro amphetamine sulfate. Elixir, 5 mg/5 cc. Bot. 12 oz. Spansule (5, 10 & 15 mg) Bot. 30s, 250s, Tab. (5 mg) Bot. 100s, 1000s.

Use: Central nervous system stimulant.

W/Amobarbital. See: Dexomyl Spansule (Smith, Kline & French)

W/Thorazine. See: Thoro Des, Tab. (Smith, Kline & French)

DEXIBAR. (Barre) d-Amphetamine phos. 15 mg./time disint. tab. Bot. 100s, 1000s

DEXIBAR-B. (Barre) Dextroamphetamine sulfate 15 mg., amobarbital 1 gr./D.A. Tab Bot. 100s, 1000s

Use: Orally (1 daily), delayed action antidepressant & sedative.

* Generic name recognized by N.F., U.S.P. or A.M.A.

DIXIN. (Burroughs Wellcome) Dextrin 75%, maltose 24%, moisture 0.75%, ash 0.25%. Container, 12 oz., 3 lb.

Use: Infant feeding formulas.

DEX-OB. (Tully Pharmaceuticals) Tab. 1D mg.—d-Amphetamine sulf. 2 mg. in the coating and 8 mg. in the compressed core. Tab. 1S mg.—d-Amphetamine sulf. 5 mg in the coating and 10 mg. in the compressed core. Bot. 30s, 100s

Use: Controlled release tablets in the treatment of obesity

DEXOVAL. N.N.D. 1962. (Yale) d-Desoxyephedrine NCL Tab. 2.5 mg., 5 mg. Bot. 100s, 1000s.

Use: Central nervous system stimulant.

***DEXOXADROL HYDROCHLORIDE.** (Cutter) d-2-[2,2-diphenyl-1,3-dioxolan-4-yl] piperidine hydrochloride. See: Belone.

Use: Antidepressant

***DEXPANTHENOL.** N.N.D. 1962. Panthanol, dextro pantothenyl alcohol.

See: Cozyme (Travenol)

Hopan (Warren-Teed)

Motilyn (Abbott)

DEXSTIM. (Central) Methamphetamine HCl. Tab. 5 mg., 10 mg. Bot. 100s, 500s, 1000s. Vial 20 mg./cc. 10 cc.

Use: Elevates mood.

DEX-SULE. (Normal) d-Amphetamine sulf. 5 mg., 10 mg. or 15 mg./Tab. Bot. 100s, 1000s

***DEXTRAN.** U.S.P., N.N.D. 1962. Inj., U.S.P. (Abbott, Baxter & Cutter) Polyoxyhydroglucose Hydrolyzed dextran 6% w/vial in isotonic Sod chloride sol. Bot. 500 cc.

See: Expander, Sol. (Commercial Solvents)

Genron, Sol. (Baxter)

Plavalex, Sol. (Wyeth).

Use: I.V., shock states.

Sol. 6% and 12% w/venaset & needle set Bot. 500 cc. Also available w/dextrose 5% —Abo—Liter 500 cc. 12%. Bot. 250 cc.

See: Expander (Com Sol Corp)

Genron (Baxter)

Plavalex (Wyeth)

Use: Blood volume expander.

***DEXTRIFIBROH.** N.N.D. 1962. The hematinic preparation of a colloidal solution of ferric hydroxide in complex with partially hydrolyzed dextrin. See: Astraler (Astra)

DEXTRI-MALTOSE. (Mead-Johnson)

No. 1: (Maltose 55%, dextrins 42%, Sod. chloride 2%, moisture 1%) Tin 1 & 5 lb.

No. 2: (Maltose 56%, dextrins 43%, moisture 1% & ascorbic acid 50 mg./oz. Tin 1 & 5 lb.

No. 3: (Maltose 55%, dextrins 41%, Pot.

- bicarb 3%, moisture 1%) Tin 1 & 5 lb
Use Nutritional
W/Vit D, low fat milk See: Dalaatum, Pow
(Mead Johnson)
- DEXTRINS** W/Dextrase, maltose See: Car
tose, Liq (Winthrop Lab)
- * **DEXTRO-AMPHETAMINE HYDROCHLO-
RIDE.** (Various Mfr) d Amphetamine HCl
{+} 2 Aminopropylbenzene
See Timed Tridex, Cop (Testogor)
d Amphetamine HCl for combinations
- * **DEXTROAMPHETAMINE PHOSPHATE,**
USP (Various Mfr) {+} 2 Aminopropyla
mine d Amphetamine phosphate See
d Amphetamine Phos for combinations
- * **DEXTROAMPHETAMINE SULFATE, USP,**
NNO 1962, Tab, USP d Amphetamine
sulfate d Methylphenylamine sulfate
See: d Amphetamine Sulfate for combina
tions
See: Adrizine, Tab (Key Corp)
Am Dex, Tab (Superior)
Amirene Tab & Vial (Normond)
Amphedrine, Tab (Vonpelt & Brown)
Amphex, Tab (Algre)
Amustoin, Tab (Key Corp)
Dadex, T D Cop (Central Pharmacol)
DAS, Tab (Stayner)
D Amfetazol, Tab (Pilman Moore)
D Ale Ph 747, Cop (Paul Money)
D Citramine (Preslan)
Dexaphel Tabs & Spencaps (Damed)
Dexedrine, Prep (Smith, Kline & French)
Dexolme, Tab (Mayer)
Dex OB Tab (Tully)
Dex Sule Tab (Narmol)
Dextrazole (Storck)
Dextra Unicelles (Miss Pharm)
Dura Dex, Cop (Bonar)
Evradox (Evran)
Hetamine (Dumas Wilson)
Medex, Cop (Medex)
Pellcaps (Kirkman)
- * **DEXTROAMPHETAMINE TANNATE.** See:
Synolan, Tab (Irwin, Neisler)
- DEXTRO A.T.A.** (Vitamix Corp) No. 1—
morning tab; d Amphetamine HCl 5 mg,
thyroid 1 gr, atropine sulfate 1/360 gr,
olain 1/4 gr/Tab No. 2—Noon tab;
d-amphetamine HCl 5 mg, thyroid 1 gr,
atropine sulfate 1/360 gr/Tab No. 3—
Afternoon tabs; d-amphetamine HCl 5 mg,
thyroid 1 gr, phenobarbital 1/4 gr/Tab
Bot 100s, 1000s
Use: Treatment of obesity
- DEXTROBAR.** (Lanett) Dextra-amphetamine
sulfate 5 mg, amobarbital 32 mg /tab Bot
300s & 1000s
Use: Central stimulant
- * Generic name recognized by N.F., U.S.P or A.M.A.
- DEXTROCEL.** (Lanett) Dextra amphetamine
sulfate 2.5 mg, methylcellulose 5 gr/tab
Bot 500s & 1000s
- DEXTRO-CHLORPHENIRAMINE MALEATE**
See: Paloramine, Repetab and Tab (Scher
ing)
- DEXTROGEN.** (Nestle) Milk w/dextrin
maltose, disodium phosphate, iron citrate
Vit A & C & thiamine Liq 14.5 oz
Use Infant feeding
- DEXTROLEN.** (Len Tag) d Amphetamine sul
fate 15 mg /Tab
Use Treatment of obesity
- * **DEXTROMETHORPHAN HYDROBROMIDE,**
NF, NND 1962 Syr, Tabs, NF d 3
Methoxy N methylmorphinan Dormethon d
Methorphan
See: Ramilar HBr, Tab, Syr (Roche)
Metharate HBr (Upjohn)
Tussin Liquid (Dumas Wilson)
- W/Acetyl p aminophenol, phenylephrine
hydrochloride, phenylpropranolamine hydro
chloride ascorbic acid, chlorpheniramine
maleate See Thorphan C P, Tabs (Bryant)
- W/Ascorbic acid, propyl para aminobenzo
ate See: Thoreze, Laz (Central)
- W/Chlorpheniramine maleate, ephedrine
HCl, phenylephrine HCl, ammonium Cl
ipecac Auidextract, ethyl alcohol See
Ouelidine Syrup (Abbott)
- W/Chlorpheniramine maleate, phenyleph
rine HCl, acetyl p aminophenol, ascorbic
acid See Thorphan, Syr (Bryant)
- W/Chlorpheniramine maleate, phenylephras
HCl, ammonium chloride See Nilfuss Syr
(Minn Pharm)
- W/Chlorpheniramine maleate, phenylephrine
HCl, phenindamine tartrate, menthol See
Melodon, Syr (Moyrand)
- W/Chlorpheniramine maleate, phenyleph
rine HCl, terpin hydrate Sepi Syntussin,
Tab (Ives Cameron)
- W/Chlorpheniramine, phenylephrine, N
acetyl p aminophenol See: Ramilar—CF
Syr (Roche Lab)
- W/Chlorprophenpyridamine maleate, phen
ylephrine HCl, aspirin, phenacetin anhy
drous caffeine, phenylpropranolamine HCl
citrus bioflavonoid complex, ascorbic acid
See: Daladyne, Tab (Damsco)
- W/Chlorpheniramine maleate, pyrilamine
maleate, phenylephrine HCl, acetamina
phen, caffeine, ascorbic acid See: Tacol
Tabs (Messingill)
- W/Glyceril guaiacolate, ammonium chloride
sodium citrate See: Pedicole Pediatric, Liq
(Lemmen)
- W/Homatropine methylbromide, amm chlor
ide See: Endotussin NN pedi Syr (Endo)

- W/Phenylephrine tartrate, phenylephrine. See: Cerose, Pediatric (Ives-Cameron)
- W/Phenylephrine HCl, chlorprophe-nyl-pyridamine maleate, pyrilamine maleate, APC, citrus bioflavonoid complex, ascorbic acid. See: Daldyne, Tab. (Domsco)
- W/Phenylephrine, N-acetyl-p-aminophenol, pyrilamine maleate, potassium guaiacolsulfonate, alcohol. See: Thormal, Liq. (Central)
- W/Phenylephrine, chlorpheniramine maleate, terpin hydrate, acetaminophen, ascorbic acid. See: Syntussin Compound, Cap. (Ives-Cameron)
- W/Phenylephrine maleate, potassium guaiacolsulfonate, sodium citrate, citric acid, chloroform. See: Thormal Syrup (Central)
- W/Pyrimidine maleate, phenylephrine HCl, ammosium chloride, citric acid. See: Pyral-dime Pediatric, Syr. (Yaepelt & Brown)
- W/Tripelnic, terpin hydrate. See: Tessaminic, Tab. (Smith-Dorsey).
- W/Terpin hydrate. See: T. H. & M., Syr. (Upjohn)
- * **DEXTROMORAMIDE TARTRATE**, di-(2-Methyl-4-morpholino-2,2-diphenyl-butyl) pyrrolidine. See: Dimorlin, (Smith, Kline & French) Palium (G. P. Harvey)
- DEXTRO-PANTOTHENYL ALCOHOL**. See: Dexpanthenol.
- DEXTRO-PROFETAMINE**. (Clark & Clark) d-Amphetamine phosphate. Tab. 5 mg., Eat. 100s, 1000s. Use: C.N.S. stimulant.
- * **DEXTRO PROPOXYPHENE HCl**, N.N.D. 1962. Alpha {d-4-Dimethylamino-1,2-diphenyl-3-methyl-2-propionybutane HCl. See: Dorvon (Lilly)
- W/Acetylsalicylic acid, pheneglycolol. See: Darvo-Tron, Pulvule. (Lilly)
- DEXTRO-RPHAN**, d-3-Hydroxy-N-methylmorphinone. See: Dextromethorphan HBr, N.N.D. 1959
- DEXTROSE**, U.S.P. [α], U.S.P. (Various Mfr.) d-Glucose. Amp. (25 Gm./50 cc.) 50 cc. W/Col. ascorbate & benzyl alcohol [α]. See: Calscarbonate, Amp. (Cale).
- W/Melycaine. See: Melycaine & Dextrose. Amp. (Lilly).
- DEXTROSE & SODIUM CHLORIDE INJECTION**, U.S.P.
- DEXTROSULE**. (Storck) d-Amphetamine sulf. 5 mg./Cap. Rat. 100s, 1000s
- DEXTROSULPHENIDAL**. (Winthrop) D-(+)-threo-2-dichloroacetamido-1-(4-methylselenophenyl)-1,3-propanediol (2). Use: Antimicrobial.
- DEXTROTEST**. (Ames) Tab. A: salicylsulphonic acid (protein-precipitating agent)
- * Generic name recognized by N.F., U.S.P. or A.M.A.

- Tab. R: copper sulfate, sodium hydroxide, sodium bicarbonate, citric acid (determination of sugar). 30-test Kit. Rat. 30 tabs. refill
- Use: Quick blood sugar estimates
- * **DEXTROTHOXINE**, D-a-Amino-b-[4-(4-hydroxy-3,5-di-iodophenoxy)-3,5-di-iodophenyl]propionic acid. Choloxin sodium salt
- DEXTRO-TUSSIN**, (Ulmer) Dextromethorphan HBr 10 mg., chlorpheniramine maleate 2 mg., phenylephrine HCl 5 mg., ammonium chloride 15 mg./5 cc. Rat. 4 ox. Use: Treatment of cough due to cold.
- DEXTRO UNICELLES**. (Hiss Pharm) Dextro-amphetamine sulfate 10 mg. or 15 mg./cap. Rat. 100s
- Use: (one o.m.) Delayed action, C.N.S. stimulant.
- DTP**. See: Diisopropyl fluorophosphate
- D.H.E.-45**. See: Dihydroergotamine, Amp. (Sandoz).
- DIACETINESE**, N.N.D. 1962, (Pfizer) Chlorpropamide 1-(p-chlorobenzene-sulfo-ethyl)-3-propylurea 100 mg./Tab. Eat. 100s; 250 mg./Tab. 60s, 250s. Use: Hypoglycemic agent.
- DIARISMUL** (Tilden) Paregoric 60 min., bismuth subsalicylate 1 gr., salol 13/5 gr., zinc phenolsulfonate 4/5 gr./fl. oz., pink or yellow, 1 pt., 1 gal. Use: (1-2 tsp. m.p.) Absorbent, soothing, astringent.
- DIACETIC ACID TEST**. See: Acetest, Tab. (Ames)
- DIACETOXYDIPHENYL**. See: Clysodrast (Barnes-Hind Barium Co)
- DIACETOXYPHETHYLOXINDOL**. See: Diacetyldihydroxyphenylisatin, Prep. (Various Mfr.)
- DIACITRIZOATE, SODIUM**. See: Hypaque Sodium, Amp. (Winthrop Labs.)
- DIACETYLDIHYDROXYPHENYLISATIN**. See: Diacetyldihydroxyphenylisatin. Succinylcholine Chloride. See: Anectine Chloride, Vial. (Burroughs Wellcome)
- DIACETYLDIHYDROXYDIPHENYL-ISATIN**. See: Diacetyldihydroxyphenylisatin.
- DIACETYLDIOXYPHENYLISATIN**, N.N.R. 1930. See: Diacetyldihydroxyphenylisatin.
- DIACETYLDIHYDROXYPHENYLISATIN**. (Chemo Pure) 3,3-Bis-(4-acetylhydroxyphenyl)oxindole. (Diacetoxydiphenylisatin, Acetphenolisatin, Diacetoxyphenylloxindol, Bis-salin, Phenylisatin) bis-(acetoxyphenyl) oxindol Oxyphenylisatin acetate. See: Acetalox, Cap. (G. F. Harvey) Acetalocin, Cap. (G. F. Harvey) Endophenolphthalein, Tab. (Rache)

- Isacen, NNR 1940 (No Mfr. currently lists)
- Isocrin Tab (Harrower)
- Nilax, Cap (Direct Lab)
- Prulel Tab (Mission)
- W/Brewer's yeast & sod carboxymethylcellulose See Zylax Tab (Glidden)
- W/Butabarbital, d amphetamine sulf See Nilax Cap (Direct Labs)
- W/Danthron See Via Law Tab (Rowell)
- W/Diactyl Sod Sulfasuccinate
- See Daxol Forte, Tab (Blair Labs)
- Evasof Tabs (Lemman)
- Nalac Cap (Dumas Wilson)
- W/Hamartapine methylbromide, mephobarbital, levamfetamine sulfate, Thyroid See Amcabese Dakaps (Amca Drug)
- W/Prune conc
- See Prubilax, Tab (Bell Pharma)
- Prunisolin Tab (Sull H & Cose)
- W/Prune conc & Sod carboxymethylcellulose
- See Bulalin Tab (Ulmer)
- Prulose Complex, Liq (Harrower)
- W/Prune conc & methylcellulose
- See Hydrolase Fortified, Syr (Upjohn)
- Prulose Tab (Harrower)
- W/Psyllium Hydrocolloid See Effersyl (Stuart)
- W/Sod carboxymethylcellulose
- See Cethytin, Tab (Ascher)
- Effergel bulk (Stuart)
- W/Sod carboxymethylcellulose, dioctyl sod sulfasuccinate
- See Dialose Plus Cap (Stuart)
- DIACETYLMORPHINE SALTS, NND 1962**
- Heroin forbidden in USA by Federal statute because of its addiction causing nature
- DIACETYLTANNIC ACID, NNR 1941** See Acetyltannic Acid (Various Mfr)
- DIACHYLON OINT** See Lead Oleate Oint
- DIACTOL, (Plessner) Vit D 2, calciferol** Cap 50,000 u Bot 100s, 500s
- Use Orally, vitamin D therapy
- DIAD, (Rand) Vit D 50,000 u from viosterol** in oil/Cap Bot 100s, 1000s
- DI-ADEMIL, (E R Squibb)** See *Hydroflumethiazide
- DI-A-DISCS (Reed & Carnrick)** Penicillin, bacitracin streptomycin, chlaramycetin aureomycin & terramycin sensitivity determination discs Set, 24 tabs, Box 12s
- Use: Diagnostic aids
- DIADOL, (Buffington)** Diallylbarbituric acid Elixir 2 gr/fl oz Bot 1 pt, 1 gal Tab 1/2 & 1 1/2 gr Bot 100s, 1000s
- Use: Orally, sedative & hypnotic.
- DIAPEN, NND 1962 (Schenley)** Diphenylpyraline HCl Tab 2 mg Bot 100s, 1000s.
- Use: (1 q 4 h) Antihistaminic
- DIAGNEX, NND 1959 (Squibb)** Quinine carboxylic resin Box of 10 tests
- Use: Test for gastric acidity
- DIAGNEX BLUE, NND 1962 (Squibb)** A carboxylic cation exchange resin in reversible combination with azure A dye Pkg 5s & 50s
- Use Determination of gastric acidity
- DIAGNEX BLUE REAGENT, (Squibb)** Ac copper sulfate solution Bot 15 cc
- Use Acidification of the urine
- DIAGNIOL** See Sodium Acetate
- DIAGNOSTIC KIT, (Ames)** Comb kit for acetone, albumin, occult blood & urine sugar
- DIAGNOSTIC AGENTS.**
- See Benodaine, Amp (Merck Sharp & Dahme)
- Cardiografin, Vial (Squibb)
- Cholografin, Amp (Squibb)
- Diognex, NND 1960 Pow (Squibb)
- Evans Blue, Amp (Worner)
- Fluorescein Sodium, Pow (Merck Sharp & Dahme)
- Histolog 50, Amp (Lilly)
- Isokon, Amp (Mallinckrodt Chem)
- Phentolamine Methanesulfonate (Ciba)
- Piperaxoa (Merck Sharp & Dahme)
- Regitine Amp & Tab (Ciba)
- Renografin Inj (Squibb)
- Toluidine Blue (Abbott)
- See also: Cholecystography Agents
- Liver Function Agents
- Kidney Function Agents
- Urography Agents
- DIAGNOSTIC AGENTS FOR URINE**
- See: Acetest (Ames)
- Albustix (Ames)
- Albustest (Ames)
- Albu Tester (Ames)
- Bumintest (Ames)
- Christix (Ames)
- Chintest (Ames)
- Denca Acetone Test (Denver)
- Denca Sugar Test (Denver)
- Dextrotest (Ames)
- Hematest (Ames)
- Ictatest (Ames)
- Ketostix (Ames)
- Occultest (Ames)
- Uroslik (Ames)
- DIAL, NNR 1951 (Ciba)** 5,5 diallylbarbituric acid Tab 1 1/2 gr Bot 100s
- Use: Orally, sedative & hypnotic
- W/Aminopyrine
- See: Barprene, Tab (Ulmer)
- Cibalgin, Amp & Tab (Ciba)

W/Urethane & monoethylurea. (Ciba) Dial ½ gr., urethane 0.4 Gm., monoethylurea 0.4 Gm./cc. Amp. 1 cc., Carton 5s; 2 cc., Carton 5s, 20s.

DIAL SOAP. (Armour) Hexachlorophene. Complexion soap, 2s. Deodorant bath soap, 2s. Shampoo, Bot. 3½ oz.
Use: Antiseptic soap.

***DIALLYLAMICOL.** Diallyl-diethylaminoethyl phenol di HCl. See: Camafarm (Parke, Davis)

DIALLYLBARBITURIC ACID, N.F. X, N.N.R. 1951; Tab. N.F. X. Allobarbital, Allobarbitone, Curral.

See: Diadol, Ellair & Tab. (Beechington).
Dial, Prep. (Ciba).

W/Acetaphenetidin & aspirin. See: Alprine, Tab. (Ulmer).

W/Acetaphenetidin, aspirin, caffeine & gelsemium tr. See: Darodol, Tab. (Beechington).

W/Aminopyrine.

See: Barprine, Tab. (Ulmer)

Cibalgine, Amp. & Tab. (Ciba)

W/Atropine selfate, Mag. carb., Col. carb. precipitated & bismuth subcarb. See: Diatropine, Tab. (Beechington).

W/Cortaeus tr., Sod. nitrite & nitroglycerin. See: Diatrogueus, Pill. (Beechington).

W/Mephensin, N-acetyl-p-aminophenol, homatropine methylbromide. See: Stri-Eze, Tab. (Beechington)

W/Nitrite Sod., cortaeus tr., sellin & nitroglycerin. See: Vosorutin, Pill. (Beechington).

W/Pat. bromide, Sod. bromide, ommenem bromide & alcohol. See: Nuradol, Liq. (Beechington).

W/Secobarbital sodium. See: Bartwo, Tab. (Ulmer)

W/Stramonium ext. & ephedrine HCl. See: Fhedorline, Pill. (Beechington).

DIALLYLBARBITURIC ACID & URETHANE INJECTION, N.F. See: Dial & Urethane Injection, N.N.R. (Ciba).

DIALMINATE. Mixture mag. carb. and (alminate) dihydroxyaluminum glycinate.

DIALOSE. See: Stuart Dialose, Cap. (Stuart)

DIALOSE PLUS. (Stuart) Diacetyl Sod selfo-succinate 100 mg., Sod. carboxymethylcellulose 400 mg., di-(acetylhydroxyphenyl)-isatin 3 mg./Cap. Bot. 100s, 1000s.

Use: Constipation.

DIAMAGMA. (Wyeth) Attopalgite 3 Gm., pectin 270 mg., in a special alumina gel/30 cc. Bot 12 oz.

Use: Treatment of diarrhea.

DIAMAL LIQUID. (Amid Labs) Neomycin sulfate 300 mg.; kaolin 6 Gm.; pectin 130 mg./oz. Bot 1 pt.

Use: Antidiarrheal

* Generic name recognized by N.F., U.S.P. or A.M.A.

DIAMERAZOLE. (Haskell) Sulfathiazole 37%, sulfadiazine 37%, sulfamerazine 26% or D.5 Gm./Tab. Bot. 100s, 1000s. Sesp. 5 cc. eq. fr. to 1 tab. Bot. 4 fl. oz., 16 fl. oz.
Use: Oroally, triple sulfonamide therapy.

DIA-MER-SULFONAMIDES, N.N.R. 1955.

Selfadiazine, Sulfamerazine

See: Aldiazol-M (Messingill)

Bi-Sulfazine, Susp., N.N.R. (Warren-Teed)

Biselfon, Susp., N.N.R. (Veltex).

Chera Seltabs. (Sulliff & Case).

Combiset-DM, Tab. (Schering)

Diamerzine, Tab., N.N.R. (Vale).

Disulfyn, Susp. & Tab., N.N.R. (Rorer).

Duo-Sulfonyl, Syr., N.N.R. (Fenk).

Duoazine, Susp. & Tab., N.N.R. (Abbott).

Eskadiamer, Tab. (Smith, Kline & French)

Gleco-Dimerzine, Susp. (Danley-Evans).

Liqua-D.M., Susp. (Ethical Pharm.)

Lezomid, Tab. (Falch)

Magmaid Sulf-Alka-Loc (Pitman-Moore)

Mer-Diozine, Susp., N.N.R. (McNeil).

Merdizul, Tab. (Bowman)

Selfadimer, Sesp. (Pitman-Moore)

Sel-Di-Mul, Tab., N.N.R. (Miller).

Sulfonamide Duplex, Savoret & Tab., N.N.R. 1953 (Lilly).

Selfonol, Susp. (National Drug)

Sulmeradine, Tab. (Physician's Drug).

So-Di-Zole, Susp. (First Texas).

Tizine, Susp. (Tilden).

W/Alkoline agents.

See: Aldiazol-M, Susp., Tab. (Messingill).

Bisulfon, Sesp. (Veltex).

Deo-Selfonyl, Sesp. (Cosmir Fenk).

Duoazine, Susp. (Abbott).

Gleco-Dimerzine, Susp. (Danley-Evans).

Magmaid Selfadimer, Sesp (Pitman-Moore).

Sel-Di-Mul, Tab. (Miller).

W/Liver conc. & folic acid. See: Lifosulfos, Tab. (Upjohn).

W/Penicillin V. See: Pen-Vee Sulfos, Tab. (Wyeth)

DIA-MER-THIA SULFONAMIDES.

Suspensions

See: Coco-Sulfonamides Triplex (Lilly).

Gleco-Seltas (Danley-Evans).

Magmaid Selca (Pitman-Moore).

Mutti-Selfo (Rossmar).

Tisel (Tilden).

Tri-Selfonyl (Fenk).

Diamerazale (Haskell).

Tablets

See: Cambodine, Tab. (Ives-Cameron)

Combiset. (Schering)

Diamerazale (Haskell).

Thi-Di-Mer (Pitman-Moore).

Tisel. (Tilden)

Triazolone, Dulcet & Tab. (Abbott).

- Tresamide (Merck Sharp & Dohme)
Tri sulfanyl (Punk)
W/Urethan See Trisulfone, Emulsion (Central)
W/Phenylazodiamino pyridine See Pyridium
Tri Sulfa (Warner Chilcott)
- DIAMERZINE**, N N R 1955 (Vale) Sulfadiazine 0.25 Gm, sulfamerazine 0.25 Gm / Tab Bot 100s, 1000s
Use Disulfonamide therapy
- DIAMET.** (Lynn) Methamphetamine HCl, 15 mg; methapyrilene HCl 60 mg / timed disintegrating cap Bot 50s, 250s Methamphetamine HCl 5 mg methapyrilene HCl 25 mg / Tab Bot 100s
Use Obesity control
- DIAMETHINE** See Dimethyl tubacurarine
- DIAMINE DIPHENICILLIN G.** See Benzethacl, Prep
- DIAMINIDE MALEATE** (Merck Sharp & Dohme) Pyrimidine maleate Tab (Blue) 25 mg, Bot 100s 1000s; (Peach) 50 mg, Bot 100s, 1000s
Use Antihistamine
- DIAMINODIPHENYLSULFONE** (Dapsone) See Avlosulfon, Tab (Ayerst)
Dapsone (Abbott)
Glucosulfone Sod (Parke Davis)
- 2,6-DIAMINO-3-PHENYLAZOPYRIDINE HYDROCHLORIDE.**
See: Phenylazo diaminopyridine hydrochloride
Use Urogenital antiseptic
- DIAMINOPROPYL TETRAMETHYLENE.** See Spermine
- DIAMOX**, N N D 1962 (Lederle) Acetazolamide 2 acetylamino 1,3,4 thiadiazole 5 sulfanamide Syr (250 mg / 5 cc) Bot 4 fl oz Tab, (250 mg) Bot 25s 100s 1000s Vial (500 mg / 5 cc) 1 V
Use Diuretic Carbonic anhydrase inhibitor
- DIAMOX SEQUELS** (Lederle) Acetazolamide 500 mg / Cap Bot 30s
Use Diuretic
- DIAMOX SODIUM**, H N D 1962 See Acetazolamide Sod
- DIAMPROMID SULFATE.** (Lederle) N (2 [(methyl)phenethylamino]propyl]propionamide sulfate
Use Analgesic
- DIAMTHAZOLE DIHYDROCHLORIDE.** N N D 1962 See: Asterol Preps (Rache)
- DIAMYCIN.** (Maurry) Dihydrastreptomycin 300 mg, kaolin 2.92 Gm pectin 0.26 Gm / oz Susp Bot 90 cc
Use: Orally, treatment of diarrheas
- DIANABOL** (Ciba) *Methandrostenedione 2.5 and 5 mg / Tab Bot 100s
Use: Anabolic
- * Generic name recognized by N.F., U.S.P. or A.M.A.
- DIANEAL** (Baxter) 1.5% Dextrose 7% Dextrose Bot 1000 ml
Use Intraperitoneally treatment of renal failure
- DIAPARENE CHLORIDE**, N N D 1962 (Breon) Methylbenzethonium chloride Pow 1 1800 Bot 3 1/2 oz, 9 oz Oint, 1 1000 Tube 1 oz, 2 oz Jar 1 lb Tab (0.09 Gm) 1 Tab added to 2 qts of water makes a 1 22000 sol lot 1 1500 Bot 4 oz
Use Surface active disinfectant
- DIAPARENE PERI-ANAL CREAM.** (Breon) Methylbenzethonium chloride 1 1000 zinc oxide, starch cod liver oil & casein Cream Tube 1 oz, 2 oz, 1 lb
Use Rectally dermatitis
- DIAPEC** (Pfizer) Neomycin, sulloguanidine methylhomatropine bromide
Use Gastrointestinal infections
- DIAPEP.** (KIR Lab) Amphetamine sulfate 2.5 mg, pyranisamine maleate 12.5 mg / tab Bot 100s
Use Anobesity
- DIAPHINE HYDROCHLORIDE** (Sohyun) [Et phenamine HCl] 3 Phenyl 2 hydroxybenzoate HCl Oint 1% (free amine) 1 oz Tube Sol 1.4% (Mandelate), Vial 10 cc Supp 10 mg (HCl) each 2 Gm See Melaphine (Sohyun)
- DIAPHRAGM MUSCLE EXTRACT.** See Carnoetan, Sol (Covendish)
- DIARRHEA THERAPY.** See Antidiarrheals
- DIASAL.** (Fougero) Salt substitute Pot chloride, glutamic acid Shaker Bot 2 oz, Bot 8 oz Pkg 1 Gm
Use: Salt free diet
- DIASONE SODIUM**, N N D 1962 (Abbott) Sulfonamide Sod (Disodium formaldehyde sulfoxylate derivative of p-diaminodiphenylsulfone) 0.15 Gm & 0.33 Gm Tab Bot 100s, 1000s
Use Orally, leprosy therapy.
- DIASTASE** See Aspergillus oryzae enzyme
W/Conca papaya enzyme See: Coripeptic Lq (Upjohn)
- W/Pepsin, rennin, hydrochloric acid & lactic acid See: Diastanal Lq (Smith Dorsey)
- DIASTASE CONCENTRATE.** See Eugein, Sol (Rexall)
- DIASYSTOL** (Rorer) Acanite root tr vera trum tr, passiflora tr, each 2 4/5 fl dr Bot 14 fl oz 5 pt, Jug 1 gal
Use: Hypertension
- DIATRAEGUS** (Buffington) Diallylbarbituric acid 1/4 gr, atrataegus tr, 2 min, Sod nr trite 1 gr, nitraglycerin 1/250 gr / Pilllet Bot 1000s
Use: (1-3 daily) Sedative w/vnsaditator
- DIATRINE HYDROCHLORIDE**, N N D 1959

(Warner Chilcott) (Methaphenilene HCl) N,N-Dimethyl-N'-alpha-phenyl-N'-phenylethylenediamine HCl Tab (50 mg) Bot. 100s, 1000s.

Use: Orally, antihistamine.

* **DIATRIZOATE METHYLGLUCAMINE**, N.N.D. 1962. See: Cardiografin, Vial (Squibb)

Renografin (Squibb)
Renovist Inj. (Squibb)

W/ladipamide methylglucamine. See: Sino-grafrin, Lq. (Squibb)

* **DIATRIZOATE SODIUM**, N.N.D. 1961. See: Hypaque Sod, Amp. (Winthrop Labs.)

DIA-TROPINE, (Burlington) Diallylbarbituric acid $\frac{1}{4}$ gr., atropine sulfate $\frac{1}{300}$ gr., Mag. carb. $2\frac{1}{2}$ gr., Cal. carb. precipitated $3\frac{1}{2}$ gr., bismuth subcarb. 1 gr./Tab. Bot. 100s, 1000s.

Use: Antacids & antispasmodics

DIATUSSIN, (Ames) *Drasera rotundifolia* & *Thymus vulgaris*, alcohol ext. Cont., Bot w/dropper 6 cc. Syr., Bot. 4 oz., 1 pt., 1 gal.

Use: (1-2 tsp. q. 3-4 h.) Cough preparation.

* **DIAYERIDINE**, (Burroughs Wellcome) 2,4-diamino-5-[3',4'-dimethoxybenzyl] pyrimidine.

Use: Antibacterial (for coccidiosis in chickens)

* **DIAZEPAM**, (Hoffmann-La Roche) 7-chloro-1,3-dihydro-1-methyl-5-phenyl-2H-1,4-benzodiazepin 2-one See: Valium (Hoffmann-La Roche, Inc)

DIAZINE CITRATE, (Massengill) Sulfadiazine 7.7 gr., Cit. citrate 8 gr./Tab. Bot. 100s, 500s, 1000s.

Use: Sulfadiazine susceptible organisms.

DIAZOLINE, (Abbott) Sulfadiazine 25 Gm., sulfathiazole 25 Gm./Tab. Bot. 100s, 1000s.

Use: Di-sulfonamide therapy.

DI-AZO-MUL, (First Texas) Sulfadiazine 40 gr./fl. oz. Bot. 1 pt. Citrated, Bot. 1 pt.

DIAZOMYCINS A, B, & C. Antibiotic obtained from *Streptomyces embolicus* Under Study.

6-DIAZO-5-OXO-L-NORLEUCINE. See D. O. N. (Parke, Davis)

* **DIAZOXIDE**, (Schering) 7-chloro-3-methyl-1,2,4-benzothiadiazine-1,1-dioxide.

DIAZPRIN, (Squibb) Sulfadiazine 5 gr., aspirin 3 gr./Water. Box 12s.

Use: Topically, sulfanamide therapy w/analgesic.

DIBASIC CALCIUM PHOSPHATE, U.S.P. See Dicalcium Phosphate

DIBENCIL. See: Benzathine Penicillin G.

* Generic name recognized by N.F., U.S.P. or A.M.A.

DIBENZIPEN, (Bristol) Dibenzylamine penicillin G 200,000 u., Pot. penicillin G 100,000 u./5 cc. Vial.

Use: Penicillin therapy.

DIBENZYLETHYLENEDIAMINE DIPENICILLIN G, U.S.P. See: Benzethacil, Prep.

DIBENZYLIN, N.N.D. 1962 (Smith, Kline & French) N-Phenoxyisopropyl-N-benzyl-beta-chloroethylamine HCl. Phenoxybenzamine. Cap. 10 mg. Bot. 100s.

Use: Long acting adrenergic blocking agent.

W/Bauwolfia serpentina & protaveratrine See: Mio-Pressin, Cap. (Smith, Kline & French)

DIBENZYL SUCCINATE W/Mephenesin, salicylamide. See: Easpmus, Tab. (M. H. Smith).

DIBESTIL, (Breon) Diethylstilbestrol dipropionate, Tab. 25 mg Bot. 100s.

Use: Orally, estrogen therapy.

DIBISTINE, (Ciba) Anilaine HCl 50 mg., pyribenzamine HCl 25 mg./tab. Bot. 100s, 1000s.

Use: (1-2 q.i.d., p.c. & h.s.) Di-antihistamine therapy.

3,3-DIBROMOSALICYLALDEHYDE.

See: Dolyde. (Hynson, Westcott & Dunning)

Use: Fungicide & antiseptic.

* **DISUCAINE HYDROCHLORIDE**, U.S.P., N.N.R. 1956; Inj., U.S.P. 2-Butoxy-N-(2-diethylaminoethyl)cinchonamide HCl. (Cinchocaine, Nuparals, Percaine)

See: Benzoin, Oint. (Breon).

Nupercaine HCl. (Ciba).

Use: Surface & spinal anesthesia.

W/Anipyrine, hydrocortisone, polymyxin B, sulfate, neomycin sulfate See: Otacort, Lq (Lemman)

W/Colistin sodium methanesulfonate, citric acid, sodium citrate. See: Caly-Mycin Injectable (Warner-Chilcott)

DIBULINE SULFATE, N.N.D. 1962. (Merck Sharp & Dahme) Dibutaline sulfate 25 mg / cc Vial 5 cc.

Use: I.M., S.C., antispasmodic

* **DIBUTOLINE SULFATE**, N.N.D. 1962. (2-Butylcarbamyl-oxyethyl)dimethyl-ethylammonium sulfate. See: Dibuline Sulfate. (Merck Sharp & Dahme)

Use: G.I., G.U. & biliary spasm; mydriatic & cycloplegic

DICALATE WAFLETS, (Suthff & Case) Dicalcium phosphate, 15 gr./Tab. Bot. 100s

Use: Calcium therapy

DICALCIUM PHOSPHATE, A.D.E. 1955 Dibasic Calcium Phosphate, U.S.P., Monocalcium Phosphate (Various Mfr.) Cap. $7\frac{1}{2}$ gr.

10 gr Tab 7½ gr 10 gr, 15 gr Wafers
15 gr

Use: Calcium supplement

See Calphate Tab (Marrell)

D C P 340, Paw (Parke, Davis)

W/Bone meal powder See: Cadal, Cap
(Ingram)

W/Bone phosphate purified See Bone Phos-
phate Laz (Upjohn)

W/Cal gluconate & Vit D (Various Mfr)

Cap Tab & Wafer See Vitamin D W/
Calcium Salts table

W/Ferrous sulfate vit B1, B2 D See
Aprivate Cap (Approved Pharm)

W/Iron & Vit D (Various Mfr)

Lilly—Pulvules Bot 100s 1000s

Premo—Cap Bot 100s 1000s

See Vitamins Prenatal table

W/Thiobromine sod acetate phenobarbital
See: Barbithea Tab (C D Smith)

W/Vitamin D (Various Mfr) See: Vitamin D
W/Calcium Salts table

W/Viosteral (Squibb) Vit D 660 u d-basis

Cal phosphate 583 mg Cal gluconate 389
mg/Wafer or 2 Cap Bot 90s

DICAL-D (Abbott) Dicalcium phosphate D 5

Gm Vit D 333 u/Cap Bot 100s 500s

1000s Wafers Dicalcium phosphate 972

mg Vit D 666 u Bot 30s 250s

Use Increase oral intake of calcium &
phosphorus

W/Iron (Abbott) D calcium phosphate D 5

Gm Vit D 333 u iron pyrophosphate
soluble 87 mg/Cap Bot 100s 500s

W/Vit C (Abbott) D calcium phosphate D 5

Gm Vit D 333 u ascorbic acid 15 mg/
Cap Bot 100s 500s 1000s

DICALDIMIH (Abbott) Dicalcium phosphate

anhy 725 mg, ferrous sulfate 50 mg thia-
mine mononitrate 3 mg riboflavin 3 mg

nicotinamide 12 mg Vit D 500 u/Cap
Bot 100s 500s, 1000s

Use: Calcium & vitamin therapy

W/Vit C (Abbott) Same formula as Dical-
dimih but with addition of 50 mg ascorbic

acid/Cap Bot 100s 500s 1000s

DICALPHOS (New Mexico Pharm) Vit D

333 u, dicalcium phosphate 2619 mg,
calcium gluconate 162 mg/cap Bot 100s

1000s

Use Dietary supplement

DICASAN (Chicago Pharmacaf) Diethyl

sodium sulfosuccinate 100 mg casanthranol

30 mg/Tab Bot 100s 1000s

Use Laxative

DICASTRON (Elder) Strychnine arsenate

1/100 gr, iron arsenate 1/50 gr, d g talin
patent 1/120 gr sports ne sulfate 1/25

* Generic name recognized by NF USP or A.M.A.

gr, cactus grandiflorus ext 1/50 gr/Tab.
Bot 100s, 500s 1000s

DI-CET, (Sanford & Son) Methylbenzethonium

chloride 24.4 Gm, Sod carb monohydrate

48.8 Gm, Sod nitrite 24.4 Gm, Irisodium

ethylenediamine tetra acetate monohydrate

2.4 Gm Pow Pkg 2.4 Gm, Box 24s

Use: Disinfection of dental instruments

DICHLORALANTIPYRINE, Dichloralphen-
azone Chloralpyrine A complex of 2 mol

chloral hydrate with 1 mol antipyrine See
Sominol, Tab (National Drug)

W/Amobarbital See: A.N.S., Tab (Kenwood
Labs)

W/Isametheptene mucate, N acetyl p amiza-
phenal See Midrin Caps (Carrick)

DICHLORALPHENAZONE See Dichloral-
antipyrine

DICHLORAMINE T, N.N.E. 1949, A.D.E.
1955 P Toluene-sulfone dichloramine

Use: (1 to 5% in chlorinated paraffin)
Antiseptic

DICHLORON, See: Mechlorethamine HCl
Sterile NF

DI-CHLORICIDE See: p Dichlorobenzene
(Merck)

***DICHLORIBONE ACETATE**, 9o 11b di-
chloro di,4 pregnadiene 17o 21 dol 320

dione 21 acetate See Diloderm (Schering)

W/Aluminum chlorhydroxy allantoinate

neamycin sulfate See Neo Diloderm Aero-
sol and Cream (Schering)

DICHLOR-MAPHARSEN (Parke, Davis) Di-
chlorophenarsine HCl Amp 0.05 Gm

DD75 Gm, Pkg 10s, 100s; 0.75 Gm, Pkg
10s

Use I.V., antisyphilitic arsenical prepara-
tion

DICHLOROBENZENE (Merck) Di chloride,
Can 1 lb

DICHLORODIPHENYL TRICHLOROETHANE,
See: Chlorophenothane USP

3,7-DICHLORO-8-HYDROXYQUINOLINE

See: Chlorquinolal (Geigy)

Sterosan, Cream & Oint (Geigy)

DICHLOR-MAPHARSEN (Clarasen, N.N.E.
1949)

***DICHLORMETHAZANONE**, 2 (3,4-dichloro-
phenyl)-3-methyl-4-metathiazanone 1 d ox-
ide

**DICHLOROPHENARSINE HYDROCHLO-
RIDE**, N.F. 3 Amino-4-hydroxyphenyl-dichloro-
arsene (Clararsen Clararsen Fontarol

Nalarsal) See Dichlor Mapharsen Amp
(Parke Davis)

DICHLOROPNE (Sindar Corp) G 4 bit
(5-chloro-2-hydroxyphenyl) methane Related

to hexachlorophene See Eez (Pfizer)

DICHLOROSALICYLANILIDES, 5,3 end

3,4'-*w*/salicylanilide, 5 chloro-salicylanilide, undecylenic acid & zinc undecylenate. See: Salundek New Oint. (Wallace & Tierman).

***DICHLORPHENAMIDE**, U.S.P., N.N.D. 1962. Tab. U.S.P. 1,3 disulfamyl-4-5-dichlorobenzene.

See: Doranide, Tab. (Merck Sharp & Dohme) Oratrol, Tab. (Alcon)

N,N-DICHLOR SULFAMIDORENZOIC ACID. See: Holazone, N.F.

DICLONE, See: Cyclonine Hydrochloride. (Pitman-Moore)

DICODETHAL ELIXIR, (Lennett) Dihydrocodeinone bitartrate 1/6 gr./fl. oz. in el. ferpin hydrate. Bot. 1 pt., 1 gal.

Use: Cough preparation.

DICODID, N.N.D. 1962 (Bilhuber-Knoll) Dihydrocodeinone bitartrate. Viol 15 gr., Tab. 5 mg., Bot. 10s, 100s, 500s. Hydrochloride: Pow., Viol 15 gr.

Use: Narcotic.

DICODRINE SYRUP, (Braun) Phenylephrine HCl 50 mg., phenylpropranolamine HCl 3.34 mg., p-ol. guaiacolate 83.34 mg., dihydrocodeinone bitartrate 1.67 mg./5 cc. Bot. fl., gal.

Use: Antitussive, antihistaminic, decongestant, expectorant.

DI-COLD, (Haug) White layer-aspirin 200 mg., phenacetin 150 mg., caffeine 30 mg.; blue layer-ascorbic acid 30 mg., phenylephrine HCl 5 mg., chlorpheniramine maleate 1 mg./Tab. Bot. 100s & 1000s.

Use: Symptoms of colds and sinus congestion.

DICOPHANE, N.N.R.

See: Chlorophenothone, U.S.P.

DDT.

DICOPHEN, (Buffington) Dihydrocodeinone bitartrate 10 mg., pyrilamine maleate 80 mg., phenylephrine 30 mg., Pot. citrate 20 Gm./oz. Syr. Bot. 1 pt. & 1 gal.

Use: Treatment of cough due to colds.

DICORIL, (Lemmon) Pyrilamine maleate 80 mg., carbopentane citrate 43.5 mg., sod. citrate 1 Gm./fl. oz. Bot. 3 fl. oz., 1 pt., 1 gal.

Use: Relief of coughs due to colds

DICORVIN, (Amfre Grant) Diethylstilbestrol 100 mg., Vit. C 667 mg., Vit B-2 10 mg./Tab Bot. 50s, 100s, 500s.

Use: Orally, 1 to 3 daily, estrogen and vitamin supplement

DICOTRATIL, (Buffington) Dihydrocodeinone bitartrate 1/6 gr., Pot. citrate 30 gr., aromatized syr., q.s./fl. oz. Syr., Bot. 1 pt., 1 gal.

* Generic name recognized by N.F., U.S.P. or A.M.A.

Use: (1 tsp. q. 3-4 h., mp) Antitussive agent.

W/Paraminyl, (Buffington) Dihydrocodeinone bitartrate 1/6 gr., paraminyl (pyranisamine maleate) 1 1/2 gr., Pot. citrate 30 gr., aromatized syr. q.d./fl. oz. Bot. 1 pt., 1 gal.

Use: (1 tsp. q. 3-4 h.) Cough sedative.

DICOUMARIN. See: Dicumarol (Various Mfr.)

DICOUMAROL. See: Dicumarol (Various Mfr.)

DICRYSTICIN, FORTIS, (Squibb) Procaine penicillin G 300,000 u., Pot. penicillin G 100,000 u., dihydrostreptomycin sulfate 0.5 Gm., streptomycin sulfate 0.5 Gm. Viol 1 dose.

DICUMAROL, N.N.D. 1962. (Wisconsin Alumni Res. Found) 3,3'-methylenebis(4-hydroxy-coumarin). (Dicoumarol, Dicoumarin, Dihydroxy-coumarin, Melitoxin)

Use: Blood anticoagulant.

W/Rhomonin, (Strassenburgh) Dicumarol 50 mg., rutin 30 mg., ascorbic acid 40 mg./Tab, Bot, 100s.

Use: Orally, anticoagulant w/vitamin P & C therapy.

W/Rutin & ascorbic acid See: Dicumarol w/rhomonin, Tab (Strassenburgh).

DICURIN PROCAINE, N.N.D. 1962. (Lilly) Merethoxylline procaine 100 mg., theophylline 50 mg./cc. Amp & 10 cc. Merethoxylline procaine 200 mg Theophylline 100 mg./cc. Amp. 2 cc.

Use: I.M., I.V., S.C. (0.5 cc. to 2 cc. daily), diuretic.

***DICYCLOMINE HYDROCHLORIDE**, N.N.D. 1962. See Betyl HCl, Amp, Cap. & Syr. (Merrell)

W/Carboxymethylcellulose Sod., sodium lauryl sulfate, casanthanol. See: Benulox. (Merrell)

W/Magnesium ox. hydrated, alum hydrated, mag. trisulfate, sodium lauryl sulf, methyl cellulose See: Kolantyl Wafers (Merrell)

DICYSTINE. See: Cystine (Various Mfr.)

DIDREX, (Upjohn) (-)-N-benzyl-N,N-dimethylphenethylamine hydrochloride. Benzphetamine HCl. 25 and 50 mg./Tab Bot 100s

Use: Appetite depressant.

DIDROXANE W/Salicylic acid. See: Karium, Oint. & Pow. (Karium)

DI-EFRIN, (Buffington) Dihydrocodeinone bitartrate 10 mg., ephedrine HCl 50 mg., ammonium Cl 64 mg., potassium guaiacal-sulfonate D51 Gm./az. Syr. Bot. 1 pt.

Use: Antitussive

***DIENESTROL**, U.S.P., N.N.D. 1962; Tab U.S.P. (Non-steroid, synthetic estrogen) (Di-

dose. Vial 150 T.B.U., Box 4s, 25s; Vial 1500 T.B.U.

Use: Dispersion & absorption of subcutaneously injected fluids.

DIGALEN, N.N.B. 1952. (Roche) Cardioactive principles of digitalis lvs. Amp. (1 U.S.P. u) 2 cc., Box 6s, 12s, 100s. Sol., oral use (1 U.S.P. u./cc) Vial 1 A. oz. Tab. $\frac{1}{2}$ U.S.P. u. Bot 100s; 1 U.S.P. u. Pkg. 4s, Bot. 100s

Use: Digitalis therapy.

DIGALLOYL OLEATE. See: Neo-A-Pol (Texas Pharm. Co)

DIGAMMACAINI. (Lilly) 1-Benzomido-1-phenyl 3-piperidinopropene.

W/Erythromycin. See: Erythromycin L.M. Solution. (Lilly)

DIGESTANT. (Canright) Pancreatin 5 $\frac{1}{2}$ gr., ox bile Est 2 gr., pepsin 5 gr., betaine HCl 1 gr./E.C. Tab Bot. 100s, 1000s

Use: Natural digestants.

DIGESTIL. (Breon) Diethylstilbestrol dipropionate Tab 25 mg., Bot. 100s

Use: Orally; estrogen therapy.

DIGESTIVE COMPOUND TABLETS. (Morton & Converse) Pancreatic triple strength 2 gr., papain 1 gr., est. ox bile 2 gr., pyridoxine HCl 100 mg., betaine HCl $\frac{1}{2}$ gr./Tab. Bot. 100s, 500s

DIGESTULES. (Caldwell & Bloor) Pepsin 200 mg., dl methionine 25 mg., whole dry pancreas 250 mg., ox bile extract 50 mg., mixed oxidized bile acids 50 mg./Cap Bot. 100s, 1000s

Use: Digestive aid

DIGICARDIUM. (Rorer) Purified prep. from digitalis lvs. Cap. Tab. 1 & 1.5 gr., Bot. 100s, 500s, 1000s.

Use: Digitalis therapy.

DIGICARDIUM W/Theobromine. (Rorer) Digisardium $\frac{1}{2}$ gr. or 1 gr., theobromine 5 gr., Bot. 100s, 500s, 1000s.

Use: Digitalis w/diuretic.

DIGICO. (Sulfriff & Case) Digitalis pow. 1 $\frac{1}{2}$ gr./Cap. Bot. 100s, 1000s

DIGICOID UNICELLES. (Miss Pharm.) Digitoxin 0.15 mg./cap. Bot. 100s.

Use: Delayed action, congestive heart failure.

DIGICOTIN. (Grant) Digitalis U.S.P. Tab. 0.1 Gm. Bot. 20s, 100s.

Use: Orally; digitalis therapy.

DIGIFOLIN, N.N.B. 1952 (Ciba) Not Available Cardioactive principle of digitalis lvs

DIGIFORBIS KAPSEALS. (Porke, Davis) Amp. 1 cc., Box 12s, 100s. 1 U.S.P. u. or 0.1 Gm. (1 $\frac{1}{2}$ gr.) digitalis U.S.P., or 1 cc. digitalis (1 $\frac{1}{2}$ gr.) U.S.P./Cap. Bot 100s, 500s Liq. (Each ss. equals 1 $\frac{1}{2}$ U.S.P. u. or 125% strength digitalis tr. U.S.P.) Bot 1 oz. Tab. (Each

* Generic name recognized by N.F., U.S.P. or A.M.A.

Tab. representing 0.1 Gm. (1 $\frac{1}{2}$ gr.) digitalis tr. U.S.P.) Bot. 50s, 100s.

Use: Cardiac conditions.

DIGIGLUSIN. (Lilly) Digitalis glycosides. Amp. 1 U.S.P. u./cc, 1 cc. Box 6s, 100s, Tab. $\frac{1}{2}$, 1, 1 $\frac{1}{2}$ U.S.P. u., Bot. 100s, 500s.

Use: Digitalis therapy.

DIGILANID, N.N.D. 1959. (Sandoz) Lanatoside A 47%, lanatoside B 16%, lanatoside C 37%.

Amp. (0.4 mg) 2 cc. I.M.; Box 6s, 100s. Tab. (0.333 mg) Bot. 50s, 100s.

Use: Digitalis therapy.

DIGILANIDE C. See: Cedilanid, Amp. & Tab. (Sandoz)

DIGISEALS. (C. F. Harvey) Digitalis leaf powdered in oil. Cap. 1 $\frac{1}{2}$ gr. equiv. to 1 U.S.P. u. Bot. 1000s.

Use: Orally; myocardial stimulant.

DIGITALIN. (Merk Sharp & Dohme) Digitalis, potency 10 times that of digitalis lvs. U.S.P. Pow., Bot. 5 gr., 15 gr., $\frac{1}{2}$ oz.

Use: Digitalis therapy.

W/Oubaine. See: Nalibaine, Liq. (Varick).

DIGITALINE NATIVELLE, N.N.B. 1950. (Varick) Chief active glycoside of digitalis purpurea (Digitoxin). Amp. (0.2 mg) 1 cc., & 2 cc. Sol. (oral use) 1:1000 (1 mg/cc) Bot. 10 cc. Tab. 0.1, 0.15, 0.2 mg. Bot. 40s, 100s, 250s, 1000s

Use: Digitalis therapy.

DIGITALINUM VEBUM. See: Digitalin, Pow. (Merk Sharp & Dohme)

* **DIGITALIS**, U.S.P. (Various Mfr) Cap., N.F., 1 gr, 1 $\frac{1}{2}$ gr. Pill $\frac{1}{2}$ gr., $\frac{3}{4}$ gr., 1 gr. Tab., U.S.P., $\frac{1}{2}$ gr., $\frac{3}{4}$ gr., 1 gr., 1 $\frac{1}{2}$ gr. Tr., N.F., 73%. Pow., U.S.P.; Inj., N.F. X

DIGITALIS OLYCOSIDES.

See: Acylanid (Sandoz).

Cedilanid, Tab (Sandoz).

Cedilanid D, Amp (Sandoz).

Digalen (Roche).

Digifolin (Ciba)

Digiglusin (Lilly).

Digilanid (Sandoz).

Digitoxin U.S.P.

Digoxin

Gitoxin

Gitoxin (White)

Lanoxin (Burroughs Wellcome)

DIGITALIS LEAF, POWDERED.

See: Digifolin, Tab. (Grant).

Digiseals, Cap. (C. F. Harvey)

Digitalin, Tab. (Upjohn).

W/Theobromine. Cal. salicylate & phenobarbital. See: Theodigital, Tab. (Drug Products).

DIGITALIS LEAF, TABLOID. (Burroughs Wellcome) Tab. 1 $\frac{1}{2}$ gr. Bot. 100s, 1000s

DIGITALIS TINCTURE W/Crotaegus fr., straphanilus fr & strychnine See: Cardol, Tab (C D Smith)

DIGITIN See: Digitonin, Paw (Merck Sharp & Dahme)

DIGITONIN, (Merck Sharp & Dahme) Digitin Bot 5 gr, 15 gr

DIGITORA, (Upjohn) Digitalis lvs paw Tab 0.85 gr equiv Ia 0.55 USP v; 128 gr equiv Ia 0.83 USP v Vial 30s

Use: Orally digitalis therapy

***DIGITOXIN**, USP NNR 1950, Int. USP; Tab USP (Unidigin) (Various Mfr) Amp (0.2 mg/cc) 1 cc Box 6s 25s 100s Cap in oil 0.1 mg 0.2 mg Box 100s, 1000s Bot 0.1 mg 0.2 mg 25s 100s

Use Digitalis therapy

See Cardigin, Tab (National Drug)

Crylodigin, Amp & Tab (Lilly)

Digicoid Unicelles (Hiss Pharm)

Digisidin, Amp & Tab (Winthrop Labs)

Digiloline Nativelle Amp & Sol (Varick)

Olucodigin, Tab (Cole)

Puradigin, Amp & Tab (Wyeth)

Unidigin, Tab (Merrell)

W/Carboxymethylcellulose & Sod See: Poxalin, Tab (Grant)

W/Gilolin See Myapurin, Tab (Palch)

W/Nitroglycerin, strychnine sulf See: Uniloxin, Liq (Tilden)

W/Strophanthin & strychnine sulfate See: Procardium, Tab (Grant)

W/Strophanthin strychnine sulfate nitroglycerin, cactus grandiflorus See: Pasonol, Liq (Tilden)

DIGITOXIN, ACETYL. See Acylanid, Tab (Sandoz)

DIGLYCOCOLL HYDROCHLORIDE. See: HCl Cap (Frost)

DIGLYCOCOLL HYDRIODIDE. See: HI Cap (Frost)

DIGLYCOCOLL HYDRIODIDE-IODINE, NND 1959

See: Bursaline, Tab (Burnham)

Glabaline, Tab (Warren Teed)

DIGLYSAL (Chemo Pura) **GLYCOL MONOSALICYLATE**.

DIGOLASE, (Boyle) Pancreatin 300 mg, teralase 10.5 mg, dehydracholic acid 25 mg polysorbate 80 10 mg /Cap Bos 100s

Use: Orally, digestive enzyme deficiencies

***DIGOXIN**, USP, NND 1962 Int., USP; Tab, USP (Burroughs Wellcome) Cryst glycoside isolated from lvs of digitalis lanata Amp (0.5 mg/cc) 1 cc (0.25 mg/cc) 2 cc Box 12s 100s Tabloid 0.25 mg & 0.5 mg, Bot 100s 500s Elixir Pediatric

* Generic name recognized by NF USP or A.M.A.

(0.05 mg/cc) Bot 2 fl oz See: Lanoxin (Burroughs Wellcome)

Use: Digitalization.

***DIHEXYVERINE HYDROCHLORIDE**, (Leeming Miles Pharm) 2 piperidoneethyl ester of bicyclohexyl 1 carboxylic acid See: Melspas (Leeming Miles Pharm)

DINYCON, (Consolidated Midland) Diphenylhydantoin 250 mg /Cap Bot 30s

Use: Antiepileptic

DINYDAN SOLUBLE. See: Diphenylhydantoin Sodium (Various Mfr)

DINYDERGOT, (Sandoz) Dihydroergolamine methanesulfonate 1 mg /Tab

Use: Treatment of migraine

DI-NYDRIN, (Broemmel) Plastic drop bot 7.5 cc

Use: Sterile ophthalmic solution

DIHYDROCHLOROTHIAZIDE.

See: Esdrez (Ciba)

Hydra Divril (Merck Sharp & Dahme)

DIHYDROCODEINE, (Mollinckrodt) Paracodin Orocode See: Ropacodin Bitartrate (Knoll)

Use: Antitussive & analgesic.

W/Pyrimine maleate di desoxyephedrine HCl, pol guaiacolsulfonate, omm chloride, lortar emetic, chloroform See: Banadene Syr (Banar)

***DIHYDROCODEINONE BITARTRATE**, N.F., NND 1962, Syr, N.P., Tab, N.P

See Codone, Tab (Lemmon)

Dicodethal, El (Lanett)

Dicodid, Tab & Vial (Bilhuber Knoll)

Mercodinone, Tab (Merrell)

Stodcodon, Syr (Stoddard)

Use Narcotic

W/Aluminum Aspirin, Acetophenelidin, Caf Feine See: Dracogesic No 3 (Cartone)

W/Ammonium chloride & chloroform See: Nullicof, Liq (Noyes)

W/Ammonium chloride, tartar emetic, Sod citrate, citric acid & chloroform See: Anacal Cough Syrup (Warren Teed)

W/APC See: Rapadyne, Tab (Knoll)

W/Asparin, acetophenelidin & caffeine See: Anadynas DHC, Tab (Brean)

W/Chlorpheniramine maleate, Disoephedrine hydrochloride See: Hasanone No 2 Sol (Maskell)

W/Chlor Trimetan maleate, Sod salicylate Sod citrate, caffeine & glyceryl guaiacolate See: Caricodin, Syr (Schering)

W/Citric acid, ammonium chloride chloroform & alcohol See: Per Sed, Liq (Columbus)

W/di Desoxyephedrine HCl & Pot Iodide See: Bransyl Syr (McNeil)

W/Dihydrocodeinone Terephthalate, Homa

W/Pyrilamine, desoxyephedrine HCl, ammonium chloride, citric acid, chloroform & alcohol See Pyra Sed, Cough Prep (Columbus)

W/Pyrilamine maleate, dl desoxyephedrine HCl, amm chloride, pal guaiacal sulf, chloroform & tartar emetic See Tusstor Syr (Tylar Labs)

W/Pyrilamine maleate, mesopin, ammonium chloride & Sod citrate See Endalussin, Syr. (Endo)

W/Pyrilamine maleate, pot guaiacal sulfonate, sod citrate See Codimal, Syr (Central Pharm)

W/Pyrilamine, Pot guaiacal sulfonate, Sod citrate & chloroform See Pyrilcod, Cough Prep (Raymer)

W/Salicylamide & caffeine See Codis, Tab (Haack)

W/Sod citrate, ammonium chloride, ephedrine HCl tartar emetic See Semcol, Sol (Massengill)

W/Sod citrate citric acid, ammonium chloride, Pot guaiacal sulfonate, chloroform & benzyl alcohol in syr vehicle See Amophan C, Syr (McNeil)

W/Sod citrate, citric acid, lulu syr & menthol See Dicosed, Cough Prep (Flint, Eaton)

W/Terpin hydrate See Draderne, Cough Prep (Pitman Moore)

W/Tri histin, pal guaiacal sulfonate NH₄Cl, sod citrate, chloroform See Trihisto Cod Liq (Eczsei)

DIHYDROCODEINONE RESIN COMPLEX. W/Phenyltoloxamine resin complex See Tussionex, Prep (Strassenburgh)

DIHYDROCODEINONE TEREPHTHALATE w/ Dihydrocodeinone Bitartrate Hamatropine Terephthalate Pentylenetetrazol See Para Hycadan (Endo)

DIHYDRODESOXYMORPHINE. See Desomorphine

DIHYDRO-DIETHYLSTILBESTROL. See Hexestral Tab & Vial (Various Mfr)

DIHYDRO DISPULATOR. (Squibb) Micro pulverized dihydrostreptomycin sulfate, 50 mg

Use: Intections of respiratory tract

DIHYDROERGOTAMINE. (D H E 45) (Sandoz) Dihydroergotamine methanesulfonate Amp [1 mg] 1 cc, Box 6s, 50s

Use: S.C., I.V., I.M.; sympathetic nervous system depressant recommended for migraine headache

W/Scopolamine HBr, phenobarbital Sod, barbital Sod & sandoptal See Plexanal Tab. (Sandoz)

* Generic name recognized by N F, USP or A M A

DIHYDROFOLLICULAR HORMONE. See Estradiol (Various Mfr).

DINYDROFOLLICULINE. See: Estradiol (Various Mfr)

OINHYDROHYDROXYCODEINONE. Oxycodone NCI (Ducodol, Eukodol, Eucodol) (No mfr currently lists)

Use: Narcotic, analgesic.

OINHYDROHYDROXYCODEINONE HYDR-CHLORIDE

W/Dihydrohydroxycodeinone terephthalate hamatropine terephthalate & pentylenetetrazol

See: Nucadan, Tab (Endo)

W/Dihydrohydroxycodeinone terephthalate hexobarbital, hamatropine terephthalate aspirin phenacetin, caffeine See Percobarb, Cap (Endo)

W/Dihydrohydroxycodeinone terephthalate hamatropine terephthalate, aspirin, acetophenetidin & caffeine

See Percobarb Demi (Endo)

***OIHYDROMORPHINONE HCl, USP** N N R 1954, Inj., USP; Tab., USP (Various Mfr)

See Dilaudid Prep N N R (Knoll)

Hymorphan, Sol & Tab, (Endo)

Semcol w/Dilaudid (Massengill)

W/Atropine sulfate See Hymorphan Atropine, Vial (Endo)

2-OIHYDROPHENOBARBITAL. See Mysoline, Prep (Winthrop Labs)

***DIHYDROSTREPTOMYCIN, USP, MND** 1962 A D R Pow (I M)

W/Atapulgitte activated, polymyxin B sulf. pectin See Polymagma, Susp (Wyeth)

W/Bicillin, streptomycin, procaine penicillin G & Pal penicillin G See Bicillimycin All Purpose, Vial (Wyeth)

W/Kaolin and pectin See Diamycin, Susp (Maurry)

W/Polymyxin B sulfate, activated atapulgitte pectin, hydrated alumina powder See Polymagma, Tab (Wyeth)

W/Procaine penicillin See Panmycin A S Liq (Philadelphia Ampaule)

W/Streptomycin See Cambistrep, Vial (Pfizer)

DINYDROSTREPTOMYCIN HYDROCHLORIDE, USP XIV

DINYDROSTREPTOMYCIN OTIC w/BRISTAMIN. (Bristol) Each cc contains 25 mg dihydrostreptomycin & 10 mg bristamin

dihydrogen citrate in glycerine vehicle Bot w/drapper 15 cc

Use: External; otitis, bacterial or fungus in origin

- *DIHYDROSTREPTOMYCIN SULFATE, U.S.P., N.N.D. 1962;** Inj., U.S.P. Oint. See: Daramicin (Smith-Dorsey)
- N/Bristamin dihydrogen citrate.** See: Dihydrostreptomycin ole w/Bristamin, Sol. (Bristol)
- N/Chloromycetin.** See: Chlorostrep, Capsules (Parke, Davis).
- N/Kaolin & pectin.** See: Streptomagma, Liq. (Wyeth).
- N/Penicillin fortified & streptomycin sulfate.** See: Cillimycin, Improved (Wyeth).
Dicyrsticin Fortis (Squibb).
- V/Penicillin G Sod & napoine HCl.** See: P-D N Dental Cones (Novacal Chem. Co.)
- V/Polymyxin B sulf., attapulgit activated, pectin, alumina gel.** See: Spensin-PS, Susp. (Ives-Cameron)
- V/Procaine penicillin G & Pot. penicillin G or Sod.** See: Penicillin preparations.
- V/Streptomycin sulfate, chlorprocaine penicillin G, penicillin G sod.** See: Cer-O-Strep, Vial. (Upjohn)
- V/Streptomycin & procaine penicillin O.** See: Crystimycin A.S., Vial (Squibb).
- DIHYDROSTREPTOMYCIN-STREPTOMYCIN SULFATE.** (Parke, Davis) Streptomycin 0.5 Gm., Dihydrostreptomycin base 0.5 Gm./1 Gm. vial. Streptomycin 2.5 Gms., dihydrostreptomycin base 0.5/5 Gm. vial. Soln. 1 Gm in 2½ cc. vial, 5 Gm. in 12½ cc. vial.
- DINYDROTACHYSTEROL, U.S.P., N.N.D. 1962;** Caps, Soln, U.S.P. 9,10-Seco-5,7,22-ergostan-3 betaol. (A.I. 10) Sol. (oil) 0.5% in veg. oil (C. Bowman) See: Hy-lakeral, Cap. & Sol (Winthrop Labs)
Tachysterol. (No Mfr. currently lists.)
- DIHYDROTOSTERONE.** Androstane-17-beta-ol-3-one. See: Stenolone.
- DIHYDROTHEELIN, N.N.D. 1962.**
See: Estradiol (Various Mfr.)
- DIHYDROXYACETONE.** See: Chromelin (Summers Labs)
- DIHYDROXY ALUMINUM AMINOACETATE, N.F., N.N.D. 1962,** Magma, N.F.; Tab, N.F. [Aluminum Dihydroxyaminoacetate, Basic Aluminum Aminoacetate]
See: Antacid.
See: Ada-Tabs, Tab. (G. F. Harvey)
Alglyn, Tab. N.N.R. (Broyten).
Alkam, Tab. (First Texas).
Alminaf, Tab. (Bristol)
Aspogen, Tab., N.N.R. (Flint, Eaton)
Dimothyn, Tab. (Flint, Eaton).
Doraxamin, Susp. & Tab., N.N.R. (Smith-Dorsey).
Rabalate, Tab., N.N.R. (Eobins)
- W/Belladonna alkaloids** See: Belglyn, Tab Broyten).
- W/Belladonna alkaloids & phenobarbital.** See: Molygn Compound, Tab. (Broyten).
- W/Carboxymethylcellulose Sod. & glycine.** See: Acichak, Tab. (Valentine).
- W/Gastric mucin & ethaverine HCl.** See: Mugastral, Tab. (G. F. Harvey)
- W/Homatropine methylbromide & phenobarbital.**
See: Alkam Compound, Tab. (First Texas)
Alziox, Magma, (Patch)
Homoldran, Tab. (Flint-Eaton)
- W/Nyascyamine sulfate, atropine sulfate, hyasine HBr, phenobarbital, kaolin & pectin.** See: Donnagel, Susp. (Eobins).
- W/Mag. carb.** See: Talmin, Tab. (Jackson-Mitchell)
- W/Methscopolamine bromide, sod. lauryl sulfate & magnesum hydraxide** See: Alu-Scap, Cap, Susp. (Lloyd, Dabney & Westerfeld)
- W/Phenobarbital.** See: Barbanate, Tab. (Bristol).
- W/Phenobarbital & atropine methyl nitrate.** See: Narvtrate A, Tab. (G. F. Harvey)
- W/Phenobarbital, homatropine methylbromide & Vit. C.** See: Dalzinate, Tab. (Patch)
- W/Prednisolone.** See: Pred Kam, Tabs. (First Texas).
- W/Roserpine, phenobarbital & glytheonate.** See: Glytheonate W/Phenobarbital & Roserpine, Tab. (Patch)
- W/Sod. salicylate, Sod. p-aminobenzoate & ascorbic acid.** See: Elpagen, Tab. (Patch)
- DINYDROXY ALUMINUM SODIUM CARBONATE.** See: Kaloids, Tab. (American Chicle).
- 1,8-DIHYDROXYANTHRANOL.** See: Anthrolin, N.F.
- 1,8-DIHYDROXYANTHRAQUINONE.** Dioxanthraquinone.
See: Danthron, N.F.
Dionone, Tab. (Klug)
Dorbane, Tab. (Schenley)
Daxidan, Cap. (Lloyd Bras)
Istixia, Prep. (Winthrop Labs)
- 2,5-DINYDROXYREHZOATE SODIUM.** Gentic acid. See: Gentic Acid Sodium. (Various Mfr.)
- 3,12-DIHYDROXYCHOLANIC ACID.** See: Desoxycholic acid.
- DINYDROXYESTRIN, N.N.D. 1962.**
See: Estradiol (Various Mfr.)
- DIHYDROXYFLUORANE.** See: Fluorescein. (Merck Sharp & Dahme)
- DIHYDROXYPHENYLISATIN.** See: Oxyphe-nisatin. For related products. See: Diacetylhydroxyphenylisatin
- DINYDROXYPHENYLOXINDOL.** See: Oxyphe-nisatin.

DIHYDROXYPROPYLTHEOPHYLLINE.

Dyphylline NND 1959 See Neophylline, Supp (Paul Maney)

W/Ethoverine HCl See Coravar Tab (Crystal)

W/Glycerine See Coralixin, Elix (Crystal)

W/Pentylentetrazol nicalinic acid See Pentorack Tab (Crystal)

DIHYPRYLONE See Sedulon, Syr (Roche)

3,5-DIIODO-4-HYDROXYPHENYLALANINE See: Di Ioda Tyrosine, Tab (Organon)

*DIIODOHYDROXYQUIN, USP NND 1962 Tab USP; Di iodo hydroxyquinoline

USP XIV (Embequin, Enterosept) 57 Di ioda 8 quinoline

See Diiodoquin NNR (Searle)

Iaquin Susp, (Abbott)

Lanadoxin, Tab (Lanett)

Maebiquin, Tab (Consol Midland)

Quinodone Creme (Dome)

Seboquin (Summers)

W/Hydrocortisone alcohol Zetar See Zetar Oin (Dermik)

Yodoxin, NNR (Glenwood Labs)

W/Hydrocortisone alcohol See: Vytone (Dermik)

W/Hydrocortisone alcohol, coal tar soln in a watermiscible base See Cor D Tar (Deslin)

W/Phenylmercuric acetate, sodium lauryl sulfate alum, papain See Baculin, Vag Pow & Tob (Amfra)

W/Silbestrol sulfodiazine, tartaric acid boric acid, etc See Gyneben vag (Bentex Phr)

W/Surfactants See: Lycinale Supp (Lloyd)

DIODOHYDROXYQUINDLINE, NND 1939 See Diiodo hydroxyquin Prep

5,7-DIIDO-8-QUINDLINDL. See: Diiodo hydroxyquin USP

DIIDDOTYROSINE. (Organon) (Iodogargoric Acid) 3.5 diiodo 4 hydroxyphenylamine

Tab 1½ gr Tube 20s, 100s

Use: (1-3 daily) Iodine therapy

DIISOBUTYL CRESOXY ETHOXY ETHYL DI-METHYL BENZYL AMMONIUM CHLORIDE (Hyamine 10 X) W/Alkyl dimethyl

benzyl ammonium chloride and di isobutyl phenoxy ethoxy ethyl dimethyl benzyl ammonium chloride See: Merphene Amp (Barry Labs)

DIISOBUTYL PHENDXY ETHOXY ETHYL DIMETHYL BENZYL AMMONIUM CHLORIDE W/Alkyl dimethyl benzyl ammonium

chloride and di isobutyl cresoxy ethoxy ethyl di methyl benzyl ammonium chloride

See: Merphene, Amp (Barry Labs)

DI-ISDPACIN. (CMC) Aminosalicylic acid

* Generic name recognized by N.F., U.S.P., or A.M.A.

buffered W/MgCO₃ and Mg glycinate in niazid 25 mg/Tab Bot 500s 1000s

Use Treatment of tuberculosis

*DIISOPROMINE HCl. N N diisopropyl 3,3-diphenylpropylamine HCl (Lab for Pharmaceutical Development, Inc)

DIISOPROPYL FLUOROPHOS. See: Floropryl, Sol (Merck Sharp & Dohme)

DIISOPROPYL FLUOROPHOSPHATE, USP NNR (DFP, Isofluraphale) See Floropryl (Merck Sharp & Dohme)

DILABIL-SODIUM INJECTION. (Brean) Sterile aq sol of Sod dehydrocholate 20% for IV use Amp 5 cc

Use: Management of biliary dysfunction

DILABIL TABLETS (Brean) Each tab contains 0.25 Gm dehydrocholic acid (3,12 triketocholonic acid) Bot 100s

Use (1-3 daily & h s) Biliary stasis not due to obstruction

DILABRON (Brean) Isoetharine

Use Bronchodilator

DILAC. (Ascher) Troinitrate phosphate 4 mg theophylline 300 mg, butalbarbital sodium 60 mg/Tab Bot 100s

Use Treatment of angina pectoris

DILAMINATE Mixture of mag carb and dihydroxy aluminum glycinate

DILANTIN (Parke, Davis) Diphenylhydantoin delayed action Kapseal 1½ gr Bot 100s

1000s Infatab ¾ gr Bot 100s In oil, Cap 1½ gr Bot 100s 300s Susp. (100 mg/4 cc) Bot 8 oz

Use (m p) Anticonvulsant in epilepsy

W/Phenobarbital, desoxyephedrine HCl See: Phelantin, Cap (Parke, Davis)

W/Phenobarbital & desoxyephedrine HCl See Phelantin, Kapseals (Parke Davis)

DILANTIN SODIUM, NND 1959 (Parke Davis) Sod 5,5 diphenylhydantoinate Kapseal ½ gr 1½ gr Bot 100s, 1000s Pow Bot 1 oz Susp 100 mg/4 cc Bot 8 oz Vial (250 mg) 5 cc

Use Antiepileptic

DILANTIN SODIUM W/PHENOBARBITAL KAPSEALS. (Parke, Davis) Dilantin Sod 1½ gr phenobarbital ¼ gr or ½ gr/ Cap Bot 100s, 1000s

Use: Anticonvulsant w/sedative

DE-LA-SOL. (Direct Labs) Germicidal detergent Bot 1 pt, 1 qt, 1 gal

Use: Anti pruritic & fungicide

DILAUDID COUGH SYRUP. (Knoll) Dilaudid HCl 1 mg, glyceryl guaiacolate 100 mg/5 cc Bot 16 oz

Use Harassing and nonproductive cough

DILAUDID HYDROCHLORIDE, NNR 1954 (Knoll) Dihydramorphinone HCl, USP

Amp.: (1/32 gr. & 1/20 gr.) 1 cc., Box 6s, 100s.

Comp. Tab.: (1/2 gr.) Bot. 8s, 40s

Hypodermic Tab.: (1/64, 1/48, 1/32, 1/20, 1/16 gr. each) Tube 20s; Bot. 100s, 500s.

Oral Tab.: (1/24 gr.) Tube 10s; Bot. 100s, 500s.

Pow.: Bot. 15 gr., 1/2 oz., 1/2 oz.

Rectal Supp.: (1/25 gr.) Box 6s, 100s.

Sterile Soln.: (1/32 gr./cc) Vial 10 cc.

Use: Analgesic.

W/Benzocaine. See: Hibtane, Loz. (Ayerst)

W/Glyceryl guaiacolate. See: Dilaudid Cough Syrup. (Knoll)

W/Sod. citrate, anilmony Pot. tartrate & chloroform. See: Dilocal, Syr. (Tablerock)

DILAUID SULFATE, (Knoll) Dihydromorphine sulfate. Vial, (2 mg./cc.) 10 cc.

DILCORON, (Winthrop Labs.) Glyceryl trinitrate 0.4 mg., pentaerythritol tetranitrate 15 mg./T.D. Tab. Bot. 100s.

Use: Angina pectoris.

DELOCAL, (Tablerock) Dilaudid HCl 1/10 gr., Sod. citrate 20 gr., anilmony Pot. tartrate 1/10 gr., chloroform 1 gr./fl. oz. Bot. 1 pt., 1 gal.

Use: (1 tsp. q. 3-4 h., p.r.n.) Cough depressant.

DILDERM, (Schering), Dichlorisone acetate and aluminum chlorhydraxyl allantoinate, 1.75 mg. each/10 Gm. foam aerosol. Dichlorisone, 8.3 mg. with propellant /50 Gm. aerosol spray, Dichlorisone, 2.5 mg./Gm. and 0.25% alum. chlorhydraxyl allantoinate/5 Gm. cream, Aerosol foam, 10 Gm. can; Aerosol, 50 Gm. can; Cream, 5 Gm., 15 Gm.

Use: Dermatitis

DILYN, (Elder) Methphenoxydial 100 mg./Tab. Bot. 60s, 100s

Use: Orally, treatment of tension states

DIMACID GEL, (Ollis Clapp) Mixture of aluminum hydroxide gel and magnesium magma hydroxide gel.

Use: Temporary relief of gastric hyperacidity.

DIMAZON, (Heilkratt) Diacetylaminoozotoluen 2%. Oint. Tube 1 oz., Jar 1/2 oz., 1/2 oz., 1 lb.

Use: Antiseptic skin therapy.

DIMENFORMON, N.N.D. 1961. (Organon) Estradiol Tabs, 1/5 mg & 1/2 mg. Bot 60s & 250s

Use: Estrogen therapy

DIMENFORMON BRNZOATE, N.N.D. 1962 (Organon) Estradiol benzoate. In oil, Amp. containing 0.166, 0.33, 1 or 1.66 mg./1 cc. Vial 10 cc. containing 0.33 or 2 mg./cc.

***DIMENYDRINATE**, U.S.P., N.N.D. 1962;

* Generic name recognized by N.F., U.S.P. or A.M.A.

Enj, Syr, Tab., U.S.P. beta-Dimethylaminoethyl benzohydryl ether 8-chlorotheophyllinate, combination of benodryl & 8-chlorotheophyllinate, dromain & dramyl. See: Dramamine (Searle)

W/d-Amphetamine sulf. See: Dramamine-D, Tab. (G. D. Searle)

***DEMIPROZAN HCl**, (Lederle Lab) 2-Methoxy-N,N dimethyl-d⁹,o-xanthenepropylamine hydrochloride.

Use: Tranquilizing agent.

***DIMERCAPROL**, U.S.P., N.N.D. 1962, A.D.R. Inj., U.S.P. See: BAL Amp. (Hynson, Westcott & Dunning)

2,3-DIMERCAPTOPROPANOL. See: Dimercaprol, U.S.P.

DE-ME-SAL, (Superior Pharm) Dipyrone 175 mg salicylamide 175 mg./Tab Bot 100s, 500s.

Use: Analgesic and antipyretic

DEMETANE, N.N.D. 1962, (Robins) *Brompheniramine maleate, parabromdylamine maleate, 1-(p-bromophenyl)-1-(2-pyridyl)-3-dimethylaminopropane maleate. Tabs. (4 mg.) Bot 100s, 500s Extend tabs. (12 mg) Bot. 100s. Elixir (2 mg /5 cc) Bot 1 pt., 1 gal. Amps. (10 mg /cc.) 1 cc. Box 6s; Vial (100 mg./cc) 2 cc.

Use: Antihistamine

W/Phenylephrine hydrochloride, phenylpropanolamine hydrochloride. See: Dimetapp Extentabs (Robins)

DEMETANE EXPECTORANT, (Robins) Brompheniramine maleate 2 mg., glyceryl guaiacolate 100 mg., phenylephrine HCl 5 mg., phenylpropanolamine HCl 5 mg./5 cc. Bot. pts.

Use: Anti-allergic.

DIMETANE EXPECTORANT-DC, (Robins) Same as Dimetane Expectorant plus codeine phosphate 10 mg /5 cc.

DIMETANE-TEN INJECTABLE, (Robins) Brompheniramine maleate 10 mg /cc, Amp 1 cc., 6s.

Use: Antihistaminic.

DIMETANE-100 INJECTABLE, (Robins) Brompheniramine maleate 100 mg /cc. Vial 2 cc.

Use: Antihistaminic

DIMETAPP ELIXIR, (Robins) Brompheniramine maleate 4 mg., phenylephrine HCl 3 mg., phenylpropanolamine HCl 5 mg./5 cc. Bot, pt., gal

Use: Antihistaminic, nasal decongestant

DIMETAPP EXTENTABS, (Robins) Brompheniramine maleate 12 mg., phenylephrine HCl 15 mg., phenylpropanolamine HCl 15 mg./Tab Bot. 100s.

Use: Antihistaminic.

DI-MET-BP, (Organon) Estradiol benzoate 1

- mg, testosterone propionate 20 mg/cc
Vial 10 cc Box 1 6s
Use: 1 M; estrogen androgen combination
- ***DIMETHADIONE** (Abbott) 5,5 Dimethyl
oxazolidine 2,4 dione
Use Anticonvulsant
- ***DIMETHAZAN** 1,3 Dimethyl 7 (2 dimethyl
aminoethyl)xanthine See Eldin (Endo)
- ***DIMETHICONE**, NND 1962 Dimethyl
siloxane polymers dimethyl polysiloxane
See Silicone Oil (Various Mfr)
- ***DIMETHINDENE MALEATE** (Ciba) 2 {1 [2
(2 dimethylamino ethyl)nden 3 yl]ethyl}
- ***DIMETHISOQUIN HYDROCHLORIDE**, N.F.,
NND 1962 Oint. Lot NF 3 Butyl 1 (2
dimethylaminoethoxy)isoquinoline HCl See
Quatane Hydrochloride Oint. & Lot
(Smith, Kline & French)
- ***DIMETHOLIZINE PHOSPHATE**, 1 (2 Methyl
oxyphenyl) 4 (3 methoxypropyl)piperazine
phosphate See Ansyl (Abbott Lab)
- DIMETHONE** (Columbus) Dipyrone 50%
aqueous sol Vial 10 cc, 30 cc
Use Analgesic
- ***DIMETHOXANATE HCl**, NND 1962
beta Dimethylaminoethoxyethyl phenolthia
zine 10 carboxylate HCl See Cothera Syr
W/Isolhipendyl HCl, phenylephrine HCl,
acetylaminophen, ammonium chloride so
dium citrate chloroform See Cothera
Compound Syr (Ayerst)
- DIMETHOXYPHENYL PENICILLIN SODIUM**
Methicillin Sodium
See Dimacillin Vial (Squibb)
Staphicillin Vial (Bristol)
- ***DIMETHYPYRINDENE MALEATE** 2 {1 [2 (2
dimethylaminoethyl) 3 indenyl] ethyl} pyr
ridine maleate See Forhistal
Use Antihistaminic
- DIMETHYLAMINOPHENAZONE**. See:
Aminopyrine (Various Mfr)
- DIMETHYLANE**, NND 1962 (National
Drug) Pramoxolane (Dimethylin) 2,2 Diiso
propyl 4 methanal 1,3 dioxalane Tab D 25
Gm Bot 100x 1000x
Use Orally (3,4,1 d) muscle relaxant
- DIMETHYLCARBAMATE** of 3 hydroxy 1
methylpyridinium bromide See Mestinon,
Tab (Rache)
- DIMETHYLHEXESTROL DIPROPIONATE**,
NND 1962 Promethestrol Dipropionate
See: Meprane Dipropionate Tab (Reed &
Carrick)
- DIMETHYL POLYSILOXANE** See Dimethi
cone
- ***DIMETHYL TUBOCURARINE CHLORIDE**,
NND 1962 See: Mecastin Chloride, Vial
(Squibb)
- * Generic name recognized by NF USP or AMA
- ***DIMETHYL-TUBOCURARINE IODIDE**, NF
NND 1962; Inj., NF See Melubine
Iodide, Vial (Lilly)
- 1,3-DIMETHYLXANTHINE**. See: Theophyl
line, Prep (Various Mfr)
- 3,7-DIMETHYLXANTHINE**. See: Theabra
mine, Prep (Various Mfr)
- DIMETNYLN**. See: Dimethylone, Tab (Na
tional Drug)
- DIMINIC TABLETS**. (Durst) Pyrimidine mole
ote 25 mg, methapyrilene HCl 25 mg/
Tab Bot 100s, 1000s
Use: (1-3 daily) Antihistamine therapy
- DIMOCILLIN** (Squibb) Sodium dimethoxy
phenyl penicillin 1 Gm, equivalent to 900
mg of 2,6 dimethoxyphenyl penicillin/
Vials 1 Gm
Use Staphylococcal infections
- DIMORLIN TARTRATE**. (Smith, Kline &
French) Dextromoramide d 1 (3 methyl-4
morpholino 2,2 diphenylbutyryl)pyrrolidine
tartrate Under Study
Use Analgesic
- DIMOTHYN**, NND 1962 (Flint Eaton)
Dihydroxy aluminum aminoacetate Tab
7½ gr Bot 100s, 500s, 1000s
Use (½ t i d) Antacid
- DINDEVAN** See Phenindione
- DINACRIN**, NND 1960
- DI-O-CHROME** (Warren Teed) Sod di
chromate & boric acid Bot 3 fl oz Sol
Bot 4 fl oz, 1 pt
Use Antiseptic
- ***DIOCTYL CALCIUM SULFOSUCCINATE**
Bis 2 ethylhexyl calcium sulfosuccinate
Calcium bis dioctyl sulfosuccinate See
Surlak (Lloyd Bros)
- DIOCTYL POT SULFOSUCCINATE**, W/Gly
cerin, pol oleate and stearate See Re
ctalad Liq (Wampole)
- W/Sorbitol See Sigmal Enema Sol (Pharma
seal)
- ***DIOCTYL SODIUM SULFOSUCCINATE**,
NND 1962, NF ADR Aerastal OT (Am
Cyanamid)
See Calace, Cap, Liq (Mead Johnson)
Constanate, Cap (Nysca Labs)
Diactlyn (Bruce)
Diamedicane Tab (Medicane,
Dioxuccin Cap [Consolidated Midland
Corp)
Diovac, Preps (Gray Pharmacol Co)
Di-Sosal (Drug Ind Co)
Doxinate, Cap, Liq (Lloyd Bros)
Doxol, Cap (Blair Labs)
Duozal, Cap (Kirkman)
Kantax, Tab (Buffington)
Kosate, Cap (Haack)
Laxinate, Tab (Mallord)

- Molofac, Cap.** (Squibb)
Regutal Tab. (Pharmaco)
Revoe Suppelles, Supp. (Webster Co)
Velmal, Cap. (Potch)
- W/Acetphenolisalin.** See: **Evasol, Tabs.** (Lemnan)
W/Acetylphenolisatin, thiomine HCl, citrus pectin cellulose complex. See: **Nion-Lax, Tab.** (Nion)
W/Bulkagen. See: **Bulkasol, Cap.** (Columbus)
W/Carboxymethylcellulose, sed. See: **Stuart Dialose, Cap.** (Stuart)
W/Calcium & sodium alginates. See: **Celginace, Tab. & Gron.** (Mead Johnson)
W/Calcium & sodium alginates, anthraquinone derivatives. See: **Combinoce, Tab. or Gron.** (Mead Johnson)
W/Casonthronal See: **Dicoxon, Tabs** (Chicago Pharm)
W/Cosmo acutifolia active principles. See: **Senokop, Cap.** (Purdue Frederick)
W/Ext. castora sagrada. See: **Diocetyl-Zem, Tab.** (Zemmer)
W/Cholic acid. See: **Aquatyl, Tab.** (Irwin, Neisler)
W/Danthron. (West-Word)
 See: **Diocetyl W/Danthron, Tab.** (West-Word)
Dorbantyl, Cap. (Schenley)
Doxon, Cap. (Lloyd Bros.)
Doxidon, Cap. (Lloyd Bros.)
Doxinate-Danthron, Cap. (Lloyd Bros.)
Doxitil, Tab (Crestmed)
- W/Dehydrocholic acid.** See: **Neolox, Tab.** (Central Pharm.)
W/Desoxycholic acid, dehydrocholic acid See: **Dexohyl, Tab.** (Amer)
W/Diacetyldihydroxyphenylisatin. See: **Doxal Forte, Tab.** (Blair Labs)
Noloc, Cap (Dumas Wilson)
- W/Ferrous fumarate** See: **Personalink, Tab.** (Person & Covey)
W/Hesperidin complex, vitamins and minerals. See: **Siatric, Tab.** (Smith-Osney)
W/Methylcellulose. See: **Hydralax, Tab** (Invenex)
W/Ox bile. See: **Bidrolar, Tab.** (Armour)
W/Peristim. See: **Peri-Colace, Cap., Syr** (Mead-Johnson)
W/Phenolphthalein. See: **Correctol, Tab.** (Pharmaco)
W/Scopolamine methylbromide, guar. gum. See: **Colactin, Cap.** (Hillard Labs)
Synacol, Cap. (Reed & Cornick)
- W/Sodium alkyl polyether sulfonate, sodium sulfacetate, Kerohydric, sulfur, salicylic acid, hexachlorophene** See: **Sebulex, Liq** (Westwood)
W/Sod carboxymethylcellulose, di-acetylhydroxyphenyl) isatin See: **Dialose Plus, Cap.** (Stuart)
- DIOCTYL-ZEM.** (Zemmer) Diocetyl sodium succinate 60 mg. ext. castora sagrada 4 mg / **Tab. Bot. 100s**
 Use: *Fecal softener and peristaltic stimulant*
- DIOCUBB.** (Tufag) d-Amphetamine suff. 5 mg / **Perle. Bot. 100s, 1000s**
 Use: *CNS stimulant*
- DI-ODINE TABLETS.** (Brawn Pharm) Potassium iodide & chlaramine-T (NF) w/other organic materials in tab. form. 22mg I₂ at pH 6 / **Tab. Bot 25 tabs** (foil wrapped)
 Use: *Disinfectant.*
- DIODONE INJECTION.** See: **Iodopyracel Injection, U.S.P.**
- DIOBOQUIN, N.N.D. 1962.** (Searle) (Diiodo-hydroxyquinoline) 5,7-Diiodo 8-quinolinal **Tab., Bot. 60s, 300s.**
 Use: (3 1/2 d., o.c. for 15 to 20 days) *Amebiasis & Trichomonas intestinalis infections*
- DIODRAST, N.N.D. 1962.** (Winthrop Labs) (Iodopyracel) 3,5-diiodo-4-pyridone-N-acetic acid diethanolamine Amp. 30 cc., 20 cc., 30 cc. (Each with) cc. sensitivity test amp.), **Box 1s, 10s, 25s.**
 Use: *I.V. (excretion) urography & retrograde pyelography.*
- DIODRAST COMPOUND SOLUTION, N.N.D. 1962.** (Winthrop Labs) (Iodopyracel Comp) Contains 40.5 per cent w/v of Diadrast. Amp. 20 cc., **Box 1s, 10s, 25s.**
 Use: *Same as Diadrast*
- DIODRAST CONCENTRATED SOLUTION 70%, N.N.D. 1962.** (Winthrop Labs) (Iodopyracel Conc) Amp. 20 cc., 50 cc. (each with 1 cc. sensitivity test amp) **Box 1s.**
 Use: *For cardiac & angiographic visualization; also may be used for cholangiography*
- DI-O-GEST.** (A. P. C.) Mag carb., Col. carb., colloidal kaolin, tricalcium phosphate, Sod bicarb., bismuth subcarb., papain, dextrose **liq, Bot. 6 fl oz., 1 pl. Pow., Tin 3 oz., 6 oz. Tab., Bot. 100s, 500s, 1000s.**
 Use: *Gastric hyperacidity.*
- DIOGYN, N.N.D. 1960** (Pfizer) Estradiol. Vial (0.25 mg/cc.) 1 cc., 10 cc; (10 mg/cc.) 1 cc., 10 cc. **Tab. (D) 125 mg., 0.25 mg., 1 mg.) Bot. 50s, 100s.**
 Use: *Estragen therapy.*
- DIOGYN-B, N.N.D. 1960.** (Pfizer) beta-Estradiol benzoate. Vial (0.33 mg/cc.) 10 cc; (1 mg/cc.) 10 cc.
 Use: *Estragen therapy.*
- DIOGYN-E, N.N.D. 1960.** (Pfizer) Ethinyl estradiol. **Tab. (0.02 & 0.05 mg.) Bot. 100s, 1000s; (0.5 mg.) Bot. 25s, 100s.**
 Use: *Estragen therapy*
- DIOGYNETS.** (Pfizer) Estradiol. **Tab 5 1 0.125 mg., D 25 mg., 1 mg Bot. 100s.**
 Use: *Estragen therapy.*

- DIOLANDRONE.** (Carnrick) Methylandrostenedial Aq. Susp., 25 mg /cc, 50 mg /cc, Vial 10 cc Hypoglossal 10 mg, 25 mg, Bot 30r, 100r Tab. 10 mg, 25 mg, Bot 30r 100r
Use Protein anabolic action of androgens w/minimal masculinizing effect
- DIOLOSTENE.** See Methylandrostenedial (Various Mfr)
- DILOXOL.** NND 1962 (Carnrick) (Mephenesin) Cap 0.5 Gm, 0.25 Gm, Bot 100r, 100r Elixir 0.1 Gm /cc, Bot 8 oz, 1 pt, 1 gal Tab 0.25 Gm, 0.5 Gm, Bot 100r, 1000r
Use Orally muscle relaxant
- DIOMICONE** (Medicone) Diocetyl sodium sulfosuccinate 50 mg /Tab Vials 50r
Use Fecal softener
- DIONEDRA COMPOUND.** (Central) Ethyl morphine HCl $\frac{1}{4}$ gr, Pat guaiacol sulfonate 4 gr, ephedrine HCl $\frac{1}{2}$ gr, cociflona tr 40 mg, wild lettuce syr 120 mg, quill ryr comp 24 min, euphorbia tr 120 min /fl oz Bot 1 pt, 1 gal
Use: (1 tsp q 3-4 h, prn) Cough preparation
- DIONICOL.** (Rorer) Ethylmorphine HCl (Dionin) $\frac{1}{4}$ gr chloralform 2 min, Pat, guaiacol sulfonate 8 gr, ammonium chloride 8 gr Sod citrate 20 gr, tartar emelic $\frac{1}{2}$ gr /fl oz Bot 1 pt
Use: (1 tsp q 3-4 h, prn) Sedative expectorant
- DIONIN.** (Merck Sharp & Dohme) Ethyl morphine HCl Pow, Bot 15 gr, $\frac{1}{2}$ oz, 1 oz
Use Orally cough depression, ocular lymphagogue
- DIONONE.** (Klug) 1,8 dihydroxyanthraquinone (danthron) 750 mg /Tab Bot 100s
Use Functional constipation
- DIONOSIL.** (Picher X-ray Corp) Pyridane N acetic acid n propyl ester 50% m susp Vial 20 cc
Use Bronchography
- DIOPHYLLIN** See Aminophylline Prep (Various Mfr)
- DIOPTERIN.** (Pteroylglutamic acid FDGA) Pteryl-alpha glutamylglutamic acid
Use Treatment of neoplastic diseases
- DIORAPIN** (Standex) Estrogenic conjugate 0.625 mg, methyl testosterone 5 mg /Tab Bot 100s
Use: Hormone therapy to promote anabolsim
- DIOSMIN.** (Barosmin) Buchu resin obtained from lvs of barosma serratifolia & allied rutaceae
- DIOSTATE D.** (Upjohn) Dicalcium phosphate
* Generic name recognized by N.F., U.S.P. or A.M.A.
- 5 gr, purified bone phosphate $2\frac{1}{2}$ gr, Vit O 777 u, w/traces of iron, copper, fluorine zinc & Mang /Tab Bot 100s, 1000s
Use: Calcium w/Vit D & mineral therapy
- DIOSTATE O FERROUS.** (Upjohn) Ferrous sulfate 1 gr, dicalcium phosphate 5 gr purified bone phosphates $2\frac{1}{2}$ gr, Vit D 333 u w/traces of copper, fluorine zinc & Mang /Tab Bot 100s 1000r
Use Calcium, Vit D & mineral therapy
- DIOSUCCIN.** (Consolidated Midland Corp) Diocetyl sod sulfosuccinate 20 mg /Cap Bot 30s, 100r 250r, 1000r
Use Fecal softener
- DIOTERPINE** (Shielfelin) Ethylmorphine HCl 4 mg, terpin hydrate 2 gr /derrertsp Bot 1 pt, 1 gal
Use (1 tsp q 3-4 h m p) Sedative expectorant
- DIOTNANE HYDROCHLORIDE, N N R 1954.** (Merrell) Dipiperdon HCl Cream 1%, Tube 1 oz, Jar 1 lb Sol 1%, Bot 60 cc, 500 cc
Use Local anesthetic therapy
- DIOTHANE OINTMENT W/Oxyquinoline Benzolate, N N R 1954** (Merrell) Contains 1% dipiperdon & 0.1% oxyquinoline benzolate in suitable oil base Tube 1 oz
Use Topical onerthia & antiseptic
- DIOTNOID SUPPOSITORIES.** (Merrell) Dipiperdon (Diothane brand) 1%, Isobutyl p-aminobenzoate 1%, urea 10%, ephedrine HCl 4 mg (1/16 gr), oxyquinoline benzolate 1 1000 Box 12s
Use Hemorrhoidal suppositories, anesthetic & antireptic
- DIOVAC CAPSULES.** (Gray Pharmacol Co) Diocetyl sod sulfosuccinate 50 mg /cap Bot 50s
Use To promote normal bowel evacuation
- DIOVAC SYRUP.** (Gray Pharmacol Co) Diocetyl sodium sulfosuccinate 20 mg /5 cc Bot 6 oz Pediatric Syr. (10 mg /5 cc) Bot 4 oz
Use To promote normal bowel evacuation
- DIOVOCYLIN** See Etradiol, Prep (Various Mfr)
- * **DIOXADROL HYDROCHLORIDE.** (Cutter) 2 (2,2-diphenyl 1,3-dioxolan 4-yl) piperidine hydrochloride See Rydar
Use Antidepressant
- DIOXINDOL** See Diacetylhydroxyphenyl natin
- DIOXYANTHRAQUINONE, 1-8.** See Danthron NF (Various Mfr)
- DIOXYANTHRANOL** See Anthralin, NF (Various Mfr)
- DIOXYANTHRAQUINONE.** See 1,8 Dihydroxyanthraquinone
- DIOXYLAMINE SUCCINATE W/Pyrimamine**

maleate, pheniramine maleate. See: Tridacemine, Tab. (Merrill)

* **DIOXYLINE PHOSPHATE**, 6,7 Dimethoxy-1-[4'-ethoxy-3'-methoxybenzyl]-2-methylisoquinoline. See: Poveril Phosphate, Tab. (Lilly)

DIPAMYCIN, (Mead-John) Vit. A, D and Inositol; in a urine repellent base / Tube 1 oz., carton 12s; jars 1 lb.

Use: Dermatologic, pediatric

DI-PARALINE HYDROCHLORIDE, NND 1962. (Abbott) [Chlorcyclizine HCl] 1-p-chlorobenzhydryl 4-methylpiperazine HCl, Tab. 25 mg, 50 mg Bot 100s, 500s, 1000s. Elix-tabs 25 & 50 mg Bot 100s

W/Celomine 2% cream.

Use: Antihistaminic drug of low toxicity. W/Tranethane HCl, colomine, zinc oxide & menthol. See: Tranolan, Lat. (Abbott)

DIPARCOL, (May & Baker) [Diethazine HCl, Casontia] 10-[2-Diethylaminoethyl]-phenothiazine.

Use: Parasympatholytic & sympatholytic. Parkinson's disease.

DIPASIC, (Ponray) Isoniazid & p-aminosalicylic acid 100 mg

Use: Treatment of tuberculosis

DIPAXIN, NND 1962. (Upjohn) Diphenylacetylidandione Dipendione. 2 Diphenylacetyl-1,3 Indandione. Tab 1 mg, 5 mg. Bot. 100

Use: (1-2 1/d) Orally, anticoagulant.

DIPHOYL. See: Picotinicamide, Prep. (Vorlove Mfr)

DIPENICILLIN G. See: Rensethacil.

DIPERODON HYDROCHLORIDE, NMR. 1954. See: Diathane Hydrochloride, Amp., Cream & Sol. (Merrell).

W/Alantoin neomycin sulfate See: Ro Derm (Lemman)

W/Cetalkonium Chloride, pyrilamine maleate, hydrocortisone, neomycin sulfate, urea. See: Tega Otic, Soln. (Ortega)

W/Furacin. See: Furacin Inserts (Eaton)

W/Hydrocortisone Alcohol. See: Pentacort. (Dalin)

W/Hydroxyquinoline Benzoate, NNR. 1954. See: Diathane W/Hydroxyquinoline Benzoate aint. (Merrell).

W/Neomycin sulfate, hydrocortisone, pyrilamine maleate, cetalkonium chloride. See: Fent O Augine Ear Drops, Liq. (Confield)

W/Neomycin sulf., hydrocortisone, pyrilamine maleate, cetalkonium chloride, urea. See: Olotrin Ear Drops (Lauisons)

Tyler, Otc, Sol. (Tyler)

W/Nitrofurazone. See: Furacin Urethral Supp. (Eaton).

* Generic name recognized by N.F., U.S.P. or A.M.A.

W/Promoxine HCl. See: Nasculo, Creme (Celacomb)

W/Zincum oxide, bithional, tyrothricin, neomycin sulf., phenyltoloxamine dihydrogen citrate, zirconium oxide. See: Allergenic, Lat. (Bryant)

DIPHASOL SOLUTION TESTOSTERONE.

(Kramers-Urbani) Synthetic testosterone 25 mg or 50 mg/cc. Vial 10 cc.

Use: 1 M₂ male hormone therapy.

* **DIPHEMANIL METHYLSULFATE**, NF, NND. 1962. Inj., Tab., NF, N,N dimethyl-4-piperidylidene-1, 1-diphenylmethane methylsulfate

See: Advelin (Schering)

Prantal Methylsulfate, Tab., Vial (Schering)

W/Undecylenic acid, salicylic acid. See: Advicia, Cream, Pow (Schering)

DIPHENIN. See: Hydroxydiphenine

* **DIPHENADIONE**, NF, NND. 1962. Tab., NF. See: Dipoxin, Tab. (Upjohn)

DIPHENAN, (Burroughs Wellcome) Not listed, p-amino-Benzylphenylcarbamate, Water D.5 Gm Tube 20s; Rot 100t.

Use: (1 s.i.d. for 7 days) Anthelmintic for pinworm infestation.

DIPHENATIL. See: Diphemonil methylsulfate

* **DIPHENIDOL**, (Smith, Kline & French) o,o-diphenyl-1-piperidinebutanol.

Use: Antiemetic.

* **DIPHENHYDRAMINE HYDROCHLORIDE**, U.S.P., NND 1962, A.D.R. Inj., Cop. U.S.P., Elixir U.S.P., 2 (Benzhydroxy)-N,N-dimethylethylamine HCl. See: Benadryl Hydrochloride, Prep. (Parke, Davis)

W/Aminophylline. See: Hydryllin, Elixs & Tab (Searle).

W/Aminophylline & rocephadrine. See: Hydrodrine, Tab. (Searle).

W/Aminophylline, Pot. Iodide, chlorofom, xugol & alcohol. See: Hydryllin Compound, Liq. (Searle).

W/Aminophylline rocephadrine HCl. See: Hydryllin W/Rocephadrine HCl, Tab. (Searle)

W/3 Chlorotheophylline. See: Dramamine (Searle)

DIPHENMETHANIL METHYLSULFATE. See: Prantal, Tab. & Vial (Schering).

* **DIPHINOXYLATE HCl**, 2,2 Diphenyl 4-(4-carboxy-4-phenyl-1-piperidino) butyranitrile ethyl ester HCl. See: Lomohil (Searle)

DIPHENTOLIN, (Massengill) Diphenylhydantoin Sod. Cop. 1½ gr. Rot. 100s, 1000s

Use: Orally; epilepsy therapy.

DIPHENTLACETYLDINDANDIONE. See: Dipaxin, Tab. (Upjohn)

DIPHENYLAN SODIUM, (Lannett) Diphenyl-

- hydantoin sodium $\frac{1}{2}$ gr & $1\frac{1}{2}$ gr Cap
 Bot 100s, 1000s
 Use Epilepsy
- DIPHENYLCARBINOL.** See: Benzhydrol
- DIPHENYLHYDANTOIN, USP Oral Susp**
 USP See Dihycon Caps (Consolidated
 Midland)
- W/Mebaral See Mebaroin, Tab (Winthrop
 Labs)
- W/Mephobarbital See: Medylon (G F
 Harvey)
- ***DIPHENYLHYDANTOIN SODIUM, USP,**
 NND 1959, Cap; USP Alepsin, Dihydan
 soluble, Diphenoin, Silantin Sod, Epanutin
 Eptain, Hydantal, Phenytoin soluble, Solan
 tain, Solantyl, Denyl Sod, Soluble Phenytoin
 (3,5 diphenyl hydantoinate, Sod)
 A.P.C.—Cap $1\frac{1}{2}$ gr, Bot 100s, 1000s
 Prema—Cap $\frac{1}{2}$ gr, $1\frac{1}{2}$ gr, Bot 50s,
 100s, 1000s; Paw, Bot 1 oz, $\frac{1}{4}$ lb, 1 lb
 Use: Anticonvulsant
 See: Denyl sod (Prema)
 Dianlin Sod (Parke, Davis)
 Diphenoin (Mossengill)
 Diphenyl Sod, Cap (Lanett)
- W/Mephobarbital See Medylon, Tab (G. F
 Harvey)
- W/Phenobarbital See Elmoloin, Cap (Elder)
- DIPHENYLISATIN.** See Diacetylhydroxy
 phenylisatin
- ***DIPHENYLPYRALINE NCI, NND 1962 4**
 Diphenyl methoxy 1 methylpiperidine HCl
 See: Cupertin (Tinsley)
 Diafen, Tab (Schenley)
 Hispril, Sponsule & Tab (Smith Kline &
 French)
- W/Chlophedianol HCl phenylephrine HCl
 glyceryl guaiacolate, alcohol See Ultramic
 (Riker)
- W/Chlophedianol HCl, phenylephrine HCl,
 glyceryl guaiacolate, acetaminaphen See:
 Ulagesic, Tabs (Riker)
- DI-PHOCAL** (Texas Pharm Co) Dicalcium
 phosphate 12.5 gr, Vit D 250 u/isp Bot
 & oz (in veg oil)
 Use: Calcium therapy for infants
- DIPHODYNE**, (Hisi) Chloroquine diphos
 phate 125 mg, salicylamide 300 mg, as
 corbic acid 50 mg/Tab Bot 100s, 500s
 1000s
 Use: Rheumatoid arthritis
- DIPHOSPHOPYRIDINE (DPH)**—Treating al
 coholism Under study
- DIPHOSPHOTHIAMIN** (Cocarboxylase)
 See: Coenzyme B, Cap & Inj (Barrows)
- ***DIPHOXAZIDE, N¹-[p hydroxy b b di
 phenylpropionyl] N² acetylhydrazine in
 ductin**
- DIPHTHERIA, (Parke, Davis) Diphtheria**
 * Generic name recognized by N F, U S P or A.M.A.
- toxoid & pertussis vaccine, aluminum phos
 phate adsorbed Vial 7.5 cc
 Use (I.M. 0.5 cc q 4-6 weeks for 3 doses)
 Active Immunization.
- DIPHTHERIA ANATOXIN.** See: Diphtheria
 Toxoid, U S P
- ***DIPHTHERIA ANTITOXIN, U S P**
 Use I.M., S.C.; protection & treatment of
 diphtheria
- DIPHTHERIA, PERTUSSIS & TETANUS.**
 See: Dip Pert Tet (Cutter)
 Inlagen (Pitman Moore)
 Tridipigen (Lilly)
 Trinovac (Merck Sharp & Dahme)
 Triogen (Parke, Davis)
- DIPHTHERIA & TETANUS TOXOIDS, N F X
 N N R 1955**
- DIPHTHERIA & TETANUS TOXOIDS, ALU
 MINUM HYDROXIDE ADSORBED, U S P,
 N N R 1955** See: Diphtheria & Tetanus
 Toxoids, Alhydrox (Cutter)
- DIPHTHERIA & TETANUS TOXOIDS, ALUM
 INUM PRECIPITATED, U S P, N N R 1955**
- DIPHTHERIA & TETANUS TOXOIDS COM
 BINED, ALUMINUM HYDROXIDE AD
 SORBED.** See: Diphtheria & Tetanus Tox
 oids, Aluminum Hydroxide Adsorbed, U S P
- DIPHTHERIA & TETANUS TOXOIDS COM
 BINED, ALUM PRECIPITATED.** See: Dipht
 heria & Tetanus Toxoids, Alum Precipi
 tated, U S P
- DIPHTHERIA & TETANUS TOXOIDS &
 PERTUSSIS VACCINE COMBINED, N N R
 1955**
 See: Dip Pert Tet, Plain (Cutter)
 Tridipigen, Fluid (Lilly)
- DIPHTHERIA & TETANUS TOXOIDS &
 PERTUSSIS VACCINE COMBINED, ALU
 MINUM HYDROXIDE ADSORBED, U S P,
 N N R 1955**
 See: Dip Pert Tet, Alhydrox (Cutter)
 Triogen (Parke, Davis)
- DIPHTHERIA & TETANUS TOXOIDS &
 PERTUSSIS VACCINE COMBINED, ALU
 MINUM PHOSPHATE-ADSORBED, U S P**
 See: Tri Immunol, Vial (Lederle)
- DIPHTHERIA & TETANUS TOXOIDS &
 PERTUSSIS VACCINE COMBINED, ALUM
 PRECIPITATED, U S P, N N R 1955**
 See: Inlagen, Vial (Pitman Moore).
 Tridipigen, Alum Precipitated, Vial (Lilly).
 Trinovac, Vial (Merck Sharp & Dahme)
 Triple Antigen, Vial (Wyeth)
- DIPHTHERIA & TETANUS TOXOIDS &
 PERTUSSIS & POLIO VACCINES, U S P**
 See: Quadrigen, Vial (Parke, Davis)
 Tetravox, Vial (Merck Sharp & Dahme)
- DIPHTHERIA YOXIN, DIAGNOSTIC, U S P**
 Schick Test

- DIPHTHERIA TOXIN, DIAGNOSTIC, INACTIVATED**, U.S.P. Schick Test Control.
- DIPHTHERIA TOXIN, INACTIVATED FOR DIAGNOSTIC TEST CONTROL**, U.S.P. See: Diphtheria Toxin, Diagnostic, Inactivated.
- DIPHTHERIA TOXIN FOR SCHICK TEST**. See: Diphtheria Toxin, Diagnostic, U.S.P.
- ***DIPHTHERIA TOXOID, ADSORBED**, U.S.P., N.N.D. 1962. Alum Precipitated (Anatoxin-Roman, Diphtheria Anatoxin).
- ***DIPHTHERIA TOXOID, ALHYDROX**, N.N.D. 1962. (Cutter) Viols 1 & 5 cc.
- DIPHTHERIA TOXOID, ALUMINUM HYDROXIDE ADSORBED**, U.S.P., N.N.D. 1962. See: Diphtheria Toxoid, Alhydrox. (Cutter) W/Pertussis vaccine. See: Diftussis Alhydrox, Vial (Cutter).
- DIPHTHERIA TOXOID PHOSPHATE ADSORBED**, N.N.D. 1962.
- DIPHTHERIA TOXOID & PERTUSSIS VACCINE**, Lilly—Vial 10 tests. See: Diph-Pertussis (Parke, Davis). Diftussis (Cutter).
- DIPHTHERIA TOXOID & PERTUSSIS VACCINE COMBINED**, N.F. X, N.N.R. 1955. See: Diftussis (Cutter).
- DIPHTHERIA TOXOID & PERTUSSIS VACCINE COMBINED, ALUMINUM HYDROXIDE ADSORBED**, U.S.P., N.N.R. 1955. See: Diftussis, Alhydrox (Cutter).
- DIPHTHERIA TOXOID & PERTUSSIS VACCINE COMBINED, ALUM PRECIPITATED**, N.N.R. 1955. National Drug—Vial 0.5 cc., 2.5 cc. Pilman-Moore—Vial 4.5 cc.
- DIP-PERT-TET**, N.N.R. 1955. (Cutter) Diphtheria & tetanus toxoids w/pertussis vaccine, combined. Vial 1.5, 7.5 & 22.5 cc. Use: (Hypodermic 0.5 cc. q. 3 or 4 weeks for 3 doses) Active immunization.
- DIP-PERT-TET, ALHYDROX**, N.N.R. 1955. (Cutter) Diphtheria & tetanus toxoids w/pertussis vaccine combined, aluminum hydroxide adsorbed. Vial 1.5 & 7.5 cc. Use: (Hypodermic 0.5 cc. q. 3 or 4 weeks for 3 doses) Active immunization.
- DI-PODII**. (Superior Pharm) Dipyron 5 gr./Tab. Bot. 100s, 500s. Use: Analgesic & antipyretic.
- DIPRAL**. (Haskell) Dipyron 62.5 mg., salicylamide 192 mg., PABA 192 mg., vit. C 20 mg., vit. B-12 2.5 mcg./Tab. Bot. 100s. Use: Analgesic.
- DIPRALON FORTE**. (Anar-Stone) N Acetyl-p-aminophenol 125 mg., dipyron 125 mg., salicylamide 125 mg., Vit. C 50 mg./Tab. Bot. 100s, 1000s. Use: Analgesic in the treatment of arthritis
- * Generic name recognized by N.F., U.S.P. or A.M.A.
- DI-PRO**. (Organon) beta-Estradiol benzoate 2.5 mg., progesterone 12.5 mg./cc. Vial 10 cc. Box 1s, 6s. Use: Estrogen-androgen therapy.
- DIPROFARN**. (Farnsworth Lab.) Dipyron, 500 mg./Tab. Bot. 100s. Amp. (500 mg./cc.) 5 cc. Use: Analgesic.
- DIPRONE**. (Myers-Carter) Dipyron 50%/cc. Vial 30 cc.
- ***DIPROTRIZOATE SODIUM**, U.S.P., N.N.D. 1962. See: Miokon, Amp. (Mollinckrodt)
- DIPSAL**. (U. S. Standard Products) Dipyron 62.5 mg., salicylamide 180.0 mg., p-aminobenzoic acid 180.0 mg., Vit. C 20.0 mg., B-12 2.5 mcg./Tab. Bot. 100s, 1000s. Use: Rheumatoid arthritis.
- DIP-TET, ADULT**, (Cutter) Diphtheria and tetanus toxin for immunization. Viols 1 cc., 5 cc.
- DIFTUSSIS**, N.N.R. 1955. (Cutter) Diphtheria Toxoid & Pertussis Vaccine Combined. Plain Vial, 1½ cc., 7½ cc. Alhydrox Vial, 1.5 cc., 7.5 cc. Use: Immunization.
- DIFTUSSIS, ALHYDROX**, N.N.R. 1955. (Cutter) Diphtheria Toxoid and Pertussis Vaccine Combined, Alum Hydroxide Adsorbed. Vial, 1.5 cc., 7.5 cc. Use: Dose 3 x 0.5 cc. Inj. active immunization.
- DIPTILIDIUM CANIUM ANTIDEN**. (Lederle) Dog tapeworm ext. Vial, 10 test & 10 controls. Use: I.D. diagnostic aid for hydatid disease in man.
- ***DIPTRIDAMOLE**. See: Personin, Tabs., Amps (Gaigy)
- DIPYRONE**, Sod. phenyl[2-methylpyrazolon-methylamino-methane sulfonate, novalgin, methylmalubrin, methampyrone, 4-Sod methanesulfonate methylamine-anilpyrine. See Barone Inj. (Barre Drug Co) Dimethone, Vial. (Columbi) Di Podii, Tab. (Superior Pharm) Dipraf, Tab. (Haskell) Diprofern (Farnsworth) Dipron, Vial. (Myers-Carter) Dpron (Minn. Pharm) Favonil, Liq. (Cortone Lab) Nyprone. (New Mexico Pharm) Key-Pyrone, Tabs. and liq. (Key Pharmacol) Methampyrone. Vial (500 mg./cc.) 30 cc. (Testagar) Narone, Tab. & Vial. (Uimer) Nartate, Tab. & Vial. (G. F. Harvey) Novaldin, Amp. & Tab. (Winthrop Labs.) Pydron, Amp. (Braun). Pyratgin Ped. Drops. (Savage)

Viperone, Vials (Vitamix Corp)

Use: Analgesic, antipyretic & antirheumatic
W/n acetyl p aminophenol, diphenine HCl
phenyltolaxamine citrate See Anfac, Caps
(Pharmafas)

W/N Acetyl p aminophenol, salicylamide
Vit C See Dipralan Forte, Tab (Amar
Stone)

W/p Aminobenzoic acid, salicylamide & Vit
B 12 See Cobasal, Tab (Rand Pharm)

W/Glucuronolactone & salicylamide See
Glu Solgin, Cap (Warren Tead)

W/Phenobarbital, aspirin, phenacetin See
Axafar, Cap (Warren Tead)

W/Salicylamide See Di Me Sal Tab (Su
perior Pharm)

W/Salicylamide, p aminobenzoic acid, me
phenesin, vit C, B 1, niacinamide &
menadiolone See Mybasal, Tab (Lloyd
Dabney & Westerfield)

W/Salicylamide, p aminobenzoic acid, Vit
C, B 12 See Dipal, Tab (U S Standard
Products) Dipral, Tab (Haskell)

W/Salicylamide, di desoxyephedrine HCl
See Detamide Tab (Mallard Inc)

DIRECT SKY BLUE 4% INJ (Wyeth) Sod
Dimethoxydiphenyl diaza bis 8 amine 1
naphthal-5,7 disulfonate Vials 5 cc

Use: Vital staining of regional lymphatics
during surgery

DIRIDONE, (Prama) Phenylaza diamino
pyridine HCl 100 mg/Tab Bot 100s
100s Available W/Trisulfa Bot 30s, 100s,
100s

Use: Urinary anesthetic and antiseptic

D' ISATIN, See: Diacetylhydroxyphenylisatin
DIPAL, HND 1962 (Riker) Orphenadine
HCl Tab (30 mg) Bot 100s, 500s

Use: Parkinsonism

DISODIUM AUROTHIOMALATE, NNR
1935 See, Myochrysin, Amp (Merck Sharp
& Dohme)

***DISODIUM EDATHAMIL**, USP See
Edathamil Disodium

DISODIUM HYDROGEN CITRATE, See:
Citralka, liq (Parke, Davis)

DISODIUM PHOSPHATE, See: Sodium Phos-
phate, USP.

DI-SODIUM SUCCINATE HERAhydrate
See: Soduxin, Amp (Brewer)

DI-SODIUM VERSEHATE, See: Edathamil
Disodium

DISOMER, NND 1962 (White s) Dextrom
pheniramine maleate; each tablet, 2 mg/
100s; Chronotabs 4 mg/100s, 6 mg/100s;
syrup 2 mg/5 cc, Bot 1 pt

Use: Antihistaminic.

DISOHATE, (Rannett) Diactyl sod. sullosuc-
* Generic name recognized by NF, USP or A.M.A.

cinato 60 mg & 100 mg /Cap Bot 100
1000s

Use Fecal softener

DISOPHRIN, (White Tabs) Disomer (dex
bromphenisamine maleate) 2 mg d iso-
ephrine sulfate, 60 mg/Tab Bot 100s

Use Nasal Congestion

***DISOPYRAMIDE**, (G D Searle) o (2 di
isopropylaminoethyl) o phenyl 2 pyridine-
acetamide

DISPADAL, (Squibb) Vit A 5000 u D 1000
u, thiamine HCl 1 mg, riboflavin 0.5 mg,
niacinamide 10 mg, ascorbic acid 50 mg/
D 6 cc Bot w/dropper 20 cc, 50 cc

Use Multivitamin supplement

DISPATABS TABS, (Person and Covey)
Vitamin A palmitate 50,000 USP u/Tab
Bot 100s 1000s

Use Vit A deficiency

DISPERDON (Standard Pharmacal) Acetyl-
carbromal 2 gr, hyoscyamine sulfate 0.1
mg, atropine sulfate 0.02 mg, hyoscin
HBr 0.006 mg/Tab Bot 50s, 100s

Use (1 2 1 d) Sedative w/antispasmodic

DISPERTAR, (Texas Pharm Co) Coal tar
1%, polyoxyethylene sorbitan monooleate
0.5% in zinc oxide paste Jars 5 oz & 1 lb

DI-SPOR, (Wintersmith Chemical) Copper
undecylenate 1%, zinc undecylenate 20%
undecylenic acid 2%, phenylmercuric ace-
tate 1:3000 talc base 76.97% Oint 1 oz,
8 oz Jar 1 lb Paw 1 oz, 1 lb

Use Fungicide

DISTAQUAINE, See: Penicillin V

DISTAROL, (Manhattan Eye Salve) Distilled
coal tar 10%, wool fat 30%, petrolatum
60%, color & perfume 1 lb, 5 lb

Use Antiseptic

DISTAROL & CALAMINE OINTMENT, (Man-
hattan Eye Salve) Distilled coal tar 10%
calamine 20%, salicylic acid 2%, lanolin
30%, petrolatum 38% 1 lb, 5 lb

Use Antiseptic

DI-STERIOD IH DIPHASOL, (Kremers Urban)
Estrone 2 mg, testosterone propionate 10
mg/cc oil Vial 10 cc, Box 1s, 6s, 12s, 25s

Use: Estrogen & androgen therapy

DI-STERIOD SUSPENSION (Kremers Urban)
Estrone 2 mg, testosterone 10 mg/cc Vial
10 cc, Box 1s, 6s, 12s, 25s; Vial (Estrone
5 mg, testosterone 50 mg/cc) 10 cc

Use: Estrogen & androgen therapy

DISTREPTOCIN SULFATE, (Lilly) Strepto-
decin, streptomycin base 0.5 Gm, dihydro-
streptomycin base 0.5 Gm as sulfate salt
Amp 1 Gm base, 5 cc; 5 Gm base, 20 cc

Use: I.M. streptomycin therapy

DISTRYCIN (Squibb) Streptoduoctin, strep-
tomycin sulfate & dihydrostreptomycin sul

tate, equal parts. Viol 0.5 Gm., I Gm., 5 Gm. [amounts represent free base].

Use: Tuberculosis.

DISULAX. (Crestmed) Dioctyl sodium sulfacrylate U.S.P. 100 mg., acetophenolisathm 2 mg./Tab. Bot. 100s.

Use: Laxative.

DISULFADINE. (Kremers-Urbain) Sulfadiazine 4 gr., sulthiozole 4 gr./Tab. Bot. 100s, 1000s.

Use: Orally, sulfonamide therapy.

* **DISULFIRAM, N.N.D. 1962.** (Tetraethylthiuram Disulfide)

See: Alcophobin, Tab. (Consol. Midland Corp)

Antobuse, Tab. (Ayerst)

Nozol, Liq. (Nozol)

DISULFONAMIDE.

See: Dia-Mer-Sulfonamide

DISULFYN, N.N.R. 1955. (Rorer) Susp. Sulfadiazine, sulfamerazine of each 50 mg./cc. Bot. 2 oz., 1 pl. Tab. Sulfadiazine, sulfamerazine of each 0.25 Gm./Tab. Bot. 100s, 1000s.

Use: Disulfonamide therapy.

DI-SUP. (Taylor) Ferrus gluconate 3 gr., cal. lactate 5 gr., vit. A 824 u., D 300 u., B-1 0.34 mg., B-2 0.4 mg., C 5 mg., E 0.34 I.U., niacinamide 4 mg., cobalt chloride 0.2 mg./Tab. Bot 100s, 1000s

Use: (1-2 I.U.) Dietary supplement.

DI-THELIN. (Bowman) Theelin 2 mg. potassium theelin sulfate 1 mg./10 cc. Vials 10 cc.

Use: Deficiency of estrogens.

* **DITHIAZANINE IODIDE, U.S.P., N.N.D. 1962.** Tabs, U.S.P. 3,3' Diethylthiodicarbocyanine iodide 3-ethyl-2-[5-(3-ethyl-2-benzothiazolylidene)-1,3-pentadienyl] benzothiazolium iodide

See: Abminthic, Tab. (Pfizer)

Delvez, (Lilly)

DITHIRAZOL. (Mossengill) Sulfamerazine 0.15 Gm., sulfadiazine 0.175 Gm., sulthiozole 0.175 Gm./Tab. Bot. 100s, 500s, 1000s

DITHRANOL. See: Anthrolin, N.F. (Abbott)

DITRAN. (Lakeside) N-ethyl-3 piperidyl cyclopentylphenylglycolate HC. Under study. Use Treatment of depression.

DITUBIN, N.N.D. 1960. (Schering) (isoniazid, Isonicotinic Hydrazide) Tab. 50 mg. Bot. 100s, 1000s.

Use: (1 q.i.d., m.p.) Antituberculosis therapy.

DIUCARDYN SODIUM. (Ayerst) Mercaptopurine Sod. Vial 1.4 Gm. with 10 cc. diluent. 4.2 Gm. without diluent.

Use: Diuretic.

* Generic name recognized by N.F., U.S.P. or A.M.A.

DIUMERIN. (C. D. Smith) Mercurophylline 280 mg., theophylline 20 mg./2 cc. Amp. 2 cc. Box 5s, 25s, 100s.

Use: Diuretic.

DIUPRES. (Merck Sharp & Dohme) Diuril 250 mg. or 500 mg., reserpine 0.125 mg./Tab. Bot. 100s, 1000s.

Use: Hypotensive.

DIURALETS. (Estro) Phenobarbital ¼ gr., Cal. lactate 1½ gr., theobromine Sod. salicylate 3 gr./Tab. Bot. 1000s.

Use: Hypertension.

DIURBITAL. (Amira-Grant) Theobromine Sod. salicylate 3 gr., Cal. lactate 1½ gr., phenobarbital ¼ gr./E.C. Tab. Bot. 100s, 500s, 1000s

Use: (1-3 daily) Diuretic.

DIURETICS.

See: Acetazolamide.

Ademol, Tab. (Squibb).

Aldactone, Tab (Searle)

Amisametrodine.

Benzhydroflumethiazide

Carbonic Anhydrase Inhibitors

Cordrase, Tab. (Upjohn).

Daranide, Tab. (Merck Sharp & Dohme)

Diamox, Tab. (Lederle)

Dichlorphenamide.

Esidrix (Ciba)

Ethexolamide

Flumethiazide.

Hydrochlorothiazide.

Hydrodiuril (Merck Sharp & Dohme).

Hydroflumethiazide.

Methazolamide.

Mercurial diuretics

NoClex, Tab (Robins)

Naturein (Squibb)

Neptozene, Tab. (Lederle)

Oretic, Tab. (Abbott).

Rolicton, Tab. (Searle).

Saluron (Bristol)

Spironolactone.

DIURETIC Inj. (Ortega) Mersalyl, USP, 100 mg; theophylline USP, 50 mg./cc. Vial 10 cc

Use: Diuresis, especially in cardiac edema

DIURIL, N.N.D. 1962. (Merck Sharp & Dohme)

Chlorothiazide, 6-chloro-7-sulfamyl-1, 2, 4-benzothiazodiazine-1, 1-dioxide Tab. 0.5 Gm., D 25 Gm. Bot. 100s, 1000s. Also vials 20 cc

"LYOVAC" N.N.D. 1960. [0.5 Gm] Vial 20 cc

Use: Diuretic therapy.

W/Reserpine. See: Diupres, Tab. (Merck Sharp & Dohme)

DIURIL SYRUP. (Merck Sharp & Dohme)

- Chlorothiazide 0.25 Gm / 5 cc Bot 8 oz
Use Edema and hypertension
- DIURNAL, NND 1962 (Upjohn)** Procaine penicillin G 300M u/cc Vial 5 & 10 doses Fortified, Pot penicillin G 100M u added Vial 1, 5, 6, 10 doses w/disposable syringe, 300M & 600M u/cc 1 cc
Use: I M; penicillin therapy
- DIURNAL-PENICILLIN, 3DD M., NND 1962 (Upjohn)** Procaine penicillin G 300 M u/cc Vial 5 & 10 doses
Use: I M.
- DIURNAL-PENICILLIN, FORTIFIED, 4DD M., NND 1962 (Upjohn)** Procaine penicillin G 300M u & Sod penicillin G 100M u/cc Vial 1, 5, 10 doses
Use: I M.
- DIURNAL-PENICILLIN, FORTIFIED, 5DD M., NND 1962 (Upjohn)** Procaine penicillin G 300M u & Pot penicillin G 200M u/cc Vial 1, 5, 10 doses
Use: I M.
- DIURNAL-PENICILLIN, READIMIXED, 300 M., NND 1962 (Upjohn)** Procaine penicillin G 300M u/cc Available as 1 cc cart w/metal syr 5s & 25s; sterile disp. 1 cc cart syri sterile disp 1 cc plasket syr; Vial 10 cc
Use: I M.
- DIURNAL-PENICILLIN, READIMIXED, 500 M., NND 1962 (Upjohn)** Procaine penicillin G 500 M u/cc Vial 10 cc
Use: I M.
- DIURNAL-PENICILLIN, READIMIXED, 6DD M., NND 1962 (Upjohn)** Procaine penicillin G 600M u/cc Sterile disp 1 cc plasket syr
Use: I M.
- DIURNAL-PENICILLIN, READIMIXED, 1000 M., NND 1962 (Upjohn)** Procaine penicillin G 1000M u/17 cc Sterile disp 17 cc plasket syr
Use: I M.
- DIUROBES (Biber)** Allylisabutylbarbituric acid 30 mg, methamphetamine sulfate 5 mg, carbamide 500 mg/Tab Bot 100s, 1000s
Use: Treatment at obesity
- DIURODAN (Knoll)** Theocalcin 7 1/4 gr, Pot sulfocyanate 1 1/2 gr/Tab Tube 20s
Use: Hypertension
- DIURSAL (G F. Harvey)** Mersalyl 100 mg, theophylline 50 mg/cc, Amp 2 cc, Box 12s, 25s, 100s
Use: Diuretic
- DIUTENSEN (Irwin, Neisler)** Cryptenamine (as the tannate salts) 2 mg, methyldiothiazide 2.5 mg/Tab Bot 100s
Use: Hypertension
- DIUTENSEN-R. (Irwin, Neisler)** Cryptenamine (as the tannate salts) 1 mg, methyldiothiazide 2.5 mg, reserpine 0.1 mg/Tab Bot 100s
Use: Hypertension
- DIVINYL OXIDE (Vinyl Ether, Divinyl Ether)** N N R 1957, A D R See: Vinethene (Merck Sharp & Dahme)
Use: Inhalation anesthetic
- DIVIRON (Mollard)** Dicalcium phosphate 5 gr, ironate 2 gr, Vit D 334 u, B 1 1 mg, B 6 0.5 mg, folic acid 0.25 mg, w/minerals/Tab Bot 100s, 500s, 1000s
Use: Dietary supplement
- DMCT. (Lederle)** Demethylchlartetracycline See Declomycin, Caps (Lederle)
- DOAK-OIL (Doak)** Tar distillate 2%, in a lanolin, mineral oil suspension Pl Bot 8 oz
Use: Dermatologic
- DORELL'S SOLUTION, A D R See: Sodium Borate Solution Compound (Various Mfr)**
- DOCA ACETATE, NND 1962 (Organon)** Desoxycorticosterone acetate Amp (5 mg/cc) 1 cc, Box 6s, 50s Vial (5 mg/cc) 10 cc, Box 1s, 6s
Use: Adrenocortical insufficiency
- DOCHEMA (Ives Cameron)** Ferrous sulfate essic. 129 mg, folic acid 0.5 mg, Vit B 12 4 mcg, ascorbic acid 50 mg, insoluble liver fraction 192 mg/Cap Bot 100s
Use: Treatment of most anemias
- DOCIBIN (Walker Labs)** Vit B 12 Drop (1 mcg/drop) Bot w/dropper 15 cc, 60 cc, 1 pl
Inj: (30 mcg/cc) Vial 5 cc
Tab: (10 mcg) Bot 25s, 100s, 1000s; (25 mcg) Bot 100s, 1000s; Saluble (10, 25 & 50 mcg) Bot 100s, 1000s; (1000 mcg) Box 10s
Use: Orally I M; hematopoietic
- DOCRASERPINE (Docra)** Reserpine alkaloid 0.25 mg/Tab Bot 100s 1000s
Use: Tranquihizer
- DODECABEE, NND 1962 (Miller)** Vit B 12 cryst Vial (100 mcg/cc, 1000 mcg/cc) 10 cc
Use (m p) Pernicious anemia
- DODECAVITE (U S Vitamin)** Vit B 12 Amp: (15 mcg/cc) 1 cc Box 6s, 25s 100s Vial 5 & 10 cc
T S drop (30 mcg/cc) Bot w/dropper 15 cc & 60 cc
Sol. (30 mcg/cc) Bot w/dropper 15 & 60 cc
Tab (25 mcg) Bot 30s 100s
Vial (50 100 1000 mcg/cc) 5 cc 10 cc
Use: Orally, I M; hematopoietic
- DODEKROID (Riker Labs)** Vit B 12 10 mcg w/thyroid substance 10 mg/Tab Bot 50s
Use: Orally (1 3 daily); growth failure

DODERCIL. (Wynlit) Doderlein bacilli (viable) lyophilized 200,000 min./Vial 5 cc.

Use: Vaginitis.

* **DODERLEIN BACILLI**. See: Dodercil (Wynlit)

DODEX, N.N.D. 1962. (Organon) Vit. B-12. Amp. (15 mcg./cc.) 1 cc., Box 6s.

Vial: (30, 100, 1000 mcg./cc.) 5 cc

Tab.: W/activator (2 mcg. Vit. B-12, 200 mg. pyloric substance) Bot. 100s, 1000s
w/activator forte (5 mcg. Vit. B-12, 500 mg. pyloric substance, 1 mg. folic acid) Bot. 100s, 1000s.

Drop: W/ferrous gluconate (Vit. B-12 10 mcg., ferrous gluconate 218 mg./cc.) Bot. w/dropper 15 cc.

Use: Hematopoietic.

DODEX ADD DROP. (Organon) Vit. A 5000 u., D 1000 u., B-1 1 mg., B-2 0.25 mg., nicotinamide 10 mg., pantothenic acid 2 mg., Vit. B-6 1 mg./O 6 cc. Bot. 15 cc.

Use: Multivitamin therapy.

D-O-E. (Bran) *d*-Desoxyephedrine HCl. Elixir 2.5 mg./5 cc. Bot. 4 oz., 1 pt., 1 gal. Tab. 2.5 mg. Bot. 100s, 250s, 1000s.

Use: Orally; central nervous system stimulant.

DOFOCYTE-M. (Smith, Miller & Patch) Liver injection 10 mcg, folic acid 5 mg., Vit. B-12 50 mcg./cc. Vial 10 cc. Box 6s, 25s, 50s

Use: Parenteral; anemic therapy.

DOFOPAN. (Smith, Miller & Patch) Vit. B-12 25 mcg, folic acid 5 mg, liver inf 2 mcg./cc. Vial 10 cc., Box 1s, 6s, 25s

Use: Parenteral; anemic therapy

DOFORAL. (Smith, Miller & Patch) Vit. B-1 2 mg, B-2 2 mg., C 50 mg, B-6 1 mg, B-12 10 mcg, folic acid 1.2 mg., liver 120 mg., ferrous sulfate exsic. 200 mg., niacinamide 18 mg, Cal. pantothenate 2 mg, copper sulfate 4 mg./Tab. Bot 100s, 500s, 1000s

Use: Hematinic.

DOLAMIN. (Harvey Lab.) Ammonium sulf. 0.75% with sod. chlor. and benzyl etc. Amp. 10 cc.

Use: Various neuralgias

DOLANTIN. See: Meperidine Hydrochloride, U.S.P. (Winthrop Labs.)

DOLATE. (Crestmed) *N*-acetyl-*p*-aminophenol 5 gr., phenobarbital 1/4 gr., atropine sulfate 1/400 gr./Tab Bot. 100s.

Use: Pain, fever, sedation

DOLCIN. (Dalcin Corp.) Aspirin 3.7 gr., Cal. succinate 2.8 gr./Tab. Bot. 100s, 200s, 500s.

Use: (1-2 q.i.d.) Analgesic, antipyretic.

DOLICHOS BIFLORUS. See: *Anti-A₂ Techn*

(Purified), (Myland).

* Generic name recognized by N.F., U.S.P. or A.M.A.

DOLITRONE. (Merrell) MRD-125, 5-ethyl-6-phenyl-meta-thiazone-2,4-dione.

Use: Systemic anesthetic & analgesic.

DOLODYNE. (Domsco) Phenylephrine HCl 4 mg., chlorpheniramine maleate 2 mg., d-methorphan HBr 7.5 mg. (outer coat), aspirin 180 mg., phenacolin 100 mg., anhydrous caffeine 15 mg., phenylephrine HCl 5 mg., pyrilamine maleate 12 mg., d-methorphan HBr 7.5 mg., citrus bioflavonoid complex 25 mg., ascorbic acid 60 mg., phenylpropanolamine HCl 12 mg. (core)/Tab. Bot. 12s, 36s.

Use: Cold symptoms & cough control.

DOLOPAR. (Burlington) Paraminyl (Pyrilamine maleate) 1/2 gr., salicylamide 3 1/2 gr., acetophenetidin 2 1/2 gr., caffeine 1/4 gr./Tab. Bot. 100s, 1000s.

Use: Antihistaminic, analgesic & antipyretic.

DOLOPAR FORTE. (Burlington's) Pyrilamine maleate, 1/2 gr.; salicylamine, 3 gr.; acetophenetidin, 2 1/2 gr.; caffeine, 1/4 gr.; phenylephrine HCl, 5 mg./Tab. Bot. 100s, 1000s

Use: Relief of common cold, rhinitis, hay fever.

DOLOPHINE HYDROCHLORIDE, N.N.D. 1962. (Lilly) Methadone HCl. Amp. 10 mg., 1 cc. (10 mg./cc.) 20 cc. Syr. 10 mg./20 cc. 1 pt., 1 gal. Tab. 5 mg., 7.5 mg., 10 mg. Bot. 100s, 1000s

Use: Narcotic, analgesic & antitussive.

DOLORAL. (Wally Pharm.) Colchicine salicylate 0.1 mg., phenobarbital 8 mg., Sod. p-aminobenzoate 15 mg., Vial B-1 25 mg., aspirin 225 mg.-Tab. Bot. 100s, 500s.

Use: 2 q.i.d.; treatment of gout, arthritis & rheumatism.

DOLORO COMPOUND. (Burlington) Acetophenetidin 2 1/2 gr., aspirin 3 1/2 gr., caffeine 1/4 gr., *N*-acetyl-*p*-aminophenol 1/2 gr./Tab. Bot. 100s, 1000s

Use: Orally, analgesic

DOLOSAL. See: Meperidine Hydrochloride (Demeral HCl) (Winthrop Labs.)

DOLOXTNE. (Lilly) Dextropropoxyphene hydrochloride 32 mg., 65 mg./Tab D-Comp: dextropropoxyphene hydrochloride 32 mg., acetophenetidin 162 mg., acetylsalicylic acid 227 mg., caffeine 32.4 mg./Tab.

Use: Analgesic.

DOLVANOL. See: Meperidine Hydrochloride (Demeral HCl) (Winthrop Labs.)

DOMAPAL. (Hoag) Amobarbital 30 mg., dextroamphetamine sulfate 5 mg./Tab Bot. 100s.

Use: Obesity control.

DOMEORO. (Dome Chemicals) Aluminum sulfate & Cal. acetate when added to

water gives therapeutic effect of Buraw's Sol. 1.20 Bot 4 oz
 Sol. 1.20 Bot 4 oz
 Oint.: 4 oz., 1 lb., 5 lb
 Pow.: 4 oz., 13 oz., 4 lb
 Pkg.: 2.2 Gm., 12s., 100s
 Tab.: Bot 12s., 100s., 500s., 1000s.; Tab
 offer., Bot 12s., 100s., 500s., 1000s
 Use: Externally; inflammatory skin conditions

DOMEBORO OTIC. (Dome) Acetic acid 2% in an aluminum acetate solution buffered to pH of 5.0 Plas Bot w/otic tip 2 oz
 Use: Otitis externa

DOMEFORM. (Dome) Domatorm has been changed to Domeform

DOME-PASTE BANDAGE. (Dome Chemicals) Zinc oxide, glycerine & gelatine bandage Pkg Resh salored 4" x 10 yd Impregnated gauze bandage

Use: Treatment of leg ulcers

DOMERINE MEDICATED SHAMPOO (Dome) Salicylic acid 2%, allantoin 0.2%, Bithional 1%, incorporated in vehicle combining surfactant wetting agents with protein condensate (pH 5.3) Bot 4 oz., Pls

Use: Sebarrheic conditions of the scalp

DOMESTROL. See: Diethylstilbestrol, Prep (Various Mfr)

DOM F. See: Merbromin (Mercurochrome)

(Myron, Westcott & Dunning)

DOMICONE. (Dome Chemicals) Silicone (Dimethylpolysiloxane) Cream 20%, Tube 1 oz., Jar 1 lb

Use: Skin protective

DOMINE.

See: Pruraine, Oint (Rand)

DOMIPHEN BROMIDE Dodecyl dimethyl (7 phenosethyl) ammonium bromide See: Brodosol Bromide (Ciba)

W/Benzosone See Brodosol, las (Ciba)

DOMOFORM. (Dome Chemicals) Iodochlorhydroxyquin 3%, cream in Acid Mantle base, oint in Domolene base Tube 1 oz., Jar 1 lb

Use: Eczematous skin conditions

DOMOFORM-HC. (Dome Chemicals) Hydrocortisone alcohol 5 mg/Gm., Iodochlorhydroxyquin 3%, in Acid Mantle vehicle Cream tube 1/2 oz., 1 oz.; Jar Bot 1/2 oz.

Use: Treatment of skin infections

DOMOFORM-HC FORTE. (Dome Chemicals) Hydrocortisone alcohol 10 mg/Gm., Iodochlorhydroxyquin 3% Acid Mantle vehicle Cream, tube 1/2 oz., 1 oz.; Jar Bot 1/2 oz.

Use: Eczema

DOMOGYN (Dome Chemicals) Buffered acid douche pH 4.2 Pkg 4 Gm., 12s., 30s

Use: Vag acid douche cleanser

DOMOL BATH EMOLLIENT. (Dome) Isopropyl myristate and isopropyl sebacate w/ non ionic emulsifiers and hypoallergenic agents, in a mineral oil base Bath Emollient containers 8 fl oz

Use: Scalp skin conditions

DOMOLENE-HC OINT. (Dome) Micronized hydrocortisone alcohol 1% in special bland emollient base Tube 1/2 oz., 1 oz., 2 oz

Use: Chronic eczematous dry skin

DONABARB (Elder) Phenobarbital 1/4 gr Belladonna ext 1/6 gr/Tab Bot 100s, 500s, 1000s

Use: Sedative and antispasmodic

DONASIDA TABLETS & ELIXIR. (Ives-Cameron) Each tsp contains Belladonna alkaloids 0.2 mg as a sulfated ext, phenobarbital 0.25 gr, Vit B 1) 5 mg, B 2) 2 mg, B 6) 0.5 mg, B 12) 5 mcg, niacinamide 10 mg, in sherry flavored wine base Each tab contains identical amounts of phenobarbital & B complex factors In addition 0.25 gr of specially powdered ext belladonna USP Elixir 1 pt Tab 100s

Use: Nervous tension & nutritional debility

DONCAL. (Lemar) Hyoscyamine sulfate 0.1037 mg, atropine sulfate 0.0194 mg, hyoscyne hydrobromide 0.0065 mg, phenobarbital 16.2 mg, glycine 0.15 Gm, calcium carbonate 0.35 Gm/tab Bot 100s

Use: Treatment of hypermotility & smooth muscle spasms

DONCARB. (Crestmed) Phenobarbital 8 mg hyoscyamine sulfate 0.1 mg, hyoscyne HBr 0.008 mg, atropine sulfate 0.02 mg, calcium carbonate 225 mg/Tab Bot 100s

Use: Antispasmodic

DONDRIEL. (Whitehall Labs) Caromphen ethanedisulfonate 10 mg, phenylephrine hydrochloride 5 mg, phenindamine tartrate 7.5 mg/Tab Bot 24s

Use: Antihistive

DONNA EXTENTABS (Robins) Hyoscyamine sulfate 0.3111 mg, atropine sulfate 0.0527 mg, hyoscyne HBr 0.0195 mg/tab Bot 100s, 500s

Use: Antispasmodic

DONNAFID Jr. (Jenkins Labs) Ephedrine HCl 1/16 gr., belladonna ext 1/20 gr., ethacrylamide 1/2 gr/tab Bot 100s, 1000s

Use: Coryza, rhinitis, asthmatic conditions

DONNAGEL. (Robins) Hyoscyamine sulfate 0.1037 mg, atropine sulfate 0.0194 mg, hyoscyne HBr 0.0065 mg, phenobarbital 16.2 mg, kaolin 6 Gm, pectin 142.8 mg, dihydroxy aluminum aminoacetate 0.5 Gm/oz Sosp, Bot 6 oz

Use: (2 dr stat & 1 dr p r n) Dorrkes

DONNAGEL W/NEOMYCIN (Robins) Neo-

* Generic name recognized by N.F., U.S.P. or A.M.A.

- mycin base 210 mg., others as in Donnagel/oz. Bot. 6 oz.
Use: Enteritis and bacterial causes of diarrhea
- DONNAGEL-PG.** (Robins) Po. opium 24 mg., kaolin 6 Gm., pectin 142.8 mg., hyoscyamine sulfate 0.1037 mg., atropine sulfate 0.0194 mg., hyoscyne HBr 0.0065 mg., phenobarbital 16.2 mg /30 cc. Bot. 6 oz.
Use: Control diarrhea.
- DONNAGESIC EXTENTABS.** (Robins) No. 1—Equiv. to one Donnatal Tab. plus codeine phosphate $\frac{1}{4}$ gr./tab. Bot. 30s.
No. 2—Equiv. to one Donnatal Tab. plus codeine phosphate $1\frac{1}{2}$ gr./tab. Bot. 30s.
Use: long-acting analgesic.
- DONNALATE.** (Robins) Dihydroxy aluminum aminoacetate 0.5 Gm., phenobarbital 8.1 mg., hyoscyamine sulfate 0.052 mg., atropine sulfate 0.01 mg., hyoscyne HBr 0.003 mg /Tab Bot. 100s, 500s.
Use: (1 l.l.d.) Antispasmodic & antacid.
- DONNATAL.** (Robins) Tab., Cap. or 5 cc. of Elixir contains hyoscyamine sulfate 0.1037 mg., atropine sulfate 0.0194 mg., hyoscyne HBr 0.0065 mg., phenobarbital 16.2 mg., Cap. & Tab. Bot. 100s, 500s, 1000s. Elixir 1 pt., 1 gal.
Use: Sedative & antispasmodic
W/Codeine phosphate. See: Donnagesic Extentabs. (Robins)
- DONNATAL EXTENTABS.** (Robins) Hyoscyamine sulfate 0.3111 mg., atropine sulfate 0.0362 mg., hyoscyne HBr 0.0195 mg., phenobarbital 48.6 mg /tab Bot 100s, 500s.
Use: Antispasmodic & sedative
- DONNATAL #2.** (Robins) Phenobarbital $\frac{1}{2}$ gr., hyoscyamine sulfate 0.1037 mg., atropine sulfate 0.0194 mg., hyoscyne HBr 0.0065 mg /Tab. Bot. 100s, 1000s.
Use: (1 or 2 t.l.d.) Visceral spasm.
- DONNATAL PLUS.** (Robins) Donnatal formula plus B-complex factors thiamine HCl 3 mg., riboflavin 2 mg., nicotinamide 10 mg., Cal. pantothenate 2 mg., pyridoxine 0.5 mg /Tab. or 5 cc. Bot. 100s, 1000s Elix. Bot. 1 pt., 1 gal
- DONNAZYME.** (Robins) Hyoscyamine sulf. 0.0518 mg., atropine sulf. 0.0097 mg., hyoscyne HBr 0.0033 mg., phenobarbital 8.1 mg., pepsin 150 mg in outer layer, and pancreatin 300 mg, bile salts 150 mg in core. Tab Bot. 100s, 500s
Use: Digestant therapy.
- DONORAL.** (Lemmon) Chloral hydrate 1.6 Gm., Hyoscyamine sulfate 0.8 mg., atropine sulfate 0.16 mg., hyoscyne hydrobromide 0.048 mg /fl. oz. Bot. 3 fl. oz., 1 pt., 1 gal.
Use: Management of spastic disorders
- DONPHEN.** (Lemmon) Phenobarbital 15 mg., hyoscyamine sulfate 0.1 mg., atropine sulfate 0.02 mg., hyoscyne hydrobromide 0.006 mg /Cap or Tab. Bot. 100s
Use: Treatment of motion sickness
- DOPBESE.** (Philadelphia Labs) Allylisabutylobarbituric acid 30 mg., methamphetamine sulfate 5 mg., urea 500 mg /Tab.
Use: For control of obesity.
- DOPHENCO.** (Blue Line) Opium pow. 1/20 gr., ipecac pow. 1/20 gr., acetophenetidin $1\frac{1}{2}$ gr., atropine sulf. 1/500 gr., acetylsalicylic acid 3 gr., camphor monobromated $\frac{1}{2}$ gr., caffeine $\frac{1}{2}$ gr./Cap. Bot. 100s, 1000s
Use: Analgesic antipyretic
- DORAL.** (Massengill) Vit. D. Perle 50,000 u. Bot. 100s, 500s.
Use: Vit. D therapy.
- DORANA.** (Ives-Cameron) Skin respiratory factor 2,000 u., shark liver oil 3%, phenylmercuric nitrate 1:10,000 in an emollient base/oz. Oint. Tube 1 oz., Suppos. pkg. 12s
Use: Rectally, regression of hemorrhoids, relief of associated pain, and pruritus ani
- DORANTAMIN.** N N R, 1951. (Smith-Dorsey) Pyrilamine maleate Tab 50 mg. Bot. 100s, 1000s
Use: (1-2 t.l.d.) Antihistamine therapy.
- DORAXAMIN.** N N D 1959 (Smith Dorsey) Dihydroxy aluminum aminoacetate, Gel 0.5 Gm./5 cc. Bot. 1 pt. Tab 0.5 Gm. Bot. 100s, 500s, 1000s.
Use: Gastric antacid.
- DORBANE.** (Shen Lab) Danthron, 1.8 Dihydroxyanthraquinone. Conlet 0.075 Gm. Tube 20s. Tab. 0.150 Gm. Bot. 100s.
Use: Constipation.
- DORBANE SUSPENSION.** (Shen Lab) Danthron 37.5 mg /5 cc. Bot. 1 pt.
Use: Laxative.
- DORBANTYL.** (Shen Lab) Diocetyl sod. sulfosuccinate 50 mg., danthron 25 mg /Cap Bot. 30s, 250s Susp 5 cc. Is equiv. to 1 Cap. Bot. 150 cc.
Use: Fecal softener and laxative
- DORBANTYL FORTE.** (Shen Lab) Diocetyl sod sulfosuccinate 100 mg., dorbane 50 mg /Cap. Bot 30s, 100s, 250s
Use: Fecal softener and laxative
- DORCOL.** (Smith Dorsey) Phenyltolerant d. hydrogen citrate 12.5 mg., w/emmonium chloride, ipecac fl. ext., Sod. citrate & menthol/5 cc. Bot. 1 pt., 1 gal.
Use: (1 t.p. q. 3-4 h., p.r.n.) Antihistamine cough prep
- DORCOSTRIN.** (Smith Dorsey) Desoxyaceticosterone acetate 5 mg., benzyl alcohol

- 3%, persic oil q s / 10 cc Vial 10 cc
Use: Adrenal cortical insufficiency
- DORESTRO** (Smith Dorsey) Estragenic substance
Aq susp: Vial (1 mg, 10,000 IU/cc) 10 cc rub cap; (2 mg, 20,000 IU/cc) 10 cc rub cap; (5 mg, 50,000 IU/cc) 10 cc rub cap
In Persic oil Amp (1 mg, 10,000 IU/cc) 1 cc Vial (1 mg, 10,000 IU/cc) 10 cc rub cap; (2 mg, 10,000 IU/cc) 10 cc rub cap
- DORICO SOLURLE**. See: Hexobarbital Soluble (Epsal Sodium) (Winthrop Labs)
- DORIDEN**, N N O 1962 (Ciba) Glutethimide alpha ethyl alpha phenyl-glutarimide Tab (0.125 Gm) Bot 100s Tab (0.25 & 0.5 Gm) Cap (0.5 Gm) Bot 100s 500s, 1000s
Use Sedative
- DORMETHAN**, N N D 1959 See: Dextromethorphan HBr
- DORMIN** (Dormin, Inc) Methapyrilene 25 mg / Cap Bot 12s, 36s, 72s
Use (12 hr) Antihistaminic & insomnio
- DORMIRAL**, See Phenobarbital, Prep (Various Mfr)
- DORMISON** (Schering) Methylparafynol Cap 250 mg, 500 mg Bot 100s Liq 250 mg / 5 cc Bot 1 pt
Use Hypnotic
- DORMITOL COMPOUND** (Ruffington) Phenobarbital 2 gr, strontium bromide 8 gr, alcohol 20% / fl oz Bot 1 pt, 1 gal
Use (13 hr) Sedative & hypnotic
- DORMONAL**, See: Barbitol, Prep (Various Mfr)
- DORNAVAC FOR INHALATION**, N N D 1962 (Merck Sharp & Dohme) Desoxyribonuclease, derived from beef pancreas Pancreatic dornase lyophilized 100,000 u / Vial Vial 2 cc w/diluent
Use: Inhalation, lysis of pus and reduction in viscosity
- DORNVAL** (Maltbie Labs) Amphetamine 100 mg, or 200 mg / Tab Bot 100s 500s
Use Anxiolytic and tension states
- DORNVAL 400** (Maltbie Labs) Amphetamine aminopyridine 400 mg / Tab Bot 100s 500s
Use Anxiolytic and tension
- DORODOL**. (Buffington) Diallylbarbituric acid ¼ gr, acetophenetidin 2½ gr acetylsalicylic acid 3½ gr, caffeine ¼ gr gel semium tr 1 min / Tab Bot 100s 1000s
Use: (14 daily) Sedative, antipyretic & analgesic
- DORODOL-DEX**. (Buffington) Diallylbarbituric acid ¼ gr, d desoxyephedrine HCl 2.5 mg, acetophenetidin 2½ gr aspirin 3½ gr, caffeine ¼ gr, gelsemium tr 1 min. / Tab Bot 100s
Use: 1 q 4 hr; menstrual tension & depression
- DORSACAIN**, N N D 1962 (Smith Dorsey) 2-Orethyl-aminoethyl 4-amino 3-butoxy benzoate NCl Sol 0.4% Bot ½ oz
Use: Ophthalmic anesthetic
- DORSALLIN**. (Smith Dorsey)
Aq susp: Vial (Procaine penicillin G 300M u / dose) 10 doses; Vial, Fortified (Procaine penicillin G 300M u, Pal penicillin G 100M u / dose) 10 dose
In sesame oils With 2% aluminum monostearate, Vial (Procaine penicillin G 300M u / dose) 10 dose
Paws (Pot penicillin G Ruf 1200M u) Bot 2 oz; w/Sulfadiazine, sulfamerazine & sulfamethazine 0.167 Gm each & Pal penicillin G 100M u / dr
Tab: (Pot penicillin G Buf 100M u) Bot 12s, 100s; Soluble, effor (Pot penicillin G 100M u) Bot 24s; Procaine penicillin G 100M u, Bot 24s; w/Triple sulfo (Pot penicillin G 100M u & sulfadiazine, sulfamerazine sulfamethazine, 0.11 Gm each) Bot 25s
- DORSALLIN NOSE DROPS**, (Smith Dorsey) Not listed Cryst Pot penicillin G 6700 u desoxyephedrine ½% Bot ½ oz
- DORSALLIN TROCHES WITH BENZOCAINE**, (Smith Dorsey) Cryst Pal penicillin G 5000 u, benzocaine 5 mg / Troche Bot 30s
- DORSALLIN TROCHES CHEWING**, (Smith Dorsey) Cryst Procaine penicillin G 10,000 u Bot 10s
- DORSAPHYLLIN**, N N D 1959 (Smith Dorsey) Elixir, Theophylline Sod glycinate 1 Gm, equiv to Theophylline 0.5 Gm / fl oz Bot 1 pt, 1 gal Supp, Theophylline Sod glycinate 0.78 Gm, equiv to Theophylline 0.39 Gm / Supp Rox 12s Tab, Theophylline Sod glycinate 0.325 Gm equiv to Theophylline 0.162 Gm / Tab Bot 100s
Use Diuretic coronary vasodilator
- DORSULFAS**, N N R. 1955 Not listed (Smith Dorsey) Susp, 33 mg each of sulfacetamide, sulfadiazine & sulfamerazine / cc Bot 1 pt Tab, 0.67 Gm each of sulfacetamide sulfadiazine & sulfamerazine / Tab Bot 100s 1000s
Use: Orally triple sulfonamide therapy
- DORTYL** (Merck Sharp & Dohme) (Carbachol Carbamylcholine Chloride) Amp 0.25 mg T cc Tab 2 mg Bot 25s
Use: Orally, S C; parasympathetic agent
- DOUCHE POWDER**. (Lanfene) Chloralhycol

menthol, methylsalicylate, cossia oil, lactose & lactic acid. Pow. 5 os., 10 ex.
 Use: Vaginal cleanser.

DOVERINE. (Kremers, Urban) Dovers Pow.
 Tabs. Bot. 100s, 1000s.
 Use: Opiate for colds.

DOVER'S POWDER. Ipecac 1 part, opium 1 part, lactose 8 parts.
 Use: Analgesic sedative & diaphoretic.

W/Acetophenetidin, atropine sulfate, aspirin, camphor, caffeine & Sod. sulfate dried. See: Davlum, Cop. (Hance).

W/Acetophenetidin, aspirin, camphor, monobromated & caffeine. See: Ovesen, Tab. (Warren-Teed).

W/Acetophenetidin, aspirin, caffeine & camphor. See: Horvadyne No. 3, Cop. (G. F. Horvay)

W/Acetophenetidin, camphor, aspirin, caffeine & atropine sulfate.
 See: Analgestine, Cop. (Mollard).
 Ovasen, Cop. & Tab. (Warren-Teed).
 Aspldyne Comp., Cop. & Tab. (Lemmon).

W/Acetophenetidin, salicylamide, rocephedrine HCl, pyrilamine maleate & ascorbic acid. See: Cenamol, Cop. (Central)

W/A.P.C., atropine sulf., camphor. See: Aspldyne Comp., Tab. (Lemmon)

W/Aspirin, acetophenetidin, camphor monobromated, caffeine, atropine sulfate. See: Asco (Suttiff & Case)

W/Atropine sulfate, acetophenetidin & aspirin. See: Axopine, Tab. (Warren-Teed).

W/Aspirin, acetophenetidin & caffeine. See: Acetyl-Cof-Phen, Cop. (C. D. Smith).

W/Atropine sulfate, acetophenetidin, aspirin, gelsemium fr. & camphor monobromated. See: Afro-Salco, Cop. (Pitmon-Moore).

W/Atropine sulf., A.P.C., sod.-sulf. See: Dasin, Cop. (Messengill)

W/Atropine sulfate, Ipecac, aconite ext., capsicum, podophyllin, acetophenetidin & camphor. See: C.C.G. No. 2, Tab. (Noyes).

W/Atropine sulf., phenacetin, aspirin, gelsemium fr. 2 min., camphor monobromated. See: Afro-Salco, Cop. (Pitmon-Moore)

W/Pyranilamine maleate, acetophenetidin, aspirin, camphor, caffeine & atropine sulfate. See: Alercop, Cop. (Ethical Pharm.).

W/Pyrilamine maleate, rocephedrine HCl, aspirin, phenacetin. See: Asphomal D, Tab. (Central Pharm.)

W/Thenylpyromine HCl. See: Thorysa, Cop. (Warren-Teed).

DOVIAL. (Winthrop Labs) Methamphetamine
 * Generic name recognized by N.F., U.S.P. or A.M.A.

HCl 2.5 mg., thyraclin 10 mg., mebaral 30 mg./Tab. Bot. 100s.
 Use: Control obesity.

DOVIUM. (Hanse) Daver paw. 1/2 gr., acetophenetidin 1 1/2 gr., atropine sulfate 1/500 gr., aspirin 2 gr., camphor 1/4 gr., caffeine 1/2 gr., Sod. sulfate dried 7 1/2 gr./Cop. Bot. 100s, 1000s.
 Use: (1-2 q. 4 h.) Analgesic & antipyretic.

DOX&N. (Lloyd Bros) (Old name Doxinate with Danthron) (Diacyl sod sulfasuccinate 60 mg., danthron 50 mg./Cop. Bot. 30s, 100s.
 Use: Fecal softener

* **DOXAPRAM HYDROCHLORIDE.** (Robins) 1-Ethyl-4-(2-morpholinoethyl)-3,3-diphenyl-2-pyrrolidinone HCl hydrate.
 Use: CNS stimulant.

DOXIGEST. (Brean) Pancreatin 262.5 mg., hemicellulose 25 mg., belfaine HCl 65 mg., papain 15 mg., ketocholonic acid 12.5 mg., & deoxycholic acid 32.5 mg Enteric-coated Tabs. Bot. 100s
 Use: Treatment of maldigestion syndromes

DOXIDAN. (Lloyd Bros.) Danthron 50 mg., calcium bis-(diacyl sulfasuccinate) 60 mg./Cop. Bot. 30s, 100s.
 Use: laxative.

DOXIN&TE. N.N.D. 1962. (Lloyd Bros) Diacyl sodium sulfasuccinate, Aerosol OT. Cop. 60 mg. Bot. 30s, 100s. 240 mg. Bot. 15s, 100s tiq. (5%) Bot. 60 cc.
 Use: Orally (1-2 caps. or 1 cc. dolly), a wetting agent that keeps the stool soft.

DOXINATE-DANTHRON. (Lloyd Bros) Diacyl sod. sulfasuccinate 60 mg., danthron 50 mg./Cop. Bot. 30s, 100s. Name changed to Doxon.
 Use: Orally, laxative

DOXISUL. (Creitmed) Diacyl sodium sulfasuccinate 80 mg., danthron 40 mg/Tab. Bot. 100s.
 Use: Fecal softener.

DOXITOL. (Nisan) Pyridoxine HCl 5 mg., rutin 20 mg., ascorbic acid 100 mg., in base of protein hydrolysate/Tab. Bot. 100s.
 Use: (2 t.i.d.) Radiation sickness.

DOXTONE. (Haack) Pyridoxine HCl 20 mg., phenobarbital 16 mg/Tab. Bot. 50s, 100s.
 Use: (1 Tab t.i.d. or q.i.d.) For treatment of motion sickness.

DOXO. (Mallard) Deoxycorticosterone acetate 5 mg./cc. Vial 10 cc.
 Use: I.M., adrenal cortical insufficiency.

DOXOL. (Blair Labs.) Diacyl sod. sulfasuccinate 100 mg/Tab. Box 50s.
 Use: Treatment of constipation.

- DDXDTNE.** (Testagar) Desoxycholate ester
one acetate 5 mg/cc
Use Addison's disease Restores electro-
lyte balance
- DDXYCHDL-AS** (Brean) Daxychol X w/phe-
nobarbital 8 mg, atropine sulfate D 16 mg,
hyoscyamine HBr 0.16 mg/Tab Bot 100s,
1000s
Use (1 2 T i d, p c) Hydrocholeretic, spas-
molytic & sedative
- DDXYCHDL-K.** (Brean) Ketacholonic acid
3 gr, desoxycholic acid 1 gr/Tab Bot
100s, 1000s
Use (1 2 T i d, p c) Stimulate bile flow
- DDXYFED.** NND 1962 (Raymer) Meth-
amphetamine HCl, d desoxyephedrine HCl
Elix 20 mg/fl oz Bot 1 pt, 1 gal Tab
2.5 mg 5 mg Bot 100s 500s 1000s
Use: Orally central nervous system stimu-
lant
- W/Phenobarbital** (Raymer) Methamphetamine
HCl 25 mg, phenobarbital ¼ gr Tab Bot
100s 1000s
Use Stimulant w/sedative
- *DOXYLAMINE SUCCINATE, USP, NND**
1962 Tab USP 2 [alpha (2 d methyl-
amino ethoxy) alpha methylbenzyl]pyridine
succinate Available Amp, Cap, Cream,
Elixr, Lotion, Dphthal Dint, Solution,
Syrup, Tablet, See Decapryn Succinate
Syr, Tab (Merrell)
Use: (12 & 25 mg) Antihistamine
- W/APC** See: Decapryn Succinate w/APC,
Tab (Merrell)
- W/1 N Ethylephedrine HCl & theophylline**
aminasubutanol See Nathapryn Syr (Mer-
rell)
- W/Mercadol** See: Mercadol w/Decapryn
Syr (Merrell)
- W/Pyrimamine maleate phenstamine maleate**
See: Tridecamine Tab (National)
- W/Sod salicylate & caffeine** See: Decapryn
Compound, Syr (Merrell)
- W/Teprin hydrate & narcotine** See: Consol
Syrp (Merrell)
- DDZAR, NND 1962** (Tulag) See Metha-
pyriline HCl
- D P-A INJECTION** (Hiss) Pantotheryl alca-
hol 250 mg/cc Vial 10 cc
Use: I.M., vitamin therapy in the treatment
of intestinal atony and distention
- OPRDN** (Munn Pharm) Dipyrone 500 mg/
ml Vial 10 cc
Use Inj analgesic
- DRAMAL** (Klug) Phenyltoloxamine d,hydra-
gen citrate 50 mg/Tab Bot 100s
Use: Antihistamine
- DRAMAMINE, NND 1962** (Searle) beta
D-methylaminoethyl benzohydryl ether B
e Gener. c name recognized by NF, USP or AMA
chlorotheophyllinate (Dimenhydrinate
USP; Dramatin, Dramyl) Amp 250 mg
Box 6s, 25s, 100s; Elix 12.5 mg/dr Bot
1 pt Tab 50 mg Bot 12s, 100s, 1000s
Use Motion sickness
- DRAMAMINE-D.** (Searle) Dimenhydrinate
50 mg, d amphetamine sulf 5 mg/Tab
Bot 100s, 1000s
Use Antinauseant with central stimulant
- DRAMARIN.** See Dramamine, Prep (Searle)
- DRAMCILLIN, NND 1962** (White) Peni-
cillin G 100M u buffered w/Sod
citrate/5 cc Bot 60 cc
Use Orally; for penicillin susceptible or-
ganisms
- DRAMCILLIN WITH TRIPLE SULFONA-
MIDE,** (White) Penicillin G 100M u
sulfadiazine 0.167 Gm, sulfamerazine
0.167 Gm, sulfacetamide 0.167 Gm/5 cc
Bot 60 cc
Use Orally; penicillin & sulfonamids
therapy
- DRAMCILLIN-25D,** (White) Penicillin G
250M u, buffered w/Sod citrate/5 cc
Bot 60 cc
Use Orally; for penicillin susceptible or-
ganisms
- DRAMCILLIN 250 WITH TRIPLE SULFONA-
MIDE SUSP,** (White) Penicillin G
250 000 u, w/sulfadiazine sulfamerazine
& sulfacetamide D 167 Gm each/5 cc Bot
60 cc
- DRAMCILLIN-30D,** (White) Penicillin G
300 000 u/5 cc Susp 60 cc
- DRAMCILLIN SDD,** (White) Penicillin G
500 000 u/5 cc Bot 60 cc
- DRAMCILLIN-S,** NND 1962 (White) Pa-
lasstrum phenethicillin 125 mg/tsp pow-
form Bot 30 cc, 60 cc
Use Oral penicillin therapy
- DRAMYL** See Dramamine, Prep (Searle)
- DRE-NEL** (Warren Teed) Nux vomica FE
DD24 cc, hydrats 0.75 Gm, resorcinol 0.25
Gm w/digestive enzymes/30 cc Bot 1 pt-
1 gal
Use: Digestive and stomachic
- DRIED ALUMINUM HYDROXIDE GEL, USP**
See Aluminum Hydroxide Gel, Dried
- DRIED YEAST, USP, NNR 1952** See
Yeast, Dried
- DRILITDL S DLUTIDN** (Smith Kline & French)
Isotonic aq sol containing methoprene
HCl 0.2%, gramicidin D 0.05%, polymyxin
B sulfate 500 u, paredrine HBr 1%/cc-
Bot w/dropper ½ fl oz Plastic squeeze
bot containing ½ fl oz
Use: Respiratory tract infections
- DRINACET.** (Columbus) Phenylpropanolamine
HCl 15 mg, aspirin 230 mg, acetophenet-

- din 160 mg, caffeine 15 mg /Cap. Bot. 100s, 500s, 1000s
 Use: Anesthetic, antilussive, bronchodilator
- DRINALFA**, N.N.D. 1962. (Squibb) Methamphetamine HCl. (Each cc. contains 20 mg of d-dextroamphetamine HCl in sterile, aq sol.) Tab. 5 mg Bot. 100s, 250s. Vial 10 cc.
 Use: Stimulant.
- DRISDOL**, N.N.D. 1961. (Winthrop Labs.) Pure cryst. vit. D-2 [calciferol] from ergosterol. (Cryst. Vit. D-2) Sol. 5 cc., 10 cc., 50 cc., each accompanied by special dropper delivering 250 vit. D u./drop.
 Use: Rickets.
- DRISDOL 50,000 UNIT CAPSULES**. (Winthrop Labs.) Each 3 min. Cap. contains 25 mg. pure cryst. vit. D-2, equiv. to 5000 u. Cap. 50,000 u., Bot. 50s, 500s.
 Use: Vit. D therapy.
- DRISDOL W/VITAMIN A**. (Winthrop Labs.) Oily sol. contains 10,000 u. vit. D-2 & 50,000 u. vit. A/Gm. Sol. (water dispersible) in oil, Bot. 10 cc., 50 cc. Cap., Bot. 100s, 1000s.
 Use: Vit. A & O therapy.
- DRITOXEN**. (Walker) Methapyrilene HCl 1%, phenol 1%, zinc oxide 10%, zinc sulfate 0.5%, menthol 0.5%, benzethonium chloride 0.1% in washable, greaseless base Oint. Tube 1 oz. Jar ½ lb., 1 lb., 6 lb.
 Use: Fungus infections w/antihistamine.
- * DROCABIL**, N.F. Acetarsene salt of orecoline. See: Nemuro, Tab. (Winthrop Labs.)
 Use: Antihelmintic in animals.
- * DROCODE**. See: Dihydrocodeine.
- DROCOGESIC** No. 3. (Carrlone) Dihydrocodeine bitartrate 30 mg., aluminum aspirin 287 mg., acetophenetidin 150 mg., caffeine 30 mg /Tab. Bot. 50s, 100s.
 Use: Analgesic.
- DROCOLATE SINGLES**. (Lloyd Bros.) Tinfoil wrapped 4.5/cc. dose. Box 100s.
 Use: Cough.
- DRODEINE**. (Pitman-Moore) Dihydrocodeine bitartrate 1.67 mg., terpin hydrate 0.25 Gm./5 cc. Bot. 1 pt., 1 gal.
 Use: (1-2 tsp. p.r.n. m.p.) Narcotic cough preparation.
- DROLBAN** Inf. (Lilly) * Dramostanolone propionate 50 mg./cc. in sesame oil amp 10 cc.
 Use: Metastatic carcinoma of the breast
- DROMORAN HYDROBROMIDE**, N.N.R. 1953. (Roche Lab.) Racemorphan HBr, Methorphanon, di-3-Hydroxy-N methyl-morphinan HBr. Amp. (5 mg /cc.) 1 cc., Box 12s, 100s; Vial 10 cc. Tab. 5 mg H.T. Box 20s, 100s; O.T. Box 100s.
 Use: Narcotic analgesic similar to morphine.
- * DROMORAN**, LEVO. Levorphanol, N.N.D. 1959. Levorphan, N.N.R. 1954. See: Levor-Dromoran Tartrate, Amp., Tab. (Hoffmann-La Roche).
- * DROMOSTANOLONE PROPIONATE**. (Eli Lilly) 2 α -methylandrostan-17(β)-ol-3-one propionate. See: Drolban (Lilly)
 Use: Palliative treatment breast cancer.
- ORONACTIN**. (Merck Sharp & Dohme) Dexamethasone 0.25 mg., cyproheptadine HCl 4 mg /Tab. Bot. 100s
 Use: Antipruritic and anti-inflammatory.
- OROP CHALK**. (Various Mfr.) Cot. carb., prepared.
- OROPICILLIN**, N.N.D. 1962. (White) Pot. penicillin G 50M u. buffered w/Sod. citrate/20 drops. Bot. w/dropper 9 cc.
 Use: Orally, penicillin-susceptible organisms.
- OROPLEX**. (Massengill) Thiamine HCl 4 mg., riboflavin 1 mg., nicotinamide 40 mg., pyridoxine HCl 0.5 mg., ferrous sulfate 50 mg., liver fraction 400 mg. B-12 3.3 mg./cc. Sol. w/dropper 1 oz.
 Use: Anemia.
- OROPSPRIN**, N.N.D. 1962. (M. R. Thompson) Salicylamide 1 gr./cc. Liq. Bot. 1, 2, and 16 oz.
 Use: Orally, analgesic, antipyretic
- OSGERA ROTUNDIFOLIA AND THYMUS VULGARIS EXTRACT**. See: Diatessin, Drop & Syr. (Bischoff).
- DROWZ**. (Persen & Covey) Methapyrilene fumarate 25 mg., scopolamine HBr 0.125 mg., aluminum aspirin 100 mg /Tab. Bot. 25s.
 Use: Analgesic & sedative.
- DROXOLAN**. (Breon) 3,12-Dihydroxycholanolic acid Tab. 1½ gr. Bot. 100s.
 Uses: (1 q Ld.) Hydrocholeretic.
- DRXYPROPINE**. (Registered in Great Britain) 1-[2-(2-Hydroxyethoxy)ethyl]-4-phenyl-4-propionylpiperidine
- DRYCO**. (Borden) Vit. A 2500 u., D 400 u., B-1 0.43 mg., B-2 2.2 mg. added to spray-dried whole & skim milk/qt. Can 1 lb., 2½ lb.
 Use: Infant food.
- DRYTEX**. (C & M) Salicylic acid 2%, methylbenzethonium chloride 0.1%. Sol., Bot. 6 fl. oz.
 Use: Acne and seborrhea
- DRYVAX**. (Wyeth) Prepared calf lymph w/polyoxin B, streptomycin, & neomycin —when reconstituted, glycerin 50%, phenol 0.25%/Pkg. 1 10-dose vial, 1 diluent syringe w/sterile needle & 10 sterile scarifying needles.
 Use: Vaccination against smallpox.
- DS-OTICO**. (Bristol) Dihydrostreptomycin

* Generic name recognized by N.F., U.S.P. or A.M.A.

- base as sulfate 2.5 mg phenyltoloxamine citrate 10 mg, sodium citrate 5 mg glycerin q.s./ml Bot ½ oz w/ dropper
Use Treatment of infectious external otitis
- DST** See: Dihydrostreptomycin, Prep (Various Mfr)
- DT3 (DEXTRO 3 3 3' TRIIODOTHYRONINE)**, (Smith Kline French) Under study
- DUADACIN** (Lloyd Bros) Phenylephrine NCI 5 mg, chlorpheniramine maleate 1 mg, pyrilamine maleate 12.5 mg, caffeine 30 mg, ascorbic acid 50 mg, phenacetin 130 mg, salicylamide 200 mg / Cap Bot 20s 100s
Use: Cold syndrome
- DUALAR** (Armour) Benzcarbazimine benzyl N bis(ethyleneimido)phosphorocarbamate
Use Cancer therapy
- DUATRIC**, (Dumas Wilson) Ethinyl estradiol 0.01 mg, methyltestosterone 2 mg / Tab Bot 100s 1000s
Use: Treatment of menopause, osteoporosis and endocrine disorders
- DUATRIC PLUS** (Dumas Wilson) New name for Votric
- *DUAZOMYCIN**, (Pfizer) Antibiotic isolated from broth filtrates of *Streptomyces ombrofaciens*
Use Ant neoplastic agent
- DU-BIOTIC INTRANASAL**, (White) Hydrocortisone 0.2 mg, neomycin sulf 3.5 mg, gramicidin 0.05 mg, phenylephrine NCI 2.5 mg/cc Bot plastic spray and dropper bottle 13 cc
Use Intranasal antibiotic w/decongestant
- DU-BIOTIC TROCHES**, (White) Bacitracin 200 u, neomycin sulfate equiv to base 3.5 mg, propyl p aminobenzoate 2 mg / Troche Vial 15s & 100s
Use (1 q 3 4 h) Infections of throat
- DUCOBEE** (Breen) Cryst Vit B 12 In sterile isotonic sol Vial (1000 mcg/cc) 5 cc, 10 cc
Use IM, Vit B 12 therapy
- DUCOBEE-HY** (Breen) Hydroxocobalamin 1000 mcg/cc Vial 5 cc
Use Vitamin B 12 therapy
- DUCREY VACCINE**, (Lederle) Vial D2 cc, 5 cc
Use: ID; diagnostic aid in test for chancroid
- DUENITE** (Superior Pharm) Phenobarbital ½ gr, pentobarbital sod 1½ gr / Eupule Bot 50s
Use: (1 2 pr n) Sedative
- *DUGOS CRUDE CRYSTALS** See: Tyrothricin Prep (Various Mfr)
- DULCIVITE** (Lloyd Dabney & Westerfield)
* Generic name recognized by NF USP or A.M.A.
- New name for Bonavites; no change in formula, size or price
- DULCOLAX**, NND 1962 (Geigy) Bisacodyl bis(p acetoxypheyl) 2 pyridylmethane Tab EC 5 mg Box 6s, Bot 100s Supp 10 mg Box 6s 48s
Use: Constipation
- DUMOGHAN** (Squibb) Ethinyl estradiol 0.008 mg, methyl testosterone 4 mg with 11 Vits & 6 minerals / Tab Bot 60s, 250s
Use Geriatric therapy
- DUMONE**, (Squibb) Methyltestosterone 40 mg, ethinyl estradiol 0.008 mg / Tab Bot 100s, 1000s
Use: Androgen estrogen therapy
- DUNAGUENT**, (Ascher) Propesin (propyl p aminobenzoate) 10%, benzyl alcohol 4% urea 10%, para chloro-meto xylenol 0.8% Oint Tube w/applicator 1 oz, Jar 1 lb
Use: Topically anesthetic.
- DUNOKOL** (Cole Chem) Ox bile ext 3 gr-aur vomico ext ½ gr, Sod bicarb ½ gr-Cel phosphate 2 gr / Tab Bot 100s
Use (1-3 tid) Choleric & chologogue
- DUO-AOUA-DRIN** (McKesson & Robbins) Tyrothricin 1.5 mg, neomycin sulfate 5 mg, benzocaine 10 mg / lozenge Vial 12s
Use Treatment of mouth and throat irritation
- DUOBROPNEN**, (Massengill) Phenobarbital ½ gr, sod bromide 10 gr / 3 cc Elix Bot 1 pt, 1 gal
Use: Sedative and hypnotic
- DUOCNOL** (Plessner) Bile salts 2 gr, capsicum oleoresin & peppermint oil / Tab Bot 100s, 500s, 1000s
Use (1 2 tid) Choleric & chologogue
- DUO-C.V.P.** (U.S. Vitamin) Citrus bioflavonoid comp 200 mg, ascorbic acid 200 mg / Cap Bot 50s, 100s
Use (1 3 daily) Capillary tone & resistance W/Vitamin K (U.S. Vit Pharm) Citrus bioflavonoid compounds 200 mg C 200 mg, Vit K 132 mg / Cap Bot 50s
Use Anthemorrhagic
- DUODENAL ELECTROLYTE SOLUTION** (Don Baxter) Sod chloride 0.51 Gm, pot chloride 0.09 Gm, sod lactate 0.56 Gm, dexrose 10.0 Gm / 100 cc Bot 1000 cc
Use IV, Replace electrolytes
- DUODENAL SUBSTANCE**,
W/Bile salts, Sod bicarb & phenolphthalein
See: Chologic, Tab (Central)
W/Pancreatin, ox bile ext & papain
See: Karazyme, Tab (Karmel Co)
Miller zyme, Tab (Miller)
- DUODENUM POWDER** W/Metopine, olu

minum hydroxide & Mag. trisilicate. See Melracin, Tab. (Sirasenburgh)

DUODENUM WHOLE, DESICCATED & DEFATTED. See Crymanose, Tab. (Irwin-Neister).

W/Ferrous gluconate, vit. B-12. See: Bitric-sic, Cap. (Elder)

DUODON. (Kirkman) Belladonna paw. ext. $\frac{1}{8}$ gr., Sod. pentobarbital $\frac{3}{16}$ gr., phenobarbital $\frac{3}{16}$ gr./Tab. Bot. 100s, 300s, 1000s.

Use: Orally; antispasmodic & sedative.

DUO-ESTRO. (Central) Estrogenic sub. 2 mg., pot. estrone sulf. 1 mg./cc. Vials 10 cc.

Use: Estrogen therapy (0.25 to 1 cc. once or twice a week).

DUOGEN SUSPENSION. (Smith, Miller & Patch) Testosterone 25 mg., estrone 2 mg./cc. Vial 10 cc., 15 cc.

Use: Estrogen & androgen therapy.

DUOGBAFIN (Squibb) Iodipamide methylglucosamine and diatrizoate, Vials 50 cc.

Use: I.V., Bona and biliary tracts visualization.

DUOHORM. (Elder) Estrogenic substance 20,000 u., testosterone 25 mg., thiamine 20 mg./cc. Vial 10 cc.

Use: Estrogen & androgen therapy.

DUOHORM-F. (Canfield) Natural estrogenic substances 10 mg., gonadotropic hormone, 125 rat units (1 rat unit = 1 Fevald-Hisaw unit), Vial 5 cc.

Use: Will prevent biological depression of testicles or ovaries.

DUOHORM-M. (Canfield) Gonadotropic hormone, 125 rat units (1 rat unit = Fevald-Hisaw unit), testosterone 125 mg Vial 5 cc.

Use: Will prevent biological depression of testicles or ovaries.

DUOLVITE DROPS. (Ives Cameron) "A", Vit. A 10,000 u., D 2000 u., ascorbic acid 125 mg., mixed tocopherols 3 mg./cc. "B", Vit. B-12 10 mcg., thiamine HCl 3 mg., riboflavin 1 mg., pyridoxine HCl 1.6 mg., pantothenal 3 mg., niacinamide 20 mg./cc. Pkg. 2/15 cc. Bot.

Use: Growth promoting vitamin therapy.

DUOMIN. (Rorer) Ferrous sulfate 3 gr., Vit. B-1 0.3 mg., B 2 0.025 mg./Tab. Bot. 100s, 500s, 1000s.

Use: Anemias.

DUOMYCIN. See: Aureomycin, Prep. (Lederle)

DU-ORIA. (Ascher) Reserpine 0.25 mg., methamphetamine HCl 10 mg./tab. timed-released, Bot. 60s, 500s

Use: (1 daily) Tranquillizer.

DUOSCORB. (E. S. Miller) Procaine and Sod. ascorbates equiv. to 1 Gm. of ascorbic acid/amp. Box 12s, 25s, 100s.

DUOSOL. (Kirkman) Diactyl sod. sulfosuccinate 65 mg or 120 mg./Cap Bot. 100s

Use: Fecal softener

DUOSTERONE. (Roussel) Anhydroxyprogesterone 10 mg., ethinyl estradiol 0.01 mg./Tab. Bot. 10s, 25s, 100s.

Use: Orally; corpus luteum combined w/ estrogee.

DUO-STREP. (Merck Sharp & Dohme) Streptaducin, dihydrastrepilamycin sulfate & streptomycin sulfate in equal mix. Vial 1 & 5 Gm.

Use: Combined therapy

DUO-SULFANYL, NNR 1955 (Casimir Funk) Sulfadiazine, sulfamerazine 0.25 Gm. each/5 cc. Syr., Bot. 4 oz., 1 pt., 1 gal.

Use: Orally; disulfonamide therapy.

DUOSULFON. (Haskell) Oint., Sulfanilamide 12.5%, sulfathiazole 12.5%, Tube 1 oz. Susp., Plain, sulfanilamide 6.25%, sulfathiazole 6.25%, Bot. 1 fl. oz., w/ephedrine, sulfanilamide 6.25%, sulfathiazole 6.25%, ephedrine sulfate 1%. Bot. 1 fl. oz., 2 fl. oz., 6 fl. oz.

DUOTAL. (Marian) Phenobarbital $1\frac{1}{2}$ gr., phenobarbital Sod $1\frac{1}{2}$ gr./Tab. Bot. 100s.

Use: (1 m.p.) Sedative.

DUOFAL, NNR, 1937. See Guaiacol Carbonate (Various Mfrs.)

DUO-TBAL. (Fint, Eaton) Folic acid 5 mg., Vit. B-12 15 mcg./cc. Vial 10 cc., Box 6s

Use: I.M.; hemopoietic.

DUOTIN. (Klug) Duodenal sulistame $1\frac{1}{2}$ gr., aluminum hydroxide $1\frac{1}{2}$ gr., magnesium trisilicate $2\frac{1}{2}$ gr., homatropine methyl bromide 1/200 gr., whole pancreas $1\frac{1}{2}$ gr., amylose pancreatin N.F. 1 gr./Tab. Box 100s

Use: Ulcer therapy

DUOTBATE. (Marian) Pentaerythritol tetranitrate 30 mg./Timed release Cap Bot. 60s

Use: Treatment of angina pectoris

DUOTBATE-45 W/PHENOBARBITAL. (Marian) Pentaerythritol tetranitrate (PETN), 45 mg.; phenobarbital, 45 mg./Cap Bot. 60s

Use: Angina pectoris.

DUOTRONE. (U S Standard) Na. S: Ethinyl estradiol 0.02 mg., methyl testosterone 5 mg./Tab Bot 30s, 100s Na. 10: Ethinyl estradiol 0.04 mg., methyltestosterone 10 mg./Tab. Bot. 30s, 100s.

Use: Androgen & estrogen therapy.

DUOVIRON. (Haskell) Ferrous sulfate, a.s.c. 3 gr., thiamine HCl 1 mg., riboflavin 0.125 mg./Tab. Bot. 100s, 500s, 1000s.

Use: Hematinic.

DUOZINE, NNR 1955 (Abbot) (Dulcet Tab.) Sulfadiazine 0.075 Gm., sulfamerazine 0.075 Gm./Dulcet, Bot 100s, sulfadia-

- zine 0.15 Gm sulfamerazine 0.15 Gm /
 Dulcet Bat 100s sulfadiazine D25 Gm
 sulfamerazine 0.25 Gm / Tab Bat 100s
 1000s sulfadiazine 0.15 Gm sulfamerazine
 0.15 Gm Sod citrate 1.5 Gm / 5 cc Susp
 Bat 1 pt 1 gal
- DUPONOL** See Guidinal type Detergents
 (Sod Lauryl Sulfate) Various Mfrs.
- DUPHASTON** (Philips Roxane) Dydrogester
 one 5 mg / Tab Bat 20s Formerly iso
 gregnenone
 Use Orally prevent abortion
- DURABOLIN** (Organon) Nandrolone phen
 propionate 25 mg / cc in sesame oil Amp
 1 cc Box 3s Vial 5 cc See also Decca
 Durabolin
 Use Anabolic agent
- DURABOLIN-SD** (Organon) Nandrolone
 phenpropionate 50 mg in sterile sesame
 oil/10% benzyl alcohol / Vials 2 cc
 Use Prolonged anabolic stimulation
- DURACILLIN, ADR 1955** (Lilly) Procaine
 penicillin G
 Aq inj (300M u dose) Vial 1 11 10 dose
 in oil (300M u / cc) Vial 1 & 10 cc
 Disp syr 1 cc
 Ophth Oint (5M u / Gm) 1/2 oz
 Tab buffered (50M & 100M u) Bat 12s
 36s 100s 250s 1000s; (250M u) Bat 12s
 100s
 Traches (10M u) Bat 24s 100s
- DURACILLIN A S** (Lilly) Procaine penicillin G
 Disp syr 1, 5 & 10 cc; 600 000 u / Cartrid
 1s 25s 100s 300 000 u 1 000 000 u
- DURACILLIN A S W/Dihydrostreptomycin**
 (Lilly) Procaine penicillin G 300M u, dihy
 drostreptomycin base 0.3 Gm / Cartrid Bat
 25s 100s
- DURACILLIN FA** (Lilly) Procaine penicillin
 G 750M u Sod penicillin G 250M u / dose
 Vial 1 dose 10 dose
- DURACILLIN FORTIFIED** (Lilly) Procaine
 penicillin G 300M u, Pot or Sod penicillin
 G 100M u / cc Vial 1 dose, disp syringe
 1 cc & 10 cc; aq inj 400M u / dose Vial
 1 dose 10 dose
- DURACTON** (Nordic & chem cols) Corti
 cotrapin 40 u carboxymethylcellulose 8.3
 mg glucose 50 mg phenol 5 mg water
 for inject an cc Multidose vials 1 cc 5 cc
 Use ACTH therapy
- DURA-DEX.** (Banar) Dextroamphetamine sul
 fate 15 mg / S R Cap Bat 100s
 Use Control of appetite
- DURADORM** (Lincoln) Secobarbital 50 mg
 butobarbital 30 mg, phenobarbital 50 mg /
 Tab Bat 100s
 Use Orally (1 HS) sedative and hypnotic
- DURAFOAM VAGINAL TABLETS** (Durex)
 * Generic name recognized by N F U S P or A M A
- p Triisopropylphenoxypolyethoxyethanol 5%
 in a foaming tablet Pkg 12s
 Use: To prevent conception by chemical
 and physical means
- DURA-TAB 5 M** (Wynn Pharmacol Corp)
 Dextroamphetamine sulfate—Tab 10 mg
 or 15 mg Bat 30s 100s 250s
 Homatal—Homatrapine methylbromide 1/4
 gr phenobarbital 1 gr / tab Bat 30s
 100s, 250s
 Use: Sustained medication, antispasmodic
 w/sedative
- DURA-TAB 5 M AMINOPHYLLINE** (Wynn)
 Aminophylline 4 1/2 gr / Tab Bat 30s 100s
 Use Antiasthmatic diuretic
- DURIBEX** (Durst) Thiamine HCl 100 mg,
 nicotinamide 100 mg, pantothenic acid 10
 mg, riboflavin 1 mg pyridoxine HCl 1
 mg / cc Vial 10 cc
 Use Vit B complex
- DURYCIN** (Lilly) Procaine penicillin G 300
 000 u Sod penicillin G (buffered) 100M
 u dihydrostreptomycin sulfate 0.3 Gm /
 Vial Amp 1 dose Pkg 1s 25s, 100s
 Use IM; penicillin w/dihydrostreptomycin
 therapy
- DURYCIN A S** (Lilly) Procaine penicillin G
 400 000 u w/dihydrostreptomycin sulfate
 0.3 Gm / 2 cc Procaine penicillin G 300 000
 u, dihydrostreptomycin sulfate 0.3 Gm / 2
 cc Vial 10 cc 5 dose
 Use IM; penicillin & streptomycin therapy
- DURYCIN FA** (Lilly) Procaine penicillin G
 300 000 u, Sod penicillin G 100 000 u,
 dihydrostreptomycin sulfate 1 Gm / dose
 Procaine penicillin G 400,000 u Dihydro
 streptomycin sulfate 0.5 Gm Vial 1 dose
 Pkg 25s 100s
- DUSOTAL** (G F Harvey) Sod amobarbital
 3/4 gr, sod secobarbital 3/4 gr / cap Bat
 500s 1000s
 Use Sedative
- * **DUSTING POWDER, ABSORBABLE, U.S.P.**
 NND 1962 Starch derivative dusting
 pow See Bio Sorb Pow (Ethicon)
- DUSTING POWDER, SURGICAL** See B F I
 Powder (Merck Sharp & Dohme)
- DUTCH DROPS** Oil of Turpentine Sulfurated
- DUTCH OIL** Oil of Turpentine Sulfurated
- DUVA DOUCHE CONCENTRATE** (Embur)
 Lactic acid zinc sulfacarbolate thymol
 eucalyptol, menthol dextrose Bat 8 oz
 1 pt, 1 gal
 Use Vag nat cleanser
- D-VATINE** (Smith Darsey) Vit D, synthetic
 Cap 50 000 u Bat 100s 1000s
 Use Orally rickets & hypocalcemic tetany

D-VITAMIN CAPSULES. (Squibb) Vit. D-2 (Calciferol) Cap. 50,000 u. Bot. 100s.

Use Orally; rickets & tetony.

DVB 1D. (Mayrand) Vit. B-1 1 mg., B-2 1.2 mg., B-6 0.25 mg., niacinamide 10 mg., DVB 10, d-amphetamine sulfate 10 mg., butabarbital 60 mg./DVB 15, d-amphetamine sulfate 15 mg., butabarbital 60 mg. Bot. 100r.

Use: Mental and emotional distress.

D-V-R. (Horton & Converse) Vit. A 12,000 u., D 1000 u., C 100 mg., B-1 5 mg., B-2 5 mg., B-6 1 mg., niacinamide 25 mg., cal. pantothenate 6 mg., E 1 IU, B-12 2 mcg., folic acid 100 mcg., choline bitartrate 40 mg., inositol 20 mg., biotin 4 mcg., iron 15 mg., yeast ext. 60 mg., liver desic. 60 mg w/ minerals/2 Tabr. Bot. 100s, 250s, 500s, 1000s. Ure Vitomin and mineral supplement.

D-W CAPSULES. (Dumas-Wilson) Vit. D 400 u., B-1 2 mg., B-2 2 mg., dibaric Col. phosphate 11 gr., ferrous gluconate 3 gr./Cap. Bot. 100s, 1000s.

Use: Vitamin & mineral supplement in pregnancy & lactation.

DYCLONE, N.N.D. 1962. (Pitman-Moore) Dyclonine 4-n-Butoxy-benz-(1-piperidyl) propiophenone HCl.

Cream—1%, Tube 1 oz.

Sol.—0.5%, Bot. 1 & 8 oz.

Use: Topically, anesthetic & antiseptic.

See: Neo Polycin Otic, drops (Pitman-Moore)

***DYCLONINE HCl,** N.N.D. 1962. See: Dyclone, Cream, Sol. (Pitman-Moore)

W/Neomycin sulfate, polymyxin B sulfate, hydrocortisone acetate. See: Neo polycin HC Otic (Pitman-Moore)

DYCOMENE. (Honce) Dihydrocodeinone bitartrate 1/6 gr., pyrilamine maleate 1 1/7 gr./fl. oz. Bot. 3 oz., 1 gal.

Use: (1-2 tsp. q. 3-4 h) Sedative for coughs due to cold.

***DYDROGESTERONE.** (Philips Roxone) 9 β 10a-pregna-4,6-diene-3,20 dione. See Duphaston

DYES.

See: Antreptic, Dyer.

Diagnostic Agents.

DYFLOS. See: Dinorpropyl fluoraphosphate

DYLATE. (Spencer Lab) Clonidine

Use: Coronary vasodilator

DYLEPHRIN. (Irwin, Neirler) Epinephrine

HCl 2.5%, atropine sulfate 0.5 Gm./100 cc.

Aerial sol., Bot. 15 cc. & 30 cc.

Ure: Vaporized for orthmatic therapy.

DYMELOZ. (Eli Lilly & Co) Acetabexamide.

Use: Blood sugar lowering compound

DYNACAL LIQUID. (McNeil) Dicalcium phos-

* Generic name recognized by N.F., U.S.P. or A.M.A.

phate 6.5 Gm., Cal. 1.5 Gm., Phos. 1.15 Gm., Vit. D-3 2400 u./fl. oz. Bot. 12 fl. oz.

Use: Calclem & Vit. D therapy.

DYNACORYL. See: Nikethamide, Prep (Various Mfr)

DYNARSAN. See: Acelarrone, Tab. (Abbott)

DYNATROPIN, N.N.D. 1960

DYNOLEN. (Merrell) Betaine HCl 167 mg., glycine 167 mg., folic acid 0.85 mg., Vit. B-6 1 mg., B-12 5 mcg., Intrinsic factor conc. 333 mg., Vit. C 50 mg., ferrous sulfate 200 mg./Cap. Bot. 100r.

Use: (1 t i d) Anemiar.

***DYPHYLLINE,** N.N.D. 1962, 7-(2,3-Dihydroxypropyl) Theophylline. Hyphylline. See: Neothylline, Preps (Paul Maney)

DYPRIN. (Lincoln Labs.) dl-methionine 0.25 Gm./Cap. Bot. 30s.

Use: Eliminate acidosis and ammoniacal urine and dioper torh.

DYSCO TABLETS. (Abbott) Ephedrine HCl 1/2 gr., acetophenetidin 1 1/2 gr./Tab. Bot. 100s, 1000s.

Use: Analgesic w/vasoconstrictor.

DYSMENGESIC. (Walden) Pentobarbital sodium 30 mg., prophepyridomine molecule 15 mg., desferoamphetamine sulfate 2 mg., hyoscine hydrobromide 0.05 mg., octylsalicylic acid 300 mg./Tab. Bot. 50s.

Use: Premenstrual tension and dysmenorrhea

EAR DROPS. (Sutrin & Case) Phenol 0.50%, menthol 2.00%, benzocaine 1%, alcohol 16.83%, glycerin 79.39%. Ulp. Dropper Bot. 1/2 oz., 1/2 oz., pl

EARTHNUIT OIL. (Peonut Oil.)

EAST INDIAN SANDALWOOD W/Kova-Kova resins. See: Gonaron, Cap (Riedel).

EBICOL. (M. R. Thompson) Thiamine HCl 1 mg., riboflavin 2 mg., niacinamide 10 mg., choline citrophosphate, equiv. to choline 410 mg., inorital 200 mg., liver fraction #1 250 mg., yeast concentrate 250 mg., ace bran 250 mg., honey 250 mg., Pot acetate 100 mg., alcohol 10%/5 cc.

Bot. 8 oz., 1 gal.

Use: Ergotropic & Vit. B complex.

E-B-Z COMPOUND. (Cole Chem. Co) Ergat 10%, benzocaine 1%, zinc oxide, balsam peru, phenol, pine needles oil, cartor oil.

Oint. Tube 1 oz.

Use: Topically; antreptic & protective.

ECBOLINE. See: Ergotaxine. (Na Mir. currently lists)

***ECHOETHIOPHATE IODIDE,** N.N.D. 1962

2 Diethoxyphosphinythiaethyl-trimethylammonium iodide See: Phosphatine iodide [Campbell]

ECOFROL. (Breen) d alpha Tocopherol (30

- mg /Cap as the acetate Cap 30 mg, 100 mg } Bot 100s 500s 1000s 200 mg Bot 100s Inj (200 mg /ml) Vial 10 cc
 Use Orally; Vitamin E therapy
- ECOLID, NND 1962** (Ciba) Chlorisonda mine chloride Tab (10, 25 mg, 50 mg) Bot 100s
 Amp [5 mg] 1 ml, Box 5s
 Use Orally, antihypertensive agent
- ECON** (Cale Chem Co) d Tacopherol acetate (equiv in biological activity to 50 mg, international standard Vit E) Bot 100s, 500s 1000s
 Use Orally Vitamin E therapy
- ECON-O-GENE** (Econ O Lab) Douche powder 3 oz
- ECOTRIN** (Smith, Kline & French) Aspirin 5 gr /duentric coated tab Bot 100s
 Use Analgesic
- *ECTYLUREA, NND 1962** See 2 Ethyl cis cralanylurea
 See Levonil Tab (Upjohn)
 Nostyn, Tab (Ames)
- W/Ethaxolamide medroxyprogesterone
 See Cytron, Tab (Upjohn)
- W/Hexocyclium methylsulfate See: Tralcyon, Tab (Abbott)
- W/Medroxyprogesterone acetate ethazol amide See Cytron (Upjohn)
- W/Trol See Tralcyon Fillob (Abbott)
- ECZOL OINTMENT** (Smith, Muller & Patch) Ammoniated mercury, zinc oxide, Sod hyposulfite salicylic acid ichthammol rusci oil cod liver oil Tube 1 oz Jar 1 lb
 Use Topically antiseptic
- E O SYRUP** (Tilden) Ethymorphine NCl $\frac{1}{4}$ gr, chloratam $\frac{1}{8}$ min, ephedrine sulfate 1 gr Pot guaiacol sulfonate 10 gr squill flex 2 min, ipocac flex 2 min, sanguinaria flex 4 min, menthol 1/16 gr benzocaine 1/5 gr, talv 1 gr /fl oz Bot 1 pt, 1 gal
 Use (1 2 tsp q 3 h) Expectoant, sedative
- *EDATHAMIL** Ethylene diammetetraacetic acid
 See Nullapans (General Aniline)
 Sequestrene (Altrae Chem)
 Versene (Berswarth Chem, Du Pont)
- *EDATHAMIL CALCIUM-DISODIUM, NND 1962** Calcium Disad Ethylenediamine tetraacetate See: Calcium Disodium Versenate Amp & Tab (Riker)
- W/Piperazine NND 1959 See Perin, Syr (Endo)
- *EDATHAMIL DISODIUM, USP** Disad salt of ethylene diamine tetraacetic acid See Endrate Disadum (Abbott)
 Prachelate Salm (Croakes Barnes)
- * Generic name recognized by NF USP or A.M.A.
- EDITIC, See: Ethylenediamine Tetraacetic Acid**
- EDITHAMIL, See: Edalhamil**
- EDIOL** (Schenley) Coconut oil 50% & sucrose 12.5% Bot 16 fl oz
 Use Nutritional supplement
- EDRISAL** (Smith Kline & French) Benzodrine sulfate 2.5 mg, acetylsalicylic acid 2 $\frac{1}{2}$ gr phenacetin 2 $\frac{1}{2}$ gr /Tab Bot 50s, 500s
 Uses (1 q 3-4 h) Relieves pain & the depression state
- W/Codaine (Smith, Kline & French) Codeine sulfate $\frac{1}{2}$ gr or $\frac{1}{4}$ gr, benzodrine sulfate 2.5 mg, acetylsalicylic acid 2.5 gr, phenacetin 2.5 gr /tab Bot 50s
- *EDROPNIUM CHLORIDE, USP, NND 1962** Inj USP Dimethylethyl (3 Hydroxyphenyl) ammonium chloride See Tension Chloride, Vial (Roche Lab)
- EDYA** See Edalhamil
- EENT-O AUGINE** (Canfield) Neomycin sulfate 6 mg hydrocortisone 5 mg, pyril amine maleate 0.6%, cetalkonium chloride 0.05%, dipiperdon HCl 1.0%/cc Liquid Bot plastic dropper 5 cc
 Use: Treatment of otitis media
- EEZ ACNECARE**, (Pfizer) Tyrothricin, 0.5% hexachlorophene 2.5%, sulfur, 8% resorcinol, 2% zinc oxide 10% alcohol, 9.3% /Aerosol Bot 2 oz
 Use Treatment of symptoms of acne pimples & other skin blemishes
- EEZ SPRAY POWDER**, (Pfizer) Dichlorophens Aerosol Spray Cons
 Use Athlete's Foot
- EPA** (Columbus) Soybean oil 50%, soybean lecithin 300 mg, pyridoxine HCl 6 mg /15 cc Bot 1 pt
 Use Hypercholesteremia in atherosclerosis
- EFACIN** (Persan & Covey) Niacin 500 mg /Tab Tab 100s
 Use In elevated blood cholesterol levels
- EFEDRON HYDROCHLORIDE** (Hart) Ephedrine NCl 0.6% & chlorbutanol 0.5% w/Sad chloride, menthol & cinnamon oil in a water soluble jelly base Nasal tipped collapsible tubes 20 Gm
 Use Shrinkage of nasal mucous membranes
- EFEMIST** (Hart) Ephedrine HCl 1%, chlorbutanol 0.5%, w/menthol, eucalyptol & sucrose in physiological salt sol Bot 1 oz 8 oz, 1 pt
 Use Relief of acute rhinitis, coryza, sinusitis, eustachitis
- E-FEROL**, (Testagar) dl Tacopherol acetate Cap (30 mg) Bot 100s, 500s 1000s &

- Cop (100 mg) Bot. 25s, 50s, 100s, 500s, 1000, 200 & 400 I.U. Bot. 100s, 500s. Vial (200 mg/cc) 10 cc.
 Use: Vitamin E therapy.
- E-FEROL SUCCINATE.** (Testogor) d-alpha Tocopherol acid succinate, equiv. to vit. E 200 I.U./tab. Bot. 100s, 500r, 1000s.
 Use: Orally, vitamin E therapy.
- EFFERGEL.** (Stuart) Sodium carboxymethylcellulose 2 Gm., di-(acetylhydroxyphenyl) isatin. 6 mg./7 Gm. Pow. Bot. 4 & 9 oz.
 Use: Laxative.
- EFFERSYL.** (Stuart) Psyllium 3 Gm., di-(acetylhydroxyphenyl) iratin 3 mg./tspt. Bot. 4 & 9 oz.
 Use: Laxative.
- EFFERSYLLIUM.** (Stuart) Psyllium hydrocolloid with sucrose, potassium bicarbonate, citric acid, artificial sweeteners 3 Gm./rounded tsp. (7 Gm) Bot 9 oz.
 Use: Constipation
- EFFERVESCENT GRANULES.** (Reed & Carnrick) Sod. bicarb., citric acid, Cal. lactate, Pot. bicarb., Sod. phosphate, Mag. gluconate, sugar. Bot. 1 oz., 2 1/2 oz.
 Use: Orally, antacid.
- EFFERVESCENT LAXATIVE COMPOUND.** See: Hepatone, Pow. (Merck Sharp & Dohme)
- EFOCAINE.** (Fougere) Not listed Procaine 1%, procaine HCl 0.25% & butyl p-aminobenzoate 5% in polyethylene glycol 300 2%, propylene glycol 7E% & water Vial 20 cc.
 Use: I.M., S.C., anesthesia analgesia for prolonged action.
- EFO-NOL.** (Warren-Teed) Ephedrine HCl 1/6 gr., w/loraz emetic, squill, senega, cocillano, euphorbia & wild lettuce/5 cc. Bot. 1 pt., 1 gal.
 Use: {t-2 lrp. q. 3 h. p r n} Cough preparation.
- EFRACIN.** (Petson & Covey) Niacin 0.5 Gm / Tab
- EFRICEL-10%.** (Professional Pharm) Phenylephrine HCl, 10% in methylene blue Bot. 5 cc. 1/2% & 2.5% Bot. 15 cc.
 Use: Produce pupil dilatation.
- EFRICON EXPECTORANT.** (Lannett) Dihydrocodeinone bitartrate 18 mg, phenylephrine HCl 5 mg, chlorpheniramine maleate 2 mg, ammonium chloride 90 mg. Pot. guaiaacal suff 90 mg., Sod citrate 60 mg / 5 cc. Bot 1 pt.
 Use: Antihistaminic nasal decongestant.
- EFROXINE HYDROCHLORIDE, N N D. 1967.** (Maltbie) Methamphetamine HCl; Elixir 20 mg / fl. oz. Bot 1 pt., 1 gal.; Tab. (5 mg) Bot. 100r, 500r, 1000r. w/phenobarbital, Bot. 100r, 500r.
- Use Orally; central nervous system stimulant & appetite depressant.
- EGESTEROL-STRONGER.** (Tartagor) Progesterone 5 mg, Vit. E 10 mg. Vial 10 cc. & progesterone 25 mg., Vit. E 50 mg /cc. Vial 10 cc.
 Use: Recurrent abortions & threatened abortions.
- EGRAINE.** A protein binder from oatr. See prep. Treidal (Organon).
 W/Mag. trisulfate, aluminum hydroxide, Cal. carb., Mag. carb. & regional See: Treidal, Tab. (Organon)
- E-HART.** (Hart) di-alpha-Tocopherol acetate 200 mg /cc. Vial 10 cc.
- EHRLICH 594.** See: Acetarrone, Tab. (Abbott).
- EHRLICH 606.** See: Arspenamine (Variour Mfr.).
- EIGHT VITAMIN DROPS.** (Philadelphia Ampoule) Vit. A 5000 u., D-2 1000 u., B-1 1 mg, B-2 1 mg., B-6 1 mg, d-pantothal equivalent to 11.6 mg d-calc pantothenate, niacinamide 10 mg, Vit. C 50 mg / 0.6 cc. Eol. 50 cc.
 Use: Vitamin therapy in children
- EKANS.** (Maag) Rauwolfia serpentina 100 mg./tab. Bot. 100s.
 Use: (2-3 daily) Hypertension
- EKOMINE.** (Lloyd) Atropine methyl nitrate Tab. (1/60 gr.) Eol. 100s, 500s, 1000s.
 Use: Orally; antispasmodic.
- W/Phenobarbital.** (Lloyd) Atropine methyl nitrate 1/60 gr., phenobarbital 1/4 gr./tab. Bot. 100s, 500r, 1000s
 Use Orally; antispasmodic & sedative.
- EKZEBROL.** (Tosco) Strontium bromide 1 Gm./10 cc. saline. Amp. 10 cc. Box 6.
 Use: I.V., antipruritic.
- EL-ACORTO GEL.** (Breon) Pituitary adrenocorticotrophic hormone (ACTH) in gelatin. Corticotropin. Vial (40 u. or 80 u./cc) 5 cc.
 Use: I.M., corticotropin therapy.
- ELASE.** (Parke, Davil)
 Use: Fibrinolytic enzyme for debridement of ulcers, wounds, etc.
- ELASE GINT. & PWD. FOR SOL.** (Parke, Davil) Fibrinolysin 25 u, deoxyribonuclease 15,000 u /Vial or lyophilized pwd May be reconstituted with 10 cc of isotonic NaCl Sol Gint.s Fibrinolysin 30 u., deoxyribonuclease 20,000 u /30 Gm. Tube 10 Gm. Vials 30 cc Tube pkg of 6.
 Use: Pretealytic Enzyme Combination in gynecologic complications
- ELAVIL.** (Merck Sharp & Dohme) Amitriptyline HCl/Tab. 10 mg., 25 mg., Bot. 100s; Vial (10 mg/cc) 10 cc.
 Use: Antidepressant therapy
- ELDEC.** (Parke, Davil) Methyl tartartrate

167 mg, theelin 0.167 mg, lakadiastase 20 mg, pancreatin 1333 mg Vt B 12 w/intrinsic factor conc 0.033 u, folic acid 0.1 mg, Vit C 333 mg ferrous sulf 167 mg Vit A 1667 u B 1 0.67 mg B 2 0.67 mg B 6 0.5 mg nicotinamide 167 mg choline bitartrate 6.67 mg sod pan lothenate 5 mg, iodine 0.05 mg, cal carbonate 667 mg lysine HCl 667 mg dl methionine 167 mg /Kapreol Bot 100s
Use: Hormone, vitamin and mineral supplement

ELDEMOL (Elder) Stearyl alcohol white petrolatum; acetylated lanolin lanolin alcohol extract sorbitan sesquioleate; polyoxy 40 stearate propylene glycol methyl paraben purified water gm
Use: Dermatores

ELDI-PEN (P B Elder) Procaine penicillin G 300,000 u, Pot penicillin G 100,000 u /vial Vial 10 cc
Use: Respiratory penicillin therapy

ELDOCAINE (Elder) Procaine hydrochloride 2%, Epinephrine 1:50,000 Vial 30 cc

ELDONAL, (Canright) Belladonna alk, pentobarb ¼ gr, ½ gr or ¾ gr Tabs & El
Use: Sedative spasmolytic

ELDOVITE (Canright) Vitamins 17 and Minerals 10 Tabs
Use: Vitamin & mineral supplement

ELE-B (U S Vitamin) Vit B 1 25 mg B 2 10 mg B 6 15 mg B 12 5 mcg niacinamide 75 mg cal pantothenate 75 mg choline 50 mg, inositol 15 mg, citrus bio flavanoid compound 20 mg, dentic liver 125 mg, yeast 25 mg /Cap Bot 50s, 100s
Use: Dietary supplement

ELECTROLYTE THERAPY,
See: Gastric Electrolyte Soln (Baxter)
K.M.C. (Ingram)
KNL Solution (Cutter)
Lactated Pot Soln USP
Lactol Sol (M E Thompson)
Lytren (Mead Johnson)
Polycol (Cutter)
Tis u sol (Baxter)

ELECTROCORTIN See: Aldosterone

ELEVENS, (Fuller) Vitamins -11 and minerals -12 Bot 100s

ELIDIN (Endo) See: Dimethozan

ELIPTIN, NND 1962 (Ciba) Aminoglutathimide alpha (p-aminophenyl) alpha ethyl glutarimide 125 mg and 250 mg /Tab Bot 100s
Use: Anticonvulsant in grand mot and petit mal

ELIXIR B-O-PHOS, See: B G Phos Elixir (Merck Sharp & Dahme)

ELIXIR METHENAMINE, SALICYLATE &
Generic name recognized by N.E., U.S.P. or A.M.A.

CITRATE COMPOUND See: Methenamine Elixir (Merck Sharp & Dahme)

ELIXIR PHENOBARBITAL, See: Liquid Elixir (Merck Sharp & Dahme)

ELIXIR RNUBARB ALKALINE, N.F. See: Neutralizing (Merck Sharp & Dahme)

ELIXIR SAW PALMETTO & HYDRANGEA COMPOUND See: Lithiated Sorghum Compound (Merck Sharp & Dahme)

ELIXIR STRONTIUM & COLCHICINE COMPOUND See: Panalgyn Liq (Merck Sharp & Dahme)

ELIXIR VIT B COMPLEX W/GLYCEROPHOSPHATE See: B G Phos Elixir (Merck Sharp & Dahme)

ELIXODYNE (Lemman) N acetyl p-aminophenol 120 mg alcohol 10%/5 cc Bot 3 fl oz 1 pt 1 gal
Use: Reduce fever and pain in infants and children

ELIXOPHYLLIN (Sherman) Theophylline 80 mg alcohol sugar free base/15 cc Bot 1 pt, 2 pt

Use: Diuretic for original patients

ELIXSED (Chicago Pharmacal) Barbital Sod 4 gr, strontium bromide, Pot bromide 8 Sod bromide 20 gr each & hyoscyamine HBr 1/32 gr/or Bot 1 pt 1 gal
Use: Sedative & mildly hypnotic

ELKOSIN, NND 1962 (Ciba) Sulfonamide Sulfisomidine N¹ (2,6 dimethyl 4 pyrimidyl) sulfanilamide Tob (0.5 Gm) Bot 100s 500s, 1000s; Susp in Syr 250 mg per fl oz (4 cc) Bot 1 pt

Use: Sulfonamide susceptible organisms

ELLAGIC ACID See: Gallogen Tab (Mason Gilg)

ELMALOIN W/PENOBARBITAL, (Elder) Di phenylhydantoin sod 0.1 Gm, phenobarbital 15 mg /Cap Bot 100s 500s
Use: Sedative

ELON See: p-Methylaminophenol (No list currently lists)

ELORINE CHLORIDE, NND 1962 (Lilly) Tricyclamide chloride 100 mg /Fulvule Bot 100s, 1000s 50 mg Bot 100s, 1000s
Use: Anticholinergic in the treatment of peptic ulcer

W/Amytal See: Co Elarine, Tob (Lilly)

ELPATEC, (Patch) Pot penicillin G 250 M.u. acetaphenetidin 100 mg, salicylamide 100 mg, phenyltoloxamine dihydrogen citrate 25 mg /Tab Bot 24s 100s

Use: (1 tid) Respiratory infections

ELPAGEN (Smith Miller & Patch) Sod salicylate 5 gr, Sod p-aminobenzoate 3 gr, salicylamide ½ gr, Vit C 30 mg, d hy droxyaluminum aminoacetate ½ gr /Tab Bot 100s
Use: Antacid w/analgesic salicylates

ELPAGEN W/CODEINE. (Smith, Miller & Patch) Sod. salicylate 5 gr., Sod. p-aminobenzoate 3 gr., Vit. C 30 mg., dihydroxy-aluminum aminoacetate $\frac{1}{2}$ gr., salicylamide $\frac{1}{2}$ gr., codeine phosphate $\frac{1}{4}$ gr./Tab. Bot. 60s.

Use: 2 t.i.d.; rheumatic conditions.

ELPRECAL. (Conright) Combination of 11 Vits., 12 minerals, choline inositol/Cap. Bots. 100s.

ELSERPINE. (Conright) Reserpine D.25 mg / Tab. Bots. 100s.

Use: Hypotensive-ironquillizing agent.

ELVANOL. (Dupont) Polyvinyl Alcohol

EMAGRIN FORTE. (Clapp) Phenylephrine HCl 5 mg., atropine sulfate 1/1000 gr., acetophenetidin 1 gr., aspirin 2 gr., caffeine $\frac{1}{2}$ gr., salicylamide 1 gr./Tab. Bot. 25s, 125s, 500s.

Use: Common cold.

EMBECHINE. Aliphatic chloroethylamine.

Use: Treatment of cancer.

EMBUR 90. (Embura) Amino acids, oral, Tab. (7 $\frac{1}{2}$ gr.) Bot. 100s

Use: (6-12 daily) Nutritional supplement.

EMOEE MARGARINE. (Pitman-Moore) Each 100 Gm. provides 52 Gm. unsaturated fatty acids w/each pound fortified w/Vit. A 15,000 u. and Vit. D 2000 u. Can 1 lb. Use: To reduce blood cholesterol levels.

EMERACOL. (Upjohn) Codeine phosphate 1/12 gr. w/Pol. guaicol sulfanate, chloroform, ammonium chloride, antimony & Pol. lactate/5 cc. Bot. 1 pt., 1 gal.

Use: (1-2 tsp. q. 3-4 h. p.r.n.) Cough preparation.

EMESD. (Columbus) Vit. B-6 10 mg., betaine HCl 400 mg., inulose 11.2 Gm., dextrose 11.2 Gm./fl.oz. Bot. 3 oz.

Use: Antinausea.

EMETINE BISMUTH IODIDE, N.N.R. 1936.

Use: Same as Emetine HCl.

***EMETINE HYDROCHLORIDE, U.S.P. 1ej, U.S.P.**

Lilly—Amps. $\frac{1}{2}$ and 1 gr./cc. pkg. 6s and 12s.

Lilly—Enseols.

Upjohn—Amps. $\frac{1}{2}$ and 1 gr./cc. pkg. 5s.

Use: Amebicide.

EMETROL. (Kinney) A phosphated carb. sol. Bot. 3 oz., 1 pt.

Use: Nausea & vomiting.

EMIVAN. (U.S. Vitamin) Vanillic acid diethylamide. *Ethamivan. Parenteral, Amps., 100 mg / 2 cc 5s, 25s, 100s; 500 mg / 10 cc. 1s, 5s, 25s; Tabs. 20 mg 100s, 1000s.

Use: Respiratory stimulant.

EMKO VAGINAL FOAM. (Emko Co) Noxyl

* Generic name recognized by N.F., U.S.P. or A.M.A.

phenoxy polyoxyethylene ethanal 8%, benzethonium chloride 0.2%. Kit.

Use: Prevention of pregnancy.

EMOCIN. (Burroughs-Wellcome) Not listed. Aspirin 2 gr. w/slippy elm/loz. Bot. 20s. Use: Orally; analgesic.

EMOLLIENTINE. (Parke, Davis) Aluminum hydroxide, zinc oxide, zinc sulfoacarbolate, phenol, phenylmercuric lactate. Tube 1 $\frac{1}{2}$ oz. Jar 1 lb., 5 lb.

Use: Emollient in minor burns.

EMP INJECTION. (United Labs) Liver inj. 10 mcg., Vit. B-12 20 mcg., folic acid 5 mg., Vit. B-1 10 mg./cc. Vial 10 cc.

Use: Anemias.

EMPIRAL. (Burroughs-Wellcome) Phenobarbital $\frac{1}{4}$ gr., acetophenetidin 2 $\frac{1}{2}$ gr., aspirin 3 $\frac{1}{2}$ gr./Tab. Bot. 100s, 1000s.

Use: (1 q. 3-4 h.) Sedative & analgesic.

EMPIRIN. See Acetylsalicylic Acid (Various Mfr)

EMPIRIN COMPOUND TABLET. (Burroughs Wellcome) Acetylsalicylic acid 3 $\frac{1}{2}$ gr., acetophenetidin 2 $\frac{1}{2}$ gr., caffeine $\frac{1}{2}$ gr./Tab. Bot. 12s, 25s, 100s, 500s, 1000s.

Use: Orally; analgesic & antipyretic.

W/Codeine phosphate. (Burroughs Wellcome) In addition to Empirin comp. No. 1 codeine phosphate $\frac{1}{2}$ gr., Bot. 100s, 500s. No. 2 codeine phosphate $\frac{1}{4}$ gr., Bot. 100s, 500s. No. 3 codeine phosphate $\frac{1}{2}$ gr., Bot. 100s, 500s. No. 4 codeine phosphate 1 gr., Bot. 100s, 500s.

Use: Orally; analgesic & antipyretic.

EMPLEX. (Minnesota Pharmacol) Vit. B-1 10 mg., B-2 5 mg., niacinamide 100 mg., calcium pantothenate 5 mg./ss. Vial 10 cc

Use: B complex deficiencies.

EMPRAZIL. (Burroughs Wellcome) Pseudoephedrine HCl 20 mg., chlorcyclizine HCl 15 mg., acetophenetidin 150 mg., aspirin 200 mg., caffeine 30 mg./Tab. Bot. 100s, 1000s.

Use: Treatment of the common cold

EMPRAZIL-C. (Burroughs Wellcome) Codeine phosphate 15 mg., pseudoephedrine HCl 20 mg., chlorcyclizine HCl 15 mg., acetophenetidin 150 mg., aspirin 200 mg., caffeine 30 mg./Tab. Bot. 24s, 100s.

Use: Treatment of colds.

EMULSEROL. (Upjohn) Mineral oil w/Vit. A 27,000 u./100 cc. Plain, emulsion, Bot. 12 oz., 24 oz. w/cascara, Bot. 12 oz., 24 oz.

Use: Orally; laxative.

***EMULSION OINTMENT BASE (O/W).** (Modified Hydrophilic Ointment) See: Dermabase (Marcella Pharmaceuticals)

EMULSION BASE. (Almay) Fatty acid esters,

spermaceti, cetyl alcohol, propylene glycol
Oint 4 oz, 1, 5, 25 lb
Use Ointment base

***EMYLAMATE**. 1 Ethyl 1 methylpropyl carbamate See Striatran, Tab (Merck Sharp & Dahme)

ENARAX. (Roerig) Oxyphenyclimine NCI 5 mg or 10 mg, hydroxyzine HCl 25 mg / Tab Bot 60s
Use Ant cholinergic

ENBIN, NNR 1955 (Witham Cooper)
Benzyl benzoate 0.113 Gm, chlorophenothane 10 mg & ethylaminobenzoate 20 mg / cc Emulsion, Bot 90 cc
Use Scabicide w/local anesthetic

EN-CEBRIN-F. (Lilly) Vit A 4000 u, B 1 30 mg B 2 20 mg, nicotinamide 100 mg, B 6 20 mg folic acid 10 mg, B 12 activity equivalent to 1/2 oral u, calcium pantothenate 50 mg, C 500 mg, synthetic D 400 u K 0.5 mg, rutin 100 mg, phosphorus free calcium carbonate 6000 mg, plus seven other minerals / Pulvule Bot 100s 500s
Use Prenatal vitamin & mineral supplement

ENCOTUSSIN-C. (Endo) Codeine phosphate 10 mg hamatropine methylbromide 0.5 mg pyrilamine maleate 125 mg, ammanium Cl 60 mg, sodium citrate 85 mg / 5 cc 4 oz
Use Relief of cough due to colds

ENCYNEX. (Anglo French) Acetophenetidin 2 1/2 gr w/Sod salicylate, paminobenzoic acid & Sod bicarb / Tab Bot 50s
Use (1 2 q 3 h) Analgesic & antipyretic

END-A-KOFF. (Columbus) D Methorphan HBr 5 mg chlorpheniramine maleate 1 mg, glyceryl guaiacolate 20 mg / Tab Bot 24s 100s 1000s
Use: Treatment of cough due to cold symptoms

ENDOARSAN (Endo) Sod cacodylate 1 Gm, mercury bismuthide 5 mg, sod iodide 10 mg, organic phosphorus 0.6 mg / 5 cc Amp 5 cc Box 6s 25s & 100s
Use Herpes zoster

***ENDOBENZILINE BROMIDE**. N N [dimethyl] aminomethyl a [bicyclo[2.2.1] 5 heptenyl] mandelate methyl bromide See Ulcyn (Pitman Moore)

ENDOCAINE. (Endo) Pyrrocaine
Use Local anesthetic

ENDOFER (Endo) Ferrous sulf 12 Gm thiamine HCl 10 mg, riboflavin 4 mg nicotinamide 30 mg, Cal pantothenate 5 mg, liver concentrate 0.5 Gm, Vit B 12 10 mcg / fl oz Bot 8 oz, 1 pt. 1 gal
Use Anemias

ENDOFERARSAN (Endo) Iron cacodylate
* Generic name recognized by NF, USP or A.M.A.

65 mg, Sod cacodylate 100 mg / 5 cc Amp 5 cc Box 6s, 25s, 100s
Use: Anemias

ENDOGEN-A. (Endo) 50 mg solubilizer casein, 5 mg colloidal sulfur, 60 mg pep lone / 2 cc Amp. 2 cc, Box 12s, 25s, 100s
Use: Arthritis

ENDOGLOBIN. (Endo) Ferrous sulfate exsic 0.2 Gm, Vit B-12 2 mcg, thiamine HCl 1 mg, riboflavin 2 mg, nicotinic acid 10 mg, liver fraction, secondary pow 0.2 Gm / Tab Bot 100s 1000s
Use Iron & vitamin therapy

ENDOGLOBIN-C. (Endo) Ferrous sulfate exsic 0.2 Gm, Vit B 12 2 mcg, thiamine NCI 1 mg, riboflavin 2 mg, nicotinic acid 10 mg, ascorbic acid 50 mg, liver fraction secondary pow 0.2 Gm / Tab Bot 100s 1000s
Use Anemias
W/folic acid (Endo) Ferrous sulfate exsic 0.2 Gm, thiamine HCl 1 mg, riboflavin 2 mg, nicotinic acid 10 mg, ascorbic acid 50 mg, folic acid 1 mg, liver fraction secondary pow, 0.2 Gm, cobalamin 2 mcg / Tab Bot 100s, 1000s
Use Anemias

ENDOGLOBIN FORTE (Endo) Celachrom 1/2 oral NF unit, B 1 10 mg, ferrous fumarate 500 mg, Vit C 150 mg, E 2 6 mg B 6 3 mg nicotinamide 30 mg, pantothenol 10 mg / Tab Bot 100s, 1000s
Use Iron deficiency anemia

ENDOJODIN. See Enlodan (Winthrop Lab)

ENDOLAC. (Endo) Vit A 7500 u, D 450 u, B 1 3 mg B 2 3 mg, B 6 1.5 mg, E 12 3 mcg, C 90 mg, nicotinamide 30 mg, Cal pantothenate 15 mg, folic acid 0.75 mg w/Cal lactate, Cal carbonate ferrous sulfate, Pot iodide copper sulfate cobalt sulfate, Mag sulfate, Mang sulfate ammonium molybdate, Pot sulfate, zinc sulfate / 3 Tab Bot 100s, 1000s
Use (1 1 d) Nutritional supplement

ENDO-MAGSAL. (Endo) Aluminum hydroxide 6 1/2%, Mag trisilicate 7 1/2 gr / fl oz Bot 12 oz Tab, Bot 50s 100s, 1000s
Use Antacid

***ENDOMYCIN**. A new antibiotic obtained from cultures of *Streptomyces endus*

ENDOPHENOLPNTNALEIN (Roche Lab)
Diacetyldioxyphenylisatin Isocan Bisatin See: Diacetylhydroxyphenylisatin Prep (Vorlies Mir)
Use, laxative

ENDOTNYRIN. (Nelson Bayer) Standardized thyroglobulin ext Tab (1/4, 1/2 & 1 gr) Bot 100s, 500s (1 1/2 gr) Bot 100s 500s 1000s

Use: Myxedema, cretinism, hypothyroid states.

2NDOTUSSIN-NN Pedf. Syr. (2nd) d-methorphan HBr 5 mg., homatropine methylbromide 0.15 mg., ammonium chloride 60 mg., sodium citrate 40 mg./Tsp. Bot. 4 oz.

Use: Control of unproductive cough

INDBATZ DISODIUM. (Abbott) 2eothamil disod., Disodium salt of ethylenediamine tetra-acetic acid 3 Gm./Amp Box 5x-20 cc. amp.

Use: Chelating agent for post-agric calcification, angina pectoris & scleroderma.

ENDURON. (Abbott) Methylclothiazide 2.5 or 5 mg./Tab. Bot. 100s, 1000s.

Use: Diuretic.

ENDUBONYL. (Abbott) Methylclothiazide 5 mg., deserpidine 0.25 mg./Tab. Bot. 100s, 1000s.

Use: Antihypertensive.

2NDUBONYL PASTE. (Abbott) Methylclothiazide 5 mg., deserpidine 0.5 mg./Tab. Bot. 100s, 1000s.

Use: Antihypertensive.

ENELONZ. (Funk, Cosmir) Pregnenolone 100 mg./cc. of aq. susp. & pregnenolone acetate 100 mg./Tab. Vial 10 cc., Tab. Bot. 24s, 100s.

Use: Rheumatoid arthritis.

ENEMOL. (Cutler) Sodium biphosphate 16 Gm., sod. phosphate 6 Gm., methyl paraben and propyl paraben/4½ oz. Plastic disp. tube with six in. tip

Use: Routine enemas

EN2SCOB. (Arlington-Funk) Pregnenolone acetate 100 mg., ascorbic acid 333 mg./Tab. Bot. 24s, 100s.

Use: Rheumatoid arthritis.

2N2VAC INS2RTS. (Storck) Pot. bicarbonate and lactic acid in a special nonirritating base. Supp. Box 12s.

Use: Insert to start bowel movement.

2NFAMIL. (Mead Johnson) A 1500 units, D 400 units, C 50 mg., thiamine 0.4 mg., riboflavin 1 mg., niacinamide 4 mg., pyridoxine 0.3 mg., Co-pantothenate 2 mg., E 5 mg., B-12 1 mcg., inositol 300 mg., choline 85 mg., Fe 1.4 mg., Cu 0.39 mg., I 65 mcg./13 oz. can liquid, 1 lb. can pwd.

Use: Infant feeding.

ENFAMIL WITH IRON. (Mead Johnson)

Use: Infant feeding.

ENFORBEC, A D.R. 1955. (Harrower) Thiamin HCl 10 mg., riboflavin 5 mg., niacinamide 100 mg., pyridoxine HCl 1 mg., Cal. pantothenate 10 mg., folic acid 0.1 mg., ascorbic acid 100 mg., plus natural B complex

from liver B fraction and yeast/Tab. Bot. 100s, 1000s.

Use: Orally water-soluble vitamins.

2N GRAN. (Squibb) Vit. A 5,000 u., D 500 u., thiamine mononitrate 3 mg., riboflavin 3 mg., pyridoxine HCl 2 mg., vit. B-12 2 mcg., K 0.5 mg., folic acid 0.25 mg., niacinamide 20 mg., calcium pantothenate 5 mg., ascorbic acid 75 mg., cal. 150 mg., iron 10 mg., iodine 0.15 mg., pot. 5 mg., copper 1 mg., mag. 6 mg., mang. 1 mg., zinc 1.5 mg./Tab. Sor 10s, 250s, 1000s. Tab. Term-Pak. 27Cs

Use: 1 Tab daily. Vitamin-mineral supplement.

ENGBAN BABY DROPS. (Squibb) Vit. A 5000 u., D 1000 u., C 70 mg., B-1 1.2 mg., B-2 2 mg., niacinamide 12 mg., B-6 2 mg., d-pantothelol 5 mg., B-12 6 mcg./0.6 cc. Pediatric drops, Bot. 15 cc., 50 cc.

Use: Vitamin supplement

ENISYL W/IRON. (Luka) L-lysine monohydrochloride 100 mg., ferrous fumarate 50 mg., Vit. B-6 0.5 mg./Tab. Bot. 100s.

Use: Iron deficiency anemia.

2NISYL-V. (Luka) Vit. A 2000 U.S.P. u., Vit. D 200 U.S.P. u., Vit. B-1 10 mg., Vit. B-2 10 mg., Vit. B-6 40 mg., niacinamide 50 mg., calcium pantothenate 2.5 mg., Vit. C 340 mg., Vit. B-12 20 mcg., calcium 100 mg., L-lysine monohydrochloride 200 mg., ferrous fumarate 100 mg./Tab. Bot. 100s.

Use: Iron deficiency anemia of pregnancy.

ENKID2. (Brewer) Enteric-coated tab. contains 0.5 & 1 Gm. of Pot. iodide. Bot. 100s, 500s, 1000s.

Use: Pot. iodide therapy.

ENOFEN. (Kremers Urban) Aminophylline 1½ gr., phenobarbital ¼ gr., ephedrine ½ gr./Tab. Bot. 100s, 500s, 1000s.

Use: Asthma relief.

ENOVID. (Searle) Norethynodrel [17-alpha-ethynyl-17-hydroxy-5(10)-estren-3-one] and Ethynylestradiol 3-methyl ether. (The latter present 0.15 mg./10 mg. Tab) Tab. 10 mg. Bot. 50s, Norethynodrel 5 mg., & ethynylestradiol 3-methyl ether 0.075 mg./Tab. Vial 20s.

Use: Control of many menstrual disorders

2NRUMAY. (Tilden) Pyriminole maleate 25 mg., caffeine 32 mg./Tab. Bot. 100s, 1000s.

Use: Antihistamine.

2NSOLB2C. (Harrower) Each 30-cc. vial contains: Thiamine HCl 150 mg., riboflavin 45 mg., pyridoxine HCl 25 mg., Cal. pantothenate 250 mg., niacinamide 1875 mg., gentiolic acid 2% (as preservative) 30-cc vials with or without diluent. Box 1s, 25s.

Use: Vitamin B complex.

ENSOLBEC-C. (Harrower) Each 20 cc. vial

contains Thiamine HCl 500 mg, riboflavin 50 mg, pyridoxine HCl 100 mg, Sod pantothenate 100 mg, niacinamide 500 mg ascorbic acid 1000 mg, gentisic acid 2% (as preservative) Vial 20 cc with diluent

Use Water soluble vitamins

ENSOLBEC-C 500 In Horro matic vial (Hor rows) Each 2 cc vial contains Thiamine HCl 100 mg, riboflavin 10 mg, pyridoxine HCl 20 mg, Sod pantothenate 20 mg niacinamide 100 mg, ascorbic acid 500 mg Horro matic vial Box 1s, 25s

Use Water soluble vitamins

ENTAB. (Schilling) Hesperidin 10 mg, Vit C 100 mg, salicylamide 2 gr, phenylephrine 5 mg, pyrrolamine lactate 10 mg, caffeine citrate 15 mg/Tab Bot 18s, 100s

Use Symptomatic relief of common cold

ENTACYL SUSP. (British Drug Houses) Piperazine adipate 600 mg /5 cc Syr Bot 4 16 & 80 oz

Use Pinworm

ENTACARB (Reed Cornrick) Col carb, Mag carb, Sod bicarb, Pot bicarb, kaolin bismuth subcarb Pow 3 oz, 1 lb, 5 lb

Use Orally; antacid

ENTERA-VITA N 2. (Ethical Pharm) Ferrus gluconate 3 gr, folic acid 0.5 mg, B 6 33 mg, Vit A 5000 u, O 1000 u, B-1 1.5 mg, B 2 1.5 mg, C 15 mg, B 12 0.5 mcg, niacinamide 20 mg, cal pantothenate 1 mg, cal lactate 5 gr, cu 1 mg, mang 1 mg, molybd 0.2 mg, pot 5 mg, zn 12 mg, mag 0.2 mg, A 0034 mg / Tab Bot 100s

Use (1 to 6 daily) Dietary supplement

ENTIREX. (Person & Covey) ladachlorhydroxyquin 167 mg, methylcellulose 167 mg, aluminum hydroxide 160 mg, atropine sulf 0.0067 mg, hyosine HBr 0.003 mg, hyoscyamine sulf 0.0367 mg/Tab Bot 100s

Use Diarrhea

ENTEROSULPON (Consol Midland) Phthalyl sulfacetamide 0.5 Gm /tab Bot 50s, 100s, 250s, 1000s

Use: Sulfonamide therapy

ENTERO-VIOFORM. (Ciba) ladachlorhydroxyquin Entero-Tab 250 mg, Bot 20s, 100s

ENTODON. (Winthrop Labs) Propisodal Entodoin, Endopodin, 2 Hydroxylmethyl ene bis (trimethyl-ammonium iodide) 20% Amp 2 cc, Box 10s

Use: (I V, I.M., subcu 2 cc) Iodine therapy

ENTODOIDIN. See: Entodon (Winthrop Labs)

ENTOMIST. (Waisen Teed) Neomycin sulfate 0.1%. phenylephrine HCl 0.25% thymyl pyramine HCl 0.1%. methylbenzethonium

chloride 0.01% in a buffered solution Bot spray 5/8 oz

Use Antibiotic & decongestant

ENTOQUEL. (White) Thihexinal methylbromide 5 mg /5 cc Syr Bot 6 oz

Use Diarrheal

ENTORAL PULVULES (Lily) Pneumococci 40 000 million, streptococci 15 000 million H influenzae 2500 million, M. catarrhalis 2500 million/Cop Bot 20s, 60s

Use Orally, cold vaccine

ENTOZYME (Robins) 250 mg pepsin, 300 mg pancreatin & 150 mg bile salts/Tab Bot 100s, 500s

Use (1 2 pc) Digestant

ENTROMONE, NND 1960 (Endo) Chorionic gonadotropin Vial (500 rat u/cc when diluted w/accompanying vial containing 9 cc of saline sol)

Use I.M., chorionic gonadotropin therapy

ENTRON S-M. (Whittier) Vit D, 50 000 u activity, salicylamide 162 mg, mephensin 125 mg/Cop Bot 100s

Use For acute & chronic rheumatoid arthritis

ENTROSYL. (Wirth) Oral suspension Bot 3 oz

ENZACTIN, NND 1962 (Ayerst) Triacetin (glyceryl triacetate) 250 mg /Gm Spray Bot 3 oz Pow Pkg 1 1/2 oz Cream Tube 1 oz

Use Prevention and treatment of superficial dermatophytoses, particularly athlete's foot and ringworm of the scalp

ENZON (Breon) Chymaltrypsin 5000 u/cc in sesame oil Vial 5 cc

Use I.M., anti-inflammatory

ENZIFLUR. (Ayerst) Sod fluoride 2.21 mg Vit C 30 mg Vit D 400 u/1oz Bot 100s

Use Prevention of dental caries

ENZIN (Minnesota Labs) Pepsin 100 mg pancreatin 400 mg, iron bile salts 100 mg-liver concentrate w/stomach substance 150 mg Vit B 1 1 mg B 2 0.75 mg, B-6 1 mg, C 50 mg/Tab Bot 100s

Use Gastric digestant

ENZOBARR (K & M Labs) Amylolytic enzyme 30 mg proteolytic enzyme 10 mg cellulolytic enzyme 3 mg, phenobarb sol 1/8 gr/Tab Bot 50s

Use Digestive enzyme

ENZOBILE (Mallard) Pancreatin 200 mg, mycozyme diastase 100 mg, pepsin 100 mg, hog bile desiccated 200 mg, ext aux vomica 8 mg/Tab Bot 100s, 1000s

Use: Pancreatic deficiency

ENZO-CAL. (Crookes Barnes) Cream containing calamine 5%, zinc oxide 5% & benzocaine 3% Tube 1 1/2 oz Jar 1 lb

Use: Minor irritations of the skin

ENZODASE, LYOPHILIZED. (Squibb) Hyaluronidase. Vial 150 or 1500 I.U.

Use: S.C.; to promote diffusion into the tissues.

ENZYDYNE-B. (Endocrine) Digestive enzymes with B complex. Tab., Bot. 100s

Use: Vit. B complex & digestive.

ENZYLAC. (Sulff & Case) Isolated pepsin 100 gr./fl. oz. liq. Bot. gal

ENZYMES, N N D. 1960

See: Adenosine. 5-Monophosphoric acid (Various Mfr.).

Alidase (Searle).

Alpha Chymar (Armour)

Aspergillus Oryzae Enzyme (Various Mfr.).

Cabaden (Rand).

Cholinesterase (Various Mfr.).

Chymar, Vial. (Armour Labs)

Colozyme (Organon)

Cytochrome-C (Wyeth).

Dioctase (Various Mfr.).

Diffusin (Hyaluronidase) (Ortho)

Dornovac (Merck Sharp & Dahme)

Enzodase (Squibb).

Ficln. (No Mfr. lists)

beta-Glucuronidase (Warner-Chilcatt).

Histaminase (Various Mfr.).

Hyaluronidase (Various Mfr.).

Hyozyme (Abbott).

Inerase (Neisler)

Inverase. (No Mfr. lists.)

Kelodase (Warner-Chilcatt).

Mall Extract (Barchardt).

My-B-Den (Bischoff).

Neurinase (Faugera)

Pancreatin (Various Mfr)

Popain. (No Mfr. lists)

Papayo, Carico. (No Mfr. lists.)

Pepsin (Various Mfr.)

Protamide (Sherman).

Rennin (Various Mfr.).

Streptokinase-Streptodarnase (Lederle).

Taka-Dioctase (Parke, Davis).

Tryptar Trypsin (Armour)

Varidase (Lederle).

Wydose (Wyeth)

ENZYMET. (Lloyd, Dabney & Westerfield)

Pancreatin 300 mg., pepsin 200 mg., taurocholic acid 40 mg./Tab. Bot. 77.

Use: Indigestion.

ENZYPAN. (Norgine) Peptic potency equiv. to pepsin 0.009 Gm., amylolytic potency equiv. to pancreatin 0.276 Gm., proteolytic potency equiv. to pancreatin 0.372 Gm., lipolytic potency present, as bile desic 0.056 Gm./Tab. Vial 40s, 120s, 500s.

Use: Digestive enzyme deficiencies.

EPHED-ORGANIDIN. (Wampole) Ephedrine

* Generic name recognized by N.F., U.S.P. or A.M.A.

sulfate 24 mg., organidin 60 mg., methapyrene hydrochloride 50 mg./Tab Bot 50s.

Use: Respiratory diseases.

EPHEDRA W/Pat. guaiacalsulfonate, sanguinaria, squill, thyme, menthol, lactic acid, Sod. benzoate. See: Fedrang, liq. (Cole Chem. Co.).

EPHEDREMAL. (G. F. Harvey) Ephedrine sulfate $\frac{1}{2}$ gr., pyrilamine maleate 25 mg./tab. Bot. 1000s, 5000s.

Use: (1 q. 3 h) Antihistamine w/vasoconstrictor.

***EPHEDRINE, N.F., N.N.R. 1950;** Spray, N.F. X; Compound spray, N.F. X 1-1-Phenyl-2-methylamine-propanol.

Abbott—Inhalant. Ephedrine 1% in light mineral oil. Bot. 1 fl. oz., 4 fl. oz., 1 pt., 1 gal.

Burroughs Wellcome—Isotonic sol. Ephedrine 1% in modified lacte's sol. Bot. 1 fl. oz., 1 pt.

Lilly—Compound inhalant. Ephedrine 1% w/menthol, thyme oil in light liq mineral oil. Bot. 1 fl. oz., $\frac{1}{2}$ pt., 1 pt., 1 gal.

Lilly—Plain, inhalant. Ephedrine 1% w/cinnamic aldehyde, benzaldehyde in light liq. petrolatum. Bot. 1 fl. oz., $\frac{1}{2}$ pt., 1 pt., 1 gal.

Mallinckrodt—Pow. hydrate $\frac{1}{2}$ gr., 1 oz.

Merck—Pow. hydrate. Bot. $\frac{1}{2}$ oz., 1 oz., 4 oz., 1 lb.

Parke, Davis—Compound, inhalant. Ephedrine 1% w/camphor, menthol, geraniol in liq. petrolatum. Bot. 1 fl. oz., 1 pt.

Parke, Davis—Plain, inhalant. Ephedrine 1% in liq. petrolatum. Bot. 1 fl. oz., 1 pt. See: Monadrin, liq. (Endo).

Racephedrine HCl (Various Mfr.).

EPHEDRINE & AMYTAL. (Lilly) Ephedrine sulfate $\frac{1}{2}$ gr. & amytal $\frac{1}{4}$ gr./Tab. Bot 100s, 500s. Pulvules contain three times the above.

EPHEDRINE COMPOUND W/Merthiolate. (Lilly) In addition to plain, contains menthol, camphor & thyme oil. Bot. 1 fl. oz.,

EPHEDRINE ETHYLENEDIAMINE W/Theobromine sod. acetate, caffeine, and acetophenetidin. See: Kraton, Tab. (Field Labs)

EPHEDRINE GLYCEROLE. (E. L. Patch) Ephedrine HCl gr. $\frac{1}{2}$ /fl. oz. in a base of Irish moss & glycerin. Bot. 4 fl. oz., 1 pt., 1 gal. Use: Orally; cough.

EPHEDRINE HYDROCHLORIDE, N.F., N.N.E. 1957; Cap., N.F. Tab., N.F. (Various Mfr.); Cryst Box $\frac{1}{2}$ oz., $\frac{1}{4}$ oz., 1 oz., 4 oz.

Use: Sympathomimetic agent W/Phenobarbital, theophylline, belladonna leaves, pyrilamine maleate, potassium io-

dide, magnesium trisilicate See Theofed Pediatric, Tab (Sutliff & Case)

W/Prednisone, theophylline, phenobarbital See Deltasmyl (Roussel)

EPHEDRINE HYDROCHLORIDE NASAL

JELLY. See Eledron Hydrochloride, Oint (Hart)

EPHEDRINE HYDROCHLORIDE, RACEMIC,

NNR 1957, NND 1959 See: Ephelotin, Tab (Merck Sharp & Dahme)

EPHEDRINE LACTATE W/Dextrose, phenyl mercuric lactate, tyrothricin, cetyl trimethyl ammonium See Gluco Thricil, Sol. (Parke, Davis).

EPHEDRINE & MERCAPTOLIDE, SOLUTION

(Upjohn) Ephedrine HCl 10%, orthoxy droxypheylmercuric chloride (mercaptolide) 1:5000 Bot 1 pt

Use Nasal mucous membranes

EPHEDRINE & NEMBUTAL (Abbott) Ephedrine HCl $\frac{3}{4}$ gr., nembutal gr $\frac{3}{8}$ /Cap Bot 100s, 500s

Use: Orally; sedative w/antispasmodic

EPHEDRINE & NEMBUTAL CAPSULES, NO. 2. See Nembu Fedrin Cap (Abbott)

EPHEDRINE-NOVOCAIN SOLUTION. (Winthrop Labs) A clear, colorless solution, each cc containing 50 mg ephedrine HCl & 10 mg novocain HCl Amp 1 cc & 2 cc, Box 10s, 100s

Use: Spinal anesthesia

EPHEDRINE PLAIN W/Merthiolate (Lilly)

Ephedrine 1%, merthiolate 1:5000 in oil base Bot 1 fl oz, $\frac{1}{4}$ pt, 1 pt, 1 gal

Ephedrine Product Combinations on pp 281, 282, 283 and 284.

EPHEDRINE-RELATED COMPOUNDS.

See: Ephedamine (Various Mfr)

Aramine (Merck Sharp & Dahme)

Aranthral (Knoll)

Benzedrex (Smith, Kline & French)

Sutanetrin (Winthrop Labs)

Clapone (Lilly).

Cyclopentamine . HCl (Lilly).

Dexedrine Sulfate (Smith, Kline & French)

Ethylnorepinephrine (Winthrop Labs)

Parthone (Lilly).

Hydroxyamphetamine . HBr (Smith, Kline & French)

Mephentermine (Wyeth).

Metaraminol (Merck Sharp & Dahme)

Methamphetamine . HCl (Various Mfr.)

Methoxamine . HCl (Burroughs Wellcome).

Methoxyphenamine . HCl (Upjohn)

Methylaminoheptane (Knoll)

Methylhexanamine (Lilly).

Naphazoline . HCl (Ciba).

Nethamine . HCl (Merrell).

Nethalol (Merrell).

Oenethyl (Knoll)

Orthaxine (Upjohn)

Paredrine (Smith, Kline & French)

Phenylpropanolamine . HCl (Merck Sharp & Dahme)

Phenylpropylmethylamine (Merrell).

Privine (Ciba)

Propadrine (Merck Sharp & Dahme)

Propylhexadrine (Smith, Kline & French).

Tuaminoheptone (Lilly).

Vasoxyl (Burroughs Wellcome)

Vanadrine (Merrell)

Wyamine (Wyeth).

EPHEDRINE & SECONAL SODIUM (Lilly)

Ephedrine sulfate $\frac{3}{4}$ gr., & seconal Sod $\frac{3}{4}$ gr./ensole or pulvule Pulvule Bot 100s 500s

Use Nocturnal asthma

*EPHEDRINE SULFATE, USP, NNE 1950

Cap, USP; Inj, USP; Jelly, NF₄ Sol, NF; Syrup, NF, Tab, NF

See AFKO, Syr (APC)

Ephedsol, Liq (Lonnell)

Isorefrol, Isotonic Solution (Blue Line)

W/Phenobarbital See: Ephedrol, Cap (Sutliff & Case)

EPHEDRINE SULFATE 25 mg W/Benadryl

HCl 50 mg (Parke, Davis) 8al Kapsenit 100s, 1000s

EPHEDRINE TANNATE W/Corbetapentane

tannate, phenylephrine tannate, chlorpheniramine tannate, pyrilamine tannate See Rynatan Expectorant, Liq (Irwin, Neisler)

W/corbetapentane tannate, chlorpheniramine tannate, phenylephrine tannate See Rynatuss Tabule (Irwin, Neisler)

EPHEDRITAL (Sherman) Phenobarbital $\frac{1}{4}$ gr, ephedrine sulfate $\frac{3}{8}$ gr/Tab Bot 100s 500s

Use: (1 q 3-4 h, mod praesc) Vasoconstrictor w/sedative

EPHEDROL (Lilly) Codeine sulfate 1/6 gr, ma huang, Pot guaiaecalsulfonate, Tolu balsam, menthol, squill/5 cc Bot 1 pt, 1 gal

Use (1 tsp q 3-4 h, mod praesc) Cough

EPHEDROMAL (Blue Line) Carbramal 5 gr, ephedrine HCl $\frac{3}{8}$ gr/Cop Bot 100s

Use (1 2 tid) Sedative & antiasthmatic

EPNEDROYAL (Sutliff & Case) Ephedrine sulf $\frac{3}{8}$ gr, phenobarbital $\frac{3}{4}$ gr/Cop Bot 100s, 1000s

Use Vasoconstrictor w/sedative

EPHEDSOL (Lonnell) Ephedrine sulfate 1% (pink) and 3% (green) in normal saline Bot 1 pt, 1 gal

Use: Oral ephedrine therapy

*1-EPHENAMINE PENICILLIN G See Cephenamine Vial (Commercial Solvents).

* Generic name recognized by NF, USP or AMA

Ephedrine Product Combinations

| Ephedrine | Others | Manufacturer | Brand |
|-----------------|--|-----------------|--|
| (Sulf.) 0.5% | Allantoin 0.4%, sulfanilamide 2%, telfthiazole 1% | National Drug | Allantamide liq. w/Ephedrine, liq. Oint. |
| (Sulf.) 3/4 gr. | Allyl-phenylbarbitol 3/4 gr. | Massengill | Alphed, Cap. |
| (Sulf.) 3/4 gr. | Aludrine 10 mg., phenobarbitol 1/4 gr., theophylline 2 gr. | Leeming | Nephenolin, Tab. |
| (HCl) 3/4 gr. | Aminophylline 2 gr., amobarbitol 3/4 gr. | Hoack | A. E. A., Tab. |
| (HCl) 3/4 gr. | Aminophylline 2 gr., omylat 3/4 gr. | Lilly | Amaset, Cap. |
| (Sulf.) 1/4 gr. | Aminophylline 2 gr., phenobarbitol 1/4 gr. | Harvey | Ampofed, Cap. |
| (Sulf.) 3/4 gr. | Aminophylline 1 1/2 gr., phenobarbitol 1/4 gr. | C. D. Smith | Amsdarb, Tab. |
| (Sulf.) 3/4 gr. | Aminophylline 1 1/2 gr., pheobarbitol 1/4 gr. | Smith-Dorsey | Dorsamin, Tab. |
| (HCl) 1/12 gr. | Aminophylline 1 gr., pheobarbitol 1/4 gr., ethyl aminobenzoate 1/4 gr., aluminum hydroxide 1/2 gr. | Irwin, Neisler | Dainite Pediatric, Tab. |
| (Sulf.) 1/2 gr. | Aminophylline 1 1/2 gr., phenobarbitol 1/2 gr., epharbita, f. ext., 1 1/2 gr. | Warree-Teed | Eu-Phed-Amie, Tab. |
| (HCl) 1/4 gr. | Aminophylline 2 gr., pheobarbitol 1/2 gr., Pot. iodide 3 gr. | Foythress | Mudrane, Tab. |
| (Sulf.) 24 mg. | Aminophylline 65 mg., phenobarbitol 8 mg., pyrilamine maleate 12.5 mg. | Raymer | Barbhist, Tab. |
| (Sulf.) 3/4 gr. | Aminophylline 1 1/2 gr., phenobarbitol 1/2 gr., reybosed 1 gr. | Raymer | Rayphyllin, Cap. |
| (HCl) 1/4 gr. | Aspirin 7 1/2 gr. | Wallace | Caryphedrine, Tab. |
| (Sulf.) 10 mg. | Aspirin 3 gr., phenobarbitol 10 mg., hyoscyanus ext. 10 mg., Col. gluconate 100 mg. | Plessner | Hyalite, Tab. |
| 0.1% | Antipyrine 0.4%, borax, boric acid | Wyeth | Callorium w/Ephedrine, Sol. |
| (HCl) 1/4 gr. | Atropine sulfate 1/250 gr., chlorpheiramine maleate 12 mg. | Marsh-Emery | Histacan, Tab. |
| (Sulf.) 0.1% | Belladonna, bis oxyiod., bis. suberb., balsam Peru, boric acid | Wyeth | Wyenalds, Supp. |
| (Sulf.) 3/4 gr. | Belladonna fr. 0.5 cc., phenobarbitol 1/2 gr. | Haskell | Solphedriarb, Tab. |
| (Sulf.) 1/4 gr. | Belladonna 1/10 gr., phenobarbitol 1/2 gr., stramonium 1 gr. | Chemica | Acapnold, Tab. |
| (HCl) 1/16 gr. | Belladonna ext. 1/20 gr., salicylamide 1/4 gr. | Jenkins Labs | Dannafed Jr., Tab. |
| 0.125% | Benzocaine 4.5%, menthol 0.5%, phenol 1.75% | Malloe | Rectaligan, liq. |
| (HCl) 1/4 gr. | Benzocaine, Amm. chl., Col. creosote, Pot. guaiacofsulfate | Breco | Amanidin, Tab. |
| 0.5% | Benzocaine 0.5%, bis. tubert., 3.3%, acetanilid 1%, eucalypt., menthol | Abbott | Nasalene, No. 1, Oint. |
| 0.1% | Benzocaine 5%, menth 10% | Warner-Chilcote | Mucigel, Jel. |
| (HCl) 0.1% | Benzocaine 5%, Ichthammol 1%, oxyquinoline sulf. 1% | Medicone | Derma-Medicone, Oint. |
| 10 mg. | Carbaplenone tennote 60 mg., chlorpheiramine tennate 5 mg., phenylephrine tennate 10 mg. | Irwin, Neisler | Rynotus, Tabs. |
| (HCl) 3/4 gr. | Carbromal 5 gr. | Blue Line | Ephedramel, Cap. |

| Ephedrine | Others | Manufacturer | Brand |
|--|--|--|--|
| (Sulf) 1% 1% | Chlorbutanol, dextrose | Abbott | Isobalant, Sol |
| (HCl) 0.6% | Chlorbutanol, gluconic acid | Illy | I Sedrin Plain Sol |
| (HCl) 1% | Chlorbutanol 0.5%, menthol, cinnamon oil | Hart | Efedron Oint |
| (Sulf) 60 mg | Chlorbutanol 0.5%, menthol, eucalypt | Hart | Efemist, Liq |
| (Sulf) 12 mg | Chlorpheniramine maleate 12 mg, codeine phosphate 60 mg | Lemman | Forcold, Liq / oz |
| (Sulf) 1/4 gr / oz | Chlorpheniramine maleate 1 mg, glyceryl guaiacolate 50 mg, theophylline 15 mg, phenobarbital 4 mg | Breen | Brantotab Elix / 5 cc |
| (HCl) 5 mg | Cod phos 1 gr / oz, benzylate 7.47 min / oz, mannitol 45.3 gr / oz, sorbitol 180 gr / oz | Od Peacock | Ben Co Fed, Liq |
| (HCl) 1/4 gr (HCl) 25 mg | Oxizime/hesperon HBr 10 mg, chlorpheniramine maleate 2 mg, phenylephrine HCl 5 mg, w/ammonium chloride ipetoc | Abbott | Ouelidine Syr / 5 cc |
| (HCl) 0.5 gr | Diallylbarb 1/2 gr, siramamum ext 1/4 gr | Buffington | Phedrine Pilllet |
| (Sulf) 5 mg | Dihydrocodeinone bitartrate 10 mg, nembuatal sod 25 mg, calcium iodide 9/10 mg / A oz | Abbott | Colcidrine Syr |
| (HCl) 1/2 gr (Sulf) 12 mg | Dihydrocodeinone bitart 1/4 gr, Pot guaiacolsulf 8 gr, Sod cit 20 gr, citric acid 5 gr | Mallard | Codplex, Liq |
| (HCl) 3/4 gr (Sulf) 6.25 mg (HCl) 1/2 gr / oz (HCl) 1/4 gr, (Sulf) 12 mg | Dihydrocodeinone bitartrate 1.25 mg, pyrimidine maleate 10 mg, phenylpyrimine HCl 5 mg | Zemmer | Zem Histinone Syr / dr |
| (HCl) 25 mg, (Sulf) 0.3 gr (HCl) 3/4 gr | Glucophylline 2 1/2 gr, neonol 3/4 gr | Abbott | Phylladrine Cap |
| (HCl) 1/2 gr / oz | Hydroxizine HCl 2.5 mg, theophylline 32.5 mg | Zoering | Morax Syr / 5 cc |
| (HCl) 1/4 gr, (Sulf) 12 mg | Irish moss, glycenn | Folch | Ephedrine Glycerole, liq |
| (HCl) 25 mg, (Sulf) 0.3 gr (HCl) 3/4 gr | Labello, Col, iodabehen, sarsaparilla ext | Breen | Lobiodrin, Tab |
| (HCl) 1/2 gr | Luminal 6 mg, isupret 2.5 mg, theophylline 45 mg, & pot iodide 130 mg / 15 cc | Winthrop | Isupret Compound |
| (Sulf) 3/4 gr | Mephobarbital 25 mg, aminophylline 125 mg, int histin 25 mg | Receal | Asmosen, Tab |
| (HCl) 1/2 gr / oz, (HCl) 25 mg (Sulf) 1/4 gr | Methapyrimine HCl 3/4 gr, theophylline 1 1/4 gr | Mastengill | Theohist, Tab |
| (HCl) 1/2 gr | Methylbromotrapin sulf. 1/4 gr, buloben 1 mg, Sod brom 10.5 gr | Kohlenberg | X L Seo Rem, Vial |
| (Sulf) 1/2 gr | Morphine sulf 1/4 gr, fobeline inf 4 gr, Pot guaiacolsulf, Sod cit 20 gr, menthol | Mallard | Phedrocol, Liq |
| (HCl) 1/4 gr, (HCl) 1/2 gr / oz, (HCl) 25 mg (Sulf) 1/4 gr | Phenacetin 1 1/2 gr, Phenacetin 3 gr, phenobarbital 1/4 gr Pentobarbital Sod 1/2 gr / oz, Cal iod 8 gr / oz Phenobarbital 15 mg Phenobarbital 1/4 gr | Abbott Mallard Flint, Eaton Abbott Sherman | Dystoc Tab Phendrin, Cap Pentaphedrine, liq Ephetal Tab Ephedital, Tab |

Ephedrine Product Combinations

| Ephedrine | Others | Manufacturer | Brand |
|----------------|---|-----------------|---|
| (Sulf) | Phenobarbital | (Various Mfr.) | Ephedrine sulf. & Phenobarbital Cap. N.F. |
| ½ gr. | Phenobarbital ¼ gr., aminophylline 1½ gr. | Kremers Urban | Enafen, Tab. |
| (HCl) ¾ gr. | Phenobarbital ¼ gr. | Mallard | Ephobarb, Cap. |
| (HCl) ¾ gr. | Phenobarbital ¼ gr., atalaph. 3 gr. | Rorer | Pnedretal, Tab. |
| (Sulf) 24 mg. | Phenobarbital 8 mg., aminoph. 65 mg., pyrilamine maleate 12.5 mg. | Mallard | Phenadrin, Cap. |
| | | Roemer | Barbohisl, Tab. |
| (HCl) 26 mg. | Phenobarbital 26 mg., aspirin 130 mg., vit. C. 100 mg. | Arlington | Artcaps, Cap. |
| (Sulf) ¾ gr. | Phenobarbital ½ gr., belladonna tr. 0.5 cc. | Haskell | Belphegriobarb, Tab. |
| (Sulf) ¼ gr. | Phenobarbital ¾ gr., belladonna 1/10 gr., stramonium 1 gr. | Chemico | Acapnoid, Tab. |
| (Sulf) ½ gr. | Phenobarbital Sod. ½ gr., euphorbia pil. ext. 1½ gr. | Warren-Teed | Eu.Phed-lital, Tab. |
| (HCl) ½ gr. | Phenobarbital ½ gr., lobelia ext. ½ gr. | Cole | Barbeline, Tab. |
| (Sulf) ¼ gr. | Phenobarbital ½ gr., ergonidine 10 mg. | Wampole | Sedaryl, liq. |
| (Sulf) ¾ gr. | Phenobarbital ¼ gr., Pot. iod. 2½ gr., Col. lactate 2½ gr. | Lemman | Novelene, Tab. |
| ¾ gr. | Phenobarbital Sod. ¼ gr., Pot. iod. 2½ gr., Col. lactate 2½ gr., pyrilamine 20 mg. | Prof. Drug | Histo-Novolene, Tab. |
| (HCl) ½ gr. | Phenobarbital ½ gr., theobromine 5 gr. | Yenselt & Brown | Theobarb w/Ephed., Cap. |
| (Sulf) ½ gr. | Phenobarbital Sod. ½ gr., theophylline 2 gr. | Cole Chem. Co. | Asmilyn, Tab. |
| (Sulf) ½ gr. | Phenobarbit ½ gr., theophylline 2 gr. | Cole | Asmilyn, Tab. |
| (HCl) 48 mg. | Phenobarbital 25 mg., theophylline 180 mg | Warner-Chilkott | Tadrol-SA, Tab. |
| (HCl) ½ gr. | Phenobarbital ¼ gr., theophylline 1½ gr. | E. S. Miller | Thermostat, Tab. |
| (HCl) ¾ gr. | Phenobarbit ½ gr., theophylline 2 gr. | Chicago | Barbaphine, Tab. |
| ¼ gr. | Phenobarbital ½ gr., theophylline 1 gr. | Medicone | Meditabs., Tab. |
| (HCl) ¾ gr. | Phenobarbital ¾ gr., theophylline 1½ gr. | Ende | Teprin, Tab. |
| (HCl) 3/16 gr. | Phenobarbital 3/16 gr., theophylline calcium salicylate 1 gr., Pot. iodide 2½ gr. | Knoll | Quadrinol. Susp./5 cc. |
| (Sulf) ¾ gr. | Phenobarbital ¼ gr., theophylline sod. glycinate 5 gr. | First Texas | Glyozan-EP, Tab. |
| (Sulf) 10 mg. | Phenylpropanolamine HCl 10 mg., phenylephrine HCl 10 mg., N-acetyl p-amino-phenol 300 mg., chlorpropenylpyridamine maleate 4 mg. | Bernhart | Pyrelt, Tab. |
| (Sulf) 5 mg. | Phenyltolaxamine dihydrogen citrate 6.25 mg., glyceryl guaio-colate 50 mg., pyrilamine maleate 6.25 mg., meth-dia-mer sulfonamide 0.166 of each | Tilden | Sultustin, Susp./5 cc. |
| (HCl) 1/16 gr. | Pot. iod., quebracha tr. | McNeil | Brocodin, liq. |

Ephedrine Product Combinations

| Ephedrine | Others | Manufacturer | Brand |
|---|---|-----------------------------|-----------------------------|
| (Sulf) 3/8 gr | Prednisolone 0.5 mg, theophylline 2 gr., phenobarbital 1/8 gr | Vitamin | Prednisophine, Tab |
| (Sulf) 10 mg | Pyrimidine maleate 20 mg, aspirin 2 1/2 gr., acetoph 2 1/2 gr. | Haskell | Cofetrin, Tab |
| (free HCl) 1/12 gr | Pyrimidine maleate 1/5 gr., aspirin 4 gr., acetoph 2 gr | Central | Asphamol, Tab |
| (free HCl) 1/12 gr | Pyrimidine maleate 1/5 gr., aspirin 4 gr., acetoph 2 gr., Dover's pow 24 mg | Central | Asphamol D, Tab |
| (Sulf) 3/8 gr, (HCl) 5 mg | Pyrimidine maleate 25 mg | Harvey | Ephedremal, Tab. |
| (Sulf) 5 mg | Pyrimidine maleate 12.5 mg, amm chl 1 1/2 gr., caffeine cit rate 1/2 gr. | Am Ph Co | Fedrahist, Liq |
| (Sulf) 2.5 mg | Pyrimidine maleate 12.5 mg, Cal cresol sulfonate 150 mg, chloroform 5 mg | Schenley | Crephax, Liq |
| 3/8 gr | Pyrimidine maleate 3 125 mg, phenylethylamine dihydrogen citrate 3 125 mg, sulfo meth dia mar 83 mg, glyceryl guaia colate 25 mg | Tilden | Sulfussin, Tab |
| (HCl) 1/16 gr, (Sulf) 1% (HCl) 1/4 gr (Sulf) 1% (Sulf) 0.5% 0.5% (HCl) 1/6 gr, (HCl) 25 mg 25 mg 3/8 gr 24 mg | Pyrimidine 20 mg, phenobarbital 50d 1/2 gr., Pot iod 2 1/2 gr., Cal lactate 2 1/2 gr. Quebrocha tr., Pot iod Silver prot mild 5% Sironium bromide 24 gr., Hf 16 min., aspidosperma tr 240 min Sulfanilamide 6.25%, sulfathiazole 6.25% Sulfanilamide 2%, sulfathiazole 1%, allantoin 0.4% Tannic acid 1 1/2%, procaine buf 5%, phenol aum. Tartar emetic, cocifeno, senega, etc. Thenylene HCl 50 mg Theophylline, 130 mg; hydroxyzine HCl (Atrax), 10 mg /Tab Theophylline 2 gr., isoproterenol 5 mg., secobarbital 1/8 gr Theophylline, 100 mg; thenyldiamine HCl, 10 mg; glyceryl guaiacolate, 100 mg; phenobarbital, 2 mg /tablet Theophylline 130 mg phenobarbital 8 mg pow ext bello donna 5 mg, pyrimidine maleate 25 mg, Pot iodide 65 mg, magnesium trisilicate 130 mg | Prof. Drug | Hista Novaleine, Liq |
| (Sulf) 24 mg | Thyme ext | McNeil | Brocodin, Liq. |
| (Sulf) 1.3% | Tyrothricin 0.02%, chlorbutanol 0.5%, menthol 0.03% | Hart | Neo-Efemist, Liq |
| | | Natick | Aspidrine, Tab |
| | | Haskell | Desulfon Sulp w/Ephed., Liq |
| | | National Drug | Allantamide Liq w/Ephed. |
| | | Rarer | Butfrinonic, Oint. |
| | | Warren Teed | Elo Nol, Liq |
| | | Abbott | Thenyfred, Tab. |
| | | Roering | Marax |
| | | Hart | Aluted, Tab |
| | | Breen | Brontolab |
| | | Suffiff & Case | Theated, Tab |
| | | Am Ph Co, Suffiff & Case | Alko, Syr Korizol, Liq |

- pianomida) penicillanic acid See: I Maxi
 pen (Roerig & Co)
- EPI-PHIL CREAM & LOTION** (Greco) Non
 sensitizing oil of octa dodecanol in a
 sodium diacetate buffered washable base
 Cream 1 oz, tubes 4 oz jars 4 oz, 1 lb
 5 lb Lotions Bot 4 oz pts. 1/2 gal
 Use Dermatitis
- *EPIPROPIDINE** 1,1' di(2,3 epoxypropyl)
 4,4 bipiperidine See Eponate (Eli Lilly)
- EPIRENAN**. See Epinephrine (Various Mfr)
- EPISCORB INNALANT**. (Paschall) Rac epi
 nephrine 2%, protected by ascorbic acid
 Bot 1/4 oz, 1/2 oz, 1 oz
 Use: Symptomatic relief of bronchial asthma
- EPISCORB OINTMENT**. (Paschall) Rac epi
 nephrine 2.5%, ascorbic acid benzocaine
 1%, benzyl alcohol 1%, wintergreen oil in
 a base of carbowax Tube 1/4 oz, 1 oz
 Use: Allergic manifestations
- *EPITHIAZIDE** (Pfizer) 3 [(2,2,2 trifluoro
 ethylthio)methyl] 6 chloro 3,4 dihydro 2H 1,
 2,4 benzothiadiazine 7 sulfonamide 1, 1
 dioxide
 Use Hypotensive and diuretic
- EPITRATE**. (Ophthalmos) 1 Epinephrine bi
 tartrate 2% Drop Bot 7.5 cc
 Use Glaucoma
- EPONATE**. (Eli Lilly) Epi-propidine
 Use: Anticancer compound
- EPOXYAMINOXYTROPINE TROPATE HY-
 DROBROMIDE**. See Genoscapolamine NBr
 (Lobica Debrulle)
- W/Pentobarbital Sod, phenobarbital, ami
 naxytropine trapate HCl, procaine NCl, Vit
 B-6 & aluminum hydrate gel, dried See
 Prametic, Tab (Harrower)**
- EPOXYPROPIDINE**. (Lilly) Under study
 Use: Cancer therapy
- EPOXYTROPINE TROPATE METHYLBRO-
 MIDE** See: Methscapolamine Bromide (Var
 lous Mfr.)
- EPY**. (Barnes Hind) Stabilized free base
 complex of Epinephrine USP 1% in
 incorporated in a sterile isotonic ophthalmic
 vehicle with sodium bisulfite 0.3% and
 asine sulfate 0.01%/Bot 7.5 cc
 Use: Open angle glaucoma
- EPRAGEN** (Lilly) Acetaphenetidin 3 1/2 gr
 amylol 3/4 gr, ephedrine HCl 1/2 gr, ASA
 2 gr/Pulvule Bot 100s, 500s
 Use: (1-3 daily) Antispasmodic sedative &
 anti pyretic
- EPROLIN**. (Lilly) Alpha Tocopherol, Gelseals
 (50 mg, 100 mg) Bot 100s, 500s
 Use Vitamin E therapy
- EPSILAN** (Warren Teed) d Alpha Tocopherol
 acetate in a Amp [200 mg/cc] 1 cc &
 15 cc, Box 12s, 25s
- * Generic name recognized by N.F., U.S.P. or A.M.A.
- Use: 1 M; hyaluronidase inhibitor for col
 ogen diseases
- EPSILAN-M**. (Warren Teed) d Alpha Tocoph
 erol acetate Vit E Cap (100 I U) Bot 100s;
 (30 I U) Bot 100s, 1000s
 Use Orally; hyaluronidase inhibitor for col
 ogen diseases
- EPSILAN PHOSPHATE**. (Warren Teed) d
 Alpha tocopherol monosodium phosphate
 m eq sal Amp (100 mg) 2 cc Box 12s
 25s
 Use: 1 M; vitamin E therapy
- E-P-T AQUEOUS** (New Mexico Pharm)
 a Estradiol 12 mg testosterone 25 mg
 progesterone 25 mg, flimerosal 1 10 000/
 cc Vials 10 cc
- EPTOIN**. See: Diphenylhydantoin Sodium
 (Various Mfr.)
- EQUAGESIC**. (Wyeth) Meprobamate 150 mg,
 ethoheptozina citrate 75 mg, acetylsalicylic
 acid 250 mg/Tab Bot 50s
 Use Relief of pain accompanied by anxiety
 and tension
- EQUALYSEN**. (Wyeth) Meprobamate 200
 mg pentolinium tartrate 20 mg/Tab Bot
 50s
 Use Hypertension
- EQUANIL**, NND 1961 (Wyeth) Mepro
 bamate 2 Methyl-2 n propyl 1,3 propane
 dial dicarbamate Tab 200 mg, 400 mg
 Bot 50s Susp 200 mg/5 cc Bot 4 oz
 1 pt Wyseals 400 mg Bot 50s
 Use (1 lid) Sedative and tranquilizing
 agent
- EQUANIL SUSPENSION**. (Wyeth) Meprobo
 mate 200 mg, fruit flavored base/5 cc
 4 oz
 Use: Sedative and tranquilizing agent
- EQUANIL L-A**. (Wyeth Labs) Meprobamate
 400 mg/Cap Bot 50s
 Use: Relieve anxiety and tension
- TOUANITRATE**. (Wyeth) Meprobamate 200
 mg, pentaerythritol tetranitrate 10 mg or
 20 mg/Tab Bot 50s
 Use: Prophylaxis of angina pectoris
- EQUINE ENCEPHALOMYELITIS (EASTERN)
 DIAONOSTIC ANTIGEN**. (Lederle) 5 cc
 vial 20 tests Nat for human administra
 tion—diagnostic agent only
 Uses: For the complement fixation test ap
 plicable to the detection of antibodies et
 on aid in the diagnosis
- EQUINE ESTROGENS W/Lutrexin** See: Tres
 Inest Tab (Hynson, Wescott, & Dunning)
- EQUINE GONADOTROPIN**. See: Gonadin,
 Vial (Cutter)
- EQUINE PITUITARY GONADOTROPIC
 HORMONE**. See: APG Horm, Inj (Canfield)
- EQUINEX**. (Ayerst) Highly purified pregnant

mores' serum gonadotropin. One cubic centimeter containing 5000 I.U. in dry form and one 10-cc. vial sterile diluent containing 0.5% phenol.

Use I.M., I.V.; hypo-ovarianism & testicular hypofunction.

EQUIPHYSIN. (G. F. Horvey) Anterior pituitary gonadotropin 125 fevald-hisaw e./Vial. Vial 5 cc.

Use: Gonadal hypofunction

ERASEN. (Amfre-Grant) Phenobarbital 16 mg, Vit B-1 5 mg./Tab. Bot. 100s

Use: Sedative w/Vit. B-1 therapy

ERCAFITAL. (Blue line) Ergotamine tartrate 0.5 mg., caffeine 50 mg., phenobarbital 7.5 mg./Tab. Bot. 100s, 1000s

Use: Cranial artery vasoconstrictor

ERGOPIOL W/Savin. (M. H. Smith) 0.65 Gm. pow. ergot ext, 1/2 min. savin oil, 3 min. distilled opiol (M. H. S. Special) in castor oil/cap. Pkg. 20 Cap.

Use: Uterine stimulant.

ERGOCALCIFEROL, N N D. 1961.

See: Calciferol, U.S.P.

ERGOCORNINE. (Various Mfr.) Ergot alkaloids.

Use: Peripheral vascular disorders.

ERGOCRISTINE. (Various Mfr.) Ergot alkaloid

Use: Vascular disorders.

ERGOCRYPTINE. (Various Mfr.) Ergot alkaloid.

Use: Peripheral vascular disorders.

ERGOMAR. (Nordson) Ergotamine tartrate 2 mg./Tab. Pkg. 12s.

Use: Migraine and vascular headaches.

ERGOMETRINE MALEATE, U.S.P., N N D 1961. See: Ergonovine (Various Mfr.)

ERGONOVINE. (Various Mfr.) Ergobasine, eroklinine, ergometrine, ergostetrine, ergotocine. See: Ergonovine Maleate.

ERGONOVINE MALEATE, U.S.P., N N D. 1961; Inj., U.S.P.; Tab., U.S.P.

See: Ergotole Maleate, N.N.R. (Billy).

Margovine (Thompson)

(Burroughs Wellcome) Amp. (0.2 mg) 1 cc., Box 100s; Tab. (0.2 mg) Bot. 500s

Use: I.V., I.M., orally, oxytocic.

W/Chloralhydratase. See Tace W/Ergonovine, Cap. (Merrell)

W/Ergotaxine ethanesulfonate. See Ergotora, Tab. (Upjohn).

ERGONOVINE TARTRATE, N N R. 1950

See: Basergin, Amp., Tab. (Sandoz).

W/Ergotamine tartrate. See Neo-Gynergen, Amp., Tab. (Sandoz).

ERGOPHENE. (Upjohn) Ergot ext. 1D 1/2 gr., phenol & zinc oxide/oz. Oint. Tube 1/2 oz., 1 oz. Jar 1 lb.

Use: Topically; antiseptic.

* Generic name recognized by N.F., U.S.P. or A.M.A.

ERGOSTEROL, ACTIVATED OR IRRADIATED. See: Calciferol, U.S.P.

ERGOSTERON W/Sodium salicylate. (Aldrich)

Vit. D 50,000 u., Sod. salicylate 5 gr./Cap. Bot. 100s.

Use: (I.I.D.) Antiarthritic.

ERGOSTETRINE. See: Ergonovine. (Various Mfr.)

***ERGOTAMINE TARTRATE, U.S.P., N N R.** 1950; Inj., U.S.P.; Tab., U.S.P. (Various Mfr.) (Femergin).

See: Gynergen, Amp., Tab. (Sandoz)

Ergomar, Tab. (Nordson)

Medihaler-Ergotamine (Riker)

W/Belladonna alkaloids, acetylphenethidin & caffeine. See: Wigraine, Tab. (Organon).

W/Belladonna alk. & phenobarbital. See: Bellergot, Tab. (Sandoz).

W/Caffeine, N N R.

See: Cafetan, Tab. (Vitamine)

Cofergot, Supp. & Tab. (Sandoz)

Cofetrate, Tab. (Vitomin)

W/Caffeine, chloral hydrate, scopolamine aminoxide HBr. See: Rectoloid Migraine, Dis. Applicators (Wampole Labs)

W/Caffeine, homotropine methylbromide.

See: Ergotofropin (Cole)

W/Caffeine, hyoscyamine sulfate & pentobarbital. See: Pentergot, Rectol insert. (Bentex Pharm.).

W/Caffeine, phenobarbital. See: Ergofito, Tab (Blue line)

W/Cyclizine HCl, caffeine.

See: Migrol Tab. (Burroughs Wellcome)

W/Ergonovine tartrate. See: Neo-Gynergen, Amp., Tab. (Sandoz).

W/Mareline, caffeine See: Migrol Forte, Tabs (Burroughs Wellcome)

ERGOTATROPIN. (Cole) Ergotamine tartrate 1 mg., caffeine 100 mg., homotropine methylbromide 1 mg./Tab. Bot. 100s.

Use: Migraine.

ERGOT EXTRACT.

W/Castor oil. See: Ergoval. (Mottin Smith)

W/Phenol & zinc oxide. See: Ergophene, Oint. (Upjohn).

W/Savin oil & opiol. See: Ergopiol w/Savin. (C. D. Smith).

ERGOT FLUID EXTRACT, N.F. (Merck Sharp & Dohme) 1 cc. is equiv. to 1 Gm. of ergot (alcohol 40%). Bot. 4 oz., 1 pt.

ERGOTIDINE. See: Histamine (Various Mfr.)

ERGOTININE CITRATE. See: Vasomotor Disorders.

ERGOTOCINE. See: Ergonovine (Various Mfr.)

ERGOTOLE. (Merck Sharp & Dohme) Aq ergot ext. Bot. 1 oz.

Use: To stimulate uterine contractions.

- ERGOTORA.** (Upjohn) Ergotaxine ethane sulfonate 12 mg., ergonovine maleate 0.15 mg/Tab Bot 250s, 1000s
Use Oxytocic
- ERGOTOXINE ETNANESULFONATE.** (Burroughs Wellcome) Pow 0.1 Gm Reference Standard
W/Ergonovine maleate See: Ergotora, Tab (Upjohn)
- ERGOTOXINE, NISTAMINE, TYRAMINE**
See Ergutin, Amp, Liq (Burroughs Wellcome)
- ERGOTRATE.** See Ergonovine (Various Mfr)
- ERGOTRATE-N.** See: Ergonovine (Various Mfr)
- ERGOTRATE MALEATE, NND 1961** (Lilly)
Ergonovine maleate, USP Amp (0.2 mg) 1 cc, Pkg 6s, 100s; Tab (1/320 gr/tab) Bot 25s, 100s, 500s, 1000s
Use: 1 V, 1 M, orally; oxytocic
- ERGOT-RELATED PRODUCTS.**
See: Basergin, Amp (Sandoz)
Catergat, Supp, Tab (Sandoz)
Dihydroergotamine, Amp (Sandoz)
Ergoaplof, Cap (M H Smith)
Ergonovine (Various Mfr)
Ergot Aseptic, Amp (Parke, Davis)
Ergotamine (Various Mfr)
Ergotora, Tab (Upjohn)
Ergotrate (Various Mfr)
Ergoval, Cap (Martin Smith)
Oxyergen, Amp, Tab (Sandoz)
Hydergine, Amp, Tab (Sandoz)
Methergine, Amp, Tab (Sandoz)
Neo Gynergen, Amp, Tab (Sandoz)
Nigraine, Supp & Tab (Organon)
- ERGOVAL.** (Martin H Smith) Ergot ext 125 mg in castor oil 8 min/ Cap Bot 12s
Use (1 2 1 d) Ergot therapy
- ERIODICTIN** See Vitamin P & Rutin
- ERIODICTYON, NF, NNR 1937; Flexl, NF; aromatic syrup, NF** See: Vitamin P & Rutin
- ERIDIONE, NND 1962** (Eric, Kirk & Gray)
Phenindione 50 mg/Tab Bot 100s, 1000s
Use Anticoagulant
- ERION** See: Quinacrine Hydrochloride (Ata brine Hydrochloride) (Winthrop Labs)
- E-R-O** (First Texas) Antipyrine 19 gr., anesthesin 5 gr., glycerin q s/fl oz Bot w/ dropper 1/2 oz
Use Pain in simple earache
- EROUA.** (Fellows) Sod salicylate 5 gr., merodiane (Vit K analogue) 0.125 mg/ E.C. Tab & double strength Tab, Bot 100s, 1000s; Liq, Bot 4 fl oz
Use: (1-4 daily) Antipyretic & analgesic
* Generic name recognized by NF, USP or A.M.A.
- ERTRON.** (Whittier) Activated ergosterol (Vit D) Cap (50,000 u) Bot 50s, 100s, 500s; Amp (500,000 u/cc) Box 6s
Use: Orally, 1 M; vitamin D therapy
- ERTRON S-M, Cap.** (Whittier) Vit D 50,000 u, salicylamide 162 mg., mephensazin 125 mg/ Cap Bot 100s
Use: Analgesic, relaxant
- ERUCYTE.** (Smith Miller & Patch) Ferrous fumarate 200 mg., ascorbic acid 125 mg., Vit B 12 2 mcg/ Tab Bot 100s
Use Iron deficiency anemia
- ERUCYTE NEOTABS.** (Smith, Miller & Patch)
Ferrous fumarate 100 mg., ascorbic acid 50 mg., Vit B 12 5 mcg/ Tab Bot 100s, 1000s.
Use: Iron deficiency anemia
- ERYCYTOL.** See: Vitamin B 12
- ERYSIPELAS VACCINE.** (Sherman) Streptococcus erysipalatis 300 million, Staphylococcus albus 500 million/cc Vial 12.5 cc, 20 cc See: Coley's Mixture (Parke, Davis)
Use: Prophylaxis & treatment of erysipelas
- *ERYTHRITYL TETRANITRATE, Tab, NF X; NNR 1950** (Various Mfr) Erythritol erythrol tetranitrate, nitraerythritol, tetranidin, tetranitral
See: Cordilate, Tab (Burroughs Wellcome)
Use (1 2 daily) Coronary vasodilator
W/Sod nitrite, mannitol nitrate, ammonium hippurate See: Vasodilator Compound Tab (Burroughs Wellcome)
- ERYTHROCIN.** (Abbott) Erythromycin Tab film sealed (0.2 Gm & 0.2 Gm) Bot 25s, 100s; Oint (1%) Tube 1 oz
Use: Against gram positive & certain gram negative organisms
W/Sulfis See: Erythromid, Fimtabs (Abbott)
- ERYTHROCIN-I.M. IN ABBJECT.** (Abbott)
Erythromycin ethyl succinate 50 mg dissolved in polyethylene glycol with butamben 2% (Butessin Abbott), also 100 mg/ml Abject disposable syringe w/needle Box 5s, 25s, 100s
Use: Infections caused by organisms sensitive to erythromycin
- ERYTHROCIN ETNYL SUCCINATE** (Abbott) (As ester) (50 mg/cc) Amp 2 cc, 10 cc
Use: 1 M Erythromycin
- ERYTHROCIN ETHYLSUCCINATE SUSP.** (Abbott) Erythromycin ethylsuccinate 200 mg/ 5 cc Oral pediatric Susp Bot 60 cc 1 pt
Use: Infections due to erythromycin sensitive organisms
- *ERYTHROCIN LACTOBIONATE, NND 1962** (Abbott) Erythromycin lactobionate (Equiv to 1 Gm of erythromycin) Vial 30 cc; also Vial (equiv to 300 mg of erythromycin) 10 cc soln

- Use: I.V., I.M., organisms sensitive to erythromycin.
- ERYTHROCIN-NEOMYCIN OINT.** (Abbott)
Erythromycin 10 mg., neomycin sulfate 5 mg./Gm. Tube ½ oz.
Use: Dermatitis, impetigo, furunculosis, granuloma pyogenicum & acne.
- ERYTHROCIN PEDIATRIC**, N.N.D. 1962 (Abbott) Erythromycin stearate equiv. to erythromycin 100 mg / 5 cc. Bot. 75 cc. late salt of the propionic acid ester of erythromycin. See: Ilosone (Eli Lilly)
- ERYTHROCIN STEARATE**, N.N.D. 1961. (Abbott) Oral susp. (100 mg / 5 cc. cinn. flavor) Bot. 75 cc, Tab. {100 & 250 mg} Bot. 25 & 100s.
- ERYTHROCIN STEARATE w/Sulfo**,
See: Erythramid, Tab. (Abbott)
- ERYTHROGRAN**, (Upjohn) Protection coated erythromycin 50 mg./packet. Pkg. 18s.
Use: Infections susceptible to erythromycin.
- ERYTHROGUENT OINTMENT**, (Upjohn) Erythromycin 5 mg., methylparaben 0.2 mg., butyl-2-hydroxybenzoate 1.8 mg /Gm. Tube ½ oz.
Use: Against gram-positive organisms.
- ERYTHROMID**, (Abbott) Erythrocin 75 mg., sulfadiazine sod., sulfamerazine sod., and sulfamethazine of each 111 mg./Pilltab. Bot. 25s, 100s
- ERYTHRO-MYCIQUENT**, (Upjohn) Erythromycin 5 mg., neomycin sulfate 5 mg., methylparaben 0.2 mg., butyl-p-hydroxybenzoate 1.8 mg./Gm. Tube ½ oz.
Use: Erythromycin & neomycin therapy.
- ***ERYTHROMYCIN**, U.S.P., N.N.D. 1962. A.D.R. An antibiotic from *Streptomyces erythreus* Inj., oral Susp. Tab., U.S.P. (Upjohn) Tab. 100 mg., Bot. 25s, 250 mg peach color, Bot. 25s, 100s.
See: Erythrocin, N.N.R. (Abbott) Tab. 100 mg. Erythrogran, Granule (Upjohn). Erythrogent, Oint. (Upjohn). Ilotycin, N.N.D. (Lilly) Tab. 100 mg
- W/Digoxinocaine. See: Ilotycin I.M. Solution. (Lilly)
- W/Meth-Dia-Mer sulfonamides. See: Erythrosulfa, Granules (Upjohn)
- W/Methylparaben, butyl-2-hydroxybenzoate.
See: Erythrogent, Oint. (Upjohn).
- W/Neomycin sulfate, methylparaben, butyl-p-hydroxybenzoate. See: Erythro-Myciquent, Oint. (Upjohn).
- W/Neomycin sulfate. See: Erythrocin Neomycin, Oint. (Abbott).
- ***ERYTHROMYCIN ESTOLATE**, N.N.D. 1962 Lauryl sulfate salt of the propionic acid ester of erythromycin See: Ilosone (Eli Lilly)
- * Generic name recognized by N.F., U.S.P. or A.M.A.
- ***ERYTHROMYCIN ETHYL CARRONATE**, U.S.P., N.N.D. 1962. See: Ilotycin Pediatric (Lilly)
- ***ERYTHROMYCIN ETHYL SUCCINATE**, U.S.P. See: Erythrocin Ethyl Succinate (Abbott)
- ***ERYTHROMYCIN GLUCOHEPTONATE**, U.S.P., N.N.D. 1962. See: Ilotycin Glucoheptonate, Amp. I.V. (Lilly)
- W/Polymyxin-B sulfate & benzocaine. See: Ilotycin Otic (Lilly).
- ***ERYTHROMYCIN LACTOBIONATE**, U.S.P., N.N.D. 1962. See: Erythrocin Lactobionate (Abbott)
- ***ERYTHROMYCIN PROPIONATE**, U.S.P., Cop. U.S.P.
- ERYTHROMYCIN PROPIONATE LAURYL SULFATE**, N.N.D. 1962. Erythromycin Estralate. See: Ilosone.
- ***ERYTHROMYCIN STEARATE**, U.S.P., N.N.D. 1962, A.D.R. See: Erythrocin Pediatric, Susp & Tab (Abbott)
- W/Meth-Dia-Mer sulfonamides. See: Erythrocin Sulfas, Tab. (Abbott).
- ERYTHROSULFA**, (Upjohn) Erythromycin 100 mg., sulfadiazine, sulfamerazine & sulfamethazine 0083 Gm. each/Tab Bot. 50s, 500s
Use: (1-2 q. 4 h.) Against gram-positive bacteria.
- ERYTHROSULFA GRANULES**, (Upjohn) Erythromycin 50 mg., sulfamethazine, sulfadiazine, sulfamerazine of each 83.3 mg./pkg. Granule pkg. 18s.
Use: Antibiotic & triple sulfonamide therapy.
- ERYTHROTIN**, See: Vitamin B-12.
- E5-A-CORT**. (Dome) Vit. A 100,000 u., estrone 56,000 u., hydrocortisone 5 mg or 10 mg./Gm. CREME Tube ½ oz. Lot Bot. ½s, 2 oz.
Use: Senile vulvovaginitis, anogenital pruritus.
- E5-A-CREME**. (Dome) Estrone 56,000 u., Vit. A 100M u./oz. pH 5. Creme 1 oz., Jar 1 lb.
Use: Topically estrogen therapy.
- E5-A-MYCIN CREME**. (Dome) Neomycin base 100 mg., estrone 56,000 u., Vit. A 100,000 u./oz. Tube 1, 2, 4 oz. Jar 1 lb.
Use: Treatment of acne infections.
- ESCHATIN**. (Parke, Davis) 200 mcg. Cortisone/cc. Vial 10 cc., 50 cc.
Use: Cortical hormone insufficiency.
- ESDAVITE PEARLS**. (Merck Sharp & Dohme) Vit. A palmitate 5000 u., D 1000 u., thiamine HCl 1.5 mg., riboflavin 2 mg., ascorbic acid 37.5 mg., niacinamide 10 mg., calc. pantothenate 1 mg., vit. B-12 *cryst.* 1 mcg., pyridoxine HCl 0.25 mg., w/choline di-

- hydrogen citrate 10 mg/pearl Bot 100s
250s, 1000s
Use Dietary supplement
- ESOOHE** (Lloyd, Dobney and Westerfield)
Dienestrol 0.25 mg methyltestosterone 5
mg/Tab Bot 100s Vial 10 cc Caps TD
Bot 100s
Use Androgen estrogen therapy in
geriatrics
- ESERIHE** (physostigmine as alkaloid, salicyl
ate or sulfate salt) See: Physostigmine salt
- ESERIHE AMIHOXIDE.** See Geneserine (Ha
Mfr currently lists)
- ESIORIX, NNO 1962** (Ciba) Hydrochloro
thiazide 25 mg or 50 mg/tab Bot 100s,
1000s
Use Diuretic and antihypertensive
W/Apresaline See Apresaline Esidrix (Ciba)
- ESIORIX-K.** (Ciba) Hydrochlorothiazide 25
mg potassium chloride 500 mg/Tab Bot
100s Also Esidrix 50 mg, pot chloride
1000 mg/Tab Bot 100s
Use Diuretic hypotensive
- ESIDRIX-SERPASIL** (Ciba) Esidrix 25 mg,
serpasil 0.1 mg/Tab Bot 100s
Use Control of high blood pressure
- ESJAYDIOL** (Testogor) Methylandrostenodial
(methandriol) Vial (25 mg, 50 mg/cc)
Susp 10 cc
Use: (IM 10-40 mg daily) Anabolic nitro-
gen retention
- ESKABARB.** (Smith, Kline & French) Pheno
barbital Spansule (1 gr & 1½ gr) Bot 30s
250s
Use: Sustained action sedative
- ESKACILLIN** (Smith, Kline & French) Esku
cillin 100, Penicillin O Pot 100M u/5 cc
Bot 2 fl oz, Eskacillin 250 Penicillin G
Procaine 250M u/5 cc Bot 2 fl oz
Use: Orally; penicillin sensitive organisms
- ESKACILLIN-100-SULFAS.** (Smith, Kline &
French) Penicillin G Pot 100M u, sulfa
diazine 0.167 Gm, sulfamerazine 0.167
Gm, sulfamethazine 0.167 Gm/5 cc Bot
2 fl oz
Use: Orally; penicillin with triple sulfona
mide therapy
- ESKACILLIN-250-SULFAS** (Smith Kline &
French) Penicillin G procaine 250 000 u
w/sulfadiazine, sulfamerazine sulfametha
zine 0.167 Gm of each/5 cc Bot 2 fl oz
Use: (1-2 tsp 11 d) Penicillin & sulfonamide
therapy
- ESKADIAZINE.** (Smith, Kline & French) Sulta
diazine 0.5 Gm/5 cc Susp. Bot 12 fl oz
Use: Orally; sulfadiazine therapy
- ESKAPHEN B ELIXIR & TABLETS.** (Smith
Kline & French) Each 5 cc of the elixir &
each tab contains phenobarbital ¼ gr &
- thiamine HCl 5 mg Elixir 12 fl oz Bot.
Tab, Bot 50s
Use 12-6 Tab or tsp daily) For tense
nervous patients
- ESKAR.** (Merck Sharp & Dahme) Trichlor
acetate acid ointment Jar ¼ oz
Use For removal of warts & corns
- ESKASERP** (Smith Kline & French) Eser
pine Spansule 0.25 mg & 0.50 mg Bot
30s 250s
Use Sedative & tranquilizing agent
- ESKATROL** (Smith, Kline & French) Oexo
amphetamine sulfate 15 mg, prochlorper
azine as the dimaleate 7.5 mg/spanule
Bot 30s, 250s
Use Treatment of overweight
- ESKAY'S "NEURO PHOSPHATES"** (Smith
Kline & French) Strychnine glycerophosphate
1/64 gr, sod glycerophosphate 2 gr, cal
glycerophosphate 2 gr, phosphoric acid
75% 17 min, alcohol 10%/2 fl dr Bot
12 fl oz
Uses Tonic
- "ESKAY'S PENTAPLEX,"** (Smith, Kline &
French) Vit B 1 225 mg, B 2 2 mg, B 6
0.5 mg, niacin 20 mg, pantothenic acid
15 mg/6 fl dr Bot 12 oz
Use Orally (2 dr 11 d) Vitamin B complex
deficiency
- ESOMID CHLORIDE, NND 1959** (Ciba)
Hexamethylene bis trimethylammonium di
chloride, hexamethonium chloride Tab
(250 mg) Bot 100s, Syr each tsp 250 mg/
4 cc Bot 1 pt
Use Indicated in hypertension
- ESOPRIHE.** (Ulmer) Physostigmine salicylate
1 mg & homatropine methylbromide 2
mg/cc or Tab Vials 10 cc Bot 100s &
500s
Use: (IM 1 cc or 2 Tabs) Muscle spasm
in arthritis
- ESORB.** (Wyeth) d Alpha tocopherol 50 mg
sorbitan fatty acid ester derivative 0.226
Gm/Cap Bot 100s
Use Vitamin E therapy
- ESPIHOL.** (Lloyd) di Amphetamine phosphate
1 mg, atropine methyl nitrate 1 mg, as
pirin 3½ gr, acetophenetidin 2½ gr/Tab
Bot 100s 500s 1000s
Use (1-3 11 d) Analgesic w/antispasmodic
- ESPROHATE.** (Chicago Pharmacal) Testos
terone propionate 25 mg, progesterone 25
mg, alpha estradiol 0.28 mg (equiv to
not less than 200 000 IU/1 cc) Vial 10 cc
Use Functional uterine bleeding
- ESPRORONE.** (Chicago Pharmacal) Beta
estradiol 20,000 u, progesterone 10 mg/
cc Vial 10 cc In oil 30 cc
Use: Estrogen & corpus luteum therapy
- ESROMIOTIN.** (Crookes Barnes) Stabilized

solution of physostigmine salicylate from $\frac{1}{2}$ to $\frac{1}{2}$ %. Vials 7.5 cc. and dropperettes.

Use: Miotic

ESSENAMINE. (Winthrop Labs) Protein derived from lactalbumin. Pkg. 7½ oz., 14 oz., 5 lb., 25 lb.

Use: Protein therapy.

ESTAN. (White) Dienestrol 0.25 mg., methyltestosterone 5 mg./Tab. Bot. 30s, 100s.

Use: Orally; active estrogen & androgen therapy.

ESTEED. (Warren-Teed) Ethinyl estradiol 0.05 mg./Tab. Bot. 30s, 100s, 500s.

Use: Menopause.

W/Methyltestosterone. See: Testeed, Tab. (Warren-Teed).

ESTERCOL. (Toash) Mercury oleate 0.45%, phenol 0.5%, salicylic acid 3%, coal tar crude 1.2%. Emulsion, Bot. 4 oz.

Use: Dermatitis.

ESTERIL. (Ensley-Scott) Prednisone 25 mg., phenylalanine dihydrogen citrate 12.5 mg., chlorpheniramine maleate 1 mg., ephedrine sulfate 12.5 mg./Tab. Bot. 50s.

Use: Treatment of allergic disorders.

ESTILBEN. See Diethylstilbestrol Dipropionate. (Various Mfrs.)

ESTINYL. N.N.D. 1962. (Schering) Ethinyl estradiol, Tab., coated 100 μ g. & 0.05 mg.) Bot. 100s, 250s, 1000s; Tab. (0.5 mg.) Bot. 30s, 100s.

Use: Orally; estrogen therapy.

ESTIVIN. (Schieffelin) A specially processed aq. infusion of Rosa gallica (rose petals). Dropak 0.25 fl. oz. plastic container; Bot. w/dropper 0.25 fl. oz.

Use: To reduce congestion of lacrimal caruncle glands.

ESTOLATE. Erythromycin See: Erythromycin propionate Lauryl Sulfate.

ESTOMUL Tabs. & liquid, (Riker) Orphenadrine HCl 25 mg., bismuth aluminate 25 mg., magnesium oxide 45 mg., aluminum hydroxide and magnesium carbonate as a co precipitate 500 mg./Tab. Bot. Liq. 12 oz. bottles. Tabs. 100s.

Use: Treatment of peptic ulcer.

ESTOPEN. See: Benzylpenicillin 2-diethylaminoethyl ester Hl

ESTOPHEROL. (Pitman-Moore) Estrogens D 625 mg., d tocopherol acetate 10 mg., phenobarbital 5.4 mg./Tab. Bot. 100s, 1000s.

Use: Menopausal symptoms

ESTORALS. (Key Corp.) Ethinyl estradiol 0.02 mg./Tab. Bot. 100s

ESTRACORP. (Sutliff & Case) Ethinyl estradiol 0.025 mg., methamphetamine HCl 40 mg., d amphetamine sulfate 2.0 mg., pentobarbital eod. 30.0 mg., Vit. C 50.0 mg., B-1 1.0 mg., B-2 1.0 mg., B-6 0.1 mg., Vit. B₁₂ 0.2 mcgm., Cal. pantothenate 1.0 mg., niacinamide 5.0 mg./Cap. Bot. 100s, 1000s

ESTRA-D. (New Mexico Pharm) Alpha estradiol (100,000 IU) 1 mg., carboxymethyl cellulose 0.2%, Tween 80 0.4%, benzyl alcohol 1.5%/cc. Liq. Vials 30 cc.

ESTRADIOL. N.F., N.N.O. 1962. Aq. Susp., N.F.; Pellets, N.F.; Tab., N.F. Beta-estradiol, agofollin, dihydroxyestrin, dihydrotheelin, gynergan, gynaestryl. The form now known to be physiologically active is the "beta" form rather than the "alpha" that was designated in U.S.P.

Use: Estrogen-replacement therapy.

See: Almedial, Tab., Vial (Meyer).

Altrad, Vial, Aq. (Ascher).

Aquadiol, Vial, Aq. (National Drug).

Aquagen, Vial, Aq. (Lemmon).

Aqual, Vial, Aq. (Estro).

Baridol, Aq. (Barry)

Dimenforman, Vial, Aq., Tab., N.N.R. (Organon)

Diogyn, Vial, Aq., N.N.R. (Pfizer)

Diogynets, Tab. (Pfizer).

Estraldine, Vial, Aq. (Rorer).

Femestral, Vial, Aq. (Kirk).

Femogen, Aq. (Testagor)

Macrodiol, Vial, Aq. (Kirk & Gotham).

Micsodiol, Vial, Aq. (Gold Leaf).

Nardiol, Buccal. (Normand)

Ovacylin, Tab., N.N.R. (Ciba)

Pregynon, Vial Aq., Pellets, Tab., N.N.R. (Schering)

Syndiol, Vial, Aq. (Central).

W/Estrogenic substance, propylene glycol, benzyl alcohol, chlorbutanol. See: Mikrogen, Vial (Durs).

W/Estrone.

See: Estragenol, Amp. & Vial in oil (Chemical).

Met-Estrin, Amp. in oil (Metro)

W/Estrone, benzyl alcohol in peanut oil.

See: Ovadiol, Vial (C. D. Smith)

W/Estrone & progesterone in oil.

See: Almestrone, Vial (Meyer).

Pracycla, Vial (Metro).

W/Estrone, progesterone, benzyl alcohol in peanut oil. See: Ovagest, Vial (C. D. Smith)

W/Methyltestosterone.

See: Di-Met, Tab. (Organon).

Gynetone, Tab., Vial (Schering).

Testaldiol, Tab. (Meyer)

W/Methyltestosterone, pregnenolone, monobasic amphetamine phosphate. See: Longeon, Cap. (Northwest Labs.).

W/Methyltestosterone & Vit. B complex. See: Combogenic, Tab. (C. D. Smith)

* Generic name recognized by N.F., U.S.P. or A.M.A.

W/Procaine HCl See Almadial, Vial, Aq, Tab (Meyer)

W/Progesterone

Aq Susp

See Aquapriol, Vial (Estra)

Eiprone, Vial (Chicago Pharmacol)

Myasol Tergesten (Ulmer)

Neo Femotrone, Vial (Testagar)

Prastriadiol Vial (Kirk)

In Oil

See Corpa Gen, Vial (Barry)

W/Progesterone, methyltestosterone See: Trihormonal, Tab (Key Corp)

W/Progesterone & testosterone aq susp See Tergesten Vial (Ulmer)

Triganaq Vial (Central Pharm)

Triharm, Vial (Ascher)

Tri Hormogen, Vial (Mallord)

W/Progesterone testosterone procaine HCl, procaine base See Hormo Triad, Vial (Bell Pharmacol)

W/Progesterone, testosterone propionate See Trigestondriol, Vial (Kirk)

W/Testosterone

See: Aquadiol w/Testosterone, Vial, Aq (National Drug)

Combondrets Tab (Pfizer)

Testaldol, Vial, Aq (Meyer)

W/Testosterone & chlorbutanol in cotton seed oil See Depo Testadiol, Vial (Upjohn)

W/Testosterone procaine HCl, aq susp See: Testregyn Vial (Ascher)

W/Testosterone propionate in oil See: Bi Ger Mona, Vial (Kirk)

Testriol, Vial (G F Horvey)

W/Thiamine niacin Vit B₆, riboflavin & Cal pantothenate See: Estrabex, Tab (C D Smith)

*ESTRADIOL BENZOATE, USP NND 1961 Inj USP (Various Mfr) Beta estradiol benzoate, benzogynaestril benzostrolal ditallucin, tallicarman, tallidrin See:

In Oil

Dimenlarman benzoate, NNR (Organon)

Diogyn B, Vial, NNR (Pfizer)

Estrabev E, Vial (Key Corp)

Ovocyclin Benzoate, Vial, NNR (Ciba)

Progynan B, Vial (Schering)

Saleitra, Vial (Merck Sharp & Dahme)

W/D ethylthiobital See: Nu Estro, Tabs (Gately)

W/Progesterone in oil

See: Di Pro, Vial (Organon)

Fasigyn, Vial (Pfizer)

Pro-Estrabev E, Vial (Key Corp)

Promestron, Vial (Schering)

W/Testosterone propionate

* Generic name recognized by N.P., U.S.P. or A.M.A.

Aq Susp.

See: Estrabev E, Vial (Key Corp)

Testradiol, Vial (Metro)

In Oil

See Cambandrin, Vial (Pfizer)

Di Met BP, Vial (Organon)

Gynstans, Vial (Schering)

Mal O Fem, Vial (Testagar)

Testadiol, Vial (Metro)

W/Testosterone Propionate in Oil See Estromol Impravad, Vial (Key Corp)

W/Testosterone propionate, progesterone See Pro Mal O Fem, In Oil (Testagar)

Trihormonal, Vial (Key Corp)

*ESTRADIOL CYCLOPENTYLPROPIONATE, NF NND 1962 Inj NF Estradiol 17 beta(3 cyclopentyl)propionate See Depo-estradiol cyclopentylpropionate Vial (Upjohn)

W/Testosterone cyclopentylpropionate See Depo Testadiol (Upjohn)

*ESTRADIOL DIPROPIONATE, U.S.P., NND 1962 Inj U.S.P. Ovocyclin D propionate, NND 1958 (Ciba)

Progynon D P (Schering)

Oil Inj, U.S.P.

See: Dimenformon Dipropionate (Organon)

ESTRADIOL, ETHINYL, NND 1962 See Ethinyl Estradiol

ESTRADIOL MONOBENZOATE, NND 1960 See Estradiol Benzoate

ESTRADIOL PHOSPHATE, See Estrodurin Amp (Ayerst)

ESTRADIOL, TRIMETHYL ACETIC ACID ESTER, See: Estrate, Vial (Lakeside)

W/Procaine HCl, propylene glycol aqueous diluent See Femestron AD (Moore Kk)

W/Progesterone See: Estrolate w/progesterone Vial (Lakeside)

*ESTRADIOL VALERATE, NND 1962 See: Defestrogen Vial (Squibb)

Lastrogen (Key)

Reposa E Vial (Canfield)

W/Hydroxy progesterone caproate

See Deluteval, Amp (Squibb)

W/Hydroxyprogesterone caproate See: De luteval 2X, Vials (Squibb)

W/Testosterone enanthate

See Defadumane, Vial (Squibb)

Lastomone (Key)

Reposa TE Vial (Canfield)

ESTRADURIN (Ayerst) Polyestradiol phosphate Amp (40 mg) 2 cc

Use: IM, treatment of prostatic carcinoma and estrogen deficiency states

ESTRALDINE (Rorer) Estradiol Vial (10 000 IU/cc & 20 000 IU/cc) 10 cc, Vial Susp (20,000 IU/cc) 10 cc

Use: IM, estrogen therapy

ESTRA-PLEX. (Rawell) Hexestral D1 mg/

Gm. of base adjusted to pH 4 with lactic acid. Cream, tube w/applicator 2 oz.

Use: Estrogenic vaginal therapy.

ESTRAZONE, (New Mexico Pharm) Estrogenic material 10,000 u./Tab. Bot. 100s.

ESTREX INJ. (Rier) Estrone (2.0 mg) 20,000 I.U., testosterone 5 mg., Vit C 100 mg., B-1 25 mg., B-2 5 mg., B-6 5 mg., niacinamide 25 mg., pantothenol 10 mg./cc. Vial (multiple dose) 20 cc.

Use: Estrogenic therapy.

ESTRIFOL, N.N.D. 1962 (Prama) Estrogenic substances conjugated. Tab. 1.25 mg, Bot. 100s.

ESTRIN. See: Estrone

ESTRIOL, N.N.D. 1958, Cap. (Abbott). See Theolol, N.N.R. (Parke, Davis)

ESTROBARB No. 1, (Cole Chem. Co) Phenobarbital ¼ gr., hexoestrol 1 mg./Tab. Bot. 100s, 500s, 1000s.

Use: (I.I.D.) Estrogen w/sedative

ESTROBARB No. 2, (Cole Chem. Co) Phenobarbital ¼ gr., hexoestrol 3 mg./Tab. Bot. 100s, 500s, 1000s.

Use: (I.M.P.) Estrogen w/sedative.

ESTROBINE DP. See: Diethylstilbestrol Dipropionate (Various Mfrs.)

ESTROBEV E AQUEOUS, (Key Corp.) Estradiol Benzoate in Aq. Susp. Vial (20,000 I.U./cc.) 30 cc.

ESTROBEV E IN OIL, (Key Corp.) Estradiol Benzoate in sesame oil. Vial (20,000 I.U./cc.) 30 cc., (100,000 I.U./cc.) 10 cc.

ESTROCASE, (Estra) Aq. sol. of 50 mg. of E hydrolyzed proteins/cc. Vial. 1 cc., 30 cc. Use: Protein therapy.

ESTROEN, (Estra) Sol. in oil of a concentrate of natural estrogen consisting principally of estrone. Amp. (10,000 I.U./cc.) 1 cc., Vial 10 cc., 20 cc., 30 cc., Amp. (20,000 I.U./cc.) 1 cc., Vial 10 cc., 20 cc., 30 cc.

Use: Estrogen therapy.

ESTROFOL, (Prama) Natural estrogenic substance. Aq. Vial (10,000 I.U./cc. & 20,000 I.U./cc.) 10 cc. In oil, Vial (10,000 I.U./cc. & 20,000 I.U./cc.) 10 cc. Tab. (1.25 mg) Bot. 100s, 1000s.

Use: Menopausal symptoms.

ESTROGEL, (Lantean) Estrogenic substances, principally estrone in Isopropyl alcohol base. Tube w/applicator (20,000 I.U., 50,000 I.U. & 100,000 I.U.) 10 cc.

Use: Estrogen therapy.

ESTROGENS EQUINE, (Premark) See: PMS-200, Tab. (Ayerst)

***ESTROGENIC SUBSTANCES, CONJUGATED**, U.S.P., N.N.D. 1962. Tab. U.S.P. (Water-soluble) An amorphous mixture of the naturally occurring water-soluble con-

* Generic name recognized by N.F., U.S.P. or A.M.A.

jugated (ester form) estrogens obtained from the urine of pregnant mares. Cream.

See: Premarin, (Ayerst)

Premarkin H-C, Vaginal Cream, (Ayerst)

Intravenous. See: Premarin, (Ayerst)

Liquid. See: Premarin, Liq. (Ayerst)

Lozion. See: Premarin Lozion, (Ayerst)

Tablets.

See: Amnertragen, (Squibb)

Amoryn, (Plasner)

Estrogen Tab (Barre)

Cogenol, Tab. (National Drug)

Conestron, (Wyeth)

Estrifol, (Prama)

Glyestrin (First Texas)

Kanogen, (Lilly).

Premark, (Ayerst)

Theogen, Tabs (Sig-Ebs)

Ultragen, Tab (Ulmer)

W/di-desoxyephedrine HCl

See: Hormesteral, Tab. (Miller)

W/Hydrocortisone. See: Premarin-HC, Tube (Ayerst)

W/Meprobamate. See: Milprem, Tab. (Wollock)

W/Methyltestosterone, d-desoxyephedrine HCl, vitamins & minerals. See: Vitrimin, Tablets (Paul Maney).

W/Methyltestosterone, rutin, inositol, methionine, vitamins and minerals. See: Via-Gerich, Tab. (Rawell).

W/Phenobarbital. See: Premarin w/phenobarbital, Tab. (Ayerst).

W/Testosterone. See: Kertas, Amp. (Buffington)

ESTROGENIC SUBSTANCES MIXED, May be a crystalline or an amorphous mixture of the naturally occurring estrogens obtained from the urine of pregnant mares.

Aq. Susp.

See:

Aquasension, Vial (Pitman-Moore).

Aque-Strone, Vial (Kirby).

Coestrone, Vial (Barry)

Dorestro, Vial (Smith-Dorsey).

Estrafol, Vial (Prama)

Estromone, Vial (Endo).

Estronoid, Vial (Sherman).

Estrorion, Amp., Vial. (Pitman-Moore)

Estrorin, Amp. (Warren-Teed).

Estrugenone, Vial (Kramerr-Urban).

Estrusal, Vial (C. D. Smith).

Famestrone, Vial (Kirby).

Gothestrone, Amp., Vial (Gatham).

Hormertrin, Vial (Miller).

Lanestrin, Vial (Lanett)

Myasol Ultragen (Ulmer)

Neo-Amniotin, Vial (Squibb).

W/Progesterone (Breon) Estrone 25 mg, progesterone 25 mg/cc Vial 10 cc

Use (IM 0.5 to 1 cc daily) For amenorrhea menorrhagia & habitual abortion

W/Progesterone & estradiol in sesame oil

See: Alamestrone, Vial (Meyer)

Pracyclo Vial (Metro)

W/Progesterone testosterone & procaine

HCl See Kar Tep Aq Susp (Karmel Co)

W Testosterone

See And Est Vial (Ulmer)

Androsterone Vial, Tab (Lincoln)

Androvar n Vial (Warren Teed)

Aneitra Vial (Mallard)

Angen Vial (Eotan)

Di Steroid, Amp (Kremers Urban)

Duagen Vial (C D Smith)

Hormestrin T, Vial (Miller)

Mol O Fem Aq (Testogor)

Myosal and Est (Ulmer)

Testosterone w/Estrone (Ayerst)

Testosterone progesterone

See: Pelrone Vial (Kremers Urban)

P E T, Vial (Maury)

Pro Mol O Fem, Aq Susp (Testogor)

Termone, Vial (Metro)

W/Vil A See Es A Cream (Dome)

ESTRONE PELLETS (Roussel) Estrone Amp

(50 mg) w/needle applicator Box 1s, 6s

Use (1 by implantation) Menopausal symptoms

ESTRONE SULFATE, PIPERAZINE, NNR

See Sulestrex Piperazine (Abbott)

ESTRONE SULFATE POTASSIUM W/ESTRONE See Hormogen R A, Vial (Mallard)

ESTRONE SULFATE, SODIUM

See: Almescon Tab (Meyer)

Konogen, Tab (Lilly)

Linestoral, Tab (Lincoln)

Maretin Tab (Durst)

W/Homatropine methylbromide, phenobarbital & thyroal See Mena Sed Tab (Columbi)

W/Vitamins See: Femestron Compound

Tab (Kirk)

ESTRONID (Sherman) Estrogens mixed

Aq Susp: Vial (20 000 u/cc) 10 cc In

Oil Amp (10 000 u/cc) 1 cc 10 cc; Vial

(20 000 u/cc) 10 cc

Use: Estrogen therapy

ESTROPHEROL (Pitman Moore) Estrone con-

jugated 0.625 mg, d-alpha tocopherol ace-

late 10 mg & phenobarbital 5.4 mg/Tab

Bot 100s

Use: Estrogen therapy w/sedative

ESTROSED (Chicago Pharmacal) Reserpine

0.1 mg, ethinyl estradiol 0.01 mg/Tab

Bot 100s, 1000s

Use: Orally anxiety or tension neurases

ESTRO-SERP. (Ortega) Ethinyl estradiol 0.01

mg, reserpine 0.1 mg/Tab Bot 100s, 500s

Use: Treatment of menopausal symptoms

ESTRO-STEN (New Mexico Pharm) Estro-

genic substances 2 mg, methyl and o-

stenediol 15 mg, carboxymethyl cellulose

0.2%, Tween 80 0.4%/cc Liq Vials 10 cc

ESTROSYN. (Burlington) Stilbestrol Tab

EC (0.1 mg, 0.5 mg, 1 mg, 5 mg) Bot

100s, 1000s

Use: Orally; estrogenic therapy

ESTROTEIN. (Estra) 5% sol of hydralized

casein with 10 mg of sulfur in the form of

Sad mono & poly sulfides, sulfate & thio-

sulfate w/benzyl alcohol 1% Amp 2 cc

Vial 30 cc

Use: Nonspecific protein therapy

ESTRO-TESTONE (New Mexico Pharm)

Estrogenic substances 20M IU 2 mg, tes-

tosterone 10 mg, benzyl alcohol 2%/cc

Vials 10 cc

ESTROVARIN, NNR 1950 (Warren Teed)

Estrogens mixed Aq · Amp (20 000 u/cc)

1 cc, 12s 25s 100s Vial 15 cc In Oil

Amp (10 000 u/cc) 1 cc, 12s, 25s 100s

Vial 15 cc, 1s

Use IM; estrogen therapy

ESTRUGENONE, NND 1958 (Kremers-

Urban) Estrone

Aq · Amp (2 mg/cc) 1 cc 25s 100s Vial

2, 6, 10, 25 cc Amp (5 mg/cc) 1 cc 6s

25s 100s Vials 5 & 10 cc

In Oil Amp (1 mg/cc) 1 cc, 25s 100s

Vial 10 cc

Use IM; estrogen therapy

ESTRUSOL, NND 1958 (Smith, Miller &

Patch) Estrone

Parent: all (1 mg/cc) Vial 30 cc

Susp · aq. (2 mg/cc) Vial 10 cc & 15 cc

(5 mg/cc) Vial 15 cc

Tab · (0.1 mg) Bot 100s 1000s

Use Estrogen therapy

ETALATE (Parke Davis) A sterile 5% sol of

ethylamine oleate with 2% benzyl alcohol

Amp 2 cc, Box 12s 100s; Amp 5 cc, Box

6s

Use Sclerosing varicose veins

ETAMON CHLORIDE, NND 1962 (Parke

Davis) Tetraethylammonium chloride Vial

(0.1 Gm/cc) 20 cc

Use Circulatory diseases of the extremities

ETHACAIN. (Seydel Chem Co) Benzocaine

benzoate 2.5%, oxyquinoline benzoate

0.25%, glycol ether in petrolatum 2.5%

Oint, Tube 1 oz Jar 1 lb

Use: Burns sunburn eczema pruritus ani

or vulvae

ETHACREO (Merck Sharp & Dahme) Chloro-

form 60 mg, creosote 0.12 cc, terpin hyd

- 0.12 Gm., cal. glycerophos. 0.25 Gm., sod. glycerophos. 0.25 Gm./30 cc. Sol. 1 oz.
Use: (1-2 tsp. q 3 h.) Cough
- ETHALYL ELIXIR.** (Premo) Sod. pentobarbital 1 gr., Sod. phenobarbital 1 gr./fl. oz. Bot. 8 oz. 1 pt., 1 gal.
Use: (1-2 tsp. p.r.n., m.p.) Sedative & hypnotic.
- ETHALYL SODIUM CAPSULES.** (Premo) Sod. phenobarbital 1 gr., Sod. pentobarbital 1 gr./Cap. Bot. 100s, 1000s.
Use: (1 q. 3-4 h., p.r.n.) Sedative & hypnotic.
- ***ETHAMBUTOL HCl.** (Lederle) N,N'-bis(4-hydroxymethylpropyl)ethyl-enediamine diHCl. See: *Myambutol Hydrochloride* (Lederle)
Use: Antitubercular.
- ETHAMICORT.** See: *Hydrocortamole*.
- ***ETHAMIVAN.** N,N-diethylvanillamide. See: *Emivan* (U.S. Vitamin)
- ETHAHACOL.** (Blue Line) Ethyl carbamate 4 gr., ammonium chloride, Sod. benzoate & sugar/5 cc. Sol. 1 pt., 1 gal.
Use (1-2 tsp q. 3-4 h.) Cough
- ETHANOL.** A.D.R. 1955. (Various Mfr) (Alcohol, anhydrous).
- ***ETHASULFATE SODIUM.** (Abbott) Sod 2-Ethyl-1-hexanol sulfate.
- ETHAVERINE.** (Chema Pura) Bivinal, perparin, perperine, preparein. 6,7-Diethoxy-1-(3,4-diethoxybenzyl) isoquinoline HCl.
Use Antispasmodic.
- ETHAVERINE HYDROCHLORIDE.** The ethyl analogue of papaverine HCl, 6,7-diethoxy-1-(3,4-diethoxybenzyl) isoquinoline HCl. Biquinol HCl, Preparin HCl, Perperine HCl. The tetraethyl homolog of papaverine, is 2 to 4 times more active and less than half as toxic as the parent drug. This compound is not a narcotic.
See: Ethiquinol, Tab. & Vial. (Uimer)
Harverine (New Mexico Pharm.)
Neopavrin A (Savage)
Papaverine Hydrochloride. (Lincoln)
Verina (Testogor)
(Lincoln) Tab. (1/2 gr.) Bot. 100s, 300s. Yield (15 mg./cc.) 10 cc.
Use. (i.v. 15 to 100 mg.) Antispasmodic.
W/Belladonna ext. & phenyltoloxamine. See: *Tri-Synar*, Tab. (Armour).
W/Betaine HCl, Mephenesin. See: *Mepherin*, Tab. (G. F. Harvey)
W/Dihydroxy aluminum aminoacetate & gastric mucin. See: *Mugastrel*, Tab (G. F. Harvey)
W/Dihydroxypropyl theophylline. See: *Caravar*, Tab (Crystal)
- * Generic name recognized by N.F., U.S.P. or A.M.A.
- W/Glyceryl guaiacal ether & aspirin See: *Thevasin*, Tab. (Lincoln)
- W/Isopropylarterenol, theophylline, butobarbital. See: *Benedrin*, Tab. (Bene)
- W/Mannitol hexanitrate & phenobarbital. See: *Hexaverine*, Tab. (Lincoln).
- W/Mephenesin, betaine HCl. See: *Mepherin*, Tab. (G. F. Harvey)
- W/Neothylline. See: *Neavar*, Tab. (Paul Maney)
- W/Pentaerythritol tetranitrate
See: *Papavital*, Tab. (Kenwood Labs)
Pertal, Tab. (Uimer)
- W/Phenyltoloxamine & belladonna ext. pow. See: *Tri-Synar*, Tab. (Armour).
- W/Theophylline Sod. glycinate. See: *Thaverine*, Tab (Pitman-Moore).
- ***ETHCHLORVYNOL**, N.F., NND 1962 β -chlorovinyl ethyl ethinyl carbinal See: *Flacidyl*, Cap. N.F. (Abbott)
- ***ETHIBENECID.** p-Diethylsulfamoylbenzoic acid. *Uretim*.
- ***ETHINZAMIDE.** o-ethoxybenzamide.
- ETHER-3H-OIL.** (Brewer) Ether 50% in all. Amp. 2 cc. Box 12s, 100s.
Use: I.M., treatment of bronchial asthma.
- ETHER, SPIRIT OF NITROUS.** (Squibb) Ethyl nitrite spirit N.F. Bot. 1 pt.
Use: Diuretic & diaphoretic.
- ETHIXAL.** (Hiss) 900 calorie food supplement with dextroamphetamine and amobarbital. Pow. Bot. 8 oz.
Use: Obesity.
- ***ETHINAMATE**, N.F., NND. 1962 Valmidate, 1-Ethinylcyclohexyl carbamate. Tab N.F. See: *Valmid*, Tab (Lilly)
- ETHINORAL.** (Smith, Miller & Patch) Ethinyl estradiol 0.05 mg /Tab. Bot 1000s.
Use: (1 t.d.) Estrogen therapy
- ***ETHINYL ESTRADIOL**, U.S.P., NND 1962, Tab., U.S.P 17-Ethinyl-3,17-estradiol.
See: *Choo-O-Gen*, Tab. (Keene)
Diogyn-E, Tab (Pfizer)
Esteed, Tab (Warren-Teed).
Estinyl, Lq & Tab (Schering).
Ethinoral, Tab (C. D. Smith).
Eticylol, Tab & (inguets, N.N.R. (Ciba)
Estoral, Tab. (Key Corp)
Inestra, Tab. (Merck Sharp & Dohme)
Lynoral, Elixir & Tab. (Organon)
Navestral, Tab. (Rarex)
Oradial, Tab (Yanpelt & Brown)
Orestrolyn, Tab. & Elixir. (McNeil).
Palonyl, Tab (Palmedica)
- W/Anhydroprogesterone See: *Predictabs*, (Drug Specialties-Prane)
Progesteroid (Brown Pharmaceutical)
- W/Anhydroxyprogesterone. See: *Pro-Duo-sterone*, Tab. (Rovsief)

- W/Ethisterone See Duasterone Tab (Roux
set)
- W/Ethisterone See Pregnancy Test Tablets
(West Ward)
- W/Ethisterone, thyroid ext., hesperidin Vit
B₁, Vit C, & glutamic acid See Proges-
traid, Tab (Brown Pharmaceutical)
- W/Fluoxymesterone See Haladrin, Tab
(Upjohn)
- W/Halofestine Vit A D thiamine mono-
nitrate riboflavin pyridoxine HCl cyana-
cobalamin, ascorbic acid nicotinamide cal-
cium pantothenate folic acid, calcium phos-
phate d basic See Hormozyme, Tab
(Upjohn)
- W/Methyltestosterone
See: A E, Tab (Warner Chilcott)
Duatric, Tab (Dumas Wilson)
Dumans Tab (Squibb)
Duatrons Tab (US Standard)
Femadren Linguets (Ciba)
Gynetone Repetab (Schering)
Para Testrogen Tabs (Para)
Testrolx Elix (Chicago Pharm)
- W/Methyl testosterone, d-amphetamine sul-
fate, Geval formula See: Gevrestin, Cap
(Lederle)
- W/Methyltestosterone, d-amphetamine sulf-
ate and vitamins and minerals See: Yealric
Cap (Dumas Wilson)
- W/Methyltestosterone, d-amphetamine sulf-
ate, nicotinic acid, & Vit C See: Stimogen, Tab
(Lemman)
- W/Methyltestosterone, metrozol vitamins
and minerals See: Senazol, Cap (Tablet
rock)
- W/Methyltestosterone pentylenetetrazol, nic-
otinic acid and dicalcium glutamate See:
Senatric, Tab (Dumas Wilson)
- W/Methyltestosterone, pentylenetetrazol,
nicotinic acid Vit C, thyroid See: Matri-
cine Tab (Ascher)
- W/Methyltestosterone, pentylenetetrazol nic-
otinic acid tetrahydrotyranine See: Tetra-
force Tab (E S Miller)
- W/Methyltestosterone, prelad See: Plestran,
Tab (Warner Chilcott)
- W/Methyltestosterone, B₁₂ folin, B₁, B₂ B₆
B₁₂ nicotamide dicalcium phosphate
See: Plancic Cap (Ciba)
- W/Methyl testosterone thyroid extract as-
corbic acid calcium lactate d calcium
phosphate vit A D E See: Forastee Tabs
(Brown)
- W/Methyltestosterone vitamins & minerals
See: Duorgan Tab (Squibb)
Forastee Tabs (Brown)
Hormo Plus in Caps (Vitarine)
Neobon Tab (Roerig)
- * Generic name recognized by N.F., U.S.P. or A.M.A.
- W/Norethindrane acetate See Gestest Tab
(Squibb)
- W/Pentylenetetrazol, methyltestosterone
strontium salicylate See Analeptone Ana-
bolic (Reed & Carnrick)
- W/Phenobarbital Sod See: Sedanyl, Liq
(C D Smith)
- W/Reserpine
See Estrosed, Tab (Chicago Pharmacal)
Estra serp (Ortega)
Sergynol, Tab (Ascher)
- W/Thyroxine methyltestosterone, d-ampheta-
mine sulf & vitamins See: Neabon, Liq
(Roerig)
- W/Vitamins See Vi Estry, Cap (Arling
ton Funk)
- W/Vitamins & minerals See Gevrine, Cap-
(Lederle)
- ETHINYL (17)-19-NORTESTOSTERONE.**
See Norlutin, Tab (Parke, Davis)
- ETHIODAN** See Iophendylate USP
- ETHIODIZED OIL** See Ethiodol (Fougera)
- ETHIODOL** (Fougera) Fatty acid ethyl ester
of poppy seed oil w/37% iodine Amp 10
cc, Box 2s
Use: Diagnostic agent, roentgenography
- * **ETHIONAMIDE**, 2 Ethylthioisonicotinamide
See Trecator (Ives Cameron Co, Inc)
- * **ETHISTERONE**, USP Tab USP 17-alpha-
ethinyl 17 beta hydroxy 3 oxoandrostene-4
See Anhydrohydroxyprogesterone USP
XIII (Various Mfr)
- Also See: Lucortem, Tab (Merck Sharp &
Dohme)
Lutocyl, Tab & Linguets (Ciba)
Lutoral, Tab (Schieffelin)
Ora lutin, Tab (Parke Davis)
Pronone, Tab (Schering)
Prodroxan, Tab (Smith, Dorsey)
Progestab, Tab (Massengill)
Progestal, Tab (Organon)
Prone, Tab (Barre)
Syngestratabs, Tab (Pfizer)
Troisane, Tab (Abbott)
- W/Ammanium chloride d-amphetamine sul-
fate, levo-hyoscyamine hydrabramide See
Gymens Tab (Calumbus)
- W/Chlorophenpyridamine maleate &
bromotheophylline, salicylamide See C
Compound (Schering)
- W/Ethinyl estradiol
See Duasterone, Tab (Roussel)
Estrafuleum, Tabs and Susp (Lincoln)
Pred tabs, Tab (Drug Specialties Prodel)
Pregnancy Test Tablets (West Ward)
Progestrail (Brown Pharmaceutical)
- W/Ethinyl estradiol thyroid ext., hesperid-
in Vit B-1, Vit C, & glutamic acid See Pro-
gestroid, Tab (Brown Pharmaceutical)

- W/Hesperidin complex, ascorbic acid, Vit. K analogue, di-alpha tocopherol acetate. See: Nugesoral, Tab. (Organon)
- W/Phenobarbital & Thiamine. See: Surdon, Tab. (Organon).
- ETHNINE**, [Lab. for Pharmaceutical Development, Inc.] See: *Phalcaline
- ETHOBRAL**, (Wyeth) Secobarbital Sod. 50 mg., butabarbital Sod. 30 mg. & phenobarbital 50 mg./cap. Bot. 100s, 1000s. Also available in half-strength.
Use: (1 h.s.) Sedative.
- ETHOCEL**, (Dow Chem) Ethyl Cellulose
- ETHOCAINE**, See: Procaine Hydrochloride (Various Mfrs.)
- ETHOCHLORVYNOL**, β -Chlorovinyl ethyl ethynyl carbinol. See: Placidyl, Cap. (Abbott)
- ETHODRYL**, See: Diethylcarbamazine Citrate.
- ***ETHOHEPTAZINE CITRATE**, N.N.D. 1962. 3-Methyl-4-carboethoxy-4-phenyl hexamethylenimine citrate. See: Zactane Tab (Wyeth)
- W/Acetylsalicylic Acid. See: Zoclirin, Tab. (Wyeth)
- W/Meprabamate, acetylsalicylic acid See: Equegest, Tab. (Wyeth)
- ***ETHOHEXADIOL**, U.S.P. Ethyl hexanediol, 2-ethylhexane-1,3-diol, Rutgers 612. Used in Comp. Dimethyl Phthalate, N.F.
Use: Insect repellent.
- ETHOLATE**, (Metro) Monoethanolamine oleate. Vial (5%) 30 cc.
Use: Sclerosing agent.
- ***ETHOMOXANE HCl**, (Eli Lilly) dl-8 ethoxy-2-n-butylaminomethyl-1,4-benzodioxane HCl. Use: Tranquillizer.
- ***ETHOPROPAZINE HYDROCHLORIDE**, N.N.D. 1962. Lysivane See: Parsidal, Tab. (Warner-Chilcott)
- ***ETHOSUXIMIDE**, 2-Ethyl-2-methylsuccinimide See: Zorontin (Parke, Davis)
- ***ETHOTOIN**, N.N.D. 1962. 3-Ethyl-5-phenyl hydantoin. See: Peganone, Tab. (Abbott)
- ETHOVAN**, See: Ethyl Vanillin, N.F.
- ***ETHOXAZENE HYDROCHLORIDE**, 2,4-diamino-4'-ethoxyazobenzene hydrochloride. See: Serenium (Squibb)
Use: Urinary analgesic
- ***ETHOXZOLAMIDE**, N.N.D. 1962. 6-Ethoxybenzothiazole-2-sulfonamide. See: Cordrase, Tab. (Upjohn)
- W/Ectylurea, medroxyprogesterone. See: Cytran, Tab (Upjohn)
- ETHOUINOL**, (Ulmer) Ethaverine HCl, Tab. 30 mg. Bot. 100s. Vial (15 mg./cc) 10 cc.
Use: Antispasmodic.
- ***ETHYBEHTROPINE BROMIDE**, (Sandoz)
* Generic name recognized by N.F., U.S.P. or A.M.A.
- 3 (d-phenylmethoxy)-8-ethylnoretopane hydrobromide.
Use: Anticholinergic.
- ETHYLAMINE OLEATE**, See: Etalate (Parke, Davis)
- ***ETHYL AMIHOENZOATE**, N.F., N.N.R. 1949; A.D.R. 1955; Oint., N.F., A.D.R. 1954
Anesthetin, anesthone, benzocaine, parathesin See: Benzocaine.
Use: Surface anesthetic.
- ***ETHYL BISCOUMCETATE**, N.N.D. 1962. 3,3'-Carboxymethylene bis (4-hydroxycoumarin)ethyl ester. See: Tramexon, Tab N.N.R. (Geigy)
- ETHYL BROMIDE**, (Various Mfrs.) Bromoethane.
Use: Inhalation anesthetic.
- 3-ETHYL-5-N-RUTYRARRRITURIC ACID**. See: Butethal, N.F.
- ETHYL CARBAMATE**, U.S.P. See: Urethone, Cap. (Lilly).
- W/Ammonium chloride, Sod. benzoate & sugar. See: Ethonacal, Cough Preparation (Blue Line)
- ***ETHYL CARTRIZOATE**, (Winthrop Labs) Ethyl (3,5-diacetamido-2,4,6-trimodobenzoyloxy)acetate.
Use: Bronchographic agent.
- ETHYL CHAULMOODRATE**. See: Chaulmestrol, Amp. (Winthrop Labs) Moogrol, liq. (Burroughs Wellcome).
Use: Hansen's disease, sarcoidosis.
- ***ETHYL CHLORIDE**, N.F., N.N.R. 1949, A.D.R. 1955.
Bio America—Spray pack, 100 Gm.
Stratford-Cook—Amp. 5 cc. Tube 30, 40, 60, 80, 100 Gm.
Use: Local anesthetic.
- 2-ETHYLCROTONYLUREA**. See: Levani, Tab. (Upjohn)
Nostyn, Tab. (Ames)
- ETHYL DIBUNATE**, (Nat. Drug Co) Ethyl 3,6-dialkylbutyl-naphthalene-1 (2)-sulfonate
- ETHYL DIODOBRASSIDE**, Iodobrossid. See: Lipodine, Tab. (Dwight)
Lipiodine, Tab. (Ciba)
- ETHYLDIMETHYLAMMONIUM**, See: Ambutanum Bromide
- ***ETHYLENE**, N.F., N.N.R. 1950, A.D.R. 1955. (Various Mfrs.) Ethene.
Use: Inhalation anesthetic.
- ETHYLENEDIAMINE DIHYDROCHLORIDE**. See: Chlor-Ethamine, Tab. (Pitman Moore)
- ETHYLENEDIAMINE DIHYDRIODIDE** W/Ammonium chloride, squill fr.
See: Cheronine, Cough Prep. (Pitman Moore)
Iod-Ethamine, El. & Tab. (Pitman Moore)
W/Phenobarbital, Cal. theobromine-Cal. so-

- lyclate See Xanclate Tab (Pitman Moore)
- W/Sod salicylate See: Ioclyate, Tab (Pitman Moore)
- W/Theobramine See Xanamine Tab (Pitman Moore)
- W/Theobramine phenobarbital See: Xanaphen Tab (Pitman Moore)
- ETHYLENEDIAMINE DIHYDROCHLORIDE.**
See Chlor Ethamine, Tab (Pitman Moore)
- *ETHYLENEDIAMINE SOLUTION, USP**
(67% w/v)
Use Solvent (Aminophylline Inj)
- ETHYLENEDIAMINETETRAACETIC ACID**
See Edathamil EDTA
- ETHYLENEDIAMINE TETRAACETIC ACID-DISODIUM SALT.** See Endrale Disodium, Amp (Abbott)
- 1-H-ETHYLEPHEDRINE HYDROCHLORIDE**
See Nethamine Hydrochloride, Prep (Merrell)
- Use Branchadilator & vasoconstrictor
- *ETHYLESTRENOL** (Organon, Inc) 17 α ethyl 17 β hydroxy estr 4 ene
See Orgobalin
Uses Anabolic steroid
- 2-ETHYLHEXANE-1,3-DIOL** See Ethohexadiol, USP
- ETHYLHYDROCUPREINE HYDROCHLORIDE, NNR 1946** See: Optochin, Oint (Mesco)
- ETHYL IODIDE** (Merck Sharp & Dohme) Bot 1 oz, $\frac{1}{4}$ lb 1 lb
Use Inhalation in iodide therapy
- ETHYL IODOPHENYLUNDDECYLATE, USP, NND 1962** Iophendylate
See Iophendylate Injection USP
Mulsapaque Amp (Lafayette)
Pantopaque NNR (Lafayette)
- 5-ETHYL-5-ISOAMYL BARBITURIC ACID**
See Amobarbital USP
- ETHYL OXYLATE** W/Ethyl malonate ethyl alcohol See Kaogamin Box (Chatham)
- *ETHYLMORPHINE HYDROCHLORIDE, NF, NND 1962** See Diamin, Pow (Merck Sharp & Dohme)
Use Narcotic
- W/Chloroform, ammonium chloride terpin hydrate syr solu, wild cherry syr See Sedaca Liq (Strassenburgh)
- W/Chloroform, Pot guaiacal eulfanate ammonium chloride, Sod citrate tartar emetic See: Dionical Liq (Rorer)
- W/Chloroform ephedrin sulfote Pot guaiacal sulfanate squill fl ext, ipecac fl ext sanguinaria fl ext, menthol benzocainn solu See E D Syrup Liq (Tilden)
- W/Chloroform, labelex tarlar emetic See: Calsecat, Elix (Massengill)
- * Generic name recognized by HF, USP or AMA
- W/Cociflora fr, euphorbia fr, wild lettuce syr, cascara sagrada ext & menthol See Cocis Thal, Syr (Buffington)
- W/Euphorbia pilulifera fr, squill comp syr-wild lettuce syr, menthol, alcohol See Casanyl, Syr (Parke, Davis)
- W/Pot citrate, citric acid, Pot guaiacal sulfanate See: Cidical Cough Mixture (Upjohn)
- W/Pot guaiacal sulfanate cas evac ipecac flexit citric acid sod citrate See Citronin Liq (Parke Davis)
- W/Pot guaiacal eulfanate, ephedrine HCl, cociflora fr, wild lettuce syr, squill comp syr, euphorbia fr See Dionedro Compound, Liq, Syr (Central)
- W/Sod citrate, citric acid, Pot guaiacal eulfanate, cas evac ipecac flexit & alcohol See Citronin, Cough Preparation (Parke Davis)
- W/Terpin hydrate Semi Diaterpine Liq (Schieffelin)
- W/Terpin hydrate, Col glycerophes, Sod glycerophes, creosote carb See Terpalone Liq (Strassenburgh)
- W/Terpin hydrate, creosote, eucalyptol See Krea Dionin Terpin Compound, Liq (Tilden)
- ETHYLNOREPINEPHRINE**
See Butanefrine Hydrochloride, Amp, Vial (Winthrop Labs)
Bronkephyrine HCl, Vial (Breon)
- ETHYLPAVERINE HCl.** See Ethovarine HCl
- W/Phenobarbital pentaerythritol tetrantrate See Papavatal LA w/Phenobarbital, Capl (Kenwood)
- ETHYL SALICYLATE CARBONATE** W/Bromisovalum, acetophenetidin & caffeine See Acetabra Pediatub No 1 (Columbus Pharm)
- 5-ETHYL-SEC-BUTYLBARBITURIC ACID**
See Butobarbital
- ETHYLSTIBAMINE, USP, NNR 1950** Astibil enastibosan
Use Anismany therapy
- dl- α -ETHYLTRYPTAMINE** See Manase (Upjohn)
- ETHYL VANILLATE.** (Squisbb) Ethyl hydroxy methoxy benzoate Cap (0.5 Gm) Bot 500s
Use (1 s d) For histoplasmosis & coccidioidomycosis
- *ETHYL VANILLIN, NF 3** Ethoxy 4 hydroxy benzaldehyde
Use Flavor
- *ETHYNOBIOL DIACETATE** [G D Sealey] 19 nar 17a pregn 4 en 20 yne 3 β 17 β diol diacetate [7 α ethynyl 4 estrene 3 β 17 β diol acetate
Uses Progestin

ETHYNYLESTRADIOL. See: Ethinyl Estradiol, U.S.P. (Various Mfr.)

ETHYNYLESTRADIOL 3-METHYL ETHER. See: Enovid (Searle)

ETICYLOL. N.N.D. 1962. (Ciba) Ethinyl estradiol. Tab. (0.02 mg. & 0.05 mg.) Bot. 100s. Linguets (0.5 mg.) Bot. 100s. Use: (0.05 mg. i.i.d.) Estrogen therapy.

E-TOPLEX. (U.S. Vitamin) Alpha-tocopherol. Cap. (30, 100, 150 mg.) Bot. 100s, 500s, 1000. Drops (50 mg./cc.) Bot. 15 cc., 60 cc. Use: Orally; Vit. E therapy.

ETOVAL. See: Bulethol, N.F. (Various Mfr.)

ETRYNIT. (Breon) Propyl nitrate
Use: Coronary vasodilator.

* **ETRYPTAMINE ACETATE.** (Upjohn) 3-(2-aminobutyl)indol acetate. See: Manase.

EUBASIN. See: Sulfopyridine (Various Mfr.)

EU-BEN-SAL. (Tilden) Benzoic and salicylic acids, thymol, pine oil, petrolatum. Jar 1 oz., lb.
Use: Treatment of athlete's foot

EUBERTONE. (First Texas) Ovarian substance desic. 3 gr., thyroid ext. 1/8 gr., gland's moss 2 gr., Col. glycerophos. 2 gr., arsenous acid 1/40 gr., strychnine sulfate 1/120 gr./Cap. Bot. 100s, 500s, 1000s.
Use: Ovarian substance & thyroid.

EUCADRYL. (Crookes-Bornes) Phenylephrine HCl 5%, eucatropine HCl 5% in buffered methylcellulose 1%. Vials 7.5 cc. and dropperettes.
Use: Mydriasis

EUCAPINE SYRUP. (Lonnell) Ammonium chloride 10 gr., pot. guaiacal sulf. 8 gr., chloroform 2 min. in white pine & wild cherry syrup. Bot. 1 pt., 1 gal.
Use: Cough expectorant.

* **EUCATROPINE HYDROCHLORIDE, U.S.P., N.N.R. 1950.** (M. R. Thompson) Cryst., Bot. 1, 5 & 25 Gm.

See: Euphthalmine, Cryst. (Warner-Chilcott)
Isopne-Eucatropine 3% (Alcon).

W/Phenylephrine
See: Eucodryl (Crookes-Bornes) Mydriatin (Ophthalmus, Inc)

EUCEMA. (Tilden) Zemecide, milk of magnesia, eupinol (12 1/2%) lot. Bot. pts., gals.
Use: Emollient, antipruritic.

EUCODAL. See: Dihydrohydroxycodainone, Prep. (No Mfr. currently lists)

EUCORAN. See: Nikelhomide (Various Mfr.)

EUCUPIN DINYDROCHLORIDE. (White Labs.) Isoomyl-hydrocupreine dihydrochloride
Oint.: Tube (with applicator tip) 1 oz., Jar 1 lb.

* Generic name recognized by N.F., U.S.P. or A.M.A.

Sol. (Eucupin D₂HCl 0.2%, procaine HCl 1%) Vial 10 & 30 cc.

Sol. in oil: (Eucupin base 0.1%, ethyl aminobenzoate 3%, benzyl alcohol 5%, sweet almond oil) Amp. 5 cc., Box 6s, 24s, 96s. Vial 10 cc.

Supp.: (Eucupin D₂HCl 1%, benzocaine 2%, bismuth subgallate 5%) Box 12s.

Use: Local anesthetic.

W/Zalamine. See: Otodyne, Liq. (White Labs.)

EUDICAINE. (Rexall) Benzocaine 3%, titanium dioxide, menthol, balsam peru & phenyl mercuric acetate. Oint. Tube 1 oz.

Use: Topically; local anesthetic & skin protective.

EUFILAVINE. See: Acriflavine (Various Mfr.)

EUGESTIN. (Rexall) Diastase concentrate 20 gr./oz. Bot. 1 pt., 1 gal.

Use: To promote digestion

EUGLYCIN. (Upjohn) Metohexonide. Under study.

Use: Hypoglycemic agent.

EUKADOL. See: Dihydrohydroxycodainone, Preps. (No Mfr. currently lists)

EULCIN. (Staff Research Labs.) Methscopolamine bromide 2.5 mg., butobarbital Sod 10 mg., aluminum hydroxide gel dried 250 mg., Mag. trisilicate 250 mg./tab. Bot. 100s. Use: 1-2 h.s.; gastric & duodenal ulcer therapy

EULISSIN. See: Decamethonium iodide

EUMYDRIN. (Winthrop Labs.) Atropine methylnitrate. Tab. (1 mg.) Bot. 23s, 100s.

Use: Antispasmodic biliary & intestinal colic.

EUNERYL. See: Phenobarbital (Various Mfr.)

EUPHAGIN. (Winthrop Labs) Anesthesin D1 Gm. w/Sod. borate & menthol/Tab Bot. 100s.

Use: Orally; local anesthetic

EUPHASID. (Schenley) d-Desoxyephedrine HCl 25 mg. & acetylcarbromal D.26 Gm./Tab. Bot. 100s, 1000s.

Use: To reduce the appetite; sedative action.

EUPHASED-S. (Schenley) d-Desoxyephedrine HCl 5 mg., acetylcarbromal D.26 Gm./tab. Bot. 100s, 1000s.

Use: Anabesity.

EU-PHED-AMIN. (Warren-Teed) Aminophylline 1 1/2 gr., ephedrine sulfate 1/2 gr., phenobarbital 1/2 gr., euphorbia pilulifera ext. 1 1/2 gr./Tab. Bot. 100s, 500s, 1000s. Use: (1-2 i.i.d.) Bronchial asthma.

EU-PHED-ITAL. (Warren-Teed) Phenobarbital Sod 1/2 gr., ephedrine sulfate 1/2 gr. eu-

phorbia pilulifera ext 1½ gr/Tab Bot 100s
 Use (1 t i d) Antiothmatic w/sedative

EUPHENEX. (Lloyd Dabney & Westerfield) Phenyltoloxamine dihydrogen citrate 25 mg, salicylamide 100 mg, acetyl p omino phenol 200 mg, ascorbic acid 50 mg citrus bioflavonoids 50 mg, caffeine 15 mg, racemic amphetamine sulfate 2 mg/Tab Bot 100s
 Use Treatment of the common cold

EUPHORBIA COMPOUND (Lilly; Merck Sharp & Dohme) Euphorbia pilulifera fl ext 1.5 cc, lobelia tr 22 cc nitroglycerin spirit 0.29 cc, Sod iodide 1.04 Gm, Sod bromide 1.04 Gm, alcohol 24%/30 cc Bot 1 pt, 1 gal
 Use (1 2 tsp p r n) Sedative & expectorant

EUPHORBIA & SQUILL COMPOUND SYR-UP. (Merck Sharp & Dohme) Chloroform 0.12 Gm, euphorbia 8 cc, squill flexil 0.2 cc, ipecac 0.18 Gm, Jamaica dogwood 1 D4 Gm, menthol 8 mg/30 cc Bot 1 pt, 1 gal
 Use Expectorant & sedative

EUPHTALMINE, NNR 1930 (Worner Chl colt) Eucatropine HCl Cryst, Rot D 5 Gm, 1 Gm, 5 Gm, 25 Gm
 Use Topically; mydriatic

EUPINOL UNGUENT. (Tilden) Eupinol resorcin ichthammol oil birch tar, alum camphor Oint Jar 1 lb
 Use: Ant septic antiphlogistic

EU-PRO-CAMPH. (Brean) Eucalyptol 0.15 Gm, camphor 0.05 Gm, menthol 0.06 Gm Pracoine base 0.04 Gm, in sesame oil/2 cc Amp 2 cc, Box 12s, 25s 100s
 Use Local anesthetic

EUQUININE, NNR 1949 Quinine ethyl carbonate
 Mallinckradt—Pow, Bot 1 oz, Canister 5 oz
 Merck—Pow, Canister ½ oz, 1 oz
 Use Orally; antimalarial & antipyretic

EURAX CREAM, NND 1962 (Geigy) Cro taminon 10% N ethyl o cratonaloluide in a vanishing cream base Tube 2D Gm, 60 Gm, Jar 1 lb
 Use: Antipruritic & scabicide

EURAX LOTION, NND 1962 (Geigy) Cro taminon 10% Bot 2 oz, 6 oz 1 pt, 1 gal
 Use: Relief from itching in skin disorders

EURESOL, NNR 1950 (Knoll) (Retorcinol monoacetate) Meta hydroxyphenyl acetate Rot 1 oz, 8 oz
 Use: Topically; antipruritic & antipruritic

EURESOL PRO CAPILLIS, NNR 1949 (Knoll) This product is identical with euresol, except that it is pleasantly per
 * Generic name recognized by N F, U S P or A M A

fumed and contains 6% isopropyl alcohol Bot 1 oz, 8 oz, 1 lb

EUTONYL. (Abbott Lab) *Pargyline Hydrochloride
 Use Antihypertensive antidepressant

EVAC-U-GEN. (Walker) Sod salicylate ¼ gr, bismuth subcorb ½ gr, bismuth sub gallate ¼ gr yellow phenolphthalein/Tab Pkg 24s Bot 100s
 Use Laxative

EVANARD (Brean) Vit D 50 000 u, deoxy cholic acid 32.4 mg, activity equiv to dl alpha tocopherol 50 mg/Cop Bot 100s 1000s
 Use: (1 3 daily) For arthritis & liver dysfunction

***EVANS BLUE,** USP NND 1962 In USP (Worner Chilcott) Pure diazo dye Amp (0.5%) 5 cc
 Use (1 V 25 mg) Diagnostic aid

EVASOF (Lemmon) Diethyl sod um sulfat succinate 50 mg acetophenolisatin 2 mg/Tab Bot 100s
 Use Constipation

E-VATINE. (Smith Dorsey) dl Alpha tosopherol acetate 200 mg, chlorobutanol 0.5% in perse oil/cc Vial 10 cc, Cop (50 mg) Bot 100s, 1000s
 Use Vit E therapy

EVICYL. (Winthrop Labs) Each tablet contains 65 mg of Evipal & D 49 Gm of acetyl salicylic acid Tab, Bot 100s, 500s
 Use (1 2 q 3 4 h) Analgesic & sedative

EVIDORN. (Winthrop Labs) Each tablet or capsule contains 4 gr of Evipal & 1½ gr of Phenodorn Col Tab Tube 10s Bot 100s 500s; Cap, Bot 100s, 500s
 Use: (1 q 4 h p r n) Sedative

EVIPAL (Winthrop Labs) (Hexobarbital) N methylcyclohexenylmethyl barbituric acid Tab (4 gr) Bot 100s
 Use Sedative

EVIPAL SODIUM, NND 1961 (Winthrop Labs) Hexobarbital Sod Amp (0.5 Gm) w/ 5 cc of distilled water, Amp (1 Gm) Vial (5 Gm) for rectal use only
 Use Anesthetic for short operations
 W/Aspirin See Evicyl Tab (Winthrop Labs)
 W/Phenodorn Cal See Evidorn, Cap & Tab (Winthrop Labs)

EVIPHEROL. (Kirk) dl Alpha tocopherol Cap. (50 mg) Bot 5Ds, 100s, 1000s, Vial (200 mg/cc in oil) 1D cc
 Use: Vitamin E therapy

E-VITUM (ives Cameron) dl Alpha tocopherol acetate Cap (70 I U) Bot 100s, 1000s
 Use: Vitamin E therapy

EVROCILLIN. (Ewan) No 1 Pat penicillin G

- 100,000 u./5 cc; No. 2 Pot. penicillin G
250,000 u./5 cc. Bot. 2 oz.
Use: Penicillin therapy.
- EVRODEX TEMPULES.** (Evron) d-Amphetamine sulfate, Tab. 10 or 15 mg Bot. 100s.
Use: Antiobesity.
- EVRODEX-PLUS TEMPULES.** (Evron) d-Amphetamine sulfate 10 mg., amobarbital 60 mg./cap. or d-Amphetamine sulfate 15 mg., amobarbital 100 mg./cap. Bot. 100s.
Use: Antiobesity.
- EVRONAL SODIUM, NND 1961.** (Evron) Secobarbital Sod. Cap. (0.1 Gm) Bot. 100s
Use: Sedative.
- EVROVITE.** (Evron) Vitamins multiple. Cap., Bot. 100s.
Use: Dietary supplement
- EXACAPS.** (Approved Pharm. Corp) Vit. A 5000 u., D 500 u., B-1 2.5 mg., B-2 2.5 mg., B-6 0.5 mg., C 37.5 mg., niacinamide 20 mg., Cal. pantothenate 5 mg., folic acid 15 mg., Vit. B-12 1 mcg./Cap. Bot. 100s, 1000s.
Use: Multivitamin therapy.
- EXACHOL.** (Hart) Dehydrocholic acid 130 mg., desoxycholic acid 65 mg./Tab Bot. 100s.
Use: Biliary tract derangements.
- EX-CALORIC WAFERS.** (Eastern Research) Sod carboxymethylcellulose 181 mg., methylcellulose 272 mg./Wafer, Bot 100s, 500s, 5000s.
Use: (1-4 daily) Bulk laxative
- EXCEDIN.** (Bristol Myers) Solicylamide 2 gr., acetophenetidin 2 1/4 gr., caffeine 1 gr./Tab.
Use: Analgesic.
- EXERIN EXPECTORANT.** (Hoag) Pyrilamine maleate 80 mg., Sod. citrate 10 gr., ammonium chloride 10 gr., tartar emetic 1/2 gr., chloroform 2 min., alcohol 5%, menthol & mint flavor/30 cc. Bot. 1 pl.
Use: (1-2 tsp. q. 4 h) Expectorant.
- EXODYNE.** (Hart) Acetophenetidin 2 gr., acetylsalicylic acid 4 grs., caffeine 1/2 gr., tincture gelsemium 3 minims/Tab. Bot. 100s, 500s, 1000s.
Use: Anodyne, antipyretic, antispasmodic and sedative.
- EXONAL.** (Hart) Dehydrocholic acid 0.24 Gm. & dioctyl sod. sulfosuccinate 50 mg./Tab. Bot. 100s
Use: Constipation and hydrocholeretic
- EXORBIN.** (Ayers) Polyamine methylene resin, NND. 1959, Tab. 250 mg Bot 100s.
Paw. 1 Gm. Box 50
Use: Gastric hyperacidity.
- EXPANDEX, NND 1962.** (Commercial Solvents) Dextran 6% sol Bot. 500s.
Use: I.V.; to expand plasma volume.
- EXPASMUS.** (M. H. Smith) Dibenzyl succinate 125 mg., mephenesin 250 mg., solicylamide 700 mg./Tab. Bot. 100s.
Use: (2 q. 4 h) Antispasmodic, analgesic.
- EXPECTORANT, N.F.** (Marck Sharp & Dahme) Opium camphorated liq. 17.5 cc., senega fldst. 3.5 cc., squill 3.5 cc., ammonium carb. 1.8 Gm., alcohol 12%/100 cc. Bot 1 pt., 1 gal.
Use: (1-2 tsp) Expectorant.
- EXPICTO SYRUP.** (Paul Money) Dihydrocodeine bitartrate 1/27 gr., Sod. citrate 2 1/2 gr., ammonium chloride 1/2 gr., Pot. guaiaaculfonate 1 gr./dr. Bot. 1 pt., 1 gal.
Use: (1 tsp. q. 3-4 h, p.r.n) Cough due to colds.
- EXTOSEN.** (Squibb) Prophepryridamine maleate 25 mg., ammonium chloride 500 mg., ephedrine sulfate 75 mg., alcohol 5 1/2 %/fl. oz. Bot. 1 pt., 1 gal.
Use: (1-2 tsp. q. 3 h.) Cough due to colds.
- EXTRALIN, NNR, 1949.** (Lilly) Liver-stomach conc. equiv. to 240 Gm. of fresh liver/12 Cap. Bot 84s, 500s.
Use: Pernicious anemias.
- EXTRALIN B.** (Lilly) Liver-stomach conc. 500 mg., Vit. B-1 1 mg., Vit. B-2 0.25 mg., Pulv. Bot. 84s, 500s.
Use: Pernicious anemias.
- EXTRALIN F.** (Lilly) Stomach conc. w/intrinsic factor 220 mg., Vit B-12 15 mcg., folic acid 2 mg./Pulv. Bot. 30s, 500s.
Use: Pernicious anemias.
- EXUL** (Yorktown Products Co) Nupra (non-hormonal, nonsteroidic extract of beef organs, liver, brain, adrenals) 2000 Standard Chick units, dehydrated milk (55% fat) 14 Gm., thiamin HCl 0.6 mg., niacinamide 1.2 mg., ferrous gluconate 8.7 mg., cocoa, coconut oil, casein, sucrose, ethyl vanillin malt ext./Tab. (20 Gm) Tins, 5 Tabs. Carton, 50s
Use: (5 tab. daily) Gastric or duodenal ulcer
- EYE-MD.** (Winthrop Labs) A sol. containing zinc sulfate, boric acid & berberine sulfate. Bot. 1/2 fl. oz.
Use: Miliar irritations of the eyes.
- FACTIMIN.** (Rorer) Thiamine HCl 0.5 mg., cobalfavin 1 mg., niacinamide 6.6 mg., Cal. pantothenate 0.2 mg., pyridoxine HCl 0.1 mg., ferrous sulfate 3 gr./Tab. Bot. 100s, 500s, 1000s.
Use: Anemias.
- FALCO DIHYDROSTREPTOMYCIN SULFATE.** (Fallek Prods) Dihydrostreptomycin sulfate 10 Gm/Vial.
- FALGOS.** (American Ferment) Aspirin 3 gr., acetophenetidin 2 gr., caffeine 1/2 gr., w/ aluminum hydroxide & Mag hydroxide/Tab Bot. 15s, 40s.
Use: Analgesic.

FALVIN. (Lederle) Vit B 12 W/Autinnic in intrinsic factor concentrate 1 u, ferrous fumarate 274 mg, ascorbic acid 75 mg, folic acid 1 mg/Colp Bot 60s 500s
Use Hematinic

FARNOQUINONE. 2 Difornesyl 3 methyl 1,4 naphthoquinone

FASIGYN IN OIL. (Pfizer) Estradiol benzoate 2.5 mg, progesterone 125 mg/cc Vial 10 cc

Use Estrogen w/progesterone therapy

FATTY ACIDS, UNSATURATED

See Butrcaps (Stanley Drugs)

Fats Unsaturated

Linoleterol, Oint (Chicago Pharm)

Undecylenic Acid (Various Mlx)

FAT, UNSATURATED

See Arcofac (Armour)

Atheroxin, Lrq (Gray Pharm)

EFA (Columbus)

F Vitagen, Cap (Endocrine)

Linodaxin, Cap & Liq (Pfizer)

Lufa (US Vit Corp)

Saff (Abbott)

Saff La Safflower Oil (Safflower Prod Co)

Saffola (Saffola Labs)

F C A H (First Texas) Thénylpyromine HCl 25 mg, acetaphenetid n 2½ gr, salicylamide 2½ gr, atropine sulfate 1/500 gr, colchicine salicylated 1/700 gr/Cop Bot 100s 500s 1000s

Use Ant histaminic for common cold

F-CORTEF, NND 1962 See Cortel, F, Oint (Upjohn)

FE-1. (Marsh Emarh Labs) Ferrous fumarate 330 mg thiamine chloride 5 mg/Cop Bot 100s

Use Hemat nrc

FEBAL1. (Westerfield) Ferrous sulfate 250 mg, Vit B 1 08 mg, Vit B 2 0.5 mg, niacinamide 18 mg/dr Bot 4 oz, 1 pt, 1 gal

Use [2-4 tsp daily] Hematnrc

FEBEX B-12 (Kendall) Vit B 12 3 mcg, ferrous sulfate 10 mg, folic acid 0.5 mg, thiamine monontrate 15 mg, riboflavin 18 mg, niacinamide 20 mg, pyridoxine HCl 0.5 mg, panthenol 1 mg (equiv to 116 mg Cal panthenate), whole liver pow 130 mg, ascorbic acid 50 mg/Tab Bot 100s

Use: Hematnrc tonic

FERRO-GESIC. (First Texas) Acetaminophen 15 gr, caffeine 2 gr, sod salicylate 50 gr/A oz Bot 1 pt 1 gal

Use Analgesic—antipyretic

FERROLIN. (Buffington) N Acetyl p amlna phenol 125 mg, alcohol 14%/5 cc Bot 1 pt, 1 gal

Use: Antipyretic and analgesic

FEARB. (Ayerst) Ferrous carbonate Cap (3 gr/Cap) Bot 100s, 1000s

Use [3 daily] Iron deficiency anemias

FEDOFYL TABS & LIQ (Cambridge) Neothylline 100 mg, ephedrine sulfate 25 mg, butobarbital 30 mg/Tab or 10 cc Bot 100s, pls

Use Asthma

FEDRANIST. (Amer Pharm) Pyrilamine maleate 125 mg, ephedrine HCl 5 mg, ammonium chloride 1½ gr, caffeine citrate ½ gr/5 cc Bot 3 oz

Use: (1 tsp p r n) Antihistaminic, antispasmodic, expectorant, cough syr

FEDRANG. (Cole Chem Co) Ephedra 12 gr, Pot guaracolsulfanate 8 gr, sanguinaria ½ gr, squill 1 gr, thyme ½ gr, menthol ½ gr, lactic acid 3 min, Sod benzoate/ff oz of syr Bot 1 pt

Use: (12 tsp q 3-4 h) Cough

FEDRAZIL. (Burroughs Wellcome) Pseudoephedrine Hydrochloride 30 mg, Chlorzoxizone Hydrochloride 25 mg/Tab Bot 24s 100s

Use Respiratory tract congestion

FEE-B-CEE-PLUS (New Eng Pharm Co) Vit A 2500 u, B 1 1 mg, B 2 15 mg, B 6 0.1 mg, B 12 0.5 mcg, C 20 mg, D 250 u, folic acid 1 mg, niacinamide 10 mg, Cal panthenate 1 mg, Rutin 10 mg/Tab- plus 11 minerals and isopropraptic agents Bot 100s

Use Dietary supplement

FEGLUMIN. (Elder) Ferrous gluconate 5 gr, Thiamine HCl 1 mg/Tab Bot 100s

Use: (2 t i d, p c) Iron deficiencies

FELIRON (Rond) Vit B 1 2 mg, Vit B 2 2 mg, Cal panthenate 1 mg, niacinamide 20 mg, iron & ammonium citrate 125 mg, copper 1 mg, Vit B 6 0.2 mg, secondary liver traction 180 mg/Cap Bot 100s 1000s

Use Vitamin therapy

FELLO-SED. (Fellows) Chloral hydrate 0.5 Gm, Cal bromide 0.5 Gm, atropine sulfate 0.125 mg/4 cc Bot 8 oz

Use (1 tsp t i d & h s) Sedative & hypnotic

FELLO'RAL CAPSULES OR LIQUID (Fellows) Chloral hydrate 250 mg, hyoscyamine 0.100 mg, atropine 0.020 mg, hyosine 0.008 mg/Cap or 5 cc Bot 100s, Lrq 8 oz

Use: (1) Sedative & antispasmodic

FELSOL. (Amer Felsol) Antipyrine 0.87 Gm, caffeine citrate 0.1 Gm, iodopyrine 0.03 Gm, labeline sulfate 0.1 mg/Gm Pow Pkg (1 Gm) Box 15s 90s

Use: (1 Pkg t r d w/water p c) Analgesic & antasthmatic

FELSULES. (Fellows) Chloral hydrate 0.25

- Gm./Cap. Bot. 100s; 0.5 Gm./Cap. Bot. 50s
 Use: Sedative
- FEMACRINE INJ.** (Endocrine) Estradiol and natl. estrogen each 0.1 mg./cc. Vials 30 cc.; each 0.25 mg./cc. Vials 30 cc.; each 0.5 mg./cc. Vials 10 cc.
 Use: I.M., menopausal syndrome
- FEMACRINE-D.** (Endocrine) Dienestrol 0.5 mg., hexestrol 1 mg., Vit. E 15 I.U., thiamine HCl 5 mg., ascorbic acid 50 mg./Cap. Bot. 30s, 100s.
 Use: (1-3 daily) Menopausal symptoms
- FEMANDREN LINGUETS.** (Ciba) Ethinyl estradiol 0.02 mg., methyltestosterone 5 mg./Linguet, Bot. 30s, 100s.
 Use: Menopause, osteoporosis.
- FEMERGIN.** See: Ergotamine Tartrate (Various Mfr.).
- FEMESTRAL.** (Kirk) Estradiol (0.11 mg./cc.) Amp. 1 cc., Vial 30 cc.; (0.55 mg./cc. 10 cc. & 0.28 mg./cc.) Vial 30 cc.
 Use: I.M., estrogen therapy.
- FEMESTRONE.** (Kirk) Estrogens mixed, Parenteral, Amp. (10,000 u./cc.) 1 cc., 12s, 25s, 100s, Vial 10 cc., 30 cc. Vial (20,000 u./cc.) 10 cc., 30 cc., Tab. (1.25 mg.) 30s, 100s.
 Use: Orally & I.M., estrogen therapy.
- FEMESTRONE-AD.** (Moore Kirk Labs) Estradiol 3-trimethylacetate 5 mg., procaine HCl 5 mg., propylene glycol aqueous 24%/sc. Vial 10 cc.
 Use: Uterine bleeding.
- FEMESTRONE COMPOUND TABLETS.** (Kirk) Sod. estrane sulfate 1.25 mg., vit. B-1 3 mg., B-2 2 mg., niacinamide 10 mg., B 6 1 mg., calcium pantothenate 5 mg., C 25 mg., D 500 u./tab. Bot. 30s, 100s.
 Use: Treatment of menopausal syndrome.
- FEMIDYN.** See: Estrone (Various Mfr.).
- FEMOGEN.** (Testagar) Estradiol.
 Aq. Susp.—(0.2 mg./sc.) Vial 30 ss. Natural—(20,000 u., 30,000 u./cc.) 10 cc., 30 cc. In sesame oil—(0.1, 0.2, 0.5 mg./cc.) Vial 10 cc., 30 cc. Natural—(10,000 u., 20,000 u./sc.) 10 cc., 30 cc.
 Tabs.—1.25 mg. & 0.625 mg. conj. estrogen. Bot. 100s, 1000s.
 Use: Estrogen therapy.
- FEMOTRONE.** (Testagar) 10,000 I.U. of estrogen (as estradiols) & 10 mg. of progesterone in vegetable oil/cc. Vial 10 cc.
 Use: Primary & secondary amenorrhea.
- FEMSERT CREAM.** (Pharmafac) Dihydrodazyquinoline 100 mg., sulladiazine 250 mg., sullathiazole 250 mg., stilbestrol 0.7 mg., sodium lauryl sulfate 7 mg., beta lactose 10 mg., buffered to pH 4.5/5 Gm Tube 3 oz.
 Use: Trichomonas vaginalis.
- FINARSONE.** See: Carbarsona (Various Mfr.).
- FENATRATE.** (Sherman) Iron 0.084 Gm., Vit. B-1 1 mg., sucrose 0.130 Gm./Tab. Iron 1.35 Gm., Vit. B-1 8 mg., sucrose 14.8 Gm./A. oz. Tab., Bot. 100s, 500s, 1000s; Syr. 3 oz., 1 pt.
 Use: Iron deficiencies
- FENBANE.** (Walker Corp.) Phenobarbital 1/2 gr., hyoscyamus ext. 1/6 gr./Tab. Bot. 100s.
 Use: (1 q. 3 h.) Sedative & antispasmodic.
- FEND.** (Mine Safety Appliances)
 A-2—Water soluble cream which forms a physical barrier to water insoluble irritants. Tube 3 oz., Jar 1 lb.
 I-2—Water insoluble cream which forms a physical barrier to water soluble irritants. Tube 3 oz., Jar 1 lb.
 S-2—A silicone cream which forms a barrier against a combination of water soluble and water insoluble irritants. Tube 3 oz., Jar 1 lb.
 X—Industrial cold cream which rubs well into the skin and serves as a skin conditioner. Tube 3 oz., Jar 1 lb.
 Use: Protective creams against irritants of industrial and household origin.
- FENDOL.** (Bullington) Salicylamide 3 1/2 gr., acetophenetidin 2 1/2 gr., acetaminophen 2 gr., phenylephrine HCl 5 mg./Tab. Bot. 100s.
 Use: Analgesic, relief of the common cold symptoms.
- FENDON.** (A.P.C.) Acetyl p-aminophenol 5 gr./Tab. Bot. 36s.
 Use: Analgesic.
- FENOBEX.** (Strassenburgh) Phenobarbital Sod. 15 mg., thiamine HCl 1.8 mg., riboflavin 2.7 mg., niacin 32.7 mg., yeast conc. 25%, malt syr. 18%/15 sc. Bot. 1 pt., 1 gal.
 Use: (1 tsp. tid.) Sedative & hypnotic w/ Vit. B complex.
- FENOXYDYNE COMPOUND.** (Breon) Chlorphenylamine citrate 25 mg., acetophenetidin 320 mg., caffeine 32 mg./Tab. Bot. 25s, 100s, 500s, 1000s.
 Use: (1-3 daily) Common cold.
- FENYRIPOL HYDROCHLORIDE.** (Irwin, Neiser) *o*-(2-pyrimidinylaminoethyl)benzyl alcohol hydrochloride.
 Use: Skeletal muscle relaxant.
- FENTA.** (Superior Pharm) Tablets. Bots. 1000s
- FENT-O AUGINE EAR DROPS.** (Confield) Neomycin sulfate 6 mg., hydrocortisone 5 mg., pyrilamine maleate 0.6%, cetalkonium

FENT-O OPHTHINE EYE DROPS

306

- chloride 0.05%, dipredon HCl 1% /cc
Plastic drop bot 5 cc
Use Treatment of infections
- FENT-O OPHTHINE EYE DROPS (Confield)**
Polymyxin B sulfate 16,500 u, hydrocortisone 5 mg, neomycin sulfate 5 mg /cc Bot 5 cc
Use Inflammation
- FENTON W/VITAMIN C.** (Winthrop) Ferrous gluconate 300 mg, Vit B 12 5 mcg, C 50 mg, dried pyloric mucosa 25 mg /Tab
Use Anemias
- FENTROPINE.** (Smith, Miles & Patch) Propylfenit citrate 60 mg, hamotropine methylbromide 25 mg /Tab Bot 100s 1000s
Use (1 2 1 1 d) Antispasmodic
- W/Butobarbital (C D Smith) Propylfenit estrate** 60 mg, hamotropine methylbromide 25 mg, butobarbital sod 75 mg /tab Bot 100s 1000s
Use (1 2 1 1 d) Antispasmodic w/sedative
- FENTROPINE INJECTION.** (Smith, Miller, Patch) Propylfenit HCl 450 mg, hamotropine methylbromide 25 mg, urethane 0.1 Gm /cc Ampuls 1 cc Box 5s
- FEOPLEX** (Walker) Thiamine HCl 1 mg, riboflavin 1 mg, pyridoxine HCl 0.3 mg, Cal pantothenate 1 mg, niacinamide 10 mg, ferrous sulfate exsic 200 mg /Cap Bot 50s 500s 1000s; Drops, 30 cc, 1 pl
Use Nutritional anemias
- FEOPLEX B-12 DROPS** (Walker) Vit B 12 10 mcg, thiamine HCl 2 mg, B 2 2 D mg B 6 0.6 mg Cal pantothenate 20 mg, niacinamide 20 mg, iron & ammonium citrates (green) 400 mg /cc Dropper Bot 30 cc
Use Tonic hematopoietic
- FEOSOL ELIXIR** (Smith, Kline & French) Each 2 fl dr supplies 5 gr ferrous sulfate Bot 12 oz
Use (1 tip 1 1 d) Iron deficiency anemias
- FEOSOL PLUS.** (Smith, Kline & French) Ferrous sulfate exsic 200 mg, desic lives 325 mg, Vit B 12 (activity equiv) 17 mcg, folic acid 0.4 mg, thiamine HCl 2 mg, riboflavin 2 mg, nicotinic acid 10 mg, pyridoxine HCl 1 mg, ascorbic acid 50 mg, pantothenic acid 2 mg /Cap Bot 100s
Use Metabolic deficiencies
- FEOSOL SPANSULES** (Smith, Kline & French)—excitiated $FeSO_4$ —150 mg /Tab Bot 30s & 250s
Use Sustained release action of Ferrous sulfate in the treatment of iron deficiency anemias
- FEOSOL TABLETS** (Smith, Kline & French) Exsic ferrous sulfate 3 gr /Tab Bot 100s 1000s
Use Iron therapy
- FEOSORB.** (Smith, Kline & French)
Use: Iron vitamin preparation
- FEOSTAT.** (Lloyd, Dabney & Westerfield) Ferrous fumarate 60 mg, or 200 mg / Chewable Tab Bot 100s
Use: Iron deficiency anemia
- FEOSTAT SUSPENSION** (Lloyd Dabney & Westerfield) Ferrous fumarate 60 mg /5 cc 5 oz w/eolibrated droppers
Use: Anemia
- FEOSTIM.** (Lloyd, Dabney & Westerfield) Vit B 12 5 mcg, leucous fumarate 60 mg /chewable tab
Use: Treatment of iron deficiency anemia
- FEPISEM** (Mastengill) Ephedrine sulfate $\frac{1}{2}$ gr, acetophenetidin 3 gr, pentobarbital sodium $\frac{1}{4}$ gr atropine sulfate 1/150 gr /Cap Bot 100s, 1000s
Use Bronchial asthma, coryza and chronic sinusitis
- FER-A-BIN.** (Paul Maney) Ferrous sulfate 5 gr, thiamine HCl 1 mg /Tab Bot 100s 1000s
Use Iron therapy
- FERAD.** (Burroughs Wellcome) Ferrous sulfate anhyd 3 gr, Sod carbonate anhyd 2 gr /Tab Bot 100s, 300s, 1000s
Use Hematinic
- FERAD-B.** (Burroughs Wellcome) Ferrous sulfate anhyd 0.195 Gm, thiamine HCl 1.0 mg, riboflavin 1.0 mg, niacinamide 7.5 mg, pyridoxine HCl 0.25 mg, Cal pantothenate 2.5 mg /Tab Bot 100s, 300s, 1000s
Use (3 daily) Anemia due to insufficient iron
- FERANCEE.** (Stuart) Ferrous fumarate 200 mg, Vit C 150 mg /Softab Bot 100s
Use Treatment of anemia
- FERAPLEX.** (Diamond) Polysaccharide-iron complex solution containing equivalent of 100 mg elemental iron /10 cc Bot 12 oz Also with B 1 & B 12 and also with Mult vitamins
Use Anemia
- FER-ARSENIC.** (Metropolitan) Ferric chloride $1\frac{1}{2}$ gr, Sod dimethylarsenate 5 gr w/ chlorides of Cal, Pot & Sod /10 cc Val 10 cc
Use Iron deficiency
- FER-B-PLEX.** (Paul Maney) Ferrous sulfate 5 gr, Vit B 1 5 mg Vit B 2 2 mg, pantothenic acid 186 gamma Vit B 6 125 gamma mo niacinamide 5 mg, liver fraction 2 $\frac{1}{2}$ gr /Tab Bot 100s, 1000s
Use Iron & vitamin therapy
- FERBRISOL.** (Tilden) Acetophenetidin 16 gr acetanilid 10 Gm, total 4 gr caffeine 2 gr /8 oz Bot 8 oz
Use (1 2 3p 1 1 d) Antipyretic & analgesic
- FER-B-TELVE.** (Paul Maney) Ferrous glu

- conote 300 mg., thiamine HCl 1 mg., Vit. B-12 conc. 5 mcg., desic. stomach conc. 100 mg./Tab. Bot. 100s, 1000s.
Use: Anemias.
- FER-DONA.** (Fres-Cameron) Ferrous sulf. dS mg., cap. sulf. 0.333 mg., Vit. B-1 0.33 mg., B 2 0.5 mg., niacinamide 333 mg., pepsin 19.5 mg., liver fraction, frac. #2 227 mg., desiccated liver 227 mg./Cap. Bot. 100s.
Use: Hematinic vitamin.
- FER-DONA ELIXIR.** (Fres-Cameron) Thiamine HCl 4 mg., riboflavin 6 mg., niacinamide 40 mg., iron, as green iron, & ammonium citrates 200 mg., copper, as inherent cupric compounds, 2.8 mg., soluble liver fraction & secondary anemia fraction 6R gr./R. oz. Rot. 12 oz.
- FERGON.** (Winthrop Labs.) Pure ferrous gluconate stabilized to maintain a minimum of ferric ions, containing not less than 11.5% of iron, of which approx. 95% is in the ferrous state. Tab. of 2½ gr., Bot. 100s; Tab. of 5 gr., Bot. of 100s, 500s, 1000s. Elixr (6%), Rot. 16 fl oz, 1 gal.
Use: Anemias & Iron therapy.
- FERGON-C.** (Winthrop Labs.) Fergon 450 mg., ascorbic acid 200 mg./Cap. Bot. 100s
Use: (1 l.i.d) anemia
- FERGON COMPOUND ELIXIR.** (Winthrop Labs.) Fergon 0.25 Gm. vit. B-1 5 mg., vit. B-2 2 mg., vit. B-6 1 mg., vit. B-12 10 mcg., folic acid 0.5 mg., nicotinamide 10 mg., panthelol 0.94 mg./5 cc. Bot. 1 pt., 1 gal.
Use: Dietary supplement.
- FERGON PLUS IMPROVED.** (Winthrop Labs.) Ferrous gluconate 500 mg., vit. B-12 w/intrinsic factor conc. ½ u., folic acid 1.5 mg., ascorbic acid 75 mg./caplet. Rot. 100s, 500s.
Use: Hematinic.
- FER HEPARUM B-1.** (Torqian) Vit. B-1 3 mg., ferrous gluconate 35 mg., liver conc. Iron 25 Gm. liver/cc. Amp. 1 cc.
Use: 1 M. (1 or 2 cc. weekly) Anemias.
- FERRI-CAC.** (Smith, Miller & Patch) Iron cacodylate 30 mg., Sod. cacodylate 120 mg., strychnine sulfate 7 mg., Sod. glycerophosphate 120 mg./2 cc Vials 50 cc. 100 cc.
Use: Tonic.
- FERRIDEXATE.** (U.S. Standard) d Desoxyephedrine HCl 5 mg., ferrous gluconate 2 gr., ascorbic acid 35 mg./Tab. Rot. 100s, 1000s.
Use: (1 l.i.d., a c.) Obesity.
- FER-IN-SOL.** (Mead Johnson) Sol. of ferrous sulfate 125 mg., ferrous sulfate (25 mg. iron)/cc. Bot. w/dropper 15 cc., 50 cc.
Use: Iron-deficiency anemias.
- FERRISORB TABLET.** (Reid) Ferrous fumarate 200 mg., ascorbic acid 120 mg., pyridoxine HCl 6 mg./Tab. Bot. 100s.
Use: Iron deficiency.
- FERMALOX.** (Rorer) Iron sulfate 200 mg., maalox 200 mg./Tab. Bot. 100s.
Use: 1 Tab. b.i.d; hematinic.
- FERMATIN.** (Rorer) Ferrous sulf. 200 mg., folic acid 1.5 mg., Vit. B-12 7.5 mcg., C 75 mg., d-sorbitol 150 mg., maalox 200 mg./Cap. Bot. 100s, 500s.
Use: Treatment of anemias.
- FERNISOLOONE.** (Ferndale) Prednisolone 5 mg./Tab. Rot. 100s, 1000s.
Use: Steroid hormone.
- FERNISONE CREAM 1%.** (Ferndale) Hydrocortisone acetate 1%. Tube ½ oz.
Use: Dermatological.
- FERO-GRADUMET.** (Abbott) Ferrous sulfate 525 mg./Tab Bot. 100s
Use: Iron-deficiency anemias.
- FERO-GRADUMET TARS.** (Abbott) Ferrous sulf. U.S.P. 525 mg. in controlled release form./Tab. Rot. 100s, 1000s.
Use: Hematinic.
- FERRALYN.** (Loanett) Ferrous sulf. 3 gr./tab. E.C. Bot. 1000s, 5000s. Green & red tabs.
- FEBBATED LIVIB W/Folic Acid.** (Upjohn) Ferrous sulfate 3.6 gr., Vit. B-1 2 mg., Vit. B-2 2 mg., niacinamide 10 mg., folic acid 2 mg., Vit. B-12 2 mcg., liver conc. 7 gr./Tab. Bot. 100s.
Use: (1 t.i.d) Hematinic.
- FERRATED LIVER CONCENTRATE.** (Upjohn) Ferrous sulfate 33/5 gr., liver conc. 7 gr., Vit. B-12 2 mcg., thiamine HCl 0.5 mg., riboflavin 1 mg., nicotinamide 5 mg./tab. Bot. 100s, 500s.
Use: Hematinic.
- FERRI-RETALIN.** (Billy) Ferric citrochloride r. 4 cc., Thiamine HCl 14.5 mg./R. oz. Bot. 1 pt., 1 gal.
Use: Secondary anemias.
- FERRIC AMMONIUM SULFATE.** (Various Mfr.)
Use: Astringent & styptic.
- FERRIC AMMONIUM TARTRATE.** (Various Mfr.)
Use: Iron deficiency.
- FERRIC CACODYLATE, N.F. X.** (Various Mfr.)
Use: Leukemias & iron deficiency.
- FERRIC CACODYLATE—FARNSWORTH FERRIC CACODYLATE.** Amp. 5 cc., 12s.
- FERRIC-R Jr.** (Jenkins Labs) Vit. B-12 5 mcg., B-1 2 mg., C 30 mg., ferric glycerophosphate 50 mg./tab. Bot. 100s, 1000s.
Use: Nutritional deficiencies.
- FERRIC CHLORIDE, A.D.R.** Sol., N.F.; Tr., N.F. (Various Mfr.)
Use: Astringent for skin disorders.

* Generic name recognized by N.F., U.S.P. or A.M.A.

FERRIC "PEPTONATE." (Various Mfr) See: Iron Peptonized

FERRIC QUININE CITRATE, "GREEN" (Various Mfr)

Use For iron deficiency

FERRIC QUININE & STRYCHNINE CITRATE. (Various Mfr)

Use As a tonic

FERRIC SUBSULFATE SOLUTION, N.F., A.D.R. (Various Mfr)

Use Local use on the skin

FERRIDYL (Ascher) Sod cocodylate 320 mg, ferric chloride 150 mg /vial Vial 10 cc

Use Anemias

FERRIDYL-B (Ascher) Ferric chloride 1½ gr, Sod cocodylate 2½ gr, Vit B 12 50 mcg /5 cc Amp 5 cc, Box 6s, 25s

Use Ant anemia therapy

FERRI-FOLEX. (Ascher) Vit A 5000 u, D 500 u, B 1 2 mg, B 2 2 mg, C 30 mg, niacinamide 10 mg, pyridoxine HCl 0.5 mg, Col pantothenate 3 mg, iron 10 mg, stomach 50 mg, liver 150 mg, yeast 100 mg w/Col, Phos, Mag, cobalt, copper & zinc/Tab Bot 100s

Use Vitamin & mineral therapy

FERRI-FOLEX FORTE. (Ascher) Vit A 5000 u, D 500 u, B 1 2 mg, B 2 2 mg, C 100 mg, niacinamide 15 mg, Vit B 6 0.1 mg, B 12 3 mcg, folic acid 1 mg, Col pantothenate 1 mg, iron 100 mg, stomach 50 mg, liver 150 mg, yeast 100 mg, w/copper, Mang, zinc, cobalt/Tab Bot 100s, 1000s

Use Multivitamin w/minerals

FERRILEX LIQUID (Smith Ossey) liver fraction 550 mg, thiamine HCl 10 mg, riboflavin 10 mg, pyridoxine HCl 2 mg, Col pantothenate 5 mg, niacinamide 50 mg, glycerite of red bone marrow 8 gr, iron peptonate 8 gr cobalt sulf 30 mg, choline 65 mg, and inositol 30 mg /fl oz Bot 1 pt 1 gal

Use Secondary anemias

FERRIPLEX FORTE. (Myers Carter) Liver int 0.1 cc, iron peptonized 59 mg, Vit B 2 0.3 mg, B 12 30 mcg, B 6 0.3 mg, niacinamide 50 mg, folic acid 2 mg /2 cc Vial 30 cc

Use Secondary anemias

FERRITRINSIG, TAB (Upjohn) Vit B 12 & intrinsic factor conc ½ v, ferrous sulfate 300 mg, desic liver 100 mg, folic acid 2 mg, ascorbic acid 50 mg, Vit B 1 2 mg, B 2 2 mg, niacinamide 10 mg /Tab Bot 50s, 500s, 1000s

Use Nutritional anemias

FERRO-ARSEN (Breon) Each 5 cc contains Sod cocodylate 0.16 Gm, ferric chloride

* Generic name recognized by N.F., U.S.F. or A.M.A.

0.075 Gm, chlorides of Col, Pot & Sod Amp, 5 cc, 10 cc, Box 6s, 25s

Use Secondary anemias

FERRO-B. (Pitman Moore) Iron & ammonium citrate 45 gr, Mang citrate soluble 2 gr, thiamine HCl 1 B mg, yeast ext equiv to brewer's yeast (dried) 40 gr /fl oz Bot 1 pt, 1 gal Tabs Bot 100s, 1000s

Use Nutritional & secondary anemias

FERRO-BETALIN (Lilly) Ferrous sulfate 5 gr, Vit B 1 1 mg /Tab Bot 100s, 1000s

Use: (1 1 d) Hematinic

FERRO-BEVEX. (Melfa Labs) Vit B 12 10 mcg, B 1 10 mg, B 6 10 mg, B 2 0.1 mg, niacinamide 25 mg, pantothenate 10 mg, ferrous lactate 20 mg, liver (Vit B 12 active) 10 mcg /cc Vial 10 cc

Use Vitamin B complex w/liver & iron therapy

FERRO-B IMPROVED. (Pitman Moore) Iron & ammonium citrate 5.625 gr, Mang citrate soluble 0.25 gr, thiamine HCl 0.25 mg, yeast ext equiv to Brewer's Yeast (dried) 5 gr /Tab Bot 100s, 500s, 1000s

Use Nutritional & secondary anemias

FERRO-C. (Physiological Lab) Ferrous ascorbate Amp (20 mg) 2 cc, Box 10s, 25s

Use (3 gr 1 d) Hematinic

FERROCHOLATE. See Ferrocholine

***FERROCHOLINATE**, NND 1962 Ferrocholate Ferrocholine A chelate prepared by reacting equimolar quantities of freshly precipitated ferric hydroxide with choline dihydrogen citrate

See Chel Iron (Kinney)

FERROCHOLINE, NND 1960 See Ferrocholate

Use (3 gr 1 d) Hematinic

FERROCOL CAPSULES. (Mulford) 20% iron dispersed in or combined with a protective colloid made by the hydrolysis of casein Amp 25 cc Faw 20 Gm, 100 Gm Cap (5 gr) Bot 60s, 500s, 1000s

Use Iron deficiency anemias

FERRO-C TABLETS. (Physiological Lab) Ferrous ascorbate Tab (3 gr) Bot 100s

Use (1 1 d) Anemias

FERRO-CUPRIN. (Kremers Urban) Iron cocodylate 2½ gr, copper cocodylate 1/15 gr, Sod cocodylate 5 gr /10 cc Amp 5 & 10 cc, Box 25s

Use Hematinic

FERRO-DESICOL. (Forke, Davis) Ferrous sulfate 2 gr, desical 4 gr /Kopseal Bot 100s

Use: (1-2 qid) Anemias

FERRO DROPS (Forke Davis) Aqueous iron preparation containing ferrous lactate

equiv. to 25 mg. of iron/cc. Bot. w/dropper 15 & 50 cc.

Use: Iron-deficiency anemias.

FERROGALT. (Teitagar) Ferrus sulf. 0.2 Gm., cobalt chloride 15 mg./E.C. Tab. Bot. 100s.

Use: Hematinic agent.

FERRO-GEL. (Maurry) Ferric ammonium citrate 100 mg./cc. in gelatine base. Vial 10 cc.

Use: Iron deficiency.

* **FERROGLYCINE SULFATE COMPLEX.** See Ferronord (Nordmark)

FERROLIP, N.N.D. 1962. (Flint, Eaton) Ferrocholine (Iron choline citrate) 120 mg./Tab. Bot. 100s. Syr. (120 mg of elemental iron/oz.) Bot. 1 pt., 1 gal. Pediatric Drops (25 mg of elemental iron/cc. Sqs. Bot. 30 cc.

Use: Hematinic.

FERROLIP-OB. (Flint, Eaton) Tricalcium citrate 600 mg., Cal. gluconate 300 mg., iron choline citrate 150 mg., Vit. B-1 3 mg., B-2 3 mg., niacinamide 30 mg., Cal. pantothenate 10 mg., Vit. B-6 10 mg., C 200 mg., folic acid 0.5 mg., Vit. B-12 w/intrinsic factor 1 u., Vit. A 5000 u., D 500 u./3 Tab. Bot. 100s, 1000s.

Use: Orally (1-3 daily); hematinic in pregnancy & lactation.

FERROLIP PLUS. (Flint-Eaton) liver desic. 100 mg, iron choline citrate 200 mg, Vit. B-12 w/intrinsic factor conc. 1/2 unit, B-1 2 mg, B-2 1 mg, B-6 0.5 mg, C 50 mg, folic acid 0.5 mg./Cap. Bot. 100s, 1000s.

Use: (1-2 t.i.d.) Hematinic.

FERROLIP-PLUS LIQUID. (Flint, Eaton) liver fraction #1 200 mg., iron choline citrate 200 mg., folic acid 0.5 mg., Vit. B-12 10 mcg., B-1 2 mg., B-2 1 mg, B-6 0.5 mg./5 cc. Bot. B oz., 1 gal.

Use: Anemias.

FERROLIVEX. (New Mexico Pharm.) Vit. B-1 1.5 mg, B-2 2 mg, B-6 0.1 mg, calcium pantothenate 1 mg, niacinamide 5 mg, folic acid 0.5 mg, ferrus sulfate 26 mg, liver fraction 2.67 mg., yeast hydrolysate 1 mg./Tab. Bot. 100s, 1000s

Use: Iron deficiency anemias and vitamin deficiency

FERROMALT. (Borchardt) Ferrus sulfate 195 mg, copper sulfate 0.5 mg., malt soup extract 150 mg./Tab. Bot. 100s.

Use: Iron deficiency anemia.

FERRO-MANDETS. (Lederle) Ferrus fumarate-iron 20 mg, Vit. C. 50 mg./chew tab. Bot. 60s.

Use: Hematinic

FERROMANG W/Vitamins. (Drug Products)

* Generic name recognised by N.F. U.S.P. or A.M.A.

Iron & Mang. peptonized 1 Gm., thiamine HCl 3 mg., riboflavin 3 mg., butaben 0.05%/fl. oz. Ferrus gluconate 1 gr., Mang. hypophosphite 1/4 gr., thiamine HCl 1 mg., riboflavin 2 mg./Cap. Liq., Bot. B fl. oz., 1 pt. Pulvoids, Bot. 100s, 1000s.

Use: Hematinic.

FERROMIN. (McNeil) Ferrus carb. 200 mg., thiamine HCl 0.167 mg., riboflavin 0.333 mg., niacinamide 1.67 mg./Tab. Bot. 100s, 1000s

Use: Secondary anemias.

W/Liver Conc. (McNeil) Ferrus sulfate 3 gr., liver conc. 2 1/2 gr., thiamine HCl 0.3 mg./Cap. Bot. 100s, 1000s

Use: Anemias.

FERROMIN COMPOUND. (McNeil) Iron & ammonium citrates 12 gr., thiamine HCl 500 I.U., red bone marrow ext., malt conc./fl. oz. Bot. 1 pt., 1 gal.

Use: Hematinic.

FERRONORD. (Nordmark Pharm. Labs) Ferroglycine Sulfate Complex. Glycamine iron. Ferrus iron 40 mg./tab. Bot. 100s. Liq (40 mg of elemental iron/cc.) Bot. 60 cc.

Use: (2 b.i.d. between meals) Iron therapy.

FERRONORD PROGRAVA. (Nordson) Ferroglycine sulfate complex 125 mg (20 mg. elemental iron), Vit. A 4000 u, D 400 u, B₁ 2 mg., B₂ 1 mg., niacinamide 10 mg, B₆ 2 mg, B₁₂ 1/6 U.S.P. oral unit (with intrinsic factor concentrate), C 30 mg., K 0.5 mg, folic acid 0.2 mg, and seven minerals/Cap. Bot. 90s

Use: Iron-vitamin-mineral supplement.

FERROPHYL. (Lakeside) Ferrus sulfate extic. 200 mg., Sod. Pot. copper chlorophyllin 25 mg., Vit. B-12 2 mcg./Tab. Bot. 50s

Use: Hypochromic or secondary anemias.

FERROPLEXATE. (U.S. Standard) Thiamine HCl 1 mg., riboflavin 1 mg., ascorbic acid 30 mg, niacinamide 10 mg, Cal. pantothenate 1 mg., ferrus sulfate dried 51 mg, liver desic. 25 mg./Tab. Bot. 100s, 1000s.

Use: Anemias.

FERROPYRIN. (Knoll) Phenyl dimethylpyrazoloferric chloride Pow. 1:5. Bot. 1 oz.

Use: Hematinic.

FERROSENICUM. (Parke, Davis) Iron 3 gr., Mang. 1/2 gr., arsenic peptonate 4/25 gr./fl. oz. Bot. 1 pt., 1 gal.

Use: Hematinic.

FERRO-SEQUELS. (Lederle) Ferrus fumarate 150 mg, diethyl sod sulfosuccinate 100 mg./Cap T.D. Bot 30s, 1000s

Use: Anemia.

FERROSPAN. (Myers Carter) Ferrus fuma-

rate 200 mg (equiv to 65 mg iron)
d glucitol for absorption/Tab Bol 100s
Use Iron deficiency anemias

FERRO-THERON (Tilden) Ferrous sulfate 24 gr/A oz liq Bol pls, gols
Use Nutritional iron tonic in hypochromic anemias

FERROTONE-R, (Horton & Converse) Ferrous gluconate 3 gr, Vit B 1 3 mg, B 2 2 mg, B 6 0.5 mg, niacinamide 6 mg, pantothenyl alcohol 6 mg w/wheat germ ext. In orange flavored syr /15 cc Rat 1 pt
Use Hematinic, B complex therapy

FERROUS ASCORBATE. See: Ferro C, Tab, Amp (Physiological Lab)

FERROUS BROMIDE (Various Mfr)
Use: In chorea & tuberculous cervical adenitis

FERROUS CARBONATE MASS, NF X Pills, NF Vallets mass (Various Mfr) See: Fercarb Cap (Ayerst)
Use: For iron deficiency

W/Cuprous carbonate See Cupron, Cap (Ayerst)

FERROUS CARRONATE, SACCHARATED, NF X (Various Mfr)
Use For iron deficiency

FERROUS COMPOUND, (Kirk) Iron cacodylate 1/6 gr, Sod glycerophosphate 1 1/2 gr, strychnine sulfate 1/200 gr/cc Amp 1 cc, Box 12s, 100s
Use Hematinic

***FERROUS FUMARATE**, NND 1962
Tab 200 mg Bol 100s, 1000s

See C Ron Fediatic (Rowell)
Ferro Mandets (Lederle)
Ferro Sequels (Lederle)
Ferraspan, Tab (Myers Carter)
Fumasorb (Marion)
Fumiran (D rect, Standex)
Hematate (Kirkman)
Ircan (Lakeside)
Fontrinsic Tab (Brown Pharm)
Prematinic (Ingram Pharm)
Toleran (Mallinckradt)

W/Ascorbic Acid
See Cevrate Tab (Physicians Prod)
Cytoferin, Tabs (Ayerst)
Furan Suspension (Metra Med)
Palaran (Smith, Miller Patch)
W/Vit C copper sulfate, etc See Rocyte Tabs & Inj (Sig)

W/Ascorbic Acid Vit A, D, purified powdered bone See: C Ron Prenatal, Tabs (Rowell)

W/Ascorbic Acid, Vit B-12 See: Eruocyte, Tab (Miller)

W/Diethyl sodium sulfasuccinate See Per sanatinic, Tabs (Person & Covey)

* Generic name recognized by NF, USP or AMA

W/Thiamine See: Arne Timesules Caps (Hoag)

W/Vitamin B 12, thiamine HCl, Ilyune d sorbital See: Crystlivite Fe Sorbettes Tab (Hass)

***FERROUS GLUCONATE**, USF Tab, USF See: Fergon Preparations (Winthrop Labs)
Irox, Tab (Cole Chem Co)

Nionate, Tab (Nion)
Ray Gluciron, Tab (Raymer)
Use: Iron deficiency

W/Ascorbic acid, thiamine mononitrate See Gluco Ferrum w/C (Vanpell & Brown)

W/Cobalt & Vit B 12 & B 1 See Hemo Vital Liquid (Smith)

W/Polyoxyethylene glucitan monolaurate See: Simran, Cap (Merrell)

W/Vit C See C Ron Tab (Rowell)

FERROUS IODIDE, (Various Mfr)
Use In chronic tuberculosis

FERROUS IODIDE SYRUP, NF (Various Mfr)
Use In chronic tuberculosis

FERROUS LACTATE, NNR 1949 (Various Mfr)
Use For iron deficiency

***FERROUS SULFATE**, USP, Exsic, USF Syr, USF XV; Elixir USF, Tab USF See: Feosol, Spansule, Tab (Smith, Kline & French)

Ferrotlyn, Tab EC (Lonnell)
Ferro Theron, Liq (Tilden)
Irosul, Tab (Haskell)

Sulferrous, Pill (Chicago Pharm)
W/Cobalt chloride See: Rancovite, Prop (Hoyd)

W/Dicalcium phosphate & Vit D See Calc D Ferrous, Tab (Irwin Neisler)

W/Glutamic acid HCl See: Feroma, Cap (Sharp & Dohme)

W/HCl See Acidrin Tab (Walker)
W/Vitamins See: Vitamin combinations

W/Vitamins B (Squibb) Ferrous sulfate, exsic 3 1/2 gr, riboflavin 1 mg thiamine HCl 1 mg, niacinamide 6 mg/Tab Bol 100s, 1000s

Use: Treatment of secondary anemias
FERRO-ZEM PLUS ELIXIR (Zemmer) Iron phosphate 720 mg, thiamine HCl 8 mg riboflavin 8 mg, ascorbic acid 8 mg/A oz Bol pls

Use: Tonic hematinic
FERRUGINOUS COMPOUND, (Mallard) I cacodylate 1/6 gr, Sod glycerophosphate 1 1/2 gr, strychnine cacodylate 1/20 gr, cacodylic acid 0.003 Gm/cc Amp 1 cc, Box 12s, 50s, Vial 30 cc

Use: Hematinic
FER S-B (Dumas Wilson) Ferrous sulfate

- 3 gr., yeast cont. 2 gr./Tab. Bot. 100s.
Use: (1 t.i.d.) Hematinic.
- FERUBEX PALATAS.** (Central) Ferrous peptonate 130 mg., Vit. B-12 2 mg., B1 0.5 mg., B-2 0.5 mg., niacinamide 3 mg., pantothenic acid 0.5 mg., Vit. B 6 0.3 mg., folic acid 0.25 mg./Tab. Bot. 100s, 500s, 1000s.
Use: (1 daily) Anemias.
- FERVENULIN.** (Upjohn) An antibiotic produced by streptomycetes *fervens*.
Use: Trichomonas infections
- FERYL.** (Central) Ferrous sulfate 136.1 mg., Vit B-1 0.333 mg., liver concentrate 194.4 mg./Cap. Bot. 100s, 500s, 1000s
Use: Anemia due to iron deficiency
- FESTAL.** (Lloyd Brothers) Lipase 10 W.U., amylase 10 W.U., protease 17 W.U., hemicellulose 50 mg., bile, constituents 25 mg./Tab Bot. 100s
Use: Enzymes for digestion.
- FETAMIN.** (Mission) d Melthomphetamine HCl 50 mg., pentobarbital sodium 20 mg., Vit. A acetate 2500 u., Vit. D-2 250 u., C 10 mg., B-1 20 mg., B 2 20 mg., B 3 50 mg., B-5 1.0 mg., B 6 10 mg., ferrous gluconate 65 mg., iron 7.5 mg., calcium lactate 270 mg., calcium 35 mg., copper (as sulfate) 0.15 mg., manganese (as citrate soluble) 0.25 mg., zinc (as oxide) 0.08 mg., potassium (as chloride) 50 mg., magnesium (as carbonate) 25 mg./Tab Bot. 100s.
Ure: (1 t.i.d.) Dietary supplement in control of obesity
- FEVERALL.** (Medics) Dipyrone. Tabs. (500 mg) Bot. 100s; Elixir (500 mg/5 cc) Bot 1 pt., Inj (500 mg/cc) Vial 10 cc, 30 cc.
Use: Analgesic
- FEVONIL.** (Carrlone) Dipyrone 0.5 Gm./5 cc. or Tab Bot 1 pt. Tabs. Bot 100s.
Use: Analgesic
- FEXEROL.** (Ferler Co) Calcium hypochlorite containing active chlorine 3% to 4%. Pow. Bot. 5 Gm. Box 3s
Use: Dilute before using, in moniliasis and trichomoniasis
- FIBRIN INDEX THROMBIN.** (Ortho) Amps., Pkg-12s
Use: Diagnostic aid for critical fibrinogen index
- FIBRIN FOAM.** N.N.R. 1954. Sterile human fibrin in yellowish rectangular spongelike masses. Vials. See: Fibrin Foam & Thrombin (Human). (Cutter).
Use: Topical coagulant to oozing surfaces.
- FIBRIN FOAM & THROMBIN.** N.N.R. 1954, A.D.R. (Cutter) Fibrin foam 250 mg. w/5 cc. vial of thrombin, Pkg. 1s
Use: Hemostatic.
- * Generic name recognized by N.F., U.S.P. or A.M.A.
- FIBRIN HYDROLYSATE.** See: Aminosal, Sal. (Abbott)
- * **FIBRINOGEN (HUMAN).** U.S.P., N.N.D. 1962. Partially purified fibrinogen prepared by fractionation from normal human plasma. See: Fibrinogen, Vial (Merrell).
Farenogen, N.N.R. (t.v.), Vial. (Cutter)
- FIBRINOGEN HUMAN, IRRADIATED.** (Merck Sharp & Dahme) Prepared from normal human plasma. 2 Gm/Bot. with injection outfit.
Use: Hypofibrinogenemia
- * **FIBRINOLYSIN (human).** N.N.D. 1962. Plasmin. An enzyme obtained from human plasma by conversion of profibrinolysin with streptokinase to fibrinolysin
See Actose (Orlha)
Thrombolysin Lyovac (Merck Sharp & Dahme)
- W/Desoxyribonuclease. See: Elase, prep. (Parke, Davis)
- FIBROGEN (ORAL).** N.N.R. 1944. (Merrell) Tissue fibrinogen, Vial (1.5%) 3 cc., Box 4s.
Use: Orally; hemostatic.
- FIDALAC.** (Beard) Calcium lactate 500 mg., Vit. D₃ 400 U.S.P. units, ferrous fumarate 200 mg., C 50 mg./Tab Bot 100s, 1000s
Use: Calcium-iron deficiencies.
- FILISON.** (Lederle) Vit. A 4000 u., D 400 u., B-1 3 mg., B-6 1 mg., niacinamide 10 mg., B 2 2 mg., B-12 W/Int. loc. 1/6 oral u., C 50 mg., K 0.5 mg., folic acid 1 mg., ferrous fumarate 90 mg., intrinsic factor 5 mg., iron 30 mg., Ruarine 0.015 mg., copper 0.15 mg., iodine 0.01 mg., pal. 0.835 mg., mang. 0.05 mg., mag 0.15 mg., malyb 0.025 mg., zinc 0.085 mg., cal. carbonate 375 mg./Cap. Bot. 100s
Use: Supplement to the diet during pregnancy and lactation
- FILMERINE.** (Novocal Chem) Sod. nitrite conc. soln.
Ure Antirust agent
- FINAC LOTION.** (C & M Pharnacal) Sullur 2%, hyamine 10% 0.1%, in a flesh-tane base 2 fl. oz.
Use: Acne therapy
- PINE-SUL.** (Kendall) Sulfacetamide 250 mg., mandelic acid 136 mg., methenamine 125 mg./tab. Bot. 100s, 500s, liq. Bot. 8 oz
Use: 4 t.i.d., p.c.s. urinary tract infection.
- FIORINAL.** (Sandoz) (Sandoptal) Isobutyl-ethyl barbituric acid $\frac{3}{4}$ gr., caffeine $\frac{3}{8}$ gr., aspirin 3 gr., acetophenetidin 2 gr./Tab. Bot. 100s, 1000s
Use (1 q. 3-4 h. m.p) Sedative & analgesic
- FIROLYPTOL** (Tilden) White pine 55 gr.,

wild cherry 5½ gr eucalyptol 2 min, cotton seed oil ½ fl oz/fl oz Bot pts

Use Expectorant eutrophic
FIROLYPTOL & KREOSOTE. (Tilden) Kreosote 4 min, white pin 55 gr, wild cherry 5½ gr eucalyptol 2 min, cotton seed oil ½ fl oz/fl oz Bot pts
 Use Expectorant eutrophic
FIRON, NND 1962 (G A Beard) Ferrous fumarate 200 mg/Tab Bot 100s
 Use Iron deficiency

FIRWEIN (Tilden) Sodium bromide 2½ gr/fl oz with white pine and wild cherry Bot pts gals
 Use Sedative expectorant
FI-TEST. (Hyland) Anti body to human fibrinogen and polystyrene particles
 Use Determination of fibrinogen deficiency in blood

FIVE-B-VITAMINS (Raymer) Thiamine HCl 2 mg riboflavin 1 mg, pyridoxine HCl 0.1 mg, Cal pantothenate 0.25 mg, niacin 10 mg dried brewer's yeast 325 mg/Cap Bot 100s 500s, 1000s
 Use Dietary supplement

FIZRIN POWDER (Glenbrook) Aspirin, sodium bicarbonate sodium carbonate, citric acid/pack 1 dose foil packets
 Use Analgesic antacid

FLAGYL (G D Searle) Metronidazole
 Use Trichomonocidal agent
FLAMOTIDE. (Philo Amp) Riboflavin 5 phosphate Sodium 50 mg/cc Vial 10 cc
 Use Ariboflavinolic lesions

LAHITHIN (Toblerack) Glutamic acid HCl 5 gr, thiamine ½ gr riboflavin 0.67 mg niacin 5 mg/Cap Bot 100s 500s
 Use: (1 q 3 h) Achlorhydria or hypochlorhydria

LAV-A-D (Kirkman) A new name for Vitamin
LAVASCORB (Kirkman) Ascorbic acid 100 mg/Tab Bot 100s

FLAVAXIN (Winthrop Labs) Syn riboflavin Tab, (1 mg) Bot 25s, 500s, Tab (5 mg) Bot 25s
 Use Vitamin B therapy

FLAVAXIN SOLUBLE (Winthrop Labs) The double salt of riboflavin Sod Sod tetraborate buffered with Sod tetraborate & boric acid in the instantly soluble (naphthoid) form suitable for subcutaneous, intramuscular & intravenous administration Amp (5 mg) Box 25s

FLAVIHIST. (Amiab) Neomycin, tyrothricin cetyl dimethyl benzyl ammonium chloride benzocaine/lozenge Vial 12s
 Use: Minor throat irritations

FLAVINE See Acriflavine Hydrachloride (Various Mfr)

FLAVINOL-C. (Walker) Hesperidin complex 200 mg, ascorbic acid 200 mg/Cap Bot 24s, 100s
 Use: Capillary fragility and the common cold

FLAYISAL (Columbus) Pot para amino-benzoate 5 gr, pot salicylate 5 gr, hesperidin concentrate 50 mg, Vit C 50 mg/Tab Bot 100s, 1000s
 Use: Rheumatoid diseases

FLAVO-C. (Damsta) Hesperidin 200 mg, ascorbic acid 200 mg/Tab Bot 100s
 Use: Respiratory infections

FLAVOCILLIN ORAL (Philo Ampoule) Buffered crystalline penicillin G potassium powder 3 000 000 u Pwd (60 cc/250 000 u/5 cc)
 Use Pediatrics and patients who require fluid medication

FLAVOLUTAN. See Progesterone (Various Mfr)

FLAVONOID COMPOUNDS
 See: Bio Flavonoid Cpd
 Vit ?

FLAVOSERP. (Blue Line) Reserpine 0.25 mg, hesperidin 100 mg, ascorbic acid 100 mg/Cap Bot 100s, 1000s
 Use Antihypertensive tranquilizer

FLAVUROL. See Merbreamin (Hynson Westcott & Dunning; Prema)

FLAXEDIL TRIETHIODIDE, NND 1962 (Lederle) Gallamine triethiodide salt 100 mg/cc amp 1 cc Vial 200 mg in 10 cc
 Use Similar to curare Relax skeletal muscle

FLEET ENEMA (C B Fleet) Sod biphosphate 16 Gm, Sod phosphate 6 Gm/100 cc Bot w/rectal tube 4½ oz Pediatric size 2½ oz
 Use Laxative

FLEET OIL RETENTION ENEMA (C B Fleet) Mineral Oil 4¼ fl oz in an unbreakable vinyl squeeze bottle
 Use To soften stools

FLEET THEOPHYLLINE RECTAL UNIT (Fleet) Unit delivers of theophylline 0.25 Gm Aqueous Soft Pack 6s
 Use Rectally relief of acute bronchial asthma

FLEX-O-CASE PASEE (Ulmer) Zinc Gelatin USP Jar 1 lb
 Use Gelatin paste for Unna's boot

FLIKTENA (Rorer) Boric acid benzoic acid betanaphthol, zinc oxide salicylic acid & phenol 0.5% Jar 25 Gm 4 oz, 1 lb
 Use Fungicide

FLO-CILLIN AQUEOUS (Bristol) Cryst procaine penicillin G in aq susp One dose vials 3 000 000 u, & also in Brist O Malt

- Disposal Syr., 600,000 & 1,000,000 u. per dose.
 Use: In repository penicillin therapy.
- FLO-CILLIN "96," A D B.** (Bristol) 300,000 u. per cc. of crystalline procaine penicillin G in oil, jelled with 2% aluminum monostearate as a dispersing agent. Viol 1 cc. & 10 cc.
 Use: Repository penicillin therapy.
- FLO-CILLIN "96" FORTIFIED.** (Bristol) 100,000 u. of crystallin Pot. penicillin G plus 300,000 u. of procaine penicillin G in oil containing 2% aluminum monostearate. Sesame oil, 10-dose vial; peanut oil, 10-dose vial.
- FLORAMINE.** (Lemmon) Phenobarbital 15 mg., hyoscyamus est. B mg., passiflora 60 mg., valerian 60 mg., thiomine HC 1 mg./Tab. Bot. 100s
 Use: Nervous Insomnia
- ***FLORANTYRONE,** NND 1962. gamma-Fluoranthun-B-, 1 gamma-oxibutyric acid. See Zanol, Tab (Searle)
- FLORAQUIN.** (Searle) 100 mg. of diadoquin (dilatohydroxyquinoline) per vaginal tablet or per each 1.4 Gm. of the powder, together with lactose, dextrose & boric acid. Tab., Pkg. 24s; Paw., Bot. 3 oz., 8 oz.
 Use: Vaginal Trichomonas & vaginalis vaginitis.
- FLORIDINE.** (Drug Specialties) Sodium fluoride 2.21 mg., Vit. B-6 3 mg., C 30 mg., D 400 u./Lozenge. Bot. 100s.
 Use: Dental caries prevention
- FLORINEF ACETATE,** NND 1962. (Squibb) Fluorohydrocortisone Oint. (0.1 & 0.2%) Tube 5 & 20 Gm
 Use: Many types of dermatitis & skin inflammations.
- FLORINEF TABLETS,** NND 1962. (Squibb) Fludrocortisone acetate Tab. 0.1 mg. Bot. 100s Tab. 1.0 mg Bot. 30s.
 Use: (0.1 to 0.3 mg. daily) Treatment of Addison's disease and adrenal hyperplasia
- FLORINEF NEMISUCCINATE.** (Squibb) Fludrocortisone hemisuccinate eq to 1 mg fludrocortisone acetate/ss. (0.1%) Ophthalmic solution. Bot 2.5 ss.
 See: Florinef-S, Ophth Sol.
 Use: Inflammation
- FLORINEF-S.** (Squibb) Lot.: Fludrocortisone acetate 0.05% & 0.1% combined with neomycin 2.5 mg., gramicidin 0.25 mg./cc. Squeeze bottle 15 cc. Oint. or Cream: Fludrocortisone acetate 0.1% combined with neomycin 2.5 mg., & gramicidin 0.25 mg./Gm. Tube 5 & 20 Gm.
 Ophth. Sol.: Fludrocortisone hemisuccinate 1.0 mg., neomycin 2.5 mg., gramicidin 0.25 mg./cc. Bot. dropper 2.5 cc.
 Ophth. Susp.—Fludrocortisone acetate 0.1%, neomycin 2.5 mg., gramicidin 0.25 mg./Gm. Bot. w/dropper 5 cc.
 Ophth. Oint.—Fludrocortisone acetate 0.1%, neomycin 2.5 mg., gramicidin 0.25 mg./Gm. Tube 3.6 Gm. w/ophth. tip.
 Use: Inhibits inflammatory reactions to bacterial, allergenic & chemical agents.
- FLOROPHYL,** NND. 1962. (Merk Sharp & Dohme) Diisopropyl fluorophosphate. Isofluorophosphate, U.S.P. (DFF) Ophthalmic use only, 5 cc. of an 0.1% sol. in peanut oil Oph. Oint. 0.025%, Tube 3.5 Gm.
 Use: Miotic, a potent antagonist of cholinesterase.
- FLOROTIC.** (Squibb) Fludrocortisone acetate 10 mg., neomycin base as sulfate 3.5 mg., polymyxin B sulf., 10,000 u., nystatin (Mycostatin) 100,000 u./cc. Plastic dropper 5 cc.
 Use: Inflammatory conditions of the ear.
- FLOURAN DROPS.** (Cambridge) Fluoride ion 0.1 mg./drop of liq. 1 oz. also vit. C 30 mg & D 400 U.S.P. u/drops. Bot. 1 oz. plastic droppers
 Use: Dental caries prevention
- FLIOBAN LOZENGES.** (Cambridge) Fluoride ion 1 mg./loz. vit C 30 mg & D 400 U.S.P. u/loz. Bot. 100s.
 Use: Dental caries prevention
- FLUAGEL.** (Bieeon) Alum-num hydroxide equiv to 4% aluminum oxide. Bot. 10 oz., Comp. Tab. aluminum hydroxide w/Mag. trisilicate. Bot. 100s, 500s, 1000s.
 Use: Orally; antacid.
- FLUCAPS.** (Walker) Pyriminamine moleale 20 mg., quinine sulf. 30 mg., aluminum hydroxide, APC (300, 120 & 30 mg)/Cap. Bot. 24s, 100s
 Use: Influenza symptoms
- ***FLUDROCORTISONE ACETATE,** NND. 1962. 9 alpha Fluorohydrocortisone.
 See: Alflorane Acetate, NNR (Merk Sharp & Dohme)
 Cortel F, NNR, Oint. (Upjohn)
 Florinef Acetate, NNR. (Squibb)
 W/Neomycin & gramicidin See: Florinef S Lot. & Oint. (Squibb).
 W/Neomycin base as sulfate, polymyxin B sulf., nystatin (Mycostatin) See: Florotic, plastic dropper. (Squibb)
 W/Nystatin, neomycin sulf., gramicidin. See: Myconef Ointment (Squibb)
 W/Spectrocin. See: Florinef-S, Preps. (Squibb)
FLUDROCORTISONE HEMISUCCINATE. See: Florinef hemisuccinate sal'n. (Squibb)

- *FLUFENAMIC ACID** (Parke Davis)N (a a a trifluoro m tolyl)anthranic acid
Use Anti-inflammatory agent
- FLUITRAN**, (Schering) Trichlormethiazide 2 mg or 4 mg /draque
Use Diuretic
- *FLUMETHASONE**, (Upjohn) 6a 9a difluoro 16a methylprednisolone
Use Corticoid
- *FLUMETHIAZIDE**, NND 1962 6 Trifluoro methyl 7 sulfamyl 1 2 4 benzothiazine 1 1 dioxide
See Ademel Tab (Squibb)
Routrox (Squibb)
Use Diuretic
- *FLUNISOLIDE ACETATE**. See Fluoxolone
- *FLUOCINOLONE ACETONIDE** 6a 9a difluoro 16a hydroxyprednisolone 16 17 acetate See Synalar (Synlex)
- FLUORAC**, (US Vitamin) Fluoride 0.5 mg Vit B 6 D2 mg, C 75 mg A 5000 u D 500u /chewable Tab Bot 100s
Use Fluoride and vitamin supplement
- FLUORESCIN** (Merck Sharp & Dahme) Resorcinolphthalein Pow, Bot 1/2 oz, 1 oz
Use Diagnostic aid
- FLUORESCIN ISOTHIOCYANATE**, (Borden Chem Co—Dajac Lab)
Use: Antigen antibody reaction Rabies diagnosis
- *FLUORESCIN SODIUM**, USF NNR 1950; Ophth Sol, USF (Merck Sharp & Dahme) Soluble fluorescein Bot 1/2 oz 1 oz 1/4 lb See Fluor I Strip (Ophthalmus Inc)
Use 2% sol in diagnosis of eye conditions
- FLUORESCIN SODIUM IV** See Fluorescite (Kirk)
- FLUORESCIN SODIUM 2% SOLUTION** (Alcon) Drop Trainer 15 cc Dropperettes 2%/1 cc (Iso-Sol); 0.5% Bot 5 cc
Use Diagnostic agent in ophthalmology
- FLUORESCIN SODIUM 2% SOLUTION**, (Crookes Barnes) Vials 15 cc and dropperettes
Use Diagnostic agent in ophthalmology
- FLUORESCITE**, (Kirk) Fluorescein as Sod salt 5% sol Amp (10 cc) Box 6s 25s, 100s
Use: Diagnostic aid
- FLUORIDE CALCIUM**. See: Ce De Flor, Tab (Hoack)
- FLUORIDE PREP**. See: Cadrosan, Tab (Merck Sharp & Dahme)
- FLUORIDE SODIUM**.
See: Fluorol, Sol (Novocol)
Fundo Vite (F), Drops (Hoyt)
Keridum, Tab (Odora)
- FLUORIDE THERAPY**.
See Colic gen, Cap (Boyle)
* Generic name recognized by N.F., U.S.P. or A.M.A.
- Ce Ce Fira Tab (Hoack)
Enzifur Laz (Ayerst)
Flouran (Cambridge)
Fluro Laz (Kirkman)
Flursol Sol (Novocol)
Fundo Vite (Hoyt)
Les Cav (Crookes Barnes)
Paly Vi Flor (Mead Johnson)
Tri Vi Flor (Mead Johnson)
- FLUOR I STRIP**, (Ophthalmus) Fluorescein sod 9.3 mg on paper
Use Ophthalmic diagnostic agent
- FLUOROCORTISONE** (Currently not available)
- FLUOROHYDROCORTISONE ACETATE**, 9a fluoro hydrocortisone See Fludrocortisone Acetate
- FLUOROMAR** (Ohio Chemical & Surgical Equipment Co) Fluoroxene
Use Inhalation anesthetic
- FLURO-METHANE SPRAY** (Gebauer) Di chlorodifluoromethane 15% trichlorofluoromethane 85%/Liq Bot Dispensed 4 oz
Use Counterirritant
- *FLUOROMETHOLONE**, NND 1962 6 methyl 9 fluoro prednisolone See Oxylone (Upjohn)
- 6a-FLUOROPREDNISOLONE** See Alphadrol Tabs (Upjohn)
- FLUOROURACIL** (Roche) Fluorouracil 50 mg/cc 10 cc amp 50 mg/cc/12s
Use Malignancies
- FLUOTHANE**, NND 1962 (Ayerst) Halothane Trifluoro Chlorobromoethane 2 bromo 2 chloro 1 1 1 trifluoroethane Bot 125 cc
Use General anesthetic agent
- FLUOXALONATE**, (Eli Lilly) See Fluoxalate 6a fluoro 16a hydroxyprednisolone 16 17 acetate 21 acetate Changed from Fluoxalone
- FLUOXALONE**, (Eli Lilly) 6a fluoro 16a hydroxyprednisolone 16 17 acetate 21 acetate Changed to Fluoxalate
- *FLUOXYMESTERONE**, NND 1962 9 alpha fluoro 11 beta hydroxy 17 alpha methyl testosterone
See: Halatestin, Tab (Upjohn)
Oro Testryl, Tab (Squibb)
Ultandren, Tab (Ciba)
- W/Ethinyl estradiol** See: Haladrin Tab (Upjohn)
- W/Ethinyl estradiol, Vitamins** See Hormozyme Tab (Upjohn)
- *FLUPEROLONE ACETATE** (Pfizer) 9a fluoro 21 methylprednisolone 9a fluoro 11 beta 21 trihydroxy 21 methylpregna 1 4 d ene 3 20 diene See: Methrol
Use: Corticosteroid

FLUPNENAZINE HYDROCHLORIDE, N N D. 1962. 1-(2-Hydroxyethyl)-4-[3-(2-trifluoromethyl-10-Phenothiazinyl-propyl)]-piperazine di HCl.

See: Permitt, Tab (White Lab.)

Prolixin, Tab & El. (Squibb)

Troncin, Tab. (Schering)

* **FLUPREDNISOLONE.** (Upjohn) a) 6 α fluoro-11 β ,17 α ,21-trihydroxypregna-1,4-diene 3, dione; b) 6 α fluoroprednisolone. See: Alphadrol (Upjohn)

FLURA DROPS. (Kirkman) Sodium fluoride 2.21 mg / 4 drops. (4 drops q.s. one quart provides 1 part per million of fluoride ion.) Bot. 2 oz.

Use: Prevention of dental caries.

FLURA-LOZ. (Kirkman) Sodium fluoride 2.21 mg / lozenge. Bot. 100s

Use: Dental caries

FLURA-LOZ PLUS. (Kirkman) Sodium fluoride 2.21 mg, Vit. C 30 mg, D 400 u / Lozenge Bot. 100s

Use: Dental caries.

* **FLURANDRENOLONE.** 6-Fluoro 16-hydroxy-hydrocortisone 16,17-acetonide. See: Cortran (Lilly)

FLURA-PREN. (Kirkman) Vit A 4000 u, D 400 u, B-1 3 mg, B 2 2 mg, B 6 10 mg, niacinamide 10 mg, Vit. C 100 mg, K 0.5 mg, calcium carbonate 625 mg, ferrous fumarate 125 mg, Vit. B-12 w/intrinsic factor $\frac{1}{2}$ oral u., sodium fluoride 2.21 mg / Tab Bot. 100s.

Use: Nutritional supplement

* **FLUROTHYL.** (Ohio Chem. and Surg Eq) bis (2,2,2-trifluoroethyl) ether. See: Indoklon (Ohio Chem. and Surg Eq)

* **FLUROXENE.** 1,1,1-trifluoroethyl vinyl ether. See: Fluoromor (Ohio Chemical & Surgical Equipment Co)

FLURSOL. A.D.R. (Novocot) Sod. fluoride 2% sol. Bot. 2 oz.

Use: Topically; dental caries.

FLUTRA. (Schering) Trichlormethiazide 2 mg or 4 mg / Tab.

Use: Diuretic.

FOILLE. (Carbisulphail) Benzyl alcohol, benzoicaine, oxyquinoline base, sulfur, in a bland vegetable oil base. liq Bot. 1 $\frac{1}{2}$ oz, 4 oz, 1 pt., 1 gal. Oint. Tube 1 $\frac{1}{4}$ oz. Jar 1 lb, 6 lb Spray liq Con 3 & 10 oz.

Use: Topically, burn therapy, antiseptic and protective.

FOLABIN. (Paul Money) Folic acid 5 mg, Vit. B-12 25 mcg, liver 5 mcg, of Vit. B-12 / 10 cc. Vial. 10 cc.

Use: Hematinic.

FOLACIN. N N D. 1962. See: Folic Acid, USP

* Generic name recognized by N.F., U.S.P. or A.M.A.

FOLACINE. See: Folic acid

FOLATE, SODIUM, N N D. 1962. See: Folvite, Sod (Lederle)

FOLBESTYN. (Lederle) Thiamine 10 mg, niacinamide 50 mg, riboflavin 5 mg, ascorbic acid 175 mg, folic acid 5 mg, Vit. B-12 5 mcg / Tab Bot. 50s, 500s. Thiamine HCl 10 mg, riboflavin 10 mg, niacinamide 75 mg, Sod. pantothenate 10 mg., pyridoxine HCl 5 mg., Vit. B-12 15 mcg, folic acid 3 mg, ascorbic acid 300 mg./2 cc. vial. Parenteral 1-dose vial with diluent 1 & 25 viol pkg

Use: Supplement to diets

FOLBEXIL. (Chicago Pharmacal) Vit. B-12 5 mcg, folic acid 1 mg., glutamic acid HCl 30 mg / Tab. Bot. 30s, 100s, 500s, 1000s.

Use: (1 t.i.d.) Hematopoeitic.

FOLIXEL. (Elder) Folic acid 5 mg, Vit. B 12 50 mcg, liver inj. q.s./cc. Vial 10 cc.

Use: Hematinic.

FOLI-B-12. (New Mexico Pharm) Cyanocobalamin 10 mcg, folic acid 10 mg, B 12 crystalline 100 mcg / cc Vial 10 cc

FOLS-BRUMIN. (Philadelphia Ampoule Labs) Liver inj. refined 5 mcg, B-12 30 mcg, folic acid 5 mg./cc Vial 10 cc.

Use: Vitamin B complex therapy.

FOLS-BRUMIN FORTIFIED. (Key Corp) Liv inj. refined 10 mcg, B-12 60 mcg, folic acid 10 mg./cc. Vial 10 cc

Use: Vitamin B complex therapy.

* **FOLIC ACID, USP, N N D.** 1962, A.D.R. Cop., U.S.P.; Inj., U.S.P.; Tab USP Pterylglutamic acid Vitamin Bc. See: Folvite, (Lederle), N N R. 1956.

Use: (5 to 10 mg. daily) Vitamin therapy W/Cyanocobalamin, methamphetamine HCl, niacinamide. See: Rutalox (Central)

W/Estroate, cholinn bitartrate & Vit. B-12 See Menotrape, Tab. (Lakeside)

W/Iron. See: Hemaboloids w/Folic Acid, Tab. (Arlington).

W/Liver Conc. & Disulfo. See: Ifosulfas, Tab. (Upjohn).

W/Vit. B 12.

See: B Folvite, Cap. (Abbott)

Dodecfol, Tab. (U S Vitamin Corp)

Duo-Tal, Vial (Flint, Eaton).

Rubrofolin, Cap., Vial (Squibb)

W/Vit. B 12. See: Normocytin, Tab. (Lederle)

W/Vitamin multiple. See: Vitamin preparations

FOLIC ACID ANTAGONISTS.

See: Aminopterin (Lederle)

Methotrexate (Lederle)

Terapterin Sod. (Lederle)

- FOLIC ACID SALTS** See: Folvite 30d, Vial (Lederle)
- FOLICUPRIN** (Kremers Urban) Folic acid 2 mg, ferrous gluconate 260 mg, copper gluconate 15 mg, ascorbic acid 50 mg / Tab Bot 100s, 1000s
Use: Macrocytic anemias
- FOLIDOZIN** (Meyer) Folic acid 5 mg, Vit B 12 40 mcg, benzyl alcohol 2%, phenol 0.5%, liver inj qs (10 mcg Vit B 12 activity before mixing/cc) Vial 10 cc
Use: Anemias
- FOLIDOZIN R.** (Meyer) Choline 100 mg, folic acid 5 mg, Vit B 12 40 mcg, benzyl alcohol 2%, liver injection 10 mcg qs 1 cc / 1 cc Vial 10 cc
Use: Supportive therapy in the treatment of anemias
- FOLINIC ACID-SYNTHETIC FACTOR.** See: Leucovorin Amp (Lederle)
- FOLIVER** (National Drug) Folic acid 5 mg, w/liver 20 mg/cc Vial 10 cc
Use: Hematinic
- FOLIVER.** (Kirk) Vit B 12 30 mcg, folic acid 5 mg, liver inj 5 mcg/cc Vial 10 cc
Use: Hematinic
- FOLLESTROL** (Blue Line) Aq susp, Vial (estrone 20,000 IU/cc & 50,000 IU/cc) 10 cc
Use: Estrogen therapy
- FOLLICORMON** See: Estradiol Benzoate (Various Mfr)
- FOLLICULAR HORMONE HYDRATE** See: Estril, Cap (Abbott)
- FOLLICULAR HORMONES.** See: Estrone (Various Mfr)
- FOLLICULIN** See: Estrone (Various Mfr)
- FOLLUTEIN, NND 1960** (Squibb) Chorionic gonadotropin, NNR 1954 Vial 1000 IU, 5000 IU, 10,000 IU, w/amp of diluent 10 cc
Use: Cryptorchidism
- FOLVITE, NND 1962** (Lederle) Folic acid Elixir (5 mg / 5 cc) Bot 4 fl oz
Sol (15 mg / 5 cc) 10 cc Vial
Tab: (5 mg / Tab) Bot 100s, 1000s; (20 mg / Tab) Bot 25s
Use: Megaloblastic anemias
- FOLVITE SODIUM** (Lederle) Sodium Folate Vial (15 mg / 5 cc) 10 cc
- FOLYRON.** (Lederle) Folic acid 17 mg, ferrous sulfate exsic 3 gr / Cap Elixir (folic acid 0.89 mg, ferrous gluconate 28 mg / 5 cc) Bot 16 fl oz, 1 gal Tab (folic acid 17 mg, ferrous sulfate exsic 3 gr / Tab) Bot 100s, 500s, 1000s
Ule (f 2 t i d w/meals) Dietary supplement
- FOLYRON-G** (Lederle) Ferrous gluconate
* Generic name recognized by NF USP or AMA
- 494 mg, folic acid 1.7 mg Vit C 50 mg / Cap Bot 100s, 500s, 1000s
Use: (1 2 i d) Iron deficient anemias
- FONAMIDE-3** (Morton & Converse) Sulfamethazine sulfadiazine & sulfamerazine of each 0.167 Gm / dr Bot 4 oz, 1 pt
Use: Triple sulfonamide therapy
- FONATOL.** See: Diethylstilbestrol (Various Mfr)
- FONTARSOL.** See: Dichlorophenarsine Hydrochloride (Maphorsin—Parke Davis)
- FORCOLO** (Iemmon) Codeine phosphate 60 mg, chlorpheniramine maleate 12 mg, ephedrine sulfate 60 mg sodium citrate 1 Gm menthol 3 mg / 30 cc 3 fl oz
Use: Expectorant
- FOR-OYNE** (First Texas) Aspirin 5 gr, colchicine salicylate 1/700 gr, acetophenetidin 1 gr caffeine 1/2 gr / Cap or Tab Bot 100s, 500s, 1000s Also available w/codeine phosphate 1/2 gr / Tab
Use: (1 2 q 3 h) Analgesic & antipyretic & anodyne
- FORHISTAL.** (Ciba) Dimethylpyridene maleate di 2 { 1 [2 (2 dimethylaminoethyl) 3 indanyl] ethyl } pyridine maleate
Kontabs—2.5 mg (orange) Bot 100s
Tab—1 mg (pale orange scored) Bot 100s
Syrup (pink)—1 mg Forhistal maleate per 5 ml tsp Bot 4 fl oz
Pediatric Drops (pink)—0.5 mg Forhistal maleate per 0.6 ml Bot of 1 fl oz w/droppers 0.3 or 0.6 ml
Use: Allergy and pruritus
- FORMALDEHYDE PASTILS.** Paraformaldehyde Tab (98%) Bot 100s
Use: Vaporize by heating for sterilization
- * FORMALDENYDE SOLUTION (37%).** USP, NNB 1949 ADS
Use: For poison try, fungus infections of the skin, hyperhidrosis & as an astringent w/Tannin See Yonnamaf (Myer Drug)
- FORMALIN, NNB 1949, ADS** See: Formaldehyde Solution (Various Mfr)
- FORMASAL** (First Texas) Aspirin 7 gr, colchicine salicylate 1/500 gr / Cap or Tab Bot 100s, 500s, 1000s
Use: Analgesic & antipyretic
- FORMASAL & CAMPHOR COMPOUND** (First Texas) Aspirin 2 1/2 gr, acetophenetidin 2 1/2 gr, Gelsemium 1/5 gr, colchicine salicylate & camphor manabromide / Cap Bot 100s, 500s, 1000s
W/Antihistamine Same as plain but pyronis amine maleate 25 mg / Cap added Bot 100s, 500s, 1000s
W/Codeine phosphate: 1/4 gr Cap, Bot 100s, 500s; 1/2 gr Cap Bot 100s, 500s
Use: Orally; analgesic & antipyretic

FORMATONE. (Ives-Cameron) Vit. B-1 12.5 mg., B-12 100 mcg., in a wine base containing sorbitol/5 cc. Bot. 16 fl. oz.

Use: Stimulate appetite.

FORMATRIX. (Ayerst) Conjugated estrogens equine (Premarin) 1.25 mg., methyltestosterone 10 mg., ascorbic acid 400 mg./Tab. Bot. 60s, 500s

Use: Treatment of osteoporosis.

FORMIC ACID W/Silicic acid. See: Ray-Formosil, Amp. (Raymer)

W/Silicic acid, cobra venom. See: Nyloxin, Amp. (Hynson, Wescott & Dunning)

FORMIN. N.N.R. 1950. (Merck Sharp & Dohme) (Methenamine) (Hexamethylenetetramine) Pow., Bot. 1/2 lb., Drum 5 lb.

Use: Orally; urinary antiseptic

FORMITOL. (Patch) Formaldehyde, citric acid, tannic acid, hyoscyamus, ammonium chloride, guaiac, benzoic acid, menthol, methyl salicylate & eucalyptus oil. Flg. 20s, Bot. 50s, 1000s.

Use: Throat irritations.

FORMOHYDRION. (Upjohn) Methenamine 5 gr., Sod. biphosphate 20 gr./tsp. Bot. 4 oz., 8 oz. Tob. (5 gr. & 7 1/2 gr) Bot. 100s, 1000s.

Use: (1-2 t.i.d.) Urinary antiseptic

FORMO-QUINOCAINE. (Fornsworth Lab) Formic acid .05%, dibucaine HCl 0.5%, quinine HBr. Vials 100 cc.

Use: See sting therapy for arthritic.

FORMULASE. (Kremers Urban) Proteolytic enzyme 5 mg., amylolytic enzyme 10 mg., lipolytic enzyme 20 mg./Tab. Box 60s.

Use: Digestive disorders.

FORNOTROPERIA. (Fornsworth Lab) Chorionic gonadotropin with Vit B-12. Vials (2500, 5000 and 10,000 IU 1 cc) 10 cc

Use: Gonadotropin therapy.

FORSTEO. (Brown) Ethinyl estradiol 0.02 mg., methyl testosterone 4 mg., thyroid ext. 10 mg., Vit. C 300 mg., D 1000 u., A 3000 u., E 2 iu, calcium lactate 250 mg., dicalcium phosphate 125 mg./Tab. Bot. 100s

Use: Osteoporotic

FORTE-LAC-B. (Consolidated Midland) Vit D 500 IU, C 100 mg., calcium 250 mg., iron 50 mg./Tab. Bot. 100s, 250s.

Use: Dietary supplement during pregnancy and lactation.

FORTE-LAC-C. (Consolidated Midland) Folic acid .3 mg., B-12 2 mcg., calcium pantothenate 5 mg., pyridoxine HCl 3 mg., niacinamide 15 mg., B-2 3 mg., B-1 2 mg., D 500 IU, A 6000 IU, C 100 mg., Ca 250 mg., Fe 50 mg./Tab. Bot. 100s, 250s.

Use: Dietary supplement during pregnancy and lactation.

FORTESPAN. (Smith, Kline & French) B-1 6 mg., B-2 6 mg., B-6 6 mg., B-12 6 mcg., nicotinamide 60 mg., A 15,000 U.S.F. units, D 1000 U.S.F. units, C 150 mg./spanule Bot. 50s, 100s, 1000s.

Use: Vitamin supplementation in sustained release form

FORTHANE, INHALER, NND 1962. (Lilly) 2 Amino 4 methylhexone in individual containers.

Use: Allergic rhinitis & head colds.

FORTICILLIN. (Premo) Penicillin G procaine 300M u., penicillin G. Sod. 100M u./cc. Vial 1 cc., 5 cc., 10 cc.

Use: Fortified penicillin therapy.

FORTICON. (Consolidated Midland Corp) Vit. A 5000 u., D 500 u., C 37.5 mg., B-1 1.5 mg., B-2 1.5 mg., B-6 0.1 mg., B-12 1 mcg., E 0.17 IU, niacinamide 5 mg., calcium pantothenate 1 mg., dicalcium phosphate 50 mg., iron 7.5 mg., cobalt 0.1 mg., copper 0.25 mg., iodine 0.08 mg., manganese 0.81 mg., zinc 0.23 mg., molybdenum 0.4 mg., choline bitartrate 50 mg., inositol 25 mg., methionine 50 mg./Vial sphere. Bot. 30s, 100s, 250s

Use: Multiple vitamin deficiencies

FORTICON LACTATE. (Consol. Midland) Vit. A 5000 u., D 200 u., B-1 2.5 mg., B-2 2.5 mg., B-6 1.0 mg., B-12 1 mcg., Cal. pantothenate 1 mg., niacinamide 5 mg., Vit. C 50 mg., E 0.17 u., choline bitartrate 50 mg., inositol 25 mg., methionine 50 mg., cal lactate 250 mg., w/iron, cobalt, copper, iodine, Mang, molybdenum, zinc/sphere Bot. 30s, 100s, 250s, 1000s.

Use: Treatment of multiple vitamin deficiencies.

FORTIFIED. (Dumas-Wilson) Primary yeast 7 1/2 gr., B-1 1.8 mg., B-2 0.7 mg., niacin 3.5 mg./Tab. Bot. 100s

Use: Dietary supplement

FORTIFIED YEAST EXTRACT. (Kinney) Thiamine HCl 2 mg., riboflavin 3 mg., niacinamide 20 mg., cal. pantothenate 1 mg., pyridoxine HCl 0.5 mg./5 cc. Bot. 4 oz., 1 pt

FORTI-PLEX. (Plessner) Thiamine HCl 5 mg., riboflavin 5 mg., pyridoxine HCl 1 mg., Cal. pantothenate 1 mg., niacinamide 20 mg., yeast & liver conc. base/Tab. Bot. 100s, 500s, 1000s.

Use: Dietary supplement.

FORTIZYME. (Breon) Alpha amylase 10 mg./Tab. Bot. 4s

Use: Inflammation.

FORTY PLEX. (Philadelphia Ampoule Labs) Vit. B-12 30 mcg., B-1 20 mg., B-2 3 mg., B-6 1 mg., niacinamide 100 mg., cal. pantothenate 3 mg., choline chloride 20 mg.

- inositol 10 mg, liver in] crude 2 mcg/cc
Vial 25 cc
Use: Geriatric therapy
- FOSFREE** (Mission Pharmacol Co) Calcium lactate 1000 mg, calcium gluconate 1000 mg calcium carbonate 1200 mg, calcium 703 mg Vit 0 2 600 u, A acetate 6000 u, E (as mixed Tocopherols) 50 IU, C 150 mg K 10 mg B 6 10 mg B 1 80 mg, Vit B 2 B mg B 3 40 mg d calcium pantothenate 20 mg ferrous gluconate 320 mg, iron 37 mg B 12 (crystalline an sesin) 10 mcgm copper (as sulfate) 0.6 mg, manganese (as citrate soluble) 10 mg zinc (as oxide) 0.32 mg, potassium (as chloride) 20 mg magnesium (as carbonate) 10 mg
Use: Prevention and relief of leg cramps; vitamin and mineral deficiencies
- FOSTEX CAKE and CREAM** (Westwood Pharmacol) Soapless cleanser and wetting agents w/hexachlorophene 1%, sulfur 2% salicylic acid 2% Pkg 1 cake Cream Jars 4.5 oz
Use: Treatment of fungus infections
- FOSTRIL** (Westwood) Polyoxyethylene lauryl ether 6% hexachlorophene 1%, sulfur 2% 1oz Tubes 1 1/2 oz
Use: Acne
- FOSTRIL HC** (Westwood) Polyoxyethylene lauryl ether 6%, hydrocarizone 1/4%, hexachlorophene 1%, sulfur 2% Tubes 25 Gm
Use: Acne
- FOURALCO** (Jenkins Labs) Salicylic acid 1/2 gr, sod bicarbonate 3 gr, mag salicylate 2 gr, strontium salicylate 2 gr, aceto phenol 1/4 gr w/methyl salicylate, pancreatin diatose/Tab Bot 100s
Uses (2 q 3 h) Treatment of rheumatic conditions
- FOWLER'S SOLUTION, ADB 1955** See: Potassium Arsenite Solution (Various Mfr)
- FOXALIN** (Grant) Digifoxin USP 01 mg w/Sod carboxymethylcellulose/Tab Bot 100s, 1000s
Use: Orally digitalis glycoside for rapid digitalization
- FOXGLOVE.** See Digitalis USP
- FOXIXILLIN PWD FOR SOLN** (Fox) Buffered penicillin G 250 000 u in cherry flavored vehicle/5 cc Bot 2 oz
Use: Penicillin sensitive organisms
- FRAGIKAP-I.** (Ferndale) Vit C 200 mg hesperidin methyl chalcane 50 mg hesperidin complex 150 mg, menadione [Vit K] 1.34 mg/Tab Bot 100s
Use: Capillary fragility
- FRANOL** (Winthrop Labs) Tab containing bethedon 1/2 gr, luminal 1/2 gr, & theophylline 2 gr Tab, Bot 100s
* Generic name recognized by NF, USP or AMA
- Use: (1) d) Bronchial asthma & hay fever
- FREI ANTIGEN.** See: Lygromum, Vial (Squibb)
- FRENOLICIN.** (Ederle) Antibiotic isolated from *Streptomyces fradiae*
Use: Inhibits fungi and gram positive bacteria
- FRENOSIL.** (Wallau) Mammary gland 1.5 Gm, corpus luteum 0.08 Gm/cc Bot 8 cc 1.6 cc Tabs Bot 30s 60s
Use: Hormone therapy
- FRENUQUEL, NND 1961** (Merrell) Alpha (4 Piperidyl) benzhydrol HCl Azacyclonol Tab 20 mg or 100 mg Bot 100s Amp (5 mg/cc) 20 cc
Use: Treatment of acute schizophrenia
- FRIEDLANDER VACCINE** (Sherman) Pneumococci 100 million streptococci 100 million, *N catarrhale* 200 million *Staph aureus* and *aureus* 400 million & friedlander 300 million/cc Vial 12.5 & 20 cc
Use: Respiratory vaccine for the treatment of upper respiratory infections
- * **FRUCTOSE, NF Inj NF NND 1962**
See: FreeLabs (Pfanzstiel)
Levugen, Sol (Mead Johnson)
- FRUCTOSE, Sol** (Abbott) Levulose 10% Sol Bot 1000 cc
Use: Parenteral nutrition
- FRUTABS** (Pfanzstiel) Fructose tablets
Use: Quick energy source for athlete workers, etc
- F-3-1** (Myers Carter) Anterior pit gonadotropin hormone 125 RU/5 cc Vial 10 cc
- FUADIN, NND 1959** (Winthrop Labs) Sod antimony III bis catechol 2,4 disulphonate in 6.3% aq sol (Stibophen) Amp 3.5 cc & 5 cc, Box 10s
Use: Considered to be a specific agent in bilharziasis (schistosomiasis) Also used for granuloma inguinale
- * **FUCHSIN, BASIC, NF** Basic Fuchsin is a mixture of rosaniline and pararosaniline HCl Basic Magenta
Use: Ingredient in Carbo Fuchsin Sol NF
- FUGILLIN** (Fumagillin) (Upjohn)
- FULVICIN, NND 1962** (Schering) *Griofulven 125 mg 250 mg 500 mg/Tab Bot 60s, 250s
Use: Oral, fungal infections of the skin, hair, and nails
- * **FUMAGILLIN, NND 1962** Fumagillin is a crystalline antibiotic produced during the growth of a strain of *Aspergillus fumigatus*
See: Fumadif (Abbott)
Fugilin (Upjohn)
- FUMASORB** (Marion) Ferrus fumarate 200 mg, sorbitol 200 mg/Tab Bot 100s 1000s
Use: Anemias
- FUMEX** (CMC) Silver acetate 6 mg, coen

zyme 0.025 mg, NH_4Cl 25 mg /Loz. Bot. 30s.

Use: To break the smoking habit.

EUMIDIL, N.N.D. 1962. (Abbott) Tob. (10 mg) Bot. 30s.

Use: Intestinal amebiasis.

FUMIRON, N.N.D. 1962. (Standex) Ferrous fumarate 324 mg /Tabs. Bot. 100s.

Use: Iron deficiency.

FUNDA-VITE. (Hoyt) Vit. D 400 u, C 30 mg /0.6 ml. Bot. 30 ml. Pediatric Drops 50 ml bot w/calibrated dropper lozits, 100s.

Use: Vitamin supplement.

FUNDA-VITE (E). (Hoyt) Vit. D 400 u, C 30 mg, fluoride (as sod. fluoride) 0.5 mg / 0.6 ml. Bot. w/dropper 30 ml.

Use: Vitamin supplement and safeguard against future dental caries.

FUNDA-VITE (E) /A. (Hoyt) Vit. A 3000 u, D 400 u, C 60 mg, fluoride 0.5 mg /0.6 cc. Bot. w/calibrated drop 50 cc.

Use: Dietary supplement.

EUNGACETIN, N.N.D. 1962. (G. F. Harvey) Tricelatin (glyceryl trilcelate) 25% in a water miscible base Tube 1 oz., Jar 1 lb liq. (30%) Bot. 1 oz. Paw. (33 1/2%) Shaker Pkg. 1 1/2 oz.

Use: Topically, fungus infections.

EUNGOICIDES.

See: Asteral (Roche)

Basic Fuchsin, N.F.

Benzolic Acid (Various Mfr.)

Caprylate, Ca, Cu, Na, Zn (Various Mfr.)

Copyric Acid (Various Mfr.)

Coparofinate (Ira-Par) (Medical Chem.)

Cresalin. (Merck Sharp & Dohme)

Dolyde (Hynson, Wescott & Dunning)

Dermycin. (Chol-Yon)

Dibromsalicylaldehyde (Hynson, Wescott & Dunning)

Didroxane (Sarnoy)

Furaspor Cream (Eaton).

Leningallal (Bilhber-Knoll).

Metacresylacetate. (Merck Sharp & Dohme)

Nitrofurfuryl Methyl Ether (Eaton).

p Nitrophenol. Sodium iodate. (Chol-Yon)

Phenylmercuric Cpd. (Various Mfr.)

o Phenylphenol (Smith-Dorsey).

Propionate. Sodium (Various Mfr.)

Propionic Acid (Various Mfr.)

Salicylic Acid (Various Mfr.)

Salicylanilide (Various Mfr.)

Shibamidine isethionate (Merrell)

Thymol (Various Mfr.)

Triacetylpyrogallal (Bilhber-Knoll).

Undecylenate, Cu, K, Zn (Various Mfr.)

Undecylenic Acid (Various Mfr.)

* **EUNGIMYCIN**. (Warner-Lambert) Antibiotic

* Generic name recognized by N.F., U.S.P. or A.M.A.

atic of the heptane type elaborated by *Streptomyces coelicolor* var. *aminophilus*. Use: Antifungal antibiotic.

EUNGIZONE, N.N.D. 1962. (Squibb) Amphotericin B, derived from a previously undescribed species of *Streptomyces*, 50 mg of amphotericin B actively combined with approximately 41 mg of sod. desoxycholate with sod. phosphate as a buffer /Vial Lyophilized 20 cc.

Use: I.V. Mycotic infections.

FUNGIZONE LOTION. (Squibb) Amphotericin B 3%, Ithioerosol, lithium dioxide/cc, Plastic Sq. Bot. 30 cc.

Use: Monilial infections.

FURACIN, N.N.D. 1962. (Eaton) Nitrofurazone, N.F., 5-nitro-2-furafdehyde semicarbazone. Tube 1 oz.

Use: Topical, antibacterial agent.

W/Phenylephrine HCl. See: Furacin Nasal Sol. (Eaton)

W/Diethylstilbestrol. See: Furesinol, Supp (Eaton)

FURACIN CREAM, (Eaton) Furacin D.2%, a water miscible, self-emulsifying cream. Tube 3 oz. w/(vaginal) applicator.

Use: Bacterial vaginitis where cream is desired; topical antibacterial agent.

"**FURACIN EAR SOL., IMPROVED**," See: Furacin Otc. (Eaton)

FURACIN-HC CREAM, (Eaton) Nitrofurazone 0.2%, hydrocortisone acetate in a vanishing cream base 1.0% Tubes 5 Gm., 20 Gm. Use: Inflammation.

FURACIN-HC URETHRAL SUPPOSITORIES, (Eaton) Nitrofurazone 0.2%, hydrocortisone acetate 1%, diprostan HCl 2% in water-dispersible bases of glyceryl monolaurate and sorbitol monostearate/1.3 Gm. Box 12s.

Use: Antibacterial, anti-inflammatory, anesthetic.

FURACIN INSERTS. (Eaton) Furacin 0.2%, diprostan HCl 2%. Inserts. Box 12s.

Use: Bacterial urethritis, vulvovaginitis, rectal lesions, etc.

FURACIN NASAL W/PHENYLEPHRINE, (Eaton) Furacin 0.02%, phenylephrine HCl 0.25% in an aqueous sol Plastic Atomizer 15 cc.

FURACIN OPHTHALMIC LIQUID. (Eaton) Furacin 0.02% dissolved in isotonic aq. sol. Dropper, Bot. 1/2 fl. oz.

Use: External ophthalmic bacterial infections.

FURACIN OPHTHALMIC OINTMENT, N.N.D. 1954. (Eaton) Furacin 1% in a petrolatum type base. Tube 1/2 oz.

Use External ophthalmic bacterial infections

FURACIN OTIC. (Eaton) Furacin Ear Sol, Improved Nitrofurazone 0.2%, nitfurazime 0.375% dipiperdon HCl 2% in water soluble non drying, hygroscopic polyethylene glycol/Bot w/dropper 15 cc

Use Otitis externa, otomycosis, otitis media when eardrum is perforated

FURACIN SENSI-DISCS. (Baltimore Biol Lab) Concentration 100 mcg

FURACIN SOLUBLE DRESSING, N F (Eaton) 0.2% dissolved in a water soluble non drying ointmentlike base of polyethylene glycols Jar 5 oz, 1 lb, 5 lb Tube 1 oz, 2 oz

Use Topical application in surface infections

FURACIN SOLUBLE POWDER. (Eaton) 0.2% dissolved in Carbowax to form a cream colored water soluble pow Vial 1/2 oz

Use: When antibacterial powder is desired, as for wounds, external otitis & vaginitis

FURACIN SOLUTION, N N R (Eaton) 0.2% dissolved in water miscible liq of polyethylene glycols, wetting agent & water Bot 2 fl oz, 1 pt

Use: For use when a liquid vehicle is more convenient than Furacin Solution Dressing

FURACIN TOPICAL CREAM (Eaton) Nitrofurazone 0.2% in water miscible base of glycerin cetyl alcohol liquid petrolatum polyethylene glycol ether complex of higher fatty alcohols methylparaben propylparaben water/Tube 1 oz

Use Dermatologic antibacterial, topical

FURACIN VAGINAL SUPPOSITORIES (Eaton) 0.3% dissolved in water miscible base Box 12s

Use: Bacterial cervicitis & vaginitis

FURADANTIN, N N D 1962 (Eaton) N (5 Nitro 2 furfurylidene) 1-aminohydantoin (Nifurazone) Tab [50 mg] Bot 25s, 100s 500s (100 mg) Bot 100s 500s Hospital use 1000s

Use (1 2) Urinary infections

FURADANTIN ORAL SUSPENSION (Eaton) Nifurazone Bot [5 mg./cc] 2 oz

Use: Urinary antibacterial agent

FURADANTIN SENSI-DISCS (Baltimore Biol Lab) 100 mcg ea

FURADANTIN SODIUM. (Eaton) 1 (5 nitro furfuryl deneamino) hydantoin sodium 180 mg/Vials 20 cc

Use Gen urinary infection

FURADROXYL (Eaton) See Nifuroxime

***FURALAZINE HYDROCHLORIDE** (Nederle)

*Generic name recognized by N.F., U.S.P. or A.M.A.

3 amino 6 [2 (5 nitro 2 furyl)vinyl] aziridine hydrochloride

Use Anti microbial compound

***FURALTADONE, N N D 1962** Withdraw Nov 1961 5 morpholinomethyl 3-(5 nitro-furfurylideneamino) 2 oxazolidinone Sec Altafur, Tab (Eaton)

FURAMAZONE (Eaton) An ant bacterial nitrofurazone 5 nitro 2 furaldehyde semioxazone Nifuraldezane

Use Veterinary

FURAMAZONE SENSI-DISCS (Baltimore Biol Lab) 100 mcg each

FURASOL CREAM (Eaton) 1% Nitrofurfuryl methyl ether in a white water washable odorless nonstaining vanishing cream Jar 1 oz

Use Fungicide sporicide and bactericide

FURATADONE, (Eaton) See *Furmethanol

***FURAZOLIDONE, N F N N D 1962** N-(5-nitro 2 furfurylidene) 3 amino 2-oxazolone See Furaxone Tab (Eaton)

Tricofuron Vaginal (Eaton)

W/Nifuroxime See Tricofuron (Eaton)

FURESTROL. (Eaton) Furacin 0.2%, diethylstilbestrol 0.1 mg & dipiperdon HCl 2% in water miscible base/Urethrol Supp Bas 12s

Use Senile urethritis

FURETHIDINE 1 (2 tetrahydrofurfuryloxyethyl) 4 phenylpiperidine 4 carboxylic acid ethyl ester

FURFURYLUREA (National Drug) 1 (p tolyl sulfonyl) 3 furfurylurea

Use Hypoglycemic

***FURMETHONOL** (Eaton) 5 (4 morpholinyl methyl) 3 (5 nitro 2 furfurylideneamino) 2 oxazolidone See Furaladone

***FURMETNOXADONE** (Eaton) 5 methyl 3-(5 nitro 2 furfurylideneamino) 2 oxazolidone

FURON (Metro Med) Ferrus fumarate 2 mg, ascorbic acid 3 mg/Drop Pediatric suspension

Use Iron deficiency anemia

FUROXONE, N N D 1962 N N R (Eaton) Furazolidone, N 5 nitro 2 furfurylidene 3 amino 2 oxazolidone Tab (100 mg Bot 20s 100s Liq (50 mg/15 cc W/kaolⁿ & pectin) Bot 240 cc See Tricofuron T.M. Vaginal (Eaton Lab)

Use Antimicrobial agent

FUROXONE SENSI-DISCS (Baltimore Biol Lab) 100 mcg each

FUSINOL (Kendall) Boric acid phenol resorcin fuchsin, acetone, isopropyl alcohol & propylene glycol Bot 1 oz, 4 oz

Use Topically; fungus of the skin

FUZENE (Priman Moore) A polyethylene

- glycol diester oint. base of liq & white petrolatum, glycowax 3932, distearate & dilaurate at polyethylene glycol 400.
- VITAGEN**, (Endocrine) Unsaturated free fatty acids of linseed oil, linoleic and linolenic acids 168 mg./Cap. Bot. 100s
Use: (2 t i d) Various skin conditions; allergies, dermatitis, etc.
- G-11**, (Sindar Corp) (Hexachlorophene) Pow. form for mfg. See: Hexachlorophene, U.S.P.
- G-4**, See: Dichlorophene.
- GADOMENT OINTMENT**, (Smith, Miller & Patch) Cod liver oil 70%, phenol D.375% W/benzoin ext., & zinc oxide. Tube 1½ oz., 5 oz. Jar 1 lb.
Use: Topically, dermatitis of the skin.
- GALACTOSE INJECTION**, (Horton & Converse) Galactose 25% & 50% in sterile distilled water. Vial 50 cc.
Use: Diagnostic agent
- GALATEST**, (Denver Chemical) Vial 3½ dr.
Use: Diagnostic aid for detection of sugar in urine.
- GAL-BEN-ZOL, OINT**, (Ulmer) Gallic acid, benzocaine, zinc oxide, phenol Tube 1 oz
Use: Topically, antiseptic and protective.
- GALEN-B**, (White) Thiamine 4.5 mg., riboflavin 4.5 mg., niacin & niacinamide 60 mg., pyridoxine 4.5 mg., pantothenic acid 12 mg., inositol 225 mg., choline 300 mg., biotin 0.03 mg./30 cc. Bot. 8 oz., 1 pt., 1 gal.
Use: Vitamin B deficiencies.
- GALEN FORTIFIED**, (White) Thiamine 10 mg., riboflavin 5 mg., niacin & niacinamide 60 mg., pantothenic acid 12 mg., pyridoxine HCl 4.5 mg., inositol 112 mg., choline 150 mg., biotin 0.015 mg., w/iron, Mang. & rice bran/fl. oz. Bot. 8 fl. oz., 1 pt., 1 gal.
Use: Vitamin therapy.
- * **GALLAMINE TRIIODIDE**, N.N.D. 1962. Tri-diethylaminoethoxybenzene triethyliodide. See: Flaxedil, Vial (Lederle)
- GALLIADINE**, (Astor) Methamphetamine HCl 5 mg., caffeine 30 mg./Tab. Bot. 100s, 1000s.
Use: (1 t.i.d., m p) Obesity & depressed states.
- GALLOCHROME**. See: Merbramin (Various Mfr).
- GALLOGEN**, (Massengill) Diethanol-amine salt of mono-d-camphoric acid ester p-tolylmethyl carbinol 75 mg./Tab. 100s, 1000s. Also available as Compound Tabs. Bot. 100s.
Use: (1 t.i.d. for 3-4 wks) Increase bile flow.
- GALLOTANNIC ACID**. See: Tannic Acid. Preps.

* Generic name recognized by N.F., U.S.P. or A.M.A.

- * **GAMMA BENZENE HEXACHLORIDE**, U.S.P., N.N.D. 1962. See: Benzene Hexachloride, U.S.P.
- GAMMACORTEN**, N.N.D. 1962. (Ciba) Dexamethasone D.75 mg./Tab. Bot. 50s, 250s
Use: Corticosteroid therapy.
- GAMMA GLOBULIN**, (Breon) Vial 10 cc.
- GAMMA GLOBULIN**, (Cutter; Lederle; Pitman-Moore) Labeled "Poliomyelitis Immune Globulin (Human)" Vial 2 & 10 cc.
Use: Prevention of paralytic polio, measles & infectious hepatitis.
- GAMMA GLOBULIN**, (Hyland Labs.) Pooled normal human plasma containing concentrated antibody-bearing gamma globulin fraction. Vial 2 & 10 cc.
Use: Treatment of measles & poliomyelitis.
- GAMMA GLOBULIN**, (Parke, Davis) Pooled normal human plasma containing concentrated antibody-bearing gamma globulin fraction. Vial 2 & 10 cc.
Use: Treatment of measles & poliomyelitis.
- GAMOPHEN**, N.N.D. 1962, A.D.R. (Ethicon Suture) Soap containing 2% hexachlorophene Cake 127.5 Gm., liq. 3 oz., 12 oz., Gal.
Use: Local anti-infective.
- GANATREX ELIXIR**, (Merrell) Vit. A 1667 u., D 167 u., E 0.6 u., folic acid 0.6 mg., Vit. B-1 3.3 mg., B-2 1.7 mg., B-6 0.3 mg., B-12 3.3 mcg., niacinamide 17 mg., inositol 33.3 mg., choline 33.3 mg., w/Cal. glycerophosphate, iodine & minerals/dr. Bot. 1 pt., 1 gal.
Use: (2-3 tsp. t i d) Metabolic supplement
- GANGLIONIC BLOCKING AGENTS**.
See: Ansoletysen (Wyeth).
Ariponid (Roche Lab)
Benodaine HCl, Amp. (Merck Sharp & Dohme)
Bisrium Bromide (Squibb).
Bisrium Chloride High Potency (Squibb).
Chlorisondamine Citrate.
Dibenzylamine HCl, Cap. (Smith, Kline & French).
Ecolid. (Ciba)
Evamide Chloride (Ciba).
Etaman Chloride, Vial (Parke, Davis).
Hexalin Chloride (McNeil).
Hexamethanum Chloride & Bromide (Various Mfr).
Hexametan Chloride (Burroughs Wellcome).
Hydergine, Amp. Tab. (Sandoz).
Inversine, Tab. (Merck Sharp & Dohme)
Mecamylamine HCl.
Methium Chloride (Warner-Chitcott).
Pentolinum Tartrate.
Priscoline HCl, Elixir, Tab., Vial (Ciba).
Regisine, Amp., Tab. (Ciba).

- Tetraethylammonium Chloride (McNeil & Parke Davis)
Trimelhaphan
- GANONE**, (Endo) Tab, Greens d methamphetamine HCl 3 mg, ketocholanic acids 130 mg, thiamine HCl 1 mg/Tab; Orange d methamphetamine HCl 6 mg, ketocholanic acids 130 mg, thiamine HCl 1 mg/Tab Bot 100s, 1000s
Use Management of obesity
- GANTANOL** (Roche Lab) Sulfamethoxazole 0.5 Gm/Tabs Bot 50s 300s
W/Phenazopyridine See Azo Gantanol
Tob (Roche)
Use Antibacterial
- GANTRICILLIN-100**, (Roche Lab) Pot penicillin G 100 000 u, gantrisin 0.5 Gm/Tab Bot 24s 100s, 500s
Use (1 2 q 4 h) Sulfonamide & penicillin therapy
- GANTRICILLIN-200** (Roche) Gantrisin acetyl 0.5 Gm, Pot penicillin G 200 000 u / 5 cc Susp, Bot 2 oz when prepared as directed
Use Penicillin Sulfam therapy
- GANTRICILLIN 300**, (Roche) Pot penicillin G 300 000 u, gantrisin 0.5 Gm/Tab Bot 24s, 100s, 500s
Use: Penicillin with sulfonamide therapy
- GANTRIMYCIN**, (Roche) Gantrisin 333 mg, oleandomycin phosph 75 mg/Tab Bot 50s
Use: Double spectrum antibacterial
- GANTRISIN, USP** NND 1962 ADR (Roche) Sulfisoxazole 3,4 dimethyl 5 sulfonamide/sulfazole
Tab (0.5 Gm) Bot 100s 500s 1000s
Amp (5 & 10 cc) containing 2 & 4 Gm gantrisin, respectively, in ready to use solution in the form of gantrisin diethanolamine) Box 6s 25s
Pediatric Susp (0.5 Gm / tsp) Bot 4 oz, 1 pt
Pow (Not Sterilized) Bot 4 & 16 oz
Syr 0.5 Gm / tsp 15 cc) Bot 4 & 16 oz
Use: Antibacterial sulfonamide therapy
W/Oleandomycin See: Gantrimycin, Tab (Roche)
- W/Phenylaza diamino pyridine HCl See Azo Gantrisin, Tab (Roche)
W/Pot penicillin G See: Gantricyllin (Roche)
W/Pot penicillin G & Acet Dia Mer sulfamides See: Trilonacil, Tab (Firm, Eaton)
- GANTRISIN ACETYL**, NND 1962 (Roche) Acetyl Sulfisoxazole Susp Pediatric (100 mg/cc) Bot 1 pt; Syrup (100 mg/cc) Bot 1 pt See: Lipo Gantrisin, Bot (Roche)
Use: Sulfonamide therapy.
- GANTRISIN, AZO**, See: Azo Gantrisin, Tab (Roche)
- GANTRISIN DIETHANOLAMINE EAR SOLUTION**, NND 1962 (Roche) Gantrisin 4%, urea 10%, chlorobutanol 3% Sol Bot ½ oz
Uses: Ear Infections
- GANTRISIN DIETHANOLAMINE NASAL SOLUTION**, NND 1961 (Roche) 4% Gantrisin diethanolamine & ¼% phenylephrine HCl Vial w/dropper 1 oz; Bot 1 pt
Use Local antibacterial & decongestive action
- GANTRISIN DIETHANOLAMINE OPHTHALMIC SOLUTION & OINTMENT**, NND 1961 (Roche) Both gantrisin ophthalmic oint & sol are stable antibacterial prep containing 4% gantrisin in the form of gantrisin diethanolamine Ophth Oint Tube ½, 1 oz; Sol Vial 5 cc, 15 cc
Use Antibacterial, sulfonamide therapy
- GANTRISIN PEDIATRIC SUSPENSION**, NND 1961 (Roche) Gantrisin 0.5 Gm/di Bot 4 fl oz, 1 pt
Use Sulfonamide therapy
- GANTRISIN VAGINAL CREAM** (Roche) Cream (10%) Tube 3 oz W/Applicator W/18 applicators or W/a applicator
- GARDENAL** See Phenobarbital (Various Mfr)
- GARDINOL TYPE DETERGENTS** Aural
Cyclopon, Dreff, Drene Dupanol Lissopal
Maprofix, Modinal Orvus, Sandopan
Sodipan
Use Detergents
- GARDOL** Sod Laurayl Sarcosinate
- GARGALINE**, (Blue Line) Phenol 7½ m iodine, Pot Iodide zinc sulfate hydraline HCl & Ionic acid/oz Bot 1 pt
Use Throat & mouth infections
- GARLIC**, Allium
Use Intestinal antispasmodic
Guardian—Cap & Tab Bot 100s 1000s
Levitt—6 min Cap Bot 100s 1000s
Testagar—Cap Bot 100s, 1000s
See: Allimin (Van Patten)
- GARLIC CONCENTRATE** W/Parsley Concentrate See: Allimin, Tab (Van Patten)
- GARTRONE**, (Eric, Kirk & Gray) Veratrum Viride Alk 250 mg/100 cc Vials
Use 1M 1.2 cc; 1Y 0.2 cc in 5% glucose
- GAS GANGRENE ANTITOXIN**, NF X NNR 1952
Lederle—(Polyvalent) Vial 1 dose
Lilly—(Therapeutic) Vial 1 dose
National Drug—(Trivalent) Vial 1 dose
Parke, Davis—(Trivalent) Vial 1 dose
Wyeth—(Polyvalent) Vial 1 dose
- GASTRALME**, (Meyer) Cal carb 0.7 Gm, aluminum hydroxide gel 0.2 Gm, Mag trisilicate 0.1 Gm/Tab Bot 40s
Use: Antacid
- GASTRALME LIQUID**, (Meyer Chem) Cal carb 0.7 Gm, Mag trisilicate 0.1 Gm,

- aluminum hydroxide gel, dried, 0.2 Gm./8 cc. Bot. 1 pt.
 Use: (2 tsp.) Prolonged reduction of hyperacidity.
- GAST-REX.** (Paul Money) Gastric mectin 120 mg., aluminum hydroxide gel 240 mg., magnesium trisilicate 360 mg., hamatropine methylbromide 0.2 mg./tab. Bot. 100s, 1000s.
 Use: Gastric disorders.
- GASTRIC ELECTROLYTE SOLUTION.** (Don Baxter) Sod. chloride 0.37 Gm., pot. chloride 0.13 Gm., amm. chloride 0.37 Gm., dextrose 10 Gm./100 cc. Bot. 1000 cc.
 Use: I.V., replace electrolytes
- GASTRIC MUCIN, N.N.D. 1959.**
 Willan—Gran Bot. 8 oz.
 Winthrop—Gran. Bot. 8 oz.
 Use: Orally, ulcers.
- W/Aluminum hydroxide gel dried, hamatropine methylbromide & Mag. trisilicate.
 See: Mucosa-Seth, Tab. (Harrower).
 Trimucalon, Tab. (Winthrop Labs.)
- W/Aluminum hydroxide gel & Mag. trisilicate.
 See: Mucatin, liq. & Tab. (Harrower).
 Trimucalon, Tab. (Winthrop Labs.)
- N/Aluminum hydroxide gel, magnesium trisilicate, hamatropine methylbromide. See: Gast Rex, Tab. (Money)
- N/Dihydroxy aluminum olnineacetate, ethaverine HCl. See: Mugastral, Tab. (G. F. Harvey)
- N/Magnesium glycinate and aluminum hydroxide. See: Mucogel, Tab. (Barrows Chemical Co)
- N/Pentobarbital Sod., hamatropine methylbromide, aluminum hydroxide & Mag. trisilicate See: Antased, Tab. (Lemmon).
- ASTROGRAFIN, N.N.D. 1962.** (Squibb) Radiopaque medium. Bot. 4 fl. oz.
 Use: Roentgenographic study of the gastrointestinal tract.
- ASTROLIC.** (Mallard) Phenobarbital $\frac{1}{4}$ gr., belladonna lvs. $\frac{1}{2}$ gr., benzocaine 2 gr./Tab. Bot. 100s, 500s, 1000s.
 Use: (1-2 t.i.d.) Anadyne, sedative & antispasmodic.
- ASTRON** W/Vit. B-12. (Winthrop Labs.) Stomach mucosa bag ext, w/Vit. B-12 7 $\frac{1}{2}$ mcg./oz. Bot. 6 oz., 22 oz.
 Use: (1-2 tsp. p.c.) Promote digestion.
- B. S.** (G. F. Harvey) Oxidized bile acid 1 $\frac{1}{2}$ gr., atropine methyl nitrate 1/60 gr. ext of ox & hog bile 1 $\frac{1}{2}$ gr., phenobarbital $\frac{1}{4}$ gr./tab. Bot. 100s, 1000s.
 Use: Biliary stasis & insufficiency.
- BEL-A-NEUTRIN.** (Walker) Hamatropine methylbromide 1/48 gr., aluminum hydroxide

- ide gel, dried, 2 gr., Mag. trisilicate 3 gr., Mag. hydroxide 1 gr., pectin $\frac{1}{2}$ gr./Tab. Bot. 100s, 500s, 1000s.
 Use: Hyperacidity, in the treatment of gastric & duodenal ulcers.
- GELATIN, U.S.P., Glycerinated, U.S.P.**
 See: Pharmagel A & B.
 Use: Gelatin foam or sponge w/thrombin as absorbable local hemostatic; 5% sol. has been used as plasma substitute in shock.
 See: Plazmoid (Upjohn)
- GELATIN COMPOUND, PHENOLIZED, N.N.D. 1951.** (Merck Sharp & Dahme) Gelatin 14%, phenol 1 $\frac{1}{2}$ %, zinc oxide 5 $\frac{1}{2}$ %, w/Glycerin & water. Jar 1 lb., 5 lb.
 Use: Topically, antiseptic protective.
- GELATIN-EPINEPHRINE.** (Rorer) Epinephrine HCl 1:500 in aq. gelatin base. Amp. 1 cc., Box 12s; Vial 10 cc.
 Use: S.C., I.M., bronchial antispasmodic.
- GELATIN FILM, ABSORBABLE, N.N.D. 1962.** See: Gelfilm (Upjohn)
- GELATIN IN ISOTONIC SOLUTION OF SODIUM CHLORIDE.** See: Plazmoid, liq. (Upjohn)
- GELATIN SOLUTION, SPECIAL INTRAVENOUS, N.N.D. 1962.** 6% sterile isotonic gelatin sol
 Knox Gelatin Co. Bot. 300 cc. containing 60 mg. gelatin/cc.
 Use: Blood volume expander
- GELATIN SPONGE, ABSORBABLE, U.S.P., N.N.D. 1962, A.D.R.** See: Gelfoam (Upjohn)
- GELAZYME.** (Bebbs) Activate aluminum hydroxide gel 4 gr., Mag. trisilicate 7 $\frac{1}{2}$ gr. & pancreatin triplex $\frac{1}{2}$ gr./Tab. Bot. 100s.
 Use: (1-3 t.i.d.) Gastric hyperacidity.
- GEL-ETS TRIASYN B.** (Brewer) Vit. B-1 2 mg., B-2 3 mg., nicotinamide 20 mg./cap. Bot. 100s, 500s, 1000s.
 Use: Vitamin B therapy.
- GELFILM, N.N.D. 1961.** (Upjohn) Sterile non-antigenic absorbable water-insoluble gelatin film. Box 6 films, 100 mm. X 125 mm. X 0.16 mm. Also available as Ophth. Sterile 25 X 50 mm. Box 6s
 Use: Aid in surgical closure.
- GELFOAM, N.N.D. 1961, A.D.R.** (Upjohn)
 Size 12—3 mm.—20 X 60 mm. (12 sq. cm.) X 3 mm. Boxes of 4 sponges in ind. envelopes
 Size 12—7 mm.—20 X 60 mm. (12 sq. cm.) X 7 mm. Boxes of 4 sponges in ind. envelopes, and jars of 4 sponges
 Size 50—62.5 X 80 mm. (50 sq. cm.) X 10 mm. Boxes of 4 sponges in ind. envelopes
 Size 100—80 X 125 mm. (100 sq. cm.) X

¹ Generic name recognized by N.F., U.S.P. or A.M.A.

10 mm Boxes of 6 sponges in ind envelopes
 Size 200—80 X 250 mm (200 sq cm) X
 10 mm Boxes of 6 sponges in ind envelopes
 Compressed size 100 (intended primarily for application in the dry state) 12.5 X 8 cm Boxes of 6 sponges in ind envelopes
 Packs size 2 cm (designed particularly for nasal packing) 2 X 40 cm Boxes of 6 sponges in ind bottles
 Packs size 6 cm (for large surface areas and packing cavities) 6 X 40 cm Boxes of 6 sponges in ind envelopes
 Prostatectomy cones (for use with Foley catheter)
 Size 13 cm (approx 13 cm in diameter) and
 Size 18 cm (approx 18 cm) Both in boxes of 6 cones in ind envelopes
 Use: Hemostasis in neurosurgery, otolaryngology, bone surgery, malignancy, dental surgery and abdominal surgery
GELFOAM COMPRESSED (Upjohn) Size 100 (12.5 cm X 8 cm) Box 6s, 36s, 72s, 144s
 Use Topically hemostatic
GELFOAM DENTAL PACK (Upjohn) 20 mm X 20 mm X 7 mm Jar 15 sponges Size 2 10 mm X 20 mm (2 sq cm) Jars 15 sponges
 Use: Control bleeding
GELFOAM PACKS (Upjohn) {6 cm X 40 cm} Box 6s, 36s, 72s, 144s
 Use Hemostatic
GELFOAM POWDER (Upjohn) Con 1 & 10 Gm Sterile or nonsterile
 Use: Orally, hemostatic.
GELFOAM PROSTATECTOMY CONES (Upjohn) 13 cm in diameter & 18 cm in diameter Box 6s, 36s, 72s, 144s
 Use: Hemostatic
GELFOAM TUMOR DIAGNOSIS KIT (Upjohn) 2 sterile Gelfoam biopsy swabs & 1 30 cc. bot of fixative sol (Formaldehyde Sol 1:10)/kit
 Use: Gladrone technic of tumor diagnosis.
GELINA VAGINAL GEL (Lemmon Pharm co) 9 aminopropyl na HCl 0.20%, allantoin 0.75%, cetalkonium chloride 0.05%, lactose 3.0%, lactic acid 3.0% /ml Tube 75 Gm /w/applicator
 Use: Vag nitis therapy
GELOCAST (Duke Lab) Zinc oxide, calamine and gelatin bandage 10 yd X 4 in
 Use: Unna's cast dressing r
GELPIRIN (Schuermann Jones) Aspirin caffeine and acetophenetidin buffered wth gelminate/Tab Bot 100s
 Use: Analgesic

GELSEMIUM, N F IX. (Various Mfr) Pkg 1 oz
 Use: For neuralgia
 W/Aspirin & caffeine in propylene glycol
 See Anogill, Liq (Massengill)
 W/APC See APC Combinations
 W/Atropine sulfate, strychnine sulfate and podophyllin, camphor monobromated, aspirin & phenacetin See Aspacom, tea (Smith Oarsey)
 W/Colchicine salicylate, aspirin, camphor monobromated & acetophenetidin See Formasal & Camphor Comp, Cap (First Texas)
 W/Phenobarbital & caffeine See Phac Tab (Cale Chem Co)
GELSODYNE TABLETS (Blue Line) Aspirin 3 1/2 gr, acetophenetidin 2 1/2 gr, caffeine 1/2 gr, gelsemium tr 1 min /Tab Bot 100s
 Use: (1 3 1 d) Analgesic & antipyretic
GELU-CILLIN (Warner Chilcott) Pot penicillie G, cryst 50 000 & 100 000 u buffered w/aluminum hydroxide & col carb /Tab. Pkg 12s, 100s
 Use: Orally, penicillin therapy
GELU-CILLIN TROCHES (Warner Chilcott) Pot penicillin G 5000 u Bot 25s 1000s
 Use: Orally, penicillin susceptible organisms of throat & mouth
GELUMINA (APC) Aluminum hydroxide colloidal Susp Bot 8 fl oz 1 pt, 1 gal. Tab Bot 100s 1000s Available w/Mog Irrisilicate
 Use: Gastric antacid
GELUSIL (Warner Chilcott) Mog Irrisilicate 7 1/2 gr, aluminum hydroxide 4 gr /tab or dr susp Bot 6 oz, 12 oz; Tab Bot 50s 100s 1000s
 Use: Gastric hyperacidity
GELUSIL FLAVOR-PACK (Warner Chilcott) Aluminum hydroxide 4 gr, magnesium trisilicate 7 1/2 gr per 5 cc Bot 12 oz w/separate packages of flavoring agent.
 Use: Gastric hyperacidity
GELUSIL IAC. (Warner Chilcott) Magnesium trisilicate 30 gr, aluminum hydroxide 16 gr w/high protein milk solids/16 Gm Bot 320 Gm
 Use: Treatment of hyperacidity
GEMINIMYCIN A AND B (Pfizer) Antibiotics
 Use: Inhibits many gram positive bacteria
GEMIVITE (Ulmer) Combination of 11 Vit-12 minerals, choline, inositol/Cop Bot 60s
GEMONIL, N N O 1962 (Abbott) Metharbital (5,5 Diethyl 1 methyl barbiturate acid) Tab 01 Gm Bot 100s, 1000s
 Use: (1 1 1 d) Grand mal, petit mal & myoclonic epilepsy & in mixed types of seizures
GINATRATE (General Drug & Chem) Pen

erythritol tetranitrate, 10 or 20 mg./Tab. Bot. 100s, 500s.

Use: Vasodilator.

* **GENATROPINE NCL.** Atropine N-oxide HCl. Aminasytropine Tropate HCl. See: Xtra (Xitrum Lab.)

GENIREX. (Nevama Products) Nicotinic acid, caffeine, calcium succinate, thiamine, nboflavin Tablets Bot. 10s.

Use: Vitamin & energizer.

GENITONE. (Merrill) Viburnum prunifolium 4.15 Gm., passiflora 5.2 Gm., pulsatilla 1.04 Gm., senecio aureum 2.6 Gm., aromatic cordial, q. s. ad 30 cc., alcohol 12%/30 cc. Bot. 1 pt.

Use: (1-2 tsp. t.i.d.) Uterine sedative.

GENOPHYLLIN. See: Aminophylline (Various Mfrs.)

GENOSCOLAMINE HBr. (Lobico-Debrulle) Scopolamine amineoxide HBr, scopolamine-N-oxide HBr, epaxaminooxytropine tropate HBr, Pellets: 0.5 mg. Box 60s. See: Scopodex (Lobico-Debrulle)

Use: Orally, hypnotic & sedative.

W/Ergotamine tartrate, caffeine, chloral hydrate. See: Rectal Migraine, Dis. Applicators (Wempole Lab.)

W/Mephenesin, mephobarbital & betaine HCl. See: Stophenesin, Tab. (Bayer Nelson)

GENSALATE SODIUM. (Parroy) Sod. gentisate. (Sod. salt of 2,5-dihydroxybenzoic acid) Tab. 0.5 Gm.

Use: (2 q. 3 h.) Analgesic & antipyretic.

GENTAMIN. (American Pharm.) Sod. gentisate 2½ gr., salicylamide 2½ gr./tab. Bot. 100s, 1000s.

Use: Analgesic.

GENTARTH. (Raymer) Each enteric-coated tab. contains 100 mg. Sod. gentisate, 325 mg. mysal (representing 43% salicylic acid & 3% iodine in Cal. Sod. phosphate buffer combination) & 130 mg. succinic acid. Bot. 100s, 500s, 1000s.

Use: (1-2 t.i.d. or q.i.d.) Rheumatic diseases especially arthritis.

GENTASOL. (Gold Leaf) Sod. gentisate. Tab. 0.5 Gm. Bot. 50s, 100s, 500s, 1000s.

Use: (4 q. 3 h.) Anti-rheumatic & analgesic.

GENTERSAL. (Ortho) Gentian violet 0.05%, alkyl-dimethylbenzylammonium chloride 0.05%, absorbable cream base. Tube 2½ oz. w/disposable applicator.

Use: Vaginal moniliais.

GENTEXIN. (Crookes Barnes) Sod. gentisate 200 mg., acetaphenatidin 200 mg., di-dexoxyephedrine HCl 2 mg./Tab. Bot. 50s, 500s, 1000s.

Use: (1-3 q. 3-4 h.) Relieves pain of rheumatism & brightens mood.

* Generic name recognized by N.F., U.S.P. or A.M.A.

GEN-THERA. (Gen. Drug & Chem) Vit. B-1 15 mg., B 2 & mg., C 50 mg., calcium pantothenate 5 mg., B 6 0.5 mg., niacinamide 30 mg., desiccated whole liver 175 mg., yeast conc. 75 mg., ferrous gluconate 30 mg., choline dihydrogen citrate 20 mg., inositol 25 mg., methionine 25 mg., B-12 3 mcg., folic acid 0.1 mg./Cap. Bot. 100s, 500s.

Use: Water soluble vitamin deficiencies.

GENTIA JEL. (Westwood) 0.2% Gentian violet, 3% lactic acid, 1% acetic acid. 1-dose applicators. Pkg. 12s.

Use: Vaginal moniliais.

GENTIAN E.V., Suppberries. (Webster) Gentian violet 0.25%, benzocaine 2.0%. Suppberries, Jar 14s.

Use: Orally; Enterobiasis semicirculari.

GENTIAN VIOLET, U.S.P., N.N.R. 1952, A.D.R. 1955. Methylrosaniline chloride. Available. Alcoholic sol., aq. sol., I.C. Tab., Pow., Supp., Tab. Bismuth Violet.

See: Gintersol. (Ortho)

Pyoktanin. (Merck)

W/Acridine & brilliant green. See: Dymixal, Pow. (McNeil)

W/Benzocaine. See: Gentian E.V., Suppberries (Webster).

W/Preservatives and inert materials. See: G.V.S. Aerosol Foam (Sarage)

W/Surfactants. See: Hyva, Vag. Tab. (Holland-Rentas)

GENTIAN VIOLET SUPPRIES. (Webster) Gentian violet 0.2%, lactic acid 0.3%, acetic acid 1% in Neocera base. Jar 12s.

Use: Vaginal fungicide.

GENTISATE SODIUM. 5-Hydroxysalicylate sod., 2,5-Dihydroxy benzoate sodium

See: Casale Sod., Tab. (Sulliff & Case)

Gensalate Sodium, Pow. & Tab (Parroy)

Gentisol, Tab. (Gold Leaf)

Gentam, Tab. (Crookes-Barnes)

Na-Gent, Pow. & Tab. (Raymer)

W/Salicylamide. See: Gentamin, Tab. (American Pharm.)

W/Succinic Acid & Ray-Sol. See: Gentarth, Tab. (Raymer)

GENTISIC ACID. 5-Hydroxysalicylic acid.

W/Vitamins. See: Dar-Plex. (Darwin)

GENTLAX GRANULES. (Purdue-Frederick) Sennasides A and B 15 mg., guar gum 1 Gm., polygalacturonic acid 100 mg./Tsp Pkg 3 oz., 6 oz.

Use: Treatment of constipation.

GENTRAN, N.N.O. 1962 (Don Baxter) Gentrin 6% in saline. Bot. 250 cc., 500 cc.

Use: Plasma substitute.

GENTRAN IN TRAVERT. (Baxter) Gentian

GERICAPS. (Sherman) Choline dihydrogen citrate equiv. 500 mg., inositol 200 mg., rutin 20 mg., ascorbic acid 12.5 mg., Vit. A 1000 u., thiamine HCl 1 mg., riboflavin 1 mg., pyridoxine HCl 0.25 mg., niacinamide 4 mg., Cal. pantothenate 1 mg./Cap. Bot. 100s.

Use: (1-3 daily) Cholesterol metabolism therapy.

GERICOLE. (Sherman) Choline dihydrogen citrate 250 mg., inositol 100 mg., ascorbic acid 75 mg., thiamine HCl 12.5 mg., riboflavin 1.25 mg., pyridoxine HCl 0.62 mg., niacinamide 12.5 mg., Cal. pantothenate 2.5 mg./5 cc. Syr. Bot. 3 oz., 1 pl.

Use: (1 trp. 1 d) Vitamin supplement.

GERI-DEIMAL. (Boyle) Vit. B-12 w/intrinsic factor 1/10 u., Vit. B-12 2 mcg., B-1 6 mg., B-2 6 mg., B-6 0.5 mg., cal. pantothenate 5 mg., niacinamide 100 mg., folic acid 0.34 mg., biotin 10 mg., Vit. A 6000 u., D 600 u., E 5 u., K 1 mg., C 75 mg., hesperidin 25 mg., choline bitartrate 200 mg., inositol 25 mg., betaine HCl 100 mg., pancreatin 97.2 mg., pepsin 97.2 mg., bile acids 25 mg w/cal, iron, copper, iodine, manganese, pot. magnesium, zinc/2 Tabs. Bot 100s, 1000s

Use: Geriatric vitamin and mineral supplement

GERI-DEIMAL-N. (Boyle) Methyl testosterone, 50 mg; ethyl estradiol, 0.01 mg; B-12 with intrinsic factor conc. 1/15 oral unit; amylose, 75 mg; pepsinase, 30 mg; B-1, 30 mg; B-2, 20 mg; B-6, 1.0 mg; panthenol, 50 mg; niacinamide, 500 mg; l lysine monohydrochloride, 2000 mg; A, 5000 USP units; D, 500 USP units; E, 25 mg; C, 750 mg; choline bitartrate, 1500 mg; inositol, 250 mg; betaine HCl, 1000 mg; pancreatin, 97.2 mg; Fe, 200 mg; Cu, 0.67 mg; I, 0.15 mg; Mn, 20 mg; Mg, 100 mg; Zn, 15 mg; Mo, 0.1 mg / Tab Bot. 100r, 1000r

Use: Support in nutritional and hormonal problems of aged and convalescents

GERIFORY. (Amer. Pharm. Co) Vit. A 28,000 USP. u., B-1 14 mg., B-2 14 mg., B-6 210 mg., C 210 mg., niacinamide 180 mg., iron, Cal., Phos., iodine, Cal. pantothenate, folic acid, Vit. E, Vit. B-12, choline bitartrate, inositol, dl-methionine, rutin, biotin, cobalt, copper, Mag., Mang., Pot. zinc/Cap. Bot. 100r.

Use: Supplementary nutritional needs

GERIFORY ELIXIR. (A.P.C.) Thiamine HCl 5 mg., riboflavin 5 mg., iron 100 mg., iodine 1 mg., niacinamide 50 mg., pyridoxine HCl 1 mg., Vit. B-12 2 mcg., panthenol 10 mg., choline chloride 200 mg.

inositol 100 mg., dl-methionine 100 mg., cobalt acetate 0.2 mg., copper acetate 2 mg., Mag. chloride 2 mg., Mang. citrate soluble 2 mg., zinc acetate 2 mg., w/Vit. B complex factors naturally occurring in 0.9 Gm. yeast ext./Cap. Bot. 12 oz.

Use: Geriatric & general nutritional fortifier.

GERIJECT INJECTION. (Kirk) Vit. B-12 30 mcg., B-1 20 mg., B-2 1 mg., niacinamide 100 mg., B-6 2 mg., liver inj 1 mcg., ferric citrate 12 mg., choline chloride 20 mg., inositol 10 mg., dl-methionine 5 mg., pantothenol 5 mg., benzyl alcohol 2%/cc. Vitol 10 cc.

Use: I.M., 1 cc. weekly. Treatment of the geriatric patient.

GERIJECT TABLETS. (Kirk) Vit. A 2500 U.S. D 250 u., B-1 1 mg., B-2 1.5 mg., B-6 0.1 mg., folic acid 0.1 mg., cal. pantothenate 1 mg., E 0.25 mg., rutin 10 mg., inositol 10 mg., choline bitartrate 30 mg., dl-methionine 10 mg., w/Mrneral/tab. Bot. 100s.

Use: Treatment of the geriatric patient.

GERI-K. (Kirkman) Vitamin (13), mineral (10), dl-methionine/cap. Bot. 100s, 1000s

GERILAC. (Sorden) Whale milk, skim milk, brewer's yeast, riboflavin concentrate, Vit. A & D concentrate, ergosterol (Vit. D), ascorbic acid, niacinamide, monosodium phosphate & iron citrate, Can 1 lb.

Use: Dietary supplement

GERILETS. (Abbott) Thiamine mononitrate 5 mg., riboflavin 5 mg., pyridoxine HCl 1 mg., niacinamide 20 mg., cal. pantothenate 5 mg., Vit. A 5000 u., Vit. D 500 u., Vit. E 10 IU, Bevidoral 1/2 u., ferrous sulf 75 mg., folic acid 0.25 mg., ascorbic acid 50 mg., Quertine 12.5 mg., betaine HCl 50 mg., inositol 50 mg., Desoxy HCl 1 mg., Solalex 0.3 mg., methyltestosterone 2.5 mg./Frimtab. Bot. 25s, 100r, 250s

GERI LIQUID. (Lakeride) Each trp. (5 ml) niacin 75 mg., glycine 750 mg in cherry wine base Bot. 8 fl oz, 16 oz

Use: Poor peripheral circulation.

GERINEED. (Hontan Drug) Liquid. Bot 16 oz.

GERI PLEX. (Parke, Davly) Vit. A 5000 u., B-1 5 mg., B-2 5 mg., B-12 2 mcg., niacinamide 15 mg., Vit. C 50 mg., choline dihydrogen citrate 20 mg., mixed tocopherols 10 mg., laka diartare 1 gr., rutin 25 mg., ferrous sulfate 30 mg., copper sulfate 4 mg., Mang sulfate 4 mg., zinc sulfate 2 mg., dicalcium phosphate (anhydrous) 200 mg./Cap Bot 30r, 100r, 500s

Use: Vitamin mineral combination.

GERISAL. (Geriatric) Sod. salicylate 5 gr., mephensin 3 gr., secobarbital 1/2 gr., phenobarbital 1/2 gr., ascorbic acid 5/6 gr.,

- aluminum hydroxide gel 1 gr/Tab Bot 100s 1000s
 Use: (1 2 pc) Management of arthritis & allied disorders
- GERITAG.** (Tutag) Methyl testosterone 2 mg, ethinyl estradiol 0.01 mg, ferrous sulf 50 mg rutin 10 mg, Vit A 5000 u, 0 400 u, E 11 U, C 30 mg, B 12 1 mcg, B 1 2 mg, B 2 2 mg, B 6 0.3 mg, cal pantothenate 3 mg, niacinamide 20 mg, choline bitartrate 40 mg methionine 20 mg inositol 20 mg, malyb 0.5 mg, ca 0.1 mg, cu 0.2 mg, mang 1 mg, mag 5 mg, iodine 0.15 mg, pot 2 mg, zinc 1 mg / Cap Bot 100s 500s, 1000s Vial 10 cc
 Use Orally, I.M., dietary supplement in geriatrics
- GERITINIC** (Geriatric) Vit A 5000 u, 0 500 u, B 1 3 mg, B 2 3 mg, B 6 0.1 mg, B 12 5 mcg, C 60 mg, niacinamide 10 mg, folic acid 0.1 mg, inositol 20 mg choline bitartrate 30 mg, methionine 24 mg amino acetic acid 30 mg, Cal pantothenate 1 mg, desic liver 22.4 mg, conc yeast 22.4 mg, glutamic acid HCl 120 mg, Pot acetate 30 mg, cobalt chloride 15 mg, ferrous sulfate 195 mg, copper, zinc, cobalt & Mang phylins/Cop Bot 42s 500s
 Use (1 1 d) Nutritional anemias
- GERITOL JUNIOR** (Pharmaceuticals) Vitamins A C O, B complex including B 12, plus iron Bot Tob 40s liq Bot 8 oz.
- GERITONE** (Myers-Carter) Testosterone 10 mg, estrone 1 mg, with B vits / 2 cc Vial 20 cc
- GERITONIC LIQUID** (Geriatric) Thiamine chloride 1 mg, riboflavin 1 mg, Vit B 12 3 mcg, niacinamide 10 mg, Pyridoxine HCl 0.1 mg, choline as dihydrogen citrate 20 mg, inositol 20 mg, cobalt as chloride 0.1 mg, Mag as sulfate 1 mg, iodine as Pot iod 0.02 mg, iron as ammonium citrate 20 mg, Mang as sulfate 0.5 mg, strychnine sulfate 0.3 mg, Cal as glycerophosphate 10 mg, Phos as Cal glycerophosphate 8 mg, Pot as sulfate 1 mg, aminoacetic acid 60 mg, yeast conc 125 mg, liver fraction I 125 mg, molybdenum as Sod molybdate / 5 cc Bot 8 oz. 1 gal
 Use Nutritional deficiencies
- GERITROPIN** (U S Standard) Tricheline citrate 4.15 Gm (equiv to choline base 3 Gm) inositol 750 mg Vit B 12 (cryst) 5 mcg / 15 cc Bot 1 pt, 1 gal's Tabocap Bot 100s, 500s 1000s
 Use Lipid metabolism
- GERI-VITE** (Rexall) Vit B 1 10 mg, Vit B 2 5 mg, niacinamide 50 mg, Vit C 50 mg Vit B 12 1 mcg, folic acid 0.2 mg, Vit B 6 0.35 mg, methionine 50 mg, choline dihydrogen citrate 20 mg, inositol 20 mg, pantothenic acid 5 mg, dried yeast 100 mg, liver desic 50 mg / Cap Bot. 100s
 Use: Dietary supplement
- GERIX ELIXIR.** (Abbott) Vit B 1 1 mg, B 2 1 mg, B 6 0.33 mg, B 12 1 mcg, iron cinamide 333 mg, choline 33 mg, w/iron 35 mg inositol 333 mg, betaine HCl 333 mg / dr Bot 1 pt
 Use: Hematinic
- GERIZYME.** (Upjohn) Liver conc 75 mg ferrous gluconate 43.17 mg, Cal glycerophosphate 100 mg, B 1 3.3 mg B 2 3.3 mg, B 6 1 mg, nicotinamide 333 mg d pantothenyl alcohol 33 mg B 12 3.3 mcg, inositol 100 mg choline dihydrogen citrate 100 mg, cobalt 0.33 mg, copper 0.33 mg, mang 0.33 mg, mag 2 mg molybdenum 0.066 mg zinc 0.5 mg, pot 2 mg / 15 cc Bot 12 oz, 1 pt
 Use Geriatric tonic
- GERMA-MEDICA, NNR 1953, ADR** (Huntington) Soap containing 1% hexachlorophene (anhy basis) Bot 378 L
 Use: Surgical soap w/anti-infective agent
- GERMANIN** See: Suramin Sodium (Naphthide Sod) (Winthrop Labs)
- GERMICIN.** (Consolidated Mid) Benzalkonium chloride 12.8% Bot 1 pt 50% Bot 1 gal
 Use (diluted) Antiseptic
- GERMOL-E.** (Ayerst) Tocopherols natural mixed Cap 34 mg Bot 100s
 Use Vitamin E therapy
- GER-O-FOAM.** (Geriatric) Methylsilylate benzocaine and volatile oils Aerosol can 4 oz
 Use Anesthetic analgesic
- GERONE.** (Pitman Moore) Dextro amphetamine sulfate 2 mg, thiamine HCl 2 mg nicotinamide 10 mg, riboflavin 0.5 mg pyridoxine HCl 0.5 mg, Cal pantothenate 1 mg / 5 cc or Tab. Elixir, Bot 8 oz, Tab Bot 100s, 1000s
 Use (2 tsp or Tab 1 d) Cerebral stimulant w/vitamin B therapy
- GERONIAZOL.** (Calumbus) Pentylene tetrazol 100 mg, nicotinic acid 50 mg / Tab or 1 cc Bot 100s, 500s, 1000s Inj Vial 10 cc Liq 5 cc eq 1 tab pt
 Uses: (1 or 2 1 s d) Stimulant for depressed states
- GERONIAZOL TT** (Temporal) (Calumbus Ph Cal) Pentylene tetrazol 300 mg nicotinic acid 150 mg / Tab Bot 42s
 Use Geriatric
- GESIK** (Schilling) N acetyl p aminophenol 240 mg acetophenetidin 60 mg caffeine 15 mg B 1 10 mg / Tab Bot 30s 100s
 Use Analgesic

GESTANIN. (Organon Inc) See: Allylestrenol.

GESTATABS. (White Lab) Vit. A 6000 u., D 600 u., K (menadione) 2 mg., B-12 activity equiv. to 2 mcg., folic acid 1 mg., ascorbic acid 100 mg., thiamine mononitrate 3 mg., riboflavin 5 mg., pyridoxine HCl 1.5 mg., Col. pantothenate 10 mg., nicotinamide 30 mg., mal-l-iron ferrous sulfate 120 mg., molybdenum oxide 1.8 mg., Cal. 380 mg./Tab. Bot. 60s, 1000s.
Use: (1 daily) Prenatal supplement.

GESTEROL. (Testagar) Progesterone. Aq. Susp. Vial (25 mg./cc. & 50 mg./cc.) 10 cc./In oil: Vial (5 mg., 10 mg. or 25 mg./cc.) 10 cc.
Use: I.M., corpus luteum hormone therapy.

GESTEST. (Squibb) Norethindrone acetate 2.5 mg., ethinyl estradiol 0.05 mg./Tab. Box 16s.
Use: Pregnancy diagnosis.

GEVITONE. (Crestmed) Pentylentetrozol 50 mg., methyltestosterone 1 mg., ethinyl estradiol 2 mcg., B-12 1 mcg., nicotinic acid 50 mg., ferrous sulfate 50 mg., hesperidin 50 mg., B-1 1 mg., B-2 1 mg., B-6 0.5 mg., C 50 mg./Tab. Bot. 100s.
Use: Anabolic and cerebral stimulant.

GEVRABON. (Lederle) Thiamine HCl 5 mg., riboflavin 2.5 mg., Vit. B-12 1 mcg., niacinamide 50 mg., Vit. B-6 1 mg., pantothenic acid 10 mg., choline 100 mg., inositol 100 mg., Cal., Phos., Iodine, Pot., Mag., zinc, Mang., Iron/30 cc. Bot. 16 fl. oz., 1 gal.
Use: Vitamin & mineral deficiencies.

GEVRAL. (Lederle) Vit. A 5000 u., D 500 u., B-12 1/15 oral u., B-1 5 mg., B-2 5 mg., niacinamide 15 mg., folic acid 1 mg., Vit. B-6 0.5 mg., Col. pantothenate 5 mg., choline bitartrate 30 mg., inositol 50 mg., Vit. C 50 mg., Vit. E 10 u., rutie 25 mg., Iron 10 mg., w/Iodine, lysine, Cal., Phos., barce, copper, fluorine, Mang. Mag., Pot. & zinc/Cap. Bot. 30s, 100s, 250s, 1000s.
Use: (1 daily) Vitamin & mineral supplement.

GEVRAI PROTEIN. (Lederle) Vit. A 2500 u., D 250 u., thiamine 2.5 mg., riboflavin 2.5 mg., niacinamide 7.5 mg., pyridoxine HCl 0.25 mg., Cal. pantothenate 2.5 mg., folic acid 0.5 mg., Vit. B-12 1/15 oral u., choline dihydrogen citrate 500 mg., inositol 250 mg., ascorbic acid 25 mg., rutin 12.5 mg., Vit. E 5 I.U., lysine 170 Gm., Cal. 414 mg., Phos. 60.9 mg., Cal. caseinate 21.0 Gm., ferrous fumarate 15 mg., fluorine 0.05 mg., copper 0.5 mg., iodine 0.1 mg., Pot. 2.5 mg., Mang. 0.5 mg., zinc 0.25 mg., Mag. 0.5 mg., barce 0.05 mg., Carb. 7.0 Gm., calories 105.3, total protein 60.0%.

Sod. 1.8 mg., tol 2.0%/30 Gm. Jar 1/2 lb., Can 5 lb.
Use: (2 tbsp. daily) Dietary supplement.

GEVRAL-T. (Lederle) Vit. A 25,000 u., D 1000 u., B-12 1/3 oral u., B-1 10 mg., B-2 10 mg., B-6 2 mg., E 5 u., K 2 mg., C 150 mg., cot. pantothenate 5 mg., niacinamide 100 mg., folic acid 1 mg., choline bit., inositol, lysine, rutie, and 12 micerals/Cap. Bot. 100s.
Use: Severe vitamin deficiencies

GEVBESTIN. (Lederle) Ethinyl estradiol, 0.01 mg.; methyl testosterone, 2.5 mg.; d-Amphetamine sulfate, 2.5 mg. plus a strengthened Geval formula/Tab. Bot. 100s.
Use: Provide essential vitamins and minerals as well as estrogen-androgen therapy.

GEVRINE. (Lederle) Ethinyl estradiol 0.01 mg., methyl testosterone 2.5 mg., Vit. A 5000 u., D 500 u., B-12 1/15 oral u., B-1 5 mg., B-2 5 mg., niacinamide 15 mg., Vit. B-6 0.5 mg., folic acid 1 mg., Col. pantothenate 5 mg., choline bitartrate 50 mg., inositol 50 mg., Vit. C 50 mg., E 10 I.U., rutin 25 mg., w/Iron, Iodine, Cal., Phos., fluorine, copper, Pot., Mang., zinc, Mag., boron/Cap. Bot. 100s, 1000s.
Use: Orally (1 daily) estrogen-androgen therapy w/nutritional supplement.

GEXANE. N.N.D. 1962 (Strosenburgh) Gamma benzene hexachloride. Liq. 1%. Bot. 2 oz., 16 oz., Oint. 1%, Tube 1/2 oz., Jar 1 lb.
Use: Topically, scabicide & pediculicide.

GG-TEST KIT. (Hyload) Kit, 40 tests.
Use: Estimating gamma globulin levels.

G.H.P. CARBAMIDE. (Internat. Pharm. Corp.) Urea peroxide 4%, B-hydroxyquinoline 0.1%, glycerin anhyd. qs Bot. 1 aed 4 ac
Use: Topical antiseptic

G.H.P. OTIC. (International Pharmaceutical) New name Oebrix. Glycerite of hydrogen peroxide carbamide w/tetracaine HCl. Liq. Bot. w/dropper 15 cc
Use: Bactericide, fungicide, and analgesic in the treatment of ear infections

GIASOL SYRUP. (Utter) Pot. gualacol sulfonate 8 gr., amm. chlor. 8 gr., sod. citrate 20 gr., tartar. emetic 1/12 gr., cascara fluid 5 min., chloroform 1 min./R oz. 1 pt & 1 gal.
Also with Codeine 1 gr./R oz.
Use: Cough.

GISPASIM. (Amfre-Grant) Phenobarbital 14 mg., magnesium hydroxymagnacetate 250 mg., hydroxyamine sulf. 0.052 mg., hyacinine HBr 0.003 mg., atropine sulf. 0.01 mg./Tab. Bot. 100s
Use: Anspasmodic with sedative

GITALIGIN. N.N.D. 1960. (White) Gitalin

- (amorphous) a cardio active glycoside derived from *digitalis purpurea* Tob: 0.5 mg of gitalin (amorphous) Bot 30s, 100s, 1000s, Drops: (0.05 mg/2 drops) Bot w/cali broted dropper 30 cc Amp: (0.5 mg/cc) 5 cc Box 3s, 12s
 Use: Orally, digitalis therapy
- ***GITALIN (AMORPHOUS)**, USP NND 1960 Tabs, USP See Gitaligin, Tab (White)
- W/Digitoxin See Myapurin, Tab (Patch)
- G. I. TRAN.** (Kendall) d Amphetamine sulfate 5 mg, amobarbital 32 mg, belladonna ext 16 mg/Tab Bot 100s, 500s.
 Use: (1 q 3-4 h) Gastrointestinal disorders
- GLANOBOLIN.** See: Estrone (Various Mfr)
- GLARB** (Cale Chem Co) Theobromine 3½ gr, Cal gluconate 1½ gr, phenobarbital ¼ gr Tab Bot 100s
 Use: (1 1 1 d) Hypertension
- GLARUBIN**, NND 1962 (Mossengill) Glauconin (glycoside from *Simarouba glouca*) 50 mg/Tab Bot 40s
 Use: Intestinal amebiasis
- GLAUBER'S SALT.** See Sodium Sulfate (Various Mfr)
- ***GLAUCARUBIN**, NND 1962 Crystalline glycoside from *Simarouba glouca* aubl *Simarouba* See Glarubin, Tab (Mossengill)
- GLAUCON Ophth Soln** (Houg Drug) d epinephrine HCl, 2% benzalkonium chloride 0.02%; sodium chloride 0.2%; sodium metabisulfite, 0.3% (as preservative) Bot 10 ml
 Use: Glaucoma, miatic therapy
- GLICOMP.** (Eisrol) Sol of Sod glycerophosphate 0.2 Gm, iron cacodylate 0.02 Gm, strychnine sulfate 0.001 Gm Amp 1 cc, Box 6s, 12s, 25s, 100s
 Use: Hematinic
- GLIDDA-CAL.** (Glidden) Vit A 5000 u, D 400 u, C 70 mg, B 1 2 mg, B 2 2 mg, K 0.5 mg, B 6 4 mg, niacinamide 10 mg, ferrous fumarate 90 mg, iron 30 mg, folic acid 0.5 mg, B 12 3 mcg, copper 0.3 mg, cabalt sulfate 0.66 mg, iodine 0.02 mg, magnesium 0.3 mg, zinc 0.17 mg, potassium 1.67 mg malybdenum 0.5 mg, potassium 1 mg, calcium carbonate 600 mg / chewable tab Bot 100s
 Use: Dietary supplement
- GLIDDATINIC.** (Glidden) Ferrous fumarate 90 mg Vit B 12 7.5 mcg, folic acid 0.5 mg, Vit C 50 mg / chewable Tab Bot 50s
 Use: Anemias
- GLIM.** See: Gardinol Type Detergente (Various Mfr)
- * Generic name recognized by NF, USP or A.M.A.
- GLISOTAL.** (Metropolitan Labs) Iron cacodylate ½ gr, Sod glycerophos 1½ gr, strychnine glycerophos 1/65 gr, Sod cacodylate ¼ gr/cc Vial 1 cc, 2 cc, Box 10s, 100s
 Use: Hematinic
- GLOBALINE**, NNR 1954 (Maltbie Labs) Tetra glycine hydraperladide contains 39.5% to 42.6% active iodine eq to 8 mg of iodine/Tab 20 mg Bot 50s
 Use: Purification of drinking water, 1 tab per pl for 3 minutes
- ***GLOBIN INSULIN W/ZINC**, USP, NND 1962 Glabin zinc insulin, insulin W/globin & zinc chloride added
 Burroughs Wellcome—Vial (40 u/cc) 10 cc, (80 u/cc) 10 cc
 Squibb—Vial (40 u/cc) 10 cc, (80 u/cc) 10 cc
 Use: S.C., insulin therapy
- See: Insulin Globin Zinc, injection, USP
- GLOBOTRIN.** (Smith, Miller & Patch) Ferrous lactate 195 mg, Vit B 12 w/intinsic fac for conc 0.5 u, thiamine HCl 2.5 mg, ascorbic acid 50 mg, betaine HCl 60 mg, methylcellulose 32.5 mg/Tab Bot 60s, 500s
 Use: (2 1 1 d, a c) Iron deficiency & pernicious anemias
- GLOBULIN, ANTINEMOPNIC,** USP XY Antihemophilic Globulin, Human
- GLOBULIN, GAMMA.**
 See Gamma Globulin (Various Mfr)
 Poliomyelitis Immune Globulin, Human Vial (Merck Sharp & Dohme)
- ***GLOBULIN, IMMUNE SERUM (HUMAN)**, USP NND 1962
 Celler—Vial 2 & 10 cc
 Lederle—Vial 2 cc
 Pitman Moore—2 & 10 cc
 Merck Sharp & Dohme—2 & 10 cc
 Use: IM, measles prophylactic & polio
- GLOBULIN, POLIOMYELITIS IMMUNE.** (Human) See Poliomyelitis Immune Globulin (Human), NND 1961
- GLOVARIAN.** (Schiefelin) Estrogenic substance natural 500 u & ovary ext 2 gr / Pill Also w/thyroid, Bot 36s, 100s
 Use: Orally, estrogen therapy
- GLOX.** (Cale) Phenolphthalein ½ gr, aloin ½ gr, ax bile ext ½ gr, cascara sagrada ext ½ gr/Tab Bot 100s
 Use: Laxative
- ***GLUCAGON**, NND 1962 (Lilly) Glucagon HCl 1 mg or 10 mg w/diluent, soln contains glycerin 1.6% w/phenol 0.2% as a preservative Vials
 Use: Hypoglycemic shock
- GLUCALBROM** (Kirk) Cal gluconate 0.5

Gm., strontium bromide 0.5 Gm., dextrose 1 Gm. Vial 10 cc., Box 6s, 25s, 100s.

Use: Sedative w/calcium therapy.

d-GLUCITOL [*d-Sorbitol*]/*Nomatropine* methylbromide. See: ProBitogol, liq. (Purdue Frederick)

GLUCO-CALCIUM. (Lilly) Col. salts of sugar acids. Vial 30 cc., Box 25s

Use: Calcium therapy.

GLUCO-CORTICOIDS. See: Cortical Hormone Products.

GLUCODIGIN. (Cole) Digitoxin 0.2 mg./Tab. Bot. 100s, 500s, 1000s.

Use: Oral digitalization.

GLUCODOX. (Kirkman) Calcium gluconate 5 gr., ferrous gluconate 2.5 gr., vit. B-6 2.5 mg., D 333 u., cobalt $\frac{1}{2}$ mg./cop. Bot. 100s, 1000s.

Use: (1-3 daily) Cobalt therapy.

GLUCO-PEDRIN. (Parke, Davis) Isotonic sol. of ephedrine (as the lactate) 1%, chlorzane 0.5%, w/menthol in aq. dextrose base. Drapper-Cop 1 oz. & 1 pt. Plastic spray bot. 15 cc.

W/Phemeral (benzethonium chloride 1:5000) & w/sulfathiazole & gluco-thricil (Tyrothricin 1:5000) Bot. 1 oz.

Use: Relief of congested mucous membrane w/antiseptic.

GLUCO-FERRUM. (Yonpelt-Brown) Ferrous gluconate 0.3 Gm., Vit. B-1 0.6 mg./Tab. Ferrous gluconate 0.18 Gm., Vit. B-1 0.6 mg./fl. oz. Elixir, Bot. 8 fl. oz., Tab., Bot. 100s, 500s, 1000s.

Use: Hemofinic.

GLUCO-FERRUM W/C. (Yonpelt & Brown) Ferrous gluconate 4.6 gr., B-1 1 mg., C 100 mg./Cop. Bot. 100s

Use: Hemofinic.

GLUCO-MYCIN. (Commercial Solvents) Glucuronolactone 400 mg., streptomycin sulfate 100 mg./Tab. Bot. 50s.

Use: Orally, streptomycin therapy.

GLUCONIC ACID SALTS.

See: Calcium Gluconate

Ferrous Gluconate

Magnesium Gluconate

Potassium Gluconate

GLUCOPHYLLINE, N.N.D. 1962. (Abbott) Theophylline methylglucamine. Amp. (0.366 Gm.) 10 cc. Box 6s, 25s; Tab. E.C. (0.15 Gm.) Bot. 100s, 1000s; (0.15 Gm. & 0.3 Gm.) Bot. 100s, 1000s. Supp. (0.5 Gm.) Box 12s, 100s.

Use: Diuretic & antispasmodic.

W/Ephedrine & Neonal Cop. See: Phylodrine Capsules (Abbott).

GLUCOPHYLLINE & NEMBUTAL, TABLETS.

*Generic name recognized by N.F., U.S.P. or A.M.A.

(Abbott) Glucophylline 150 mg., nembutal Sod. 15 mg./Tab. Bot. 100s, 1000s.

GLUCOPHYLLINE W/NEMBUTAL SUPPOSITORIES. (Abbott) Theophylline 0.25 Gm., methyl glucamine 0.25 Gm. & nembutal 0.2 Gm./Supp. Box 12s.

Use: Diuretic & antispasmodic.

GLUCOPHYLLINE & PHENOBARBITAL TABLETS. (Abbott) Glucophylline 0.15 Gm., phenobarbital 30 mg./Tab. Bot. 100s, 1000s.

GLUCORON. (Lemmon) Ferrous gluconate 250 mg., Vit. B-1 1 mg., B-2 0.5 mg., manganese 2 mg., copper 1 mg., cobalt 0.1 mg./Tab. Bot. 100s, 1000s. Also available with Vit. B-12 3 mcg./Tab. of above formula

Use: Anemias.

GLUCORONIDASE, BETA. See: Ketodase, liq. (Warner-Chilcott).

GLUCOSAMINE. W/Tetracycline HCl, nystalin See: Terrastatin (Pfizer)

W/Tetracycline. See: Tetracyc, Cap. (Pfizer)

W/Oxytetracycline. See: Terramycin (Pfizer)

W/Oxytetracycline, buclizine HCl, phenacetin, salicylamide, caffeine. See: Terracyclin, Caps (Pfizer)

W/Oxytetracycline, nystalin. See: Terrastatin, Caps (Pfizer)

W/Triacetyl Oleandomycin. See: Signemycin (Pfizer)

***GLUCOSE**, liq., U.S.P., N.N.R. 1943. (Various Mfr.) Cerebrate, Dextrose. See: Cortose (Winthrop Labs.)

Use: 5-50% sol. as nutrient for acute hepatitis and dehydrated; to increase blood volume.

GLUCOSE TEST. See: Cambistix (Ames)

GLUCO-SULPADIAXINE. (Danley-Evans) Sulfadiazine 6.48 Gm., Sod. lactate & glucose 21.6 Gm./100 cc. of susp. Bot. 1 pt.

Use: Sulfonamide susceptible infections.

GLUCO-SULPAMERAZINE. (Danley-Evans) Sulfamerazine 6.48 Gm., Sod. lactate & glucose 21.6 Gm./100 cc. of susp. Bot. 1 pt.

Use: Sulfonamide susceptible infections.

GLUCO-SULPANILAMIDE. (Danley-Evans) Sulfanilamide 0.065 Gm. 1 gr./cc., Sod. lactate 13.8%, Pot. citrate 4.9%. Bot. 6 fl. oz.

Use: Sulfanilamide susceptible infections.

GLUCOSULPAS. (Danley-Evans) Sulfadiazine 3.7 Gm., sulfathiazole 3.7 Gm., sulfamerazine 2.6 Gm. w/Sod. lactate/100 cc. of susp. Bot. 1 pt.

Use: Triple sulfonamide for sulfonamide susceptible infections

GLUCO-SULPATHAZOLE. (Danley-Evans) Not listed. Sulfathiazole 6.48 Gm., sod.

- lactate 21.6 Gm, pot citrate & glucose 6.4R Gm/100 cc Bot 1 pt, 1 gal
Use Sulfanamide susceptible infections
- * **GLUCOSULFONE SODIUM**, NND 1962
p'-Diaminodiphenylsulfane N,N'-di(dextrose sodium sulfonate) See Promin, Amp (Parke, Davis)
- GLUCO-THRICIL** (Parke, Davis) Tyrothalcin 15000, ephedrine lactate 1% in isotonic dextrose sol Bot/dropper, 1 oz, 1 pt
Use: Typically, vasoconstrictor & antibiotic nasal solution
- * **GLUCUROLACTONE**, NND 1962 Gamma lactone of glucuronic acid See: Glucurone, Tab (Commercial Solvents)
- W/Dipyrrone & salicylamide See Glu Salgin Cap (Warren Teed)
- W/Salicylamide See Glu Sol, Tab (Warren Teed)
- W/Streptomycin sulfate See: Gluco Mycm, Tab (Commercial Solvents)
- GLUCURONE**, NND 1962 (Reed & Carnrick) Glucuronolactone Tab 0.5 Gm Bot 50s
Use Orally, rheumatoid & other forms of arthritis
- GLUCURONIDASE**, RETA See Ketodase (Warner Chilcott)
- GLUFERATE**, (Wyeth) Ferrous sulfate 2 gr, glutamic acid HCl 200 mg/Tab Rot 100s, 1000s
Use: (2 t i d) Iron therapy for nutritional anemias
- GLUKOR**, (Research Supplies) Chorionic gonadotropin 200 IU, Vit B1 25 mg, glutamic acid 52.5 ppm, procaine HCl 1%, chlorbutanol 0.5%/cc Vial 10 cc, 25 cc
Use: IM (1-2 cc weekly); androgen deficiency, uterine disorders
- GLU-SAL** (Warren Teed) Glucuronalactone 0.2 Gm, salicylamide 0.4 Gm/Tab Rot 100s, 500s
Use (2-3 q i d) Analgesic & antiarthritic
- GLU-SALGIN**, (Warren Teed) Glucuronalactone 2½ gr, dipyrrone 1½ gr, salicylamide 5 gr/Cap Bot 100s
Use: (1-2 t i d) Analgesic
- GLUSATE**, (United Lab) Glutamic acid Tab 0.5 Gm, Bot. 100s, 1000s.
Use: Orally, antiepileptic in reduction of petit mal attacks
- GLUSIDE**, See: Saccharin.
- GLUTALATE**, (Dumas Wilson) Pot p-Aminobenzoate 225 mg, salicylamide 225 mg, pot salicylate 165 mg, dicalcium glutamate 75 mg, Vit C 20 mg/Cap Bot 100s
Use: Sodium free treatment of arthritis
- * Generic name recognized by N F U S P or A M A
- GLUTALATE W/PREDNISON**, (Dumas Wilson) Prednisone 2 mg, ascorbic acid 20 mg, dicalcium glutamate 75 mg, potassium salicylate 165 mg, salicylamide 225 mg, potassium p-aminobenzoate 225 mg./Cap Bot 100s, 1000s
Use Arthritis
- GLUTAMATE SODIUM**, NND 1961
See: Glutavite, Pow. (Gray Pharm Co)
Glutavene (Gray Pharm Co)
- GLUTAMATE SODIUM 20% INJECTION** (Bruce) Monosodium glutamate 200 mg/cc Vial 60 cc
Use: Hepatitis
- GLUTAMATE THERAPY**.
See: Glutavene (Gray Pharm Co)
Glutavene K (Gray Pharm Co)
- GLUTAMIC ACID**. See: Glusate, Tab (United Labs)
- W/Methyltestosterone, thyroid ext, Vit B1 See Android, Tab (Brown Pharm)
- W/Reserpine & estrogenic substance. See Glutrone, Vial, (Jeffery)
- W/Sodium citrate See: Glutamcol, Pow (Walker)
Use Anticonvulsant
- * **GLUTAMIC ACID HYDROCHLORIDE**, NF Cap, NF Acidogen, aciglutamin, glutasin
Use: The hydrochloride for achlorhydria & hypochlorhydria
See Acidoride, Cap (Abbott)
Acidothyn, Cap (Plint, Eaton)
Acidulin, Pulvules (Lilly)
Aclor, Cap (Cole Chem Co)
Gastaloric, Cap, & Tab (Warren Teed)
Glutan HCl, Cap (Lederle)
Hydronic, Cap (Upjohn)
Munamic, Cap (Pitman Moore)
- W/Cellulose, Pepsin, Pancreatin NF, Ox Bile Extract See: Kanulose, Tab (Smith Dorsey)
- W/Mephenesin
See Mephate, Cap (Robins)
Tolamic Cap (Kremers Urban)
- W/Mephenesin, phenobarbital & 1 hyoscyamine HBr. See: Kused, Cap (Kremers Urban)
- W/Pepsin See: Glutasin, Cap (McNeil)
- W/Vitamins See: Flanithin, Cap (Table-rock)
- W/Vitamin B preparation See: Actamin Cap (Massengill)
- W/Vitamin B1, riboflavin & niacinamide See: Acid V, Cap (Merck Sharp & Dohme)
- W/Vit B12 & folic acid See: Fatbexil Tab (Chicago Pharmacal)
- W/Vit B12, Intrinsic factor, iron oxide saccharated & folic acid See: Antemia Tab (Chicago)
- GLUTAMIC ACID SALTS**

See: Ammonium Glutamate

Calcium Glutamate (Various Mfr.)

Gustamate (Arlington)

GLUTAMICOL. (Walker) Glutamic acid 3 Gm. w/Sad. citrate/dr. Pow. 10 ox.

Use: Orally, antiepileptic in reduction of petit mal attacks.

GLUTAN HCl. (Lederle) Glutamic acid HCl 5 gr., equiv. In hydrochloric acid content to approximately 10 min. of diluted hydrochloric acid U.S.P./Cap. Bot. 100s, 500s, 1000s.

Use: Orally, achlorhydria & hypochlorhydria.

GLUTATHIONE, SODIUM. See: Triplide, Vial (Schwarz).

GLUTAVENE, N N D. 1961. (Crookes Barnes) Sodium L-glutamate eq. to 25 Gm. of L-glutamic acid/100 ml. Vials 100 ml.

Use: Reduction of blood ammonia level associated with hepatic disease

GLUTAVENE-K. (Crookes-Barnes) Potassium and sod. L-glutamate eq. to 25 Gm. of L-glutamic acid/100 ml. Vials 100 ml.

Use: Reduction of blood ammonia level associated with hepatic disease

L-GLUTAVITE. (Crookes-Barnes) Levoglutamate monosod. 3.48 Gm., niacin 25 mg., Vit. B-1 0.6 mg., B-2 0.8 mg., B-6 0.7 mg., B-12 2.0 mcg., ferrous sulfate 11 mg./packet or tsp. Box 40s. Bottle (50 level teaspoonfuls) Caps. 100s.

Use: Cerebral tonic in geriatric therapy.

GLUTAZYME CAPSULES. (Purdue Frederick) Monosodium L-Glutamate 500 mg., niacin 15 mg., Vit. B-1 1 mg., B-2 1 mg., C 20 mg., cal. pantothenate 2 mg., Vit. B-6 0.34 mg., B-12 1 mcg., A 2000 u., ferrous sulf. 12 mg., cal. carbonate 100 mg./Cap. Bot. 100s.

Use: Nutritional supplement.

GLUTAZYME POWDER. (Purdue Frederick) Monosodium L-Glutamate 3.812 Gm., niacin 16.4 mg., Vit. B-1 1.08 mg., B-12 1.08 mcg., C 21.6 mg., B-2 1.08 mg., B-6 0.36 mg., cal. pantothenate 2.16 mg., Vit. A 2180 u., ferrous sulf. 13.08 mg., cal. carbonate 108.8 mg./Tsp.

Use: Nutritional supplement.

GLUTEST. (Research Supplies) Testosterone 10 mg., thiamine HCl 25 mg., glutamic acid 52.5 ppm, procaine HCl 1%, benzyl alc 2%/cc. Vial 10 cc., 25 cc.

Use: I.M. (1 cc. weekly); female climacteric.

* **GLUTETHIMIDE, N F., N N D.** 1962. Alpha-ethyl alpha phenyl-glutarimide. Tabs., N.F. See: Douden, Tab. (Ciba)

* Generic name recognized by N.F., U.S.P. or A.M.A.

GLURONE. (Jeffery) Reserpine 0.1 mg., estrogenic substance 2 mg., glutamic acid 200 mg./cc. Vial 10 & 30 cc.

Use: Menopausal disturbances.

GLYAL WAFERS. (Fay) Aluminum hydroxide-magnesium carbonate 400 mg., glycine 100 mg./chewable wafer/wafer. Bot. 100s.

Use: Antacid.

GLYBROM. (Polch) Pyrabrom 50 mg./Tab. Bot. 100s, 500s, 1000s.

Use: Orally, premenstrual tension & molar sickness.

GEYCAMINE. (Minnesota Pharm. Labs., Inc.) Nymagnate 0.5 Gms./Tab Bot. 100s, 500s. Liq. 0.5 Gm./tsp. Bot. 1 pt.

Use: Peptic ulcers, hyperacidity.

GLYCAMINE-AM. (Barre) Hydroxy aluminum magnesium aminoacetate. Tab. 7 1/2 gr., Bot. 100s, 500s, 1000s.

Use: 1-3 cc. & h.s.) Antacid.

GEYCANA. (Schleffelin) Strychnine glycerophosphate 1/16 gr., Sod. glycerophosphate 8 gr., Cal. glycerophosphate 2 gr./fl. oz. Elixir, Bot. 8 fl. oz., 1 gal.

Use: Tonic.

* **GLYCARNINE IRON.** See: Ferronard (Nordmark)

GLYCARSAN. (Endo) Iron cacodylate 20 mg., sod. cacodylate 16 mg., strychnine sulf. 0.8 mg., sod. glycerophosphate 0.1 Gm./cc. Amp. 1 cc. Box 12s & 100s

Use: Anemia and neuroasthenia.

GLYCERITE OF HYDROGEN PEROXIDE. (Int'l Pharm. Corp.) Hydrogen peroxide 2.5%, 8-hydroxyquinoline 0.1%, glycerol q.s. ad 120 cc. Bot. 4 oz.

Use: Antiseptic.

GLYCIRITE OF HYDROGEN PEROXIDE WITH CARBAMIDE. (Internat. Pharm. Corp.) Hydrogen peroxide 1.5%, ureo (Carbamide) 2.5%, 8-hydroxyquinoline 0.1%, anhy. glycerol. Bot. w/dropper 30 cc.

Use: Antiseptic.

GEYCERYL GUAIACOLATE. Methphenoxy-dol, 3 [o-methoxyphenoxy]propanediol-1,2. Guayanesin Guayanesin. Glyceril Guaiacol Ether.

See: Dilyn (Elder)

Glycotuss (Vale)

W/N-Acetyl-p-aminaphenal. See: Rystal, Syr. (Grail)

W/N-Acetyl-p-aminaphenal, phenylpropanolamine, noscapine. See: Caryz, Tab (Moran)

W/Brompheniramine maleate, phenylephrine HCl, phenylpropanolamine HCl. See: Dimethane Expectorant (Robins)

W/Chlorpheniramine maleate, Phenylpropanolamine HCl, Phenylephrine HCl. See: Parathane, Syr. (Grail)

- W/Dextromethorphan HBr See: Glycotree dM (Yale)
- W/Dihydrocodeinone, Chlor Trimeton, Sod salicylate, Sod citrate, caffeine See: Cordia, Syr. (Schering).
- W/Dihydrocodeinone bitartrate chlorpheniramine maleate, phenindamine tartrate, phenylpropranolamine HCl See: Nolamine Expectorant Syr (G W Carrick)
- W/Dihydrocodeinone bitartrate, methopirylene fumarate, Sod citrate See: Neo Codonyl, Syr (C. B Kendall)
- W/Dilaudid HCl See: Dilaudid Cough Syrup (Knoll)
- W/Ephedrine sulfate, theophylline, thendiamine HCl, phenobarbital See: Bronkafab (Brees)
- W/Ethaverine See: Thevasin (Lincoln)
- W/d Isopropylamine HCl, chlorpheniramine maleate, dihydrocodeinone bitartrate, sodium citrate, citric acid, chloroform See: Hozanone No 2, Tab (Haskell)
- W/d Methorphan HBr, chlorpheniramine maleate See: Golden Cough Syrup, liq (Stoner)
- W/Noscapine, citric acid See: Noscatene (Burlington)
- W/Noscapinealkaloid phenylpropranolamine, acetaminophen, chloroform, AR See: Coryx Suspension, liq (Morion)
- W/Brompheniramine maleate phenylephrine, phenylpropranolamine See: Dimetone Expectorant, liq (Robins)
- W/Phenylephrine HCl, phenylpropranolamine HCl codeine phos, brompheniramine maleate See: Dimetone Expectorant DC (Robins)
- W/Phenylpropranolamine HCl, isopropylterrenal HCl See: Broastodyl Pediatob 17. (Columbus)
- W/Phenylpropranolamine HCl, Pheniramine maleate See: Darathon, Sol (Graft)
- W/Phenylpropranolamine HCl, phenyltolazamine d hydrogen citrate, pyriminamine maleate, sec'rylamide See: Respal, Syr (Tilden)
- W/Phenyltolazamine d hydrogen citrate See: Guafalussin, Syr (White).
- W/Phenyltolazamine d hydrogen citrate pyriminamine maleate meth-dia mar sulfonamide and ephedrine sulfate See: Sulfasin, Susp (Tilden)
- W/Pyronal maleate & phenylephrine HCl See: Guafalit Tab (Columbus).
- W/Pyronal maleate phenylephrine HCl, sec'rylamide acetan acetos See: Guafalussin, liq (Columbus)
- W/Terbutaline HCl, pseudoephedrine HCl, codeine phosphate See: Asted C. Syr (Burlington-Walton)
- OLTCBYL GUAIACOLATE CARBAMATE**
* Generic name recognized by I.F.U.S.P. or A.M.A.
- Methocarbamol. See: Robaxin, Tab (Eobin)
- GLYCERYL GUAIACOL ETHER**, See: Glyceryl Gualacolate
- * **GLYCERYL MONOSTEARATE**, NF (Various Mfr) Monostearin
- GLYCERYL TRIACETATE**, NND 1962 NF X, ADR 1955 (Various Mfr) Triacetin
See: Enzocin, Oint (Ayerst)
Fungacetin, Oint, liq & Pow (G F Harvey)
Vanay, Cream (Ayerst)
- GLYCERYL TRINITRATE OINTMENT**, Sem Nitrol, Oint. (Kremers Urban)
- * **GLYCERYL TRINITRATE TABLETS**, USP
See: Nitraglyn (Key Corp)
Nitrol. (Kremers Urban)
- W/Pentaerythritol tetranitrate See: Dilcora Tab (Winthrop Labs)
- W/Phenobarbital Sod, triple bromides hyoscyamus & Pot nitrate, See: Nitabrom, Tab (Yilo)
- GLYCINE**, NND 1962 (Various Mfr) See: Aminoacetic Acid, NF
- W/Ammonium gluconate, ammonium salicylate, caffeine, terpin hydrate, benzalkonium chloride See: Mortussin Special, liq (M. R Thompson)
- W/Calcium Carbonate, NNR 1934 See: Titorac, liq, Tab & Pow, (Schenley)
- W/Nicotinic acid. See: Nicatron Ellair (Maug Drug)
- W/Sod salicylate, chlorphenylpyridamine maleate See: Corlinh liq (Schering)
- GLYCINE, ALUMINUM SALT**, See: Dihydroxyaluminum Aminoacetate (Various Mfr)
- OLYCINE HYDROCHLORIDE**, (Various Mfr)
See: Glyco-HCl, Cap (Burnham)
Use: Recommended for ophthalmia
- OLYCINE HYDRIODIDE**, See: Glyco-HL Cap. (Burnham)
- OLYCINE SOLUTION**, (Abbott) Sol 1000 diluted to a volume of 10 L w/sterile distilled water yields a 1.5% sol of glycine.
Use: Urologic Irrigation
- OLTCOBEN**, (Tilden) Codeine sulfate 1 gram, antimony & Pot tartrate 1/25 gr, Pot guaiacolsulfonate 8 gr, ammonium chloride 8 gr, chloroform 1/8 ml / R as Sol. 4 oz., 1 pt., 1 gal
Use: (1 tsp. q 3 h) Expectorant, bronchial sedative
- * **OLYCOBIARSOL**, USP, NND 1962, Tab
Bismuth Glycylarsosulfate, NPLB 1934
Chemo Pura, Pow for Mfr
See: Ambebian, Tab (Cansol Almond Corp)
Brazlin, Cream, Veg Supp (Broom)
Miltab, Tab & Veg Supp (Winthrop Labs)

W/Chloroquine diphosphate. See: Milbisa
w/Aralen Diphosphate, Tab. (Winthrop
Labs.)

Uses: (250-500 mg. i.i.d.) Amebiasis. Tricho-
monas vaginalis, Monilia albicans.

GLYCOCOLL, N.N.R. 1941. Glycine.

See: Aminoacetic Acid (Various Mfrs.)

Glycolixir (Squibb).

W/l lysine. See: Lixacol Elixir (Lanett)

OLYCOCOLL HYDROCHLORIDE. See: Glyco-
HCl, Cap. (Burnham).

OLYCOCOLL HYDRIODIDE. See: Glyco-HI
(Burnham)

GLYCOCYAMINE, Guanidoacetic acid.

W/Betaine. See: Betasyamine. (Internat.
Minerals & Chem.)

GLYCO-HCl. (Burnham) Diglycine hydrochlo-
ride. 1 Cap. equivalent to 10 min. of diluted

Uses: (1-2 i.i.d.) Achylarhydria & hypochlor-
HCl. Bot. 50s, 100s, 500s.

GLYCO-HI. (Burnham) Glycocol hydriodic
acid. Cap., Bot. 50s, 100s, 500s.

Uses: Iodine therapy.

GLYCOLIXIR. (Squibb) Elixir of glycololl
1.82 Gm. (28 gr.) glycololl/½ fl. oz. (one
tbsp.) Alcohol 12%. Bot. 1 pt., 1 gal.

Uses: Promote the development of protein.

***OLYCOHENYLATED BROMIDE**.

See: Mepenzolate Methylbromide.

***GLYCOPYRROLATE**, 1-Methyl-3-pyrrolidyl
α-phenylcyclopentaneglycolate methobro-
mide. See: Robinul (Robins)

GLYCORAL. (Schleffelin) Adrenal cortex
gland ext. 2 gr./Pill. Bot. 50s, 225s.

Uses: Orally, adrenal cortex therapy.

GLYCOTAURO, N.N.R. 1949. (Hynson, Wes-
cott & Dunning) Ox bile salts 1.2 gr./E.C.
Tab. or Cap. Tab. Bot. 72s, 500s, 1000s;

Cap. Bot. 36s, 500s; w/cascara ½ gr. &
phenolphthalein ½ gr., E.C. Tab., Bot. 50s,
500s, 1000s

Uses: (2-4 daily) Choleric & chologogue.

GLYCOTUSS. (Vale) Glyceryl guaiacolate
100 mg./Tab or 5 cc. Syr. pts Tabs. 100s.

Uses: Expectarant.

GLYCOTUSS-DM. (Vale) Glyceryl guaiacolate
100 mg., dexamethorphan HBr 10 mg./
Tab or 5 cc. Tab 100s. Syr. pts.

Uses: Expectarant/antitussive.

GLYCYRRHIZA, U.S.P. Elixir, N.F. Ext.,
N.F. Ext. Pure, U.S.P. Flex., U.S.P. Syr.,
U.S.P. (Merck Sharp & Dahme) Licarice root.

GLYESTRIN. (First Texas) Estragline con-
jugates (equine) 125 mg./tab. Bot. 100s,
500s.

Uses: Estrogen insufficiency.

GLYKERON. (M. H. Smith) Codeine phos-
phate 32 mg., Hyoscyamus flex. 0.25 cc.,

* Generic name recognized by N.F., U.S.P. or A.M.A.

ammonium chloride 650 mg., Sod. citrate
650 mg., citric acid 325 mg./30 cc. Bot.
4 oz., 1 pt., 1 gal.

Uses: (2-3 tsp. q. 3 h.) Expectarant, cough
depressant.

GLYLENE. (Flint-Eaton) Glylene 5 gr., brand
of theophylline-Sod. Glycinate equiv. to
theophylline 2½ gr. Tab. Bot. 100s, 500s,
1000s.

Uses: (1-2 q. 3-4 h.) Antispasmodic, par-
ticularly of bronchiales, coronary vasodila-
tor & myocardial stimulant.

OLYLENE WITH PHENOBARBITAL. (Flint-
Eaton) Glylene 5 gr., phenobarbital ¼
gr./Tab. Bot. 100s, 500s, 1000s

Uses: (1-2 q. 3-4 h.) Sedative & antispas-
modic.

GLYMOL. See: Petrolatum Liquid (Various
Mfrs.)

GLYNAZAN, N.N.D. 1961. (First Texas) The-
ophylline-Sod. glycinate. Elixir, Bot. 1 pt. &
1 gal., Pow., Bot. ¼ & 1 lb., Syr., 35 mg/
cc., Bot. 1 pt. & 1 gal., Tab., Bot. 2.5 & 5
Gm.

Uses: Cardiac therapy. Same as aminophyl-
line.

OLYNAZAN-EP. (First Texas) Theophylline
sodium glycinate, 5 gr., ephedrine sulfate,
¾ gr., phenobarbital, ¼ gr./Tab. Bot. 100s

Uses: Bronchial asthma

GLYOTOL. (U. S. Standard) Mephenesin.
Tab. 0.5 Gm. Bot. 100s, 1000s.

Uses: (2 i.i.d.) Muscle relaxant.

GLY-OXIDE. (International Pharmaceutical
Corp.) Carbamide peroxide 10% in flavored
anhydrous glycerol Plas. bot 2 oz. & ½ oz.

Uses: Oral infections.

GLYOXYLDIUREIDE. See: Allantoin (National
Drug).

GLYPECTUS. (Schleffelin) Chloroform, white
pine, cherry, squill, senega, ipecac, son-
geinaria, honey & eucalyptol. Bot. 1 pt.,
1 gal.

Uses: (1 tsp. q. 3 h.) Cough preparation.

***GLYPINAMIDE**. (Upjohn) 1-(p-chloro-
phenylsulfonyl)-3-(hexahydro-1H-azepin-1-
yl)urea.

Uses: Oral hypoglycemic agent.

GLYROL. (Smith, Miller & Patch) Ipecac,
ammonium chloride, cinnamic acid, cinn-
amic aldehyde, glycerin & aromatics. Syr.,
Bot. 4 R. oz., 1 pt., 1 gal

Uses: (1-2 tsp. p.r.n.) Expectarant.

GLYSENNID. (Sandax) Sennosides A & B 12
mg./Tab. Bot. 40s, 250s, 1000s.

Uses: Orally, laxative.

GLYSUFED. (Massengill) Ephedrine 1%.

- menthal, chlorobutanol isotonic lactose sol
Bot w/dropper 1/2 oz, 1 pt, 1 gal
Use Nasal vasoconstrictor
- GLY TABS** (Deseret) Amino acetic acid 100 mg calcium carbonate USP 300 mg, aluminum hydroxide, magnesium carbonate 200 mg/Tab Bot 100s
Use Antacid
- GLYTHEONATE, NND 1962** (Smith, Moller & Patch) Theophylline Sod glycinate Syr. 5 gr/5 cc Bot 1 pt, 1 gal; Tob 5 gr Bot 100s; Supp 12 gr Box 12s
Use Antispasmodic & vasodilator
- W/Phenobarbital (Patch) Glytheonate 5 gr, phenobarbital 1/4 gr/Tab Bot 100s, 500s
Use (1 2 11 d) Antispasmodic, vasodilator with sedative
- W/Phenobarbital & raphedrine (Patch) Glytheonate 5 gr, phenobarbital 1/4 gr, raphedrine HCl 3/4 gr/Tab Bot 100s, 500s
Use: (1 q 3 h, m p) Antispasmodic, vasodilator & sedative
- W/Phenobarbital & Reserpine (Patch) Glytheonate 325 mg, phenobarbital 10 mg, reserpine 0.1 mg dihydroxy aluminum aminoacetate 325 mg/tab Bot 100s
Use Orally, antispasmodic & tranquilizing agent
- W/Phenobarbital & rutin (Patch) Glytheonate 5 gr, phenobarbital 1/4 gr, rutin 20 mg/Tab Bot 100s, 500s
Use: (1 q 3 4 h, m p) Antispasmodic & vasodilator with vitamin P factor
- GLYTINIC**, (Boyle) Ferrous gluconate 135 gr, aminoacetic acid 13 Gm, Vit B 12 10 mcg, B 1 75 mg B 2 7.5 mg, B 6 225 mg, niacinamide 45 mg, panthenol 65 mg, liver fraction 1 H.F. 3 gr, cobalt 0.05 mg, & mong 5 mg/4 tab or 2 tsp Bot 100s & 1 pt
- GLYVITA**, (Massengill) Sod glycerophosphate 4 gr, Cal glycerophosphate 4 gr, Pot glycerophosphate 3/4 gr thiamine HCl 5 mg, riboflavin 2 mg, pyridoxine HCl DS 5 mg, pantothenic acid 1 mg, nicotinic acid 30 mg/A oz Bot 1 pt, 1 gal
Use: Tonic
- GOAT'S MILK**.
Cutter—Can 8 oz
Jackson—Pkg 14 oz
King—Cap, Bot 75s; Pow 7 oz
See: Dale Goat Milk (Cutter)
Use: Dietary supplement
- GOBROMOL**, (Endo) Gomenol 100 mg guaiacal 100 mg, eucalyptal 80 mg, lada farm 20 mg, camphar 50 mg in olive oil solution Amp 2 cc, Box 12s, 25s, 100s
Use: (I.M.) (Liquefying expectorant
- * Generic name recognized by NF USP or A.M.A.
- GOLD BROMIDE**. See: Gold Tribromide (Various Mir)
- GOLD (Au¹⁹⁸) Colloid, Radio** See: Radio Gold (Au¹⁹⁸) Colloid
- GOLD, COLLOIDAL**.
See: Colloidaurum, Amp (Kahlenberg)
Collosol Aurum, Sol, Amp (Crookes Barnes)
Rexaurum, Amp (Kahlenberg)
- GOLD COMPOUNDS**
See: Auracein, Amp (Christina)
Aural Sulfide, Sol, Vial (Hille)
Auralake, Vial (Lakeside)
Aurathiooglucose, Amp, NNR (Schering)
Aurathio glycander, Vial, NNR (Endo)
Gold Bromide (Merck Sharp & Dohme)
Gold Sodium Thiomalate NNR (Merck Sharp & Dohme)
Gold Sodium Thiosulfate (Various Mir)
Gold Thioglucose, Vial NNR (Schering)
Lauron, Vial (Endo)
Myochrysin, Amp (Merck Sharp & Dohme)
Rexaurum, Sol, Amp (Kahlenberg)
Solgonal, Amp (Schering)
- GOLDEN COUGH SYRUP** (Stayne) d metharphan hydrobromide 7.5 mg glyceryl guaiacolate 25 mg, chlorpheniramine maleate 1 mg/5 cc Syr 4 oz
Use Antitussive
- GOLD NAPHTHYL SULFHYDRYL DERIVATIVE**. See Auracein, Amp Christina
- * **GOLD SODIUM THIOMALATE, NF NND 1962**; Inj, NF See: Myochrysin Amp (Merck Sharp & Dohme)
- * **GOLD SODIUM THIOSULFATE, NF**; Sterile NF; NNR 1951 Auricidine Auracidⁿ Aurolin, Aurapin, Aurasan, Novacrysin Sulfactosol & Thiachrysin
Abbott—Amp 10, 25, 50, 100 mg; Bot 6s, 25s
Merck—Pow, Amp 10 mg, 25 mg; 50 mg, Bot 100 mg
Miller—Cryst Flacon 50 mg; Box 6s 50s
Use: Rheumatoid arthritis
- W/Sod thiosulfate & Sod bisulfite See Auralake, Amp (Lakeside)
- GOLD SULFIDE, COLLOIDAL**. See: Aural Sulfide (Hille)
- GOLD SUSPENSION**. See: Collosol Aurum Amp (Crookes Barnes)
- GOLD THIOGLUCOSE**. See: Aurathiooglucose
- GOLD TRIBROMIDE**.
See: Aubromid, Ellixir (McNeil)
Auri Tussin Liq (Zemmer)
Aurabram (Prof Pharm Co)
Tribromaura, Liq (Chicago Pharm)
- GOMENOL AQUEOUS** (Medics Pharmaceutical) Gomenol saturated metharphanated potassium guaiacal sulfanate 40 mg

sodium iodide 50 mg., sodium bisulfite 0.1%, guaiacol saturated, eucalyptal saturated, benzyl alcohol 2%/cc. Inj.—Amps. 2 cc. Multiple dose vials 30 cc.
Use: Expectorant.

GOMENOL PREPARATIONS.

See: Gombromal (Endo)

Gomicol, Vial (Kirk)
Gomoleaf, Vial (Gold Leaf)
Guacamphal, Vial (Metropolitan)
Komerlin, Vial (Lincoln)
Tego-Nole, Vial (Ortega)

GOMENOL COMPOUND. (New Mexico Pharm.) Gomenol 0.1 Gm., guaiacol 0.1 Gm., eucalyptal 0.08 Gm., iodolerm 0.02 Gm., camphor 0.05 Gm./2 cc. Vial 30 cc.

GOMICOL. (Kirk) Gomenol 0.10 Gm., guaiacol 0.10 Gm., iodolerm 0.02 Gm., camphor 0.05 Gm., eucalyptal 0.10 Gm., In oil, Amp. 2 cc., Box 12s, 25s, 100s, Vial 30 cc.

Use: (I.M.) Expectorant

GOMOLEAF. (Gold Leaf) Gomenol 0.12 Gm., iodolerm 0.02 Gm., guaiacol 0.10 Gm., camphor 0.05 Gm., eucalyptal 0.10 Gm., sesame oil q.s./2 cc. Amp. 2 cc., Vial 10 cc., 30 cc.

Use: (I.M.) Expectorant in bronchitis & in whooping cough

GONACRINE. See: Acrifovine (Various Mfr.)

GONADIN. (Cutter) Equine gonadotropin dried. Vial (500 u.) 1 cc.; (5000 u.) 10 cc.
Use: Functional uterine bleeding.

GONADOGEN, STERILE. (Upjohn) Purified & biologically standardized prep. of gonadotropic hormone obtained from serum of pregnant mares. Tab. Vial (10 Cortland-Nelson u., 20 C-N u., 50 C-N u., 250 C-N u.)
Use: (I.M., I.V., S.E.), gonadotropin therapy.

GONADOTROLIN. (Lincoln) Chorionic gonadotropin. Vial (500 u./cc.) 10 cc.

Use: Functional uterine bleeding in the female and depressed sexual development in males.

GONADO-TRONE. (Smith, Miller & Patch) Gonadotropin, pituitary oel. lobe. Vial (1000 Mouse Units) 10 cc. R/C w/diluent, Box 1s, 6s, 25s.

Use: Hypopituitarism w/associated sexual underdevelopment.

GONADOTROPIC SUBSTANCE. See: Gonadotropie Chorionic.

GONADOTROPIN, PITUITARY ANT. LOBE. Extracted from anterior lobe of equine pituitaries (not pregnant mare urine) (1 rat unit = 1 Fernald-Hisaw unit).

See: Aedatrape, Vial (Ewell)
Biotropin, Vial (Biophysics)
Climactie, Vial (Myers-Carter).
Equiphysic (G. F. Harvey)

* Generic name recognized by N.F., U.S.P. or A.M.A.

F-S-L, Vial (Myers-Carter)
Gonado-Trone, Amp. (Miller)

Tropiate, Vial (Mallard)

* **GONADOTROPIN, CHORIONIC, U.S.P.** 10 I.U., U.S.P. Numae Pregnancy Urine.

Uses: In the female: Chronic cystic mastitis, functional sterility, dysmenorrhea, premenstrual tension, & threatened abortion. In the male: Cryptorchidism, hypogonadism, dwarfism, impotency, euresis.

See: Anpitrane, Vial. (Gold Leaf)

Antotrin, Vial. (Parke, Davis)

A.P.L., Vial (Ayers).

Apestrin, Vial (Harrower).

Apitor, Vial. (Ethro)

Charoald, Vial (C. D. Smith).

Chargon, Vial. (Massengill)

Choroginin, Amp. (Lakeside).

Entromone, Vial (Endo).

Fallulele, Vial (Squibb).

Gestazol-Dry, Vial (National Drug).

Glentropin, Vial (Ingram)

Gonadotropin, Vial (Lincoln).

Harvatropin, Vial (G. N. Harvey)

Karotrin, (No Mfr. currently lists.)

Kuorin, Vial (Kremers-Urban).

Prenturon, Vial (Schering).

Pregnyl, Amp. (Organon).

Biogon, Vial (Breon).

Tropigon, Vial (Schleifelin).

W/Vit. B-1, glutamic acid, procaine HCl.

See: Glakar, Vial. (Research Supplies)

* **GONADOTROPIN SERUM, U.S.P. Pregnant Mare's Serum.**

See: Anteron, Vial (Schering).

Gonadin, Vial (Cutter)

Gonadogen, Vial (Upjohn)

GONIOSCOPIC FLUID. (Crookes-Barnes)

Methylcellulose 2.5%/Lacrival 5 cc

Use: With gonioscopic prisms.

GONOCOCCUS ANTIGEN. (Lilly) UBA-60 combined. Vial 5 cc.

GONOCOCCUS VACCINE. (Sherman) Combined: Gonococci, Strep., Staph. albus & aureus, E. coli, C. pseudodiphthericum. Vial 12.5 cc., 20 cc.

GOTHESTRONE. (Gotham) Estrogens mixed, eq. 20,000 u./cc. Amp. 1 cc., 10s, 100u Vial 10 cc., 30 cc.

Use: Estrogen therapy.

GOULARD'S EXTRACT. See: Lead Subacetate Solution (Various Mfr.)

GOYL. See: Acetarsone, Tab. (Abbott).

G.P.C. (Newman) Glycero-phosphates of Cal., Sod., Pot., Iron & Mang. w/Vit. B-1. Bot. 1 pt.

Use: Stimulants of appetite

GRABEXINE. (United Lab) Pyridoxine 100 mg., thiamine 100 mg./cc. Vial 10 cc.

- *GRAMICIDIN, NF, NNR 1951 See Gramoderm Oint (Schering)
- W/Fludrocortisone acetate & neomycin See Flanef S, Lot & Oint (Squibb)
- W/Hydrocortisone, neomycin sulfate, polymyxin B sulfate, phenylephrine & povidone HBr. See: Trisacort, Spraypak. (Smith, Kline & French)
- W/Neomycin See: Bia Cuta, Oint (Columbus)
- W/Neomycin sulfate, hydrocortisone, phenylephrine HCl See Du Biatic Intranasal (White)
- W/Neomycin sulfate, polymyxin B sulfate See Nea Polycin Ophthalmic Solution (Pitman Moore)
- W/Neomycin sulfate & polymyxin B sulfate See Neosporin, Sol (Burroughs Wellcome)
- W/Neomycin sulf, polymyxin B sulf, benzocaine See: Tricidin, Oint (Boyle)
- W/Neomycin sulfate, propyl paminobenzoate See: Orabiotic Gum Tab (White)
- W/Neomycin sulfate, thonzylamine HCl, phenylephrine HCl & thonzonium bromide See Biomydrin, Sol (Nepero)
- W/Neomycin sulf, thonzylamine HCl, thonzonium bromide See: Biomydrin Otic (Nepero)
- W/Neomycin sulf, thonzylamine HCl, thonzonium bromide, phenylephrine HCl and hydrocortisone See: Biomydrin F W/Hydrocortisone (Nepero)
- W/Neomycin, triamcinolone acetate See Kenalog S, Cream, Oint (Squibb)
- W/Neomycin, triamcinolone, nystatin See: Mycalog Cream (Squibb)
- W/Nystatin, fludrocortisone acetate, neomycin sulf See: Mycanef Ointment (Squibb)
- W/Phenylephrine HCl Chlor Trimeton glyconate See: Coricidin Nasal Mist, Soln (Schering)
- W/Phenylephrine HCl, pyrilamine maleate, cetyl dimethylbenzylammonium chloride See: Biohisdin, Sol (Columbus)
- W/Polymyxin B sulfate, benzocaine, cetalkonium chloride, chlorpheniramine maleate, tyrrathricin See: Ramex Traches (American Pharmaceutical)
- W/Polymyxin B sulf, methoxamine HCl See Aerodrin Sol (Burroughs Wellcome)
- W/Polymyxin B sulfate, quaternary HCl & cetyl dimethyl benzylammonium chloride See Driazets, Loz (Smith, Kline & French)
- GRAMODERM. (Schering) Gramicidin 0.25 mg/Gm Oint Tube 20 Gm
Use: Topical, antibiotic ointment
- GRAN. (Taylor Labs) Trypsin conc, 1 mg; Peru balsam, 72.4 mg; & castor oil/cc Aerial container, 5.5 oz
Use: Treatment for decubital ulcers, burns, abscesses, & other wounded surfaces
- GRANULESTIN. (Assoy Concentrates) Concentrate of purified soy phospholipids 80% comprising equal parts of lecithin, cephalin & lipositol Jar 9 oz
Use: Vitamin supplement
- GRATUS STROPHANTHIN. See: Quabain (Various Mfr)
- GRAUDA. (Raxall) Vit A 6000 u, D 1200 u thiamine HCl 9 mg, riboflavin 6 mg, ascorbic acid 111 mg, niacinamide 30 mg, folic acid 21 mg, mixed tocopherols 37 mg, dicalcium phosphate (anhy) 2607 mg, ferrous gluconate 135 mg/Cap Bot 100s, 300s, 1000s
- GRAVIDOX. (Lederle) Pyridoxine HCl & thiamine HCl Sol (50 mg of each) Vial 10 cc; Tab (20 mg of each) Bot 50s
Use: I.M., IV & orally, to relieve excessive nausea and vomiting of pregnancy
- GRAVA-VITE-P. (Hoyt) Fluorine (from 221 mg sodium fluoride) 1 mg elemental iron (from 135 mg), ferrous sulfate exsiccated 40 mg, calcium (from 625 mg calcium carbonate) 250 mg vit A 6000 u, D 400 u B 1 13 mg, B 2 2 mg, B 12 2 mcg C 100 mg, niacinamide 20 mg calcium pantothenate 5 mg/Tab Bot 100s
Use: Prenatal nutritional support
- GRAVOCIBUS. (Canfield) Vit A 4500 USP u, D 900 USP u E 0075 IU, & 1 1125 mg B 2 1 875 mg, B 6 0 075 mg, C 75 mg d calcium pantothenate 0.9 mg, niacinamide 11.25 mg, B 12 0 075 mcg, Ca 561 mg, Mg 225 mg, Fe 45 mg, P 0 075 mg I 0 111 mg, Mo 0 075 mg Cu 1 125 mg Mn 2 25 mg, Zn 2 25 mg, Co 0 225 mg lolic acid 0 0501 mg/Tab Bot 100s 500s 1000s
Use: Prenatal vitamin mineral combination
- GRAVOCIBUS-P. (Canfield) Vits A 2000 u; D 270 u, B 1 1 mg, B 2 1 2 mg, niacinamide 10 mg, B 6 2 mg, Ca pantothenate 1 mg, cyanocobalamin 2 mcg C 33.3 mg; Ca 250 mg, P 194 mg Fe 5 mg, Mn 0 01 mg, K 1 mg Zn 0 15 mg/Cap Bot 100s 500s, 1000s
Use: Prenatal vitamin mineral combination
- GRAVOCIBUS-P-F. (Canfield) Vits A 3000 u, D 300 u, B 1 3 mg, B 2 2 mg, niacinamide 10 mg, B 6 3 mg, Ca pantothenate 2 mg, cyanocobalamin 3 mcg, C 37.5 mg; Ca 150 mg, Fe 5 mg, Mg 1 mg, Mn 0 1 mg, K 1 mg, Zn D 15 mg/Cap Bot 100s 500s, 1000s
Use: Prenatal vitamin mineral combination
- GRAY POWDER. See Mercury with Chalk (Various Mfr)

GRECORT & NEO-GRECORT CREAM & LOT. (Greco) Greco Cort Cream: $\frac{1}{4}$, $\frac{1}{2}$, 1 or 3% and lotion $\frac{1}{4}$, $\frac{1}{2}$ or 1% microfine hydrocortisone. Neo-Grecort: $\frac{1}{4}$, $\frac{1}{2}$ or 1% and lotion $\frac{1}{4}$, $\frac{1}{2}$ or 1% microfine hydrocortisone with $\frac{1}{4}$ % neomycin sulfate added. Cream: Tube 1 oz.; Jar 4 oz.; 1 lb. Lotion: Bot. 1 oz.; 4 oz.; pts.

Use: Dermatitis.

GREENIES. (Central) Acetylsalicylic acid $3\frac{1}{2}$ gr., acetophenetidin $2\frac{1}{2}$ gr., caffeine, anhyd. $\frac{1}{2}$ gr./Tab. Bot. 100s, 1000s.

Use: Urter Analgesic.

GRIFULVIN, N.N.D. 1962. (McNeil) * Griseofulvin 250 mg scored tabs. Bot. 16 & 100s, 500 mg/Tab. Bot 20r. 100s. Susp. (250 mg/3 cc.) Bot. 4 oz.

Use: Oral. Fungus infection of the skin, hair, and nails.

GRIFULVIN V. (McNeil) Griseofulvin microcrystalline 125 mg., 250 mg/Tab. Bot. 30s, 100s, 500 mg/Tab. Bot. 20s, 100s.

Use: Ringworm infections.

GRISACTIN, (Ayerst Lab) Griseofulvin 125 mg/Tab. Bot. 100s, 500s.

Use: Ringworm infections.

* **GRISEOFULVIN, N.N.D. 1962.** An antibiotic. 7-chloro-4,6-dimethoxycoumaran-2-one-2-spiro-1'-(2'-methoxy-6'-methylcyclohex-2'-en-4'-one)

See: Fulvicin-U/F (Schering)

Grifulvin Suspension (McNeil)

Grisactin (Ayerst Lab)

Grifulvin V (McNeil)

* **GRISEOFULVIN, N.N.D. 1962.** (Ayerst) 250 mg Bot 30s, 100s & 500 mg Tab. Bot 50s.

Use: Oral fungistat for a variety of dermatomycoses.

GROWTH (SOMATOTROPIC) HORMONE.

See: Antultrin Growth (Farke, Davis).

Leutropin (Barry).

G-STROPHANTHIN, N.N.R. 1949. See: Quabain. (Various Mfrs.)

GUACAMPHOL. (Metropolitan Eebs.) Gamenal 0.10 Gm., gualacol 0.10 Gm., eucalyptol 0.08 Gm., iodotorm 0.02 Gm., camphor 0.05 Gm./2 cc. Amp. 2 cc., Box 10s, 100s. Vial 30 cc.

Use: (I.M.) Expectant

GUAIACODYL. (Meyer) Concentrated mixture of Guaiacol 5% and cacodylic acid $2\frac{1}{2}$ %. Vial 10 cc., 30 cc.

Use: Liquefying expectorant.

GUAIACOL, N.F. X, N.N.R. 1937, A.D.R. (Various Mfrs.) Methylcatechol. Use: Chronic infections of the respiratory tract.

* Generic name recognized by N.F., U.S.P. or A.M.A.

GUAIACOL CARBONATE, N.N.R. 1937. (Various Mfrs.) (Duotal).

Use: Expectorant.

GUAIACOL GLYCERYL ETHER. See: Glyceryl Guaiacolate

GUAIJA-COMPH. (Smith-Dorsey) Guaiacol, methyl salicylate, camphor, menthol, oil of Siberian fir, wool fat, petrolatum. Jar 2 oz., 1 lb., 5 lb.; child's strength, Jar 2 oz. Use: Counterirritant.

GUAIAFAGE. (Meyer) Guaiacol 25 mg, cacodylic acid 12.5 mg, foreign protein 20 mg., benzyl alcohol 1%/cc. In aqueous vehicle. Vial 10 cc., 30 cc.

Use: Upper respiratory infection.

GUAIAGESIC. (Columbus) Phenylephrine HCl 5 mg., pyrilamine maleate 6.25 mg., glyceryl guaiacolate 50 mg., salicylamide 60 mg., acetaminophen 60 mg./5 cc. Bot. 3 oz., pt.

Use: Pediatric cold therapy.

GUAIAMIST. (Columbus) Phenylephrine HCl 5 mg or 10 mg, pyrilamine maleate 12.5 mg, glyceryl guaiacolate 100 mg/Tab. Bot. 100s, 1000s. Available w/Narcolline 3 mg/Cap. Bot 100s.

Use: (1 q.t.d.) Antihistaminic.

GUAIAMIST DX. (Columbus) Dihydrocodeinone bitartrate 166 mg, phenylephrine hydrochloride 10 mg, pyrilamine maleate 12.5 mg, glyceryl guaiacolate 50 mg, ammonium chloride 50 mg, chloroform 9.2 mg, l-methol 1 mg/5 ml. Bot. 1 pts, 1 gal.

Use: Treatment of the common cold.

GUAIAMIST TT. (Columbus) Phenylephrine HCl 20 mg, Chlorpheniramine maleate 4 mg, glyceryl guaiacolate 300 mg/Temp. controls-trms-controlled Bot. 50s.

Use: Respiratory Infections

GUAIAMIN CREAM. (Lonnell) Methyl salicylate, guaiacol, menthol in greaseless base. Jar 4 oz. & 1 lb.

GUAIANESIN. See: Glyceryl Guaiacolate

GUAIASYL ELIXIR. (Meyer) Paragoric 5 cc., elixir phenobarbital 3.5 cc., terpin hydrate 0.5 Gm., Pot. gualacol sulfonate 1 Gm., comp. squill syr. 2 cc., creosote 0.1 cc., trest. glycyrrhiza 5.5 cc./fl. oz. Bot. 1 pt, 1 gal.

Use: Sedative with antispasmodic.

* **GUAIIPHENESIN.** 3-a-Methoxyphenoxypropene-1,2-diol. Guaiacol glycerol ether. Resphenyl. See: Glyceryl Guaiacolate.

GUAMIDE. See: Sulfoguanidine. (Various Mfrs.)

* **GUANETHIDINE SULFATE.** [2-(actohydro-1-acetyl)ethylguanidine sulfate. See: Isonin (Ciba)]

Use: Ganglionic blocking agent.

GUANIPINE. (Davies, Rose) Guanidine NCI
 Tab 125 mg Bot 50s
 Use: Parasympathetic stimulant

***GUAR CELLUPECTINOID.** W/Magnesium
 oxide aluminum hydroxide gel, magnesium
 trisilicate See Belvis, Tabs (Carrick)

GUAR GUM. W/Sennasides A and B, poly
 galacturonic acid See Gentles Granules
 (Purdue Frederick)

GUAYANESIN See: Glyceril Guaracolate
 (Various Mfr)

GUIA-CAMPH (Smith Darsey) Guaracol,
 methyl salicylate, camphor, menthol, pine
 needle oil Ointment Jors Reg 2 oz, Ib
 Childrens 2 oz

GUI-ACUA. (Tulag) Guaracol 50 mg,
 cocodylic acid 25 mg, benzyl alcohol 0.06
 cc, chlorbutanol 5 mg intracorne 0.5%/
 cc Vial 20 cc
 Use: Expectorant and antitussive

GUIAMIDE. (Tulag) Sulfoguanidine 0.5 Gm,
 kaolin 2 Gm, pectin 40 mg/5 cc Liq Bot
 4 oz, pl, gal
 Use: Antibacterial and intestinal adsorbent
 in diarrhea

GUIATISE (Warner) Pot guaicol sulfanole
 8 4/5 gr, iron hypophosphite 1 2/5 gr,
 Mang hypophosphite 1 gr, Cal hypophos-
 phite 8 gr/A oz Bot 8 A oz, 1 pl., 1 gal
 Use (1 2 tsp q 3 h pr n) Expectorant

GUIODEC. (New Mexico Pharm) Camphor
 60 mg, Eucalyptol 80 mg, Guaracol 100
 mg, Iodafarm 15 mg, Gamonal 10 mg,
 Benzyl Alcohol 2% / 2 cc Vials 30 cc

GUNCOTTON, SOLUBLE. See Pyroxylan
 USP

GUSTALAC. (Genetric) Cal carbonate 300
 mg, defatted skim milk pow 200 mg/
 Tab Bot 100s, 250s, 1000s
 Use: Antacid in the treatment of peptic
 ulcers

GUSTAMATE (Arlington) Not listed Mono
 ammonium glutamate, plus glycine & glu-
 tamic acid in salt shaker type dispensers
 containing 1 oz
 Use: Salt substitute

GUSTASE PLUS. (Geriatric) Phenobarbital 8
 mg, hamatrapine methylbromide 2.5 mg,
 geriare 30 mg, genpratare 6 mg, gerr
 cellulare 2 mg/Tab Bot 42s, 100s
 Use: Antispasmodic and a sedative

GUSTAZYME (Geriatric) Amylolytic enzyme
 30 mg, proteolytic enzyme 6 mg, cellu-
 lolytic enzyme 2 mg/Tab Bot 42s, 100s
 500s
 Use: Gastrointestinal enzymes

G-VITAMIN See: Riboflavin (Various Mfr)

G.V.S. AEROSOL FOAM. (Savage) Gentran
 violet 1.35%/Gm preservatives and inert

* Generic name recognized by N.F. USP or A.M.A.

materials 1/2 oz w/reusable plas applic
 2 oz w/4 reusable plas applic
 Use: Vaginitis

GWJALATE. (Breen) Codeine phosphat 1/1
 gr, phenylephrine HCl 6 1/4 mg, Pot guaico
 col sulfanole, ammonium chloride, menthol
 chloroform & alcohol/5 cc Bot 1 pl, 1 gal
 Use: (1 tsp q 3 h) Narcotic & antihib-
 itamine cough preparation

GYN. (Grant) Mag gluconate 0.5 Gm/Tab.
 Bot 100s
 Use: (1 2 tid) Antispasmodic in treatment
 of primary dysmenorrhea

GYNALGOS. (Wyeth) Aminopyrine 20 gr,
 Pot bromide 20 gr/A oz Bot 1 qf
 Use: Analgesic & antipyretic sedative

GYNAPLEX. (Rowell) Tyrothricin 0.05%
 zinc caprylate 5%, zinc propionate 5% in
 water miscible cream adjusted to pH 4.5
 Tube 3 oz
 Use: Vaginal cream

GYNBEN. (Barre) Dried dihydroxyquinolone
 100 mg, tartaric acid 1/10 gr, boric acid
 2 gr, stilbestrol 0.1 mg, sulfadiazine 0.5
 Gm, kaolin 11 gr, dextrose 11 gr, lactose
 11 gr/insert Box 24s
 Use: Vaginal insert for the treatment of
 vaginal infections

GYNERGEN. NNR 1950 (Sandoz) Ergob-
 amine tartrate Amp (0.25 mg) 0.5 cc,
 Box 6s, 50s, (0.5 mg) 1 cc, Box 6s 50s
 Tab. (1 mg) Bot 100s, 500s, 1000s
 Use: SC, S.I., I.M., orally, sympatholytic
 agent & oxytocic

GYNERGON. See: Estradiol (Various Mfr.)

GYNETONE. (Schering) Oil sol estrad of
 benzoate 1 mg, testosterone propionate
 20 mg/cc Vial 10 cc, Box 1s 6s Tab.
 estradiol 1 mg, methyltestosterone 10 mg/
 Tab Bot 30s, 100s
 Use: I.M., orally, androgen estrogen therapy

GYNETONE REPETABS. (Schering) Ethinyl
 estradiol 0.02 mg & 0.04 mg, methyltes-
 tosterone 5 mg & 10 mg in 2 layers that
 divide dose inner coat for delayed action
 & outer coat for immediate absorption
 Tab 02, Bot 30s 100s, 04 Bot 30s
 100s
 Use: Orally, androgen estrogen therapy

GYNOESTRYL. (Standard) Estrogen mixed
 in oil Amp 10 000 u/cc, Box 6r, 100r
 25 000 u/cc Box 3s Vial 30 cc Amp
 50 000 u/cc 1 cc Box 3s
 Use: I.M., estrogen therapy

HAEMOL. (Irwin, Nersler) Liver, desic (de
 fatted) 3 gr, ferrous sulfate, asic 2 gr,
 thiamine mononitrate 3 mg, riboflavin 1

- mg., niacinamide 9 mg./Cap. Bot. 100s, 500s, 1000s.
Use: Anti-anemia therapy.
- HAIMASED.** (Tilden) Each 100 cc. contains 4.4 Gm. Sod. thioeyanate (20 gr. to fl. oz.) Bot. 1 pt., 1 gal.
Use: (1 tsp. t.i.d., p.c.) Hypertension.
- HAKRITON.** (Hoac) Phenylpropanolamine HCl 10 mg., d pseudoephedrine HCl 10 mg., pyrilamine maleate 5 mg./Tab. Bot. 100s, 1000s.
Use: Nasal congestion.
- HALABAR.** (Carrick) Butobarbital 16 mg., mephenesin 300 mg./Tab. Bot. 100s, 1000s.
Use: (1 p.c., t.i.d. & h.s.) Sedative.
- HALABAR No. 2.** (Carrick) Butobarbital 28 mg., mephenesin 300 mg./Tab. Bot. 100s.
Use: Sedative.
- HALDRONE.** (Eli Lilly) Paramethasone Acetate.
Use: Anti-inflammatory, anti-allergic glucocorticoid steroid.
- ***HALAZONE, N.F., N.N.B. 1952;** Tab., N.F. (Abbott) N,N-dichlor sulfamidobenzolic acid. Tab. 4 mg. Bot. 100s, 1000s.
Use: Disinfectant.
- HALAZYME.** (Halco) Amylolytic enzyme 30 mg., proteolytic enzyme 10 mg., cellulolytic enzyme 2 mg., aluminum hydroxide gel 60 mg./Tab. Bot. 30s, 100s
Use: Indigestion.
- HALDRONE.** (Lilly) Paramethasone acetate 1 mg., 2 mg./Tab. Bot. 30s, 100s.
Use: Steroid therapy, anti-inflammatory agent
- HALEY'S M.O.** (Phillips) Mineral oil & milk of magnesia. Bot. 8 oz., 1 pt., 1 qt. Also flavored.
Use: Orally, laxative.
- HALIBUT LIVER OIL, N.F., Cap., N.F.**
Use: Vitamin A & D deficiencies.
- W/Viosterol, N.N.R. 1935, A.D.R. 1935.
- HALIMIDE.** (Bard-Parker) Dodecylbeazyl trimethyl ammonium chloride 25.6%. Bot. 4 oz.
Use: (Diluted 1 oz. to gal. of water or etc.) Disinfection of instruments
- HALINONE.** (U.S. Vitamin & Pharmaceutical) Bromindione.
Use: Anticoagulant.
- HALIVER OIL.** See: Halibut Liver Oil (Various Mfr.).
- HALIVER OIL WITH VIOSTEROL, N.N.R. 1955, A.D.R. 1955.** See: Halibut Liver Oil with Viosterol.
- HALODRIN.** (Upjohn) Fluoxymesterone 1 mg., ethinyl estradiol 0.02 mg./Tab. Bot. 100s
Use: Hormone preparation.
- * Generic name recognized by N.F., U.S.P. or A.M.A.
- HALOPERIDOL.** (Searle) 1-[3-p-fluorobenzyloxypropyl]-4-p-chlorophenyl-4-hydrozypiperidine.
Use: Ataractic.
- ***NALOPROGESTERONE.** 6a-fluoro-17a-bromoprogesterone. See: Prohalone (Ayerst Lab)
- HALOTESTIN, N.N.D. 1962.** (Upjohn) Fluoxymesterone, 9-alpha-fluoro-11-beta-hydroxy-17-methyltestosterone, Tab. 2 mg. 100s and 5 mg. & 10 mg. Bot. 50s.
Use: (5 to 10 mg. do.) Androgenic and anabolic agent
- W/Ethinyl estradiol, Vit. A, D, thiamine mononitrate, riboflavin, pyridoxine HCl, cyanocobalamin, ascorbic acid, nicotinamide, calcium pantothenate, folic acid, calcium phosphate, dibasic. See: Hormozyme, Tab. (Upjohn)
- ***HALOTNANE, N.N.D. 1962.** 1,1,1-Trifluoro-2-bromo-2-chloroethane. See: Fluothane (Ayerst Lab)
- HALOXSIL.** (Holler) Methylpolysiloxane 40 mg./Tab. Bot. 100s.
Use: Gastro-intestinal.
- HALPHTOL.** (Holler) Pentacerythritol tetranitrate 80 mg./Tab. Bot. 100s.
Use: Coronary vasodilatation
- HALTRAN.** (Holler) Amobarbital 1/2 gr., butobarbital 1/2 gr./Tab. Bot. 100s.
Use: Sedative.
- HALVASTIM.** (Holler) Niacin 150 mg., niacinamide 50 mg., ascorbic acid 50 mg., riboflavin 5 mg., thiamin HCl 10 mg., pyridoxine HCl 2 mg., Vit. B-12 activity 3 mcg., calcium pantothenate 5 mg., tolazoline HCl 10 mg./Tab. Bot. 100s.
Use: Vitamin deficiencies and vascular disorders.
- NALVICORT.** (Holler) Prednisone 1 mg., salicylamide 200 mg., hesperidin conc 30 mg., ascorbic acid 30 mg., Vit. K 0.3 mg., alum. hydrox. gel. 30 mg./Tab. Bot. 100s
Use: Rheumatic disorders.
- NAMA.** N-hydroxy-aluminum magnesium aminoacetate. See: Glycamine-A.M. (Victor M. Hermelin)
- NAMAMELIS, EXTRACT** W/Butelin & hydroxide ext. See: Butelin Compound, Supp. (Abbott).
- W/Zinc oxide. See: Hazeline, Soap (Burgess Wellcome).
- NANSEN'S DISEASE.**
See: Chaulmoogra Oil (Various Mfr.).
Diazene, Tab. (Abbott).
Ethinyl Chaulmoograte (Various Mfr.).
Isosalazid (Various Mfr.).
Promie, Amp. (Parke, Davis).
Promizole, Tab. (Parke, Davis).

protein 50 mg, folic acid 3 mg /Tab Bot 50s

Use Orally hematinic

W/Liver conc (Arlington) Alcohol 17%, iron (chemically combined with protein 120 mg liver conc 500 mg, sugar, glycerin, flavor ing qs /fl oz Bot 12 fl oz, 1 gal Tab iron (chemically combined with protein) 35 mg, liver conc 100 mg Bot 60s

Use Hematinic

HEMACETS (Walker) Ferrous fumarate 100 mg, ascorbic acid 50 mg, Vit B 12 2 mcg, pyridoxine HCl 0.5 mg /Candisphere Bot 100s, 1000s

Use: Treatment of iron deficiency anemia

HEMACIL (Endocrine Co) Vit B-12 10 mcg C 50 mg, folic acid 1 mg, liver NF 200 mg, stomach pow 100 mg ferrous sulfate 194 mg /Cap Bot 50s, 100s

Use: All anemias

HEMACIN (Walker) Liver stomach conc 300 mg, Vit B 12 15 mcg, folic acid 1 mg, ascorbic acid 75 mg, ferrous sulfate exsic 300 mg, copper, cobalt, Mang, molybde num, zinc, Mag /Cap Bot 60s

Use (1 daily) Nutritional anemias

HEMACOMBIN (Rand) Vit B 12 5 mcg, folic acid 2 mg, Vit C 75 mg, B-1 3 mg B 2 2.5 mg, ferrous gluconate 136 mg, desic liver 100 mg, Vit B 6 1.5 mg, nicotinamide 10 mg /Tab Bot 100s

Use (1-3 daily) Hematinic

HEMACOMBIN ELIXIR (Rand) Vit B 12 4 mcg, iron ammonium citrate 200 mg, B 1 1 mg, B 2 0.5 mg, sod glycerophosphate 1 gr, manganese glycerophosphate 2 mg pot glycerophosphate 0.5 gr /5 cc Bot pt, gal

Use: Anemias

HEMACOMBIN-P (Rand) Vit B 1 3 mg B 2 2.5 mg, B 6 1.5 mg, B 12 5 mcg folic acid 2 mg, ascorbic acid 75 mg, ferrous gluconate 136 mg, desiccated liver 100 mg, niacinamide 10 mg, dextro amphetamine sulfate 1.5 mg /Tab Bot 100s, 1000s

Use: Vitamine supplement W/CNS stimulant

HEMACOMBIN-T (Rand) Vit B 1 3 mg, B 2 2.5 mg, B 6 1.5 mg, B 12 5 mcg, folic acid 2 mg, ascorbic acid 75 mg, ferrous gluconate 136 mg, desiccated liver 100 mg, niacinamide 10 mg, reserpine 0.05 mg /Tab Bot 100s, 1000s

Use: Vitamin supplement

HEMAFER (Stuart) Formerly known as Stuart Hematinic w/Folic Acid and Vit B-12

HEMAFOL-12 (Normand Pharm) Crude liver, vit B 12 (equiv to 2 mcg cyano

cobalamin) folic acid 5 mg, Vit B 12, 30 mcg /cc Vial 10 cc

Use Periculous anemia

HEMAFOLATE (Canright) Ferrous glucooate 293 mg, liver fraction II 250 mg gastric sub 100 mg, Vit C 50 mg, B-12 10 mcg, folic acid 0.3 mg /Tab Bot 100s

Use Hematinic agent

HEMALIQUID (Ulmer) Ferrus s H 10 Gm, vit B 1 B mg B 2 B mg, nicotinamide 32 mg, yeast conc 1.5 Gm, liver fraction No 1 2 Gm /fl oz

Bot 1 pt, 1 gal

Use: (1-3 tsp daily) Hypochromic anemia

HEMALIRONE CAPSULES (Blue Line) Ferrous sulfate 1 gr, liver conc 5 gr, cascara ext 1/4 gr /Cap Bot 100s

Use (2 tsd) Anemias

HEMALIRONE LIQUID (Blue Line) Ferrous sulfate 15 gr, liver fraction 27.5 gr /2 dr Bot 1 pt, 1 gal

Use: (1-2 tsp daily) Hematinic

HEMANEED (Hanson Drug) liquid Bot 4 oz, 8 oz

HEMAPOIETIC AGENTS

See Lipotropic Preparations

Liver Products

Iron Products

Vitamins

HEMAREX (Estraf) Liver inj 10 mg, iron 19 mg, niacinamide 30 mg, riboflavin 0.3 mg, pyridoxine HCl 0.3 mg /2 cc Amp 2 cc Vial 10 cc, 30 cc

Use Nutritional & secondary anemias

HEMAREX CIRCLETS (Estraf) Liver fraction No 2 3 gr ferrous sulfate exsic 3 gr, Vit B 1 1 mg, B 2 0.66 mg, niacinamide 10 mg, Vit C 50 mg /Cap Bot 1000s 5000s

HEMASTIX (Ames) A cellulose strip, im pregnated with a peroxide and ortho tolidine Bot 50s

Use Detection hematuria and hemo globinuria

HEMATIST (Ames) Strontium peroxide o Tolidie, tartaric acid & Cal acetate Tab Bot 60s, 500s

Use Diagnostic aid for the test of blood in the feces, urine & body fluids

HEMATINIC (Canright) Ferrous gluconate 183 mg, liver desic 200 mg, Vit B 12 1 mcg, folic acid 0.2 mg copper gluconate 0.3 mg, cobalt gluconate 0.12 mg, Vit C 25 mg, B 1 3.3 mg /Tab Bot 100s 1000s

Use Anemias

HEMATINIC FORTIFIED (Stuart) Vit B 1 10 mg, Vit B 2 10 mg, niacinamide 150 mg, Cal pantothenate 10 mg, Vit B 6 2 mg, Vit C 300 mg, folic acid 51 mg ferrous

gluconate 18 gr., liver 1200 mg., gastric subs. 600 mg., Vit. B-12 60 mcg., copper sulfate 15 mg./6 Tab. Bot. 100s, 1000s.
Use: Hematinic.

HEMATIHC PLASTULES. (Wyeth) Ferrous sulfate 5 gr., yeast conc. 0.75 gr./Cap. Bot. 100s.

Use: (1 t.i.d.) Hematinic.

W/Liver conc. (Wyeth) Ferrous sulfate 162 mg., liver conc. 170 mg., yeast/Plastule. Bot. 50s, 100s, 1000s.

Use: Hematinic.

HEMATIHCs.

See: Iron Products.

Ferric Compounds.

Ferrous Compounds.

Liver Products.

Vitamin B-12.

Vitamin Products.

HEMATOCRIN. (Harrower) Ferrous gluconate 194 mg., liver desic. 130 mg., thiamine HCl 5 mg., riboflavin 2 mg., niacinamide 10 mg., pyridoxine HCl 0.1 mg., Cal. pantothenate 0.5 mg., folic acid 0.34 mg., ascorbic acid 30 mg./Cap. Bot. 100s, 500s.

Use: (1-2 t.i.d.) Iron-deficiency and nutritional anemias.

W/B-12. (Harrower) Vit. B-12 cryst. 10 mcg., gastric substance 150 mg., folic acid 1 mg., liver desic. 150 mg., ferrous gluconate 194 mg., ascorbic acid 50 mg./Tab. Bot. 100s, 500s.

Use: (1-2 t.i.d.) Anti-anemia therapy.

HEMATOHE B. (Chicago Pharmacal) Contains glycerophosphate of Cal., berberine, strychnine sulfate, nux vomica, hypophosphites & cell-tissue salts, Bot. 6 oz., 1 pt., 1 gal.

Use: Hematinic.

HEMATOVALS. (Ulmer) Ferrous sulf. 4.5 gr., cobalt sulf. 2 mg., liver 110 mg., vit. B-12 1 mcg., B-1 1 mg., B-2 1 mg., B-6 0.25 mg., C 16.66 mg., folic acid 0.25 mg., cal. pantothenate 0.25 mg., niacinamide 3.3 mg./Cap. Bot. 100s, 500s, 1000s.

Use: (1-3 t.i.d.) Hematinic.

HEMATRHSIC. (Invenex) Ferrous gluconate 325 mg., liver conc. 250 mg., vit. B-12 w/intrinsic factor 1/5 u., B-12 15 mcg., B-1 5 mg., B-2 5 mg., B-6 1 mg., C 40 mg., folic acid 1 mg., niacinamide 25 mg./Cap. Bot. 50s, 100s, 500s.

Use: (1 t.i.d.) Hematinic.

HEMATRHSIC. (Rexoll) No. 10 Ferrous gluconate 200 mg., Vit. B-12 10 mcg., Vit. C 50 mg., folic acid 0.33 mg., liver 100 mg., stomach 100 mg./Cap. Bot. 100s, Ho. 25. Ferrous gluconate 200 mg., Vit. B-12 25 mcg., Vit. C 50 mg., folic acid 1.67 mg.,

liver 100 mg., stomach 100 mg./Cap. Bot. 100s.

Use: (1 t.i.d.) Hematinic.

HEMBRO. (Farrest) Minerals. Tab. Bot. 50s, 100s, 500s.

W/Liver concentrate. Bot. 50s, 100s, 500s.

Use: Secondary anemia.

HEMIACIDRH. See: Renocidin (Guardion Chem.).

HEMICIN. (Walker) Liver-stomach concentrate with Intrinsic factor 300 mg., Vit. B-12 15 mcg., folic acid 1 mg., Vit. C 75 mg., ferrous sulfate exsic. 300 mg., copper 1 mg., cobalt 0.2 mg., Mang. 1 mg., molybdenum 0.2 mg., zinc 1 mg., Mag. 1 mg./cap. Bot. 60s.

Use: 1 b.i.d.; anemias.

HEMISINE. See: Epinephrine (Various Mfr.).

HEMISUCCHOXYPREGNENOLOHE,

DELTA-5- See: Panzalone (Daak)

HEMO. (Borden) Vit. A 4000 International u., B-1 333 International u., B-2 2 mg., D 400 International u., niacinamide 10 mg., iron 14.7 mg., calcium 376 mg., phosphorus 288 mg./1½ oz. dry chocolate-flavored malted drink.

Use: Nutritional supplement.

HEMOCHROMIH. (Carnrick) Liver fraction 0.162 Gm., ferrous sulfate exsic. 0.162 Gm. Bot. 100s, 1000s.

Use: Secondary anemias.

W/B Complex and vitamin C. (Carnrick) Ferrous sulfate exsic. 195 mg., liver fraction 163 mg., Vit. B-1 2 mg., Vit. B-2 3 mg., nicotinamide 20 mg., Cal. pantothenate 1 mg., Vit. C 25 mg./Tab. Bot. 100s, 1000s.

Use: Secondary anemia.

HEMOCOAVIT. (Wynhit) Vit. C, and bioflavonoids plus Vit. K. Tabs. Bot. 50s, 100s, 1000s. Vial 10 cc.

Use: Antihemorrhagic

HEMOCCULT. (Schieffelin). Electrophoresis filter paper impregnated with guaiac gum and a developer solution/Kit. 100-test kit.

Use: Test for occult blood in feces and urine.

HEMODAL. See: Menadione, U.S.P.

HEMO-GEHIN. (Winthrop Labs) Ferrous gluconate 0.133 Gm., liver conc. 0.10 Gm., thiamine HCl 0.25 mg., riboflavin 0.50 mg., niacinamide 3.50 mg., pyridoxine HCl 0.05 mg., cal. pantothenate 0.05 mg./cap. Bot. 100s, 500s, 1000s.

Use: Hematinic with vitamin V complex.

HEMOGLOBIN. (Wilson) Ferruhemoglobin. Pow., Bot. 1 oz.

Use: Anemias.

HEMOLATUM. (First Texas) Benzocaine 2%, tannic acid 3%, gum camphor 4%, phenol 1%, Ichthammol 6%, zinc oxide 1½%.

sulfur precipitated 6% Tube 1 oz for 1 lb

Use: Soothing & antiseptic for hemorrhoids

HEMOLETS, (Warren Teed) Stramonium alk., phenol, ichthammal, horse chestnut, homo melis & benzocaine/Supp Boxes 12s

Use Hemorrhoid suppository

HEMOLIV-100, (Kirk) Crude liver injection 2 mcg, vit B 12 100 mcg / 2 cc Vial 10 cc Also available with vit B-12 250 or 500 mcg

Use (I.M. 1 or 2 cc daily) Pernicious anemia

HEMOMIN, NND 1962 (Kirk) Vit B 12 cryit Vial (30, 50 mcg / cc) 10 cc, 30 cc; (100, 1000 mcg / cc) 10 cc; (2000 mcg / cc) 5 cc

Use I.M., hematinic

NEMOMIN COMPOUND (Kirk) Vit B 1 3 mg, B 2 3 mg, B-6 0.5 mg, cal pantothenate 5 mg, niacinamide 20 mg, B 12 5 mcg, liver fraction No 2 60 mg, yeast dried 60 mg / cap Bot 100s, 1000s

Use: Orally hematopoietic with B complex

HEMOMIN-GEL, (Kirk) Vit B 12 200 mcg / cc Vial 10 cc

Use: I.M. prolonged-action, vitamin B 12 therapy

HEMONUTRON, (Nion) Ferrous gluconate 0.19 Gm, liver desic 96 mg, brewers yeast 32 mg, thiamine HCl 0.5 mg, riboflavin 0.5 mg niacinamide 3.5 mg / Tab Bot 100s, 500s Vial 10 cc

Use: (1-3 daily) Hypochromic macrocytic anemias

HEMONUTRON PLUS, (Nion) Vit B 12 5 mcg, folic acid 1 mg, ferrous gluconate 4 gr, ascorbic acid 50 mg, thiamine HCl 1 mg, riboflavin 2 mg, pyridoxine HCl D 5 mg, Cal pantothenate 0.5 mg, niacinamide 10 mg / Tab Bot 100s

Use (2 t.i.d., p.c) Nutritional macrocytic anemias

HEMO-PAK, NND 1962, ADR (Johnson & Johnson) Oxidized cellulose Absorbable surgical dressings, Tube 3" X 3" gauze pad, 14" X 2" gauze strip, 6" X 2" cotton pad Hemostatic absorbable gauze packing strip 14" X 2", 1/2" X 5", 1/2" X 2 1/2" yd, 3" X 3" gauze pad Hemostatic absorbable cotton 6" X 2", sterile packing strip 2" X 3" yd Packed 12s

NEMOPHILUS INFLUENZAE ANTISERUM. See Influenzae Serum, Vial (Squibb)

HEMOPHILUS INFLUENZAE VACCINE, MIXED (Merck Sharp & Dohme) Vial 5 cc 20 cc

Use For the prevention of complications with upper respiratory infections

HEMORATE (Kirkman) Ferrous fumarate 200

mg, ascorbic acid 100 mg / Tab Bot 100s, 1000s

Use: Iron deficiency anemia

HEMOSTASIN, (United Lab) Lymphoid tissue ext Vial 10 cc, 1s, 6s

Use: S.C., hemostatic

HEMOSTATICS, LOCAL

See: Absorbable Gelatin Sponge

Calamine Chloride (Merck Sharp & Dohme)

Fibrin Foam (Cutler)

Gelfilm (Upjohn)

Gelfoam (Upjohn)

Oxycel (Parke, Davis)

Oxidized Cellulose

Solution of Brain Extract

Salusponge (Parroy)

Stypven (Burroughs Wellcome)

Styptican (Merck Sharp & Dohme)

Thrombin (Parke, Davis)

Thrombol (Merck Sharp & Dohme)

HEMOSTATICS, SYSTEMIC

See: Adrenosem (Mastengill)

Adrenostat (Organon)

A.M.F., Gran (Armour)

Anti Hemophilic Serum

Blutene (Abbott)

Carbazochrome Salicylate

Ceanothyn (Flint, Eaton)

Fibrogen (Merrell)

Koogamin (Chatham)

Premarin 1V (Ayerst)

Thromboplastin (Various Mfr)

HEMOSTATIN See: Epinephrine (Various Mfr)

HEMOSULES, (Warner-Chilcott) Ferrous sulfate, dried 162 mg, thiamine HCl 1 mg, riboflavin 1 mg, ascorbic acid 15 mg, nicotinic acid 4 mg, pyridoxine HCl D 5 mg, 6 Panthelol 0.47 mg, folic acid D 2 mg, vit B 12, cryit 1 mcg, liver fraction #2 162 mg / cap Bot 100s, 250s 1000s

Use: (1-3 daily) Nutritional deficiencies

NEMOTIN CAPSULES, (Rowell) Liver 5 gr, ferrous gluconate 5 gr, thiamine HCl 2 mg, riboflavin 2 mg, niacinamide 10 mg, folic acid 1.5 mg, Vit B 12 (activity equiv) 5 mcg / Cap Bot 100s 1000s

Use (1-2 t.i.d.) Nutritional anemias

NEMOTON, NND 1962 (Carrione) See: Ferrous fumarate

HEMOVITE, (Cowley) Cap Bot 100s, 500s, 1000s Tab Bot 100s, 500s, 1000s

Use: (1-2 t.i.d.) Hematinic

HEMO-VITOL (Smith, Mutter & Patch) Ferrous sulfate, exsic 200 mg, Vit B 12 activity 5 mcg, ascorbic acid 37.6 mg, folic acid 1 mg, liver desic 200 mg, gastric mucosa desic 100 mg / Tab Bot 100s 1000s

Use (1-2 t.i.d., m.p) Hematinic

HEMO-VITOL LIQUID. (C. O. Smith) Ferrous gluconate 100 mg., Vit. B-12 10 mcg., thiamine HCl 2 mg., cobalt 2.5 mg./5 cc. Bot. 1 pt.

Use: (2 tsp. p.c., t.i.d.) Anemias.

HEMO-VITONIN. (Buffington) Cap. Liver conc. 3 gr., colloidal iron peptonate 3 gr., Vit. B-1 1.5 mg., B-2 2 mg., B-6 0.1 mg., niacinamide 20 mg., Cal. pantothenate 1 mg./Cap. Bot. 84s, 500s. Liq. Liver conc. equiv. to 50 Gm. fresh liver, Vit. B-12 10 mcg., B-1 3 mg., B-2 2 mg., B-6 0.22 mg., pantothenic acid 1.2 mg., niacinamide 20 mg., folic acid 1 mg., colloidal iron peptonate 6.5 gr./fl. oz. Bot. 8 oz., 1 gal. Tab. Whole dried liver 3 gr., ferrous sulfate exsic. 3 gr., Vit. B-1 1.5 mg., B-2 2 mg., B-6 0.25 mg., Cal. pantothenate 1 mg., niacinamide 20 mg./Tab. Bot. 100s, 1000s.

Use: Secondary anemia.

HEMO-VITONIN PEDIATRIC. (Buffington) (Vit. B-1) 1 mg., B-2 1 mg., B-6 1 mg., B-12 5 mcg., niacinamide 6 mg., Cal. pantothenate 1 mg., folic acid 0.33 mg., iron peptonate 65 mg./5 cc. Bot. 1 pt.

Use: 1 dr. t.i.d.; anemias.

HEMO-VITONIN SUPER. (Buffington) Vit. B-1 2 mg., B-2 2 mg., C 50 mg., B-12 w/intrinsic factor conc. 1/2 u., B-12 5 mcg., liver desic. 100 mg., ferrous sulfate exsic. 150 mg., niacinamide 20 mg., folic acid 0.25 mg./tab. Bot. 100s.

Use: 1 t.i.d.; anemias.

NENASPHEN. (Drug Products) Acetaphenetidin 0.13 Gm., hyoscyamus pow. ext. 11 mg., aspirin 0.13 Gm., caffeine 16 mg., gelsemium pow. ext. 2.2 mg., w/aromatics/Cap. Bot. 100s, 1000s.

Use: (1-2 q. 3 h) Analgesic & antipyretic.

HINBANE. See: Hyoscyamus (Various Mfr.)

HEPA-B. (Pitman-Moore) Thiamine HCl, not less than 10 mg., liver sol., purified 10 U.S.P. u./cc. Vial, 10 cc.

Use: Pernicious anemias.

HEPACIDS. (Gold leaf) (liver fraction 20.2 Gm., ferrous sulfate exsic. 0.2 Gm., thiamine HCl 1.5 mg., riboflavin 2.5 mg., niacinamide 15 mg., ascorbic acid 50 mg./Tab. Bot. 50s, 100s, 1000s.

Use: (1-2 q. 4 h.) Hypochromic anemia.

HEPACOL. (Vitamax) Heparin sod. 2500 u., Vit. B-12, folic acid & niacin/cc. Vial 10 cc.

HEPA-DISICOL KAPSIALS. (Parke, Davis) di-Methionine 150 mg., choline bitartrate 200 mg., inositol 50 mg. & desicol (desicc. whole fresh bile) 150 mg./Cap. Bot. 100s, 1000s.

Use: (1-2 t.i.d.) For liver disease states.

HEPAFOLIN. (Gold leaf) Liver inj. 2 U.S.P.

u., folic acid 5 mg., Vit. B-12 15 mcg./cc. Amp. 1 cc., Box 6s, 25s, 100s. Vial 10 cc., Box 1s.

Use: Sprue & other macrocytic anemias.

HEPAFOLIN-X. (Gold leaf) Liver inj. 10 U.S.P. u., folic acid 5 mg., Vit. B-12 30 mcg./cc. Vial 10 cc.

HEPALEAF. (Gold leaf) Vit. A 2500 u., D 200 u., B-1 1 mg., B-2 1.5 mg., C 37.5 mg., niacinamide 10 mg./Cap. Bot. 100s, 1000s.

Use: Multivitamin therapy.

HIPAMIN. (Gold leaf) Vit. B-12 20 mcg., liver fraction #1 3 gr., iron & ammonium citrate 18 gr., Vit. B-1 10 mg., B-2 4 mg., nicotinamide 20 mg./fl. oz. Elixir. Bot. 6 oz., 12 oz.

Use: Hematinic in secondary anemias.

HEPAMINE. (Myers-Carter) Heparin 2500 u., choline 100 mg., Vit. B-12 15 mcg., folic acid 2 mg., niacinamide 50 mg./cc. Vial 10 cc.

HEPAMINE. (Ball Pharmacol) Choline chloride 75 mg., methionine 25 mg., B-1 25 mg., B-2 2 mg., niacinamide 100 mg., B-12 15 mcg., liver inj. 19 mcg./cc. Vial 10 cc.

Use: I.M., dietary supplement.

HEPALEXIN. (Gold leaf) (liver inj.) 100 mg., Vit. B-1 2 mg., B-2 0.3 mg., B-6 0.3 mg., niacinamide 50 mg./2 cc. Amp. 2 cc., Box 6s, 25s, 100s. Vial 10 cc., 30 cc.

Use: Nutritional & secondary anemias.

HEPARIN, N N D. 1960, A.O.R. 1955.

Harvey Labs.—Vial (10 mg./cc.) 10 cc.

Sherman—Vial (1000 u./cc) 10 cc.

HEPARIN ANTAGONIST.

See: Blytens (Abbott).

Polybrene, Amp. (Abbott)

Protamine Sulfate (Various Mfr.)

HEPARIN IN ABBJECT FORM. (Abbott) Heparin sodium 20,000 u./cc. In disposable syringe. 1 cc. Abboject disposable syringe. Box 5s.

Use: Anticoagulant therapy.

HEPARIN IN PITKIN MENSTRUUM. (Worner) Amp. 2 cc., 3 cc., Box 1s, 6s. Also w/ vasoconstrictors.

Use: Thrombosis.

HEPARIN IN SALINE. (Abbott) Isotonic saline 30 cc., heparin sod. 15 mg., or 1500 u., isotonic saline 30 cc., and heparin sod. 18 mg./cc. Bot. 500 ml. Abbo-Voc or Non-Vac.

***HEPARIN POTASSIUM, N N D. 1962.** See: Clarin, Sub. Tab (Leeming)

HEPARIN REPOSITORY INJECTION. (Led-erle) 20,000 U.S.P. u. (approx 200 mg./cc.) 1 cc. disposable syr. Pkg. 1s, 5s.

Use: (S.C., 20,000 to 40,000 u.) Prevention & treatment of thrombosis & embolism.

* Generic name recognized by N.F., U.S.P. or A.M.A.

***HEPARIN SODIUM, USP, NND 1962;**
In USP

Use: I.M., I.V. or S.C., anticoagulant in prevention & treatment of thrombosis or embolism (Note: protamine sulfate is antidote)

See Clarin, Sublingual (Leeming)

Dopa Heparin, Vial (Upjohn)

Hed Heparin, Vial (Hed)

Heparinate, Sol (Cy Labs)

Hepathram, Amp (Testagar)

Lipo Heparin, Vial (Darwin)

Liquemine, Vial (Organon)

Metarin, Vial (Bell Ph)

Panheprin (Abball)

W/Choline chloride, Vit B 12, folic acid & niacinamide

See: Hep Nine B, Vial (Columbus)

Hepamine, Vial (Myers Carter)

Hepavite, Inj (Mallard)

W/Choline chloride, B 12, B 6, inositol, dl methionine See: Hepatropic, vial (Central)

W/Choline chloride, Vit B 12 folic acid niacinamide, liver injection See Ro Co Lin (Paul Maney)

W/Neomycin sulfate polymyxin B sulfate hydrocortisone See: Oerme Parin, Oint (Broemmel)

W/Polymyxin B sulfate neomycin, hydrocortisone See Bro Parin susp (Broemmel Pharm)

HEPARINATE SOLUTION (Cy Labs) Sod heparin 10 mg, mutopolysaccharides 1% / cc Vial 10 cc

Use I.M., anticoagulant

HEPATHROM. (Testagar) Heparin sodium 1000 u equals 10 mg supplied: (1000 u/cc) Vials 10 cc; (5000 u/cc) Amps 1 cc, Vials 10 cc; (10,000 u/cc) Amps 1 cc Vial 4 cc; (20,000 u/cc) Vial 4 cc

HEPATINIC (McNeil) Ferrous sulfate 167 mg, liver conc 333 mg, Vit B-12 1 mcg, thiamine HCl 1 mg, riboflavin 2 mg, niacinamide 10 mg / 5 cc Each tab represents 5 cc of the elixir Elixir Bot 1 pt Tab, Bot 50s 100s

Use: (2 tsp or 2 Tab p.c. t.i.d.) Secondary anemias

HEPATONE (Merk Sharp & Dahme) Contains sod phosphate, lithium citrate, phenol phthalein & saccharin, sod bicarb, citric & tartaric acids Bot 5 oz

Use: Effervescent laxative compound

HEPATOPLEX. (G F Harvey) Thiamine HCl 500 mg, riboflavin 20 mg, pyridoxine HCl 5 mg, cal pantothenate 5 mg, niacinamide 1250 mg, Vit B 12 (cryst) 50 mcg, liver USP (at time of preparation) 50 w/Vial

*Generic name recognized by N.F. USP or A.M.A.

Pkg 2 elafs, 1 al dried material, other of diluent

Use: Anemias

HEPATROPIC. (Central) Heparin Sod 30 mg (3000 u) Vit B 6 5 mg, B 12 50 mcg, choline chloride 100 mg, inositol 50 mg, dl methionine 25 mg / cc Vials 10 cc

Use: Anticoagulant

HEPAVITE. (Mallard) Heparin Sod (2500 u, 25 mg, choline chloride 100 mg, Vit B 12 50 mcg, lactic acid 2 mg, nicotinamide 50 mg, benzyl alcohol 1.5% / cc Vial 10 cc

Use: Coronary vascular disease

HEPAZYN. (Endocrine) Betaine choline 150 mg, inositol 25 mg, dehydracetic acid 30 mg, ax bile 90 mg, liver 100 mg / Tab. Bot 100s

Use (1 2 t i d) Anemias

HEPBETA. (Columbus) Liver Injection crude 1 mcg, Vit B 12 5 mcg, folic acid 2 mg, Vit B 1 10 mg, niacinamide 50 mg, Vit B 2 0.1 mg Vit B 6 0.1 mg / cc. Vial 10 cc

Use I.M., anemias

HEPCOVITE. (Endo) Vit B 12 Amp. (15 mcg / cc) 1 cc Box 12s, 25s 100s; Vial (30, 100, 1000 mcg / cc) 5 cc, 10 cc. Drops (30, 25 mcg / cc.) Bot w/dropper 15 cc, 30 cc Tab (50 mcg), Bot 25s, 100s

Use I.M., orally; hematopoietic

HEP-FER-B CAPSULES. (Pitman Moore) Liver conc #1 #2 4 gr, ferrous sulfate 3 gr, thiamine HCl 0.15 mg / Cap Bot 100s 500s, 1000s

Use Anemias

HEPFERPLEX. (Gold Leaf) Liver inj 10 u / cc. 100 mg, iron hydroxide colloidal 191 mg, riboflavin 0.3 mg, niacinamide 50 mg, pyridoxine HCl 0.3 mg / 2 cc Amp 2 cc Vial 10 cc, 30 cc

Use: Nutritional deficiencies

HEPICEBRIN GELSEALS (Lilly) Thiamine chloride 2 mg, riboflavin 3 mg, ascarbic acid 75 mg, nicotinamide 20 mg, vit A 5000 u, D 400 u / cap Pkg 100s, 1000s

Use (1 t i d) Vitamin deficiencies

HEP-IRON W/Vit B (Norgine Labs) Ferrous citrate 10 gr, vit B 1 2 mg, B 2 2 mg, niacinamide 14 mg, Vit B 12 10 mcg, liver fraction #2 20 gr / fl oz Bot 8 oz, 1 pt, 1 gal

Use (4 6 tsp t i d) Hematinic

HEP-IRON DROPS. (Norgine Labs) Vit B 12 25 mcg, lactic acid 0.25 mg, iron and ammon citrate 120 mg / cc Bot w/dropper 15 cc, 30 cc

Use Hematinic

HEP-IRON ELIXIR (Norgine) Liver Traction 2 650 mg, ferric ammonium citrate 1 Gm

Thiamine hydrochloride 5 mg., riboflavin 3 mg., niacinamide 20 mg., Vit. B-12 15 mcg., Vit. B-6 1 mg., panthenol 2 mg., choline chloride 20 mg., inositol 10 mg., copper sulfate U.S.P. 1.6 mg., cobalt sulfate 1 mg., manganese citrate soluble 30 mg., alcohol content 12% by volume/fl. oz. Bot. 8, 16 oz.

Use: Hypochromic anemias.

HEPIVIN. (Filt, Eaton) Liver, desic. 3½ gr., ferrous sulfate 2 gr., Vit. B-1 1 mg., B-2 2 mg., B-6 0.5 mg., B-12 1 mcg., niacinamide 5 mg., Cal. pantothenate 0.5 mg. & intrinsic factor from dried hog duodenal mucosa/Cop. Bot. 100s, 500s, 1000s.

Use: (3 t.i.d.) Hematopoietic & hematinic w/vitamin B complex.

HEP-NINE B. (Columbus) Heparin Sod. 25 mg., choline chloride 100 mg., Vit. B-12 15 mcg., folic acid 2 mg., niacinamide 50 mg./cc. Vial 10 cc.

Use: (i.m., 1-2 cc. weekly) Heparin & lipotropic therapy.

HEP-NINE B FORTIS. (Columbus) Heparin sod. 50 mg., choline chloride 100 mg., Vit. B-12 50 mcg./cc. Vial 10 cc.

Use: Lipotropic therapy

HEPOVEX COMPOUND. (Lilly) Liver-stomach ext. 2.2 Gm., ferrous sulfate anhy. 0.4 Gm., thiamine HCl 0.6 mg., riboflavin 0.2 mg./3 Gm. Bot. 65 Gm. or 260 Gm.

Use: Anemias.

HEPOJEL. (Prema) Heparin sodium in gelatin 20,000 u./cc. Vial 2 cc.

Use: Anticoagulant.

HEPRON TABLETS. (Norgine Labs.) Liver fraction #2 600 mg., gastric substance w/ intrinsic factor 180 mg., vit. B-12 10 mcg., ferrous sulf. exsic. 600 mg., vit. A 10,000 u., D 1000 u., B-1 3 mg., B-2 5 mg., C 90 mg., niacinamide 30 mg./6 tabs. Bot. 50s, 100s, 500s.

Use: (1-2 t.i.d.) Hematinic.

***HEPTABARBITAL.** N.N.D. 1962. (Geigy) 5-(1-Cyclohepten-1-yl)-5-ethylbarbituric acid. See: Medamin (Geigy)

HEPTERYL-12. (Kremers-Urban) Vit. B-12 100 mcg., folic acid 5 mg., purified liver ext. 10 u./cc. Amp. 1 cc. Vial 10 cc. & 30 cc.

Use: Pernicious anemias.

HEPTERYL-12 (CRUDE). (Kremers-Urban) Vitamin B-12 2 mcg., folic acid 5 mg., in crude liver ext./cc. Vial 10 cc.

Use: Pernicious anemias

HEPTOFEB. (Endo) Vit. B-12 cryst. 15 mcg., thiamine HCl 10 mg., riboflavin 5 mg., nicotinamide 15 mg., liver deriv. 10 mg./cc. Amp. 1 cc., 10 cc.

Use: Anemias.

* Generic name recognized by N.F., U.S.P. or A.M.A.

HEPTOFEB. (Endo) Liver deriv. 62.5 mg., iron peptonate 12 mg./cc. Amp. 1 cc. Box 12s, 25s, 100s

Use: I.M. Secondary anemia

***HEPTOLAMIDE.** (Upjohn) 1-cycloheptyl-3-p-tolylsulfonyleurea

HEPTRON. (Lakeside) Liver Inj. purified 2 u., iron (as iron peptonate) 5 mg., copper (as copper sulfate) 0.14 mg., thiamine HCl 3.3 mg./cc. Vial 10 cc.

Use: Anemias.

HEPTRON. (Superior) Heparin sod. 25 mg., choline chloride 100 mg., vit. B-12 crystalline 50 mcg., folic acid 2 mg., niacinamide 50 mg./cc. Vial 10 cc.

Use: Atherosclerosis.

HEPTUNA. (Roerig) Ferrous sulfate 4.5 gr., Vit. A 5000 u., D 500 u., B-1 2 mg., B-2 2 mg., B-6 0.1 mg., Cal. pantothenate 0.333 mg., niacinamide 10 mg./Cap. Bot. 50s, 100s.

Use: Secondary anemias.

HEPTUNA PLUS. (Roerig) Vit. A 5000 u., D 500 u., B-1 2 mg., B-2 2 mg., B-6 0.1 mg., niacinamide 10 mg., calcium pantothenate 0.33 mg., B-12 5 mcg., folic acid 0.33 mg., C 50 mg., liver des. 150 mg., ferrous sulf. dried 199 mg., cobalt 0.1 mg., copper 1 mg., molybdenum 0.2 mg., calcium 37.4 mg., iodine 0.05 mg., manganese 0.33 mg., mg. 2 mg., phosphorus 29 mg., pot. 1.7 mg., zinc 0.4 mg./cap. Bot. 100s.

Use: Nutritional deficiency.

W/Folic acid. (Roerig) In addition to above contains folic acid 1.7 mg. Cap. Bot. 100s.

HEPTUNA PLUS IMPROVED. (Roerig) Iron 100 mg. (ferrous fumarate), Desiccated liver N.F. 50 mg., folic acid 0.5 mg., Vit. C 75 mg., Vit. B₁₂ oral Concentrate 5 mcg., Non-inhibitory intrinsic factor concentrate w/ activity equivalent to ½ U.S.P. oral unit, Vit. A 2500 u., D 250 u., B₁ 3 mg., B₂ 2 mg., B₆ 2 mg., calcium pantothenate 1 mg., Niacinamide 15 mg., and ten minerals/Cap. Bot. 100.

Use: Microcytic & Macrocytic anemias

HERMESOL. (Kohlenberg) Colloidal mercury biolate 2.5%. Sol., Bot. 1 fl. oz., 2 fl. oz., 16 fl. oz.

Use: Germicide, lungicide.

HEBON. See: Diacetylmorphine. (No Mfr. currently lists)

HESCOR. (Haug Drug Co.) Ascorbic acid 100 mg., hesperidin purified 100 mg./tab. Bot. 100s.

Use: Orally (2 t.i.d.), to maintain normal capillary function.

HESPER-C. (National Drug) Bioflavonoid hesperidin 100 mg., Vit. C 100 mg./Tab. or

dr Bot 100s, 1000s Liq Bot 4 oz, 12 oz
Use (3-4 t i d) Habitual abortion, rheuma-
toid arthritis diabetes hypertension.

HESPER-C AGIATRIC (National) Hesperidin
complex 100 mg, ascorbic acid 100 mg,
pancreatin 100 mg, dl methionine 100 mg
choline 125 mg, inositol 75 mg, diastase
20 mg/Cap Bot 100s
Use Geriatric

HESPER-C BITABS (National Drug) Hes-
peridin complex 200 mg and ascorbic acid
200 mg/Tab Bot 100s
Use (1 tab 3 or 4 times daily) Prevention
of capillary damage

HESPER-C PRENATAL (National Drug) Hes-
peridin complex 100 mg, Vit C 100 mg-
A 1000 u D 200 u, R1 125 mg, B 2
075 mg, nicotinamide 5 mg, Vit B 12
075 mcg folic acid 0.05 mg, Vit B 6 1.67
mg, col pantothinate 1 mg ferrous glu-
conate 21.6 mg cal carbonate 208.25 mg
copper sulfate 2 mg, pot iodine 0.065
mg/Cap Bot 100s, 500s
Use (2 t i d) Prenatal therapy

HESPERCONE (Testagar) Hesperidin Methyl
Chalcone 50 mg Vit C 50 mg/Tab Bot
100s, 1000s
Use Protected against capillary damage

HESPERGESIC (National Drug) Hesperidin
complex 100 mg, ascorbic acid 100 mg
guayanesin 25 mg, pyrilamine maleate
12.5 mg, methapyrilene fumarate 12.5 mg,
salicylamide 200 mg, ephedrine HCl 6.25
mg/Cap With or without codeine phos-
phate 6 mg/Tab Bot 30s, 100s
Use Treatment of the common cold

HESPERIDIN See Vitamin P; also Rutin
Use Capillary fragility and permeability

W/Ascorbic acid
See Caplaid Tab (Person & Covey)
Cevanoid, Tab (Kirkman)
Citric, Tab (Blue Line)
Flavinal C, Tab (Walker)
Hescar, Tab (Haug Drug Co)
Hesper C, Tab (National)
Hesperidin C, Cap & Tab (National Drug)
Hesperidin W/C, Cap (West Ward)
Inabex HC, Cap (Tilden)

**W/Adrenochrome semicarbazone, ascorbic
acid, sod menadiol diphosphate** See:
Adrestat, Cap, Laz (Organon)

W/Ascorbic acid See: Cebnoids, Cap
(Morton & Converse)

**W/Ascorbic acid choline bitartrate, dl
methionine, inositol pancreatin** See:
Hesper C Agiatric Cap (National)

**W/Ascorbic acid, guayanesin pyrilamine
maleate methapyrilene fumarate salicyl-
amide, ephedrine HCl, and codeine phos-**

phate See: Hespergesic, Cap (National
Drug)

**W/Ascorbic acid, phenyltoloxamine, aspirin,
acetophenetidin, caffeine.** See: Corycio,
Tab (Standard)

**W/Ascorbic acid, prophenpyridamine male-
ate, thienpyramine fumarate, pyrilamine
maleate salicylamide & caffeine.** See:
Tripac Cap (Person & Covey)

W/Ascorbic acid, rutin See: Ascorutin Bia,
(Arnar Stone)

**W/Ascorbic acid, salicylamide, pyrilamine
maleate, phenacetin, caffeine** See: Hesp-
ron, Cap (CMC)

**W/Diactyl sod sulfosuccinate, vitamins and
minerals** See: Sulfic, Tab (Smith Dorsey)

**W/Ethisterone, Vit K analogue, di-alpha
tocopherol acetate, ascorbic acid** See: Nu-
gesteral Tab (Organon)

**W/Pentylentetrazol, nicotinic acid, Vit C,
See Niatex, Tab (Mallory)**

**W/Phenacetin, salicylamide, caffeine, as-
corbic acid, pyrilamine maleate** See: Bical,
Cap (Bryant Pharmaceutical Corp)

**W/Phenylephrine HCl, chlorpheniramine male-
ate Vit C, aspirin** See: Calrex, Cap
(Rowell)

**W/Phenylephrine HCl, prophenpyridamine
maleate methapyrilene HCl, pyrilamine
maleate vit C, salicylamide acetophene-
lidin, caffeine** See: Bayle Citra, Caps
(Boyle)

W/Piperidolate HCl, ascorbic acid See: Doc-
in OB, Tab (Lakeside)

**W/Pal para-amino benzoate, pot salicylate,
Vit C** See: Flavisal, Tab (Columbus)

**W/Prednisone, salicylamide ascorbic acid,
Vit K, alum hydrax gel** See: Holvicort,
Tab (Haller)

**W/Pyrilamine maleate, thienpyramine fu-
marate, chlorprophenpyridamine maleate
vitamin C, phenylpropanolamine HCl, sali-
cyclamide caffeine anhydrous** See: Tripac
DC Tab (Person & Covey)

W/Reterpine, ascorbic acid See: Flavoserp,
Cap (Blue Line)

W/Vitamins See: Preablin, Tab (Person &
Covey)

HESPERIDIN-C. (National Drug) Hesperidin
50 mg, ascorbic acid 50 mg/Cap or Tab
Cap, Bot 100s, 500s, 1000s Tab, Bot
100s, 500s, 1000s

Use (1 2 q 4 h) Capillary fragility or
permeability

HESPERIDIN W/C. (West Ward) Hesperidin
100 mg ascorbic acid 100 mg/cap Bot
100s, 1000s

Use: Capillary fragility or permeability

HESPERIDIN RIOFLAVONOID W/Salicyl-
amide, acetophenetidin, caffeine prophen-

- pyridomine maleate & ascorbic acid. See Caryban, Cap. (Roerig)
- HESPERIDIN CHALCONE SODIUM.** See Pev-Gram (Ingram).
- W/Ascorbic acid. See: Pecegram, Cap. (Ingram).
- HESPERIDIN METHYL CHALCONE.** (Lilly)
Pulvules 50 mg. Bot. 100s, 500s.
Use: Deficiency of vitamin P.
W/Ascorbic acid. See: Carperidin, Cap. (Dumas-Wilson)
- W/Diethylstilbestrol, ascorbic acid, and menadiol. See: Nulabart, Tab. (Dumas-Wilson)
- W/Dihydrocadinone bitartrate, Vit. C, pheniramine maleate, pyrilomine maleate, pot citrate. See: Citra Forte. (Boyle)
- W/Pipradol HCl, vitamins, lipotropics, iron, essential minerals. See: Peptolin, liq. (Walker Labs.)
- W/Vitamin C. (Lilly) Hesperidin methyl chalcone 50 mg. & ascorbic acid 100 mg./Pulvule. Pulvule, Bot. 100s, 500s.
Use: [1-2 i.l.d.] Capillary defects caused by deficiency of vitamins P & C.
See: Hespercane, Tab. (Testagar)
- W/Citrus bioflavonoids complex, Vit. C & mucopolysaccharides. See: Hesperinate-C, Cap. (Cy Labs.)
- HESPERINATE-C.** (Cy Labs.) Hesperidin methyl chalcone 10 mg, citrus bioflavonoids complex (Vit. P complex) 25 mg, Vit. C 50 mg., mucopolysaccharides 10 mg./Cap. Bot. 50s.
Use: [1 i.l.d.] In diseases associated with capillary fragility.
- HESPER-VITE.** (Lowitt Labs) Vit. C 200 mg, A 5000 u., D 500 u., B-1 3 mg., B-2 3 mg., B-12 5 mcg., B-6 2 mg, hesperidin complex 200 mg, niacinamide 20 mg., calcium pantothenate 5 mg., iodine 15 mg, potassium 5 mg., copper 1 mg, manganese 6 mg., zinc 1.5 mg., menadione 1 mg./Tab Bot. 60s.
Use: Vitamin and mineral supplement.
- HESPYRON.** (CMC) Pyrilamine maleate 25 mg, sacilylamide 3 1/2 gr., phenacetyl 2 1/2 gr., caffeine 1/2 gr., ascorbic acid 50 mg, hesperidin purified 50 mg./Cap. Bot. 30s, 100s, 250s, 1000r
Use: Common cold.
- HETAMINE.** (Dumas-Wiltron) d-amphetamine sul. 5 mg./Tab. Bot. 100s, 1000s
- HETERIL.** (Harrar) Non-specific antigen from sheep erythrocyte & spleen. Vial 10 cc.
Use: Non-specific therapy.
- HETOXIN.** (Warner-Chilcott) Hydroxyquinoline benzoate 0.3%, benzoic acid 0.2%, boric acid 3%, talcum diluent. Container, 3 oz.
Use: Antireptic dusting powder.
- HETRAZAN.** U.S.P., N.N.D. 1962. (Lederle) (Diethylcarbamazine) 1-Diethylcarbamyl-4-methylpiperazine dihydrogen citrate. Syr. 120 mg./5 cc. Bot. 1 pt. Tab. 50 mg Bot. 100s.
Use: Anthelmintic, specific for certain parasites, including flukes & ascarides.
- NEVITRON TABLETS.** (United Lab.) Ferrous gluconate 3 gr., Vit. B-12 2 mcg., B-1 4 mg., B-2 2 mg., C 20 mg., niacinamide 10 mg., liver fraction # 2 5 gr., copper sulfate 2.5 mg./Tab. Bot. 100s.
Use: [1-2 t.i.d.] Hematinic.
- NEXA-RETALIN.** (Lilly) Pyridazine HCl, Amp. (50 mg./cc.) 1 cc., Vial 5 cc., (100 mg./cc.) 5 cc. & 10 cc. Tab. 10 mg. Bot. 100s; 25 mg., 50 mg. Bot. 10s, 100s.
Use: Vitamin B-6 therapy.
- HEXARORATE WITH ATROPINE AND SOD. RENZOATE.** (Warren-Teed) Atropine sulf. 0.12 mg, methenamine 0.2 Gm., sod. benzoate 0.12 Gm., & sod. borate 0.12 Gm./Tab. Bot. 1000s
Use: Urinary antiseptic
- HEXACHLOROCYCLOHEXANE, N.N.D. 1962**
See: Benzene Hexachloride, U.S.P.
- *NEXACHLOROPHENE, U.S.P., N.N.D. 1962.**
A.D.R. Bis[2-hydroxy-3,5,6-trichlorophenyl]-methane
See: Almederm, Cream (Meyer Chem.)
Aveeno Baby Pow. (Faugero)
Dial Soap. (Armour)
Fostri, Lot. (Washwood)
G-11 (Sindar)
Gamaphen (Ethicon)
Germa-Medica (Huntington)
Hex-O-Son (Retart Ph. Co)
Septisol (Yertal)
Surgi-Cen (Central)
- W/Alantain, Coal Tar Extract. See: Sebical Cream (Reed & Carnrick)
- W/Hydrocortisone free alcohol, tyrothricin. See: Acnestat (Dale)
- W/Menthol, glycerin & Isopropyl alcohol. See: Appliderm, Lot. (White Labr.)
- W/Oatmeal Colloidal, kaolin, and sulfur. See: Aveeno, Lot. Pow. (Aveeno Corp)
- W/Phenol, Resorcinol, Oil of Tor rectified, Zinc Oxide. See: Retafen. (Vanpett & Brown)
- W/Polyoxyethylene Lauryl ether, Hydrocortisone, Sulfur. See: Fostri HC. (Wertwood)
- W/Prednisolone. See: Lipolone, Cream, (Spir)
- W/Pyridazine HCl, pantothenyl alcohol &

menthol See: VI Cuta, Skin Cream (Columbus)

W/Salicylic acid, sulfur, sodium diacetyl sulfoxuccinate, sodium alkyl polyether sulfonate, sodium sulfacetate, Kerahydric See: Sebulex, Lq (Westwood)

W/Silicone See: Perfection, Lot (Suliff & Case)

W/Silicone cod liver oil & zinc oxide See: Hollandex Silicane Oint (Holland Rankos)

W/Sod octylphenoxymethyl ether sulfonate See: phisoHex Lq (Winthrop Labs)

W/Sulfur, alcohol, zinc, titanium and iron oxides, koolin and talc See: Liquimat (Texas)

W/Sulfur resorcinol isopropyl alcohol See: Cenac lot (Central)

W/Sulfur, salicylic acid See: Sebacin (Summers)

W/Tyrosinacin sulfur resorcinol, zinc oxide, alcohol See: Eez Acne-care, aerosol (Pfizer)

W/Zinc undecylenate, undecylenic acid See: Undex Improved (Vonpelt & Brown)

***HEXACHLOROPHENE LIQUID SOAP, USP**

See Gamophen Surgical Soap (Ethicon Suture)

Hexachlorophen Germa Medica Liquid Surgical Soap (Huntington Lab)

Hexachlorophen Liquid Soap (J I Holcomb Lab)

Hex-O San Surgical Soap (Retort Pharm Co)

phisoHex (Winthrop Lab)

Sepsisol with Hexachlorophene (Vestol Inc)

Surgi Cen Liquid Soap (Central)

HEXACIDE, (Central) Zinc undecylenate 8%, undecylenic acid 2% hexachlorocyclohexane 1% methylbenzethonium chloride 2% Oint 1 oz, 1 lb

Use: Topically, scabicide & pediculicide

HEXACOSE (Lincoln) Mixture of C-6 alcohols derived from oxidation of tetraosane— $C_{24}H_{50}$ See: Hexathricin Aeropak (Lincoln)

HEXACYCLONATE SODIUM Sodium 3,3-pentamethylene 4-hydroxybutyrate Treatment of alcoholism Under study

HEXADECADROL See: Decadron (Merck Sharp & Dahme)

HEXADENOL See: Hexacase (Lincoln)

HEXADERM-H C CREME (Federal Pharmaceutical) Hydrocarbons free alcohol 0.5%, hexachlorophene 0.5%, menthol 0.125% Tube 1/2 oz

Use: Acute dermatosis

HEXADIENOL See: Hexacase (Lincoln)

*Generic name recognized by N F USP or A.M.A.

***HEXADILINE**, (Merrill) 2(2,2 Dicyclo Hexyl vinyl) piperidine

Use: Coronary dilator

***HEXADIMETHRINE BROMIDE**, N N D 1962 1,3 dimethyl 1,5 dioxoundecamethyl

see methobromide See: Polybrene AMP (Abbott)

HEXADROL (Organon) Dexamethasone D.75 mg / Tab Bot 100s

Use: Rheumatoid arthritis

***HEXAFLUORENIUM BROMIDE**, Hexo-methylene bis [9 fluorenyldimethylammonium] dibromide See: Myloxen (Irwin, Neisler)

HEXAFLUORODIETHYL ETHER. See: Indoxon

HEXAFORM, (Central) Methenamine 5 gr / Tab Bot 100s, 1000s

Use: urinary antiseptic

HEXAGELS, (Walker Vitamins) Vit A 2,500 u, B 1 1 mg, B 2 1.5 mg, C 37 mg, D 200 u, nicotinamide 10 mg / Cap Bot 100s, 1000s

Use: (1 2 t.l.d.) Dietary supplement

HEXA-GERM, (Huntington) Hexachlorophene 3%, In Iodofin and pet base Lq Bot 1 gal

Use: Skin antiseptic cleanser

HEXAHYDROXYCYCLOHEXANE. See: Inatol, Prep (Various Mfr)

HEXAKOSE, Mixture of tetraosanes & oxidation products See: Hexathricin, Aeropak (Lincoln)

W/Benzethonium chloride, p chloro-m-xylene ethyl p aminobenzoate & tyrosinacin See: Hexathricin, Aeropak (Lincoln)

HEXALOID (Lilly) Methenamine 5 gr, 7 1/2 gr Pkg 100s 1000s

Use: (1 2 q 3 h) Urinary antiseptic

HEXALOL, (Central) Methenamine 30 mg, methylene blue 6 mg, atropine sulfate 0.03 mg, hyoscyamine 0.12 mg, benzoic acid 300 mg, salal 300 mg / Tab Bot 100s, 1000s

Use: Genito-urinary emollient and anodyne

***HEXAMARIUM BROMIDE** (Hexomethyl ene bis(3 pyridyl N methylcarbamate) dimethyl bromide

HEXAMETHONIUM W/Rauwiloid See: Rauwiloid w/hexamethonium, Tab (Riker)

HEXAMETHONIUM BROMIDE, N N D 1958 Vegoflysen Hexamethylene (bistrimethyl ammonium) bromide See: Bistilium Bromide (Squibb)

***HEXAMETHONIUM CHLORIDE**, N N D 1962 Hexamethylene (bistrimethyl ammonium) chloride

See: Bistrium Chloride High Potency N N D (Squibb)

Exonid Chloride N N D (Ciba)

- Hexhex Chloride, N.N.D.** (Kellih-Victor)
Methium Chloride, N.N.D. [Warner-Chilcott]
 W/Reserpine. See: Hexatina, Tab. (Testagar)
HEXAMETHYLENAMINE, U.S.P., N.N.S. 1949.
 See: Methenamine (Various Mfr.).
- HEXAMETHYLENAMINE MANDELATE, N.N.D. 1961.** See: Mandelamine (Warner-Chilcott)
- HEXAMETHYLENAMINE COMPOUND ELIXIR.** See: Farmatrapin (Abbott).
- HEXAMETHYLENAMINE SILVER.** See: Argalol, Oint. (Daho)
- HEXAMETHYLENETETRAMINE, U.S.P., N.N.R. 1957.** See: Methenamine, U.S.P., N.N.R. Hexamethylenetetramine Mandelate, N.N.D. 1961.
- HEXAMETHYLPARAROSANILINE CHLORIDE.** See: Bismuth Violet, Oint. (Tablerock)
- HEXAMETHYLOSANILINS CHLORIDE.** See: Gentian Violet.
- HEXAMINE.** See: Methenamine (Various Mfr.).
- HEXANASTAB.** See: Hexobarbital Soluble (Eliol Sed) (Winthrop Labs.)
- HEXANICOTOL.** (Phladelphia Lab) Nicotinic acid and Inositol/200 mg. eac. 200 mg cap. Bot. 30s, 100s
 Use: For ecna.
- HEXANITROL.** (Drug Products) Mannitol hexanitrate 32 mg./Cap. Bot. 100s, 1000s. Also available w/phenobarbital.
 Use: Hypertension.
- HEXAPHOS.** (Central Pharm.) Methenamine 4 gr. w/disodium hydrogen phosphate, Mag. oxide & citric acid/isp. Pow., Pkg. 2½ oz.
 Use: Urinary antiseptic.
- ***NEXAPRADOL HCl.** (Mead Johnson) α-[1-aminohexyl]benzhydrol HCl.
 Use: Central nervous system stimulant.
- HEXA-RUTALIN.** (Columbus) Mannitol hexanitrate 25 mg., theophylline 45 mg., Vit. C 20 mg., butabarbital cap. 15 mg., Resperidin conc. 20 mg./Cap. Bot. 100s, 500s, 1000s
 Use: (1-2 daily) Vasodilator w/sedative.
- HEXARUTAN.** (C. B. Kendall) Aminophylline 50 mg., mannitol hexanitrate 16 mg., rufin 20 mg., phenobarbital 16 mg./Tab. Bot. 100s, 500s.
 Use: Cardiovascular disease associated with hypertension.
- HEXATNAL SODIUM.** See: Ortal Sodium, Cap. (Parke, Davis).
- HEXATHRICIN AEROPAK.** (Lincoln) Hexacose 8 Gm., benzethonium chloride 80 mg., p-chloro-m-xylene 800 mg., ethyl p-aminobenzoate 1.6 Gm., tyrothricin 40 mg./6 oz. Aeropak 3 oz., 6 oz.
 Use: Topically, burns, pyoderms & fungal infections.
- NEXATHRICIN Otic.** (Lincoln) Tyrothricin 10 mg., benzocaine 200 mg., p-chloro-m-xylene 400 mg., benzethonium chloride 10 mg./10 cc. Bot. 15 cc., 25 cc., 1 pt.
 Use: Fungal infections of ear.
- HEXATINA.** (Testagar) Hexamethonium chloride 125 mg., reserpine 0.1 mg./tab. Bot. 100s, 1000s.
 Use: Treatment of hypertension.
- HEXATONE.** (Upjohn) Methenamine 2 gr., Sod. salicylate 1½ gr., Sod. benzoate 1½ gr., sobal 2½ gr., carn silk 2½ gr., triticum 2½ gr., buchu 2½ gr./5 cc. Elixir, 1 pt., 1 gal.
 Use: (1-2 tsp. q. 4 h.) Urinary antiseptic.
- NEXAYERINE.** (Lincoln) Ethaverine HCl 25 mg., mannitol hexanitrate 20 mg., phenobarbital ½ gr./Tab. Bot. 100s, 1000s.
 Use: (1-2 q. 4 h.) Antispasmodic & vasodilator.
- NEXAVIBEX.** (Parke, Davis) Pyridoxine HCl, Amp. (50 mg.) 1 cc., Box 6s, 25s, Kapsals 10 mg. Bot. 20s, 100s. Vial (100 mg./cc.) 5 cc.
 Use: Vitamin B 6 therapy.
- ***NEXAVITAMIN, N.F., N.N.R. 1955.** Vit. A 5000 u., B-1 2 mg., B-2 3 mg., C 75 mg., D 400 u., nicotinamide 20 mg./Tab. or Cap. Bot. 100s, 1000s.
 Merrell—Tab. Bot. 100s, 1000s.
 Pfizer—Tab. Bot. 100s, 1000s.
 Fream—Tab. Bot. 100s, 1000s.
 Walker—Cap. Bot. 100s, 1000s.
 See: Avicap (Burroughs Wellcome)
 Mepcevirin (Lilly)
 Polyvitin (Winthrop Lab.)
- NEXAVITAMIN TABLETS.** Holt N.P. Strength. (Merrell) Vit. A 2500 u., B-1 1 mg., B-2 1.5 mg., C 37 mg., D 200 u., nicotinamide 10 mg./Tab. Bot. 100s
- ***NEXCARBACHOLINE BROMIDE.** 1,6-Nexamethylenebiscarbaminaylcholine bromide.
- NIXENE-DL.** See: Hexacose (Lincoln).
- NIXENOL.** See: Hexacose (Lincoln).
- NEXESTROPEN-1** (Merrell) Hexestrol 1 mg., phenobarbital 15 mg./Tab. Bot. 100s, 1000s.
 Use: Estrogen therapy
- NEXESTROPEN-3.** (Merrell) Hexestrol 3 mg., phenobarbital 15 mg./Tab. Bot. 100s, 1000s.
 Use: Estrogen therapy.
- ***NEXESTROL, N.F., N.N.D. 1962;** Inj., N.F.; Tab., N.F. (Dihydro-diethylstilbestrol, Hexaestrol) Mesa-3,4-di(p-hydroxyphenyl)-n-hexane.

* Generic name recognized by N.F., U.S.P. or A.M.A.

HEXETHAL SODIUM

Lantene—Tab 0.25 mg, 1 mg, 3 mg,
Bot 100s 1000s

Massengill—Tab 3 mg, Bot 100s 500s
1000s

Merrill—Tab 1 mg, 3 mg, Bot 100s
1000s

See: Extra Plex Vag Cream (Rowell)
Use (200 mcg to 3 mg i.M.) Estrogen
therapy

W/Phenobarbital

See Hexestrolen, Tab (Merrill)

Hexital Tab (Ortho)

Hexypheen Tab (Lantene)

W/Phenobarbital (Massengill) Hexestrol 3
mg phenobarbital 1/2 gr Tab Bot 100s
1000s

W/Pentobarbital scopolamine HBr See: Tal
Estomine Tab (Walker Corp)

W/Sulfacetamide sulfa meth dia mer See:
Vagi Plex, Cream (Rowell)

HEXETHAL SODIUM, N.N.R. 1954 A.D.R.
(Mabral) Sod ethyl hexylbarbiturate See:
Oral Sodium (Parke Davis)
Use Sedative & hypnotic

***HEXETIDINE**, N.N.D. 1962 5 Amino 1,3
bis (beta ethylhexyl) 5 methylhexahydro
pyrimidine Triacil See Sterisal Vag Gel
(Warner Chilcott)

HEXITAL TABLETS (Ortho) 3 mg of the
synthetic estrogen Hexestrol combined with
20 mg (1/2 gr) phenobarbital/Tab Bot
100s 1000s 5000s

Use Estrogenic & sedative

HEXOBARBITAL, A.D.R.

See Evipal (Winthrop Labs)

Sombulex Tab (Schenley)

W/Dihydrohydroxycodone HCl, dihydro
hydroxycodone Ierephthalate homotro
pine Ierephthalate, aspirin phenacetin
caffeine See Percobarb Cap (Endo)

W/Dihydrohydroxycodone Ierephthalate,
dihydrohydroxycodone HCl homotropine
Ierephthalate aspirin, phenacetin caffeine
See Percobarb Dem (Endo)

***HEXOBARBITAL SODIUM**, N.F. N.N.D.
1961; Sterile N.F. (Cyclonal Sod, Dorco
Soluble Evipan Sod Hexanostab Hexo
barbitone Sod, Methenexyl Sod) N methyl
5 cyclohexenyl 5 methylbarbiturate See
Evipal Sodium Prep (Winthrop Labs)

HEXOBABBITONE SODIUM See: Evipal
Sodium, Prep (Winthrop Labs)

***HEXOCYCIUM METHYLSULFATE**, N.N.D.
1962 N (beta Cyclohexyl beta hydroxy
beta phenylethyl) N methylpiperazine di
methylsulfate See: Trol Filmtabs & Gradu
mets (Abbott)

W/Etylurea See Trolcyon Tab (Abbott)

* Generic name recognized by N.F. U.S.P. or A.M.A.

354

HEXOESTROL. See: Hexestrol (Various Mfr.).

***HEXOPROPYIMATE**, (Mead Johnson) 1
propargyl 1 cyclohexyl carbamate

HEXOPYRROLATE (Robins) 1 methyl 3
pyrrolidyl a phenyl cyclohexaneglycolate
methobromide

Use Anticholinergic

***HEXOPYRROLONIUM BROMIDE**; See: Hexo
pyrrolate

HEX-O-SAN, N.N.D. 1959 (Retort Ph Co)
Soap containing 2% hexachlorophene (an
hy bases) Can 379 L & 189 L; Drum
567 L 1135 L, 208 L L

Use: A soap with local anti infective agent

HEXODE (Mallard) Hexamethylenimine 3
gr, Sod acid phosphate 3 gr, hyoscyamus
ext 0.125 gr, triticum ext 0.5 gr, adiphe
nine HCl 0.154 gr/Tab Bot 100s 500s
1000s

Use (1 2 q 3 4 h) Urinary antiseptic, seda
tive & relaxant

***HEXYLCALINE HYDROCHLORIDE**, N.F.
N.N.D. 1962, Inj, Soln, N.F. 1 Cyclohexyl
amino-2 propylbenzoate HCl See Cycloine
HCl (Merck Sharp & Dohme)
Use Spinof & local anesthetic

***HEXYLRESORCINOL**, U.S.P., N.N.R. 1934;
Pill, U.S.P. (Various Mfr.)

See Caprotol, Cap Jelly & in oil (Merck
Sharp & Dohme)

Crystoids Anthelmintic (Merck Sharp &
Dohme)

ST 37 Sol & Tooth Paste (Merck Sharp &
Dohme)

Secrets (Merck Sharp & Dohme)

W/Benzocaine & benzoic acid See: Benzo
Hexyl Loz (Rowell)

W/Benzocaine, camphor, menthol See: Au
ristil (Blue Line)

W/Tannic acid jelly See: Hexylton Jelly
(Merck Sharp & Dohme)

HI-A-VITA (Brewer) Vitamin A Capsules
50 000 u Bot 25s, 100s, 1000s
Amp (50,000 u/cc) 1 cc, Box 6s, 12s
100s

Amp. (100 000 u/cc) 2 cc, Box 6s, 12s
100s

Use Vitamin A deficiency

HI-BISTROL (Bayle) Diethylstilbestrol Tab
25 mg Bot 100s, 1000s

Use: (1 t i d) Estrogen therapy

HIBITANE LOZENGES, (Ayerst) Chlorhexi
dine dihydrochloride (16 d: 4 chlorophe
nyl diguamidohezone di HCl) 5.0 mg Benz
ocaine 25 mg /Loz Pkg 12s

Use: Moderate infections or irritations of
the mouth and throat

HI-CAPS (Frost) Diglycolol hydriodide

- Equiv. of 4 cc. Syr. HI/Cap. Bot. 50s, 100s, 500s
 Use: Hydriodic acid therapy.
- HI-COMPLEX.** (Kirk) Vit. B-1 30 mg., niacinamide 50 mg., Vit. B-6 2 mg., riboflavin 0.5 mg./cc. Amp. 10 cc., 30 cc.
 Use: Vitamin B complex.
- HI-COMPLEX, FORTE.** (Kirk) Vit. B-1 100 mg., riboflavin 1 mg., niacinamide 120 mg., pyridoxine 4 mg., Cal. pantothenate 4 mg., Vit. B-12 15 mcg./cc. Amp. 1 cc., Box 12s, 25s; Vial 10 cc., 30 cc.
 Use: Vitamin B complex therapy.
- HICOMPLEX.** (Blue Line) Thiamine HCl 0.5 mg., riboflavin 1 mg./Tab. Bot. 100s, 1000s.
 Uses Vitamin B-1 & B-2 deficiencies.
- HI-DERATOL.** (Brewer) Vit. D, activated ergosterol. Amp. (200,000 u./cc.) 1 cc., 6s, 12s, 100s (400,000 u./cc.) 2 cc., 6s, 12s, 100s (600,000 u./cc.) 1 cc., 6s, 12s, 100s. Cap. 200,000 u. Bot. 50s, 100s.
 Use: Vitamin D therapy.
- HIDEX W/Iron.** (Winthrop Labs) Oestrins, dextrose, maltose w/Iron. Pow., Can 1 lb.
 Use: Infant food.
- HI-DOXIN.** (Dumas-Wilson) Tab. Bot. 100s, 1000s.
- HIESTROHE.** See: Estrone (Various Mfr.).
- HIGADOCE FORTE ELIXIR.** (Burlington) Vit. B-12 25 mcg., liver conc. 2 Gm., folic acid 2 mg., thiamine chloride 10 mg., riboflavin 5 mg., pyridoxine HCl 1 mg., Cal. pantothenate 2 mg., niacinamide 50 mg., iron peptonized 0.5 Gm., alcohol 12%/fl. oz. Bot. 1 pt., 1 gal.
- HIGADOCE FORTE TABLETS.** (Burlington) Vit. B-12 10 mcg., folic acid 0.5 mg., ascorbic acid 50 mg., stomach sub. desicc., including pepsin portion 50 mg., whole dried liver 150 mg., ferrous sulfate exsic. 100 mg./Tab. Bot. 100s, 1000s.
- HINKLE PILL FORMULA, ORIGINAL.** See: Cascara Compound w/strychnine (Merck Sharp & Dohme)
- HINKLE'S PILL FORMULA, N.F.** See: Atain, Belladonna, Cascara & Podophyllum Tab (Merrell).
- W/Strychnine.** See: Atain, Belladonna & Podophyllum W/Strychnine (Merrell).
- HINKLE'S PILLS.** See: Cascara Compound (Merck Sharp & Dohme)
- HINTON INDICATOR.** (Lederle) Cardiolipin, lecithin & cholesterol in an alcoholic sol. Amp. 5 cc.
 Use: Diagnostic agent for syphilis.
- HIOHEX CHLORIDE, N.D. 1959.** (Keith-Victor) Hexamethanum chloride. Tab (125 & 250 mg.) Bot. 24s.
- HIPERNINE.** [Westerfield] Nitroglycerin 1/100 gr., Sod. nitrate 2 gr. w/lobelia & pilocarpine/Tab. Bot. 100s, 1000s.
 Use: Orally, vasodilator.
- HIPPUSAN, N.N.R. 1954.** (Mollinckradt) o-tolohippurate, Sod. Am. Sol. 25 cc. Pow., Bot. 12 Gm., 100 Gm., 500 Gm.
 Use: Radiopaque medium for urography.
- HI-PRO.** (Jackson-Mitchell) High protein & low fat milk. Pow., Can 1 & 2 1/2 lb.
 Use: Infant food.
- HISCAPHEH.** (Ulmer) Aspirin 4 gr., phenacetin 2 gr., ephedrine HCl 1/12 gr., pyrilamine maleate 1/5 gr./Tab. Bot. 100s, 500s, 1000s.
 Use: Analgesic & antihistaminic.
- HISDIN.** (Massengill) Semikan HCl (methapyrilamine HCl) 50 mg., semoxydrine HCl (methamphetamine HCl) 2.5 mg./Tab. Bot. 100s, 500s, 1000s.
 Uses Antihistamine effect.
- HISPRIIL, N.N.D. 1962.** (Smith, Kline & French) Diphenylpyraline HCl Spansule 5 mg. Bot. 30s, Tab. 2 mg. Bot. 50s.
 Use: Antihistamine.
- HISTABAR "Plus" & "Plain".** (Medco) Pylamine maleate, 25 mg.; metapyrilamine HCl, 25 mg.; phenpyridamine maleate, 10 mg./Cap. Bot. 30s, 100s. "Plus" has dextroamphetamine HCl, 6 mg. added.
 Uses Manifestations in allergic states.
- HISTABORT.** (Lincoln) Anisapradamine. Tab. 25 mg., 50 mg. Bot. 100s, 1000s.
 Use: Antihistamine.
- HISTABORT FORTIFIED.** (Lincoln) Pylamine maleate. (Anisapradamine) 30 mg., lisompheta (amphetamine sulfate) 2 1/2 mg./tab. Bot. 100s, 1000s.
 Use: Antihistamine w/control nervous system stimulant.
- HISTACAP.** (Ethical Pharm) Pylamine maleate 75 mg./Timed disintegration tab Bot. 30s
 Use: Antihistaminic
- HISTA-CARB APC.** (Caritane) Chlorpheniramine maleate 2 mg., aspirin 230 mg., phenacetin 160 mg., caffeine 30 mg./Tab. Bot. 100s.
 Use: Relief of cold symptoms
- HISTACHIOR.** (Vitamins Corp) Chlorpheniramine maleate 4 mg./Tab & Cap Bot. 200s, 1000s Also with APS, also timecaps 8 mg. or 12 mg.
- HISTA-CLOPANE.** (Lilly) Pulvules: Histadyl 25 mg., etopone HCl 12.5 mg./Pulv. Bot. 100s, 1000s Sol; Histadyl 0.5% & etopone HCl 0.5%. Bot. 1 fl. oz., 1 pt.
 Uses Allergic symptoms of hay fever
- HISTACOMP SYRUP.** (Approved Pharm) Thienpyramine HCl 80 mg., ammonium chloride 10 gr., Sod. citrate 5 gr., antimony

- Pot tartrate 1/24 gr, menthol & aromatics q s / fl oz Sol, Syr 4 oz, 8 oz, 1 gal
Use Allergic & asthmatic coughs
- HISTACOMP TABLETS.** (Approved Pharm)
Pyrimilamine maleate 25 mg, aspirin 3 1/2 gr phenacetin 2 1/2 gr, caffeine 1/2 gr / Tab Bot 12s 30s, 100s
Use (1 q 3 h) Allergic symptoms & the common cold
- HISTACON** (Marsh Emory Labs) Chlorpheniramine maleate 12 mg, ephedrine hydrochloride 1/4 gr atropine sulfate 1/250 gr / Tab Bot 100s
Use Antihistaminic
- HIST-A-CORT-E CREME** (Dome) Vit A 100 000 u estrane 36 000 u / oz, pyrimilamine maleate 20 mg / Gm hydrocortisone (free alcohol) 5 mg / Gm, in acid mantle creme Creme Tube 0.5% or 1% hydrocortisone 1/2 1 oz lot 1/2 % Bot 2 oz
Use: Anti-inflammatory, antipruritic
- HISTA-DERFULE** (Cale) Thenylpyramine HCl 25 mg, ipecac & opium pow 1/4 gr, acetophenetidin 1 1/2 gr, atropine sulfate 1/500 gr, camphor monobromated 1/2 gr, salt cyclamide 2 gr, caffeine 1/2 gr / Cap Bot 100s, 500s, 1000s Available in half strength
Use: Analgesic
- HISTADYL HYDROCHLORIDE, NND 1962** (Lilly) (Methapyrilene Hydrochloride)
Amp: (Free base 20 mg / cc) 1 cc Box 6s, 100s, 10 cc
Cream: Salt 2% HCl Tube 1 oz; Jar 1 lb, 3 lb
Enterals: Free base 50 mg Bot 100s, 500s, 1000s
Let: Free base 2% Bot. 4 oz, 1 pl
Nasal Salt: Free base 0.5% Bot 1 oz, 1 pl
Ophth. Oint: Free base 0.5% Tube 1/2 oz
Pulvules: 25 mg, 50 mg, 100 mg, free base Bot 100s, 1000s
Syr: Fumerate salt 6 mg / cc Bot 1 pt, 1 gal
Use: Antihistaminic.
- W/Calamine & zinc oxide See: Enzo-Col AH (Crookes).
W/Clonidine HCl See: Hista Clonidine, Cap, Sol (Lilly).
W/Clonidine HCl & pyranil See: Co Pyranil, Preps (Lilly)
W/Sulfacaine & titanium dioxide See: Sur-tadyl, Lotion (Lilly)
- HISTADUR DURA-TABS.** (Wynn) Chlorpheniramine maleate 8 mg / Tab Bot 30s 100s 250s 12 mg / Tab Bot 30s, 100s 250s
Use: Relief of allergic conditions
- HISTADYL & A.S.A. COMPOUND.** (Lilly) Histadyl 25 mg, acetophenetidin 2 1/2 gr, acetylsalicylic acid 3 1/2 gr, caffeine 1/2 gr / Pulvule Bot 100s, 1000s
- HISTADYL E.C.** (Lilly) Cadeine phosphate 1 gr, ephedrine HCl 1/2 gr, thenylpyramine fumarate 1 1/2 gr, ammonium chlorido 10 gr, chloralform 1 min w/menthol & flavor R oz Bot 1 pt, 1 gal
Use (1 tsp q 3 4 h) Coughs & colds
- HISTADYL & EPHEDRINE HCl.** (Lilly) Histadyl 25 mg ephedrine HCl 8 mg / Pulvule Bot 100s, 1000s
Use (1 2 q 3 d h) Hay fever & allergic rhinitis
- HISTADYL & SURFACAIN** (Lilly) Histadyl 2% & Sulfacaine 0.5% Cream 1 oz, 1/2 lb, 1 lb, 5 lb
Use Combined antihistaminic & anesthetic effect
- HISTADYL & SURFACAIN LOTION** (Lilly) Histadyl 2 Gm, sulfacaine 0.5 Gm, zinc oxide 8 Gm, calamine 8 Gm / 100 cc Bot 4 oz, 1 pt
Use: Antihistaminic w/local anesthetic
- HISTATED.** (Cale) Thenylpyramine HCl 25 mg ephedrine HCl 8 mg, ascorbic acid 10 mg / Cap Bot 100s, 500s, 1000s
Use: Antihistaminic
- HISTALME DURACAP5** (Meyer) Pyrimilamine maleate 75 mg, hyoscyamine HBr 0.12 mg, hyoscyamine HBr 0.024 mg, d-amphetamine sulfate 2.3 mg / TD cap Bot 30s, 100s
Use: (1 2 fld) Motion sickness and antihistaminic
- HISTALOG, NND 1962** (Lilly) 3 Beta aminoethylpyrazole diHCl Betazole HCl Ampul 50 mg / cc Pkg 6s
Use: I.M., S.C.; diagnostic aid for carcinoma at the stomach or peptic ulcer
- HISTALOG-SD** mg (Lilly) Betazole HCl 3 Beta amino ethylpyrazole dihydrochloride 50 mg / cc Amp 1 cc
Use: (50 200 mg I.M.) Diagnostic for gastric secretory capabilities
- HISTALOH.** (Hance) Pyrimilamine maleate 25 mg / Tab Folders 1Bs; Bot 50s, 1000s
Use: Antihistaminic
- HISTAMINASE.** Diamine oxidase See: Torontol, Tab (Winthrop Labs)
- HISTAMINE.** 2 (4 imidazole) ethylamine
Use: Diagnostic agent
- W/Ergotaxine & tyramine See: Ernufin Amp & liq (Burroughs Wellcome)
- HISTAMINE ACID PHOSPHATE, U.S.P.** See: Histamine Phosphate, U.S.P. (Various Mfr)
- HISTAMINE DIHYDROCHLORIDE W/Acetyl** glycol salicylic acid ester, methyl salicylate menthol & thymol See: Imadyl, Oint (Roche)
W/Methyl nicotinate, glycol monosalicylate

capsicum oleoresin. See: Rubiguent, Cream (Wyleh).

***HISTAMINE PHOSPHATE, U.S.P., N.N.R. 1957;** Inj. U.S.P. beta-Iminozolyethylamine-acid phosphate (Histamine acid phosphate) Burroughs Wellcome—Amp. (1 mg./cc.) 1 cc.

Parke, Davis—Amp. 1:1000/1 cc. Box 12s, 100s.

See: Histapon, Viol. (Endo)

Use: S.C., I.V., allergy therapy & diagnostic aid.

HISTAMINE PHOSPHATE SOLUTION. (Don Baxter) Histamine phosphate 2.75 mg. in 250 cc. of 0.9% sod. chloride solution

Use: I.V., therapeutic agent

HISTAR. (Sherman) Pyramisamine maleate 25 mg./Tab. Bot. 100s, 1000s, 6000s

Use: Antihistaminic.

HISTANTIN. See: Chlorcyclizine

HISTA-PAC. (Hance) Pyrilamine maleate 25 mg., aspirin 3½ gr., acetophenelidin 2½ gr., caffeine ½ gr./Tab. or Cap. Bot. 12s, 100s, 1000s.

Use: (1 l.l.d.) Antihistamine-analgesic.

HISTA-PACQ. (Horton & Converse) Histapyran (Pyrilamine maleate) 25 mg., phenacetin 2 gr., aspirin 3 gr., caffeine ½ gr., quinine sulfate ½ gr./Tab. Bot. 18s.

Use: (1 l.l.d. or q. 3 h) Cold symptoms.

HISTAPAH. (Lynn) Chlorphenirpyridomine maleate, 4. mg + phenylpropanolamine HCl 25 mg + vit. C, 50 mg + acetylsalicylic acid, 5 gr/Tab. Bot. 50's

Use: Common cold and allergies.

HISTAPAH TIMED. (Lynn) Chlorphenirpyridomine maleate, 10 mg; phenylpropanolamine HCl, 50 mg /timed disintegrating cap. Bot. 50s, 250s.

Use: Common cold and allergies.

HISTAPON. (Endo) Repository Injection of histamine phosphate 2.75 mg./cc. in peanut oil and aychotesteral derivatives. Vials 5 cc. Box 6s.

Use: (I.M. on graduated schedule, 0.05 cc. to 0.5 cc) Multiple sclerosis and acute allergies.

HISTAPYRAN. (Horton & Converse) Pyrilamine maleate 25 mg. or 50 mg/Tab. Bot 100s, 500s. Inj. (10 mg./cc.) Vial 10 cc.

Use: Antihistamine therapy

HISTAPYRAN COMPOUND. (Horton & Converse) Pyrilamine maleate 25 mg., aspirin 3½ gr., phenacetin 2½ gr., caffeine ½ gr./Tab. Bot. 1000s

Use: Antihistamine therapy and treatment of the common cold

HISTAOIN. (Taylor) Pyrilamine maleate 25 mg., salicylamide 3 gr., phenacetin 2 gr.

* Generic name recognized by N.F., U.S.P. or A.M.A.

caffeine ½ gr., atropine sulf. 1/600 gr., guanine sulf ½ gr., ascorbic acid 25 mg/ Cap. Bot. 100s, 1000s

Use: Analgesic-antihistaminic, in the treatment of the common cold

HISTAR. (Reed & Cornrick) Pyrilamine maleate 2% w/special coal tar ext. (Tarbanis Brand) 5% in a hydrophilic base. Jar. 2 oz., 1 lb.

Use: Allergic skin therapy.

HISTASCORB. (Nalick) Pyrilamine maleate 25 mg., Sod. ascorbate 175 mg., iodine 5 mcg., Vit. R-1 7 mg., B-2 2 mg., niacin 5 mg/Tab. Bot. 56s, 500s, 1000s

Use: (1 q.l.d.) Hay fever, urticaria, pruritus & rhinitis.

HISTASCORRINE. (Torign) Histidine HCl 4%, ascorbic acid 500 mg/5 cc. Amp. 5 cc., Box 12s, 25s, 100s.

Use: Peptic & duodenal ulcers & peripheral vascular diseases.

HISTA-TRANE. (Cole) Tyrothricin 2 mg., benzocaine 5 mg., phenylpyramine HCl 5 mg./lozenge. Bot. 100s.

Use: Antibiotic & antihistaminic throat lozenges.

HISTA-VADRIN. (First Texas) Phenylpropanolamine HCl 40 mg., chlorpheniramine maleate 4 mg., methoppyriline HCl 40 mg., phenylephrine HCl 5 mg./Tab. Bot. 100s, 1000s.

Use: Antihistaminic-Decongestion.

***HISTIDINE MONOHYDROCHLORIDE, N.F. X₁ Inj., N.F.** beta-4-imidazolyl-1-amino propionic acid HCl.

Endo—Amp. (0.2 Gm.) 5 cc., Box 6s, 25s, 100s.

Kremers-Urban—Amp. (0.2 Gm.) 5 cc., Box 25s.

Lakeside—Amp. (0.2 Gm.) 5 cc., Box 25s.

Miller—Amp. (0.2 Gm.) 5 cc., Box 6s, 50s.

National Drug—Amp. (0.2 Gm.) 5 cc., Box 6s, 12s, 25s, 100s.

C. D. Smith—Amp. (0.2 Gm.) 5 cc., Box 5s, 25s, 100s.

Use: I.M., peptic & jejunal ulcers.

I-NISTIDINE MONOHYDROCHLORIDE.

See: Larastidin, Amp. (Roche).

W/Ascorbic acid. See: Histascorbine, Amp. (Torign).

W/Choline, methionine, tryptophane, and B complex vitamins. See: Amvitaf, Amp. (Walker)

HISTONEX. (Strassenburgh) Phenyltoloxamine resin 25 mg. or 50 mg/cap. Bot. 100s.

Use: Antihistaminic.

HISTITRIN. (Vitarine) Chlorpheniramine mol-

cate 8 or 12 mg /Cap T D Bot 30s, 100s
Tab 4 mg Bot 100s, 1000s
Use Antihistamine

HISTIVITE-D (Vitarin) Dextromethorphan
HBr 30 mg, chlorafarm 2 min, methapyril
ene fumarate 75 mg ammonium Cl 8 gr,
ephedrine sulfate $\frac{3}{4}$ gr, menthol alcohol
(by volume) 4.8% /oz Bot 4 oz
Use Coughs due to colds

HISTO-CLOPAHE. (Lilly) Histodyl 25 mg,
clopone HCl 12.5 mg /Cap Bot 100s
Use (1 1 d) Antihistamine

HISTONIUM (Columbus) Zirconium oxide
3%, phenyltolazamine dihydrogen citrate
1%, colamine 3% camphor 0.15%, phenyl
1%, benzocaine 1% Bot 2 oz
Use: Poison ivy and poison oak

HISTOPLASMIN (Parke Davis) Histoplasma
capsulatum filtrate Vial (0.01 cc) w/diluent
Use: Diagnostic aid in testing for histo-
plasmiasis

HIST-OXA-MINE (Walker Casp) Methapy-
riene HCl 25 mg, d desoxyephedrine HCl
1.25 mg /Tab Bot 100s
Use: Antihistaminic with central nervous
system stimulant

HISTOSTAB * Antazoline

HISTUSSIN (Meyer) Codeine phosphate 10.9
mg, psyllium maleate 12.5 mg, Pot
guaiacac sulfonate 169 mg, hyoscine HBr
0.005 mg, hyoscyamine HBr 0.025 mg,
chlorafarm 0.333 min, alcohol 10%, men-
thol 0.62 mg /dr Bot 3 oz, 1 pt, 1 gal
Use: [1 tsp q 3 h] Expectorant

HI-TEST (Mylord) Test reagent combining
ant body to human fibrinogen with polysty-
rene particles. Reacts specifically with
human fibrinogen. Only 1 drop of whole
blood is necessary for test. Kits of 6 tests
Use: Rapid slide determination of fibrinogen
blood deficiency

HI-VI-TOL (Patch) Fish liver oil containing
Vit A 5000 u, D 500 u /Cap Bot 30s,
100s, 1000s 1/4 Vit A 30,000 u, D 3000
u /Gm Bot 10 cc, 30 cc
Use: Orally, dietary supplement

H-M-C, No. 1 HYPDERMIC TABLETS (Ab-
bott) Hyoscine HBr 1/100 gr, morphine
HBr $\frac{1}{4}$ gr, cactus ext 1/60 gr /Tab Tube
20s, Bot 100s, 500s

H-M-C, No. 2 HYPDERMIC TABLETS (Ab-
bott) Hyoscine HBr 1/200 gr, morphine
HBr $\frac{1}{4}$ gr cactus ext 1/120 gr /Tab Tube
20s, Bot 100s, 500s, 1000s

HOLADIN, (No Mfr currently lists) Pancre-
atic gland est

Use: Pancreatic deficiencies

HOLAGOGUM HERB TEA (Overseas) Ex-
tracts of peppermint and fennel cara

* Generic name recognized by NF USP or A.M.A.

way, curcuma and calamus 2 X 110 Gm
containers

Use: Treatment of flatulence

HOLLANDEX SILICOHE OINTMENT, (Hal-
land Kantos) Silicone, Norwegian cod liver
oil zinc oxide hexachlorophene & Iodine
Tube 1 oz, 2 $\frac{1}{4}$ oz Jars 1 lb

Use: Topically antiseptic & protective

HOLOCAINE HYDROCHLORIDE, N N R
1950 Phenacaine HCl

Abbott—Ophth Oint 2% Tube $\frac{1}{2}$ oz

Lilly—Ophth Oint 2% Tube $\frac{1}{4}$ oz.

Winthrop Labs—Pow, Vial 5 Gm Bot 1
oz

Use: Topically local anesthetic

HOLOCAINE & EPINEPHRINE, (Abbott) Holo-
caine HCl 1%, epinephrine HCl 2%
Ophth Oint Tube $\frac{1}{2}$ oz

Use: Topically, local anesthetic

HOLOCAINE & MEYAPHEN, (Abbott) Holo-
caine HCl 2%, metophen 13000 Ophth
Oint Tube $\frac{1}{2}$ oz

Use: Topically local anesthetic with anti-
septic

HOMACHOL, (Lemmon) Sod pentobarbital
8 mg homatropine methylbromide 2.5 mg,
dehydrocholic acid 60 mg, ox bile ext
0.15 Gm /Tab Bot 100s

Use: Biliary dysfunction

HOMADONNA (Van Pelt & Brown) Pheno-
barbital $\frac{1}{2}$ gr, homatropine methylbromide
1/26 gr Tab or tsp Elix Bot 1 pt; Tab
Bot 100s, 500s, 1000s

Use: Orally, sedative with antispasmodic

HOMADRIH, (Crabbe Barnes) Homatropine
HBr 5% hydroxyamphetamine HBr 1%/
lacrival 15 cc

Use: Mydriasis Cycloplegia.

HOMAGEHETS ADORAL (Massengill) Vit A
acetate 50,000 u /Tab Bot 30s

Use: Vitamin A deficiencies

HOMAGEHETS GERIATRIC (Massengill) Vit
A 5000 u, D 3 1000 u, B 1 3 mg, B 2 2
mg, B 6 1 mg, nicotinamide 10 mg, cal
pantothenate 2 mg, ascorbic acid 37.5 mg;
Vit B 12 2 mcg, cal 10 mg, cobalt, copper,
iron, radine, pot, mag, mang, molyb,
zinc /Tab Bot 50s

Use: Geriatric

HOMAGEHETS-PEDIATRIC (Massengill) Vit
A 5000 u, D 3 1000 u, B 1 2 mg, B 2 2
mg, C 37.5 mg, B 12 2 mcg, B 6 1 mg,
nicotinamide 10 mg, cal pantothenate 2
mg /Tab Bot 40s

Use: Dietary supplement

HOMAGEHETS-PREHATAL (Massengill) Vit
A 2500 u, D 3 250 u, B-1 1 mg, B 2 1 mg,
B-6 1 mg, B 12 2 mcg, C 25 mg, nicotina

vide 5 mg., iron 10 mg., cal. carbonate 188 mg./Tab. Bot. 100s.

Use: Dietary supplement.

HOMAGENETS-THERAPEUTIC. (Massengill)

Vit. A 12,500 u., D-3 1000 u., B-1 5 mg., B-2 5 mg., B-6 0.5 mg., B-12 2.5 mcg., C 37.5 mg., nicotinamide 25 mg./Tab. Bot. 50s.

Use: Vitamin deficiencies.

HOMAPHINE. (C. D. Smith) Phenobarbital 15 mg., homatropine methylbromide 1/150 gr./Tab Bot. 1000s.

Use: Orally, sedative with antispasmodic

HOMAPIH. (Mission) Homatropine methyl-

bromide, No. 1; Tab. 5 mg., No. 2; Tab.

2.5 mg. w/phenobarbital 1/4 gr. No. 3;

Tab. 5 mg. w/phenobarbital 1/4 gr. No.

4; Homatropine methylbromide 10 mg.,

phenobarbital 1/4 gr. Bot. 100s, 500s, 1000s

Use: Orally, antispasmodic.

* **HOMARYLAMINE HYDROCHLORIDE.** N-

Methyl-3,4-methylenedioxyphenethylamine

HCl. W/Zinc bacitracin, lysothricin, neomy-

cin sulf., benzocaine. See: Pentazels Troches,

Vial. (Merck Sharp & Dohme)

HOMATAL. Homatropine Methylbromide 1/4

gr., Phenobarbital 1 gr./Tab. S.M. Bot. 30s,

100s, 250s. See: Dura-Tab S.M.

HOMATHYN. (Flint, Eaton) Homatropine

methylbromide 2.5 mg. (1/25 gr.), pheno-

barbital 16 mg. (1/4 gr.)/Tab. Bot. 100s,

500s, 1000s.

Use: Antispasmodic & sedative.

HOMATRAL. (Central) Phenobarbital 6 mg.,

homatropine methylbromide 0.4 mg./Pala-

tabs. Bot. 100s, 1000s

Use: Antispasmodic and sedative

HOM-ATRO-BARB. (Filtan-Moore) Not

listed Phenobarbital 16 mg., homatropine

methylbromide 2.5 mg./Tab. Bot. 100s, 500s.

Use: Orally, sedative with antispasmodic.

HOMATROMIDE. (Blue Line) Homatropine

methylbromide. Tab. 2.5 mg. Bot. 100s,

1000s.

Use: Orally, antispasmodic.

HOMATROPHEN. (Central) Phenobarbital 20

mg., homatropine methylbromide 2.5 mg./

Tab. Bot. 100s, 1000s

Use: Antispasmodic and sedative

HOMATROPINE HYDROBROMIDE, U.S.P.

(Various Mfr.).

Use: 1-3% for mydriatic & cycloplegia.

HOMATROPINE HYDROBROMIDE. (Crookes-

Barnes) 2% and 3% solution/Bot. 4 cc.

and 15 cc.

Use: Mydriasis.

W/Hydroxyamphetamine HBr. See: Homad-

rin, Oph. Soln. (Crookes Barnes)

* Generic name recognized by N.F., U.S.P. or A.M.A.

HOMATROPINE HYDROCHLORIDE, N.N.R.
1950.

Alcon—(2% sol.) Isopto-dropper. Bot.

15 cc.

Burroughs Wellcome—Tabloid aphth. 1/40

gr. Tube 12s.

Lilly—Tabloid aphth. 1/2 gr. Tube 10s.

Mallinckrodt—Cryst. Bot. 5 gr., 15 gr.,

1 oz., 18 oz.

Merck—Cryst. Vial 5 gr., 15 gr.

Parke, Davis—Tab. aphth. 1/2 gr. Tube

10s. Bot. 100s.

Use: Topically, mydriatic & cycloplegic.

HOMATROPINE HYDROCHLORIDE & CO-

CAINE HYDROCHLORIDE. Tabloid, ophth.

1/50 gr. each. Tube 25s. (Burroughs Well

come).

Use: Topically, local anesthetic with

mydriatic.

* **HOMATROPINE METHYLBROMIDE, U.S.P.,**

N.N.R. 1957; Tab., U.S.P. (Novatropine,

Methatropin, Comatropine)

Amer. Drug Prods.—Tab. 1/24 gr., 1/12

gr. Bot. 1000s.

Mallinckrodt—Pow. Bot. 1/2 oz., 1 oz., 4 oz.

Merck—Pow. Bot. 1 oz., 5 oz.

See: Homapln, Tab. (Mission).

Homatromide, Tab. (Blue Line).

Malcotran, Tab. (Maltale)

Metapin, Elix., Pow. & Tab., N.N.R. (Endo).

Novatrin, Tab., N.N.R. (Campbell).

Sed-Tems, Tab. (Lemman)

Sethyl, Tab. (Harrower).

W/Codeine phosphate, pyrilamine maleate,

ammonium Cl, sodium citrate. See: Endo-

tussin-C (Endo)

Homatropine Methylbromide Combina-

tions on pp. 360, 361, 362 and 363;

Homatropine Methylbromide & Pheno-

barbital Combinations on pp. 363 and

364.

HOMATROPINE TEREPHTHALATE W/Di-

hydroxycodolone, dihydrohydroxycode-

lone terephthalate & pentylene-tetra-

zol. See: Nucodan, Tab. (Endo)

W/Dihydrohydroxycodolone HCl, dihydro-

hydroxycodolone terephthalate, hexobar-

bitol, aspirin, phenocelin, caffeine. See:

Percobarb, Cap. (Endo)

HOMATROPINE TEREPHTHALATE W/Di-

hydrocodolone Bitartrate, Dihydrocodol-

one Terephthalate, Pentylene-tetra-

zol. See: Para Nucodan, (Endo)

HOMESAL. (Lemman) Mephenesin 0.2 Gm.,

Solycylamide 0.15 Gm., Homatropine meth-

ylbromide 2 mg./Tab. Bot. 100s

Use: Central ad pain and spasm

HOMICEBRIN. (Lilly) Thiamine chloride 1

mg., riboflavin 1.2 mg., Vit. B-12 (activity

Homatropine Methylobromide Combinations

| Homatropine Methylobromide | Others | Manufacturer | Brand |
|----------------------------|--|------------------|--------------------------------|
| 0.125 mg. | Bromisovalerylurea 300 mg., hyoscyamine HBr 0.125 mg., scopolamine HBr 0.0065 mg. | Tablerock Labs. | Tadana, Tabs. |
| 2.5 mg. | Butabarbital sodium 10 mg. | Zemmer | Butathyl Tablet or Elksir |
| 1.25 mg. | Butazolidin 100 mg., aluminum hydroxide 100 mg., mag. trisilicate 150 mg. | Geigy | Butazolidin Alka, Cop. |
| 2.5 mg. | Butisal sod. 10 mg., mag. hydroxide 0.3 Gm. | McNeil | Bunesia, Tab. |
| 1/100 gr. | Dehydrocholic acid 2 gr. | Blue Line | Atracholin, Tab. |
| 5 mg. | Dehydrocholic acid 250 mg. | Malibie | Chelon V, Tab. |
| 2.5 mg. | Dehydrocholic acid 4 gr. | Lakeside | Chalotropin, Tab. |
| 2.5 mg. | Dehydrocholic acid 250 mg., phenobarbital 9 mg. | Malibie | Chelon HMB, Tab. |
| 2.5 mg. | Dehydrocholic acid 60 mg., ac bile ext. 150 mg., pentobarbital sod. 8 mg. | Lemman | Hamachol, Tab. |
| 1.2 mg. | Diadol 1/4 gr., mephenesin 5 gr., N-acetyl-p-aminophenol 5 gr. | Buffington's | Steril-Ere, Tab. |
| 1.5 mg. | Dihydrocodeinone bitartrate 5 mg. | Endo | Hycodan, Syr./5 cc. or Tab. |
| 1.5 mg. | Dihydrocodeinone bitartrate 5 mg., chlorpheniramine maleate 2 mg., phenylephrine HCl 10 mg., N-acetyl-p-aminophenol 250 mg., caffeine 30 mg. | Endo | Hycomine Compound, Tab. |
| 1.5 mg. | Dihydrocodeinone bitartrate 5 mg., pyrilamine maleate 12.5 mg., amn. chloride 60 mg., sod. citrate 8.5 mg. | Endo | Hycomine, Syr./5 cc. |
| 1 mg. | Ergotamine tartrate 1 mg., caffeine 100 mg. | Cole | Ergolotropin, Tab. |
| 0.13 mg. | Estrone sulfate Sod. 0.42 mg., thyroid 8 mg., phenobarbital 16 mg. | Columbus | Men-Sed, Tab. |
| 1 mg. | d-Glucose 4.5 Gm., | Purdue Frederick | Probiogel/5 cc., liq. |
| 0.5 mg. | Glycine 180 mg., col. carbonate 420 mg. | Schenley | Titrolac-SP, Tab. |
| 2.5 mg. | Hydrocarisone acetate 5 mg., sod. salicylate 250 mg., mephenesin 250 mg. | Crookes-Barnes | Mephalac W/Hydrocarisone, Tab. |
| 1.512 mg./cc. | Hyoscyamine HBr 0.283 mg./cc. | Sherman | Navodanno, Sol. |
| 0.2 mg. | Magnesium trisilicate 360 mg., aluminum hydroxide gel 240 mg., gastric mucin 120 mg. | Money | Gast-Rex Tablets |
| 1.25 mg. | Magnesium trisilicate 150 mg., aluminum hydroxide gel 100 mg., prednisone 1.25 mg., phenylbutazone 50 mg. | Geigy | Sterozolidin, Cop. |
| 2.5 mg. | Man-'al hexanitrate 30 mg., veratrum viride 75 mg., Rauwolfia serpentina whole root 50 mg. | Crookes-Barnes | Rau-Parlanol, Tab. |
| 1/50 mg. | Mephenesin 4 gr., physostigmine salicylate 1/120 gr. | Columbus | Mephylin, Tab. |

Homotropine Methylobromide Combinations

| Homotropine Methylobromide | Others | Manufacturer | Brand |
|-------------------------------|--|-----------------------|---------------------------------|
| 2.5 mg. | Profenil citrate 60 mg., butobarbital sod. 7.5 mg. | C. D. Smith | Fentropine, Tab. |
| 2.5 mg. | | C. D. Smith | Fentropine W/autobarbital, Tab. |
| 2.5 mg. | Profenil HCl 45 mg., urethane 100 mg. | Smith, Miller & Patch | Fentropine Inj./cc. |
| 1.2 mg. | Pyrimilone maleate 30 mg., pamabrom 50 mg., hyoscyamine sulf. 0.1 mg., scopolamine HBr 0.02 mg., methamphetamine HCl 1.5 mg. | Central | Neoparbel, Tab. |
| 0.5 mg. | Reserpine, 0.1 mg., amn. chloride 500 mg., caffeine 30 mg. | Haack | Amiens, Tab. |
| 1/20 gr. | Secobarbital Sod. 1/2 gr., Cal. carb. 3 1/2 gr., charcoal 3 1/2 gr. | Roswell | Via-Alta-Phen, Cap. |
| 2.5 mg. | Sucetylulofathazole 3 Gm., Koolin 3 Gm., pectin 0.45 Gm. | Chicago Pharm. | Sulfased, Susp./Oz. |
| 2.5 mg. | Thyroid 1/2 gr., w/vitamins | Kenwood | InferTab-F, Tab. |
| 0.5 mg. | W/Vit. B complex | Purdue Frederick | Pre-Mens, Tab. |

Homatropine Methylobromide & Phenobarbital Combinations

| Homatropine Methylobromide | Phenobarbital | Others | Manufacturer | Brand |
|-------------------------------|---------------|--------------------------|-----------------|-----------------------------|
| 5 mg. | 16.2 mg. | | Chicago Pharm. | Bromaspaz, Tab. |
| 1/4 gr. | 3 mg. | | Klinney | Coactyn, Sol. |
| 10 mg. | 1 gr. | | Wynn | Duro-Tab S.M., Tab. |
| 1/26 gr. | 1/4 gr. | | Vanpelt & Brown | Hamadonna, Elixir & Tab. |
| 5 mg. | 1/4 gr. | | Filini, Eaton | Hamathyn, Tab. |
| 0.5 mg. | 8 mg. | | Maitbie | Malcatron w/Phenob., Elixir |
| 2.5 mg. | 20 mg. | | Central | Hamatropen, Tab. |
| 10 mg. | 65 mg. | | Ballard | Matrafen Dispellets |
| 2.5 mg. or 5 mg. | 1/4 gr. | | Endo | Mesopin-PB, Tab. |
| 1/50 gr. | 1/4 gr. | | C. D. Smith | Hamaphine, Tab. |
| 1.3 mg. | 16 mg. | | Campbell | Navatrin Compound |
| | | | | Fediatric/cc. |
| 2.5 mg. | 1/4 gr. | | Pitman-Moore | Hom-Altra-Barb, Tab. |
| 5 mg. | 45 mg. | | Brewer | Spatial, Tab. |
| 1/50 gr. | 1/6 gr. | Aluminum hydroxide 5 gr. | Irwin, Neisler | Hamatropol, Tab. |

Homatropine Methylbromide & Phenobarbital Combinations

| Homatropine Methylbromide | Phenobarbital | Others | Manufacturer | Brand |
|---------------------------|---------------|---|------------------|-------------------------|
| 1/50 gr | 1/4 gr | Alum num hydroxide 5 gr | Hart | Hartgel H M B Tab |
| 1/24 gr | 1/4 gr | Aluminum hydroxide 7 1/2 gr. Mag trisilicate 2 1/2 gr | Bassmar | Alased, Tab |
| 2.5 mg | 8 mg | Aluminum hydroxide gel 129.6 mg. Mag trisilicate 194.4 mg | Blue Line | Neutratabs, Tab |
| 2 mg | 1/6 gr | d Amphetamine sulfate 2 mg mepheneese 300 mg | Adams | Relaxamine, Tab |
| 2.5 mg | 8 mg | Dehydrocholic acid 250 mg | Malibie | Cholan HMB Tab. |
| 2.5 mg | 8 mg | Dehydrocholic acid 250 mg | Chicago Pharm | Hytraphen Tab |
| 1/48 gr | 1/8 gr | Dehydrocholic acid 2 gr. ox bile 2 gr | Nargine Labs | Bilamide Tab |
| 1 mg | 8 mg | Dihydroxy aluminum aminoacetate 0.5 Gm | Flint Eaton | Hamaldran, Tab |
| 1/100 gr | 1/4 gr | Dihydroxy aluminum aminoacetate 7 1/2 gr | First Texas | Alkam Comp. Tab |
| 1/50 gr | 1/8 gr | C 25 mg | Patch | Delzinate Tab |
| 0.13 mg | 1/4 gr | Estrone sulfate Sod 0.42 mg thyroid 8 mg | Columbus | Mena Sed Tab |
| 0.567 mg | 1/4 gr | Hyocyanine HBr 0.106 mg. hyosceae HBr 0.007 mg | Sherman | Navadannal, Tab |
| 0.0447 mg | 1/4 gr | Hyocyanine HBr 0.54 mg. scopalamine HBr 0.0144 mg thiamine HCl 3 mg | Paul Maney | Sedlin Unoday Cap |
| 5 mg | 1/4 gr | Kaolin 300 mg | Malibie | Lusyn Tab |
| 2.5 mg | 1/4 gr. | Mannitol hexanitrate 30 mg. veratrum viride 100 mg | Crookes Barnes | Pertenal, Tab |
| 1 mg | 8 mg | Mannitol hexanitrate 16 mg. veratrum viride 20 mg | Stoddard | Verante, Tab |
| 1.2 mg | 8 mg | Mepheneese D 3 Gm hyoscyamine sulfate 0.1 mg. scopalamine HBr 0.02 mg | Central | Mydanna, Tab |
| 1 mg | 8 mg | Methylcellulose 500 mg | Chilcatt | Airalase, Tab |
| 1/50 gr | 1/8 gr | Ox bile 1.5 gr. oxidized bile acids 1.5 gr. steapum 1/4 gr | Columbus | Oxochalin, Tab |
| 2.5 mg | 8 mg | Pentobarbital Sod 8 mg | Barer | Precibus, Tab |
| 1/480 gr | 1/4 gr | Pentobarbital Sod 1/16 gr | Walker | Sedadrapt, Liq |
| 1/480 gr | 1/4 gr | Pentobarbital Sod 1/16 gr | Walker | Sedaplex, Elixir & Tab. |
| 1/480 gr | 1/4 gr | Pentobarbital Sod 1/16 gr | Jackson Mitchell | Salsed Pediatric, Liq |
| 2 1/2 mg | 7 1/2 mg | Pentobarbital 7 1/2 mg | Meyer | Almased, Elixir |

- equivalent) 3 mcg., ascorbic acid 60 mg., Vit. A 3000 u., Vit. D 1000 u./5 cc. Bot. 60 cc., 120 cc., 1 pt.
 Use: Vitamin deficiencies.
- * **HOMOCHLORCYCLIZINE.** (Abbott) 1-(p-chlorobenzhydryl)-4-methylhomopiperazine.
- HOMOCILLIN.** (Rexall) Procaine penicillin G 300,000 u., Pot. penicillin G 100,000 u./dose. Vial 1 dose.
 Use: Penicillin therapy.
- HOMOPENT.** (Lemmon) Pentobarbital Sod. ¼ gr., homatropine methylbromide 1/25 gr., Mag. trisilicate 5 gr./Tab. Bot. 100s
 Use: Gastric antacid & antispasmodic.
- * **HOMOSULPANILAMIDE HCl.**
 See: Sulfamylon HCl (Winthrop Labs.)
 Sulfbenzamide HCl (Flint, Eaton)
- HOMVITE.** (Carrone) Thiamine hydrochloride 5 mg., pyridoxine hydrochloride 5 mg., nicotinic acid 20 mg., vit. B-12 25 mcg., L-lysine hydrochloride 300 mg./15 cc.
 Use: Geriatric nutritional supplement.
- HONEY CILLIN 400.** (Physicians Products) Penicillin G, buffered, 4,800,000 u./60 cc. 400,000 u./5 cc. Bot. 60 cc.
 Use: Penicillin therapy.
- HORMALE.** (Key Corp.) Testosterone propionate in oil. Vial (25 mg./cc.), (50 mg./cc.) 10 cc.
 Use: Sublingual (5-20 mg) androgen therapy.
- HORMALE LINGUALS.** (Key Corp.) Methyltestosterone 10 mg. Bot. 100s, 500s.
 Use: Sublingual (5-20 mg) androgen therapy.
- HORMESTRAL.** (Smith, Miller & Patch) Estrogenic substances-conjugated 1.25 mg., di-desoxyephedrine HCl 5 mg./Tab. Bot. 30s, 100s, 1000s.
 Use: (1 i.l.d.) Premenstrual tension.
- HORMESTRIN.** (Smith, Miller & Patch) Estrogens mixed, aq (20,000 u./cc.) Vial 10 cc., 30 cc. Box 1s, 6s, 25s
 Use: Estrogen therapy.
- HORMESTRIN-T.** (Smith, Miller & Patch) Estrone 20,000 u., testosterone 25 mg./cc. Vial aq. sol. 10 cc. Box 1s, 6s, 25s
 Use: i.M., estrogen & androgen therapy.
- HORMOFOLLIN.** See: Estrone (Various Mfr.).
- HORMOGEN R-A.** (Mallard) Estrone 2 mg., pot. estrone sulfate 1 mg. w/thimerosal and benzyl alcohol/cc. Vial 10 cc.
 Use: i.M., estrogen therapy
- HORMOPLETE.** (Key) Testosterone 10 mg., estrane 1 mg., Vit. B-1 50 mg., B-2 2 mg., B-6 5 mg., panthanol 10 mg., niacinamide 100 mg., inositol 25 mg., choline chloride 25 mg., dl-Methionine 25 mg., procaine hydrochloride 1%, benzyl alcohol 2%/2 cc. Vial 10 cc., 30 cc.
 Use: Androgen-estrogen therapy.
- * Generic name recognized by N.F., U.S.P. or A.M.A.
- HORMO-PLUVIMIN.** (Vitorine) Ethinyl estradiol 0.01 mg., methyl testosterone 2.5 mg., Vit. A 5000 u., D 1000 u., B-12 3 mcg., B-1 5 mg., B-2 20 mg., B-6 1 mg., C 50 mg., niacinamide 20 mg., calcium pantothenate 5 mg., choline bitartrate 50 mg., inositol 50 mg., L-lysine HCl 50 mg., Vit. E 10 I.U., rutin 25 mg., calcium 93 mg., iron 1 mg., copper 1 mg., potassium 1.5 mg., manganese 1.5 mg., zinc 7.5 mg. Vit. B-12 w/intrinsic factor concentrate 1/15 u., lemon bioflavonoid complex 10 mg., biotin 2 mcg., PABA 10 mg./Cop. Bot. 30s, 100s.
 Use: Geriatric supplement.
- NORMOTONE T.** (Cornrick) Estrogens w/thyroid. Tab. 1000 u. w/thyroid 1/20 gr. Bot. 100s, 1000s, 2000 u. w/thyroid 1/20 gr. Bot. 100s, 1000s, 5000 u. w/thyroid 1/10 gr. Bot. 100s, 1000s, 10,000 u. w/thyroid 1/10 gr. Bot. 100s, 1000s.
 Use: Orally, estrogen therapy.
- NORMO-TBIAD.** (Bell Pharmacol) Estradiol 1.2 mg., progesterone 25 mg., testosterone 25 mg., procaine HCl 1%, procaine base 1%/cc. Vial 10 cc.
 Use: Triple hormone therapy.
- HOUGASE.** (Haug) Enzyme concentrate from fresh beef pancreas 50,000 Haug Units/Tab. Bot. 50s.
 Use: Mild inflammatory conditions.
- HOUSE DUST CONCENTRATE.** (Endo) Allergic ext., purified. Diagnostic: 1 cc. vial conc. 1:200. Treatment sets: 10 cc. vial conc. 1:40,000, 1:4000, 1:400, 1:40. Maintenance treatment Pkg.: 1 cc. of 1:40. Bulk treatment Pkg.: 10 cc. of 1:40 therapeutic conc.
 Use: Specific desensitization.
- HOUSE DUST EXTRACT.**
 Abbott—Vial 1:50,000, 1:5000, 1:500.
 Parke, Davis—Pkg. 15 dose.
 Wyeth—Tube 1 cc., Vial 10 cc.
- H.P. ACTHAR GEL.** (Armour) Corticotropin highly purified. Disp. Cart. (20 & 40 u./cc.) 1 cc., Vial 5 cc., (40 u./cc.) 1 cc., Vial 5 cc., (80 u./cc.) Vial 5 cc.
 Use: i.M., stimulation of adrenal cortex & diseases related to it.
- H.P.S. SIXTY.** (Smith-Dorsey) Protein 60%, fat 1.5%, carbohydrate 27%, minerals 6.7%, moisture 5%, caloric equiv./Gm. 3.6. Can 1 lb., 4 lb.
 Use: Feeding.
- H.P.V.** (Robins) Vit. B-1 3 mg., B-2 2.5 mg., A 5000 u., D 500 u., C 30 mg., niacinamide 20 mg., pantothenic acid 1 mg./Cop. Bot. 100s, 500s.
 Use: (1-2 daily) Multivitamin therapy.

H-R LUBRICATING JELLY, (Holland Ranios)

Tube 2, 4½ oz

Use Lubricating jelly

H-STADUR (Wynn) Chlorpheniramine maleate 8 mg or 12 mg / Tab Bot 30s 100s

Use Antihistaminic

HUMACORT, (Parke, Davis) Chloramphenicol 10 mg, paramamycin 5 mg, hydrocortisone acetate 10 mg per Gm In ½ oz tubes

Use For treatment of dermatoses

HUMATIN (Parke, Davis) *Paramamycin sulfate 250 mg of paramamycin base as the sulfate Kapsel Pkg 16s

Use Intestinal amebiasis

HUMATIN SENSITIVITY DISCS, (Parke Davis) Paramamycin 30 mcg / paper disc Vial 25s

Use Testing sensitivity of microorganisms to antibiotic

HUMATIN SYRUP (Parke Davis) Paramamycin 125 mg / 5 cc

Use Antibiotic

HUMATIN SYRUP PEDIATRIC (Parke Davis) Paramamycin 125 mg / 5 cc Bot 60 cc

Use Enteric infections and diarrheas

HUMAN BLOOD CELLS DRIED. See Lyocyte Pow (Merck Sharp & Dohme)**HUMAN MEASLES IMMUNE SERUM, U S P** N N R 1955 See Globulin, Immune Serum (Human) (Various Mfr)**HUMORSOL**, N N D 1962 (Merck Sharp & Dohme) Demecarium bromide 0.25% sol 5 cc dropper vial

Use: Treatment of chronic glaucoma

HVC (Hayden's Viburnum Compound) (New York Pharmaceutical) Liq containing viburnum apulus discocorea villosa, prickly ash berries, aromatics & sufficient alcohol to release resins of the crude drugs 4 oz, 16 oz Bot

Use: Relieves smooth muscle spasms

HYACIDE, A D R (Powells) Benzethonium chloride 0.1% with sod nitrite 0.55% Bot 1 oz.

Use: Germicidal solution

NYADRINE (Searle) Diphenhydramine 37.5 mg, aminophyllin 150 mg, racephedrine 25 mg / Tab Bot 100s, 1000s

Use: (2 t i d) Bronchial asthma, hay fever & allergic disorders

HYALEX, (Bayer) Magnesium salicylate 4 gr, magnesium p aminobenzoate 2½ gr, Vit A 1500 u, Vit D 500 u, Vit E 3 u, Vit B 12 2 mcg, magnesium ascorbate 30 mg, calcium pantothenate 5 mg, zinc D7 mg, magnesium oxide 5 mg, hesperidin complex 10 mg / Tab Bot 100s

Use: Arthritis, bursitis neuritis & rheumatism

HYBEPHEN (Massengill) Phenobarbital ¼ gr, belladonna tr 5 min, hyoscyamus tr 15 min, B 1 3 mg / 5 cc or Tab Elixir, Bot 1 pt, 1 gal Tab, Bot 100s, 500s

Use: Orally, spasmodic

HYBURQUIN CREAM (Texas Pharm) Hydrocortisone ½%, iodochlorhydroxyquin 3% I C O 5% Buraw's solution 2½%, vanishing cream base ½ oz tube 2 oz jar

Use: Eczematous skin diseases

HYCETIN (Vampelt Brown) Not listed Phenobarbital ¼ gr, aspirin 2½ gr, acetaphen

*Generic name recognized by H F U S P or A M A

NYALIDASE. See: Hyaluronidase (Various Mfr)***NYALURONIDASE INJ., U S P, N N D** 1962 Inj U S P (Hyalidase, Hydase)

See: Alidase, N N R, Amp (Searle)

Diffusin, Vial (Ortho)

Enzodase, N N R, Vial (Squibb)

Hyazyme N N D, Vial (Abbott)

Infiltrase, Vial (Armaur)

Wydase Vial (Wyeth)

Use (150 u) Hypodermoclyses & promote diffusion

HYAMAGHATE. Hydraxy Aluminum Magnesium Aminoacetate (Sodium free) See Glycamine and Glycamine Liquid (Minnesota Pharm Labs Inc)**HYAMINE 10-X**, (Rohm & Haas) Diisobutyl cresoxy ethoxy ethyl dimethyl benzyl ammonium chloride See: Merphene, Amp (Barry Lab)**HYAMINE 1622**, (Rohm & Haas) See: Benzethonium Chloride U S P**HYAMINE 2389** (Rohm & Haas) Methyl dodecyl benzyl trimethyl ammonium chloride

Use Disinfection and deodorization

HYAMINE 35DD (Rohm & Haas) See: Benzethonium Chloride U S P**HYAMINE 3528** (Rohm and Haas) Trimethyl octadecyl, ammonium pentachloro phenate W/Salicylanilide See: Quatrasol, Ont (Doms)

W/Tyrolthricin & benzocaine See: Semels Loz (Massengill)

HYASORB PENICILLIN, (Key Corp) Penicillin G Pot 250 000 u / Tab Bot 36s Granules, Box 18 pkgs

Use: Penicillin therapy, prolonged release

HYAZYME, N N D 1962 (Abbott) Hyaluranidase Each vial contains sterile lyophilized hyaluronidase 150 turbidity reducing (TR) units ready to be dissolved in 1 cc of water for injection or in other infusion solution**HYBEPHEN** (Massengill) Phenobarbital ¼ gr, belladonna tr 5 min, hyoscyamus tr 15 min, B 1 3 mg / 5 cc or Tab Elixir, Bot 1 pt, 1 gal Tab, Bot 100s, 500s

Use: Orally, spasmodic

HYBURQUIN CREAM (Texas Pharm) Hydrocortisone ½%, iodochlorhydroxyquin 3% I C O 5% Buraw's solution 2½%, vanishing cream base ½ oz tube 2 oz jar

Use: Eczematous skin diseases

HYCETIN (Vampelt Brown) Not listed Phenobarbital ¼ gr, aspirin 2½ gr, acetaphen

*Generic name recognized by H F U S P or A M A

elidn 2½ gr., hyoscyamine alkaloids 0.025 mg./Tab. Bot. 100s, 500s.

Use: Orally, sedative & analgesic.

HYCIDARON, (Warren-Teed) Ferrous gluconate 100 mg., Vit. B-1 1 mg., glutamic acid 300 mg./tab. Bot. 100s, 1000s.

Use: Hematinic.

HYCLOTRITE, N.N.R. 1950. Sod. hypochlorite sol. N.F.

HYCODAH, N.N.R. 1955 (Endo) Dihydrocodeinone bitartrate 5 mg., homatropine methylbromide 1.5 mg./5 cc. or Tab. Syr Bot. 1 pl., 1 gal., Tab. Bot. 100s, 500s, 1000s. Pow. for compoundings Bot 1, 5, 10 Gm.

Use: Antiflusive.

HYCOLA, (Columbus) Aluminum hydroxide gel, dried. Tab. 7½ gr., Bot. 100s, 1000s.

Use: Orally, gastric antacid.

HYCOMINE, (Endo) Dihydrocodeinone bitartrate 5 mg., homatropine methylbromide 1.5 mg., pyrilamine maleate 12.5 mg., phenylephrine HCl 10 mg., amm. chloride 60 mg., sod citrate 85 mg./5 cc. Syr. Bot. 1 pl., 1 gal.

Use: Orally, antitussive and antihistaminic.

HYCOMINE COMPOUND, (Endo) Dihydrocodeinone bitartrate 5 mg., homatropine methylbromide 1.5 mg., chlorpheniramine maleate 2 mg., phenylephrine HCl 10 mg., N-acetyl-p-aminophenol 250 mg., caffeine 30 mg./Tab. Bot. 100s, 500s, 1000s.

Use: Antihistaminic for treatment of respiratory tract infections.

HYCOMINE TAB., (Endo) Dihydrocodeinone bitartrate 5 mg., homatropine methylbromide 1.5 mg., chlorpheniramine maleate 2 mg., phenylephrine HCl 10 mg., N-acetyl-p-aminophenol 250 mg., caffeine 00 mg./Tab. Bot. 100s, 500s, 1000s.

Use: Cold therapy.

HY-CORTEFRIN, (Allergon) Hydrocortisone 0.12% & 0.5% in suspension with phenylephrine HCl. 0.12%/15 cc., 0.5%/7.5 cc.

Use: Ophthalmic.

HYCORTOLE, N.N.D. 1962. (Premo) Hydrocortisone. Creams: 0.5%, 1%, 2.5% Tube 5 Gm., 15 Gm.; Lot. r 1% Bot. Squeeze 15 cc.; Oint. 1%, 2.5% Tube 5 Gm., 15 Gm.

HYCORTOLE ACETATE, (Premo) Hydrocortisone acetate, saline suspension 25 mg./cc Vial 5 cc. For Intra-Articular use only Ophthalmic Susp. (0.5% or 2.5%) Bot w/dropper 5 cc

HYDANTAL, (Sandoz) Mesantoin 1½ gr., phenobarbital ½ gr./tab. Bot. 100s, 500s, 1000s.

Use: (1 q. 3 h., m.p.) Anticonvulsant & anti-epileptic.

* Generic name recognized by N.F., U.S.P. or A.M.A.

HYDAHTOIH DERIVATIVES.

See: Dilantin (Parke, Davis).

Diphenylhydantoin Sod., U.S.P.

Etholoin

Mesantoin (Sandoz).

Phenatoin.

HYDASE. See: Hyaluronidase (Various Mfr.).

HYDELTRA, N.N.D. 1962. (Merck Sharp & Dohme) Prednisolone. Tab. 2.5 mg. & 5 mg. Bot. 100s, 500s, 1000s. Tab. 1 mg. Bot. 100s, 1000s.

Use: Initial dose 20-30 mg. daily; maintenance dose 5-20 mg. daily, anti-inflammatory.

HYDELTRA-T.B.A., N.N.D. 1962. (Merck Sharp & Dohme) Prednisolone tertiarybutylacetate 20 mg./cc. Aq. Susp. Vial 5 cc. Use: Rheumatic, arthritis

HYDELTRACIN TOPICAL LOTION, (Merck Sharp & Dohme) Prednisolone 5 mg., neomycin sulfate 5 mg./cc. Bot. Plastic Squeeze 15 cc.

Use: Topically t.i.d., skin inflammations.

HYDELTRASOL, N.N.D. 1962. (Merck Sharp & Dohme) Prednisolone 21-phosphate sod Topical Lot. (5 mg./cc) Plastic Bot. 15 cc. Topical Oint. (5 mg./Gm.) Tube 1/6, ½ oz. Ophth. Sol. (5 mg./cc.) Dropper Bot. 2.5 cc., 5 cc.

Use: Inflammatory disorders of the eye and skin.

HYDELTRACIN, N.N.D. 1962. (Merck Sharp & Dohme) Prednisolone 21-phosphate sod Topical Lot. (5 mg./cc) Plastic Bot. 15 cc. Ophth. Sol. (5 mg./cc.) Dropper Bot. 2.5 cc., 5 cc.

Use: Inflammatory disorders of the eye and skin.

HYDERGINE, (Sandoz) Dihydroergocornine 0.1 mg., dihydroergocristine 0.1 mg., dihydroergokryptine 0.1 mg./cc. Amp. 1 cc, Box 6s, 12s, 50s, 100s.

Use: (I.M.) in peripheral vascular disease.

HYDERGINE, S.L. TABLETS, (Sandoz) Equiproportional mixture of dihydroergocornine, dihydroergocristine & dihydroergokryptine as the methanesulfonates. Tab. S.L. 0.5 mg. Bot. 100s, 500s, 1000s.

Use: (4-6 daily) Hypertension & peripheral vascular disease.

HYDRABAMINE PENICILLIN V, N.N.D. 1960. N,N'-bis (dihydroabietyl) ethylenediamine salt of Penicillin V. See: Compacillin-V, Oral Susp. (Abbott)

* **HYDRABAMINE PHENOXYMETHYL PENICILLIN**, N.N.D. 1962. See: Penicillin V Hydrobamine.

HYDRACILLIN, (Premo) "C" Cartridges: Procaine penicillin G 300,000 u. w/procaine HCl 2%. 1 cc., 10s, 100s "600" Cartridges Procaine penicillin G 600,000 u. w/procaine HCl 2%. 1s, 10s, 100s "AMM" Cartridges: Procaine penicillin G 1,000,000 u. w/procaine HCl 2%. 1s, 10s, 100s. Tab. Pot

Homatropine Methylbromide & Phenobarbital Combinations

| Homatropine Methylbromide | Phenobarbital | Others | Manufacturer | Brand |
|---------------------------|---------------|---|------------------|-------------------------|
| 1/50 gr | 1/4 gr | Aluminum hydroxide 5 gr | Hart | Hartgel H M B, Tab |
| 1/24 gr | 1/4 gr | Aluminum hydroxide 7 1/2 gr, Mag trisilicate 2 1/2 gr | Rossmar | Alased Tab |
| 2.5 mg | 8 mg | Alumseum hydroxide gel 129.6 mg, Mag trisilicate 194.4 mg | Blue Line | Neutratabs, Tab |
| 2 mg | 1/6 gr | d Amphetamine sulfate 2 mg | Adams | Relaxamine Tab |
| 2.5 mg | 8 mg | Dehydrocholic acid 250 mg | Mallinckrodt | Cholan HMB, Tab |
| 2.5 mg | 8 mg | Dehydrocholic acid 250 mg | Chicago Pharm | Hydrophen, Tab. |
| 1/48 gr | 1/4 gr | Dehydrocholic acid 2 gr, ox bile 2 gr | Norgine Labs | Bilamide Tab. |
| 1 mg | 8 mg | Dihydroxy aluminum aminoacetate 0.5 Gm | Flint, Eaton | Homaldron, Tab. |
| 1/100 gr | 1/4 gr | Dihydroxy aluminum aminoacetate 7 1/2 gr | First Texas | Alkam Comp. Tab |
| 1/50 gr | 1/4 gr | Dihydroxy aluminum aminoacetate 7 1/2 gr, Vit C 25 mg | Patch | Oalzinatc, Tab. |
| 0.13 mg | 1/4 gr | Estrone sulfate Sod 0.42 mg, thyroid 8 mg | Columbus | Meno-Sed Tab |
| 0.567 mg | 1/4 gr | Hyoscyamine HBr 0.106 mg, hyosine HBr 0.007 mg | Sherman | Novadonnal, Tab |
| 0.0447 mg | 1/4 gr | Hyoscyamine HBr 0.54 mg, scopolamine HBr 0.0144 mg, thiamine HCl 3 mg | Paul Maney | Sedlie Unoday, Cap |
| 5 mg | 1/4 gr | Kaolin 300 mg | Mallinckrodt | Lusyn, Tab. |
| 2.5 mg | 1/4 gr | Mannitol hexacetate 30 mg, veratrum viride 100 mg | Crochet Barnes | Perenal, Tab. |
| 1 mg | 8 mg | Mannitol hexacetate 16 mg, veratrum viride 20 mg | Stoddard | Verante, Tab |
| 1.2 mg | 8 mg | Mephnesin 0.3 Gm, hyoscyamine sulfate 0.1 mg, scopolamine HBr 0.02 mg | Central | Mydonna, Tab. |
| 1 mg | 8 mg | Methylcellulose 300 mg | Chickitt | Atralase, Tab |
| 1/50 gr | 1/4 gr | Ox bile 1.5 gr, acidized bile acids 1.5 gr, steapsin 3/4 gr | Columbus | Oxachalin, Tab. |
| 2.5 mg | 8 mg | Pentobarbital Sod 8 mg | Rorer | Precibus, Tab. |
| 1/480 gr | 1/4 gr | Pentobarbital Sod 1/16 gr | Walker | Sedadrops, Elix |
| 1/480 gr | 1/4 gr | Pentobarbital Sod 1/16 gr | Walker | Sedaplex, Elixir & Tab. |
| 1/480 gr | 1/4 gr | Pentobarbital Sod 1/16 gr | Jackson Mitchell | Salsed, Pedalatric, Uq |
| 2 1/2 mg | 7 1/2 mg | Pentobarbital 7 1/2 mg | Meyer | Almesed, Elixir |

- equivalent) 3 mcg., ascorbic acid 60 mg., Vit. A 3000 u., Vit. D 1000 u./5 cc. Bot. 60 cc., 120 cc., 1 pt.
Use: Vitamin deficiencies.
- ***HOMOCHLORCYCLIZINE**, (Abbott) 1-(p-chlorobenzhydryl)-4-methylpiperazine.
HOMOCILLIN, (Rexall) Procaine penicillin G 300,000 u., Pot. penicillin G 100,000 u./dose. Vial 1 dose.
Use: Penicillin therapy.
- HOMOPENT**, (Lemmon) Pentobarbital Sod. $\frac{1}{4}$ gr., homotropine methylbromide 1/25 gr., Mag. Trisilicate 5 gr./Tab. Bot. 100s
Use: Gastric antacid & antispasmodic.
- ***HOMOSULFANILAMIDE HCl**.
See: Sulfamylon HCl (Winthrop Labs)
Sulfbenzamide HCl (Flint, Eaton)
- HOMVITE**, (Cartone) Thiamine hydrochloride 5 mg., pyridoxine hydrochloride 5 mg., nicotinic acid 20 mg., vit. B-12 25 mcg., L-lysine hydrochloride 300 mg./15 cc.
Use: Geriatric nutritional supplement.
- HONEY CILLIN 400**, (Physicians Products) Penicillin G, buffered, 4,800,000 u./60 cc. 400,000 u/5 cc. Bot. 60 cc.
Use: Penicillin therapy.
- HORMALE**, (Key Corp.) Testosterone propionate in oil. Vial (25 mg./cc.), (50 mg./cc.) 10 cc.
HORMALE LINGUALS, (Key Corp.) Methyltestosterone 10 mg. Bot. 100s, 500s.
Use: Sublingual (5-20 mg.) androgen therapy.
- HORMESTRAL**, (Smith, Miller & Patch) Estrogenic substances-conjugated 1.25 mg., di-desoxyephedrine HCl 5 mg./Tab. Bot. 30s, 100s, 1000s.
Use: (1 t.i.d.) Premenstrual tension.
- HORMESTRIN**, (Smith, Miller & Patch) Estrogens mixed, aq. (20,000 u./cc.) Vial 10 cc., 30 cc. Box 1s, 6s, 25s
Use: Estrogen therapy.
- HORMESTRIN-T**, (Smith, Miller & Patch) Estrone 20,000 u., testosterone 25 mg./cc. Vial aq. sol. 10 cc. Box 1s, 6s, 25s
Use: I.M., estrogen & androgen therapy.
- HORMOFOLLIN**. See: Estrane (Various Mfr)
HORMOGEN R-A, (Mallard) Estrone 2 mg., pot. estroee sulfate 1 mg w/thimerosal and benzyl alcohol/cc. Vial 10 cc.
Use: I.M., estrogen therapy
- HORMOPLETE**, (Key) Testosterone 10 mg., estrone 1 mg. Vit. B-1 50 mg., B-2 2 mg., B-6 5 mg., panthenol 10 mg., niacinamide 100 mg., inositol 25 mg., choline chloride 25 mg., dl-Methionine 25 mg., procaine hydrochloride 1%, benzyl alcohol 2%/2 cc. Vial 10 cc., 30 cc.
Use: Androgen-estrogen therapy.
- ***HORMO-PLUVIMIN**, (Vitarine) Ethinyl estradiol 0.01 mg., methyl testosterone 25 mg., Vit. A 5000 u., D 1000 u., B-12 3 mcg., B-1 5 mg., B-2 20 mg., B-6 1 mg., C 50 mg., niacinamide 20 mg., calcium pantotheate 5 mg., choline bitartrate 50 mg., inositol 50 mg., l-lysine HCl 50 mg., Vit. E 10 I.U., rutin 25 mg., calcium 93 mg., iron .1 mg., copper 1 mg., potassium 1.5 mg., magnesium 1.5 mg., zinc 7.5 mg., Vit B-12 w/intrinsic factor concentrate 1/15 u., lensae bioflavonoid complex 10 mg., biotin 2 mcg., PABA 10 mg./Cap. Bot. 30s, 100s.
Use: Geriatric supplement
- NORMOTONE T**, (Carrick) Estrogens w/thyroid. Tab. 1000 u. w/thyroid 1/20 gr. Bot. 100s, 1000s, 2000 u. w/thyroid 1/20 gr. Bot. 100s, 1000s, 5000 u. w/thyroid 1/10 gr. Bot. 100s, 1000s, 10,000 u. w/thyroid 1/10 gr. Bot. 100s, 1000s.
Use: Orally, estrogen therapy.
- NORMO-TRIAD**, (Bell Pharmacal) Estradiol 1.2 mg., progesterone 25 mg., testosterone 25 mg., procaine HCl 1%, procaine base 1%/cc. Vial 10 cc.
Use: Triple hormone therapy.
- NOUOASE**, (Haug) Enzyme concentrate from fresh beef pancreas 50,000 Haug Units/Tab. Bot. 50s.
Use: Mild inflammatory conditions.
- NOUSE DUST CONCENTRATE**, (Endo) Allergic ext., purified. Diagnostic: 1 cc. vial conc. 1-200. Treatment set: 10 cc. vial conc. 1:40,000, 1:4000, 1:400, 1:40. Maintenance treatment Pkg: 1 cc. of 1:40. Bulk treatment Pkg: 10 cc. of 1:40 therapeutic conc.
Use: Specific desensitization.
- NOUSE DUST EXTRACT**.
Abbott—Vial 1,50,000, 1,500, 1,500.
Fork, Davis—Pkg. 15 dose.
Wyeth—Tube 1 cc., Vial 10 cc.
- H.P. ACTHAR GEL**, (Armour) Corticotropin highly purified. Disp. Cart. (20 & 40 u./cc.) 1 cc., Vial 5 cc., (40 u./cc.) 1 cc., Vial 5 cc., (80 u./cc.) Vial 5 cc.
Use: I.M., stimulation of adrenal cortex & diseases related to it.
- N.P.S. SIXTY**, (Smith-Dorsey) Protein 60%, fat 1.5%, carbohydrate 27%, minerals 6.3%, moisture 5%, caloric equiv./Gm. 3.6. Can 1 lb., 4 lb.
Uses: Feeding.
- N.P.V.** (Robins) Vit. B-1 3 mg., B-2 2.5 mg., A 5000 u., D 500 u., C 30 mg., niacinamide 20 mg., pantotheic acid 1 mg./Cap. Bot. 100s, 500s.
Use: (1-2 daily) Multivitamin therapy.

penicillin G buffered w/Cal cost 50 000 u, 100 000 u, 200,000 u, 250 000 u, 500,000 u, 1,000 000 u Bot 12s, 100s Vial, procaine penicillin G 300 000 u/cc 10 cc

HYDRAL. (Person & Covey) Chloral hydrate Cap (3¼ & 7½ gr) Bot 100s

Use Sedative

***HYDRALAZINE HCl, NF, NND 1962** 1 Hydroxynaphthalazine HCl Inj Tab, NF See Apresoline, Amp & Tab (Ciba)

W/Reserpine See Serpassil Apresoline Tab (Ciba)

W/Reserpine hydrochlorothiazide (Esidrix) See SER AP ES Tab (Ciba)

HYDRAMIN (Nion) Amino acids approx 12 Gm, thiamine HCl 3 mg, riboflavin 6 mg niacinamide 60 mg, Vit C 75 mg/lip Can 6 oz 1 lb 2 oz.

Use Malnutrition in pregnancy & lactation

***HYDRARGAPHEN.** (Mead Johnson) Phenyl mercury 2,2' dinaphthylmethane 3:3' disulphonate Penotrans

HYDRASOL CONCENTRATED. (Blue Line) Hydrostine HCl 11 gr, zinc sulfate, zinc phenolsulfonate, boric acid, cresol/oz Sol Bot 4 oz 1 pt

Use: Topically antiseptic

HYDRASTINE ALKALOID (Mallinckrodt) Bot 5 gr, ¼ oz, 1 oz

HYDRASTINE HYDROCHLORIDE (Mallinckrodt) Pow, Bot 5 gr, ¼ oz, 1 oz

Use: Uterine hemostatic

W/Ichthammol, iodine, phenol, glycerite of baroglycerin & eucalyptol See Ichtholdine, Liq (Merck Sharp & Dohme)

HYDRASTIS, EXTRACT, NF, W/Butesin & hamamelis ext See: Butesin Compound (Abbott)

HYDRELT CREAM (Elder) Aluminum acetate sol 20 mg, iodochlorohydroxyquin 30 mg hydrocortisone 5 mg each Gm in water miscible base ½ oz tube

Use Dermatoses

HYDREPEL-A. (Hygiene Research Inc) Plasticized combinations of ethyl cellulose & castor oil in mixed propanols Cream, Pkg 1 oz, 2½ oz

Use: Skin protective against water solutions & water borne irritants

HYDREPEL-S (Hygiene Research Inc) Plasticized combinations of ethyl cellulose & castor oil in mixed propanols & a silicone Cream, Pkg 1 oz, 2½ oz

Use: Protective against water solutions & water borne irritants

W/Hydrocortisone acetate by NF, USP or A.M.A.

tate 2% in a buffered aq base Vial 5 cc
Use: Hydrocortisone therapy

HYDRIODIC ACID. (Various Mfr) Hydrogen iodide About 70% Bot 25 Gm, 100 Gm
Use: Iodide therapy

***HYDRIODIC ACID SYRUP, NF** See: Hyadin, Syr (Gardner)

HYDRIODIC ACID THERAPY.

See: Aminoacetic Acid HI

Hyadin, Syr (Gardner)

Glyco HI, Cap (Burnham)

HYDRIONIC. (Upjohn) Glutamic acid HCl Cap 5 gr Bot 100s, 500s, 1000s

Use: (1 2 1 d) Achlorhydria & hypochlorhydria

HYDROBALM. (Merck Sharp & Dohme) Hydrocortisone 0.5% in calamine cream Tube 5 Gm, 15 Gm & 1 oz

Use Topically, allergic dermatoses

HYDROBALM 0.5% TOPICAL LOTION

(Merck Sharp & Dohme) Hydrocortisone 0.5% in calamine lotion Bot 15 & 30 cc
Use Skin conditions

HYDRO-BILEIN TABLETS (Abbott) Bilein (Abbott's dried & purified ox bile) 2 gr, dehydrocholic acid 2 gr/Tab Bot 100s, 1000s

Use: (1-2 q d) Bile therapy

HYDROCHLORATE. Same as Hydrochloride

HYDROCHLORIC ACID, USP, A DR (Various Mfr) (Muriatic Acid) Absolute 38% Diluted 10%, NF, A DR 1954

Use: Well diluted, achlorhydria

HYDROCHLORIC ACID THERAPY.

See Betaine HCl (Various Mfr)

Glutamic Acid HCl (Various Mfr)

Glycine HCl (Various Mfr)

***HYDROCHLOROTHIAZIDE, USP, NND 1962;** Tab, USP 6 chloro 3,4 dihydro 7 sulfamoylbenzo 1,2,4 thiazolidine dioxide 11

See: Aquarius, Tab (Frossi)

Esidix (Ciba)

Hydro Diurnl (Merck Sharp & Dohme)

Oretic Tab (Abbott)

W/Butabarbital See Butizide Prestab (McNeil)

W/Butabarbital reserpine See Butiserpazide Prestab (McNeil)

W/Meprobamate See Cyclex, Tab (Merck Sharp & Dohme)

W/Pentaerythritol tetranitrate See Perithiaside SA, Tabs (Warner Chilcott)

W/Reserpine

See: Hydropres Tab (Merck Sharp & Dohme)

Serpassil Esidrix Tab (Ciba)

W/Reserpine, Hydralazine HCl See: SER AP ES, Tab (Ciba)

W/Reserpine potassium chloride

- See: Hydropres-KA, Tab. (Merck Sharp & Dohme)
 Salupres, Tab. (Merck Sharp & Dohme)
 W/Syrasingopine. See: Singaverp-Esdirix. (Ciba)
- HYDROCHOLERETICS.**
 See: Bile Salts (Various Mfr.).
 Cholic Acid (Various Mfr.).
 Dehydrocholic Acid (Various Mfr.).
 Desoxycholic Acid (Various Mfr.).
 Ketocholic Acid (Various Mfr.).
 Ox Bile (Various Mfr.).
- HYDROCIL** (Fuller) Psyllium mucilloid, karaya, vit. B-1, dextrase. Cont. 4 oz., 1 lb. Use: Laxative.
- HYDROCIL FORTIFIED.** (Fuller) Hydrocil with acetphenolisatin 2 mg/5 Gm. Bot. 4 oz., 1 lb.
- HYDROCIL FORTIFIED CAPS.** (Fuller) Psyllium mucilloid 1.925 gr., methylcellulose 5.775 gr., acetphenolisatin 3 mg./Cap. Bot. 30s, 100s. Use: Laxative.
- HYDROCIN.** (Philadelphia Ampoule) Hydrocortisone acetate 50 mg., neomycin sulfate 5 mg./cc. Vial 5 cc. Box 25s, 100s. Use: Antibiotic/hormone preparation.
- HYDROCODONE.** See: Dihydrocodeinone. (Various Mfr.)
- *HYDROCORTAMATE, HCl, N.N.D. 1962.**
 17-Hydroxycorticosterone-21-diethylaminoacetate HCl.
 See: Magnacort. (Pfizer)
 Ulicort (Ulmer)
- W/Neomycin sulfate
 See: Neo-Ulicort, Oint. (Ulmer)
 Vio-Cort, Cream. (Rowell)
- W/Neomycin, proteolytic enzymes. See: Chymar (Armour Pharmaceutical)
- *HYDROCORTISONE, U.S.P., N.N.D. 1962.**
 Oint., Lot., Tab., U.S.P.
 Tab., U.S.P.
- See: Cort-Dome, Cream. (Dome Chemical)
 Cortef. (Upjohn)
 Cortilan, Cr. (Schering)
 Cortispray, Liq. (Walker)
 Cortril (Pfizer).
 Demafene-HC, Oint. (Dome)
 Heb-Cort (Barnes-Hind)
 Hycortale, N.N.R., Prep. (Prema)
 Hydrocortone. (Merck Sharp & Dohme)
 Hytone Lotion (Dermik)
 Oplef, Sol. (Upjohn)
 Otosone-P, Liq. (Braemmel)
 Torcortin, Oint. (Reed & Carrick)
 Texacort Lotion 25 (Texas Pharm)
 Topicort, Spray. (Rausiel)
- W/Acetyl-p-aminophenol, Flexin. See: Flexilan-HC, Tab. (McNeil)
- W/Allantoin, tar, and soybean complex gel
 See: Altara-Cort, Gel (Dome)
- W/Aluminum acetate solution, iodochlorohydroxyquin See: Hydrelt Cream (Elder)
- W/p-Amino benzoic acid, aspirin, ascorbic acid. See: Pabrin-AC, Cap. (Smith-Dorsey)
- W/Ascorbic acid. See: Cortalex, Tab. (Upjohn)
- W/Aspirin. See: Hydradyne, Tab. (Merck Sharp & Dohme)
- W/Aspirin, aluminum hydroxide gel. See: Aceto-Cort, Tab. (Mallard)
- W/Bacitracin, neomycin sulfate See: Bacimycin W/Hydrocortisone, Oint. (Walker)
- W/Bacitracin, tyrothricin, neomycin sulf., benzocaine. See: Hydrazets, Trache. (Merck Sharp & Dohme)
- W/Calamine Lotion. See: Hydrobalm 0.5% Topical Lotion. (Merck Sharp & Dohme)
- W/Chloromycetin. See: Chloromycetin-Hydrocortisone, Ophth. Oint. (Parke, Davis)
- W/Chloroxylenol. See: Sepliderm-HC, Cream (Aveeno)
- W/Coaf tar, allantoin. See: Alphosyl-HC, Lot. (Reed & Carrick)
- W/Coaf tar sol. in acid Manile vehicle. See: Carbo-Cort, Cream, Lot. (Dome)
- W/Coaf Tar Ext. See: Tarcortin, Cream. (Reed & Carrick)
- W/Coaf Tar Solution, diiodohydroxyquinoline See: Car-Tar-Quin, Creme. (Dome)
- W/Desitin ointment formula. See: Desitin-HC Oint. (Desitin)
- W/di-desoxyephedrine hydrochloride, pyrilamine maleate, antipyrine. See: Rhinalgon HC (Daha)
- W/Diiodohydroxyquinoline, coal tar soln., in a water-miscible base. See: Car-D-Tar (Desitin)
- W/Diperodon HCl See: Pentacort, (Dolin)
- W/Estrogens (Premarin). See: Premarin-HC, Tube. (Ayerst)
- W/Gramicidin, neomycin sulfate, polymyxin-B sulfate, phenylephrine & paredrine HBr. See: Trisacort, Spraypak. (Smith, Kline & French)
- W/Ichthammol. See: Ixal, Creme. (Dole)
- W/Iodochlorohydroxyquin.
 See: Demaform-HC, Cream, Lot. (Dome Chemicals)
 Lidaform-HC Oint. (Dome)
 Vytone (Dermik)
- W/Iodochlorohydroxyquin See: Vioform-Hydrocortisone, Lotion. (Ciba)
- W/Iodochlorohydroxyquin, LCD See: Cortarin Cream—1/4%, Tube (C & M Pharmacal)
- W/Iodochlorohydroxyquin, LCD, Buraw's solution. See: Hyburquin Cream. (Texas Pharm)
- W/Diiodohydroxyquinoline, Zetar. See: Zetar-Quin (Dermik)

* Generic name recognized by N.F., U.S.P. or A.M.A.

- W/Adochlarhydroxyquin, neomycin sulfate See Nafsa Carl Oint (Rowell)
- W/Mephenesin, sod salicylate homatropine methylbromide See Mephasol with Hydrocortisone Tob (Crookes Barnes)
- W/Methylandrostenol, Vit B 12 See: Vis tabolic, Tob (Organon)
- W/Neomycin See Neo Carl Dame, Cream (Dame)
- W/Neomycin, polymyxin B sulf, chlorpheniramine maleate phenylephrine HCl See: Polycort Sol (Lemman)
- W/Neomycin polymyxin B sulfate heparin Sod See Bro Parin susp (Braemmel Pharm)
- W/Neomycin sulfate See: Heb Carl N 1/4% Lotion (Barnes Hind)
- W/Neomycin sulfate See: Neo Tarcortin Oint (Reed & Carrick)
- W/Neomycin sulfate See: Protef, Supp (Upjohn)
- W/Neomycin sulf, gramicidin, phenylephrine HCl See Du Biotic Intra-nasal (White)
- W/Neomycin sulf, phenylephrine HCl See: Medihaler Phen Vial (Riker)
- W/Neomycin sulf, polymyxin B sulf See: Cortisporin Otic Drops (Burroughs Wellcome)
- W/Neomycin sulfate pyrilamine maleate cetalkonium chloride, dipiperdon HCl urea See Tega Otic Soln (Ortega)
- W/Neomycin sulfate pyrilamine maleate cetalkonium chloride dipiperdon HCl See Fent O Augine Ear Drops Liq (Confield)
- W/Neomycin sulfate, pyrilamine maleate cetalkonium chloride, dipiperdon HCl urea See Otatrin Ear Drops (Louvissans)
- W/Neomycin sulf pyrilamine maleate cetalkonium chloride dipiperdon HCl urea See Tyler Otic Sol (Tyler)
- W/Nystatin neomycin sulfate See Neo Nysta Carl (Dame)
- W/Pantothényl See: Pantho F, Cream (U S Vitamin)
- W/Phenol menthol chlorbutanol See Pruristal Cream (Dole)
- W/Phenylephrine HCl See Hy cartelzin Liq (Allergan)
- W/Phenylephrine methylcellulose See: Isopto Sterafin Soln (Alcon)
- W/Polymyxin B sulfate See Cortamixin Vial (Braemmel)
- W/Polymyxin B sulfate bacitracin & neomycin sulfate See: Cortisporin Ophth Oint (Burroughs Wellcome)
- W/Polymyxin B sulfate neomycin sulfate See: Ocutrin Eye Drops (Louvissans)
- W/Polymyxin B sulfate neomycin sulfate Steramine Ophthalmic, Liq (Mayrand)
- W/Polymyxin B sulfate neomycin sulfate
- antipyrine dibucaine HCl See: Otocort Liq (Lemman)
- W/Polymyxin B sulfate, neomycin sulfate heparin sodium See: Dermo-Porin, Oint (Braemmel)
- W/Polymyxin B sulfate, neomycin, heparin sodium See Bro Parin H, Vial (Braemmel)
- W/Polymyxin B sulfate, neomycin, pramoxine hydrochloride See: Auracort, Drops (Columbus)
- W/Polyoxyethylene Lauryl Ether, Hexachlorophene, Sulfur See Fosiril HC, (Westwood)
- W/Pot salicylate, Pot p-aminobenzoate & Vit C See Neocylate H, Tob (Central)
- Pobalate HC, Tob (Robins)
- W/Pyrimine maleate, pheniramine maleate phenylpropranolamine HCl See: Triaminic HC, Tob (Smith Dorsey)
- W/Salicylamide, p-aminobenzoic acid, Vit C See: Artamide HC, Cap (Wampale)
- Sylopar HC, Tob (Burlington)
- W/Steratan See: Sterosan Hydrocortisone Oint (Geigy)
- W/Sulfanilamide sodium dimethylacryl, benzocaine, menthol See Reclolyt HC, Rectisris (Mallan/Daho)
- W/Sulfur colloidal salicylic acid, resorcinol See Theracort Lot (C & M Pharm)
- W/Superiah See Neo Superiah S, Cream (Tailby Nazon)
- W/Terramycin See Terra Cortin Topical Aerosol Spray (Pfizer)
- W/Tetracycline HCl See: Achramycin W/Hydrocortisone Oint & Ophth Oint (Lederle)
- W/Thenyldiamine HCl, Neomycin sulf, polymyxin B sulf, phenylephrine HCl See Biosynephrine (Winthrop Labs)
- W/Thonzonium bromide, neomycin sulfate gramicidin, thanzylamine HCl & phenylephrine HCl See: Biomydrin F W/Hydrocortisone Sol (Nepera)
- W/Triaminic formula See Triaminic HC, Tob (Smith Dorsey)
- W/Trichlisonium chloride See Triburan HC Ointment (Roche Labs)
- W/Tyrosin hexachlorophene See Acne sol (Dole)
- W/Tyrosin, phenylephrine HCl, thenylpyrimine HCl See Pyracort, Liq (Lemman)
- W/Vit A, estrone, pyrilamine maleate See Hial A Carl, Creme (Dame)

*HYDROCORTISONE ACETATE, U.S.P., NND 1962 Oint In U.S.P., Sterile Susp Ophth Oint Ophth Susp, U.S.P. 17 hydrocortisone 21-acetate comp F See: Anusol HC, Supp (Warner-Chilcott)

* Generic name recognized by NF U.S.P. or A.M.A.

- Cartef Acetate, Ophth. Oint. & Tob., N.N.R. (Upjohn).
- Cortril Acetate, Aq. Susp., Oint. & Tob., N.N.R. (Pfizer).
- Hydrin-2, Viol. (Broemmal)
- Hydrocortone Acetate, N.N.R. (Merck Sharp & Dohme)
- Isopto-Hydrocortisone, Ophth. Susp. (Alcon Lab.)
- Triamel-HC (Fuller)
- W/Bacitracin, neomycin sulf., polymyxin B sulf. See: Neopolycin-HC, Oint., Ophth. Oint. (Pitman-Moore)
- W/Benzocaine, ephedrine HCl, oxyquinoline sulfate, zinc oxide, menthol, balsam of Peru. See: Medicone-HC, Supp. (Medicone)
- W/Benzocaine, oxyquinoline sulfate, ephedrine HCl, menthol, ichthammol, zinc oxide. See: Derma Medicone-HC (Medicone).
- W/Chloramycetin. See: Chloramycetin-Hydrocortone, Ophth. (Parke, Davis).
- W/Chloramycetin, polymyxin B sulfate. See: Ophthacort, ointment. (Parke, Davis)
- W/Chloramphenicol, paramoxycin, in a liquid petrolatum base. See: Humacort Ointment (Parke, Davis)
- W/Cod liver oil, lanolin, zinc oxide, bismuth subgallate, balsam peru. See: Desitin HC Supp. (Desitin Chem.)
- W/Celistin base, neomycin base, thonzonium bromide, polysorbate 80, acetic acid, sodium acetate. See: Caly-Mycin Otic
- W/Neomycin and Hydrocortisone (Warner-Chilcott)
- W/Mephenesin and salicylamide. See: Artican, Tab. (Walker Lab.)
- W/Neomycin.
- See: Cortamycin, Oint. (Bryant)
- Hydrotropic Oph. Susp. (Merck Sharp & Dohme)
- W/Neomycin sulfate.
- See: Hydrocin, Liq. (Philadelphia Ampoule)
- Hydrocortisone-Neomycin, inf. (Philadelphia)
- Neo Cortef, Oint. (Upjohn)
- Scinocort, Vial (Testogor)
- W/Neomycin, bacitracin. See: Hydroderm, Top. Oint. (Merck Sharp & Dohme)
- W/Neomycin sulfate, dyclanine HCl, polymyxin B sulfate. See: Neo polycin HC Otic (Pitman-Moore)
- W/Paradrine HBr, phenylephedrine HCl & thimerosal. See: Vosocort, Sol., Spraypak (Smith, Kline & French).
- W/Phenylephrine & chlorobutanol. See: Op-Hydrin, Susp. (Broemmal)
- W/Phenylephrine HCl, neomycin sulfate. See: Neo-Cortef, Nasal Spray (Upjohn).
- W/Polymyxin B sulfate, and Neomycin sulfate. See: Isopto-P.H.N., Ophth. Sol. (Alcon)
- W/Prednisone, prednisolone, neomycin sulfate. See: Pansteroid, Oint. (Kenwood)
- W/Propodrine HCl and neomycin sulf. See: Hydrospray Nasal Susp. (Merck Sharp & Dohme)
- W/Salicylamide, mephenesin. See: Articon, Tab. (Walker)
- W/Selenium disulfide. See: Se'sunef, Oint. (Abbott)
- W/Sod. salicylate, mephenesin, homatropine methylobromide. See: Mephasol with Hydrocortisone, Tab. (Ames)
- W/Terramycin. See: Terra-Cortril Oint. (Pfizer).
- W/Terramycin HCl. See: Cortril Acetate w/Terramycin, Ophth. Susp. (Pfizer).
- W/Tetracycline HCl & phenylephrine HCl. See: Achromycin Nasal Suspension, (Lederle)
- W/Tyrosine, phenylmercuric acetate, 9 aminoacridine HCl, polysorbate-80, urea, lactose. See: Aquocort, Vog Supp. (Webster)
- W/Xylocaine. See: Lidafarm-HC (Dome)
- HYDROCORTISONE CREAM** (free alcohol)
See: Cortilan Cream 1% (Schering).
- *HYDROCORTISONE CYCLOPENTYLPROPIONATE**, N.N.D. 1962.
See: Cortel Fluid, Susp. (Upjohn)
Cortef Oral Suspension, (Upjohn)
- HYDROCORTISONE DIETHYLAMINOACETATE HCl**. See: Hydrocortamate
- W/Neomycin, Undecylenic acid in a petrolatum base. See: Padiacort ointment, (Padiacort)
- W/Neomycin sulfate. See: Neo-ulcort (Ulmar)
- HYDROCORTISONE I. V.** See: Solu-Cortef, Vial (Upjohn)
- HYDROCORTISONE-NEOMYCIN**. (Philadelphia Amp.) Hydrocortisone acetate 5D mg., neomycin sulfate 5 mg./cc. Vial 5 cc. Box 25s, 100s.
Use. Rheumatoid and osteoarthritis and post-traumatic bursitis.
- HYDROCORTISONE-NEOMYCIN OINTMENT**. (Maney) Neomycin sulfate 3.5 mg., Hydrocortisone acetate 1D mg./Gm. Ophth. or top. Tube 1/2 oz.
Use: Control of dermatitis.
- HYDROCORTISONE PHOSPHATE**. See: Hydrocortone Phosphate (Merck Sharp & Dohme)
- *HYDROCORTISONE SODIUM SUCCINATE**, U.S.P., N.N.D. 1962. (I.V. form) inf., U.S.P. See: Solu-Cortef, Vial (Upjohn)
- *HYDROCORTISONE TERTIARY-BUTYLACETATE**. See: Hydrocortone-T.B.A. Supp. (Merck Sharp & Dohme)
- HYDROCORTONE**. See: Hydrocortisone, U.S.P., N.N.D. 1962.

*Generic name recognized by N.F., U.S.P. or A.M.A.

W/Calamine cream See: Hydrobalm, Cream (Merck Sharp & Dahme)

W/Propadrine HCl & neomycin sulfate See Hydro spray, Nasol Suspension (Merck Sharp & Dahme)

HYDROCORTONE ACETATE, NND 1962 (Merck Sharp & Dahme) 17 Hydroxycorticosterone 21 acetate Kendall's Comp F Dental Oint. 2.5% Tube 5 Gm; Oint 1D mg or 25 mg/Gm Tube 5, 15 & 30 Gm; Ophth Oint 15 mg/Gm Tube 3.5 Gm

Ophth Susp 5 mg or 25 mg/cc Bot w/ dropper 5 cc r Saline Susp 25 mg/cc & 50 mg/cc Vial 5 cc Tab 5 Bot 50s 10 Bot 50s, 100s & 20 mg, Bot 25s, 100s Use Rheumatoid & osteo arthrrhs

W/Neomycin See Hydroptic Ophth Susp (Merck Sharp & Dahme)

HYDROCORTONE ACETATE, TOPICAL OINTMENT, NND 1958 (Merck Sharp & Dahme) 10 or 25 mg/Gm Tube 5 Gm

HYDROCORTONE, INFUSION CONCENTRATE, (Merck Sharp & Dahme) Hydrocortisone Amp (100 mg) 20 cc Use: To be diluted w/dextrose or physiologic saline sol for surgical shock, drug reactions & acute allergic emergencies

HYDROCORTONE 0.5%, 1% & 2.5% LOTION TOPICAL, (Merck Sharp & Dahme) Hydrocortisone Plastic bot 15 cc Use: (Apply 2 or 3 times daily) Pruritus or dermatitis

HYDROCORTONE TABLETS (Merck Sharp & Dahme) Tab (5 mg) Bot 50s, (10 & 20 mg) Bot 50s 100s 500s

HYDROCORTONE PHOSPHATE Inj (Merck Sharp & Dahme) Hydrocortisone (as hydrocortisone 21 phosphate disodium salt) 50 mg/cc Vials 2 cc Use Adrenocortical Steroid

HYDROCORTONE-T B A, SUSPENSION, (Merck Sharp & Dahme) Hydrocortisone tertiarybutylacetate Vial (25 mg/cc) 5 cc Use: Intrasyneovial, hydrocortisone therapy

HYDRDDERM OINT, TOPICAL, (Merck Sharp & Dahme) Oint 1% (Hydrocortisone 1D mg, neomycin 3.5 mg, bacitracin 1000 u/Gm) Tube 1/6 oz & 1/2 oz Oint 2.5% (Hydrocortisone 25 mg, neomycin 3.5 mg, bacitracin 1000 u/Gm) Tube 1/6 oz & 1/2 oz

HYDRODIURIL, NND 1962 (Merck Sharp & Dahme) Hydrochlorothiazide 25 mg, 50 mg/Tab Bot 100s, 1000s

Use: Edema congestive heart failure renal disease, acute and chronic glomerulonephritis and nephrosis

HYDRDDIURIL-KA (Merck Sharp & Dahme) Hydrochlorothiazide 25 mg, and 50 mg. *Generic name recognized by NF, USP or A.M.A.

potassium chloride 572 mg/Tab Bot 100s Use Diuretic and antihypertensive

HYDRODYNE, (Merck Sharp & Dahme) Hydrocortisone 2.5 mg, aspirin 300 mg/Tab Bot 100s, 500s, 1000s

Use: Analgesic w/hydrocortisone therapy

***HYDROFLUMETHIAZIDE** 3,4 dihydro 6 tri fluoromethyl 7 sulfamoylbenza 1,2,4 thiazine 1,1 dioxide See: Saluron Tab & Syr (Bristol)

W/Reserpine, proloaveratine A See: Salutensin Tab (Bristol)

HYDROGEN, (Breon) 10% Aluminum hydroxide Susp Jar 10 oz r Tab Box 50s

Use: Gastric antacid

HYDROGEN DIOXIDE, U.S.P., A.D.R. See: Hydrogen Peroxide, U.S.P. (Merck Sharp & Dahme)

HYDROGEN PEROXIDE, See: G.H.P. Carbamide (Internal Pharm Corp)

HYDROGEN PEROXIDE SOLUTION, U.S.P., A.D.R. (Various Mfr) (3%) See: Neaxyn Rorer

Use Antiseptic

HYDROGEN PEROXIDE SOLUTION 3D%, A.D.R. Perhydrol Hydrogen peroxide Bot 1/4 lb, 1/2 lb, 1 lb See Superoxol (Merck Sharp & Dahme)

Use Dentistry & preparing the 3% solution

HYDROLAMINS (Lewal) Amino acids in a polyethylene glycol ointment base Tube 5 Gm 20 Gm

Use: Antipruritic

HYDROLAX (Invenex) Diacetyl sod succinate 1/2 gr, methylcellulose 500 mg/tab Bot 100s

Use: Orally (12 tabs daily), a swelling agent that keeps the stool soft

HYDROLAX PEDIATRIC, (Invenex) Diacetyl sod succinate 1% Liq Bot 30 cc

Use: Fecal softener

HYDROLOSE, FORTIFIED, SYRUP, (Upjohn) Diacetylhydroxyphenylisatin 15 mg, prune pow 148 Gm, methylcellulose 5.91 Gm./fl oz Bot 6 oz & 12 oz

Use Bulk producing laxative

HYDROLOSE, NND 1962 (Upjohn) Each 5 cc/1 Gm methylcellulose Bot 12 oz Tabs 500 mg Bot 100s

Use: Orally, bulk laxative

HYDROLYSIN (Merck Sharp & Dahme) Fibrinolysin (human) 50 000 u/vial with 10 cc water

Use: Intravascular blood clots (Dissolve in 1D cc water dilute to 250 cc for IV Inj)

***HYDROMADINONE ACETATE,** (Eli Lilly) 6a-chloro 17a-acetoxypregesterone acetate

Use: Progestational compound

- * **HYDROMORPHONE**. Dihydromorphinone Dilaudid hydrochloride.
- HYDROPNEH**. (Amtra) *o*-Phenylphenol mercuric nitrate. Oint. 1 oz., 4 oz., 1 lb.
Use: Topically, antiseptic ointment.
- HYDROPHILIC OINTMENT, U.S.P.** Stearyl alcohol, white petrolatum, propylene glycol, Sod. lauryl sulfate & water.
Morgan Sobolal Prod. Co.—16 oz., Schiefelin—1 lb., 5 lb.
See: Plashbase. (Squibb)
Polysorb. (Faugera)
- HYDROPRES**. (Merck Sharp & Dahme) Hydrochlorothiazide 50 mg., reserpine 0.125 mg./Tab. Also hydrochlorothiazide 25 mg., reserpine 0.125 mg./Tab. Bot. 100s & 1000s.
Use: (1 or 2 a day) Hypertension therapy.
- HYDROPRES-KA'-25**. (Merck Sharp & Dahme) Hydrochlorothiazide 25 mg., reserpine 0.125 mg., potassium chloride 572 mg./sugar-coated brown tab. Bot. 100s.
Use: Hypertension.
- HYDROPRES-KA'-5D**. (Merck Sharp & Dahme) Hydrochlorothiazide 50 mg., reserpine 0.125 mg., potassium chloride 572 mg./yellow tablet. Bot. 100s.
Use: Hypertension.
- HYDROQUIHONE MONOBENZYL ETHER**.
See: Benzoquin, Oint. (Elder)
- HYDROPTIC OPHTHALMIC SUSPENSION**. (Merck Sharp & Dahme) Hydrocortisone acetate 0.5% or 2.5% with neomycin. Susp. Dropper bottles 5 cc.
Use: Inflammation of the eye.
- HYDROSAL**. (Hydrosal Co.) A colloidal susp. of aluminum acetate 5%. Bot. 4 oz., 8 oz., 16 oz. and 1 gal. Oint. Jars ½ oz., 2 oz. and 1 lb.
- HYDROSPRAY, NASAL SUSPENSION**. (Merck Sharp & Dahme) Hydrocortone 1.0 mg., phenylephrine HCl 2.5 mg., propadrine HCl 7.5 mg., neomycin sulfate 75 mg./15 cc Bot. plastic spray 15 cc.
Use: Management of acute and chronic allergic disorders of the nose.
- HYDROSULPHOSOL** (Lientz) Calcium hydrol. An alkaline (pH 10.2) soln. of calcium pentathionate, sulfide, polysulfide & thiosulfate. It contains about 0.237 Gm. of the —SH (sulfhydryl) ion per 100 cc. Oint., Jar 1 oz., 1 lb.; Oraf, Vial 1 fl. oz.; Ophth. Oil, Vial 1 fl. oz., 3 fl. dr.; Sal., Bot 5 cc., 1 R. oz., 2 fl. oz., 4 fl. oz., 8 fl. oz.
Use: (Oral & topical). Various ocular conditions.
- HYDRO-TAR**. (Schiefelin) Two strength Hydrocortisone 0.5% or 1% with liquor
* Generic name recognized by NF, U.S.P. or A.M.A.
- carbowis detergent 3.83%/cream. Tube 15 Gm.
Use: Dermatoses.
- HYDROTEX**. (Texas Pharmacal) Absorption oint. base 1 lb., 5 lb.
Use: Ointment base.
- HYDROUS WOOL FAT, U.S.P., N.N.R. 1939, A.D.R. 1955**. See: Lanolin. (Various Mfr.)
- HYDROXAL** (Blue Line) Aluminum hydroxide susp. Bot. 1 pt., 1 gal.
Use: Orally, antacid.
- * **HYDROXINDASATE HCl**. (Merck Sharp & Dahme) 5-acetoxy-3-(2-aminoethyl)-1-(*p*-methoxybenzyl)-2-methylindole HCl.
- * **HYDROXYCOBALAMIN**. α -(5,6-dimethylbenzimidazolyl)hydroxocobamide. See: AlphaREDISOL (Merck & Co., Inc.)
- HYDROXY ALUMINUM MAGNESIUM AMINOACETATE**. See: Glycamine-AM, Tab. (Mermelin)
- * **HYDROXYAMPHETAMINE HYDROBROMIDE, U.S.P., N.N.D. 1962**. Ophth. Soln., U.S.P. See: Paredrine Hydrobromide (Smith, Kline & French)
- 2-HYDROXYBENZAMIDE**. See: Salicylamide, N.N.D. 195E
- * **HYDROXYCHLOROQUINE SULFATE, U.S.P., N.N.D. 1962**; Tab, U.S.P. 7-Chloro-4-[4-(*N*-ethyl-*N*-beta-hydroxyethylamino)-1-methylbutylamino]quinoline sulfate. See: Plaque nil, Tab. (Winthrop Labs.)
Labs)
W/Chloroquine, quinacrine, See: Triquine, Tab. (Winthrop Labs)
- 17-HYDROXYCORTICOSTERONE**, See: Hydrocortisone Preparations (Merck Sharp & Dahme)
- HYDROXYDIONE SODIUM**. See: Viadri, Vial (Pfizer)
- * **HYDROXYDIONE SODIUM SUCCINATE, N.N.D. 1962**. See: Viadri (Pfizer)
- HYDROXYDIPHENINE HCl**. (Chemo Pura) Dimethylaminoethyl-diphenyl-hydroxyacetate HCl. (Diphemin) Similar to Adiphenine
- * **HYDROXINDASOL HCl**. 5-hydroxy-1-(*p*-methoxybenzyl)-2-methyltryptamine HCl. See: Presid (Merck)
- * **HYDROXYCOBALAMIN**. (Merck & Co) α -(5,6-dimethylbenzimidazolyl)-hydroxocobamide. See: alphaRedial.
- HYDROXYISOINDOLIN**. Under study.
Use: Hypertension.
- HYDROXY MAGNESIUM AMINOACETATE, W/Mepapyritens HCl, Scopolamine HBr, benzocaine**. See: Banousea, Tab. (Grant)
- HYDROXYMYCIN**. An antibiotic substance obtained from cultures of *Sireptomycetes paucisporogenes*.
- * **HYDROXYPHENAMATE**. 2-hydroxy-2-

phenylbutyl carbamate See: Lisica Tabs (Armour)

o-HYDROXYPHENYLMERCURIC CHLORIDE, ADR W/Thymal, cassia oil, boric acid & kaolin, NNR 1952 See: Myaleto, Pow (Abbott)

***HYDROXYPROGESTERONE CAPROATE**, NND 1962
See Delafutin, Vial (Squibb)
Pradox (Upjohn)

W/Estradiol valerate See Deluleval Amp (Squibb)

R-HYDROXYQUINOLINE SULFATE (Mal Inckrad) See Oxyquinoline Sulfate (Varr aus Mfr)
Use: As an antiseptic

S-HYDROXY SALICYLIC ACID. See: Genlissic Acid (Various Mfr)

***HYDROXYSTENOZOLE**, [Winthrop Lab] 17b hydroxy 17a methylandro-4-eno [32 c] pyrazole

***HYDROXYSTILAMIDINE ISETHIONATE**, U.S.P., NND 1962 (Merrell) 4,4 Oxamidino 2 hydroxystilbene di(beta hydroxy ethane sulfonate) Amp (225 mg) 20 cc, Box 1s, 5s See also Stilbamidine Isethionate (Merrell)
Use: I.V. 225 mg, may be repeated at 24 hour intervals; blastomycosis, leishmaniasis & Indian kala azar

***HYDROXYZINE HCl**, NF NND 1962 Tab, N.P. 1 (p Chlorobenzhydryl) 4 [2 hydroxyethoxyethyl] piperazine di HCl
See: Atarax, Tab (Roerig)
Vistoril Stereject, Parenteral (Pfizer)

W/Ephedrine sulf
See: Marax Syr (Roerig)
Marax Tab

W/Oxyphenyclimine HCl See: Enarax, Tab (Roerig)

W/Pentaerythritol tetranitrate See: Cartrax, Tab (Roerig)

W/Prednisolone See: Ataraxid, Tab (Pfizer)

W/Theophylline ephedrine sulfate See: Marax, Tabs (Roerig)

***HYDROXYZINE PAMOATE**, NND 1962 1,1' methylene bis (2 hydroxy 3 naphthalene carboxylic acid salt of 1-(p chlorobenzhydryl) 4 [2 (2 hydroxyethoxyethyl) piperazine
See: Vitoril, Susp (Pfizer)

HYDROZITS (Merck Sharp & Dohme) Hydrocortisone 5 mg bacitracin 50 u, tyrothricin 1 mg, neomycin sulf 5 mg, benzocaine 5 mg /trache Bot 12s
Use: Throat and mouth infections

HYDRYLIN, (Searle) Diphenhydramine 25 mg, aminophylline 100 mg /Tab or R cc.

*Generic name recognized by NF, U.S.P. or A.M.A.

of Elixir Elixir, Bot 1 pt, 1 gal, Tab, Bot 100s, 1000s
Use: Orally, antihistaminic

HYDRYLIN COMPOUND, (Searle) Aminophylline 25 mg, diphenhydramine 6.25 mg, Pot iodide 15 mg, chlorofarm B mg & sugar & alcohol/4 cc Bot 1 pt, 1 gal
Use: Orally, antihistaminic

HYDRYLIN W/RACEPHEDRINE HYDROCHLORIDE, (Searle) Diphenhydramine 25 mg, aminophylline 100 mg, racephedrine HCl 25 mg /Eab Bot 100s, 1000s
Use: Antihistaminic with vasoconstrictor

HYEALIN, (Plessner) Ephedrine sulfate 10 mg, aspirin 3 gr, phenobarbital 10 mg Cal gluconate 100 mg, hyoscyamus ext pow 10 mg /Tab Bot 100s, 1000s
Use (I I d) Sympathomimetic

HYELAVIN, NND 1962 (Endo) Methylol riboflavin Vial (10 mg /cc) 10 cc
Use: I.V., I.M., vitamin B 2 therapy

HYEFEM DOUCHE POWDER, (Bilhuber Knoll) Benzalkonium chloride, potassium alum with buffers, aromatics and excipients Jar 4 oz, 10 oz
Use: Astringent and caudic vaginal douche

HYGIENIC POWDER. See: Bo Car Al, Pow (Merck Sharp & Dohme)

HYGLOX, (Lorre) Oxyquinoline sulf 50 mg /Tab Vials 12s
Use: (1 per qt makes 12000) Vaginal douche

HYGROMYCIN, (Lilly) Antibiotic (from *S. hygroscopicus*) used for the control of worms in hogs Available as Hygromix

HYGROTON (Geigy) 3 Hydroxy 3 (4 chloro 3 sulfamylphenyl) phthalimidine Chlorothalidone 100 mg /Tab Bot 100
Use: Hypertension and diuretic

HYXINONE, NND 1962 (Abbott) Menadiolone Sod bisulfite Amp (25 mg) 0.5 cc, Box 6s, 25s, 100s (5 mg) 1 cc, Box 6s, 25s, 100s (10 mg) 1 cc, Box 6s, 25s, 100s (7.2 mg) 10 cc, Box 3s, 25s, 100s Tab. 5 mg Bot 100s, 1000s
Use: Vitamin K therapy

HYXOLEX, (Chicago Pharm) Dehydracholic acid 250 mg /Tab Bot 100s, 1000s
Use: Hydrocholeretic

HYLAND LIQUID PLASMA (Hyland) Normal human blood plasma 300 cc liq u cit rated, ready for use
Use: I.V.

NYMORFAN-ATROPINE (Endo) Dihydromorphinone NCl 2 mg, atropine sulfate D 25 mg /cc Vial 10 cc Box 3s
Use: Analgesic with antispasmodic

NYMORFAN HYDROCHLORIDE, (Endo) Dihydromorphinone HCl Tab 2.5 mg Tube

20s, Bot. 100s Inj. {2 mg./cc} Vial 10 cc.
Use: Orally & S.C., analgesic & cough
sedative.

HYMOSA. (Walker Pharm.) Sod. salicylate
12 gr., Sod. citrate 8 gr., Pot. iodide 4 gr.,
caffeine citrate 8 gr., colchicum ext. 5 gr.,
cimicifuga fr. 48 min./oz. Bot. 1 gal.

Use: Salicylate therapy.

HYODIH. (Gardner) Iodine 0.85 gr., os
H₂O 5 cc. Bot. 4 fl. oz., 8 fl. oz.

Use: Iodine therapy.

HYOSCINE ALKALOID W/Chloral hydrate,
atropine, hyoscyamine. See: Chloradonna,
Cap. (Ulmer)

HYOSCINE AMINOXIDE HBr.

See: Scalamine, Tab. T.D. (Nysco Labs.)

Scalen Unicelloc. (Miss Pharm.)

HYOSCINE HYDROBROMIDE, U.S.P., N.N.R.

1948. Scopolamine HBr. See: Alcon Isopto
0.25% Sol. Bot. w/dropper 15 cc.

See: Tables for combinations.

W/Atropine and hyoscyamine.

See: Atropine sulfate table.

Belladonna alkaloids.

HYOSCINE HYDROBROMIDE. (Crookes-
Barnes) 0.2% and 0.5%/Bot. 4 cc.

Use: Mydriasis.

Hyosine Hydrobromide Combination
on pp. 376, 377, end 378.

HYOSCINE-HYOSCYAMINE-ATROPINE.

See: Atropine w/hyoscyamine w/hyosine.

HYOSCINE METHOBROMIDE. See: Meth-
scopolamine Bromide

HYOSCYAMINE. *Levo form of atropine.*

See: Bellafoline preps. (Sandoz)

Levsin preps. (Kremers-Urban)

W/Butobarbital. See: Levamine, Tabs., Caps.
(Meyer)

W/Chloral hydrate, atropine, hyosine. See:
Chloradonna, Cap. (Ulmer)

W/Methenamine, Cal. mandelate & atropine
sulfate. See: Urolate, Tab. (Borchardt)

W/Phenobarbital, aspirin & acetophenetidin.
See: Hycetin, Tab. (Vanpel & Brown)

W/Phenobarbital, atropine sulfate, hyosine
HBr, activated charcoal. See: Barcole-B,
Tabs. (Haag)

HYOSCYAMINE-ATROPINE-HYOSCINE.

See: Atropine w/hyoscyamine w/hyosine.

HYOSCYAMINE CAMPHORATE W/Scopola-
mine camphorate. See: Vesano, Tab.
(Schering).

***HYOSCYAMINE HYDROBROMIDE, N.F.**
(Various Mfr.) Darvane HBr. See: Table for
combinations.

W/Atropine, hyosine HBr, phenobarbital.
See: Atropine Sulfate Tablets.

W/Atropine, phenobarbital & scopolamine
HBr. See: Atropine Sulfate Tablets.

* Generic name recognized by N.F., U.S.P. or A.M.A.

W/Atropine sulfate. See: Atropine Sulfate
Tables.

W/Bromisovaleryurea, homatropine methyl-
bromide & scopolamine HBr. See: Tadano,
Tab. (Tablerock)

W/Dehydrocholic acid, mag trisilicate, alu-
minum hydroside. See: Barmal, Cap. (Paul
Maney)

W/Dehydrocholic acid, Prolase-300, Mylase-
100, hyosine HBr, atropine sulfate. See:
Almezyme, Cap. (Meyer)

W/Dihydrocodeinane bitartrate, orthoxin
HCl & Sod. citrate. See: Orthoxin Liq.
(Upjohn).

W/Ethisterone, ammonium chloride, d-am-
phetamine sulfate. See: Gymens, Tab. (Co-
lumbus)

W/Homatropine methylbromide. See: Nava-
donna, Sol. (Sherman).

W/Mephenesin, glutamic acid & pheno-
barbital. See: Kused, Cap. (Kremers-Urban).

W/Phenobarbital. See: Scyophen, Tab. &
Elix. (Kremers-Urban).

W/Phenobarbital, atropine sulfate & Mag
trisilicate. See: Belnesium, Tab. (Haskell).

W/Phenobarbital, homatropine methylbro-
mide & hyosine HBr. See: Navadonnol,
Tab. (Sherman).

W/Phenylsodiuminapyridine HBr, atropine
sulfate, scopolamine HBr. See: Urogestic,
Tab. (Edwards)

W/Phyostigmine salicylate. See: Phyotro-
mine H, Amp. & Vial (Kremers-Urban).

W/Troxigmine methyl sulfate & benzyl al-
cohol. See: Kelagen, Vial (Paul Maney)

W/Pyrilamine maleate & hyosine HBr. See:
Almahist Syrup (Meyer Chemical).

W/Pyrilamine maleate, hyosine HBr, d-
amphetamine sulfate. See: Histalme Dura-
caps. (Meyer)

W/Pyrilamine maleate, hyosine HBr, op-
pirin, acetophenetidin & caffeine. See:
Almegetic, Tab. (Meyer Chemical).

W/Pyrilamine maleate, hyosine HBr &
d-amphetamine sulfate. See: Almahist, Cap.
(Meyer).

W/Scopolamine HBr, aluminum hydroxide,
mag. hydroside, bulkagen. See: Hycodros,
Tab. (Columbus)

HYOSCYAMINE HYDROCHLORIDE. (Vari-
ous Mfr.)

HYOSCYAMINE MALEATE. See: Bellafoline
(Sandoz)

HYOSCYAMINE SALTS W/Atropine salts.
See: Atropine W/Hyoscyamine.

***HYOSCYAMINE SULFATE, N.F.** (Various
Mfr.).

See: Donnzyme (Robins)

Levsin Sulfate, Tab. (Kremers-Urban)

Hyoscine Hydrobromide Combinations

[376]

| Hyoscine HBr | Others | Manufacturer | Brand |
|--------------|---|--------------------|----------------------------|
| 0.075 mg | N Acetyl p-aminophenol 300 mg, butobarbital sod 10 mg | Denab | Pallacin Tab |
| 0.65 mg | Aluminum hydroxide 400 mg, mag hydroxide 100 mg, l hyoscyamine HBr 0.125 mg, bulkogen 50 mg | Calumbus | Hyocdrax Tab |
| 0.02 mg | Atropine methylnitrate 1 mg, mephobarbital 65 mg, homatropine methylbromide 5 mg | Nysca Labs | Spascaps Tab T D |
| 0.02 mg | Atropine methylnitrate 1 mg, mephobarbital 65 mg, homatropine methylbromide 5 mg | Hiss Pharm | Trascine Un celled Cap |
| 0.0065 mg | Atropine sulf 0.0194 mg, dehydrated acid 250 mg, Hyoscyamine sulf 0.1037 mg, phenobarbital 16.2 mg | Whittier | Bellabul, Tab |
| 0.0086 mg | Atropine sulf 0.0258 mg, hyoscyamine sulf 0.1384 mg, acetyl p-aminophenol 2.5 gr, salicylamide 2.5 gr, aspirin 2.5 gr, d-amphetamine sulf 5 mg, butobarbital 1/2 gr | Stuart | Stuart Prolar 8 Tab |
| 0.0065 mg | Atropine sulfate 0.0194 mg, hyoscyamine sulfate 0.1037 mg, buta-barbital Sod 16.2 mg | Zemmer | Zemarine Elix / dr |
| 0.0065 mg | Atropine sulfate 0.02 mg, hyoscyamine HBr 0.1 mg, dehydrated acid 100 mg, prolose 300.30 mg, mylose 100.30 mg | Meyer | Almeszyme, Cap |
| 0.0193 mg | Atropine sulf 0.06 mg, hyoscyamine sulf 0.31 mg, phenobarbital 48.6 mg | Raberis | Meinal Tab |
| 0.0065 mg | Atropine sulf 0.0194 mg, hyoscyamine sulf 0.1037 mg, pheno-barbital 16 mg | Dumas Wilson | Anspas, Tab or Elix / 5 cc |
| 0.0032 mg | Atropine sulf 0.0097 mg, hyoscyamine sulf 0.0318 mg, pheno-barbital 8.1 mg, pepsin 150 mg, pancreatin 300 mg, bile salts 150 mg | Robins | Dennazyme Tab |
| 0.0074 mg | Atropine sulfate 0.025 mg, Hyoscyamine sulfate 0.1286 mg, W/Phenobarbital 16 mg in core | VanPelt & Brown | Barbidanna D A, Tab |
| 0.22 mg | Atropine sulf 0.16 mg, phenobarb 32 mg | Ives Cameron | Campamal Tab |
| 0.00325 mg | Atropine sulf 0.0097 mg, phenobarbital 8.1 mg, hyoscyamine sulf 0.0519 mg, aluminum hydroxide magnesium carbonate 5 gr | Bentan | Almadan Tab |
| 0.0065 mg | Atropine sulf 0.0195 mg, sulfacetamide 500 mg, phenylazodi-aminopyridine HCl 50 mg, hyoscyamine sulf 0.1040 mg | Vitamix | Pyloscetamide Tab |
| 0.0065 mg | Atropine sulf 0.0195 mg, sulfacetamide 500 mg, phenylazodi-aminopyridine 50 mg, hyoscyamine sulf 0.104 mg | Richard & Williams | Uricin Tab |
| 0.325 mg | Benadryl 25 mg | Park, Davis | Benacine Tab |
| 0.5 mg | Benztacine 100 mg, methapyridene HCl 10 mg, mag hydroxy-aminoacetate 100 mg | Grant | Banoxase, Tab |

Hyoscine Hydrobromide Combinations

| Hyoscine HBr | Other | Manufacturer | Brand |
|--------------|--|-----------------|------------------------------|
| 1/400 gr. | Pentobarbital ½ gr. | Walker Corp | Pentacaine, Tab |
| 1/200 gr. | Pentobarbital ½ gr., hexestrol 3 mg | Walker Corp | Tal Ertamine, Tab |
| 0.0195 mg | Pentobarbital sod 48.6 mg., hyoscyamine sulf 0.3113 mg., atropine sulf 0.0582 mg | Columbus | Belkoids TT, Tab |
| 0.08 mg | Phenobarbital 15 mg., barbitol Sod 45 mg., sandoptal 25 mg dihydroergalamine 0.16 mg | Sandoz | Plexonal Tab. |
| 0.22 mg | Phenobarbital 32 mg., atropine sulfate 0.16 mg | Ives Cameron | Compamal, Tab. |
| 1/500 gr. | Phenobarbital ½ gr., atropine sulfate 1/2000 gr | Haack | Cymital, Tab |
| 0.2 mg | Phenobarbital 15 mg., atropine sulfate 0.1 mg., benzocaine 100 mg., Vit B complex | Winthrop Labs | Apalamine, Tab |
| | Phenobarbital See: Atropine—Hyoscine—Hyoscyamine Combina tions | | |
| 0.4 mg | Phenyltolazamine dihydrogen citrate 25 mg., nlocinamide 25 mg., benzocaine 100 mg | Tutag | Hyloxine, Tab |
| 0.4 mg | Phenyltolazamine dihydrogen citrate 25 mg., pyridoxine HCl 10 mg methamphetamine HCl 1.25 mg | Lenman | San neus, Tab |
| 0.15 mg | Pheniramine 15 mg | Nhan | Nio Piracene, Tab |
| 0.15 mg | Pheniramine maleate 10 mg w/aluminum hydroxide gel & ethyl aminobenzoate | Davis & Sly | Dapren, Tab |
| 0.008 mg | Pyrimamine maleate 25 mg., hyoscyamine HBr 0.04 mg | Meyer | Almehist, Syr |
| 0.016 mg. | Pyrimamine maleate 50 mg., hyoscyamine HBr 0.08 mg., d amphet amine sulfate 1.5 mg | Meyer | Almehist, Cap |
| 0.008 mg | Pyrimamine maleate 25 mg., hyoscyamine HBr 0.04 mg A.P.C | Meyer | Almegestat, Cap. |
| 0.0042 mg | Pyrimamine maleate 12.5 mg., salicylamide 3½ gr., acetophenetidin 2½ gr., caffeine ½ gr., hyoscyamine sulf 0.0702 mg., atropine sulf 0.0140 mg | VanPelt & Brown | Pyr Aspra, Tab. |
| 0.006 mg. | Reserpine 0.1 mg., hyoscyamine sulf 0.1 mg., atropine sulf 0.2 mg | Lenman | Redona, Tab or Elix./5 cc |
| 0.0074 mg | Reserpine 0.1 mg., hyoscyamine sulf 0.1286 mg., atropine sulf 0.025 mg | VanPelt & Brown | Raudonna, Tab |
| 0.0035 mg | Sulfacetamide 250 mg., phenylazo-diamine pyridine HCl 50 mg., atropine sulf 0.0097 mg., hyoscyamine sulf. 0.05185 mg | Columbus | Sulfid B A, Tab |

- Levsinex, Tab. (Kremers-Urban)
- W/Acetophenolidin, aspirin, phenobarbital, prophenpyridamine maleate, phenylephrine HCl. See: Phenaphen Plus, Tab. (Robins)
- W/N-acetyl-p-aminophenol, salicylamide, acetylsalicylic acid, d-amphetamine sulf., butobarbital, atropine sulf., hyoscine HBr. See: Stuart Prolar-B, Tab. (Stuart)
- W/Acetylcarbromal, atropine sulf. & hyoscine HBr
See: Disperdon, Tab. (Standard)
- W/d-Amphetamine sulfate, phenacelin, salicylamide, amobarbital. See: Sedamosal, Cap. (Finley)
- W/Atropine, methamphetamine, phenyl salicylate, methylene blue, benzoic acid. See: Urisep-tin, Tab. (Blaine)
- W/Atropine sulf. & scopolamine HBr. See: Tables under Atropine Sulfate
- W/Atropine sulf., scopolamine HBr, buclizine HCl, Vit. B-6. See: Stuart Buclodia, Tab. (Stuart)
- W/Atropine sulfate, hyoscine HBr & phenobarbital. See: Tables under Atropine Sulfate.
- W/Atropine sulfate, phenobarbital. See: Tables under Atropine Sulfate.
- W/Atropine sulfate, hyoscine, phenobarbital. See: Tables under Atropine Sulfate.
- W/Dehydracetic acid, atropine sulfate, hyoscine hydrobromide, phenobarbital. See: Bellabil, Tabs. (Whittier)
- W/Homatropine methylbromide, pyrilamine maleate, pamabrom, scopolamine HBr, methamphetamine HCl. See: Neaporbol, Tab. (Central)
- W/Hyoscine HBr, atropine sulfate, sulfacetamide, phenylazodiaminopyridine. See: Uricla, Tab. (Richard & Williams)
- W/Methocarbamol, phenacelin, acetylsalicylic acid, phenobarbital. See: Robaxisol-PH, Tab. (Robins)
- W/Pentobarbital sodium, Atropine sulfate, Hyoscine HBr. See: Relokoids TF, Tab. (Columbus)
- W/Phenobarbital. See: Levsinex-Phenobarbital.
- W/Phenobarbital, atropine sulf., hyoscine hydrobromide. See: Bar-Cy-A-Tab, Tab. (Sul-liff & Case)
- W/Phenobarbital, atropine SO₄, hyoscine HBr. See: Barbidanna-CR, Tabs. (Vampell & Brown)
- W/Pyrilamine maleate, salicylamide, acetophenetidin, caffeine, atropine sulfate, hyoscine hydrobromide. See: Pyr-Aspra. (VanPelt & Brown)
- W/Reserpine, atropine sulf., hyoscine hydrobromide. See: Redona, Tab., Eliz. (Lemmon)
- W/Secobarbital. See: Levsin Sulfate W/Secobarbital, Tab. (Kramer-Urban)
- W/Sulfacetamide, phenylazo-diamino-pyridine HCl, atropine sulf., hyoscine HBr. See: Sulhd B-A, Tab. (Columbus)
- HYOSCYAMUS EXTRACT, N.F.**
Lilly—Bot. 1 oz., 4 oz.
- W/Acet-Dio-Mer Sulfonamide. See: Suro-mate, Tab. (Patch)
- W/Acetophenetidin, aspirin, caffeine & gelsemium ext. pow. See: Henasphen, Cap. (Drug Products).
- W/Aminopyrine, acetophenetidin & caffeine. See: Amidaphen, Pulv. (Lilly)
Andofive, Cap. (Massengill)
- W/Arbutin, chromium sulfate, triticum ext. & echinacea ext. See: Pasta-Cole, Tab. (Cafe Chem. Co.)
- W/Aspirin & acetophenetidin. See: Hosac, Tab. (Haskell).
- W/Aspirin, nux vomica ext., Gelsemium tr. & cascara sagrada ext. See: A.A.S. Compound, Tab. (Tilden).
- W/Atropine sulfate, hexamethylenamine, solol & benzoic acid. See: Cystarrhal, Tab. (Buffington).
- W/Barbital, chamanil & passiflora. See: Insomnilol, Tab. (Buffington).
- W/Barbital, passion flower ext. & valerian ext. See: Neuroids, Tab. (Buffington; Tilden).
- W/Chloral hydrate & pot. bromide.
See: Bromide-Chloral Comp. (Merck Sharp & Dohme)
Bromidia Liq. (Bottle)
Hypno-Bromic, Sol. (Wampole)
- W/Cocaine phosphate. See: Glykeron, cough preparation (M. M. Smith).
- W/Hexamethylenamine, Sod. acid phosphate, triticum ext. & adiphenine HCl. See: Hexosed, Tab. (Mallard).
- W/Kaolin. See: Maalin No. 2 (The Alphaden Co.)
- W/Paragaric, Ipecac ayr., licorice ext. See: Crupic, Tab. (Jenkins Labs.)
- W/Sulfacetamide, sulfamethylthiadiazole. See: Uronamide, Tab. (Flint, Eaton)
- HYOSCYAMUS FLUID EXTRACT.** Each 100 cc. yields not less than 0.035 Gm. & not more than 0.045 Gm. of hyoscyamus alkaloids.
Lilly—Bot. 1 pt., 1 gal.
Sharp & Dahme—Bot. 1 pt., 1 gal.
- W/Pot. bromide. See: U.S.T., liq. (Tilden).
- Hyoscyamus Products & Phenobarbital Combinations** on pp. 38D and 38I.
- HYOSCYAMUS TINCTURE, N.F.**
W/Phenobarbital, Sod. bromide & Pot. bromide. See: Sedalca, Tab. (Mallard).
- W/Sod. bromide, strontium bromide, ammonium bromide & valerian tr. See: Neuro-bromyde, liq. (Rorer).
- HYOSID.** (Walker Corp) Phenobarbital ½

| Hyoscyamus | Phenobarbital | Others | Manufacturer | Brand |
|----------------------|---------------|---|-----------------|--------------------------|
| Ext., 1/6 gr, | 1/2 gr | | Walker | Fenbane, Tab |
| Ext., 1/6 gr, | 1/2 gr | | C D Smith | Phenobane, Tab |
| Ext., 1/6 gr | 1/2 gr | | Cole Chem Co | Phenocymus, Tab |
| HBr, 1/500 gr, | 1/4 gr | | Kremers Urban | Scyaphen, Elixir & Tab |
| 1/2 gr | 1/4 gr | Amniopyrine 2 1/2 gr | G F Harvey | Amifabs, Tab |
| Ext., 1/6 gr | 1/4 gr | Aspirin 5 gr | Rorer | Lumaspirin w/Hyoscy, Tab |
| 1/6 gr | 1/4 gr | Aspirin 5 gr | Massengill | Andasus, Cap |
| Alisaloids, 0.025 mg | 1/4 gr | Aspirin 2 1/2 gr acetophenethidin 2 1/2 gr | VanPelt & Brown | Hyoclin, Tab |
| Ext., 1/6 gr | 1/4 gr. | Aspirin 2 1/2 gr, acetophenethidin 2 1/2 gr | Haskell | Hasamal, Tab |
| Ext., 1/6 gr | 1/4 gr | Aspirin 3 gr, acetophenethidin 2 gr | Frast | Acetaphen Tab |
| Ext., 1/6 gr | 1/4 gr | Atipurin 3 gr, phenacetin 3 gr | Bentex | Anoben, Cap |
| Ext., 1/6 gr | 1/4 gr | Aspirin 2 1/2 gr, phenacetin 1 1/2 gr | Massengill | Analdyne Cap |
| Ext., 1/16 gr | 1/2 gr Sod | Atropine sulfate 1/1000 gr, Mag trisilic 7 gr, Sod bicarb 2 1/2 gr | Lemman | Aitmag Tab |
| 0.1266 mg | 16 mg | Atropine sulf 0.025 mg hyoscyne HBr 0.0074 mg | VanPelt & Brown | Barbidanna O A, Tab |
| Ext., 1/2 gr | 1/4 gr | Belladonna ext 1/2 gr | Metra | Spasticon Tab |
| Ext., 1/2 gr, | 1/2 gr | Camphor cypripedium, valerian | Noyes | Restaphen, Tab |
| Ext., 10 mg | 10 mg | Ephedrine sulfate 10 mg, aspirin 3 gr, Cal gluconate 100 mg | Piessner | Hyefalin, Tab |
| Tr, | 1/16 gr Sod | Glyceryl trinit 1/100 gr, Pot nitrate bromides | Vita | Nitrobrom, Tab |
| 0.54 mg, | 3/4 gr. | Homatropine methylbromide 0.0447 mg sco polamine HBr 0.0144 mg, thiamine HCl 3 mg | Paul Maney | Sedlin Unoday, Cap |
| Ext., 1/2 gr | 1/4 gr. | Mag hydroxide 7 1/2 gr Mag oxide 7 1/2 gr, kaolin 22 gr, agar, karaya, pepsin, pepsin, parmit | Alphaden | Maalin No 3L, Tab |
| Ext., 1/25 gr, | 1/4 gr, | Passiflora 1 gr, valerian 1 gr. | Buffington | Lumiflora Cpd, Tab |
| Ext., 8 mg | 15 mg | Passiflora 1 gr, valerian 1 gr | Drug Prod | Sed O El, Cap |
| 1/2 gr | 1/4 gr | Passiflora ext 1/4 gr, valerian ext 1/4 gr | National Drug | Sed A Nat, Tab |
| Ext., 1/4 gr | 1/2 gr | Passiflora 1/4 gr, valerian 1/4 gr, camphor monobromate 1/4 gr | Vale | Nervofase, Cap |

Hyoscyamus Products & Phenobarbital Combinations

| Hyoscyamus | Phenobarbital | Others | Manufacturer | Brand |
|--------------|---------------|---|---------------|-------------------|
| Ext., ½ gr. | ¼ gr. | Passiflora fr. 1 min., chamomilla 1 min., pulsor-tillo fr. 2 min., Sod. bicarb. ½ gr. | C. D. Smith | Passi-Barb., Tab. |
| Ext., ½ gr. | ½ gr. | Passiflora, valerian, camphor, sumbul | Tallby-Nelson | Repasans, Tab. |
| Tr., 10 min. | ¼ gr. | Sod. bromide 4 gr., Pot. bromide 3½ gr. | Mallard | Sedalta, Tab. |
| Ext., ½ gr. | ½ gr. | Salicylamide 3 gr., dl-desoxyephedrine HCl 1/20 gr. | Lehman | Aspifedrine, Tab. |
| Tr., 10 min. | ¼ gr. | Thyroid ¼ gr. | Mallard | Sedocrine, Tab. |

gr., aminopyrine 1 gr., hyoscyamus ext. 1/6 gr./Tab. Bot. 100s.

Use: (1 q 3 h) Antispasmodic & sedative.

HYOSED ELIXIR. (Klug) Phenobarbital 1d.2 mg., hyoscyamine sulfate 0.1037 mg., atropine sulfate 0.0194 mg., hyoscine HBr 0.0065 mg./isp. Bot. 4 oz.

Use: Sedative

HYOSED T.B. (Klug) Hyoscyamine HBr 0.256 mg., hyoscine HBr 0.0144 mg., atropine sulfate 0.0480 mg., phenobarbital 32.0 mg /Timed disintegration tab. Bot. 100s.

Use: Sedative

HYTOLE-12 SYRUP. (Merck Sharp & Dohme) Vit B-12 cryst. 25 mcg., ferrous sulfate 1.3 Gm., thiamine HCl 10 mg., riboflavin 2 mg., pyridoxine HCl 1 mg., pantothenol (equiv. to 2.32 mg. cal. pantothenate) 2 mg., niacinamide 30 mg., choline chloride 20 mg., folic acid 5 mg., liver fraction #2, N.F. 1 Gm./30 cc. Bot. 1 pt., 1 gal.

Use: Hematinic.

HYPAN. (Hoack) Aspirin 5 gr., precipitated calcium carbonate 1½ gr., dried aluminum hydroxide gel 1 hr., anhydrous citric acid ½ gr./Tab

Use: Analgesic.

HYPANALDYNE. (Chicago Pharmacal) Phenobarbital ½ gr., hyoscyamine sulfate 0.04 mg., atropine sulfate 0.00225 mg., scopolamine HBr 0.06 mg./Tab. Bot. 100s, 500s, 1000s.

Use: (1-2 m.p.) Antispasmodic.

HYPACUE-M, 75%. (Winthrop Labs.) 15 Gm. Vials. 20 cc 37.5 Gm. Vials 50 cc.

HYPACUE-M 90%. (Winthrop Labs) Sed salt of n-methylglucamine salt of 3,5-diacetamido-2,4,6-triiodobenzoic acid 45 Gm / 50 cc. Vial

Use: X-ray contrast medium for blood vessels of heart and thorax

HYPACUE SODIUM, N.N.D. 1962. (Winthrop Labs) Diatrizoate Sod., Sod 3,5-diacetamido-2,4,6-triiodobenzoate. Amp. (50% sol) 30 cc., Box 1s, 10s, 25s. Vial (90%) 50 cc. Oral Paw. Cons 250 Gm.

Use (I.V. slowly); radiopaque medium for excretory urography.

HYPACOTIN INFECTION. (Cutter) Gamma globulin 165 mg/cc. Amp. 1.5, 4.5 cc. Use: I.M., prevention and treatment of mumps.

HYPERCILLIN. (Cutter) Procaine penicillin G 300,000 u/cc. Vial 10 cc
Use: I.M.; penicillin therapy.

- HYPERCILLIN PROLONGED** (Cutter) Procaine penicillin G 300 000 u in oil w/2% aluminum monostearate/cc of susp Vial 1 cc., 10 cc
Use I.M., prolonged penicillin therapy
- HYPERLOID.** (Person & Covey) Rauwolfia serpentina whole root Tob 2 mg Bot 100s
Use: Hypotensive & tranquilizing agent
- HYPER-RU** (Sutliff Case) Na 1 Rutin 5 mg, Pot thiocyanate 1½ gr/5 cc; Na 2 Rutin 5 mg, Pot Thiocyanate 3 gr/Tab; Bot 100s, 1000s
Use: Orally capillary permeability therapy
- HYPER-SED.** (Sutliff & Case) Pot thiocyanate 97 mg, phenobarbital 32.5 mg/Tab Bot 100s 1000s
Use: Sedative
- HYPERTENCIN** (CMC) Mannitol hexanitrate 32 mg/Tab Bot 30s, 100s, 250s, 1000s
Use: Hypertension
- HYPERTENCIN MILD.** (CMC) Mannitol hexanitrate 32 mg, phenobarbital 16 mg/Tab Bot 30s, 100s, 250s, 1000s
Use: Hypertension
- HYPERTENSIN-CIBA** (Ciba) Angiotensin Amide 2.5 mg/Vial as pow Box 5s
Use: Hypertensive agent for treatment of shock and circulatory collapse
- HYPERTENSION DIAGNOSIS.**
See: Benodolol HCl, Amp (Merck Sharp & Dohme)
Regline, Amp, Tob (Ciba)
- HYPERTENSION THERAPY.**
See: Ansoleson (Wyeth)
Aprexoline, Amp, Tob (Ciba)
Artanod (Haffmann La Roche)
Dibenzyline HCl, Cop (Smith, Kline & French).
Ecolid, Tab (Ciba)
Hesperidin Methyl Chalcane (Various Mfr).
Hexamethanum Chloride & Bromide (Various Mfr).
Hyderygine, Amp, Tob (Sandoz)
Inversine, Tab (Merck Sharp & Dohme)
Potassium Thiocyanate
Priscoline HCl, Preps (Ciba).
Protoveratrine A & B
Bouwaiffo Serpentina (Various Mfr)
Besperine (Various Mfr)
Eutin (Various Mfr).
Tetraethylammonium Chloride (Various Mfr).
Vasodilators.
Veralbo, Vial, Tob (Pitman Moore).
Veratrum Albo
- Veratrum Viride
Verifaid, Preps (Riker)
- HYPERTENS N** (Ciba) Angiotensin amide 2.5 mg, mannitol 4.7 mg, thiomersal 0.1 mg as lyophilized powder/5 cc 2.5 mg/Vial 5s
Use: Vascular collapse
- HYPERTET INJECTION** (Cutter) Tetanus immune globulin of human origin. Vial 1500 unit
Use: For immediate injection after injury
- HYPERTHYROIDISM** See: Antithyroid agents
- HYPERTUSSIS SERUM** (Cutter) Prepared from the gamma globulin fraction of serum Vial 2.5 cc
Use: I.M., immunization
- HYPER-VISERA LIQUID.** (Lloyd) Viscum album 25 Gr, mactocantha 19 gr/5 cc Bot liq 1 pt, 1 gal
Use: Orally, vasodilator
- HYPER-VISERA TABLETS.** (Lloyd) W/Rutin ½ gr, viscum album 25 gr, crotaegus 1/6 gr/Tab Bot 100s, 500s, 1000s
Use: Orally, vasodilator
- HYPYER-X.** (Tully) Reserpine 0.1 mg theobromine 325 mg, phenobarbital 10 mg/Tab Bot 100s
Use: Anxiety and hypertension
- HYPYER-ZEM NO. 2.** (Zemmer) Sod nitrite 1 gr, Sod nitrate 2 gr w/echinacea & ezebile/Tab Bot 100s, 1000s
Use: Orally vasodilator
- HYPNYLINE.** NNR 1957 Dyphylline NND 1958 (7 Dihydroxy propyltheophylline) See: Neophylline, Tab (Paul Money)
- HYPNALDYNE.** (Chicago Pharmacal Co) Phenobarbital ¼ gr, hyoscyamine sulfate 0.0016 gr, atropine sulfate 0.00029 gr scopolamine HBr D 001 gr/tab Bot 100s 500s, 1000s
Use: (1 2 prn) Sedative w/antispasmodic
- HYPNO-BROMIC.** (Wampole) Chloral hydrate 16 gr hyoscyamus ext 10 mg, Pot bromide 8 gr/5 cc Bot 16 oz
Use: Orally, sedative
- HYPNOGENE** See: Borbital (Various Mfr)
- HYPNORM, CAP.** (Massengill) An E.C. pill in each cap of 5 mg methamphetamine HCl admixed with secobarbital Sod 30 mg, phenobarbital 15 mg/Cap Bot 100s
Use: Hypnotic with central nervous system stimulant to be released in 8 hours
- HYPNOTICS.** See: Sedatives
- "HYPO."** See: Sodium Thiosulfate (Various Mfr).

HYPO-ALLERGIC MILK. (Wyeth) Pow., less allergenic by prolonged thermal processing, Con 1 lb. Liq., less allergenic by prolonged thermal processing. Coe 14.5 fl. oz.

HYPOBETA-20. (Merck Sharp & Dohme) Vit. B-1 10 mg., B-2 2 mg., B-6 1 mg., niacinamide 100 e.g., pantothenol 4 mg., liver inj. 20 mcg., B-12/cc. 0.05 cc., B-12 cryst. 20 mcg./cc. Vol 10 cc.

Use: I.M. (1-2 cc. daily) Vitamin B-complex therapy.

HYPOCAL. (Benton) Methopramine HCl 5 mg., pentobarbital 20 mg., Vit. A 3000 U.S.P. u., D 200 U.S.P. u., B-1 1 mg., B-2 1 mg., niacinamide 10 mg., C 100 mg., iron 15 mg./Tab. Bot. 100s.

Use: Obesity when dietary supplement is desired.

HYPOCHLORITE PREPS.

See: Antiformin

Clorapactin (Guordion)

Dakin's Soln.

Pexeral (Fesler Co)

Hyclorite, N.N.R. 1950

Zonite (Chemway Corp.)

HYPOGLYCEMIC AGENTS.

See: Chlorpropamide

DBI (U. S. Vit.)

Dibinase (Pfizer)

Orinose (Upjohn)

Ozinase (Upjohn)

Phenformin

Talbutamide

HYPOSULFITE. See: Thiosulfate.

HYPOTAN. (E. Fougere) Chloral hydrate $\frac{3}{4}$ gr., methylacetylcholine 1/32 gr., bromocholine bromide 1/12 gr./tab. Bot. 44s.

Use: Orally, sedative.

HYPOTENSIN. (Cy Lab)

Formula A: Pentolinium 25 mg. of base & 60 mg. of bitartrate, apresoline 20 mg., serpassil 0.2 mg., prostigmine 15 mg.

Formula B: Pentolinium 45 mg., apresoline 50 mg., serpassil 0.2 mg., prostigmine 15 mg.

Formula C: Pentolinium 80 mg., apresoline 90 mg., serpassil 0.2 mg., prostigmine 15 mg.

HYPOVITE. (Kirk) Vit. A 10,000 u., D 2000 u., B-1 10 mg., B-2 1 mg., B-6 2 mg., B-12 10 mcg., C 50 mg., niacinamide 30 mg / 2 cc. Amp. 2 cc.

Use: Vitamin supplement.

HYPRONE. (New Mexico Pharm.) Dipyrone 50% /cc. Liq. Vials, 30 cc.

HYPROTIGEN, SOLUTION 6%. N.N.D. 1962. (Oon Baxter) Protein hydrolysates, as enzymatic hydrolysate of casein containing

amino acids & polypeptides. Sol. Bot. 1000 cc.

Use: I.V., amino nitrogen therapy.

W/Dextrose 5%. Bot. 500 cc., 1000 cc.

Use: I.V.

HYPTAN. (Wampole) Dual release. Outside, phenyltoloxamine citrate 25 mg., secobarbital 60 mg.—core, phenyltoloxamine citrate 75 mg /Tab. Bot. 100s.

Use: Insomnia.

HYROCAIN. (Amer. Pharm.) Pyrilamine maleate 10 mg., tyrothricin 0.5 mg., benzocaine 10 mg./Gm. Oint. Tube 1 oz.

Use: Topically, antihistaminic w/antibiotic

HYRYE. (Ours) Riboflavin-5-phosphate Sod. Tab. 10 mg. Bot. 100s; Viol (50 mg./cc.) 10 cc.

Use: Dermatitis & Vit. B-2 deficiency.

HYSLIPTINE, A.D.R. (Warren-Teed) Methylbenzethonium chloride 12.8%. Bot. 1 oz., 1 pt., 1 gal.

Use: Disinfection of surgical instruments

HYSOBEL. (Zemmer) d-Desoxyephedrine HCl 5 mg., methylcellulose 2½ gr., thyroid 15 mg., phenobarbital 8 mg./Tab. Bot. 100s, 500s, 1000s.

Use (1 & 2) Antilethargy.

HYSSEH-AC. (Conright) d-Amphetamine sulfate 12 mg., pentobarbital sodium 65 mg / tab. Bot. 100s.

Use: Appetite control.

HYSTAL. (First Texas) Pentobarbital 18.125 mg., hyoscyamine sulfate 0.1296 mg., atropine sulfate 0.02425 mg., hyoscine HBr 0.00812 mg., thiamine HCl 12.5 mg./Tab. or dr. Liq. Bot. 1 pt., 1 gal. Tab. Bot. 100s, 500s, 1000s.

Use: (1 tsp. or 1 Tab. q 3 h.) Antispasmodic, sedative & hypnotic.

HYTAKEROL, N N O. 1962. (Winthrop Labs) Dihydrotachysterol, A.T. 70. Cap., 0.625 mg. Bot. 50s, 500s. Sol. in oil, 1.25 mg /cc. Bot. 15 cc.

Use: Orally, tetany therapy.

HYTROHA. (Pitman-Moore) Hyoscyamine HBr 0.18 mg., atropine 0.015 mg., scopolamine HBr 0.0048 mg., phenobarbital ¼ gr./Tab. or 5 cc. elixir. Elixir, Bot. 1 pt., 1 gal. Tab., Bot. 100s, 1000s.

Use: Orally, antispasmodic & sedative.

HYTONE LOTION. (Oermik) Hydrocarthans alcohol ½%, ¼%. Bot. 1 oz.

Use: Dermatologic.

HYTRAST. (E. Fougere) 50% w/v of combined iodine, 3,5 diiodopyridone-4—30.5% w/v. N-[2,3 propyldiol]-3,5-diiodopyridone

—44.60% w/v sodium carbamethylcellulose methyl, propyl parabens, monosodium phosphate, water/25 ml vial of 20 ml susp

Use For Bronchography

HYTROPHEN, (Chicago Pharmacal) Dehydrocholic acid 250 mg, homotropine methyl bromide 2.5 mg, phenobarbital 8 mg/tab Bot 100s, 1000s

Use (1 2 1 1 d p c) Hydracholeretic & antispasmodic w/sedative

HY-VAL-FLOBA WITH PHENOSABBITAL (Warren Teed) Phenobarbital 15 mg, hyoscyamus ext 15 mg, valerian ext 60 mg, passiflora ext 60 mg/Tab Bot 1000s

Use: Sedative

HYVANOL, (Walker) Vit A 4000 u, D 400 u, B-1 2 mg, B-2 2 mg, B-6 0.1 mg, C 37.5 mg, niacinamide 20 mg, cal pantothenate 1 mg, l cysteine HCl 8 mg, urea 75 mg, l tyrosine 60 mg, choline dihydrogen citrate 30 mg, glutamic acid 60 mg, liver conc 25 mg, yeast dried 50 mg/cap Bot 25s, 100s

Use: (1 2 1 1 d) Nutritional deficiencies

HYVA VAG. TABS, (Holland Ranlos) Methylrosaniline Cl, 1 mg, alkyldimethylbenzyl ammonium chloride, 0.5 mg, polyoxyethyl enenonylphenol 10 mg, polyethylene glycol tert dodecylthioether, 5 mg in a gel forming base/Tab Boxes 12s & 36s

Use Antimycotic

HY-VIDOX, (Vilaine) Pyridoxine HCl 10 & 25 mg/Tab Bot 100s 1000s

IBEBET, (Abbott) Ferrous sulfate 525 mg Vit B 12 25 mcg B 1 6 mg, B 2 6 mg B 6 5 mg, nicotinamide 30 mg, calcium pantothenate 10 mg, Vit C 150 mg/Tab Bot 60s

Use Iron deficiency anemias

IBEBIN CAPSULES, (Abbott) Ferric ammonium citrate 5 gr, thiamine HCl 0.375 mg, riboflavin 0.125 mg, nicotinamide 5 mg liver conc NF & liver fraction #2 NF 4 gr (0.25 Gm) (Containing the Vit B complex/Cop) Bot 100s, 500s

Use: (3 daily) Hematinic In anemias

IBEBIN FERROUS CAPSULES (Abbott) Ferrous sulfate, anhyd 2 gr (0.13 Gm) thiamine HCl 0.375 mg, riboflavin 0.125 mg nicotinamide 5 mg, liver conc, NF & liver fraction #2, NF 7 gr (0.455 Gm)/Cop Bot 100s, 250s, 1000s

Use: (1 daily) Hematinic

*Generic name recognized by NF, USP or A.M.A.

IBERIS AMARA. See: Anti M Lectin (Hyland)

IBEROL, (Abbott) Hematinic, differs from IBEROL-F in the omission of folic acid 1 mg, contains Vit B 12 12.5 mcg, ferrous sulfate 525 mg, liver fraction No 2 100 mg, Vit C 75 mg, B 1 3 mg, B 2 3 mg, nicotinamide 15 mg, B 6 1.5 mg, calcium pantothenate 3 mg

IBEROL-F FILMTABS (Abbott) Vit B 12 w/ intrinsic factor 1 oral u, iron 210 mg, liver fraction No 2 200 mg, folic acid 2 mg, Vit C 150 mg, B 1 6 mg, B 2 6 mg, nicotinamide 30 mg, Vit B-6 3 mg, cal pantothenate 6 mg/2 Filmtabs Bot 100s 500s, 1000s

Use: (2 daily) Anemias

IBEROL-F TABLETS, (Abbott) Ferrous sulfate 0.35 Gm, Vit B 12 (as Vit B 12 conc) 10 mcg, folic acid 12 mg, ascorbic acid 50 mg, stomach liver digest 0.5 Gm, thiamine mononitrate 2 mg, riboflavin 2 mg, nicotinamide 10 mg, pyridoxine HCl 1 mg, pantothenic acid (as Cal pantothenate) 2 mg/Tab Bot 100s, 500s 1000s

Use (3 daily) Anemias

ICHTHALBIN, (Knoll) Ichthyol albuminate is an organic sulfur comp Pow 1 oz Tab (5 gr) Bot 1000s

Use: Colarrhal conditions of the intestines

***ICHTHAMMOL**, NF NNR 1944 Oint, NF (Hirathal) Ichthyinate (Isarol) Oint NF (10%) Tube 1 oz, Jar 1 lb (Various Mfr) Oint (20%) Tube 1 oz (Merck Sharp & Dohme)

See Ichthymol Oint (Mollinckradt)

Ichthyol Oint (Schering)

Use: Mild antiseptic in skin disorders

W/Iodine Tr, boroglycerin, phenol hydrastine HCl See: Ioda Ichthol, Liq (Ulmer)

W/Iodine, phenol, hydrastine HCl, glycerite of boroglycerin eucalyptol See Ichthaldine, Liq (Merck Sharp & Dohme)

W/Naftalan, catamine petrolatum See Naftal, Paste (Rowell)

W/Phenol benzocaine, balsam peru, alum exsic, eads oil, eucalyptus oil See: Aiucaene, Oint (Jenkins Labs)

W/Phenol iodine zinc phenolsulfonate & boric acid Supp See: Vaginaids, Supp (Merck Sharp & Dohme)

ICHTHOLDINE, (Merck Sharp & Dohme) Ichthammol 4 gr, iodine 0.06 gr, phenol 4 gr, hydrastine HCl 2 mg, glycerite of baro-

glyceric 9 gr., eucalyptol 1/20 min./oz.
Bot. 1 pt., 1 gal.

Use: Mild antiseptic.

ICHTHYMALL. (Mallinckrodt) Sulfated oil obtained from distillation of bituminous shale. Bot. 1 oz., 1/4 lb., 1 lb., 5 lb.

Use: Mild antiseptic for skin diseases.

ICHTHYNATE, N.N.R. 1944. See Ichthammal (Mallinckrodt).

ICHTHYOL, N.N.R. 1944. (Scherling) Substances classified as Ichthammal. Conc., Bot. 4 oz. Oint. (10% & 20%), Tube 1 oz., Jar 1 lb.

Use: Mild antiseptic for skin diseases.

ICHTHYOL ALBUMINATED. See Ichthalbumin, Pow. & Tab. (Knall)

ICOPERSUL. (New Mexico Pharm.) Iron Cadmate 1 gr., Copper Sulfate 1/100 gr., Sodium Chloride 0.9%, Chlorobutanol 0.5%/2 cc. Vials 30 cc.

ICTOTEST. (Ames) p-Nitrobenzene diazonium p-toluene sulfonate. Reagent Tab., Bot. 90s.

Use: Test for urine bilirubin in the detection of hepatic disorders.

IDARON. (Marion) Polysaccharide complex 50 mg., thiamine mononitrate 5 mg., vit. B-12 5 mcg., B-6 2 mg./O.6 cc., Bot. 8 oz. Use: Anemia.

* **IDOXURIDINE.** (Smith, Kline & French) 2'-deoxy-5-iodouridine. See: Staxil.

* **IDURIDINE.** (Smith, Kline & French) 2'-Deoxy-5-iodauridine. Use: Antiviral agent.

IGEGAL CA 630. (Antara Chemicals) Octylphenoxy polyethoxyethanol, N.F.

Use: Ingredient in nitrofurazone solution.

IGOMOL. (Drug Products) Iodized oil 50 mg., gomenol 100 mg., guaiacol 100 mg., camphor 50 mg., menthol 50 mg., eucalyptol 50 mg., chlorobutanol 30 mg., olive oil/2 cc. Boxes 12s, 25s, 100s.

Use: (1 tsp. p.r.n.) Expectorant for coughs due to colds.

ILBAN. (Smith-Barney) Liver ext. 1:20, ferrous gluconate extic. 5 gr., Vit. B-1 2 mg., B-2 2 mg., B-6 0.2 mg., niacinamide 5 mg., Cal. pantothenate 0.2 mg./Cap. Bot. 100s, 1000s.

Use: (3 daily) Hematiocel.

ILETIN, N.N.D. 1961. (Insulin) (Lilly)

N.P.H. Vial (40 & 80 u./cc.) 10 cc.

Plafin Vial (0, 80 & 100 & 500 u./cc.) 10 cc.

Protamine zinc Vial (40 & 80 u./cc.) 10 cc.

* Generic name recognized by N.F., U.S.P. or A.M.A.

Zinc Insulin Vial (40 & 80 u./cc.) 10 cc.

Lente Vial (40 & 80 u./cc.) 10 cc.

Semi-Lente Vial (40 & 80 u./cc.) 10 cc.

Ultra-Lente Vial (40 & 80 u./cc.) 10 cc.

Use: Diabetes.

ILIDAR PHOSPHATE, N.N.D. 1962. (Bache)

Azopetine phosphate, 6-allyl-6,7-dihydro-5H-dibenz(c,e)azepine phosphate. Tab. 25 mg. Bot. 100s & 500s.

Use: (I.D.) Vascular diseases (Raynaud's Disease); quaternary vasodilator; vasospastic disorders.

ILOCALM. (Warren-Teed) d-calcium pantothenate 25 mg., methscopolamine nitrate 2.5 mg., mephobarbital 30 mg./Tab., Bot. 100s, 500s

Use: Anticholinergic sedative.

ILOMEL. (Warren-Teed) d-calcium pantothenate 25 mg., calcium carbonate 350 mg., milk solids/Tbsp. Pow. 9 1/2 oz.

Use: Peptic ulcer.

ILOPAN, N.N.D. 1962. (Warren-Teed) * Dex-pantonal. Pantothényl alcohol 250 mg./cc. Amp 2 cc. Box 12s, 25s, and Vial 10 cc.

Use: Intestinal atony and distention

ILOPAN-CHOLINE. (Warren-Teed) Ilopan 50 mg., choline bitartrate 25 mg./Tab. Bot. 100s, 500s.

Use: Hypodynamic states of gastrointestinal tract.

ILOSONE, N.N.D. 1962. (Eli Lilly) See: * Erythromycin Estolate.

ILOSONE, N.N.D. 1961. (Lilly) Erythromycin Estolate 250 mg./Cap. and 125 mg./Cap. Bot. 24s, 100s.

Use: Bacterial infections

250 mg. Bot. 24s & 100s.

W/Sulfadiazine, sulfamerazine, sulfamethoxazine. See: Ilosone Sulta, Tab. (Lilly)

ILOSONE. (Lilly) Erythromycin Estolate. (Erythromycin base) 125 mg/5 cc. Susp. Bot. 60 cc. Drops. (5 mg. of base/drop) Bot. 10 cc.

Use: Most common bacterial infections.

ILOSONE SULFA. (Lilly) Erythromycin Estolate 125 mg., sulfameth-dia-mer each 167 mg./5 ml. Bot. 60 cc.

Use: Bacterial infections.

ILOSONE SULFA. (Lilly) Erythromycin Estolate 125 mg., sulfadiazine 167 mg., sulfamerazine 167 mg., sulfamethazine 167 mg./Tab. Bot. 24s, 100s.

Use: Mixed or refractory bacterial infections

ILOTYCIN, N.N.D. 1962 (Erythromycin)

ILOTYICIN ETHYL CARBONATE DROPS AND PEDIATRIC 386

- (Lilly) Olet. (10 mg/Gm) Tube ½ oz, 1 oz Jar 1 lb Ophth Olet. (5 mg/Gm) Tube ½ oz Tab. EC (100 mg) Bot 36s, 100s (250 mg) Bot 24s 100s
 Use Antibiotic therapy against gram positive & gram negative organisms
- ILOTYICIN ETHYL CARBONATE DROPS AND PEDIATRIC, NND 1962** (Lilly)
 Ilotycin ethyl carb 1 Gm in dry flavored mixture when mixed 100 mg/cc Bot dropper 10 cc, 200 mg/cc Pediatric Bot. 60 cc
 Use Orally; acute bacterial infection
- ILOTYICIN GLUCOHEPTONATE DENTAL CONES.** (Lilly) Ilotycin G 5 mg/cone
 Use: Locally after tooth extraction and debridement
- ILOTYICIN GLUCOHEPTONATE I.V., NND 1961** (Erythromycin glucoheptonate) (Lilly)
 Amp 1 V (250 mg) 20 cc (500 mg) 30 cc (1 Gm) 50 cc
 Use 1 V; antibiotic therapy
- ILOTYICIN OTIC.** (Lilly) Ilotycin glucoheptonate 25 mg in 1 Bot One Bot diluent containing polymyxin B sulfate 50 000 u benzocaine 5%/5 cc propylene glycol
 Use Ear infections
- ILOTYICIN, PEDIATRIC, 200.** (Lilly) Ilotycin (Erythramycin) ethyl carbonate crystalline 24 Gm/60 cc Bot or 200 mg/5 cc Bot 60 cc (as dry pow) cherry cinnamon flavor
 Use Systemic infections
- ILOTYICIN-SULFA, PEDIATRIC.** (Lilly) Ilotycin (Erythromycin) ethyl carbonate 24 Gm sulfadiazine 2 Gm, sulfamerazine 2 Gm and sulfamethoxine 2 Gm/60 cc (dry pow) or 5 cc equal to Ilotycin EC 200 mg and sulfa 167 mg each
 Use: Infections requiring antibiotic and sulfa
- ILOTYICIN-SULFA TABLETS** (Lilly) Ilotycin crystalline 75 mg sulfadiazine sulfamerazine & sulfamethoxine 111 mg of each/Tab Bot 36s, 100s, 500s
 Use: Antibiotic w/triple sulfa
- I.L.X. W/Vit B 12 & Folic Acid** (Kenwood)
 Vit B 12 10 mcg, folic acid 0.85 mg ferrous gluconate 5 gr liver fraction No 2 2 gr, thiamine HCl 2 mg, riboflavin 2 mg nicotinamide 20 mg/Tab Bot 100s
 Use: (11 d) Hematinic
- I.L.X. ELIXIR.** (Kenwood) Ferric & ammonium citrate 27 gr, liver fraction #145 gr Vit B-12 30 mcg, B 1 6 mg B 2 6 mg niacin 30 mg/1½ oz Bot 12 oz
 Use: Anemias
- IMADYL UNCTION.** (Roche) Histamine diHCl 1%, acetyl glycolsalicylic ester 5% methyl salicylate 15%, menthal (not USP) 10% & thymal 1% Tube 1½ oz, Jar 1 lb
 Use: Local vasodilation & analgesic
- IMBICOLL**
 W/Cascara (Upjohn) A stercula gum w/cascara sagrada 15 min/dr. Bot 4 oz, 8 oz., 16 oz
 Use: Bulk producing laxative
- W/Vit B 1** (Upjohn) Stercula gum w/Vit B 1 1 mg./dr Bot 4 oz, 8 oz, 1 pt
 Use: Bulk producing laxative
- IMIFERON, NND 1960** (Lakeside Labs)
 An iron dextran complex (50 mg iron/cc) Amp 2 cc, Box 10s, 5 cc Box 4s Vial 10 cc
 Use 1M Anemias
- *IMINOPHENIMIDE**, (Lederle Lab) 3 Ethyl 3 phenyl 2,6 piperazinedione
 Use Hypnotic agent
- *IMIPRAMINE HCl, NND 1962** 5-(3 Di methylaminopropyl) 10,11-dihydro 5H dibenz (b, f) azepine HCl See Tofranil (Geigy)
- IMMOLIN** (Julius Schmid) Tube 75 Gm
 Use: Contraceptive
- IMMU-G** (Parke, Davis)
 Use Immune Globulin
- IMMUNE SERUM GLOBULIN, U.S.P., NND 1962** Gamma globulin fraction of normal human plasma
 Cutter—Vial (160 mg/cc) 2 cc, 10 cc
 Lederle—Vial (16%) 2 cc, 10 cc
 Pitman Moore—Vial (160 mg/cc) 2 cc, 10 cc
 Merck Sharp & Dohme—Vial (160 mg/cc) 2 cc, 10 cc
 Use (15 mg/lb of body weight) Modification of active measles
- IMMUNE SERUM GLOBULIN-HUMAN, U.S.P., NND 1962** See Globulin Immune Serum, USP
- IMMUNE SERUMS**
 See: Anth Hemophilus Influenzae Type B Serum (Rabbit)
 Human Measles Immune Serum
 Human Scarlet Fever Immune Serum
 Immune Serum Globulin (Human)
- IMMUNE SERUM (ANIMAL).**
 See Antipertussis Serum (Wyeth)
 Anthrax Serum (Lederle)
 Botulism Antitoxin (Lederle)
 Diphtheria Antitoxin USP
 Gas Gangrene Antitoxin NF

*Generic name recognized by NF USP or A.M.A.

- Tetanus Antitoxin U.S.P.
Tetanus Antitoxin—Gas Gangrene Combined N.F.
- IMMUNE SERUM (NUMAN).**
See: Antimumps Serum (Hyland)
Antipertussis Serum
Hypertussis (Cutter)
Poliomyelitis Immune Globulin (Various Mfg.)
- IMMUNOVAC ORAL VACCINE TABLETS.**
(Parke, Davis) Antigenic product combining the immunizing properties of whole bacterial vaccines & soluble ecto-antigens known as immunogens. Bot. 20s, 100s
Use: Prophylaxis of certain bacterial infections.
- IMMUNOVAC RESPIRATORY VACCINE.**
(Parke, Davis) A distinctive antigenic product combining the immunizing properties of whole bacterial vaccines & the soluble ecto-antigens known as immunogens. Vial 10 cc.
Use: Infections of the upper respiratory tract.
- IMPOTEX SOLUTION.** (Whittaker) Chorionic gonadotropin, thiamine HCl, L-glutamic acid, chlorbutanol, procaine HCl. Two 5 cc. vials (powd. & diluent)
Use: Impotence in the male.
- IMPREGON LIO.** (Fleming) Tetrachlorosalicylonilide, 2%. Bot. 4 oz.
Use: Diaper soak.
- IMPREGON OINT.** (Fleming). Tetrachlorosalicylonilide, 5 mg; neomycin base, 2.5 mg./Gm. Tube 20 Gm.
Use: Diaper rash.
- IMPROVED POTENSORS PORTE.** See: Potensors Forte, Improved, Tab. (Tailby-Nason)
- INACTIVATED DIAGNOSTIC DIPHTHERIA TOXIN, U.S.P.** See Diphtheria Toxin, Inactivated Diagnostic.
- INCNOLIP.** (Flint, Eaton) Choline 20%, inositol 1 Gm./15 cc. Bot. 1 pt., 1 gal.
Use: Nutritional deficiencies.
- INCORPOHIST.** (Blue Line) Methoppyrilene HCl 10 mg., pyrilamine maleate 10 mg., prophepyridamine maleate 5 mg./Tab. Bot. 100s, 1000s
Use: Triple antihistaminic
- INCORPOSUL, N.N.R. 1953.** (Acet-Dia-Mer-Sulfonamides) (Blue Line) 0.167 Gm. each of sulfacetamide, sulfadiazine & sulfamerazine/Tsp. or Tab. Susp., Bot. 1 pt., Tab., Bot. 100s.
Use: Orally, triple sulfa therapy
* Generic name recognized by N.F., U.S.P. or A.M.A.
- INCREMIN.** (Lederle) L-Lysine 300 mg., Vit. B-12 25 mcg., B-1 10 mg., B-6 5 mg., alcohol & cherry flavor/cc. or Tab. Bot. 15 cc. Plastic Squeeze or Tab. Bot. 30s.
Use: (10-20 drops daily) Appetite stimulant
- INCREMIN W/IRON.** (Lederle) L-Lysine HCl 300 mg., ferric pyrophosphate 280 mg., Vit. B-1 10 mg., B-6 5 mg., B-12 25 mcg./5 ml. Syr. Bot. 4 oz., 1 pt.
Use: An appetite stimulant.
- INCRESOL W/Vitamin B-1.** (G. W. Carrick) A sol. of gentian, cardamom & foraxocum, d.6 mg. of thiamine HCl/fl. oz. Bot. 1 pt., 1 gal.
Use: Vitamin B-1 deficiencies.
- INDALONE.** See: Butopyronoxyl (Various Mfr.)
- INDIAN GUM.** See: Karaya Gum
- INDIGO CARMINE.** Indigotin disulfonate Sod.
Assn. Physicians Lab.—Amp. (0.8% aq. sol.) 3 cc., 5 cc., Box 12s, 25s.
Nynson—Amp. (0.8% aq. sol.) 5 cc., Box 10s, 100s.
Rorer—Amp. (0.8% aq. sol.) 5 cc., Box 6s, 12s, 25s, 100s.
Use: I.V., I.M. diagnostic aid.
- INDIOTINDISULFONATE SODIUM, U.S.P.** Inj., U.S.P. See: Indigo Carmine, Amp. (Various Mfr.)
- INDOKLON.** (Ohio Chem. and Surg Eq) Hexafluorodethyl ether. J.A.M.A. 188, 1555, 1958 flurathyl.
Use: Shock inducing agent.
- *INDDMETHACIN.** (Merck Sharp & Dohme) 1-(p-chlorobenzoyl)-5-methoxy-2-methylindole-3-acetic acid.
Use: Anti-inflammatory agent.
- INDON, N.N.D. 1961.** (Parke, Davis) (Phen-Indione) Tab. (50 mg) Bot. 100s, 1000s.
Use: Anticoagulant.
- *INDOPINE HCl.** (Irwin, Neiser) 4-[3-indolylethyl]-1-phenethylpiperidine HCl.
Use: Analgesic and/or tranquilizer.
- INESTRA.** (Ethiny Estradiol) (Merck Sharp & Dohme) Tab. (0.02 mg. & 0.05 mg) Bot. 100s, 1000s.
Use: Oral estrogen therapy.
- INFACAPS.** (Lanett) Vitamin A 3000 u., D 800 u./cap. Bot. 1000s, 5000s.
- INPACE.** (Ives-Cameron) 100 mg. of ascorbic acid/cc. Bot. 10 cc., 60 cc.
Use: Vitamin C therapy.
- INFA-CONCEMIN WITH B-12 ACTIVITY.** (Marrell) Thiamine HCl 0.8 mg., riboflavin

- 0.8 mg, niacinamide 5 mg, pyridoxine HCl 0.5 mg, ferrous sulfate (½ gr) 37.5 mg, Vit B12 activity 2 mcg w/slice bran ext/cc Bot w/dropper 30 cc, Bot 2 oz
 Use: Nutritional & iron deficiency anemias
- INFADROPS LIQ.** (Sig. Labs) Vit A 5000 u, D 1000 u, B1 1 mg, B2 1 mg, B6 1 mg, C 50 mg, nicotinamide 10 mg, panthenol 3 mg/D6 cc Bot 30 cc
 Use: Vitamin deficiencies in infants
- INFAGEN, NNR 1955** (Pitman Moore) Diphtheria A tetanus toxoids & pertussis vaccine, combined, alum precip Vial 7½ cc
 Use: Immunization (3 in) of 0.5 cc)
- INFANTOL, PLAIN.** (First Texas) Pectin 4 gr, zinc phenolsulphonate 1 3/5 gr, phenyl salicylate 2 1/5 gr, bismuth subsalicylate 8 gr Irish moss ext q s/n oz Bot 1 pt, 1 gal
 Use: Diarrhea
- W/Paregoric (First Texas) Pectin 4 gr, paregoric 60 min, zinc phenolsulphonate 1 3/5 gr phenol salicylate 2 1/5 gr, bismuth subsalicylate 8 gr, Irish moss ext q s/n oz Bot 1 pt, 1 gal
 Use: Diarrhea
- INFANTOVIT DROPS** (Consol Midland) Vit A 6000 u, D 1200 u, B1 1.8 mg, B2 0.6 mg, C 60 mg, niacinamide 3 mg, Vit B6 0.3 mg, Cal pantothenate 12 mg/O.6 cc Bot 15 30 2 60 cc
 Use: Dietary supplement
- INFANTOVIT PLUS SPHERE.** (Consol Midland) Vit A 5000 u, D 1000 u, C 60 mg, B1 1 mg, B2 1.5 mg, B6 1.0 mg, Cal pantothenate 3 mg, nicotinamide 10 mg, B-12 1 mg, folic acid 0.25 mg/sphere Bot 30s 100s, 250s 1000s
 Use: Dietary supplement
- INFANTUSSIS** (Blue Line) Pot citrate, squill fl ext, ipecac syr, talu syr, honey & alcohol Bot 1 pt 1 gal
 Use (1 2 tsp p r n) Cough
- INFAZYME** (Merrill) Protein hydrolysate 5 Gm thiamine HCl 3 mg, riboflavin 2 mg, niacinamide 12 mg, pyridoxine HCl 1 mg, opatinized iron, N.F., 135 mg, rice bran ext supplied additional B complex factors/15 cc Bot 12 oz
 Use: Multiple vitamin therapy
- INFERTAB-P** (Kenwood) Thyroid ½ gr, hamotropine methyl bromide 25 mg, supplemented with vitamins/Tab Bot 60s
 Use: Treatment of female infertility
- INFERTAB-M.** (Kenwood) Thyroid ½ gr supplemented with vitamins/Tab Bot 60s
 Use: Treatment of male infertility
- INFILTRASE.** (Armour) Hyaluronidase Vial (150 turbidity reducing u) 1 cc; (1500 turbidity reducing u) 10 cc.
 Uses: 3 C, 1 M; rapid dispersal of fluids into tissues
- INFLOLAC BACTERIAL VACCINE.** (U.S. Standard) Hemophilus influenzae 1000 million, Streptococcus hemolyticus 500 million, Str viridans 500 million, Micrococcus catarrhalis 200 million, Pneumococcus Types I II & III 3000 million in casein menstruum/cc Vial 10 cc, 30 cc.
 Use: Treatment of acute & chronic rhinitis sinusitis mastoiditis & influenza
- INFLUENZA COMBINED VACCINE.** (U.S. Standard) Bacillus influenzae 1000 million, Streptococcus hemolyticus 500 million, Str viridans 500 million, Micrococcus catarrhalis 200 million, Pneumococcus Types I, II & III 3000 million/cc Vial 20 cc
 Use: Treatment of acute & chronic rhinitis sinusitis, mastoiditis & influenza
- INFLUENZA TYPE B SERUM (RABBIT)**
ANTI-HEMOPHILUS, NND 1959 See: Anti Hemophilus influenzae Type B Serum (Babbit)
- INFLUENZA MIXED VACCINE.**
 Lilly—Vial 5 cc, 20 cc.
 Pitman Moore—Vial 20 cc.
 U.S. Standard Prods.—Vial 20 cc.
 Use: Treatment of rhinitis, sinusitis & influenza
- INFLUENZA-PNEUMONIA VACCINE.** (Parke Davis) Pneumococcus Types I, II, III, V, VII, VIII & XIV 15,000 million, Streptococcus hemo & non hemo 2500 million, Hemophilus influenzae 500 million, Staphylococcus aureus 500 million/cc Vial 20 cc
 Use: Treatment of infections produced by the above organisms
- INFLUENZA-PNEUMONIA VACCINE.** (Pitman Moore) Pneumococcus Types I, II, III, IV, VII, VIII & XIV 3000 million Streptococci hemo & non hemo & viridans 1000 million, Hemophilus influenzae 500 million X pneumoniae 500 million, Staphylococcus aureus 500 million, M catarrhalis 500 million/cc Vial 20 cc
 Use: Stimulation of active immunity
- INFLUENZA SERUM.** (Squibb) A sterile, refined & conc antiserum obtained by immunizing rabbits w/Hemophilus influenzae Type B potency Vial 25 mg, agglutinin antibody nitrogen, equiv to not less than 25,000 provisional units
 Use: Influenzal meningitis due to Hemophilus influenzae Type B

- ***INFLUENZA VIRUS VACCINE MONOVALENT-TYPE A-ASIAN STRAIN, N.N.D. 1962.** (Various Mfr.) The sterile suspension of formaldehyde-killed influenza virus, type A, Asian strain grown in the extra-embryonic fluid of chick eggs. Vials 10 cc.
Use: Immunization (1 cc. I.M.)
- ***INFLUENZA VIRUS VACCINE, POLYVALENT, N.N.D. 1962.** (Types A & B)
Iederle—Vial 10 cc.
Lilly—Vial 1 cc.
National Drug—Vial T cc., 5 cc.
Pitman-Moore—Vial 1 cc., 5 cc.
Merck Sharp & Dohme—Vial T cc., 10 cc.
Use: Immunization.
- INFOSE.** (Sorden) Apple juice conc. Eq. Bot. T pt.
Use: Infant food.
- INFRON PEDIATRIC.** (Whittier) Vit. D 100,000/u. Cap. Cap., Bot. 6s.
Use: Orally; vitamin D therapy
- INFUSION CONCENTRATE HYDROCORTONE.** See: Hydrocortone, Infusion Concentrate, Amp. (Merck Sharp & Dohme)
- INNISTON.** (Union Pharm.) 10 mg 1-phenyl-1-(2-pyridyl)-3-dimethylamino-propane. (Propylphenylamine/Tab. Bot. 36s.
Use: Allergy & colds.
- INN TABLETS, N.N.D. 1962.** (Lilly) (Isoniazid) Isonicotinyl hydrazide, Isonicotinic acid hydrazide. Tab. (100 mg) Bot. 100s, 1000s.
Use: (3-5 mg./Kg. of body weight) Tuberculosis.
- INJECTION DIRECT SKY BLUE 4%.** (Wyeth) Amp. 5 cc.
Use: Vital staining of lymphatic tissue.
- INOEX-CAL.** (Tilden) Calcium carbonate 200 mg., ferrous sulf. 5 mg., nicotia 3 mg., Vit. B-1 0.3 mg., B-2 0.3 mg., B-6 2 mg., C 25 mg., D 133 u./Tab. Bot. 100s.
Use: Dietary supplement.
- INOEX-NC.** (Tilden) Hesperidin 200 mg., ascorbic acid 200 mg./Cap. Bot. 50s, 100s
Use: Prenatal supplement for maintenance of capillary integrity.
- INOPHYLLINE.** See: Aminophylline (Various Mfr.).
- INOSIT.** See: Inositol, N.F.
- ***INOSITOL, N.F., A.D.R. Tab., N.F. X (Bios T, Hexahydroxycyclohexane, Inosil, Dambase)**
Use: Alterations in lipid metabolism.
W/Betain-choline, dehydrathalic acid, ox bile, liver-B. See: Hepozyme, Tab. (Endocrine).
- W/Choline dihydrogen citrate, polysorbate 80. See: Manichol, liq. (Ives-Cameron)
- * Generic name recognized by N.F., U.S.P. or A.M.A.
- W/Choline gluconate. See: Sirnositol, liq. (Commercial Solvents).
- W/Choline. See: Wychol, liq. (Wyeth).
- W/Choline & pyridoxine HCl. See: Vasculat, liq. (Schenley).
- W/Choline bitartrate & dl-methionine. See: Chalathol, Cap. (Nysco)
Tipocaps, Cap. (Lakeside)
- W/Rutic, methyltestosterone, conjugated estrogen (equine), methionine, vitamins and minerals. See: Via-Geric-N, Tab. (Rowell)
- W/Vitamin, Multiple. See: Vitamin preparations.
- W/Vit. B-12 & trichovine citrate. See: Lipoliquid, liq. (Lakeside).
- ***INOSITOL NEXANITRATE.** See: Talanate, Tab. (Reed & Carnrick)
- W/Phenobarbital. See: Talanate W/Phenobarbital, Tab. (Reed & Carnrick).
- INOSITOL NEXAPHOSPHATE** (phylic acid)
W/Calcium & iron, salts. See: Calciphos, Tab. (Knoll)
- INOSITOL NEXAPHOSPHORIC ACID CAL & MAG. SALT.** See: Phytin (Ciba)
- ***INOSITOL NIACINATE.** Meso-Inositol hexanicathinate. See: Inoadil (Winthrop Labs)
- INOSOL B W/INVERT SUGAR 10%.** (Abbott) Bot. 250 ml., 500 ml., 1000 ml.
- INOTON.** (Fougere) Ammonium sulphalchthylolate, Hamamelis ext., titanium comp., zinc oxide, Sod. borate. Oint., Tube 50 Gm., Jar 500 Gm. Pow. (ammonium sulphalchthylolate, Hamamelis ext., Sod. borate, zinc oxide, Col. carb & Mag. silicate) Bot. 4 oz.
Use: Topically; antiseptic.
- INPERSOL SOL. & EQUIP.** (Abbott) Sodium lactate, anhydrous 500 mg., sodium chloride U.S.P. 560 mg., calcium chloride, hexahydrate 15 mg., dextrose U.S.P. 1.5 Gm. or 7 Gm./100 ml., 1000 ml. Abbaliter bottles, packed in 6s, and disposable administration set, packed in 8s.
Use: Peritoneal Dialytic.
- INSOBEL TD.** (Denab) Butobarbital sod. 30 mg., ext. of belladonna 30 mg./T.D. Tab. Bot. 100s
Use: Sedative and antispasmodic.
- INSOLAT.** (Denab) Butobarbital sod. 15 mg., or 30 mg./Tab. Bot. 1000s.
Use: Sedative.
- INSOMNITAL.** (Bumington) Barbital 1 gr., hyoscyamus ext. 1/10 gr., chamomile T/10 gr., passiflora 1/10 gr./Tab. Bot. 100s, 1000s.
Use: (1-2 h.s.) Mild sedation.
- INSTA-PEP.** (Drug Research) A vitamin-mi-

- eral formula w/o 'self starter' (Dynamol)/
Tab Bat 25s, 50s
- INSTA-PERM.** (Paul B Elder) Potassium permanganate 4.86 gr In efferv base/Tab Bat 100s, Vial 16s
Use: For Dermatoses
- ***INSULIN, USP, NND 1962**
Burroughs Wellcome—See: Globin Insulin Lilly—See: Iletin & Lente Insulin
Merck Sharp & Dahme—NPH, Vial (40 u/cc & 80 u/cc) 10 cc Plain, Vial (40 u/cc, 80 u/cc & 100 u/cc) 10 cc Protamine Zinc Vial (40 u/cc & 80 u/cc) 10 cc
Squibb—See also: Globin Insulin NPN, Vial (40 u/cc & 80 u/cc) 10 cc Plain, Vial (40 u/cc, 80 u/cc & 100 u/cc) 10 cc Protamine Zinc, Vial (40 u/cc & 80 u/cc) 10 cc Zinc Insulin, Vial (40 u/cc & 80 u/cc) 10 cc
- ***INSULIN, GLOBIN ZINC, SUSP, USP**
NND 1962 Globin insulin w/zinc globin zinc insulin See: Lente Iletin Susp (Lilly)
- ***INSULIN INJECTION, USP** Insulin Insulin HCl
- INSULIN INJECTION, ISOPHANE, USP,**
NND 1962 Isophane insulin See: NPN insulin (Lilly)
- ***INSULIN, ISOPHANE SUSP, USP** NND 1962
See NPN Iletin Susp (Lilly)
Protamine Zinc and Iletin Susp (Lilly)
- INSULIN, PROTAMINE ZINC, INJECTION, USP**
- ***INSULIN, ZINC, SUSP, USP** See Lente Insulin NND 1962 Vial (Lilly)
- ***INTERFERON** A protein formed by the interaction of animal cells with viruses capable of conferring on animal cells resistance to virus infection
- INTOCOSTRIN, NND 1962** (Squibb) Chondrodendron tomentosum ext 1 cc Is equiv to 20 u Physiologically equiv to 3 mg tubocurarine chloride pentahydrate Vial (20 u/cc) 10 cc
Use: IV; anticonvulsant
- INTRACANE HYDROCHLORIDE, NND 1959** (Squibb) Parthoxycoaine, Diethoxyn β Diethylaminoethyl p ethoxybenzoate NCl Amp (sterile crys) 1 Gm Bat (monodried crys) 1 oz, 1 lb In oil (2% & 5%) Bat 30 cc Oint in plastibase (2% & 5%) Tube 1 oz Jar 1 lb
Use: Following types of anesthesia regional spinal topical dental or endotracheal
W/Polymyxin B sulfate hydrocortisone neomycin sulfate alcohol glycerine propylene glycol See Intramycin Otic (Philadelphia Labs)
- *Generic name recognized by NP USP or AMA
- INTRADERM TYROTRICIN.** (Wallace Labs)
Tyrothricin 0.1% Oint, Tube 1 oz
Use: Pyogenic skin infections
- INTRADERM TYROTRICIN SOLUTION.**
(Wallace) Tyrothricin 1 mg/cc. Bat. 120 cc
Use: Antibiotic penetrating skin preparation
- INTRANEPTOL.** (Lederle) Whole liver 33.3 Gm/cc Vial 10 cc
Use: IV,—dilute before using; chronic infectious hepatitis (virus hepatitis) & cirrhosis of liver
- INTRAMYCIN OTIC.** (Philadelphia Labs)
Polymyxin B sulfate 0.65 mg, (2000 u) hydrocortisone 100 mg, tetracaine HCl 5.0 mg neomycin sulfate 5.0 mg, alcohol 3%, glycerine 50%, propylene glycol 50% base/cc Drop Bat 7.5 cc
Use: Infection otitis externa
- INTRAVAL SODIUM.** See: Pentothal Sodium (Abbott)
- INTRAVITE.** (Myers Carter) Vit B 1 50 mg, B 2 5 mg, B 6 5 mg, C 50 mg, niacin amide 75 mg, pantothenol 6 mg/cc. Vial 10 cc
- INTRAVITE C.** (Philadelphia Ampule Labs)
Vit B 1 50 mg B 2 5 mg, niacinamide 75 mg, B 6 5 mg, cal pantothenate 6 mg ascorbic acid 50 mg/cc Vial 10 cc
Use: Vitamin B complex w/vit C
- INTRIBEX KAPSEALS.** (Parke, Davis) Vit B 12 7.5 mg w/intrinsic factor $\frac{1}{2}$ u, liver stomach conc 200 mg, folic acid 1 mg, ferrous sulfate 375 mg, Vit C 75 mg/Kapsel Bat 100s, 500s
Use: Anemia
- INTRINASE, NND 1962** (Upjohn) Vit B 12 w/intrinsic factor $\frac{1}{2}$ oral u/Tab Bat 30s
Use: Anemias
- INTRINSIC FACTOR, USP**
See: Stomach Products
Vit B 12 W/Intrinsic Factor
- INTRON CT** (Deseret) Intrinsic factor conc w/B 12, $\frac{1}{2}$ USP anti anemia oral unit, ascorbic acid 75 mg, liver stomach conc 200 mg, ferrous sulf exsic 300 mg, folic acid 1 mg, B 12 (cobalamin conc NF) 75 mcg/Tab Bat 100s
Use: Hematonic
- INTROMYCIN** (Pitman Moore) Streptomycin sulfate 15 mg, neomycin sulfate 7.5 mg in carob pow/Gm Pow, Jar 75 Gm
Use: Treatment of diarrhea
- INULIN** (US Standard Prods) (Dahlin) A starchlike carbohydrate Amp 10 cc, Box 100s 50 cc, Box 10s, 25s, 50s, 100s (Warner Chilcott) Sol 10% Bat 50 cc
Use: Diagnostic aid, kidney function, glomerular filtration rate
- INVERSINE, NND 1962** (Merck Sharp &

Dohme) Mecamylamine HCl, 3-Methylamino-isocamphane HCl. Tab. 2.5 mg., 10 mg., Bot. 100s.

Use: Hypertension.

INVERT SUGAR, N.N.R. 1955.

See: Emetrol, liq. (Kinney)

Travert, Sol. (Don Boxter)

Verclystyl Inlusloe, Sol. (Abbott)

INVERT SUGAR, N.N.R. 1955. (Abbott) An equal mixture of dextrose & levulose. Rot. (5% in saline or water) 1000 cc., (10% in water or saline) 1000 cc.

Use: Carbohydrate therapy.

See: Travert, N.N.R. (Don Boxter)

INVERT SUGAR, N.N.R. 1955. (Lilly) Mixture of dextrose & levulose 7.5 Gm., sucrose 0.15 Gm., Sod. tartrate 0.023 Gm./10 cc. Amp. 10 cc., Box 6s, 25s, 100s.

Use: Varicose veins.

IOCAPRAL (Wiethrop Labs.) Theobromine 5 gr., mebaral ½ gr. & col. iodide 2 gr., di- and triethaalamines. Tab., Bot. 100s, 500s

Use: (1-2 t.i.d.) Hypertensioe.

IOCLIDE, (Clyp-Adams) Polyethoxypropoxy-polyethoxyethanol-Iodine complex 15.8%, eonylphenoxypropylhexyethaol-Iodine complex 15.2%, hydrogee chloride 0.20%, inert ingredients 68.8%. Bot. 26 ml., 1 pt., 1 qt.

Use: Germicide, disinfectant.

IOCORT & IOCORTAR CREAM. (Greco)

Iocort. ¼% cream contains microflee hydrocortisone ¼%, Iodochlorhydroxyquie 3%

½% cream contains ½% hydrocortisone

and 3% Iodochlorhydroxyquin. Iocortar: ¼

& ½% creams contain some with 3% coal

tar sol. added. Tube 1 oz.; jar 4 oz.

Use: Dermatitis.

IOCORTAR 1% OINTMENT. (Greco Phar-

macol) Microphine hydrocortisone 1%,

Iodochlorhydroxyquie 3%, solution of coal

tar 3% petrolatum base./Oint. Tube

1 oz.

Use: Dermatologic

IOCYLATE, (Pitman-Moore) Ethylenediamine

dihl ¼ gr., Sod. salicylate 7½ gr./Tab.

Bot. 100s, 500s, 1000s.

Use: (1-2 q.i.d.) Analgesic & antipyretic.

IOD-ETHAMINE, (Pitman-Moore) Ethylenedi-

amine dihl. Elixir (16 gr./fl. dr.) Rot. 1 pt.,

1 gal. Tab. (2 gr.) Bot. 100s, 500s, 1000s (5

gr.) Bot. 100s, 500s, 1000s.

Use: Iodine therapy.

IODEX. (Menley & James) Iodine in organic

combination. Oint., Tube 1 oz., Jar 4 oz.,

1 lb. Pessaries (vaginol) for vaginitis, 6s.

Supp. (rectal), 12s. Soap for skin care.

Use: Cell stimulant.

W/Methyl salicylate, (Meeley & James) lo-

* Generic name recognized by N.F., U.S.P. or A.M.A.

dex w/5% methyl salicylate (syn.) Tube

1 oz., Jar 4 oz., 1 lb.

Use: Iodex plus plus.

IODINATED CASTOR OIL. See: Kiodlee,

Cap. (Gallia).

IODINATED PEPTONE W/Sod. Iodide. See

Iodoloke-S, Amp. & Vial (Lakeside).

IODINATED PROTEIN W/Sod. Iodide. See:

Protide, Vial. (Paul Money)

W/Colloidal proteie. See: Criodie, Vial.

(Crookes).

***IODINE, U.S.P., A.D.R.; Amp., N.F.; Oint.,**

N.F.; Sol., N.F.; Sol. phenalated, N.F.; Tr.

stroeg, N.F.; Sol. stroeg, U.S.P.; Tr., U.S.P.

W/Colloidal proteie. See: Criodin, Vial

(Crookes)

IODINE & RORIC ACID POWDER. (Endo)

Mild (0.44% iodine) 1 oz., Bot. Strong

(0.88% iodine) 1 oz., Bot.

Use: Antiseptic & deodorant.

IODINE CACODYLATE, COLLOIDAL. See:

Cacodye, Amp. (Res. Medications Inc.).

IODINE COMBINATION. See: Calcidree

Prep. (Abbott).

IODINE DIGLYCOCOLL NI, N.N.D. 1959.

See: Bursoline, Tab. (Eurehom Soluble Iodine).

IODINE, GLYCERINATED TINCTURE. See:

Thalogen, liq. (United Lab.).

IODINE-IODOPHOR.

See: Iodine (Iodine Co.)

Betadise (Carbie & Conwell)

Wescodyne (West Disinfecting Co.)

IODINE, ORGANIC PROTEIN BOUND. See:

Amand's Solution (Lesming).

IODINE POVIDONE, N.N.D. 1959.

IODINE PRODUCTS, ANTI-INFECTIVE.

See: Anayodine (Eischoff).

Betadise (Carbin & Coewell)

Bursoline (Burnham Soluble Iodine).

Chiofian (Vorlaus Mfr.).

Diglycocol Hl.

Diadoquin (Searle).

Diado-Hydroxyquinolise (Various Mfr.).

Globalise (Wallace-Tieran)

Heliogen, Tab. (Penick)

Iodise (Iodine Pharm.)

Quinoxyl (Burrroughs Wellcome).

Viorm (Ciba).

Wescodyne (West Disinfecting Co.)

IODINE PRODUCTS, DIAGNOSTIC.

See-

Iosaluble

Chloriodized Oil.

Ethyl Iodaphenylundecylate (Lafayette).

Iodeikon (Mollinckrodt).

Iodized Oil (Fougera).

Iodoaliphoeic Acid (Various Mfr.).

Iodobrassid (Various Mfr.).

Iodochloral (Searle)
 Iodopanoic Acid (Winthrop Labs)
 Lipiodol (Fougera)
 Lipiodine (Ciba)
 Pantopaque (Lafayette)
 Priodax (Schering)
 Skiodon (Winthrop Labs)
 Telepaque (Winthrop Labs)
 Umbradol (Winthrop Labs)
 Veripaque (Winthrop Labs)
 Soluble
 Diadrast (Winthrop Labs)
 Hippuran (Mallinckrodt)
 Iodohippurate Sodium (Mallinckrodt)
 Iodopyrocet (Winthrop Labs)
 Iodophthalein Sodium (Various Mfr)
 Methiodal Sodium (Winthrop Labs)
 Neo Iopax (Schering)
 Sodium Acelrizoate (Mallinckrodt)
 Sodium Iodomethamate (Schering)
 Urokon Sodium (Mallinckrodt)

IODINE PRODUCTS, NUTRITIONAL
 See: Amend's Solution (Leeming)
 Calcidin (Abbott)
 Calcium Iodobehenate (Various Mfr)
 Criodin, Vial (Crookes)
 Di Iodo Tyrosine (Organon)
 Entodon (Winthrop Labs)
 Glyco HI, Cap (Burnham)
 Hyadin, liq (Coardner)
 Hydriodic Acid Diluted (Various Mfr)
 Iod Ethimine (Pitman Moore)
 Iodin Ration (Paul Money)
 Iodobrassid (Various Mfr)
 Linodin (Central)
 Lipodine (R E Dwight)
 Lipiodine, Tab (Ciba)
 Lugol, Cap (Various Mfr)
 Organidin (Wampole)
 Orgaphen (Wampole)
 Oridine (Lilly)
 Potassium Iodide (Various Mfr)
 Riodine (Gallia)
 Sajodin (Winthrop Labs)
 Stearadine (Parke Davis)
 Strong Iodine Solution (Various Mfr)
 Syniodin, Tab (Blue Line)
 Thyrolagen (United Lab)

IODINE RATION (Paul Money) Iodine D I mg / Tab Bot 1000s
 Use: Dietary source of Iodine

IODINE SOLUBLE. See: Burnham Soluble Iodine (Burnham Soluble Iodine)

***IODIPAMIDE, N F**
***IODIPAMIDE METHYLGLUCAMINE, U S P, N N D** 1962 N N Odipyl bis(3 amine 2 4 6 Iriodobenzoic acid) Also sodium salt inj See: Cholografin Methylglucamine Amp (Squibb)

W/Diatrizoate methylglucamine See: Sino grafin liq (Squibb)
***IODIPAMIDE SODIUM INJ, N F, N N D** 1962 See: Chlorografin Sod (Squibb)
IODIZED AMMONIUM OLEATE. See: Iodized Petrogen, liq (Wyeth)
IODIZED CALCIUM CREOSOTATE W/Terpin hydrate Hyascyamus ext, strychnine sulfate & phosphorus See: Cresradine, Tab. & liq (Raymer)
***IODIZED OIL, U S P, N N R** 1950 (Fougera) A vegetable oil containing not less than 38% and not more than 42% of organically combined iodine
IODIZED PETROGLN. (Wyeth) Iodized ammonium oleate emulsion Total Iodine 5%. Also available as 10%. Bot, 3 fl oz. Use: Iodine therapy
IODIZED POPPY-SEED OIL, N N R 1950. See: Lipiodol Amp. (Fougera)
***IODOLPHIONIC ACID, N F, N N D** 1962; Tab, N F Alpha phenyl-alpha(4 hydroxy 3 5 diiodophenyl) propionic acid Bihselecton, dikol, pheniodol See: Perfectochol, Tab (Lafayette) Priodax, Tab., N N R (Schering)
O-IODOBENZOIC ACID. W/Triethanolamine dibucorne base in a neutral hydrophilic base See: Pro Nasyi (Progonosyl Co)
iodo BISMUTHATE OF QUININE. (Fougera) Quinine bismuth iodide containing approximately 19% bismuth Amp (4 cc & 3 cc) Box 6s, 25s
 Use: Antisyphilitic & bismuth therapy
IODOBRESSID, N N R 1953 (Ethyl Diiodobrossidole)
 See: Lipodine, Tab (Dwight) Lipiodine Tab (Ciba).
iodo BROMIDE AND CALCIUM COMPOUND. (Tilden) Sodium Bromide 2 1/2 gr, potassium iodide 8 gr / fl oz Bot pts, gals Use: Metabolic adjuvant
iodo BROMIDE AND CALCIUM COMPOUND WITH MERCURY. (Tilden) Sodium bromide 2 1/2 gr, potassium iodide 8 gr, mercury bichloride D D 8 gr / fl oz Bot pts gals
 Use: Metabolic adjuvant
iodochlorhydroxyquinoline
 See: Iodochlorhydroxyquin
***iodochlorhydroxyquin, U S P, N N R** 1952; Comp Pow, Tab Oint, U S P Entrokin, Enteroquinol, Entero Viaform, Nio form Quinamide Rometin
 See: Domalarm, cream, oint (Dome) Ounofarm, Oint (C & M Pharmacal) Oun O Creme (Walker) Viaform (Cibal)

* Generic name recognized by N F, U S P or A M A

- W/Aluminum acetate solution, hydrocortisone. See Hydrelt Cream, (Elder).
- W/Bacitracin-methylene disalicylate. See: Anameba, Tabs. (Chicago)
- W/Bismuth subcarb. See: Bismucin, Tab. (Ciba).
- W/Hydrocortisone.
See: Damaform-HC, Cream, Lot. (Dome)
Viatorm-Hydrocortisone, Lotion. (Ciba)
- W/Hydrocortisone alcohol, acetic acid, Xylocaine. See: tidoform-HC Otic (Dome)
- W/Hydrocortisone, LCD, Burrow's Solution
See: Hyburquin Cream (Texas Pharm)
- W/Hydrocortisone alcohol, neomycin sulfate. See: Nalta-Cort, Oint. (Rowell)
- W/Nystatin. See: Nystoform (Dome)
- W/Prednisolone. See: Iodo-Lipalane, Cream, (Spir)
- IODOPROPYLDENEGLYCEROL.** W/Theophylline. See: Theo-Organidin
- ODOCHLOROL, N.N.D. 1962.** (Seate) Chlorodized radiopaque oil contains 26.5% of iodine & 7.5% of chlorine organically combined w/highly refined peanut oil. Bot. 20 cc.
Use: Roentgenographic study.
- IODOCORT CRIME.** (Alan) Hydrocortisone acetate 25 mg., salicylic acid 100 mg., Iodochlorhydroxyquin 50 mg., menthol 12.5 mg /5 Gm. Tube 5 Gm., 20 Gm.
Use: Dermatologic.
- ***IODOFORM, N.F., A.D.R.** (Various Mfr.). Triiodomethane. Pow., Bot. 1/8 oz., 1 oz., 1/4 lb., 1 lb.
Use: For dressing wounds.
- W/Antimony triiodide. See: Camitrol, Amp. (C. D. Smith)
- ODOFORMINE.** See: Siamine, Cap. (Pitman-Moore)
- ODOGEN.** (Metro) A clear sol. of iodized milk protein in a 2% sol. of Sod. iodide. Each cc. contains protein 0.05 Gm., Sod. iodide 0.02 Gm., iodine 0.005 Gm. Vial, 30 cc
Use: (I.M.); induces liquefaction of mucous secretions.
- ODOGUICAN.** (Estra) Gemenal 0.12 Gm., guaiaacal 0.1 Gm., iodalarm 20 mg., camphor 50 mg., eucalyptal 0.1 Gm., in corn oil/2 cc. Amp. 2 cc., Vial 30 cc.
Use: Expectorant in bronchitis.
- ODOHIPPURATE, SODIUM, N.N.R. 1954.**
See: Hippuran, Amp. (Mallinckrodt)
Medapaque (Bell Craig)
- 7-ODO-8-HYDROXYQUINOLINE-5-SULFONATE, SODIUM.** See: Chiniaton, U.S.P.
- W/Purified casein. See: Activin, Amp. & Vial (Bischoff).
- ODO-ICHTHOL.** (Ulmer) Ichthammol 10 gr.,
* Generic name recognized by N.F., U.S.P. or A.M.A.
- tr. iodine 9.3 min., baroglycerin 20 gr., phenol 3.5 gr., hydrastine HCl 1/32 gr. in a glycerol and aromatic vehicle/fl. oz. Bot. 1 pt.
Use: Mucous membrane antiseptic.
- IODOLAKE-S.** (Iakeside) Sod. iodide 80 mg., iodinated peptone 40 mg./2 cc. Vial (30 cc.) Box 1s, 6s & (60 cc.) Box 25s
Use: Nonspecific iodide therapy.
- ODO-LIPALONE.** (Spir) Prednisolone 0.25%, Iodochlorhydroxyquin 3%, in emollient base Tube 5 Gm., 1/2 oz., 1 oz
Use: Oermatoses.
- ODOMETHAMATE, SODIUM, U.S.P., N.N.R. 1954.** (Inj. U.S.P. See: Neo-lapax, Amp. (Schering).
- ODO-NIACIN.** (Cole Chem. Co.) Pot. iodide 125 mg., niacinamide HCl 25 mg./Tab. Bot. 100s. Also Amps. 5 cc.
Use: (2 p.c. I.I.D.) Iodine therapy.
- ODO-NIACIN INJECTION.** (Cole Chem. Co.) Niacinamide HI 50 mg., sod. iodide 20 mg., niacinamide 20 mg., dextrose 5%, chlorobutanol 0.45%/5 cc. Amp. 5 cc., Box 10s.
Use: Iodine therapy.
- ODOPANOIC ACID, U.S.P., N.N.D. 1961.** Tab., U.S.P. See: Telepaque, Tab. (Winthrop Labs.)
- ODOPHENE.** See: Iodophthalein. (Various Mfr.)
- ODOPHINE SODIUM.** See: Iodophthalein Sodium (Various Mfr.)
- ODOPHOR.**
See: Betadine (Corbin & Corwell)
Iodide (Clay-Adams)
Iodine (Iodine Co.)
Wescodyne (West Disinfecting Co.)
- ***ODOPHTHALEIN SODIUM, N.F., N.N.R. 1950.** Tetraiodophenolphthalein Sod., tetraiodophthalein Sod., tetraiodolein Sod. (Antinastin, Cholepulvis, Chalumbin, Foriod, Iodaphene, Iodarayal, Nosaphene Sod., Opacin, Photabiline, Piliophen, Radiatetron)
Merck—Pow., Bot. 3.5 Gm., 25 Gm., 100 Gm., 500 Gm.
Use: Radiopaque medium for cholecystography.
- IODOPROPYLDENE GLYCEROL.** See: Organidin (Wampole)
- IODOPROT.** (Harvey) Sod. iodide 10 mg with protein/cc. Vial 10 cc.
- ODO-PROTEIN.** (New Mexico Pharm.) Foreign protein 0.04 Gm., sodium iodide 0.08 Gm., sodium bisulfate 1%, chlorobutanol 0.5%/2 cc. Vials 30 cc
- ODOTEIN.** (Myers-Carter) Sod. iodide 0.6 Gm., protein 0.04 Gm./2 cc. Vial 30 cc.

IODOPIRACET INJ.

394

***IODOPIRACET INJ**, NF, NND 1958
(Diatrast Diadone Iopyracil Neo Methia
dal Neo Skiodan Nesylom Oporenel Per
Abradil Per Radiographical Pycelosi Pylem
brin Uradine)

Use Radiopaque medium
See: Diadrast Amp (Winthrop Labs)

***IODOPIRACET COMPOUND**, NND 1962
See Diadrast Compound (Winthrop Labs)

***IODOPIRACET CONCENTRATED**, NND
1962 See Diadrast Concentrated (Winthrop
Labs)

IODOPIRINE. (No Mtr currently lists) Anti
pyrine iodide
Use Iodides & analgesic
W/Antipyrimo, caffeine See Felsol (Am
Felsol)

IODO-TAC, (Cetoc Drug) Expectoant 16 oz
IODOTHIOURACIL. See Itrumil, Tab (Ciba)

IODOTOPE, NND 1962 (Squibb) Sodium
Radio iodide-131 Oral and parenteral
forms

IODOXYL. See Sodium Iodomethamate
USP

IONAL. (Strassenburgh) Codeine 30 mg
combined with methyl a tolylquinazalone
resin/cap
Use Analgesic

IONAMIN, NND 1962 (Strassenburgh)
Phentermine Phenyl tert butylamine os resin
complex 15 mg ond 30 mg/Cap Bot
100s 400s
Use Obesity

IONAZE (Eli Lilly) See Propazalamide

ION-EXCHANGE RESINS.
See: Palyomine Methylene Resins, NND
1961

Resins, Potassium Removing
Resins Sodium Removing

IONIL (Owens Labs) Salicylic acid 2%
benzalkonium Cl 0.2%, alcohol 13%/cc
Plus Bot w/disp cap 4 oz
Use: Anti seborrheic

IONLEX, (Fellows) Ferrous gluconate 3.3 gr
liver est 2 gr, thiamine HCl 3 mg, ribo-
flavin 5 mg, niacinamide 30 mg, pyridox-
ine 1 mg, Cal pantothenate 1 mg, folc
acid 1 mg, choline (calculated as chloridel
5 mg, Inositol 10 mg, ascorbic acid 30
mg/Tab Bot 100s 1000s
Use: Hematinic

IONOSOL MB W/Dextrose 5% (Abbott)
Dextrose 5 Gm, mag chlor 14.2 mg, pot
chlor 141 mg, manopat phosph 15 mg,
monosod phosph 21.4 mg, sod lact 260
mg/100 cc Bot 150 250 & 500 cc
Use: IV Dehydration

IONOSOL D-CM (Abbott) Sod chloride 516
mg, Pot chloride 29.4 min, Cal chloride
* Generic name recognized by NF, USP or AMA

368 mg, Mag chlorida 30 mg, Sod lac
tate anhy, 560 mg/1000 cc Bot 1000 cc
Available w/dextrose 5%, Bot 500s, 1000s
Use Replacement fluid therapy

IONOSOL D MODIFIED SOLUTION. (Ab-
bott) Sod Chloride 1.169 Gm, Pot Chloride
2.683 Gm, Cal chloride 255 mg, mag
chloride 133 mg, sod lactate 6.724 Gm,
Invert sugar 100 Gm/1000 cc
Use Parenteral fluid therapy

ION-O-TRATE SOLUTIONS. (Abbott)
Ammonium chloride—(Amm chloride 5.35
Gm (100 mEq), water for injection to
make 20 cc) Vial 20 cc, Box 25s

Dextrose—(Dextrose 20 Gm, water for in-
jection to make 40 cc) Vial 40 cc, Box 25s

Potassium chloride—(Pot chloride 1.49 Gm
(20 mEq), water for injection to make 20
cc) Vial 20 cc, Box 25s

Potassium chloride—(Pot chloride 2.98 Gm
(40 mEq), water for injection to make 20
cc) Vial 20 cc, Box 25s

Potassium phosphate—(Pot phosphate, di-
basic 3.06 Gm, pot phosphate, monobasic
670 mg, water for injection to make 20
cc) Vial 20 cc, Box 25s

Sodium chloride—(Sod chloride 2.92 Gm
(50 mEq), water for injection to make 20
cc) Vial 20 cc, Box 25s

Sodium chloride—(Sod chloride 5.84 Gm
(100 mEq) water for injection to make 40
cc) Vial 40 cc, Box 25s

Sodium lactate—(Sod lactate 5.6 Gm (50
mEq) water for injection to make 20 cc)
Vial 20 cc, Box 25s

Sodium lactate—(Sod lactate 11.2 Gm (100
mEq), water for injection to make 40 cc)
Vial 40 cc, Box 25s

***IOPANOIC ACID**, USP, NND 1962
Tab USP See: Telepaque, Tab (Win-
throp Labs)

***IOPHENDYLATE INJECTION**, USP,
NND 1962 Ethiodan, Myodil See Ethyl
iodophenyl endocylate inj

***IOPHENOXIC ACID**, USP, NND 1962
Tab USP a (2.4.6 triado 3 hydroxybenzyl)
butyric acid See: Teridax, Tab (Schering)

IOPREP ANTISEPTIC. (Johnson & Johnson)
Iodine 1%, non ionic surfactants 10%,
de ionized water 89%/liq Bot 16 oz
gals

Use Topical antiseptic
IOQUIN SUSP (Abbott) 10% w/v diado
hydroxyquin USP Bot 120 ml

Use: Seborrheic dermatitis of the scalp con-
trol of common dandruff

IOSOL (Morse Labs) Thymol Iodide Oint
1/2 oz
Use: Topically antiseptic.

- IOSTHIPENDYL HYDROCHLORIDE.** See: Theruhistin, Tab. & Syr. (Ayerst)
- IOXAN, PHTHOLATID.** (Serriff & Case) Iodotannic S&C 2.25%, phenol 1.00%, glycerin 7.50%, methyl D 0.3%, aqueous base. Sol 1 pt., 1 gal.
Use: Antiseptic, astringent and local anesthetic
- *IOTHALAMIC ACID.** (Malinchrodt) 3-acetamido-2,4,6-triiodo-N-methylsophthalamic acid
See: Conroy
(meglumine iothalamate 60%)
Angio-Conroy
(sodium iothalamate 20%)
- *IOTHIOURACIL SODIUM, N.N.D. 1962.**
Sod salt of 5-Iodo-2-thiouracil. See: Iru-mil Sodium, Tab (Ciba)
- IOZANTHAL.** (Mastell) Theobromine 5 gr., phenobarbitol $\frac{1}{2}$ gr., ethyl dilodabromistate 1 gr./Tab. Bot. 50s, 250s, 1000s.
Use: (1 I.L.D.) Sedative w/diuretic.
- *IPECAC, U.S.P.** See: Alcresta Tablets of Ipecac (Lilly).
- W/Opium pow., acetophenetidin, atropine sulf., acetylsalicylic acid, camphor monobromated, caffeine. See: Dophenco, Cap. (Blue line)
- W/Phenolphthalein, alole, belladonna lvs. ext. & styeheline sulfate. See: Alothal, Tab. (Mallard).
- IPESANDRINE.** (Saedox) Opium alkaloids 108 mg., ameline HCl 2.2 mg., w/ephedrine HCl, balsam tolu, cherry laurel water & alcohol/oz. Bot. 1 pt., 1 qt., 1 gal.
Use: (1 tsp. q. 3 h., p.r.n.) Narcotic cough preparation.
- IPRAL CALCIUM, N.N.R. 1951.** (Squibb) (Proborbital Cal.) 5-Ethyl-5-isopropylbarbiturate. Cal. Tab. ($\frac{1}{2}$ gr. A 2 gr) Bot. 100s, 1000s.
Use: ($\frac{1}{2}$ gr. to 2 gr., m.p.) Sedative.
- IPRAL SODIUM, N.N.R. 1955.** (Squibb) (Proborbital Sod) Ethylisopropylbarbituric acid. Elixir (1 gr./dr.) Bot. 1 pt., 2 Tab. (4 gr.) Bot. 100s.
Use: Sedative A hypnotic.
- *IPRONIAZID, N.N.D. 1961.** See: Marsilid, Tab (Roche)
- IPSATOL.** (Davies, Rose) Squill, Ipecac, eucalyptus, ammonium chloride A Infu balsam. Bot. 3 oz.
Use: (1-2 tsp. q. 3 h) Cough preparation.
- IRON, N.N.D. 1962.** (Lakeside) Ferrous fumarate 200 mg./Tab. Bot. 100s.
Use: Hematinic.
- IRILIN.** (Tilden) Methyl salicylate, menthol, camphor, oil mustard. Bot. pts., gots.
Use: Counter-irritant liniment.
- IRISIN.** A polysaccharide found in several species of Iris. See: Herb, Tab. (Cole Pharmacy)
- IRO-B.** (Maltbie) Elixir (ferrous sulfate 3 gr., Vit. B-1 0.4 mg./dr.) Bot. 1 pt., 1 gal. Tab. (ferrous sulfate 3 gr., Vit. B-1 0.3 mg., B-2 0.5 mg./Tab) Bot. 100s, 500s.
Use: Hematinic.
- IROCAINE.** See: Procaine Hydrochloride (Various Mfr.)
- IRONCINE.** (Reed & Carnrick) Iron Sod. malate 200 mg., copper sulfate 4 mg., liver desic. 200 mg., thiamine HCl 0.17 mg., Vit. D 67 u., Vit. B-12 actively equiv. to 1 mcg./Tab. Bot. 100s, 500s, 1000s.
Use: Anemias due to iron deficiency.
- IRONEP-B.** (Lanett) liver fraction 27 gr., iron and ammonium citrate 3 gr., vitamin B-1 2 mg., B-2 1 mg., B-6 0.5 mg., cal. pantothenate 1 mg., nicotinamide 10 mg./3 cap. Bot. 100s, 1000s.
Use: Iron deficiency.
- IRONIC.** (Hetro Med) Ferrous fumarate 120 mg., ciferavin 1 mg., thiamine HCl 1 mg., pyridoxine HCl 1 mg., Vit. B-12 activity 5 mcg., ascorbic acid 30 mg., polygalacturonic acid 65 mg., desiccated liver 65 mg., folic acid 0.1 mg., cobaltous chloride 0.1 mg., citric acid. 30 mg./Cap. Bot. 100s, 1000s.
Use: Treatment of iron deficiency anemia
- IRONIN-O.** (Mission) Ferrous gluconate 1000 mg., iron 116 mg Vit. B-12 (crystalline, or resin) 60 mcgm., C 120 mg., A acetate 15,000 u., D-2 1500 u., R-1 60 mg., B-2 60 mg., B-6 3.0 mg., R 3 3.0 mg., B 3 60 mg., calcium lactate 376 mg., calcium carbonate 210 mg., calcium 159 mg., copper (as sulfate) 0.45 mg., manganese (as citrate soluble) 0.75 mg., zinc (as oxide) 0.24 mg., potassium (as chloride) 15 mg., magnesium (as carbonate) 7.5 mg./3 Tab. Bot. 100s.
Use: Hematinic nutrient
- IRON, N.F.**
W/Calcium. See: Ricalol, Tab. (Ortho)
W/Copper. See: Calron, Tab. (Abbott)
W/Copper A Vit. B-12. See: Calron w/vitamin B-12 (Abbott)
W/Folic acid. See: Hemabolids w/folic Acid, Tab. (Arlington)
W/Polyoxyethylene glucite manalaurate (Socagen). See: Simroe, Cap. (Merrell)
W/Vitamins mixed. See: Vitamin preparations.
- IRONATE.** (Wyeth) Ferrous sulfate, dried, 227 mg., copper 1 mg., Vit. B-1 5 mg., B-2 2 mg., B-6 1 mg., B-12 5 mcg., C 75 mg., folic acid 0.33 mg., Cal. pantothenate 1

- mg, niacinamide 20 mg, liver desic, NF, 175 mg /Cap Bot 100s
 Use: (1 t i d) Nutritional anemias
- IRONATE.** (Mallard) Represents iron 50 mg as ferrous sulfate, ferrous gluconate & Sod ferric pyrophosphate
- IRONATE-B.** (Mallard) Ironate 5 gr, w/not less than Vit B 1 1.5 mg, B 2 0.5 mg, niacinamide 5 mg /Tab Bot 100s 500s, 1000s
 Use: Hematinic
- IRONATE-B PLUS** (Mallard) Ironate 5 gr Vit B 1 5 mg, B 2 5 mg, B 12 (cryst) 5 mcg, C 50 mg, folic acid 0.5 mg, nicotinamide 10 mg, gastric mucosa desic 75 mg w/cobalt & mang sulf /Tab Bot 100s 500s, 1000s
 Use: Hematinic
- IRON BILE SALTS.** See Bilron, Pulvules (Lilly)
- IRON CACODYLATE, NF VII**
 Merck Sharp & Dohme—Vial 1 gr /5 cc, Box 25s
 Parke, Davis—Vial 1 gr /5 cc Box 25s
 Ingram—Vial 1 gr /2 cc & 2½ gr /5 cc Box 12s, 25s, 100s
 Use: IY
 W/Cal cacodylate 6 gr (Ingram) Iron Cacodylate 1 gr /5 cc Vial, Box 12s, 25s, 100s
 W/Copper cacodylate, Sod cacodylate See Ferro Cuprin, Amp (Kremers Urban)
 W/Sod cacodylate strychnine and sod glycerophos See: Glisotal, Amp (Metro palatin Lab)
 W/Strychnine cacodylate 1/65 gr Iron cacodylate 1 gr /2 cc Vial, Box 12s, 25s 100s (Ingram)
 W/Strychnine and sod glycerophos See Gl comp Amp (Estral)
- IRON CACODYLATE COMPOUND.** (Ingram) Cacodylates of Iron ½ gr, Sod 2 gr strychnine 1/32 gr, Sod glycerophosphate 1½ gr /2 cc Vial, Box 12s, 25s, 100s
- IRON CARBONATE COMPLEX.** See Polyferase (Pitman Moore)
- IRON CHOLINE CITRATE COMPLEX.** See: Chel Iron, Tab (Kinney)
 Ferrolip (Flint, Eaton)
- IRON-DEXTRAN INJ., USF, NND 1960** (Not available) See: Interon, Amp (Lek side)
- IRON & FOLIC ACID** (National Drug) Ferrus sulfate essic 0.2 Gm, folic acid 17 mg /Cap Bot 100s 500s
 Use Orally; hematinic.
- IRON PEPTONIZED.** See: Saleron, Tab (Messengier)
- IRON PROTEIN COMPLEX.** See Ovaferzin (Barnes)
- IRON, QUININE & STRYCNINE, ELIXIR, NF** (Merck Sharp & Dohme) Ferric citrochloride 12.5 cc, quinine HCl 0.8 Gm, strychnine sulfate 17.5 mg, alcohol 25%/100 cc Bot 1 pt, 1 gal
- IRON & QUININE WITH STRYCNINE PHOSPHATE, LIGHT COLOR, NOT NF** (Merck Sharp & Dohme) Ferric citrochloride tr 5 cc, quinine HCl 0.45 Gm, strychnine phosphate 20 mg, alcohol 15%/100 cc Bot 1 gal
- IRON SULFATE W/Maalox** See: Fermalox Tab (Rorer)
- IRON-VITAE,** (Kirk) Liver, iron, and vitamins Vial 30 cc
 Use: Hematinic
- IRONYL.** (Physiological Chem) Iron salt of yeast adenylic acid (equiv to 4.4 mg iron & 28 mg adenosine 2 monophosphate) Amp 1 cc, Box 10s, 25s, 100s Tab 324 mg Bot 12s, 25s, 250s
 Use: Nematonic
- IROPHOS-D.** (Lonnert) Dicalcium phosphate anh 330 mg, ferrous sulf 30 mg, vit D 333 u /cap Bot 500s, 1000s
 Use: Nematonic
- IRUSOL** (Naskell)
 Tab: (Ferrus sulfate 3.5 gr) Bot 100s 500s, 1000s
 W/Vit B 1: (Ferrus sulfate 3.5 gr, Vit B 1 8 mg./dr) Bot 1 pt, 1 gal
 Fortified: (Ferrus sulfate 5 gr, Vit B 12 4 mcg, folic acid 0.5 mg, Vit C 30 mg, B 1 5 mg, B 2 3 mg, niacinamide 30 mg, Vit B-6 0.67 mg, Cal pantothenate 2 mg /tab) Bot 100s, 500s
 Use: Nematonic
- IRUSOL-C.** (Haskell) Ferrus sulfate 5 gr, ascorbic acid 75 mg /tab Bot 100s
 Use: Anemias
- IROETHERON B** (Tilden) Cal gluconate 2 gr, ferrous sulfate 3 gr, w/Vit B 1 0.24 mg /Tab Bot 100s, 500s
 Use: (2 t i d) Dietary supplement
- IROETHERON B-PLUS.** (Tilden) Ferrus sulf 3 gr, mactn 3 mg, Vit B 1 0.3 mg, C 25 mg /Tab Bot 100s
 Use: Treatment of anemias
- IROX.** (Cafe Chem Co) Ferrus gluconate Tab (5 gr) Bot 100s
 Use: Hematinic
- IRRADIATED 7-DEHYDROCHOLESTEROL.** See: Activated 7 Dehydrocholesterol, USF
- IRRADIATED ERGOSTEROL.** See: Calciterol USF
- IRRADOL-A.** (Parke, Davis) Haliver oil other fish oils & irradiated ergosterol combined w/other vitamins & iron & malt ext (non-diatolic) Vit A 18,000 u, D 2600 u, B 1 4

* Generic name recognized by NF, U.S.P. or A.M.A.

mg., B-2 2 mg., nicotinamide 40 mg., iron & ammonium citrate, green, 4 gr., malt ext. (nondiastolic) q.s./fl. oz., Jar 1 lb., 2½ lb. (hospital size).

Use: Nutritional therapy.

IR-VATINE. (Smith-Dorsey) Ferrous lactate 5 gr., thiamine HCl 2 mg./Tab., Bot. 100s. 1000s.

Use: Anemias.

IRVITOL. (Rorer) Ferrous sulfate 20 gr., thiamine HCl 6 mg./fl. oz., Elixir, Bot. 14 oz., 1 gal.

Use: Anemias.

IRWINONE. (Irwin, Neisler) Vit. B-1 18 mg., B-2 12 mg., niacinamide 60 mg., cal. pantothenate 30 mg., rice bran ext. 72 gr., fresh liver 120 gr., dry yeast 18 gr./fl. oz. Bot 1 pt., 1 gal.

Use: Dietary supplement

ISACEN. N.N.R. 1940. See: Diacetylhydroxyphenylisatin, Prep. (Various Mfrs.)

ISADOXOL. (Blair Labs) Diacetyl sod. sulfosuccinate 100 mg., diacetylhydroxyphenylisatin 1 mg./Tab., Bot. 50s.

Use: Constipation.

ISAMIN-C. (Arlington-Funk) Pyrilamine maleate 25 mg., ascorbic acid 100 mg., caffeine 16 mg./Tab., Bot. 50s, 100s; Syr., Bot. 4 oz., 16 oz.

Use: Antihistaminic.

ISATIN. See: Diacetylhydroxyphenylisatin.

I-SEDRINE COMPOUND. (Lilly) Ephedrine 1% combined w/glucanonic acid, merthiolate 1:5000 in an aromatized sol. made isotonic w/dextrose. Bot. 1 oz., 1 pt.

Use: Shrinkage of nasal mucous membranes.

I-SEDRIN PLAIN. (Lilly) Ephedrine alkaloid 1% combined w/glucanonic acid w/chlorobutanol 0.5%. Bot. 1 oz., 1 pt.

Use: Nasal decongestant.

ISMELIN. N.N.D. 1962 (Ciba) *Guarethidine sulfate, 10 mg. and 25 mg./Tab. Bot. 100s.

Use: (1 tab. daily) Hypertensives.

***ISOAMINILE.** (Robins) 4-Dimethylamino-2-isopropyl-2-phenylvaleranoltrile.

Use: Antitussive.

ISOAMYLHYDROCUPREINE DIHYDROCHLORIDE. See: Eucupie Dihydrochloride. Prep. (Rare-Galen).

ISOAMYL NITRITE. See: Amyl Nitrite. U.S.P. (Various Mfrs.)

ISOAMYNE. See: Amphetamine (Various Mfrs.)

ISOBARB. (U. S. Standard) Pentobarbital Sod. outer coat ¾ gr., Pentobarbital Sod inner core ¾ gr./Tab. Box 100s. 1000s.

Use: (1 h.s.) Sedative.

* Generic name recognized by N.F., U.S.P. or A.M.A.

***ISOBORNYL THIOCYANOACETATE, TECHNICAL.** N.N.D. 1962.

See: Bornate, N.N.R. (Wyeth).

Thioite. (No Mfr. currently lists)

Use: Pediculicide.

ISO-BROVITE. (Braemmel) Bot. 100s, 1000s.

Use: Special dietary supplement. Vit. A 5000 u., Vit. D 400 u., Vit. C 600 mg., Vit. B-1 10.0 mg, Vit. B-2 5.0 mg, Vit. B-6 1.0 mg, Vit. B-12 2.0 mcg, Folic Acid 0.5 mg, Calcium Pantothenate 100 mg, Niacin 100 mg., Choline Citrate 300 mg, Inositol 300 mg., Dry Yeast Powder 240.0 mg.

ISOBUTYLALYL BARBITURIC ACID, N.N.R. 1952. See: Allylbarbituric Acid, N.F. Sandoptal, Tab. (Sandoz)

W/Acetophenetidin, acetylsalicylic acid, caffeine. See: Fiorinal, Cap. (Sandoz)

W/Acetophenetidin, aluminum aspirin, caffeine. See: Alatal, Tab. (L. L. Drug)

ISOBUTYL p-AMINO BENZOATE, (No Mfr. currently lists.) (Cyclafarm) (Isacaine).

Use: As a surface anesthetic.

W/Bismuth subnitride, zinc oxide, bismuth subcarb. & boric acid. See: Proclafarm, Supp. (Testagor).

W/Procaine base & benzyl alcohol. See: Proclacaine, Amp. (Columbus).

W/Tetracaine HCl & benzalkonium chloride. See: Isotraine, Oint. (Columbus).

W/Tyrothricin, tetracaine HCl & menthol. See: Tyrosotraine, Loz. (Columbus).

W/Tyrothricin & benzalkonium chloride. See: Cyclo-Thrycin, Loz. (Hance).

ISOCAINE. See: Isobutyl p-aminobenzoate.

ISOCAINE LACTATE. See: Dimethylecaine lactate.

***ISOCARBOXAZID.** 1-Benzyl-2-[5-methyl-3-isoxazolyl-carbonyl]hydrazine. See: Marplan, Tab. (Rache Lab.)

ISOCLOL. (Haskell) Chlorprophenpyridamine maleate 4 mg., d-isopropylamine HCl 25 mg./Tab. Bot. 100s.

Use: Colds, sinusitis, hayfever, etc

ISOCLOL CAPS. (Arner-Stone) Chlorpheniramine maleate 10 mg, d-isopropylamine HCl 65 mg./Cap. Bot. 50s, 250s.

Use: Relief of hay fever.

ISOCOCAINE. See: Pseudococaine. (No Mfr. currently lists)

ISOCRIN PRULETS. (Warner-Chilcott) Oxyphezzin acetate. Diacetylhydroxyphenylisatin 5 mg./Tab., Bot. 50s.

Use: As a laxative.

ISODINE. N.N.D. 1959. (Isidine Pharmacol Corp.) Iodine & polyvinylpyrrolidone (PVP) Povidone-Iodine. Bot. ½ oz. Available as gargle, aetz. & loat pow.

- Use Topical antiseptic (No sting & non toxic)
- ISOEPHEDRINE HCl** d Isoephedrine HCl
See Pseudoephedrine HCl
- W/Chlorpheniramine maleate, dihydrocodeinone bitartrate See *Hasanone Na 2 Sal* (Hoske'l)
- W/Chlorpheniramine maleate, salicylamide N acetyl p aminophenol See *Isogastic Tab* (Arnar Stone)
- W/Chlorpropranpyridamine maleate See *Isoclar Tab* (Hoskell)
- W/Theophylline phenyltoloxamine See *Aztec Tabs* (Arnar Stone)
- ISOEPHEDRINE**, W/Dexbrompheniramine maleate See *Disaphrin Tab* (White)
- ***ISOETHARINE** (Brean) a β isopropylamino propylprotocatechuy alcohol See *Dilabron*
- ***ISOETHARINE METHANESULPONATE**, W/Phenylephrine HCl thenyldiamine, tream propellan See *Brankometer* (Brean)
- ISOFEDROL**, (Blue Line) Ephedrine sulfate 1% in an isalonic sol Bat 1 oz, 1 pt.
Use Nasal vasoconstrictor
- ***ISOFUCROPHATE**, NND 1962 Ophth Oint Ophth Sol USP Diisopropyl fluoro phosphate See *Flarapryl Oint & Soln* (Merck Sharp & Dohme)
- ISOGESIC**, (Arnar Stone) d Isoephedrine HCl 25 mg chlorpheniramine maleate 2 mg salicylamine 150 mg, N acetyl p amino phenol 150 mg /Tab Bat 100s
Use: Common cold
- ISOHIST IMPROVED OPNTH SOL** (Broom-mal) Chlorpheniramine maleate 0.10%, phenylephrine HCl 0.125% and methyl cellulose 0.5% 15 cc plastic dropper bottle
Use Allergic and chronic conjunctivitis
- ISO-IODEIKON**, NNR 1951 See *Phen-tethothalein Sodium* (No Mfr currently lists)
- ISOLEUCINE**, DL Isoleucine
Merck—Bat 5 Gm, 10 Gm
Winthrop—Bat 25 Gm
Use Nutrient
- ISOLYN**, (Abbott) (Isoniazid) Isonicotinic acid hydrazide Tab (50 mg) Bat 100s 1000s
Use Streptomycin resistant tuberculosis
- ISOLYTE SOLUTION** (Don Baxter) Sod acetate 0.64 Gm, sod chloride 0.30 Gm, pot chloride 0.075 Gm, sod citrate 0.079 Gm, cal chloride 0.037 Gm, mag chloride hexahydrate 0.031 Gm /100 cc Flask 250 cc 1000 cc Also available w/Dextrose 5% Flask 1000 cc
Use: IV, balanced electrolyte
- ISOMEPROBAMATE**, See: *Carsoprodol*
- ISOMETH**, (Lloyd, Dabney and Westerfield)
Isomethiene tartrate 2 Methylamino 6 methyl 5 heptene tartrate 100 mg/cc Vial 10 ml
Uses Antispasmodic
- ***ISOMETNAZINE HCl**, (Winthrop Labs) 10-(2 dimethylamino 1 methylethyl)phenanthroline HCl
Use: Antinuseant, antihistaminic, and sedative
- ***ISOMETHEPTENE HCl**, NND 1962 Methyl isoocetyl amine mucate or tartrate Also available as Hydrochloride
See: *Isometh*, Vial (Lloyd, Dabney & Westerfield)
Isanyl, Vial (Bell Pharm)
Octin Mucate, Prep (Knoll)
- W/N acetyl p-aminophenol, dichloralphenazone See *Miorin*, Caps (Cornrick)
- W/Acetyl p-aminophenol, salicylamide, caffeine See *Nygron* (Dumas Wilson)
- ISOMYN**, See *Amphetamine* (Various Mfr.)
- ***ISONIAZID**, USP, NND, 1962; Tab, USP Isonicotinic acid hydrazide, isonicotiny hydrazide
See: *Armozide*, NNR (Armour)
Colinoxin, NNR (Pfizer)
Dinacrin, Tab (Winthrop Labs)
Disubun, NNR (Schering)
INN, NNR (Lilly)
Isolyn (Abbott)
Isonozid, NNR (Walker)
Isonico, Tab (Kirkman)
Isonizide (Various Mfr.)
Isoniazid (Lanett)
Niadrin, Tab, NNR (Endo)
Nisonyl, Tab, NNR (Porko, Davis)
Nicozide, NNR (Prama)
Nidaton (Organon)
Nydrozid, NNR (Squibb)
Pyrazidin, NNR (Nepero)
Rimfon, NNR (Hoffmann La Roche)
Teebacoin, Tab (Consol Mid)
Tisin, NNR (U S Vitamin)
Tyvid, NNR (Merrell)
Zinadan, NNR (K Victor)
Zanazide (G P Harvey)
- W/p Aminosulicylate Sod
See *Dipasic* (Ponray) Under study
Isopacin, Tab (Consol Midland)
- W/Pyridoxine HCl See *PI N Tab* (Lanett)
- W/Seromycin See *Seromycin W/Isoniazid Pulv* (Lilly)
- ISONICO**, (Kirkman) Isoniazid 50 mg & 100 mg /Tab Bat 100s, 1000s 5000s
Use: Tuberculosis
- ISONICOTINIC ACID HYDRAZIDE**, NND 1962 See *Isoniazid* USP
- ISONICOTINYL HYDRAZIDE** See *Isoniazid*
- ISONIPECAINE HYDROCHLORIDE**, NNR 1954 See *Meperidine Hydrochloride*, USP

- ISONORIN SULFATE**, N.N.D. 1962. (Smith, Miller & Potch) Isopropylarterenal sulfate. Sol (1 200) Bot. 15 cc. Tab (10 mg) Eot. 100s.
Use: Orally (1-2 m.p.); vasoconstrictor & central nervous system stimulant.
- ISONYL**, (Bell Pharm.) Isometheptene tartrate 100 mg. Vial 10 cc.
Use: I.M., treatment of smooth muscle spasm.
- ISOPACIN**, (Consol. Midland) p-Aminosalicylate Sod. 500 mg., isoniazid 12.5 mg./tab. Bot. 500s., 1000s.
Use: Treatment of tuberculosis.
- ISO-PAR**, (Coparaffinate) N.N.D. 1962. (Medicol Chem.) A mixture of water-insoluble isoparaffinic acids. Oint. (coparaffinate 17%, titanium dioxide 4%) Jar 14 Gm., 28.5 Gm., 114 Gm., 454 Gm.
Use: Topically; antifungal agent.
- ISOPENTAQUINE**. See: Pentaquine Phosphate (Abbott).
Use: Curative & prophylactic in malaria.
- ***ISOPNANE INSULIN**, U.S.P., N.N.D. 1962. See: NPH Insulin (Lilly).
NPH Insulin, (Squibb)
- ISOPHRIN HYDROCHLORIDE**, N.N.R. 1954. (Broemmel) Phenylephrine HCl, Nasal Sol. (0.25%) Bot. 1 oz., 4 oz., 1 pt. Ophth. sol. (0.125%) Bot. 15 cc., 118.3 cc.
- ISOPREGNOLONE**.
See: Duphaston, Tabs. (Phillips Roxone).
Dydrogesterone
- ISOPRENALINE**, See: Isoproterenol NCI
- ISOPHYLLIN**, (Maney) Isoproterenol 10 mg. in outer coating; in inner core racephedrine HCl 16 mg., phenobarbital E mg., neothylline 130 mg./Tab. Eot. 100s., 1000s.
Use: Control of bronchospasm in chronic asthma.
- ISOPRO**, (Philadelphia Ampoule) Isopropylarterenal 0.5%. Bot. 10 cc., 30 cc.
Use: Symptomatic treatment of bronchial spasm.
- ISOPRO NP**, (Philadelphia Ampoule) Isopropylarterenal 0.4%, phenylephrine HCl 2%, propylene glycol 10%. Bot. 10 cc.
Use: Bronchial asthma and emphysema.
- ***ISOPROPAMIDE IODIDE**, N.N.D. 1962. (3-Carbamoyl-3,3-diphenylpropyl) diisopropylmethyl ammonium iodide. See: Darbid, Tab. (Smith, Kline & French)
W/Phenylpropranolamine HCl, chlorpheniridine maleate. See: Ornade, Span. (Smith, Kline & French)
- ***ISOPROPHENAMINE** NCI. 1-(O-chlorophenyl)-2-isopropylaminoethanol HCl
* Generic name recognized by N.F., U.S.P. or A.M.A.
- ISOPROPYDRIN W/CALCIUM IODIDE**. (Abbott) Isoprenoline 0.06%, calcium iodide 3%. Syr
Use: Bronchial asthma
- ISOPROPYLARTERENOL HYDROCHLORIDE**, U.S.P., N.N.D. 1962 See: Isoproterenol Hydrochloride, U.S.P.
Use: Asthma, vasoconstrictor & allergic states.
- ISOPROPYLARTERENOL SULFATE**, N.N.D. 1962. See: Isoproterenol Sulf.
- ISOPROPYL-NORADRENALINE** NCI. See: Isoproterenol HCl, U.S.P.
- ISOPROTERENOL**. See: Medihaler-ISO, Inhaler, (Riker)
W/Racephedrine HCl, phenobarbital, neothylline. See: Asophylline, Tab. (Cristol)
W/Theophylline, ephedrine & secobarbital. See: Aluted, Tab. (Hart)
- ***ISOPROTERENOL HYDROCHLORIDE**, U.S.P., N.N.D. 1962; Inhalation, U.S.P.; Tab, U.S.P.; Isopropylarterenal HCl, N.N.R. 1954; alpha-Isopropylaminomethyl protocatechuyal alcohol HCl, Oleudrin.
See: Aleudrin. (National Drug)
Aludrin, N.N.R. (Lilly)
Asdrin, Tab. (Cofe)
Isopro, liq. (Philadelphia Amp.)
Isuprel, N.N.R. (Winthrop Labs.)
Use: Asthma, vasoconstrictor & in allergic states. (Subcu. 10 mg.; Inhalation 0.25 mg.)
W/Ethaverine NCI, theophylline, butobarbital. See: Eendrin, Tab. (Bene)
W/Phenobarbital, benzylephedrine HCl, theophylline. See: Isuprel-Franol. (Winthrop)
W/Phenylephrine bitartrate. See: Medihaler-Duo, (Riker).
W/Phenylephrine HCl, propylene glycol See: Isopro NP, liq. (Philadelphia Ampoule).
W/Phenylpropranolamine HCl, glyceryl guaiacolate. See: Eranodyl Padiotab 17. (Columbus).
W/Theophylline, ephedrine sulfate, phenobarbital. See: Nephentalin, Tab. (Leeming)
- ***ISOPROFRENOL SULFATE**, N.F., N.N.D. 1962. Inhal. N.F.
See: Duraldrin, Tab. (Durst)
Isanorin, Liq., N.N.R. (C. D. Smith)
Norisadrine, Liq., N.N.R. (Abbott)
W/Phenylephrine HCl. See: Nebu-Prel. (Mahan)
- ISOPTO-ALKALINE**, (Alcon) 1% Methyl cellulose sterile ophth. sol. Dropper Bot. 15 cc.
Use: Ophthalmic emollient & lubricant
- ISOPTO-ATROPINE**, (Alcon) Atropine sulfate 0.5% and 1.0% in methyl cellulose solution. Drop-tainer S, 15 cc.
- ISOPTO-CARBACHOL**, (Alcon) Carbachol U.S.P. 0.75%, 1.5%, or 3%. In a sterile

- buffered solution of 1% methylcellulose
Drop Tainer 15 cc
Use: Glaucoma contraindications corneal
injuries
- ISOPTO-CARPINE.** (Alcon) Pilocarpine HCl
in stable sterile 0.5% methyl cellulose sol
w/isotonicity of lacrimal fluid & adjusted
to pH 6.4 Bot Drop Tainer (0.25%, 0.5%,
1%, 2%, 3%, 4% & 6% pilocarpine HCl)
Drop tainer 15 cc
- ISOPTO-CETAMIDE.** (Alcon) Sod sulfaceta-
mide 15% in buffered pH 7.4, 0.5 methyl
cellulose sol Dropper Bot 15 cc & Drop
Tainer 15 cc
Use: Conjunctival infections
- ISOPTO-CETAPRED DROPS** (Alcon) Sulfo-
cetamide sod USP 10% prednisolone
USP 0.25% methylcellulose (4000 cps)
USP 0.5% in a sterile buffered and stable
suspension Plas Bot 5 cc
Use: Chronic and allergic blepharitis and
conjunctivitis
- ISOPTO-CORTISONE.** (Alcon) Cortisone ace-
tate 0.5% or 0.25% in a methyl cellulose
sol Susp Ophth Drop tainer (0.3%) 5 cc
(0.25%) 15 cc
Use: Eye infections
- ISOPTO-ESERINE.** (Alcon) Eserine salicylate
0.25% & 0.5% in stable sterile 0.5%
methyl cellulose sol preserved against con-
tamination & w/adjusted pH & tonicity
Drop tainer Bot 15 cc
Use: Treatment & control of glaucoma
- ISOPTO-EUCATROPINE 3%.** (Alcon) Euca-
tropine HCl 3% ophth w/methylcellulose
Drop tainer 15 cc
- ISOPTO-FRIN** (Alcon) 1/4% Phenylephrine
HCl in a methyl cellulose solution Drop
tainer Bot 15 cc
- ISOPTO-HOMATROPINE.** (Alcon) Homatra-
pine HBr 2.0% and 5.0% in methyl cellu-
lose solution Drop tainer 15 cc
- ISOPTO-HYDROCORTISONE.** (Alcon Labs)
Hydrocortisone acetate 0.5% or 2.5%. Susp
Ophth Drop-tainer 5 cc.
Use: Eye infections
- ISOPTO-HYOSCINE** (Alcon) Hyoscine HBr
0.25% in a methyl cellulose solution Drop-
tainer 15 cc.
- ISOPTO-MYDRAPRED DROPS.** (Alcon) Atro-
pine sulf USP 1%, prednisolone USP
0.25%, methylcellulose 0.5% in a sterile,
buffered, isotonic suspension Plas Bot
5 cc.
Use: Granulomatous and non-granulomatous
uveitis
- ISOPTO-P-ES.** (Alcon) Pilocarpine 2%, eser-
ine salicylate 0.25%, sterile buffered, iso-
tonic
- ISOPTO-P-H-N.** 0.5% or 1.5% (Alcon) Hy-
drocortisone acetate 0.5% or 1.5%, with
polymyxin B sulf 16,250 u and neomycin
sulf 5 mg/cc Drop tainer 5 cc
- ISOPTO-P-N-P.** (Alcon) Phenylephrine
0.12%, neomycin sulfate 5 mg, polymyxin
R sulfate 16,250 u, isotonic vehicle con-
taining 0.5% methylcellulose Bot 5 cc
Use: Mild ocular infections
- ISOPTO-PLAIN.** (Alcon) Methylcellulose
0.5% Drop tainer 15 cc
Use: Ophthalmic lubricant and emollient
- ISOPTO-PHENYLEPHRINE.** (Alcon) Phenyl-
ephrine 2.0% and 10% in a methyl cellu-
lose solution Drop tainer 15 cc 10% also
available in 5 cc
- ISOPTO-PREDNISOLONE.** (Alcon) Predniso-
lone 0.25%, phenylephrine HCl 0.12%,
methylcellulose 0.5% 5 cc plastic Drop
Tainer dispenser
Use: Treatment of inflammatory and allerg-
ic eye disorders
- ISOPTO-STEROFRIN** (Alcon) Hydrocortisone
USP 0.5%, phenylephrine 0.12%, isotonic
solution of 0.5% methylcellulose Drop
tainer 5 cc
Use: Ocular allergies and inflammatory
conditions in absence of infection
- ISORDIL** (Ives Cameron) Isosorbide dinitrate
1,4,3,6 dranhydro-sorbitol 2,5 dinitrate 10
mg/Tab Bot 100s Also 5 mg sublingual
tabs Bot 100s
Use: Control of angina pectoris and coro-
nary insufficiency
- ISORDIL TEMBIDS** (Ives Cameron) Isosorb-
ide dinitrate 40 mg/Tab Bot 100s
Use: For Angina pectoris
- ISORDIL WITH PHENOBARRITAL.** (Ives
Cameron) Isosorbide dinitrate 10 mg,
phenobarbital 15 mg/Tab, Bot 100s
Use: Angina pectoris
- ISOSORBIDE DINITRATE.**
See: Isordil, Tab (Ives Cameron)
Isordil Tembids (Ives Cameron)
W/Phenobarbital See Isordil w/Pheno-
barbital, Tabs (Ives Cameron)
- ISO-SYNECAINE.** (Warren Teed) Phenyleph-
rine HCl 0.03% & procaine HCl 2% Vials
10 cc
Use: In] local anesthetic
- ISOTHIMPENDYL HCl.** See: Theruhistin, Prop-
(Ayerst)
- W/Aspirin, phenacetin, l phenylephrine HCl
ascorbic acid Sear Kryl, Tab (Ayerst).
W/Dmethaxanate HCl, l phenylephrine HCl,
acetaminophen, ammonium chloride, Sod.**

- citrate, chloroform. See: Cothera Compound, Liq (Ayers)
- ISO-THRICIN.** [Approved Pharm.] Tyrothricin 2 mg., benzocaine 5 mg., cetyl dimethyl benzyl ammonium chloride 0.1% /oz. Bot. 14s, 1000s.
Use: Antibiotic-anesthetic-analgesic lozenges
- ISOTOGEN.** [Broemmel] Sulfabenzomide 1%, hexachlorophene 1%, thimerosal 1-4000, & Tween 80 1%, benzocaine 1% in a polyethylene glycol base. Bot. ½ oz.
Use: Ear infections.
- ISOTOPES RADIOACTIVE, N.N.D. 1960.**
- ISOTRAINE.** [Columbas] Tetracaine HCl ½%, Isocaline 3% w/benzalkonium chloride. Cream, Tube 1 oz., Jar 1 lb.
Use: Topically; local anesthetic.
- ***ISOXSUPRINE HCl, N.N.D. 1962** 1-(p-Hydroxy phenyl)-2-(1'-methyl-2'-phenoxy ethylamino) propanol-1 HCl. See: Vasodilan (Mead Johnson)
- ISTIZIN.** [Winthrop Labs] (Donthron, N.F.) 1,8-Dihydroxyanthraquinone.
Bouises: (½ oz.) Box 24s.
Pow.: Bot. 1 & 5 lb.
Tab.: (2½ gr) Tube 30s.
- ISUPREL COMPOUND.** (Winthrop Labs) Phenobarbital 6 mg., Isopropylarterenol HCl 2.5 mg., ephedrine sulfate 12 mg., theophylline 45 mg., potassium iodide 150 mg., alcohol 19% / 15 cc. Sol. 16 fl. oz., Gal.
Use: Treatment of bronchial asthma.
- ISUPREL-FRANOL.** (Winthrop Labs) Outer layer of 10 mg Isopropylarterenol HCl, middle layer, and inner core of "Fronol" (8 mg. phenobarbital, 32 mg. benzylephedrine HCl, 130 mg. theophylline) / Tab. Bot. 100s.
Use: Bronchial asthma.
- ISUPREL-FRANOL MILD.** (Winthrop Labs) Outer layer of 5 mg. isopropylarterenol HCl, middle layer, and inner core of "Fronol" (8 mg. phenobarbital, 32 mg. benzylephedrine HCl, 130 mg. theophylline) / Tab. Bot. 100s.
Use: Acute asthma.
- ISUPREL HYDROCHLORIDE, N.N.D. 1962.** (Winthrop Labs) Isopropylarterenol HCl. Amp. (1:5000) 1 cc., Box 5s. Tab. (sublingual) (10 mg. & 1.5 mg) Bot. 50s, 500s.
Use: Bronchial asthma.
- ISUPREL MISTOMETER.** (Winthrop Labs) 200 Inhalations of 125 mcg. of isopropylarterenol HCl, 10 cc.
Use: Bronchial asthma.
- ISUPRENE.** See: Isopropylarterenol
- ITRUMIL, N.N.D. 1962.** (Ciba) Sod. 5-iodo-2-thiouracil. Tab. (50 mg) Bot. 100s, 1000s
Use: Antithyroid action.
- IVOKO, N.N.R. 1952.** (Pitman-Moore) Ivy & oak ext. 1 mg./cc. Pkg. 2 treatments.
Use: Poison Ivy & poison oak
- IVOTOX.** (Bruce Parenterals) Calamine 1%, zirconium carbonate 2%, benzocaine 1%, isopropyl alcohol 8%, methylparaben 0.08%, propylparaben 0.02% w/colloidal magnesium aluminum silicate Lallan, Plastic Bot. 4 oz.
Uses: Treatment of poison oak and poison ivy dermatitis
- IVY-EZE.** (Lloyd, Dabney & Westerfield) Zirconium oxide 5%, phenyltoloxamine dihydrogen citrate 1%, oxyquinoline sulfate 0.1%. Lot, Bot. 4 oz., 1 pt., 1 gal.
Use: Poison ivy, oak and sumac dermatitis therapy.
- IVYOAK.** (Sherman) Sterilin oil sol. of the rhus group, Viol 5 cc.
Use: Prophylaxis & therapy in poison ivy & poison oak dermatitis.
- IVY & OAK EXTRACT.**
See: Birchtox, Amp. (Centrol).
Ivoko, Amp. (Pitman-Moore).
Ivyoak, Viol (Sherman).
Ivyol, Amp. (Merck Sharp & Dohme)
- IVYOL, N.N.R. 1943.** (Merck Sharp & Dohme) Ivy & oak ext. 1:1000 in olive oil. Viol 0.5 cc., 1s, 5s.
Use: Poison ivy & poison oak.
- IXOL.** (Dale) Ichthammol 5%, hydrocortisone 0.5%. Creme 1 oz., 3 oz.
Use: Eczema.
- JACILIN.** (Jamleson Pharms) Pot. penicillin G. 50,000 u./Tab. Bot. 24s.
Use: Orally; penicillin therapy.
- JACOBSON'S SOLUTION.** (Fougero) Benzyl cinnamate 1.6%, ethyl cinnamate 0.5%, benzyl alcohol 2% in olive oil. Amp. 1 cc., Box 12s, 50s. Also w/Vit. A Amp. 1 cc Box 12s.
Use: I.M. (1 cc. daily); congestion, infiltration & inflammatory lesions of middle ear.
- JALOVIS.** See: Hyaluronidase [Various Mfr]
- JAPAN AGAR.** See: Agar [Various Mfr]
- JAPAN GELATIN.** See: Agar [Various Mfr]
- JAPAN ISINGLASS.** See: Agar [Various Mfr]
- JAYRON PLUS INTRINSIC FACTOR.** (Viobin) Specially processed raw liver and duodenum. Pwd. 4 oz. Bot., Tab. 7½ gr. Bot 100s, 500s.
Use: Hematopoietic.
- JECO-B.** (Pitman-Moore) Thiomine HCl 10 mg., riboflavin 5.4 mg., nicotinamide 24 mg., Cal. pantothenate 24 mg., pyridoxine

- HCl 27 mg, liver conc eq to liver, 40 Gm, rice bron conc. 2 cc, aromoliz & diluent, q s / fl oz Bot 1 pt, 1 gal
Use: Vitamin B deficiencies
- JECULIN CAPS.** (Upjohn) Ferrous & ammonium citrate 1 gr, liver conc. 65 gr, red bone marrow ext 1 25 min / Cap Bot 84s, 200s, 500s
Use: (2 3 1 d) Anemias
- JECULIN LIQUID.** (Upjohn) Ferrous & ammonium citrate 8 gr, liver conc 52 gr, red bone marrow ext 10 min, & orange juice / oz Bot 12 oz, 24 oz
Use (3 tsp daily) Hematinic
- JEFBON** (Pitman Moore) Polyterose Iron carbohydrate complex 100 mg / 5 cc or Tab Elixir Bot 8 oz Tabs Bot 60s
Use Treatment of iron deficiency anemia
- JEFRONIC ELIXIR** (Pitman Moore) Cyanocobalamin 25 mcg, thiamine mononitrate 10 mg, riboflavin 0.5 mg, pyridoxine HCl 10 mg, with Jelron (polyterose) alcohol 10% sugar content 25% / 5 ml Bot 6 oz
Use Appetite stimulant and hemolinic
- JENETTES.** (Bristol) Pot penicillin G 10 M.u Jenette (Trache) Pkg 10s
Use: Penicillin sensitive organisms
- JEN-I-SOL** (F E Young) Benzocaine and benzalkonium chloride in lubricant base Tube 1 1/2 oz & Jar 1 lb
- JERIAT.** (Brewer) Iron and amm citrate 6 gr, liver conc 12 gr, vit B 1 10 mg, B 2 10 mg, B 6 1 mg, B 12 6 mcg, niacinamide 20 mg, tricholine citrate 150 mg folic acid 2 mg, pantothenol 4 mg / fl oz Bot 1 pt 1 gal
Use: Treatment of iron deficiency anemias
- JODA-SAL** (Suliff & Case) Sod silicylate 325 mg, Pot iodide 72 mg / Tab Bot 1000s
Use: Analgesic
- JOYCIN No 1 and No 2** (Tracy Drug) Tablets Bot 45s 100s
- JUNIPER TAB OINTMENT.** (Almay Div) Cade oil 4%, Pot stearate, stearam mono glycerol, cetyl alcohol, propylene glycol Oint Tube 1 oz, Jar 4 oz, 1 lb
Use: Topically antipruritic
- JUNIPLEX.** (Chicago Pharm) Vit B 1 2 mg, B 2 4 mg, B-6 4 mg, B 12 30 mcg, niacinamide 10 mg, Cal pantothenate 4 mg, cinchona 129 mg, iron oxide, saccharated 1.55 Gm, Mang glycerophosphate 10 mg, copper gluconate 3.6 mg, nux vomica D 6 mg / oz Bot 8 oz, 1 pt, 1 gal
Use: Vitamin B complex deficiency
- JUNITAB.** (Marcelle) 40% jumper tar (cade oil U.S.P.) in bland water miscible emulsion base Bot 8 fl oz
Use Tar both
- JUNITONIC SYRUP.** (Chicago Pharmacol) Nux vomica D 01 gr, iron oxide saccharated 24 gr, cinchona 0.2 gr, thiamine HCl 225 mg / oz Bot 1 pt, 1 gal
Use Orally, hematinic
- JUVENAL LIQUID.** (Vitamin Corp) Vit A 4000 u, D 400 u, B-1 125 mg, B-2 1 mg, B-6 0.3 mg, niacinamide 10 mg, Vit B-12 3 mcg, iron & ammonium citrate 210 mg, Cal glycerophosphate 10 mg, Mang glycerophosphate 3 mg, cobalt sulfite 0.2 mg / 3 dr Bot 1 pt
Use: Vitamin & iron supplement
- JUVENALS.** (Vitamin Corp) Vit A 4000 u, B 1 1 mg, B 2 1 mg, C 20 mg, D 400 u / Cop Bot 100s, 250s, 1000s
Use: Multiple vitamin therapy
- JUVOCAINE.** Semi Procaine Hydrochloride (Various Mfr.)
- K34** Hexachlorophene
- KADALEX.** (Don Baxter) Potassium chloride 0.2%, dextrose solution 5% Bot 1000 cc, Pkg 6s
Use: IV, treatment of potassium deficiency
- KAERGONA** See: Menadiolone (Various Mfr.)
- KANN ANTIIDIN** (Lederle & Parke, Davis) Kahn precipitation test Viol 10 cc, 25 cc
Use Test for syphilis
- KANN PRECIPITATION TEST.** See: Kahn Antigen (Lederle & Parke, Davis)
- KALACET.** (Donley Evans) Phthylsulfacetamide 0.2 Gm w/Sod carboxymethylcellulose, inorganic salts & glucose / dr Bot 1 pt
Use: Intestinal infections
- KALINIL** (National) Cationic resin, 80% hydrogen, 20% sodium
Use Under study for removal of potassium
- KALIBON.** (National) Pol guaricol sulfonate 1 1/2 gr w/tartar emetic & chloralorm / 5 cc Bot 4 oz, 16 oz, 1 gal
Use Orally, cough preparation
- KAINATAL** (Paul Maney) Bone meal 420 mg, ferrous sulf 200 mg, Vit A 1000 IU, D 500 IU, C 25 mg, B 1 1 mg, B 2 1 mg, niacinamide W/Minerals / Tab Bot 100s, 500s, 1000s
Use Dietary supplement
- KALOX SUSPENSION** (Martin Phillips) Balanced colloidal suspension of calcium carbonate in magnesium and aluminum hydroxide gel / tsp Bot 12 oz
Use: Antacid
- KALPEC** (Wyeth) Kaolin 45 gr, pectin 4 gr in special alumina gel / fl oz Bot 12 fl oz - 1 gal
Use Central of diarrhea
- KALPEN** (Cale) Phenobarbital 1/6 gr, kamatrapine methylbromide 1/24 gr, oil

- minum hydroxide 3 gr., colloidal kaolin 3 gr./Tab. Bot. 100s, 500s, 1000s
 Use: (1 q. 3-4 h) Spasmodic gastro-intestinal disorders.
- KALPIN.** (Walloce) Sod. alginate & Col. alginate complex. Jar 3¼ oz.
 Use: Orally (1 tsp. w/water) Bulk laxative.
- KAMADBOX-JEL.** (Massengill) Aluminum hydroxide gel 5 gr., Mag. trisilicate 10 gr./5 cc. Bot. 1 pt.
 Use: Orally (1-4 dr. p.r.n.) Gastric antacid
- KAMADBOX POWDER.** (Massengill) Aluminum hydroxide gel, dried 25%, Mag. trisilicate 50%, kaolin 25%. Bot. 4 oz., 1 lb.
 Use Orally (1-2 dr. p.r.n.) Gastric antacid.
- KAMADROX TABLETS.** (Massengill) Mag. trisilicate 4 gr., aluminum hydroxide gel, dried 2 gr., colloidal kaolin 2 gr./Tab. Bot. 100s, 500s, 1000s.
 Use: Orally (2 p.r.n.) Antacid.
- KAMELIN.** (Lincoln) Gualacol 100 mg., Iodoform 20 mg., eucalyptol 80 mg., camphor 50 mg./2 cc. Vial 30 cc.
 Use: I.M., expectorant
- *KANAMYCIN SULFATE CAPS., INJ.** U.S.P., U.S.P., N.N.D. 1962. An antibiotic obtained from *Streptomyces kanamyceticus*. See: Kantrax. (Bristol)
- KANONE.** (Massengill) Merodione 2 mg./Fert. Bot. 100s, 500s
 Use: Prevention of hemorrhagic disease
- KANTREX.** N.N.D. 1962. (Bristol) Kanamycin sulf. Cap. (0.5 Gm) Bot. 20s, 100s; Vial (0.5 Gm/2 cc) or (1.0 Gm/3 cc)
 Use: Infections from gram-positive and gram-negative bacteria.
- KANTREX SENSITIVITY DISCS.** (Bristol) 10, or 30 mcg. Bot. 50s.
 Use: Sensitivity, kantrex-susceptible microorganisms.
- KANULASE.** (Smith-Dorsey) Cellulase (Dorsey) 320 u., pepsin 150 mg., glutamic acid HCl 200 mg., pancreatin 500 mg., ox bile extract 100 mg./Tab Bot. 50s.
 Use: Gastro-intestinal.
- KANUMODIC.** (Dorsey) Pentobarbital 8 mg., methscopolamine nitrate 2 mg., pepsin 150 mg., glutamic acid HCl 200 mg., cellulase (Dorsey) 9.14 mg., ox bile ext. 100 mg., pancreatin NF 500 mg./Tab. Bot. 50s.
 Use: Antispasmodic.
- KAOCABB.** (Rorer) Belladonna ext ¼%, bismuth subcarb. 16%, Sod bicarb. 25%, colloidal kaolin 58%/dr. Paw-dispensing container.
 Use: Gastric hyperacidity
- KAO-GALLATE.** (Pitman-Moore) Bismuth subgallate 40 gr., kaolin 80 gr., benzoic acid 9/10 gr. w/mucilage & aromatics/fl. oz. Susp., Bot. 4 fl. oz., 1 pt., 1 gal.
 Use: Gastro-enteritis.
- KAO-LACTOS B.** (Crookes) Kaolin, lactose, thiamine, riboflavin, nicotinic acid & nicotinic acid amide. Pow. 4 oz., 16 oz.
 Use: Orally, intestinal disturbances.
- KAOLIN, N.F.** (Various Mfr.)
 Use: Absorbent for diarrhea.
W/Alumina Gel.
 See: Duterra, Vaginal Lot. (Wyeth)
 Kaomagma, Susp. (Wyeth)
- W/Aluminum hydroxide dried gel, magnesium trisilicate, methyl cellulose, hamatropine methylbromide.** See: Maginal, Tab. (Sutliff & Cose)
- W/Aluminum hydroxide, phenobarbital & adiphenine HCl.** See: Alltrale, Tab. (Malkard)
- W/Bismuth subgallate, Ir. of opium, ginger.** See: Bi-O-Kal, Liq. (Ulmer)
- W/Chlorophyllin, Cap.** (Warren-Teed)
- W/Dihydrostreptomycin and pectin.** See: Dimycin, Susp. (Mourry)
- W/Magnesium trisilicate, aluminum hydroxide.**
 See: Kao-Magnin, Syr. (Sutliff & Cose)
 Kathmojel, Tabs. (Mason)
- W/Methyl cellulose.** See: Kaolase, Susp. (Brewer)
- W/Neomycin sulfate, pectin.** See: Neopectate, Syr. (Tyler)
- W/Opium Pow., pectin, hyoscyamine sulfate, atropine sulfate, hyoscyne HBr and phenobarbital.** See: Danmogel-PG, Susp. (A. H. Robbins)
- W/Pectin, N.F.**
 See: Kaopectate, Liq (Upjohn)
 Kapek, Liq. (Garde)
 Ka-Pak (Am. Phr. Co.)
 Keolin, Lio. (Blue Line)
 Paecin, Liq. (Massengill)
 Pecta-Kalin, Liq. (Paul Maney)
 Pectafold, Liq. (G. F. Harvey)
 Paktamolr, Liq (Warren-Teed).
- W/Phthalylsulfacalimide & pectin** See: Thalin, Susp. (Abbott)
- KAOLIN-BISMUTH PECTIN COMPOUND.** (Pitman-Moore) Kaolin 80 gr., pectin 4 gr., milk of bismuth 77 min., benzoic acid 0.9 gr./oz. Pkg. 4 oz., 12 oz., 1 gal.
 Use: Orally, intestinal disturbances.
- KAULOSE.** (Brewer) Methyl cellulose 1.35 Gm., colloidal kaolin 1.5 Gm./15 cc. Susp. 6 oz., Bot. 1 pt.
 Use: (4 to 8 dr. q. 4 hr.) Diarrhea.
- KAOMAGMA.** (Wyeth) Kaolin in alumine gel. Bot. 12 fl. oz., 6 fl. oz.
 Use: Control of diarrhea.

* Generic name recognized by N.F., U.S.P. or A.M.A.

W/Mineral oil (Wyeth) Kaolin re alumina gel w/mineral oil Bot 12 fl oz

W/Pectin See: Kalpec, Susp (Wyeth)

KAO-MAGNIN. (Sutliff & Case) Aluminum hydroxide 19 Gm, magnesium trisilicate 0.65 Gm, kaolin 4.5 Gm /fl oz Susp 1 pt, 1 gal

Use: Reduces stomach acidity

KAOMIN (Lilly) Bismuth subcarb, Mag hydroxide, mucilage vegetable, kaolin, sucrose & vaellin Pow 4 oz, 1 lb

Use: Orally, intestinal disturbances

KAOMUTH (Elder) delected Kaolin 40 gr, bismuth subcarb 40 gr, pectin 10 gr/oz, 8 oz, 16 oz, 1 gal

Use Orally, intestinal disturbances

KAOMYCIN. (Upjohn) See Kaopectate with Neomycin

KAON ELIXIR (Warren Teed) Pot glucoate equal to 250 mg of elemental Pot /5 cc w/aromatics 1 pt, 1 gal

Use: Potassium deficiency

KAOPATCH (Smith, Muller & Patch) Kaolin 12%, mineral oil 40% w/Irish moss Emulsion 1 pt

Use Orally intestinal disturbance

KAOPECTATE, new name Kaomycin (Upjohn) Kaolin 90 gr, w/pectin 2 gr/oz Bot 6 oz 10 oz 1 gal

Use Orally intestinal disturbances

W/Neomycin new name Kaomycin (Upjohn) Neomycin sulfote 4 1/2 gr kaolin 90 gr, pectin 2 gr/fl oz Bot 6 oz 1 pt

Use Antibacterial action of neomycin

KAOPHYL. (Warren Teed) Chlorophyll ext 60 mg, kaolin 5 1/2 gr /Cap Bot 30s

Use: Orally, deodorize undesirable odors

KAORESIN (National Drug) Polyamine methylene resin 750 mg, sodium aluminum silicate 750 mg magnesium aluminum silicate 187 mg, kaolin 3 Gm pectin 300 mg/oz Susp Bot 4 fl oz

Use: Treatment of diarrhea

KAPHEBEL. (Elder) Phenobarbital 1/2 gr belladonna root 1/4 gr, colloidal kaolin 7 1/2 gr /Tab Bot 100s, 1000s

Use: (1 2 q 4 h) Sedative & antispasmodic

KAPIGAM (Reid) Colloidal Lactose 90 gr, pectin 3 gr, aluminum hydroxide 10 gr cake syrup/5 cc Bot pint

Use: Treatment of diarrhea

KAPIGAM SYRUP. (Reid Labs) Colloidal kaolin 90 gr, pectin 3 gr, aluminum hydroxide 10 gr cake syrup qs /isp Bot 1 pt

Use: Anti-diarrheal

KAPILIN See: Menadione (Various Mfr)

KAPPADIONE, NND 1962 (Lilly) (Menadione Sodium Diphosphate, USP) 2 Methyl 1,4 naphthoquinone diphos

phate tetrasodium (Vitamin K action) Amp (5 mg or 10 mg) 1 cc, PLg 6s, 25s, 100s
Use: Vitamin K therapy

KAPPAXIN. (Winthrop Labs) Pure synthetic menadione 2-methyl-1,4-naphthoquinone Cap 1 Gm in oil, Bot 25s Sol 1 mg/cc Bot 10 cc

Use: Synthetic vitamin K analogue

KAPRYLEX. (Strassenburgh) Resin caprylate 250 mg /Cap Bot 100s

Use: (2-4 qid) Intestinal coelidiosis (mansoniasis)

KARABIM. (Breese) Karaya, kaolin, thiamine Also w/cascara Pow 1/2 oz, 1 oz, 5 lb

Use: Orally, laxative

KARANAL. (Smith Darsey) Pheobarbital 1/4 gr, benzocaine 1/4 gr, Cal carb 3 gr, Sod bicarb 3 gr, papain 1 gr, pan creatin 1 gr, charcoal 5 gr /Tab Bot 100s 1000s

Use: Gastric antacids

KARAYA GUM, A D R (Penick) Indian Gum Can 1/2 lb 1 lb

Use Sterculia gum, NF Used in Tri Cathart (Prof Lab)

W/Apple pectin, Viol B 1 See: Vitapectose w/Karaya (Strassenburgh)

W/Vit B See: Betokar Pow (Eedo)

W/Vit B 1, Kaolin See: Karobim, Pow (Breese)

KARAYA GUM-AROMATIZED. (Mossengill) Can 1/2 lb, 5 lb

Use: Bulk laxative

KAREON. See: Menadione (Various Mfr)

KARIOIUM (Lorvic Corp) Sod fluoride 2.21 mg /Tab Bot 180s, 1000s 1 tab eq 1 mg fluoride Tab Bot 780s 1000s

Use: Dental caries

KARIOIUM LIQ. (Lorvic) Each 8 drops yield 1 mg fluoride ion from sodium fluoride

2 oz plastic bot

Use: Control of dental caries

KARTRYL. (Direct Lab) Bromisovalum 1 gr, carbromat 3 gr thiamine mononitrate 3 gr /Tab Bot 100s, 500s, 1000s

Use: Anxiety and tension

KASAGRA. (Winthrop Labs) Cascara Adxt, nonalcoholic Bot 1 pt

Use: Orally, laxative

KASILAC. (United Lab) Plantago ovata sterculia pectin & lactase Pow 14 oz

Use: Orally, laxative

KATHMAJEL. (Masan) Magnesium trisilicate 0.25 Gm, dried aluminum hydroxide gel D13 Gm, colloidal kaolin D13 Gm /Tab Bot 50s

Use: Treatment of hyperacidity

KATONIUM (Winthrop Labs) Stryanate resin, sulfonic exchange resin, Pow in fl

- dividual pkg. 15 Gm. corl. 21 pkg. Bot. 1 lb., 5 lb.
 Use: Edematous states.
- KATRASUL.** (Chicago) Sulfadiazine 0.167 Gm., sulfamerazine 0.167 Gm., sulfamethazine 0.056 Gm., sulfacetamide 0.111 Gm./Tab. or dr. Tab., Bot. 100s, 1000s. Susp., Bot. 4 oz., 1 pt., 1 gal.
 Use: Sulfonamide therapy.
- KAVITON.** See: Menadiene, U.S.P.
- KAYEXALATE.** (Winthrop Labs) Sod. polystyrene sulfonate. Pkg. 1 lb
 Use: Hyperkalemia as seen in oliguria or anuria due to acute tubular necrosis.
- KAYKLOT.** (Raymer) Menadiene, Tab. 1 mg. Bot. 100s, 500s, 1000s
 Use: Orally, reduce clotting time of blood.
- KAYOUNONE.** (Abbott) See: Menadiene, U.S.P.
- K.B.P.** (Cole) Kaolin 40 gr., bismuth subcarb. 40 gr., pectin 2 gr., glycerin 29 min./oz. Bot. 1 pt
 Use: Orally, intestinal disturbances.
- K-CILLIN.** (Bio-Same) Pot. penicillin G 300,000 u., or 600,000 u./5 cc. in oil. Bot. 2 oz.
 Use: Orally, penicillin therapy.
- K-CILLIN SULFA.** (Maynard) Penicillin G 250,000 U., sulfadiazine 0.167 Gm., sulfamerazine 0.16 Gm., sulfacetamide 0.167 Gm./5 cc. Bot. 60 cc.
 Use: Antibiotic.
- K-C TABLET.** (Paul Maney) Menadiene 10 mg., ascorbic acid 500 mg./Tab., lemon flavored. Bot. 100s, 1000s.
 Use: (1 daily) Anticoagulant in surgical cases.
- KE.** See: Cortisone (Various Mfr.).
- KECTIL SUSPENSION.** (Brital) Dihydrostreptomycin sulfate 50 mg., sulfoguanidine 250 mg., sulfadiazine 250 mg., aminopentamide hydrogen sulfate 0.033 mg., bismuth subcarbonate 250 mg., kaolin 500 mg., pectin 25 mg./5 cc. Sol. 5 fl. oz.
 Use: Relief of acute diarrheal and gastrointestinal episodes.
- KEEP-A-WAKE.** (Stayner) Vit. B-1 5 mg., B-2 2 mg., B-12 activity (cyanocobalamin conc. 1.5 mCg), niacinamide 10 mg., caffeine citrate 2½ gr./Tab. Sol. 24s.
 Use: Ward off drowsiness & fatigue.
- KELATSATE.** (S. J. Tutag & Co) Iron 30 mg., cobalt 0.1 mg., manganese 1.0 mg., zinc 1.25 mg., magnesium 6.0 mg., Vit. B-1 1.5 mg., Vit. B-2 1.2 mg., Vit. B-12 6.0 mcg., niacinamide 10 mg., panthelal 10 mg./5 cc. Bot. pts.
 Use: Iron deficiency anemia and appetite stimulant.
- KILINE.** N.N.R. 1949. (Merck) Ethyl chloride, U.S.P.
- KILOIN.** (Kelco) Algin
- KILLIN.** See: Kheilin (Various Mfr.).
- KIMADSHN.** N.N.O. 1962 (Burroughs Wellcome) Procyclidine HCl (1-cyclohexyl-1-phenyl-3-pyrrolidinopropan-1-ol). Tab. 5 mg., Bot. 100s, 1000s
 Use: (Orally, 2.5 mg. t.i.d.) parkinsonism.
- KIMITHAL.** Thiotharbital. 3-allyl-5-cyclohex-2-enyl-7-thiobarbitic acid
- KINACOST.** N.N.D. 1962. (Squibb) Triamcinolone 1 mg./Tab. Sol. 50s, 500s. 2 mg./Tab. Bot. 50s. 4 mg./Tab. Bot. 30s, 100s. Syr. 4 mg./5 cc. Bot. 4 oz.
 Use: Antiallergic, antirheumatic and anti-inflammatory.
- KINACOST DIACETATE SYRUP.** (Squibb) Triamcinolone diacetate equiv. to 4 mg. triamcinolone; buffered with sodium citrate and sodium phosphate/5 cc. Sol. 120 cc.
 Use: Antirheumatic agent.
- KINAKION.** (Marlett Lane Home of Johns Hopkins Hospital) Vit. K-1 oxide.
 Use: Vit K-induced kernicterus.
- KINALOG.** N.N.D. 1962. (Squibb) Triamcinolone acetonide (16,17-acetonide, 9- α -fluorohydrocortisone-16,17-acetonide). 0.1% cream tubes 3 & 15 Gm. tot. sq. bot. 15 cc. Oint. tube 5 & 15 Gm. Parenteral (10 mg./cc) Vial 5 cc. Spray (50 Gm. or 150 Gm.)
 Use: Topically (10 to 20 times as active as hydrocortisone for inflammation).
- KINALOG-S.** (Squibb) Triamcinolone acetonide 1 mg., neomycin base 2.5 mg., gramicidin 0.25 mg./cc. Bot. 7.5 cc. Cream or Oint. (some formula per Gm) tubes 5 Gm., 15 Gm.
 Use: Anti-inflammatory, anti-bacterial.
- KINALOG IN ORABASE.** (Squibb) Triamcinolone acetonide 1 mg. in arabase vehicle/Gm. Tube 5 Gm.
 Use: Stomatitis.
- KENCEL.** (Kendall) Sod. carboxymethylcellulose 0.5 Gm., ox bile ext. 0.03 Gm./Tab. Bot. 100s.
 Use: Orally, bulk laxative.
- KENDALL'S "COMPOUND A."** See: 11-Dehydrocorticosterone (Various Mfr.).
- KENDALL'S "COMPOUND B."** See: Corticosterone (Various Mfr.).
- KENDALL'S "COMPOUND E."** See: Cortisone (Various Mfr.)
- KENDALL'S "COMPOUND F."** See: 17-Hydroxycorticosterone (Various Mfr.)
- KENDALL'S "DESOXY COMPOUND B."** See: Desoxycorticosterone (Various Mfr.).
- KENHEX.** (Kendall) Cream: Hexachlorophene, hydroxy-quinoline, quaternary ammonium

- camp & sulfacetamide Tube 3 oz with or without applicator
- Vaginal Tabs** Sulfacetamide 50 mg, phenylmercuric acetate 3 mg, 9-amino acridine HCl 2 mg, benzalkonium chloride 2 mg, in a hydrophilic lactose base containing soluble chlorophyllins and buffered with succinic acid/supp Box 24s
Use Vaginal anti-infective
- KENPECTIN SUSP.** (Kenwood) Purified kaolin 90 gr aluminum hydroxide 10 gr, pectin 2 gr/fl oz Bot 8 oz
Use Diarrhea
- KENPECTIN-P SUSP.** (Kenwood) Poregore 1 dram purified kaolin 90 gr, aluminum hydroxide 10 gr pectin 3 gr/fl oz Bot 4 oz
Use: Diarrhea
- KENWOOD THERAPEUTIC LIQUID** (Kenwood) Vit A 30,000 u, D 3000 u, B1 12 mg B2 6 mg, niacinamide 120 mg, Vit C 62 mg, Cal pantothenate 12 mg, Vit C 300 mg, E 30 mg, w/Cal, Phos, Mag Mang & Pot/fl oz Bot 6 & 12 fl oz
Use (1-4 dr daily) Nutritional supplement
- KEOTIN** (Blue Line) Kaolin 80 gr, pectin 8 gr/oz Bot 1 pt, 1 gal
Use Orally, intestinal disturbances
- KEPHALIN** (Wilson) Cephalin, NNR 1949
Use Local hemostatic
- KERALAC.** (Salem Pharm) Chloroxyl, X 0.5% In nail lacquer/Liq Bot ½ oz
Use Treatment of fungus infections of the nails
- KERAMIN INJECTION** (Campbell Pharmaceutical Co) (Steinberg's Solution) Vit A 50,000 USP u/cc, di a tocopherol 0.4%, butyl hydroxyanisole 0.03%, polysorbate EO 1%, chlorbutanol 0.5%, citrate phosphate buffer 0.25%, water for injection q.s. 100%; Amp ½ cc Box 10s Vial 22 cc
Use: Verrucae of all kinds and painful and resistant seed corns
- KERAMOL** (Dumas Wilson) 2% Salicylic acid in an emollient vehicle Jar 1 oz
Use: Topically, antipruritic
- KERID DROPS** (Blair Labs) Triethanolamine polypeptide oleate condensate 10% in propylene glycol with chlorbutanol 0.5% Bot 7 cc
Use: Ear wax remover
- KEROCAINE.** See: Procaïne Hydrochloride (Various Mfr.)
- KERODEX.** (Ayerst)
No 51; Water miscible, Tube 4 oz, Jar 1 & 10 lb
No 55; Water miscible, Jar 1 & 10 lb
No 5E; Water miscible Jar 1 & 10 lb
* Generic name recognized by NF USP or A.M.A.
- No 71; Water repellent, Tube 4 oz, Jar 1 & 10 lb
Use: Skin protective
- KEROHYDRIC W/Sod** dioctyl sulfosuccinate, Sod alkyl polyether sulfonate, Sod sulfacetate, sulfur salicylic acid, hexa chlorophene See: Sebulex, liq (Westwood)
- KESTOS.** (Buffington) Natural estrogenic hormones 2.5 mg, testosterone 25 mg/cc Vial 10 cc
Use 1 M, estrogen & androgen therapy
- KETOCHOL.** (Searle) Oxidized or keto form of all 4 bile acids present in human bile Tab Bot 500s 1000s
Use: Inflammatory conditions of gallbladder
- KETOCHOLANIC ACID,** 90% oxidized cholic acid of natural bile acids See: Dehydrocholic acid
- KETODASE.** (Warner Chilcott) Beef liver beta glucuronidase buffered in acetate to pH 5 Bot 50 & 200 cc (5000 Fishman u/cc)
Use: Determination of steroids in urine & blood
- KETODESTRIN** See: Estrone (Various Mfr.)
- * **KETOHEXAZINE,** 4,6 Diethyl 3(2H) pyridazinone (Lederle)
- KETOHYDROXYESTRADIENE.** See: Estrone
- KETOHYDROXYESTRIN.** See: Estrone (Various Mfr.)
- KETONIL.** (Merck Sharp & Dohme) Protein Hydrolysates A mixture of amino acids and minerals Bot 1 lb
Use: Treatment of phenylketonuria
- KETOSOX.** (Ascher) Ketocholeic acid 3 gr; desoxycholic acid 1 gr/Tab Bot 100s 500s, 1000s
Use Bile acid therapy
- KEYOSTIX** (Ames) Sod Nitraprusside, sod phosphate glycine Slick test for ketones in urine (measures oetaloacetic acid) Bot 90s
Use Urine ketone estimation
- KEYOBARS** (Flint Eaton) Oxidized mixed bile acids 3¼ gr, phenobarbital ½ gr, homatropine methylbromide 1/60 gr/Tab Bot 100s, 500s, 1000s
Use: Antispasmodic & sedative
- KEYADON** (Merrill) Thalidamide N phthalylglutamic acidamide
Use: Potentiated analgesic
- KEWELL SHAMPOO.** (Reed & Carrick) Bot 2, 16 oz
Use: Pediculosis (head and crab lice) only
- KEY-PRED.** (Key Pharmacal) Parenteral, Vial 5 cc
- KEY-PYRONE.** (Key) Dipyrone 0.5 Gm/Tab. Bot 100s; Liq tsp Bot pts; inj cc Amp 10 cc 30 cc
Use: Analgesic—antipyretic
- KHAROPHEN** See: Acetarsone (Abbott)

KHELISEM. (Massengill) Khellin $\frac{3}{8}$ gr. (25 mg./Tab. Bot. 100s

Use: Effect on coronary arteries.

* **KHELLIN.** 5,8-dimethoxy-2-methyl-4',5'-furo-6,7-chromone.

See: Ammivin, Vial, Tab. (National Drug)

Khellsem, Tab. (Massengill).

Khellloyd, Tab. (Lloyd)

Methatrome, Tab. (Physicians Drug)

Visommin, Tab. (Physicians Drug)

Use: Coronary vasodilator.

W/Phenobarbital. See: Khellsem w/Phenobarbital, Tab. (Massengill)

See: Khellloyd w/Phenobarbital, Tab. (Lloyd)

KHELLOYD. (Lloyd) Khellin, Tab. 50 mg. Bot. 50s, 250s, 1000s.

Use: Coronary vasodilator.

W/Phenobarbital. (Lloyd) Khellin 50 mg., phenobarbital $\frac{1}{4}$ gr./Tab. Bot. 50s, 250s, 1000s.

Use: Coronary vasodilator.

KIDNEY FUNCTION AGENTS,

See: Hippuran, Amp. (Mallinckrodt)

Inulin (U. S. Standard Prods.)

Indigo Carmine Sol. (Various Mfr.)

Iodohippurate, Sod.

Monnitol Sol. (Merck Sharp & Dohme)

Medopoque, Amp. (Bell-Craig)

Methylene Blue (Various Mfr.)

Phenolsulfonphthalein (Various Mfr.)

Sodium p-Aminohippurate (Merck Sharp & Dohme).

KI-NI. (Irwin, Neisler) Potassium Iodide 10 gr./Tab. Bot. 50s.

Use: Liquefies secretions and promotes drainage of the upper respiratory tract.

KINATE. Hexahydrotetra hydroxybenzoate salt, quinic acid salt. See: Maxukol, Amp. (Breon)

KINAVOSYL. N.N.D. 1959. (Schanley)

Mephenesin, Elixir 0.1 Gm./cc. Bot. 16 oz., 1 gal. Tab. 0.25 Gm. & 0.50 Gm. Bot. 100s, 1000s.

Use (0.50 Gm. i.i.d.) Muscle relaxant.

KIOPHYLLIN. (Searle) Aminophylline 2½ gr., phenobarbital $\frac{1}{4}$ gr., Pot. iodide 2 gr./Tab. Bot. Tab. E.C., or Plain 100s, 1000s.

Use: Orally (1 i.i.d.) Diuretic w/sedative.

KIPYRONE. (Kirk) Dipyrone 1 Gm./2 cc. Vial 30 cc.

Use: Analgesic in types of rheumatism

KIRKAINE. (Kirk) Procaine 2%, epinephrine 1:25,000. Vial 30 cc.

Use: Local anesthetic.

KIRKAVIN. (Kirk) Casein purified. Vial 1 cc., Box 6s, 12s, 100s. Vial 25 cc., Box 1s.

Use: S.C., nonspecific protein therapy.

KIRKOBEE CAPSULES. (Kirk)

No. 1—Amphetamine sulf. 5 mg., thyroid

* Generic name recognized by N.F., U.S.P. or A.J.A.

1 gr., atropine sulf. 1/360 gr., aloin $\frac{1}{4}$ gr./cap.

No. 2—Amphetamine sulf. 5 mg., thyroid 1 gr., atropine sulf. 1/360 gr./cap.

No. 3—Amphetamine sulf. 5 mg., thyroid 1 gr., phenobarbital $\frac{1}{4}$ gr./cap. Bot. 100s, 1000s.

Use: Timed disintegration, treatment of obesity.

KIRKOBEE TABLETS. (Kirk)

No. 1—Amphetamine sulf. 5 mg., thyroid $\frac{1}{2}$ gr., atropine sulf. 1/360 gr., aloin $\frac{1}{4}$ gr./tab.

No. 2—Amphetamine sulf. 5 mg., thyroid $\frac{1}{2}$ gr., atropine sulf. 1/360 gr./tab.

No. 3—Amphetamine sulf. 5 mg., thyroid $\frac{1}{2}$ gr., phenobarbital $\frac{1}{4}$ gr./tab. Bot. 100s, 1000s.

Use: Timed disintegration, treatment of obesity.

KIRKOLAC. (Kirk) Milk protein defatted susp. Vial 5 cc., Box 6s, 25s, 100s.

Use: S.C., nonspecific protein therapy.

KIRKOPLEX. (Kirk) Vit. B-1 30 mg., riboflavin 0.5 mg., nicotinamide 50 mg., Vit. B-6 2 mg., liver injectable (10 u.) 0.2 cc./cc. Vial 10, 25 cc.

Use: Vitamin B deficiencies.

KIRKSTIOMINE. (Kirk) Neostigmine 1:2000 & 1:1000 sol. Vial 1 cc., 10 cc.

Use: Parasympathomimetic agent.

KITINE. (Superior Pharm) Reserpine 0.1 mg. or 0.25 mg./Tab. Bot. 100s, 500s.

Use: Hypertension.

KIJM. (Borden) Whole milk w/minerals Pow., Can 1 lb., 5 lb.

Use: Infant food.

KIOTONE. (Metro) Sesame oil sol. of Mandiame. Amp. (2 mg. in oil) 1 cc., 10s, 100s (10 mg. in oil) 1 cc., 10s, 100s

Use: Prevention & treatment of hemorrhage.

KIARON LOTION. (Dermik) Salicylic acid 2%, sulfur precipitate 5%, 3'4'6'-trichlorophenol 5%. In a hydro-alcoholic lotion Bot 2 oz. plas. sq.

Use: Seborrhea and related conditions.

K-1-SEA-REM. (Kahlenberg) Sod. bromide 10½ gr., ephedrine HCl $\frac{1}{4}$ gr., methylbromtropin maleate (MAB) $\frac{1}{2}$ gr., n-butyl p-hydroxybenzoate 1 mg./2 dr. Vial 2 dr., Box 4s.

Use: (1 Vial p.r.n.) Travel sickness.

K.M.C. (Ingram) Pot. chloride 1.8 Gm., Mag. chloride 0.5 Gm., Cal. chloride 0.6 Gm., phenol red 0.0001 Gm., water q.s./10 cc. Amp. 10 cc.

Use: L.V., dilute to 1000 cc.

KNL SOLUTION. (Cutter) Pot. chloride 0.26 Gm., Sod. chloride 0.6 Gm., Sod. lactate

only 0.59 Gm /100 cc Soft flasks 500 cc.
 Use: Treatment of severe diarrhea

KOAGAMIN (Chatham) Aq sol of oxalic & malonic acids w/0.25% phenol Vial 10 cc diaphragm stoppered
 Use (IM or IV) To mobilize prothrombin for rapid blood coagulation

KOAGAMIN SUBLINGUAL. (Chatham) Ethyl oxylate 0.0324 Gm, ethyl malonate D0154 Gm ethyl alcohol 35.76%/4 minim unit Box 6 single dose units
 Use Administered to the sublingual veins behind the anterior teeth to accelerate hemostasis

KODOVEN CAPSULES (Flint Eaton) Dover's Pow NF 1/4 gr (equiv to opium 0.025 gr) acetophenetidin 2 gr, Sod citrate 2 gr Pot guaiacol sulfonate 2 gr/Cap Bot 100s, 500s, 1000s
 Use Expectorant & sedative

KOFREDENE. (Abbott) Chloroform, ammonium chloride, ipecac, menthol tartar emetic, honey & cherry syrup Also available w/codeine Bot 1 pt 1 gal
 Use: Orally cough preparation

KOGLUCOID (Parroy) Active principle of Rauwolfia serpentina Tab 50 mg 100 mg, or 250 mg Bot 100s
 Use Hypotensive & tranquilizing agent

KOLAG. (Sulliff & Case) Carcara sagrada 120 gr, lappo 36 gr, taraxacum 48 gr, leptandro 32 gr fringe tree bark 36 gr, Sod phosphate 40 gr, citric acid 8 gr / 8 oz Bot 1 pt 1 gal
 Use (1 tsp p.c. fld) For chronic constipation

KOLANTYL GEL. (Merrell) Bentyll HCl 5 mg, aluminum hydroxide gel, dried 400 mg Mag oxide 200 mg, Sod lauryl sulfate 25 mg methylcellulose 100 mg/1D cc Bot 6s 12 oz

KOLANTYL (TABLETS). (Merrell) Bentyll HCl 5 mg, aluminum hydroxide gel, dried 400 mg, Mag oxide heavy 200 mg, Sod lauryl sulfate 25 mg, methylcellulose 100 mg/Tab Bot 100s, 1000s
 Use: (2 q 3 h) Antispasmodic to control peptic ulcers

KOLANTYL WAFERS (Merrell) Dicyclomine HCl 25 mg, magnesium ox hydrated 100 mg, alum hydrax 240 mg, magn trisilicate 90 mg, sodium lauryl sulf 12.5 mg, methylcellulose 250 mg/Water Bot 32s 100s
 Use: Peptic ulcer therapy

KOLIUM CHLORIDE, ADR (Webster Co) Methyl benzethanium chloride 10%, sod borate 44.75%, sod nitrite 45%, and inert ingredients Ptg 10 Gm
 Use: Powder made for dilution with water

to provide 1:1000 sol of methyl benzethanium chloride

KOLKLOT. (Cole) Menadione Cap (2 mg) Bot 100s
 Use Orally, reduce clotting time of blood

KOLMER ANTIGEN. (Parke, Davis) Vial 5 cc
 Use: Fixation test in the diagnosis of suspected syphilitic sera

KOLOYD. (Lloyd) Aluminum aminoacetate 0.5 Gm, methyl atropine nitrate 1/60 gr / Tab Also available w/phenobarbital 1/4 gr/Tab Bot 100s, 500s, 1000s
 Use (1 p.c. & hs) Antacid with antispasmodic action

KOLPIX (Dane) A—Crude coal tar from coke over 2% w/starch & zinc oxide Oint Tube 1 oz, Jar 2 oz, 4 oz, 1 lb, 5 lb D—Crude coal tar from vertical retort 2%, w/starch & zinc oxide Oint Tube 1 oz, Jar 2 oz, 4 oz, 1 lb 5 lb
 Use: Skin irritations

KOLYTYPE. (Ortho) Polyvalent screening antiserum O111 B4, O55 B5, O26 B6, O127 B8 Vials 2 cc
 Use Diagnostic aid, identification of enteropathogenic E coli

KONAKION, NND 1962 (Roche) Phytonadone synthetic vitamin K₁+2 methyl 3 phytyl 1,4 naphthoquinone Cap 5 mg Bot 50s, Amp 1/2 cc, 1 mg Box 12s, 100s Amp 10 mg 1 cc 25 mg 25 cc
 Use Prevention and treatment of hypoprothrombinemia

KONOREMUL, NNR 1949 (Smith Miller & Patch) Plains Mineral oil w/Irish moss emulsion Bot 1 pt W/Phenolphthalein 147 gr/100 cc Bot 1 pt W/Cascara ext, Bot 14 oz
 Use Laxative

KONDRETTABS. (Smith Miller & Patch) Methylcellulose 7 1/2 gr & Irish moss 38 gr Tab 100s, 500s
 Use Orally, laxative

KONLAX. (Buffington) Dioctyl sod sulfosuccinate 50 mg /Tab specially coated Bot 100s
 Use: Fecal softener

KONOGEN, NND 1958 (Lilly) Estrogenic substances, conjugated Tab 0.625 mg, 1.25 mg, 2.5 mg Bot 100s, 1000s
 Use: Estrogen therapy

KONSYL, NND 1961 (Burton Parsons) Plantago ovata coating Pow, Can 6 oz, 12 oz
 Use: Orally bulk laxative

KORA-12 W/LIVER (Karmel) Vit B 12 30 mcg, folic acid 5 mg, liver inj 5 mcg./cc Vial 10 cc
 Use: IM, anemias

- KORACITRIN.** (Kormel) Citrus bioflavonoid 200 mg./tab. Bot. 100s.
Use: Treatment of increased capillary permeability.
- KORACOL INJ.** (Kormel) Eucalyptol 100 mg., gomenol 100 mg., guaifacel 100 mg., camphor 60 mg., menthol 50 mg., chlorobutanol 30 mg., olive oil q.s./2 cc. Vial 30 cc. Use: Expectorant.
- KORATE.** (Kormel) Phenol 2.75%, iodine & pot. iodide 3.1%, menthol 0.55%, guaifacel 1.5%, in a base of glycerin and alcohol. Bot. 1 oz., 4 oz.
Use: Bactericidal, fungicidal & local anesthetic.
- KORATONE.** (Kormel) Vit. B-12 10 mcg., folic acid 1 mg., B-1 2 mg., B-2 4 mg., ferrous sulfote exsic. 200 mg., ascorbic acid 50 mg., liver desic. 200 mg., stomach mucosa 100 mg./tab. Bot. 100s.
Use: Orally (1 t.i.d.), anemias.
- KORATONE LIQ.** (Kormel) B-12 4 mcg., liver fraction 1 300 mg., B-1 4 mg., B-2 2 mg., B-6 1 mg., calcium pantothenate 2 mg., niacinamide 50 mg., inositol 100 mg., choline hydrochloride 100 mg., l-lysine monohydrochloride 100 mg., bealene HCl 100 mg., Fe 20 mg./30 cc. Bot. 8 oz.
Use: Geriatric.
- KORAVITE.** (Kormel) Vit. A 5000 u., D 1000 u., E 2 mg., B-1 3 mg., B-2 3 mg., B-6 1 mg., C 90 mg., B-12 5 mcg., niacinamide 25 mg., pantothenol 1 mg., folic acid 1 mg., choline dihydrogen citrate 20 mg., inositol 20 mg., methionine 20 mg., w/trace minerals/2 tab. Bot. 100s.
Use: Dietary supplement.
- KORAZYME.** (Kormel) Ox bile ext. 130 mg., pancreatin 130 mg., papain 65 mg., duodenal substances detotted & desic. 65 mg./tab. Bot. 100s.
Use: Choleretic & choleagogue.
- KOREP.** (Kormel) Progesterone 25 mg., estrone 2.5 mg., procaine HCl 0.5% & thimerosal 0.01%/cc. Vial 10 cc.
Use: I.M., estrogen & androgen therapy.
- KORICIDE.** (Kormel) Nitromersol 37 mg., isopropyl alcohol extracts from coal tar 2.22 Gm., quilloja 0.74 Gm., benzoin 2.96 Gm., isopropyl alcohol 80-85%/fl oz. Bot. 1 oz.
Use: Topically, fungus infections.
- KORIUM.** (Korium) Didrozone (dichlorophene) 2%, salicylic acid 3%. Cream 1 oz., 4 oz., 1 lb. Pow. 2 oz.
Use: Topically, fungicide.
- KORIZOL.** (Sutliff & Case) Tyrothricin 0.02%, phenylephrine HCl 0.25%, cetylalkonium chloride 1:2000, menthol 0.03%. Bot. w/dropper 20 cc., 1 pt.
Use: Acute & chronic sinusitis.
- KORMACORAL.** (Kormel) Vit. B-12 4 mcg., stomach mucosa desic. 200 mg./tab. Bot. 90s.
Use: Orally, anemias.
- KOROMEX A.** (Holland Rontas).
Use: SpERMAtocidic agent.
- KOROMEX JELLY.** N.N.R. 1935. (Holland-Rontas) Boric acid 2%, oxyquinoline benzoate 0.02% & phenylmercuric acetate 0.02%, in a suitable jelly base or cream. Cream, Tube 3 oz. with or without measured dose applicator & large economy refill, 5 oz.
Use: Spermicide, contraceptive & vaginal lubricant.
- KORPARIN PLUS LIQ.** (Kormel) Heparin sodium U.S.P. 2500 U.S.P. u., B-12 50 mcg., folic acid 2 mg., choline chloride 100 mg., niacinamide 30 mg./cc. Vial 10 cc.
Use: Heparin-lipotropic.
- KORPAZINE LIQ.** (Kormel) Piperazine citrate equivalent to 100 mg. piperazine hexohydrate/cc. Bot. 4 oz.
Use: Pinworm and roundworm infection.
- KOR-TEP.** (Kormel) Progesterone 25 mg., estrone 6 mg., testosterone 25 mg., procrone HCl 1%, methylparaben 0.135%, propylparaben 0.015%/cc. Vial 10 cc.
Use: I.M., estrogen & androgen therapy.
- KOSATE.** (Maack) Diactyl sod. sulfosuccinate 60 mg., or 100 mg./Cap. Bot. 100s.
Use: Fecal softener.
- KOUGH GREME.** (Central) Syp. pl., Gal.
- K-PHOS.** (Ingram) Pot. phosphate 2.0 Gm., Pot. biphosphate 0.4 Gm., phenol red 0.0001 Gm., water q.s./10 cc. Amp. 10 cc. Use: I.V., diluted to 1000 cc.
- K.P.G. '400'.** (Physicians) Penicillin G potassium buffered 400,000 u./Tab Bot. 100s.
Use: Oral penicillin therapy.
- K-PREDNE DOME.** (Dome Chemicals) Prednisolone 1 and 5 mg., pot. chloride 3 gr./Tab. 1 mg. Bot. 30s, 100s; 5 mg. Bot. 30s.
Use: Corticosteroid therapy.
- KRADON.** (Field Labs.) Ephedrine ethylenediamine 10.6 mg., theobramine sod. acetate 25 mg., caffeine 50 mg., acetophenetidin 160 mg./tab. Bot. 60s.
Use: Pre-menstrual tension
- KREMPLIX.** (Kremers-Urban) Cap.: Thiamine HCl 5 mg., riboflavin 5 mg., niacinamide 50 mg., pyridoxine HCl 5 mg., Col. pantothenate 20 mg., dried brewer's yeast 250 mg./Cap. Bot. 100s, 500s, 1000s. [Inj.] Thiamine HCl 10 mg., riboflavin 0.5 mg., niacinamide 10 mg., pyridoxine HCl 5 mg., Col. pantothenate 5 mg./cc. Vial 30 cc. Tab.: Thiamine HCl 2 mg., riboflavin 2 mg., niacinamide 10 mg., pyridoxine HCl 2 mg., Col. pantothenate 5 mg., dried

- brewers yeast 250 mg / Tab Bot 100s, 500s 1000s
- KREMPLEX-B** (Kremers Urban) Thiamine NCI 100 mg, niacinamide 50 mg, riboflavin 2 mg pyridoxine HCl 2 mg, Cal pantothenate 5 mg / cc Vial 10 cc, 30 cc
Use: Vitamin B complex therapy
- KREO-DIONIN-TERPIN COMPOUND** (Tilden) Ethylmorphine HCl $\frac{1}{4}$ gr, terpin hydrate 2 gr, kreasate (Creasate) 2 min eucalyptal 1 min / fl oz
Use (1 tsp q 3 h) Expectorant & onltussive
- KRES-LUMIN**, (Winthrop Labs) Cal cresol sulfonates 6%, luminal $\frac{1}{16}$ gr / dr Syr, Bot 4 fl oz, 8 fl oz, 1 gal
Use (1 2 tsp) Sedative & expectorant
- KRONOHIST**, (Ferdale) Chlorpheniramine maleate 4 mg pyrilamine maleate 25 mg, phenylephrine HCl 10 mg Vit C 150 mg / Tab Bot 100s
Use Antihistaminic decongestant
- KRYL**, (Ayerst) Terbutaline 4 mg spirin 230 mg, phenacetin 160 mg, phenylephrine HCl 5 mg, Vit C 100 mg / Tab Bot 100s, 1000s
Use Allergic rhinitis
- K-SYRUP** (Walker) Potassium (as lactate citrate and acetate) 75 mEq 8 1 1 mg 8 2 0.5 mg 8 6 0.5 mg niacinamide 35 mg / 5 ml Syr Bot 1 pt
Use Treatment of hypokalemia
- KUDELATE** (Ulmer) Copper undecylenate 6%, undecylenic acid 3% w/dioctyl sod sulfosuccinate Ointment Jar 1 oz, 1 lb
Use Fungicidal ointment
- KUDROX** (Kremers Urban) Susp of aluminum hydroxide gel and magnesium hydroxide Bot 12 oz Also available in double strength
Use Antacid
- KUDROX (DOUBLE STRENGTH) TABLETS** (Kremers Urban) Aluminum hydroxide magnesium carbonate 0.75 Gm / Tab Bot 60s
Use Antacid
- KUSED**, (Kremers Urban) Mephenesin 250 mg, glutamic acid 62.5 mg phenobarbital 7.5 mg, 1 hyoscyamine HBr 0.0625 mg / Cap Bot 100s 500s, 1000s
Use: (1 2 q 3-4 h) Sedative with antispasmodic
- KUTABS**, (Kremers Urban) #2 Amphetamine phos 5 mg, thyroid 30 mg, atropine sulf 0.18 mg
#1 w/cascara sag 30 mg / tab
#3 w/phenobarb 15 mg / tab
Bot 100s
Use: Obesity
- KUTAPRESSIN**, (Kremers Urban) Selective vasoconstrictor from liver Amp 2 cc Vial 10 cc, 20 cc.
Use: Subcutaneous, I.M. & topically; in wide variety of skin disorders from oile gles to pruritus
- KUTRASE** (Kremers Urban) Amylolytic enzyme 30 mg, proteolytic enzyme 6 mg, lipolytic enzyme 75 mg, cellulolytic enzyme 2 mg, phenyltolaxamine citrate 15 mg, 1 hyoscyamine 0.0625 mg / Cap Bot 50s
Use Indigestion
- KUTROL**, (Parke, Davis) Uroenterane, also known as uroanthelone is extract of pregnancy urine which has been found to be effective by oral route in treatment of patients with crateriform peptic ulcer (duodenal and jejunal types) Each Kapsel (No. 4 Cap) contains 75 mg Bot 100s
Use: Treatment of peptic ulcer
- KU-ZYME**, (Kremers Urban) Amylolytic enzyme 30 mg proteolytic enzyme 6 mg, d sorbitol qs / Cap Bot 50s
Use To promote digestion
- KWELL**, NND 1959 (Reed & Cornick) Gamma benzene hexachloride Lot 1% Bot 2 oz Oint 1% Jar 2 oz, 1 lb Sham poo Bot 2 oz, 1 pt
Use: Topically, scabies & pediculosis
- KYLATRATE** (Kyle Labs) Tablets 20 mg Bot 100s
- KYLATUSS** (Kyle Labs) Liquid Bot pl
- KYLECILLIN** (Kyle Labs) Potassium G Penicillin Crystalline, Buffered 200 000 u / Tab Bot 50s
Use Penicillin therapy
- KYNEX**, NND 1962 (Lederle) Sulfamethoxypyridazine 0.5 Gm / Tab Bot 24s, 100s, 1000s
Use long acting sulfanamide therapy
- KYNEX ACETYL PEDIATRIC SUSPENSION**, NND 1962 (Lederle) N Acetyl sulfamethoxypyridazine 250 mg / 5 cc Bot 4 oz & 1 pt
Use Sulfanamide therapy
- KYNEX ACETYL PEDI. DROPS**, NND 1962 (Lederle) Sulfamethoxypyridazine 125 mg, methylparaben 0.8 mg, propylparaben 0.2 mg / cc 10 cc squeeze bottle
Use Sulfanamide sensitive infections
- K-4 LUBRICATING JELLY**, (Johnson & Johnson) Boric acid, glycerin, chandrus, tragacanth Tube 2 oz, 4 1/2 oz.
Use Surgical lubricant
- LAC-BISMO**, NNB 1949 (Hart) Bismuth magma Bot 1 pt, 1 gal
Use Intestinal discomfort
- LACOTEIN**, (Christina) Protein digest 5% w/preservatives Vial 30 cc, 100 cc.
Use: I.M., nonspecific protein therapy

- LACBIL ARTIFICIAL TEARS.** (Allergon) Methylcellulose, polysorbate 80, gelatin, boric acid in isotonic solution w/chlorbutanol 0.5%. Drop bot. 15 cc.
- LACTALBUMIN HYDROLYSATE.** See: Amino-acid, Pow. (Notianol Drug).
- LACTALBUMIN SUSPENSION.** See: Aolan, Amp. (Duke).
- LACTATED POTASSIC SALINE, U.S.P., N.H.R. 1955.** (Don Baxter) Darrow's solution. Pot. chloride 260 mg., Sod. chloride 400 mg., Sod. lactate 590 mg./100 cc. Bot. Vocoliter 150, 500 cc.
Use: Treatment of dehydration & acidosis.
- LACTATED BINGER'S INJ., U.S.P.** See: Hortmann's Soln. (Cutter, Dan Baxter & Rorer)
- LACTAZINE DELTABS.** (M. R. Thompson) Sulfadiazine 83 mg., sulfamerazine 83 mg., sulfamethazine 83 mg., magnesium lactate 324 mg./Tab. Bot. 50s, 250s.
Use: Sulfonamide therapy.
- LACTHAZOL.** (Trifoil) Topical Susp. Bot. 1 oz., pt.
- LACTIC ACID MILK No. 2.** (Mead Johnson) Whole milk acidified with lactic acid, homogenized and spray dried. Cons 1 lb.
Use: Nutrient.
- LACTICATED PEPSIN.** (Merck Sharp & Dohme) Pepsin (N.F.) 16 gr., alcohol 15%/30 cc. Bot. 1 pt., 1 gal.
- LACTIKOL CREAM, N.H.R. 1955.** (Durez) Glyceryl monoricinateate, lactic acid, Sod. lauryl sulfate, stearic acid, glyceryl mono-stearate, glycerin. Tube 1 oz., 2 oz. & 3 oz.
- LACTIKOL JELLY, N.H.R. 1955.** (Durez) Lactic acid, glyceryl monoricinateate, oxy-quinoline sulfate, Sod. lauryl sulfate, karaya, tragaconth, glycerin, butyl p-hydroxybenzoate. Jelly, Tube 2 oz., 3 oz., 4 oz., with or without applicator.
Use: Contraceptive.
- LACTINEX.** (Hynson, Wescott & Dunning) Lactobacillus acidophilus & bulgaricus mixed culture. Tab., Bot. 50s.
Use: Orally (2-4 i.i.d.), to combat intestinal disturbances.
- LACTINEX GRANULES.** (Hynson, Wescott & Dunning) Pkg. 1 Gm., 12i.
Use: Diarrhea.
- LACTOBACILLUS ACIDOPHILUS & BULGARICUS MIXED CULTURE.** See: Lactinex, Tab. (Hynson, Wescott & Dunning)
- LACTOBACILLUS ACIDOPHILUS, VIABLE CULTURE.** See: Bu-Tabs. (Hynson, Wescott & Dunning) Lactinex (Hynson, Wescott & Dunning) Neo-Cuttal, Susp. (Arlington).
- LACTOBACILLUS RULGARICUS.** See: Bu-Tabs. (Hynson, Wescott & Dunning)
- * Generic name recognized by N.F., U.S.P. or A.M.A.
- LACTOFLAVIN.** See: Riboflavin, U.S.P., N.N.B., A.D.R.
- LACTOFLOTT.** (White) Vit. A acetate 3750 u., D 1000 u., R-1 0.75 mg., B-2 1.25 mg., niacinamide 7.5 mg., Vit. B-6 0.75 mg., Cal. pantothenate 7.5 mg., Vit. B-12 2.5 mcg., folic acid 0.25 mg., Vit. C 75 mg., l-lysine 500 mg., iron 7.5 mg., Cal. 130 mg./2.3 Gm. Pow. Bot. 46 Gm.
Use: Vitamin & mineral supplement.
- LACTOGEN.** (Nestle) Vit. A 1000 u., B-1 0.13 mg., C 6 mg., in modified cow's milk w/milk fat, sugar & iron/32 oz. Pow., Pkg. 4 oz., 1 lb.
Use: Infant feeding.
- LACTOGENIC FACTOR.** See: Luteotrophin, Viol. (Squibb)
- LACTOL.** (M. R. Thompson) Sod. lactate 1.35 Gm., Pot., Cal. & Mag. lactates 0.15 Gm./dr. Bot. 16 fl. oz.
Use: To heal on ulcerous condition.
- LACTO-SALICYLATE.** (M. R. Thompson) Sod. salicylate 5 gr., Sod. lactate 1.5 Gm./5 cc. Syr., Bot. 1 pt., 1 gal.
Use: (1 tsp. q. 3 h.) Analgesic & antipyretic.
- * **LACTOSE, U.S.P.** Milk sugar.
W/Dextrin, psyllium seed conc. See: LA Formulo, Liq. (Burton, Parsons)
- LACTOSE, BETA.** See: Beta lactose.
- LACTUM.** (Mead Johnson) Protein 4.0%, fat 5.6%, carbohydrate—from milk 7.0%, dextrimaltose 8.0%, Cal. 0.22%, Phos. 0.2%, iron 0.0002%, water 70%. Can 13 fl. oz.
- LADAFORM-HC.** (Dome) Xylocaine 3%, hydrocortisone alcohol 0.5%/Gm. Bot. Cream ½ oz., lot 1 oz.
Use: Treatment of infection and inflammation.
- LADD'S PASTE.** See: Aluminum Paste, U.S.P.
- LA. FORMULA.** (Burton, Parsons) Lactose 37.5%, dextrin 12.5%, psyllium seed conc. 50%. Bot. 7 oz., 14 oz.
Use: (1 tsp. in water a.c. or p.c.) Laxative.
- LA.J.** (Toblox) Lactic acid, boric acid, amino acetic acid, & methyl & propyl p-hydroxybenzoate, glycerin, & tragaconth. Jelly, Tube 3 oz., with or without applicator.
Use: Contraceptive.
- LAKOBIE.** (Pharm. Dist.) Amphetamine HCl 5 mg., 10 mg. or 15 mg., phenobarbital ½ gr., ¼ gr., w/thyroid ½ gr. In each Cap. Bot. 100s.
Use: (1 a.c.) Obesity.
- LAMO.** (Tailby-Nor-an) Lanolin skin cream. 1 oz., 4 oz. Jar 1 lb.
Use: Topically, skin protective.
- LANABROM ELIXIR.** (Lanett) 60 gr. of

- combined bromides of sod, pot, strontium, ammonium/fl oz Bot 1 pt, 1 gal
- LANABURN OINT.** (Lanett) Aluminum basic acetate, phenol, zinc oxide boric acid, ich thammol, eucalyptol Jar 1 & 5 lb
- LANAFORM** (Doak) Lanolin & veg oil Emulsion 4 oz, 1 lb, 5 lb
Use Topically, skin protective
- LANAMINS** (Lanett) Combination of vitamins (11) minerals (11)/cap Bot 100s 1000s
- LANASCORBIN** (Lanett) Ruth 20 mg ascorbic acid 100 mg /tab Bot 100s, 1000s
Use Capillary permeability
- LANATOSIDE A, LANATOSIDE B A LANATOSIDE C, MIXTURE, NNR 1951** See Digilanid (Sandoz)
- ***LANATOSIDE C, NF NNR 1951, Int. USP, Tab, NF** See Cedilanid (Sandoz)
- LANAURINE**, (Lanett) Antipyrine & benzoic acid in glycerin Bot 4 oz
Use: Otitis media
- LANAVITE** (Lanett) Vitamins A 5000 u, D 1000 u B 1 1.5 mg, B 2 2 mg, B 6 0.1 mg, C 37.5 mg, B 12 0.1 mcg niacinamide 20 mg cal pantothenate 1 mg /cap Bot 500s 1000s
- LANAZINE** (Lanett) Desoxyephedrine HCl racemic 5 mg & 10 mg /tab Bot 500s, 1000s
- LANESTA GEL**, (Brean) 7 Chloro 4 indanol 0.10% sod chloride 10%, Sod lauryl sulfate 0.2% ricinoleic acid 1%, in a glycerin tragacanth base Tube 3 oz w/ app or 3 oz refill
Use Conception control
- LANESTRIN** (Lanett) Estragenic seb natural in aq susp 20000 u/cc Vials 10 cc & 30 cc
- LANIAZID**, (Lanett) Isoniazid 50 mg & 100 mg /tab Bot 500s, 1000s, 5000s
Use: Tuberculosis
- LANNATES ELIXIR** (Lanett) Glycerophos photos of strychnine 1/64 gr, sod 2 gr, cal 2 gr & phosphoric acid in wine/fl oz Bot 1 pt, 1 gal Also with B-125D w/lsp
Use Tonic.
- LANODOXIN** (Lanett) Diadohydroxyquin 210 mg /tab Bot 500s 1000s
Use Amebiasis & *Trichomonas infestans* infections
- LANOKALIN** (Lanett) Phenobarbital 15 mg, homatropine MBr 5 mg, colloidal kaolin 300 mg /Tab Bot 100s
Use Antacid absorbant
- LANOPLEX**, (Lanett) Vitamins B-1 215 mg B 2 2 mg B-6 0.5 mg cal pantothenate 1 mg, niacinamide 20 mg /cap Bot 500s, 1000s, 5000s
- LANOPLEX OERITRIC**, (Lanett) Vitamins B 1 20 mg, B 2 3 mg, B 6 1 mg, B 12 30 mcg, niacinamide 100 mg, cal, pantothenate 3 mg, choline chloride 20 mg, inositol 10 mg, liver 2 mcg /cc Vials 10 cc
- LANOPLEX INJECTION**, (Lanett) Vitamins B 1 100 mg, B 2 1 mg, B 6 2 mg, niacinamide 50 mg, cal pantothenate 10 mg /cc Vials 10 cc & 30 cc
- LANOSTABIL** See Digilanid (Sandoz)
- LANOTAL PILLS** (Lanett) Phenolphthalein 1/2 gr, aloin 1/2 gr, ipecac 1/15 gr, bella donna ext 1/12 gr /pill Bot 1000s, 5000s
- LANOXIN, NND 1962** (Burroughs Well come) Digoxin Tab, 0.25 mg, Bot 100s, 500s, 1000s; 0.5 mg, Bot 100s, 1000s Amp (0.25 mg /cc) 2 cc Pediatric Elixir (0.05 mg /cc) Bot 2 fl oz
- LANTENE JELLY, NNR 1955** (Lanett) Ricinoleic acid, hexylresorcinol, Sod borate, chlorothymol, tragacanth, starch, hydrochloric acid & Cal hydroxide Tube 1 1/2 oz - 3 oz, with or without applicator
Use: Contraceptive
- LANTRISUL**, (Lanett) Sulfamerazine, sulfadiazine, sulfamethoxazole 2 1/2 gr each /tab Bot 500s, 1000s
Use Triple sulfa therapy
- LANUM, ADR 1955** See Lanolin (Various Mfr)
- LANZETS**, (Lanett) Tyrothricin 1 mg, benzocaine 5 mg /oz Bot 500s, 1000s
- LAPACTIC PILLS**, (Merck Sharp & Dohme) Aloin 15 mg, specac 4 mg, belladonna ext 5 mg /pill Bot 100s
Use: Orally, laxative
- LARGACTIL** See Tharazine, Amp, Tab (Smith Kline & French)
- LARGON** (Wyeth) Propriazone HCl Amps (20 mg /cc) 1 cc, 2 cc Pkg 25s
Use Tranquilizing agent
- LAROCAINE HYDROCHLORIDE, NNR 1946** (Roche Lab) deleted p Aminobenzoic acid ester of 2,2 dimethyl 3 diethylamino propanol HCl
Use: Anesthetic
- LARODON**, (Haffmann La Roche) 4 Isopropyl antipyrine
Use Analgesic
- LARYIGAN** (Doho) Val oils, antipyrine pyrimidine molecule, methyl rasaniline, glyc erine alcohol Atomizer bot 1 oz & 1 pt
Use: Throat & nasal spray
- LASSAR'S PASTE** See Zinc Oxide Paste USP (Various Mfr)
- LASTOMONE**, (Key) Estradiol valerate 4 mg, testosterone enanthate 90 mg, w/0.5% chlorobutanol in sesame oil/cc Vial 5 cc
Use Estrogen testosterone therapy

* Generic name recognized by NF USP or A.M.A.

- LASTROGEN.** (Key) Estradiol valerate 10 mg/cc. in sesame oil w/0.5% chlorobutanol. Parenteral, Vial 5 cc.
Use: Estrogen therapy.
- LATEX-TRICHINA REAGENT.** (Tyland) Test reagent combines polystyrene latex and an extract of *Trichinella spiralis* and has been processed to give rapid, color-cut reactions w/serum specimens containing antibodies to trichina./12-test kit containing 0.5 cc. reagent & 28 capillary pipettes.
Use: Diagnostic reagent.
- LATHANOL LAL, A.D.R.** (National Aniline Corp.) Sod. alkyl sulfacetate. Sod. lauryl sulfacetate, Pow.
Use: Detergent and foaming agent
- LATRODOL.** (Toilby-Nason) Mephensin 200 mg., nicotinic acid 25 mg., belladonna ext 5 mg./Tab. or dr. Bot. 100s, 300s. Liq. 1 pt., 1 gal. Also w/phenobarb. ¼ gr. Tab. Bot. 100s.
Use: [1-2 i.l.d.] Rheumatic & arthritic conditions.
- LAUDEXIUM METHYLSULFATE.** Decamethylene- α , ω -bis-[1-13',4'-dimethoxybenzyl]-1,2,3,4-tetrahydro-6,7-dimethoxy-2-methylisouquinolinium methosulphate. See: Laudolisin Methylsulfate (Fougera).
- LAUDOLISSIN.** (Fougera) Laudexium methylsulfate. A synthetic muscle relaxant chemically closely related to d tubocurarine. Vial (6 mg/cc.) 10 cc.
Use: I.V. muscle relaxant.
- LAUGHLEN REAGENT.** (Lederle) Diagnosis of syphilis. Vial 3 cc. (120 tests)
- LAURIUM.** (Whittier) Ferrous gluconate 300 mg., liver conc. 200 mg., folic acid 1 mg., thiamine HCl 2 mg., riboflavin 1 mg., niacinamide 10 mg., ascorbic acid 15 mg./Cap. Bot. 100s.
Use: [1-2 daily] Hypochromic & nutritional anemias.
- LAURO-RORACIC EYE BATH IMPROVED.** (Clapp) Borax, berberin HCl, boric acid, sodium chloride in camphar. Bot. R or w/cup, 8 oz w/out cup, ½ oz. Drop Bot. Use: Optic.
- LAURON, N.N.D. 1962.** (Endo) Aurethiagliconide, ourathiagliocolanilid susp in oil. Vial (50 mg./cc.) 5 cc., 10 cc.; (150 mg./cc.) 10 cc.
Use: Gold therapy & antiarthritic.
- LAURYL SULFOACETATE.** See: lowto, Cahn & Liq. (Wertward).
- LAVABO.** (Superior Pharm) 1-Amphetamine sulfate 5 mg./Tab Bot. 100r, 500s.
Use: Treatment of obesity.
- LAVATAR.** (Doak) Coal tar dirtifate 33.3% in a shampoo base. Liq. Bot. 4 oz., 1 pt., 1 gal.
Use: Treatment of dermatoses, and psoriasis of the scalp
- LAVEMA POW.** (Winthrop Labs) Oxyphe-nisatin 3 Gm., edothami Sod. 3 Gm./Pkg of Pow. Box 100r.
Use: Dissolve Pkg. in 1-2 qts. of water for enema.
- LAVEMA COMPOUND DISPOSABLE EN-EMA KIT.** (Winthrop) Oxyphe-nisatin (Lav-ema) 20 mg., tyloxopol (oxyethylated ter-troty octylphenolpolymethylene polymer; Supermono) 135 mg., w/methylparaben 0.005%, propylparaben 0.015%, in propy-lene glycol 9 cc., water q s /180 cc.
Use: Facal impaction.
- LAXINATE.** (Mottard) Diethyl sod. sulfosuc-cinate 60 mg./Tab. Bot. 100s, 1000s
Use: Facal softener
- LAYOR CARANG.** See: Agor (Various Mfr.).
- LAZAGEL.** (Squibb) Zinc oxide 20% in plastibase. Oint., Tube 1 oz. Jar 1 lb.
Use: Topically, protective.
- L.C.D.** See: Coal Tar Soln., U.S.P.
- L.C.D. COMPOUND OINTMENT.** (Almay) Coal tar sol. 5.83% in lanolin, menthol & vanishing cream base. Tube 1 oz.; Jar 1 lb., 5 lb.
Use: Topically, in dermatitides.
- L-CILLIN AQUEOUS 250 & TABS.** (Leyden) Crystalline penicillin G potassium 200,000 units/Tab 250,000 units/5 cc. Tab. Bot. 100s. Liq. Bot. 60 cc.
Use: Penicillin-susceptible infections
- L-CILLIN AQUEOUS 250 with THREE SULFONAMIDES.** (Leyden) Crystalline penicillin G potassium 250,000 units; three sul-fonamides 0.5 Gm., plus buffer/5 cc. Bot. 60 cc.
Use: Antibiotic
- LE 80/40.** (Ross Labs) Modified milk for infant feeding Can 1 lb.
- LEAD OLEATE OINTMENT, N.F. VII** (Ulmer) Diachylon Oint. Jar 1 lb
- LEBRUN'S.** (Fougera) Eucalyptol 0.12 Gm., carbolic acid 0.02 Gm., chlorophyll 0.1 Gm., sulfuric ether 0.05 Gm., galatol 0.05 Gm., camphor 0.05 Gm., olive oil. Amp. 1 cc., Box 12s.
Use: I.M., expectorant
- LECITHIN.**
Cavendish—Tab. (½ gr.) Bot. 100s, 500s, 1000s.
Dumas-Wilson—Granul. Bot. 1 pt.
See: Aclatin, Cap. (Arroc. Concentrator)
Colastin, Pow (Lawman)
W/Dextrose anhydrous, cottonseed oil, oxy-nithylenoxypropylene polymer. See: Lipa-mul L.V. (Upjohn).
W/Thyroid, adrenal substance & spermin.
See: Lotans, Tab. (Warterheld).

LECITHIN COMPLEX CAPSULES

414

W/Vitamins See Aletin, Cap [Associated Concentrates]

LECITHIN COMPLEX CAPSULES. (Victor Lemon Co) Soybean lecithin 8 gr with natural choline inositol, sterole tocopherols, unsatd fatty acids/cap Bot 100s
Use: (1 or 2 pc) Hypercholesterolemia

LECITHIN COMPLEX PLUS. (Victor Lemon Co) Soybean phosphatides including choline inositol, and cephalin; protein hydrolysates and citrus bioflavonoids Bot 8 oz paw
Use: (1 tsp p c t i d) Hypercholesterolemia

LECITHIN-MULTIPLY PERLES (Victor Lemon Co) Soybean lecithin 8 gr, vit A 3000 u D 180 u, B 1 0.6 mg B 2 1.2 mg, B 6 1.8 mg, cal pantothenate 12 mg/perles Bot 100s
Use: (1 t i d) Hypercholesterolemia

LECITHIN, SOYA OIL-FREE "RG." (Glidden Co) Gran 8 oz pkg
Use: (1 tsp equals 7.5 Gm, 1 3 tsp t i d) deranged lipid & cholesterol metabolism

LECODRINE (Lemman) Dihydrocodeinone bitartrate 30 mg, chlorphenamine 6 mg Phenylpropylamine HCl 60 mg, sad citrate 1 Gm/A oz Bot 3 oz 1 pt, gal
Use: Upper respiratory congestion

LECTAL (M R Thompson) Syrup containing sad pot cal, and mag for inducement of electrolyte balance Bot pl gal

LEDERCILLIN, ADR 1955 (Lederle)
Oint.—Penicillin G procaine 1000 u/Gm Tube 1 oz Also 10 000 u/Gm
Tab.—Penicillin G procaine 50 000 u & 100 000 u Bot 100s 1000s 250 000 u Bot 12s
Traches.—Penicillin G procaine 5000 u Bot 25s, 250s

LEDERM LOTION (Scott Lee) 4 oz

LEDERPLEX CAPSULES (Lederle) Vit B 1 2 mg B 2 2 mg, B 6 0.2 mg, B 12 1 mcg niacinamide 10 mg Cal pantothenate 3 mg, liver 414 mg choline 20 mg inositol 10 mg folic acid 2 mg/Cap Bot 100s, 250s 500s
Use: (1 t i d) Anemias

LEDERPLEX LIQUID (Lederle) Vit B 1 2 mg B 2 2 mg niacinamide 10 mg pantothenic acid 2 mg Vit B 6 0.2 mg B 12 5 mcg liver 470 mg, folic acid 0.2 mg inositol 10 mg, choline 20 mg/dr Bot 4 oz, 8 oz 12 oz, 1 gal
Use: Hematinic w/vitamin B complex

LEDERPLEX PARENTERAL (Lederle) Vit B 1 10 mg, B 2 0.5 mg, niacinamide 10 mg, pantothenic acid 5 mg, Vit B 6 1 mg, B-12 5 mcg/cc Vial 10 cc
Use: Vitamin B complex

LEDERPLEX TABLETS. (Lederle) Vit B-1 2 mg, B 2 2 mg, B-6 0.1 mg, B 12 1 mcg, niacinamide 10 mg, Cal pantothenate 3 mg, folic acid 0.2 mg, liver 250 mg/Tab Bot 50s, 100s, 250s, 500s
Use: (1 t i d) Anemias

LEDINAC. (Lederle) Thiamine 1 mg, riboflavin 1.5 mg, niacinamide 3.5 mg, pantothenic acid 1.25 mg, pyridoxine HCl 0.15 mg, biotin 9 mcg, inositol 23 mg, choline 60 mg, Vit B 12 30 mcg, folic acid 1.5 mg, iron Cal, Phos/30 Gm Jar 1/2 lb, Can 5 lb
Use: Orally, vitamin supplement

LEMASCORB. (Ascher) Ascorbic acid 100 mg/Tab Bot 100s, 500s, 1000s
Use: Vitamin C deficiency

LEMBROSE (Wyeth) Methacholine chloride 0.25% menthol 5.5%, thymol 0.1%, eucalyptol 1%, methyl salicylate 11%. Tube 1 oz, Jar 1 lb
Use: To produce erythema by local dilation of arterioles & capillaries

LEMISERP. (Lemman) Reserpine 0.1 mg, 0.25 mg/Tab Bot 100s Sal (0.25 mg/5 cc) Bot 3 oz, 1 pt, 1 gal
Use: Treatment of hypertension

LEMISERP TY-MED (Lemman) Reserpine 0.5 mg/Tab Bot 100s
Use: Hypertension

LENETRAN TABS. (Lakeside) Mephenoqualone 400 mg/Tab Bot 50s, 500s
Use: Tranquilizer

LENIC (Crookes Barnes) Linoleic and Linolenic acids 500 mg, oleic acid 175 mg; Arochudonic, Pentaenoic and Hexaenoic Acids—100 mg With saturated fatty acids 100 mg, mixed tocopherols 1 mg
Use: Decrease and control of blood cholesterol level

LENIC HP (High Potency) (Crookes Barnes) Tetraenoic pentaenoic, hexaenoic 350 mg linoleic 300 mg/Cap Bot 100s
Use: With low fat diet to decrease and control blood cholesterol levels
Use: Supplement

LENIC W/NIACIN (Crookes Barnes) Tetraenoic pentaenoic, hexaenoic 90 mg, linoleic acid 450 mg niacin 100 mg/Cap Bot 50s
Use: With low fat diet to decrease and control blood cholesterol levels

LENICET See: Aluminum Acetate, Basic [Various Mfr]

LENTE ILETIN, NND 1962 (Lilly) Zinc Insulin Vial (40 u/cc) U-40 10 cc, (80 u/cc) U 80, 10 cc
Use: Insulin therapy

- *LENTE INSULIN, N.D. 1962. Susp. of zinc insulin cryst. & amorphous insulin.
See: lente insulin (Lilly).
- LENTIN, (Merck) Carbachol. Vial (1:1000) 20 cc. (1:10,000) 10 cc.
- LENTOPEN, N.D. 1962. (Wyeth) Penicillin G procaine 300,000 u./cc. in peanut oil w/2% aluminum monostearate. Tubex w/sterile needle. Vial 10 cc.
Use: Prolonged penicillin therapy.
- LENTOPEN PECTINATED. (Wyeth) Penicillin G procaine 300,000 u. w/penicillin G Pot. 100,000 u. in oil w/aluminum stearate/cc. Tubex w/needle 1 cc., Vial 10 cc.
- LENTOVET. (Wyeth) Penicillin G procaine 300,000 u. in oil/cc. Tubex w/needle, Vial 10 cc.
- LENTOVET PECTINATED. (Wyeth) Penicillin G procaine 300,000 u., penicillin G Pot. 100,000 u. in oil w/aluminum stearate/cc. Tubex w/needle 1 cc., Vial 10 cc.
- LENTOVET 600 SUSPENSION. (Wyeth) Penicillin G procaine 600,000 u./cc. Aq. inj., 1 Tubex w/needle.
- LEOTIC EAR DROPS, (Scott-Lee) 5 cc.
- LEPAZOL. See: Pentylentetrazol, U.S.P.
- LEPIFRIN OPHTH. SOL. (Allergan) Levoprenaline 2% and benzalkonium chloride 1:10,000 as the preservative./Sol. Bot. 15 cc. plastic dropper.
Use: Topical sympathomimetic amine.
- LEPROSY THERAPY. See: Hansen's Disease.
- LERGINE. See: Tricyclamol Chloride
- LERITINE HCl AND PHOSPHATE, N.D. 1962. (Merck Sharp & Dohme) Anileridine Ethyl 1-(4-aminophenethyl)-4-phenylisopiperolate. Tab. (25 mg) Bot. 100s, 500s; Inj. (25 mg./cc) Amp. 1 cc., 2 cc., Box 25s, Vial 30 cc.
Use: Analgesic.
- LERJA. (Burns) Phenitamine maleate 25 mg., pyrilamine maleate 25 mg., phenylpropanolamine HCl 50 mg./tsp; Phenitamine maleate 6.25 mg., pyrilamine maleate 6.25 mg., phenylpropanolamine HCl 12.50 mg./mg., phenylpropanolamine HCl 12.50 mg./mg. Bot. 100s, Bot. pt.
Use: Hay fever, asthma, sinusitis.
- LERJA SYR. (Burns) Dexamethorphan HBr 10 mg., phenitamine maleate 6.25 mg., pyrilamine maleate 6.25 mg., phenylpropanolamine HCl 12.5 mg., ammonium chlorate 60 mg., sodium citrate 120 mg., chloroform 10 mg., menthol 0.75 mg./tsp. Bot. ph. Use: Cough due to common cold
- LES-CAV. (Crookes-Barnes) Fluoride ion 1 mg (from sodium fluoride)/0.6 cc. 60 cc. disp bottle
Use: Reduction in dental caries.
- LESCOPINE BROMIDE, N.D. 1962. (Lincoln) Methscopolamine bromide 25 mg./Tab. Bot. 100s
Use: Antispasmodic & anticholinergic.
- LES-I-THIN. (National Lecithin) Linoleic and linolenic acids. Bot. 6 oz., 12 oz.
Use: Anticholesteral.
- LISOFAC. (Wyeth) Casein & lactalbumin, hydrolyzed rice flour, sucrose, dibasic Col. phosphate, Pot. carb., Mag. oxide, thiamine HCl, riboflavin, niacinamide & vanillin. Can 1 lb.
Use: Dietary supplement.
- LESTEMP. (Zeid Labs) N acetyl-p-aminophenol 120 mg, alcohol 10%/5 cc.
Use: Analgesic and antipyretic.
- LE-TEST. (Hyland Labs) Latex-nucleoprotein reagent 2 cc., positive control serum 0.5 cc., negative control serum 0.5 cc., 60 capillary pipettes & 2 ceramic ring glass slides./Kits with enough material for 50 screening or control tests.
Use: Detecting anti-nuclear factors.
- LETHDRONE, (Burroughs Wellcome) Nalorphine, n-allyl normorphine hydrobromide. Amp. (5 mg./cc.) 2 cc.
Use: Antidote for morphine-type drugs.
- LEUCOVORIN CALCIUM, N.D. 1962. (Lederle) Folinic acid S.F. Formyl tetrahydropteroylglutamic acid, a derivative of folic acid as calcium salt. Amp (3 mg./cc.) 1 cc.
Use: Antagonist of aminopterin & other folic-acid antagonists.
- LEUKEMIA AGENTS.
See: Aminopterin, Tob. (Lederle)
Cytosan (Mead Johnson)
Leukeran (Burroughs Wellcome)
6-Mercaptopurine (Burroughs Wellcome)
Methotrexate (Lederle)
Mustargen HCl (Merck Sharp & Dohme)
Myleran. (Burroughs Wellcome)
Purinethal, Tab. (Burroughs Wellcome)
Teroplerin, Amp. (Lederle)
Thio-Tepa, Pow. (Lederle)
Triethylene Melamine (Lederle)
Urethane, Pulvules. (Lilly)
- LEUKERAN, N.D. 1962 (Burroughs Wellcome) Chlorambucil, C.B. 1348. p-(Di-2-chlorethyl) aminophenylbutyric acid. Tab. 2 mg, Bot. 50s
Use: Orally. [D.1 to D.2 mg./Kg. body weight daily for 3 to 6 weeks] for the treatment of chronic lymphocytic leukemia and malignant lymphomas
- LEUKORAY. (Graft) Dinadobydroxyquinoline, sorbic acid, a quaternary ammonium cpd, sulfadiazine, stilbestrol, detergent and wetting agents, buffered to pH 4. Vaginal Tabs. 20s incl. appl.
Use: Trichomonas.

LEVALLOPHAN, 1-3 Hydroxy N allyl morphinan See: Lofan, Vial (Roche)

***LEVALLOPHAN TARTRATE**, USP [N] USP

See: Lofan Pediatric Ampules (Roche)

LEVAMINE (Meyer) Butabarbital 15 mg, 1 hyoscyamine 0.15 mg / Tabs butabarbital 45 mg, 1 hyoscyamine 4.5 mg / Cap Bot 100s

Use: Nervous indigestion

***LEVAMPHETAMINE SUCCINATE** (Isomer) 1 phenyl 2 aminopropene succinate Amphetamine, levo 1- α methylphenethylamine succinate See: Cydril (Tutag)

***LEVAMPHETAMINE** levo Amphetamine Levamfetamine

See: Ad Nil Tab (Medics)

Amphetamine Tab (Van Pelt Brown)

Lababa Tab (Superior)

Levanor Tab (Nordmark)

***LEVARTERENOL BITARTRATE**, USP, NNO 1962, ADR levo Norepinephrine levo norepinephrine, 3,4 dihydroxyphenyl 2 aminoethanol [N], USP See: levophed Sol (Winthrop Labs)

LEVAHIL, NNO 1962 (Upjohn) Ecliyurea 300 mg / Tab Bot 50s, 500s

Use: Apprehension and tension

LEVETAMINE (Abbott) 1 desoxyephedrine

LEVO-AMPHETAMINE (Isomer) alpha 2 phenylaminopropene succinate See: Lev amphetamine

LEVO-DROMORAN TARTRATE, HND 1962 (Roche) Levorphanol tartrate, HND 1959 Levorphan Tartrate NHR 1954; levo

3 hydroxy α methylmorphinan Amp (2 mg / cc) 1 cc Box 12s, 100s Vial (2 mg / cc) 10 cc Tab (2 mg) Bot 25s, 100s, 1000s

Use: Narcotic, analgesic

***LEVO-EPINEPHRINE BITARTRATE**. See: Lyophrin (Alcan)

LEVO-GLUTAVITE. See: 1 Glutavite, Pow (Gray Pharm)

LEVOMEPROMAZINE () 10-(3 dimethyl amino 2 methylpropyl) 2 methoxypheno thiazine

LEVONOS (Nordmark) levo Amphetamine 1 Phenyl 2 aminopropene alginate 5 mg / Tab Bot 100s

Use: Appetite control

***LEVONOBDEFRIN**, NF 1-1 (3,4 Dihydroxyphenyl)-2 aminopropenal See: Cobe frin (Winthrop Labs)

LEVOPHED, NND 1962 ADR (Winthrop Labs) Levarterenol bitartrate, levoarterenol 1 nor epinephrine 1 (3,4 Dihydroxyphenyl) 2 aminoethanol Sol 0.2%, Amp 4 cc Bot 10s Sol 0.02%, Amp 2 cc., Box 5s

*Generic name recognized by HF USP or AMA

Use: To maintain blood pressure in acute hypotensive states.

LEVOPHENACYLMORPHINAN. See: ()-3-hydroxy N phenacylmorphinan

LEVO-4-PHENYL-4-(CHLOROBENZYL)-5-OXOHENANOIC ACID. See: Caprochlorone (Merck Sharp & Dahme)

LEVOPROPOXYPHENE. (Lilly) See: Navrad

***LEVOPROPOXYPHENE NAPSYLATE**. (Eli Lilly Co) a 1,4 dimethylamino 1,2 diphenyl 3 methyl 2 butanol propionate 2 naphthalene sulfonate hydrate See: Navrad Napsylate (Lilly)

LEVOBENINE. See: Epinephrine (Various Mfr)

LEVOBPHAN TARTRATE, HND 1962 See: Levorphanol Tartrate, NND 1961

***LEVOBPHANOL TARTRATE**, HF, NNO 1962 [N] Tabs NF See: Levo Oramoran Tartrate, Amp, Tab (Roche)

LEVOTHYROXINE SODIUM, HND 1962 See: Synthroid Sodium (Travenol)

***LEVOXADROL HYDROCHLORIDE**. (Cutter) 1,2 (2,2 diphenyl 1,3 dioxolan-4 yl)piperidine hydrochloride

Use: Antidepressant

LEVSIN SULFATE. (Kremers Urban) 1 Hyoscyamine sulfate Tab: 0.125 mg & 0.25 mg Bot 100s, 500s, 1000s; Elixirs: 0.125 mg / Tsp Bot 1 pt, 1 gal; Injectable: 0.25 mg / cc Amp 1 cc 25s, Vial 10 cc and 30 cc

Use: Anticholinergic

W/Secobarbital (Kremers Urban) 1 Hyoscyamine sulfate 0.25 mg, secobarbital 15 mg / Tab Bot 100s, 500s, 1000s

Use: Anticholinergic with sedative

LEVSIN-C. (Kremers Urban) Tab Levsin sulfate 0.25 mg and sodium ascorbate 50 mg Bot 100s, 500s, 1000s Bulk, 5000s

Use: Peptic ulcer

LEVSIN PB (Kremers Urban) Levsin sulf and phenobarbital Tab 0.125 mg & 15 mg Elixir Levsin sulf 0.125 mg and phenobarb 15 mg / Tsp Bot 1 pt, 1 gal; Injectable Levsin sulf 0.25 mg and phenobarbital 15 mg 1 cc, Amp 1 cc 25s, Vial 10 cc and 30 cc

Use: Anticholinergic and sedative

LEVSIN-SB (Kremers Urban) Tab Levsin sulfate 0.25 mg and secobarbital 15 mg Bot 100s, 500s, 1000s

Use: Anticholinergic and sedative

LEVSINEX. (Kremers Urban) 1 Hyoscyamine sulf 0.375 mg / long action Tab Bot 100s

Use: Anticholinergic

LEVSINEX/PHENOBARBITAL. (Kremers Ur-

LIBETOPLEX-B (New Mexico Pharm) Liver injection (equiv to B 12 10 mcg) 100 mg, ferric citrate 10 mg, thiamin HCl 10 mg, riboflavin 0.5 mg, niacinamide 50 mg, calcium pantothenate 1 mg, pyridoxine HCl 0.5 mg/cc Vial 30 cc

LIBEXRON-F, (C. D. Smith) Ferrous sulfate, 194 mg, ascorbic acid 25 mg, riboflavin 2 mg, thiamine HCl 2 mg, niacinamide 10 mg, pyridoxine HCl 1 mg, Calcium pantothenate 1 mg, folic acid 2 mg, liver conc 250 mg, Vit. B 12 2 mcg/Tab Bot 100s 1000s

Use B complex & Iron therapy in secondary anemia

LIB TABLETS, (Warren-Teed) Ferrous gluconate 2 gr, liver, whole, 4 gr, Vit B 1 3 mg B 2 1 mg, niacinamide 5 mg/Tab Bot 100s

Use (2 3 t i d) Anemias

LIBRAX (Roche) (Quarzan) Clidinium bromide 25 mg, chloridiazepoxide 5 mg (librium)/Cop Bot 50s 500s

Use Gastrointestinal disorders

LIBRIUM (Roche) Chloridiazepoxide 7 Chloro-2 methylamino 5 phenyl 3H 1 4 benzodiazepine 4 oxide HCl Methaminodiazepoxide 5 mg or 10 mg 25 mg/Cop Bot 50s 500s

Use Relief of anxiety, apprehension and nervous tension

LIBRIUM INJECTABLE (Roche) Librium Amp (dry filled) 100 mg Special IM Diluent 2 cc for IM 1.5% benzyl alcohol, 4% polysorbate 80 20% propylene glycol 5 cc, Box 6s 25s

Use Relief of anxiety and tension

LICURON-B, (Lokeside) Liver fraction #2 195 mg, ferrous sulfate, dried 84 mg, copper sulfate 2.5 mg, thiamine HCl 0.33 mg, riboflavin 0.67 mg, niacinamide 3.33 mg/Tab Bot 100s, 500s, 1000s

Use: Anemias

LIDAFORM-HC CREME & LOTION, (Dome) Xylacaine 30 mg, iodochlorhydroxyquin 30 mg, Hydrocortisone alcohol 5 mg/Gm Creme Tube 1/4 oz, Lotion 1 oz 1 pt

Use: Inflammation

LIDAFORM-HC OTIC, (Dome) Hydrocortisone alcohol 10 mg, iodochlorhydroxyquin 30 mg, acetic acid 2%, Xylacaine 30 mg/cc Bot 10 cc

Use: Treatment of infection associated with otitis externa, otitis media, otomycosis

LIDA-MANTLE CREME, (Dome) 3% lidocaine HCl in Acid Mantle vehicle Tube 1/4 oz, 1 oz 1 lb Jar

Use: Topical anesthetic

LIDA-MANTLE-HC, (Dome) Lida Mantle

*Generic name recognized by NF, USP or A.M.A.

formula plus hydrocortisone alcohol 1/4% & 1/2% in Acid Mantle base Tube 1/4 oz, 1 oz

Use Anti-inflammatory

LIDA-MYCIN CREME, (Dome) 3% lidocaine HCl, neomycin 0.5% Tube 1/4 oz, 1 oz, 1 lb Jar

Use: Analgesic, Anti-Infective Topical

***LIDOCAINE HYDROCHLORIDE INJ**, USP NND 1962 ADR; Inj USP Alpha diethylamino 2,6 aceto xylicide HCl See Xylacaine Hydrochloride (Astro)

W/Epinephrine See: Xylacaine W/Epinephrine (Astro)

W/Dezomethasone See: Decadron W/Xylacaine, (Merck Sharp & Dohme)

LIDOSPORIN OTIC SOLN (Burroughs Wellcome) Polymyxin B Sulf 10000 u, lidocaine HCl 50 mg/cc propylene glycol Bot 10 cc

Use Ear drops

LI-FER-B-12, (New Mexico Pharm) Ferrous sulphate 1361 mg, thiamine HCl 1 mg, riboflavin 2 mg, liver desiccated 200 mg, B 12 1 mcg/Cop Bot 100s, 1000s

Use Dietary supplement

LIFIC-B-12, (Ascher) Cryst Vit B 12 5 mcg, folic acid 2.5 mg, liver inj equiv to cyanocobalamin 5 mcg/cc; Cryst Vit B 12 15 mcg, folic acid 5 mg, liver inj equiv to cyanocobalamin 10 mcg/cc; Cryst Vit B-12 50 mcg, folic acid 10 mg, liver inj equiv to cyanocobalamin 10 mcg/cc

LIFIC-B-12 "100" (Ascher) Crystalline Vitamin B 12 100 mcg, folic acid 10 mg, liver injection, equivalent to cyanocobalamin 10 mcg, phenol, os preservative 0.5%/ml

Use Anemias

LIFO-B-12 (New Mexico Pharm) Vit B-12 activity (equiv to 5 mcg B 12) 5 mcg, folic acid 5 mg, B 12 25 mcg/cc Vial 10 cc

LIFO-DEX (New Mexico Pharm) Liver Injection 10 mcgm, desoxyephedrine 5 mg, Vit B 12 50 mcgm, niacinamide 75 mg, folic acid 10 mg, phenol 0.5%/cc Vials 10 cc

LIF-O-GEN (Lif O Gen Inc) USP Oxygen (7 gal) at 400 p.s.i.g. (70° F) Unit complete with disposable mask

Use First aid treatment

LIFO-HEPARIN, (New Mexico Pharm) Heparin sodium 25 mg, choline chloride 100 mg, Vit B 12 50 mcg, folic acid 2 mg, niacinamide 50 mg, benzyl alcohol 1.5%/cc Vials 10 cc

LIFOLBEX, (Central) Cryst Vit B 12 5 mcg, folic acid 2.5 mg, liver inj 5 u/cc Vial 10 cc

- Vit B 1 1.5 mg, D 500 u /Cap or Tab Bot 180s 500s
Use Orally treatment of psoriasis
- LIPAN IMPROVED** (Spirit) 400 mg of 5x NP whole hog pancreas equiv to 2000 mg pancreat n NF plus pyridoxine NCI 3 mg /tab or Cap Bot 180s 500s
- ***LIPANCREATIN** A preparation of hog pancreas containing a reliably high content of steapsin (pancreatic lipase) with adequate amounts of trypsin, amylase and other pancreatic enzymes See Cotazym (Organon Inc)
- LIPASE W/Amylase Protease** See Trulase Tabs (Smith Miller & Patch)
- LIPIODOL**, NNR 1950 (Fougera) Iodized poppy seed oil Amp (40% Iodine) 5 cc, 10 cc Vial 20 cc
Use: Radiopaque
- W/Sulfonamide See: Visciodol Susp (Fougera)
- LIPIODOL RADIOLOGIQUE ASCENDANT**, NNR 1950
- LIPO-ADRENAL CORTEX, STERILE SOLUTION**, NNR 1950 (Upjohn) Each cc contains biological activity equiv to 1 mg (10 glycogen u) of 17 hydroxycorticosterone Vial 1 cc, 5 cc
Use: I.M., adrenal cortex therapy
- LIPO-B** (United Lab) Vit B 1 2 mg, B 2 4 mg niacin 20 mg, pantothenic acid 2 mg, Vit B 6 2 mg, inositol 150 mg, choline dihydrogen citrate 300 mg /Tab Bot 100s 1000s
Use: (2 t i d) Lipotropic
- LIPICAL** (G P Harvey) Choline bitartrate 0.2 Gm, dl methionine 0.1 Gm inositol 0.1 Gm Vit B 12 2 mcg /Cap Bot 100s 1000s
Use: 3 w th meals; vitamin lipotropic
- LIPOCAPS** (Lakeside) Choline bitartrate 450 mg, dl methionine 150 mg inositol 100 mg /Cap Bot 100s
Use: (1 Cap t i d) Lipotropic therapy
- LIPPOCHOLINE**, NND 1961
- LIPO-COBEX** (New Mexico Pharm) Choline chloride 75 mg, dl methionine 25 mg, Thiamine HCl 25 mg, riboflavin 2 mg, nicotinic acid 100 mg Vit B 12 15 mcgm, liver inject 10 mcgm, phenol 0.5%/cc Vials 10 cc
- LIPO-DIAZINE**, (Donley Evans) Sulfadiazine 77 gr /5 cc Susp, Bot 4 ox. 1 pt
Use: Orally, sulfonamide therapy
- LIPODINE**, (Dwight) Iodobrassid Tab (4½ gr) Bot 100s 500s 1000s
Use: Iodine therapy
- LIPOFLAVONOID** (Smith Miller & Patch) Choline bitartrate 700 mg, inositol 334 mg dl methionine 84 mg, cabotamin
- * Generic name recognized by N.F., U.S.P. or A.M.A.
- concentrate 5 mcg, Vits B 1 1 mg, B-2 1 mg, niacinamide 10 mg, B 6 1 mg panthanal 1 mg, C 300 mg, leman bio flavonoid complex 300 mg /3 caps Bot 100s 500s
Use Nutritional supplement
- LIPO GANTRISIN**, NND 1962 (Roche Lab) Gantrisin acetyl 20% in a palatable, readily digestible vegetable oil emulsion Bot 4 & 16 oz
Use: Systemic and urinary tract infections
- LIPOGEST**, (Breen) Inositol 43 mg, methionine 85 mg, choline dihydrogen citrate 220 mg /Cap Bot 100s, 1000s
Use Lipotropic
- LIPO-NEPIN**, (Darwin) Heparin Sod Vial (200 mg, 20,000 u /cc) 2 cc; (100 mg, 10,000 u /cc) 4 cc, 5 cc; (50 mg, 5000 u /cc) 10 cc; (10 mg, 1000 u /cc) 10 cc.
- LIPO-NEPLEX**, (U.S. Vitamin) Liver 11.5 Gm, Vit B 12 & B 12b 0.5 mcg, choline 10 mg, Inositol 5 mg, Thiamine HCl 3 mg, riboflavin 3 mg, niacinamide 10 mg, pyridoxine NCI (B 6) 0.3 mg, Cal pantotheante 3 mg /Cap Bot 100s, 500s, 1000s
Use: Multiple vitamin B deficiencies
- LIPO-HORMESTRIN**, (Smith, Miller & Patch) Natural estrogens in oil Vial (20,000 u /cc) 10 cc, 30 cc Box 1s, 6s 25s
Use: I.M., estrogen therapy
- LIP-OID**, NND 1962 (Tulog) See Estradiol
- LIPOIODINE**, NNR 1953 (Ciba) Ethyl di iodobrassidate Iodobrassid 41% Iodine in organic combination Tab (4½ gr) Bot 30s, 100s
Use Orally, iodine therapy
- LIPO-LEVAZINE** (Hiss) Sulfadiazine 0.5 Gm /5 cc Bot pt
Use Sulta therapy
- LIPOLIQID**, (Lakeside) Each 15 cc contains tricholine citrate 3.75 Gm (equiv Ia 9.15 Gm of choline dihydrogen citrate), Vit B 12 4.20 mcg & inositol 75 mg Bot 16 oz
Use Lipotropic therapy
- LIPO-LUTIN**, NND 1962 (Parke Davis) An oil sol of progesterone 2 u (2 mg) /cc, Amp 1 cc, 5 u (5 mg) /cc, Amp 1 cc, Vial 10 cc 10 u (10 mg) /cc, Amp 1 cc, 25 u (25 mg) /cc Amp 1 cc, Box 10s, Vial 10 cc 50 u (50 mg) /cc Vial 5 cc
Use: I.M. for deficiency of progesterone
- LIPOLYTIC ENZYME**, W/Proteolytic enzyme amylolytic enzyme See Formulose Tabs (Kremers Urban)
- LIPOMIC**, (Chicago) Inositol 25 mg, methionine 50 mg, choline dihydrogen citrate 123 mg, liver conc 60 mg, Vit B 12 1 mcg /Tab Bot 100s, 1000s
Use: Lipotropic.

LIPOMIC INJECTION. (Chicago Pharm)
Heparin sod. 25 mg., choline chloride 50
mg., inositol 50 mg., Vit. B-12 100 mcg.,
folic acid 2 mg., nicotinamide 50 mg.,
benzyl alc. 1.5%/cc. Vial 10 cc.
Use: I.M. for atherosclerosis

LIPO-MISTRONE. (Smith, Miller & Patch)
Natural estrogens 2.5 mg., progesterone 25
mg. In oil/cc. Vial 10 cc., Box 1s, 6s, 25s
Use: I.M., long-acting hormone deficiency

LIPOMUL-ORAL. (Upjohn) Corn oil 10 Gm /
15 cc. w/antioxidants, emulsifying agents,
cyclamate sod., & preservative. Bot 1 pt.
Use: (1-3 R. oz. t.i.d., p.c.) Underweight in
the malnourished.

LIPOMUL I.V., N.N.D. 1962. (Upjohn) Col-
linsed oil 15 Gm., dextrose anhydrous 4
Gm., lecithin 1.2 Gm., oxyethyleneoxy-
propylene polymer 0.3 Gm./100 cc. Bot 500
cc./15% w/administration unit.

Use: Nutrient.

LIPO-NICIN. (Brown) Nicotinic acid 250 mg.,
niacinamide 150 mg., C 100 mg., B-1 25
mg., B-2 2 mg., B-6 3 mg./Tab Bot. 100s.
Use: Hypercholesteremia, cerebral sclerosis.

LIPO-NICIN PTZ. (Brown) Pentamethylene
tetrazole 100 mg., nicotinic acid 100 mg.,
Vit. C 100 mg., l-glutamic acid 50 mg.,
niacinamide 5 mg., riboflavin 2 mg., d-
pantholanyl alcohol 5 mg., In a port wine
vehicle, alcohol 20%/5 cc. Bot pts.
Use: Vasodilator.

LIPO-PALSUL. (Polmedica) Sulfadiazine,
sulfamerazine, sulfamethiazine 0.5 Gm.
total/Cap. Bot. 30s. Tsp. Emulsion, Ptt.
Use: Treatment of infections.

LIPOPHILATE. (Brewer) Choline bitartrate
200 mg., inositol 100 mg., methionine 150
mg., liver ext. (1-50) 80 mg./Cap. Bot. 100s
Use: Liver dysfunction

LIPOPLETE. (Hart) Vit. A 125,000 u., D 1000
u., B-1 5 mg., B-2 2.5 mg., B-6 0.5 mg.,
C 75 mg., niacinamide 40 mg., col. pantho-
thenate 4 mg., Vit. E 2 I.U., B-12 1 mcg.,
folic acid 0.5 mg., dl-methionine 10 mg.,
inositol 15 mg., choline bitartrate 31.4 mg.,
col. phosphorus, iron, cobalt, copper, mang.,
iodine, molybdenum, pot., zinc, mg./Cap-
Bot. 100s, 500s.
Use: Dietary supplement.

LIPOPROTEIN POWDER. (Upjohn) Pkg. 1/2
& 1 lb.
Use: 2 tbsp. with milk I.t.d. debilitated
patients.

LIPO-SAL. (Danley-Evans) Sodium salicylate
327 mg., vegetable oil/5 cc. Bot. pts.

Use: Antirheumatic.

LIPO-SULFALOID. (Lloyd Dabney & Wester-

feld) Sulfameth-dia-mer 167 mg./5 cc.
Bot. pts.

Use: Sulfonamide therapy.

LIPOTAIN CAPSULES. (Stuart) Betaine 333
mg., choline 35 mg., desic. liver N.F. 35
mg., Vit. B-12 2 mcg./Cap. Bot. 100s, 1000s
Use: Disturbed fat metabolism.

LIPOTAIN LIQUID. (Stuart) Betaine 3 Gm.,
choline 210 mg., liver fraction N.F. 210
mg., Vit. B-12 12 mcg./5 cc. Bot. 1 pt.,
1 gal.

Use: Disturbed fat metabolism.

LIPO-TEX. (Bentex) Sulfadiazine 0.167 gm.,
sulfamerazine 0.167 gm., sulfacetamide
0.167 gm., emulsified vegetable oil 50%,
sucrose 12.5%/5 cc Bot 4 oz., pt.

Use: Sulfonamide therapy.

LIPOTHYN CAPSULES. (Flint, Eaton) Choline
bitartrate 400 mg., methionine 100 mg.,
inositol 100 mg., Vit. B-12 2 mcg., alpha-
tocopherol acetate 5 mg./Cap. Bot. 100s,
500s, 1000s.

Use: Lipotropic agents & dietary supple-
ments.

LIPOTINIC. (Wampole) Vit. B-1 3.34 mg.,
B-2 3.34 mg., B-6 0.67 mg., niacinamide
33.34 mg., Col. panthothenate 6.7 mg., Vit.
C 100 mg., choline dihydrogen citrate 33.34
mg., inositol 33.34 mg., methionine 3.34
mg., folic acid 0.5 mg., Vit. B-12 1.34 mcg.,
iron 16.67 mg., liver-stomach 100 mg./cap
Bot. 100s.

Use: 1 daily; dietary supplement.

LIPOTRASI. (G. F. Harvey) Choline bitor-
trate 0.2 Gm., dl-methionine 0.1 Gm., inositol
0.1 Gm., Vit. B-12 2 mcg./Cap. Bot
100s, 1000s

Use: Correction of liver dysfunction and
abnormal fat metabolism

LIPOTRIAD CAPSULES. (Smith, Miller &
Patch) Choline bitartrate 700 mg., inositol
334 mg., dl-methionine 84 mg., Vit. B-12
activity 5 mcg., thiamine HCl 1 mg., ribo-
flavin 1 mg., niacinamide 10 mg., pyri-
doxine HCl 1 mg., panthothenol 1 mg./3 Cap
Bot. 100s, 1000s.

Use: Lipotropic agents w/vitamin supple-
ment.

LIPOTRIAD LIQUID. (Smith, Miller & Patch)
Tris choline citrate 460 mg., inositol 334 mg.,
dl-methionine 84 mg., Vit. B-12 5 mcg.,
riboflavin 1 mg., thiamine HCl 1 mg., nia-
cinamide 10 mg., pyridoxine HCl 1 mg.,
panthothenol 1 mg./5 cc. Bot. 4 oz., 1 pt.

Use: Lipotropic agents w/vitamin supple-
ment.

LIPO-TRIAZINE. (Danley-Evans) Sulfametha-
zine, sulfadiazine & sulfamerazine of equal
parts, 77 gr. al total sulfonamides/5 cc. or

- Cap Bot 4 oz, 1 pt Caps Bot 30s, 250s
Use Sulfonamide therapy
- LIPO TRI S** (Hiss) Sulfo metho diomer 0 167 Gm / 3 cc Bot pts
Use Sulfa therapy
- LIPO-TRISULTEX.** (Bentex) Sulfadiazine 0 167 Gm, Sulfamerazine 0 167 Gm, Sulfacetamide 0 167 Gm, Emulsion of vegetable oil 50%, sucrose 12.5%/5 cc Bot 4 oz 1 pt
Use Triple sulfonamide therapy
- LIPOTROPIC.** (Lokeside) Choline dihydrogen citrate 200 mg dl methionine 100 mg, Inositol 100 mg / Cap Bot 100s
Use: Cirrhosis diabetes mellitus
- LIPOTROPIC PREPARATIONS.**
See Choline Products (Various Mfr)
Cytellin, Susp (Lilly)
Betaine Products (Various Mfr)
Inositol Products (Various Mfr)
Lecithin (Various Mfr)
Methionine Products (Various Mfr)
- LIPOTROPIN.** (Vascular) Choline dihydrogen citrate 422 mg, inositol 134 mg, pyridoxine HCl 2 mg / Tab Bot 100s, 1000s
Use Treatment of disturbed fat metabolism
- LIPOTROPLEX.** (Lincoln) Methionine 150 mg, choline 100 mg, inositol 100 mg, Vit B 1 3 mg, B 2 3 mg, B 6 20 mg, Cal pantothenate 3 mg, niacinamide 20 mg, brewers yeast qs / Tab Bot 100s, 1000s
- LIPROTEIN POWDER.** (Upjohn) Protein 45 Gm, fat 35 Gm, carbohydrate 14 Gm, yeast 15 Gm, liver concentrate 2 Gm, B 1 5 mg, B 2 5 mg B 6 2 mg, cal pantothenate 10 mg, nicotinamide 30 mg, folic acid 2 mg B 12 5 mcg, w/Co 0.91% No D 0.9B% Cl 0.7B% K 0.91%, P 0.65% / 100 Gm Pkg ½ 1, & 5 lb
Use High protein & caloric intake
- LIQUACILLIN** (Lilly) Penicillin G Pot 3 600,000 u When mixed as directed, each 5 cc will contain 200,000 u Bot 2 oz
Use: Penicillin susceptible organisms
- LIQUA COMPLEX.** (Ethical Pharm) Sulfadiazine 2.5 gr, sulfamerazine 2.5 gr, sulfamethoxime 2.5 gr / dr Bot 1 pt
Use: Triple sulfonamide therapy
- LIQUADIAZINE** (Ethical Pharm) Sulfadiazine 7.7 gr, Sod lactate 7.7 gr / tsp Bot 1 pt
Use: Sulfadiazine therapy
- LIQUA-D.M.** (Ethical Pharm) Sulfadiazine 3.8 gr, sulfamerazine 3.8 gr, sod lactate 7.7 gr / dr Bot 1 pt
Use: Dio Mer sulfonamide therapy
- LIQUAEMIN SODIUM, NND 1962** (Organon) Heparin sod aq sol Vial (200 mg, 20,000 u/cc) 2 cc; (50 mg, 5000 u/cc)
- 1 cc, 10 cc; (100 mg, 10 000 u/cc) 4 cc; (10 mg, 1000 u/cc) 10 cc; (20,000 u w/gelatin/2 cc) 2 cc Disposable syringe 1 cc cartridge providing 20,000 u Box 1s.
Use: (I/M and I/V) Anticoagulant
- LIQUAMAR.** (Organon) 3 (1' Phenylpropyl)-4 hydroxycoumarin * Phenprocoumon 3 mg / Tab Bot 100s, 1000s
Use: Anticoagulant
- LIQUAPEN** (Pfizer) Penicillin G Pot. 250,000 u / 5 cc, dry Bot 50 cc
Use: Orally, penicillin therapy
- LIQUA-SWEET.** (Forward House) (Matheson) Mixture of saccharine & sorbitol Plastic squeeze bot 1 oz,
Use Noncaloric sweetening agent
- LIQUI-CEE.** (U S Standard) Ascorbic acid 1 Gm / 5 cc Bot 4 oz
Use Ascorbic acid therapy
- LIQUO CITRALKI.** (Parke, Davis) Disodium hydrogen citrate 15 gr / dr Bot 8 oz, 1 gal
Use: Orally, antacid and to increase the alkali reserve
- LIQUO COGEL** (Lilly) Methylcellulose 9 Gm / 100 cc Bot 8 oz, 16 oz
Use: Orally, laxative
- LIQUO GERMICIDAL DETERGENT.** (Parke Davis) Susp of alkylamine hydrachlorides containing phemeral Bot 16 oz, 1 gal
Use: Cleansing & germicidal properties
- LIQUO LIVER EXTRACT.** See: Liver Sallv lion, U S P
- LIQUO PETROLATUM EMULSION, NNR 1949** See: Petrolatum Liq
- LIQUIMAT.** (Texas Pharmacol Co) Sulfur 5%, hexachlorophene 0.25%, alcohol 22%, zinc, titanium and iron oxides, kaolin and talc Plastic Bot 2 fl oz
Use Acne and other skin ailments
- LIQUIPIX CREAM** (Dermatologicals, Inc) Liquor carbonis detergens, 3% in water washable base Jar 2 oz, 1 lb
Use Dry, chronic dermatoses
- LIQUIPRIN.** (Johnson & Johnson) Salicylo mide 2½ gr / 2.5 cc Dropper bot 50 cc
Use Analgesic
- LIQUITAL** (Merck Sharp & Dohme) Each 30 cc (1 fl oz) contains 0.12 Gm (2 gr) of phenobarbital Differs from elixir phenobarbital U S P & should not be dispensed when the U S P product is specified Bot 1 gal
Use Sedative
- LIQUOID METHA-MERDIAZINE** See: Metha Merdiazine Susp (McNeil)
- LIQUOR CARBONIS OETERGENS.** See: Cool Yar Solution U S P (Various Mfr)
- LIQUOR DIASTOS** (Merck Sharp & Dohme) Pepsin D 52 Gm, papain 0.7 Gm, rennin 50 mg, nitrohydrochloric acid, dilute D.3
- * Generic name recognized by N F, U S P or A M A

- cc., lactic acid. 0.03 cc., nux vomica 27 mg./30 cc. Bot. 1 pl.
 Use: Digestant.
- LIRIMIN CAPSULES.** (Merck Sharp & Dohme)
 Ferrous sulfate exsic. 0.30 Gm., liver-yeast conc. 0.15 Gm., thiamine HCl 0.25 mg., riboflavin 0.50 mg./Cap. Bot. 100s.
- Use: Nutritional deficiency.
- LIRON-B.** (Breon) Liver conc. N.F. 0.29 Gm., ferrous sulfate exsic. 0.065 Gm., thiamine HCl 0.33 mg., riboflavin 0.67 mg., nicotinamide 3.33 mg./Cap. Bot. 100s, 500s, 1000s.
- Use: Secondary anemias.
- LIRON-FORTE.** (Breon) Liver conc. N.F. 240 mg., ferrous sulfate exsic. 200 mg., thiamine HCl 2 mg., riboflavin 2 mg., nicotinamide 10 mg., folic acid 1.7 mg./Cap. Bot. 100s, 500s, 1000s.
- Use: Hematinic.
- LISCOR-B INJECTION.** (E. S. Miller) Thiamine HCl 100 mg., ascorbic acid 100 mg., choline chloride 100 mg., methionine 50 mg., riboflavin 5 mg., pyridoxine HCl 5 mg., d-pantothenyl alcohol 5 mg., niacinamide 75 mg./2 cc. Vial 10 cc.
- LISSEPHEN.** N.N.D. 1959. (Abbott) Mephenesin .25 Gm./Tab. Bot. 100s, 1000s.
- LISTACORT.** (Testogor) Prednisone 5 mg./Tab. Bot. 100s. Rheumatoid arthritis and other rheumatic disorders.
- LISTICA.** (Armour Pharm.) Hydroxyphenamate. 200 mg./tab. Bot. 50s
- Use: Anxiety and tension states.
- LITHIATED SORGHUM COMPOUND.** (Merck Sharp & Dohme) Hydro-alcoholic ext. of broom-corn seed, corn silk, saw palmetto berries & hydrangea & lithium benzoate 1.3 Gm. & lithium citrate 2.2 Gm. Bot. 1 pl., 1 gal.
- Use: Diuretic preparation.
- LITHITROL.** (Columbus) Salicylamide 250 mg., methenamine mandelate 250 mg./Tab. Bot. 100s, 1000s w/pyridoxine Tab. Bot. 100s, 1000s
- Use: Orally, (2 t.i.d.) urinary antiseptic
- LITHITROL W/PYRIDOXINE.** (Columbus) Pyridoxine HCl 166 mg., salicylamide 250 mg., methenamine mandelate 250 mg./Tab. Bot. 100s.
- Use: Urinary disorders.
- LITHOPLEX.** (Kremers-Urban) Thiamine HCl 10 mg., riboflavin 0.2 mg., niacinamide 1 mg., liver inj. 10 u./cc. Vial 10 cc., 20 cc.
- Use: Anemias.
- LITRISON.** (Hoffmann-La Roche) dl-Methionine 150 mg., choline dihydrogen citrate 180 mg., thiamine HCl 3 mg., riboflavin 3 mg., panthanal 4.5 mg., pyridoxine HCl 3 mg., niacinamide 9 mg., folic acid 0.6 mg., biotin 0.15 mg., Vit. B-12 0.001 mg., A palmitate 4500 u., E acetate 4.5 mg./Cap. Bot. 100s, 500s.
- Use: Functional impairment of liver.
- LIT-U-PEPT.** (Overseas) New name for Vit-U-Pept.
- LITWELFO.** (C. D. Smith) Liver injection 20 mcg., Vit. B-12 30 mcg., folic acid 5 mg., phenol 0.5%/cc. Vial 10 cc.
- Use: Anemias
- LIVER B.** Liver desic. See: Hepazyn, Tob. (Endocrine)
- LIVER-B-VITAMIN.** (Sherman) Riboflavin 0.5 mg., niacinamide 20 mg., thiamine HCl 20 mg., liver inj., crude 2 u./cc. Vial 30 cc.
- Use: Hematinic.
- * **LIVER CONCENTRATE.** N.F. See: Intrahepitol, Vial (Lederle)
- W/Folic acid & disulfa See: Lifasulfas, Tab. (Upjohn)
- * **LIVER, DESICCATED.** N.F. Desic. liver substance
- * **LIVER EXTRACT.** N.F. Dry liver ext w/Vit. B-12 & folic acid.
- See: Anplex, Amp. (Chicago Pharm.). Anplexol, Vial (Chicago Pharm.). Anplexul, Vial (Chicago Pharm.).
- LIVER EXTRACT OBTAINED FROM PREGNANT COWS.** See: Biohepulin, Vial. (International Biochemical)
- LIVER-FERROUS CAPSULETTES.** (Armour) Not listed. Thiamine HCl (B-1) 1 mg., riboflavin (B-2) 2 mg., niacinamide 10 mg., ferrous sulfate 0.24 Gm., liver conc. (N.F.) 0.5 Gm./cap. Bot. 100s, 500s.
- Use: Hypochromic anemia.
- * **LIVER FRACTION 1.** N.F. Soluble liver fraction.
- * **LIVER FRACTION 2.** N.F.
- LIVER FUNCTION AGENTS.**
- See: Bromsulphalein, Amp. (Hynson, Westcott and Dunning)
 Iodaphthalein (Vorhaus Mfr.)
 Sod. Benzoate Seln.
 Sulfabromophthalein Sodium, Amp., U.S.P. (Gotham)
- * **LIVER INJECTION.** N.F. Liver ext. for parenteral use (Various Mfr.)
- See: Bioxiver, Vial (Breon).
 Fols Brunin, Amp. (Philadelphia Amp. Labs)
- * **LIVER INJECTION CRUDE.** N.F. (Various Mfr.)
- See: Campolan, Amp. (Winthrop Labs)
 Fernoemon PL (Organon)
 W/Folwits (Lederle)
 Cyanocobalamin 2 mg. & folic acid 2 mg./cc. Vial 10 cc.
 W/Vit. B-12. See: Cru-B-12, Vial (Thompson).
- * Generic name recognized by N.F., U.S.P. or A.M.A.

- LIVER & IRON TONIC.** (Sherman) Liver desic 200 mg, ferrous sulfate exsic 136 mg, thiamine 1 mg, riboflavin 2 mg, Vit B 12 conc 1 mcg/Cap Bot 100s, 500s, 1000s
 Use: Secondary anemias
- LIVERPLEXIR-B-12.** (Lincoln) Thiamine HCl 20 mg, niacinamide 80 mg, riboflavin 2 mg, Sad pantothenate 2 mg, pyridoxine 4 mg, Vit B 12 10 mcg, liver 2 w/2 cc Vial 30 cc
- LIVERPLEXIR W/VIT. B-T2.** (Lincoln) Liver desic 250 mg, ferrous sulfate exsic 136 2 mg, copper sulfate 3 mg, thiamine HCl 15 mg, riboflavin 1 mg, niacinamide 5 mg, methionine 2 mg, Vit B 12 5 mcg/Tab Bot 100s.
- LIVER, RAW** See Joyran plus Intrinsic Factor (Viobin)
- ***LIVER SOLUTION, NF** Liq liver ext (Armour) Sol, oral, Bot 8 oz, 1 pt
- LIVER VASODILATOR PRINCIPLE.** See: Cortunon, Amp & Cap (Cortunon)
- ***LIVER WITH STOMACH, NF** Cap NF W/Thiamine (C D Smith) liver 10 w, 6 mg of thiamine HCl/cc Vial 15 cc
 Use Anemias
- LIVER VASOCONSTRICTOR** See Kutapresin, Vial (Kremers Urban)
- LIVER, WHOLE DESICCATED.** See: Desiver, Tab (US Vitamin Corp)
- LIVER & YEAST CAPSULES.** (Armour) Thiamine HCl 0.34 mg, riboflavin 0.67 mg, niacinamide 5 mg, yeast conc 0.15 Gm liver fraction #2 0.5 Gm/Cap Bot 100s, 500s
 Use: Anemia
- LIVER & YEAST WITH IRON TABLETS.** (Armour) Thiamine HCl 0.34 mg, riboflavin (B 2) 0.67 mg, niacinamide 5 mg, ferrous sulfate 0.166 Gm, yeast conc 0.15 Gm liver fraction Na 2 (NF) 0.5 Gm/Tab Bot 100s, 500s
 Use: Anemia
- LIVER & YEAST LIQUID.** (Armour) Yeast digest of liver fraction Na 1 NF 4.5 Gm thiamine HCl 3 mg, riboflavin 5 mg, liq yeast ext Na 3 0.6 cc/fl oz Bot 8 oz, 1 gal
 Use: Anemia
- LIVERGRAN** (Rawl) Desiccated whole liver 9 Gm, Vit B 1 18 mg, B 2 36 mg, niacinamide 90 mg, choline bitartrate 216 mg, B 6 3.6 mg, calcium pantothenate 3.6 mg, inositol 90 mg, biotin 6 mcg, Vit B 12 5.4 mcg, methionine 198 mg, arginine 242 mg, cystine 72 mg, glutamic acid 675 mg, histidine 99 mg, isoleucine 333 mg, leucine 495 mg, lysine 297 mg, phenylalanine 189 mg, threonine 333 mg, tryptophane 45 mg, tyrosine 180 mg, valine 306 mg/3 tsp Bot gran 3 oz, 15 oz
 Use: Nutritional supplement
- LIV-FER-B IMPROVED.** (Sutliff & Cose) Liver fraction Na 1 0.6 Gm, ferrous gluconate 0.6 Gm, mang gluconate 0.12 Gm, copper gluconate 0.01 Gm, cobalt gluconate 0.03 Gm, zinc sulfate 0.01 Gm, Vit B 1 2 mg, B 2 2 mg, B 12 2 mcg, folic acid 0.4 mg, sad citrate 0.6 Gm/fl oz Elix 1 pt, 1 gal
 Use Hematinic
- LIV-FER-BLEK.** (Sutliff Case) Ferrous gluconate 0.194 Gm, liver conc 0.228 Gm, thiamine HCl 1 mg, riboflavin 1 mg, niacin 10 mg, brewers yeast 0.194 Gm/Tab Bot 100s, 1000s
 Use Iron deficiency anemias
- LIVIBRON** (Parke, Davis) Liq liver conc equiv to 10 Gm fresh liver, Vit B 1 5 mg, B 2 2 mg, ferrous sulfate 12 gr, Mang citrate 1/4 gr, alcohol 12%/fl oz Bot 8 oz, 1 gal
 Kapsbal, No 1 Cap contains Liver conc equiv to 2.5 Gm fresh liver Vit B 1 1.25 mg, B 2 0.5 mg, B 12 1 mcg, ferrous sulfate 3 gr, Mang citrate (soluble) 1/16 gr. Bot 100s, 1000s
 Use Nutrient tonic & hematinic
- LIVIFOL INJECTABLE** (Smith Dorsey) B 12 25 mcg, folic acid 5 mg, purified sol of liver 2 mcg/cc Vial 10 cc
 Use Anemias
- LIVIPÉPTONE.** (Borchardt) Liver fraction No 1 NF. 28 Gm, iron & Mang peptones 2 Gm, Vit B 1 10 mg, B 2 5 mg, B 12 like activity 10 mcg, B 6 1 mg, panthelol 5 mg, niacinamide 25 mg, choline 33 mg, alcohol 10%/fl oz Syr Bot 8 fl oz, 1 gal
 Use: Hypochromic anemias
- LIVIRON FORTIFIED** (Philadelphia Ampule Labs) Vit B 12 10 mcg, niacinamide 25 mg, B 2 1 mg, B 6 1 mg, iron peptonate 30 mg, liver lnj refined 1 mcg, B 1 20 mg/cc Vial 30 cc
 Use: Fortified vitamin therapy
- LIVITAMIN** (Massengill) Liver desic 400 mg, iron peptonized 147 mg, thiamine HCl 3 mg, riboflavin 3 mg, niacinamide 10 mg, pyridoxine HCl 0.5 mg, Cal pantothenate 2 mg, Vit B 12 5 mcg, folic acid 1 mg/Cap Bot 100s, 500s, 1000s
 Use: Anemias
- W/Intrinsic factor** (Massengill) Desicc liver 400 mg, iron peptonized 147 mg Vit B 1 3 mg, B 2 3 mg, niacinamide 10 mg, Vit B 12 5 mcg, B 6 0.5 mg, Cal pantothenate

- 2 mg., folic acid 1 mg., intrinsic factor 1/2 cc./Cap. Bot. 100s
 Use: Pernicious anemia & geriatrics.
- W/IRON.** (Massengill) Iron peptonized N.F. 6 1/2 gr., mag. citrate 2 1/2 gr., liver fraction No. 1 2 Gm., rice bron ext. 1 Gm., thiamine HCl 10 mg., riboflavin 10 mg., niacinamide 50 mg., pyridoxine HCl 1 mg., pantothenic acid 5 mg., Vit. B-12 20 mcg., inositol 30 mg., choline 60 mg./30 cc. Bot. 8 oz., 1 pt., 1 gal.
 Use: General hematinic.
- LIVITAMIN LIQUID.** (Massengill) Liver conc. equiv. to fresh liver 2 oz., Vit. B-1 3 mg., B-2 1 mg., nicotinamide 25 mg., Vit. B-6 1 mg., pantothenic acid 5 mg., Iron & Mang. peptonized 1.94 Gm./oz. Bot. 8 oz.
 Use: Hematinic w/vitamin supplement.
- LIVITOL** (Consolidated Mid.) Vitamine A 2500 u., D 250 u., B-12 3 mcg., C 37.5 mg., folic acid 0.5 mg., Iron 15 mg., cobalt 0.1 mg., copper 0.25 mg., liver fr. 250 mg., liver pow 25 mg., gastric mucosa 75 mg./cap. Bot. 30s, 100s, 250s, 1000s, 5000s.
 Use: Anemias.
- LIVITOL SYRUP.** (Consol. Midland) Liver crude 21.7 Gm., ferrous sulfate 440 mg., Vit. B-1 3.3 mg., B-2 0.67 mg., B-6 0.33 mg., cal. pantothenate 0.67 mg., nicotinamide 10.10 mg., choline chloride 6.70 mg., tauric acid 1.7 mg./10 cc. Bot. 100 cc.
 Use: Orally, hematinic.
- LIVOTON.** (Estra) Crude liver 2 u., thiamine HCl 100 mg., riboflavin 0.5 mg., pyridoxine HCl 5 mg., Cal. pantothenate 5 mg., niacinamide 100 mg./cc. Amp. 1 cc., Vial 10 cc., 30 cc.
 Use: Vitamin B complex.
- LIVREX-2.** (Consolidated Mid) Liver ext. crude equal to 2 mcg. B-12/cc. Vials 10 cc.
- LIVREX-10.** (Consolidated Mid) Liver ext. refined equal to 2 mcg. B-12/cc. Vials 10 cc.
- LIVREX-20.** (Consolidated Mid) Liver Inf. equiv. to Vit. B-12 20 mcg./cc. Vial 10 cc.
- LIVREX-BB.** (Consolidated Mid.) Liver ext. refined equal to B-12 10 mcg., vit. B-1 5 mg., B-12 5 mcg./cc. Vials 10 cc.
- LIVREX-BP.** (Consolidated Mid.) Liver ext. crude equal to 2 mcg. B-12, vit. B-12 15 mcg., folic acid 5 mg./cc. Vials 10 cc.
- LIXA-BETA.** (Warner-Chilcott) Thiamine HCl 1 mg./dr. Bot. 8 fl. oz., 1 pt., 1 gal.
 Use: Palatable Vit. B-1 therapy.
- LIXAMINOL ELIXIR.** (Ferndale) Aminophylline 259 mg with alcohol, 30% by volume/ Tsp Bot. pts
 Use: Caranary vasodilator, diuretic.
- LIXAMINOL AT.** (Ferndale) Aminophylline 259 mg, alcohol 4.5 cc., dextromethorphan hydrobromide 20 mg. in a sugar-free
 * Generic name recognized by N.F., U.S.P. or A.M.A.
- pleasantly flavored elixir/5 cc. Bot. 1 pt.
 Use: Acute and chronic asthma
- LIXATONE.** (Buffington) Liver fraction No. 1 750 mg, betaine HCl 180 mg., choline 180 mg., leucitol 180 mg., L-lysine monohydrochloride 300 mg., Vit. B-12 30 mcg, B-1 18 mg., B-2 18 mg., niacinamide 72 mg, B-6 3 mg., cal. pantothenate 12 mg., folic acid 1.5 mg./fl. oz. Bot. 8 oz., 1 gal.
 Use: Dietary supplement
- LIXOCOL ELIXIR.** (Lannett) Glycocol 28 gr., L-lysine 100 mg/Tsp. Bot. 1 pt.
 Use: Anorexia, muscular weakness and amino acid supplement.
- LIXOIL** (Lixoil Labs) Sulfonated fatty oils and one or more esters of higher fatty acids. Bot. 1 pt.
 Use: Dermatologic.
- LIXOPHEN.** (Massengill) Phenobarbital 1/2 gr. (20 mg./5 cc) Bot. 1 pt., 1 gal.
 Use: Sedative & hypnotic.
- LLD FACTOR.** See: Vitamin B-12, Prep. (Yorkous Mir.)
- L-MAXIPEN.** (Roerig) * Epiphenethicillin.
 Use: Antibiotic.
- LOBANA HOSPITAL LOTION.** (Ulmer) Stearic acid, lanolin, menthol.
 Use: Skin lotion.
- LOBELIA EXT.** W/Ephedrine HCl, calcium iodobehenate. See: Lobladrin, Tab. (Breon)
- LOBELIA EXT.** W/Phenobarb., ephedrine HCl. See: Barbelline, Tab. (Cole)
- LOBELIA, HCl.** See: Labelin, Amp. (Blschoff).
- LOBELINE.** (Sandoz) Amp. (1/20 gr. & 3/20 gr) 1 cc.
 Use: (M. or I.V.) respiratory stimulant.
- LOBELINE SULFATE.** See: No-Kolin, Tab. (Walker Pharm.).
- W/Methapyritene hydrochloride.** See: Sitabs, Tab. (Cenright)
- LOBIDINE.** (Searle) Pot. iodide, chloroform, camphorated tr. of Jamaica dogwood, cajuput oil, lobelia ext. & stinging ext. Syr. Bot. 1 pt., 1 gal.
 Use: (1-2 tsp. as indicated) Coughs.
- LOBIODRIN.** (Breon) Ephedrine HCl 1/4 gr., lobelia ext. pow., Cal. iodobehenate & sarsaparilla ext. pow./Tab. Bot. 100s, 500s, 1000s.
 Use: (1 q. 3 h) Cough preparation.
- LOFINALAC.** (Mead Johnson) Protein hydrolyzates with most of phenylalanine removed. Pwd. 2 1/2 lb.
 Use: Phenylketonuria.
- LOMOTIL** (Searle) Atropine sulfate 0.025 mg., diphenoxylate HCl 2.5 mg/Tab. Bot. 100s.
 Use: Diarrhea associated with acute infections.
- LOHALAC.** (Mead Johnson) Lactose 38%,

- fol 28%, casein 27%, Col phosphate monobasic Pot carb, Cal hydroxide Cal chloride Col carb, Pot chloride, Mag oxide Vit A conc, riboflavin, niacinamide, thiamine HCl Artificial color Tin 1 lb, 4 lb
- LONGEVAN.** (Northwest Labs) Alpha estradiol 0.5 mg, methyl testosterone 5 mg, pregnenolone 5 mg, monobasic ampheta mine phosphate 5 mg / Cap Bot 30s, 100s
- LONGIFENE.** See Buchizine
- LONTABS** (Ciba) Pyribenzomine hydrochloride 100 mg / long acting Tab Bot 100s, 1000s
Use For protection against allergic reactions
- LOOKASINE** (National Drug) Boric acid, zinc sulfate zinc phenolsulfonate, phenol, Pot alum, Sod salicylate menthol eucalyptol thymol & methyl salicylate Pow 4 oz, 1 pt
Use Vaginal cleanser
- LORA** (Wallace Labs) Chlorhexadrol
Use Hypnotic
- LORAKON.** (Baybank) Benzalkonium chloride 1:4000 & glycerin in an isotonic solution B oz, 1 pt
Use Topically, antiseptic
- LORFAN** (Rache) Levallorphan tartrate 1:3 Hydroxy Nallylmorphinan Amp (1 mg / cc) 1 cc Box 6s
Use IV or IM, overdosage of narcotics
- LORFAN PEDIATRIC AMPULES** (Rache) Levallorphan tartrate 0.05 mg / cc Amp 1 cc Box 6s, 25s
Use: Treatment of narcotic induced respiratory depression
- LORINAL** (Ainar Stone) Chloral hydrate Caps (7 1/2 gr) Bot 100s, Caps (15 gr) Bot 50s Drops (graduated dropper to deliver 3 1/4 gr or 7 1/2 gr) Bot 15 cc
Use Sedative
- LORINGO-TABS** (Cole) Benzocaine 1/20 gr tartar emetic 1/200 gr ext licorice 1 gr, menthol 1/100 gr, camphor 1/100 gr, cinnamon oil 1/150 min, peppermint oil 1 min / 50 Tab Bot 100s
Use: Sedative expectorant
- LOROPHYN SUPPOSITORIES & JELLY.** NNR 1955 (Eaton)
Supp Phenylmercuric acetate 0.02%, methylbenzethonium chloride and methyl paraben in a water miscible base Box 12s
Jelly (Phenylmercuric acetate 0.05%, polyethylene glycol of monoiso octyl phenyl ether 0.3%, methyl p hydroxy benzoate 0.05%, Sod borate 3%, in a special jelly base) Tube 3 1/4 oz
Use: Contraceptive
- LOROTRIDOL.** (Hilton Davis Chem) Bithionol 1:1000 conc. In 5% castor oil soap
Use: Germicidal soap
- LORYL** (Krameris Uibon) Chloral hydrate 337.5 mg phenyltoloxamine citrate 37.5 mg / Cap Bot 50s
Use: Hypnotic sedative
- LOTEN INCOTE.** (Paul Money) Rauwolfia serpens alkaloids 0.75 mg, Neethylline 100 mg, rutin 20 mg / Tob Bot 100s, 1000s
Use: Hypertension
- LOTIOBLANC.** (Arnor Stone) Zinc sulf, sulfured potash Plastic Bot 2 oz Bot 4 oz
Use: Acne, seborrheic dermatitis
- LOTOCREME.** (Abbott) Lanolin, olive oil glycerin, menthol Emulsion, B oz
Use: Topically, body rubs
- LOTONE.** (Westerfield) Thyroid 1/12 gr, odenol substance 1/4 gr, spermine 1 gr, leathan 3/4 gr / Tob Bot 100s, 500s, 1000s
Use: (I I d) Thyroid therapy
- LOTUSATE.** NND 1962 (Winthrop Labs) Tolbutol 5 allyl 5 sec butyl barbituric acid Cap (30, 50 & 120 mg) Bot 100s 1000s
Use: (1/2 to 2 gr HS) Sedative & hypnotic w/ Demerol HCl See Demerol w/ Lotusate, Tab (Winthrop Labs)
- LOUBARB** (Lauisons) Butobarbital sodium 30 mg / 5 cc Elixir Containers 1 pt, 1 gal
Use Sedative barbiturate
- LOUCARBATE** (Lauisons) Corbetapentane citrate 12.5 mg, salicylamide 3 1/2 gr, phenocelin 1 1/4 gr, caffeine 1/2 gr / Cap Bols 25s 100s Tabs 25 mg Bot 100s
Use Non narcotic cough suppression analgesic compound
- *LOUCEVERIN CALCIUM** Calcium 5 formyl 5.678 tetrahydroteroylglutamate
- LOUCORT-N NASAL SPRAY.** (Lauisons) 0.5% saln hydrocortisone acetate 0.5% neomycin sulfate 0.5% 1.5% hydrocortisone acetate 1.5%, neomycin sulfate 0.5% / cc 0.5% soln 15 cc squeeze bot 1.5% saln 10 cc squeeze bot
Use Steroid antibiotic therapy
- LOUKAPEC-N** (Lauisons) Neomycin sulfate 300 mg kaolin 60 Gm, pectin 130 mg, suspended with methylcellulose / fl oz Bot pts
Use Oarhea
- LOUPRED** (Lauisons) Prednisone 0.75 mgm, salicylamide (5 gr) 325 mgm, ascorbic acid 20 mgm aluminum hydroxide 75 mgm in a suitable buffered base / Tab Bols 100s 1000s
Use Corticoid salicylate therapy for dermaloses arthritis fibrositis and related disorders
- LOUPRED-N LOTION.** (Lauisons) Prednisone

- lane 5 mg, neomycin sulfate 5 mg/cc. 15 cc. plastic squeeze bottle.
 Use: An anti-infective, anti-inflammatory lotion.
- LOUPRED-N OINT.** (Lousonar) Prednisolone 5 mg, neomycin sulfate 5 mg./gr. Tubex 10 gr.
 Use: Topical treatment of allergic dermatoses.
- LOWEBARS CTR.** (Lowe) dextro-amphetamine sulfate 15 mg, amobarbital 90 mg./Cap. Bot. 100s, 250s.
 Use: Oberity and treatment of alcoholism and narcolepsy.
- LOWEBESE CTR.** (Lowe) dextro-amphetamine 15 mg., thyroid 3 gr., atropine 1/4 gr., atropine sulfate 1/180 gr., phenobarbital 1/4 gr./Cap. Bot. 100s, 250s.
 Use: Control of obesity.
- LOWEDIX CTR.** (Lowe) d-amphetamine sulfate 15 mg /Cap. Bot. 100s, 250s.
 Use: Oberity control.
- LOWESERP.** (Lowe) Reserpine 0.25 mg /Tab. Bot. 100s, 500s, 1000s.
 Use: Hypertension and anxiety.
- LOWETRATE.** (Lowe) Pentoerythritol tetranitrate 30 mg /Cap. Bot. 100s, 250s.
 Use: To prevent attacks in angina pectoris.
- LOWETRATE CTR.** (Lowe) Pentoerythritol tetranitrate 30 mg /Cap. Bot. 100s, 250s
 Use: Prevention of attacks in angina pectoris.
- LOWETRATE-S CTR.** (Lowe) Pentoerythritol tetranitrate 30 mg, secobarbital 50 mg /Cap. Bot. 100s, 250s.
 Use: Prevention of attacks in angina pectoris, short acting sedative.
- LOWILA.** (Westwood) Sod. lauryl sulfacetate, Cake 4 oz., liq., 1 pt. Emollient, Tube 62.5 Gm.
 Use: Topically, skin cleanser.
- LOZILLES.** (White) A lozenge containing 2 mg. tyrothricin & 2 mg. propolis. Vial 15 Lozilles.
 Use: Infections of mouth & throat.
- LOZI-TARS.** (Hoyt) Sod. fluoride equiv. to 1.0 mg. of fluoride/lozenge. Bot. 100s, 1000s.
 Use: Prevention of dental caries
- LUASMIN.** (Brewer) Theophylline Sod. acetate 3 gr., ephedrine sulfate 1/2 gr., Sod. phenobarbital 1/2 gr./Cap. or E.C. Tab. Also available in half strength. Cap., Bot. 30s, 100s, 1000s. Tab., Bot. 30s, 100s, 1000s.
 Use: Relief of bronchial asthma & other allergic states.
- LUAFAX.** (Burroughs Wellcome) Surgical lubricant, sterile, water-soluble, non-staining
 * Generic name recognized by N.F., U.S.P. or A.M.A.
- tube 2 oz., 5 oz., 5 Gm. single use tube 4Bs.
 Use: Surgical lubricant.
- LUBASPORIN.** (Burroughs-Wellcome) Polymyxin B sulfate 5000 u, benzalkonium Cl 330 mcg/Gm Tube 5 Gm, 12r w/opplr
 Use: Antiseptic lubricant.
- LUBATH.** (Texor Pharmacol) A dispersible oil for the bath. Bot. 8 oz., 1 pt.
- EURISMIN.** (Maltbie) Homatropine methylbromide 1/4 gr., colloidal kaolin 60 gr., bismuth subcarb. 30 gr., pectin 5 gr./fl. oz. Bot. 4 oz., 1 pt., 1 gal.
 Use: Diarrhea, dysentery.
- LUBRIDERM.** (Texas Pharm.) Lanolin (oxycholestrin), earbital, cetyl alcohol, mineral oil & cholestrin. Emultran R ex., & New Pump Bot. 8 oz., 16 oz.
 Use: Topically, protective.
- LUBRIDERM-AC.** (Texas Pharm.) Emulsion of oxycholestrin, mineral oil, sorbitol, cetyl alcohol, cetyl sulfate, stearyl sulfate & lauryl sulfate. Bot. 1/2 & 1 pt.
 Use: Topically; protective.
- LUBRIDERM CREAM.** (Texas) Refined derivatives of lanolin, sorbitol, cetyl alcohol, petrolatum blend & purified water in heavy white cream./jar 4 oz.
 Use: Relief from dry scaly skin.
- LUCAINE HYDROCHLORIDE.** (Maltbie) Pridocaine HCl, Amp. 20 & 30 mg. Box 12s
 Use: Spinal anesthesia.
- * LUCANTHONE HCl.** (Burroughs Wellcome) 1-β-diethylaminoethylamine-4 methyl thioxanthone HCl 200 mg /Tab. Bot. 30s.
 Use: Oral treatment of schistosomiasis
- LUCIDON.** (Lloyd, Dabney & Westerfeld) Pentylene tetrazol 100 mg, niacin 50 mg, dienestrol 0.05 mg., methyltestosterone 1 mg /Tab. Bot. 100s.
 Use: Anabolic stimulant.
- LUCORTEUM, NND.** 1961. (March Sharp & Dahms) Progesterone. Vial (23 mg/cc.) 5 cc.
 Use: Progestational therapy.
- LUTA.** (U.S. Vitamin Corp.) Unsaturated fatty acid 378 mg, pyridazine HCl 2 mg, choline bitart. 233 mg., dl methionine 110 mg., inositol 40 mg., desic. liver 87 mg., Vit. B-12 1 mcg and Vit. E 3.5 u./Cap. Bot 500s, 1000s
 Use: (1) (Ld) Atherosclerosis
- LUGOL CAPS.** (Burnham) Iodine 40 mg./Cap. Bot. 21s, 100s
 Use: Iodine therapy.
- LUGOL'S SOLUTION, STRONG IODINE SOLUTION, U.S.P.**
 See: Cape-Lugol's, Cap. (Free)
 Lugol Caps. (Burnham)
- LUKESTRA SOLUTION.** (March Sharp &

Oshme) Estradiol benzoate 1.67 mg, progesterone 25 mg, testosterone propionate 25 mg/cc Vial 10 cc, rubber capped Vial 5 cc & 10 cc rubber capped

Use: Hormonal therapy

LULLAMIN DROPS (Reed & Carnrick)
Methoprylone HCl 16 mg/cc Bot. 25 cc

Use: Pediatric sedative

LUMASPIRIN W/HYOSCYAMUS. (Rorer)
Aspirin 5 gr, phenobarbital 1/4 gr, hyoscyamus ext 1/4 gr/Tab Also available w/cadaverin Bot 100s

Use (1 qid) Analgesic & sedative

LUMBRICIDE (Massengill) Santonia 1/6 gr, spigelia 2.6 gr, frangula 80 mg, senna 80 mg/5 cc Syr, Bot 1 pt, 1 gal

Use: Orally (12 dr tid for 3 days) anthelmintic

LUMIFLORA COMPOUND (Buffington) Phenobarbital 1/4 gr, hyoscyamus ext 1/25 gr, passiflora 1 gr, valerian 1 gr/Tab Bot 100s 1000s

Use Sedation

LUMINAL, NNR 1949, ADR (Winthrop Labs) Phenobarbital Phenylethylbarbituric acid

Cap (1 1/2 gr) Bot 100s

Ellixir (1/4 gr/4 cc) Bot 12 fl oz, 1 gal

Powr Bot 1/2 & 4 oz

Tab (1/4 & 1/2 gr) Bot 100s, 1000s (1 1/2 gr) Bot 50s 100s, 1000s

Use Long acting sedative & hypnotic

W/Aspirin See Lumalgin Tab (Winthrop Labs)

W/Atropine sulfate scopolamine HBr & Vit B complex See Apalamin Tab (Winthrop Labs)

W/Col tressal sulfonates See Kres Lumin, Syr (Winthrop Labs)

W/Bephedrin & theophylline See: Franol Tab (Winthrop Labs)

W/Isuprel ephedrine sulfate theophylline potassium iodide, alcohol See Isuprel Compound (Winthrop Labs)

W/Scopolamine atropine sulfate benzocaine pyridoxine nicotinamide, riboflavin See: Apalamin Tab (Winthrop Labs)

W/Theobromine & rauwolfia serpentina See Theaminol RS Tab (Winthrop Labs)

LUMINAL OVOIDS. (Winthrop Labs) Phenobarbital Ovoids (1/4 gr yellows 1/2 gr light green 1 1/2 gr dark green) Bot 100s 1000s

LUMINAL SODIUM, NNE 1952 ADR 1955 (Winthrop Labs) Phenobarbital Sodium Ampine (5 gr) 2 cc. Box 5s 100s Amp (2 gr & 5 gr) Box 5s 25s 100s Tab (1/4 gr, 1/2 gr & 1 1/2 gr) Bot 50s 1000s Tab, NY (1 gr) Bot 50s 500s

Use: Sedative & hypnotic

LUMINAL SODIUM IN PROPYLENE GLYCOL (Winthrop Labs) Amp. (130 mg./cc.) Box 100s

Use: I.M.

LUNARGEN. (Lilly) Mild protein silver, colloidal Pow, Pkg 1 oz, 4 oz, 1 lb, 5 lb, Cap (0.389 Gm) Bot 50s

Use: Topically, silver antiseptic

LUNOSOL (LIQUID), NNR 1950 (Hille) Colloidal silver chloride liq, Bot w/drop per 1/2 fl oz, 2 fl oz; Bot 1 fl oz, 4 fl oz. 1 pt Oint, Tube 1/2 oz, 1/2 oz

Use: Topically, silver antiseptic

LURIDE. (Hoyl) Sod fluoride equiv to 0.1 mg al fluoride/drop Plastic dropper bottle 40 ml

Use Prevention of dental caries

LUSYN. (Mallinckrodt) Homatropine methylbromide 5 mg, alukalin (activated kaolin) 300 mg, & phenobarbital 15 mg/Tab Bot 100s 500s, 1000s

Use: Gastro-intestinal disorders

LUSYN IMPROVED (Mallinckrodt) Homatropine methylbromide 5 mg, phenobarbital 1/4 gr, alukalin 300 mg/Tab Bot 100s, 500s 1000s

Use (1 2 tid) Antispasmodic & sedative

LUTEIN, NNR 1931 (Hynson, Westcott & Dunning) Corpus luteum 2 dg/cc Amp 1 cc, 100s 3 cc, 100s

Use I.M. corpus luteum therapy

LUTEOGAN See Progesterone (Various Mfr)

LUTEOSAN. See Progesterone (Various Mfr)

LUTEOTROPHIN. (Squibb) Prolactin A highly purified, freeze dried luteotropic hormone conc from sheep pituitary glands Vial 5 cc, 1000 IU w/sterile diluent

Use: I.M., functional uterine bleeding to stimulate lactation

LUT-EST-EL (Elder) Estrogen natural 2 mg progesterone 20 mg/cc Aq Susp Vial 10 cc

Use I.M. estrogen w/corpus luteum therapy

LUTENIZING NORMONE. W/Chronic Gonadotropin See Kuarin Vial (Kremers Urban)

LUTORYL, NND 1959

LUTOCYLIOL (Ciba) Ethisterone Tab (5 mg-10 mg, 25 mg) Bot 20s, 40s, 100s Linglets (10 mg) Bot 30s 100s

Use: Orally corpus luteum therapy

LUTREN See: Progesterone (Various Mfr)

LUTREXIN, NND 1962 (Hynson Westcott & Dunning) Lututrin, water soluble protein like uterine relaxing factor obtained from ovary Tab (2000 u) Bot 25s

Use: (2-4 q 3-4 h) premenstrual tension dysmenorrhea & threatened abortion

- W/Equine estrogens. See: *Trexinest*, Tab. (Hynson, Wescall & Dunning)
- LUTROMONE**, N.N.D. 1962. [Endo] Progesterone.
Amp: (1 & 2 I.U.) 1 cc., Box 6s, 12s, 25s, 100s; (5 I.U.) 1 cc., Box 3s, 6s, 12s, 25s, 100s; (10 I.U.) 1 cc., Box 3s, 6s, 12s, 25s Vial. (1 & 2 I.U.) 10 cc; (5, 10 & 25 I.U./cc) 5 cc.
Use: I.M., *carpus luteum* therapy.
- LUTUTRIN**, N.N.D. 1962. See: *Lutrexin* (Hynson, Wescall & Dunning)
- LYBETA**. (Columbus) L-lysine monohydrochloride 300 mg., Vit. B-12 25 mcg., B-1 10 mg., B-6 5 mg./3 tab. Bot. 100s.
Use: Dietary supplement
- LYCOAN**. (Schwarz) Adenine 5' phosphoric acid 20 mg./Tab. Bot. 20s, 50s Amp. 1 cc. Box 12s.
Use: Nucleic acid deficiency.
- LYCINATE VAGINAL**. (Lloyd Bros) Di-iodo-hydroxyquin 100 mg. sod. lauryl sulfate 5 mg., dioctyl sodium sulfosuccinate 5 mg., aluminum potassium sulfate 14 mg., lactose 380 mg., dextrose 650 mg./Tab. Bot. 50s w/applicator.
Use: Inserted high in vaginal vault, morning and night for treatment of leukorrhoea
- LYCINE HYDROCHLORIDE**. See: *Betaine Hydrochloride* (Various Mfrs.)
- LYCOL ELIXIR**. (Lonnell) Glycolal 28 gr., lysine 100 mg./16 cc. Bot. 1 pt.
Use: Anorexia.
- LYCOPLEX**. (United Lab) liver inj. crude U.S.P. 1 mcg., Vit. B-12 10 mcg., Cal. pantothenate 10 mg., Vit. B-6 10 mcg./cc. Vial 10 cc., Box 1s, 6s,
Use: Treatment of Anemias.
- LYCORAL**. (Fellows) Chloral hydrate 10 gr./dr. liq. Bot. 1 pt., 1 gal.
Use: Sedative and hypnotic
- LYGEL VAGINAL JELLY**, N.N.R. 1955 (Lehn & Fink) Tube 92 Gm. Tube with applicator
- LYGENES VAGINAL SUPPOSITORIES**, N.N.R. 1955. (Lehn & Fink) Zinc phenolsulfonate, oxyquinoline benzoate, phenol derivatives, Cocoa butter base Box 12s
Use: Contraceptive
- LYGRANUM C.F. TEST**. (Squibb) Antigen prepared from the causative organisms (*Miyagawanella lymphogranulomatis*) of lymphogranuloma venereum. Vial 5 cc. & amount sufficient for testing about 50 sera.
Use: Diagnosis of lymphogranuloma venereum.
- LYGRANUM S.T.** (Squibb) Purified & potent antigen prepared from lymphogranuloma venereum organisms grown in chick embryos. Vial: Single test & in 10 test pkg

* Generic name recognized by N.F., U.S.P. or A.M.A.

- Use: Skin test diagnosis of lymphogranuloma venereum.
- LYMPHOID TISSUE EXTRACT**. See: *Hemostasin*, Vial (United Lab).
- *LYNDESTRENOL**. (Organon, Inc) Ethinyl oestrol.
Use: Pre-gestational and ovulation inhibiting action.
- LYNORAL**, N.N.D. 1962 (Organon) Ethinyl estradiol, Elixir (0.03 mg./4 cc.) Bot. 4 as., 1 pt., 1 gal. Tab. (0.01 mg.) Bot. 100s, 1000s; (0.05 mg.) Bot. 30s, 60s, 250s, 1000s.
Use: Oral estrogen
- LYO B-C (W/B-12)**. (Merck Sharp & Dohme) Vit. B-12 1 mg./cc. w/above ingredients, Vial 7 cc. With or without diluent
- LYO B-C (W/DILUENT)**. (Merck Sharp & Dohme) Thiamine HCl 100 mg., riboflavin 10 mg., pyridoxine HCl 10 mg., panthelol (equiv. to 58 mg.), cal. pantothenate 50 mg., niacinamide 250 mg., ascorbic acid 200 mg./7 cc. Also available without diluent, Vial 7 cc.
Use: Vitamin B complex & ascorbic acid deficiency states.
- LYO B-C FORTE (W/DILUENT)**. (Merck Sharp & Dohme) Thiamine HCl 100 mg., riboflavin 10 mg., pyridoxine HCl 10 mg., panthelol (equiv. to 58 mg.), cal. pantothenate 50 mg., niacinamide 500 mg., ascorbic acid 500 mg./7 cc. Also available without diluent.
- LYOCYTE**. (Merck Sharp & Dohme) Human blood cells dried Pow. Bot. 6 Gm.
Use: Aid to tissue growth & regeneration
- LYOPHRIN**. (Alcon Labs, Inc) 1-Epinephrine bitartrate 2%, sodium bisulfite 0.2%, chlorobutanol, 0.15%, boric acid, diadium ethylene diaminetetra-acetate, distilled water./Vial sterile lyophilized 1 epinephrine bitartrate (100 mg.), Bot. 5 cc. Lyophrin diluent.
Use: Chronic simple glaucoma.
- LYOVAC DIURIL**, N.N.D. 1962.
- LYOVAC NORMAL HUMAN PLASMA**. (Merck Sharp & Dohme) Irradiated normal human plasma. Vial 2.5 cc.
Use: Ophthalmic.
- LYSIDIN**. Methyl glyoxalidin. See: *Lyanthine* (Gallio)
- LYSIDOX**. (Gray) L-lysine HCl 250 mg., pyridoxine HCl 2 mg., milk powder defatted 12 Gm./pkg. Pow. Pkg 15s.
Use: Nutritional supplement.
- LYSI-GERIATRIC CAPS.** (Nyco Lab.) L-lysine HCl 50 mg. with 8 vitamins, 8 minerals, rutin 10 mg., liver 50 mg., yeast 15 mg. & fipatropic agents/2 tabs. Bot. 1000

LYSINE (Various Mfr)
Use Nutrient

LYSINE DROPS (Walker) 200 mg/cc Bot 30 cc

L-LYSINE W/Glycocal

See Lixacol Elixir (Lannett)

Lycal Elixir (Lannett)

L-LYSINE HCl W/Ferrous fumarate, Vit B 6
See Enisyl w/Iron Tabs (Luka)

L-LYSINE HCl W/Pyridoxine HCl & milk powder defatted See Lysidox Pow (Gray)

LYSINE THERAPY. See Stimavit, Tab (Roerig)

LYSIVANE. See Parsidal Tab (Warner Chilcott)

***LYSOSTAPHIN** (Mead Johnson) Antibiotic derived from *Staphylococcus staphylolyticus*

Use Antibiotic

LYTECA SYRUP (Lloyd Dabney & Westler field) N acetyl p aminophenol 120 mg / 5 cc Bot pts

Use Analgesic & antipyretic

LYTECA TABS (Lloyd Dabney & W) Acetaminophen 300 mg / Tab Bot 100s

Use Analgesic

LYTEERS (Barnes Hind) Isotonic viscous liquid adjusted to pH of tears Plas Bot 2 oz

Use Ophic

LYTREN (Mead Johnson) A complete electrolyte mixture: Dextri maltose 90.4% Sod citrate 3.6%, Pot chloride 1.9%, Sod phosphate 1.8%, Cal lactate 0.8% Sod chloride 0.8%, citric acid 0.4% Mag sulfate 0.3% Con 8 oz

Use Prep sol 1 tbsps to 4 oz water; give orally enough to meet fluid requirements

LYXANTHINE. (Gallia) Sod iodopropion sulfonate 12% lysidin (methyl glyoxalidin) bitartrate 9%, Cal gluconate 12%, Sod bicarb 37%, tartaric acid 15%, citric acid 6%, Sugar 9% Effer granules, Bot 60 Gm

Use: Arthritis gout & neuritis

MAALOX. (Rorer) Colloidal susp of Mag & aluminum hydroxides Susp Bot 12 oz Tab Bot 100s Double strength tab Bot 50s & 250s

Use: Gastric antacid

W/Aspirin See: Ascriptin Tab (Rorer)

W/Iron sulfate See: Fermalox, Tab (Rorer)

MAALOX NO 2. (Rorer) New double strength tablet Bot 50s, 250s

Use: Antacid

MASUTONE. (Reed & Cornrick) Mephenesin 250 mg, d-amphetamine phosphate (dibasic) 2.5 mg, butabarbital 8 mg / Tab Bot 50s 100s Elixir Bot 8 oz

Use: (1 lid) Muscle relaxant w/sedative

* Generic name recognized by N.F., U.S.P. or A.M.A.

MACRODIOL

Kirk—Estradiol 0.55 mg/cc Vial 10 cc 30 cc

Gatham—Estradiol Vial (0.22 mg, 0.55 mg/cc) 30 cc

MACROGESTIN. (Kirk) Progesterone 25 mg / cc Vial 10 cc

Use Progesterone therapy

MACROGOL 400 See Polyethylene 400

MACROGOL 400D. See Polyethylene glycol 4000

MACRON TABLETS (Abbott) Ferrous sulfates 0.35 Gm thiamine HCl 2 mg, riboflavin 2 mg, nicotinamide 10 mg pyridoxine HCl 1 mg pantothenic acid (as Cal pantothenate) 2 mg ascorbic acid 50 mg, liver conc NF 0.5 Gm / Tab Bot 100s, 500s 1000s

Use Secondary & nutritional anemias

MADRIBON, NND 1962 (Eahe lab) Sulfadimethoxine 2.4 dimethoxy 6 sulfamidamide 1.3 diazine 0.5 Gm / Tab Bot 30s 100s 250s 1000s Pediatric Susp 0.250 Gm / 5 cc Bot 4 oz, 1 pt Pediatric Drops (250 mg / cc) plastic conta ner 10 cc

Use (2 doses then 1 q 24 hours) Bacterial infections

W/Theophorin tartrate N acetyl p aminophenol caffeine See Madricidin Cap (Eahe)

MADRIBON CHEWABLE TABLETS (Rahe Lab) 2.4 Dimethoxy 6 sulfamidamide 1.3 diazine 250 mg / Tab 0.25 Gm Bot 30s 100s

Use Bacterial infections

MADRICIDIN (Rahe) Madribon 125 mg theophorin tartrate 10 mg, N acetyl p aminophenol 120 mg caffeine 30 mg / Cap Bot 100s 1000s

Use Therapy for the common cold

MADRIDOID, NND 1962 (Eahe) 2.4 Dimethoxy 6 sulfamidamide 1.3 diazine 125 mg / Cap Bot 100s 1000s

Use Upper respiratory infections

MAFENIDE HYDROCHLORIDE (SULFABENZAMINE). W/Streptomycin See: Sulfamylan Sol (Winthrop Labs)

MAGALIN. (Kendall) Mag gluconate 7 1/2 gr, phenobarbital 16 mg dl desoxyephedrine HCl 5 mg / Tab Bot 100s

Use: (1 lid) Premenstrual tension

MAGANHIST. (Marlo) Pyrilamine maleate Tab Bot 50s 100s 1000s

Use: Antihistaminic

MAGCYL (Elder) see old name Magnocyl

MAGINAL (Sutthiff & Case) Aluminum hydroxide dried gel 2 gr, magnesium trisilicate 5 gr, kaolin 2 gr, methyl cellulose

- $1\frac{1}{2}$ gr., homatropine methylbromide 1/60 gr./Tab. Bot. 100s, 1000s
Use: Antacid
- MAGKAOLUM.** (Elder) Mag. trisilicate 4 gr., aluminum hydroxide 2 gr., kaolin 2 gr./Tab. Bot. 100s, 1000s.
Use: Orally (2-4 p.r.n.), antacid.
- MAGMA ALZINOX.** (Patch) Dihydroxy aluminum amino acetate 7.7 gr./5 cc. Bot. B oz.
Use: Orally, gastric antacid.
- MAGMASIL.** (Leeming) Mag. trisilicate 20%. Susp. Bot. 12 oz.
Use: Orally, antacid.
- MAGMOID SULCO.** (Pitman-Moore) Sulfamerazine 1 Gm., sulfadiazine 1 Gm., sulfathiazole 1 Gm./fl. oz. Susp. Bot. 2 fl. oz., 12 fl. oz., 1 gal.
Use: Triple sulfonamide therapy.
- MAGNACILLIN.** (Standard Pharm.) Peccoine penicillin O 300,000 u., Pot. penicillin G 100,000 u./dose. Vial 1 dose.
Use: Penicillin therapy.
- MAGNACORT, N N D.** 1962. (Pfizer) (Hydrocortisone) Hydrocortisone diethylamineacetate HCl 0.5%. Oint. Tube 1/6 oz., 1/2 oz.
Use: Topically, skin inflammations
- W/Neomycin.** See: Neo-Magnecort (Pfizer)
- MAGNAMYCIN, N N D.** 1960. (Pfizer) Carbomycin. Tab. 100 mg. Bot. 25s, 100s; 250 mg. Bot. 16s.
Use: Orally, penicillin-resistant gram-positive infections.
- MAGNEPHYLLINE.** (Sherman) Theophylline 79%. Mag. 5.36%. Tab. $1\frac{1}{2}$ gr., 3 gr. Bot. 100s, 500s, 1000s.
Use: (1-2 i.i.d. or q.i.d.) Theophylline therapy.
- W/Phenobarbital.** (Sherman) Magephylline $1\frac{1}{2}$ gr., w/phenobarbital $\frac{1}{4}$ gr., $\frac{1}{2}$ gr./Tab. Bot. 100s, 500s, 1000s.
Use: (1-2 i.i.d.) Diuretic.
- *MAGNESIA MAGMA, A D R, U S P.** Milk of Magnesia. See: Magnesium Hydroxide, Prep.
- MAGNESIUM ACETYSALICYLATE.** (Apyron, Magnesperin, Magisal, Novacetyl.) See: Yimmary
Use: Like acetylsalicylic acid.
- MAGNESIUM ALUMINATE HYDRATE.** See: Riapan, Susp. (Ayerst)
- MAGNESIUM ALUMINUM HYDROXIDE.** See: Maalox, Susp. (Ries))
- MAGNESIUM ALUMINUM HYDROXIDE AMINOACETATE.** See: Glycamine-AM, Tab. (Hermelin)
- MAGNESIUM ASPARTATE.** W/Potassium aspartate. See: Spartox, Tabs (Wyeth).
- MAGNESIUM CARBONATE** W/Amino acetic acid, aluminum hydroxide, calcium carbonate See: Glytabs (Dereist)
- MAGNESIUM GLUCONATE.** See: Almaro, Tab. (Durst).
GYN, Tab. (Grant)
Menesia, Tab. (Pitman-Moore).
Relaxin, Tab. (Dumas-Wilson).
W/Phenobarbital & dl-desoxyephedrine HCl. See: Magalin, Tab. (Kendall).
- MAGNESIUM GLYCINATE** W/Gastric mucin & aluminum hydroxide See: Mucagel, Tab. (Barrows Chemical Co)
- *MAGNESIUM HYDROXIDE, N F; Tab., N F** See Magnesia Magma, U.S.P.
W/Aluminum hydroxide
See: Almgah (Hoag)
Aludrox, Susp. (Wyeth)
Bidrox, Susp. (McNeil)
Creamlets, Tabs. (Winthrop Labs)
Maalox, Susp. (Ries)
W/Aluminum hydroxide, scopolamine HBr, l-hyoscyamine HBr, bulkagen. See: Hydroxox, Tab. (Columbus)
W/Aluminum hydroxide, scopolamine methylnitrate, 1-piperidine ethanol benzilate HCl. See: Madutrol, Tab (Reed & Carnrick).
W/Carboxymethylcellulose Sod. See: Turicum, Susp. (Whittier).
W/Danthron, prvne concentrate powder. See: Danthro-Lax, Tab. (Molard)
W/Dicyclomine HCl, alum. hydroz, magn. trisilicate, sodium lauryl sulf., methylcellulose See: Kalantyl Waters (Merrell)
W/Dihydroxyaluminum aminoacetate, meth-scopolamine bromide & sod. lauryl sulfate. See: Alu-Scop, Cap., Susp. (Lloyd, Dabney & Westerfield)
W/Mineral oil.
See: Haley's M.O., liq. (Phillips).
Magaleum, liq. (Blue line)
W/Phenolphthalein, yellow. See: Colovac, Tab. (McNeil)
W/Scopolamine & aluminum hydroxide See: Ampyrox, Tab #293. (Elder)
- *MAGNESIUM OXIDE, U S P.** W/Acacia & veg. mucinoid. See: Acarasil, Pow. (Sherman).
- W/Aluminum hydroxide glycine. See: phtan, Tab. (Sandaz)
W/Guar cellupectinoid, aluminum hydroxide gel, magnesium trisilicate. See: Belvis, Tabs. (Cornick)
- *MAGNESIUM PEMOLINE.** 2-imino-5-phenyl-4-oxazolidinone magnesium See: Cylert
- *MAGNESIUM PHOSPHATE, TRIBASIC, N F., N N R** 1947. Tab. N F.
- MAGNESIUM SULFATE, U S P.; Inj., N F.**

- Epsom salt (Various Mfrs) Pow 1 lb Sol
 Abbott—10% Amp 20 cc (50%) 2 cc
 Baxter—10% Vial 10 cc, 20 cc
 Ingram—50% Vial 2 cc
 Lilly—10% Amp 20 cc; 25% Amp 2 cc;
 25% Amp 10 cc; 50% Amp 2 cc
 Merrell—50% Amp 2 cc
 Parke, Davis—25% Amp 10 cc; 50%
 Amp 2 cc See Osmopak
- *MAGNESIUM TRISILICATE, USP, NH&
 1947, Tab, USP (Various Mfr)
 See Chooz Tab (Pharmacia Inc)
 Magmasil, Susp (Leeming)
 Salisal, Tab (Elder)
 Trimax, Pow (Winthrop Labs)
 Trinesium, Tab (Abbott)
 Tri Sil, Tab (Warren Teed)
 Trisamin, Cap (Lilly)
- W/Aluminum hydroxide
 See: Alinase, Susp & Tab (Haskell)
 Almag Tab & Cap (Drug Products)
 A M T, Susp & Tab (Wyeth)
 Bidiox Susp (McNeil)
 Ca Lu Gel M T, Susp (Ulmer)
 Fluogel Camp, Tab (Beech)
 Gelusil, Susp & Tab (Warner)
 Kamodiox, Jel (Massengill)
 Magal, Susp & Tab (Endo)
 Malcogel (Upjohn)
 Malcolabs (Upjohn)
 Mardazine, Tab (G F Harvey)
 Milk of Trinesium, Susp (Abbott)
 Neutrolan Composite Tab (Cansel Mid
 land)
 Silmagel, Tab (Lannett)
 Tri Cremalate, Liq & Tab (Winthrop Labs)
 Tridroxal Liq (Warren Teed)
 Trigelma Sesp & Tab (Burrington)
 Tisax, Sesp & Tab (Bee Line)
 Trisogel, Susp & Pulv (Lilly)
- W/Aluminum hydroxide & ascorbic acid
 See Cimadrox, Pow & Tab (Massengill)
- W/Aluminum hydroxide gel dried, predni
 sone phenylbutazone hamotrapine methyl
 bromide See Stiazolidin Cap (Geigy)
- W/Aluminum hydroxide gel & Col carb
 See: Alcarval Tab (Wellace)
 Gastralme Liq (Mayer Chem)
- W/Aluminum hydroxide chlorophyllin
 See Chloragel, Tab (Walker)
 Stawex Tab (Rchlyn)
- W/Aluminum hydroxide & gastric mucin
 See Almagusin NH&
 Mucatin, Liq & Tab (Harraver)
 Trimucatan, Tab (Winthrop Labs)
- W/Aluminum hydroxide gastric mucin
 hamotrapine methylbromide See: Gast Rex
 Tab (Money)
- * Generic name recognized by NF, USP or A.M.A.
- W/Aluminum hydroxide dried gel & kaolin
 coloidal
 See: Kamodiox, Tab & Pow (Massengill)
 Koa Magnin, Syr (Sutliff & Case)
 Silmocal, Cap, Tab & Pow (Amfco)
- W/Aluminum hydroxide dried gel kaolin
 methyl cellulose, hamotrapine methylbromide
 See Maginal, Tab (Sutliff & Case)
- W/Aluminum hydroxide & okra See Chlor
 xum Mucinoid (Rystan)
- W/Aluminum hydroxide & pectin See
 Alumapex Tab (Lemmon)
- W/Aluminum hydroxide phenylbutazone
 prednisone hamotrapine methyl bromide
 See Stiazolidin Cap (Geigy)
- W/Aluminum hydroxide, reganal esquire
 Cal carb & mag carb See Trevidal Tab
 (Organon)
- W/Aluminum hydroxide & reduced iron See
 Alinase Tab & Susp (Haskell)
- W/Ascorbic acid See Trismagnal, Tab
 (Patch)
- W/Aspirin, carlisono acetate See: Cortisol
 Tab (Lincoln)
- W/Atropine sulfate & phenobarbital See:
 Malrobarb, Tab (Smith Miller & Patch)
 Triphen Tab (Lannett)
- W/Atropine methyl nitrate, aluminum hy
 droxide duodenum pow See Melacin
 Tab (Strosenburgh)
- W/Belladonna alkaloids aluminum hydroxide
 benzocaine & phenobarbital
 See Alkaspamol Tab (Walker)
 Al Si Col, Tab (Smith Dorsey)
- W/Belladonna ext & aluminum hydroxide
 See Traldin, Tab (Raries)
- W/Belladonna ext & charcoal
 See Belladonna Trisilicate & Charcoal, Tab
 (Raymer)
 Trisidanno Tab (Rarer)
- W/Betabarbital Sod, methacapalamine bromide
 & aluminum hydroxide gel dried See:
 Eelcin, Tab (Staff Research Labs)
- W/Betabarbital, aminophylline phenylpro
 panalamine HCl, chlorpheniramine maleate
 aluminum hydroxide See: Amocal, Tabs
 (Vale)
- W/Betazolodin (phenylbutazone) aluminum
 hydroxide, hamotrapine methylbromide
 See Rutaxolidin Alka Cap (Geigy)
- W/Col carb See Traco-Discs (Neoca)
- W/Col carb & Mag carb
 See: Calinase Tab (Haskell)
 Jests, Tab (Jests Inc)
 Neutralizer Water (Upjohn)
 Parluguan, Trache (Panray)
 Tricarb, Tab (Raymer)
 Tricarbs, Tab (G F Harvey)
 Temx Tab (Howe Co)

- W/Homatropine MB & Aluminum hydroxide.
See: Trigelma, HMB (Burlington)
Tropasil, Tab. (Crookes-Barnes)
- W/Homatropine methylbromide, aluminum hydroxide gel, dried, Mag. hydroxide & pectin. See: Gel-A-Nutrin, Tab. (Walker)
- W/Homatropine methylbromide, aluminum hydroxide & pectin. See: Alucen, Tab. (Centrol)
- W/Homatropine methylbromide, aluminum hydroxide, phenobarbital.
See: Alased, Tab. (Nargine)
Neutrolabs, Tab. (Blue Line)
- W/Homatropine MB, & pentobarb. See: Homopent, Tab. (Lemman)
- W/Methylcellulose, d-amphetamine sulfate, phenobarbital & vitamins. See: Nexarin, Cap. (Smith-Dorsey).
- W/Methscopolamine bromide, phenobarb. col. carb., mag. oxide. See: Scopamol, Tab. (Miller)
- W/Metrolase, duodenum pow. & aluminum hydroxide. See: Metrocin, Tab. (Strosenburgh).
- W/Neojel. See: Neo-Gel Susp. & Tab. (Diamond)
- W/Phenobarbital sod., homatropine methylbromide, gastric mucin, aluminum hydroxide. See: Antased, Tab. (Lemman)
- W/Phenobarbital & atropine sulfate.
See: Trilaf, Tab. (Warren-Teed)
Trilaphen, Tab. (Lannett)
- W/Phenobarb. atropine sulf., & hyoscyamine see: Atromag, Tab. (Lemman)
- W/Phenobarb. atropine sulf. & hyoscyamine. See: Silalaid, Tab. (Van Peft & Brown)
- W/Phenobarbital, theophylline, ephedrine HCl, belladonna leaves, pyrilamine maleate, potassium iodide See: Theated-Pediatric, Tab. (Suliff & Case)
- W/Prednisalone & aluminum hydroxide gel.
See: Co-Hydeltro, Tab. (Merck Sharp & Dohme)
- W/Prednisone & aluminum hydroxide gel.
See: Co-Deltro, Tab. (Merck Sharp & Dohme)
- W/Scopolamine methylbromide & aluminum hydroxide gel. See: Trigelma w/Mescamina, Tab. (Burlington)
- W/Sippyloc, aluminum hydroxide, vit. B-1, B-2, B 6, C, Col. pantathenate, niacinamide, menadione. See: Sippyplex, Pow. (Furdue Frederick)
- MAGNICAPS.** (Plessner) Vit. A 25,000 u., 1000 u., thiamine HCl 10 mg., riboflavin 5 mg., ascorbic acid 150 mg., niacinamide 150 mg./Cap. Bot 100s, 1000s
Use: Vitamin deficiencies
- MAGNICAPS-M.** (Plessner) Vit. A 25,000 u., D 1000 u., B-1 10 mg., B-2 5 mg., B-6 1 mg., B-12 5 mcg., C 150 mg., niacinamide 100 mg., with trace minerals/Cap. Bot. 100s, 500s, 1000s
Use: Therapeutic multivitamins with B-12 and minerals
- MAGNOCYL.** (Elder) new name Magcyl Propethyleneoxides 250 mg./Cap. Bot 100s.
Use: Stool softener.
- MAGOLEUM.** (Blue Line) Mineral oil 50%, milk of Mag. 50%. Bot. 8 oz.
Use: Orally, laxative.
- MAGSAL.** (Endo) Aluminum hydroxide 10 gr., Mag trisilicate 1½ gr./Tab. or dr. Tab. Bot. 100s, Liq., Gel., 12 oz. 1 gal.
Use: Orally, antacid.
- MAGSOBBAL.** (U. S. Standard) Aluminum hydroxide gel dried 400 mg., Mag. trisilicate 400 mg., Mag. oxide heavy 35 mg./Lax. Bot. 100s, 1000s. Also avail. in susp.
Use: Hyperacid stomach.
- MALAGBIDE.** See: Acetarsona, Tab. (Abbott)
- MALCOGEL.** (Upjohn) Aluminum hydroxide 3½ gr., Mag trisilicate 7½ gr./R. dr. Bot. 12 oz., 32 oz.
Use: Orally, antacid
- MALCOTABS.** (Upjohn) Aluminum hydroxide 10 gr., Mag. trisilicate 5 gr./Tab. Bot. 100s, 500s, 1000s.
Use: Orally, antacid.
- MALCOTABS W/BELLADONNA.** (Upjohn) Malcotabs ½ strength w/belladonna exl. 1/16 gr./Tab. Bot. 100s, 500s, 1000s.
Use: Orally, antispasmodic w/antacid.
- MALCOTRAN.** (Maltbie) Homatropine methylbromide. Tab. 10 mg. Bot. 100s, 500s, 1000s.
Use: [1 q. 3 h] Anticholinergic & ganglionic blocking agent.
- W/Phenobarbital. (Maltbie) Malcotran 10 mg., phenobarbital 8 mg./Tab. Bot. 50s, 500s, 1000s.
Use: Psychosomatic patient.
- MALCOTRAN W/PHENOBARBITAL PEDI-ATRIC ELIXIR.** (Maltbie) Homatropine methylbromide 2.5 mg., phenobarbital 8 mg./5 cc. Bot. 4 oz., 1 pt., 1 gal.
- MALESTRONE.** [N.N.O. 1962. (Kirk) Testosterone aq. susp. Vial. {25 mg/cc} 10 cc; {50 mg/cc} 10 cc
Use: I.M., androgen therapy.
- MALESTRONE TABLETS, N.N.R.** (Kirk) Methyl testosterone. Tabs. 10 mg., 25 mg., Bot. 1000s
Use: Hormone therapy.
- MALGLYN COMPOUND.** (Brayten) Dihydroxy aluminum aminoacetate 0.5 Gm., belladonna alkaloids (as sulfates) 0.162 mg., phenobarbital 16.2 mg./Tab. Bot. 100s.
Use: [1-2 t.i.d.] Peptic ulcer.
- MALGLYN COMPOUND MAGMA.** (Brayten)

- Each 5 cc is equiv to 1 Maltglyc Compound Tablet Bot B oz
- MALIRUATIN** (Smith Dorsey) Vit A 20000 u D 2000 u, thiamine HCl 2 mg, riboflavin 2 mg niacinamide 40 mg, iron & ammonium citrates 4 gr /fl oz Bot 24 oz, 48 oz
Use Detary supplement
- MALLOPNE** (Mallinckrodt) 3 phenylazo 2,6 diaminopyridine monohydrochloride Pow, Bot 1 Gm 5 Gm Soluble discs 0.1 Gm, Bot 50s, Tube 10s Tab 0.1 Gm, Bot 25s 50r 500r
Use Orally; irrit llation & topically urinary antiseptic
- MAL-O-FEM** (Testagar) Aq—Testosterone 25 mg estrane 2 mg/cc Vial 10 cc 5m Ori—Testosterone propionate 25 mg estradiol benzoate 1 mg/cc Vial 10 cc
Use IM, estrogen and androgen therapy
- MALONE INJ** (Testagar) Testosterone Aqueous—[25 mg/cc] 10 cc 30 cc; [50 mg/cc] 10 cc; [100 mg/cc] 10 cc
Use IM male hormone therapy
- MALONE TABLETS**, (Testagar) Methyl testosterone 10 mg/tab Bot 100s 1000s
Use Orally male hormone therapy
- MALONAL**, Seer Barbitol (Various Mfr)
- MALONIC ACID W/Oxalic acid inj** See Kaogamin, Vial (Chatham)
- MALRYN** (Bernhoff Labs) Tabs 100s
- MALT EXTRACT, NF** (Borchardt) Malted barley ext containing maltose dextrins protein & minerals w/Vit B complex Plain Bot 1 pt W/Cod liver oil Bot 1 pt
Use Nutrient
- MALT EXTRACT & NYPOPHOSPHITES COMPOUND** Seer Arcadal Liq (Merck Sharp & Dahme)
- MALTINE PLAIN** (Warner Chilcott) Malted barley ext Bot 1 pt 1 gal
Use Detary supplement
- W/Cascara Sagrada** (Warner Chilcott) Cascara sagrada ext 2.9 Gm/fl oz Bot T pt 1 gal
- W/Cod liver oil** (Warner Chilcott) Maltine 70% cod liver oil 30% Bot T pt 1 qt 1 gal
- W/Creosote** (Warner Chilcott) Creosote D 25 cc/fl oz Bot 1 pt 1 gal
- W/Iron** (Warner Chilcott) Iron pyrophosphate 3.9 Gm/fl oz Bot 12 fl oz 1 gal
- W/Vitamin conc** (Warner Chilcott) Vit A 10000 u D 1000 u, B 1.3 mg B 2.4 mg niacinamide 40 mg pantothenic acid D 35 mg, dicalcium phosphate TT Gm/fl oz Bot 12 fl oz, 1 gal
- MALTO-FERRO** (Tilden) Iron and ammonium citrates 80 gr/fl oz Bot pts, gals
Use: Treat secondary anemias
- MALTOSE W/DEXTRINS**. See: Alerdex, Pow (Wyeth)
- MALTRATE**, (Mallard) Mannitol hexanitrate Bot 100s, 500s 1000s
W/Phenobarbital (Mallard) Mannitol hexanitrate 1/2 gr, phenobarbital 1/4 gr/Tab. Bot 100s, 500s, 1000s
Use: Vasodilator w/sedative
- MALT SOUP EXTRACT, NNR 1932** (Borchardt) Non drastic malt ext neutralized w/Pot carb laxative modifier or milk for constipated babies Effective in constipation in pre school children & elderly patients Liq, Bot 8 oz, 1 pt Pow, dry, Can 1 lb, Chocolate flavor, B oz
Use Orally laxative and pruritus ani
- MALT-C** (Lincoln Labs) Sodium ascorbate 75 mg, magnesium trisilicate 200 mg, aluminum hydroxide gel, dried 100 mg, homaloprine methyl bromide 2.5 mg, sodium lauryl sulfate 20 mg/Cop Bot 100s
Use Pepsic ulcer and hyperacidity
- MALUMEL**, (Rawell) Aluminum phosphate 200 mg magnesium glycinate 200 mg/Tab Bot 100s 1000s
Use Antacid
- MAMLENE OINT.** (Ulmer) Bismuth subnitrate castor oil, peru balsam Tube 1 oz
Use Nipple ointment
- MAMMARY GLAND W/Carpus luteum** Seer Frensil, Liq (Wallow)
- MAMMOL OINTMENT** (Abbott) Bismuth subnitrate 40% castor oil 30%, wool fat 22%, ceresin wax 7%, balsam peru 1% Tube 1/2 oz Box 12s
Use: Dressing for nipples of nursing mothers
- MAMOLIENT** (Smith Dorsey) Bismuth subnitrate 40%, balsam peru 1%, benzoin T% Tube 2 oz
Use Dressing for cracked nipples of nursing mothers
- MANADRIN**, (Endo) Isotonic sol of ephedrine Bot w/dropper 1 fl oz, Bot 1 pt, T gal
Use Vasoconstrictor
- MANAKALI** (Trident) Sod salicylate 40 gr, Pot iodide 8 gr, calchicum carb fl ext 8 mm manaca fl ext 20 mg, gelsemium fl ext 8 mm/fl oz Bot 1 pt 1 gal Also available w/haud calchicum
Use: Pain in rheumatism, arthritis & gout
- MANARTAL** (Abbott) Nat lsted Mannitol hexanitrate 30 mg phenobarbital 15 mg/Tab Bot 100s 1000s
Use (1 2 tr d) Vasodilator w/sedative
- MANDECIOLOR** (Vanpel & Brown) Ammonium mandelate Sod mandelate ammonium chloride Elixir, Bot 8 oz, 1 gal
Use: Orally, urinary antiseptic.

MANDELAMINE, N N D. 1962. (Warner-Chilcott) Enteric coated Tab. contains D 25 Gm (3½ gr) methanamine mandelate. Bot. 120s, 500s, 1000s. Susp. (250 mg / 5 cc.) (sesame oil) Bot. 4 oz.

Use: (3-4 t.i.d.) Urinary antiseptic.

MANDELAMINE "HAFGRAMS." (Warner-Chilcott) Methanamine mandelate 0.5 Gm./Tab. Bot. 100s, 500s, 1000s.

Use: Urinary antiseptic.

MANDELAMINE "UNOGRAM." (Warner-Chilcott) Methanamine mandelate 1 Gm./Tab. Bot. 100s

MANDELIC ACID, N F., N.N.R.

Mallinckrodt—Pow., Bot. ¼ lb., 1 lb., 5 lb.

Merck—Pow., Bot. 1 oz., ½ lb., 1 lb.

Use: Orally, urinary antiseptic.

W/Sulfacetamide & methanamine. See: Finestul, Tab. (Kendall).

MANDELIC ACID & AMMONIUM CHLORIDE, (Lilly) Monoethanolamine mandelate 5 gr., ammonium chloride 3½ gr./Enseal. Bot. 100s, 500s, 1000s.

Use: Orally, urinary antiseptic.

MANDELIC ACID ELIXIR, (Lilly) Ammonium mandelate 8 Gm./fl. oz. Elixir, Bot. 1 pt.

Use: Orally, urinary antiseptic.

MANDELIC ACID AS MONOETHANOLAMINE W/AMMONIUM CHLORIDE. See: Mandelic Acid & Ammonium Chloride, Enseal (Lilly)

MANDELIC ACID SALTS.

See: Ammonium mandelate.

Calcium mandelate.

Mandeliclar (Vompelt & Brown)

MANDELYTROPEINE. See Hamatropine (Various Mfr.).

MANDEX. (Vale) Methanamine mandelate 250 mg, salicylamide 120 mg, extract belladonna 5 mg./Tab. Bot. 100s.

Use: Urinary antiseptic.

MANEXIN. (Massengill) Mannitol hexanitrate ½ gr./Tab. Bot. 100r, 1000s

Use: (1-2 t.i.d.) Hypertension & far angina pectoris.

MANEXINE. (Cole) Phenobarbital 16 mg., mannitol hexanitrate 16 mg, theobromine 162 mg./Tab. Bot. 100r, 300r, 1000s.

Use: (1 q.i.d.) Hypotenrive.

MANGALAC. (Endo) Milk protein, w/Mang. butyrate. Vial 1 cc., 2 cc., 25 cc.

Use: I.M. (1 cc. daily), nonspecific protein therapy.

MANGANESE BUTYRATE W/Carsein. See Mangalac, Amp. (Endo)

MANGANESE HYDROXIDE COLLOIDAL. See: Colomang, Amp. & Vial (Crookes-Barnes)

MANHEXIN. (Normand) Mannitol hexanitrate ½ gr./Tab. Bot. 100s, 500s, 1000s

Also available with phenobarbital ¼ gr./Tab.

Use: Vasodilator

MANIBEE AMPULS. (Endo) Thiamine HCl 25 mg., nicotinamide 50 mg., Cal. pantothenate 10 mg., pyridoxine HCl 5 mg., riboflavin eqv 10 mg./cc Amp. 1 cc Box 12s, 25r Vial 10 cc

Use: (1-2 cc.). Vit. B therapy.

MANIBEE-C. (Endo) Vit. B-1 25 mg., B-2 10 mg., nicotinamide 50 mg., Vit. B-6 5 mg., Cal. pantothenate 10 mg., Sod. ascorbate 500 mg./dose. Pkg. 6r, 12s, 100r.

Use: I.V., Vit. B complex w/Vit C.

MANIBEE ELIXIR. (Endo) Vit. B-1 20 mg., B-2 6 mg., B-6 3 mg., nicotinamide 250 mg., pantothenol 15 mg., B-12 15 mcg./fl. oz. Bot 8 oz., 1 pt, 1 gal.

Use: (1-3 t.p. daily) Vitamin deficiencies.

MANIBEE TABLETS. (Endo) Vit. B-1 2 mg., B-2 1 mg., nicotinamide 10 mg., Cal. pantothenate 1 mg., Vit. B-6 0.3 mg., brewer's yeast 0.3 Gm./Tab. Bot. 100s, 1000s

Use: (2 or more daily) Vit. B complex

MANICOLE. (Cole) Mannitol hexanitrate ½ gr./Tab. Bot. 100s, 500s, 1000s.

Use: (1-2 t.i.d.) Essential hypertension.

MANICOLE W/PHENOBARBITAL. (Cole) Phenobarbital ¼ gr., mannitol hexanitrate ½ gr./Tab. Bot. 100r, 500s, 1000s.

Use: (1-2 t.i.d.) Essential hypertension.

MANIRON. (Bowman) Ferrus fumarate 3 mg./Tab. Bot. 100s, 500s, 1000s, 5000s

Use: Anemias.

MANITE. (Dumas-Wilson) Mannitol hexanitrate Tab. ½ gr. Bot. 100s, 1000s. W/Phenobarbital ¼ gr./Tab., Bot. 100s, 1000r.

Use: Orally (1 t.i.d.) Vasodilator.

MANITOLIN. (Blue Line) Mannitol hexanitrate ¼ gr., phenobarbital ¼ gr., rutin 20 mg./Tab. Bot. 100s.

Use: Orally (1-2 q. 3 h.) Vasodilator.

MANNA SUGAR. N N D. 1962 See Mannitol

MANNCOR. (Chicago Pharm) Mannitol hexanitrate 30 mg. (½ gr.) phenobarbital 15 mg. (¼ gr.), rutin 50 mg. (¾ gr.), ascorbic acid 50 mg. (¾ gr.)/Tab. Bot. 100r, 1000s.

Use: (1 q.i.d.) Hypertension & capillary fragility.

MANNECAPS CAPS. (Manne Lab) Thyroid 3 gr., amphetamine sulfate 15 mg., atropine sulfate 2/360 gr., olain ¼ gr., phenobarbital ¼ gr./Cap. Bot. 100r, 500r and 1000s.

Use: Anorectic

MANNEX. (Flint Eaton) Mannitol hexanitrate 32 mg. (½ gr.)/Tab. Bot. 100s, 500r, 1000s.

Use: Angina pectoris & hypertension.

MANNEX W/PHENOBARBITAL. (Flint-

- Eaton) Mannitol hexanitrate 32 mg ($\frac{1}{2}$ gr), phenobarbital 16 mg ($\frac{1}{4}$ gr)/Tab Bot 100s, 500s, 1000s
 Use: Sedative & vasodilator in essential hypertension
- MANNITE**, NND 1962 See Mannitol
- *MANNITOL**, NF NND 1962 Inj, NF (Manna Sugar, O Mannitol Mannite) See Mannitol Solution (Merck Sharp & Dahme)
- *MANNITOL HEXANITRATE**, NND 1962 See Hexanitrol, Cap (Drug Products)
- Hypertensin Tab (CMC)
 Manexin, Tab (Massengill)
 Manhexin, Tab (Normand)
 Manicale, Tab (Cole)
 Manite, Tab (Dumas Wilson)
 Mannex, Tab (Flint, Eaton)
 Maxitate, Tab (Strassenburgh)
 Nitranol, Tab (Merrell)
 Use Orally, coronary vasodilator
- W/Ascorbic acid & rutin See Monrucee Tab (Cole Chem Co)
- W/Nitraglycerine See Maxitate w/nitro glycerine, Tab (Strassenburgh)
- W/Phenobarbital (Squibb Raymer) Maenital hexonitrate $\frac{1}{2}$ gr, phenobarbital $\frac{1}{4}$ gr/Tab Bot 100s, 1000s
 Use: Orally, coronary vasodilator w/sedative
- W/Rauwolfia serpentina alkaloids See: Nitranol R S, Tab (Merrell)
- W/Rauwolfia serpentina Veratrum Viride ext, homotropine methylbromide See Roupertal, Tab (Crookes Barnes)
- W/Reserpine, rutin ascorbic acid See Ruxhexotal W/Reserpine Tab (Lemnae)
- W/Reserpine, theophylline, rutin & ascorbic acid See: Neo Semhyten, Cap (Massengill)
- W/Rutin & delvinal See Siotic Tab (Merck Sharp & Dahme)
- W/Sad nitrite & rutin See Mantrin Tab (Tilden)
- W/Veratrum viride, rutin & rauwolfia serpentina See: RauRa, Encale (Paul Maney)
- Mannitol Hexanitrate w/Phenobarbital** on pp. 437 and 438
- MANNITOL SOLUTION**, NND 1961 (Merck Sharp & Dahme) Amp (25%) 50 cc Box 1s
 Use: I.V. to determine kidney function
- MANOLA**. (Manala Ca) Thiamine HCl 10 mg, riboflavin 3 mg, niacinamide 2.5 mg, iron citrate 3 gr conc salts at glycerophosphates at Sod, Fat & Cal 12 min in palatable aromatic menstruum & 15% nicotinal/n oz Bot 6 ax, 1 gal
 Use: Deficient nutrition
- MANOPHTHLLIN** (Geriatric) Mannitol hexonitrate $\frac{1}{2}$ gr, aminophyllin $1\frac{1}{2}$ gr. Pot
 * Generic name recognized by NF USP or A.M.A.
- Iodide 1 gr, secobarbital $\frac{1}{2}$ gr, phenobarbital $\frac{1}{2}$ gr, rutin $1/6$ gr, ascorbic acid $\frac{1}{2}$ gr, aluminum hydroxide gel 1 gr/Tab Bot 100s, 1000s
 Use: (1-2 ac, t i d, & h s) Vasodilator
- MANRUCAL** (Cole) Phenobarbital 16 mg, mannitol hexanitrate 32 mg, rutin 15 mg, ascorbic acid 25 mg/Tab Bot 100s, 500s, 1000s.
 Use (1 t i d) Essential hypertension
- MANRUCEE**, (Cole) Mannitol hexanitrate $\frac{1}{2}$ gr, ascorbic acid 25 mg, rutin 15 mg/Tab Bot 100s
 Use: (1 2 t i d) Vasodilator.
- MANRUTAL** (G F. Harvey) Mannitol hexonitrate $\frac{1}{2}$ gr, veratrum viride 100 mg, phenobarbital $\frac{1}{2}$ gr, rutin 10 mg./tab Bot 1000s, 5000s
 Use (1 t i d) Vasodilator
- MANRUTAL**, (Normand) Mannitol hexonitrate $\frac{1}{2}$ gr, rutin $1/6$ gr, phenobarbital $\frac{1}{4}$ gr/Tab Bot 100s, 500s, 1000s
 Use Hypertension
- MANTOMIDE**, (Winthrop Lab) See: Chlorbetamide
- MANTOUX TEST** See Tuberculin, Old USP (Various Mfr)
- MANTRIN** (Tilden) Mannitol hexonitrate $\frac{1}{2}$ gr, rutin $\frac{3}{8}$ gr, phenobarbital $\frac{1}{4}$ gr/tab Bot 100s, 1000s Also available without rutin
 Use: (1 t i d) Hypertension
- MANVENE**, (G D Searle) 3 methoxy 16a methyl 1,3,5,10 estratriene 16B 17B diol
 Use Treatment of prostatic cancer
- MANVERTIN** (Sulff & Case) Phenobarbital 10 mg, veratrum viride 50 mg, mannitol hexonitrate 30 mg, sad nitrite 50 mg rutin 20 mg/Cap Bot 100s, 1000s
 Use Coronary vasodilator w/sedative
- MAOLIN NO. 1**, (Alphaden) Mag oxide $7\frac{1}{2}$ gr mag hydroxide $7\frac{1}{2}$ gr, kaolin 22 gr/3 Gm Jar $3\frac{1}{2}$ ax
 Use Antacid & laxative
- MAOLIN NO. 2**, (Alphaden) Hyascyamus ext $\frac{1}{2}$ gr kaolin 22 gr/3 Gm Jar $3\frac{1}{2}$ ax
 Use: Antidiarrhea
- MAOLIN NO 3 1** (Alphaden) Phenobarbital $\frac{1}{4}$ gr, hyascyamus ext $\frac{1}{2}$ gr., mag hydroxide $7\frac{1}{2}$ gr, mag oxide $7\frac{1}{2}$ gr, kaolin 22 gr, agar agar, gum karaya, pepsin & peppermint/tab Bot 60s, 200s, 500s
 Use: Antispasmodic
- MAOLIN NO. 3 PLAIN** (Alphaden) Phenobarbital $\frac{1}{4}$ gr, hyascyamus ext $\frac{1}{2}$ gr., mag oxide 4 gr, mag hydroxide 4 gr., kaolin 22 gr/3 Gm Jar $3\frac{1}{2}$ ax
- MAOLIN NO. 4**, (Alphaden) Mag oxide 4

Mannitol Hexanitrate w/Phenobarbital

| Mannitol Hexanitrate | Pheno-barbital | Others | Manufacturer | Brand |
|----------------------|----------------|---|------------------|----------------------------------|
| ½ gr. | ¼ gr. | | Mallard | Maftrate W/Phenobarbital, Tab. |
| ½ gr. | ¼ gr. | | Strasenburgh | Maxitate W/Phenobarbital, Tab. |
| ½ gr. | ¼ gr. | | Narmond | Manhasin W/Phenobarbital, Tab. |
| ½ gr. | ¼ gr. | | Massengill | Mexal, Tab. |
| ½ gr. | ¼ gr. | | Sutcliffe & Case | Caso-Hex, Cap. |
| ½ gr. | ¼ gr. | | Abbott | Manartal, Tab. |
| 32 mg. | ¼ gr. | | Arice | Mannes W/Phenobarbital, Tab. |
| 32 mg. | 16 mg. | | C.M.C. | Hypertendin Mild, Tab. |
| ½ gr. | ¼ gr. | | Dumas-Wilson | Manite W/Phenobarbital, Tab. |
| ½ gr. | ¼ gr. | | Breen | Nibarb, Tab. |
| ½ gr. | ¼ gr. | | Merrell | Nitranital W/Phenobarbital, Tab. |
| ½ gr. | ¼ gr. | | Smith-Dorsey | Trabar, Tab. |
| 20 mg. | ¼ gr. | Ethaverine 25 mg. | Linsch | Hazaverine, Tab. |
| ½ gr. | ¼ gr. | Rutin 20 mg. | Plessner | Ru Nitral, Tab. |
| ½ gr. | ¼ gr. | Rutin ¼ gr. | Tilden | Mantrin, Tab. |
| ½ gr. | ¼ gr. | Rutin 20 mg. | Blue Line | Manitalin, Tab. |
| ½ gr. | ¼ gr. | Rutin 10 mg., veratrum viride 100 mg. | Harvey | Manrutal, Tab. |
| 25 mg. | ¼ gr. | Rutin 10 mg., aminophylline 45 mg. | Calumbas | Hexa Rutalin, Cap. |
| ½ gr. | ¼ gr. | Rutin 1/6 gr., aminophylline 1 ½ gr., ascorbic acid, ½ gr., secobarbital ¾ gr., Pot. iodide 1 gr., albumin hydroses 1 gr. | Geriatric | Manaphyllin, Tab. |
| 30 mg. | ¼ gr. | Rutin 15 mg., ascorbic acid 20 mg. | Strasenburgh | Maxitate W/Rhomnatin, Tab. |
| 30 mg. | ¼ gr. | Rutin 15 mg., ascorbic acid 20 mg., Va. B-12 | Strasenburgh | Maxitate W/Rhomna-B-12, Tab. |
| 30 mg. | 15 mg. | Rutin 20 mg., ascorbic acid 10 mg. | Lemmon | Ruhezatal, Tab. |
| ½ gr. | ¼ gr. | Rutin 20 mg., theophylline 1.5 gr. | Plessner | Ru-Nitral W/Theophylline, Tab. |

Mannitol Hexamitrate w/Phenobarbital

[438]

| Mannitol Hexamitrate | Phenobarbital | Others | Manufacturer | Brand |
|----------------------|---------------|--|-----------------|------------------------------------|
| ½ gr. | ½ gr. | Rutin 10 mg, veratrum viride 100 mg | G F Harvey | Mannitol Tab |
| ½ gr. | ¼ gr. | Rutin ½ gr, veratrum viride 2/5 gr | F Trout | Vermantin Tab |
| 30 mg | ¼ gr. | Rutin 10 mg, veratrum viride 100 mg | Rand | Verutal Tab |
| 30 mg | ¼ gr. | Rutin 20 mg, veratrum viride 100 mg | Paul Money | Vir I Tin Tab |
| 30 mg | 10 mg | Rutin 20 mg, veratrum viride 50 mg, sod nitrite 50 mg | Suffitt & Case | Monverlin, Cop |
| ¼ gr. | ¼ gr. | Rutin 20 mg, homatropine methylbromide 1 mg | Staddord | Vernite Tab |
| ¼ gr. | ½ gr. | Theobromine 2½ gr | Smith Dorsey | Teoline, Tab |
| ¼ gr. | ¼ gr. | Theobromine 2½ gr, rutin ¼ gr, Vit C 20 mg | Smith Dorsey | Teoline R C |
| ½ gr. | ¼ gr. | Theophylline Sod glycerate 162 mg | Ascher | Nitraphyl, Tab |
| ¼ gr. | ¼ gr. | Theophylline Sod glycerate 2½ gr | Central | Syntrate Tab |
| 15 mg | ¼ gr. | Theophylline Sod glycerate 130 mg | Bobst | Thea Nitral, Tab |
| ½ gr. | ¼ gr. | Thiamine HCl 3 mg | Welter | Thea Mannital W/Phenobarbital, Tab |
| ¼ gr. | ¼ gr. | Veratrum viride 32 mg, rutin 20 mg, choline methyl tartrate 100 mg | Mallord | Amvera, Tab |
| ½ gr. | ¼ gr. | Veratrum viride 1 gr | Merrill | Nitranital, P V, Tab |
| 30 mg | ¼ gr. | Veratrum viride 100 mg, homatropine methylbromide 2.5 mg | Crockett Barnes | Pertenol, Tab |

- gr., Mag. hydroxide 4 gr., kaolin 22 gr./
3 Gm. Jar 3½ oz.
Use: Antacid.
- MAPHARSEN**, N.N.R. 1955. (Parke, Davis)
Oxaphenarsine HCl, U.S.P. 3-Amino-4-hy-
droxyphenyl arsineoxide HCl Amp. 0.04
Gm., 0.06 Gm Box 10s; 0.6 Gm. Box 10s
Use: I.V., antisyphilitic.
- ***MAPHENIDE**. p-Sulphamoylbenzylamine
hydrochloride. See: Sulfbenzamine HCl.
- MAPROFIX**. See: Gordinal Type Detergents
(Various Mfr.).
- MARAX SYRUP**, (Roerig) Hydroxyzine HCl
25 mg., ephedrine sulfate 6.25 mg.,
theophylline 32.5 mg., alcohol 5%/5 cc
Bal pt.
Use: Allergic conditions.
- MARAX TAB.** (Roerig) Hydroxyzine HCl 10
mg., ephedrine sul. 25 mg., theophylline
130 mg./Tab. Bot. 100s.
Use: Bronchodilator.
- MARDIL**, (Warner-Chilcott) See: *Phenazine
Sulfate.
- MAREDOX**, (Burroughs Wellcome) Marezine
50 mg., pyridoxine HCl 50 mg./Tab. Bot.
25s, 100s.
Use: Treatment of nausea & vomiting of
pregnancy.
- MARIZINE HYDROCHLORIDE**, N.N.D. 1962.
(Burroughs Wellcome) Cyclizine HCl, Tab
(50 mg.) Tube 10s, Bot. 100s, 1000s. Supp.
(100 mg.) Box 12s. Pediatric Supp. (50 mg.)
Box 12s.
Use: (1 t.i.d.) Motion sickness & antihista-
mine.
- W/Ergotamine tartrate, caffeine. See: Migral
Forte, Tabs (Burroughs Wellcome).
- W/Pyridoxine HCl. See: Maredox, Tab. (Bur-
roughs Wellcome)
- MAREZINE LACTATE**, N.N.D. 1962. (Bur-
roughs Wellcome) Cyclizing lactate. Amp.
(50 mg./cc) 1 cc., Box 12s, 100s.
Use: Antihistaminic.
- MARPANIL**. See: Sulfbenzamine HCl
- MARGEL**, (Thompson) Balanced formulation
of selected aluminum hydroxide, magnes-
ium hydroxide, magnesium trisilicate and
magnesium carbonate. Susp. Bot. of 12 oz.
Tubs. Boxes of 96.
Use: Gastric hyperacidity.
- MARGELATE**, (Thompson) Aluminum hydrox-
ide and mag. aluminum silicate gels w/0.5
Gm. mag. trisilicate/5 cc. Bot. 1 gal.
Use: Gastric and duodenal ulcer manage-
ment.
- MARGONOVINE**, (Thompson) Ergonovine
maleate 0.2 mg./Tab. or ml. Bot. 100s.
Amp. 1 cc. Box 25s, 100s.
- MARIA SOAKING SOL.** (Santo) Benzalko-
nium chloride 1.50,000, chlorbutanol 0.15%,
in detergent action vehicle. Bot. 4 oz
Use: Lens soaking solution.
- MARLA WETTING SOL.** (Santo) Benzalko-
nium chloride 1:50,000 and chlorbutanol
0.15% in detergent action vehicle. Bot.
4 oz
Use: Lens wetting solution.
- MAR-PHEDACO**, (Strassenburgh) Codeine
sulfate 1 gr., ephedra 30 gr., Cal. guaiacol
sulfonate 5 gr./R. ox. Syr., Bot. 1 pt., 1 gal.
Use: (1 tsp. q. 3 h.) Sedative expectorant.
- MARPLAN**, (Roche) Isocarboxamid 10 mg/
tab. Bot. 100s & 1000s.
Use: Amine oxidase regulator, Angina pec-
toris and mental depression
- MARROW MALT**, (Smith, Miller & Patch)
Red bone marrow ext. 30 Gm., thiamine HCl
1 mg., riboflavin 2 mg., niacinamide 10 mg.,
Vit. D 400 IU, ferrous sulfate ¼ gr., Cal.
glycerophosphate 4 gr., sherry wine 20%,
malt & aromatics, alcohol content 4%/R.
oz. Bot. 4 oz., 1 pt., 1 gal.
Use: Hematinic.
- MARBHMALLOW**. See: Althea (Various
Mfr.).
- MARSILID**, N.N.D. 1961. (Roche) Iproniazid
Phosphate of 1-isanicotiny-2-isopropylhy-
drazine 10, 25 or 50 mg./Tab. Tab., Bot.
100s, 1000s.
Use: Antituberculous agent and mental en-
ergizer
- MARTBATE-45**, (Marsh-Emery Labs) Penta-
erythritol tetranitrate 45 mg./Cap. Bot.
100s
Use: Angina pectoris.
- MARTBATE-45 WITH PHENOBARBITAL**
(Marsh-Emery Labs) Pentaerythritol tetra-
nitrate 45 mg, phenobarbital 45 mg./Cap.
Bot. 100s
Use: Angina pectoris.
- MARTUSSIN SPECIAL** (M. R. Thompson)
Terpin hydrate 200 mg, caffeine 50 mg,
ammonium salicylate 200 mg., ammonium
gluconate 50 mg, glycine 100 mg, benzal-
konium chloride 0.05%/5 cc. Bot. 4 oz.,
pls.
Use: Cough preparation for diabetic or
hypertensive persons.
- MASINATE**, N.N.D. 1962. (Schiefelin)
Testosterone Propionate 1n) 10, 25, 50
mcg/ccs 1n) 100, 250, 500 mg, 1 Gm/
1D cc. Tabs. buccal 5, 10 mg
- MASOBABE No. 1 & No. 2**, (Mason) No 1
—Phenobarbital ¼ gr., niacinamide 0.2
mg./Tab. No 2—Phenobarbital ½ gr.,
niacinamide 0.2 mg./Tab
Use: Sedative and hypnotic.
- MASODONNA**, (Mason) Hyoscyamine sulfate

0.1037 mg, hyoscine hydrobromide 0.0065 mg atropine sulfate 0.0194 mg, phenobarbital 162 mg / Tab Bot 100s
 Use Sedative and spasmolytic

MASSENGILL POWDER (Massengill) Baric acid ammonium alum, berberine sulfate, phenol, thymol, menthol & eucalyptol. Pow, Box 3 oz, 6 oz, 1 lb
 Use: (1 dr in 1 pt water) Vaginal cleanser

MASSE NIPPLE CREAM (Orith) Aflonain & 9 amino-acridine in absorbent cream base
 Tube 1 oz
 Use: Bacteriostatic & healing effect on nipple

MATCO ANALGESIC. (Matco) Acetaminophen 200 mg salicylamide 200 mg, caffeine 100 mg / Tab Bot 30s
 Use Analgesic

MATERCAPS. (G F Harvey) Calcium lactate 616 mg, Vit A 8000 u, D 800 u, B 1 6 mg, B 2 4 mg, niacin 80 mg, C 66.6 mg, E 2 u, cal pantothenate 2 mg, B 12 2 mcg, folic acid 0.2 mg, choline bitartrate 62.8 mg, dl methionine 20 mg, inositol 30 mg, w/iron, mang, pot, pot iodide, mag, Mo, Zn, Co, Cu/2 cap Bot 90s
 Use: Dietary supplement in pregnancy

MATERNAVITE. (Elder) Vit A 5000 u, D 1000 u, B 1 3 mg, B 2 2.5 mg, niacinamide 20 mg, Vit B 6 0.75 mg, cal pantothenate 5 mg, folic acid 0.34 mg, Vit B 12 1.5 mcg, C 50 mg, E 3 u, cal 215 mg, phosph 166 mg, iron 13.4 mg, copper 1 mg, cobalt 0.1 mg, iodine 0.1 mg, mag 7.5 mg, mang 1.5 mg, molybd 0.4 mg, potassium 5 mg, zinc 1.4 mg / Cap Bot 100s, 500s, 1000s
 Use: Vitamin and mineral supplement

MATROFEN DISPELLETS. (Ballard) Homatropine methylbromide 10 mg, phenobarbital 65 mg / cap Bot 60s 250s
 Use 1 morning & at bedtime; antispasmodic, sedative

MATROMYCIN, NND 1962 (Pfizer) An antibiotic elaborated by a strain of Streptomyces antibioticus Oleandomycin phosphate 250 mg / Cap Bot 16s Intravenous Vial 500 mg dry powder diluent to be added

Use: Gram positive organisms (staphylococci)
 W/Tetracycline See: Signemycin (Pfizer)

MATURON. (Ayerst) Amylase 3000 u, protease 12000 u, tryptic activity 500 u, inositol 10 mg dl methionine 5 mg dehydrocholic acid 40 mg. Vitamins A 2500 USP u, D 250 USP u, C 25 mg B 1 1.5 mg, B 2 1 mg, B 6 0.5 mg pantothenate 35 mg, nicotinamide 7.5 mg, rutin

*Generic name recognized by N.F., U.S.P. or A.M.A.

12.5 mg, E 2.5 IU Minerals: Co 35 mg, P 27 mg, I 25 mg, Fe 12.5 mg., Mn 0.15 mg, K 0.85 mg, Zn 0.2 mg, Mg 2.5 mg / Tab Bot 100s, 1000s

Use: Enzyme Vitamin Compound
M.A.V. ELIXIR. (Taylor Labs) Iron saccharated 200 mg with vit B complex glycerophosphates of cal, & mang with Ca Cu and molybdate/oz Bot 1 pt, 1 gal
 Uses Dietary supplement

MAXIDEX. (Alcon Lab) Dexamethasone in a 0.5% methylcellulose vehicle/5 cc
 Use Eye conditions

MAXILETS. (Abbott) (Some as Dayamineral, Tab) Vit A 10,000 u, D 1000 u, B 1 5 mg, B 2 5 mg, nicotinamide 25 mg, Vit B-6 2 mg, B 12 2 mcg, folic acid 0.1 mg, pantothenic acid 5 mg, Vit C 100 mg, Iron 15 mg, copper 1 mg, iodine 0.15 mg, cobalt 0.1 mg, Mang 1 mg, Mag 5 mg, Pot 5 mg, zinc 1.5 mg, molybdenum 0.2 mg / Tab Bot 100s
 Use: Vitamin & mineral supplement

MAXIPEN, NND 1962 (Roerig) *Phenethicifen pot Tab 125 mg, Bot 36s 250 mg Bot 24s, 100s Sal (125 mg/5 cc) 60 cc

MAXIPLATE. (Hart) Vit A 25,000 u, D 1000 u, B 1 10 mg, B-2 5 mg, B 6 1 mg, B 12 5 mcg, niacinamide 100 mg, cal 106 mg, phosphorus 82 mg, iron 10 mg, cobalt 0.1 mg, copper 1 mg, mang 1 mg, iodine 0.1 mg, molybdenum 0.18 mg, pot 49 mg, zinc 1.4 mg, mag 5.6 mg / Cap Bot 100s
 Use: Therapeutic multiple vitamin with minerals therapy

MAXITATE. (Strassenburgh) Mannitol hexanitrate 1/2 gr, Bot 100s, 1000s
 Use (1 to 3 d) Coronary vasodilator,
 W/Nitroglycerine (Strassenburgh) Mannitol hexanitrate 1/2 gr, nitroglycerine 1/100 gr Bot 100s, 1000s
 Use (1 to 3 d or p r n) Coronary vasodilator

W/phenobarbital (Strassenburgh) Mannitol hexanitrate 1/2 gr, phenobarbital 1/4 gr / Tab Bot 100s, 1000s Also w/phenobarbital 1/2 gr
 Use: Orally (1 q 3 h), Vasodilator w/sedative

W/Rauwolfia (Strassenburgh) Maxitole 30 mg, rutin 30 mg, rauwolfia (whole root) 30 mg / Tab Bot 100s, 1000s
 Use: (2 to 3 daily) Severe hypertension

W/Rhamnobarbital (Strassenburgh) Maxitole 30 mg, rutin 15 mg, ascorbic acid 20 mg, phenobarbital 15 mg / Tab Bot 100s 1000s

W/Rhamnolsin (Strassenburgh) Maxitole 30

- mg., rutin 15 mg., ascorbic acid 20 mg./
Tab. Bot. 100s, 1000s.
Use: Essential hypertension.
- MAXITON.** See: Amphetamine (Various
Mfr.).
- MAXUKAL** (Breon) A calcium kinate [cal-
cium quinolate] gluconate. Calcium 50 mg./
cc. Amp. 2 cc., Box 25s
Use: I.M., calcium therapy
- MAZZINI ANTIGEN, CHOLESTERINIZED.**
(Lilly) Mazzini flocculation test for serodiag-
nosis of syphilis. Vial 30 cc.
- M.D.D.** (G. F. Harvey) Vit. A 1333 u., D
133 u., B-1 0.333 mg., B-2 0.667 mg., C
76.65 mg., niacinamide 3.3 mg., dicalcium
phosphate anhyd. 210 mg., ferrous sulfate
12 mg./Cap. Bot. 100s, 1000s
Use: Dietary supplement during pregnancy
- M.D.M. TABLETS** (Pitman-Moore) Sulfameth-
azine 0.983 Gm., sulfadiazine 0.083 gr.,
sulfamerazine 0.083 gr., Sod. citrate 0.25
gr., aromatics & diluent, q.s./Tab. Bot. 100s,
1000s.
Use: Sulfonamide therapy.
- MDM SULFA-SPENSION LIQUID.** (Martin-
Phillips) Sulfamerazine 0.165 Gm., sulfa-
diazine 0.165 Gm., sulfamethazine 0.165
Gm./Tsp. Bot. Pts.
Use: Anti-infective.
- MEADININ.** Mixture of Amidin & Amidin
alk. of Ammi Majus linn. See: Biomoidia-
alk. (Bioproducts Oregon)
- MEASLES CONVALESCENT SERUM, N.N.R.**
See: Measles Immune Human Serum.
- MEASLES IMMUNE HUMAN SERUM, N.F.,
N.N.D.** 1959. (Jr. League Blood Center, Mil-
waukee) Vial 5 cc., 10 cc. (Michael Reese
Research Found.) Vial 5 cc., 7.5 cc., 20 cc.
Use: I.M., prevent or modify measles.
- MEASLES PROPHYLACTIC SERUM, N.N.D.**
1958. See: Immune Serum Globulin (Human)
- MEBARAL, N.N.D.** 1962. (Winthrop Labs.)
(Mephobarbital) N-methyl-5,5-diethyl phenyl
barbituric acid. Tab. ½ gr. Bot. 100s,
1000s; 1½ gr. Bot. 100s, 500s, 1000s; 3 gr.
Bot. 100s, 500s.
Use: Sedative.
W/Diphenylhydantoin.
See: Mebaroin, Tab. (Winthrop Labs.)
Medylan, Tab. (G. F. Harvey)
- W/Reserpine.** See: Reserpine w/Mebaral,
Tab. (Winthrop Lab.)
- W/Theobramine & Col. iodide ditiethanol-
amine.** See: Iacopal, Tab. (Winthrop Labs.)
- MEBAROIN.** (Winthrop Labs.) New name
Mebarin. Mebaral 90 mg., diphenylhydant-
oin 60 mg./Tab. Tab. scored, Bot. 100s,
1000s.
- Use: Epilepsy (grand mal, petit mal) & its
variants.
- MEBROIN.** (Winthrop Labs.) New name for
Mebaroin.
- MeBuSol-C.** (Ethical Specialties) Mephenein
400 mg., butabarbital 16 mg., salicylamide
100 mg., Vit. C 50 mg./Cap. Bot. 100s,
500s, 1000s.
Use: I.I.J.d., muscular disorders & tension
states.
- MEBUTAL.** (Medco) Butabarbital sodium 30
mg., 100 mg./Tab
Use: Sedative and hypnotic.
- * **MEBUTAMATE.** 2-sec-butyl-2-methyl-1,3-
propanediol dicarbamate. See: Capla (Wal-
lace Lab)
- * **MECAMYLAMINE HYDROCHLORIDE,**
U.S.P., N.N.D. 1962. Tab. U.S.P. 3-Methyl-
aminoisocamphane HCl. N-Methyl-di-isobar-
nylamine HCl. See: Invertine, Tab. (Merck
Sharp & Dohme)
- * **MECHLORETHAMINE HYDROCHLORIDE,**
U.S.P., N.N.D. 1962. Inj. U.S.P. Trifluorol,
N.F. X Dichloron. See: Mustargen, Vial
(Merck Sharp & Dohme)
- METHOL TABLETS** (Mann Labs) di-Methlonine
100 mg.; choline 50 mg.; Vit. B-1 3 mg.;
B-2 2 mg.; C 25 mg.; niacinamide 20 mg.;
panthothenic acid 2 mg.; brewer's yeast 25
gr./Tab. Bot. 100s, 1000s.
Use: Metabolic tonic.
- METHOLIN HCl.** See: Methacholine Chloride,
U.S.P.
- METHOLYL BROMIDE, N.N.D.** 1962. (Merck
Sharp & Dohme) Methacholine Bromide Tab
0.2 Gm. Box 24s Bot. 500s.
Use: Orally, parasympathomimetic agent.
- METHOLYL CHLORIDE, N.N.D.** 1961. (Merck
Sharp & Dohme) Methacholine Chloride.
Amp. 25 mg. Box 6s. Bot. 1 Gm., 10 Gm.
Use: S.C. only, parasympathomimetic agent.
W/Menthol, thymol, eucalyptol & methyl
salicylate. See: Lembrase, Oint. (Wyeth).
- * **MECILIZINE HYDROCHLORIDE, U.S.P.,
N.N.D.** 1962. Tab. U.S.P. 1-p-chlorobenzhy-
dryl-4-m-methyl-benzylpiperazine 2 HCl.
See: Bonadettes, Tab. (Pfizer)
Bonine Hydrochloride, Tab (Pfizer)
W/Nicotinic acid See: Antivert, liq. Syr.,
Tab. (Roerig)
- W/Pyridoxine HCl.** See: Bonadaxin, Amps,
Tab. (Roerig)
- MECODRIN.** See: Amphetamine (Various
Mfr.).
- MECOSTRIN CHLORIDE, N.N.D.** 1962.
(Squibb) Sterile, aq. isotonic sol. of di-
methyl tubocurarine chloride, each cc. con-
taining 1 mg. anhy. salt. Vial 10 cc.
Use: Relax skeletal muscle.

* Generic name recognized by N.F., U.S.P. or A.M.A.

- MEDACHE** (Organon) Phenyltoloxamine dihydrogen citrate 44 mg salicylomid 150 mg, acetophenetidin 150 mg, caffeine 32 mg / Tab Bot 100s
Use Tension headache
- MEDAQINT** (Upjohn) Methylprednisolone 15 mg aspirin 300 mg, calcium carbonate 200 mg / Tab Bot 50s
Use Anti-inflammatory & analgesic
- MEDALAC** (Medco) Glycine 390 mg, calcium carbonate 800 mg / Tab Vial 20s, Bot 100s 500s
Use Hyperacidity
- MEDAPRIN** (Upjohn) Methylprednisolone 1 mg, acetylsalicylic acid 300 mg, calcium carbonate 200 mg / Tab Bot 100s
Use Antirheumatic
- MEDA-TEX** (Sutcliffe & Case) Ethinyl estradiol 0.01 mg, methyltestosterone 20 mg Vial A 5000 u, O 500 u, E 2 IU C 30 mg B 1 3 mg B 2 2 mg, B 6 0.5 mg, B 12 1 mcgm niacinamide 20 mg calcium pantothenate 3 mg, folic acid 0.5 mg, choline bitartrate 50 mg inositol 10 mg Fe 60 mg Cu 1 mg, Co 0.2 mg Mn 3 mg iodine 0.15 mg Mo 0.2 mg, Zn 1 mg Mg 5 mg, Ca 48 mg, P 37 mg / 3 Tabs Bot 100s 1000s
- MEDATINIC** (Medco) Nicotinic acid 500 mg / Tab Bot 100s
Use Treatment of hypercholesterolemia
- MEDATUSSIN SYRUP** (Medco) Dihydrocodeinone bitartrate 1/6 gr phenylephrine HCl 30 mg, pyrilamine maleate 50 mg, potassium guaifenesinate 8 gr ammonium chloride 8 gr, tartar emetic 1/12 gr, chloroform 2 min / fl oz Bot / pt / gal
Use Common cold bronchial spasmolysis
- MEDIX** (Medco) O amphetamine sulfate 10 mg or 15 mg / Tab
Use Appetite depressant
- MEDIATRIC CAPS A TABS** (Ayerst) Conjugated estrogens equine (Premarin) 0.25 mg, methyltestosterone 2.5 mg, vit C 100 mg, thiamine mononitrate 10 mg, vit B 12 2.5 mcg, intrinsic factor concentrate 8 mg, ferrous sulfate basic 30 mg, B 2 5 mg, B 6 3 mg, nicotinamide 50 mg, Cot pantothenate 20 mg, d-desoxyephedrine HCl 1 mg / cap & tab Bot 30s 100s, 1000s Tabs Bot 100s
Use: Estrogen, androgen & vitamins for emotional uplift
- MEDIATRIC LIQUID** (Ayerst) Premarin D 25 mg, methyltestosterone 2.5 mg, Vit B-1 5 mg, B 12 1.5 mcg, d-desoxyephedrine HCl 1 mg / 15 cc Bot 1 pt, 1 gal
Use: Treatment of aging patients
- MEDI-BAND** (Purpac) Neomycin in a plastic film forming base Aerosol spray
Use Treatment of minor cuts, scratches burns
- MEDICATED FOOT POWDER** (Upjohn) Benzoic acid 2.5%, salicylic acid 2.5%, chlorothymol 0.04%, boric acid 10%, zinc oxide 2%, camphor 0.1% cinnamaldehyde 0.08% Shaker top tube 1 1/2 oz, 8 oz
Use Treatment and prevention of athlete's foot
- MEDICATED THROAT DISCS** (Parke Davis) Chloroform 1/2 min, w/capsicum, licorice cubeb, peppermint, anise & linseed / disc Bot 60s
Use Orally, throat & mouth infections
- MEDICONE DRESSING** (Medicone) Anesthin, cod liver oil, zinc oxide, hydroxy quinoline Tube 1 oz, 3 oz; Jar 1 lb
Use Antiseptic skin preparation
- MEDICONE-HC RECTAL** (Medicone) Hydrocortisone acetate 10 mg, benzocaine 2 gr, ephedrine HCl 1/16 gr, oxyquinoline sulfate 1/4 gr, zinc oxide 3 gr menthol 1/7 gr balsam peru 1 gr in a cocoa butter base / supp Box 12s
Use Anorectal inflammation, pruritus and hemorrhoids
- MEDIHALER-DUO** (Riker) Isoproterenol HCl 40 mg phenylephrine bitartrate 60 mg / cc in an inert propellant Oral adapter and 15 cc vial
Use Treatment of asthma
- MEDIHALER-EPI** (Riker) Epinephrine bitartrate 70 mg / cc solution Pkg Medihaler in inert propellant 10 cc Stainless steel vial 10 cc Vial 30 cc
Use Inhalation bronchial asthma therapy
- MEDIHALER-ERGOTAMINE** (Riker) Ergotamine tartrate 90 mg / cc 0.36 mg is delivered with each inhalation 25 cc stainless steel vial and plastic mouthpiece adapter
Use Recurrent type headache including migrainous headache
- MEDIHALER-ISO**, N N O 1962 (Riker) Isoproterenol 20 mg / cc solution Pkg Medihaler Stainless steel vial 10 cc Vial 30 cc
Use: Inhalation, bronchial asthma therapy
- MEDIHALER-NITRO** (Riker) Not listed Octyl nitrite aerosol 1% Container 10 cc
Use: Angina pectoris and antispasmodic
- MEDIHALER-PHEN** (Riker) Phenylephrine HCl 0.25%, neomycin sulfate 0.10%, hydrocortisone 0.04% in inert propellant Stainless steel vial 10 cc
Use: Relief of nasal congestion and inflammations
- MEDINAL**, N N R. 1949 (Warner) Barbitol Sod Elaxr 0.13 Gm / 5 cc. Bot 6 fl oz

- 1 gal. Tab. 0.32 Gm. Box 12s; Bot. 100s, 500s.
 Use: (5 gr. b.i.d.) Long-acting hypnotic.
- MEDITABS.** (Medicone) Ephedrine HCl $\frac{1}{4}$ gr., phenobarbital $\frac{1}{8}$ gr., theophylline 1 gr./Tab. Bot. 30s, 100s.
 Use: Orally (1 q. 3 h.), antioesthmic w/ diuretic.
- MEDITRATE TABS.** (Medicines, Inc) Pentacrythral telranitrate 20 mg./Tab. Bot. 100s, 500s, 5000s.
 Use: Coronary artery disease.
- MEDITUSSIN.** (Palmedico) Dihydrocodeinone bitartrate 10 mg., ammonium chloride 520 mg., potassium guaiacolsulfonate 520 mg., chlorofarm 2 min., pyrilamine moleate 50 mg., phenylpropanolamine HCl 50 mg., dextoxyphedrine HCl 2 mg., tartar emetic 5 mg./30 cc. Syrup, phs., gals.
 Use: Cough control.
- MEDOMIN.** N.N.D. 1962. (Geigy) Cycloheptanylethyl barbituric acid, Heptobarbital Tab. 200 mg. Bot. 50s, 250s, 1000s.
 Use: Sedative.
- MEDOPAQUE.** (Bell-Craig) Medopaque H, sterile sol., Sod. ortho-iodohippurate 45%, Sod. carboxymethylcellulose 1.83%, w/stabilizer & preservatives. Vial 10 cc. Medopaque U, sterile sol., Sod. ortho-iodohippurate 45%, Sod. carboxymethylcellulose 2.10%, w/stabilizer & preservatives. Vial 30 cc.
 Use: Radiopacity in diagnosis.
- MEDROL.** N.N.D. 1962. (Upjohn) Methylprednisolone [6-methyl-delta-1-hydrocortisone) 2 & 4 mg./Tab. Bot. 30s, 100s, 2 mg. Bot. 30s; 16 mg., Bot. 50s.
 Use: [$\frac{1}{2}$ dose of prednisolone] An anti-inflammatory corticosteroid
- W/Othaxine.** (Upjohn) Methylprednisolone 2 mg., Orthaxine 75 mg./Tab. Bot. 30s, 100s.
 Use: Bronchial asthma.
- MEDROL ACETATE.** N.N.D. 1962. (Upjohn) Methylprednisolone acetate 0.25% and 1% in veriderm base. Tube 5 Gm., 20 Gm.
 Use: Allergic dermatoses.
- MEDROL W/ORTHOXINE.** (Upjohn) Methylprednisolone 2 mg., methoxyphenamine HCl 75 mg./Tab. Bot. 30s, 100s, 500s, 1000s.
 Use: Asthma and allergic rhinitis.
- MEDROL ACETATE, VERIDERM.** (Upjohn) Medrol 2.5 mg., 10 mg. Methylparaben 4 mg. Butyl p-hydroxybenzoate 3 mg. .25% & 1%. 5 Gm. & 20 Gm. Tubes.
 Use: Allergic dermatoses and inflammatory skin diseases.
- MEDROL-DEPO.** (Upjohn) A sustained depo
 * Generic name recognized by N.F., U.S.F. or A.M.A.
- effect methylprednisolone acetate 40 mg./cc. Vials 1 and 5 cc.
 Use: Anti-inflammatory therapy.
- MEDROL MEDULES.** (Upjohn) Methylprednisolone 4 mg./medule Bot. 30s, 100s.
 Use: Anti-inflammatory therapy
- MEDROL-NEO.** (Upjohn) Veriderm; Medrol acetate 2.5 mg., neomycin sulfate 5 mg./Gm. Tubes 5 Gm.
 Eye-Ear Drops; not listed. Medrol 1 mg., neomycin sulfate 5 mg./Gm. Bot. 5 cc.
 Eye-Ear Oint.; not listed. Medrol 1 mg., neomycin sulfate 1 mg./Gm. Tube 5 Gm.
- MIDROL-SOLU.** (Upjohn) Intravenous and intramuscular methylprednisolone 40 mg./cc. Mix-a-vial 1 cc
- ***MEDROXYPROGESTERONE ACETATE.** N.N.D. 1962. 17a-acetoxy-6a methylpregn-4-ene-3,20 dione.
 See: Depo-Provera, Vials (Upjohn)
 Provera, Tab (Upjohn)
 Provera-Testpac (Upjohn)
 W/Ethoxzolamide, cetylurea. See: Cytron, Tab. (Upjohn)
- MEDYLAN.** (G. F. Harvey) Not listed. Mephobarbital 1 gr., diphenylhydantoin sod. 1 gr./tab. Bot. 100s, 1000s.
 Use: Treatment of epileptic seizures.
- ***MEFENAMIC ACID.** (Parke, Davis) N-2,3-xylanthranilic acid.
 Use: Anti-inflammatory agent.
- MEGA-B.** (Smith, Miller & Patch) Vit. B-1 100 mg., B-2 5 mg., B-6 50 mg., B-12 25 mcg., pantothen 10 mg., niacinamide 75 mg./cc. Vial 10 cc., Box 1s, 6s, 25s
 Use: I.M., I.V., 1 cc. daily, Vit. B complex.
- MEGIMIDE.** N.N.D. 1962. (Abbott) Bemegride (50 mg.) Amp. 10 cc. Vial (5 mg./cc.) 30 cc.
 Use: C.N.S. stimulant to counteract barbiturate depression.
- MEGIMIDE.** N.N.D. 1961. 3-Methyl-3-ethyl glutarimide. (Bemegride). See: Mikedimide.
- MIGLUMINE IOTHALAMATE.** (Mallinckrodt) See: Conray.
- MEINAL.** (Roberts) Phenobarbital 48.6 mg., hyoscyamine sulfate 0.31 mg., etrapine sulfate D06 mg., hyacin hydrobromide D0195 mg./tab.
 Use: Treatment of hypermatility, smooth muscle spasms, etc.
- MEJALIN.** (Mead Johnson) Vit. B-1 1 mg., B-2 1 mg., B-6 0.2 mg., B-12 3 mcg., niacinamide 10 mg., pantothenic acid 1 mg., choline 50 mg., inositol 20 mg., folic acid 0.2 mg., biotin 0.02 mg., p-aminobenzoic acid 0.5 mg., liver fraction 300 mg., iron 7.5 mg./5 cc. or Cap. Cap. Bot. 100s, 500s. Liq. Bot. 12 oz.
 Use: Dietary supplement.

MELABUTOXYCAINE HCl

444

- ***MELABUTOXYCAINE HCl**, NF 2 Diethyl aminoethyl 3 amino 2 butoxybenzoate HCl w/Epinephrine Inj, NF
- MELACTIN** (Squibb) Thiamine HCl 40 mg riboflavin 20 mg, niacinamide 600 mg ascorbic acid 600 mg /lb Bot 1 lb
Use: Dietary supplement
- MELANEX**, (Lilly) Methohexamide Under study
Use: Hypoglycemic agent
- MELCO** (Sulliff & Case) Cocilono tr 40 min, morphine cult 1/4 gr, cyr equil comp pound 24 min, menthol 2/25 gr/8 oz Syr 1 pt, 1 gal
- MELITOXIN** See: Dicumarol (Various Mfr)
- MELLARIL**, NND 1962 (Sandoz) Thioridazine HCl 2 Methylmercapto 10 [2[N methyl 2 piperidyl]ethyl] phenothiazine HCl Tob 10 mg, 25 mg, 50 mg, 100 mg, 200 mg Bot 100c
Concentrate As HCl salt 30 mg/cc Bot 4 fl oz available to hospital only
Use: Tranquilizer
- MELLIN'S FOOD** (Mellin) Dried ext of wheat flour, malted barley & wheat bran w/Pot bicarb protein 10.3%, lol 0.2%, carbohydrate 79.6%, w/Cal, Phos, iron & carotene Bot 10 oz
Use: Dietary supplement
- MELLOSE** Methylcellulose
- MELMAC**, (Davis & Geck) Plastic resin Can 9.74 oz, Cert 6s
Use: Orthopedic
- MELODAN** (Mayrand) Dexamethorphan 60 mg, chlorpheniramine maleate 10 mg, phenylephrine HCl 45 mg, phenindamine tartrate 20 mg, menthol 1 mg/8 oz Bot pt
Use: Cough syrup
- MELONEX**, (Lilly) Methohexamide
Use: Oral anesthetic
- MELOXINE**, NND 1962 (Upjohn) Methoxalen 10 mg /Tab Bot 28s
Use: Sun protectant, tanning aid
- MELOZITS** (Merck Sharp & Dohme) Methylcellulose 1.5 Gm /water, Bot 25s
Use: (1 2 a.c.) Chronic constipation & obesity
- ***MELPHALAN** L3 [p[bis(2 chloroethyl) amino]phenyl]alanine or Lp[bis(chloroethyl)amino]phenylalanine
- MELSAM W/MORPHINE**, (Sulliff & Case) Morphine sulfate 16 mg, chloroform 0.093 cc, ammonium chloride 1.56 Gm, menthol 10 mg, sanguinaria 0.13 Gm, guindelia 1.3 Gm, cenega D 39 Gm, yerba santa 1.3 Gm/A oz Liq Bot pt, gal
- MELSAFIN**, (Sahyun Labc) Name changed from Daphne B phenamine HCl
*Generic: some recognized by NF, USP or A.M.A.
- Orel 1% (free amine) 1 oz Tube Sol 1.4% (Mandelate) Vial 10 cc, Supp 10 mg (HCl) each 2 Gm
Use: Local anesthetic
- MELULOSE** (Tri State) Methylcellulose 0.5 Gm /Tob Bot 100s, 1000s
- MELVARON**, (Lilly) Vit A synthetic 4500 u O synthetic 450 u, thiamine chloride 0.25 mg, riboflavin 0.5 mg, iron & ammonium citrate green 1/2 gr, nondiabetic malt syrup 30 min/fl dr Pkg 1 lb
Use: Vitamin supplement
- MENACYL**, A OR (Lakeside) Aspirin 5 1/2 gr, ascorbic acid 1/2 gr, menadiolone 0.33 mg /Tab Bot 100c, 1000c
Use: Orally (1-2 tid) analgesic & anti rheumatic
- ***MENADIOL SODIUM DIPHOSPHATE**, USP, NND 1962 Inj, Tabs USP 2 Methyl 1 4 naphthohydroquinone diphosphoric ester tetrasodium salt
See: Koppadione, Amp (Lilly)
Synkayvite, Amp & Tab, NNR (Roche)
Thylakoy, Amp & Tab (Squibb)
Use: Vitamin K therapy (Orally or I.M. 4 to 75 mg)
W/Adrenochrome, cemicarbozone, ascorbic acid, heparidin See: Adreclol, Cap, Loz (Organon)
- W/Vit C See: Synkayvite C, Loz & Drop (Roche)
- ***MINADIONE**, USP ADR Cap, NF1 Inj NF; Tab, USP 2 Methyl 1 4 naphthoquinone (Menaphthone, Dondamon K, Aquinone, Aquokay Menoquinone)
Use: Orally & I.M., Vit K therapy
See: Xanone, Perls (Massengill)
Kappoxin, Cap & Sol (Winthrop Labs)
Kayklot, Tab (Raymer)
Kayquinone (Abbott)
Klaltone, Amp (Metho)
Kalklot, Cap (Cole Chem Co)
Thylaqunone (Squibb)
W/Ascorbic acid See: K C, Tab (Paul Maney)
W/Ascorbic acid, Cal lactate & rutin See: Rudonate C, Tab (Accher)
W/Ascorbic acid, calcium, PABA, cod calicylate See: Be Em Cates, (Paul Maney)
W/Ascorbic acid & citrus bioflavonoid See: Coresidin, Cap (Dumas Wilson)
Viflavo, Cap (Columbus)
W/Acpiin & ascorbic acid
See: A-C-K, Tab (G F Horvey)
Asplrite, Cap (Merck Sharp & Dohme)
Menacyl, Tab (Lakeside)
W/Bioflavonoid citrus compound, ascorbic acid See: C.V.P. W/Vitamin K, Syr, Tab (US Vitamin)

- W/Diethylstilbestrol, hesperidin methyl chalcone, and ascorbic acid. See Nulabart, Tab. (Dumas-Wilson)
- W/Sod. salicylate. See: Erqua, Tab. & Liq. (Fellows).
- MENADIONE DIPHOSPHATE SODIUM.** See: Menadiol Sodium Diphosphate.
- ***MENADIONE SODIUM BISULFITE, U.S.P.** N.N.D. 1962, A.D.R.; Inj., U.S.P. 2-Methyl-1,4-naphthoquinone sod bisulfite. See: Hykinane, Amp. & Vial (Abbott)
- Use: Vit. K therapy.
- MENAGEN.** (Parke, Davis) Oral estrogens. (Mixture of highly purified estragenic factors of natural equine origin. Each capsule contains 5000 u. or 10,000 I.U. of ketohydroxyestratriene.) Bot. 100s, 1000s.
- Use: Orally, menopausal symptoms.
- W/Methyltestosterone. (Parke, Davis) Estrogenic activity of 10,000 I.U. ketohydroxyestratriene, & methyltestosterone 10 mg./Cap., Bot. 100s.
- See: Theolondrol. (Parke, Davis)
- Use: Climacteric symptoms in either sex.
- MENAPHTHENE OR MENAPHTHONE, N.N.D.** 1962. See: Menadione (Various Mfr.)
- MENAOQUINONE.** See: Menadione (Various Mfr.)
- MENESIA.** (Pitman-Moore) Mag. gluconate 0.5 Gm./Tab. Bot. 100s, 1000s.
- Use: (2-3 I.I.d.) Antispasmodic.
- MENFORMON.** (Organon) Estrogens in all. Dosules, 2000 u., 5000 u. Box 25s. Vial 10,000 u., 20,000 u./cc) 10 cc., 25 cc.
- Use: Estrogen therapy.
- MENIC.** (Geriatric) Pentylene tetrazole 100 mg., nicotinic acid 50 mg./tab. Bot. 100s, 500s.
- Use: Stimulant in the treatment of senility.
- MENOCRIN.** (Harrowe) Thyroid ext. $\frac{1}{8}$ gr. dienestral 1/640 gr./Tab. Bot. 100s.
- Use: Menopausal & menstrual disorders.
- MENOTROPE.** (Lakeside) Estrafate 0.33 mg., thalame bitartrate 80 mg., Vit. B-12 1.25 mg., folic acid 0.46 mg./Tab. Bot. 100s.
- Use: Orally (1 to 2 daily), estrogen w/vitamin therapy.
- MENSALIN.** (Walker Vitamin) Pyrabram. Tab. 50 mg. Bot. 30s, 100s, 1000s
- Use: (1 I.I.d.) Premenstrual tension & motion sickness.
- MENTHOFAX.** (Burroughs Wellcome) Methyl salicylate 40% w/menthol, camphor & eucalyptol. Tube 1 oz.
- Use: Rubefacient-analgesic ointment.
- MENTHYL VALERATE.** (Navacal) Validol.
- Use: Sedative.
- MEONINE, N.N.D.** 1960. (Ives-Cameron) dl-Methionine. Pow., Bot. 50 Gm. Tab. 0.5 Gm Bot. 100s, 1000s
- Use (1-3 Gm. daily) Nutrient.
- MEOVITE.** (Ives-Cameron) dl-Methionine 250 mg., thiamine HCl 5 mg., riboflavin 2.5 mg., niacinamide 25 mg./Cap. Bot. 100s, 1000s
- Use: Liver damage.
- MEOVITE-E.** (Ives-Cameron) dl-Methionine 125 mg., thiamine HCl 2.5 mg., riboflavin 1.25 mg., niacinamide 12.5 mg., d-alpha tocopherol 50 mg./Cap. Bot. 100s.
- Use: Liver damage.
- MEPARACRINE HYDROCHLORIDE, N.N.D.** 1961 See Quinacrine Hydrochloride, U.S.P., N.N.R.
- MEPARFYNOL, 2-ethinylbutanal-2.** See: Dar-mison (Schering)
- MEPAVLON.** See: Meprabamate
- ***MEPAZINE ACETATE & HCl, N.N.D.** 1962 10-[1-Methyl-3-(piperidyl)methyl]phenothiazine acetate or HCl 5-50. Pacotal (Warner-Chilcott)
- MEPENZOLETE BROMIDE, N.N.D.** 1962. Glycophenylate bromide. 1-methyl-3-piperidyl benzilate methylbromide. See: Centil (Lakeside)
- MEPERGAN.** (Wyeth) Equal amounts of promethazine and meperidine. Vial 25 mg/10 cc., 50 mg/10 cc Tubex 25 mg/2 cc., 50 mg/1 cc.
- Use: Analgesic and sedative.
- MEPERGAN FORTIS.** (Wyeth) Meperidine HCl 50 mg., promethazine HCl 25 mg/Tab. Bot. 30s.
- Use: Analgesic and sedative.
- ***MEPERIDINE HYDROCHLORIDE, U.S.P.** N.N.D. 1962, A.D.R.; Inj., U.S.P.; Tab. U.S.P. Ethyl 1-methyl-4-phenylpiperidine-4-carboxylate hydrochloride. (Dolantol, Dolantin, Dolosal, Dalvonal, Endalate, Isaniprocaine & Pethidine) See: Demeral Hydrochloride. (Winthrop Labs). Inj. (Wyeth) Tubex 50 mg., 75 mg., 100 mg.
- Use: Analgesic.
- See: Pro-Mepergan (Barry)
- W/Lofusate. See: Demeral w/Lofusate (Winthrop Labs.)
- W/Promethazine HCl. See: Mepergan, Amp. (Wyeth)
- MEPHAMIDE.** (Columbus) Mephenesin $3\frac{1}{2}$ gr., salicylomid 2 $\frac{1}{2}$ gr./Tab. Bot. 100s, 1000s
- Use: Management of pain spasm and tension
- MEPHATE.** (Whittier) Mephenesin 0.25 Gm., glutamic acid HCl 0.30 Gm./Cap. Bot. 100s, 500s.
- Use: Muscle relaxant.

* Generic name recognized by N.F., U.S.P. or A.M.A.

- MEPHATINE** (Mallard) Mephenesin 0.25 Gm, salicylamide 0.25 Gm, hamatropine methylbromide 1 mg/Tab Bot 100s, 1000s
Also available w/Vit C
Use: Antispasmodic
- MEPHEDAN**, (Doniels) Mephenesin 250 mg & 500 mg/Tab Bot 100s, 1000s Also pow
- MEPHELOR** (Taylor) Mephenesin 500 mg/Tab Bot 100s, 1000s
- ***MEPHENESIN**, NF, NND 1962; Cap NF; Tab NF (Various Mfr.)
Use: (1 Gm tid or qid) Skeletal muscle relaxant.
- See: Avasyl, Tab (Schenley)
Daseral, Tab, NNR (Evran)
Dialaxal, Tab, Cap & Elixir, NNR (Carnick)
Glyatol, Tab (U S Standard)
Kinovosyl, Elixir & Tab, NNR (Schenley)
Mephedan, Tab (Doniels)
Mephelor, Tab (Taylor)
Mepheral, Tab, NNR (Bryant)
Mephson, Tab, NNR (Tutag)
Myanesin (B Organic)
Myalax, Tab (Key Corp)
Myoxane, Tab, NNR (Asches)
Oranixan, Tab, NNR (Organon)
Prolax Tab, NNR (Cole Chem Co)
Prolaxin, Tab (Lannell)
Relaxant, Tab (Hart)
Sinan, Tab, NNR (Warren Teed)
Spartalaxin, Tab (Spartan)
Spasmolyn Tab, NNR (Heun)
Telhart, Tab (Hart)
Thaxid I, Tab (Normond)
Telansin Tab, NNR (Physicians Drug)
Tefax, Tab & Elixir (Sutliff & Case)
Tefasate Tab, NNR (Brewer)
Tolaxyn NNR (Delta)
Talseral, Cap, Elixir & Tab, NNR (Squibb)
Talsil, Tab (Pharmazeal)
Tolulox Elixir & Tab, NNR (Miller)
Talyripax Tab (Chicago Pharm)
Walco-Nein, Tab (Walker)
- W/Acetylcarbromal reserpine See: Amril Tab (Leeming)
- W/p Aminobenzoic acid salicylamide & Vit C See: Mybaral Tab (Lloyd & Dabney)
- W/p Aminobenzoic acid, salicylamide vit C B-1, niacinamide & menadiolone See: Mybaral Tab (Lloyd, Dabney & Westfield)
- W/Ammanium salicylate & p-aminobenzoic acid See: Sal Phenesin, Tab (Marion)
- W/d Amphetamine sulfate See: Mephedexin Tab (Columbus)
- W/Ascorbic acid prednisolone, salicylamide See: Sal meph/Prednisolone, Tab (Kremers Urban)
- W/Ascorbic acid, salicylamide, calcium p aminobenzoate See: Cousalin, Tab (Grant)
- W/Betaine HCl, ethaverline HCl See: Mepharen, Tab (G F Horvay)
- W/Butobarbital See: Talbuzem, Tab (Zemmer)
- W/Butobarbital, salicylamide, Vit C See: MeBuSal C, Cap (Ethical Specialties)
- W/Calcium glutamate See: Talamic, Improved Cap (Kremers Urban)
- W/Carbromal
See: Carbak, Tab (Hoack Loas)
Carbanesin, Tab (F Traut)
- W/Chloral hydrate & d amphetamine sulfate See: Samatol, Cap (Person & Covey)
- W/Chloral hydrate & clonidine HBr See: Somarsal, Cap (Calumbu)
- W/Cadeina sulfate See: Talseral w/Cadeine Tab (Squibb)
- W/Diadal N acetyl p aminophenol homotropine methyl bromide See: Steri Eze, Tab (Ruffingtan s)
- W/Dibenzyl succinate & salicylamide See: Expasmus, Tab (M H Smith)
- W/Dihydrocortisone bitartrate, Pot citrate & pyrimidine malate See: Tosonan (Organon)
- W/Glutamic acid HCl
See: Mephate, Cap (Robins)
Tolamic (Kremers Urban)
- W/Hamatropine methylbromide See: Prolax w/Hamatropine Methyl Bromide (Cole Chem Co)
- W/Hydrocortisone acetate, salicylamide See: Articon Tab (Walker)
- W/Hydrocortisone acetate sod salicylate hamatropine methylbromide See: Mephosal w/Hydrocortisone, Tab (Craikes Barnes)
- W/Mephobarbital See: Ceretol B (Barre)
- W/Nicotinic acid & belladonna ext See: Latrodol, Tab & Liq (Tailby Nasan)
- W/Pentobarbital See: Nebralin Tab (Smith Dorsey)
- W/Physostigmine salicylate & hamatropine methylbromide
See: Mephylin (Columbus)
Trimitis, Tab (Bell Pharmacal)
- W/Reserpine See: Relaxerpine, Tab (Adami)
- W/Salicylamide
See: Ceretol A (Barre)
Mephamide Tab (Columbus)
Pendron, Tab (Whittier)
Tolagesic, Tab (Miller)
- W/Salicylamide ascorbic acid See: Sal meph Forte, Tab (Kremers Urban)
- W/Salicylamide & ascorbic acid See: Salimeph C, Tab (Kremers Urban).

*Generic name recognized by NF, USP or A.M.A.

- W/Salicylamide, ascorbic acid, prednisolone. See: Solimeph Forte with Prednisolone, Tab. (Kremers-Urban)
- W/Salicylamide & butobarbital Sod. See: Bonasin, Tab. (Kendall)
- W/Salicylamide, homatropine methylbromide
See Homesol, Tab. (Lemman)
Mephatine, Tab. (Mallord)
- W/Salicylamide, dibenzyl succinate. See: Exspasmus, Tab. (M. H. Smith)
- W/Salicylamide, pentobarbital, phenobarbital, caffeine, acetophenetidin. See: Sopena-MBT. (Thompson)
- W/Salicylamide, pot. p-aminobenzoate, ascorbic acid. See: Hoxsol, Tab. (Hoack)
- W/Salicylamide & Vit. D. See: Ertron S-M, Cap. (Whittier)
- W/Sod. salicylate & homatropine methylbromide. See: Mephasol, Tab. & Elixir (Crookes-Barnes)
- Mephenesin & Borbiturate Combinations on p. 448.**
- ***MEPHENESIN CARBAMATE, N.N.D. 1962.**
See: Tolserom, Tab. & Susp. (Squibb)
- ***MEPHENOXALONE, 5-(p-Methoxyphenoxymethyl)-2-oxazolidinone.**
See: Lenetron Tabs (Lakeside)
Tranpase Tablets (Whittier)
Tropidone (Lederle)
- ***MEPHENTERMINE, N.N.D. 1962.** See: Wyamine Inhal. (Wyeth)
- ***MEPHENTERMINE SULFATE, U.S.P., Inj., U.S.P., N.N.D. 1962** N, alpha-alpha-Tri-methylphenethylamine sulfate, Mephine. See: Wyamine Sulfate (Wyeth)
Use: Vasoconstrictor and nasal decongestant. Also I.V. or I.M. 20 to 80 mg.
- W/Penicillin V, salicylamide, promethazine HCl, phenacetin. See: Pen-Vee-Cidin, Cap. (Wyeth)
- W/Phenacetin, aspirin, promethazine HCl. See: Synalgos, Cap. (Ives-Cameron)
- ***MEPHENYTOIN, 3-Methyl-5 ethyl-5-phenylhydantoin.** See: Mesantoin (Sandos)
- MEPHERIN, [G. F. Harvey]** Mephenesin 300 mg., betaine HCl 60 mg., ethoverine HCl 6 mg./Tab. Bat. 100s, 1000s
Use: Muscle relaxant.
- MEPHEROL, N.N.D. 1962.** (Bryant) Mephenesin, Tab. 250 & 500 mg. Bat. 100s, 1000s
Use: Skeletal muscle relaxant.
- MEPHINE.** See: Mephentermine
- ***MEPHOBARBITAL, U.S.P., N.N.D. 1962:** Tab. U.S.P.
See: Mebaral, Tab. (Winthrop Labs) 5-Ethyl-5-phenyl-N-methyl-barbituric acid.
Harvey—Mephobarbital 1½ gr./cap. Bat. 100s.
- W/Acetophenetidin, homatropine methylbromide, levo-amphetamine sulfate, thyroid. See: Amcobese Dokaps. (Amco Drug)
- W/Aminophyllin, ephedrine HCl, tri-hist. See: Asmason, Tab. (Beetz)
- W/d-Amphetamine phosphate, Vit. A, D, B-1, B-2, C, calcium pantothenate, dextrose. See: Via-Dextose, Tabs. (Rowell)
- W/Dextroamphetamine HCl, phenacetin & aspirin. See: Mepha-Dex, Tab. (Rowell)
- W/d-Desoxyephedrine HCl & pentobarbital. See: Mepha-d, Tab. (Walker)
- W/Diphenylhydantoin sod. See: Medylan, Tab. (G. F. Harvey)
- W/Homatropine methyl bromide, atropine methyl nitrate & hyoscyne HBr. See: Spascaps, Tab. T.D. (Nysco Labs.)
- W/Homatropine methylbromide, atropine methyl nitrate & hyoscyne HBr. See: Trascine Unicelles. (Hiss Pharm.)
- W/kl, aminophyllin, ephedrine HCl, tri-hist. See: Neo-Asmason, Cap. (Beetz)
- W/Mephenesin, epoxyaminoxypropine tropate HBr & betaine HCl. See: Scophenesin, Tab. (Harrower)
- W/Mephenesin, hyoscyne HBr. See: Tritanquil, Tab. (Meyer)
- W/Methscopolamine HBr. See: Almethine, Cap. (Meyer Chemical)
- W/Pantothsnole Cal, methscopolamine nitrate. See: Localm, Tabs. (Warren-Teed)
- W/Pentaerythritol tetranitrate, phenobarbital. See: Rith-A-Tol, Tab. (Walker Corp)
- W/Penthenate bromide. See: Monodral-Mebaral, Tab. (Winthrop Labs.)
- W/Scopolamine methyl nitrate. See: Methine, Tab. (Rowell)
- MEPHO-d.** (Walker) d-Desoxyephedrine HCl 5 mg., pentobarbital ¾ gr., mephobarbital ¾ gr./Tab. Bat. 500s, 1000s.
Use: T.F.d. emotional distress.
- MEPHO-DEX.** (Rowell) Dextro-amphetamine HCl 2.5 mg., mephobarbital 16.2 mg., phenacetin 162 mg., aspirin 5 gr./Tab. Bat. 50s.
Use: 1-2 q. 3 h. analgesic with central nervous system stimulant.
- MEPHOSAL** (Crookes-Barnes)
Tab. (Mephenesin 125 mg, Sod salicylate 125 mg, homatropine methylbromide 2.25 mg) Bat. 50s, 100s, 500s.
Use: Muscle spasm & pain.
- MEPHOSAL W/HYDROCORTISONE.**
(Crookes-Barnes) Hydracortisone acetate 5 mg., sod. salicylate 250 mg, mephenesin 250 mg., homatropine methylbromide 2.5 mg/Tab. Bat. 50s.
Use: Rheumatoid arthritis.

* Generic name recognized by N.P., U.S.P. or A.M.A.

Mephensin & Barbiturate Combinations

[448]

| Mephensin | Barbiturate | Others | Manufacturer | Brand |
|-----------|----------------------|---|----------------|-----------------------------|
| 250 mg | Amobarbital 25 mg | 3 (o-methoxyphenoxyl) propionated 12.50 mg | Tufag | Sonazar Tab |
| 400 mg | Amobarbital 8 mg | Hamatropine methylbromide 0.6 mg, physostigmine sulfate 0.25 mg, salicylamide 130 mg | Testogor | Myamphetane Tab. |
| 300 mg | Butobarbital 16 mg | d Amphetamine 2.5 mg | Carrick | Helobar, Tab. |
| 250 mg | Butobarbital 8 mg | d Amphetamine salt 2.5 mg | Maney | Butadex, Tab. |
| 250 mg | Butobarbital 8 mg | d Amphetamine phosphate 3 mg | Crystal | Crystadex, Tab. |
| 300 mg | Butobarbital 10 mg | | Cole | Mustax, Tab |
| 0.5 Gm | Mephobarbital 30 mg | Genacopolamine HBr 1/640 gr, benzoin HCl 1 gr | Olis Lab | Sedesin, Tab |
| 3 1/2 gr | Mephobarbital 1/2 gr | Hyoscine HBr 0.25 mg | Harrower | Scophenesin, Tab. |
| 400 mg | Mephobarbital 10 mg | | Meyer | Tritanqu 1, Tab |
| 425 mg | Pentobarbital 90 mg | | Smith Dorsey | Nebralin, Tab |
| 0.5 Gm | Pentobarbital 100 mg | | Invenex | Talaphen Pentobarbital, Tab |
| 250 mg | Phenobarbital 7.5 mg | Hyoscyamine HBr 0.0625 mg, glutamic acid HCl 62.5 mg | Kremers Urban | Kused, Cap |
| 0.3 Gm | Phenobarbital 8 mg | Hamatropine methylbromide 1.2 mg, hyoscyamine sulfate 0.1 mg, scopolamine HBr 0.02 mg | Central | Mydonna, Tab. |
| 300 mg | Phenobarbital 1/6 gr | Hamatropine methylbromide 2 mg, d-amphetamine sulfate 2 mg | Adams | Relaxamine, Tab |
| 375 mg | Phenobarbital 1/4 gr | Hamatropine methylbromide 1.5 mg, phenacetin 2.5 gr, salicylamide 2.5 gr | Adams | Relaxamine, A P |
| 6 gr | Secobarbital 1/2 gr | | Knack | Haxsen Tab |
| 400 mg | Secobarbital 30 mg | Hamatropine methylbromide 1/2 gr | Crackes Barnes | Secanesin, Tab |
| 5 gr | Secobarbital 1/4 gr | Phenobarbital 1/2 gr, Sod salicylate 5 gr, Vit C 5/6 gr aluminum hydroxide 1 gr | Rawell | Via Phen, Cap |
| 3 gr | Secobarbital 1/4 gr | | Geriatric | Gerisal, Tab |

- MEPNSON, N.N.D.** 1962. (Tulag) Mephenesin. Elixir 0.1 Gm./cc. Rot. 4 fl. oz., 1 pt., 1 gal., Tab. 0.5 Gm. Bot. 100s, 1000s.
- MEPNYSIN.** (Columbus) Mephenesin 4 gr., physostigmine solicylate 1/120 gr., homatropine methylbromide 1/50 mg./Tab. Bot. 100s, 500s, 1000s.
Use: (2 q. 4 h.) Muscle relaxant.
- MEPNTON, N.N.D.** 1962. (Merck Sharp & Dahme) Emulsion of vit. K-1 (2-methyl-3-phytyl-1,4-naphthoquinone). Amp. 50 mg/cc. of vit. K-1. Box 6s. Emulsion Amp. (10 mg/cc.) 1 cc. Tab. 5 mg., Bot. 100s.
Use: I.V., Vit. K therapy.
- MEPIBEN.** (SchenLabs. Pharm.) Methylpiperidyl benzhydryl ether.
Use: Antihistaminic.
- *MEPIPERHENIOL BROMIDE.** 1-(3-hydroxy 5-methyl-4-phenylhexyl)-1-methyl piperidium bromide. See: Dorstine Bromide (Merck Sharp & Dahme)
- *MEPIVACAINE HCl, di-N-Methylpipercolic acid-2,6-dimethylamillide HCl.** See: Carbocaine HCl (Winthrop Labs)
- MEPRANE DIPROPIONATE, N.N.D.** 1962. (Reed & Carnrick) Promethesial dipropionate 1 mg./Tab. Box 30s, 100s, 500s, 1000s.
Use: (1 i.i.d.) Oral estrogen.
- MEPRANE-PHENORARBITAL.** (Reed & Carnrick) Meprane dipropionate 1 mg., phenobarbital 16 mg./Tab. Bot. 100s, 500s.
Use (1 i.i.d.) Estrogen therapy w/sedative.
- *MEPROBAMATE, U.S.P., N.N.D.** 1962; Tabs, U.S.P. (2-Methyl-2-n-propyl-1,3-propanedial dicarbamate.
See: Equanil, Susp. & Tab. (Wyeth)
Meprasan, Tab. (Wallace)
Mepratabs, Tab. (Wallace)
Miltown, Tab. (Wallace Labs)
- W/d-Amphetamine sulfate.** See: Bamadex, Tab. (Lederle)
- W/Benactazine HCl.** See: Depral, Tab. (Wallace Lab.)
- W/Decodran.** See: Decobamate, Tab. (Merck Sharp & Dahme)
- W/Dexamethasone.** See: Decobamate, Tab. (Merck Sharp & Dahme)
- W/Dextroamphetamine.** See: Bamadex Sequels, Caps. (Lederle)
- W/Estragens conjugated.** See: Milprem, Tab. (Wallace)
- W/Ethasheptazine citrate, acetylsalicylic acid.** See: Equagesic, Tab. (Wyeth)
- W/Hydrachlarothiazide.** See: Cyclex, Tab (Merck Sharp & Dahme)
- W/Pentaerythritol tetranitrate.** See: Equanil-trate (Wyeth)
- W/Pentaerythritol tetranitrate.** See: Milttrate, Tab. (Wallace Labs)
- *Generic name recognized by N.F., U.S.P. or A.M.A.**
- W/Pentolinium tartrate.** See: Equalysen, Tab. (Wyeth)
- W/Prednisolone, dried aluminum hydroxide gel.** See: Mepralone, Tab. (Merck Sharp & Dahme)
- W/Premarin.** See: PMB-200. (Ayerst)
- W/Promazine NCl.** See: Prozine, Cap. (Wyeth)
- W/Tridihexethyl iodide.**
See: Miltpath, Tab. (Wallace)
Pothibamate, Tab. (Lederle)
- MEPROBAMATE, N-ISOPROPYL**
See: Corisopradal
- MEPROLONE #1.** (Merck Sharp & Dahme) Prednisolone 1.0 mg., meprobamate 200 mg., dried aluminum hydroxide gel 200 mg./Tab. Bot. 100s
Use: (1-2 t.i.d.) Treatment of rheumatoid arthritis
- MEPROLONE #2.** (Merck Sharp & Dahme) Prednisolone 2.0 mg., meprobamate 200 mg., dried aluminum hydroxide gel 200 mg./Tab. Bot. 100s
Use: (1-2 t.i.d.) Treatment of rheumatoid arthritis
- MEPROLONE #3.** (Merck Sharp & Dahme) Prednisolone 5 mg., meprobamate 400 mg., aluminum hydroxide gel 200 mg./Tab. Vial 30s
- MEPROSPAN, N.N.D.** 1962. (Wallace Lab) Meprobamate in form of coated pellets which release drug continuously for 10-12 hours. 200 mg or 400 mg./Cap. Bot. 30s.
Use: Anxiety and tension states.
- MEPROTABS, N.N.D.** 1962. (Wallace Lab) Meprobamate 400 mg./Coated tab. Bot. 50s.
Use: For relief of anxiety and tension.
- MEPTYLCAINE HYDROCHLORIDE, N.P., A.D.R.** See: Oracaine Hydrochloride (Orendent)
- MEPYRAMINE MALEATE.** See: Pyrimamine Maleate (Various Mfr.)
- *MEPYRAPONE.** (Ciba) 2-methyl-1,2-di-3-pyridyl-1-propanone. See: Metaprone.
- MER-29, N.N.D.** 1962. (Merrell) Withdrawn. Triperanol 250 mg Capsules. Bot. 30s.
Use Cardiovascular. Anticholinester therapy.
- *MERACTINOMYCIN. ACTINOMYCIN D.** (Merck and Co., Inc.)
Use: Antitumor agent.
- *MERAGIDONE SODIUM.** The sodium salt of anhydro-N (beta-methoxy-gamma-hydroxymethylmercuri)propyl-2-pyrrolidone-5-carboxylic acid—theophylline. See: Merdraxone Sodium (Winthrop Lab)
- *MERALLURIDE SODIUM, U.S.P., N.N.D.** 1962; Inj, U.S.P. See: Mercurhydrin, Amp. & Supp. (Lakeside)

W/Ascorbic acid See Mercurhydrin w/Ascorbic Acid (Lakeside)

MERATRAN HCl, NND 1962 (Merrell) Pipradrol HCl alpha (2 piperidyl) benzhydryl HCl Tab (1 mg) Bot 100s, 1000s (2.5 mg) Bot 100s, 1000s

Use 1-2 tid, relief of emotional & mental depression

W/Vitamins & minerals See: Alertonic, El (Merrell)

MIRATRAN W/RESERPINE (Merrell) Pipradrol HCl 1 mg, reserpine 0.25 mg/Tab Bot 100s, 1000s

Use: 1 tid; treatment of tension & anxiety

MIRRAK, NND 1962 Withdrawn (Schief felin) (Acetamercol) [2 acetoxymercuro-4 diisobutylphenol] Tr (1000) colored & stainless, Bot 1 fl oz, 4 fl oz, 1 pt, 1 gal

Use: Externally; antiseptic.

***MERRROMIN**, NF, NNR 1950 (Promo) Sol, NF; Surgical Sol, NF (Asceptichrome, Chromogyre, Cynochrome, Flavural, Oolochrome, Mercurocol, Mercurophage Mercurochrome Plonochrome)

Promo—Disodium 2,7 dibrom-4 hydroxymercurifluorescein Cryst, Bot 10 Gm, 50 Gm, 100 Gm, 500 Gm, 1000 Gm Sol 2%, Bot, 1/2 R oz, 1 R oz, 1 pt, 1 gal

Surgical sol 2%, Bot 1 pt, 1 gal

See: Mercurochrome (Hynson)

Use: Topically antiseptic

MERCAPTOARSINOL Arsthinol See: Balar sen, Tab & Amp (Endo)

***MERCAPTOMIRIN SODIUM**, USP, NND 1962 Disodium salt of N(3 carboxymethylthiomercuro 2 methoxypropyl) comphoramid acid

See: Dluordyn Sodium, Vial (Ayerst)

Thiomerin Sodium, NNR, Vial (Wyeth)

***MERCAPTOPYRINE**, USP, NND 1962 6-Mercaptopyrine Tabs, USP See: Purvina thol, Tab (Burroughs Wellcome)

MERCARROLID Hydroxy phenylmercuric chloride See: Phenacoline w/Mercarrolid (Upjohn).

W/Ephedrine HCl See: Ephedrine & Mercarrolid, Sol. (Upjohn).

MERCAZOLE, See: Methimazole

MIRCENT, (Maltible) Undecylenic acid 5%, phenylmercuric oleate 1:800 Oint 1 oz, 1 lb.

Use: Topically antiseptic & fungicidal

MERCOCEROLS, NNR 1955, ADR See: Mercresin, Stainless or Tr (Upjohn)

MERCODINONE, (Merrell) Dihydrocadinone bisulfate Tab (10 mg) Bot 100s, 500s

Use (5 mg q 4 h, p.r.n) Analgesic & cough sedative

* Generic name recognized by NF, USP or A.M.A.

MERCODOL, (Merrell) Codeine phosphate 600 mg, phenylephrine hydrochloride 300 mg, nethamine HCl 600 mg, sodium citrate 12 Gm /30 cc Bot 1 pt, 1 gal

Use 1 lsp q 3 h & p.r.n at night Anti-tussive & expectorant

MERCODOL W/DECAPRYN SUCCINATE (Merrell) Codeine phosphate 600 mg, phenylephrine HCl 300 mg, nethamine HCl 600 mg, sodium citrate 12 Gm, decapryn succinate 360 mg/30 cc Bot 1 pt 1 gal

Use (1 lsp q 3 h, p.r.n) Antitussive, expectorant, & antihistamine bronchodilator

MERCORISIN, NNR 1955, ADR (Upjohn) Sec—amyltricrosols 0.1%, a hydroxyphenylmercuric chloride 0.1%; Tr 0.2%, Bot 2 fl oz, 4 fl oz 1 pt, 1 gal; Stainless 0.2%, Bot 4 fl oz, 1 gal

Use Topically, antiseptic

MERCUHYDRIN, NND 1958 (Lakeside) Meralluride Supp 0.6 Gm w/sod bicarb

See: Ro 12s

Use Rectal, diuretic

MERCUHYDRIN W/ASCORIC ACID TABLETS, (Lakeside) Meralluride 60 mg (195 mg Hg) ascorbic acid 100 mg/Tab Bot 100 red sugar coated tab

Use (1-2 daily) Diuretic

MERCUHYDRIN SODIUM, NND 1962 (Lakeside) Sod salt of meralluride (methoxymercuripropylsacetylurea) w/theophylline Each cc contains 39 mg organically combined mercury & 48 mg theophylline

Amp 1 cc, 2 cc, Box 12s, 25s, 100s; Vial 10 cc, Box 6s, 25s, 100s

Use (5C or 1V, or 1M, 1 or 2 cc) Diuretic

***MERCUMATILIN**, NND 1962 B-(2'-Methoxy 3'-hydroxymercuripropyl)coumarin 3 carboxylic acid See: Cumerilin, Tab (Endo)

***MERCUMATILIN SODIUM**, NND 1962 See: Cumerilin, Amp (Endo)

MERCUPROCYL, (Barrow) Mersoyl 100 mg, theophylline 50 mg, procaine barate 0.5% /cc Amp 2 cc Box 100s Vial 10 cc

Use: Mercurol diuretic

MERCUPURIN, NNR 1946 See: Mercurophylline Injection (Various Mfr)

MERCURANINE, See: Merbromin (Promo)

MERCURETTES, (Porke, Davis) Metallic mercury 50 gr Briquette, Box 6s 30s, 100s

Use: Topical, syphilis

MERCURIAL ANTISYPHILITICS, See: Mercuric Oleate (Mallinckrodt)

Mercuric Salicylate (Mallinckrodt)

Mercuric Succinimide (Endo)

Mercuric Sulfide (Mille)

MERCURIAL DIURETICS.

- See Cumerilin (Endo).
 Dicurin, Amp. (Lilly)
 Diocardyn Sodium (Ayerst).
 Diurmerin (C. D. Smith).
 Diursol, Amp. (Horvey)
 Maralluride (Lokaride).
 Mercaptomerin Sodium (Various Mfr.).
 Mercurhydrin (Lakeside).
 Mercumaitin Sodium (Endo).
 Mercupurin
 Mercurophylline (Various Mfr.).
 Mercuzonthin Sod. (Campbell)
 Mersaline (Vitamin)
 Merrolyl (Kirk).
 Mersolyn, Amp. (Kirk)
 Neohydrin. (Lakeside)
 Neomersyl, Amp. (Central)
 Salyrgan
 Salyrgan-Theophylline (Winthrop Labs.)
 Thiomerin Sodium (Wyeth).

MERCURIAL-THEOPHYLLINE. See Cumerilin Sodium (Endo).**MERCURIC IODIDE.** W/Antipyrine, benzocaine. See Auralecaine. (Ethical Pharm.)**MERCURIC IODIDE, POTASSIUM, N.N.R. 1950.****MERCURIC OLEATE, N.F. (Marsengill) Oleate of mercury.**

- Use: Parasitic skin diseases.
 W/Coal tar crude, salicylic acid, phenol & p-nitrophenol.

See: Prasal, Emulsion (Torch).**Estercal, Emulsion (less p-Nitrophenol) (Torch).****W/Cresol. See: Mercroal, Sol. (Texas).****MERCURIC SALICYLATE.** (Mallinckrodt) Mercury suboilcylate.

- Use: Parasitic & fungous skin diseases.

MERCURIN. (Campbell) Sod. salt of β -methoxy- γ -hydroxymethyl propylamide of camphoronic acid. Trimethylcyclopentenedicarboxylic acid. Combined with theophylline is Mercurophylline Sod., U.S.P.**MERCURITAL.** (A. J. Parker) Mersaly acid

- 60 mg., theobromine cal. equiv. to 325 gr. of theobromine/Tab. Bot. 100s, 1000s

Use: Diuretic**MERCUROCHROME, N.N.R. 1950.** (Hynson, Wescott & Dunning) Disodium 2,7-dibrom-4-hydroxy-mercurifluorescein. (Merbramin)

- Pow. Bot. 10 & 100 Gm.
 Sol. (2%) Bot. $\frac{1}{4}$, $\frac{1}{2}$ & 1 fl. oz.

Surgical sol. (2%) Bot. 2B, 110, 220 & 440 cc., 1 gal.

- Tab. (0.3 Gm.) Bot. 30s, 100s, 500s.
 Use: Antiseptic.

MERCUROCOL. See Merbromin (Premo)**MERCUROME.** See Merbromin (Premo).**MERCUROPHEN.** (Merck Sharp & Dohme) Sod. hydroxymethyl-*o*-nitrophenolate. Pow., Vial 10 gr.

- Use: Topically, antiseptic.

MERCUROPHYLLIN D-U. (Kremers-Urban) Mercurophylline Inj., U.S.P. Amp. 2 cc., Box 10s, 25s, 5 cc., Box 5s, 25s.

- Use: I.M., I.V., diuretic.

MERCUROPHYLLINE, N.P.; Inj., Tab., N.P., N.N.D. 1959.

- A Sod. salt. See below.

MERCUROPHYLLINE SODIUM, N.N.D. 1959; Sod. salt of (beta-methoxygamma-hydroxymethylmercuri)propylamide of camphoronic acid

- and theophylline Inj., U.S.P.; Tab., U.S.P. Amp. 2 cc., Box 6s, 25s, 100s. Tab. Bot. 5Ds, 100s, 1000s.

See: Mercupurin. (Various Mfr.)

- Mercuzonthin, N.N.R. (Campbell).
 Mercurophyllin D-U (Kremers-Urban).

W/Ascorbic acid.

- See Mercuroscorb, Tab. (U. S. Vitamin)
 Methyllyne C, Pulvaid (Drug Products).

W/Theophylline. See: Diurmerin, Tab. (C. D. Smith).**MERCUROSCORB.** (U. S. Vitamin) Ascorbic acid 100 mg., mercurophylline 74 mg./Tab.

- Bot. 50s, 100s, 1000s.
 Use: Orally (1-2 daily), diuretic.

MERCURY RIIODIDE. W/Sodium Cocodylate, sodium iodide, phosphorus. See Endoorsen. (Endo)**MERCURY COMPOUNDS.**

- See: Antiseptics, Mercury.
 Diuretics, Mercury.

MERCURY CYANIDE.

- Mallinckrodt—Bot. $\frac{1}{4}$ lb., 1 lb., 5 lb.
 Merck—Pow., Bot. $\frac{1}{4}$ lb., 1 lb.

Use: Topically, antisyphilitic & antiseptic.**MERCURY SUCCINIMIDE, Inj. (Endo) Amp. 0.011 Gm./1 cc., Box 12s, 100s.**

- Use: (I.M. & S.E.) Antisyphilitic.

MERCURY SULFIDE. (Hille) Mersulfon, Mersulfol. Sol. 1%, Bot. 1, 4, 12 oz.; 2%, Bot. 30 cc., 60 cc.

- Use: Orally & I.M. (2 cc. b. weekly), antisyphilitic.

MERCUZANTHIN SODIUM, N.N.D. 1959. (Campbell) Mercurophylline Sod. Inj., U.S.P.

- Amp. 1 cc., Box 5s, 25s, 100s; 2 cc., Box 10s, 25s, 100s. Tab. 100 mg. Bot. 100s, 500s

Use: Oral, I.V. & I.M., diuretic**MERDISUL, N.N.R. 1955** (Bowman Bras) Sulfadiazine & sulfamerazine of each 250 mg./Tab. Bot. 100s.

- Use: Sulfonamide therapy.

MERDOXIN. (G. P. Harvey) Aluminum hydroxide 0.162 Gm., mag. trisilicate 0.162

Gm, peppermint oil qs/tab Bot 100s, 1000s

Use: Functional gastric disturbances

MERDROXONE SODIUM (Winthrop Lab)

See Meragidone Sodium

***MERTHOXYLLINE PROCAINE, NND**

1962 See D curin Procaine Amp (Lilly)

MERITENE, (Oiefene) Vitamin & mineral food supplement Con 1 lb

MERLENATE, (Chicago Pharm) Undecylensic acid 5%, phenylmercuric nitrate 1:1500 Oint Tube 1 oz Jars 1 lb Pow shaker 2 oz

Use: Topically fungicidal

MERODICEIN W/Saligenin See: Thonits, Loz (Hynson Wescott & Dunning)

MERODICEIN SOLUTION, (Hynson Wescott & Dunning) Sod um Merale n Monohydraxy mercurid adoresorcinsulfonaphtholein Sod 0.2% sol Bot 2 & 4 fl oz

Use: Topically, antiseptic

W/Saligenin See Thonits Loz (Hynson Wescott & Dunning)

MERPECTODEL, (Poithress) Phenyl mercuric nitrate (1:24,000 conc) in pectin jelly Tube 3 1/2 oz with or without applicator

Use: Antibacterial agent

MERPHENE (Barry Labs) Hyamine 10-X Inert ingred ents 87% Amp 10 cc to make (1:100 dilution) 1000 cc Pkg 5s clear or tinted

Use: Antiseptic

MERPHENYL NITRATE, NND 1961 (Hamilton Labs) Phenylmercuric nitrate basic Oint 1:1500 Tube 25.3 Gm Sol 1:1500 Bot 473 cc., 3.75 L

Use: Externally, antiseptic

MERPHENYL PICRATE WITH PICRIC ACID, NNR 1952 (Hamilton Labs) Phenylmercuric picrate 0.5%, picric acid 12% acetone 10%, alcohol 50%, water 33.3% Tr, Bot 59.2 cc.

Use: Externally, disinfectant

MERPHYLLINE, (Myers Carter) Mersalyl sod 100 mg, theophylline 50 mg./cc Vial 10 cc

MERSALINE (Vistamiz) Mersalyl 200 mg & theophylline 100 mg./2 cc. Vial 10 cc., 30 cc.

Use: I.M. or I.V., diuretic

***MERSALYL, NP** (Winthrop Labs; Kirks Harvey) Mersalyl Sod, Salyrgan Sod (ortho-gamma hydroxymercuri beta methoxypropylcarbonyl)phenoxy acetate W/Theophylline & betaine See: Neamersyl Vial (Central).

MERSALYL ACID W/Theobromine colcliam See: Mercuryl, Tab (A. J Parker)

MERSALYL SODIUM & THEOPHYLLINE,

*Generic name recognized by N.P., U.S.P. or A.M.A.

NND 1959 See: Mersalyl & Theophylline Injection, U.S.P.

***MERSALYL AND THEOPHYLLINE INJECTION, NF NND 1958** (Metro) Mersalyl D2 Gm, theophylline 0.1 Gm, water for inj qs/2 cc Amp 2 cc Box 10s, 100s

See: Diursol, Amp (Harvey)

Merphylline (Myers Carter)

Mersalyl, NNR (Kirk)

Merthiphyl (New Mexico Pharm)

Salyrgan Theophylline, NNR (Winthrop Labs)

Use: Diuretic

W/Procaine borate See Mercuprocy Inj (Sarraws)

MERSALYN, NND 1959 (Kirk) Mersalyl 5 Theophylline Inj, USP (Amp (0.1 Gm/cc) 2 cc

Use: I.V. & I.M., diuretic

MERSULFON See: Mercury Sulfide, Vial (H He)

MERTESTATE, NND 1962 (Breon) Testosterone 100 mg/cc Susp Vial (multiple dose) 10 cc

Use: I.M., androgen therapy

MERTHIOLATE, NNR 1950, ADR 1955 (Lilly) Merthiolate Sod Thimerosal, NF Thiomersalate

Aerosol 6 oz plastic squeeze bot 3/4 oz

Cream (1:1000) Tube 1 oz

Glycerite (1:1000) Bot 1 pt, 1 gal

Jelly: (1:1000) Jar 1 oz

Oint (1:1000) Jar 1 oz, 1 & 5 lb

Ophth Oint: (1:5000) Tube 1/2 oz

Soln 0.1%

Tr: 0.1% 1 pt 1 gal

Nasal Soln: 0.02% w/Ephedrine sulfate

1% Bot 30 cc

Use: Antiseptic

MERTNIPHYL (New Mexico Pharm) Mersalyl 200 mg, theophylline 100 mg/2 cc Vials 10 cc

MERTNYLLINE (Drug Products) Mercuraphylline Inj Amp 2 cc, Box 12s, 25s 100s

Use: I.V. & I.M., diuretic

MERTRICONE, ADR (Rorer) Phenylmercuric borate, Sod pentachlorophenate Bot 1 pt, 1 gal

Use: Disinfection of surgical instruments

MERYALDIN (Formerly Proloxin), (Lan-nett) Maphenacoin Tabs D25 Gm Bot 250s 1000s Tabs 0.5 Gm Bot 250s, 500s, 1000s

Use: Muscle relaxant

MERYTHROL (Chemico) Mercury salicylphenylformic acid Sol 1:2500 Bot 4 fl oz Surgical Ir 1:1000, 1 pt Tr Bot 4 fl oz., 1 pt 1:2400 Bot 4 fl oz, 1 pt Stainless 1:1000, Bot 4 fl oz, 1 pt, 1 gal

Use: Topically, antiseptic

- MESANTOIN.** (Sandoz) Mephénytáin) Phenontáin, N-methyl-5,5-phenyl ethylhydantáin. Tab. 0.1 Gm. Bot. 100s, 500s, 1000s.
Use: (1-2 t.i.d. until dosage is adjusted) Anti-epileptic & anticonvulsant.
W/Phenobarbital. See: Hydantal, Tab. (Sandoz).
- MESCOMINE.** See: Methscopolamine bromide.
- MESOPIN.** (Endo) Homatrapine methylbromide Elixir 5 mg./5 cc. Bot. 1 pl., 1 gal. Pow. Bot. 1 Gm. Tab. 5 mg Bot. 100s, 500s, 1000s.
Use: 2.5 mg. q. 4 h. antispasmodic.
W/Hycodon bitartrate, pyrilamine maleate, ammonium chloride & Sod. citrate. See: Endotussin, Syr. (Endo).
- MESOPIN-PB.** (Endo) Homatrapine methylbromide 5 mg, phenobarbital 15 mg./5 cc. or Tab. Elixir, Bot. 1 pl., 1 gal. Tab., Bot. 100s, 500s, 1000s.
Use: (1 tsp. t.i.d.) Antispasmodic w/sedative.
- MESTENEDIOL.** See: Methylandrostenediol.
- MESTILBOL, N.N.R. 1954;** Inj., Tab. See: Monamestril, Amp. & Tab. (Wallace & Tiernan).
- MESTINON.** Bromide, N.N.D. 1962. (Boche) Dimethylcarbamate of 3-hydroxy-1-methylpyridinium bromide. Pyridastigmin 60 mg / Tab. B elixir 60 mg./Tsp. Timespan Tab. 180 mg Bot. 100s.
Use: Myasthenia gravis.
- *MESTRANOL.** (Lilly) 17 α -ethynyl-3-methoxy-1,3,5(10)-estratrien-17 β -ol.
Use: Estrogenic compound.
- MESULFIN.** (Ayerst) Sulfamethizole 250 mg, methenamine mandelate 250 mg / Tab Bot. 100s.
Use: Urinary tract infections.
- *METABUTETNAMINE HYDROCHLORIDE,** N.F. 2-Isobutylaminoethyl m-aminobenzoate HCl.
W/Epinephrine, Inj., N.F.
- METACARPEN HYDROCHLORIDE.** See: Meirin. (Galzy)
- META-CINE.** (Brayten) Citric acid, lactose, papain, methyl salicylate, menthol, eucalyptol & chlorobutanol. Pow., Pkg. 4 oz., 8 oz.
Use: Vaginal antiseptic & cleanser.
- METACORDALONE.** See: Prednisolone. (Various Mfr.)
- METACORTALONE.** (Schering) See: Maticortalone (Schering)
- METACORTANDRACIN.** See: Prednisone, Tab. (Various Mfr.)
- METACORTANDRALONE.** See: Maticortalone (Schering)
- METACORTIN.** (Schering) See: Maticorten (Schering)
- METACRESYLACETATE.** See: Cresolin, Liq. (Merck Sharp & Dahme)
- *METAGLYCODOL.** (Lilly) 2-m-chlorophenyl-3-methyl-2,3-butanediol.
Use: Central nervous system depressant.
- METADEE.** (Merck Sharp & Dahme) Vit. D-2 600,000 u./cc. Amp. 1 cc., Box 1s, 6s, 25s, 100s.
Use: Antirachitic agent.
- *METANEXAMIDE.** N-Cyclohexyl-N'-(3-amino-4-methylbenzene sulfonyl) urea.
Under study.
See: Euglycin (Upjohn)
Metanex (Lilly)
Use: Hypoglycemic agent.
- METANYDRIN.** (Lakeside) Trichlormethoxide, 2 or 4 mg / Tab. Bot. 100s, 1000s.
Use: Diuretic.
- METAKOL.** (Durex) Sod. lauryl sulfate 1.8%, boric acid 0.5% w/glycerin, Tragaconth & koraya. Tube 4.3 oz.
Use: Contraceptive.
- METALEX.** (Storck) Nicotinic acid 50 mg., pentylsulfatrazole 100 mg./5 cc. Bot. 1 pl.
Use: Geriatric patients.
- METALID.** (Columbus) Acetyl-p-amino-phenol 5 gr./Tab. Bot. 1000s
Use: Analgesic
- METAMINE, N.N.D. 1962.** (Leeming) Trinitrate Phosphate. Triethanolamine trinitrate biphosphate. Aminetrats Phosphate, N.N.D. 1959. Tab. 2 mg, Vial 50s.
Use: Angina pectoris.
- W/Subobarbital.** (Leeming) Aminotrate phos 2 mg, Subobarbital 15 mg./Tab. Bot. 50s.
Use: Angina pectoris (1 q.d.)
- METAMINE SUSTAINED.** (Leeming) Aminotrate Phos. 10 mg /sustained Tab. Bot. 50s.
Use: Angina pectoris (1 q. 12 hr.)
- W/Reserpine.** (Leeming) Aminotrate Phos. 10 mg., reserpine 0.1 mg./Sust. Tab. Bot. 50s.
Use: Angina pectoris (2 Tab. daily).
- W/Subobarbital.** (Leeming) Aminotrate phos. 10 mg, butobarbital 50 mg /Tab Bot 50s.
Use: Angina pectoris (2 tab. daily).
- METAMUCIL, N.N.D. 1952.** (Searle) Psyllium hydrophilic mucilloid w/dextrose. 4 oz., 8 oz., 16 oz.
Use: Orally; laxative.
- METAMUCIL INSTANT MIX.** (G. D. Searle) Psyllium hydrophilic mucilloid with citric acid and sodium bicarbonate A powder which when combined with water, forms an effervescent, flavored liquid. Contains of 16 & 30 packets of 5 & 4 Gm.
Use: Therapy for constipation.

* Generic name recognized by N.F., U.S.P. or A.M.A.

METANDREN, NND 1962 (Ciba) Methyltestosterone Linguet 5 mg white, 10 mg yellow Bot 30s, 100s, 500s Tob 10 mg, Bot 30s, 100s, 1000s Tab (25 mg) Bot 15s, 100s

Use: Androgen therapy

METANITE, (Drug Products) Atropine methyl nitrate Tab 1/120 gr, 1/60 gr. Bot 100s, 500s

Use: Orally, antispasmodic

W/Phenobarbital (Drug Products) Atropine methyl nitrate 1/60 gr, phenobarbital 1/4 gr/Tab, Bot 100s, 500s

Use: Orally, antispasmodic w/sedative

METAPHEN, NNR 1950, ADR (Abbott) Anhydride of 4 nitro 3 hydroxy mercuric-ortho cresol (Nitromersol NF)

Disinfecting sol (1500) Bot 1 pt, 1 gal

First-aid sol (1500) Bot 20 cc

Sol (12500) Bot 12 fl oz, 1 gal

Tst (1200) tinted or untinted Bot 1 & 4 R oz, 1 pt, 1 gal

Use: Antiseptic

W/Allantoin See Allantoin w/Metaphen, Cream (Abbott)

W/Cod liver oil See Cod Liver Oil w/Metaphen, Oint (Abbott)

W/Ephedrine See: Metaphedrin Elq (Abbott)

W/Holocaine HCl See Holocaine & Metaphen, Ophth oint (Abbott)

METAPHEN IN COLLODION (Abbott) Metaphen (Nitromersol) 1:500 In flexible colloid Bot 1 fl oz

METAPHENYLBARRITURIC ACID, See: Mebaral, (Winthrop Labs)

METAPHYLLIN, NNR 1935 See: Aminophylline (Various Mfr)

***METARAMINOL BITARTRATE, NF, NND 1962;** Int, NP See Aramine Bitartrate Sol & Int (Merck Sharp & Dohme)

METARIN, (Bell Pharm) Heparin sod 2500 u, Vit B 12 50 mcg, folic acid 2 mg, choline chloride 100 mg, and inosinamide 50 mg/cc Vial 10 cc

Use: IM, impaired lipid metabolism Anti-coagulant

METASERTS, (Brayten) Phenyl mercuric acetate 3 mg, tyrothricin 0.3 mg succinic acid sod lauryl sulfate lactase & papaine/Insert Bot 100s

Use: Vaginal inserts, antibacterial

METASNAL (Strial) Colorado shale tar 1.7% in washable petrolatum base Tube 2 oz, 4 oz, Jar 1 lb Lotion for 18% with zinc oxide and talc Bot 2 oz, 4 oz

Use: Dermatologic conditions

METASONE TRISULS (Wilfred) Prednisone

*Generic name recognized by N.F. USP or A.M.A.

5 mg for release in 6 to 10 hrs/Cop Bot 50s

Use: Anti Rheumatic/Anil Inflammatory

METASPAS, (Leeming Miles Pharm Inc) Dihexyverine Hydrachloride

Use: Anticholinergic

METATENSIN, (Lakeside) Trichlormethiazide 2 mg and 4 mg, each containing reserpine 0.1 mg/Tab Bot 100s

Use: Hypertension

METATONE, (Parke, Davis) Strychnine glycerophosphate 1/25 gr, VII B 1 4 mg, Col glycerophosphate 4 gr, Pot glycerophosphate 4 gr, Sod glycerophosphate 2 gr, Mang glycerophosphate 1/2 gr, alcohol Bot 12 fl oz, 1 gal

Use: Tonic

***METAXALONE, (Robins)** 5 (3,5 dimethyl phenoxyethyl) 2 oxazolidinone See Ske-laxin (Robins)

Use: Skeletal muscle relaxant, tranquilizer

***METAZOCINE 1,2,3,4,5,6 hexahydro 8 hydroxy-3,6,11 trimethyl 2,6 methano 3 beza zocine**

METCARAPHEN HYDROCHLORIDE, 2-di ethylaminoethyl 1-(3',4'-dimethylphenyl)N-clopanonecarboxylate HCl See Nefrin Hydrochloride (Geigy)

METED SHAMPOO/CLEANSER, (Texas) Tri chloro-carbanilide 15 salicylic acid 2% sulfur 3% as liquid 4 oz Bot

Use: Shampoo and cleanser

MET-ESTRIN, (Metro) Sesame oil sol estro-diol & estrone Amp (0.133 mg/cc) 1 cc, 10 cc; (0.267 mg/cc) 1 cc, 10 cc; (0.667 mg/cc) 10 cc

Use: Estrogen therapy

MET-ESTROL, (Metro) Sesame oil sol di-ethylstilbestrol Vial (1 mg/cc) 30 cc; (5 mg/cc) 10 cc; (25 mg/cc) 10 cc AMP 1 mg/cc Box 10s, 100s

Use: Estrogen therapy

***METNACHOLINE BROMIDE, NF, NND 1962** Tabs, NF Mechalin bromide See Mecholyl Bromide (Merck Sharp & Dohme)

METHACHOLINE CHLORIDE, NF, NND 1961, Cop NF X Sterile NF (Acetyl β methylcholine chloride, Methiacyetyl choline Amechal See Mecholyl Chloride, AMP (Merck Sharp & Dohme)

Use: (IM 10 to 60 mg) Parasympatho-mimetic

W/Camphor menthal & methyl salicylate See: Serin, Oint (McKesson & Robbins)

W/Chloral hyd & bromocholine bromide See: Nypoton (E. Faugera)

W/Methyl salicylate menthal, thymal See: Arthralgen, Oint (Whittier)

METHACHOLINE CHLORIDE OINTMENT.

See: Lembrose (Wyeth).

***METHACYCLINE.** (Pfizer Lab.) 6-deoxy-6-demethyl-6-methylene-5-oxytetracycline.

Use: Antibiotic.

***METHADONE HYDROCHLORIDE, U.S.P.,**

N.N.D. 1962. (Various Mfr.) Inj., U.S.P., Tab., U.S.P., 6-Dimethylamino-4,4-diphenyl-3-heptanone hydrochloride. Amidon HCl, Butalgin, Diaminon HCl, Hoechst 1082D, Midanone, Physeptane HCl, Palamidon HCl Available: Elixir, pow., soln. for inj., syr., tablet.

See: Adonan Hydrochloride, N.N.R. (Winthrop Labs.)

Althase, Syr. (Wyeth).

Dolaphine Hydrochloride (Lilly).

Use: (Orally or subcu, 5 to 10 mg) Analgesic.

W/Propadrine HCl, Pot. citrate & diluted phosphoric acid, See: Methajade, Syr.

(Merck Sharp & Dahme)

METHADROME. (Physicians Drug) Kheilin.

Tab. 20 mg., Bot. 100s, 500s; 50 mg., Bot. 50s, 100s, 500s.

Use: (20 mg. b.i.d. until dosage is adjusted) Coronary vasodilator.

METHAFROME. (Physicians Drug) Kheilin.

Tab. (20 mg.) Bot. 100s, 500s, 1000s; (50 mg.) Bot. 50s, 100s, 1000s.

Use: Asthma & angina pectoris.

METHAJADE. (Merck Sharp & Dahme)

Methadone HCl 10 mg., Propadrine (Phenylpropanolamine HCl) 0.12 Gm., pot. citrate 1.2 Gm., diluted phosphoric acid 4.5 cc./30 cc. Bot. 1 pt., 1 gal.

Use: (1 tsp. q. 3 h.) Sedative expectorant.

METHAKOTE. (Borden's) Protein hydrolysate

consisting of l-leucine, l-isoleucine, l-methionine, l-phenylalanine, l-tyrosine added

methianine and cysteine, benzethonium chloride in Dermabase. Tube 1½ oz., 3 oz.

Use: Therapy and prevention of diaper rash at all degrees of severity.

METHALGEN. (Schuemann-Jones) Cream. 4

oz.

Use: Counter-irritant.

***METHALLENSTRIL.** N.N.D. 1962. 3-(6-Methoxy-2-naphthyl)-2,2-dimethylpentanoic acid.

See: Vallastril, Tab. (Searle)

METHALUTIN. (Parke, Davis) See: Normethandrone.**METHA-MERDIAZINE SUSPENSION, N.N.R.**

1955. (McNeil) Metha-dia-men-sulfanamide, sulfamethazine 15 gr., sulfamerazine 15

gr., sulfadiazine 15 gr./oz. Susp., Bot. 1 pt.

Use: Orally, triple sulfonamide therapy

METHAMINODIAZEPOXIDE. *Chlardi-

zepamide NCl. 7-chloro-2-methylamino-5-

*Generic name recognized by N.F., U.S.P. or A.M.A.

phenyl-3H-1,4-benzodiazepine-4-oxide, HCl.

See: Librium (Roche)

METHAMOCOL.** See: Aranthol (E. Bilhuber)METHAMPHAZONE.** 4-Amino-6-methyl-2-phenyl-3-pyridazone.***METHAMPHETAMINE HYDROCHLORIDE,**

N.F., N.N.D. 1962; Tab. N.F., Elixir 20 mg./fl. oz. Bot. 1 pt. Tab. 5 mg. Bot. 100s, 1000s.

See: d-Desoxyephedrine.

Use: Orally, central nervous system stimulant.

W/Phenobarbital. See: Ambar (Robles)

W/Pentobarbital, Vit. C, B-1, B-2, niacin. See: Obadrin-LA, Tabs. (Massengill).

dl-METHAMPHETAMINE HCl, N.F. See: dl-Desoxyephedrine HCl**METHAMPHETAMINE.** (Lemmon) Methamphetamine HCl 15 mg./Tab. timed-release

Bot. 100s

METHAMPHETAMINE SULFATE. W/Allylisobutylbarbituric acid, urea See: Diabese,

Tabs. (Philadelphia Labs.)

W/Allylisobutylbarbituric acid, carbamide. See: Diurobers, Tabs. (Biber).

METHAMPHIN TABLETS. (Rorer) dl-Desoxyephedrine HCl, Tab. 5 mg. Bot. 100s, 1000s.

Use: (1-2 in the morning) Central nervous system stimulant.

METHAMPYRONE. See: Dipyrone, Prop.

(Various Mfr.)

***METHANDRIOL.** See: Methylandrostenedial.

(Various Mfr.)

METHANDROLAN. (Lanett) Methylandrostenedial 50 mg./cc. aq. susp.

Use: I.M. Anabolic nitrogen therapy.

METHANDROSTENOLONE,** 17 α -methyl-17 β -hydroxyandrostano-1,4-dien-3-one. See: Dicaobal (Ciba)METHANTHELIN BROMIDE, N.F., N.N.D.**

1962; Tab. N.F. See: Bantline, Amp. & Tab. (Searle)

Use: (Orally, i.v., I.M. 25 to 100 mg) Parasympatholytic.

***METHAPHENILENE HYDROCHLORIDE,**

N.F. X, N.N.D. 1959. N,N'-Dimethyl-N-phenyl-N-(2-thenyl) ethylenediamine HCl.

Tab. N.F. X See: Diatrine Hydrochloride, Tab. (Warner)

METH-APHOR. (Borden) Protein hydrolysate

(l-leucine, l-isoleucine, l-methionine, l-phenylalanine, l-tyrosine); methianine, camphor, benzethonium chloride, in Dermabase

vehicle/Oint Tube 1½ oz.

Use: Dermatologic, amino acid preparation.

METHAPYRILENE FUMARATE.

W/Chlorpheniramine maleate, phenylpropenolamine HCl, Vit. C. See: Allerec, Tabs (Pharmocrat)

W/Dextromethorphan HBr, chloroform, amonium Cl ephedrine sulfate, menthol, alcohol See: Miltivite D, (Vitorine)

W/Pyrimine maleate See: Bihistol, Tab (Sutlif & Case)

W/Pyrimine maleate caffeine, eucapree phenylephrine, acetylsalicylic acid, N-acetyl p amino-phenol See: Tussapap, Tab (Sutlif & Case)

W/Pyrimine maleate, salicylamide & d amphetamine phosphate See: Amsake, Cap (Irwin Neisler)

W/Scopolamine HBr aluminum aspirin See: Drows Tabs (Person & Convey)

*METHAPYRILENE HYDROCHLORIDE, N.F., NND 1962 Tab NF NN' Dimethyl N (2 pyridyl) N (2 phenyl) ethylenediamine HCl (Thenylpyramine Terolite) Blue Use—Elixir 200 mg/fl oz Bot 1 pt, 1 gal Use: (Orally 25 to 100 mg q i d) Aeti histamine

See: Allergin, Tab (Myers Carter)

Barhist, Tab (Barry)

Capathyn, Tab (Flint, Eaton)

Cohisine Tab (Pitman Moore)

Caryzal, Oint (Tilden)

Darmin Tab (Tason Drug)

Histadyl, Preo: (Lilly)

Histafed, Cap (Cale Chem Co)

Lullamin, Drap (Reed & Carnrick)

Methaxylene Tab (McNeil)

Pentryl, Tab (Warner Chilcott)

Pyralhyn, Cap (Flint, Eaton)

Semikon, Cream Tab, NNR (Mosseegill)

Sannicaps (Amer Pharm)

Thenylene, Tab NNR (Abbott)

Tem Histine Ophth Oint (Zemmer)

Use Orally antihistamine

Methapyrilees Hydrochloride Combina-
tions on pp 457, 458 and 459

*METHAQUALONE HCl 2 methyl 3 a tolyl
4(3H) quinoxalinone HCl See Tuazole
(Straenburgh Co)

METHARAMINOL. See Aramine (Merck
Sharp & Dahme)

*METHARRITAL, NF NND 1962 Tabs
NF (5.5 dl ethyl 1 methyl barbituric acid
See Gemoni Tab (Abbott)

METH-ASEPTIC (Borden) Methionine benze
thanium chloride zinc lactate/26 Gm
Powder Pkg 12 envelopes

Use Acute dermatitis, exudative and
chronic eczema

*METHASTYRIDOHE 2 2 dimethyl 5 styryl
4 oxazolidinone (Merck & Co, Inc)

METHA-TAR. (Borden) Protein hydrolysate
(leucine l isoleucine l methionine l phen
ylalanine l tyrosine) methionine, benze
thanium chloride liqur carbons deter

* Generic name recognized by N.F., U.S.P. or A.M.A.

geestin in Dermabase/Oint Tube 1 1/2 oz
Use: Dermatalagic; amino acid preparation

METH-ATRO-MINE, (Breen) Methenamine 5
gr, w/atropine sulfate, hyascyamine, ben
zoic acid, salal & methylene blue/Tab Bot
100s, 1000s

Use: Orally (1 q 3-4 h); urinary infections.

*METHAZOLAMIDE, NND 1962 5 Acetyl
Imino 4-methyl Δ^2 -1,3,4 thiazolaine 2
ulfonamide See: Neptozane (Lederle)

METHCOLATE. (Ascher) Choline bitartrate
0.350 Gm, dl methanline D 167 Gm./Tab
Bot 100s, 1000s

Use: Lipotropic agent

METH-DIA-MER SULFA TABLETS, U.S.P.,
N.N.R. 1955 Trisulfapyrimidines, Tab,
U.S.P.

See: M.D.M., Tab (Pitman Moore)

Metha Merdiazine, Tab, N.N.R. (McNeil)

Meth Dia Mer, Tab (Premo)

Mullazine, Tab, N.N.R. (Kremers Urban)

Neotresamide, Tab (Merck Sharp &
Dahme)

Nealrisul (Ascher)

Nealrixine, Tab (Lilly)

Neotrizine, Savoret (Lilly)

Palatraz, Pediatabs (Columbus)

Ray Tri Mides, Tab, N.N.R. (Raymer)

Sulfaloid, Tab, N.N.R. (Lloyd & Dabney)

Sulfo Ter (American Ph Co)

Sulfo In-azine, Tab, N.N.R. (Thompson)

Sulfo Tria Tab (Rexall)

Sulfatryl Granules, N.N.R. (Wampole)

Sulfase, Tab (Wyeth)

Terfonyl, Tab, N.N.R. (Squibb)

Tersulfas, Tab, N.N.R. (Boyle)

Therasulfo (Mallard)

Thi Di Mer (Pitman Moore)

Triazine (U S Standard)

Tri Azo-Tab (Texas)

Trilaeamide, Tab, N.N.R. (Vanpelit)

Trimethazine Tab (Normand)

Trionamide, Tab, N.N.R. (Flint, Eaton)

Triprazine, Tab, N.N.R. (Eaton)

Triple Sulfa, Tab (Various Mfr)

Triple Sulfoid (Vale)

Trisem, Tab (Massengill)

Trisudone, Tab (G F Harvey)

Trisul (Haug)

Tri Sulfameth, Tab, N.N.R. (Funk)

Trisulfazine, Palatab & Tab, N.N.R.
(Central)

Trisulfil (Approved Pharm)

Triazine, Dulcet & Tab, N.N.R. (Abbott)

METH-DIA-MER SULFONAMIDES.

W/Pot penicillin G

See Table under Penicillin G Potassium

W/Erythracin See: Erythramid, FilmTab
(Abbott)

Methapyrilene Hydrochloride Combinations

| Methapyrilene HCl | Others | Manufacturer | Brand |
|---------------------|---|---------------------------------|----------------------------|
| 0.1% | Neamycin sulf 0.1%, phenylephrine HCl 0.25 %, methylbenzothionium chlor 0.01% | Warren Teed | Entamist, Sol |
| Fumarate 12.5 mg | Noxapine 5 mg, APAP 195 mg phenylephrine HCl 5mg, pyrilamine maleate 12.5 mg, caffeine 32.5 mg | Sutliff & Case | Tussapap, Tab |
| 0.2% | Phenylephrine HCl ¼ %, methylbenzethonium chlor 0.01% | Warren Teed | Allermist, Sol |
| Fumarate 30 mg | Phenylephrine HCl 25 mg pyribanase maleate 30 mg, glyceryl guanacolate 250 mg | Sutliff & Case | Branchahist Neo, Sol / oz. |
| 0.25% | Phenylephrine HCl 0.125%, zinc sulfate 0.25%, boric acid 2% | Lemman | Optaphed, Eye drops & Lq |
| Fumarate 8.3 mg | Phenyltoloxamine dihydrogen citrate 6.3 mg, pyrilamine maleate 8.3 mg, aspirin 3 gr, acetophenetidin 2 gr, caffeine ½ gr | Walker Corp | Trist A Lin, Tab |
| 0.2% | Polymyxin B sulf 50 u, grammidine 0.005%, parodrine HBr 1% | Smith, Kline & French Zovell | Driotal Soln |
| 12.5 mg | Procaine HCl 250 mg, vit C 12.5 mg | Merck Sharp & Dahme | Vig Caine, Cap |
| 10 mg | Propadrine HCl 15 mg, pot cit 150 mg, chloroform 15 mg | Rawell | Praktuss/cc |
| 12.5 mg | Pyrilamine maleate 12.5 mg | Durst | Duo-Pyril, Cap |
| 25 mg | Pyrilamine maleate 25 mg | Zemmar | Diminist, Tab |
| 5 mg | Pyrilamine maleate 10 mg, ephedrine sulf 5 mg, dihydro cadéinate bitartrate 1.25 mg | | Zam Histinann, Syr / dr. |
| 5 mg | Pyrilamine maleate 5 mg, phenramine maleate 5 mg | Columbus | Pedrahist/5 cc. |
| 25 mg | Pyrilamine maleate 25 mg, phenramine maleate 10 mg, d om phetamine HCl 6 mg | Testagar | Pymadex Timed, Cap. |
| 8.33 mg | Pyrilamine maleate 8.33 mg, phenylephrine HCl 10 mg, pro phenpyridamine maleate 6.25 mg | Boyle | Citra H P, Cap |
| 8 mg | Pyrilamine maleate 8 mg, phenylephrine HCl 10 mg, phenramine maleate 6 mg, aspirin 240 mg phenacetin 120 mg, caffeine 30 mg, Vit C 50 mg, hesperidin purified 50 mg | Crestmed | Darbacin, Cap |
| 8 mg | Pyrilamine maleate 10 mg, phenylpropanolamine 25 mg, prophenpyridamine maleate 5 mg | Paul Maney | Hislene, Tab. |
| 10 mg | Pyrilamine maleate 10 mg, phenyltoloxamine dihyd cit, 10 mg | Ascher | Triahistin, Tab |
| 15 mg | Pyrilamine maleate 15 mg, phenyltoloxamine dihyd cit 15 mg | Lemman | Trilamine, Tab |
| 10 mg | Pyrilamine maleate 10 mg, prophepyridamine maleate 5 mg | Blue Line | Incorporahist, Tab |
| Fumarate 12 mg | Pyrilamine maleate 12 mg, salicylamide 150 mg, d amphetamine phos. 1 mg | Irwin, Nelsler | Amsalin, Cap. |
| 25 mg | Salicylamide 2½ gr, calcichone sal 1/700 gr, atropine sulf 1/500 gr, phenacetin 2½ gr. | First Texas | F C A H, Cap |

- W/Pyrimidine maleate, phenyltoloxamine dihydrogen citrate, glyceryl guaiacolate & ephedrine sulfate See Sulfussin, Susp (Tilden)
- W/Tetracycline HCl See: Polycycline W/Triple Sulfonamides, Susp (Bristol)
- METH-DIA-MER SULFONAMIDES SYRUP, NNR 1955**
- See: Syrasulfa (Upjohn)
Tri Sulfameth (Funk)
- W/Pat penicillin G
See Darsallin Triple Sulfas, Liq (Smith Dorley)
Eikacillin Sulfas, Liq (Smith Kline & French)
- Pentid Sulfas, Tab (Squibb)
Phenazine, Tab (Elder)
- Sulfa Readicillin, Susp (Upjohn)
Sulfa Sugracillin, Sol (Upjohn)
Sulfa Sugracillin Fortified (Upjohn)
Sulfasal w/Penicillin Liq (National Drug)
Tetracillin, Liq (Schenley)
Trifacillin, Liq (Vanpelt & Brown)
True Cillin Dulcet Tab (Abbott)
- W/Sulfacetimide
See: Delamets Tab (Durst)
Ketrasul, Tab & Susp (Chicago Pharm)
- W/Sulfoguanidine See: Oquadet, Diplot (Eder)
- W/Sulfathiazole & Sod citrate See: Tal Par, Susp (Tilden)
- ***METHOILAZINE HCl, NND 1962** See Tacaryl Tab or Syr (Mead Johnson)
- METHEDRINS HYDROCHLORIDE, NND 1962** (Burroughs Wellcome) Methamphetamine HCl Amp (20 mg/cc) 1 cc 12s 100s Tab 5 mg, Bot 100s 1000s
Use: Central nervous system stimulant
- ***METHENAMINE, NF, NNR 1949**; Tab, NF (Hexamethyleneamine Cystamin, Cystagen, Hexamine, Hexamethylenetetramine) See: Formin, Powder, Tab (Merck Sharp & Dahme)
Hexaform, Tab (Central)
Hexaloids, Tab (Lilly)
Uritane (Parke, Davis)
Uratrapin, Tab. (Warner)
- W/Acetaminosalicylic acid as sod salt See: Salihexin, Cap, Amp (Abbott)
- W/Ammanium chloride See: Uro-Chlor, Tab (Vanpelt & Brown).
- W/Ammanium chloride, zea mays, hyoscyamus fr & triticum See: Hexa Chloride Compound, Sol (Pitman Moore).
- W/Atropine sulfate, hyoscyamine, benzoic acid salol & methylene blue See: Meth-Alo-Mine, Tab (Breon).
- W/Atropine sulf, hyoscyamus ext, salol,
* Generic name recognized by N.F., U.S.P. or A.M.A.
- benzoic acid See Cystorrhal, Tab (Buffington)
- W/Atropine sulf, lithium benzoate, corn silk, broom corn seed, saw palmetto berries hydrangea triticum, asparagus root See: Parmatropin, Liq (Abbott)
- W/Atropine sulfate, Sod benzoate & cocca & ext See Urolax, Liq (Blue Line)
- W/Buchu, uva ursi, hops, pareira brava, hyoscyamus & Pat acetate See: Cystadyne, Pow (Tyree)
- W/Cal mandelate & ammonium acid phosphate See Prahydrian, Tab (Upjohn)
- W/Cal mandelate, atropine sulfate & hyoscyamine alkaloid See Urolate Tab (Borchardt)
- W/Calcium mandelate, phenylloxamidopyridine HCl & sod phosphate monobasic See Urtrial, Cap (Central)
- W/Cholic acid See Felamine, Tab (Sandos)
- W/Cinchophen & cacodylate See Cin Co Mine E, Vial (Kirk)
- W/Dissodium hydrogen phosphate, Mag oxide & citric acid See Hexaphos (Central)
- W/Hyoscyamus ext, atropine sulfate, salol & benzoic acid See Cystorrhal, Tab (Buffington)
- W/Lithium benzoate & atropine See Formotropin, Liq (Abbott)
- W/Lithium benzoate, Sod benzoate & piperazine See Aseptural, Liq & Pow (Wallou)
- W/Methylene blue atropine sulfate, hyoscyamine, benzoic acid salol See: Hexalol Tab (Central)
- W/Phenyl salicylate methylene blue, benzoic acid, atropine hyoscyamine sulfate See Ursepha, Tab (Blaine)
- W/Resorcinol & boraxmin (diasmin) See: Eucharform, Tab (Strassenburgh)
- W/Pat citrate, ammonium salicylate, cocca sagrada, ipecac, caffeine citrate, padaphyllin & camphar See: Methenas, Liq (Merck Sharp & Dahme)
- W/Sod acid phosphate, ext hyoscyamus ext triticum & adiphenine HCl See: Hexoset Tab (Mallard)
- W/Sod biphosphate, Tab, N.F.
Abbott—Tab 5 gr at each Bot 100s 1000s
Lilly—Tab No 1, 5 gr of each Bot 100s 1000s; No 2, 7½ gr of each Bot 100s 1000s
Miller—Tab 5 gr of each Bot 500s 1000s
Merck Sharp & Dahme—Tab 0.3 Gm & 0.5 Gm of each Bot 100s, 1000s
See: Formahydrian, Liq & Pow (Upjohn)
- W/Sod salicylate, Sod, benzoate salol

- corn silk, Iriticum & buchu. See: Hexafona, Elixir (Upjohn).
- W/Sulfacetamide & mandelic acid. See: Fine-Sul, Tab. (Kendall).
- W/Sulfo-Salicylic acid. See: Hexafel, Tab. (Riedel)
- METHANAMINE ANHYDROMETHYLENE CITRATE.** (Formanol, Urapurgol, Urotropin new.) See: Helmitol. (Winthrop Labs.)
Use: Urinary antiseptic.
- METHANAMINE BISMUTH IODIDE.** See: Orbimin, Cap. (Central).
- ***METHANAMINE HIPPURATE.** (Riker) A 1:1 complex of methanamine and hippuric acid
Use: Urinary antiseptic.
- ***METHANAMINE MANDELATE, U.S.P.** N.N.D. 1962; Tab., U.S.P. Tab. (250 mg) (West-Ward)
See: Mandelamine. (Warner-Chilcott)
Mandelamine "Hafgrams," & "Unograms" Tab (Warner-Chilcott)
- W/d-Methionine, phenylazadiaminopyridine HCl, homatrapine methylbromide. See: Azo-Uroquid-A, E.C. Tab. (Beach)
- W/d-Methionine, salicylamide. See: Uroquid-Amide, Tab. (Beach)
- W/Phenylazo-diamino-pyridine. See: Azo-Mandelamine, Tabs. (Warner-Chilcott)
- W/Phenylazo-diamino-pyridine HCl, di-methionine. See: Azolole, Tabs. (Amfre-Grant)
- W/Salicylamide See: Lithitol, Tab. (Columbus)
Use: (Orally 250 mg. to 1 Gm) Urinary anti-infective.
- W/Salicylamide, ext. belladonna See: Mandes, Tabs. (Vale).
- W/Salicylamide, pyridoxine HCl. See: Lithiral w/Pyridoxine, Tab. (Columbus)
- W/Sulfacetamide. See: Praklor, Tabs. (Lloyd, Dobney & Westerfield)
- W/Sulfacetamide, homatrapine methylbromide. See: Septamide, Tabs. (Lemmon)
- W/Sulfacetamide, acid sodium phosphate. See: Uraman, Tab. (Superior Pharm)
- W/Sulfacetamide, phenylsodiaminopyridine HCl, acid sodium phosphate See: Uraman w/Azo, Tab. (Superior Pharm.)
- W/Sulfamethazole. See: Mesulfin, Tabs. (Ayerst).
- METHANAMINE TETRAIODIDE.** Iodoformine. See: Siamine, Cap. (Pitman-Moore).
Use: Iodide therapy.
- W/Sod. salicylate. See: Siamine W/Sodium Salicylate, Cap. (Pitman-Moore)
- METHANAMINE UNDECYLENATE.** (Nordson)
Under study.
- METHINDELATE.** (Carrtons) Methanamine
* Generic name recognized by N.F., U.S.P. or A.M.A.
- mandelate 0.25 Gm. or 0.5 Gm./Tab. Bot 100s.
- Use: Urinary antiseptic.
- METHENEXYL SODIUM.** See: Evipal Sodium. (Winthrop Labs.)
- METHENOS.** (Merck Sharp & Dahme) Methenamin 0.5 Gm., pat. citrate 0.52 Gm., ammonium salicylate 0.25 Gm., cascara sagrada 1.56 Gm., ipecac 60 mg., caffeine citrated 60 mg., padophyllin 15 mg., camphor 15 mg., alcohol 9%/30 cc. Bot. 1 gal.
Use: (1-2 tsp. in water, p.r.n) Analgesic & laxative.
- METHEPONEX.** (Eawl) Vit. B-1 1 mg., B-2 2 mg., Cal. pantothenate 0.2 mg., Vit. B-6 0.2 mg., niacin 5 mg., Vit. B-12 0.3 mcg., methionine 100 mg., inositol 15 mg., choline 30 mg., liver 450 mg., biotin 0.6 mcg. & amino acids 0.6 mg./Cap. Bot. 100s, 500s
Use: Uptropic.
- METMERGINE MALEATE, N.N.D. 1962.** (Sandoz) Methylerganovine maleate, Amp. 102 mg./cc.) 1 cc., Box 6s, 12s, 50s, 100s. 1/4. 0.25 mg./cc. Bot. 100 cc., 100 cc. Tab. 0.25 mg. Bot. 10s, 100s, 500s, 1000s.
Use: Orally, I.V. & I.M., oxytocic.
- METHESCULETIN, DISODIUM SULFONATE.** (Chemo Pura) Disodium 4-methylcaumarin-6,7-dihydroxysulfonate. A bioflavonoidlike compound
- METHESTROL.** See: Pramethestrol
- METHETHARIMIDE, BEMEGIBIDE, 3-Methyl-3-ethyl glutarimide.**
See: Meglmid (Abbott)
Mibedimide (Panray)
- ***METHETAIN.** (Sandoz Pharm) 3-ethyl-1-methyl-5-phenylhydantoin.
Use: Anticonvulsant.
- METHIACIL, N.N.D. 1962.** (Schwarz Lab) Methylthiuracil. Tab. 50 mg. 100s, 1000s
Use: Antithyroid therapy.
- METHIADOXINE.** (Lincoln) Thiamine HCl 1000 mg., pyridoxine HCl 1000 mg., dimethionine 500 mg., choline chloride 1000 mg., gentisic acid ethanamide 2.5%, water for inj./20 cc. Vial 20 cc.
Use: Uptropic & vitamin therapy.
- ***METHICILLIN SODIUM.** Sod. 2,6-Dimethoxyphenyl penicillin. See: Dimethoxyphenyl penicillin sod
- METHILAZINE HCl.** 10-[1-Methyl-3-pyrrolidicyl] methyl] phenothiazine HCl
- ***METHIMAZOLE, U.S.P., N.N.D. 1962; Tab. U.S.P.** 1-Methyl-2-mercaptimidazole. Mercazole. See: Tapazole, Tab. (Lilly)
Use: Thyroid inhibitor. (5 to 20 mg)
- METHINE.** (Rowell) Methscopolamine nitrat 2.5 mg, mephobarbital 32 mg./Tab. Bot 100s, 1000s

- Use: Anticholinergic in the treatment of peptic ulcer
- METHIOCON** (Consolidated Mid) Methionine 0.5 Gm /tab Bot 30s, 100s, 250s, 1000s
Use: Lipotropic agent
- METHIODAL SODIUM**, NF NNR 1952 Inj, NF Sod Iodomethanesulfonate (Abro dit Radiographol O agnorenol) See Skiodan (Winthrop Lab.)
Use: Radiopaque medium for urinary tract
W/Neomycin sulfate See Retropaque Vial (Winthrop Labs)
- ***METHIOPRAZINE HCl** dl 10 (3 di methylamino 2 methylpropyl) 2 methylthio-phenothiazine HCl (Smith Kline & French Lab.)
- ***METHIOHINE**, NF NND 1962 Cap NF, Tab, NF dl Methionine alpha amino gamma methylmercaptobutyric acid
See: Allora (Dome)
Amurex Cap (Reid)
Dyprin Cap (Lincoln)
Meonine, Tab, NNR (Wyeth)
Methiocon Tab (Consol Mid)
Metiane, Tab, NNR (Labica Debrusile)
Oradash, Cap (Adams)
Pedometh, Cap (Durst)
W/d Amphetamine, choline bitartrate, phe nobarbital & vitamins See: Obolip Cap (Lakeside)
W/Ascorbic acid See Metarbic (Elder)
W/Betaine, choline, inositol, Vit B-12 liver See: Beta Methischol, Syr (U S Vitamin)
W/Choline bitartrate & Inositol See: Lipocaps Cap (Lakeside)
W/Choline dihydrogen citrate
See: Chamene Cap (Dumas Wilson)
Methalate, Tab (Archer)
W/Choline d hydrogen citrate & Inositol
See: Lipotropic, Cap (Lakeside)
W/Dehydrocholic acid, choline bitartrate & betaine
See: Cholitropic, Tab (Buffington)
W/Inositol, rutin, methyltestosterone, conjugated estrogen (equine) vitamins and minerals See: Via-Geric H, Tab (Rowell)
W/Phenylazo diamino pyridine HCl, meth enamine mandelate See: Azolate Tabs (Amire Grani)
- METHIOHINE, ACETYL**. See: Acetylmethionine (Chemo Para)
- METHIOPLEX STABLE SOLUTION**. (Lincoln)
Methionine 50 mg., thiamine HCl 100 mg., niacinamide 200 mg., riboflavin 4 mg., choline 100 mg., pyridoxine HCl 4 mg., pan theneol 4 mg., benzyl alcohol 1%, distilled
- * Generic name recognized by N.E., U.S.P. or A.M.A.
- water qs /2 cc Vial 30 cc; Lyophilized 5 cc, 30 cc Tab, Bot 100s
Use: Lipotropic & vitamin therapy
- METHISCHOL**. (U.S. Vitamin) Choline dihydrogen citrate 2.5 Gm, dl methionine 1 Gm, inositol 0.75 Gm, Vit B 12 9 mcg, liver conc & desic liver 0.78 Gm /3 dr or 9 Cap Cap, Bot 100s, 250s, 1000s Syr Bot 1 pt, 1 gal
Use: Cirrhosis of liver
- METHISCHOL INJECTABLE**. (U.S. Vitamin)
Choline chloride 100 mg, inositol 50 mg., dl methionine 25 mg, Vit B 12 3 mg /cc Ampul 2 & 10 cc
Use: I.M.; lipotropics
- METH-I-SOL** (Ophthalmus Inc) Methylcellulose 1/2% sol Bot dropper 15 cc
Use: Eye emollient
- ***METHITURAL SODIUM**, NND 1959 See Neraval, Vial (Schering)
- METHIUM CHLORIDE**, NND 1962 (Warner Chilcott) Hexamethonium chloride Tab 125 & 250 mg, Bot 100s, 500s
Use: Hypertension
- W/Reserpine (Warner Chilcott) Methium 125 mg, reserpine D 125 mg & Methium 250 mg reserpine 0.125 mg Tab, Bot 100s 500s
Use: 1 q s d; hypertension
- METHIUM-RAUWOLFIA**, TAB. (Warner Chilcott) On trial; hypertension
- ***METHOCARBAMOL**, NND 1962 (2 hydroxy 3 a methoxyphenoxypropyl) carbamate Carbamate of glyceryl guaiacolate
See: Robaxin Tabs (Robins)
- W/Aspirin See Robaxisol Tab (Robins)
- W/phenacetin acetylsalicylic acid, hyoscyamine sulfate phenobarbital See Robaxisol PH Tab (Robins)
- METHOCEL**, NND 1962 (Coviel) Methyl cellulose Bot 4 oz, 1 lb
W/d Amphetamine phosphate & dicalcium phosphate See: Amphetacel Tab (Lemmon)
- METHOCHLORIDE**. See Triburan (Roche Lab.)
- ***METHOHEXITAL SODIUM**, NND 1962 Alpha (dl) 5-allyl 5 (1 methyl 2 pentynyl) 1 methylbarbituric sod See: Brevital (Lilly)
- METHOIN MEPHENYTOIN**. See: Meantoin (Sandoz)
- METHOLANUM OINTMENT**. (Sutliff & Case)
Methyl salicylate 22%, menthol 5%, lanolin anhydrous 50% petrolatum wax Jcr 1 lb
- ***METHOPHOLINE HCl**. 1 (p chlorophenethyl) 2 methyl 6,7 dimethoxy 1,2,3,4 tetrahydroisoquinoline HCl (Hoffmann LaRoche)
Use: Analgesic

- ***METHOPYRAPONE**. Su-4885. 2-Methyl-1,2-di-3-pyridyl-1-propanone. See: Metapirone (Ciba)
- METHORATE**. See: Dextramethorphan HBr, N.N.D. 1959.
- d-METHORPHAN HBr**. See: Dextramethorphan HBr, N.N.D. 1959.
- W/Glycerol guaiacolate, potassium citrate, W/Glycerol guaiacolate, potassium citrate, citric acid, Ipecac fluid extract. See: Sarcbutuss, (Dalin).
- W/Glycerol guaiacolate, ammonium Cl, pyrilamine maleate, phenylephrine HCl, sodium citrate. See: Carrusin-NN, (Cortane)
- METHORPHINAN**. *Racemorphan. See: Dramoran (Schering)
- i-METHORPHINAN LEVORPHANOL**, N.N.D. 1959. See: Leva-Dramoran tartrate (Roche).
- METHOSTAN**, (Schering) Methylandrostenedial, Tab. 25 mg., Bot. 30s, 100s. Vial, aq. sup. (50 mg./cc.), 10 cc.
- Use: Anabolic steroid.
- ***METHOTREXATE**, U.S.P. (Lederle) Tabs U.S.P. Amethopterin. 4-Amino-N-methyl pteroylglutamic acid Tab (2.5 mg) Bot. 100s.
- Use: Leukemia in children.
- METHOTREXATE SODIUM**, (Lederle) 4 Amino-N¹⁰-methyl-pteroylglutamic acid sodium 5 mg. Vial 5 mg.
- Use: Leukemia therapy.
- ***METHOTRIMEPRAZINE**, (Lederle) 2-methoxy-10-(3-dimethylamino-2-methyl-propyl)phenothiazine.
- W/levo-mepromazine
- Use: Tranquillizer, non-addicting analgesic.
- METHOXA-DOME**, (Dome) 8-Methoxyptoralen 10 mg./Cap. Bot. 50s, 100s.
- Use: Idiopathic vitiligo.
- ***METHOXAMINE HYDROCHLORIDE**, U.S.P., N.N.D. 1962; [n.], U.S.P. beta Hydroxy-beta-(2,5-dimethoxyphenyl)-isopropylamine HCl. (I.V. & I.M. 2.5 to 5 mg) Sympathomimetic.
- See: Valoxy, Amp. (Burroughs Wellcome)
- W/Polymyxin B sulf., gramicidin. See: Aerodin, Sol. (Burroughs Wellcome)
- ***METHOXSALEN**, N.N.D. 1962. 8-Methoxyptoralen. 8-Hydroxy-4',5',6,7-furcoumarin. 8-methoxyptoralen, ammidin, xanthoxin.
- See: Biomaidin, Tab., Amp. (BioProducts)
- Meloxine, Tab. (Upjohn)
- Methoxa-Dome, Cap (Dome)
- Oxioralen, Cap., Lot. (Elder)
- METHOXYCELLULOSE SODIUM**. (Mixture of M.C. and sod. C.M.C.) W/Methamphetamine HCl. See: Obetamine, Tab. (Sly)
- ***METHOXYFLURANE**. (Dow Chem. Co.)
- * Generic name recognized by N.F., U.S.P. or A.M.A.
- (1,1-dichloro-2,2-difluoro-2-methoxyethane)
- See: Penthrone (Abbott)
- Use: General Inhalation anesthetic.
- ***METHOXYPHENAMINE HYDROCHLORIDE**, N.N.D. 1962. beta-(o-Methoxyphenyl) isopropylmethylamine HCl. See: Orthoxine Hydrochloride (Upjohn)
- W/Aminaphylline. See: Orthoxine & Aminaphylline, Cap. (Upjohn).
- W/APC & pyrrolizate. See: Pyrraxate, Cap. (Upjohn).
- W/Chlorpheniramine maleate, acetophenetidin, acetylsalicylic acid, caffeine. See: Pyrraxate, (Upjohn)
- W/Dihydrocodeinone bitartrate, hyocyanine HBr & Sod. citrate. See: Orthoxical, Syr. (Upjohn).
- W/Medrol. See: Medrol W/Orthoxine, Tab. (Upjohn)
- 3-(o-METHOXYPHENOXY)-prepnedial-8,2**. See: Glycerol Guaiacolate.
- o-METHOXYPHENYL GLYCERYL ETHER**. See: Glycerol Guaiacolate.
- ***METHOXYPRIMAZINE MALEATE**, N.N.D. 1962. See: Tentone, Tab. (Lederle)
- 8-METHOXYPSORALEN**. See: Methoxsalen, N.N.D. 1959.
- METHRAL** (For the United Kingdom) (Pfizer) Fluperolone Acetate
- ***METHSCOPOLAMINE BROMIDE**, N.N.D. 1962. Epoxytropine tropate methylbromide, scopolamine methylbromide.
- See: Ampyrox, Tab. #276. (Elder)
- Escopine Bromide, Tab. (Lincoln)
- Nutrop, Cap. (Marin)
- Pamine Bromide. (Upjohn)
- Proscamide, Cap. (E. S. Miles)
- Scoline, Tab. (Kendall)
- Tropone, Vial (Myers-Carter)
- W/Aluminum hydroxide & magnesium hydroxide. See: Ampyrox, Tab. #293. (Elder)
- W/Amobarbital. See: Scoline-Amobarbital, Tab. (Kendall)
- W/Butobarbital. See: Ampyrox, Elixir. (Elder)
- W/Butobarbital sod. See: Ampyrox, Tab. #281. (Elder)
- W/Butobarbital Sod., aluminum hydroxide gel dried & Mag. trisilicate. See: Eulcin, Tab. (Staff Research Labs).
- W/Dihydroxyaluminum aminoacetate, sod. lauryl sulfate & magnesium hydroxide. See: Ato-Scop, Cap., Sup. (Lloyd, Dabney & Westerfield)
- W/Guar gum, dioctyl sod. sulfosuccinate. See: Synacol, Cap. (Reed & Carnick)
- W/Magnesium trisilicate & aluminum hydroxide gel. See: Trigelma w/Mescamine, Tab (Burlington)

W/Mephobarbital See Almelhine, Cap (Meyer Chemical)

W/Neomycin sulfate See Biosal M (Upjohn)

W/Phenobarbital See Prescamite B, Cap (E S Miller)

W/Phenobarbital, cal carb, mag trisilicate mag oxide See Scopamal, Tab (Miller)

W/Tuazole See Akalan T, Tab (Strassenburgh)

***METHSCOPOLAMINE NITRATE, H N D**

1962 Scopalamine Methyl Nitrate Prep (Various Mfr) Mercomine

See: Skopalate, Cap, Tab & Vial (Strassenburgh)

Skapyl, Drops (Pharmacia Lab)

Vircape, T D Cap (Lynn)

W/Aluminum hydroxide, Mag carb & sod dodecyl sulf See: Skopalate P.A.M., Cap (Strassenburgh)

W/Aluminum hydroxide magnesium carbonate See Alucen Tabs (Central)

W/Bulbarbital atropine methyl nitrate See: Seplate Tabs (Central)

W/Mephobarbital See Methine, Tab (Rowell)

W/Pantothenate cal mephobarbital See Illocalm, Tabs (Warren Teed)

W/Pentobarbital, pepin glutamic acid HCl, cellulose, ex bile extract pancreatin See: Konumadic Tabs (Dorsey)

W/Phenobarbital See Scopamate (Deseret)

W/Piperidine ethanol benzilate HCl, magnesium hydroxide, aluminum hydroxide See: Modulral, Tab (Reed & Carnrick)

***METHSUXIMIDE, N N D 1962 N Methyl**

alpha, alpha methylphenylsuccinimide See Celontin Kapsels (Parke, Davis)

METHULOSE, BUFOPTO, (Professional

Pharm Co) Methylcellulose 1% and benzalkonium chloride Bal 15 cc, 30 cc

Ure: Tear replacement and ophthalmic vehicle

***METHYLCLOTRIAZIDE 6 chlora 3 chloro-**

methyl 2 methyl 7 sulfamyl 3 4 dihydro-T, 2,4 benzothiazidine 1,1 dioxide See Enduron, Tab (Abbott)

W/Cryptenamine See: Diutensan Tabs (Irwin, Neiler)

W/Cryptenamine, rerepine See: Diutensan R, Tabs (Irwin, Neiler)

METHYLACETYLCOLINE. See: Methacholine

METHYLAMINOHEPTANE HYDROCHLORIDE. See: Oenethyl HCl (Knall)

METHYL AMINOPHENYL SULFONAMIDE. See: Sulfbensamine

METHYLAMPHETAMINE HYDROCHLORIDE & SULFATE. See: Desoxyephedrine HCl & Sulfate, Prep

METHYLANDROSTENEDIOL, 17-alpha meth-

yl Δ^8 -androstene 3 beta, 17 beta diol (DI

* Generic name recognized by N.F., U.S.P. or A.M.A.

olasteen Mestenediol, Methanabal, *Me Ibandrial)

Seer Andradial, Tab (Carnrick)

Drolandrone, Tab & Susp. (Carnrick)

Drasene (Ascher)

Esjaydiol, Vial (Testagar)

Melhandrolan, Vial (Lannett)

Methostan, Tab & Vial (Schering)

Nabadiol, Vial (Brean)

Stenediol, Tab & Vial (Organon)

Stembel, Vial, Tab (Bell Pharmaceutical)

W/dl Amphetamine sulf See Amladiol, Tab (Columbus)

W/d Amphetamine sulfate & vitamins See Gerovoidr, Tab (Columbus)

W/dl Desoxyephedrine HCl & estrogenic substances conjugated See: Oromeritral, Tab (Miller)

W/Estrogenic substances, See Neamestrin Vial (Miller)

W/Hydrocortisone, Vial B 12 See: Vistobolil, Tab (Organon)

W/Prednisone, magnesium trisilicate, aluminum hydroxide gel, calcium carbonate magnesium carbonate, guar gum, eat binder (Egraine) See: Stensone, Tab (Organon)

W/Vitamins & minerals

See: Covislan, Tab (Organon)

Gerovoids (Columbus)

METHYL ATROPINE NITRATE. See: Atropine

Methyl Nitrate (Various Mfr)

***METHYLBENZETHONIUM CHLORIDE, USP N N D 1962 Oint USP ADR**

Benzyl dimethyl 2 [2 (p 1,1,3 3 tetra methyl butyl cresoxy)ethoxy] ethyl ammonium chloride (p Tertiary octyl cresoxy ethoxy ethyl dimethyl benzyl ammonium chloride) Chemo Pura, Bulk Paw

See: Ammarid Diaper Rinse (Kinney)

Bactine (Miles)

Diaparene Chloride, H N R (Homemakers)

Hyseptine, Sal (Warren Teed)

Use: Bactericide

W/Neomycin sulfate, phenylephrine HCl & thienylpyramine NCl. See: Entamirl, Sal (Warren-Teed)

W/Phenol, acetanilid, zinc oxide, calamine & eucalyptal. See: Tolain, Oint (Warren Teed)

W/Phenylephrine HCl & thienylpyramine HCl See: Allermir, Sal (Warren Teed).

W/Phenylmercuric acetate, 9 aminoacridine HCl, succinic acid, allantain, lactate See: Vagtab Vag Tab (Ascher)

W/Phenylmercuric acetate, 9-aminoacridine HCl, tyrothricin, urea, succinic acid, lactose

See: Trisert Vaginal, Tab., Paw (Ulmer)

- METHYLBROMTROPIN MANDELATE.** See: Hamotropine Methylbromide
- ***METHYLCELULOSE, U.S.P., N.N.D. 1962, A.D.R.** Cellulose methyl ether, Mellose. Haack—Tab. 0.5 Gm. Bol. 100s, 1000s. Lannett—Tab. 0.5 Gm. Bol. 500s, 1000s. West Ward—Tab. 0.5 Gm. Bol. 500s, 1000-
- See: Bufopto Methalose, Liq. (Professional Pharm.)
- Bulkoid, Tab. (U.S. Standard)
- Cellagran (Tilden)
- Cellathyl, Tab., N.N.B. (Warner-Chilkott).
- Cellumeth, Tab. (Chicago Pharm.)
- Cologel, Liq. (Lilly)
- Gonioscopic Fluid (Crookes-Barnes)
- Hydroflose, Liq. (Upjohn)
- Isopto-Alkaline, Liq. (Alcon)
- Isopto-Pain. (Alcon)
- Lacril, Soln. (Abbott)
- Melozets, Wafer. (Merck Sharp & Dohme)
- Meth-I-Sol (Ophthalmus Inc.)
- Methocel (Dow)
- Melulose (Tri-State)
- Methulose (Prof. Pharm.)
- Methylose, Tab. (Rowell).
- Nicel (Irwin, Neisler)
- Premocel, Tab. (Promo)
- Syncelose, Tab., N.N.R. (Blue Line)
- Tearisol (Iso-Sol)
- Valocel, Tab. (Valentine).
- Uses: (1 to 4 Gm.) Bulk laxative.
- METHYLDINDYDOMOPHINONE HYDRO-CHLORIDE, N.N.D. 1958.** See: Metapon, Cap. (Parke, Davis).
- 2-METHYLDIHYDROTOSTERONE.** See: Adroyd (Parke, Davis)
- ***METHYLDOPA, Levo** 3-(3,4-dihydroxyphenyl)-2-methylalanine. See: Aldomet (Merck and Co., Inc)
- ***METHYLDOPATE NCI,** (Merck Sharp & Dohme) Ethyl ester of levo-3-(3,4-dihydroxyphenyl)-2-methylalanine HCl.
- Uses: Anti-hypertensive agent.
- 3,3'-METHYLENEBIS(4-HYDROXYCOUMABIN).**
- See: Bishydroxycoumarin, U.S.P.
- Dicamarol. (Various Mfr.)
- ***METHYLENE BLUE, U.S.P., N.N.B. 1943.** Inj. U.S.P. (Various Mfr) Methylthionine chloride.
- See: Hexalol (Central)
- Urised (Chicago)
- Uses: Urinary antiseptic.
- ***METHYLEGONOVINE MALEATE, N.F., N.N.D. 1962, Inj., Tab. N.F.** See: Methergine, Amp. & Tab. (Sandax)
- METHYLETHYLAMINO-PHENYLPROPANOL**
- ***METHYLGLUCAMINE DIATRIZOATE, N.N.D. 1962.** Inj. U.S.P. A water-soluble radiopaque iodine cpd. N-methylglucosamine salt of Diatrizoate.
- Uses: Cardiografin
- Gastrografin
- Benografin, Inj. (Squibb)
- ***METHYLGLUCAMINE DIATRIZOATE, U.S.P., N.N.D. 1962.** w/Iodipamide. See: Duografin, Liq. (Squibb).
- W/Sodium diatrizoate, sodium citrate, disodium ethylenediamine tetra-acetate dihydrate, methylparaben, propylparaben See: Benovist (Squibb).
- ***METHYLGLUCAMINE IODIPAMIDE** Inj., U.S.P., N.N.D. 1962. Cholografin Methylglucosamine Inj. (Squibb)
- METHYLGLUCAMINE THEOPHYLLINE** N.N.D. 1962.
- See: Glucophylline (Abbott)
- ***METHYLHEXANEAMINE, N.N.D. 1962.** See: Forthane Inhaler. (Lilly)
- METHYLISO-OCTENYLAMINE.** See: Isomethoplene, N.N.D. 1958.
- 2-METHYL-1,4-NAPHTHOQUINONE.** See: Menadione, U.S.P. (Various Mfr.)
- METHYL NICOTINATE** W/Histamine dihydrochloride, glycol monosalicylate & copicum oleoresin. See: Rubiguent, Cream (Wyeth).
- ***METHYLOL RIBOFLAVIN, N.N.D. 1962.** See: Hyflovin, Amp. (Endo)
- METHYLOSE.** (Rowell) Methylcellulose, Tab. 0.5 Gm. Bol. 100s, 500s, 1000s.
- Uses: (1-2 l.i.d. w/glass of water) Bulk laxative.
- ***METHYLPARABEN, U.S.P., A.D.R.** (Various Mfr.) Methyl p-hydroxybenzoate Methyl Chemasep.
- Uses: Preservative.
- ***METHYLPARAFYNOL** 3-Methyl-pentyl-1-ol-3. Obllivon, Samnesin. See: Dormison, Cap. (Schering)
- METHYLPARAFYNOL CARBAMATE.** See: Obllivon-C. (Brit. Schering)
- METHYLPENTENOL.** See: Methylparafynol.
- METHYLPENTYNOL CARBAMATE.** See: Obllivon-C. (Brit. Schering)
- METHYLPHENETHYLAMINE.** 1-alpha-methylphenethylamine. See: Amphetamine.
- ***METHYLPNINIDATE HCl, N.N.D. 1962.** See: Ritalin (Ciba)
- METHYLPHENIDYLACETATE NCI.** Methyl 1-phenyl-2-piperidylacetate. See: Methylphenidate HCl.
- METHYLPHENOBARBITAL.** N-Methyl-5-ethyl-5-phenylbarbituric Acid. See: Mebaral
- d-METHYLPHENYLAMINE SULFATE.** See: Dextroamphetamine Sulfate, U.S.P. (Various Mfr.)
- * Generic name recognized by N.F., U.S.P. or A.M.A.

- METHYL PHENYLETHYLHYDANTOIN.** See Mesantoin, Tab (Sandoz)
- 3-METHYL-3-PHENYL-4-ISOXAZOLYL PENICILLIN.** See Prostaglin, caps (Bristol)
- 2-METHYL-3-PNYTYL-1,4-NAPHTHO-QUINONE.** See Kanakion (Roche) Phytanadione, USP
- *METHYL POLYSILOXANE.**
Haloxsil (Haller)
See: Mylicon, Tab (Stuart)
W/Belladonna extract butobarbital sodium
See Sibel Tabs (Plough)
Siloin Tablets (Plough)
Silicone
- *METHYLPREDNISOLONE, NND 1962 6**
methyl della 1 hydrocortisone
See Medrol, Tab (Upjohn)
Depo Medrol (Upjohn)
Wyacosi Tab (Wyeth)
W/Acetylsalicylic acid calcium carbonate
See Medadent Tabs (Upjohn)
W/ASA calcium carbonate See Medaprin, Tab (Upjohn)
W/Neomycin sulfate chlorbutanol See: Neo Medrol Soln (Upjohn)
W/Orthoxine See Medrol w/Orthoxine, Tab (Upjohn)
- *METHYLPREDNISOLONE ACETATE, NND 1962 6**
Alpha methylprednisolone 21 acetate
See: Depo Medrol, Inj (Upjohn)
Medrol Acetate (Upjohn)
- *METHYLPREDNISOLONE SOD SUCCINATE, NND 1962 6**
Solu Medrol Mix O Vial (Upjohn)
- *METNYPROMAZINE, 10** (3 dimethyl aminopropyl) 2 methylphenothiazine
- METNYLPYRIMAL** See: Sulfamerazine (Various Mfr)
- *METNYLROSANILINE CHLORIDE, USP**
ADR; Sol USP, Tab, USP (Crystal Violet, Methyl Violet)
Use: (Oral 10 to 60 mg tid; externally 1% soln) Anthelmintic & anti infective
See: Gentian Violet (Various Mfr)
- W/Alkyl dimethylbenzylammonium chloride polyoxyethyleneonylphenol, polyethylene glycol tert didecylthioether See Nyva, Tab (Holland Ranlos Co.)
- *METHYL SALICYLATE, USP, ADR** (Various Mfr) Oil of wintergreen
W/Chloral hydrate & menthol
See Chloral Methylol, Oint (Ulmer)
Chloro Salicylate, Oint (Kremers Urban)
W/Calchicine See Calchi Sol, Cap (Huxley)
W/Menthol, camphor See: Analbalm, Liq (Central Pharm)
- *Generic name recognized by NF USP or AMA
- W/Menthol See: Analgesic Balm (Various Mfr)
- METHYL SULFANIL AMIDOISOXAZOLE**
Sulfamethoxazole See: Gantanol (Rache Lab)
- *METHYLTESTOSTERONE, USP, NND 1962**
Tab USP; Buccal, Tab, USP 17
Methyltestosterone (Anerton, Glosso-Steron dryl, Testovison)
See: Dumogron (Squibb)
Normal, Lingual (Key Corp)
Malesrone, Tab (Kirk)
Malogen, Tab (Testagen)
Masenone, Tab (Schleffelin)
Metandren, Tab (Ciba)
Metrone, Tab (Baire)
M T Mucorelles (White)
Neo Nambriol M, Tab (Organon)
Ovelon M Tab (Schering)
Synandrets, Tab (Pfizer)
Synandrolabs, Tab (Pfizer)
Testobase, Disks & Vial (Merck Sharp & Dahme)
Testora, Tab (Chicago Pharm)
- W/D enestrol
See: Eidone D Loy, Caps (Lloyd, Dabney & Westerfield)
Di Enest, Tab (Kremers Urban)
- W/Dienestrol, reserpine See Testerone, Tab (Lloyd, Dabney and Westerfield)
- W/Dienestrol pentobarbital sod d-amphetamine sulf See: Ampetone, Tab (Kremers Urban)
- W/Diethylstilbestrol
See: Melystil, Tab (Chicago Pharm)
- W/Diethylstilbestrol & reserpine See: Tylon dril, Tab (Lilly)
- W/Estoad See: Testead, Tab (Warren Teed)
- W/Estradiol
See: Di Met, Tab (Organon)
Glynetone, Tab (Schering)
Testaldiol, Tab (Meyer)
- W/Estradiol, pregnenolone & amphetamine phosphate monobasic See: Longevan, Cap (Northwest)
- W/Estradiol & Vit B complex See Comba genic, Tab (C D Smith)
- W/Estrogens See Theelandrol Oral, Cap (Parke, Davis)
- W/Estragenic conjugate See Diorapin, Tab (Standex)
- W/Ethrayl estradiol, d amphetamine sulfate
Gevral formula See Gevralin Cap (Lederle)
- W/Estragenic substance conjugated, d desoxyephedrine HCl, vitamins & minerals
See: Vitramin, Tabsules (Paul Money)
- W/Estragen conjugated (equine) uterin, in

- asitol, methionine, vitamins and minerals.
See: Vio-Geric-H, Tab. (Rowell).
- W/esthane sulfate pat., 1-thyroxine sodium.
See: Vi-Testin, (Sovage).
- W/Ethinyl Estradiol.
See: A-E, Tab. (Warner-Chilcott)
- Duatic, Tab. (Dumas-Wilson)
- Dumane, Tab. (Squibb)
- Duatrane, Tab. (U.S. Standard)
- Femandien linguets (Ciba).
- Gynelone, Repetab. (Schering)
- W/Ethinyl estradiol. See: Para Testrogen, Tabs (Para)
- Testolix, Elix. (Chicago Pharm.)
- W/Ethinyl estradiol, d-amphetamine sulf., nicotinic acid & Vit. C. See: Stimagen, Tab. (Lemmon)
- W/Ethinyl estradiol, metrazol, vitamins, and minerals. See: Senozal, Cap. (Table Rock)
- W/Ethinyl estradiol & multivitamins.
See: Neoban, Cap. (Roerig)
- V-Estandra, Cap. (Funk).
- W/Ethinyl estradiol, Vit. A, D, B-12, thio-mine monanilate, riboflavin, niacinamide, pyridoxine HCl, calcium pantothenate, choline, bitartrate, inositol, ascorbic acid, 1-lysine HCl, Vit. E, rutin, calcium, phosphorus, iron, iodine, copper, potassium, manganese, zinc, magnesium, Vit. B-12 manganese, zinc, magnesium, Vit. B-12 w/intrinsic factor concentrate, lemon bio-flavonoid. See: Harmo-Pluvimin, Caps. (Vitaline)
- W/Ethinyl estradiol, pentylenetetrazol, nicotinic acid, and dicalcium glutamate. See: Senozal, Tab. (Dumas-Wilson)
- W/Ethinyl estradiol, proloid. See: Plestron, Tab. (Warner-Chilcott)
- W/Ethinyl estradiol, thyroid extract, ascorbic acid, calcium lactate, dicalcium phosphate, vit. A, D, E. See: Forstee, Tabs (Brown)
- W/Glutamic acid, thyroid ext. Vit. B-1. See: Andiol, Tab. (Brown Pharmaceutical)
- W/Pentylenetetrazol, ethinyl estradiol, strontium salicylate. See: Analeptane-Anabolic. (Reed & Carnick)
- W/Pentylenetetrazol, niacin, dienestrol. See: Lucidon, Tab. (Lloyd, Dabney & Westerfield)
- W/Pentylenetetrazol, nicotinic acid, Vit. C, ethinyl estradiol, thyroid. See: Niatricrine, Tab. (Asche)
- W/Pentylenetetrazol, nicotinic acid, tetra-foda tyranine, ethinyl estradiol. See: Tetra-Forse, Tab. (E. S. Miller)
- W/Premarin. See: Premarin W/Methyltetra-sterone (Ayerst).
- W/Premarin, d-desoxyephedrine HCl & vitamin. See: Mediatric, liq. (Ayerst).
- *Genesis name recognized by N.F., U.S.P. or A.M.A.
- W/Progesterone, Estradiol. See: Tithormonal, Tab. (Key Corp.)
- W/Rauwolfia serpentina. See: Andio-Walfia (Ortego)
- W/Ritalin, ethinyl estradiol, B-1, B-2, B-6, B-12, nicotinamide, dicalcium phosphate. See: Ritonix, Cap. (Ciba).
- W/Sulestrex piperazine. See: Sulestrex MT, No. 1, 2 & 3 (Abbott).
- W/Tace. See: Tace With Androgen, Cap. (Merrell)
- W/Thyroxine, ethinyl estradiol, d-amphetamine sulf. & vitamins. See: Neoban, liq (Roerig)
- W/Vitamins. See: Testacoids Improved, Tab. (Reed & Carnick)
- W/Vitamins & minerals. See: Gavrine, Cap. (Lederle)
- W/Vitamins multiple. See: Vi-Andro, Cap. (Funk).
- W/Yohimbine HCl. See: Andriobine, Tab. (G. F. Horvey)
- W/Yohimbine HCl, nux vomica. See: Alrodex, Caps. (Layton Labs.)
- METHYLTHIOBROMINE.** See: Coffeine. (Various Mfrs.)
- ***METHYLTHIOURACIL, U.S.P., N.N.D. 1962**
4 Methyl-2 thiourasil Tab., U.S.P. (Anti-bason, Methioclil)
Della—Pow. Bol. 100 Gm.
Physicians Drug & Supply—Tab. 50 mg.
Bot. 100s, 500s, 1000s.
Use: (Orally 150 to 250 mg. daily) Thyroid inhibition.
See: Methioclil, Tab., N.N.R. (Schwarz)
Muracil, Tab. (Organon)
Thimesil, Tab., N.N.R. (Physicians' Drug)
- METHYL VIOLET.** See: Gentian Violet.
- ***METHYPRYLON, N.F., N.N.D. 1962, J-3**
Diethyl-5-methyl-2,4-piperidinedione. Elixir, Tab. N.F. See: Natudar, Elixir & Tab. (Roche) W/Rubber cap. Tab. 1½ gr. Bot 100s.
- METHYROID.** (Lakeside) Thyroid substance, U.S.P. 30 mg. thiamine HCl 3 mg., riboflavin 1 mg., choline dihydrogen citrate 300 mg./Tab. Bot. 100s.
Use: (1-3 daily) Thyroid therapy w/vitamin supplement.
- ***METHYSERGIDE MALEATE, N-[1-(hydroxymethyl)propyl]-1-methyl-D-lysergamide bi-maleate** See: Sanserl (Sandoz)
- METICORTELONE, N.N.D. 1962.** (Schering)
Metacortandracin & prednisolone. Tab. 1.0, 2.5 & 5 mg Bot. 30s, 100s. Sterile Sol. powder (Prednisolone sod. hemisuccinate 50 mg/5 cc) Vial, Box 1s, 25s.
Use: Rheumatoid arthritis.

W/Chlor trimeton gluconate See: Metrelon, Nasal Spray (Schering)

METICORTELONE ACETATE, (Schering) Prednisolone acetate, Susp 25 mg/cc Vial 5 cc
Use: Intra articular Injection

METICORTEN, NND 1962 (Schering) Prednisone, Tab 10, 25 & 5 mg, Bot 30s, 100s

Use: Rheumatoid arthritis
W/Chlor trimeton maleate, ascorbic acid See: Metrelon, Tab (Schering)

METI-DERM, NND 1962 (Schering) Prednisolone 5 mg/Gm of cream Tube 10 Gm & 25 Gm Aerosol (50 mg) Can 150 Gm
Use Topically, anti-inflammatory

W/Neomycin (Schering) Prednisolone 5 mg neomycin sulf 5 mg/Gm Tube 10 Gm, & 25 Gm Aerosol Cont 50, 150 Gm

METIMYO OINT, W/Neomycin (Schering) Prednisolone acetate 5 mg, sulfacetamide sod 100 mg, neomycin sulf 25 mg/Gm Tube ½ oz w/applicator tip

Use: Inflammatory eye infections
METIMYO OPHTH, SUSP (Schering) Prednisolone acetate 5 mg, sulfacetamide 10%/cc Bot dropper 5 cc

Use: Inflammatory eye infections
METIOGRAN, (Lobica Debrulle) Methionine Bot 30 Gm

Use: (1 tsp daily) Dermatitis
METIONE, NND 1962 (Lobica Debrulle) Methionine Envelope, 2 Gm Box 20s
Use: Dietary supplement

METOL, (New York Quinine) p Methylamino phenol sulfate Bot 1 oz, ¼ lb, 1 lb

METOPYRONE (Ciba) Metyrapone Methopirapone 250 mg/Tab or methopirapone ditartrate 100 mg/10 cc amp Tab Bot 18s 10 cc amp/2s

Use: Diagnosis of suspected hypopituitarism

METOPON HYDROCHLORIDE, NND 1958

Methylidihydramorphinane HCl Parke Davis—Cap 3 mg Bot 100s
Sharp & Dahme—Cap 3 mg Bot 100s
Use: Analgesic

METOQUINE See Quinacrine Hydrochloride (Atabrine HCl Winthrop Labs)

METORBIC, (Elder) Ascorbic acid 30 mg, methionine 200 mg/Cap Cap Bot 100s

Use: Fortify wound healing
METOSO, (Abbott) Metaphen 12000 sol Sol, Bot 8 fl oz, 1 pt 1 gal

Use: Antiseptic gargle

METRANIL, (Meyer) Pentaerythritol tetranitrate 20 mg/Tab Bot 100s Caps 80 mg Bot 60s

Use: Vasodilation

METRANIL-AM DURACAP, (Meyer) Pentaerythritol tetranitrate 30 mg, amobarbital 50 mg/Cap Bot 100s

Use: Vasodilation depression

METRAZOL, N N B 1950 (Knoll) Penta methylenetetrazol, Pentylene tetrazol, USP Amps 1 cc, 1½ gr, Box 6s, 100s; 3 cc, 4½ gr, Box 5s.

Pow: Bot 5, 25, 100 & 250 Gm
Sterile sol: 10%, 30 cc & 100 cc, rubber capped vial

Tab: 1½ gr, Bot 100s 500s

Use: Analeptic & antidote to the barbiturates

W/Barbiturates See: Tetbarbols (Strong & Cobb)

W/Methyl testosterone, ethinyl estradiol, vitamins, and minerals See: Senazol, Cap (Table Rock)

W/Vitamins See: Vito Metrazol, Elix (Knoll)

W/Vit B1 See: Metrazol Liquidum, Liq (Knoll)

METRAZOL LIQUIDUM (Knoll) Metrazol 1½ gr, Vit B1 1 mg, in flavored wine elixir/5 cc Bot 1 pt
Use: Central nervous system stimulant

METRECAL (Mead Johnson) Protein 70 Gm, fat 20 Gm, carbohydrate 110 Gm, minerals 145 Gm, Vit A 5000 u, D 400 u, C 100 mg, B1 2 mg, B2 3 mg, niacin 18 Gm, iron 15 mg, iodine 150 mcg, pyridoxine HCl 2 mg, folic acid 0.5 mg

calcium pantothenate 10 mg, vitamin E 10 u, B12 2 mcg, potassium 35 Gm sodium 0.9 Gm copper 1.5 mg, manganese 2 mg, zinc 5 mg/½ lb powder in a 3 can Starter Pak Also liquid and wafers

Use: Food supplement

Use: Mix with water for weight control diet

METRETON, (Schering)

Tab—Meticorten 25 mg, chlor trimeton maleate 20 mg, ascorbic acid 75 mg/Tab Bot 30s, 100s

Nasal Spray—Meticortelone acetate 20 mg, chlor trimeton gluconate 30 mg/cc Bot 15 cc

Ophth Susp—Meticortelone acetate 2 mg, chlor trimeton gluconate 3 mg/cc Dropper Bot 5 cc

Use: Caricaid antihistamine therapy

METROCAPS, (Strassenburgh) Metyrapone 1/60 gr, aspirin 3½ gr, acetaphenetidin 2½ gr, caffeine ¼ gr/Cap Bot 100s, 500s 1000s

Use: Orally (12 q 3 h) analgesic & anti-rheumatic

METROCRINE, (Metropalton Labs) Thyroid 1 gr, adrenal cortex fresh gland 3 gr anterior pituitary fresh gland 5 gr, ovarian

- substance, fresh gland 15 gr., posterior pituitary, U.S.P. 1½ ml./2 cc. Viol 2 cc. Use: Thyroid therapy.
- METRONE.** (Barre) Methyltestosterone. Tab. 10 mg, 25 mg. Bot. 100s.
- Use: Male gonadal replacement therapy.
- **METRONIDAZOLE.** 1-(2-hydroxyethyl)-2-methyl-5-nitroimidazole. See: Flagyl (G. D. Searle)
- METROPECTIN.** (Strassenburgh) Aluminum hydroxide 2 gr., metropies 1 mg., pectin 1 gr., osmakaolin 7½ gr., beta-lactose 1 gr./Tab. Bot. 100s, 500s, 1000s.
- Use: Orally (1-2 q. 3 h.), gastric hyperacidity.
- METROPINE.** (Strassenburgh) (Atropine Methylnitrate, Methyl Atropine Nitrate.)
- Elxir: (1 mg./4 cc) Bot. 1 pt., 1 gal. w/ sropalamine, Bot. 1 pt., 1 gal.
- Tab.: (1 mg) Bot. 100s, 500s, 1000s; w/ phenobarbital, ¼ gr., Bot. 100s, 500s, 1000s.
- Viol. (2 mg./cc.) 10 cc., 6s.
- Use: Antispasmodic.
- W/Alapectase. See: Alapectase W/Metropies, Tab. (Strassenburgh).
- W/Aluminum hydroxide, beta-lactose, pectin & osmakaolin. See: Metropectin, Tab. (Strassenburgh)
- W/di-Amphetamine phosphate, acetyl-p-aminophenol, salicylamide, phenyltolaxamine dihydrogen citrate. See: Acogest, Cap. (Strassenburgh)
- W/Amphetamine, sod. CMC. See: Biphetocel, Tab. (Strassenburgh)
- W/A.P.C. See: Metrocaps, Cap. (Strassenburgh)
- W/Duodenum paw., aluminum hydroxide & Mag trisilicate. See: Metracel, Tab. (Strassenburgh).
- METUBINE IODIDE, N.N.D. 1962.** (Lilly) Dimethyltubocurarine iodide. Amp. (2 mg./cc.) 20 cc. rubber-stoppered amp.; (1 mg./cc.) 10 cc. rubber-stoppered amp.
- Use: Skeletal-muscle relaxant.
- **METUREDEPA.** (Armour) ethyl[bis(2,2-dimethyl-aziridinyl)phosphinyl]-carbamate. See: Turtac
- METYCAINE HYDROCHLORIDE, N.N.R. 1955, A.D.R.** (Lilly) (Piperacaine HCl, Neohesin HCl) di-gamma-methylpiperidinopropyl benzoate HCl.
- Amp. (rubber stoppered) for regional & local infiltration anesthesia. (1%) 30 cc., Pkg. 6s, 25s, 100s.
- Amp. for spinal anesthesia: (1.5%) 5 cc., Pkg. 6s, 25s, 100s.
- Amp. (rubber stoppered) for caudal anesthesia: (1.5%) 200 cc.
- Generic name recognized by N.F., U.S.P. or A.M.A.
- Amp. (rubber stoppered) for regional and local infiltration anesthesia: (2%) 30 cc., Pkg. 25s.
- Amp. for epinal anesthesia: (5%) (150 mg / 3 cc.) Pkg. 6s, 25s, 100s.
- Amp. for epinal, regional, infiltration & topical anesthesia: (20%) 5 cc., Pkg. 6s, 25s, 100s.
- Olar.: (5%) Tube ½ oz.
- Ophth. aint.: (4%) Tube ¼ oz.
- Paw.: Bot. ½ & 4 oz.
- Sol.: (2%) Bot. 1 oz.
- Tab.: (0.15 Gm) Bot. 100s
- Use: Local anesthetic.
- W/Dextrose & epinephrine. (Lilly) Amp., Metycaine HCl 1e 5% dextrose w/epinephrine HCl 1:2500, 1 cc., Pkg. 6s, 25s, 100s.
- Use: Saddle block & minimal spinal anesthesia.
- W/Merthiolate. (Lilly) Ophth. aint. 4%, w/ Merthiolate 1:5000 Tube ½ oz. Supp. 5%, w/Merthiolate 1:5000 Box 12s.
- Use: Antiseptic & local anesthetic.
- W/Zinc oxide. (Lilly) Metycaine 125 mg., ballodonna ext. 12.5 mg., w/zinc oxide, bismuth subcarb./Supp. Box 12s.
- Use: Rectally antiseptic.
- **METTRAPONE.** (Ciba) 2-methyl-1,2-di-3-pyridyl-1-propanone. See: Metapirone.
- METYSTIL.** (Chicago Pharm.) Methyltestosterone 5 mg, diethylstilbestrol 0.25 mg./Tab. Bot. 30s, 100s, 1000s.
- Use: Estrogen & androgen therapy.
- MEVATINIC-C.** (Bentlich, Inc.) Exsiccated ferrous sulf. 200 mg, cabalamin (B-12 activity) 5 mcg, liver stomach cont. 50 mg, folic acid 0.5 mg., ascorbic acid 225 mg., hesperidine 30 mg., copper 0.33 mg, cobalt 0.1 mg., molybdenum 0.1 mg., zinc 0.4 mg./Tab. Bot. 100s.
- Use: Anemias.
- MEVTON.** (Tablet Rock) B-1 2.4 mg., B-6 1 mg., Vitamin B-12 16 mcg, lysine monohydrochloride 300 mg, metrazole 200 mg., C 100 mg, niacinamide 30 mg., B-2 4 mg., folic acid 0.5 mg., panthenol 6 mg., cobalt 0.2 mg., zinc 1 mg., iodine 0.5 mg., magnesium 4 mg., manganese 2 mg., alcohol 10% in a D-sorbitol base/or. Bot. 1 pt., gal.
- Use: Dietary supplement.
- MEKAL.** (Massengill) Mannitol hexantrate ½ gr., phenobarbital ¼ gr./Tab. Bot. 100s, 500s, 1000s
- Use: Prolonged vascular dilatation
- MEZO.** (Warren-Tee) Phenol 4½%, campher 3%, benzocaine 0.04%, zinc oxide

- 37.5%, menthol 0.15%. Tube 1½ oz; Jar 12½ oz, 25 oz, 7½ lb
 Use: Antiseptic ointment
- MIADONE.** See: Methadone
- MIADYNE.** (Hart) Niacin ¾ gr, aspirin 4 gr, acetophenetidin 2 gr, caffeine ½ gr / Tab Bot 100s, 500s, 1000s
 Use (2 q 3 h) Analgesic, sedative & vasodilator
- MIAQUIN.** See Camoquin (Parke, Davis)
- MICAJAH WAFER** (Micojah) Medicated wafers containing alum, aluminum sulfate Sod sulfate & boric acid Bot 25s
 Use: Intervaginal antibacterial
- MIC-CERRIN.** (Lilly) Thiamine mononitrate 10 mg, riboflavin 5 mg, pyridoxine HCl 2 mg, pantothenic acid (as Cal pantothenate) 10 mg, nicotinamide 30 mg, Vit B 12 (activity equiv) 3 mcg, folic acid 0.1 mg, ascorbic acid 100 mg, vit E 5 mg, vit A 10 000 u, D 1000 u Also containing iron, copper, iodine, cobalt, boron, molybdenum, pot zinc/Tab Pkg 60s, 100s, 1000s
 Use: Vitamin & mineral deficiencies
- MIC-CERIN T.** (Lilly) Vit B 1 15 mg, B 2 10 mg, B 6 2 mg, pantothenic acid 10 mg, nicotinamide 100 mg, B 12 7.5 mcg, folic acid 0.2 mg, ascorbic acid 150 mg, alphatocopherol 5 mg, Vit A 25 000 u, D 1000 u, iron 15 mg, copper 1 mg, iodine 0.15 mg, cobalt 0.1 mg, boron 0.1 mg, manganese 1 mg, magnesium 5 mg, molybdenum 0.2 mg, potassium 5 mg, zinc 1.5 mg/Tab Bot 30s, 100s, 1000s
 Use: Therapeutic vitamin minerals
- MICOFUR.** (Anti S-Nitro-2 Furoldoxime) Nitrofurantoin See: Tricofuran Prep (Eaton)
- MICRAFORM SULFATHIAZOLE SUSPENSION 20%.** (Smith, Kline & French) Susp of sulfathiazole 20% w/thimerosal 1 20 000 Bot 25 cc
 Use: Pyogenic skin infections
- MICRIDILUM** (Johnson & Johnson) Phenocridine (p hexyloxyphenyl)-10 methyl acridinium chloride
 Use: Skin disorders in infants and children
- MICRIN** (Johnson & Johnson) Cetyl pyridinium chloride dequalinium acetate oil of peppermint menthol, alcohol 15%. Oral antiseptic 7 oz, 14 oz
- MICRO-ACETARS.** (Brewer) A mixture of sod nitro-prusside & amm sulfate Kits of 50 tabs, a bottle of ammonia water & two droppers Bot 500 tabs, 1000 tabs
 Use: Detection of acetone in urine
- MICRODIOL.** (Gold Leaf) Estradiol aq susp Vial (0.1 mg/cc) 10 cc., 30 cc (0.2 mg/cc) 10 cc., 30 cc (0.5 mg/cc) 10 cc., 30 cc
 Use: I.M., estragen therapy
- MICROEST.** (Mossengill) Diethylstilbestrol 0.5 mg, 1 mg, 5 mg, 25 mg/Tab 0.5 mg, 1 mg Bot 100s, 1000s 5 mg Bot 100s, 500s, 1000s 25 mg Bot 100s, 500s
- MICROSULFON.** (Consolidated Mid) Sulfadiazine 0.5 Gm /Tab Bot 30s, 100s, 250s, 1000s
 Uses: Gram positive bacteria infections
- MICROZINE SUSPENSION.** (Horton & Converse) Microcrystalline sulfadiazine 20% in aqueous suspension Bot 1 oz
 Use: Topically, sulfonamide therapy
- MICROZOLE OINTMENT** (Hartan & Converse) Microcrystalline sulfathiazole 5% in bland base Pkg 1 oz, 1 lb
 Use: Topically, sulfonamide therapy
- MICTINE.** NND 1959 (Searle) New name Mincard Aminametradine; 1 ethyl 6 aminotetrahydropyrimidinedione Tab (200 mg) uncoated, Bot 100s
 Use: Diuretic
- *MIDAMALINE HCl** N (5 chloro 2 benz imidazolylmethyl) N phenyl NN' dimethyl ethylene diamine HCl
- MIDICEL.** NND 1962 (Parke, Davis) 3 Sulfonamide 6 methoxy pyridazine Sulfomethoxy pyridazine 0.5 Gm /Tab Bot 24s 100s
 Use: Sulfonamide therapy
- MIDICEL ACETYL SUSP.** (Parke, Davis) N'-acetyl sulfomethoxy pyridazine eq to Midicel 250 mg /5 cc Bot 4 oz
 Use: Bacterial infections
- MIDICEL SENSITIVITY DISCS** (Parke, Davis) Pkg 50 mcg, 150 mcg, 300 mcg Bot 50r
 Use: Determining qualitatively the sensitivity of micro organisms to the action of the sulfonamide
- MIDRAN.** (Columbia Medical) Phenylephrine HCl, phenylpropanolamine HCl chlorpheniramine maleate salicylamide, phenacetin caffeine & Vit C Tabs Bot 24s, 60s 100s
 Use: Analgesic, antihistaminic, & decongestant
- MIDRIN** (Carrick) Isametheptene mucate 65 mg, N acetyl p aminophenol 325 mg, dichlorophenazone 100 mg /Cap Bot 50s
 Use: Prevention of migraine
- MIDOL** (Sterling Drug) Cinnamylphedrine with APC /Tab Bot 100s
 Use: Antispasmodic analgesic
- MIOESTERONE** (Dursi) Progesterone 10 IU in propylene glycol/cc. Vial 10 cc
 Use: I.M., corpus luteum therapy
- MIGRADE.** (Crystal) Salicylamide 3 gr, meth-

- yliso-octenylamide mucate 1 gr., acetaminophen 3 gr./Tab. Bot. 100s.
Use: Anti-migraine.
- MIGRAINE RECTAL SUPPOSITORIES.** (Organon) Ergotamine tartrate 1 mg., caffeine 100 mg., levorotatory belladonna alkaloids 0.1 mg., acetophenetidin 130 mg./Supp. Box 12.
Use: Migraine.
- MIGRAL.** (Burroughs Wellcome) Ergotamine tartrate 1 mg., cyclizine HCl 25 mg., caffeine 50 mg./Tab. Bot. 20s, 100s.
Use: Migraine.
- MIGRAL PORTE.** (Burroughs Wellcome) Ergotamine tartrate 2 mg., marezine 50 mg., caffeine 100 mg./Tab. Bot. 20s.
Use: Headache.
- MIKAROL.** (Dumas-Wilson) Diethylsulbestrol. Tab. 1 mg., 5 mg., 25 mg. Bot. 100s, 1000s
Use: Orally (5-25 mg. i.i.d.), estrogen therapy.
- MIKEDIMIDE.** (Parlam) Methetharimide. 3-methyl-3-ethylglutaramide. Bamegride. Megimide in Propylene glycol. Vials (30 mg/cc) 10, 60 & 120 cc.
Use: Barbiturate antidote
- MIKROGEN.** (Ourst) (Estradiol 0.20 mg., estrogenic substance 0.20 mg., propylene glycol, benzyl alcohol & chlorobutanol/cc.) Vial 10 cc.
Use: Estrogen therapy.
- MILD SILVER PROTEIN.** N.N.R. 1949. See: Silver Protein, Mild.
- MILESTROL.** (E. S. Miller) Cryst. synthetic estrogen. Tab. 5 mg., 25 mg., Bot. 100s, 500s, 1000s. Vial (25 mg./cc.) 10 cc. Box 1s, 6s, 25s
Use: Estrogen therapy.
- MILIBIS, N N O.** 1962. (Winthrop Labs.) Glycobiarsol, N.P., bismuth glycolylarsanilate. Tab. 0.25 Gm., 0.5 Gm. Bot. 25s, 500s.
Use: (2 i.i.d. for one week) Intestinal amebiasis.
- MILIBIS W/ARALEN DIPHOSPHATE TABLETS.** (Winthrop Labs) Milibis (brand of bismuth glycolylarsanilate) 0.25 Gm., Aralen (brand of chloroquine) diphosphate 75 mg./tab. Bot. 50s, 1000s.
Use: Prophylaxis & treatment of Intestinal & extra intestinal amebiasis.
- MILIBIS SUPPOSITORIES.** (Winthrop Labs.) Each supp. contains 0.25 Gm. Milibis in gelatin-glycerine base. Box 10s.
Use: Vaginitis of trichomonal, monilial, bacterial or mixed etiology.
- MILIDEX.** (Bonar) Dextroamphetamine sulfate 15 mg., amobarbital 60 mg./Cap. Bot. 100s, 1000s
Use: Appetite control.
- MILIFERIN INJECTION.** (E. S. Miller) Liver injection, U.S.P. 5 mcg., vitamin B-12, U.S.P. 25 mcg., ferric ammonium citrate, N.P. 100 mg., thiamine HCl 10 mg., riboflavin 2 mg., pyridoxine HCl 1 mg., d-pantothenyl alcohol 2 mg., niacinamide 25 mg., dibucaine HCl 0.04%/2 cc. Vial 10 cc.
MILK OF BISMUTH, N.F., N.N.R. 1941. (Various Mfr) Bismuth hydroxide & bismuth subcarb.
See: Crema-Bismuth, Susp. (Merck Sharp & Dohme)
Lac-Bisma, Susp. (Hart)
Use: Orally, intestinal disturbances.
W/Paregoric, kaolin & pectin. See: Mul-Sed, Susp. (Pitman-Moore).
- MILK, COW, MODIFIED.** See: Baker's Modified Milk. (Baker)
- MILKINOL.** (Kremers-Urban) Diacetyl sodium sulfosuccinate and liquid petrolatum specially processed for instant aqueous mixing Bot. 12 oz.
Use: Laxative.
- MILK OF MAGNESIA, A.D.R.** See: Magnesia Magma, U.S.P.
- MILK POWDER DEFATTED W/1-Lysine HCl & pyridoxine HCl.** See: Lysidox, Pow. (Gray)
- MILK PROTEIN DEFATTED SUSPENSION.** See: Kirkolac, Vial (Kirk).
W/Sulfur & peplone. See: Endogen-A, Amp (Endo).
- MILK OF TRINISIUM.** (Abbott) Aluminum hydroxide 1.25 gr., Mag trisilicate 7.5 gr./fl. dr. Bot. 12 oz.
Use: Orally (1 dr. pr. n), antacid.
- MILL-A-SED.** (E. S. Miller) Sod. phenobarbital 1 gr., Sod. pentobarbital 1 gr./Cap. Bot. 100s, 500s, 1000s
Use: Orally; sedative.
- MILLER-DRINE.** (E. S. Miller) dl Desoxyephedrine. Tab. 5 mg., 10 mg. Bot. 100s, 500s, 1000s
Use: Orally; central nervous system stimulant.
- MILLER-ZYME.** (E. S. Miller) Ext. ox bile 120 mg., pancreatin 120 mg., duodenal subst. 60 mg., papain 60 mg./Tab. Bot. 100s, 500s, 1000s
Use: Choleretic.
- MILONTIN, N N D.** 1962. (Parke, Davis) Phensuximide. N-methyl-alpha-phenyl-succinimide Kopsels 0.5 Gm. Bot. 100s
Use: (1-2 i.i.d.) Anticonvulsant.
- MIEONTIN SUSPENSION.** (Parke, Davis) Milantin 250 mg./4 cc. Bot. 1 pt.
Use: Anticonvulsant.
- MILPATH.** (Wallace) Meprobamate 400 mg. or 200 mg. trihexethyl chloride 25 mg./Tab. Bot. 50s.
Use: (1 i.i.d.) Reduces pain and promotes normal G. I. function

- MILPREM.** (Wallace) Meprobamate 400 mg, estrogens conjugated 0.4 mg / Tab Bot 60s
Use: Treatment of menopause symptoms
- MILPREM-200** (Wallace) Meprobamate 200 mg, estrogens conj 0.4 mg / Tab Bot 60s
- MILTON ANTIBACTERIAL SOLUTION.** (Walker) 1% sodium hypochlorite Bot 16 oz
Use: Sterilize nursing bottles, nipples, utensils
- MILTOWN, NND 1962** (Wallace Labs) Meprobamate 2 Methyl 2 N propyl 1,3 propanediol dicarbamate Tab 200 & 400 mg Bot 50s See also Meprasan
Use: Muscle relaxing & ironquizing agent
W/d amphetamine sulfate See Appetrol (Wallace)
- MILTRATE,** (Wallace Labs) Meprobamate 200 mg, pentaerythritol tetranitrate 10 mg / Tab Bot 50s
Use: Angina pectoris
- MITBONAL** (Searle) Cinnasizine
Use: Antihistaminic
- MILU-B-COMPLEX,** (Smith, Miles & Patch) Thiamine HCl 100 mg, B₆ 4 mg, B₁₂ 4 mg, d Pantothényl Alcohol 4 mg, Niacinamide 100 mg / 2 cc Vials 30 cc
- MIL-U-CAL,** (Miller) Cal levulinate 10% sol Amp 10 cc Box 6s, 50s 100s
Use: Calcium deficiency & allergic manifestatons
- MILURETIC** (Wallace) Meprobamate 200 mg, hydrochlorothiazide 25 mg / Tab Bot 50s
Use: Hypertension
- MINABAR.** (Minette) Tab Bot 100s
- MINABEE W/C** (Minette) Tab Bot 100s
- MINABEL.** (Minette) Tab Bot 100s
- MINACAP** (Upjohn) Vit B 12 2 mcg, thiamine HCl 2 mg, riboflavin 2 mg, pyridoxine HCl 1 mg, Vit C 333 mg, Cal pantothenate 2 mg, nicotinamide 10 mg, folic acid D 67 mg, Vit A 2000 u, D 400 u, ferrous sulfate 25 mg, Cal lactate 590 mg plus 7 minerals / Cap Bot 100s, 500s
Use: Multivitamin therapy
- MINAGEST.** (Minette) Tab Bot 100s
- MIN-AMIN.** (Necco) Vit A 5000 u, D 500 u, B 1 2 mg, B 2 3 mg, dicalcium phosphate 480 mg / 12 Gm or 3 waters Cans 5 oz, 20 oz
Use: Dietary supplement
- MINERAL-CORTICOIDS.** See: Desoxycorticosterone
- MINCARD, NND 1962** Aminametradine See: Michne, Tab (Searle)
- MINCOLATE.** (Testagar) Calcichne 1/250 gr, sod salicylate 5 gr, Vit C 15 mg / Tab Bot 100s
Use: Gout
- MINERALIZED VITAMIN.** (Walker Vitamin) Vit A 5000 u, thiamine 3 mg, riboflavin 2 mg, Cal pantothenate 1 mg, niacinamide 20 mg, ascorbic acid 30 mg, Vit D 500 u, Cal, Phos, Iron, Mag, Mang, zinc / Tab Box 50s, 250s, 500s, 1000s
Use: Dietary supplement
- MINERAL OIL.** See: Petrolatum, Liq, USP
- MINERALAX.** (Suliff & Case) Heavy mineral oil 18 cc, Phenolphthalein 260 mg, / fl oz Liq Bot pt, 12 pts, gal
- MINEROVIT.** (Consolidated Mid) Vit A 10000 u, D 1000 u, C 50 mg, B 1 3 mg, B 2 2.5 mg, B 6 0.75 mg, E 1 U, niacinamide 15 mg, calcium pantothenate 35 mg, dicalcium phosphate 170 mg, iron 15 mg, cobalt 0.1 mg, copper 1.5 mg, iodine 0.1 mg, magnesium 1.5 mg, manganese 1.5 mg, zinc 0.7 mg / vitasphere, Bot 30s 100s 1000s, 5000s
- MINIHIST,** (Ives-Cameron) Pyranisamine maleate Tab 25 mg Bot 50s
Use: (11d) Antihistaminic
- MIN-10-PHEN.** (Paul Maney) Phenobarbital 1/4 gr, neohyline 2 gr, potassium iodide 2 gr / Tab Bot 100s 1000s
Use: Congestive heart failure
- MINIVITE** (Central) Vit A 5000 u, D 1000 u, B 1 2.5 mg, B 2 2.5 mg, niacin 10 mg, Vit B 6 20 mg, Cal pantothenate 5 mg, Vit C 30 mg, B 12 1.5 mg, folic acid 0.5 mg, Vit E 2 mg, w/minerals / Cap Bot 100s, 500s
Use: Vitamin w/mineral therapy
- MINUKOL.** (Elder) Pyrimamine maleate 25 mg, dl methamphetamine HCl 2.5 mg / Tab Bot 100s, 1000s
Use: Orally (1-2 q 4 h), antihistaminic w/ central nervous system stimulant
- MIN-VATINE.** (Smith Dorsey) Vit A 5000 u, D 1000 u, B 1 3 mg, B 2 2.5 mg, B 6 65 mg, C 50 mg, B 12 1.5 mcg, folic acid D 33 mg, Vit. E 3 mg, pantothenic acid 5 mg, niacin 20 mg, dicalcium phosphate, ferrous sulfate Pot iodide, Mang-copper, cobalt, zinc, Pot, ma Bot 100s, 500s, 1000s
Use: Vitamin & mineral therapy
- MIOCEL BUEOPTO** (Professional Co) Physostigmine salicylate 1/8%, prne HCl 2% and phenyl mercuric Bot dropper 5 & 15 cc
Use: Control and treatment of and miotic
- MIKON SODIUM, NND 1962** (rodt Chem) Sod Dipratrizoate planamide 2 4 6-trisodobenzoate

- 50%, Amp. 20 cc., Box 5s, 20s, Amp. 30 cc., Box 1s, 5s, 20s.
Use: Diagnostic agent.
- MIO-PRESSIN.** [Smith, Kline & French]
No. 1: Bouwffia serpentina whole root 12.5 mg., praloveratrine 0.1 mg., phenazobenzamine HCl (Dibenzylsine) 2.5 mg./Cap. Bot. 100s.
No. 2: Bouwffia serpentina whole root 25 mg., praloveratrine 0.2 mg., phenazobenzamine HCl 5 mg./cap. Bot. 100s.
Use: Hypertension.
- MIRADON.** [Schering] Antsindione (2-p-anisyl Indanadione-1,3). 50 mg./Tab. Bot. 50s, 100s.
Use: Anticoagulant.
- MISTLETOE.** (*Viscum album*) W/Sad. nitrite.
See: *Visca Modified*, Tab. (Pitman-Moore)
- MISTRONE.** (E. S. Miller) Estrogens natural 2.5 mg., progesterone 25 mg./cc. Vial 10 cc. Box 6s, 25s
Use: Hormone deficiency.
- MITRONAL.** (G. D. Searle) Cinnarizine.
Use: Antihistaminic.
- MITY-MYCIN.** (Reid Labs) Neomycin base 3 mg., polymyxin B sulfate 4,000 u., tyrothricin 0.05%, gramicidin 0.01%/Gm. Oint. M.C. contain in addition hydrocortisone (as acetate) 5 mg./Gm. Oint
- MIXASE A.** (Armour) Proteolytic enzymes from mammalian pancreas present in Kymor.
- MIXED YOCOPHEROL ACETATE.** (Parke, Davis) Vit. E acetate. Cap. 100 mg. Bot. 100s, 1000s.
Use: Orally (100 mg daily), habitual & threatened abortion.
- MIXED VITAMIN CAPSULES.** (Mead-Johnson) Vit. A 10,000 u., D 1000 u., thiamine 3 mg., riboflavin 4 mg., niacinamide 25 mg., ascorbic acid 100 mg./Cap. Bot. 25s, 100s.
Use: Vitamin supplement.
- MIXIBO.** (Durst) Ferrrous sulfate, exsic. 100 mg., ferrous gluconate 100 mg., reduced iron 200 mg., copper gluconate 7.5 mg. Vit. B-12 1.5 mcg./Tab. Bot. 100s, 500s, 1000s.
Use: Hypochromic microcytic anemias.
- MIXTURE 6T2.** See: Dimethyl Phthalate Solution, Compounds, N.F.
- M-MINUS-S.** (Whittier) Pamobrom 50 mg., acetophenetidin 100 mg./Tab. Bot. 24s, 100s
Use (1 t.t.d.) Premenstrual tension.
- MOAN ANTIGEN.** (Mabac) Aqueous extract of *E. histolytica* added to red blood corpuscles
Use: To test for Amebic infection.
- MOBINOL.** See: Tolbutamide.
- MOBISYL.** (B. F. Ascher) Acetylsalicylic acid 0.25 Gm., para-aminobenzoic acid 0.25 Gm., ascorbic acid 25 mg., physostigmine salicylate 0.25 mg., homatropine methylbromide 0.50 mg./Tab. Bot. 100s, 1000s
Use: Analgesic muscle relaxant
- MOBIFYN.** (Elder) Acetophenetidin 1/2 gr., belladonna leaves 1/40 gr., camphor monobromated 1/10 gr., gelsemium 1/40 gr., aconite 1/40 gr., bryonia 1/20 gr./Tab. Bot. 100s, 500s, 1000s
Use: Analgesic and antispasmodic
- MOCCASIN BITE.** See: Antivenin, Snake, Polyvalent.
- MODANE.** (Warren-Teed) Danthron (1,8-dihydroxyanthraquinone) 75 mg., d-calcium pantothenate 25 mg./tab. Bot. 100s, 500s, 1000s. Liq. (1 tspful is equivalent to 1 Modane Mild, Tab.) 1 pt.
Use: Orally (1-2 h.s.), laxative
- MODANE MILD.** (Warren-Teed) Danthron 37.5 mg., d-col. pantothenate 12.5 mg./Tab. or 5 cc. Bot. 100s. Liquid also available
Use: Nutritional decongestant.
- MODERIL.** N.N.D. 1962. (Pfizer) Rescinamine Tab. (0.25 mg.), Bot 100s, 500s, (0.5 mg) Bot. 100s. Elix (0.5 mg/5 cc) Bot 1 pt.
Use: Hypertension & tension
- MODILAC.** (Gerber) Protein 0.65 Gm., fol 0.83 Gm., carbohydrate 2.47 Gm./fl. oz. Each quart after dilution contains Vitamin A 3000 u., D 600 u., C 45 mg., B-6 0.7 mg., thiamine 0.55 mg., riboflavin 1 mg., calcium 800 mg., phosphorus 605 mg Pkg. 12 oz
Use: Milk formula for infant feeding
- MODINAL.** See: Gordinol Type Detergents. (Various Mfr.)
- MODUMATE INFUSION.** (Abbott). L-arginine, 13.5 Gm; L-glutamic acid, 12 Gm. (including 0.5 Gm. excess glutamic acid) (25% (w/v) arginine glutamate)/100 ml. 100 ml sterile containers.
Use: Treatment of ammonia intoxication due to hepatic failure.
- MODUTROL.** (Reed & Carnrick) 1-Piperidine ethanol benzilate HCl 2 mg., scopolamine methylnitrate 1 mg., magnesium hydroxide 200 mg., aluminum hydroxide 200 mg./Tab. Bot 50s, 100s.
Use: Peptic ulcer.
- MOEBIQUIN.** (Consol. Midland) Dihydroxyquinoline 210 mg./tab. Bot 30s, 100s, 250s, 1000s.
Use: Intestinal amebiasis.
- M.O., HALEY'S.** See: Haley's M.O. (Phillips)

- *MOLINAZONE** (Riker) 3 morphina 1 2
3 benzotriaz n 4(3H) one
Use Analgesic
- MOL-IRON CHRONOSULES** (White) Molybdenum oxide 6 mg ferrous sulfate 390 mg/Cap Bot 30s
Use Iron deficiency anemias
- MOL-IRON DROPS**, (White) Ferrous sulfate 125 mg (elemental iron 25 mg), molybdenum oxide 2 mg, alcohol 4.75%/cc Bot w/calibrated dropper, 15 cc, 50 cc
Use: Iron deficiency anemias
- MOL-IRON E M F**, (White) Ferrous sulfate 195 mg, molybdenum oxide 3 mg, Vit B 12 activity equiv 100 mcg, gastric sub stancia 250 mg folic acid D 85 mg ascorbic acid 50 mg desic liver 100 mg/ Cap Cap, Bot 100s, 1000s
Use Anemias
- MOL-IRON PANHEMIC** (White) Ferrous sulfate 500 mg, molybdenum oxide 77 mg, Vit B 12 w/intrinsic factor concn frate 1 USP unit, Vit B 12 77 mcg, folic acid 2.5 mg Vit C 75 mg, B 1 2 mg B 2 2 mg nicotinamide 10 mg/Cap Bot 60s, 500s
Use: Pernicious anemia
- MOL-IRON PRENATAL** (White) Vit A 6000 u, D 600 u, K 2 mg, E 11 U, C 100 mg 8 1 5 mg, 8 1 2 2 mcg folic acid 0.25 mg, Vit B 2 5 mg, B 6 2 mg, panthenol 5 mg nicotinamide 30 mg, ferrous sulf 75 mg molybdenum oxide 11 mg plus 8 minerals and trace elements/Tab Bot 30s, 90s
Use Prenatal vitamin supplement
- MOL-IRON TABLETS, LIQUID** (White) Ferrous sulfate 195 mg (3 gr) molybdenum oxide 3 mg (1/20 gr)/Tab or dr Tab Bot 100s, 1000s, 5000s Liq Bot 12 fl oz 1 gal
Use Iron deficiency anemias
- W/Calcium & Vit D (White) Mol iron ferrous sulfate 195 mg, molybdenum oxide 3 mg, dicalcium phosphate 869 mg, Cal 250 mg, Vit D 200 u/Cap Cap, Bot 100s, 1000s
Use: Nutritional supplement
- W/Liver & Vitamins (White) Mol iron ferrous sulfate 195 mg, molybdenum oxide 3 mg, desic whole liver 450 mg (equiv to 1 8 Gm whole liver), Vit B 12 1 mcg, thiamine HCl 1 mg, riboflavin 1 mg, nicotinamide 5 mg, pyridoxine HCl 0.5 mg, Cal pantothenate 1 mg/Cap Cap, Bot 100s 1000s
Use: Iron deficiency anemias
- W/Vitamin C (White) Ferrous sulf 195 mg, molybdenum oxide 3 mg, ascorbic acid 75 mg/Tab Bot 100s
- *MOLOFAC**, (Squibb) Diethyl sod sulfosacinate 60 mg/Cap Bot 30s, 100s
Use Fecal softener
- MONACETYL PYROGALLOL**, (Eugallo), Pyrogallol Monoacetate)
Use: Keratolytic in chronic dermatitis
- *MONALIUM NYDRATE**, Hydrated magnesium aluminum Sca: Riopan, Tab & Susp (Ayerst Lab)
- MONASE** (Upjohn) Withdrawn Etryptamine acetate [3-(2 aminobutyl)indole] Acetate 15 mg/Tab Bot 100s
Use Management of depressive disorders
- MONA-SERTS**, (Wintersmith) Aluminum caprylate, 3 gr, phenylmercuric acetate 9.2 mg, urea, 1.0 gr, in combination w/ citric acid, boric acid lactose & dextrose/Tab Box 24s, Bot 100s
Use Infections at vaginal tract
- MONETAMINE**, (Northwest Labs) dl Amphetamine sulfate Tab, Vial 10 cc
Use: Central nervous system stimulant
- MONICHO**, (Ives Cameron) Inositol 250 mg, choline dihydrogen citrate 500 mg, polysorbate 80 500 mg/5 cc Bot 12 oz
Use: Lipotropic
- MONITAN**, (Ives Cameron) Sorethyton (20) mono oleate (Polysorbate 80) 1.5 Gm/5 cc Bot 12 oz
Use (1-2 tsp daily) Fat emulsifier
- MONO-AMINE OXIDASE INHIBITORS**
- Cotron Tab (Lekside)
Marplan Tab (Roche Lab)
Marsid Tab (Roche Lab)
Nardil, Tab (Warner Chilcott)
Niamid Tab (Pfizer)
- *MONOBENZONE**, NF NND 1962 Soln, Oint NF See Benquin Oint (Elder)
- MONOBENZYL ETHER OF HYDROQUINONE**, See Benquin Oint (Elder)
- MONOBROMISOVALERYLUREA**, NNR 1940 See Bromisovalum (Various Mfr)
- MONOBROMOSALICYL ALCOHOL** Bismosaligenin See Bromisalizal, Tab & Amp (Hynson, Wescott & Dunning)
- MONOCAINE FORMATE**, NND 1962 (Novocain) Butethamine formate Cryst Amp 50 100, 150, 200, 300, 500 mg Sol 5% Amp 2 cc
Use Spinal anesthetic
- MONOCAINE HYDROCHLORIDE**, NND 1962 ADR (Novocain) Butethamine hydrochloride Sol 1.5% w/epinephrine 1:100000 Amp 2 cc, 3 cc, 5 cc Anestube 1 cc, 2 cc, 2.5 cc, 5 cc Novomula 2.5 cc 5 cc Bot 60 cc, 120 cc
Use: Local anesthetic
- MONO-CALAFORMULA** (Eric Kirk & Gary) Calcium carbonate 500 mg, ferrous fumarate

*Generic name recognized by NF, USP or AMA

- rate 90 mg., Vit. A 5000 u., Vit. D 400 u., Vit. B-1 1 mg., Vit. B-6 1 mg., Vit. B-12 3 mcg., Vit. C 50 mg., niacinamide 10 mg., Vit. B 2 1 mg., Iodine 0.1 mg., manganese 1.5 mg., cobalt 0.1 mg., copper 1 mg., zinc 1.4 mg., magnesium 1 mg., potassium 5 mg./Tab.
- Use: Vitamin mineral supplement.
- MONOCIDENE.** (Klee) Copper sulfate 4 1/2 % / Pwd. Bot. 8 oz.
- Use: Vaginal douche for Trichomonas infections.
- MONOCILLIN.** (Schenley) Procaine penicillin G in oil 300M u./cc. with 2% alaminem monostearate. Vial, 300M u., 1 doses; 3000M u., 10 doses. Cortridge, 300M u. w/disposable syringe.
- Use: I.M.; prolonged penicillin therapy.
- MONODRAL BROMIDE, N N O. 1962.** (Winthrop Labs.) (Penthenole bromide) 2-Diethyl-hydroxyethyl-2-cyclopentyl-2-(2-thienyl) hydroxyacetatemethobromide. Cap. 5 mg. Bot. 100s. Elxlr. Penthenole bromide 0.5 mg./cc. Bot. 1 pt.
- Use: (1 Ld) Anticholinergic.
- MONODRAL W/MEBARAL.** (Winthrop Labs.) Penthenole bromide 5 mg., mephobarbital 32 mg./Tab. Bot. 100s & 1000s.
- Use: 1-2 t.i.d.; peptic ulcer therapy.
- MONOETHANOLAMINE NICOTINATE.** See: Nicotin, Amp. (Abbott)
- MONOETHANOLAMINE OLEATE.** See: Manolate, Amp. (Abbott).
- MONOIODOMETHANESULFONATE SODIUM.** See: Methiodal Sodium, U.S.P. (Winthrop Labs.)
- MONO-KAY.** (Abbott) *Phylionadione. Vit. K-1 5 mg./Tab.; 1 mg., 10 mg., 25 mg., Bot. 50s. 50 mg. In a water emulsion w/ polyorbate "EO" / Vial, Amps, 1 mg., 10 mg., 25 mg., 50 mg. Emulsion 1 mg./0.5 ml. amp. 5s, 25s.
- Use: Control of hemorrhage.
- MONOLATE.** (Abbott) Monoethanolamine oleate 5%, benzyl alcohol 2%, w/water for inj. Amp. 2 1/2 cc., Box 12s.
- Use: Anadyne & sclerosing sol.
- MONOPHEN.** (National Synthetics) 2-(4-Hydroxy-3,5-dihydroxybenzyl)-cyclohexane carboxylic acid. Cap. 0.5 Gm. Box 50s, 100s, 250s, 500s, 1000s.
- Use: Orally; cholecystography.
- MONOPHOS.** (G. F. Harvey) Amphetamine phosphate manobasic. Tab. 10 mg. Bot. 100s, 1000s
- Use: Orally; central nervous system stimulant.
- MONOSOD.** (Martin-Phillips Labs) Caps 100s, Tabs. 100s.
- MONOSODIUM PHOSPHATE.** See: Sodium Biphosphate.
- MONOSTEARIN.** (Various Mfr.). Glyceryl monostearate.
- MONOTHEAMIN.** (Lilly) (Theamin, Theophylline monoethanolamine) Enseal 3 gr. Bot. 100s. Pulvule 1 1/2 gr. Bot. 100s, 500s; 3 gr., Bot. 100s, 500s; Supp. 0.5 gr. Box 12s.
- Use: Heart stimulant, vasodilator, diuretic & antispasmodic.
- MONOTHEAMIN & AMYAL "A."** (Lilly) Theamin 3 gr., amytal 3/4 gr./Pulvule. Pulvule, Bot. 100s, 500s. Enseals: mono, 3 gr.; amytal 1/4 gr. Bot. 100s, 500s; Pulv. Bi mono 1 1/2 gr.; amytal 1/2 gr. Bot. 100s, 500s.
- Use: Heart stimulant w/sedative.
- MONOTHEAMIN & SECONAL SODIUM.** (Lilly) Theamin 0.5 Gm., Seconal Sod. 0.0325 Gm./Supp. Supp., Pkg. 12s.
- Use: Sedative & theophylline therapy.
- MONO-VACC.** (Lincoln) Device used for vaccination against smallpox. Box of 12 sterile units.
- Use: Scarifier.
- MONOXYCHLOROSENE.** A stabilized, buffered, organic hypochlorous acid derivative. See: Oxychlorosene
- MORANLY.** See: Suramin Sod. (Naphuride Sod., Winthrop Labs.)
- ***MORAZONE.** 2,3-Dimethyl-4-(3-methyl-2-phenylmorpholinomethyl)-1-phenylpyrazol-5-one.
- MOR-CAL.** (Schen Lab) Dried skim milk solids, processed fat, whole wheat flour w/Vit. B-1 50 mg., B-12 50 mcg./lb. Can 1 lb.
- Use: Dietary supplement.
- MORISTIN.** (Durst) Estrone sulfate Sod. Tab. 0.625 mg., 1.25 mg. Bot. 100s, 500s, 1000s
- Use: Orally (m pl), estrogen therapy.
- MORNIDINE.** N.N.D. 1962. (Searle) Pipamazine, Tabs. (5 mg.) Bats. 100s & 500s. Amp (5 mg/cc) 1 cc. Boxes 6 & 25s
- Use: Control nausea and vomiting
- ***MOROXYDINE.** (Smith, Kline & French) 4-morpholinocarbonylmethyl guanidine.
- Use: Antiviral agent.
- MORPHINE ACETATE.** (Mallinckrodt) Bot. 1/2 oz., 1 oz.
- W/Terpin hydrate, ammonium hypophosphite & Pot guaifacil sulfonate. See: Branch-Tuwin No. 1, Sol. (First Texas).
- W/Antimony, Pot. tartrate, ammonium chloride, coccilana tr. See: Morphinal Lig (C. D. Smith)
- MORPHINE AROMATIC.** (Meick Sharp & Dohme) Morphine HCl 1 gr., alcohol 12%/30 cc. Bot. 1 pt.

* Generic name recognized by N.F., U.S.P. or A.M.A.

MORPHINE HYDROCHLORIDE

476

- MORPHINE HYDROCHLORIDE, USP XV**
[Various Mfr] Pow, Bot 1 oz, 5 oz
Use Analgesic
- W/Chloral hyd See Papine (Bottle)
- W/Hydrocyanic acid diluted chloroform
capsicum lr
See: Chlor Anodyne (Parke Davis)
Chlorodyne (Merck Sharp & Dohme)
- W/Scopolamine comp (Merck Sharp &
Dohme) Morphine HCl T/10 gr, ethyl
morphine HCl 1/5 gr, scopolamine HBr
1/800 gr/HT Each Tab will make 5 min
Schlesinger's sol Bot 100s
- ***MORPHINE INJECTION, USP** Sterile sol
of suitable salt
- ***MORPHINE SULFATE, USP, ADR, Tab,**
USP, ADR 1955 (Various Mfr) Flake or
Pow Bot 1/2 oz, 1 oz, 5 oz, HT 1/2 gr,
1/6 gr 1/4 gr 1/2 gr, 1 gr
Use Narcotic Analgesic & sedative
- W/Atropine Sulfate, NF (Merck Sharp &
Dohme) Morphine sulfate 1/4 gr, atropine
sulfate 1/150 gr/tab Bot 100s
- W/Chloroform ammonium chloride, menthol
sanguinaria grindelia, senega yerba santa
See Melsam w/Morphine (Suliff & Case)
- W/Chloroform Sod citrate citric acid &
Pol guaiacal sulfonate See Sodacitrol
Liq (Flint, Eaton)
- W/Ephedrine HCl, labelia inflata Sod cit
rate, Pol guaiacal sulfonate, menthol &
aromatics See: Phedrolal, Liq (Mallard)
- W/Papaverine HCl See: Asymatrine Amp
(Breon)
- W/Pol guaiacal sulfonate tartar emetic
ammonium chloride citric acid Sod citrate
& with or without sugar Sees Cetrolen
Liq (Warren Teed)
- MORPHINE LIQUID (C D Smith)** Mor
phine acetate 1/4 gr antimony Pot tartrate
1/12 gr, ammonium chloride 8 gr, cocai
lona lr 40 min/r oz Bot 4 oz, 1 pt,
1 gal
Use: (1 tsp q 4 h) Narcotic cough prep
arolan
- MORRAMIN (Wyeth)** Hypophosphites of
strychnine, Iron, Mang, Cal, Sod & Pot
w/yeast, malt ext, wild cherry & alcohol
Bot 12 fl oz, 1 gal
Use: Tonic
- MORRHU-COMP. (Hort)** Cod liver oil 60%
chlorobutanol 1% benzocaine 1% in a
perfumed emulsion base Tubes 1 oz Jars
1 lb
Use: Burns wounds, ulcers, abrasions
- MORUGINT (Massengill)** Cod liver oil
Oint Jar 2 oz, 1 lb Ophth aint Tube
1/2 oz Supp 12s
Use: Topically; vitamin therapy.
- *Generic name recognized by NF USP or AMA
- MORUMIDE, (Massengill)** Sulfanilamide 10%
w/cod liver oil in sterile aint base Oint
Tube 2 oz; Jar 1 lb
Use: Topically, sulfanilamide susceptible
organisms
- MORUMIDE & PHENACAINE, (Massengill)**
Sulfanilamide 5%, phenacaine HCl 1%
w/cod liver oil conc Ophth aint Tube
1/2 oz
Use: Topically, sulfanilamide susceptible
organisms
- MORUSAH, (Massengill)** Benzocaine 2%
cod liver oil conc Tube 2 oz; Jar 1 lb-
5 lb
Use Granulation of diseased tissue
- MORUSUL, (Endo)** Sod marrhuate 5%, Sod
sulfadiazine 10% Vial multiple dose, 25
cc
Use: (0.5 cc to 5 cc) Sclerosing agent
- M-SUL (Misemer Pharm)** Tabs 100s Susp,
pt
- MOTILYH, NND 1962 (Abball)** Dexpon
thenol d Pantothonyl alcohol 250 ml/ml 2
ml ampoules in pack of 10 and 25 10 ml
multiple dose vials in pack of 1 and 6
500 mg/ml Amp 2 ml 10s
Use Restore intestinal function in ileus
enteric atony or delayed motility
- MOTION SICKNESS,**
See Benacine, Tab (Parke, Davis)
Bonadaxin, Tab (Pfizer)
Bonamine, Tab (Pfizer)
Dramamine, Amp, Elixir & Tab (Searle)
Marezine Tab, Amp & Supp (Burroughs
Wellcome)
Nausea Therapy
Nia Piracene, Tab (Nian)
Frametic, Tab (Harrower)
Scopodex Pel (De Bruielle)
Scopolamine HBr (Various Mfr)
Vasano (Schenig)
- MPP (Abball)** 1 (2 methoxyphenyl)piper
azine monohydrochloride
Use Antihypertensive
- MSC TRIAMINIC, (Darsey)** Phenylpropanol
amine HCl 50 mg pheniramine maleate
25 mg, pyrilamine maleate 25 mg, meth
scopolamine nitrate 4 mg/Timed releas
Tab Bot 50s
Use: Decongestant
- M-SED LIO (Misemer)** Pentobarbital sod um
30 mg, phenobarbital sodium 10 mg,
NaBr 300 mg, KBr 200 mg, CoBr 100 mg/
3cc Bot pts gals
Use Hypnotic Sedative
- M S M (Creomulsion Co)** Mint flavored syrup
of magnesium hydroxide 67 gr/5 cc Bot
8 fl oz
Use: Antacid, laxative

- MUCAMIDE.** (Warren-Teed) Phthalylsulfacetamide 0.5 Gm., pectin 0.1 Gm., albumen tonate 30 mg./Cap. Bot. 100s, 500s, 1000s.
Use: 2 cap. every hr. for 3 hr. then 2 every 2 hr.; diarrhea & intestinal infections.
- MUCARA.** (Ives-Cameron) Karaya gum. Plain, Bot. 6½ oz., 2 lb.; w/Cascara, Bot. 6½ oz., 2 lb.
Use: Bulk laxative.
- MUCIENUM.** (Thurston) Gastric mucin 150 mg., duodenal substance 250 mg., ascorbic acid 25 mg., bioflavonoids 30 mg./Cap. Bot. 50s 100s
Use: Treatment of peptic ulcer.
- MUCIGEL.** (Warner-Chilcote) Ephedrine 0.1%, benzocaine 5%, mucin 10%. Jelly, Tube 1 oz., w/applicator.
Use: Topically; local anesthetic.
- MUCILOID OF PSYLLIUM SEED.** See: Metamucil, Pow. (Searle)
- MUCILOSE.** (Winthrop Labs.) Bland, non-irritating hemicellulose obtained from psyllium seed (*Plantago lanifolia*). Flake, Conc., Bot. 4 oz., 1 lb.; Special, Bot. 4 oz., 1 lb.; Gran., Special, Bot. 4 oz., 1 lb. Tab., Comp., Sol. 100s, 1000s.
Use: Chronic constipation.
W/Cascara, Gran., Bot. 4 oz.
- MUCILOSE-SUPER.** (Winthrop) Mucilose, Tylaxopal, 50 mg (Superinone, a detergent), and dextrose/5 Gm. Cons., 4 oz., 16 oz.
Use: Demulcent and bulk laxative.
- MUCIN.** N.N.D. 1959. See: Gastric Mucin. (Various Mfr)
- MUCIN, VEGETABLE W/Yeast or alkalinized.** See: Plantamucin, Pow. (Elder).
- MUCOGEL.** (Barrows Chemical Co.) Gastric mucin 160 mg., mag. glycinate 250 mg., alum. hydroxide gel 250 mg./tab. Bot. 100s.
Use: Antacid.
- MUCOGEL-B.** (Barrows Chemical Co.) Gastric mucin 160 mg., magnesium glycinate 250 mg., aluminum hydroxide gel 250 mg., phenobarbital 6 mg., belladonna ext. 5 mg./tab. Bot. 100s.
Use: (1-2 q. 2 h) Treatment of hyperacidity & peptic ulcer.
- MUCOLYTIC AEROSOL SOLN.**
See: Alevole (Winthrop Labs.)
Tergemist (Abbott)
- MUCOPLEX.** (Stuart) Desiccated liver 750 mg., Vit. B-12 5 mcg., Vit. B-2 1.5 mg./Tab. Bot. 100s.
Use: Nutritional.
- MUCOPOLYSACCHARIDES.**
W/Heparin Sod. See: Heparinate, Sol. (Cy Labs.)
W/Hesperidin methyl chalcone, citrus bio-
- flavonoids complex & Vit. C. See: Hesperinate-C, Cap. (Cy Labs)
- 'MUCORETTES,' METHYLTESTOSTERONE.** (White) Methyltestosterone 10 mg./Disk. Bot. 30s, 100s, 500s
Use: Oromucosal absorption, androgenic therapy
- MUCO-SETH.** (Harrower) Aluminum hydroxide gel dried 4 gr., gastric mucin 2½ gr., hamotropine methylbromide 1/85 gr., Mogn. trisilicate 7 gr./Tab. Bot. 50s.
Use: Orally (1-2 q. 3 h), antacid w/anti-spasmodic.
- MUCOTIN.** N.N.R. 1955. (Warner-Chilcote) Each Tab. contains gastric mucin 0.16 Gm., aluminum hydroxide gel 0.25 Gm., Mogn. trisilicate 0.45 Gm. Tab. 50s, 100s.
Use: Peptic ulcer & gastric hyperacidity.
- MUCUSOL.** (Kramers-Urbon) Foreign protein 0.04 Gm., Sod. iodide 0.10 Gm., chlorbutanol 0.01 Gm./2 cc. Amp. 2 cc., Box 25s, Vial 30 cc.
Use: I.M., S.C. (2 cc. daily), encourages normal mucous secretion.
- MUDRANE.** (Poythress) Aminophylline 2 gr., phenobarbital ½ gr., ephedrine HCl ¼ gr., Pot. iodide 3 gr./Tab. Bot. 36s, 100s.
Use: (1 t.i.d.) Treatment of asthma.
- MUGASTRAL.** (G. F. Harvey) Gastric mucin 200 mg., dihydroxy aluminum aminoacetate 300 mg., ethaverine HCl 6 mg./Tab. Bot. 100s.
Use: (1-3 q.i.d.) Antacid.
- MULLSOY.** (Sorden) Water 75%, soy protein 6%, soy fat 7.5%, carbohydrate (available) 5.9%, crude fiber 0.3%, minerals (total) 2%, Cal. 0.26%, Phos. 0.22%, Iron 0.0004%/fl. oz. Tin 15½ fl. oz.
Use: Hypo-allergic infant feedings.
- MULSAVITE CAPSULES.** (Merck Sharp & Dohme) Vit. A palmitate 20,000 u., D 2000 u., thiamine HCl 5 mg., riboflavin 10 mg., pyridoxine HCl 1 mg., vit. B-12 cryst. 5 mcg., cal. pantothenate 10 mg., niacinamide 50 mg., ascorbic acid 150 mg., mixed tocopherols 10 mg./cap. Bot. 30s, 100s, 500s.
Use: (1 or more daily) Therapeutic potency multivitamins.
- MULSAVITE-P CAPSULES.** (Merck Sharp & Dohme) Vit. A palmitate 20,000 u., D 2000 u., thiamine HCl 5 mg., riboflavin 10 mg., pyridoxine HCl 1 mg., Vit. B-12 cryst. 5 mcg., Cal. pantothenate 10 mg., niacinamide 50 mg., ascorbic acid 150 mg., mixed tocopherols 10 mg., folic acid 5 mg./Cap. Bot. 30s, 500s, 1000s.
Use: Therapeutic potency multivitamins.
- MUL-SED.** (Pitman-Moore) Mix of bismuth 77 min., paregoric 170 min., kaolin 93 gr.

- pectin 4 gr/A oz Bot 4 oz, 1 pt, 1 gal.
Use: Orally (2 dr prn), antacid
- MULSOPAQUE** (Lafayette) Ethyl iodophenyl-undecylate Amp 50%, 10 cc, Box 4s
Use: Parenterally (10 20 cc), cholangiography
- MULTAZINE**, NNR 1955 (Kremers Urban)
Meth dia mersulfanamide, D167 Gm of each per 0.5 Gm/Tab Bot. 100s, 1000s
- MULTEX**. (Sutiff Case) Vit A 5000 u, D 1000 u, C 75 mg, thiamine HCl 3 mg, riboflavin 3 mg, alpha tocopherol 5 IU, nicotinamide 20 mg, Cal pantothenate 5 mg, pyridoxine HCl 0.25 mg, Vit. B 12 1 mcg/Cap Bot 100s, 500s, 1000s
Use: Dietary supplement
- MULTEX LIQUID**. (Sutiff & Case) Vit A Palmitate 5000 U Vit O 1000 U, Thiamine Mononitrate 15 mg, Riboflavin 15 mg Nicotinamide 5 mg, Vit C 50 mg, Vit B12 2 mcgm/5cc Liq Bot 4 oz 8 oz
- MULTI-BETA**. (White) Cap. Vit B 1 1 mg B 2 2 mg, B 6 0.5 mg, Cal pantothenate 0.5 mg, nicotinamide 10 mg, brewers yeast 155 mg/Cap Bot 30s, 100s, 500s Liq Vit B 1 2.5 mg, B 2 0.5 mg, B 6 0.15 mg, Cal pantothenate 0.2 mg, nicotinic acid 10 mg/cc Bot 10 cc, 25 cc, 50 cc, B fl oz
Use: Vit B complex
- MULTI-BETA-B-12 DROPS**. (White) Vit B 12 1 mcg, thiamine HCl 2.5 mg, riboflavin 0.4 mg pyridoxine HCl 0.15 mg Cal pantothenate 0.2 mg, nicotinamide 10 mg/cc Bot 15 cc, 50 cc w/dropper
Use: Deficiencies of Vit B
- MULTIBIOTIC GAUZE PADS** (Bio Rama) Absorbent gauze containing not less than 1 mg Streptomycin Cal chloride complex, 50 u Bacitracin, 1000 u polymyxin B per sq Inch Box 2"x2", 3"x3", 4"x4" gauze pads 100s
Use: Prevention of Infection
- MULTI-B-K**. (Testagar) Thiamine HCl 100 mg, riboflavin 3 mg, niacinamide 200 mg, pyridoxine 50 mg, pantothenic acid 5 mg, benzyl alcohol 2%/cc Vial 10 cc, 30 cc.
Use: Multiple Vit B deficiencies
- MULTI-B-PLEX**. (Testagar) Thiamine HCl 100 mg, riboflavin 2 mg, nicotinamide 100 mg, pantothenic acid 10 mg, pyridoxine HCl 10 mg/cc, Amp 1 cc Vial 10 cc
Use: Multiple Vit B deficiencies
- MULTICAPS THERAPEUTIC**. (Kirk) Vit A 25,000 u, D 1000 u, B 1 10 mg, B 2 5 mg, C 5 mg, niacinamide 150 mg/Cap Bot 100s
Use: Treatment of vitamin deficiencies
- MULTICEBRIN**. (Lilly) Thiamine chloride 3 mg, riboflavin 3 mg, pyridoxine HCl 15 mg (as Cal pantothenate 5 mg), nicotinamide 25 mg, Vit B-12 (activity equiv) 3 mcg, folic acid 0.1 mg, ascorbic acid 75 mg, tocopherols, natural type 10 mg, vit A 10,000 u, D 1000 u/cap Pkg 100s, 1000s
Use: Multiple vitamin deficiencies.
- MULTICEBRIN JR**. (Lilly) Vit B 1 1.5 mg, B 2 2 mg, B 6 1 mg, Cal pantothenate 25 mg, Vit B-12 3 mcg, nicotinamide 12 mg, folic acid 0.1 mg, Vit C 75 mg, E 5 mg A 5000 u, D 500 u/Gelseal Bot 60s 1000s
Use: Multivitamin therapy for children
- MULTICIBUS LIQUID** (Canfield) Iron pep tonized 420 mg, manganese citrate 158 mg Vits B 1 10 mg, B 2 10 mg, cyanocobalamin 20 mcg nicotinamide 50 mg B 6 1 mg pantothenic acid 5 mg, liver fraction No 1 1 Gm rice bran ext 1 Gm choline 60 mg, inositol 30 mg/A oz Bot 8 oz 1 pt, 1 gal
Use: Anemia
- MULTIFUGE CITRATE**, NND 1962 (Blue Line) Piperazine citrate Syr 100 mg/cc Bot 4 fl oz 1 pt Tab (500 mg) Bot 100s 1000s
Use: Vermifugal agent
- MULTIHIST**. (Smith Dorsey) Pyrilamine maleate 10 mg, prophenpyridamine maleate 10 mg, phenyltolaxamine dihydrogen citrate 10 mg/Cap Bot 100s 500s 1000s Also avail w/APC
Use: (1 1 d) Antihistaminic
- MULTIHIST SYRUP**. (Smith Dorsey) Pyrilamine maleate 5 mg, prophenpyridamine maleate 5 mg, phenyltolaxamine dihydrogen citrate 5 mg/5 cc Bot 1 pt, 1 gal
Use: Triple antihistaminic therapy
- MULTIHIST-APC** (Smith Dorsey) Pyrilamine 5 mg, pheniramine 5 mg, phenyltolaxamine 5 mg, acetophenetidin 210 mg, aspirin 150 mg caffeine 30 mg/Cap Bot 100s 500s
Use: Treatment of common cold
- MULTINIC TABS** (Cambridge) Ferrous sulf 195 mg B 1 10 mg, B 6 2 mg, B 12 15 mcg's intrinsic factor O1 USP U/Tab Bot 100s
Use: Hematinic
- MULTIPLEX**. [Prema] Vit A 5000 u, B-1 4.5 mg, B 2 2 mg, nicotinamide 20 mg, Cal pantothenate 10 mg, Vit C 30 mg, D 500 u, Cal, Phos, Iron, Mag, Mang & zinc/Tab Vial 30s Bot 50s, 100s, 250s, 1000s
Use: Dietary supplement
- MULTI-VI** (White) Vit A 5000 u, D 500 u-

- C 37.5 mg., B-1 1.5 mg., B-2 2 mg., nicotinamide 10 mg./Cap. Bot. 25s, 100s.
Use: Multivitamin therapy.
- MULTI-VI-DROPS.** (White) Vit. A 5000 u., D 1000 u., thiamine HCl 1 mg., riboflavin 0.4 mg., nicotinamide 10 mg., pyridoxine HCl 1 mg., panthenol 2 mg., ascorbic acid 50 mg./0.6 cc. Bot. 15 cc., 30 cc., 50 cc. w/droppers.
Use: Diet supplement.
- MULTI-VIMS DBOPS.** (Poul Maney) Vit. A 5000 u., D 1000 u., thiamine chloride 1 mg., riboflavin 0.5 mg., niacinamide 5 mg., pantothenic acid 2 mg., pyridoxine HCl 0.1 mg., ascorbic acid 30 mg., cryst. Vit. B-12 1 mcg./0.6 cc. Bot. 1/2 oz.
Use: Diet supplement.
- MULTIVITALIN.** (Lincoln) Vit. A 10,000 u., D 2000 u., B-1 50 mg., B-2 5 mg., B-6 10 mg., niacinamide 100 mg., panthenol 5 mg., C 500 mg., B-12 50 mcg./10 cc. Vial 10 cc., Pkg. 6s/diluent. Tabs. Bot. 100s, 1000s.
Use: Multiple vitamin therapy.
- MULTI-VITA-MIN-ERALS.** (Raymer) Vit. A 1666 u., D 166 u., thiamine HCl 0.33 mg., riboflavin 0.66 mg., ascorbic acid 10 mg., pyridoxine HCl 0.66 mg., Cal. pantothenate 0.16 mg., niacin 3.33 mg., Cal., Phos., Mag., Iron, Mang., Iodine/Tab. Bot. 100s, 500s, 1000s.
Use: Dietary supplement.
- MULTIVITAMIN INFUSION.** (U.S. Vitamin) Not listed, Vit. C 500 mg., A 10,000 u., D 1000 u., E 5 mg., B-1 50 mg., B-2 10 mg., niacinamide 100 mg., B-6 15 mg., panthenol 25 mg./10 cc. Amp. 10 cc., Box 1s, 5s, 25s, 100s. Also oval, w/Vit. C.
Use: Multivitamin therapy.
- MULTIVITAMIN INTRAVENOUS.** (Lincoln) Vit. A palmitate 10,000 u., D-2 2000 u., thiamine HCl 50 mg., riboflavin 5 mg., pyridoxine HCl 10 mg., niacinamide 100 mg., panthenol 5 mg., ascorbic acid 500 mg., Vit. B-12 15 mcg./10 cc. Amp. 10 cc., Box 25s.
Use: Therapeutic vitamin supplement.
- MULTIVITAMIN LYOPHILIZED.** (Lincoln) Vit. A 10,000 u., D 2000 u., B-1 50 mg., B-2 5 mg., B-6 10 mg., niacinamide 100 mg., panthenol 5 mg., Vit. C 500 mg., B-12 50 mcg./10 cc. Amp. 10 cc.
Use: Multivitamin therapy.
- MULTIVITAMINS.** (C. D. Smith) Vit. A 5000 u., B-1 2 mg., B-2 2 mg., C 50 mg., D 500 u., niacinamide 20 mg., panthenol 1 mg., folic acid 0.25 mg., Vit. B-12 1 mcg./Tab. Bot. 100s, 1000s.
Use: Vitamin supplement
- MULTI-VITAMINS.** (National Drug) Vit. A 5000 u., D 1000 u., B-1 3 mg., B-2 2 mg., C 37.5 mg., nicotinamide 20 mg., cal. pantothenate 5 mg., Vit. B-6 0.5 mg./Cap Bot 100s, 500s
Use: Multi-vitamin therapy
- MULZIZINE.** (Kremers-Urban) See: Trisulfapyrimidines Tabs., U.S.P.
- MULTOVOALS.** (Vitamin Corp) Vit. A 5000 u., D 1000 u., B-1 3 mg., B-2 3 mg., B-6 0.25 mg., C 50 mg., Cal. pantothenate 2 mg., niacinamide 10 mg., Vit. E (wheat germ oil) 5 mg., folic acid 0.1 mg., Vit. B-12 1 mcg./Cap. Bot. 50s, 100s, 250s, 1000s.
Use: Dietary supplement.
- MULVIDREN.** See: Stuart Mulvidren, Tab. (Stuart)
- MULVIDREN DBOPS.** (Stuart) Vit. A 3000 U.S.P. units, D 400 U.S.P. units, C 60 mg., B-1 1mg, B-2 1.2 mg, B 6 0.5 mg, B-12 1 mcg, niacinamide 10 mg/0.6 cc. Bot 30 cc.
Use: Supplement to infant feeding and diet of children
- MULVITAB.** (Reid) Vit. A 8000 u., D 800 u., B-1 3 mg., B-2 3 mg., B 6 1 mg., calcium pantothenate 5 mg., nicotinamide 20 mg., B-12 2 mcg., folic acid 0.25 mg., ascorbic acid 75 mg., E 2 mg./tab.
- MULVITOL.** (Consolidated Mid) Vitamins A 5000 u., D 500 u., B-1 2.5 mg., B-2 2.5 mg., B-6 0.5 mg., B-12 1 mcg., C 37.5 mg., cal. pantothenate 5 mg., folic acid 0.25 mg., niacinamide 20 mg./cap. Bot. 30s, 100s, 250s, 1000s, 5000s.
Use: One daily. Avitaminosis.
- MULVITOL STRONG.** (Consolidated Mid) Vitamins A 12,500 u., D 1000 u., B-1 5 mg., B-2 5 mg., B-6 2 mg., B-12 2 mcg., C 100 mg., cal. pantothenate 10 mg., folic acid 0.5 mg., niacinamide 30 mg./cap. Bot. 30c, 100s, 250s, 1000s, 5000s.
Use: One daily. Vitamin therapy.
- MUMPS IMMUNE GLOBULIN-HUMAN.** (Hyland) Concentrated mumps gamma globulin. Vial 1.5 cc., 4.5 cc.
Use: Prevention and treatment of mumps.
- MUMPS IMMUNE SERUM.** (Hyland) Homologous serum prepared from hyperimmunized adults. Vial 20 cc. w/diluent.
Use: I.M. & I.V., prevent mumps.
- *MUMPS SKIN TEST ANTIGEN, NND 1962.** (Lederle) Antigen made from allantoic fluid of chick embryos. Amp. 1 cc. (10 tests)
Use: Diagnostic aid.
Prepared from allantoic fluid of chick embryo
- *MUMPS VACCINE, NND 1962.** (Lederle)

bryos infected w/known strain of mumps virus 1 immunization 2 cc, 5 immunization 10 cc

Use SC & IM (2 inj of 1 cc each) immunization against mumps

MURACIL, NND 1959 (Organon) (Methyl thioauracil) 6 methyl 2 thioauracil Tab 50 mg Bot 100s, 1000s

Use (1 2 t i d) Antithyroid

MUREL, NND 1962 (Ayerst) Valethamate bromide 10 mg (phenylmethylvaleric acid β d ethyl aminoethyl ester bromomethylate)/Tab Bot 100s 1000s Amp (10 mg/cc) 5 cc

Use: Spasmolytic agent

MUREL W/PHENOBARBITAL, (Ayerst) Val ethamate bromide 10 mg, Phenobarbital 1/4 gr/Tab Bot 100s 1000s

Use Spasmolytic agent and sedative

MUREL-5 A, (Ayerst) Valethamate bromide 40 mg with or without 1/2 gr phenobarbital Sod/Tab Bot 100s 1000s

Use Gastrointestinal, Genitourinary and biliary spasm

MURIAMIC (Pitman Moore) Not listed Glutamic acid HCl 5 gr Bot 100s, 500s, 1000s
Use: (1 2 t i d) Achylarhydria & hypochlorhydria

MURIC ACID, See Hydrochloric Acid

MURIPSIN (Norgine) Glutamic acid HCl, 500 mg, pepsin 35 mg/Tab Bot 100s 500s
Use: Treatment of hydrochloric acid deficiencies

MURO-OTIC, (Muro Pharmacal) Polymyxin B Sulfate 10 000 u/cc w/Neomycin Sulfate 5 mg/cc, 75 cc

MUSCLE ADENYLIC ACID (Various Mfrs)
Active form of adenosine 5 manophosphate
See: Adenosine 5 manophosphate, Preps (Various Mfr)

MUSCLE RELAXANTS

See: Curare (Various Mfrs)

Dimethylane, Tab (National Drug)

Flaxedil, Vial (Lederle)

Flexin Tab (McNeil)

Inlacastin, Vial (Squibb)

Mecastin, Vial (Squibb)

Mephenesin (Various Mfr)

Meprobamate (Various Mfrs)

Metubina, Amp (Lilly)

Mytalan, Vial (Winthrop Labs)

Narflex, Tab (Riker)

Prenderal, Tab (Squibb)

Robaxin (Robins)

Smaxor (Armour)

Soma Tab (Wallace)

Rela (Schering)

Striatran (Merck Sharp & Dahme)

Succinylcholine Chloride (Various Mfr)

d Tubacurarine Chloride (Various Mfr)

MUSLAX, (Ciba) Mephenesin 300 mg, d amphetamine phosphate 3 mg, butobarbital sod 10 mg/Tab Bot 100s, 500s, 1000s
Use: Muscular relaxant, antidepressant

MUSTARAL OIL. See: Allyl Isothiocyanate NF

MUSTARGEN HYDROCHLORIDE, NND 1962 (Merck Sharp & Dahme) (Mechlorethamine HCl) Methyl bis(beta chloroethyl)amine HCl (Nitrogen Mustard) Mustine, Caralysine Vial 20 cc, each vial containing 10 mg in dry form triturated with 90 mg of anhydrous sodium chloride
Use: IV, certain forms of neoplastic diseases

MUSTEROLE LINIMENT, (Plough) Liq

Use Analgesic

MUSTINE See Mechlorethamine HCl, Sterile NF

MUTAMYCIN (Bristol) Streptoducin, streptomycin & dihydrostreptomycin in equal parts Vial, Pow 1 Gm, 5 Gm

Use: For gram negative & gram positive infections

M.V.I (US Vitamin) Vit A 10 000 u, D 1000 u, E 5 IU, C 500 mg, B 1 50 mg, B 2 10 mg, niacinamide 100 mg, B 6 15 mg, panthothenal 25 mg/10 cc Amp 10 cc

Use IV, infusion feeding

M.V.R.I. (Cold Vaccine) #4 Count (Cutter) Vial 5 cc, 20 cc

Use SC (0.5 cc), common cold & upper respiratory infections

MYADEC, (Parke, Davis) Each Cap contains Vit B 12 cryst 5 mcg, B 2 10 mg, B 6 2 mg, B 1 mononitrate 10 mg, niacinamide 100 mg, Vit C 150 mg, A 25 000 u, D 1000 u, E 5 IU, iodine 0.15 mg, Mang 1 mg, cobalt 0.1 mg, Pot 5 mg, molybdenum 0.2 mg, iron 15 mg, copper 1 mg, zinc 15 mg, Mag 6 mg, Cal 100 mg, Phos 80 mg Bot 100s 1000s

Use Nutritional aid

MYANESIN, See: Mephenesin (Various Mfrs)

MYANIL (Reed & Carnrick) Not listed Mephenesin 250 mg, salicylamide 125 mg/Tab Bot 50s, 100s

Use: Skeletal muscle relaxation & analgesic

MYBASAL (Lloyd & Dabney) Mephenesin 195 mg, paminobenzoic acid 195 mg, salicylamide 195 mg, vit C 10 mg, niacinamide 5 mg, B 1 2.5 mg, menadione 0.35 mg/tab Bot 100s

Also available with Dipyrone

Use: 2 4 q 2 4 hr analgesic & antispasmodic

MY-8-DEN (Ames) Sod Adenosine 5 manophosphate Sustained Action (gelatin base for prolonged action) 20 mg 100 mg/cc, 10 cc vials Aq sol, (20 mg, 100 mg/cc)

- 1 cc. Amp. Box 3s, 50s. Sublingual Tab. 20 mg. Bot. 20s, 50s, 500s. Supp. (100 mg) Box 6s.
 Use: Phosphorylation & in metabolism of carbohydrate, fat & protein, ulcer therapy, treatment of varicose veins and thrombophlebitis and bursts.
- MYCADRINE.** (Myers-Carter). Hydroxyamphetamine HBr; Para-(2-aminopropyl)phenol HBr, 20 mg./Tab. Bot. 50s, 250s.
 Use: Hypotension, coratid sinus syndrome and heart block.
- MYCASINE.** (Myers-Carter) Adenosine-5-monophosphoric acid 25 mg., Vit. B-12 60 mcg./cc. Vial 10 cc. Also in gel foam
- MYCATUSSIN.** (Myers-Carter). Noscapine, 10 mg.; phenylpropanolamine HCl, 12.5 mg.; chlorpheniramine maleate, 1.5 mg.; acetaminophen, 120 mg.; glyceryl guaiacolate, 5 mg.; ammonium chloride, 70 mg / 5 cc. Bot. Pls. Gals.
 Use: Relief in colds, coughs.
- MYC(FR)ADIN-N.** (Upjohn) Neomycin sulf. 0.5 Gm., nystatin 125,000 u./Tab. Bot. 20s, 100s.
 Use: Preparation of patients for intestinal surgery.
- MYC(FR)ADIN SULFATE, N.N.D. 1961.** (Upjohn) Neomycin sulfate. Few. Vial 0.5 Gm. J.M. Paw, Topical, Vial 3 Gm. and 10 Gm. Pediatric Sol. (125 mg/cc) Bot. 2 oz. 8 pl. Tab. (0.5 Gm.) Bot. 20s, 100s, 500s lot. Bot. plastic sqz. 1 oz.
 Use: Antibiotic.
- MYCIQUENT, N.N.D. 1962.** (Upjohn) Neomycin sulfate. Cream 5 mg./Gm. Tube ½ oz. Oint. 5 mg./Gm. Tube ½ oz., 1 oz. Jar 4 oz. Ophth. oint. 5 mg./Gm. Tube 1 dr.
- MYCIMIST.** (Upjohn) Neomycin sulfate 5 mg./cc. in lat. base. Lat. polyethylene spray bottle 89 cc.
 Use: Topical on skin, minor burns, cuts & abrasions.
- MYCINOL IMPROVED.** (Parson & Covey) Vitamins with trace elements, hesperidin, and safflower oil base/Cop. & Tabs. Bot. 50s, 100s.
 Use: Therapeutic vitamin and mineral supplement.
- MYCITRACIN.** (Upjohn) Bacitracin 500 u., neomycin sulfate 5 mg., polymyxin B est. 5000 u./Gm. Oint. Tube ½ & 1 oz. Ophth. Oint. Tube ½ oz.
 Use: Reinforced antibiotic therapy.
- MYCOFAN.** (Ooth) Salicylic acid 2½%, benzoic acid 2½%, boric acid, thymol & isopropyl alcohol. Bot. 4 oz., 1 lb., 5 lb.
 Use: Topically; bactericide & fungicide.
- MYCOLOG CREAM.** (Squibb) Triamcinolone acetate 1.0 mg., neomycin 2.5 mg., gramicidin 0.25 mg., nystatin 100,000 u./Gm. Tube 5 Gm., 15 Gm.
 Use: Infected or inflamed skin.
- MYCOLOG CREAM.** (Squibb) Triamcinolone acetate 1 mg., neomycin base 2.5 mg., gramicidin 0.25 mg., nystatin 100,000 u / Gm. Tube 5 Gm., 15 Gm.
 Use: Treatment of dermatoses, pruritis, inflammation and bacterial and/or monilial infections.
- MYCONEF OINTMENT.** (Squibb) Nystatin 100,000 u, hydrocortisone acetate 1.0 mg., neomycin sulf. 2.5 mg., gramicidin 0.25 mg./Gm. Oint. Tube 15 Gm.
 Use: Antibiotic aintment.
- MYCORHODIN.** (Bristol) Antibiotic produced by an unidentified streptomyces strain.
 Use: Antibacterial against Mycobacterium tuberculosis and gram-positive bacteria.
- MYCOSTATIN, N.N.D. 1961.** (Squibb) Nystatin, an antifungal antibiotic derived from *Streptomyces noursei*. Tab. (500,000 u) Bot 12s, 100s. Crs. 100,000 u./Gm. Tube 15 Gm. Oint. (100,000 u/Gm.) Tube ½ oz., 1 oz. Susp. (100,000 u./cc.) Bot. 24 doses w/dropper. Vag. Tab. 100,000 u., lactose 0.95 Gm./Tab. Bot. 15s, 100s.
 Use: 1) I.D.; Intestinal moniliasis. Inhibits or kills many species of fungi & yeasts.
 W/Fludrocortisone acetate, polymyxin B sulf., neomycin sulf. See: Flarohc. (Squibb)
 W/Neomycin sulf. See: Neomycin-Mycostatin, Tab. (Squibb)
- MYCOSTATIN DUSTING.** (Squibb) Nystatin 100,000 u. in a talc base/Gm. Pkg. ½ oz.
 Use: Topically, treatment of fungus infections
- MYCOZOL.** (Parke, Davis) Chlorotone 5%, salicylic acid 4%, mercury salicylate 4%, in oint. base. Tube 1 oz.; Jar 1 lb. Dermoment. Comb. of salicylic acid, benzoic acid, chlorotone & malachite green. In 50% isopropyl alcohol. Liq. Bot. 50 cc.
 Use: Fungicide.
- MYCOZYME.** (Paul Lewis Lab) See: Diastase.
- MYDONNA.** (Centrol) Mephensin 0.3 Gm., hyosciamine sulfate 0.1 mg., phenobarbital 8 mg., scopolamine HBr 0.02 mg., homotropine methylbromide 1.2 mg./Tab. Bot 100s, 500s.
 Use: (1-2 I.D.) Antispasmodic & sedative
- MYDRIACYL.** (Alcon) *Tropicamide 11-tropamide 0.5% or 1.0% Sterile aqueous solution. Bot 7.5 cc. and 15 cc.
 Use: Mydriatic-cycloplegic action.
- MYDRIATICS.** (Parasympatholytic agent.)
 See: Atropine Salts (Various Mfr.)

- Cyclogyl HCl (Schieffelin)
 Cyclopentolate HCl (Schieffelin)
 Eucatropine HCl (M. M. Thompson)
 Euphthalmine HCl (Warner Chilcott)
 Homatropine HCl (Various Mfr.)
 Sympathomimetic Types
 Amphetamine Sulfate 3% (Various Mfr.)
 Clonidine HCl 0.5% (Lilly)
 Ephedrine Sulfate 3% (Various Mfr.)
 Epinephrine HCl 1% (Various Mfr.)
 Neo Symphephrine HCl 2.5% (Winthrop Labs)
 Phenylephrine HCl (Various Mfr.)
MYDRIATIN, (Ophthalmus Inc) Eucatropine HCl 5% and phenylephrine HCl 5% buffered with methyl cellulose Bot dropper 7.5 cc
 Use Mydriatic
MYELADOL, (Upjohn) Super D cod liver oil 7.4 cc red bone marrow ext glycerinated 0.62 cc iron & ammonium citrates 4 gr, malt ext nandiolate qs/8 oz Bot 12 8 oz, 32 8 oz
 Use Hemolitic
MYLANTA (Stuart) Magnesium hydroxide 200 mg aluminum hydroxide, dried gel 200 mg methylpolysiloxane (activated) 20 mg/Tab 5 cc Box 100s Liquid Bot 12 oz
 Use Gastric distress
MYLASE 100 Alpha-amylase See: Diastase
W/ProLase 300 dehydrocholic acid, hyoscyamine HBr, hyoscyamine HBr, atropine sulfate See Almezyme, Cap (Meyer)
W/ProLase cellulose calcium carbonate magnesium glycerate See Zylase Tabs (Bryant)
MYLAXEN, (Irwin Neisser) Hexafluorenum bromide 20 mg/cc, Bot 10 cc
 Use Muscle relaxant
MYLERAN, NND 1962 (Burroughs Wellcome) Busulfan 1-4 Dimethanesulfonaxy butane Tab (2 mg) Bot 25s
 Use Chronic myeloid leukemia
MYLICON (Stuart) Methyl polysiloxane 40 mg/Tab Bot 100s
 Use Gastrointestinal distress
MYLICON DROPS (Stuart Ca) Methylpolysiloxane 40 mg/0.6 cc Drop Bot 30 cc
 Use: Infant colic
MYOCARDONE, (Chemical) Heart muscle ext conc Tab 1 1/2 gr Bot 100s, 400s
 Use Orally vasodilator
MYOCHRYSLINE, NND 1962 (Merck Sharp & Dohme) Gold sad thiamolate Amp (10 mg, 25 mg, 50 mg, 100 mg) 1 cc
 Use: (I.M.) Rheumatoid arthritis & lupus erythematosus
MYODIL See laphendylate USP
MYODONNA (Central) Phenobarbital 8 mg, mephenesin 0.3 Gm, homatropine methylbromide 12 mg, hyoscyamine sulfate 0.1 mg, scopolamine hydrobromide OD2 mg/Tab Bot 100s, 500s
 Use: Sedative and antispasmodic
MYODYNE, (Canfield) Ointment sulfate 1 1/2 gr niacin 25 mg, Vit E 25 I u/Tab Bot 100s
 Use: Leg cramps
MYOFLEX, (Warren Teed) Triethanolamine salicylate 10% in a vanishing cream base Tube 1 oz, Jar 1 lb
 Use: Topically analgesic
MYOLAX, (Key Corp) Mephenesin 500 mg/Tab Bot 100s, 1000s
 Use: Muscle relaxant
MYOMEPHITANE TABLETS, (Testagar) Amobarbital 8 mg, mephenesin 400 mg, physostigmine salicylate 0.25 mg, homatropine methylbromide 0.60 mg, salicylamide 130 mg/Tab Bot 100s, 1000s
 Use: (1 or 2 tid, p.c.) Antispasmodic & analgesic
MYOPHANE, (Riker) Orphenadrine hydrochloride 50 mg/Tab
 Use: Muscle spasm and parkinsonism
MYOPONE (Drug Products) dl alpha tocopherol acetate 50 mg, whole wheat germ oil 120 mg/Perlette Oint contains 35% solvent ext wheat germ oil Perlette, Bot 30s, 100s Oint, Jar 1 oz, 1 lb
 Use: Myositis & fibrositis
MYOPURIN (Polch) Digoxin 0.06 mg, gitalin O 19 mg/Tab Bot 100s
 Use: Orally, cardiac
MYORDIL, (Winthrop Lab) Coronary vasodilator Under study
MYORGAL, (Winthrop Labs) Mysuran NN' Bis(2 diethylamino-ethyl) azomide bis 2 chlorobenzyl chloride See Mytelase, Tab (Winthrop Labs)
 Use: Chlamerigic
MYOSOL AND EST, (Uimer) Testosterone 25 mg, estrane 2 mg/cc Vial 10 cc
 Use I.M., menopausal syndrome
MYOSOL PRESTOGEN (Uimer) Estrane 2.5 mg, progesterone 25 mg/cc Vial 10 cc
 Use I.M. progesterone therapy
MYOSOL PROGESTERONE, (Uimer) Progesterone 25 mg/cc Vial 10 cc
 Use I.M. progesterone therapy
MYOSOL STEROTATE, (Uimer) Testosterone 25 mg/cc Vial 10 cc
 Use I.M. androgen deficiency
MYOSOL TERGESTEN, (Uimer) Estradiol 1.2 mg, progesterone 25 mg/cc Vial 10 cc
 Use I.M. hormone therapy
MYOSOL ULTROGEN, (Uimer) Estrogenic sub 20,000 u/cc Vial 10 cc
 Use I.M. estrogenic therapy
MYOSPAX (Chicago Pharmacal) Symamale

- 200 mg, salicyltamide 210 mg., acetophenetidin 150 mg., caffeine 30 mg./Tab. Bot. 40s, 100s.
Use: Muscle disorders.
- MYOTEN**, N.N.D. 1962 (Central) See: Mephensin, N.F.
- MYOTRATE**, (Sig Labs) Pentaerythritol tetranitrate 10 mg. or 20 mg./Tab. Pentaerythritol tetranitrate 30 mg. or 45 mg./Cap. Bot. 100s, 1000s.
Use: Coronary insufficiency, angina pectoris.
- MYOXANE**, N.N.D. 1962, (Ascher) Mephensin. Tab. 0.5 Gm, Bot. 100s, 1000s.
Use: (1-2 i.i.d.) Muscle relaxant.
- MYPROZINE**, (Lederle) Pimaricin.
Use: Antifungal.
- MYRINGACAINE**, (Upjohn) Benzocaine 4%, ephedrine HCl 0.44%, ortho-hydroxyphenylmercuric chloride 0.05% in glycerin & propylene glycol, Bot. ½ oz.
Use: Topically, ear infections.
- MYRINOL**, (Smith-Dorsey) Sulfathiazole 2%, benzocaine 0.5% in propylene glycol, Urea 10%, Bot. ½ oz., 1 pt., 1 gal.
Use: Topically, ear infections.
- MYRISTYL-γ-PICOLINIUM CHLORIDE**.
See: Wel-Tone B (Broemmel)
- W/Calamine**, zinc oxide, benzocaine. See: Camellians Lotion, (Upjohn)
- MYRJ 82**, (Allac Pow, Co) Polyoxyl 40 Stearate, U.S.P.
- MYROBIE**, (Meyer)
1: di-Desoxyephedrine HCl 5 mg., thyroid 1 gr., atropine sulfate 1/360 gr., aloin ¼ gr./Tab. Bot. 100s, 1000s.
2: di-Desoxyephedrine HCl 5 mg., thyroid 1 gr., atropine sulfate 1/360 gr./Tab. Bot. 100s, 1000s.
3: di-Desoxyephedrine HCl 5 mg., thyroid 1 gr., phenobarbital ¼ gr./Tab. Bot. 100s, 1000s.
Use: Antiobesity.
- MYROPIN**, (Meyer Chem. Co.) Procaine penicillin G 300,000 u., Pot. penicillin G 100,000 u./dose. Vial.
Use: Penicillin therapy.
- MYSOLINE**, N.N.D. 1962. (Ayers) Frimidine 5-phenyl-5-ethyl-hexahydroxypyrimidine-4,6-dione, 2-dihydrophenobarbital. Tab. 50 mg or 250 mg Bot. 100s, 1000s. Sup. (250 mg./5 cc) Bot. 8 oz.
Use: Anticonvulsant.
- MYSONE**, (Columbus) Lotion, Bot. 15 cc.
Use: Topically, anti-inflammatory
- MYSTECLIN**, (Squibb) Tetracycline HCl 250 mg., nystatin 250,000 u./Cap. Bot. 16s, 100s. Susp.: Each teaspoonful is equiv. to one half-strength capsule, Bot. 2 oz. Also available in half strength.
Use: Antibiotic & antifungal therapy.
- MYSTECLIN-F CAPSULES**, (Squibb) Tetracycline phosphate complex equiv. to 250 mg. tetracycline HCl, amphotericin B 50 mg./Cap. Bot. 16s, 100s. Half-strength cap Bot. 16s.
Use: Antimicrobial therapy and antifungal prophylaxis.
- MYSTECLIN-F FOR AQUEOUS DROPS**, (Squibb) Phosphate potentiated tetracycline 100 mg., amphotericin B 20 mg./cc. Drop. Bot. 10 cc.
Use: Antibiotic and antifungal.
- MYSTECLIN-F FOR SYRUP**, (Squibb) Phosphate-potentiated tetracycline HCl 125 mg., amphotericin B 25 mg./5 cc, Bot. 60 cc., 8 oz.
Use: Antibacterial therapy and antifungal.
- MYSTECLIN-F PEDIATRIC DROPS**, (Squibb) Tetracycline equiv. to tetracycline 100 mg. w/pot metaphosphate and amphotericin B 20 mg./cc, Bot. 12 cc.
Use: Antibiotic.
- MYSTECLIN-V**, (Squibb) Tetracycline phosphate equiv. to 250 mg. of tetracycline HCl, and mycostatin 250,000 u./Cap. Bot. 16s, 100s. Also available in half strength
Use: Antibiotic and antifungal therapy
- MYSTECLIN-V PEDIATRIC DROPS**, (Squibb) Tetracycline phosphate complex equiv. to 100 mg. of tetracycline HCl, mycostatin 100M u. in an oil suspension/cc. Bot. w/ dropper 10 cc.
- MYSTECLIN-V SUSPENSION**, (Squibb) Tetracycline phosphate complex equiv. to 125 mg of tetracycline HCl, mycostatin 125M u. in an oil suspension/5 cc. Bot. 2 oz.
- MYSURAN**. See: Mytelase, Tab. (Winthrop Labs.)
- MYTELASE**, N.N.D. 1962. (Winthrop Labs) Ambenonium chloride N.N.'bis (2-diethylaminoethyl) oxamide bis-2-chlorobenzyl chloride 10 mg., 25 mg./Tab. Bot. 100s.
Use: Myasthenia gravis.
- MYTINIC**, (Bristol) Iron, Iver & Vit. B complex including B-12 in 2 dosage forms. Liq. Bot. 8 oz., 12 oz., 1 gal. Tab., Bot. 100s, 500s.
Use: Nutritional hypochromic anemias.
- MYTOLON CHLORIDE**, (Winthrop Labs)
- MYTRATE**, (Professional Pharmacol) Epinephrine bitartrate 2%, benzalkonium chloride 1 in 25,000 in sterile buffered methylcellulose solution, Dropper bot. of 15 cc.
Use: Decongestant.
Benzoquinonium, 2,5 bis-(2-diethylamino-propylamino)-benzoquinone-bis-(benzyl chloride). Vial, multiple dose 13 mg/cc.) 10 cc.
Use: Adjunct to surgical anesthesia for neuromyal block & muscular flaccidity.

M-Z SOLUTION

484

- M-Z SOLUTION.** (Crookes Barnes) Zinc sulf $\frac{1}{4}\%$, piperocaine HCl $\frac{1}{4}\%$, phenylephrine HCl 0.12% with boric acid and methylcellulose Lacrivial 15 cc
Use: Antiseptic decongestant
- NABADIAL** (Brean) (Melhandrol, Methyl Androstenediol) Vial (50 mg/cc) 10 cc
Use (25-50 mg twice weekly) Anabolic hormone
- NABCON** (Brewer) Vitamin B Complex One teaspoonful supplies the same amount of complete natural B Complex contained in thirty five $\frac{1}{2}$ Gm tablets of primary grown whole brewers yeast Bot 120 cc
Use: Vitamin B complex deficiency
- NABOCAL** (Rand) Vit A 5000 u, D 1200 u, B 1 3 mg, B 2 2.5 mg, niacin 20 mg, Vit B 6 0.65 mg, pantothenic acid 5 mg, Vit C 30 mg, B 12 2.5 mg, folic acid 0.33 mg, Vit E 3 mg, bone meal 644 mg, minerals/Cap Bot 100s, 1000s
Use: Vitamin & mineral therapy
- NABOCAL IMPROVED.** (Rand) Vit B 1 1 mg, B 2 1.5 mg, B 6 2.5 mg, B 12 2.5 mcg, C 70 mg, folic acid 0.34 mg, niacinamide 20 mg, cal pantothenate 3.5 mg, cal lactate 3 gr, ferrous gluconate 4 gr, menadione 0.5 mg, vit E mixed tocopherols 3 mg/Tab Bot 100s, 1000s
- NACETYL.** (Allison) N Acetyl p aminophenol 5 gr, butabarbital sodium $\frac{1}{4}$ gr/Tab Bot 100s, 1000s
Use: Analgesic
- NACITRA.** (McNeil) Citric acid & Sod carb monohydrate Tab 10 gr Bot 100s 1000s
Use: Increase alkali reserve
- NACLEX.** (Rabins) 3 Benzylthiomethyl 6 chloro 7 sulfamyl 1 2 4 benzalhadiazine 11 ds oxide Benzthiazide 50 mg/Tab Bot 100s
Use: Diuretic and antihypertensive
- NACTISOL** (McNeil) Bulbarbital sodium 15 mg paldine methylsulfate 4 mg/Tab Bot 100s 500s
Use: Treatment of peptic ulcer
- NACTON.** (McNeil) Paldine methylsulfate 4 mg/Tab Bot 100s 500s
Use: Anticholinergic
- NADC** (Tilden) Phenylephrine HCl 10 mg, pyrilamine maleate 25 mg/Tab Bot 100s TD Caps (Phenylephrine 15 mg, pyrilamine maleate 25 mg)
Use: Colds and allergic rhinitis
- *NAEPAIN HYDROCHLORIDE, N.F., N.N.D.** 1962 Inj See Amylsine Hydrochloride (Novocal)
- W/Dihydrostreptomycin sulfate & Sod penicillin G See: P D N Dental Cones (Novocal Chem Co)
- *Generic name recognized by N.F., U.S.P. or A.M.A.
- *NAFCILLIN** (Wyeth) 6 (2 ethoxy-1 neph thimido) penicillanic acid
Use: Antibiotic
- NA-FEIN** (Warren Pharmacol) Fluoride 1 mg/Dose Bot Tabs 100s, liq 2 oz
Use: Control of dental caries
- NAEIRON.** (Parke, Davis) Iron & Sod cit rate Kapsel 7 $\frac{1}{2}$ gr Bot 100s, 1000s
Use: Iron deficiencies
- NAFTA-CORT.** (Rowell) Iodochlorhydroxyquin 0.5%, hydrocortisone alcohol 0.1%, neomycin sulf 0.15% in a base of naftalan, Ichthammol, colamine and petrolatum Tube 7 Gm, 28 Gm
Use: Anti-inflammatory
- NAFTAL PASTE.** (Rawell) Naftalan 10%, Ichthammol 2%, calamine 44%, petrolatum 44% Jar 2 oz, 1 lb
Use: Allergic chronic dermatitis
- NAFTALAN.** (Danner) (Rawell) Mineral crude oil 91%, soap 4%, manlan wax 2%, rape oil 3%
- NAGANOL.** See: Suramin Sodium (Naphuride Sodium Winthrop Labs)
- NA-GENT.** (Raymer) (Sod gentisate, Sod 2,5 dihydroxybenzoate) Paw, Bot 50 Gm Tab 325 mg Bot 100s, 500s, 1000s
Use: (2 3 q 3 h) Arthritis
- NAGIST Timed Tabs** (Leads) Phenylpropanolamine HCl 40 mg phenylephrine HCl 10 mg phenyltolaxamine dihydrogen citrate 15 mg, chlorpheniramine maleate 5 mg/sustained action Tab Bot 50s
Use: Decongestant antihistamine
- NAGUA** (Schering) *Trichlarmethiazide
Use: Diuretic
- NALDECON** (Bristol) Phenylephrine HCl, 10 mg; phenylpropanolamine HCl, 40 mg; phenyltolaxamine citrate, 15 mg; chlorpheniramine 5 mg/Tablet; half each in gradient in inner and half in outer layer; each tsp, $\frac{1}{2}$ above amounts, Tabs Bot 50s (Syrup) 1 pt
Use: Hayfever, sinus congestion
- NALERTAN.** (Irwin, Neisler) Chlorpheniramine tartrate 8 mg, pyrilamine tartrate 25 mg, d-amphetamine tartrate 10 mg complexed/Durabond tabule Bot 30s, 100s 500s
Use: Control of allergy
- NALGISED** (Chicago Pharm) Not listed Acetaphenetidin 1 $\frac{1}{2}$ gr, aminopyrine 2 gr, caffeine $\frac{3}{4}$ gr, phenobarbital $\frac{1}{2}$ gr/Tab Bot 100s 1000s
Use: Orally (1 2 1 1 d) analgesic w/seda tive
- *NALIDIXIC ACID** (Winthrop Lab) 1 Ethyl 7 methyl 1,8 naphthyridin 4 one 3 carboxylic acid See: Negram

- NALLINE HYDROCHLORIDE**, N.H.D. 1961. (Merck Sharp & Dohme) N-allylnormorphine. Amp. 0.2 mg./cc.) 1 cc.; (5 mg./cc.) 1, 2 cc. & 10 cc.
 Uses: I.V. (5 to 40 mg.), overdosage of morphine & derivatives.
- NALORPINE HYDROCHLORIDE**, U.S.P., N.H.D. 1962; Inj., U.S.P. N-Allylnormorphine HCl.
 See: Nalline, N.N.R., Amp. (Merck Sharp & Dohme)
 Letidrona, Amp. (Burroughs Wellcome)
- NALOXONE HYDROCHLORIDE**. [Endo] N-allyl-14-hydroxy-nordihydromorphinone hydrochloride.
 Use: Narcotic antagonist.
- NALUTRON**. (Winthrop Labs.) Progesterone in sesame oil, Amp. (1 mg./cc., 2 mg./cc., 5 mg./cc.) 1 cc., Box 5s, 50s.
 Use: Habitual & threatened abortion.
- NANDROLONE PHEENPROPIONATE**, 19. 17-phenylpropionate, Nandrostefenolone phenylpropionate. See: Durabolin Inj. (Organon)
- NAMURON**. See: Cyclobarbital. (Winthrop Labs.)
- NANDROLONE DECANOATE**. Deca-Durabolin (Organon)
- NAOTIN**. (Drug Products) Sod. nicotinate. Amp. (equiv. to 10 mg. nicotinic acid/cc) 10 cc., Box 12s, 25s, 100s.
 Use: I.V. & I.M., nicotinic acid therapy.
- NAPENTAL**. (Massengill) Pentobarbital Sod. Cap. ¼ gr., Bot. 100s, 500s, 1000s; 1½ gr., Bot. 100s, 500s, 1000s. Vial (4 gr.) 3 cc. w/5 cc. diluant
 Use: Hypnotic.
- NAPENTAL (ELIXIR)**. (Massengill) Pentobarbital Sod. ½ gr./5 cc. Bot. 1 pt., 1 gal. Use: (1-2 tsp. 1 to t.i.d.) Hypnotic action.
- NAPHAZOLINE HYDROCHLORIDE**, N.F., N.H.D. 1962; Soln., N.F. 2-[1-Naphthylmethyl]-2-imidazoline HCl. Mild Sol., Strong Sol., N.F.
 See: Privity Hydrochloride (Ciba).
 Prizole HCl (DuMont)
 W/Antazoline phosphate, boric acid, phenylmercuric ocelate, sod. chloride, sod. carbonate anhydrous. See: Vasocin-A Ophthalmic, Sol. (Smith, Miller & Patch)
 W/Phenylmercuric ocelate. See: Closol, Nasal Spray (Vitamax Corp.)
 W/Prednisolone, chlorpheniramine maleate, methylcellulose. See: Vasopred, (Smith, Miller & Patch).
- NAPHTHONONE**. (Robins Co., Inc.) 2-(2-hydroxy-1-naphthyl)cyclohexanone.
 Use: Antitussive.
- B-NAPHTHYL SALICYLATE**. (Betol, Naphthosolol, Salinophthal.)
 Use: G.I. & G.U., antispasmodic.
- NAP-OL OINTMENT**. (Superior Pharm.) Dimethyl subnitrate 30%, castor oil 30%, lanolin anhy. 32%, ceresin wax 4%, balsam peru 3%, benzoic comp. tr. 1%. Oint., Jar 1 oz.
 Use: Topically, skin protective.
- NAPRIE CAPSULES**. (Marion Lab) Phenylpropanolamine HCl 25 mg., phenylephrine HCl 15 mg., pyrilamine maleate 25 mg., chlorpheniramine maleate 4 mg./Cap. Bot. 50s, 250s.
 Use: Relief of sinus and nasal congestion
- NAPRIE SYRUP** (Marion Labs) Phenylpropanolamine HCl 10 mg., phenylephrine HCl 5 mg., pyrilamine maleate 7.5 mg., chlorpheniramine maleate 1.5 mg./5 cc. Bot. 1 pt. Use: Relief of sinus and nasal congestion.
- NAPRYLATE**, N.N.R. 1955. (Strassenburgh) Sod. coprylate w/sinc coprylate. Creme, vaginal, Sod. coprylate 100 mg Zinc coprylate 50 mg./Gm. of base. Tube 90 Gm. w/applicator. Oint., Same conc. as Creme, Tube 21.25 Gm., Jar 454 Gm. Pow., Same conc. as Creme, Bot. 3543 Gm. Supp., vaginal, 3 Gm. gelatin Cap.
 Use: Antifungal agent.
- NAP-TABS**. (Sandoz) Secobarbital 65 mg., amobarbital 40 mg., phenobarbital 20 mg./Tab. Bot. 30s, 100s.
 Use: Hypnotic-sedative.
- NAQUA** (Schering) Trichlormethiazide 2 mg. or 4 mg./Tab. Bot. 100s
 Use: Diuretic/hypotensive.
 w/serpine See: Naquival (Schering)
- NAQUIVAL**. (Schering) Trichlormethiazide 2 mg. or 4 mg., serpine 0.1 mg./Tab. Bot. 100s.
 Use: Treatment of hypertension.
- NARAKON**. (Chasebrough) Benzalkonium chloride 1:3500, astringent, with or without di-desoxyephedrine hydrochloride, 1%. Nasal sol. Plain, Bot. 1 fl. oz. w/desoxyephedrine, Bot. 1 fl. oz.
 Use: Decongestant.
- NARB**. (Cole Chem.) Phenobarbital ¼ gr., Sod. nitrite 2 gr., w/Sod. glycocholate, cholanathanin & irisin/Tab. Bot. 100s.
 Use: Orally (1-2 q.i.d.) Vasodilator.
- NARCOL**. (Barre) Nascapine 10 mg., phenyltoloxamine 10 mg., NH₄Cl in a thixotropic suspension 0.2 Gm./5 cc.
 Use: Cough medicinal.
- NARCOTINE**, N.H.D. 1962. (Merck Sharp & Dohme). See: Nascapine, N.H.D. 1961.
- NARDIL**, N.H.D. 1962 (Warnar-Chilroff) phenylethyldiazine dihydrogen sulfate

* Generic name recognized by N.F., U.S.P. or A.M.A.

- Phenelzine sulfate (beta phenyl ethyl hydrazine) 15 mg /Tab; Bot 100s
 Use: Anti depressant
- NARETIC.** (Elder) Acetophenetidin 2 gr, acetylsalicylic acid 4 gr, strychnine sulfate 1/240 gr, caffeine 1/2 gr, camphor monobromate 1/2 gr, gelsemium 1/2 gr /Tab Bot 100s 1000s
 Use: (1 2 1 1 d) Analgesic
- NARKINE.** (Tilden) Chloral hydrate 44 gr, potassium bromide 44 gr /fl oz Bot pts, gals
 Use: Hypnotic, anodyne, antispasmodic
- NARCOGEN** (Tilden) Chloral hydrate 80 gr Pot bromide 80 gr, hyoscyne HBr 1/25 gr, horse-nettle, wild lettuce & celery seed/fl oz Bot 1 pt, 1 gal
 Use: Hypnotic, anodyne & antispasmodic
- NARONE.** (Ulmer) Dipyrone Tab 5 gr, Bot 100s; Vial (50%) 30 cc
 Use: Analgesic
- NARTATE** (G F Harvey) Dipyrone Tab 300 mg, Bot 100s; Vial (50%) 30 cc
 Use: Analgesic
- NASADRINE.** (Nayes) Ephedrine chlorobutanol, Sod chloride, menthol & aromatics Bot 1/2 oz
 Use: Nasal decongestant
- NASALENE, No. 1.** (Abbott) Acetaminid 1%, anesthesin 0.5%, ephedrine alkaloid 0.5%, bismitol subnitrate 3.3%, petrolatum, menthol, eucalyptol & aromatics q.s. to make 100% Tube 1/2 oz
 Use: Irritations of nasal membranes
- NASOL.** (Tilden) Ephedrine, menthol, camphor, eucalyptol, pine oil, liquid petrolatum Bot 1 oz 1 oz w/dropper, pts
 Use: Nasal decongestant
- NASO-MIST INHALER** (Bryant) 1 desoxy-ephedrine carbonate eucalyptol, menthol, thymol camphor Inhaler/16s
- NASOPHEN** (Prama) Phenylephrine hydrochloride 0.25%, 1% Bot 1 oz, 1 pt
 Use: Treatment of vasomotor rhinitis, sinusitis allergic rhinitis and as an adjuvant in the treatment of acute coryza
- NATABA.** (Glenwood) Sod p aminobenzoate Pow Bot 100 Gm, 1 lb Tab 0.5 Gm Bot 100s
 Use: Rheumatoid arthritis
- NATABEC.** (Parke, Davis) Vit A 4000 u, D 400 u, B 1 3 mg, B 2 2 mg, B-6 3 mg, C 50 mg, B 12 2 mg Intrinsic factor concentrate 5 mg, folic acid 1 mg, niacin 10 mg, cal carbonate 600 mg, ferrous sulfate 150 mg, rutin 20 mg, synkamine HCl 0.5 mg/Cap Bot 100s, 1000s
 Use: Vitamin, iron & calcium deficiencies
- NATAGEST.** (Alphabet) Vit A 4000 u, B-1 3 mg, B-2 3 mg, B-6 5 mg, B 12 3 mcg, calcium pantothenate 5 mg, folic acid 0.25 mg, niacinamide 15 mg, C 75 mg, hesperidin 25 mg, D 400 u, K 0.5 mg, calcium carbonate 600 mg, iodine 0.1 mg, iron 66 mg, copper 0.2 mg, manganese D 2 mg, magnesium 0.5 mg, malybdanum 0.15 mg, zinc 0.5 mg, fluarline 0.015 mg, cobalt D 1 mg /tab
 Use: Prenatal dietary supplement
- NATALINS BASIC.** (Mead Johnson) Iron 40 mg, calcium 250 mg, Vit C 100 mg, O 400 u /Tab Multip Pak/100s/3s
 Use: Prenatal suppository
- NATALINE COMPREHENSIVE.** (Mead Johnson) Iron, Calcium, Vit C, D plus 8 additional vitamins/Tab Multip Pak/100s/3s
 Use: Prenatal suppository
- NATABINE.** (Dumas Wilson) Tab Bot 100s, 1000s
 Use: Obstetrical Dietary Supplement
- NATELIE.** (Gross) Vit A & D, B complex, C, K, folic acid calcium pantothenate, calcium carbonate, iron, hesperidin and rutin Cap 100s
 Use: Phosphorus free vitamin supplement
- NATESTRIN.** (Upjohn) Estragenic substance Tab 1 mg Bot 100s, 500s, 1000s
 Use: Estrogen therapy
- NATIBAIN.** (Varick) Digitaline 1 part, ouabaine 2 parts 5al, Bot 10 cc
 Use: Digitalis therapy
- NAT-I-LAC.** (National Drug) Milk protein, lactose & mineral salts Vial 10 cc, 20 cc
 Use: IM (5 10 cc q 3 5 days) nonspecific protein therapy
- NATOLA.** (Parke Davis) Each Gm of liq contains Vit A 55 000 u D 11,000 u. Each Cap contains Vit A 5000 u, D 1000 u liq 50 cc Veal w/dropper cap Cap, Bot 100s, 250s
 Use: Deficiencies of Vit A & D
- NATOLONE** (National Drug) Pregnenolone Tab 50 mg, 100 mg Bot, 24s Vial (100 mg/cc) 9 cc
 Use: Orally & IM, rheumatoid arthritis
- NATOPHEROL.** (Abbott) Mixed Iacopherols Cap 25 mg Bot 100s Cap 37.5 u Bot 100s, 500s, 1000s
 Use: Orally, Vit E therapy
- NATOPHEROL ACETATE.** (Abbott) Mixed Iacopheryl acetate Cap 100 u Bot 100s 1000s In oil, Vial (200 u/cc) 10 cc
 Use, Orally & IM, Vit E therapy
- NATOREXIC** (Walker) Diethylpropion 25 mg, Vit A 2000 U.S.P. units D 200 U.S.P. units, E 1 IU, B 1 1 mg, B 2 1 mg, niacinamide 10 mg calcium pantothenate 2 mg, B 6 1 mg, C 25 mg, B 12 1 mcg, hesperidin complex 67 mg, ferrous fumarate 333 mg, dicalcium phosphate 333

- mg., Cu .065 mg., Mn .065 mg., Zn 0.065 mg., Mg .065 mg., I .0066 mg., K .83 mg./Tab. Bot. 100s.
 Use: Prevent weight gain during pregnancy and lactation.
- NATRASCORB.** (U.S. Vitamin) Sod. ascorbate.
 Amp.; Vit. C equiv. 2 cc. (100 mg./cc.) Box 5s, 25s, 100s; 5 cc. (200 mg./cc.) Box 5s, 25s, 100s; 1 cc. (500 mg.) Box 6s, 25s, 100s.
 Drops; (150 mg./cc.) Bot. 15 cc.
 Tab. (50 mg.) Bot. 100s; (100 mg.) Bot. 100s, 1000s; (250 mg.) Bot. 50s, 100s, 1000s.
 Use: Vit. C therapy.
- NATRI-C.** (Physiological Chemicals) Sod. ascorbate. Tab. 120 mg. Bot. 100s.
 Use: Vit. C therapy.
- NATRICO.** (Drug Products) Pot. nitrate 2 gr., Sod. nitrite 1 gr., nitroglycerin $\frac{1}{4}$ gr., crotaegus oxycantha $\frac{1}{4}$ gr./Pulv. Bot. 100s.
 Use: Hypertension.
- NATRIDOL.** (Warren-Teed) Codeine sulfate $\frac{1}{6}$ gr., Sod. salicylate $3\frac{5}{6}$ gr./dr. Bot. 2 oz., 1 pt., 1 gal.
 Use: Cough preparation.
- NATRINIL.** (National Drug) Cation exchange resin, carboxylic type. Box 24 individual dose packets, 10 Gm. each. Bot. 10 oz.
 Use: Withdrawal of sodium from system.
- NATRO-NAWTORN COMP.** (National Drug) Hawthorn ext. $\frac{1}{4}$ gr., pot. nitrate $2\frac{1}{2}$ gr., sod. nitrite $\frac{1}{2}$ gr., nitroglycerin $1\frac{1}{2}$ gr., powd. aconite root $2\frac{1}{2}$ gr., veratrum viride ext. $1\frac{1}{10}$ gr., sod. bicarbonate 2 gr./Tab. Bot. 100s, 1000s
 Use: Hypertension and angina pectoris
- NATURAL VITAMIN A IN OIL, N N R. 1955.**
 See: Oleovitamins A, U.S.P.
- NATURETIN, N N D. 1962** (Squibb) Bendroflumethiazide 2.5 mg or 5 mg./Tab. Bot. 100s, 1000s.
 Use: Control of edema.
- NATURETIN W/K.** (Squibb) Benzhydroflumethiazide 5 mg., potassium chloride/Tab. Bot. 100s.
 Use: Diuretic and hypotensive
- NAUCAINE.** (Taylor Labs) Procaine HCl, Sol. 10% Bot. w/dropper 60 cc., 1 pt. Tab. 120 mg. Bot. 50s, 250s.
 Use: Nausea & pain due to gastric ulcers.
- NAUDEL** (Elder) Benzocaine 100 mg., vit. B-6 25 mg., nicotinamide 25 mg., methanone 100 mg., pentobarbital sod. 15 mg./Tab. Bot. 25s, 100s, 500s.
 Use: (1 p.c.) In therapy of hyperemesis gravidarum.
- NAUDOX.** (Dumas-Wilson) Pyrilamine maleate 25 mg., pyridoxine HCl 50 mg./cap. Bot. 100s, 1000s.
 Use: (1 cap. daily on rising) Morning sickness of pregnancy.
- NAUSEA THERAPY.**
 See: Apolamine, Tab. (Winthrop Labs.)
 Banausea, Tab. (Grant)
 Banamine, Tab. (Pfizer)
 Chloratone, Cap. & Paw. (Parke, Davis)
 Chlorpromazine, Amp. & Tab. (Smith, Kline & French)
 Emetrol, Liq. (Kinney)
 Maretox
 Marezine
 Motion Sickness.
 Pyridoxine HCl (Various Mfr.).
 Vasana Tablets (Schering).
- NAVICAPS.** (Dumas-Wilson) Cal. levulinate 360 mg., ferrous sulf. 12.5 mg., Vit. A 1000 u., O 100 u., B-1 0.25 mg., B-2 0.5 mg., niacinamide 2 mg., cal. ascorbate 8.5 mg., cal. pantothenate 0.3 mg., B-6 0.1 mg., folic acid 0.1 mg., B-12 0.5 mcg. with trace elements/Cap. Bot. 100s, 1000s.
 Use: (4 daily) A phosphorus-free dietary supplement
- NAVIDRIX (NAVIDREX, *CYCLOPENTHAZIDE).** (Ciba) Cyclopenthaside, 3-cyclopentylmethyl derivative of hydrochlorothiazide
 Use: Oral diuretic.
- NAVITOL WITH VIOSTEROL.** (Squibb) Liq. Vit. A 5900 u., and D 1180 u./0.1 cc
 Vials 10 & 50 cc. Caps. Not listed Vit A 5000 u., O 1000 u., from fish liver oil & irradiated ergosterol/Cap., Bot. 100s, 250s.
 Use: (1 daily) Vit. A & D deficiencies.
- N.B.T. SPRAY.** (Novacol) Benzocaine 20%, tetracaine 1%, naepaine 1%, alcohol 45% with propellant in pressure bottle.
 Use: Topical anesthetic.
- N-CHLORO COMPOUND ANTISEPTICS**
 See: Antiseptics, N-Chloro Compounds.
- ND-8 TABS.** (Hylix) Pyrilamine maleate 16 mg.; chlorpheniramine maleate 2 mg.; phenindamine tartrate 10 mg.; phenylephrine HCl 10 mg.; ascorbic acid 50 mg./Tab. Bot. 50s, 500s.
 Use: Cold Therapy
- NEBRALIN.** (Smith-Dorsey) Pentobarbital 90 mg., mephesisin 425 mg./T.R. Tab. Bot. 50s.
 Use: Sedative with muscle relaxation.
- NEBS.** (Norwich Pharmacol) N-Acetyl-*p*-amino-phenol. Bot. 30s
 Use: Analgesic.
- NIBU-PREL** (Mahan) Isoproterenol sulf. 0.4%, phenylephrine HCl 2%, propylene glycol 10%. Uq. Vial 10 cc.
 Use: Bronchial asthma
- NEBUTABS.** (Pema) Penicillin pot. 250 =,

200 m, 100 m, or 50 m /sublingual tab
Bot 24s and 100s

NECTADON, NND 1962 (Merck Sharp & Dohme) See: Noscopine

NECOBIN. (E S Miller) Neostigmine methyl sulfate 0.6 mg, atropine sulfate 0.3 mg, Vit B 12 100 mcg/cc Vial 10 cc, Box 1s, 6s, 25s

Use SC, IM (1 cc daily), muscle relaxant

NEGATAN. (Lilly) (Negotal) Polymerized di sulfonic-dioxy dimethyl diphenyl-methane acids Sol 45% Bot 1 fl oz

Use Cervicitis & Trichomonos vaginalis vaginitis

NEGATOL See: Negoton, Sol & Supp (Lilly)

NEGRAM (Winthrop Lab) Nalidixic acid

NEKATUSSIN (Bruce) Noscopine 10 mg terpin hydrate 4.25 mg, sodium citrate 60 mg, pyrilamine maleate 7 mg, alcohol 2% /5 cc Bot pts gals

Use Coughs

NEMBUDEINE 1/4, FILMTABS. (Abbott) Codeine sulfate 15 mg, acetophenetidin 150 mg, aspirin aluminum 210 mg, nembutal 15 mg caffeine 30 mg /Tab 100s, 500s, 1000s

NEMBUDEINE 1/2, FILMTABS. (Abbott) Codeine sulfate 30 mg, aspirin aluminum 210 mg acetophenetidin 150 mg, nembutal 15 mg caffeine 30 mg /Tab Bot 100s, 500s, 1000s

NEMBUDEINE 1, FILMTABS (Abbott) Codeine sulfate 60 mg, aspirin aluminum 210 mg, acetophenetidin 150 mg, nembutal 15 mg, caffeine 30 mg /Tab Bot 100s, 500s, 1000s

NEMBUDEINE FILMTAB WITHOUT CODEINE. (Abbott) Aspirin aluminum 210 mg, acetophenetidin 150 mg, caffeine 30 mg, pentobarbital 15 mg /Tab Bot 100s, 500s, 1000s

Use Analgesic w/sedative

NEMBU-DONNA CAPSULES. (Abbott) Cap 1/2 gr Nembutal 1/2 gr, belladonna ext 1/6 gr /Cap Bot 100s, 1000s; 1/4, Nembutal 1/4 gr belladonna ext 1/6 gr /Cap Bot 100s, 1000s

Use: (1 2 11 d) Antispasmodic & sedative

NEMBU-DONNA ELIXIR (Abbott) Nembutal Sol 15 mg, belladonna ext 10 mg, total alkaloids 0.125 mg corresponding to USP belladonna tr 7 min/fl dr Bot 1 pt, 1 gal

Use: (1 2 tsp 11 d) Antispasmodic & sedative

NEMBU-GESIC (Abbott) Nembutal 30 mg aspirin 300 mg /Cap Bot 100s, 500s, 1000s

Use: Sedative and analgesic

NEMBU-SERPIN FILMTAB. (Abbott) Nembutal calcium equiv to nembutal 30 mg,

reserpine 0.25 mg /Rimtab Bot 100s, 500s

Also available in one half strength
Use: (1 h s) Sedative & tranquilizing agent

NEMBUTAL, NND 1962, ADB (Abbott) Pentobarbital Elixir 1/4 gr /5 cc. Bot 1 pt, 1 gal

W/Aspirin See: Nembu Getic, Cap (Abbott)
W/Desoxyn See: Desbutol, Cap (Abbott).
W/Ephedrine HCl See: Nembu Fedrin, Cap (Abbott)

W/Glucophylline See: Glucophylline & Nembutal, Tab (Abbott)

W/Theophylline & methyl glucamine See: Glucophylline w/Nembutal, Supp (Abbott)

NEMBUTAL & BELLADONNA CAPSULES (Abbott) Cap No 1, Nembutal Sod 1/2 gr, belladonna ext 1/6 gr Bot 100s, 1000s; No 2, Nembutal Sod 1/4 gr, belladonna ext 1/6 gr Bot 100s, 1000s

Use: Orally, sedative w/antispasmodic

NEMBUTAL & BELLADONNA ELIXIR. (Abbott) Nembutal Sod 1/2 gr, belladonna ext 1/6 gr /5 cc Bot 1 pt, 1 gal

Use: Sedative w/antispasmodic

NEMBUTAL CALCIUM, NND 1962, ADR. (Abbott) Pentobarbital Cal Enterob 3/4 gr Bot 100s, 500s; 1 1/2 gr, Bot 100s, 500s; Tab 1/2 gr, 3/4 gr, 1 1/2 gr, Bot 100s, 500s; Filmtab 1 1/2 gr

Use: Sedative & hypnotic

W/Beserpine See: Nembu Serpin, Filmtab (Abbott)

NEMBUTAL SODIUM, NNR 1952, ADR 1955 (Abbott) Pentobarbital Sod Amp: (50 mg/cc) 2 cc Box 6s, 25s; 5 cc, Box 6s, 25s, 100s

Vial: (3/4 gr /cc) 20 & 50 cc
Cap: (1/2 gr) Bot 100s, 1000s; (3/4 gr) Bot 100s, 500s, 1000s; (1 1/2 gr) Bot 25s, 100s, 500s, 1000s

Supp: (1/2, 1, 2 & 3 gr) Box 6s, 12s, 100s
Use: Sedative & hypnotic

W/Amylphylline See: Neutraphylline & Pentobarbital, Supp (Smith, Dorsey)

W/Belladonna ext See: Nembutal & Belladonna Capsules & Elixir (Abbott)

W/Dihydrocodeinone bitartrate, ephedrine HCl & cal iodide See: Colcodrine 3yr
W/Dihydrocodeinone (Abbott)

NEMISIS. (Beutlich, Inc) Amylolytic enzyme 15 mg, proteolytic enzyme 6 mg, pyridoxine HCl 50 mg, pyrilamine maleate 15 mg /Tab Bot 50s

Use: Prevent and control emesis

NEMURAL (Winthrop Labs) (Dracorbil) 4 Oxy 3 acetyl aminophenyl arsonic acid N methyltetrahydro pyridine beta-carbonic acid methyl ester, acetorzone salt of arecoline Tab 18 mg Bot 100s, 500s

- Use: Treatment of tapeworms & stomach worms in animals.
- NEO-A-FIL** (Texas Pharm Co.) Digalloyl trioleate 3% in vanishing cream. Tubes 2 oz., also stick available
Use: Sun screen
- NEO-ALYSINE (SYRUP)**, (Merrell) Sod. salicylate natural 0.3 Gm., p-aminobenzoic acid 0.3 Gm. as Sod. salt/5 cc. Bot. 1 pt., 1 gal
Use: (2-3 tsp. q. 3 h.) Rheumatic conditions.
- NEO-AMNIOTIN**, (Squibb) Aq. susp. of Estrogenic substance mixed, 1 mg. (equiv. to 10,000 I.U. estrone)/cc. 2 mg. (equiv. to 20,000 I.U. estrone)/cc. 5 mg. (equiv. to 50,000 I.U. estrone)/cc. Vial w/diaphragm cap 10 cc.
Use: Estrogen therapy.
- NEO-ANTERGAN MALEATE, N.N.D. 1962**, (Merck Sharp & Dohme) Pyrilamine maleate. Tab. 25 mg., 50 mg., Bot. 100s, 500s, 1000s.
Use: Antihistaminic.
- NEOANTIMOSAN**. See: Stibaphen (Fuadin, Amp. Winthrop Labs.)
- NEO-AQUA-DRIN**, (McKesson & Robbins) Tyrothricin 2 mg., benzocaine 10 mg./10z. Vial 12s.
Use: Irritation of mouth & throat.
- NEO-ARISTOCORT**, (Lederle) Triamcinolone acetonide 1 mg., neomycin Sulf. 5 mg./Gm. Oint. Tube 1/2 oz.
Use: Eye and ear inflammations.
- NEO-ARISTODERM**, (Lederle) Triamcinolone acetonide 15 mg., neomycin sulfate 75 mg./15 cc. Pressure Bot. 15 cc.
Use: Skin dermatoses.
- NEO-ARSOLUIN**. See: Neocarsophanamine, Amp. (Abbott).
- NEOARSPHENAMINE, N.N.B. 1949**, (Abbott) Sod. 3,3'-diamino-4,4'-dihydroxyarsenobenzeno-N-methanal sulfoxylate. (Neosalvarsan, Nauarsenobenzal) Amp. 0.3, 0.45 & 0.6 Gm., Box 5s.
Use: I.V. & I.M., antisyphilitic.
- NEO-ASMASAN**, (Reckel) Kf 250 mg., aminophyllin 125 mg., mephobarbital 30 mg., ephedrine HCl 20 mg., tri-histin 30 mg./Cap. Bot. 100s.
Use: Antispasmodic, decongestant, sedative and antihistamine.
- NEOBACIN**, (Reed & Carnrick) Neomycin sulfate 25 mg., bacitracin 2500 u./tab. Bot. 30s.
Use: Amebiasis, infectious diarrhea & dysenteries.
- NEOBACIN OINTMENT**, (Reed & Carnrick) Bacitracin 500 u., neomycin sulfate 3.5 mg./Gm. Tube 1/2 oz.
- Use: Topically, for infections due to gram-negative & gram-positive bacteria.
- NEO-SAND**, (Rowell) Aerosol spray bandage containing 5 mg. neomycin sulf. per Gm. of solids. Aerosol can 6 oz.
Use: Antibiotic bandage.
- NEOBASE**, (Berroughs Wellcome) Water in oil emulsion base, Jar 1 lb., 5 lb.
Use: Ointment base.
- NEOBIVIN**, (Walker Vitamin) Thiamine HCl 10 mg., riboflavin 5 mg., pyridoxine 1 mg., Caf. pantothenate 5 mg., nicotinamide 50 mg./Tab. Bot. 30s, 100s, 500s, 1000s. Also avail. w/Vit. C. Bot. 1000s.
Use: Vit. B deficiencies.
- NEOBILE**, (Haug) Dehydrocholic acid 50 mg., ox bile ext. 100 mg., pepsin 300 mg., pancreatin 300 mg./tab. Bot. 100s.
Use: (2-3 t.i.d.) Chronic constipation.
- NEOBILE-HP**, (Haug) Outer coating: Pepsin 1:15,000 (60 mg.), homatropine methylbromide 2.5 mg., phenobarbital 8 mg. Esteric coated core: pancreatin 60 mg., ox bile ext. 100 mg., dehydrochloric acid 50 mg. Tab. Bot. 100s.
Use: Choleric, chologogue.
- NEOBLOTIC**, (Pfizer) Neomycin sulf. 500 mg./Tab. Bot. 20s, 100s.
- NEOBLASTINE**, (Overseas) No. 1 Oxidase 12 mg., Sod. butylsulfonate 5 mg., glutathionase 10 mg./Amp. No. 2 Sod. resorcylosulfonate 10 mg., lipase 10 mg./Amp. No. 3 lipase 10 mg., glucocorticoid hormone 5 mg./Amp. Box of 36 (12 each).
Use: Cancer therapy.
- NEOBON CAPSULES**, (Roerig) Methyl testosterone 3 mg., ethinyl estradiol 0.018 mg., glutamic acid 90 mg., pancreatic substance 150 mg., liver-stomach substance 300 mg., vit. A 6000 u., D 600 u., E 15 u., cat. pantothenate 15 mg., vit. B-1 1.5 mg., B-2 1.5 mg., B-6 1.5 mg., etacnamide 150 mg., folic acid 0.3 mg., Vit. C 150 mg., B-12 3 mcg., w/Iron, molybdenum, cobalt, copper, Mang., Mag. Iodine, Pot., zinc, rutin 15 mg., L-flysine 150 mg./3 caps. Bot. 60s.
Use: 1 t.i.d. hormone & nutritional supplement.
- NEOBOVININE-20**, (Wyeth) Ferric ammonium citrate 0.8 Gm., defibrinated beef blood 56 Gm., liver fraction #1 7.6 Gm./100 cc. Bot. 8 R. oz., 12 R. oz.
Use: Hematinic.
- W/Malt & Iron**, (Wyeth) Malt ext. 38 Gm., ferric ammonium citrate 2.82 Gm., defibrinated beef blood 56 Gm., liver fraction #1 7.4 Gm./100 cc. Bot. 8 R. oz.
Use: Hematinic.
- NEO-BROMTH**, (Broyden) Bromoelate. Pemo-

- bism 2 mol wt, pyrimidine maleate 1 mol wt Tab 80 mg Bot 100s
Use (2 b i d, 5 to 7 days before menstruation) Control of premenstrual tension
- NEO-CAPOTAN** (Premo) Pyrimidine maleate
1½ gr caffeine ½ gr, phenacetin 2½ gr, aspirin 3½ gr/Tab 80 Bot 100s
Use Analgesic, antipyretic and antihistaminic
- NEOCAINE**. (Fougere) Procaine HCl Amp
Infil ones 0.50 mg, Spinal ones 50 mg 80 mg, 100 mg, 120 mg, 150 mg 200 mg & 300 mg Cont caudal ones 500 mg Box 10s
Use Spinal anesthesia
- NEOCALAMINE**. (Various Mfrs) Red ferric oxide 30 Gm, yellow ferric oxide 40 Gm, zinc oxide 930 Gm
W/Sulfolcetamide base, talc, glycerin, camphor & menthol See Neofenamide, liq (McNeil)
- W/Talc, zinc sulfate Sod thiosulfate, menthol & camphor See Calathor, lot (McNeil)
- NEO-CALCIN** (Rorer) Cal levulinate Amp
10%, 8ax 6s, 12s 25s, 100s
Use I V, calcium therapy
- NEO-CALGLUCON** (Sandoz) Cal gluconogalactogluconate Amp 20% 5 or 10 cc; 10%, 5 or 10 cc Syr 20 gr / 5 cc Bot 8 oz, 1 pt Not listed
Uses Calcium therapy
Uses Nutritional supplement
- NEOCNOLAN**, (Pilon Moore) Dehydrocholic acid 3½ gr, phenobarbital 8 mg, homotriprine methylbromide 12 mg/Tab Bot 100s, 1000s
Uses Management of dyspepsia
- NEO-CNOLEX**, (Bell Craig) Fat emulsion containing 40% w/v pure vegetable oil Bot 60 cc.
Uses Produce maximum cholecystokinetic activity in roentgen study
- NEOCIN**. (Commerce Drug) Polymyxin B sulf, neomycin, tyrothricin Lozenge. Pkg 14s
Uses Infections of mouth and throat
- ***NEOCINCNOFEN**, N.F. N.N.R. 1947s Ethyl 6-methyl 2 phenylquinoline 4 carboxylates Tab, N.F. Navataphon
Abbott—Tab 5 gr, 7½ gr Bot. 100s, 1000s
Lilly—Tab 5 gr, 7½ gr Bot 100s
Mallinckrodt—Pow, Bot 1 oz, ¼ lb, 1 lb, 5 lb
Meck—Pow, Bot 1 oz, ¼ lb, 1 lb
Uses Orally; analgesic, antipyretic & In gout
- W/Aspirin & colchicine See Neo Salicyphen, Cap (Abbott)
- * Genetic name recognized by N.F., U.S.P. or A.M.A.
- NEO-COSEFRIN**. (Cook Waite) levo-Nordefrin 1,2,4 dihydroxyphenyl 3 hydroxy 2 iso propylamine
Use Vasoconstrictor
- NEOCOD A & D OINTMENT**. (Leeds) Cod liver oil, 10% with vitamin A & D; petrolatum base Tubes ½ oz
Use Impetiginous dermatitis
- NEO-CODENYL**. (C 8 Kendall) Glyceryl guaiacolate 100 mg, dihydrocodeinone bitartrate 108 mg, methoppyrilene fumarate 50 mg, sodium citrate 1 Gm / oz Bot. 8 oz Syr
Use Antitussive
- NEO-COROVAS**. (Amire Grant) Pentaerythritol tetranitrate 30 mg / Timed cap Bot 60s 120s, 500s, 1000s
Uses Coronary vasodilator
- NEO-CORTAPAC LOTION**. (Pharmafac) Hydrocortisone alcohol 0.5%, neomycin sulfate 0.5%, in a bland lotion base with a pH of 4.5 Bot 2 oz
Use Severe skin infections
- NEO-CORT-DOME**, (Dome) Hydrocortisone ¼% ½% or 1% w/neomycin 5 mg / Gm Cream, Tube 5 Gm, ½, 1, 2, 4 oz Lot, Sqz Bot ½, 1, 2, 4 oz
Uses Anti-inflammatory
- NEO-CORT-DOME OTIC**, (Dome) Hydrocortisone alcohol 10 mg, neomycin sulfate 5 mg acetic acid 2% / cc Bot 10 cc
Use External otitis
- NEO-CORTEF CREAM**. (Upjohn) Hydrocortisone acetate 1% & 2.5% Tube 5 Gm 20 Cm
Uses Allergic & inflammatory skin conditions
- NEO-CORTEF DROPS, 0.5%**. (Upjohn) Hydrocortisone acetate 5 mg, neomycin sulfate 5 mg / cc Susp Bot 5 cc
Uses Inflammatory conditions of eye & ear
- NEO-CORTEF DROPS, 1.5%**. (Upjohn) Hydrocortisone acetate 15 mg, neomycin sulfate 5 mg / cc Susp, Bot w/dropper 2.5 cc, 5 cc
Uses Anti-inflammatory
- NEO-CORTEF, LOTION 1%**. (Upjohn) Hydrocortisone acetate 10 mg, neomycin sulfate 5 mg, methylparaben 2 mg, butyl p-hydroxybenzoate 3 mg / cc Plastic Squeeze Bot 15 & 30 cc
- NEO-CORTEF, NASAL SPRAY**. (Upjohn) Hydrocortisone acetate 15 mg at 5 mg, phenylephrine HCl 2.5 mg, neomycin sulfate 5 mg / cc Plastic spray bot (1.5%) 10 cc (0.5%) 15 cc
Uses T's & gram positive & gram negative bacterial infections of the nasal mucosa

NEO-CORTEF OINTMENT, (Upjohn) 0.5%
 —Hydrocortisone acetate 5 mg., neomycin sulfate 5 mg., in bland, nonirritating oint. base/Gm. Tube 5 & 20 Gm. 1% —Hydrocortisone acetate 10 mg., neomycin sulfate 5 mg./Gm. Oint., Tube 5 & 20 Gm. 2.5%
 —Hydrocortisone acetate 25 mg., neomycin sulfate 5 mg./Gm., Oint., Tube 5 & 20 Gm.
 Use: Broad range antibiotic w/anti-inflammatory effect.

NEO-CORTEF, EYE-EAR OINTMENT 0.5% OR 1.5%, (Upjohn) Hydrocortisone acetate 5 mg. or 15 mg., neomycin sulfate 5 mg./Gm. Tube 1 dr.
 Use: Ocular infections.

NEO-CORTEF PARENTERAL, (Upjohn) Hydrocortisone acetate 50 mg., neomycin sulfate 5 mg./cc, Vial 1 cc., 5 cc.
 Use: Antibiotic & hormone therapy.

NEO-COSTILL OINT., (Dermatologicals, Inc) Hydrocortisone 1/2%, neomycin sulfate 2%. Tube 1/2 oz.
 Use: Skin infections.

NEO-CORTIN OPHTH. DROPS, (Klug) Polymyxin B sulfate 16,500 units, neomycin sulfate 5 mg., hydrocortisone 5 mg./cc, 5 cc. plastic applicator containers.
 Use: Topical antibiotic.

NEO-CREATINE, (Creatine) dl Methionine 200 mgm., glycine N.F. 100 mgm., arginine glutamate 20 mgm., Vit B-12 10 mcgm., Vit B-1 10 mgm., papain 10 mgm./Cap Bot 100s.
 Use: Special food supplement.

NEO-CULTOL, (Arlington) Viable susp. of lactobacillus acidophilus. Jar 6 oz.
 Use: Constipation.

NEOCURTASAL, (Winthrop Labs.) Pot. chloride, pot. glutamate, glutamic acid, cal. silicate & pot. iodide. Salt shaker 2 oz. Bot 8 oz. Unit service pkg., Rox 100s, 1000s
 Use: Salt (sodium)-free diets.

NEOCURTASAL IODIZED, (Winthrop Labs) Pot. chloride, ammonium chloride, pot. formate, cal. formate, mag. citrate, starch, pot. iodide 0.01%. Pkg. same as Neocurtasal.
 Use: Same as Neocurtasal.

NEO-CUTONE, (Sutliff & Case) Pyrilamine 1%, calomel 8%, benzocaine 3%. Lot. Bot. 2 oz., 1 pt., 1 gal.
 Use: Topically, antihistaminic w/local anesthetic.

NEOCYLATE, (Central) Sod. salicylate 4 gr., p-aminobenzoic acid 4 gr., ascorbic acid 0.3 gr./Tab. Sod. salicylate 5 gr., para-aminobenzoic acid (Sod. salt) 5 gr./dr. Syr. Bot. 1 pt., 1 gal. Entab., Bot. 200s, 500s, 1000s.
 Use: Analgesic in rheumatic conditions.

W/Cadaine (Central) Sod. salicylate 4 gr., p-aminobenzoic acid 4 gr., ascorbic acid 1/2 gr., codeine phosphate 1/4 gr./Tab Bot 100s, 500s
 Use: (1-2 Tab. q. 4-6 h.) Analgesic for arthritic conditions

W/Calcichine. (Central) Sod. salicylate 4 gr., calcichine 1/250 gr., p-aminobenzoic acid 4 gr., ascorbic acid 1/2 gr./Entab. Bot 200s, 500s, 1000s.
 Use: Gout & gouty arthritis.

NEOCYLATE-CORTISONE, (Central) Cortisone acetate 5 mg., Vit. C 20 mg., p-aminobenzoate Pot. 0.32 Gm., ammonium salicylate 250 mg./Tab., E.C. Bot 50s, 100s, 200s
 Use: 2-10 daily; antiarthritic.

NEOCYLATE-NC, (Central Pharm) Hydrocortisone 4 mg., potassium salicylate 0.28 Gm., potassium para-aminobenzoate 0.5 Gm., ascorbic acid 20 mg./Tab. Bot. 50s, 500s, 1000s.

NEOCYATE (Sodium-Free), (Central) Pot. salicylate 4 1/2 gr., pot. p-aminobenzoate 5 gr., & ascorbic acid 20 mg./Tab. Bot 100s
 Use: Rheumatic fever and arthritic conditions

NEOCYLONE, (Central) Prednisolone 2 mg., pot. salicylate 0.28 Gm., ascorbic acid 20 mg., pot. p-aminobenzoate 0.30 Gm./Tab. Bot. 50s, 100s, 500s.
 Use: Rheumatoid arthritis, acute rheumatic fever and acute gout.

NEOCYTIN, (Central) Pot. salicylate 4 1/2 gr., pot. p-aminobenzoate 5 gr., ascorbic acid 1/2 gr., physostigmine salicylate 1/250 gr., homatropine methylbromide 1/120 gr./Entab. E.C. tab. Bot. 100s, 500s, 1000s
 Use: Analgesic & antispasmodic.

NETO-DECADRON, (Merck Sharp & Dohme) Dexamethasone 21-phosphate w/Neomycin self. 5 mg
 Ophth. Sale 0.1% Dropper Bot. 5 cc
 Ophth. Oint. 0.05% Tube 3.5 Gm.
 Topical Cream 0.1% Tube 5 Gm., 15 Gm.

NEO-DELTA-CORTEF EYE-EAR DROPS, (Upjohn) Prednisolone acetate 2.5 mg., neomycin sulfate 5 mg./cc. Bot. (1/4%) 5 cc.
 Use: Treatment of infections of the eye or ear.

NEO-DELTA-CORTEF EYE-EAR OINTMENT, (Upjohn) Prednisolone acetate 2.5 mg. (1/4%), or 5 mg. (1/2%) neomycin sulfate 5 mg./Gm. Tube 1/2 oz.
 Use: Eye-ear infections.

NEO-DELTA-CORTEF LOTION, (Upjohn) Delta-1 hydrocortisone acetate 5 mg., neomycin sulfate 5 mg., methylparaben 2 mg., butyl-p-hydroxybenzoate 3 mg./cc. Plastic spray bot. T5 cc.

- Use Topically treatment of skin Inflammations
- NEO-DELTA-CORTEF NASAL SPRAY.** (Upjohn) Delta 1 hydrocortisone acetate 1 mg, neomycin sulfate 5 mg, phenylephrine HCl 2.5 mg/cc Plastic spray bot 15 cc
Use Treatment of nasal inflammations & bacterial infections
- NEO-DELTA-CORTEF OINTMENT TOPICAL.** (Upjohn) Prednisolone acetate 5 mg, neomycin sulfate 5 mg, methylparaben .02 mg, butyl p hydroxybenzoate 18 mg / Gm Tube 5 Gm, 20 Gm
Use Topically, allergic dermatitis & inflammatory skin diseases
- NEO-DELTEF DROPS** (Upjohn) Prednisolone 2 mg neomycin 5 mg & chlorbutolal 5 mg/cc (solution) Dropper Bot 2.5 cc
Use Ophthalmic use
- NEODOX** (Central) Pyridoxine HCl 100 mg thiamine 100 mg/cc Vial 10 cc
Use: I.M. & I.V.; (See B 1 & B 6 therapy)
- NEO-DILODEBM,** (Schering) Diladerm plus 37.5 mg neomycin sulfate 37.5 mg / 10 Gm foam Diladerm plus B 3 mg neomycin sulfate 16.7 mg / 50 Gm aerosol spray Diladerm plus 2.5 mg neomycin sulfate, 50 mg / Gm / 5 Gm cream Aerosol foam 10 Gm can Aerosol 50 Om can Cream, tube 5 Gm, 15 Gm
Use Dermatoses when secondary infection threatens
- NEO-DOMEFORM-HC** (Dome) Neomycin sulfate 5 mg / Gm, micro dispersed hydrocortisone alcohol 1/2%, iodochlorhydroxy quin 3% in Acid Mantle vehicle / Tube 1/2 oz Jar 1 lb
Use Dermatological conditions
- NEODENAL** See Isoproterenol
- NEODOL, NND 1962** (Fizer) Withdrawn
Stanolone Vial (50 mg/cc) 10 cc multiple dose
Use: I.M.; antitumor & anabolic therapy
- NEODROPS** (Walker) Vit A 10 000 u, D 2000 u, B 1 3 mg, B 2 0.8 mg, B 6 1 mg, panthenol 3 mg, niacinamide 20 mg, ascorbic acid 75 mg, folic acid 50 mcg / cc Bot w/dropper 15 & 30 cc.
Use: (5 to 15 drops daily) Dietary supplement
- NEODYNE,** (National Drug) Terpin hydrate 1/6 gr, guaiaacal, glycerophosphates ammonium chloride, tartar emetic, creosote carb & hyoscyamus tr / 5 cc Bot 4 oz, pt., gal
Use: Orally (1 2 dr q 3 h), cough preparation
- NEO-EFEMIST,** (Hart) Ephedrine sulfate 1% mild silver protein 5% Bot 1/2 oz
Use: Antiseptic & vasoconstrictor
- NEO-FEMOTRONE.** (Testagar) Estradiol 0.5 mg, progesterone 12.5 mg / cc Vial 10 cc
Use: I.M., hormone therapy.
- NEO-FERINEX PLUS,** (Plessner) Ferrrous sulf 150 mg, liver desic. 100 mg, stomach substance 100 mg, folic acid 0.5 mg, Vit C 50 mg, B 12 with intrinsic factor conc 0.5 u, B 12 2 mcg, oyster shell calcium 3 gr / Tab Bot 50s, 100s, 500s
Use: Hematinic
- NEOFERNISONE CREAM.** (Ferndale) Hydrocortisone acetate 10 mg, neomycin sulfate 5 mg / Gm 1/2 oz tubes
Use Dermatitis
- NEOFERBUM.** (Warner Chilcott) Iron 17 gr, Manganese 1/2 gr, Vit B 1 1.5 mg, w/trace of copper / fl oz Bot 12 fl oz, 3 gal
Use (I tsp t i d, p c) Hematinic
- NEO-FLO SOL** (Professional) Boric acid sodium chloride potassium chloride and sodium carbonate preserved with benzalkonium chloride USP 115 000 120 cc plastic bellows type dispensing bottle
Use Ophthalmic irrigant
- NEOFONAMIDE,** (McNeil) Sulfacelamide 5%, urea 5%, neocollamine 15%, talc 15% glycerin camphor, menthol Bot 4 R oz, 1 pt
Use Topically; bacteriostatic
- NEOFORTIFIED.** (Dumas Wilson) Primary yeast 500 mg, B 1 1 mg, B 2 1 2 mg, nicotinamide 10 mg B 6 0.5 mg, calcium ascorbate 36 mg, iron 10 mg, sodium 0.9 mg / Tab Bot 100s
Use: Dietary supplement
- NEO-GEL** (Diamond) Magnesium trisilicate and neigel Susp Bot 12 oz Tab bot 60s
Use: Antacid
- NEOGEN SUPPOSITORIES.** (Premo) Pentobarbital 1 gr, aminophylline 4 gr, theophylline 3 gr, ephedrine sulfate 3/8 gr benzocaine 3 gr / Supp Box 12s Also available in half strength
Use Branchodilator, mild sedative
- NEOGRAY** (Mason Pharmaceuticals) Vit A 3000 u Vit D 400 u, B 1 2 mg, B 2 1 5 mg, niacinamide 10 mg, B 6 5 mg, C 25 mg, calcium pantothenate 1 mg, purified bone meal 195 mg, ferrous sulfate 120 mg, mixed tocopherols 3 mg menadiol 0.05 mg, Vit B 12 0.5 mg / Tab Bot 100s
Use: Prenatal vitamin mineral supplement
- NEO-GYNERGEN** (Sandoz) Not listed Ergotamine tartrate 0.25 mg, ergonovine maleate 0.125 mg / Tab or cc
Use: Oxytocic
- NEO-NABVOGLOBEIN** (G F Harvey) Vit B 1 0.5 mg, B 2 1 mg, Iron hypophosphite

- 8 mg., maltose, proteins, amino acids, beef, molli & yeast ext./dr. Bot. 1 pt., 1 gal.
Use: Dietary supplement.
- NEOHEMOCHROMIN.** (Carnrick) Vit. B-1 2 mg., B-2 3 mg., inositol 20 mg., pantothenic acid 1 mg., folic acid 2 mg., ferrous gluconate 5 gr., liver fraction no. 2 195 mg./Cap. Bot. 100s, 500s.
Use: Hematinic.
- NEO-HEMOPLASTIN.** (Porke, Davis) Prothrombin, thrombokinase & antithrombin. Vial 2 cc., 5 cc.
Use: Orally, I.M. & S.C.; hemostatic.
- NEOHETRAMINE, HCl, N.N.D.** 1962. (Napso) Thanzylamine HCl. Tab. (25 mg) Bot. 12s, 100s, 1000s; (50 & 100 mg) Bot. 100s, 1000s. Syr. (6.25 mg./cc.) Bot. 1 pt., 1 gal. Cream (2%) Tube 1 oz.
Uses (50 mg. to 100 mg. q.i.d.) Anti-histaminic.
- NEO-HOMBREOL, N.N.D.** 1962. (Organon) Testosterone propionate in sesame oil. Amp. (10 mg./cc., 25 mg./cc.) 1 cc. Bot. 6s, 30s. Vial (10 mg./cc., 30 mg./cc.) 10 cc., 6s; (100 mg./cc.) 5 cc., 6s.
Use: Male sex hormone.
- NEO-HOMBREOL (F), N.N.D.** 1962. (Organon) Testosterone aq. susp. Vial (25 & 50 mg./cc.) 10 cc., 50 cc.
Use: I.M. (1-2 cc weekly); endocrine therapy.
- NEO-HOMBRIOL (M), N.N.D.** 1962. (Organon) 17-Methyltestosterone Tab. 10 mg., Bot. 30s, 100s, 500s; 25 mg., Bot. 15s, 100s. Sublingual 5 mg., 10 mg., Bot. 30s, 100s.
Use: Androgen therapy.
- NEO-HYDCORTOLE OPHTHALMIC OINTMENT.** (Premo) Hydrocortisone acetate 15 mg., neomycin sulfate 5 mg., in a bland white petrolatum base. 4 gram collapsible metal tubes w/special ophthalmic applicator tip.
- NEO-HYDELTRASOL.** (Merck Sharp & Dohme) Prednisolone 21-phosphate sod. 5.0 mg., neomycin sulf. 5.0 mg./cc. Ophth. Sol. Vial 5 cc., 2.5 cc., Topical Sol. Plastic Squeeze 15 cc. Ophth. Oint. 2.5 mg., and 5 mg., respectively. Tube 3.5 Gm. Topical Oint. 5.0 mg. & 5.0 mg., respectively. Tube 5 Gm., 15 Gm.
Use: Inflammation and infection
- NEO-HYDELTRASOL NASAL SPRAY.** (Merck Sharp & Dohme) Prednisolone 21-phosphate (as sod. salt) 1.0 mg., phenylephrine HCl 2.5 mg., propadrine HCl 7.5 mg., neomycin sulf. 5.0 mg./cc. Bot. 15 cc.
Use: Allergic disorders of the nose.
- NEOHYDRIN, N.N.D.** 1962. (Lakeside) Chlormerodrin, 3-chloromercuri-2-methoxypropylurea. Tab. 183 mg. Bot. 50s, 500s.
Use: Diuretic.
- NEO-POPAX, N.N.R.** 1954. (Schering) Sod iodamethamate. Amp. 30%, 10 cc., 20 cc., 30 cc.; 75%, 10 cc., 20 cc., 50 cc.
Uses I.V. & retrograde pyelography & angiocardiology.
- NEOJIL W/Magnesium trisilicate.** See: Neo-Gel, Susp. & Tab. (Diamond)
- NEO-LACMANESE.** (Braun) Milk protein, Mang. gluconate, procaine HCl. Vial 1 cc., 30 cc.
Use: I.M. (1 cc. daily), nonspecific protein therapy.
- NEOLAX.** (Central) Diactyl sodium sulfosuccinate 50 mg., dehydrocholic acid 0.25 Gm./Tab. Bot. 30s, 100s, 500s
Use: Choleric laxative and fecal softener.
- NEO-LIDA-MANTLE-HC.** (Dome) Lidocaine 30 mg., hydrocortisone alcohol 5 mg., neomycin sulfate 5 mg., in Acid Mantle base/Gm. Tube ½ oz., 1 oz.
Use: Dermatoses.
- NEOLIN SUSPENSION, N.N.D.** 1962. (Lilly) Benzathine penicillin G, N,N'-dihydroxyethyl-enediamine penicillin G 300,000 u./5 cc. Bot. 60 cc.
Use: Orally; penicillin therapy.
- NEOLOID.** (Lederle) Castor oil, U.S.P. 36.4% (w/w) Bot. 4 oz.
Use: Orally; laxative.
- NEO-MAGNACORT.** (Pfizer) Hydrocortisone diethylaminoacetate 0.5%, neomycin 0.5%. Oint. Tube 1/6 oz., ½ oz.
Use: Topically; skin inflammations
- NEOMALENE CAPSULE.** (Smith-Dorsey) Codeine phosphate ¼ gr., pyrilamine maleate 25 mg./Cap. Bot. 100s, 500s.
Use: (1 I.I.d.) Antihistaminic w/analgesic.
- NEO-MANTLE.** (Dome) Neomycin sulfate 50 mg./Gm in "Acid Mantle" vehicle Creme. Tubes ½ oz. Jar 4 oz. Bot. 2 oz., 4 oz.
Use: Treatment of cutaneous pyodermitis, impetigo contagiosa and other pustular dermatoses.
- NEO-MATURE.** (Ayerst) Ferrous fumarate 200 mg., ascorbic acid 100 mg., folic acid 1.67 mg., Vit. B 12 10 mcg., desiccated stomach tissue 1 Gm., liver extract equivalent to 1 Gm beef liver/Cop.
Use: Treatment of anemias
- NEO-MEDROL.** (Upjohn) Methylprednisolone 1 mg., neomycin sulfate 5 mg., chlorbutanol 5 mg./5 cc. Soln. Dropper Bot. 5 ml Methylprednisolone 1 mg., neomycin sulfate 5 mg./Gm. Oint. Tube ½ oz.
Use: Treatment of eye and ear infections.
- NEO-MEDROL ACETATE, VERIDERM.** (Upjohn) Methylprednisolone acetate 2.5

- mg or 10 mg neomycin sulfate 5 mg methylparaben 4 mg butyl p hydroxybenzoate 3 mg /Gm Tube 5 Gm 20 Gm
Use Treatment of skin lesions
- NEO-MENSALIN.** (Walker) Bromoate 2 Amino 2 methyl 1 propanol 8 bromo theophyllinate (pamobrom) 50 mg pyrrolamine maleate 30 mg /Tab Bat 30s 100s 1000s Use (111d) Premenstrual tension
- NEOMYCIN BASE W/Gramicidin** See: Spectrocin Oint & Troche (Squibb)
- W/Bacitracin, neomycin sulfate & benzocaine** See: Tetraxets, Troche (Merck Sharp & Dahme)
- W/Bacitracin & polymyxin B sulfate** See Neo Polycin, Oint & Ophth Oint (Pitman Moore)
- W/Colistin base hydrocortisone acetate thonzonium bromide polysorbate 80 acetic acid sodium acetate** See Coly Mycin Otic
- W/Neomycin and Hydrocortisone** (Warner Chilcott)
- W/Dyclonine hydrochloride polymyxin B sulfate** See: Neo Polycin Otic drops (Pitman Moore)
- W/Estrane & Vit A** See: Es A Mycin Creme (Dome)
- W/Hydrocortisone acetate & gramicidin** See Florinel S Lotion & Oint (Squibb)
- W/Gramicidin** See: Spectrocin Oint & Troche (Squibb)
- W/Gramicidin** See Bio Cuto, Oint (Columbus)
- W/Gramicidin nystatin triamcinolone acetate** See Mycolog Cream (Squibb)
- W/Hydrocortisone** See Neo Cort Dame Cream (Dome)
- W/Hydrocortisone acetate** See: Cortomycin Oint (Bryant)
- W/Hydrocortisone diethylaminoacetate** See: Neo Magnacort Oint (Pfizer)
- W/Hydrocortisone & bacitracin** See Hydroderm Oint (Merck Sharp & Dahme)
- W/Hydrocortisone diethylaminoacetate HCl, undecylenic acid in a petroleum base** See: Podiacort Oint (PodiatRex Ltd)
- W/Hydrocortisone polymyxin B sulfate piroxime hydrochloride, propylene glycol glycerin** See: Auracort, Drops (Columbus)
- W/Hydrocortisone, polymyxin B sulf, chlorpheniramine maleate, phenylephrine HCl** See: Polycort, Sol (Lemman)
- W/Kaolin pectin salol, zinc sulfocarbolate** See: Neo Kotelan (Buffngton)
- W/Kaolin, pectin hyascyamine sulf, atropine sulf, hyoscine Hbr, phenobarbital** See Dannagal with Neomycin (Robins)
- W/Phenylephrine HCl, chlorphenamine, polymyxin B sulfate, benzocaine** See Biotic Loz (Lemman Pharm)
- W/Phthalylsulfathiazole** See: Neotholidine Tab (Merck Sharp & Dahme)
- W/Polycarbophil thiexinol methyrbromide** See Sorbaquel, Tabs (White)
- W/Polymyxin B sulfate heparin Sod hydrocortisone** See Bro Porin, Susp (Braemmel Pharm)
- W/Polymyxin B sulfate benzocaine, cetylpyridinium Cl** See Biotane Lozenges (Philips Roxane)
- W/Polymyxin B sulfate, heparin sod, hydrocortisone chlorbutanol in polyethylene glycol 400 base** See Bro Porin H, Vial (Braemmel)
- W/Prednisolone, sod propionate** See Otobion (White)
- W/Sulfasuxidine pectin & kaolin** See: Cremamycin (Merck Sharp & Dahme)
- W/Triamcinolone acetonide gramicidin** See Kenalog S Cream Oint (Squibb)
- W/Triamcinolone, gramicidin, nystatin** See Nycolog Cream (Squibb)
- W/Tyrosine benzocaine** See Antibiotic Throat Lozenges (Slayner)
- W/Tyrosine, prophenpyridamine maleate, benzocaine** See: Neo T Cain, Lozenget (Chicago Pharmacol)
- NEOMYCIN LOTION.** See Mycimist (Upjohn)
- NEOMYCIN-MYCOSTATIN** (Squibb) Neomycin sulf. 0.5 Gm, mycostatin 125,000 u /Tab Bat 20s, 100s
Use: To suppress enteric bacterial & fungal organisms in gastro intestinal surgery
- NEOMYCIN PALMITATE, W/Hydrocortisone mole HCl proteolytic enzymes** See Chymot (Armour)
- *NEOMYCIN SULFATE, USP NND 1962**
An antibiotic from *Streptomyces fradiae*; Oint, USP's Tab, USP (Upjohn) Paw Vial 0.5 Gm Tab 0.5 Gm Bat 20s
See Lidamycin Creme Tubes (Dome)
Mycifradin, Vial & Tab NNR (Upjohn)
Myciguent, Oint, & Ophth Oint (Upjohn)
Neobiotic, Tab (Pfizer)
Neo Mantle Creme & Lot (Dome)
Quintess (Lilly)
White's Otobiotic, Sol (White)
- W/Arnoldin, Vit A & D** See VAD, Cream (Walker)
- W/Bacitracin**
See Bacimycin, Oint, & Ophth Oint (Walker)
Bacitracin Neomycin Ophth Oint (Paul Money)
Mycitracin, Oint (Upjohn)

* Generic name recognized by N.P., U.S.P. or A.M.A.

- Neobactin, Oint. & Tab. (Commercial Solvents)
- Neotracin, Oint. (Pramo)
- W/Bacitracin & hydrocortisone.
- See: Bacimycin W/Hydrocortisone, Oint. (Walker)
- Hydroderm, Oint. (Merck Sharp & Dohme)
- W/Bacitracin & polymyxin B sulfate.
- See: Neomycin, Oint. (Burroughs Wellcome)
- Neo-Polycin Ointment. (Pitman-Moore)
- Neo-Polycin Ophth. Oint. (Pitman-Moore)
- Neosporin, Oint. & Ophth. Oint. (Burroughs Wellcome)
- Spectra Biotic, Oint. (American Pharmaceutical Co.)
- Trimizal, Oint. (Hance)
- W/Bacitracin zinc salt & polymyxin-B sulfate.
- See: Wyblotic, Troche. (Wyeth)
- W/Bacitracin zinc, polymyxin B, benzalkonium chloride. See: Biotres, Oint. (Control)
- W/Bacitracin, polymyxin B sulf., hydrocortisone ocelate. See: Neopolycin-HC, ophth. oint. (Pitman-Moore)
- W/Bacitracin & propyl p-aminobenzoate.
- See: Du-Biotic, Troche. (White)
- W/Benzocaine. See: Americaine-Neomycin, Oint. (Amar-Stone)
- W/Cortisone ocelate. See: Neosone, Oint. (Upjohn)
- W/Dexamethasone 21-phosphate. See: Neo-Decadron, Prods. (Merck Sharp & Dohme)
- W/Diloderm. See: Neo-Diloderm (Schering)
- W/Diperodon HCl, allantoin See: Ro-Derm (Lemmon)
- W/Dyclanin HCl, polymyxin B sulf., See: Neo-Polycin Otic (Pitman-Moore)
- W/Erythromycin. See: Erythracin-Neomycin Ointment (Abbott)
- W/Erythramycin, methylparaben & butyl p-hydroxybenzoate. See: Erythro-myciguent, Oint. (Upjohn)
- W/Estrone & Vit. A. See: Es-A-Mycin Cream. (Dome)
- W/Fludrocortisone, gramicidin, mycostatin.
- See: Mycanol Ointment. (Squibb)
- W/Fludrocortisone ocelate, polymyxin B sulf., nystatin (Mycostatin). See: Piarotic, plastic dropper. (Squibb)
- W/Gramicidin & polymyxin B sulfate. See: Neosporin, Sol. (Burroughs Wellcome)
- W/Gramicidin, polymyxin B sulf., benzocaine. See: Tricidin, Oint. (Boyle)
- W/Gramicidin & propyl p-aminobenzoate.
- See: Orabiotic, Gum Tab. (White)
- W/Gramicidin, thonzonium bromide, thonzylamine HCl. See: Biomydrin Otic. (Nepero)
- W/Gramicidin, thonzylamine HCl, phenylephrine HCl & thonzonium bromide. See: Biomydrin, Sol. (Nepero)
- W/Homarylamine HCl, zinc bacitracin, thiaziclin, benzocaine. See: Pentozels traches, Vial. (Merck Sharp & Dohme)
- W/Hydrocortamote.
- See: Neo-Ulcort, Oint. (Ulmer)
- Via-Cort, Cream (Rowell)
- W/Hydrocortisone.
- See: Heb-Cort N 1/4 % Lotion (Barnes-Hind)
- Protal, Supp. (Upjohn)
- W/Hydrocortisone ocelate. See: Hydrocortisone-Neomycin, Inj. (Philadelphia Ampoule)
- W/Hydrocortisone ocelate. See: Hydrocortin, Liq. (Philadelphia Ampoule)
- W/Delta-1-Hydrocortisone ocelate & phenylephrine HCl. See: Neo-Delta-Cortef Nasol Spray. (Upjohn)
- W/Hydrocortisone, gramicidin, thonzylamine HCl, phenylephrine HCl, thonzonium bromide. See: Biomydrin P with Hydrocortisone. (Nepero)
- W/Hydrocortisone. See: Neo-Torcortin, Oint. (Reed & Carrick)
- W/Hydrocortisone alcohol, tetracycline hydroxyquin. See: Naflo-Cort, Oint. (Rowell)
- W/Hydrocortisone ocelate, and polymyxin B sulfate. See: Icapia-P-H-N, Ophth. Sol. (Alcon)
- W/Hydrocortisone, gramicidin, phenylephrine HCl. See: Du-Biotic Intranasol. (White)
- W/Hydrocortisone, gramicidin, polymyxin-B sulfate, phenylephrine & prednisone HBr See: Triacort, Sproypak. (Smith, Kline & French)
- W/Hydrocortisone ocelate, prednisone, prednisolone. See: Pansteroid, Oint. (Kenwood)
- W/Hydrocortisone & propadrine HCl. See: Hydro Spray, Nasol Suspension. (Merck Sharp & Dohme)
- W/Hydrocortisone, pyrilamine maleate, catalkonium chloride, diperodon HCl, urea See: Ototrin Ear Drops (Louisiana)
- Tega-Otic, Soln. (Ortega)
- W/Hydrocortisone, pyrilamine maleate, catalkonium chloride, diperodon HCl. See: Pant-O Angina Ear Drops, Liq. (Confield)
- W/Hydrocortisone, pyrilamine maleate, catalkonium chloride, diperodon HCl, urea. See: Tyler Otic, Sol. (Tyler)
- W/Hydrocortisone, tyrothricin, bacitracin, benzocaine See: Tyroozels, Troche. (Merck Sharp & Dohme)
- W/Hydrocortisone ocelate.
- See: Hydroptic, Ophth. Susp. (Merck Sharp & Dohme)
- Neo-Cortef, Oint. & Drop. (Upjohn)
- W/Hydrocortisone & bacitracin. See: Hydroderm, Top. Oint. (Merck Sharp & Dohme)
- W/Koolin, pecin
- See: Neoplate, Syr. (Tyler)
- Pecta-Kolin with Neomycin, Liq. (Money)
- Vahmycin, Tab. (Moore)

NEOMYCIN-TRIAMCINOLONE ACETONIDE. See: Neo-Aristocort Eye-Ear (Lederle)

NEOMYCORSONE. See: Neosone, alet. (Upjohn)

NEONAL. (Abbott) Butethal 100 mg /tab. Bot. 100s
Use: Hypnotic.

NEO-NORMADRINE. (Vanpelt & Brown) Phenylephrine HCl ¼%, pyrilamine maleate ¼%, w/cetyl dimethyl benzyl ammonium chloride in buffered solution. Bot 1 & 4 oz.
Use: Antihistaminic w/vasoconstrictor.

NEO-NYSTA-CORT. (Dome) Nystatin 100,000 U., hydrocortisone 10 mg., neomycin sulfate 5 mg /Gm. Tube ½ oz.
Use: Moniliasis.

NEO-OXYLONE. (Upjohn) 0.25 mg. Fluorometholone & 5 mg. neomycin sulfate/Gm. Oint. Tube 7.5 Gm.
Use: Contact dermatitis, atopic dermatitis, neurodermatitis, anogenital pruritus, & seborrheic dermatitis.

NEOPAN. (U S Vitamin) Neomycin sulfate 5 mg., pantolhenylol 20 mg., water-miscible cream base/Gm. Cream. Jars. 2 oz., 1 lb.
Use: Dermatoses.

NEOPARSEL. (Central) Pamobrom 50 mg., pyrilamine maleate 30 mg., homatropine methylbromide 1.2 mg., hyoscynamine sulf. 0.1 mg., scopolamine HBr 0.02 mg., methamphetamine HCl 1.5 mg./Tab Bot. 30s, 100s, 500s
Use: Treatment of premenstrual tension

NEO-PARROROM. (Central) Bromateate. Pamobrom 50 mg., pyrilamine maleate 30 mg /Tab. Bot. 100s, 500s
Use (1-3 daily) Fremenstrual tension.

NEOPAS. (Kirkman) p-Aminosalicylic acid 1 Gm./tab. Bot. 100s.
Use: Tuberculosis.

NEOPASALATE. (Dumas-Wilson) p-Aminosalicylate sod. and p-aminosalicylic acid. Tab. (equiv. to Sod. PAS 7½ gr.) Bot. 100s, 1000s
Use: Treatment of tuberculosis

NEOPAVRIN-A. (Savage) Diethoxydiethoxybenzyl isoquinoline hydrochloride (ethaverine) 30 mg., d-amphetamine 3 mg /Tab. Bot. 100s, 1000s.
Use: Treatment of smooth muscle spasms.

NEOPAVRIN TABS. & ELIXIR W/Phenobarbital. (Savage) Phenobarbital ½ gr., diethoxy (diethoxybenzyl) isoquinoline HCl 30 mg./Tab. & 5 cc. elixir. Bot pts Tabs 100s, 1000s.
Use: Smooth muscle spasms.

NEOPECTATE. (Tyler) Neomycin sulfate 300 mg., kaolin 90 gr., pectin 2 gr./oz.
Use: Treatment of diarrhea

NEOPENIL. See: Benzylpenicillin 2-diethylaminoethyl ester HI

NEO-PENT. (Paul Money) Pentylene tetrazole 100 mg., nicotinic acid 50 mg., neothylline 100 mg /Tab Bot. 100s, 1000s
Use: Treatment of fatigue and senility

NEOPHRYN. See: Phenylephrine

NEOPINZINE. (Lilly) Cryst. Fat. penicillin G 150,000 or 300,000 u., sulfadiazine 0.167 Gm. Tab., Bot 24s, 100s, 1000s. Cryst. Pot. penicillin G 200,000 u., sulfadiazine 0.167 Gm., sulfamerazine 0.167 Gm., sulfa methazine 0.167 Gm. Buffered w/Sod. citrate 5 cc. Susp. 60 cc.
Use: Penicillin or sulfonamide-susceptible organisms.

NEOPLASTIC AGENTS. See: Folic acid antagonists. Leukemia agents.

NEO-POLYICIN OINTMENT. (Fitman-Moore) Neomycin 3 mg., bacitracin 400 u., polymyxin B sulfate 8000 u./Gm. Oint. In special base of Fuzene. Tube 15 Gm. Ophth. Oint. (Neomycin base 3 mg, bacitracin 500 u., polymyxin B sulfate 10,000 u./Gm.) Tube ½ oz.
Use: All types superficial skin infections.

NEO-POLYICIN OPHTHALMIC OINTMENT. (Fitman-Moore) Neomycin 3 mg., bacitracin 500 u., polymyxin B sulfate 10,000 u./Gm Oint. In enhy. lanolin-petrolatum base. Tube ½ oz.
Use: Wide variety of pathogens.

NEO-POLYICIN OPHTHALMIC SOLUTION. (Fitman-Moore) Neomycin sulfate 2.85 mg., polymyxin B sulfate 5000 u., gramicidin 0.02 mg in a sterile isotonic solution. Bot w/sterile dropper 10 cc.
Use: Treatment of ophthalmic infections.

NEO-POLYICIN OTIC. (Fitman Moore) Dyclonine hydrochloride 1% (4-n-butoxy-beta-piperidieopropiophenone hydrochloride), neomycin 2 mg, polymyxin B sulfate 4,000 u. in fuzene base/Gm. Bot 8 Gm.
Use: Treatment of pain, pruritus, or infection in otitis externa

NEO-POLYICIN-HC OTIC. (Fitman-Moore) Neomycin sulfate 2.85 mg., polymyxin B sulfate 4000 u., dyclonine hydrochloride 10 mg (1%) hydrocortisone acetate 10 mg. (1%) /Gm. Bot. 5 Gm.
Use: Otitis externa and other painful or pruritic otic conditions.

NEO-POLYICIN-HC OINTMENT. (Fitman-Moore) Neomycin sulf. 3 mg., bacitracin 400 u., polymyxin B sulf. 8000 u., hydrocortisone acetate 10 mg in fuzene/Gm. Tube 5 Gm.
Use: Topically. inflammatory skin infections

NEO-POLYICIN-HC OPHTHALMIC OINTMENT (Pitman Moore) Neomycin sulf 3 mg bacitracin 500 u, polymyxin B sulf 10 000 u, hydrocarbons acetate 10 mg / Gm Tube 1/2 oz w/applicator tip
Use Inflammatory eye infections

NEOPRONTOSIL (Winthrop Labs) Aza sulfamide Amp 2.5% 5 cc, Bal 50 cc Bal 5% 50 cc Tab 5 gr, Bot 50s 500s
Use Streptococcus infections

NEO-PROPISOL (Ayerst) Sad propionate 5% and Naphazoline HCl 0.025% in buffered saline Bot drapper 15 cc
Use Antifungal agent for eyes

NEO-PYRAPAR (Ulmer) Bromalate Pama brom 50 mg pyrilamine maleate 30 mg / Tab Bot 100s, 500s
Use (2 b i d) Premenstrual tension

NEOQUINOPHAN See Neocinchophen (Yarnus Mir)

NEO-RESULIN-F (Schiefel) Resorcinol monoacetate 1.5%, sulfur 2%, hydrocortisone 0.5%, neomycin sulf 0.5%/cream Tube 15 Gm
Use Acne therapy

NEO-RHIBAN (Kendall) Glyceril guaracolate 5 mg, phenylpropanolamine HCl 12.5 mg, methopyrilene HCl 25 mg, n acetyl p aminophenol 195 mg / Tab Bot 100s, 500s
Use Common colds

NEO-RIBOTHYN (Flint Eaton) Thiamine HCl 50 mg, riboflavin 25 mg, pyridoxine HCl 25 mg, Col pantothenate 50 mg nicotinamide 500 mg ascorbic acid 500 mg / Vial Vial 5 cc w/diluent
Use I M & I V (1 or 2 cc daily) Vit 8 complex deficiencies

NEO-RIODINE (Gallia) Sad iodopropano sulfonate 0.1 Gm Amp 5 cc, Box 10s, 50s
Use (1 2 cc daily) Iodide therapy

NEORON-C TIMEKAPS (Caldwell & Bloor) Ferrous fumarate 250 mg, Vit C 60 mg, B 1 5 mg / Cap Bot 100s 1000s
Use Iron deficiency anemias

NEO-SALICIPHEN (Abbott) Aspirin 3 gr neocinchophen 3 gr, calcichime 1/250 gr / Cap Bot 100s 1000s
Use (1 q i d) Analgesic & antirheumatic

NEOSALVARSAN, N N R 1949 See Nearsphenamine (Abbott)

NEOSED (Diamond) Acetylcarbromal 5 gr, Neojel 2 gr calcium carbonate 2 gr / Tab Bot 100s
Use Tension headaches

NEO-SEDAPNEN (C D Smith) Fenobarbital 30 mg, phenobarbital 10 mg
Sad bromide 300 mg, Pot bromide 200

mg, Col bromide 100 mg / 5 cc Bot 4 oz, 1 pt, 1 gal
Uses (1 dr. 1 i d) Sedative

NEO-SEMNYTEN (Massengill) Reserpine 0.1 mg, mannitol hexanitrate 30 mg, theophylline 100 mg, rutin 10 mg, Vit C 15 mg / Cap Bot 100s, 500s, 1000s
Use Hypertension

NEO-SILVOL, N N R 1950 (Parke, Davis) Silver iodide w/soluble gelatin base, containing 18 to 22% silver iodide in colloidal form Cap 0.389 Gm, Bot 50s Sal Bot 1 oz 4 oz Vaginal Supp 5%, Box 12s
Use Genito urinary tract infections

NEO-SKIODAN. See Iodopyracet (Diadrost, Winthrop Labs)

NEO-SLOWTEN (Smith, Miller & Patch) Phenobarbital 16.2 mg, reserpine 0.1 mg, Thiamine HCl 5 mg / Tab Bot 100s, 1000s
Use Treatment of worry apprehension, irritability emotional instability, subjective fatigue and mild degrees of hypertension

NEOSONE (Upjohn) Neomycosone Cortisone acetate 15 mg neomycin sulfate 5 mg / Gm Ophth Oint, Tube 1 dr w/ applicator tip
Use Conjunctivitis keratitis & various trauma

NEOSORB (Hoock) Aluminum hydroxide gel 4 gr, Mag trisilicate 7 gr, methylcellulose 1 gr / Tab or 1/2 dr Bot 1 pt 1 gal Tab 100s, 500s, 1000s
Use (2 q 2 4 h) Antacid & adsorbent

NEOSPORIN ANTIBIOTIC OINTMENTS (Burroughs Wellcome) Polymyxin B sulfate 5000 u, bacitracin 400 u neomycin sulfate 5 mg / Gm Oint, Tube 1/2 oz w/appl color & 1 oz Ophth Oint, Tube 1/2 oz
Use Bacterial infections

NEOSPORIN AEROSOL (Burroughs Wellcome) Aerasparin brand polymyxin B sulfate 100 000 u, neomycin sulfate 100 mg, zinc bacitracin 8000 u dispersed in inert propellant / Containers (aerosol spray) 90 Gm Approx spraying time 100 seconds
Use Bacterial skin infections

NEOSPORIN ANTIBIOTIC LOTION (Burroughs Wellcome) Polymyxin B sulfate 10 000 u neomycin sulfate 5 mg / cc Squeeze bot 20 cc
Use Topical bacterial infections

NEOSPORIN OPHTHALMIC SOLUTION (Burroughs Wellcome) Polymyxin B sulfate 5000 u, neomycin sulfate 2.5 mg, gramicidin 0.025 mg / cc Bot 10 cc
Use Topically eye infections

NEOSPORIN TOPICAL POWDER (Burroughs Wellcome) Polymyxin B sulf 5000 u, bac

- tracin zinc 400 u., neomycin 5 mg/Gm.
Bot. shaker-top 10 Gm.
Use: Topical bacterial infections
- NEOSPORIN VAGINAL SUPPOSITORIES.**
(Burroughs Wellcome) Polymyxin B sulfate
20,000 u., neomycin sulfate 5 mg, acetar-
sone 250 mg., beta-lactose qs. 1 Gm./
Supp. Box 12s.
Use: Vaginitis, cervicitis, trichomoniasis.
- NEOSTIBOSAN.** See: Ethylstibamine (Fuadon,
Amp. Winthrop Labs)
- NEOSTIGMETIN.** (New Mexico Pharm.) Neo-
stigmine methylsulfate 1:1000 10 mg.
propylparaben 0.02%, methylparabee
0.18%/cc. Vials 10 cc.
See Kirkstigmine, Amp. & Vial (Kirk).
- NEOSTIGMINE, N.N.R. 1950.** (International
Medical) Prostigmin. Amp. (1:2000) 1 cc.,
Box 100s, Vial (1:1000) 10 cc.
See Kirkstigmine, Amp. & Vial (Kirk).
- *NEOSTIGMINE BROMIDE, U.S.P., N.N.D.**
1958, Tab, U.S.P. See: Prostigmin Bromide
(Roche)
- *NEOSTIGMINE METHYLSULFATE, U.S.P.,**
N.N.D. 1958; Int., U.S.P.
See: Kirkstigmine, Vial (Kirk)
Neostigmetin (New Mexico Pharm.)
Prostigmin methylsulfate (Roche Lab)
Use: Parasympathomimetic agent,
W/Atropine sulfate & vit. B-1. See: Thio-
silgmine, Vial, (Kirk)
- W/Atropine sulfate & Vit. B-12. See
Necobin, Vial (Miller).
- W/B-1. See: Thio-silgmine, Vial (Kirk)
- NEO-STREPSAN.** See: Sulfathiazole (Various
Mfr.)
- NEOSUM ANTIBIOTIC LOT.** (Summers) Neo-
mycin undecylenate 11%, polysorbate 80,
1.25%, ocalone 14%, Isoproponol 50%,
in water. 2 fl. oz. bals.
Use: Dermatoses.
- NEO-SUPERTAN-5.** (Tailby-Nasae) Hydro-
cortisone alcohol 1%, w/Supertan equiv.
to 5% crude coal tar, methylparabee 0.1%,
propylparabee 0.05%. Jar ½ oz.
Use: Topically, treatment of dermatitis
- NEO-SYNALAR.** (Syntex) Fluocinonide ace-
tonide 0.025%, neomycin sulfate 0.5%, in
a water washable base/Gm. Tube 5 Gm-
15 Gm.
Use: Inflammatory dermatoses
- NEO-SYNEPHRINE COMPOUND COLD**
TABLETS. (Winthrop Labs) Neo-synephrine
HCl 5 mg, N-Acetyl-p-aminophenol 150
mg, thenyldiamine HCl 7.5 mg, coffeee
15 mg./Tab. Bot. 100s
Use: Treatment and management of the
common cold
- NEO-SYNEPHRINE HYDROCHLORIDE,**
N.N.R. 1954, A.D.R. (Winthrop Labs)
- * Generic name recognized by N.F., U.S.P. or A.M.A.
- Phenylephrine HCl. Amp. 0.2%, 2 cc., 10s.
Cap. 10 & 25 mg, Bot. 100s & 500s. Elixir
0.1%, Bot. 1 pt., 1 gal. Jelly ½%, Tube
¼ oz. Ophth. Emulsion ¼%, Bot. 1 oz.,
1 pt. Pediatric Nozal Spray ¼%, Squeeze
Bot. 20 cc. Sol. ¼%, Bot. 20 cc., 1 & 4 oz.,
1 pt., ½%, Bot. 20 cc., 1 oz.; 1%, Bot.
1 & 4 oz., 1 pt.
- W/Aromatic ½%, Bot. 15 cc.; ¼%, Bot.
1 oz., 1 pt.; 2½%, Bot. 15 cc.; 10%, Bot.
15 cc.
- W/Codeine phosphate, acetylsalicylic acid,
chlorpheniramine maleate, ascorbic acid
See: Win-Codin, Tabs. (Winthrop)
- W/Novocaine. (Winthrop Labs) Neo-syneph-
rine HCl 5 mg., novocaine 10 mg./cc.
Amp. 1 cc., Box 10s.
Use: Maintains blood pressure during spinal
anesthesia.
- W/Phenobarbital & theophylline. See: Ad-
nephrine Cap. (Winthrop Labs)
- W/Sulfathiazole. (Winthrop Labs) Neo-
synephrine HCl 40%, sulfathiazole 60%,
isotonic sol. Bot. 1 oz., 1 pt., 1 gal.
Use: Sulfonamide w/nasal decongestant.
- NEO-SYNEPHRINE THENFADIL.** (Winthrop
Labs.) Neo-synephrine HCl & thenfadil
HCl. Jelly 0.5% w/0.1%, Tube ½ oz Sol
0.25% w/0.1%, Bot. 1 oz., 1 pt.; 0.5%
w/0.1%, Bot. 1 oz.
Use: Decongestant & antihistaminic.
- NEO-TARCORTIN.** (Reed & Corrick) Hydro-
cortisone 0.5%, neomycin sulfate 0.35%,
coal tar ext. 5%. Oint. Tube 7 Gm, 1 oz.
Use: Treatment of eczemas or complicated
infections
- NEOTHALIDINE.** (Merck Sharp & Dahme)
Neomycin 200 mg, phenylsulfathiazole
300 mg./Tab. Not listed. Granules sulfa. 1.5
Gm., eq. 1 Gm./15 cc. Bot. 120 cc.
Use: Antibiotic & sulfonamide therapy.
- NEO-THERACORT SUSPENSION.** (C & M
Pharmaco) Neomycin sulfate equiv. to
0.35% neomycin base, hydrocortisone
0.25%, colloidal sulfur 4%, salicylic acid
1.75% in an aqueous base. Bot. 1 oz.
Use: Treatment of acne.
- NEOINTILINE, N.N.D. 1962.** (Paul Money)
Dyphylline, Myphylline, dihydroxypropyl-
theophylline. Tab. (1½ gr) Bot. 100s, 1000s.
Tab (3 gr) Bot. 100s, 1000s. Supp. (7½
gr) Box 12s, 100s. Amp. 1M. (7½ gr)
2 cc., Box 6s, 25s. Tablets also available
w/Phenobarbital ¼ gr.
Use: Orally (1-2 11 d) diuretic, vasodilator,
bronchodilator and myocardial stimulant
- W/Ethaverine HCl See: Neovar, Tab. (Paul
Maeey)

- W/Nicotinic Acid, pentylene tetrazole See: Nea Penl, Tab (Paul Maney)
- W/Pentaerythritol See: Niphylline, Tab (Paul Maney)
- W/Phenobarbital, potassium iodide See Min lo Phen Tab (Paul Maney)
- W/Racephedrine HCl, isoproterenol, phenobarbital See Asophylline, Tab (Crystall)
- W/Rauwolfia serpentina alkaloids & rutin See: Laten, Encote (Paul Maney)
- W/Veratrum viride, rutin & phenobarbital See: Vir I Phyl, Tab (Paul Maney)
- NEOTHYLLINE W/PHENOBARBITAL**, (Paul Maney) Tab (Neothylline 1½ gr, phenobarbital ¼ gr /tab) Bat 100s, 1000s Tab (Neothylline 3 gr, phenobarbital ¼ gr /tab) Bat 100s, 1000s
- NEOTHYN**, (Bentex) Isoproterenol HCl 5 mg, pentobarbital sodium 1/5 gr, Neothylline 2½ gr, ephedrine sulfate ¾ gr, butabarbital 1/6 gr, potassium iodide 5 gr /tab
Use: Prevention of asthma
- NEOTHYN ELIXIR**, (Bentex) Neothylline 2½ gr, ephedrine sulfate ¾ gr, butabarbital ¼ gr, potassium iodide 5 gr /Tab; neothylline 1¼ gr, ephedrine HCl ¼ gr, butabarbital sodium ¼ gr, calcium iodide 1¼ gr, alcohol 10% aromatic vehicle Qs / 5 cc
Use: Bronchodilation
- NEOTINIC**, See: Stuart Neotinic, Tab (Stuart)
- NEO-TOROCAL**, (Plessner) Rile salt 1 gr, cascara sagrada 1 gr, oleoresin capsicum 1/40 min, peppermint oil 1/200 min /Tab Bat 100s, 500s 1000s
Use: Choleric
- NEO-TOTABEX**, (Smith, Miller & Patch) Thiamine HCl 10 mg, riboflavin 10 mg, niacinamide 50 mg, pyridoxine HCl 1 mg, panthoenol 5 mg, ascorbic acid 100 mg, Vit B 12 5 mcg /Cap Bat 100s
Use: Vitamin deficiencies
- NEOTRACIN** (Prema) Bacitracin 500 u, neomycin sulfate 5 mg /Gm Tube ½ oz
Use: Topically skin infections
- NEOTRESAMIDE STERILE SOLUTION**, (Merck Sharp & Dohme) Sod sulfamerazine 0.6 Gm, sod sulfadiazine 12 Gm, sod sulfamethazine 12 Gm, water for inj q s 50 cc Box 1s, 6s, 25s, 100s
Use: IV inj, triple sulfonamide therapy
- NEOTRESAMIDE TABLETS**, (Merck Sharp & Dohme) Sulfamerazine 01 Gm, sulfadiazine 02 Gm, sulfamethazine 02 Gm /tab Bat 100s, 1000s
Use: Triple sulfonamide therapy
- NEO-TREYHIST TABS**, (Merit Pharmaceutical) Thelypyromine HCl mg, chlorpheniramine mol 2 mg, pyrilamine mol 10 mg, phenylephrine HCl 5 mg /Tab Bat 100s 1000s
Use: Cold Therapy
- NEOTRISUL**, (Ascher) Sulfadiazine 1.67 gr, sulfamerazine 1.67 gr., sulfamethazine 1.67 gr, Sod bicarb 5.0 gr./Tab Bat 100s 500s, 1000s
Use: Sulfonamide susceptible bacteria
- NEOTRIZINE**, (Lilly) Susp; Sulfadiazine 3.33 Gm, sulfamethazine 3.33 Gm Tab 0.5 Gm Bat 100s 1000s sulfamerazine 3.33 Gm / 100 cc Bat 1 pt
Use: Orally; triple sulfonamide therapy
- W/Pot penicillin G (Lilly) Neotrizine 0.5 Gm, Pot penicillin G 150,000 u/Tab., Rot 24s, 100s, 1000s
Use: Bacterial infections
- NEOTROL**, (Hance) Phenylephrine HCl 0.25%, pyrilamine maleate 0.2%, cetalkonium chloride 0.05%, tyrothricin 0.03% phenylmercuric acetate 1:50,000 Sol., Squeeze Bat 20 cc
Use: Nasal congestion
- NEOTUSS-PT**, (Lloyd Dabney & Westerfield) Dihydrocodeinone bitartrate 18 mg, phenyltoloxamine dihydrogen citrate 10 mg, sod citrate 215 mg, pot guaicol sulfonate 110 mg, citric acid 22 mg, racemic amphetamine sulfate 1 mg /5 cc Bat 1 pt
Use: Antitussive
- NEO-ULCORT**, (Uimer) Hydrocortisone diethylaminoacetate HCl 5 mg, neomycin sulfate 5 mg /Gm of Oint Tube ½ oz
Use: Anti-inflammatory
- NEOVACAGEN** (Merck Sharp & Dohme) Vacagen w/methapyrilene HCl 25 mg /tab Bat 20s, 100s
Use: Cold vaccine w/antihistamine
- NEO-VADRIN**, (First Texas) dl Norephedrine HCl 0.4%, phenylephrine HCl 0.15%, chlorbutanol 0.15%, benzalkonium chloride 0.005% Dropper Bat 1 oz
Use: Nasal decongestant
- NEO-VAGISOL**, (Smith Dorsey) Tyrothricin 5 mg phenylmercuric acetate 3 mg, succinic acid 12.5 mg, papain 25 mg, sodium lauryl sulfate 3 mg, lactose 0.75 Gm / Sepp Vag Box 24s
Use: Trichomoniasis
- NEOYAR**, (Paul Maney) Neothylline 100 mg, ethaverzine HCl 30 mg /Tab Bat 100s Alia w/neothylline 200 mg
Use: Orally (1 2 1 d), treatment of cardiovascular disease
- NEOVIASEPT**, See: Milibis W/Arofen (Winthrop)
- NEO-VIN SOL**, (Walker Corp) Oxyquinoline sulfate 1/5 gr w/ammonium alum, borax

- acid & Sod. chloride, menthol & thymol/
Tab., Bot. 100s.
Use: Antiseptic solution.
- NEOXYN.** (Rorer) Acetonilid, acetic acid,
benzethonium chloride, hydrogen peroxide,
propylparahydroxybenzoate, water & aromatics. Lot. 4 oz.
Use: Poison ivy, poison oak & poison sumac.
- NEOZIN, BUOPTO.** (Professional Pharm.)
Co) Phenylephrine HCl 1/2%, zinc sulf. 1/2
gr. and benzalkonium chloride/15 cc., Bot.
Use: Eye drops
- NEPENTH.** (Tilden) Blackberry bark, nutgall,
ginger, clove, cinnamon, prickly ash, bark,
nutmeg and aromatics. Bot. pls, gals.
Use: Carminative, astringent, anti-diarrheal
- NEPHENALIN.** (Leeming) n-Isopropyl ortera-
nol HCl 10 mg., theophylline 2 gr., ephed-
rine sulfate 1/2 gr., phenobarbital 1/2 gr./
Tab. Vial 20s. Bot. 100s, 1000s.
Use: Asthmatic symptoms.
- NEPHENALIN PEDIATRIC.** (Leeming) Outer
coat contains ephedrine HCl 5 mg.; inner core
contains ephedrine sulfate 3/16 gr.,
theophylline 1 1/2 gr., phenobarbital 1/2 gr./
Tab. Bot. 20s, 100s.
Use: (1 q. 4 h.) Treatment of asthma in
children.
- NEPHRIDIN.** See: Epinephrine (Various
Mfrs.).
- NEPRITIN.** (Reed & Carnrick) Theophylline
25 mg., rutin 10 mg. In excipient containing
dical. & defatted kidney substance, milk
sugar, starch & gum arabic/Tab. Bot. 80s,
500s, 1000s.
Use: Diuretic & capillary wall strengthener.
- NEPTAZAN.** NND 1962 (Lederle) Metha-
zalamide 50 mg/Tab. Bot. 100s.
Use: Carbonic anhydrase inhibitor, treat-
ment of glaucoma.
- NEPYTOL.** (Elder) Thyroid 1/5 gr., zinc
phosphide 1/10 gr., iron pyrophosphate 2
gr./Tab. Bot. 100s, 500s.
- NEBAVAL.** N.N.D. 1959. (Schering) Methi-
lural Sod. 5-[1-Methylbutyl]-5-[2-methylthio
ethyl]-2-thiobarbiturate, Vial, sterile pow.
1, 2, 5 Gm.
Use: i.v., general anesthesia
- NEBVIN.** (Hance) Sod. bromide 12 gr., pot.
bromide 12 gr., ammonium bromide 6 gr./
fl. oz. Bot. 6 oz.
Use: Sedative.
- NECACINE HCl.** NND. 1962. (Mallinck)
Chloroprocaine HCl. Vial (1%, 2% or 3%)
30 cc.
Use: Infiltration anesthesia or various re-
gional blocks.
- NECACINE W/EPINEPHRINE INJECTION.**
(Mallinck) Nesocaine 1% and 2% solutions
w/epinephrine 1:100,000 for infiltration,
field, and regional block; 3% solution w/
epinephrine 1:200,000 for caudal and epi-
dural block. Also available without epi-
nephrine. 1% and 2% vials 30 cc. multiple
dose. 3% vial 30 cc. single dose.
- NESDONAL SODIUM.** See: Thiopental
sodium (Pantothal Sod. Abbott).
- NESTBEX TABLETS.** (Fielding) Pyridoxine
HCl (B-6) 50 mg; phenyltoloxamine dihy-
drogen citrate 50 mg; scopolamine HBr
0.5 mg/Tab. Bot. 50s.
Use: Anti-emetic.
- NETHACETIN.** (Merrill) Nethamine HCl 45
mg., acetophenetidin 0.23 Gm., acetyl-
salficylic acid 0.13 Gm./tab. Bot. 100s,
1000s.
Use: (1 q. 3-4 h.) Orally; antispasmodic &
analgesic.
- NETHACOL.** (Merrill) Nethamine HCl 65
mg., Ipecac fl. ext. 0.06 cc., ammonium
chloride 0.63 Gm., menthol 8 mg./30 cc.
Syr. Bot. 1 pt., 1 gal.
Use: (1 tsp. q. 3 h) Expectorant.
- NETHAMINE HCl.** 1-N-ethylephedrine HCl,
2-methylethylamine-1-phenylpropanol-1 HCl.
W/Acetophenetidin & aspirin. See: Netha-
cetin, Tab. (Merrill)
- W/Codaine phosphate, phenylephrine HCl,
sodium citrate, dextylamine succinate.** See
Mercodol with Decapryn Syrup (Merrill)
- W/Ipecac fl. ext., ammonium chloride & men-
thol.** See: Nethacol, Syr. (Merrill)
- W/Phenobarbital & butaphyllamine.** See
Nethaphyl, Cap. (Merrill)
- NETHAPHYL.** (Merrill) Butaphyllamine 120
mg., phenobarbital 15 mg., nethamine HCl
50 mg./cap. Bot. 100s. Also available in
half strength.
Use: (1 t.i.d.) treatment of asthma.
- NETHAPRIN.** (Merrill) Nethamine HCl 25
mg., butaphyllamine 60 mg., decapryn suc-
cinate 6 mg./Cap. or 5 cc. Bot. 100s. Syr.
Bot. 6 oz., 1 pt., 1 gal.
Use: (1 t.i.d.) Bronchial asthma allergy.
- NETRIN.** (Geigy) Under study See: Metcra-
phan HCl.
Use: Anticholinergic.
- NEULIN.** (Walker Labs.) Phenobarbital 1/2 gr.,
Vit. B-1 3 mg., Cal. phosphate tribasic 1 1/2
gr./tab. Bot. 25s, 100s.
Use: Sedative.
- NEURAMIDE.** (Sherman) Protamide.
Use: Ophthalmic herpes
- NEURINASE.** (Faugera) Barbitol 3.34 Gm.,
chloroform 0.052 cc., valerian ext. 0.75
Gm./100 cc. Bot. 110 cc. Tabs. barb. 0.26
Gm., valerian ext. 0.049 Gm/Tab. Tubes
20s. Sol. Bot. 1 pt. Tab., Bot. 22s.
Use: Sedative.

NEURO-CENTRINE. (Bristol) Not listed
Phenobarbital $\frac{1}{4}$ gr, centrine hydrogen
sulf 0.25 mg, reserpine 0.05 mg/Tab
Bot 100s

Use: (1 2 11 d) Treatment of stress & ten-
sion states

NEUROIDS. (Tilden) Barbital 1 gr, hyo-
scyamus ext $\frac{1}{8}$ gr, passion flower ext $1\frac{1}{2}$
gr, valerian ext $\frac{1}{8}$ gr/Tab Bot 100s
1000s

Use: Sedative

NEURONIDIA. (Schieffelin) Barbital 2 gr/oz
Elix Bot 8 oz, 1 gal

Use: Sedative

NEUROPLEX INJ. (Ries) Vit B 1 100 mg,
B 12 500 mcg, l glutamic acid 100 p p m/
cc Vial (multiple dose) 20 cc

Use: Nerve tone in secondary neuropathoses
NEUROSIN. See: Calcium glycerophosphate
(Various Mfr)

NEUROSINE. (Dias) Sod bromide 30 gr,
Pot bromide 30 gr, ammonium bromide
30 gr, zinc bromide $\frac{3}{5}$ gr, hyoscyamus
ext $\frac{3}{40}$ gr, belladonna fl ext $\frac{3}{40}$ min,
cascara $\frac{3}{5}$ gr, humulus ext $\frac{3}{5}$ gr. In
pleasantly flavored 4% alcoholic aq
vehicle/fl oz Bot 4 & 8 oz

Use: (1 2 1sp w/water) Antispasmodic

NEUTHENIC. (Faugera) Sod glycerophos-
phate 0.1 Gm, strychnine cacodylate
0.0005 Gm, cacodylic acid 0.005 Gm/cc.
Amp 1 cc Oral use, Bot 25 Gm

Use: Tonic

NEUTRA-CORDIALS SPECIAL (Sulliff &
Cose) Sodium bicarbonate 78 Gm, cinna-
mon 78 Gm, rhubarb 156 Gm, berberis
aquifolium 49 Gm spirit peppermint/fl oz
Liq Bot pt, gal

NEUTRAL ACRIFLAVINE, NNB 1952,
ADR 1954 See: Acriflavine (Various Mfr)

NEUTRALIZER WAFER. (Upjohn) Cal carb
12 gr, Mag carb 1 gr, Mag trisilicate 2
gr/Wafer Bot 100s 500s 1000s

Use: Gastric antacid

NEUTRALIZING CORDIAL, N.F. (Merck
Sharp & Dahme) Rhubarb flex 16 cc,
hydrastis flex 0.8 cc, pot carb 16 Gm/
100 cc Bot 1 gal

Use: Laxative & antacid

NEUTRALIZING POWDER. (Roymex) Es-
s-muth subcarb, Cal carb, kofein, Mag
oxide, Mag trisilicate, Sod bicarb Bot 35
Gm

Use: Gastric hyperacidity

**NEUTRAL PROTAMINE HAGEDORN-IN-
SULIN** See: Insulin NPH (Lilly)

NEUTRALON COMPOSITE. (Consol Med
land) Aluminum hydroxide 260 mg, mag

* Generic name recognized by N.F., U.S.P. or A.M.A.

nesium trisilicate 488 mg/Tab Bot 30s,
100s, 250s, 1000s

Use: Orally, antacid

NEUTRAPEN, NND 1962 (Schen Lab) In
jectable form of purified penicillinase Vial
(800,000 u) 1 dose, Box 12s

Use: I.M., relief of allergic reactions of
penicillin

NEUTRAZYME. (Smith Dorsey) Sod lauryl
sulfate 100 mg/Supp Box 12s

Use: Rectally; pruritus ani.

NEUTROFLAVINE. See: Acriflavine (Various
Mfr)

NEUTROGENA. (Notone Co) A neutralized
soap (pH 7.5)

NEUTRON (Myers Carter) Testosterone 25
mg, estrone 2 mg/cc Vial 10 cc

NEUTROTABS. (Blue Line) Hamatrapine
methylbromide 2.5 mg, aluminum hydrox-
ide gel 129.6 mg, magnesium trisilicate
194.4 mg, phenobarbital 8 mg/Tab Bot
100s, 1000s

Use: Antispasmodic, antacid, sedative

NEVAX. (Tilden) Sodium dioctyl sulfosuc-
inate 60 mg/Cap Bot 50s, 100s, 1000s

Use: Constipation

NEVROTOSE (Vale) Phenobarbital $\frac{1}{2}$ gr,
hyoscyamus ext $\frac{1}{8}$ gr, passiflora ext $\frac{1}{8}$
gr, valerian ext $\frac{1}{8}$ gr, camphor mono-
bromate $\frac{1}{8}$ gr/Cop Bot 100s, 1000s

Use: Sedative

NEXORIN. (Smith Dorsey) d Amphetamine
sulfate 5 mg, amobarbital 15 mg, methyl
cellulose 200 mg, Mag trisilicate 200 mg,
Vit B 1 0.67 mg, R 2 0.67 mg, nicotin-
amide 6.7 mg/Cop Bot 100s 500s, 1000s

Use: (1 11 d, a c) Antiobesity

NIACIN, USP NND 1962 ADR See:
Nicotinic Acid Prep (Various Mfr)

NIACINAMIDE, NND 1958, ADR Nic-
otinic acid amide See Nicotinamide, USP
(Various Mfr)

NIACINAMIDE HI, W/Pat iodide See:
Iodo Niacin (Cole)

NIACINATE (New Mexico Pharm) Sodium
nicotinate and nicotinic acid equiv to 100
mg chlorbutanol 0.35%/2 cc Vials 20 cc

NIADRIN, NND 1959 (Endo) Isoiazid
Tab 50 mg, 100 mg Bot 100s

Use: TB therapy

***NIALAMIDE, NND 1962** Nisonicotinoyl
N (β N benzylcarboxamidoethyl) hydro-
zine See Niamid (Pfizer)

NIALEX. (Mallard) Pentylene tetrazol 50 mg,
nicotinic acid 50 mg, hesperidin complex
50 mg vit C 30 mg/Tab Bot 100s, 1000s

Use: An effective analeptic and a CNS
stimulant

- NIALIFT.** (B. F. Ascher) Pentylene-tetraazol 100 mg., nicotinic acid 50 mg., thiamine HCl 2 mg., riboflavin 2 mg., pyridoxine HCl 1 mg., calcium pantothenate 5 mg., ascorbic acid 50 mg./Tab. Bot. 100s & 500s
Use: For fatigue and debility
- NIAMID, N.N.D. 1962.** (Pfizer) N-[2-(benzyl-carbonyl)ethylomina]-isonicotinamide Niacinamide. 25 mg./Tab., 100 mg./Tab. Bot. 100s.
Use: Antidepressant.
- NIAPENT.** (Ferndole) Pentylene-tetraazole 100 mg., nicotinic acid 50 mg., glutamic acid/Tab. or 15 cc. Elixir Bot. 1 pt.; Tab. Bot. 100s.
Use: Symptoms resulting from atherosclerosis-sclerotic degeneration.
- NIATHERM.** (Podiatrex) Niacin, iodine, vitamins B-1, B-2, B 6, for podiatrists/chiropractists. Tabs. Bot. 120s.
Use: Peripheral vasodilation.
- NIATRIC.** (Ascher) Pentylene-tetraazol 100 mg., nicotinic acid 50 mg., ascorbic acid 100 mg., biotinoids 100 mg./Tab. Bot. 100s, 500s. Elix. Bot. 1 pt.
Use: For the old age syndrome, the mildly confused, forgetful, apathetic, irritable, antisocial, aging patient.
- NIATRICRINE.** (Ascher) Pentylene-tetraazol 100 mg., nicotinic acid 50 mg., Vit. C 100 mg., methyltestosterone 2 mg., ethinyl estradiol 0.01 mg., thyroid 15 mg./Tab. Bot. 100s, 500s. Elixir contains sed. l-thyroxine 24 mcg./5 cc. in place of thyroid in above formula. Bot. 1 pt.
Use: Treatment of geriatric patients.
- NIAZO.** Neatropin. Use: Urinary antiseptic
- NIBARB.** (Breon) Mannitol hexanitrate 1/2 gr., phenobarbital 1/4 gr./Tab. Bot. 100s & 1000s.
Use: (I.I.d.) Coronary vasodilator.
- NICALEX.** (Walker Labs) Aluminum nicotinate 625 mg equivalent to 500 mg nicotinic acid/Tab. Bot. 100s, 1000s.
Use: Hypercholesterolemia.
- NICAMIDE.** See: Nixethamide (Various Mfrs.)
- NICAMIN.** (Abbott) Monoethanolamine nicotinate. Amp. (25 mg./cc) 2 cc., 6s, 25s, 100s. Vial (50 mg./cc.) 20 cc.
Use: Nicotinic acid therapy.
- NICAMINDON.** See: Nicotinamide (Various Mfrs.)
- NICASAMIN 'B.'** (Rorer) Thiamine HCl 0.5 mg., riboflavin 1 mg., niacinamide 5 mg., brewer's yeast 3 gr./Tab. Bot. 100s, 500s, 1000s.
Use: Dietary supplement.
- NICEL.** (Irwin-Netsler) See: Methylcellulose.
- NICKEL PECTINATE.** See: Tamectin, Pow (Ayers).
- NICOBEE.** (Endo) Vit. B-1 25 mg., nicotinic acid 50 mg., nicotinamide 150 mg., Vit. B-12 1D mcg./fl. oz. Bot. B 3 oz., 1 pt., 1 gal. Inj. Amp 10 cc., Vial 30 cc
Use: Vit. B complex.
- NICOBION.** See Nicotinamide (Various Mfrs.)
- NICODEN.** (Lemman) Adenine-5-monophosphoric acid 50 mg., Vit. B-12 cryst. 100 mcg., nicotinic acid 20 mg., procaine HCl 1%, benzyl alcohol 1%/cc Vial 10 cc.
Use: Treatment of arthritis.
- NICOCODINE.** (Registered in Great Britain) 6-Nicotinoylcodeine
- *NICODUOXIDE.** A mixture of nicotiazone and isoniazid
- NICOLEXIN.** (Upjohn) Nicotinic acid amide 75 mg., riboflavin 1 mg., thiamine HCl 1 mg., pyridoxine HCl 0.1 mg., Cal. pantothenate 0.25 mg./Cap. Bot. 50s, 500s.
Use: Vit. B complex.
- NICO-METRAZOL.** (Knoll) Pentamethylene-tetraazol 100 mg., nicotinic acid 50 mg./5 cc or Tab. Elixir, Bot. pt & gal. Tab. Bot. 100s, 1000s
Use: Senile deterioration.
- NICONYL, N.N.D. 1962.** (Parke, Davis) (Isoniazid, Isonicotinic Acid Hydrazide) Tab. 50 mg., 100 mg., Bot 100s, 1000s
Use: Tuberculosis.
- NICORIS.** (Eitro) Niacinamide 200 mg., riboflavin 5 mg./cc. Vial 10 cc.
Use: Niacin & riboflavin deficiency
- NICOSODE.** (Chicago Pharmacol) Nicotinic acid Sod. salt 118 mg/cc Amp 1 cc., 12s, 25s, 100s. Vial 30 cc., Box 1s, 6s.
Use: I.V. & nicotinic acid therapy
- NICO-SPAN.** (Toble Rock) Nicotinic acid 400 mg./Cap Bot. 100s.
Use: Blood cholesterol.
- NICOTAMIDE.** See: Nicotinamide (Various Mfrs.)
- *NICOTHAZONE.** Nicotinylaldehyde thiosemicarbazone.
- NICOTHIAMIN.** (Abbott) Thiamine HCl 40 mg., nicotamide 400 mg. in palatable elixir base/fl. oz. Bot. 12 oz., 1 gal.
- NICOTILAMIDE.** See: Nicotinamide (Various Mfrs.)
- *NICOTINAMIDE, U.S.P., N.N.D. 1958:** Cap., N.F. Inj. U.S.P., A.D.R.; Tab. U.S.P. (Amnicotin, Dipegyl, Niacinamide, Nicamindon, Nicotamide, Nicotinamide, Nicotinic Acid Amide, Pelatin Amide)
W/Ascorbic acid. See: Dentacevin, Tab. (Walker)

* Generic name recognized by N.F., U.S.P. or A.M.A.

- W/Bromides See Braniozin, Liq & Tab (Massengill)
- W/Methamphetamine HCl, lactic acid, cyano cabalamin See Rufaldex, Vial (Central)
- W/Pentylentetrazol thiamine HCl, cyanocobalamin, alcohol See Cenalene, Liq (Central)
- W/Phenyltolaxamine dihydrogen citrate, scopalamine HBr, benzocaine See Hyloxine Tab (Tutag)
- W/Pot Iodide See Iodo Niczin, Tab (Cole Chem Co)
- W/Riboflavin See Nicriba Amp (Breon) Nira, Vial (Barry) Riboflavin & Nicotinamide, Amp (Lilly)
- W/Thiamine HCl See Nicolthiamin, Liq (Abbott)
- NICOTINATE, METNYL, W/Other hyperemia producing agents** See Rubiquent Cream (Wyeth)
- NICOTINATE MONOETHANOLAMINE, See** Nicamin, Amp (Abbott)
- ***NICOTINIC ACID, USP, NND 1962** ADR Pyridine 3 carboxylic acid Cap NF X, Inj USP; Tab, USP (Niczin) (Various Mfr)
- See: Efacin (Person & Covey) Nico-Span Caps (Table Rock)
- W/Adenosine 5 monophosphoric acid, vit B 12, procaine HCl & benzyl alcohol See Vibuden, Vial (Philadelphia Ampoule Labs)
- W/Glycine See Nicatron Elixir (Haug Drug)
- W/Inositol Hexanicotol (Philadelphia Lab)
- W/Meclizine See: Antivert, Tab (Roerig)
- W/Pentamethylentetrazol See Nico Met, Tab & Elix (Knoll)
- W/Pentylentetrazol
- See: Anoleptone, Elix (Reed & Carnrick) Arterine, Tab or Elix (Confield) Benzal Tab, Elix (Bentex) Geraniazol, Tab (Columbus Pharmacal) Menic, Tab (Geriatric)
- W/Niopent (Ferndale) Senilex, Tab (Durst) Tolec, Tab (Hoack) Vaso-tabs, Tab (Paul Maney)
- W/Pentylentetrazol, ascorbic acid, bio flavonoids See: Niatric, Tab (Ascher)
- W/Pentylentetrazol, dienestral, methyltestosterone See: Lucidon, Tab (Lloyd, Dabney & Westerfield)
- W/Pentylentetrazol, ethinyl estradiol, methyltestosterone, and dicalcium glutamate See: Senotic, Tab (Dumas Wilson)
- W/Pentylentetrazole, neothylline See: Neo-Pent Tab (Paul Maney)
- W/Pentylentetrazol, tetraiodo-tyrosine, meth
- *Generic name recognized by N.F., USP or A.M.A.
- yltestosterone, ethinyl estradiol See: Tetra Forte, Tab (E S Miller)
- NICOTINIC ACID AMIDE, NND 1958, ADR** See Nicalinamide, USP (Various Mfr)
- NICOTINIC ACID, AS SODIUM SALT, Inj USP**
- See Naotin, Amp (Drug Products) Nicasode Vial (Chicago Pharm)
- ***NICOTINYL ALCONOL** 3 pyridinemeth anol See Roniacol (Roche) Tigacol Caps (Roche)
- NICOTRON, (Haug Drug)** Nicotinic acid 75 mg, glycine 750 mg/5 cc or Tab Bot 8 oz Tab Bot 100s Use Vitamin
- NICOZIDE, NND 1962 (Fremo)** Isonicotinic acid hydrazide (isoniazid) Tab 50 mg, 100 mg Bot 100s, 1000s Use Streptomycin resistant tuberculosis
- NICOZOL, (Hart Laboratories)** Pentylene tetrazol 200 mg nicotinic acid 100 mg, alcohol 5%/5 cc Elixir 1 pt Pentylentetrazol 100 mg, nicotinic acid 50 mg Tab, or Cap -Bot 100s 1000s
- NICOZOL COMPLEX (Hart Laboratories)** Pentylentetrazol 150 mg nicotinic acid 75 mg methyl testosterone 2.5 mg, ethinyl estradiol 0.01 mg thiamine HCl 6 mg, riboflavin 3 mg pyridoxine HCl 6 mg, Vit B 12 2 mCg folic acid 0.33 mg, pantothenol 5 mg choline bitartrate 20 mg, inositol 15 mg lysine monohydrochloride 100 mg, Vit E 3 mg, iron (as ferric pyrophosphate) 15 mg magnesium 2 mg, manganese 1 mg zinc 1 mg alcohol 15%/15 cc Liq Bot 1 pt 1 gal Use Geriatric therapy
- NICRIBO (Breon)** Nicotinamide 200 mg, riboflavin 5 mg/cc Amp 1 cc, Box 12s, 25s, 100s Use IV & IG; Vit B therapy
- NIDAR, (Armour)** Secobarbital Sod 25 mg, pentobarbital Sod 25 mg butobarbital Sod 7.5 mg, phenobarbital 7.5 mg/Tab Bot 100s Use: (1 2 hs) Sedative
- W/Reserpine See: Nitensar, Tab (Armour)
- NIDATON, (Organon)** Isonicotinic acid hydrazide (Isoniazid) Tab 50 mg, Bot 100s Use: Streptomycin resistant tuberculosis
- NIDOXITAL, (Ortho)** Benzocaine 100 mg pyridoxine HCl 50 mg, nicotinamide 25 mg, dl methionine 100 mg, pentobarbital Sod 75 mg/Cap Bot 24s, 100s Use: (1 2 30 to 40 minutes a c) Nausea & vomiting of pregnancy
- ***NIDROXYZONE, 5 Nitro 2 furaldehyde-2**

- (2-hydroxyethyl)semicarbazone. See: **Ferodoxyl** (Eaton Lab)
- NIERALINE**. See: **Epinephrine** (Various Mfr.)
- NIFURALDEZONE**. See: **Furamazone** (Eaton)
- *NIFURETHAZONE**. 5 Nitro-2-furaldehyde 2-(2-dimethylaminoethyl)semicarbazone. (Eaton Labs.)
- *NIFUROXIME**, N.F. ans 5-nitro-2-furaldoxime. See: **Micolur** (Eaton)
- W/**Furozolidone**. See: **Tricofuron**. (Eaton)
- NIGINA**. (Hoag) Triethanolamine trinitrate biphosphate 2 mg./Tab. Bot. 100s.
Use: Vasodilator.
- NIGLYCON**. (CMC) Nitroglycerine 1/25 gr./tab.
Use: Treatment of angina.
- *NIHYDRAZONE**. 5-Nitro-2-furaldehyde acetylhydrazone. See: **Puriton** (Eaton Labs)
- NIKA-CAPS Ph 749**. (Paul Maney) di-Amphetamine sulfate 15 mg., thyrald 1½ gr., atropine sulfate 1/180 gr., aloin ¼ gr., phenobarbital ¼ gr./cap. Bot. 100s.
Use: 1 daily; antiobesity.
- *NIXETHAMIDE**, N.F., N N D. 1962; A. D R; Inj. N.F. {Anacardone, Cordiamid, Cormed, Dynacoryl, Eucoran, Nicamide, Pyricardy} N, N-Diethyl-3-pyridinecarboxamide
See: **Caramine**, Amp. & Bot. (Ciba)
Nikarin, Amp. & Bot. (Massengill).
Use: Central nervous system stimulant.
- NIKETNAMIDE TARTRATE**. See: **Nikarin**. Cap. (Massengill).
- NIKORIN**, N N D 1962. (Massengill) Nikethamide, Amp. 25%, 1.5 cc.; 6s, 25s, 100s Bot., Sol. 25% orally, 4 oz., 1 pt.
Use: Central nervous system stimulant.
- NIKORIN TARTRATE**. (Massengill) Nikethamide tartrate. Cap. 250 mg. Bot. 100s, 500s.
Use: Cardiovascular stimulant.
- NILEVAR**. 1962. (Seoria) Norethandrolone 17-Alpha-ethyl-17-hydroxynorandrosthenone Tab. 10 mg., Bot. 50s. Amp. 25 mg./cc. I.M., Pk. 6s, 25s, Drops Bot. 15 cc
Use: Treatment of negative nitrogen balance & protein anabolism.
- NILTUSSIN**. (Pharmaceutical Products) Dihydrocodeinone bitartrate 1.67 mg., Pot guaicol sulfanate, antimony & Pot. tartrate, squill syr./5 cc. Syr. Bot. 4 oz., 1 pt. 1 gal.
Use: Cough preparation.
- NILO**. (Cofe) Sod. nitrite 2 gr., Sod. glycocholate ½ gr., chianonithin ¼ gr., Irlin 1/10 gr./Tab. Bot. 100s
Use: Vasodilator.
- NILOX**. (Direct Labs) d-Amphetamine Sulf 15 mg., butobarbital ¼ gr., acetphenalis-
- atin** 5 mg./Cap. T. D. Bot 30s
Use: Obesity
- NILTUSS**. (Minnesota Pharm. Lab., Inc) Dexamethorphan hydrobromide 10 mg., chlorpheniramine maleate 1.25 mg., phenylephrine hydrochloride 5 mg., ammonium chloride 83 mg./isp Syr.; Bot. 4 oz., 1 pt. Chlor: Cough syrup, antitussive, vasoconstrictor, antihistaminic, expectorant.
- NIMO-TEX**. (Sutliff & Case) Vit. A 5000 u., D 500 u., E 2 I.U., B-1 3 mg., B-2 2 mg., B-6 0.5 mg., B-12 1 mcg., cal pantothenate 3 mg., folic acid 0.5 mg., choline bitartrate 50 mg., inositol 10 mg., Vit. C 30 mg., larrous sulf. 60 mg., copper 1 mg., cobalt 0.2 mg., manganese 3 mg., iodine 0.15 mg., molybdenum 0.2 mg., zinc 1 mg., magnesium 5 mg., calcium 48 mg., phosphorus 37 mg./Tab Bot. 100s, 1000s.
Use: Vitamin & mineral supplement
- NIO-A-LET**. (Nion) Vit. A. Cap. 25,000 u Bot. 100s, 500s; 50,000 u. Bot. 100s.
Use: Vit. A deficiencies.
- NIODOLIN**. (Lincoln) Foreign protein 20 mg., Sod. iodide 40 mg./cc. Vial 1 oz.
Use: (1-2 cc. daily) Nonspecific protein therapy.
- NION A**. (Nion) Vit. A acetate 25,000 u./Cap. Bot. 100s, 250s, 1000s.
Use: Vit. A deficiencies.
- NIONATE**. (Nion) Ferrous gluconate 5 gr./Tab. Bot. 100s, 1000s.
Use: Iron deficiency hypochromic anemia.
- NIONDOX**. (Nien) Ascorbic acid 100 mg., rutin 20 mg., Vit. B-6 5 mg., citrus bioflavonoids 100 mg./Tab. Bot. 100s. Also available w/vit. K.
Use: Treatment of impaired capillary fragility
- NION-LAX**. (Nien) Diocetyl sodium sulfocinate 60 mg., acetophenolisatin 5 mg., thiamine HCl 1 mg., citrus pectin cellulose complex 200 mg./tab.
Use: Laxative.
- NIO-PIRACENE**. (Nion) Scopelamine HBr 0.15 mg., prophenpyridamine 15 mg./Tab. Bot. 30s, 100s.
Use: For motion sickness.
- NIPHEN**. (Torch) p-Nitrophenol 1% solution. Bot. 1 oz., 4 oz., 1 pt.
Use: Treatment of fungus infections.
- NIPHYLLINE**. (Paul Maney) Dihydroxypropyl theophylline (Neophylline) 100 mg., penterythritol tetranitrate 10 mg./tab. Bot 100s
Use: (1-2 T.D.) Angina pectoris.
- NIPIRIN**. (Massengill) Pentobarbital ½ gr., acetylsalicylic acid 5 gr./Cap. Bot. 100s, 1000s
Use: Sedative, hypnotic, analgesic

* Generic name recognized by N.F., U.S.P. or A.M.A.

- NI-PLEX SYRUP.** (Irwin, Neisfer) Vit B 1 16 mg, B 2 0.33 mg, niacinamide 5 mg, pantothenic acid 0.25 mg, choline salt 6.25 mg/dr Bot. 1 pt, 1 gal.
Use: Vit B complex therapy
- NIRO.** (Barry) Niacinamide 200 mg, riboflavin 5 mg/cc Vial 10 cc.
Use: I.M.; Vit B therapy
- NISALIN** (National Drug) n Propylsulfonate 5%, methyl salicylate 15%, menthol 10%, thymol 1%, nicotinylsalicylic acid 1% Oint
Use: Treatment of osteoarthritis, rheumatoid arthritis and fibrositis
- NISENTIL, NND 1962** (Rache) Alphaprodine HCl 1.3 dimethyl-4 phenyl 4 propionoxy piperidine HCl Amp (40 mg) 1 cc Box 6s, 50s; (60 mg) 1 cc Box 12s, 100s Vial (60 mg) 10 cc Box 1s
Use: (4 mg q 3 h) Obstetrical analgesic
- NISULFAZOLE, NND 1962** (Breon) p Nitro sulfathiazole Susp 10%, Bot 10 fl oz
Use: Intrarectally; sulfonamide therapy
- NITCO** (Sutliff & Case) Sod nitrite 65 mg pot nitrate 130 mg paw echinacea 260 mg, paw ext chionanthus 65 mg, paw ext iris 65 mg sod glycocholate 65 mg/Tab Bot 100s, 1000s
- NITENSAR.** (Armour) Not listed Secobarbital sod 3/16 gr, pentobarbital sod 3/16 gr butobarbital sod 1/16 gr phenobarbital 1/16 gr w/Reserpine 0.1 mg/tab Bot 100s
Use: Sedative w/ tranquilizing agent
- NITE-REST.** (AFC) Methapyriene HCl 50 mg scopolamine ominoxyde HBr 0.25 mg/Cap Bot 21s
Use: Sustained action capsules for sleeplessness
- NITOPLEX.** (New Mexico Pharm) Dicalcium phosphate 0.5 Gm, terrous sulfate 0.1 Gm B 1 1 mg, B 2 1 mg O 400 u/Cap Bot 100s
Use: Vitamin & mineral supplement
- NITRAEGUS COMPOUND.** (Buffington) Nitroglycerin 1/250 gr, Sod nitrite 1 gr, crotaegus tr 5 min/Pittet Bot 100s, 1000s
Use: Vasodilator
- NITRALOX** (Smith Dorsey) Alaxoxylon 1 mg, pentaerythritol tetranitrate 10 mg/Tab Bot 100s, 500s, 1000s
Use: (1 2 q 1 d) Angina pectoris
- NITRANITOL, NND 1962** (Merrell) Mannitol hexanitrate 32 mg/Tab Bot 100s, 1000s, 5000s
Use: (1 2 q 4-6 h) Coronary vasodilator
- W/Phenobarbital** (Merrell) Mannitol hexanitrate 32 mg, phenobarbital 16 mg/Tab Bot 100s, 1000s, 5000s
Use: (1 2 q 3-4 h) Angina pectoris
- W/Phenobarbital & rutin** (Merrell) Mannitol hexanitrate 32 mg, phenobarbital 16 mg, rutin 20 mg/Tab Bot 100s, 1000s
Use: Hypertension
- W/Phenobarbital & theophylline** (Merrell) Mannitol hexanitrate 32 mg, phenobarbital 16 mg, theophylline 100 mg/Tab Bot 100s, 1000s
Use: Hypertension
- NITRANITOL P.V.** (Merrell) Mannitol hexanitrate 32 mg, phenobarbital 16 mg, veratrum viride alkaloidal fraction 1 mg/Tab Bot 100s, 1000s
Use: (2 q 1 d) Refractory cases of hypertension
- NITRANITOL R S** (Merrell) Mannitol hexanitrate 1/2 gr, rauwolfia serpentina alkaloids 0.5 mg/Tab Bot 100s, 1000s
Use: (1 2 q 1 d) Vasodilator
- NITRAPNYL.** (Ascher) Mannitol hexanitrate 32.4 mg, phenobarbital 16.2 mg, theophylline Sod glycinate 162 mg/Tab, Bot 100s, 1000s
Use: (1-2 1 d) Vasodilator
- NITRASED** (Lemmon) Nitroglycerin 0.4 mg, pentaerythritol tetranitrate 15 mg, secobarbital sod 15 mg/Tab Bot 100s
Use: Anginal attacks
- NITRAZINE PAPER** (Squibb) Phenaphthazone Sod dinitrophenyl azo naphthal di sulfide Vial 100s, w/color chart, Box 5s Also 15 fl roll
Use: Determine pH of a solution
- NITRETAMIN PHOSPHATE, NND 1962** (Squibb) Trainitrate Phosphate Aminotrate Triethanolamine trinitrate diphosphate 2 mg or 10 mg (long acting)/tab Bot 50s
Use: Vasodilator
- NITRILLIN.** (Maney) Outer layer Glycerol trinitrate 0.4 mg, in tablet core pentaerythritol tetranitrate 15 mg neophylline 100 mg/Tab Bot 100s, 1000s Also with 15 mg phenobarbital
Use: Vasodilator
- NITRINE-TDC.** (Canfield) Nitroglycerin 1/25 gr/T O Cap Bot 100s, 500s, 1000s
Use: Angina pectoris
- NITRITAL** (Lemmon) Phenobarbital 1/4 gr, Sod nitrite 2 gr, veratrum viride tr 4 min/Tab Bot 100s
Use: Vasodilator
- NITROBAR.** (McNeil) Bismuth subnitrate 5 gr, butal 1/6 gr, passiflora ext 1/2 gr, lupulus ext 1/2 gr/Tab Bot 100s
Use: (1 q 1 d) Vasodilator & sedative
- NITROCHOLATE.** (Breon) Sod nitrite 1 gr, Sod glycocholate 1/10 gr, chionanthin 1/10 gr, iris 1/24 gr, paw ext echinacea 1/15 gr/Cap Bot 1000s
Use: Vasodilator.

NITRO-CYANATES. (First Texas) Sod. sulphocyanate 16 gr., Pot. sulphocyanate 8 gr., cascara sagrada fl. ext. aromatic 90 mg., Sod. nitrite 8 gr./oz. Elixir, Bot. 1 pt, 1 gal.

Use: (1 dr. b.i.d.) Arterial hypertension.

NITROFURAN, ORALLY. See: Alfatur, Tab. (Eaton)

***NITROFURANTOIN, U.S.P., N.N.D. 1962.**
1-[5 Nitro-2-furfurylideneamino]hydronitoin.
See: Furodanin, Prep. (Eaton)

***NITROFURAZONE, N.F., N.N.D. 1962;**
Oint., N.F. Sol., N.F. 5-Nitrofurazone-2-aldehyde semicarbazone.

See: Furacin, (Eaton)

Furacin Soluble Dressing.

Furacin Solution.

OB-GYN Furacin Cream (Eaton)

Use: Bacteriostatic for surface wounds.

W/Diperodon HCl. See: Furacin Urethral, Supp. (Eaton).

NITROGEN MONOXIDE, A.D.R. (Laughing Gas, Nitrous Oxide)

Use: Inhalation anesthetic & analgesic.

NITROGEN MUSTARD. See: Mustargen HCl (Merck Sharp & Dohme)

NITROGEN MUSTARD DERIVATIVES,
N.N.D. 1961.

See: Leukemia.

Leukeron, Tab. (Burroughs-Wellcome)

Mechlorethamine HCl, Sterile, N.F.

Mustargen HCl (Merck Sharp & Dohme)

Triethylene Melamine (Lederle)

NITROGLYCERIN, U.S.P. (Various Mfr.) (Glyceryl Trinitrate, Glonoin, Nitroglycerol, Trinitrin, Trinitroglycerol) Tab. 1/200 gr., 1/150 gr., 1/100 gr.

See: Glyceryl Trinitrate Tablets, U.S.P.

Niglycan, Tab. (CMC).

Nitrine-TDC (Confield)

Nitroglyn, Tab., S.A. (Key Corp.)

Nitrolawe, Tab. (Lowe)

Use: Vasodilator.

W/Digitalis comp. RX (DA COSTA) S.C.T. (Merck Sharp & Dohme) Nitroglycerin 0.6 mg., digitalis fr. 0.2 cc., strophanthus fr. 0.06 cc., belladonna tra. D 015 cc./tab. Bot. 100s.

U.S. Vasodilator.

W/Digitoxin, strychnine sulf. See: Unitroxin, Lir, (Tilden)

W/Hawthorn ext., pot. nitrate, sod. nitrite, powd. aconite root, veratrum viride ext., sod. bicarbonate. See: Natra Hawthorn Comp., Tab. (National Drug)

W/Homatropine methylbromide, ox bile ext., Mag. sulfate. See: Odi-Late, Tab. (Warren-Teed).

* Generic name recognized by N.F., U.S.P. or A.M.A.

W/Mannitol hexanitrate. See: Maxitate w/ Nitroglycerine, Tab. (Strassenburgh).

W/Nicotinic Acid. See: Nitrovas, Tab. (Amfre-Grant)

W/Pentaerythritol tetranitrate.

See: Bitrate, Tabs. (Arca)

Nitrol, Tab. (Kremers-Urban)

Vasotrate, Caps. No. 1 (Hiss)

W/Pentaerythritol tetranitrate and reserpine. See: Pente, Tab. (Carrick)

W/Pentaerythritol tetranitrate, butobarbital. See: Buta Penite, Tabs. Caps. (Carrick)

W/Pentaerythritol tetranitrate, secobarbital sod. See: Anti-Anginol, Tabs. (Lammon)

Nitrosed, Tabs (Lammon)

W/Pentaerythritol tetranitrate, Vit. B-1. See: Pentalar, Cap. (Sullivan & Case)

W/Sod. nitrite & crotaegus. See: Diotroegus, Pillel (Buffington)

W/Sod. nitrite, crotaegus fr., rutin & diallyl barbituric acid. See: Vasorutin, Pili (Buffington)

W/Sod. nitrate, labella & pilocarpine. See: Hipernine, Tab. (Westerfield).

W/Strychnine sulfate, strophanthin, digitoxin, cactus grandiflorus. See: Pasonol, Liq (Tilden)

W/Veratrum viride, Sod. nitrite, Pot. nitrate, crotaegus fr. & Pot. nitrate. See: Crotegonite, Tab. (C. D. Smith).

W/Veratrum viride, Sod. nitrite, Pot. nitrate, aconite root. See: Nyomin, Tab. (Elder).

NITROGLYCEROL. See: Nitroglycerin (Various Mfr.).

NITROGLYN. (Key Corp.) Nitroglycerin. Tab. sustained action 1/25 gr., 1/10 gr. Bot. 50s, 500s.

NITROL OINTMENT, (Kremers-Urban) 2%. Glyceryl trinitrate in lanolin base. Tube 1, 2 oz., Jar 1 lb.

Use: Coronary & peripheral vasodilator

NITROL TABLETS. (Kremers-Urban) Nitroglycerin 0.3 mg., pentaerythritol tetranitrate 15 mg./Tab. Bot. 100s

Use: Angina pectoris.

NITROLOWE CTR. (Lowe) Nitroglycerin U.S.P. 1/25 gr./Cap. Bot. 100s, 250r.

Use: Angina pectoris, coronary hypersensitivity, vascular conditions

NITROMANNITE, N.N.R. 1946. See: Mannitol Hexanitrate (Various Mfr.).

NITROMANNITOL. See: Mannitol Hexanitrate (Various Mfr.)

***NITROMERSOL, N.F., N.N.R. 1950, A.D.R.; Sol., N.F. Tr., N.F.** See: Metaphen, Prep. (Abbott)

W/Coal tar ext., quilla ext., benzoin &

- isopropyl alcohol. See: Keracide, Sol (Kor
mel Co)
- p-NITROPHENOL.** See: Niphen, Liq (Torch)
- NITROPIN.** (Fuller) Methylotropine nitrate
1/60 gr, phenobarbital 1/4 gr/Tab Bot
100s, 500s
Use: Antispasmodic
- NITROSCLERAN** (Tosse) Sod nitrite 100
mg, Sod & Pot phosphate/1 cc Amp 1
cc, Box 6s, 30s, 100s
Use: I.M.r essential hypertension & angina
pectoris
- ***p-NITROSULFATHIAZOLE, NF, NND**
1962 2 [p Nitrophenylsulfonamide]thiazolo
See Nisulfazole, Supp (Breon)
- NITROTALANS, Na. R65.** (Paul Maney)
Pentaerythritol tetranitrate 10 mg Bot
100s 500s
Use: Angina pectoris
W/Phenobarbital—No 866 Contains 15 mg
Phenobarbital in addition
- NITROTALANS UNODAY Ph. 755.** (Paul
Maney) Pentaerythritol tetranitrate 30 mg/
tab Bot 100s, 500s
Use: Angina pectoris
W/Phenobarbital Unoday Ph 756 (Paul
Maney) Pentaerythritol tetranitrate 30 mg,
phenobarbital 45 mg/tab Bot 100s, 500s
Use: Angina pectoris
- NITROUS ETHER** See: Ethyl Nitrite (Various
Mfr)
- NITROVAS** (Amfre Grant) Nitroglycerine
1/25 gr, nicotinic acid 35 mg/long acting
Tab Bot 60s, 120s
Use: Angina pectoris
- NIVAQUINE, See: Chloroquine Diphosphate**
(Arafen Ophosphate Winthrop Labs)
- NIVIA OIL.** (Duke) Emulsion of neutral oli-
phobic hydrocarbons Bot 4 fl oz, 1 pt,
1 qt
Use: Emollient
- NIZANTOL** (Normond) Vit D 30,000 u, A
5000 u, ascorbic acid 100 mg, thiamine
HCl 1.5 mg, riboflavin 2.7 mg, niacinamide
15 mg/Tab Bot 30s, 100s 1000s
Use: Vitamin therapy
- NOCTEC.** (Squibb) Chloral hydrate Sol 7 1/2
gr./5 cc Bot 1 pt Cap 7 1/2 gr, 3 3/4 gr
Bot 100s
Use: Natural sedation
- NO-DERM** (Lemmon) Neomycin sulfate 3.5
mg, dipiperdon hydrochloride 1%, alumi-
num d hydroxy allantoinate 0.75%/ml
Plastic Bot 30 ml
Use: Control infection in minor cuts and
abrasions
- NOION Liquid** (Barnes Hind) Naniamic,
nonsudsing nonirritating hand cleanser
Bot 4 oz, 8 oz
- *Generic name recognized by NF USP or A.M.A.
- Use: Cleanser for contact lens wearers and
practitioners
- NO-KOTIN.** (Walker Pharm) Labeline sulfate
1/16 gr./tab Tube 40s
Use: As aid to those who should or wish to
stop smoking
- NOLAMINE.** (Carrick) Chlorprophenpyrida-
mine maleate 4 mg, phenindamine tartrate
24 mg, phenylpropranolamine HCl 50 mg/
Tab Bot 50s
Use: Upper respiratory and nasal conges-
tion
- NOLAMINE ELIXIR.** (Carrick) Chlorpro-
phenpyridamine maleate 1 mg, phenindo-
mine tartrate 6 mg, phenylpropranolamine
HCl 12.5 mg, sorbitol 1 Gm/5 cc Bot
pt
Use: Nasal congestion
- NOLAMINE EXPECTORANT SYRUP.** (Carr-
rick) Glyceril guaicolate 50 mg, dihydro-
codeinone bitartrate 1.7 mg, chlorphenir-
amine maleate 0.67 mg, phenindamine
tartrate 4 mg, phenylpropranolamine HCl
8.35 mg/5 cc Bot 16 oz
Use: Cough due to colds
- NOLOC** (Dumas Wilson) Diacetylhydroxy-
phenylsatin 2 mg, dioctyl sod sulfosuc-
cinate 50 mg/Cop Bot 100s 1000s
Use: Laxative
- NOLUDAR, NND 1961** (Roche) Methyl
prylon, 3,3 diethyl 5 methyl 2,4 piperidine
dione, Tab (50 & 200 mg) Bot 100s, 500s
1000s Elixir (50 mg/5 cc) Bot 1 pt, 1
gal Caps (300 mg) Bot 100s 500s
Use: Relief of nervous insomnia & daytime
tension
- ***NONAPYRIMINE.** (Burroughs Wellcome) 4
n nanylamina 7H pyrrolo [2,3-d] pyrimidine
Use: Anticonvulsant
- NONARCOL** (Buffingtons) Pyrilamine male-
ate 15 mg, phenylephrine HCl 5 mg, glyc-
eryl guaicolate 50 mg, potassium citrate
250 mg, citric acid 50 mg, chloroform 1/4
minim menthol 1 mg/5 cc Syr
Use: Expectorant
- NONSPECIFIC PROTEIN THERAPY** See
Protein, Nonspecific Therapy
- ***NORACRYMETHADOL NCI** (Eli Lilly) α -4,4-
diphenyl 6 methylamino 3 heptanol acetate
Use: Analgesic agent
- NORADRENALIN** See: Levarterenal Bitar-
trate, USP
- ***NORDEFRIN HYDROCHLORIDE, NF ADR**
1,3,4 Dihydroxynorephedrine See Cobefrin
(Winthrop Labs)
- NORDIL** (Warner Chilcott) beta Phenylethyl
hydrazine dihydrogen sulfate

NORDIOL (Normond) Estradiol 0.125 mg / Buccal. Bot. 30s, 100s, 500s, 1000s
 Use: Menopausal disturbances

dl-NOREPHEDRINE HYDROCHLORIDE
 (alpha-hydroxy-beta-aminopropylbenzene HCl)

See: Phenylpropanolamine Hydrochloride.
 (Merck Sharp & Dohme)

Propadrine HCl, (Merck Sharp & Dohme)

l-NOREPINEPHRINE BITARTRATE, U.S.P., N.N.R., A.D.R., N.N.D. 1962.

See: Levarterenol Bitartrate, U.S.P.

Levophed, Sol. (Winthrop Labs.)

***NORETHANDROLONE, N.N.D. 1962. 17-Alpha-ethyl-17-hydroxynandrostane.**

See: Nilevar, Tab. (Searle)

***NORETHINDRONE, N.N.D. 1962.** See: Norlutin, Tab. (Parke, Davis)

***NORETHINDRONE ACETATE.** See: Norlutale (Parke, Davis) W/Ethinyl estradiol See: Gestest, Tabs. (Squibb)

NORETHISTERONE. See: Norethindrone.

***NORETHYNODREL, 17-hydroxy-19-nor-17 α -pregn-5(10)-en-20-yn-3-one** See: Enovid (Searle).

NOREXIN. (Smith-Dorsey) Vit. B-1 0.67 mg., B-2 0.67 mg., niacin 6.7 mg., d-amphetamine sulfate 5 mg., phenobarbital $\frac{1}{4}$ gr., methylcellulose 200 mg., Mag. trisilicate 200 mg./Cap. Bot. 100s, 500s, 1000s.

Use: Anti-obesity.

NORFLEX. (Riker) Orphenadrine citrate 100 mg./Tab. Bot. 50s

Use: Skeletal muscle relaxant.

NORFLEX INJECTABLE. (Eiker) Orphenadrine citrate 60 mg., sodium bisulfite 2 mg., sodium chloride 5.8 mg. for injection qs 2 cc. 2 cc. amp/6s; 2 cc. amp/50s.

Use: Relief of skeletal muscle spasm.

NORISODRINE SULFATE, N.N.D. 1962 (Abbott) Isopropylarterenal sulfate. Sol. T:100, Bot. 10 cc.

Use: (Oral inhalation) Bronchial antispasmodic.

NORISODRINE SULFATE FOR ORAL INHALATION 10% or 25% IN AEROSOL CARTRIDGE, N.N.R. 1953. (Abbott) Isopropylarterenal sulfate. Vial 3 sifter cartridges.

Use: Oral Inhalation; bronchial spasm.

NORISODRINE SYR. (Abbott) Isoproterenol sulfate 1B mg., anhydrous calcium iodide 900 mg., alcohol 6%/30 ml. Bot. pl.

Use: Asthma conditions.

NORITAL. (Normond) Hyoscyamine HBr 0.12B mg., atropine sulfate 0.024 mg.

* Generic name recognized by N.F., U.S.F. or A.M.A.

hyoscyine HBr, 0.0072 mg., phenobarbital 16 mg./tab. Bot. 100s, 500s, 1000s.

Use: Antispasmodic.

NORLEYORPHANOL. (-)-3 hydroxymorphinan.

NORLUTATE. (Parke, Davis) Norethindrone acetate 5 mg./Tab. Bot. 30s.

Use: Amenorrhea, menstrual irregularity.

NORLUTIN. 1962. (Parke, Davis) Norethindrone. (17-alpha ethinyl-19-nortestosterone) 5 mg./Tab. Bot. 30s.

Use: Control of many menstrual disorders (1 to 2 tab. twice daily)

NORMACAP. (Padiatrix) Bioflavonoids, pectinates, vitamin C for podiatrists/chiropractors Tabs Bot. 90s.

Use: Abnormal capillary permeability.

NORMACID. (Stuart) Betaine HCl 440 mg., pepsin 32.4 mg., methylcellulose 110 mg./Tab. Bot. 100s.

Use: Hydrochloric acid therapy.

NORMADRINE IMPROVED. (Vanpelt & Brown) Not listed. dl-Desoxyephedrine HCl

1% isotonic sol. Bot. w/dropper, 1 fl. oz. Bot. 1 pl., 1 gal.

Use: Nasal decongestant.

NORMAL HUMAN PLASMA, U.S.P., N.N.D. 1961.

NORMAL HUMAN SERUM, U.S.P., N.N.D. 1961.

NORMAL HUMAN SERUM ALBUMIN, U.S.P., N.N.D. 1962. See: Albumisal 25% Conc. (Merck Sharp & Dohme)

NORMAL SERUM ALBUMIN (HUMAN) SAET-POOR. (Hyland Labs) Albumin fraction 25% from pooled, normal human plasma. Bot. 50 cc. Sol. (5%), Bot. 250 cc.

Use: I.V. shock therapy.

***NORMETHANDRONE.** 17- α -methyl-19-nortestosterone. See: Methalulin (Parke, Davis)

NORMIC. (Spencer) Dihydroergocornine 0.3 mg., phenyltoloxamine dihydrogen citrate 50 mg., hyoscyamine sulfate 0.1 mg./Tab.

Use: Treatment of psychophysiological disorders.

NORMOCYTIN. (Lederle) Vit. B-12. Tab. 30 mcg. Bot. 100s. Vial (30 mcg./cc) 10 cc.; (60 mcg./cc) 10 cc.

Use: I.M. & orally; hematopoietic.

NORODIN HCl, N.N.D. 1962 (Methamphetamine HCl) d-desoxyephedrine HCl Pow., Bot. 1 Gm., 5 Gm., 10 Gm., 20 Gm. Tab. (2.5 & 5 mg) Bot. 100s, 1000s.

Use: Orally; central nervous system stimulant.

NORPHRITICA. (Tilden) Sod. bromide 12 gr., fluidextract hyoscyamus 4 min., juniper berries 24 gr., liverwort 16 gr., uva ursi 16

- gr, pot ocelote 16 gr, buchu 9 gr Lithium citrate 5½ gr/fl oz Bot pts, gals
Use Diuretic urinary tract sedative
- NORPRAMIN** (Lakeside) Desipramine Hydrate
Use antidepressant
- NORTESTERONATE** 19 nortestosterone cyclopentylpropionate See Depo Nortestonate (Upjohn)
- 19-NORTESTOSTERONE-17-CYCLOPENTYLPROPIONATE** See Depo Nortestonate (Upjohn)
- ***NORTRIPTYLIN HCl** (Eli Lilly and Co) 5 (α methylaminopropylidene) dibenzo [a,d] cyclohepta [1,4] diene hydrochloride See Anenlyl
- NOSCAP**, (Tablet) Noscapine 7.5 mg chlorpheniramine maleate 1 mg, phenylephrine HCl 5 mg, N acetyl p aminophenol 150 mg, salicylamide 150 mg Vit C 20 mg/cap
Use Relief of common cold
- ***NOSCAPINE**, NF, NND 1962 (Merck Sharp & Dohme) Narcotone Noscapine Bulk powder See Nectadon (Merck Sharp & Dohme)
Use: Antitussive
- ***NOSCAPINE HCl**, NF
W/N Acetyl p aminophenol phenylpropanolamine glyceryl guaiacolate See Coryz Tab (Marion)
- W/Chlorpheniramine maleate, phenylephrine HCl, N acetyl p-aminophenol, salicylamide Vit C. See: Noscap, Cap (Tablet)
- W/Chlorpheniramine maleate phenylephrine HCl, ammonium chloride menthol See Noscamel Compound (Squibb)
- W/Glyceryl guaiacolate, alcohol See Coryz A T Susp (Marion)
- W/Glyceryl guaiacolate citric acid, alcohol See: Noscatuss Syr (Buffington)
- W/Pheniramine maleate, phenylephrine HCl chloroform menthol See: Conar Liq, Troches (Massengill)
- W/Pheniramine maleate phenylephrine HCl glyceryl guaiacolate, chloroform menthol See: Conar Expectarant Liq (Massengill)
- W/Phenylephrine acetylsalicylic acid, N acetyl p aminophenol caffeine, pyrilamine maleate, methopirylene fumarate See: Tusso-pap, Tab (Swift & Case)
- W/Phenylephrine HCl pyrilamine maleate glyceryl guaiacolate See: Guaahist W/ Narcotone, Cap (Columbus)
- W/Phenylpropanolamine HCl, chlorpheniramine maleate, acetaminophene glyceryl guaiacolate ammonium chloride See: Mycatussie, Syr (Myers-Cortez)
- * Generic name recognized by NF, USP or A.M.A.
- W/Phenylpropanolamine, glyceryl guaiacolate acetaminophene, chloroform AR See: Coryz Suspension, Liq (Marion)
- W/Phenyltoloxamine, NH₄Cl in a thixotropic suspension See Narcol, Liq (Barrs)
- W/Salicylamide phenylephrine HCl, glyceryl guaiacolate See Noscoline Tab (Chicago Pharmacol)
- W/Terpin hydrate, doxylamine succinate See: Consol, Susp (Merrill)
- W/Terpin hydrate sodium citrate, pyrilamine maleate alcohol See: Nektussin (Bruce)
- NOSCATUSS**, (Buffington) Noscapine 15 mg glyceryl guaiacolate 50 mg, citric acid 50 mg, alcohol 5%/5 cc Syr Bot 1 pt, 1 gal
Use Antitussive and expectorant action
- NOSCOLINE**, (Chicago Pharmacol) Noscapine 5 mg, salicylamide 325 mg, phenylephrine HCl 12.5 mg, glyceryl guaiacolate 12.5 mg/tab Bot 50s, 100s, 1000s
Use: Control of cough
- NOSCOMEL COMPOUND** (Squibb) Noscapine 10 mg chlorpheniramine maleate 0.66 mg phenylephrine HCl 1.66 mg, ammonium chloride 50 mg menthol 0.25 mg/ tsp Bot 3 oz
Use Antitussive
- NOSCOSED** (Chicago) Chlorpheniramine maleate 13 mg narcotine 5 mg, with terpin hydrate/5 cc Bot 4 oz, 1 pt
Use Cough therapy
- NOSE DROPS-SPRAY**, (Johnson & Johnson) Phenylephrine HCl ¼% Liq
Use Nasal decongestion
- NOSTYN**, NND 1962 (Ames) Ectylurea 2 Eryth cis cratonylurea 300 mg/Tab Bot 4s 500s
Use Apprehension and tension
- NOVADONNA**, (Sherman) Homatropine methylbromide 1.512 mg, hyoscyamine HBr 0.283 mg, hyoscine HBr 0.019 mg/cc Bot 4 oz
Use: 6 mie orally; antispasmodic
- NOVADONNAL**, (Sherman) Homatropine methylbromide 0.567 mg, phenobarbital ¼ gr, hyoscyamine HBr 0.106 mg, hyoscine HBr 0.007 mg/Tab Bot 100s, 1000s
Use: (1-2 qid) Antispasmodic w/sedative
- NOVAHISTINE-DH** (Pitman Moore) Phenylephrine hydrochloride 10 mg, chlorprophepyridamine maleate 20 mg, calcium phosphate 10 mg, chloroform 13.5 mg; 1 menthol 10 mg/Alcohol 5%; sugar content 33½% Elix Bot 1 pt, 1 gal
Use: 2 dr tid; coughs & respiratory congestion
- NOVANISTINE-OH TAB** (Pitman Moore) Phenylephrine HCl, 20 mg; chlorpheniramine

- pyridomine maleate, 4 mg.; dihydrocodeinone bitartrate, 5 mg./Tab. Bot. 100s.
 Use: Relief from cough and congestion.
- NOVANISTINE ELIXIR.** (Pitman-Moore) Phenylephrine HCl 5 mg., prophenpyridamine maleate 12.5 mg., chloralform 13.5 mg., menthol 1 mg./5 cc. Bot. 1 pt., 1 gal.
 Use: Oral; antihistamine w/sympathomimetic agent.
- NOVAHISTINE EXPECTORANT.** (Pitman-Moore) Phenylephrine hydrochloride 10 mg.; chlorophenpyridomine maleate 2 mg.; codeine phosphate 10 mg.; Glyceryl guaiacolate 100 mg.; chloralform 13.5 mg.; 1-Menthol 1.0 mg Bot. pt., gal.
- NOVAHISTINE FORTIS.** (Pitman-Moore) Phenylephrine HCl 10 mg., prophenpyridomine maleate 12.5 mg./Cap. Bot. 36s, 500s.
 Use: (2 t.l.d.) Antihistaminic.
- NOVAHISTINE L.P.** (Pitman-Moore) Phenylephrine HCl 20 mg., chlorophenpyridomine maleate 4 mg./Tab. Bot. 50s.
 Use: Common cold.
- NOVAHISTINE SINGLET.** (Pitman-Moore) Phenylephrine hydrochloride 40 mg., chlorophenpyridomine maleate 8 mg., n-acetyl-para-aminophenol 500 mg./Tab. Bot. 50s.
 Use: Antihistamine-decongestant
- NOVAHISTINE TABLETS.** (Pitman-Moore) Phenylephrine HCl 5 mg., prophenpyridomine maleate 12.5 mg./Tab. Bot. 100s, 1000s.
 Use: Sympathomimetic & antihistamine.
- W/A.P.C.** (Pitman-Moore) Phenylephrine HCl 10 mg., prophenpyridomine maleate 12.5 mg., aspirin 3½ gr., phenacetin 2½ gr., caffeine ½ gr./cap. Bot. 24s, 100s.
 Use: (2 t.l.d.) Treatment of the common cold
- W/Penicillin.** (Pitman-Moore) Phenylephrine HCl 10 mg., prophenpyridomine maleate 12.5 mg., penicillin G pot. 200,000 u/cap. Bot. 24s.
 Use: Orally, treatment of respiratory congestion.
- NOVALDIN.** (Winthrop Labs.) Dipyrone, Navalgin, Phenyl dimethylpyrazolone Methylaminomethone Sod. Sulfonate. Amp. 50%, 2 cc., Box 5s, 50s. Tab. 5 gr., Bot. 100s.
 Use: Analgesic & antirheumatic.
- NOVALINE TABLETS.** (Lemman) Phenobarbital ¼ gr., ephedrine sulfate ¾ gr., Pot. iodide 2½ gr., Col. lactate 2½ gr./Tab. Bot. 25s, 100s, 500s, 1000s.
 Use: Asthma & hay fever.
- NOVALGIN.**
 See: Dipyrone. (Various Mfr.)
- NOVAMIDON.** See: Aminopyrine (Various Mfr.).
- NOVAPAN.** (Brewer) Vit. A 5000 u., D 1000 u., C 50 mg., B-1 1 mg., B-2 0.5 mg., B-6 0.6 mg., niacinamide 17 mg., pantothenol 1.9 mg./0.6 cc. Bot. 15 & 50 cc. w/graduated dropper.
 Use: Orally, vitamin deficiencies.
- NOVAPAN GEL-ETS.** (Brewer) Vit. A 5000 u., D 1140 u., B-1 1.5 mg., B-2 2 mg., B 6 1 mg., C 30 mg., niacinamide 10 mg., col. pantothenate 3 mg, mixed tocopherals 1.33 mg./Gel-El. Bot. 100s, 500s, 1000s.
 Use: Multiple vitamin therapy.
- NOVASENOBENZOL.** N.N.E. 1936. See: Neocarsphenamine, Amp. (Abbott)
- NOVATOPHAN.** See: Neocinchaphen (Various Mfr.).
- NOVATIN.** N.N.R. (Campbell) Homatropine methylbromide, Amp. (5 mg) 1.1 cc., 5s, 100s. Pediatric cal. 1/50 gr./cc. Bot. 1 oz. Pow., Vial 15 gr. Tab. 1/24 gr., 1/12 gr., Bot. 100s, 500s.
 Use: Antispasmodic & sedative.
 W/Methylcellulose. See: Celulrapin, Liq. (Campbell)
- W/Phenobarbital.** (Campbell) Homatropine methylbromide 1/12 gr., phenobarbital ¼ gr./Tab. Bot. 40s, 250s, 1000s.
 Use: (1 q. 3-4 h. not to exceed 6 dolly) Antispasmodic & sedative
- NOVATIN COMPOUND (PEDIATRIC).** (Campbell) Homatropine methylbromide 1.3 mg., phenobarbital 16 mg/cc Bot. 1 oz.
 Use: Treatment of colic and vomiting.
- NOVATROPINE.** See: Homatropine Methylbromide (Various Mfr.).
- NOVA-TUSSA SYRUP.** (Drug Specialties) Dihydrocodeinone bitartrate 10 mg., pheniramine maleate 30 mg., pyrilamine maleate 30 mg., phenylephrine HCl 30 mg., polysium guaiacolsulfonate 500 mg./fl. oz. Bot. pts., gals.
 Use: Cough and cold therapy.
- NOVESTROL.** (Borer) Ethinyl estradiol 0.02 and D05 mg./Tab. Bot. 100s, 1000s
- NOVIPLEX CAPSULES.** (Massengill) Vit. B-1 50 mg., Vit. B-2 2.5 mg., niacinamide 25.0 mg., Vit. B-6 0.5 mg., calcium pantothenate 1 mg., Vit. B-12 1 mcg./Cap. Bot. 100s, 500s, 1000s
 Use: Vitamin B Complex
- NOVIPLEX ELIXIR.** (Massengill) Vit. B-1 3.33 mg., Vit. B-2 1 mg., Vit. B-6 0.33 mg., pantothenic acid 0.83 mg., nicotinamide 12.5 mg., Vit. B-12 2.5 mcg./5 cc. Elix. Bot. 8 oz., 1 gal.
 Use: Dietary supplement

NOVO-BASIC, (Squibb) Vit C 150 mg, B 1 5 mg, B 2 5 mg, niacinamide 50 mg, B 6 1 mg, cal pantothenate 10 mg, B 12 2 mcg, folic acid 0.15 mg / Tab Bot 60s, 180s

Use: Nutritional therapy

***NOVOBIOCIN**, Streptanivacin (Biatexin) An antibiotic from *Streptomyces niveus* and *Streptomyces spheroides*

See Albamycin (Upjohn)

Cardelmycin (Pfizer)

Cathomycin (Merck Sharp & Dohme)

Dio Oxcs (Reed & Carnrick)

W/Panmycin See Panalba, Cap (Upjohn)

***NOVOBIOCIN CALCIUM**, USP, NNO 1962 Oral Susp USP

See Albamycin, Syr (Upjohn)

Cathomycin Calcium Cap (Merck Sharp & Dohme)

W/Tetracycline HCl potassium metaphosphate See Panalba KM Drops (Upjohn)

***NOVOBIOCIN SODIUM**, USP, NNO 1962 Cap USP

See Albamycin, Cap (Upjohn)

Cathocillin, Cap (Merck Sharp & Dohme)

Cathomycin Sodium Lyovac, Vial (Merck Sharp & Dohme)

NOVOCAIN, NNR 1950, ADR (Winthrop Labs) Procaine HCl

Amp: (50, 100, 120, & 150 mg) 10s, 100s; (200, 300, 500 & 1000 mg) 10s

Pow: Bot 1 oz

Sol: (0.1%) Bot 1000 cc (1%) 2 cc, 10s, 100s; (6 cc) 50s; Bot 30 & 100 cc (2%)

3 cc, 10s, 100s; Bot 1 oz, 100 cc (10%)

2 cc, 100s (20%) 5 cc, 10s, 100s

Tab: (0.2 Gm) Bot 100s, 500s; (0.05 Gm) Bot 1000s

See Spinocaine (Winthrop Labs)

NOVOCAIN-PONTOCAINE HYDROCHLORIDE SOLUTION (Winthrop Labs) Novocain HCl 100 mg, pontocaine HCl 10 mg / 2 cc, Amp 2 cc, Box 10s

Use: Spinal anesthesia

See also Spinocaine (Winthrop Labs)

NOVOCAIN-SUPRARENIN (N.S.) (Winthrop Labs)

Amp: E (Novocain 2%, suprarenin 1:20,000) 3 cc, Box 10s, 100s K (Novocain 1%, suprarenin 1:50,000) 2 cc, Box 10s 100s;

6 cc, Box 50s T (Novocain 2%, suprarenin 1:50,000) 3 cc, Box 10s, 100s Conc. (Novocain 20%, suprarenin 1:9000) 1.5 cc, Box 10s; 5 cc, Box 100s

Sol: (Novocain 1%, suprarenin 1:100,000) Bot 1 fl.oz; (Novocain 2%, suprarenin 1:50,000) Bot 1 fl.oz

Tab: "A" (Novocain 0.125 Gm, suprarenin 0.000125 Gm) Bot 100s "B" (Novocain 0.1 Gm, suprarenin 0.00025 Gm) Bot 100s "K" (Novocain 0.06 Gm, suprarenin 0.00006 Gm) Bot 100s "T" (Novocain 0.02 Gm, suprarenin 0.00002 Gm) Bot 100s

NOVOCELL, ADR (Novacol) Oxidized cellulose w/phenyl mercuric nitrate 0.08 mg / pellet Vial 50s

Use Hemostatic in dental surgery

NOVOCILLIN, (Novacol Chem Co) Pot penicillin G 20,000 u/Gm Oint, Tube 1/2 oz

Use Vincent's infection

NOVOGRAN, (Squibb) Vit B 1 10 mg, B 2 10 mg, B 6 2 mg, C 300 mg, niacinamide 100 mg pantothenic acid 20 mg, Vit B-12 4 mg, folic acid 1.5 mg, menadione 2 mg / Cap Bot 30s, 100s, 500s

Use Vit B complex therapy

NOVOGRAN FOR SOLUTION, (Squibb) Sod ascorbate 300 mg, thiamine HCl 5 mg, riboflavin 5 mg, niacinamide 100 mg, pantothenic acid 20 mg, pyridoxine HCl 2 mg, folic acid 1.5 mg, Vit B 12 1 mcg / cc Vial 2 cc, lyophilized w/2 cc of diluent 7-dose units, Pkg 5s

Use: IV or IM

NOVOGRAN 2X FOR SOLUTION (Squibb) Twice Novogran ingredients in Vial 5 cc lyophilized w/4 cc of diluent

Use: IV or IM

NOVRAD (Lilly) Levopropoxyphene 50 mg / 5 cc Susp Bot 4 Oz 50 mg or 100 mg / Cap Bot 50s

Use Antitussive

NOVRAD W/A.S.A (Lilly) Levopropoxyphene 50 mg, acetylsalicylic acid 300 mg / Pulvule Bot 50s

Use: Antitussive & analgesic

***NOXYTHIOLIN**, N-Hydroxymethyl N'-methylthiourea

NOZAIN (Naxemo) Cyclofarm, octamer, cat bolic acid in a greaseless cream Pkg 1 1/4 oz 2 1/4 oz

Use: First aid cream

NPH Iletin, NND 1962 (Lilly) (Isophane Insulin NPM Inselin) Vial 40 u, 80 u, 10 cc

Use: S.C. management of diabetes

NPH INSULIN, USP, NND 1962 (Squibb) Vial 40 u, 80 u, 10 cc See Insulin Injection Isophane, USP

N.R.C. THERAPEUTIC VITAMINS (West Ward) Vit B 1 10 mg, B 2 10 mg, niacinamide 100 mg, Cal pantothenate 20 mg, Vit B-6 2 mg, folic acid 1.5 mg, Vit C 300 mg, B-12 4 mcg / Cap Bot 50s, 500s

Generic name recognized by N.F., USP or A.M.A.

NTZ NASAL SOLUTION. (Winthrop Labs.)
Neosynephrine 0.5%, thantadil HCl 0.1%,
zephiran chloride 1:5000. Bot. 30 cc., w/
dropper.

Use: Allergic rhinitis.

NUALETS. (Abbott) Calcium as col. lactate
100 mg., Vit. A 1333 u., D 66.6 u., B-1
0.25 mg., B-2 0.5 mg., nicotinamide 2.5
mg., C 2.5 mg., B-6 0.25 mg., B-12 0.5 mcg.,
folic acid 0.2 mg., cobalt 0.05 mg., copper
0.5 mg., iodine 0.075 mg., iron 2.5 mg.,
magn. 3 mg., mang. 0.5 mg., molybd. 0.033
mg., pot. 2.5 mg., zinc 0.6 mg./Filmstab
Bot. 100s, 500s, 1000s

Use: Dietary supplement during pregnancy
and lactation

NUBESF. (Dumas-Wilson) Amphetamine zela-
tate 5 mg., thyroid 16 mg./Tab. Bot. 100s,
1000s.

Use: Anorectic.

NUBIKIRK. (Irwin, Neisler) Vit. B-1 6 mg.,
B-2 3 mg., niacinamide 24 mg., pantothenic
acid 6 mg., yeast 2.25 Gm./15 cc. Bot.
1 pt., 1 gal.

Use: Nutritional supplement.

NUBILAX TABLETS. (Numaloxine) Dehydro-
cholic acid 2 gr., cascara sagrada pow. ext.
2 gr./Tab. Bot. 25s, 50s, 100s.

Use: Orally (2 m p.), laxative.

NUBILIC. (Hobart Lab.) Dehydrocholic acid
0.25 Gm., phenobarbital 8 mg., belladonna
pow. 8 mg./Tab. Bot. 25s, 50s, 100s.

Use: Hydrocholeretic, antispasmodic &
sedative.

NUCITE. See: Inositol (Various Mfrs.)

NUCLEIC ACID ANTAGONISTS.

See: Leukeran (Burroughs Wellcome)

Mustorgen (Merck Sharp & Dahme)

Myleran (Burroughs Wellcome)

Purinethal (Burroughs Wellcome)

Triethylene Melamine (Lederle)

NUCLOMIN. (Baylor) Vitamins and minerals.
Tab. Bot. 100s.

Use: Dietary supplement.

NUCLON. (Smith, Kline & French) Thényl-
pyromine fumarate 37.5 mg., dextroamphetamine
sulfate 1.25 mg., acetylsalicylic acid
162 mg./Cap. Bot. 500s

Use: Common cold.

NUCODAN. (Endo) Dihydrohydroxycodolone
HCl 4.5 mg., dihydrohydroxycodolone
terephthalate 0.38 mg., homatropine tereph-
thalate 0.38 mg., pentylene tetrazol 50 mg./
Tab. Bot. 100s, 1000s

Use: (1 q. 6 h.) Analgesic.

NUCORSAL. (C. B. Kendall) Sod. salicylate
5 gr., p-aminobenzoic acid 5 gr., ascorbic
acid 1/6 gr./E.C. Tab. Bot. 100s, 500s.

Use: Antirheumatic.

NUDOX. (Ulmer) Thiamine HCl 100 mg.,
pyridoxine HCl 100 mg./cc. Vial 10 cc.

Use: I.M., I.V., nausea of pregnancy, radia-
tion sickness, alcoholism.

NU-ESTRO. (Caleway) #0 Estradiol benza-
late 0.025 mg., diethylstilbestrol 0.045 mg.;
#1 Estradiol 0.5 mg., diethylstilbestrol 0.9
mg.; #2 Estradiol 2.5 mg., diethylstilbestrol
4.5 mg.; #3 estradiol 5.0 mg., diethyl-
stilbestrol 9.0 mg.; #0, #1 Bot. 100s,
1000s; #2, #3 Bot. 36s, 100s, 1000s

Use: Estrogen therapy.

NUMESTORAL. (Organon) Ethisterone 15
mg., hesperidin complex 175 mg., ascorbic
acid 175 mg., Vit. K analogue 2 mg., di-
alpha-tocopherol acetate 3.5 mg./Tab. Pkg.
30s

Use: (I t.i.d.) Treatment of abortion

NUJOL. (Plough) liq. petrolatum. Bot. 1/4 pt.,
1 pt., 1 qt., 1 gal.

Use: Laxative.

NU-KOL-TAR. (Benet) Crude coal tar 3%
or 5%, in Ruggles cream base. Cream, Jar
2 oz., 4 oz., 8 oz., 1 lb. Lot. 1%, 2%,
3%, 5%. Bot. 4 oz.

Use: Eczema, pruritis & antipruritic.

NULABORT. (Dumas-Wilson) Diethylstilbes-
trol 25 mg., hesperidin methyl chalcone 100
mg., ascorbic acid 100 mg., menadiolone
2 mg./Tab. Bot. 100s, 1000s.

Use: (1 or 2 q.i.d.) Treatment of threatened
abortion

NULACIN. (Horlick) Mag. trisilicate 3.5 gr.,
Mag. oxide 2 gr., Col. carb. 2 gr., Mag.
carb. 0.5 gr., peppermint oil q.s./Tab.
Bot. 25s.

Use: Antacid in peptic ulcer.

NULLAPONS. (General Aniline & Film Corp.)
The whole group of chelating agents related
to ethylenediaminetetraacetic acid. Pkg. ac-
cording to demand. See: Edathamil

Use: Sequestering agent.

NULLICOF. (Naves) Dihydrocodolone bitar-
trate 1/36 gr., ammonium chloride 3 1/2 gr.,
chloroform 1/6 min. in aromatized syr. of
lobelia & tolu/dr. Bot. 1 pt., 1 gal.

Use: Coughs.

NULLO. (Depree) Chlorophyll. Tab., Bot. 15s,
45s, 100s.

Use: Orally (1-2 dolly), body odors.

NUMA DURA-TABS. (Wynn) Theophylline
3 1/2 gr., ephedrine HCl 1/4 gr., butabarbital
1/2 gr./Tab. Bot. 30s, 100s, 250s.

Use: Bronchial asthma

NUMAL. (United Chemicals) Aprabarbital
Pow., Bot. 4 1/2 oz.

Use: Sedative & hypnotic.

NUMEROIDAL SUPPOSITORIES. (Hobart

- Lab) Ephedrine HCl 0.22%, benzocaine 5% Box 12s
 Use Vasoconstrictor w/ anesthetic
- NUMORPHAN**, NND 1962 (Endo) Oxy morphone (14 hydroxydihydromorphinan) HCl 1 mg/cc amps 1 cc & 0.5 cc Box 12s, 100s Vials 15 cc Box 1 & 3s 15 mg/cc Vial 10 cc Amp 1 cc, 2 cc suppository 2 mg or 5 mg Box 6s
 Use Relief of pain
- NUMOTIZINE**, Catoplasin (Hobart Lab) Guaiacal, beechwood creosote, methyl salicylate, formaldehyde, glycenn & aluminum silicate Jar 4 oz, B oz, 15 oz, 30 oz
 Use Analgesic & decongestive
- NUMOTIZINE COUGH SYR.** (Hobart Lab) Glyceril guaiacolate 5 gr, amm chlor 5 gr, sod cit 20 gr, menthol 0.4 gr/fl oz Bot 3 oz
- NUMOTIZINE**, (Hobart) Phenylephrine HCl 5 mg, acetaminophenol 150 mg, pyrilamine maleate 7.5 mg, caffeine 15 mg, vit C 15 mg/Tab Bot 25s 100s
 Use Nasal congestion
- NUMOTIZINE Nasal Spray.** (Hobart) Polymixin B sulfate 0.009% hydrocortisone 0.02%, phenylephrine HCl 0.25%, pyrilamine maleate 0.1%, benzalkonium chloride 0.1% in a phosphate buffered base with 1% sodium bisulfite and 2% Tween 20 Spray bats 15 cc
 Use: Common cold symptoms
- NUNOL** See Phenobarbital (Various Mfr)
- NUPERCAINAL CREAM** (Ciba) Nupercaine 0.5% in water washable base Tube 1 1/2 oz
 Use: Topically; local anesthetic
- NUPERCAINAL OINTMENT.** (Ciba) Nupercaine 1% in lanolin & petrolatum Tube 1 oz Jar 1 lb
 Use: Topically; local anesthetic
- NUPERCAINAL SUPPOSITORY.** (Ciba) Nupercaine 25 mg, zinc oxide, and bismuth subgallate in a cocoa butter base/Sepp Box 12s
 Use: Treatment of pruritis ani, and hemorrhoids
- NUPERCAINE HYDROCHLORIDE**, NNR 1956 (Ciba) Dibucaine HCl Amp. Sols 1,200, 2 cc, Carton 10s; 1,500, 20 cc Carton 12s; 1,400 w/dextrose 5% (heavy soln), 2 cc, Carton 10s Bot Sols 2%, 1 fl oz, 4 fl oz Tab 1 D05 Gm Tube 20s
 Use: Local anesthetic
- W/Pyribenzamine HCl combination See, Tricainol, Oint & Supp (Ciba)
 W/Quinine HCl benzyl alcohol See, Anesacaine (New Mexico Pharm)
- NUPERCAINE, PHENOL & BENZYL ALCOHOL IN OIL.** (Ciba) Nupercaine HCl 0.5%, phenol 1%, benzyl alcohol 10%. Amp 5 cc, Carton 10s
 Use Topically; proctology
- NUPORALS.** (Ciba) Dibucaine HCl 1 mg/loz, Box 15s, 100s, 500s
 Use Orally; local anesthetic
- NUPRA.** A nonharmonic, nonsteroidic extract of beef organs, liver, brain adrenals See Exual Tab (Yorklawn Prod)
- NUPRIN** (Upjohn) Sulfamoxale
 Use Antibacterial sulfonamide
- NUROBROMYDS.** (Rorer) Sod bromide 20 gr, strontium bromide 20 gr, ammonium bromide 20 gr, valerian tr 60 min, hyocyamus tr 60 min/fl oz Bot 14 fl oz, 5 pt, 1 gal
 Use: Sedative & antispasmodic
- NURODOL.** (Bullington) Dialyl borbituric acid 1/2 gr, Pot bromide 20 gr, Sod bromide 20 gr, ammonium bromide 20 gr, alcohol 10%/fl oz Bot 1 pt, 1 gal
 Use: Sedative
- NUROKARDIAC.** (Rorer) Phenobarbital 1/4 gr, nux vomica ext 1/4 gr, carbenzyl 4 gr/Tab Bot 100s, 500s, 1000s
 Use Sedative & digestant
- NU-THERA.** (Kirkman) Contains 9 vitamins & 11 minerals/cap Bot 100s, 1000s 5000s
- NUTRACHLORIC.** (Upjohn) Skimmed milk pow 48 gr, Cal carb 11 1/2 gr, Sod bicarb 1 1/4 gr/Tab Box 48s
 Use Gastric hyperacidity
- NUTRALKA.** (Endo) Bismuth subcarb 1/2 gr, Mag carb 2 gr, Cal carb 3 1/2 gr, distase 1/40 gr papain 1/20 gr, aromatics qt/Tab Bot 100s, 500s, 1000s
 Use Antacid & digestant
- NUTRAMENT LIQUID** (Edward Dalton) Protein 20 gr fat 13.3 gr, carbohydrates 50 gr and 400 calories per 12 1/2 oz can Can 12 1/2 oz
 Use Nutritional supplement
- NUTRAMIGEN.** (Mead Johnson) Dextrin maltose 45.2%, amigen 20%, corn oil 18% arrowroot starch 10%, Cal gluconate 3.5% w/Pot phosphate, Pot dibasic phosphate Cal hydroxide, Pot chloride, Mag oxide ferrous sulfate, thiamine HCl, riboflavin niacinamide, pyridoxine, Cal pantothenate folic acid, biotin, inositol, choline chloride/lb Can 1 lb
 Use: Nutritional supplement
- NUTRATE.** (United Labs) Vit A 6000 u, D 1200 u, C 90 mg, B-1 6 mg, B-2 6 mg, B-6 0.5 mg, niacinamide 30 mg, Cal pantothenate 6 mg, Vit B 12 2 mcg plus trace minerals of Iron, Cal, Phos, Mang, Mag, Iodine, copper, Vit E 6 I U, folic acid

0.33 mg./3 Tab. Bot. 100s, 300s, 1000s. Jr. Tabs. Bot. 50s

Use: Vitamin & mineral supplement.

NUTRI-BEE. (A.P.C.) Thiamine HCl 10 mg., riboflavin 10 mg., pyridoxine HCl 5 mg., folic acid 2 mg., niacinamide 50 mg., Cal. pantothenate 25 mg., liver fraction #2 200 mg., stomach conc. 200 mg., Vit. B-12 1 mcg./Cap. Bot. 100s, 1000s.

Use: Dietary supplement.

NUTRI-CAPS. (A.P.C.) Vit. A 15,000 u., D 1500 u., thiamine HCl 5 mg., riboflavin 10 mg., pyridoxine HCl 2 mg., ascorbic acid 100 mg., niacinamide 30 mg., Cal. pantothenate 10 mg., alpha-tocopherol 10 mg., folic acid 0.5 mg., Vit. B-12 2 mcg./Cap. Bot. 30s, 100s, 1000s. Also available w/ minerals.

Use: Dietary supplement.

NUTRI-DISCS. (Ortho) Dextrose 82.54%, Sod. chloride 4.86%, Pot. phosphate monobasic 0.25%, Sod. dibasic phosphate 5.93%, egg albumen 4.79%/Tab. Bot. 12s. Use: Treatment of infertility.

NUTRI-DROPS. (A.P.C.) Vit. A 5000 u., D 1000 u., thiamine HCl 1 mg., riboflavin 0.5 mg., ascorbic acid 50 mg., niacinamide 5 mg., pyridoxine HCl 1 mg., Cal. pantothenate 2 mg./0.6 cc. 50 cc. dosage dropper bot.

Use: Dietary supplement.

NUTRIENT SUPPLEMENT. See: Delcos Granules. (Merck Sharp & Dohme)

Delmor, Pow. (Merck Sharp & Dohme)

NUTRILOR. (Taylor) Vit. A 2000 u., D 200 u., B-1 0.33 mg., B-2 0.66 mg., B-6 D.033 mg., C 12.5 mg., B-12 0.033 mcg, folic acid 0.022 mg., Cal. pantothenate 0.4 mg., niacinamide 5 mg., iodine 0.05 mg., bone phosphates 83.3 mg, dicalcium phosphate 250 mg., Cal. 100 mg., Phos. 666 mg., fluorine 0.033 mg., Mag. 5 mg., ferrous sulfate 1.6 mg., molybdenum 0.033 mg., copper 0.5 mg., zinc 1.0 mg., cobalt 0.1 mg., Mang. 1 mg./Tab. Bot. 100s, 1000s.

Use: (1 t.i.d.) in pregnancy & lactation.

NUTRINIC. (Allied Labs.) Vit. B-1 1RB mg., B-2 3.75 mg., niacin 47 mg., pantothenic acid 1.88 mg., ferrous gluconate 375 mg., liver fraction (1-20) 188 mg., folic acid 0.75 mg., beet ext. (175) 2.25 Gm., Vit. C 37.5 mg./15 cc. Bot. 1 pl.

Use: Hematinic.

NUTRI-SAL. (Ortho) Glucose, Pot. chloride, Sod. chloride, Cal. gluconate. Pkg. 3 vials. Use: Vaginal cleanser.

NUTRISAN. (Recsei) Lactalbumin 65%.

* Generic name recognized by N.F., U.S.P. or A.M.A.

aminoacids 15%, minerals 18%, carboxymethylcellulose 10%. Pow., 1 lb.

Use: Protein therapy.

NUTRISUP. (Chicago Pharm) Vit. A 1000 u., B-1 1 mg., B-2 1 mg., B-6 0.5 mg., B-12 0.5 mg., nicotinamide 5 mg., folic acid 0.25 mg., Cal. pantothenate 1 mg., Vit. C 25 mg., D 200 u., E 0.5 mg., ferrous sulfate exsic., Cal. dibasic phosphate, barium, cobalt, copper, fluorine, iodine, Mag., molybdenum & Pot./Tab. Bot. 50s, 100s, 200s. Use: (2 t.i.d.) Nutritional supplement.

NUTRITIVE. (Forks, Davis) Dicalcium phosphate 900 mg., ferrous sulfate 30 mg., Vit. B-1 2 mg., B-2 2 mg., D 400 u./Cap. Bot. 100s, 1000s.

Use: Supplement diets.

NUTROP. (Marin) Methscopolamine bromide 5 mg./cap. Bot. 30s.

Use: Antispasmodic.

W/Pentobarbital. (Marin) Methscopolamine bromide 5 mg., pentobarbital 30 mg./cap. Bot. 30s.

Use: Antispasmodic w/sedative.

NUVITIN. (Kremers-Urban) Vit. B-12 50 mcg, thiamine HCl 100 mg., niacinamide 100 mg., riboflavin 0.5 mg., pyridoxine HCl 5 mg., panthenol 5 mg., liver 2.5 u., benzyl alcohol 2%, isotonic sol. of Sod. chloride q.s./cc. Vial 10 cc.

Use: Vit. B complex deficiencies.

NUX VOMICA EXTRACT. N.F. W/Ox bile ext., Sod. bicarb. & Cal. phosphate. See: Dunalol, Tab. (Cole Chem. Co.)

W/Aspirin, Hyoscyamus ext., Gelsemium tr. & cascara sagrada ext. See: A.A.S. Compound, Tab. (Tilden)

W/Caffeine theophylline comp. See: Xanthinex, Tab. (Cole Chem. Co.)

NYDRAZID. N.N.D. 1962. (Squibb) (isonicotinic acid hydrazide, isoniazid) Cap. 50 mg., 100 mg. Bot. 100s, 1000s. Tab. 50 mg., 100 mg., Bot. 100s, 1000s. Syr. 20 mg./cc. 1 pt. inj. 100 mg./cc. Vial 10 cc.

Use: Tuberculosis.

NYGRAM. (Dumas-Wilson) Ocin (isomethoptene) mucate 1 gr., acetyl-p-aminophenol 1 gr., salicylamide 2 gr., caffeine ¼ gr./Cap. Bot. 100s, 1000s.

Use: (1 to 2) Relief of migraine and tension headaches

***NYLIDRIN HYDROCHLORIDE.** N.F., N.N.D. 1962. Inj. Tab. N.F. 1-(p-Hydroxyphenyl)-2-(1-methyl-3-phenylpropylamino)propanol HCl. See: Arlidin, Prep. (Arlington-Funk)

NYLMERATE JELLY. (Hoffand-Rantos) Phenylmercuric acetate 0.02%, boric acid in special base, pH adjusted w/acetic acid to

- lob) Ephedrine HCl 0.22%, benzocaine 5% Box 12s
Use Vasconstrictor w/anesthetic
- NUMORPHAN**, NND 1962 (Enda) Oxy morphane[14 hydroxydihydromorphine] HCl 1 mg/cc amps 1 cc & 0.5 cc Box 12s, 100s Vials 15 cc Box 1 & 3s 15 mg/cc Vial 10 cc, Amp 1 cc, 2 cc suppository 2 mg or 5 mg Box 6s
Use Relief of pain
- HUMOTIZINE** Cotaplast (Hobart Lab) Guaiacol, beechwood creosote, methyl salicylate, formaldehyde glycerin & aluminum silicate Jar 4 oz, 8 oz, 15 oz, 30 oz
Use Analgesic & decongestive
- HUMOTIZINE COUGH SYR.** (Hobart Lab) Glyceryl guacolate 5 gr, amm chlor 5 gr, sod cit 20 gr, menthol 0.4 gr/fl oz Bot 3 oz
- HUMOTIZINE**, (Hobart) Phenylephrine HCl 5 mg, acetaminophen 150 mg, pyrilamine maleate 75 mg, caffeine 15 mg, vit C 15 mg/Tab Bot 25s 100s
Use Nasal congestion
- HUMOTIZINE Nasal Spray** (Hobart) Polymixin B sulfate 0.009%, hydrocortisone 0.02%, phenylephrine HCl 0.25%, pyridamine maleate 0.1%, benzalkonium chloride 0.1% in a phosphate buffered base with 1% sodium bisulfite and 2% Tween 20 Spray bats 15 cc
Use Common cold symptoms
- HUHOL** See Phenobarbital (Various Mfr)
- HUPERCALIN CREAM**, (Ciba) Nupercaine 0.5% in water washable base Tube 1 1/2 oz
Use Topically; local anesthetic
- HUPERCALIN OINTMENT** (Ciba) Nupercaine 1% in lanolin & petrolatum Tube 1 oz Jar 1 lb
Use Topically; local anesthetic
- HUPERCALIN SUPPOSITORY**, (Ciba) Nupercaine 2.5 mg, zinc oxide, and bismuth subgallate in a cocoa butter base/Supp Box 12s
Use Treatment of proctitis ani, and hemorrhoids
- HUPERCALIN HYDROCHLORIDE**, NHR 1956 (Ciba) Dibucaine HCl Amp Sol. 1:200, 2 cc, Cortan 10s 1500 20 cc Cortan 12s 1400 w/dextrose 5% (heavy soln) 2 cc Cortan 10s Bot Sol. 2%, 1 fl oz, 4 fl oz Tab: D05 Gm Tube 20s
Use: Local anesthetic
- W/Pyribenzamine HCl combination See Tricainol, Oint & Supp (Ciba)
W/Quinine HCl, benzyl alcohol See Anesacaine (New Mexico Pharm)
- HUPERCALIN, PHEOL & BENZYL ALCOHOL 1H OIL** (Ciba) Nupercaine HCl 0.5%
- phenol 1%, benzyl alcohol 10%. Amp 5 cc, Cortan 10s
Use: Topically; proctology
- NUPORALS**, (Ciba) Dibucaine HCl 1 mg/fl oz, Box 15s, 100s, 500s
Use: Orally; local anesthetic
- NUPRA**, A nonhormonic, nonsteroidic extract of beef organs liver, brain, adrenals See Exval, Tab (Yorktown Prod)
- NUPRIH** (Upjohn) Sulfamazole
Use Antibacterial sulfonamide
- NUROBROMYDS**, (Rorer) Sod bromide 20 gr, strontium bromide 20 gr, ammonium bromide 20 gr, valerian tr 60 min, hyoscyamus tr 60 min/fl oz Bot 14 fl oz 5 pt, 1 gal
Use Sedative & antispasmodic
- NURODOL**, (Buffington) Diethyl barbital acid 1/2 gr, Pa bromide 20 gr, Sod bromide 20 gr, ammonium bromide 20 gr, alcohol 10%/fl oz Bot 1 pt, 1 gal
Use Sedative
- NUROKARDIAC**, (Rorer) Phenobarbital 1/4 gr, nux vomica ext 1/2 gr, carbonyl 6 gr/Tab Bot 100s 500s, 1000s
Use Sedative & digestant
- NU-THERA**, (Kirkman) Contains 9 vitamins & 11 minerals/cap Bot 100s, 1000s 5000s
- NUTRACHLORIC** (Upjohn) Skimmed milk pow 48 gr, Cal carb 11 1/2 gr, Sod bicarb 1 1/4 gr/Tab Box 48s
Use Gastric hyperacidity
- NUTRALKA**, (Enda) Bismuth subcarb 1/2 gr, Mag carb 2 gr, Cal carb 3 1/2 gr, diastase 1/40 gr, papain 1/20 gr, aromatics q s/Tab Bot 100s, 500s, 1000s
Use Antacid & digestant
- NUTRAMENT LIQUID** (Edward Galtan) Protein 20 gr, fat 13.3 gr, carbohydrate 50 gr and 400 calories per 12 1/2 oz can Can 12 1/2 oz
Use Nutritional supplement
- NUTRAMIGEN** (Mead Johnson) Dextrin maltose 45.2%, amigen 20%, corn oil 18%, arrowroot starch 10%, Cal gluconate 3.5% w/Pot phosphate, Pot dibasic phosphate, Cal hydroxide, Pot chloride, Mag oxide, ferrous sulfate, thiamine HCl riboflavin, niacinamide, pyridoxine, Cal pantothenate, folic acid, biotin, inositol, choline chloride/flb Can 1 lb
Use Nutritional supplement
- NUTRATE**, (United Labs) Vit A 6000 u D 1200 u C 90 mg, B 1 6 mg, B 2 6 mg, B 6 D 5 mg, niacinamide 30 mg, Cal pan tothenate 6 mg, Vit B 12 2 mcg plus trace minerals of iron, Cal, Phos, Mang, Mag, sodine, copper, Vit E 6 IU, folic acid

0.33 mg./3 Tab. Bot. 100s, 300s, 1000s Jr. Tabs. Sol. 50s

Use: Vitamin & mineral supplement.

NUTRI-BEE. (A.P.C.) Thiamine HCl 10 mg., riboflavin 10 mg., pyridoxine HCl 5 mg., folic acid 2 mg., niacinamide 50 mg., Cal. pantothenate 25 mg., liver fraction #2 200 mg., stomach conc. 200 mg., Vit. B-12 1 mcg./Cap. Bot. 100s, 1000s.

Use: Dietary supplement.

NUTRI-CAPS. (A.P.C.) Vit. A 15,000 u., D 1500 u., thiamine HCl 5 mg., riboflavin 10 mg., pyridoxine HCl 2 mg., ascorbic acid 100 mg., niacinamide 30 mg., Cal. pantothenate 10 mg., alpha-tocopherol 10 mg., folic acid 0.5 mg., Vit. B-12 2 mcg./Cap. Bot. 30s, 100s, 1000s. Also available w/ minerals.

Use: Dietary supplement.

NUTRI-DISCS. (Ortho) Dextrose 82.54%, Sod. chloride 4.86%, Pot. phosphate monobasic 0.25%, Sod. dibasic phosphate 5.93%, egg albumin 4.79%/Tab. Bot. 12s.

Use: Treatment of infertility.

NUTRI-DROPS. (A.P.C.) Vit. A 5000 u., D 1000 u., thiamine HCl 1 mg., riboflavin 0.5 mg., ascorbic acid 50 mg., niacinamide 5 mg., pyridoxine HCl 1 mg., Cal. pantothenate 2 mg./0.6 cc. 50 cc. dosage drop-per bot.

Use: Dietary supplement.

NUTRIENT SUPPLEMENT.
See: Delcas Granules. (Merck Sharp & Dahme)

Delmar, Paw. (Merck Sharp & Dahme)

NUTRILOR. (Taylor) Vit. A 2000 u., D 200 u., B-1 0.33 mg., B-2 0.66 mg., B-6 0.033 mg., C 12.5 mg., B-12 0.033 mcg., folic acid 0.022 mg., Cal. pantothenate 0.4 mg., niacinamide 5 mg., iodine 0.05 mg., bone phosphates 83.3 mg., dicalcium phosphate 250 mg., Cal. 100 mg., Phos. 66.6 mg., fluorine 0.033 mg., Mag. 5 mg., ferrous sulfate 1.6 mg., molybdenum 0.033 mg., copper 0.5 mg., zinc 1.0 mg., cobalt 0.1 mg., Mang. 1 mg./Tab. Bot. 100s, 1000s.

Use (1 t.i.d.) In pregnancy & lactation.

NUTRINIC. (Allied Labs.) Vit. B-1 18.8 mg., B-2 3.75 mg., niacin 47 mg., pantothenic acid 188 mg., ferrous gluconate 375 mg., liver fraction (1:20) 188 mg., folic acid 0.75 mg., beet ext. (1:75) 2.25 Gm., Vit. C 37.5 mg./15 cc. Bot. 1 pt.

Use: Hematinic.

NUTRI-SAL. (Ortho) Glucose, Pot. chloride, Sod. chloride, Cal. gluconate. Pkg. 3 vials.
Use: Vaginal cleanser.

NUTRISAN. (Reckel) Lactalbumin 65%,
* Generic name recognized by N.F., U.S.P. or A.M.A.

aminoacids 15%, minerals 18%, carboxymethylcellulose 10%. Paw., 1 lb.

Use: Protein therapy.

NUTRISUP. (Chicago Pharm.) Vit. A 1000 u., B-1 1 mg., B-2 1 mg., B-6 0.5 mg., B-12 0.5 mg., nicotinamide 5 mg., folic acid 0.25 mg., Cal. pantothenate 1 mg., Vit. C 25 mg., D 200 u., E 0.5 mg., ferrous sulfate exsic. Col. dibasic phosphate, barium, cobalt, copper, fluorine, iodine, Mag, molybdenum & Pot./Tab. Bot. 50s, 100s, 200s.

Use: (2 t.i.d.) Nutritional supplement.

NUTRITIVE. (Parke, Davis) Dicalcium phosphate 900 mg., ferrous sulfate 30 mg., Vit. B-1 2 mg., B-2 2 mg., D 400 u./Cap. Bot. 100s, 1000s.

Use: Supplement diets.

NUTROP. (Marin) Methscopolamine bromide 5 mg./cap. Bot. 30s.

Use: Antispasmodic.

W/Pentobarbital. (Marin) Methscopolamine bromide 5 mg., pentobarbital 30 mg./cap. Bot. 30s.

Use: Antispasmodic w/sedative.

NUVITIN. (Kremers-Urban) Vit. B-12 50 mcg., thiamine HCl 100 mg., niacinamide 100 mg., riboflavin 0.5 mg., pyridoxine HCl 5 mg., pantothenol 5 mg., liver 25 u., benzyl alcohol 2%, isotonic sol. of Sod. chloride q s./cc. Vial 10 cc.

Use: Vit. B complex deficiencies.

NUX VDMICA EXTRACT, N.F. W/Ox bile ext., Sod. bicarb. & Cal. phosphate. See Dunokol, Tab. (Cole Chem. Co.)

W/Aspirin, Hyoscyamus ext., Gelsemium Ir. & castoreo sagrada est. See A.A.S. Compound, Tab. (Tilden)

W/Caffeine-theophylline comp. See Xanthinax, Tab. (Cole Chem. Co.)

HYDRAXID, N.N.D. 1962. (Squibb) Isonicotinic acid hydrazide, Isoniazid) Cap. 50 mg., 100 mg Bot. 100s, 1000s. Tab. 50 mg., 100 mg., Bot. 100s, 1000s. Syr. 20 mg./cc 1 pt. Inj 100 mg./cc. Vial 10 cc.

Use: Tuberculosis.

NYGRAM. (Dumas-Wilson) Ocin (isomethoptene) mucate 1 gr., acetyl p-aminophenol 1 gr., salicylamide 2 gr., caffeine ¼ gr./Cap. Bot. 100s, 1000s.

Uses (1 to 2) Relief of migraine and tension headaches

***NYLIDRIN HYDROCHLORIDE, N.F., N.N.D. 1962.** Inj. Tab. N.F. 1-(p-Hydroxyphenyl)-2-[1-methyl-3-phenylpropylamino]propanol HCl See Arlidin, Prep. (Arlington-Funk)

NYLMERATE JELLY. (Halland Rantol) Phenylmercuric acetate 0.02%, boric acid in special base, pH adjusted w/ acetic acid to

pH 4.0 Jelly, Tube with or without applicator 5 oz
 Use Trichomonocide & vaginal antiseptic

NYLMERATE SOLUTION CONCENTRATE. (Holland Rantos) Phenylmercuric acetate 0.2% in buffered solvent of alcohol 30%, acetone 10%, deionized water. Bot ½ pt, 1 pt
 Use: Antiseptic & trichomonocidal

NYLOXIN (Hynson, Wescott & Dunning) Cobra venom, formic acid & silicic acid Amp 3 cc, Box 10s
 Use: SC; chronic arthritis

NYMORE. (Durst) Tyrothricin 1 mg, tyrocidine ½ mg, benzocaine 5 mg /Loz Bot 100s
 Use Throat & mouth infections

NYOMIN. (Elder) Veratrum viride 2/5 gr, Sod nitrite ½ gr, Pot nitrate 2½ gr, nitroglycerin 1/250 gr, aconite root 2/5 gr /Tab Bot 100s, 1000s
 Use Hypertension

NYSCOID. (Nysco Labs) Vit B 12 cryst Tab 5 mcg Bot 100s 1000s
 Use: Vit B 12 therapy

NYSTAFORM (Dome) Nystatin 100 000 U tetrachlorhydroxyquin 10 mg /Gm Tube ½ oz
 Use Moniliosis

***NYSTATIN, USP, NND 1962** An anti-fungal antibiotic derived from cultures of *Streptomyces noursei* Oint Oral Susp, Tab USP See: Mycostatin (Squibb)
 W/Demethylchlorotetracycline See Declosolin, Caps (Lederle)
 W/Fludrocortisone acetate, neomycin sulf, gramicidin See: Myconet Ointment (Squibb)
 W/Fludrocortisone sulf, polymyxin B sulf neomycin base as sulfate See: Florotic plastic dropper (Squibb)
 W/Gramicidin, neomycin, triamcinolone nystat See: Nycalog Cream (Squibb)
 W/Hydrocortisone neomycin sulfate See: Neo Nysta Cort, (Dome)
 W/tetrachlorhydroxyquin See Nystofarm (Dome)
 W/Neomycin base, gramicidin, triamcinolone acetate See: Mycolog Cream (Squibb)
 W/Neomycin sulfate See: Mycitraadin N, Tab (Upjohn)
 W/Tetracycline
 See: Casa Tetrastatin, Caps (Pfizer)
 Mysteclin, Cap (Squibb)
 W/Tetracycline phosphate buffered See: Achromycin V, Cap (Lederle)
 W/Tetracycline phosphate See: Comycin, Cap (Upjohn)

*Generic name recognized by N.F., U.S.P. or A.M.A.

OAK & IVY EXTRACT. See: Birhutox, Amp (Central)

OATMEAL COLLOIDAL W/Koolin, sulfur, and Hexochlorophen. See: Acnoveen, Lat Pow (Aveeno Corp)

OATMEAL, GUM FRACTION. See: Aveeno, Pow. (Aveeno)

O B.-6. (Kremers Urban) Vit A 2000 u, O 200 u, B 1 0.5 mg, B 2 1 mg, B 6 1 mg, C 33.3 mg, B 12 17 mcg, niacin 5 mg, pantothenic acid 33 mg, folic acid 0.7 mg w/iron, Cal, phosphates, iodine, copper, Mang, cobalt, molybdenum, zinc, Mag & Pot /Cap Bot 90s
 Use Pregnancy & lactation

O B.E.C. (Morton) Formula A, dl Desoxyephedrine 1/12 gr, thyroid 1 gr, atropine sulfate 1/350 gr /Tab Bot 100s Modified Thyroid eliminated Tab Bot 100s B Pheno-rborbitol ¼ gr, rochelle salts 5 gr, aloin ¼ gr /Tab Bot 100s
 Use: Antiobesity

OBEDBIN. (Massengill) Semoxydrine HCl 5 mg, pentobarbitol Sod 20 mg, ascorbic acid 100 mg thiamine HCl 0.5 mg, Vit B 2 1 mg, niacin 5 mg /Tab Bot 100s, 500s, 1000s
 Use: (1 t i d o c) With reducing diets

OBEDRIN L.A. (Massengill) Methamphetamine HCl 12.5 mg, pentobarbitol 50 mg, Vit C 200 mg, B 1 mononitrate 1 mg, B 2 2 mg nicotinic acid 10 mg /Tab double layer Bot 50s
 Use Curb appetite

OBELIP. (Lakeside) Phenobarbitol 16 mg, d amphetamine sulfate 5 mg, choline bitartrate 400 mg, dl methionine 150 mg, Vit B 12 4 mcg, methylcellulose 160 mg, Mag stearate 14 18 mg /Cap Bot 50s
 Use: 1 t i d, a c, w/glass of water; anti-obesity

OBESIFORM (New Mexico Pharm) Anterior pituitary 30 gr whole ovarian 40 gr suprarenal cortex 30 gr, thyroid 2 gr posterior pituitary 3 IU, chlorbutanol 0.5%/cc Vials 30 cc

OBESILIN. (Lincoln) Methylcellulose 0.5 Gm, ascorbic acid 25 mg, dl amphetamine 5 mg, thiamine HCl 3 mg /Tab Bot 100s 1000s
 Use: 1 t i d; antiobesity

OBESITY AGENTS. See: Anti Obesity Agents (Various Mfr)

OBESITY TABLETS. (Normand)
 No. 1—Morning Tabs.—White—Amphetamine HCl 5 mg, thyroid 1 gr, atropine sulf 1/360 gr, aloin ¼ gr /Tab
 No. 2—Noon Tabs.—Pink—Amphetamine

- HCl 5 mg., thyroid 1 gr., atropine sulf. 1/360 gr./Tab.
- Na. 3—Afternoon Tab.—Gray—Amphetamine HCl 5 mg., thyroid 1 gr., phenobarbital ¼ gr./Tab. Set of 3 bottles of 1000 tab. each.**
 Use: Treatment of obesity
- OBESITY T.D. (Dumas-Wilson) d-Amphetamine HCl 15 mg., thyroid 50 mg., phenobarbital 25 mg./Cap. Bot. 100s, 1000s.**
 Use: Control and management of obesity
- OBESIDS. (Columbus) d-Amphetamine phosphate 2.50 mg., butberbital sod. 8 mg., homatropine methylbromide 0.65 mg., bullogen 250 mg., Vit. A 1000 u., Vit. D 100 u., Vit. B-1 1 mg., B-2 1.50 mg., niacinamide 10 mg./Tab. Bot. 100s, 1000s**
 Use: Control of obesity
- OBESONIL. (Lincoln) d-Amphetamine sulfate 2 mg., methylcellulose in chocolate flavored base/Tab. Bot. 100s, 1000s.**
 Use: 1 t.i.d., antiobesity.
- OBESTAT. (Lemmon) Amobarbital 60 mg., methamphetamine 10 mg., thyroid 150 mg./Timed-release Tab. Bot. 100s, 500s, 1000s.**
 Use: Obesity.
- OBESTROL. (Crestmed) Methamphetamine HCl 5 mg., phenobarbital 16 mg., C 100 mg., B-1 1 mg., niacin 10 mg., methylcellulose 150 mg./Cap. Bot. 100s.**
 Use: Obesity.
- OBES TR. (Columbus) d-Amphetamine sulfate 15 mg., thiamine, niacinamide, Vit. B-12, col. pantothenate, ascorbic acid, riboflavin, Vit. A, D/T.C. Tab. Bot. 30s**
 Use: Anorexiast.
- OBETAMINE. (Sly) Sod. methoxycellulose 300 mg., methamphetamine HCl 6 mg./Tab. Bot. 100s, 500s, 1000s.**
 Use: 1-2 t.i.d., a.c., antiobesity.
- OBETROL. (Obetrol Pharm. Co) Methamphetamine saccharate, methamphetamine HCl, amphetamine sulf., dextroamphetamine sulf. in equal parts. Tab. 10 or 20 mg Bot. 100s, 500s, 1000s**
 Use: Treatment of obesity
- OBEVITA. (Meyer) Vit. A 5000 u., D 1000 u., thiamine HCl 3 mg., riboflavin 3 mg., pyridoxine HCl 0.5 mg., ascorbic acid 40 mg., niacinamide 20 mg., Cal. pantothenate 5 mg., folic acid 0.25 mg., Vit. B-12 1 mcg.-E 2 I.U./Cap. Bot. 100s, 1000s.**
 Use: Multivitamin therapy.
- OB-GYN FURACIN CREAM. (Eaton) Nitrofurazone 0.2%. Tube w/applicator.**
 Use: Treatment of vaginal infections and to speed healing.
- OBID. (Ingram) Calcium carbonate 340 mg., ferrous sulfate 34 mg., Vit. A 4000 u., D 400 u., B-1 3 mg., B-2 3 mg., B-6 5 mg., niacin 20 mg., pantothenic acid 3 mg., C 75 mg., folic acid 0.3 mg., B-12 2 mcg./Cap. Bot. 100s, 500s**
 Use: Phosphorus and sodium-free, prenatal multivitamin-mineral therapy
- OBLIVON. See: Darmison (Schering)**
- OBLIVON-C. (Brit. Schering) Methylpentynol carbonate.**
 Use: Sedative. Under study.
- OBNATAL. (Boyle) Vit. C 100.0 mg., B-6 10.0 mg., B-12 10.0 mcg., B-1 4.0 mg., B-2 2.5 mg., niacinamide 60.0 mg., folic acid 0.3 mg., Cal. pantothenate 5.0 mg., Vit. K 1.5 mg., A 6000 U.S.P. u. (acetate), D 600 U.S.P. u., cal. lactate 2.25 Gm., ferrous sulf. 300 mg., iodine 0.15 mg., copper 1.0 mg., mag. 15.0 mg., mang. 4.5 mg., molybdenum 0.15 mg., nickel 0.15 mg., zinc 2.25 mg./3 cap. Bot. 100s, 1000s.**
 Use: 1 t.i.d., phosphorus-free prenatal nutrition.
- ORNATAL-O. (Boyle) Vitamin and Mineral tablet. Bot. 100s**
 Use: Vitamin and mineral supplement during pregnancy and lactation.
- OBOCILL COMPLEX. (Nelsler) d-Amphetamine phosphate 5 mg., nicel (methylcellulose) 100 mg., choline bitartrate 200 mg., inositol 60 mg., vit. B-1 0.8 mg., B-2 1.2 mg., niacin 8 mg./tab. Bot. 100s, 500s.**
 Use: Anti-obesity.
- OBOCILL LIQUID. (Irwin, Nelsler) Methylcellulose 9 gr., d-amphetamine phosphate 10 mg./oz. Bot. 1 pt.**
 Use: (1-2 tsp. q.e. w/glass of water) Appetite depressant.
- OBOCILL TABLETS. (Irwin, Nelsler) Methylcellulose 150 mg., d-amphetamine phosphate 5 mg./Tab. Bot. 100s, 500s, 1000s.**
 Use: (3-6 daily a c. w/glass of water) Curb appetite.
- OBOCILL-TF. (Irwin, Nelsler) Methapyrilene 25 mg., d-amphetamine phos. 5 mg., methylcellulose 150 mg./Tab. Bot. 100s, 500s, 1000s.**
 Use: Combat tension and emotional stress.
- OBOLIP. (Lakeside) Phenobarbital ¼ gr., d-amphetamine 5 mg., choline bitartrate 400 mg., dl-methanline 150 mg., Vit. B-12 4 mcg., methylcellulose 160 mg./Cap. Bot. 50s.**
 Use: 1 t.i.d., antiobesity.
- OBRECAL. (Conright) Vitamins D 400 u., B-1 1 mg., B-2 1 mg., C 10 mg., ferrous sulf. 25 mg./tab. Bot. 100s, 1000s.**
- OBRON. (Roerig) Vit. A 5000 u., D 400 u., thiamine HCl 2 mg., riboflavin 2 mg., pyridoxine HCl 0.5 mg., ascorbic acid 37.5 mg., niacinamide 20 mg., Cal. pantothenate**

3 mg, dicalcium phosphate, anhy. 770 mg, ferrous sulfate 44 mg, cobalt 0.033 mg, copper 0.33 mg, iodine 0.05 mg, Mang 0.33 mg, Mag 1 mg, molybdenum 0.07 mg, Pot. 1.7 mg, zinc 0.4 mg / Cap Bot 100s

Use (1-3 daily) Dietary supplement.

OBRON IMPROVED. (Roeng) Vit A 6000 u, D 400 u B 1 3 mg, B 2 2 mg, B 6 2 mg, niacinamide 20 mg, K 0.5 mg, calcium pantothenate 1 mg, folic acid 0.5 mg, C 50 mg, B 12 3 mcg, as tablets with non-inhibitory intrinsic factor concentrate, calcium carbonate 593 mg, ferrous fumarate 105 mg / Cap Bot 100s

Use Nutritional needs of pregnancy and lactation

OCCULTEST. (Amet) Orthotolidin, strontium peroxide, cal acetate & tartaric acid Tab, Bot 250s w/filter papers

Use: Not for internal use Diagnostic test for occult blood in urine

OCCY-CRYSTINE. (Ocy Crystine) Sod thiosulfate 1.28 Gm, Mag sulfate 1.19 Gm / 4 cc Bot 4 R oz, 8 fl oz Sod thiosulfate 0.433 Gm, Mag sulfate 0.433 Gm, Pot sulfate, Cal sulfate / Tab Bot 75s

Use: Laxative & chologogue

OCTAPLEX. (APC) Vit A 5000 u, D 1000 u C 50 mg, B 1 3 mg, B 2 3 mg, B 6 0.5 mg, Cal pantothenate 5 mg, niacinamide 20 mg, folic acid 0.25 mg, Vit B 12 1 mcg / Cap Bot 100s, 1000s.

Also available w/minerals
Use (1 cap daily) Dietary supplement

***OCTATROPINE METHYLBROMIDE.** 2 propylpentanoylmethyltropinium bromide See Valpin (Endo Lab)

OCTIN HCl, NND 1961 (Knoll) Isometheptene HCl Methylisobutylamine Amp 100 mg / cc Box 6s 100s Oral sol Octin HCl 10% Bot 10 cc

Use Relaxes smooth muscle

W/braitureal See Valactin, Tab (Knoll)

OCTIN MUCATE, NND 1962 (Knoll) Isometheptene mucate 120 mg / Tab / Bot 100s Supp 100 mg Box 6s 100s

OCTOFEN. (McKesson & Robbins) Liq 8 Hydroxyquinoline in ethyl alcohol sol Bot 1 1/2 oz, 4 oz Pow 8 Hydroxyquinoline & silica gel Bot 2 1/2 oz

Use: Fungicidal

OCTOFOLLIN, NNR 1949 See Benzestrol (Schleffelin)

OCTOGESIC. (Merit) Sod gentisate 225 mg, megensate 225 mg / Tab Bot 50s
Use Antirheumatic.

OCTOID. (Warren Teed) Vit A 5000 u D 1000 u, ascorbic acid 37.5 mg, riboflavin

* Generic name recognized by N.F., U.S.P. or A.M.A.

2 mg, thiamine HCl 3 mg, pyridoxine HCl 0.5 mg, Cal pantothenate 5 mg, nicotinamide 20 mg / Cap Bot 100s, 1000s

Use: Vitamin supplement

OCTRIPHENATE & SALICYLANILIDE. See: Sylamin, Cream (Abbott)

OCTRITE. (Hynson, Westcott & Dunning) Ncl listed Octyl nitrite 750 mg / Inhaler Inhaler, Pkg 1

Use: Coronary vasodilator.

OCTYL NITRITE.

See: Medihaler Nitro, Vial (Riker)

Octrite, Inhaler (Hynson, Westcott & Dunning)

***OCTYLPHENOXY POLYETHOXYETHANOL, NF** A mono ether of a polyethylene glycol See Igepal CA 630 (Antara Chemicals)

Use: Ingredient in Nitrofurazone Sol, NF

OC-U-MID Ophth, Sol. (Blue Line) Sodium sulfacetamide (w/v) 30% Sterile and buffered to approx pH 7.4 15 cc plastic squeeze bottle

Use Ocular infections

OCUTRIN EYE DROPS. (Lourison) Polymyxin B sulfate 16,500 u, hydrocortisone 5 mg, neomycin sulfate 5 mg / cc Dropper bot 5 cc

Use Infection of the eye

OC-U-ZIN Ophth Sol. (Blue Line) (w/v) zinc sulfate 0.25%, boric acid 2% phenyl ephrine HCl 0.12%, sodium borate with benzalkonium chloride 1:20,000 hydrous chlorobutanol 0.25% 15 cc plastic squeeze bottle

Use: Superficial irritations of the eye

ODI-LATE. (Warren-Teed) Homatropine methylbromide 1/50 gr, ox bile ext 1 gr / nitroglycerin 1/500 gr, Mag sulfate 5 gr / Tab Bot 100s, 1000s

Use (1 tsd) Choleretic & antispasmodic

OENETHYL HYDROCHLORIDE. (Knoll) Methyl ylaminoheptane HCl Amp (50 mg / cc) 1 cc, Box 6s, 100s; (50 mg / cc) 2 cc, Box 5s, 50s Vial, Sol 5%, 10 cc

Use: Vasopressor action

OENETHYL SULFATE. (Knoll) Methylaminoheptane sulfate Sol 1%, Bot 15 cc, 240 cc; 2%, Bot 60 cc

Use: Nasal vasoconstrictor

OESTERON. See Estradiol (Various Mfr.)

OESTRADIOL. See Estradiol (Various Mfr.)

OESTRASID. See Dienestrol (Various Mfr.)

OESTRIN. See Estrone (Various Mfr.)

OESTROFORM. See Estrone (Various Mfr.)

OESTROMENIN. See Diethylstilbestrol (Various Mfr.)

OESTROMON. See Diethylstilbestrol (Various Mfr.)

OESTRONE. (Meyer) Estrone. Vial (1 mg./cc.) 10 cc.

Use: Estrogen therapy.

O.H.B. (Bauflich) Salicylamide 7 gr./Tab. Bot. 100s.

Use: Analgesic and antirheumatic.

OIL ENEMA. (Pharmoseol) Light mineral oil/126 cc. in plastic tube and tip

Use: Oil retention enema

OIL RETENTION ENEMA. (Fleet) Mineral oil U.S.P., 135 cc. in plastic container with 2-in. tip

OILZO. (G. F. Harvey) Not listed. Triolein ozonide in olive oil. Bot. 1 oz., 4 oz., 1 pt.

Use: Antiseptic & deodorant.

OINTMENT BASE, WASHABLE.

See: Absorbable Base (Upsher-Smith)

Cetaphil (Texas Pharm.)

Cetyl-Stearic Base (Upsher-Smith)

Dermobase (Marcelle)

Hydrotex (Texas Pharm.)

Neobase (Burroughs Wellcome)

Omnia

Pramulgen (Robison-Wagner)

Velvachol (Texas Pharm.)

OLAC. (Mead Johnson) Non-fat milk solids 40.6%, dextrimaltose (maltose & dextrins)

31.7%, corn oil 17.5%, Cal. caseinate

10.1%, halibut liver oil 0.1%. Can 1 lb.

Liq.—Same formula as the powder. Can 13 fl. oz.

Use: Infant feeding.

OLD TUBERCULIN. N.N.R. 1950. See: Tuberculin, Old, U.S.P.

OLEAGINOUS OINTMENT BASE. See: Flastibase. (Squibb)

***OLEANDOMYCIN.** An antibiotic from *Streptomyces antibioticus*

See: Motromycin, Cap. (Pfizer)

Tao Pediatric Drops. (Roerig)

W/Gantrisin. See: Gantrimycin, Tab. (Roche)

W/Tetracycline. See: Signamycin, Susp. (Pfizer)

***OLEANDOMYCIN, TRIACETATE, N.F., N.N.D. 1962.**

See: Cyclamycin (Wyeth)

Tao, Cap. (Roerig)

W/Sulfas. See: Taamid (Roerig)

W/Tetracycline. See: Signamycin, Cap. (Pfizer)

OLEANDOMYCIN PARENTERAL, N.N.D. 1962. (Roerig) Oleandomycin phosphate 500 mg./Vial.

Use: I.V., I.M., antibiotic therapy against gram-positive organisms.

***OLEANDOMYCIN PHOSPHATE, N.F., N.N.D. 1962.** (Wyeth) Vial 500 mg. Caps. 10. N.F. See: Motromycin Phos (Pfizer)

* Generic name recognized by N.F., U.S.P. or A.M.A.

Use: Infections caused by susceptible gram-positive and gram-negative organisms.

OLEANDOMYCIN SALT OF PENICILLIN.

See: Pen-M (Pfizer) Under study

OLEIN OZONIDE. See: Ozonides of olive oil.

OLEO-BIEND VITAMIN A. (White) Cap. Vit. A 25,000 u. Bot. 25s, 100s, 1000s, 5000s.

Use: Vit. A therapy.

OLEOVITAMIN A, U.S.P., N.N.D. 1961. A D R, Cap., U.S.P. (Natural Vit. A in Oil)

Brewer—Gel-lets, Vit. A 25,000 u. Bot. 100s, 1000s; Vit. A 50,000 u. Bot. 50s, 100s, 1000s.

Ives-Cameron—Cap. 25,000 u. & 50,000 u. Bot. 100s, 1000

See: Aflaxin, Cap. (Winthrop Labs)

***OLEOVITAMIN A & D, N.F., N.N.R. 1957.** A D R. 1954, Cap. N.F.

See: A-D Percomorph, Cap. (Abbott)

Alpha-Deltalin, Cap. (Lilly)

A-Vita-D, Cap. (Fitman-Moore)

Burbot Liver Oil, Cap., liq. (Rawell)

Caritol w/Vit. D, Cap. (Wyeth)

Cod Liver Oil Conc., Cap., liq., Tab. (White)

Drisdol w/Vit. A, Cap., liq. (Winthrop Labs.)

Hepicolium, Cap., liq. (Lilly)

4-Vi-Tol, Cap., liq. (Fatch)

Notola, Cap., liq. (Forke, Davis)

Navital w/Viosterol, Cap. (Squibb)

Oleum Percomorphum w/Viosterol, Cap. (Mead Johnson)

Super-D, Cap., liq. (Upjohn)

Vad, Cap. (Cole)

Vi-Delta, Cap., liq. (Lederle)

Use: (1 daily) Vit. A & D therapy.

OLEOVITAMIN A & D, CONCENTRATED, U.S.P., N.N.R. 1955; Cap., U.S.P. Vit. A 50,000 u. & Vit. D 1150 u.

OLEOVITAMIN D, SYNTHETIC, U.S.P. See: Viosterol In Oil, N.N.R. 1952.

OLEUM PERCOMORPHUM, N.N.R. 1955. Percomorph Liver Oil. (Vit. A 60,000 u., D 8500 u./Gm.)

Amer. Pharm.—Bot. 10 & 50 cc.

Approved Pharm.—Bot. 10 & 50 cc.

Dewey—Bot. 50 cc.

Mead Johnson—Bot. 10 & 50 cc.; Cap. Bot. 50s, 250s.

OLEUM PERCOMORPHUM, CODANOL BRAND, W/OTHER FISH LIVER OILS & VIOSTEROL, N.N.R. 1955. (Amer. Pharm)

liq. 60,000 u. A & 8500 u. D Bot. 10 cc., 50 cc.

OLEUM PERCOMORPHUM W/OTHER FISH LIVER OILS & CALCIFEROL, N.N.R. 1954. A D R. 1951. (Mead Johnson) Blend of

A rich & Vit D rich liver oils of perca morph fishes w/viosterol & other fish liver oils liq Vit A 60 000 u, D 8500 u/cc Bot 10 cc, 50 cc Cop Vit A 5000 u, D 700 u/Cop Bot 50s, 250s

Use Vit A & D therapy

OLODEX. (Wolke) Chlorophyll ext fractions 100 mg/Tab Box 30s; Bot 100s, 1000s

Use: Body odors

OLOTHORB See Polysorbate 80 (Various Mfr)

OL-VITUM CAPSULES. (Ives Cameron) Vit A 5000 u, D 1000 u, C 37.5 mg, thiamine HCl 2.5 mg, riboflavin 2.5 mg, niacinamide 20 mg, Col pantothenate 5 mg, pyridoxine HCl 0.5 mg/Cop Bot 100s, 250s, 1000s
Use Dietary supplement

OL-VITUM DROP (Ives Cameron) Vit A 10,000 u, D 2000 u B 1.3 mg, B 2 0.8 mg, C 100 mg, B 6 1.6 mg, niacinamide 15 mg, natural mixed tocopherols 3 mg / cc Bot w/dropper 15 cc, 30 cc
Use Dietary supplement

OL-VITUM-M (Ives Cameron) Vit A 5000 u, D 1000 u, C 50 mg B 1 2.5 mg, B 2 2.5 mg B 6 0.5 mg, B 12 2 mcg, niacinamide 20 mg, cal pantothenate 5 mg, folic acid 0.25 mg plus iron, copper, cobalt iodine, mang, mag, molybdenum, pot and zinc/Cop Bot 100s
Use Prevent deficiencies

OMADERM. (Dorco) Diisobutyl ethoxyethyl dimethyl benzyl ammonium chloride 0.5%, benzocaine 1%, benzyl alcohol 2% in glycerin Bot 15 cc
Use: Ear infections

OMIAL (Kenwood) Hesperidin complex 50 mg, citrus bioflavonoid compound 50 mg, ascorbic acid 200 mg, calcium 100 mg, iodine 0.15 mg, copper 1 mg, potassium 0.5 mg, zinc 0.5 mg, manganese 1 mg, cobalt 0.5 mg, molybdenum 0.3 mg, iron 33.3 mg, folic acid 0.125 mg, Vit B 12 1 mcg, Vit A 5000 u, D 500 u, B 1 2 mg, B 2 2 mg, niacinamide 20 mg, Vit K 0.5 mg, B 6 2 mg/Tab Bot 100s
Use: Nutritional supplement

OMHADIH (Winthrop Labs) Comp sterile sol of protein (obtained from nonpathogenic bacteria) various animal fats & lipids derived from bile Amp 2 cc, Box 5s, 25s Vial 10 cc w/rubber diaphragm stopper, Box 1s, 25s
Use: Non-specific protein therapy

OMNI-BETA. (Warner Chilcott) Vit B 1 2 mg, B-2 2 mg, niacinamide 10 mg, panthenol 172 mg, Vit B 6 0.2 mg, choline 20 m itol 10 mg, liver fraction #2,

N.F. 0150 Gm/dr Bot 8 fl oz, 16 fl oz, 1 gal

Use: Vit B complex deficiencies

OMNI-TUSS. (Strassenburgh) Codeine 10 mg, phenyltoloxamine (as a resin) 5 mg, chlorphaniramine (as a resin) 3 mg, ephedrine 25 mg, guaicol carbonate 20 mg

Use Antitussive

OMNI-VITA. (Warner Chilcott) Vit A 5000 u, D 1000 u, C 50 mg, B 1 1 mg, B 2 2 mg, B 6 0.5 mg, B 12, crystal 1 mcg, panthenol 1 mg/Spherette Bot 50s

Use: (1-3 daily) Vitamin deficiencies

O.N.P. (Clark & Clark) Pyrilamine maleate Tab Bot 100s, 250s, 1000s

Use: Antihistaminic

ONYCHO-PHYTEX. (Wynlit) Berotannic complex (derived from 46 mg tannic acid and 29 mg boric acid) 75 mg, salicylic acid 8 mg, ethyl alcohol 56% Bot 30 cc

Use: Nail fungus infection (onychomycosis)

OPASAL (Elder) Dovers powder 15 mg, acetophenetidin 0.1 Gm, acetylsalicylic acid 0.12 Gm, atropine sulfate 0.12 mg, caffeine 8 mg, camphor monobromated 8 mg/Cop Bot 100s, 500s

Use Analgesic

OPH EYE LOTION (Winthrop Labs) Neosynephrine 0.08%, zinc sulfate 0.06%, boric acid 2.2%, w/chlorobutanol as preservative Bot 10 cc w/dropper

Use Eye lotion

OPHTHAINE HCl, NND 1962 (Squibb) Proparacaine 2 Diethylaminoethyl 3 amino-4 propoxybenzoate HCl 0.5%, chlorbutanol 0.2% glycerin 2.45%, benzalkonium chloride 110 000 Sol Bot 15 cc

Use Topical anesthesia for the eye

OP-HYDRIN. (Braemmel) Hydrocortisone acetate 0.5%, phenylephrine 0.125%, chlorbutanol 0.10%, w/methylparaben propylparaben Susp Bot 1/4 oz

Use Eye infections

OPHTHAMOL (White) Vit A & D derived from fish liver oils in oint base Tube 1/2 oz

Use: Epithelization n corneal erosions

OPHTHA PBH DROPS. (Misemer) Hydrocortisone acetate USP 5 mg, polymyxin sulfate USP 16 250 units, neomycin sulfate USP 5 mg, in a sterile ophthalmic susp 5 cc plastic drop containers

Use Ophthalmic

OPHTHA PBH OINT. (Misemer) Polymyxin B sulfate 5000 USP units, zinc bacitracin 400 USP units, neomycin sulfate 5 mg/1 Gm Tube 1/2 oz
Use Ophthalmic

gramicidin 0.25 mg, propyl paminobenzoate 2 mg/chewing gum tab Pkg 20s
Use: Antibiotic & anesthetic chewing gum

ORACAINE HYDROCHLORIDE. (Mizzy)
Meprylcalne HCl.

ORA-CEL. (First Texas) Vit A palmitate 25,000 u, D 1000 u, thiamine HCl 10 mg, riboflavin 20 mg, pyridoxine HCl 1 mg, niacinamide 50 mg, lysine 100 mg, calcium ascorbate 150 mg, calcium pantothenate 10 mg, crystalline B 12 5 mcg/Cap Bot 100s

Use: Vitamin therapy

ORADASH. (Adams) d Methanline 200 mg/Cap Bot 12s

Use: Diaper rash

ORADIOL. NND 1959

ORAGRAFIN CALCIUM (Squibb) ipodate calcium 3 Gm/Single dose packets cartons 25s 100s

Use: Radiopaque substance

ORAGRAFIN SODIUM (Squibb) 0.5 Gm ipodate sodium/B Gm packet or 6 caps
Use: Radiopaque substance

ORAHEMA (Suliff Case) Vit B 12 5 mcg, ferrous gluconate 130 mg, stomach substance 50 mg, ascorbic acid 50 mg, folic acid 0.30 mg/Cap Bot 100s, 1000s

Use: Deficiency anemias

ORAHEMA PLUS (Suliff & Case) Intrinsic factor B-12 concentrate 0.25 u, ferrous gluconate 200 mg, zinc 1 mg, manganese 5 mg, cobalt 0.5 mg, copper 0.25 mg, molybdenum D 25 mg, ascorbic acid 25 mg, folic acid 0.5 mg/Cap Bot 100s, 1000s

Use: Anemias

ORALCID. See: Acetarsone (Abbott)

ORAL PENTACRESOL. (Upjohn) Pentacresol 1:1000 sol Bot 4 oz, 1 pt, 1 gal

Use: Antiseptic mouth wash

ORALPLEX. (Rare) Thiamine HCl 1 mg, riboflavin 2 mg, niacinamide 20 mg, Cal pantothenate 12 mg, pyridoxine HCl 12 mg/A oz. Elixir, Bot 1 pt, 1 gal

Use: Dietary supplement

ORAMESTRIN-H. (Smith, Miller & Patch) Estrogenic substances conjugated D 625 mg, dl desoxyephedrine HCl 25 mg, methylandrostenedial 5 mg/Tab Bot 30s, 100s 1000s

ORAMETS. (National Drugs) Tyrothreclin 2 mg, benzacaine 5 mg/Loz, Bot 15s, 100s, 500s

Use: (1 q 4 h) Antibiotic therapy

ORANDE SPANSULE. (Smith Kline & French) Chlorphenapyridamine maleate 8 mg, phenylpropanolamine HCl 50 mg, Isepropamide 2.5 mg/Cap Bot 30s

Use: Orally, treatment of common cold

ORANIXON. NND 1962 (Organon) Me phenesin Elixir 100 mg/cc Bot B oz, 1 pt, 1 gal Tab 250 mg, Bot 100s, 500s, 1000s, 500 mg, Bot 100s, 1000s

Use (I Gm f i d or q i d) Muscle relaxant

ORANIXON COUGH COMPOUND. (Organon) New name for Tazanon (Organon)

ORAPEN LIQUID. (Schenley) Pot penicillin G 300,000 u 5 cc when diluted w/water as directed Bot (buffered penicillin 3,600,000 u) 60 cc

Use: Orally; penicillin therapy

ORAPEN TABLETS. (Schenley) Pot penicillin G cryst Tab 50,000 u, Box 12s, 100s Bot 1000s, 100,000 u, Box 12s, 100s, 500s, 1000s, 250,000 u, Box 12s, 100s, Bot 1000s, 500,000 u, Box 10s

Use: Orally; penicillin therapy

ORARSAN See Acetarsone (Abbott)

ORASEPTINE. (Suliff & Case) Zinc chloride 17½ gr, oil cinnamon 12 min, oil clove 1 min, menthol 3 gr saccharin soluble 3 gr/pl Liq pt, gal

ORA-TESTRYL (Squibb) Fluoxymesterone Tab 2 mg, 5 mg Bot 50s, 100s

Use: Anabolic agent

ORATROL. (Alcon) Dichlorphenamide 50 mg/Tab Bot 100s

Use: Oral treatment of glaucoma

ORAVALS. (Vitamin Corp) Vit A 15,000 u, D 1500 u, B 1 10 mg, B 2 6 mg, B 6 1 mg, C 100 mg, Cal pantothenate 5 mg, nla cinamide 20 mg, Vit E 5 mg, folic acid 0.25 mg, Vit B 12 2 mcg/Cap Bot 50s, 100s, 250s, 1000s

Use: (1-3 daily) Dietary supplement

ORAVAX. (Merrell) Total killed organisms 60,000 million & the soluble antigens from 8625 million/Tab Bot 20s, 50s, 1000s

Use: (1 daily for one week) Prophylactic against common cold

ORBIMIN CAPSULES. (Central) Methen amine bismuth iodide 4 gr/Cap Bot 60s, 500s, 1000s

Use: Orally (2-3 daily); antispasmodic therapy

ORBITINIC. (Schilling) Ferrus fumarate 100 mg, B 12 5 mcg, folic acid 0.5 mg, d sorbitol 500 mg/Tab Bot 100s

Use: Iron deficiency anemias

ORCNIC-OL. (Cale Chem Co) Testosterone propionate 1n a1 Vial (25 mg in sesame oil/cc) 10 cc

Use: 1 M, androgen therapy

ORENZYME. (National) Trypsin 63%, chymotrypsin 30%, ribonuclease 2%/Tab Bot 45s

Use: Oral enzymes for treatment of inflammation and edema

ORESTRALYN. NND 1960 (McNeil) Ethinyf

contains stearic acid, cetyl alcohol glycerin ricinoleic acid, 5ad lauryl sulfato, baric acid & triethanolamine Pkg A, 2½ oz of cream w/measured dose; applicator B 2½ oz printed refill; Tube only, D refill, 2½ oz plain Tube only, large size 4½ oz Printed tube

Use: Spermicidal vaginal cream

ORTHO DERMA. (G F Harvey) Emulsion of crude coal tar Liq Bot 3 oz, 1 pt

Use: Antiseptic skin preparation

ORTNOFORM, NNR 1951 ADR 1952

No mfr listed Methyl 3 amino 4 hydroxy benzoate (Orthocaine) Bot 1 oz
Use Local anesthetic

W/Sulfamerazine & sulfadiazine Tab See Sulfocaine Tablets (Brewer)

W/Tyrothricin (Columbus Pharm) Tyrothricin 0.5 mg, tetracaine HCl 0.5%, epinephrine 1/1000 sol 2%/Gm Oint Tube ½ oz
Use: Eye infections

ORTNOFORM HEW. See: Orthoform (Winthrop Labs)

ORTNO-GYHOL, NNR 1955 (Ortho)

Water dispersible spermicidal jelly having pH of 4.5 which contains ricinoleic acid, p-diisobutylphenoxypolyethoxy ethanol in vegetable gum base Pkg A, 3 oz, Tube w/measured dose applicator; B 3 oz printed refill tube only; D Refill 3 oz, plain tube only Large size 5 oz, tube only
Use: Spermicidal jelly

ORTHO-HYDROXYBENZOIC ACID, USP
See: Salicylic Acid

ORTHO-PNEHYLPHEHOL. (Dorsey) Orthophenylphenol 5% in organic solvents & castor oil 2% Sol Bot 2 oz
Use: Topically; fungicide

ORTNOPNEHYLPNEHOL

See: O Syl (Lehn and Fink)

Phenolol Liq (Squibb)

ORTHOXICOL (Upjohn) Dexamethorphan Hydrobromide 10 mg Orthoxine NCI 17 mg, Sod um Citrate 325 mg fl oz Bot 2 oz, 4 oz, pt gal
Use: [1 tsp q 3 h] Coughs

ORTHOXINE (Methoxyphenamine) W/A P C & Pyrralozate See: Pyrralozate Cop (Upjohn)

W/Methylprednisolone See: Medrol W/Orthaxine Tab (Upjohn)

ORTHOXINE & AMINOPHYLLINE (Upjohn) Methoxyphenamine 50 mg, aminophylline 100 mg/Cap Bot 100s, 500s
Use: [1 q 4-6 h] Bronchodilator

ORTHOXINE HYDROCHLORIDE, NND 1961 (Upjohn) Methoxyphenamine NCI Tab 0.1 Gm, Bot 100s, 500s Syr 50 mg/5 cc., Bot 1 pt

Use: [1 Gm to 0.2 Gm q 4 h] Bronchodilator

Use: [1 Gm to 0.2 Gm q 4 h] Bronchodilator

W/Dihydrocodeinone bitorate, hyoscyamine HBr, Sod citrate 500 Orthoxicol, Liq (Upjohn)

W/Pyrralozate, oclophenetidin, aspirin & caffeine See: Pyrralozate, Cop (Upjohn)

OR-TOPTIC SUSP (Ortego) Polymyxin B sulfate 16,500 units hydrocortisone USP 5 mg neomycin sulfate USP 5 mg/cc & cc plastic dropper bottle

Use Ophth anti bacterial/anti fungal

ORVUS See: Gardinal Type Detergents (Various Mfr)

ORY-CIN (First Texas) Tyrothricin 1 mg, chlorpheniramine maleate 2 mg, benzocaine 5 mg /Trache Box 16s, 500s
Use: Mouth and throat infections

OSARSAL See: Acetarsone (Abbott)

OS-CAL (Marion) Oyster shells pow containing 40% Cal 625 mg, Vit D 500 u/Tab Bot 100s
Use: [1 tid] Calcium & Vit D therapy

OS-FAC (Metro Med) Osteogenic factors from cultured bone Tabs Bot 100s
Use: Bone degenerative

OSITOL (Dumas Wilson) Yeast conc 150 mg, inositol 200 mg, methionine 200 mg/Cop Bot 100s 1000s
Use: Lipotropic

OSMOPAK (Irwin Nessler) Mag sulfate 58% benzocaine & brilliant green Jar 3 oz 12 oz, 24 oz Dressing avail in rolls 8 in wide 25 ft long
Use: Varicose vein therapy

OSTAMER, NND 1962 (Merrell) Polyurethane Foam Ostamer prepolymer 61 Gm Ostamer catalyst 30 Om/Kit
Use: Grafting agent for bone

OSTENSIN, NND 1962 (Wyeth) d [N methyl N [α trimethylammoniumpropyl]] 1 methyl 8 8 dimethyl 3 oxabicyclo[3.2.1] octane d methasulfate Trimehidanium methasulfate 20 mg, 40 mg/Tabs Bot 100s
Use: Antihypertensive

OSVARSAN. See: Acetarsone (Abbott)

OS-VIM (Marion) Vit A 1500 u, D 300 u- B 1 0.37 mg B 2 0.60 mg, B-6 0.03 mg, C 25 mg, B 12 0.03 mcg, niacin 38 mg-pantothenic acid 0.3 mg, folic acid 0.002 mg, Vis E 0.03 mg, oyster shell pow 469 mg, w/Mag, Iran iodine copper Mang, zinc/Tab Bot 100s
Use: Vitamin & mineral supplement

O-STL (Lehn & Fink) ortho-Phenyl phenol (ortho hydroxy diphenol) Bot 1 gal 5 gal & 10 gal Drum 50 gal
Use: Diluted to 0.5 to 3% for germicidal disinfectant

OTALOINE. (G F Harvey) Chloral hydrate 20 gr, camphor 15½ gr phenol 10½ gr

- olive oil as solvent/fl. oz. Bot. $\frac{1}{2}$ oz., 1 pt.
Use: Analgesic
- OTAMYLON EAR DROPS.** (Winthrop Labs)
Sulfomylon HCl (sulfbenzamine) 5%, benzocaine 5% in propylene glycol. Bot. w/dropper 15 cc. Also with hydrocortisone (0.02%). Bot. w/dropper 15 cc.
Use: External otitis, otitis media & furuncular otitis.
- O. T. C. BARY OINTMENT.** (Mossengill)
Oint. Tube 1 oz.
Use: Astringent and antiseptic.
- O. T. C. FUNGACETIN.** (G. P. Horvey) Oint Tube 1 oz.
Use: Fungus infections.
- OTOBIONE.** (White) Prednisolone acetate 5 mg., neomycin 3.5 mg., Sod. propionate 50 mg./cc. Bot. 5 cc.
Use: Ear infections.
- OTOBIOTIC.** (White) Neomycin sulfate 3.5 mg., Sod. propionate 50 mg. in hydroalcohol glycerin sol./cc. Bot. w/dropper 15 cc.
Use: Otitis externa & chronic otitis media.
- OTOCAINE.** (Professional Pharm. Co) Antipyrine 5%, benzocaine 1%, alcohol 5% and phenylmercuric benzoate in propylene glycol. Bot. dropper 15 cc.
Use: Ear drops
- OTOCORT.** (Lemmon) Neomycin sulfate 5 mg., polymyxin B sulfate 2000 u., hydrocortisone 0.1%, antipyrine 5%, dibucaine hydrochloride 0.25% in hydro-glycerine vehicle/cc.
Use: Treatment of otitis externa and media.
- OTOOTYNE.** (White) Zolamine 1%, eucupin dihydrochloride 0.1%, polyethylene glycol q.s. Bot. w/dropper $\frac{1}{2}$ fl. oz.
Use: Ear drops.
- OTOLGESIC.** (Blue Line) Phenol 4%, benzocaine 10%, clove 4%, menthol 4%, in non-alcoholic nonoily solvent. 50 min./dropule Pkg. 12s.
Use: Inflammation of external ear canal.
- OTOMIOE.** (White) Sulfanilamide 5%, ureo 10%, chlorobutanol 3%, glycerin q.s. Bot. w/dropper $\frac{1}{2}$ fl. oz.
Use: External ear infections.
- OTO MYCIN SOL.** (Miserer) Cetalkonium chloride 0.05%, hydrocortisone USP. 50 mg., neomycin sulfate U.S.P. 60 mg., pyrimine maleate 0.6%, dipiperdon HCl 1%, ureo 5%. 5 cc. plastic drop containers
Use: Otitis externa.
- OTORAL.** (Hort) Acetic acid 1%, benzocaine 1%, saligenin 1% in a propylene glycol base. Bot. w/droppers $\frac{1}{2}$ oz.
Use: Otitis externa and otitis media.
- OTOS-MOSAN.** (Ayers) Urea 125 mg., sod. propionate 60 mg., sulfathiazole 0.60 mg., methylparaben 20 mg., benzethonium chloride 0.25 mg., glycerol ad to 1.0 cc. Otic solution, dropper bottle 19 Gm.
Use: Acute and chronic middle ear infections.
- OTOSONE-P.** (Broemmel) Hydrocortisone 0.5% in polyethylene glycol. Bot. 5 cc.
Use: Ear infections.
- OTOTHERON.** (Tilden) Benzocaine 11%, pine oil 0.8%, menthol 1.5%, guaiacol 0.4%, chlorothymol 0.4%, oil cloves 12%, liquid phenol 2%, camphor 0.4%. Bot. $\frac{1}{4}$ oz. w/dropper.
Use: Analgesic for earache.
- OTOTRIN EAR DROPS.** (Louisons) Neomycin sulfate 6 mg., hydrocortisone 5 mg., pyrimine maleate 0.6%, cetalkonium chloride 0.05%, dipiperdon HCl 1.0%, urea 5%/cc. 5 cc. plastic dropper bals.
Use: Otitis media and otitis externa.
- OTOZOLE.** (Hart) Sulfathiazole 3%, saligenin 5%, propylene glycol, q.s. Bot. $\frac{1}{2}$ oz., 8 oz.
Use: Bacteriostatic analgesic for otitis media.
- OTRATHOL.** (Mossengill) Atropine sulfate 1/100%, antipyrine 6%, urea 10%, menthol 2%, benzocaine 1%, propylene glycol. Bot. $\frac{1}{2}$ oz. dropper, 1 pt.
Use: Inflammation of the ear
- OTRIVIN.** N.N.D. 1962 (Ciba) Xylometazoline. 2 (4'-T. Ruly-2',6'-dimethylphenylmethyl)imidazole hydrochloride Sol (0.1%) Bot. 1 oz. Spray (0.1%) Tube 15 ml. Ped spray (0.05%) Tube 15 ml.
Use: Nasal congestion.
- *OUARAIN.** U.S.P., N.N.R. 1950, [nj], U.S.P. G-Strophanthin.
Tilly—Amp. (0.5 mg.) 2 cc., Box 12s, 100s.
C. D. Smith—Amp. (0.1 mg.) 0.5 cc., Box 5s, 25s; (0.5 mg.) 2 cc., Box 5s, 25s.
Varick—Amp. (0.25 mg.) 1 cc., Box 6s.
Use: Cardiac glycoside therapy.
- W/Orgitaline.** See: Natibaine, liq. (Varick)
- OVACOIDS.** (Reed & Cornick) Estradiol 0.1 mg./Tab. Bot. 100s, 500s, 1000s
Use: (2-3 q 1 d.) Hormone therapy.
- OVAOICL.** (C. O. Smith) Estrone 1 mg., estradiol 0.1 mg., benzyl alcohol 3% in peanut oil/cc. Vial 10 cc.
- OVAGEST.** (C. D. Smith) Estrone 1 mg., estradiol 0.1 mg., progesterone 20 mg., benzyl alcohol in peanut oil 3%/cc. Vial 10 cc.
- OVARIAN EXTRACT.** Aq. ext. of whole ovaries of cattle.
Endo—Amp. (2.6 Gm.) 1 cc., Box 12s, 100s. Vial 10 cc.

Hynson—Amp (0.2 Gm) 1 cc, Box 10s
 Lilly—Amp (0.65 Gm) 1 cc, Box 6s, 12s,
 100s; (2 Gm) 1 cc, Box 12s, 100s
 National Drug—Injectosal (2.6 Gm/cc)
 25 cc
 Parke, Davis—Amp (1.17 Gm) 1 cc, Box
 12s, 100s.
 Pitman Moore—Vial (2.6 Gm/cc) 25 cc
 Use 1 M & 5 C; estragen therapy

OVARIAN HORMONE, isolated from ovary
 See Lutrexin, Tab (Hynsan, Westcott &
 Dunning)

OVARIAN RESIDUE, DESICCATED. (Parke
 Davis) Ovaries w/carpus lutea removed
 from cattle, sheep or swine, desicc Cap 5
 gr, Bot 50s, 500s Emplats 5 gr, Bot 100s
 500s Paw Bot 1 oz, 1 lb
 Use: Orally; glandular therapy

OVARIAN RESIDUE EXTRACT Est w/
 carpus lutea removed, from cattle
 Horton & Canverse—Vial 30 cc
 Parke, Davis—Emplats, EC 5 gr, Bot
 100s 500s Pow 1 oz
 Labs.—Cap 5 gr, Bot 50s

OVARIAN SUBSTANCE (Various Mfr) Whole
 ovarian substance from cattle, sheep or
 swine

OVARIAN SUBSTANCE, DESICCATED
 (Parke, Davis) Whole ovarian substance
 from cattle, sheep or swine pow Cap 5
 gr, Bot 100s 500s Emplats 5 gr Bot
 100s, 500s Tab, 5 gr, Bot 100s, 500s
 Pow, Bot 1 oz, 1 lb
 Use Orally; glandular therapy

W/Thyroid, adrenal cortex fresh gland
 anterior pituitary fresh gland & posterior
 pituitary See: Metrocaine, Amp (Metro)

OVARY EXTRACT.
 See: Ovocoids, Cap (Carnrick)
 Stilensis, liq (Wollau)
OVEEN. (Warren Teed) Dayer's pow ¼ gr,
 acetaphenetidin 1½ gr, acetylsalicylic
 acid 2 gr, camphor manabramated ¼ gr
 caffeine ½ gr./Cap Bot 100s, 1000s
 Use Colds

OVIFOLLIN See Estrane (Various Mfr)

OVOCYLIN DIPROPIONATE, NND 1962
 (Ciba) Estradial dipropionate In oil Amp
 1 mg, 2.5 mg, 5 mg, Carton 6s, 50s Vial
 (1 mg., 5 mg) 10 cc.
 Use: f.M.; estrogen therapy

OVOFIN. (Warren Teed) Dayer's pow ¼
 gr, acetaphenetidin 1½ gr, atropine
 sulfate 1/500 gr, acetylsalicylic acid 2 gr,
 camphor ½ gr, caffeine ½ gr./Cap or
 Tab. Cap, Bot. 100s, 1000s Tab, Bot
 100s, 1000s.
 Use: Cold treatment

OVOFERRIN. (Crookes Barnes) Ferrous glu
 * Generic name recognized by N.F., U.S.P. or A.M.A.

conate 260 mg (iron 30 mg) Vit B 1 1 mg
 B 2 12 mg, B 3 10 mg B 6 1 mg, B 12
 4 mcg, alcohol 18% /fl az Bot 12 fl oz
 Use Dietary supplement

OVUTEST. (Eli Lilly) See Cervical Glucos
 Indicator

OWATE. (Paul Money) Butobarbital 20 mg,
 amphetamine sulfata 5 mg, sod ascorbate
 100 mg, thiamine HCl 0.5 mg, riboflavin
 10 mg, niacinamide 5 mg /tab
 Use: (1-3 daily) Obesity

OXACHOLIN. (Columbus) Oxidized bile
 acids 1.5 gr, ox bile 1.5 gr, steapsin ¼
 gr, hamatropine methylbromide 1/50 gr/
 Tab Bot 100s 1000s
 Use (1-2 i d) increase bile flow

***OXACILLIN, SODIUM,** 5 Methyl-3 phenyl
 4 isoxazolyl penicillin
 See Sodium oxacillin
 Prostoplin (Bristol)
 Restitopen (Squibb)

***OXADIMEDINE NCI** N (2 benzoxazolyl)-
 N benzyl N'N dimethylethylenediamine
 HCl

OXALINE. (Wyeth) Oxethazone (N,N bis(N
 methyl N phenyl tertiary butylacetamide)
 beta hydroxyethylamine HCl 10 mg in alu
 minum gel/5 cc Bot 12 oz

Use Anesthetic in gastritis and indigestion
OXALINE-M (Wyeth) Oxethazone 10 mg
 in alumina gel with the equiv of ¼ tea
 spoonful of milk of magnesia liq Bot 12
 oz

Use: Treatment of gastritis and indigestion
OXALAX. (Blue Line) Oxgall desicc 1 gr,
 aloin ½ gr, podophyllin 1/6 gr, ginger
 ½ gr./Tab Bot 100s, 1000s
 Use Cathartic

OXALIC ACID w/Malonic acid Inl See:
 Koagamis, Vial (Chatham)

OXAMYCIN. (Merck Sharp & Dahme) d-4-
 amino-3 isoxaxolidane Cyclaserline See
 Seromycin (Lilly)
 Use: Pulmonary TB

***OXANDROLONE** (G D Searle and Co)
 (1) Dodecahydro 3 hydroxy 6 (hydroxy
 methyl) 3,3a 6 trimethyl 1H benz(e)indene-
 7 acetic acid, Δ-lactone (2) 17β hydroxy
 17a methyl 2 axa 5a androstan 3-one See:
 Anavas (Searle)

***OXANAMIDE, NND 1962** 2 Ethyl 3 pro-
 pyl glycidamide See: Oxiactin, Tab
 (Merrell)

***OXAZEPAM.** (Wyeth) 7 chlora 1,3-d hydra-
 3 hydroxy 5 phenyl 2H 1,4 benzodiazep n
 2-one

Use: Management of anxiety tension states
OXAZOLIDIN. (Geigy Pharm) See: Oxy
 phenbutazone

- OX BILE EXTRACT, N.P., N.N.R. 1949.** Purified oxgall. See: Bile Extract, Ox
- OX & NOG BILE EXT. W/ Atropine methyl nitrate, oxidized bile acids, phenobarbital.** See: G. B. S., Tab. (G. F. Harvey)
- OXEDRINE.** (C. Bowman) Sympalol, Synephrine, N.N.R. 1934, 1-(p-Hydroxyphenyl)-2-Methylamino-ethanol tartrate. Pow. in bins. Use: Oral or topical epinephrine-like action
- OXETHANZINE.** (Registered in Great Britain) 2 Di-(N-methyl-N phenyl tert butyl carbamoylmethyl)-aminoethanol
- * **OXETHAZINE.** (N,N-bis-(N-methyl-N-phenyl-1-butyl-acetamido) beta-hydroxyethylamine) See: Oxaine (Wyeth)
- OXIDIZED BILE ACIDS.** See: Bile Acids, Oxidized.
- OXIDIZED CELLULOSE, U.S.P., N.N.D. 1967.** A.D.R. Absorbable cellulose
See: Hemo-Pak. (Johnson & Johnson)
Oxycel, N.N.R. (Parke, Davis)
Use: Local hemostatic.
- OXINE.** See: Oxyquinoline sulfate
- OXIPHEN.** (Pitman-Moore) Sod. glycocholate $\frac{1}{4}$ gr., Sod. taurocholate $\frac{1}{4}$ gr., phenolphthalein $\frac{1}{2}$ gr., cascara comp. ext. $\frac{1}{2}$ gr., oilain $\frac{1}{4}$ gr./tab. Bot. 100s, 500s.
Use: Laxative.
- OX GALL.** See: Bile Extract, Ox
- * **OXONAZINE.** (Upjohn) 2,4-diamino-6-(diethylamino) s-triazine, N⁶oxide.
Use: Antihypertensive agent
- * **OXOPHENARISINE HYDROCHLORIDE, N.F., N.N.R. 1955** 3-Amino-4-hydroxyphenylarsinoxide HCl. See: Maphorsen, Amp (Parke, Davis)
- * **OXPHENRIDINE.** 1-(β -phenyl- β hydroxyethyl)-4 carbethoxy-4-phenylpiperidine.
- OXSORALEN, N.N.D. 1962** (Elder) Methoxsalen 8-Methoxypsoralen (Armalidin, Xanthoxalin.) Cap. (10 mg) Bot. 100s. Lot. (1%) Bot. 1 oz.
Use: 2 Cap. twice daily and/or topically idiopathic vitiligo.
- OXSORBIL.** (Ives Cameron) Dehydrocholic acid $\frac{1}{2}$ gr., desoxycholic acid $\frac{1}{2}$ gr., ox bile ext. 1 gr., sorbitan monooleate 2 $\frac{1}{2}$ gr., oleic acid 2 $\frac{3}{4}$ gr./Cap Bot. 100s.
Use: Choleresis & chologogia.
- OXSORBIL P.S.** (Ives-Cameron) Same formula as Oxorbil w/phenobarbital $\frac{1}{8}$ gr., belladonna pow. ext. $\frac{1}{8}$ gr. added/Cap. Bot. 100s.
Use: (1-2 i d.) Choleric & hydrocholeretic.
- * **OXTRIPHYLLINE, N.N.D. 1962** See: Chaledyl, Tab (Nepera)
W/glyceril guaicolate (Warner Chilcott)
See: Brandoran.

* Generic name recognized by N.P., U.S.P. or A.M.A.

- OXUCIDE.** (Brean) Piperazine citrate 500 mg./5 cc. Bot. 1 pl.
Use: Pinworm & roundworm anthelmintic.
- * **OXYBUTYNIN CHLORIDE.** (Mead Johnson) 4-diethylamino-2-butynyl phenylcyclohexaneglycolate hydrchloride
Use: Antispasmodic agent
- OXYCEL, N.N.D. 1961.** A.D.R. (Parke, Davis) Sterile oxidized cellulose. Pledgets 2 $\frac{1}{4}$ " x 1" x 1"; gauze pads 3" x 3", 8 ply; 4" x 12", 8 ply; gauze strips 5" x $\frac{1}{2}$ ", 4 ply; 2" x 18", 4 ply; 36" x $\frac{1}{2}$ ", 4 ply; 2" x 3 yds, 4 ply; Foley cones 5" diameter, 4 ply; 7" diameter, 4 ply.
Use: Hemostatic.
- * **OXYCHLOROSENE.** (Guardian Chemical Corp) Monochlorosene Hydrocarbon derivative containing fourteen carbons and hypochlorous acid. The hydrocarbon chain also has a phenyl substituent which in turn holds a sulfonic acid group
See Clorpectin XCB (Guardian Chemical Corp)
Clorpectin WCS-90 (Guardian Chemical Corp)
Worezin. (Guardian Chem.)
- OXYCODONE HCl.** See: Dihydrohydroxycodeinone HCl
- OXYDESS.** (Chlmedic) di-Desoxyephedrine HCl 8 mg./Tab. or dr. Elixir, Bot. 1 pl., 1 gal. Tab. Bot. 100s, 1000s.
Use: Central nervous system stimulant.
- OXYDESSIN.** (Chicago Pharm) di-Desoxyephedrine HCl 8 mg., Vit. A 1350 u., D 133 u., B-1 0.34 mg., B-2 0.67 mg., B-12 1 mcg., C 20 mg./tab. Bot. 100s, 1000s.
Use: Antiobesity.
- OXYDRIN.** (Grant) Methamphetamine HCl Tab. (2.5 & 5 mg.) Bot. 100s.
Use: Central nervous system stimulant.
- OXYETHYLATED TERTIARY OCTYL-PHENOL-FORMALDEHYDE POLYMER.** See: Tison WR-1339. (Winthrop Labs)
- OXYETHYLENE OXYPROPYLENE POLYMER.** See: Poloxalkol.
- W/Lecithin, dextrose anhydrous, cottonseed oil.** See: Iipomul IV. (Upjohn)
- OXYTED.** (Cole) di-Desoxyephedrine HCl 5 mg./Tab. Bot. 100s, 500s, 1000s.
Use: Central nervous system stimulant.
- OXYGEN, U.S.P.**
See: Let O Gen (Lil O Gen, Inc)
Oxy-Hale (Inhalational Equipment Co)
Oxy Swig (Oxy Swig, Inc)
- OXYGEN-CARBON DIOXIDE MIXTURES, N.N.R. 1947.**
- OXY-HALE.** (Inhalational Equipment Co)

- Pocket size cylinder of oxygen Enough for 1½ minutes Refill cartridges, pkg 6s
Use Oxygen therapy
- OXYL-IODIDE** (Lilly) Cinchophen Hb Tab 3 gr, Bot 100s, 500s Also W/Cinchophen Ensoles and Pulvules Bot 100s, 500s
Use: Gout
- OXYLONE, NND 1961** (Upjohn) Elwara metholone 0.025% in a water soluble cream base Tube 7.5 Gm
Use Dermatitis
- * **OXYMETAZOLINE HYDROCHLORIDE** (Schering) 2,6 dimethyl 4 tertiarybutyl 3 hydroxyphenyl methylimidazoline hydrochloride
Use Decongestant
- * **OXYMETHOLONE, 17 beta hydroxy 2 hydroxymethylene 17 alpha methyl 3 androstanone**
See Adroyd, Tab (Parke Davis)
Anadrol (Syntex Labs Inc)
- * **OXYMORPHONE, HCl, NND 1962** 1,4 hydroxydihydro morphinone HCl See Nemorphan (Endo)
- OXY-H-ALATOR** (Inhalational Equipment H Y) Oxygen in portable apparatus 135 l and 75 l
Use Heart and respiratory conditions
- * **OXYPTERINE HYDROCHLORIDE** (Winthrop Labs) 5,6 dimethoxy 2 methyl 3 [2 (4 phenyl 1 piperazinyl)ethyl]indole hydrochloride
- * **OXYPHENBUTAZONE, 1 (p hydroxy phenyl) 2 phenyl 4 butyl 3,5 pyrazolidine dione**
See Oxazolindin (Geigy Pharm)
Tondeonil (Geigy)
- * **OXYPHENCYCLIMINE HCl, NND 1962** 1 methyl 1,4,5,6 tetrahydro 2 pyrimidinylmethyl alpha cyclohexyl phenylglycolate HCl
See Daricon Tab (Pfizer)
Via Thene, Tab (Rowell)
- W/Hydroxyxine HCl See Enarax, Tab (Roerig)
- * **OXYPHENISATIN 3,3 bis(4 hydroxy phenyl) oxindole Dihydroxyphenylisatin Dihydroxyphenoloxindol Related to dia cetylhydroxyphenylisatin See Lavema Pow (Winthrop Labs)**
- * **OXYPHENISATIN ACETATE** (Warner Chilcott) 3,3 bis(p hydroxyphenyl)2 indolin one d acetate See Isocaric Fruitet
- * **OXYPHENONIUM BROMIDE, NND 1962** Diethyl [2 hydroxyethyl] methylammonium bromide alpha phenyl cyclohexyl glycolate See: Antrenyl Bromide (Ciba)
W/Phenobarbital See Antrenyl Bromide Phenobarbital Tab (Ciba)
- * **Generic name recognized by NF USP or A.M.A.**
- * **OXYPHENUDRINE, (Mead Johnson and Co)** α {[(p-hydroxy α methylphenethyl) amino]methyl}protocatechuyal alcohol
- OXYQUINOLINE BENZOATE, H H R 1949** (Mallinckrodt; Merck) Pkg 1 lb 8 Hydroxy quinoline
W/Chlorophyll See: Chlorothelio, Oint (Arnar Stone)
W/Phenol, alum, boric acid, etc See: Bon Du, Pow (Warren Teed)
W/Tragacanth, quince See: Broom, Lot (Brean)
- OXYQUINOLINE SULFATE, (Mallinckrodt Merck Sharp & Dahme)** Pkg 1 lb 8 Hydroxyquinoline, oxine
See: Chinosal, Tab (Jewell)
Octafen, Liq & Pow (McKesson & Robbins)
W/Anesthasin, ephedrine HCl & Ichthammol See: Derma Medicine, Oint (Medicone).
W/Boric acid, mag sulf See: Braxalin Pow (Brean)
W/Boric acid mag sulf, sod chlor See: Aguapow, Pow (Chicago Pharm)
W/Boric acid, sod chlor, sod sulfa diethyl succinate See: Certane Dauche Pow (Vogorell)
W/Hydrocortisone acetate benzocaine ephedrine HCl menthol ichthammol zinc oxide See: Derma Medicine HC (Medicone)
W/Lactic acid, zinc sulfocarbonate, citric acid & dextrose See Acidusal, Sol (Taxis).
- OXYSORALEN, 8 Hydroxy-4',5',6,7 tetracoumarin** See: Methaxalen
- OXY SWIG, (Oxy Swig Inc)** A portable unit of 56 liters of oxygen ready for emergency use
- * **OXYTETRACYCLINE, NF NND 1962**
See Terramycin (Pfizer)
- * **OXYTETRACYCLINE HYDROCHLORIDE, USP NND 1962, ADR; An antibiotic from Streptomyces rimosus Cop, Inj Ophth, USP** See Terramycin Hydrochloride (Pfizer)
W/nystatin See Terrastatin Caps (Pfizer)
W/buchazine HCl, phenacetyl salicylamide caffeine See Terracydin Caps (Pfizer)
W/Sulfamethizole phenylaxodiaminopyridine HCl See: Urobitalc, Cop (Pfizer)
W/Vitamins & stress formula See Terramycin SF, Cap & Susp (Pfizer)
- OXYTETRACYCLINE FOR ORAL SUSPENSION, USP** See Terramycin Oral Susp (Pfizer)
- OXYTETRACYCLINE-POLYMYXIN B, NNR 1955** Mixture of oxytetracycline HCl & polymyxin B sulfate See Terramycin Hydrochloride w/Polymyxin B Sulfate (Pfizer).

- OXYTOCICS.** See Ergol Prep.
- *OXYTOCIN INJECTION,** U.S.P., N.N.R. 1952
See: Pitocin, Amp. (Parke, Davis)
Syntocinon, Amp. (Sandoz)
Syntocinon Nasol Spray. (Sandoz)
- OXYTOCIN, SYNTHETIC.** See: Syntocinon, Vial. (Sandoz)
- OXYVATINE,** (Smith-Dorsey) Thiamine HCl 10 mg., riboflavin 5 mg., ascorbic acid 30 mg., nicotinic acid 25 mg./Tab. Bot. 100s, 1000s.
Use: Vitamin supplement.
- OYSTER SHELES.** See: Os-Cal. (Marion)
- OZONIDES OF OLIVE OIL.** See: Olizo, liq (G. F. Harvey)
- PAADON CAPSULES.** (Rorer) Phenobarbital 15 mg., acetylsalicylic acid 180 mg., acetylphenelidin 120 mg., belladonna sulfates 0.0625 mg./Cap. Bot. 100s, 500s.
Use: (1 q, 3 h.) Analgesic & sedative.
- PABA.** (Ives-Cameron) p-Aminobenzoic acid. Tab. 100 mg., Bot. 100s.
Use: Dietary supplement.
- PABACAL.** (Ives-Cameron) p-Aminobenzoic acid 100 mg., Cal. alpha-pantothenate 10 mg./Tab. Bot. 100s.
Use: Dietary supplement.
- PABACEE.** (Paul Money) Sod. salicylate 6 gr., ascorbic acid 20 mg., menadione 0.5 mg., Pot. p-aminobenzoate 3 gr./Tab. Bot. 100s, 1000s.
Use: Rheumatic fever.
- PABACEE W/COLCHICINE.** (Paul Money) Sod. salicylate 0.4 Gm., pot. p-aminobenzoate 0.2 Gm., ascorbic acid 20 mg., menadione 0.5 mg., colchicine/Tab. Bot. 100s, 1000s.
- PABACEE (SODIUM FREE).** (Paul Money) Ammonium salicylate 0.4 Gm., ascorbic acid 20 mg., menadione 0.5 mg., pot. p-aminobenzoate 0.2 Gm./Tab. Bot. 100s, 1000s.
- PABAGESIC.** (Lynn) Acetylsalicylic acid 5 gr., para-aminobenzoic acid 5 gr., Vit. C 50 mg., aluminum hydroxide gel 2 gr/Tab Bot. 100s, 500s.
Use: Rheumatoid and gouty arthritis.
- PABALATE.** (Robins) Sod. salicylate & Sod. p-aminobenzoate al each, 5 gr., vit. C 50 mg. in E.C. Tab., or 5 cc. chocolate flavored liq. liq. Bot. 1 pt., 1 gal. Tab. Bot. 100s, 500s.
Use: (2 tabs. or tsp. q i d.) Antirheumatic.
- PABALATE-HC.** (Robins) Hydrocortisone alcohol 2.5 mg., pot. salicylate 300 mg., pot
- p-aminobenzoate** 300 mg., ascorbic acid 50 mg./Tab. Bot. 100s.
Use: (1-2 i.i.d.) Rheumatoid arthritis.
- PABALATE-SODIUM FREE.** (Robins) Ammonium salicylate & Pot. p-aminobenzoate 5 gr. each, vit. C 50 mg./Tab. Bot. 100s, 500s.
Use: (2 tabs. q i d.) Antirheumatic.
- PABAMIDE.** (Rowell) Salicylamide 5 gr., Pot. p-aminobenzoate 5 gr./Cap. Bot. 100s.
Use: Antirheumatic.
- PABA-SALICYLATE.** (West Ward) Sodium salicylate 5 gr., p-aminobenzoate 5 gr., Vit. C/Tab. Bot. 100s.
Use: Antirheumatic.
- PABA SODIUM.** Sod. p-aminobenzoate. Ives-Cameron—Tab. 0.5 Gm., Bot. 100s, 1000s.
Wyeth—Tab. 0.5 Gm., Bot. 100s, 1000s.
Use: Antirheumatic agent.
- PABASYL.** (Ives-Cameron) Sod. p-aminobenzoate 5 gr., Sod. salicylate 5 gr., ascorbic acid 10 mg./Tab. Bot. 100s, 1000s
Use: (2 i. d.) Rheumatic conditions.
- PABASYL SODIUM-FREE.** (Ives-Cameron) Ammonium salicylate 5 gr., p-aminobenzoic acid 5 gr., Vit. C 50 mg./Tab. Bot. 100s.
Use: 1-2 i.i.d., rheumatic therapy.
- PA-BEX.** (Brewer) Vit. B-1 5 mg., B-2 5 mg., B-6 1.5 mg., Cal. pantothenate 25 mg., niacinamide 50 mg., Vit. C 150 mg./Tab. Bot. 100s, 500s, 1000s.
Use: Nutritional supplement.
- PABIBIN.** (Smith-Dorsey) p-Aminobenzoic acid 5 gr., aspirin 5 gr., ascorbic acid 50 mg./Cap. Bot. 100s, 1000s.
Use: (1-2 i.i.d.) Analgesic.
- W/Codeine** 1/4 gr./Cap. Bot. 100s, 500s, 1000s.
- PABIBIN-AC.** (Smith-Dorsey) p-Aminobenzoic acid 5 gr., aspirin 5 gr., ascorbic acid 50 mg., hydrocortisone 2.5 mg./Cap. Bot. 50s.
Also avail. as Buffered Tabs.
Use: Analgesic w/hydrocortisone therapy.
- PABLUM BABY CEREAL.** (Mead Johnson) Carbohydrate 77.1%, protein 10.5%, fat 0.3%, minerals & 4% including Cal., Phos., iron & copper, Con B oz.
Use: Infant food.
- PABLUM MIXED CEREAL.** (Mead Johnson) Wheatmeal 52%, oatmeal 18%, wheat germ 15%, cornmeal 10%, beef bone pow. 2%, Sod. chloride 1%, alfalfa leaf 1%, primary dried yeast 1% & reduced iron. Pkg. 8 oz., 1 lb.
Use: Infant food.
- PABLUM OATMEAL.** (Mead Johnson) Oatmeal 90.3%, malt syrup 4.7%, beef bone pow. 2.3%, Sod. chloride 1.5%, primary

* Generic name recognized by N.F., U.S.P. or A.M.A.

dried yeast 12% & reduced iron Pkg 8 oz

Use: Infant food

PABLUM RICE CEREAL. (Mead Johnson) Carbohydrate 81.7%, protein 6%, fat 0.2%, minerals 3.8% including Cal 0.79%, Phos 0.53%, iron 0.03%, copper 0.0006 Pkg 8 oz.

Use Infant food

PABODYNE. (Blue Line) Acetylsalicylic acid 5 gr, para-aminobenzoic acid 5 gr, ascorbic acid $\frac{3}{4}$ gr/Tab Bot 100s, 1000s

Use Management of rheumatoid disorders

PABUTOLE (Merck Sharp & Dohme) Vit A 2400 u thiamine HCl 0.5 mg, riboflavin 1 mg, vit B 0.440 u, hypophosphites of cal iron, mong, pol, sod, sod salicylate, sod citrate, alcohol/Tsp Bot 14 R oz

Use Otolary supplement

P-A-C. Prep See: A.P.C. Preparations B Empirin Preparations

P-A-C COMPOUND. (Upjohn) Acetophenetidin $2\frac{1}{2}$ gr, acetylsalicylic acid $3\frac{1}{2}$ gr, caffeine $\frac{1}{2}$ gr/Tab & Cap Bot 100s, 1000s Also avail w/codeine sulfate Cap $\frac{1}{4}$ gr, $\frac{1}{2}$ gr, Bot 100s, 500s, 1000s Tab $\frac{3}{4}$ gr, $\frac{1}{2}$ gr, Bot 100s, 500s, 1000s

Use Analgesic

P-A-C COMPOUND, Half Strength (Upjohn) Acetophenetidin $1\frac{1}{4}$ gr, acetylsalicylic acid $1\frac{3}{4}$ gr, caffeine $\frac{1}{4}$ gr/Tab Bot 1000s Also avail w/codeine sulfate $\frac{3}{4}$ gr, $\frac{1}{2}$ gr Tab Bot 100s, 500s, 1000s

Use Analgesic

W/Cyclopal (Upjohn) Cyclopentylalyl barbituric acid $\frac{3}{4}$ gr, acetophenetidin $2\frac{1}{2}$ gr, acetylsalicylic acid $3\frac{1}{2}$ gr, caffeine $\frac{1}{2}$ gr/Tab Bot 100s, 500s, 1000s

Use Analgesic

PACATAL, Acetate & HCl NND 1961 (Warner Chilcott) Mepazine, 10 (N methyl 3 piperidylmethyl) phenothiazine acetate & HCl Tab 25 mg, 50 mg Bot 100s 100 mg Bot 500s Amp 125 mg/cc 2 cc)

Use Ataractic agent

PACINOL. (Schenck) Fluphenazine 0.25 mg/Orange

Use Ataractic

PACTAMYCIN. (Upjohn) Antibiotic Uefer study

P-A-D. (Upjohn) Acetophenetidin $2\frac{1}{2}$ gr acetylsalicylic acid $3\frac{1}{2}$ gr, dl desoxyephedrine HCl $1/26$ gr/Tab Bot 100s, 500s, 1000s

Use: Antispasmodic & analgesic

PADRIHA. (Merck Sharp & Dohme) Dihydrocodeinane bitartrate 5 mg methapyrifene HCl 50 mg, d amphetamine sulfate 2.5 mg/Tab Bot 100s

Use: Analgesic & antitusive

PADROPHYLL TABLETS. (Merck Sharp & Dohme) Propadrine HCl 30 mg, theophylline 0.12 Gm /Tab Bot 100s, 1000s

Use: Asthmatic paroxysms & bronchial spasms

PADUTIN. (Wiethrop Labs) Insulin free ext from pancreatic gland which is biologically stabilized so that 1 u counteracts pressor effect from minimal hypertensive dose of epinephrine Amp 10 u, Box 10s, 50s, Tab 10 u, Bot 100s

Use: Peripheral vasodilator

PAGITANE HYDROCHLORIDE, NND 1952 (Lilly) (Cyzimine HCl) phenyl 1 cyclophenyl 3 piperidina 1 prapanal HCl Tab 1.25 mg 2.5 mg, Bot 100s, 1000s

Use: (1.25 mg t i d) Antispasmodic

PAHA. See: p Aminohippuric Acid NF

PAL COLD TABS. (Pharm of Arkansas) Chlorpheniramine maleate 2 mg, acetylsalicylic acid $2\frac{1}{2}$ gr acetophenetidin $2\frac{1}{2}$ gr phenylephrine HCl 10 mg/Tab Bot 100s

Use Cammoe cold

PALADAC. (Parke, Davis) Vit A 5000 u, D 1000 u, B 1.3 mg B 2.3 mg, B 6 1 mg, B 12 5 mcg, C 50 mg niacin 20 mg pantothenic acid 5 mg/dr Bot 4 oz 1 pt

Use: Multivitamin therapy

PALADAC W/MINERALS (Parke Davis) Vit A 4000 u O 400 u C 50 mg E 3 IU B 1.3 mg B 2.3 mg B 6 1 mg B 12 5 mcg, nicotinamide 20 mg pantothenic acid 5 mg plus calcium, phosphorus iodine pot iron and magnesium/Chewable Tab Bot 30s 100s

Use Vitamin and mineral supplement

PALADEIN (Bristol) Codeine phosphate 0.065 Gm, terpin hydrate 0.05 Gm Sod citrate 0.650 Gm Pot guaiacol sulfonate 0.510 Gm, citric acid 0.190 Gm, squill camp syr 0.540 cc, chloroform 0.09 cc, ipecac fl ext 0.06 cc, menthol 0.003 Gm, alcohol 5%/fl oz Bot 1 pt 1 gal

Use (1 tsp q 3-4 h) Coughs

PALAGREN. (C B Kendall) Phenobarbital Sod 2 gr, Sod bromide 60 gr, Pot bromide 18 gr, Cal bromide 2 gr/fl oz Bot 8 oz

Use (1 tsp t i d & 2 h s) Sedative

PALAPENT. (Bristol) Elixir pentobarbital Sod 2 gr/fl oz Bot 12 oz, 1 gal

Use Sedative

PALARON (Smith Miller & Patch) Ferrout fumarate 200 mg, ascorbic acid 125 mg/Tab Bot 100s

Use Anemias

PALATRIZINE, PEDIATABS, NNR 1955 (Columbus) Sulfadiazine sulfamerazine.

sulfamethazine 42 mg of each/tab. Bot. 100s.

Use: Triple sulfonamide therapy.

PALBAR. (Palmedica) Phenobarbital 16.2 mg, atropine sulfate 0.012 mg, scopolamine HBr 0.005 mg, hyoscyamine HBr 0.18 mg / Tab Bot. 100s, 1000s. Elixir 23% alcohol / 5 cc.

Use: Sedative-spasmodic effect.

PALCILLIN. (Phila. Ampoule) Procaine penicillin G 400,000 u. in 0.5 Gm. of streptomycin sulfate / 2 cc 10 cc. Vial

Use: Infection caused by organisms susceptible to penicillin

PALESTROL. See: Diethylstilbestrol (Various Mfrs.)

PALFIUM. (G. P. Harvey) (d-1-[3-methyl-4-morpholino-2,2-diphenyl-butyl] pyrrolidine) Dextramoramide 5 mg / Tab, Bot. 100s.

Use: Analgesic.

PALICYLATE. (Ascher) Tab. Sod. salicylate 0.25 Gm., p-aminobenzoic acid 0.25 Gm., ascorbic acid 20 mg./Tab. Bot. 100s, 1000s. liq. Sod. salicylate 0.32 Gm., p-aminobenzoic acid 0.32 Gm./dr. Bot. 1 pt., 1 gal.

Use: (1-2 q.i.d.) Antirheumatic.

PALICYMIN. (Ascher) Sod. salicylate 5 gr., p-aminobenzoic acid 5 gr./dr. Syr., Bot 1 pt.

Use: Geriatric patients

W/Colchicine. (Ascher) Sod. salicylate 3.85 gr., p-aminobenzoic acid 3.85 gr., ascorbic acid 1/2 gr., colchicine 1/250 gr./Tab. Bot 100s, 1000s.

Use: (1-2 q i d) Pain in gout.

PALICYMIN-C. (Ascher) Sod. salicylate 3.85 gr., p-aminobenzoic acid 3.85 gr., ascorbic acid 1/2 gr./Tab. Bot, 200s, 1000s.

Use: (1-2 i.i.d.) Arthritis & rheumatic fever.

PALINUM. See: Cyclobarbitol (Phanodorn. Winthrop Labs.)

PALITINE. (Ascher) Sod. salicylate 3.85 gr., p-aminobenzoic acid 3.85 gr., ascorbic acid 1/2 gr., physostigmine salicylate 1/250 gr., hamatrapine methylbromide 1/170 gr./Tab. Bot. 100s, 1000s.

Use: (1 i.i.d.) Muscular relaxation.

PALIVITE. (Sherman) Vit. B-12 10 mcg, folic acid 5 mg., liver ext. crude 2 v./1 cc. Vial 10 cc.

Use: (1-3 cc. weekly) Macrocyclic anemias.

PALLACIN. (Denob) N-acetyl p-amino-phenol 300 mg, hyoscyne hydrobromide 0.075 mg, butabarbital sodium 10 mg / Tab Bot 50s, 250s.

Use: Analgesic.

* **PALMIOROL.** N (2-hydroxyethyl)palmitamide.

* Generic name recognized by N F, U S P. or A.M.A.

PALMINATE. (Palmedica) Vit A 4000 units, C 800 units, thiamine HCl 3 mg, riboflavin 2 mg, pyridoxine HCl 6 mg, B-12 2 mcg, niacinamide 15 mg, lactic acid 0.1 mg, rutin 15 mg, C 100 mg, K 0.5 mg, E 1 mg, calcium carbonate 1000 mg, calcium lactate 300 mg, aluminum hydroxide gel 200 mg, ferrous fumarate 200 mg, Ca .3 mg, Cu .5 mg, Mo 0.25 mg, Zn .1 mg, Mn .05 mg, Mg 1 mg, K 1 mg, F 0.015 mg, I .1 mg / Tab. Bot. 60s, 1000s.

Use: Pre-natal vitamin-mineral combination
PALMINATE-F. (Palmedica) Vit. A 4000 u, D 800 U, B-1 3 mg., B-2 2 mg, B-6 6 mg, B-12 2 mcg, niacinamide 15 mg, rutin 15 mg., Vit. C 100 mg., K 0.5 mg, E 1 mg, calcium carbonate 500 mg, calcium lactate 500 mg, aluminum hydroxide gel 200 mg, ferrous fumarate 200 mg., manganese .05 mg, magnesium 1 mg, sodium fluoride 2.2 mg./Tab Bot. 60s, 300s.

Use: Dietary supplement.

PALMIRON. (Palmedica) Ferrous fumarate 300 mg, intrinsic factor concentrate 1/2 U.S.P. oral unit, Vit. B-12 12 mcg., C 130 mg, thiamine HCl 3 mg / Tab. Bot 60s, 1000s.

Use: Anemias and wherever a complete hematonic is desired.

PALMIRON-C TABS. (Palmedica) Ferrous fumarate 200 mg, ascorbic acid 120 mg, Vit. B 6 5 mg / chewable tab. Bot. 100s.

Use: As hematonic.

PALOEISE. (Palmedica) Betaine anhydrous 75 mg, d-amphetamine 15 mg, amobarbital 16 mg., mixed oxidized bile acids 194 mg./Tab. Bot. 100s.

Use: Obesity control.

PALONIST. (Palmedica) Phenylephrine hydrochloride 25 mg, chlorophenpyridine amine maleate 7.5 mg, pyrilamine maleate 25 mg, methoprylene hydrochloride 12.5 mg / S R Cap Bot. 36s, 100s.

Use: Symptomatic relief of colds

PALONAD. (Palmedica) Vit A 6000 units, D 1200 units, B-12 10 mcg, thiamine mononitrate 5 mg, riboflavin 5 mg, niacinamide 15 mg, pyridoxine HCl .5 mg, lactic acid .25 mg, calcium pantothenate 5 mg, C 66 mg, dl-tocopherol acetate 10 I.U., ferrous fumarate 30 mg, rutin 2.5 mg, I .1 mg, Cu 1 mg, Ca .1 mg., Mn 5 mg., Mg 1 mg, K 5 mg, Zn 1 mg, Mo .2 mg, F .1 mg / Tab. Bot 100s, 1000s.

Use: Vitamin-mineral supplement.

PALONYL. (Palmedica) Ethinyl estradiol 0.05 mg./Tab. Bot 100s.

Use: Menopausal syndrome, functional uterine bleedings.

PALSORB W/NEOMYCIN. (Palmedico) Activated attopulgites 330 Gm, pectin (100 grade) 0.60 Gm, sodium and potassium lactates 30 Gm, neomycin sulfate D3 Gm, glucose USP 6.60 Gm, sucrose USP 90 Gm, benzoin orid 0.12 Gm/cc Bot pf
 Use: Diarrhea

PALUDRINE. (Ayerst) Chloroquine HCl Proguanil HCl Tab 0.1 Gm, Bal 100s
 Use: Malignant tertian malaria

PALVITE. (Ehrhart Pharm) Vit A 5000 u, D 1000 u, B 1 10 mg, B 2 1 mg, niacinamide 20 mg, Vit C 75 mg, B 6 1 mg, l-lysine 60 mg, B 12 25 mcg/dr Bot 3 oz
 Use: Vitamin supplement

PAM (Barden Co) 2 Pyridine aldazine methiodide

Use: Antidote for a lethal dose of nerve gas
PAMA (Chicago Pharm) Aluminum hydroxide 4 gr Mag trisilicate 4 gr, phenobarbital 1/8 gr, atropine sulfate 1/500 gr/Tab Bot 100s, 1000s, 6000s
 Use: Antispasmodic & sedative

PAMABROM. 2 Amino 2 methylpropanol 8 bromotheophyllinate See: Pseudoephedrine (Braylen)

W/Acetoephenedin See: M Mucus S, Tab (Whittier)

W/Dextroamphetamine sulfate, carbamol, salicylamide See Pambramal, Tab (Whittier)

W/Pyridamine maleate (Bromateate)

See Neo Bramin, Tab (Brytlen)

Neo Mensalen (Walker)

Neo Parbram, Tab (Central)

Neo Pyrapar Tab (Ulmer)

Sanbram, Tab (Smith Darsey)

W/Pyridamine maleate, homotropine methyl bromide, hyoscyamine sulf, scopolamine HBr, methamphetamine HCl See Neoparbel Tab (Central)

PAMAQUINE NAPHTHOATE. (Mann Fine Chemicals) (Aminoquin, Beprachone, Gamefar, Plasmquine, Praequine, Quipenyl) See: Plasmochin Naphthoate, Prep (Winthrop Labs)

PAMCO (Kirk Labs) Pentylentetrazol 50 mg nicotinic acid 25 mg glutamic acid 60 mg/C/T Tab Bot 100s 1000s
 Use: Geriatric energizer

PAMINE BROMIDE, NND 1962 (Upjohn)

Epoxitropine tropate methylbromide Methsropalome bromide Sropalome methyl bromide Tab 2.5 mg Bot 100s

Use: (1 tab t.i.d. p.c. & h.s.) Antispasmodic

PAMINE A PHENOBARBITAL DROPS (Upjohn) Methscopolamine bromide 0.5 mg, phenobarbital 20 mg, alcohol 20%/cc Bot 15 rr

Use: Gastro Intestinal disturbances

PAMINE & PHENOBARBITAL ELIXIR. (Upjohn) Methscopolamine bromide 125 mg, phenobarbital 8 mg, alcohol 20%/5 cc Bot 1 pt, 1 gal

Use: Parasympatholytic agent (anticholinergic) for peptic ulcer

PAMINE & PHENOBARBITAL TABLETS (Upjohn) Methscopolamine bromide 2.5 mg, phenobarbital 15 mg/Tab Bot 100s

Also available in half strength

Use: Gastro Intestinal disturbances

PAMINE STERILE SOLUTION. (Upjohn) Methsropalome bromide Vial (1 mg./cc.) 1 cc Pkg 1s, 5s 25s

Use: S.C., I.M., anticholinergic

PAMINE SYRUP. (Upjohn) Methsropalome bromide 125 mg, & other gastro-intestinal therapeutic agents/5 cc Bot 4 oz

Use: Peptic ulcer therapy

PAMISYL, NND 1962 (Parke, Davis) p-Aminosalicylic acid Tab D3 Gm, O5 Gm, Bot 500s, 1000s

Use: Tuberculosis

PAMISYL SODIUM, NND 1962 (Parke Davis) p-Aminosalicylate Sod Tab 0.5 Gm, Bot 500s, 1000s

PAMISYL SODIUM STERI-VIALS. (Parke Davis) p-Aminosalicylate Sod 69 Gm, equiv to 5 Gm acid Vial 1s, 10s

Use: I.V.; tuberculosis

PANAC (Brewer) Aspirin 5 Gm, Vit C 15 mg, calcium pantothenate 1 mg/Tab Bot 100s

Use: Analgesic

PANACARE. (Lanett) Bismuth subcarb., sed bicarb, mag carb, papain, diastase/tab Bot 1000s, 5000s

Use: Antacid

PANADOL See: N-Acetyl p-aminophenol

PANADON (Gordon Labs) Penthonal 2% Cream, Pkg 1 oz, 2 oz

Use: Treatment of dermatitis

PANAFIL (Rystan) Papain pow 10%, urea 10%, chlorophyll water sol 0.5% Oint tube 1 oz, 4 oz

Use: Treatment of skin lesions

PANALBA (Upjohn) Tetracycline phosphate complex eq to HCl form 250 mg & Novobio-cin 125 mg/Cop Bot 16s 100s

Use: (1 or 2 q.i.d.) Mixed infections

PANALBA HALF STRENGTH. (Upjohn) Tetracycline HCl 125 mg novobio-cin sod 62.5 mg/Cop Bot 16s 100s

Use: Tetracycline novobio-cin therapy, also resistant prone staphylococci

PANALBA K M GRANULES (Drops) (Upjohn) Panmycin 125 mg Albamycin 62.5 mg and Potassium metaphosphate 100 mg/5 cc Bot 30 cc

Use: Antibiotic therapy

- PANALGESIC.** (Paythress) Methyl salicylate & aspirin 58%, menthol & camphor 4%, in alcohol & vegetable oil. Bot. 2 oz., ½ gal.
Use: Analgesic & counterirritant.
- PANALGYN.** (Merck Sharp & Dohme) Strontium salicylate 1.2 Gm., strontium iodide 0.52 Gm., calcichine 1.2 mg., pilocarpine HCl 8 mg., gelsemium 60 mg./30 cc. Bot. 1 pt., 1 gal.
Use: Analgesic & antirheumatic.
- PANALINS.** (Mead Johnson) Vit. B-1 2 mg., B-2 2 mg., niacinamide 20 mg., Vit. C 50 mg., Cal. pantothenate 5 mg., Vit. B-6 0.5 mg., folic acid 0.25 mg., Vit. B-12 2 mcg., A 5000 u., D 400 u./Cap. Bot. 100s, 500s
Use: (1-2 daily) Multivitamin therapy. Maintenance formula.
- PANALINS-T.** (Mead Johnson) Vit. B-1 10 mg., B-2 10 mg., niacinamide 100 mg., Cal. pantothenate 20 mg., Vit. B-6 2 mg., folic acid 1.5 mg., Vit. C 300 mg., B-12 4 mcg./Cap. Bot. 30s, 100s.
Use: (1-2 daily) Multivitamin therapy. Therapeutic formula.
- PANALIX.** (Rhinoap) N-acetyl-p-aminophenol 60 mg./O 6 cc.
Use: Antipyretic & analgesic.
- PANAMINE.** (Metropolitan Lab.) Phenylpropanolamine HCl 25 mg./Tab. (yellow) Bot. 1000s
Use: Nasal decongestant and bronchodilator
- PANAOR.** (Dewey) di-Calcium pantothenate 10 mg./Tab. Bot. 100s
- PANBIOTIC.** (Bristol) Pot. penicillin G 300,000 u., procaine penicillin G 300,000 u., dibenzylethenediamine dipenicillin G 600,000 u./Dose. Vial 1 dose, 5 dose.
Use: I.M., penicillin therapy.
- PANBLEX-C.** (Brewer) Vit. B-1 5 mg., B-2 3 mg., B-6 0.25 mg., cal. pantothenate 3 mg., niacinamide 10 mg., vit. C 50 mg., liver ext. 30 mg., Brewer's yeast 400 mg./cap. Bot. 100s, 500s. The tablets contain the same ingredients as the capsules except that it has 220 mg. of Brewer's yeast. Also available with folic acid 0.7 mg./cap. or tab.
Use: Dietary supplement.
- PANCEBRIN.** (Lilly) Thiamine chloride 10 mg., riboflavin 2 mg., pyridoxine HCl 3 mg., pantothenic acid 3 mg., niacinamide 20 mg., ascorbic acid 60 mg., alpha tocopherol 2 mg., Vit. A 10,000 u., D 1000 u./2 cc. Vial 10 cc. Box 1s, 25s.
Use: Multiple vitamin therapy.
- PANCHOLIN.** (Strassenburgh) Dehydrocholic acid 2 gr., bile ox ext 1½ gr., pancreatic sub. 3 gr./Tab. E.C. Bot. 100s
Use: Hydrocholeretic and choleretic agent
- PANCIDIN.** (Pan American) Chlorpheniramine maleate 1 mg., salicylamide 227 mg., phenacetin 100 mg., caffeine 10 mg., Vit. C 20 mg., phenylephrine HCl 5 mg./Tab. Bot. 100s.
Use: Sinus, hay fever, colds.
- PANCIOLIN FOSTE.** (Pan American) Chlorophenpyridamine maleate 4 mg., salicylamide 0.19 Gm., phenacetin 0.13 Gm., caffeine 0.03 Gm., Vit. C 0.05 Gm., methamphetamine HCl 1.25 mg./Panseal 100s.
Use: Cold.
- PAN-CONCEMIN.** (Merrell) Vit. A 5000 u., D 800 u., thiamine HCl 3 mg., riboflavin 2 mg., ascorbic acid 50 mg., niacinamide 20 mg., Cal. d-pantothenate 2 mg., pyridoxine HCl 1 mg./Tab. Bot. 100s, 1000s
Use: Multivitamin therapy.
- PANCGRAPHEN.** (Smith-Dorsey) Phenobarbital ¼ gr., Cal. carb. 3 gr., Sod. bicarb. 3 gr., pepsin 1 gr., pancreatin 1 gr./Tab. Bot. 100s, 1000s.
Use: (t.i.d.) Sedative & gastric antacid.
- PANCREATIC DORNASE, N.N.D. 1962.** See: Dornasec for Inhalation, Vial (Merck Sharp & Dohme)
- PANCREATIC EXTRACT DEPROTEINATED.** See: Depropanex, Vial. (Merck Sharp & Dohme)
Fadulin. (Winthrop Labs.)
- PANCREATIC SUBSTANCE.** Substance from fresh pancreas of hog or ox & contains enzymes amylapsin, trypsin & lipapsin. See: Stomyl. (Winthrop Labs.)
W/Bile salts & dehydrocholic acid. See: Depancol, Tab. (Warner-Chilcott)
W/Cholic acid and ketocholonic acid. See: Chobite, Tab. (Irwin-Nelsler)
W/Lipase. See: Colozym, Cap. (Organon)
W/Ox bile. See: Pancrabillin, Liq. (Reed & Carnick)
W/Ox bile ext. & dehydrocholic acid. See: Pancroid, Tab. (Strassenburgh).
W/Vit. B-1, B-2, C, niacinamide, diastase, ox bile ext., glutamic acid. See: Be-zymin, Tab. (Croskell)
- PANCREATIN, N.F.** Pancreatic enzymes obtained from hog or cattle pancreatic tissue.
Use: Digestant.
See: Panapsin, Tab. (Harrower)
Stomyl. (Winthrop Labs.)
W/Betaine HCl, ginger oleoresin, desoxycholic acid & chlorophyllin. See: Convertin, Tab. (Ascher)
W/Cellulose, pepsin, glutamic acid HCl, ox bile extract. See: Kanulase, Tab. (Smith-Dorsey)

* Generic name recognized by N.F., U.S.P. or A.M.A.

W/Ketocholonic acids, bile salts, betaine HCl See: Pradose, Tab (Miller Labs)
 W/Myczyme diastase, pepsin, hog bile desiccated, ext nux vomica See Enzobile Tab (Mallard)
 W/Ox bile ext, duodenal subst & papain See Miller Zyme, Tab (Miller)
 W/Ox bile extract, papain, duodenal sub stances defatted & desic See: Korazyme, Tab (Karmel Co)
 W/Pepsin, bile ext and papain See Pluri zyme, Tab (Harrower)
 W/Pepsin and bile salts See: Entozyme, Tab (Robins)
 W/Pepsin, dehydracholic acid, ox bile ext See: Neabile, Tab (Haug Drug Co)
 W/Pepsin, ox bile, dehydracholic acid See: Ra Bile, Tab (Rowell)
 W/Pepsin, taurocholic acid See Enzymel, Tabs (Lloyd, Dobney & Westerfield)
 W/Phenobarbital, Cal carb, Sod bicorb, papain & benzocaine See: Pancrophen Tab (Smith Dorsey)
 W/Piperidolate HCl, cellulolytic enzyme amylolytic enzyme proteolytic enzyme taurocholic acid See Doctilase Tabs (Lake side)
 W/Rhubarb See: Altkarhein, Liq (Merrell)
 W/Sod, glycocholate, Sod taurocholate Sod salicylate & Sod. bicorb See: Toblogestin, Tab (Strong)
 W/Terolase, dehydrochloric acid, polysor bote 80 See: Digolase, Cap (Boyle)
PANCREOLIN, (Reed & Carnick) Liq Fresh pancreas ext, 2.59 Gm, ox bile ext 130 mg, alcohol 16%/fl oz Bot 1 pl, 1 gal Pill Ox bile ext 907 mg, pancreas desic 2 mg/Pill Bot 100s, 500s, 1000s Use: Hepatic & pancreatic hypofunction
PANCROID, (Strosenburgh) (New name is Pancholn) Dehydracholic acid 2 gr, ox bile ext 1 1/2 gr, pancreatic substance 3 gr / Tab Bot 30s, 100s, 500s, 1000s Use: Choleraic
PANHEPRIN (Abbott) 40,000 u/ml Vial 2 ml 20,000 u/ml 1 ml disp syringe, 5 ml Vial 1 ml ampoule 10,000 u/ml 4 and 5 ml vials, 1 ml ampoule 5000 u/ml 10 ml vial 1000 u/ml 10 ml vial
PANHEPRIN IN SALINE, (Abbott) Heparin sodium 2000 u, 1800 u, 1500 u /30 ml sterile isotonic saline Non waittable con tainers 500 ml. Use: Anticoagulant in extracorporeal circula tion of blood
PANMYCIN AQUADROPS, (Upjohn) Tetra cycline 100 mg/cc Bot 10 cc Use Antibiotic therapy
PANMYCIN AQUEOUS INJECTION, (Upjohn)

Tetracycline HCl 100 mg, Mag chloride 100 mg procaine HCl 2%/2 cc Vial 1s Use: IM Antibiotic therapy
PANMYCIN HYDROCHLORIDE, NND 1961 (Upjohn) Tetracycline HCl Cap (50 & 100 mg) Bot 25s, 100s (250 mg) Bot 16s, 100s **OROPS** 100 mg/cc Bot 10 cc w/calibrated dropper **Intravenous**, Vials 100 mg, 250 mg, 500 mg Use: Adults 250-500 mg q 6 h & children 50 mg per lb body wt q 6 h; against wide range of gram positive & gram nego live bacteria
PANMYCIN W/Albamylin See Panalba Cap (Upjohn)
PANMYCIN KM SYRUP, (Upjohn) Tetracycline equivalent to 125 mg tetracycline HCl, pol melophosphate 100 mg/5 cc Bot 2 oz, pl Use: Antibiotic therapy
PANMYCIN PHOSPHATE, NND 1961 (Upjohn) Tetracycline phosphate 250 mg/Cap Bot 16s, 100s Also available in half strength Bot 25s, 100s Use: Mixed infections
PANMYCIN REAOMIX, (Upjohn) Susp tetracycline HCl 125 mg or 250 mg/5 cc; fruit flavor Bot 1 oz Bot 2 oz (125 mg/5 cc) Use: 1-2 tsp q i d; tetracycline therapy
PANMYCIN SYRUP, (Upjohn) Tetracycline HCl 125 mg/5 cc Bot 2 oz, 1 pl Use Orally antibiotic
PAN-NUTRON (Nison) Vit A ocelate 10,000 u, O 1000 u, B 1 5 mg B 2 5 mg, B 6 1 mg, B 12 2 mcg, C 100 mg, E 1 IU, niacinamide 50 mg, Col pantothenate 5 mg, Iolic acid 0.5 mg, choline dihydrogen citrate 25 mg/Tab Bot 90s, 500s Use: (1 or more daily) Vitamin deficiencies
PANOGEN, (Smith Miller & Patch) Methyl testosterone 2.5 mg, estradiol 0.05 mg, pentylene tetrazol 50 mg, ferrous sulf 50 mg, Iolic acid 0.25 mg, B 12 1 mcg choline bitartrate 105 mg, Inositol 50 mg, dl methionine 50 mg, hesperidin 50 mg, B 1 1.5 mg, B 2 2 mg, B 6 0.5 mg, C 50 mg, niacinamide 15 mg/Cap or Tab Bot 30s, 100s Use: (1 daily) Geripentics
PANOGEN LIQUID, (Smith Miller & Patch) Methyl testosterone 2.5 mg, estradiol 0.05 mg, pentylene tetrazol 50 mg, ferrous sulf 75 mg, Vit B 12 1 mcg, choline chloride 57.6 mg, Inositol 50 mg, betaine monohydrate 50 mg, hesperidin methyl chalcone 50 mg, Vit B-1 1.5 mg, B 2 20 mg niacinamide 15 mg, Vit B-6 0.5 mg, C 50 mg/1.5 cc Bot 16 oz Use: Oletary supplement

PANOVIMS. (Plessner) Vit. A 5000 u, D 1000 u., thiamine HCl 5 mg., riboflavin 3 mg., niacinamide 25 mg., pyridoxine HCl 0.5 mg., cal. pantothenate 1 mg., ascorbic acid 75 mg., vit. E 2 mg., B-12 2.5 mcg
Tab. Bot. 100s, 250s, 1000s

Use: Dietary supplement.

PANPARNIT HYDROCHLORIDE, NND
1962. (Geigy) (Caromiphon HCl). Tab. 12.5 mg., Bot. 100s, 1000s; 50 mg., Bot. 50s, 250s, 1000s.

Use: Parkinson's syndrome.

PAN-PEPTIC. (Merck Sharp & Dohme) Pepsin, lactic acid, caffeine A celery seed. Elixir, Bot. 1 pt., 1 gal.

Use: (1 dr. t.i.d.) Dietary disturbances.

PANRAY. (Panroy Corp.) Isoniazid. Bot. 100s, 1000s.

Use: Tuberculosis.

PANREXIN TP. (Pan American) Dextro amphetamine sulfate 15 mg., amobarbital 100 mg./Panseal 100s.

Use: Obesity.

PANRONE. (Panroy Corp) p-Acetylamino-benzaldehyde thiosemicarbazone (Contaban, Amthloxane.) Tab. 12.5 mg., 25 mg., Bot. 100s, 500s, 1000s.

Use: Tuberculosis.

PAN-S. See: Pansulfa, Prep. (Merrell).

PANSCOL. (Baker Labs) Salicylic acid, cholesterol, oil olein & phenol. Oint. Jars 4 oz.

Use: Treatment and management of dermatitis.

PANSTERIOD. (Kenwood) Hydrocortisone acetate 10 mg., prednisone 1.5 mg., prednisolone 1.5 mg., neomycin sulfate 5 mg./Gm. Oint. Tube. 5 Gm., 15 Gm.

Use: Dermatitis.

PANSULFA. (Merrell) Susp., sulfamerazine 1 Gm., sulfadiazine 1 Gm., sulfacetamide 1 Gm./30 cc. Bot. 12 oz., 1 gal. Tab., sulfamerazine 0.167 Gm., sulfadiazine 0.167 Gm., sulfacetamide 0.167 Gm./Tab. Bot. 100s, 1000s.

W/Penicillin. (Merrell) Cryst. Pot. penicillin G 100,000 u., sulfamerazine 0.167 Gm., sulfadiazine 0.167 Gm., sulfacetamide 0.167 Gm./Tab. Bot. 24s, 250s

PANTAREE. (Haskell)
Syr. Thiamine HCl 12 mg., riboflavin 12 mg., pyridoxine HCl 1 mg., pantothenic acid 12 mg., choline chloride 30 mg., liver conc. equiv. to 1 oz. fresh liver/l.oz. Bot. 6 fl. oz., 1 pt., 1 gal.

W/Iron, Cap. & Tab. (Haskell) Ferrous sulfate exscl. 3 gr., in addition to vitamin content of Pantabee Cap. Bot. 50s, 500s, 1000s. Tab., Bot. 100s, 500s, 1000s.

W/Iron Syr. (Haskell) Vit. B-1 6 mg., R-2

* Generic name recognized by N.F., U.S.P. or A.M.A.

6 mg., B-12 15 mg., R-6 0.6 mg., niacin 60 mg., pantothenic acid 6 mg., ferrous sulfate 9 gr., liver fraction No. 1 625 mg., tricholine xitrate 50 mg./20 cc. Syr. Bot 1 pt., 1 gal.

Use: Hematinic.

PANTABEERIOD. (Haskell) Thyroid gland dried 1/2 gr., thiamine HCl 1 mg., riboflavin 2 mg., elacinaomide 10 mg., pyridoxine HCl 0.1 mg., Cal. pantothenate 2 mg./Tab. Bot 50s, 250s, 1000s.

Use: Thyroid w/vitamins.

PANTAREIROID No. 2. (Haskell) Same vitamin content but with 1 gr. of thyroid gland dried/Tab. Bot. 50s, 250s, 500s, 1000s.

Use: Thyroid w/vitamins.

PANTASE. (Sulliff & Cass) Pepsin saccharated 260 mg., pancreatin 65 mg., aromatic powder 16 mg., dicitrate 16 mg., extracts of whole bile 11 mg./Tab Bot. 1000s

Use: Digestant and acidulant.

PANTERIC. (Parke, Davis) Pancreatin, Caps 5 gr. Bot. 100s, 1000s.
Granules Bot. 1 & 4 oz.
Tab. 5 gr. Bot. 100s, 500s, 1000s.

Use: To aid digestion.

PANTERIC COMPOUND TABLETS. (Parke, Davis) Pancreatin 10 gr., oxgall ext. 1 1/2 gr./tab Bot 100s, 500s, 1000s.

Use: Aid digestion.

* **PANTHENOL.** Alcohol corresponding to pantothenic acid Panthanol. Panthényl. Sol. Cosyme, Viol (Travenol)

D-P-A Injection (Hiss)

Ilapon, Amp. (Worren-Teed)

Motilyn, Ivy (Abb.)

Panodon, Cream (Gordon Labs.)

Panthoderm Cream (U. S. Vitamin)

W/Cholins. See: Ilapon-Choline (Worren-Teed)

W/Hydrocortisone. See: Patha-F, Cream or Pantha-Foam (U. S. Vitamin)

W/Neomycin Sulf. See: Neopan, Cream (U. S. Vitamin)

W/Pyridoxine. See: Pylipan, Oint. (Danley-Evans)

W/Pyridoxine HCl, hexachlorophene, menthol. See: Vi-Culo, Skin Cream. (Columbus)

W/Resorcinol monoacetate A sulfur. See: R.S. Panthoderm Cream (U. S. Vitamin).

W/Tyrosinase. See: Tyrolaris, Sol. (Merck Sharp & Dohme)

W/Vitamin A. See: Aquasol A W/Panthoderm, Oint. (U. S. Vitamin)

PANTHODERM CREAM. (U. S. Vitamin)

Panthanol 2% In water-miscible cream.

Jar 1 oz., 2 oz.

Use (Once or twice daily) Emollient & healing aid

PANTHO-F CREAM. (U S Vitamin) Hydrocortisone 0.2% or 1%, pantothenyl 2%
Tube 5 Gm, 20 Gm

Use Topically, dermatitis

PANTHO-FOAM. (U S Vitamin) Hydrocortisone 0.2%, Pantothenyl 2% Aerosol
Bot 1 oz, 2 oz

Use Dermatologic

PANTHOJECT (U S Vitamin) d Calcium pantothenate 250 mg /cc Vial 10 cc, 6s
Use IM Restoration of normal intestinal motility and function

PANTHOLIN. (Lilly) dl Pantothenic acid as Sod or Cal salt Tab 10 mg, Bot 100s, 500s

Use: Orally; dietary supplement

PANTHOSAN. (Elder) Vit B-1 3 mg, B 2 3 mg, B 6 0.25 mg, cal pantothenate 5 mg, nicotinamide 20 mg, liver concentrate 50 mg /Cop Bot 100s 500s, 1000s

Use Vitamin B complex

PANTOCAINE See Tetrocaine Hydrochloride (Various Mfr)

PANTOCIN. (Liberty) dl Calcium pantothenate 10 mg /Tab Bot 100s

PANTODREN-100 (Kremers Urban) Beef peptone 100 mg, methionine 100 mg, thiamine HCl 100 mg, ascorbic acid 100 mg /2 cc Amp 2 cc, Box 25s Vial 30 cc

PANTOFENICOL PLURIOQUIMICA, 10A (Paritugol) Chloramphenicol Pantothenate Complex

Use Antibiotic

PANTOPAUQUE, NND 1961 (Lalayette) lophendylate USP, ethyl iodophenylundecylate Amp 3 cc

Use: Intraoperative, Intrahepatically or orally; myelography

PANTOPAUQUE EMULSION, NND 1961 (Lalayette) 50% v/v aq emulsion of ethyl iodophenylundecylate Amp 10 cc

Use Roentgenographic visualization of certain body cavities such as sinuses, fistulous tracts and ducts and biliary tree

PANTOPON, NNR 1930 (Roche) Total alkaloids of opium as the HCl salt

Amp r (1/2 gr) 1 cc, Box 12s, 100s

Power Vial (1/2, 1 oz)

Tab Oral; (1/6 gr) Vial 20s

Tab Hypodermic (1/2 gr) Pkg 20s, 100s

Use S.C. & Orally; Analgesic

W/Papaverine HCl & atropine (atropine sulfate) See: Spasmoalgin, Tab & Amp (Roche)

PANTOTHAXIN. (Wmthrop Labs) Pantothenic acid as cal salt Tab 10 mg Bot 25s
Use: Dietary supplement

PANTOTHENIC ACID, NNR 1950, ADR
As Cal or Sod salt

See: Vitamin, preparations

Use Nonspecific protein therapy

PANTOTHENIC ACID SALTS

See Calcium Pantothenate

Sodium Pantothenate

PANTOTHENOL. See Panthenol, Prep (Various Mfr)

PANTOTHENYL ALCONOL. See Panthenol Prep (Varroas Mfr)

PANTOTHENYLOL. See Panthenol, Prep (Various Mfr)

PAN-TRANQUIL (Nysco) Mephenesin 400 mg mephobarbital 10 mg hyoscine HBr /Tab Bot 1000s

Use (11d) Tranquilizing agent

PANTRINSIC (Brown Pharm) Ferrous fumarate 200 mg /Tab Bot 100s

Use Anemia

PANTRINSIC-C (Brown Pharm) Ferrous fumarate 200 mg hesperidin 250 mg Vit B 1 5 mg B 12 2 mcg stomach with in

Intrinsic factor 100 mg C 250 mg, liver 200 mg cobalt chloride 15 mg folic acid 0.1 mg /Tab Bot 100s

Use Anemias

PANTRINSIC PN (Brown Pharm) Ferrous fumarate 100 mg, calcium lactate 250 mg, Vit A 2500 u B 1 3 mg B 2 1.5 mg B 6

0.1 mg B 12 5 mcg stomach with intrinsic factor 50 mg C 100 mg aluminum hydroxide 50 mg /Cop Bot 100s 250s

Use Prenatal therapy

PAN-VATINE (Smith Dorsey) Vit A 5000 u, D 800 u, ascorbic acid 375 mg, thiamine HCl 15 mg nicotinamide 20 mg

riboflavin 2 mg pyridoxine 0.1 mg, Cal pantothenate 1 mg, Vit E 2 mg /Cop Bot 100s, 1000s

Use Vitamin supplement

PANVITEX. (Testagel) Ferrous fumarate 150 mg, sodium molybdate 3 mg rutin 10 mg

folic acid 0.25 mg B 12 2 mcg, A 4000 USP u, D 400 USP u, B 1 2 mg B 2 2 mg, niacinamide 10 mg B-6 0.8

mg, C 50 mg Quinine 0.25 mg, I 0.15 mg, Ca 600 mg bioflavonoids 25 mg, Vit K 0.25 mg /Cop Bot 100s 1000

Use: Prenatal vitamin mineral combination

PANWARFIN, NND 1962 (Abbott) Warfarin sodium 5 mg /10 mg or 25 mg Bot 5 mg /100s 10 mg /100s 25 mg /25s

Use Intravascular thrombosis and embolism

PANWARFIN INJECTION (Abbott) Warfarin sod 75 mg w/ sod chloride 15 mg /vial Vials w/3 ml sterile water

Use Anticoagulant

PANZALONE. (Doak) delta 5 hemisuccinoyl-pregnenolone 2%, water washable cream

- base containing Bura-Sol Dook (equiv. to 3.38 mg. aluminum acetate) pH 5.5, Tubes $\frac{1}{2}$ oz., Jar 1 lb.
 Use: Dermatoses.
- PAOCIN**, (Mossengill) Colloidal kaolin 90 gr., pectin 2 gr., aromatics q.s./15 cc. Bot. 1 pt., 1 gal. Tab., Bot. 500s, 1000s.
 Use: Diarrhea, colitis & intestinal toxemias.
- PAOQUAN**, (Massengill) Sulfaguanidine $7\frac{1}{2}$ gr., colloidal kaolin 30 gr., pectin $\frac{3}{8}$ gr./5 cc. Bot. 1 pt., 1 gal.
 Use: Diarrhea & bacillary dysentery.
- W/Duodenal subst., pancreatin & ox bile ext.** See: Miller-Zyme, Tab. (Miller)
- PAPAIN**, Vegetable pepsin,
 See: Papose (Warner-Chilcott)
- W/Diastase**
 See: Caripeptic, Liq. (Upjohn)
 Coroid W/Diastase, Tab. (Amer. Ferment)
- W/Pancreatin, ginger, rhubarb.** See: Alkagests, Tab. (Blue Line)
- W/Pancreatin, ox bile ext. & duodenal substances defatted & desic.** See: Korasyme, Tab. (Kormel Co.)
- W/Urea, chlorophyll.** See: Panoff, Oint. (Ryston)
- PAPANESIN**, (Superior Pharm) Acetyl-para-aminophenol 250 mg., guayanesin 50 mg., phenylephrine HCl 2.5 mg./Tab. Bot. 100s, 500s.
 Use: Relief of the common cold, nasal congestion.
- PAPASE**, (Warner-Chilcott) Proteolytic enzyme from carica papaya 10,000 u./Tab. spearmint-flavored. Bot. 100s.
 Use: Orally or buccally, Edema and inflammation conditions.
- PAPAVATRAL**, (Kenwood Labs.) Ethaverine HCl 30 mg., pentaerythritol tetranitrate 10 mg./tab. Bot. 100s, 1000s.
 Use: Angina pectoris.
- PAPAVATRAL L.A.** (Kenwood) Ethylpapaverine HCl 30 mg., pentaerythritol tetranitrate 50 mg./Cap. Bot. 30s, 250s. Available w/Phenobarbital 45 mg./Cap
 Use: Angina pectoris.
- *PAPAVERINE HYDROCHLORIDE, U.S.P., N.N.R. 1947;** Inj., U.S.P. Tab. U.S.P.
W/Cocaine sulfate. See: Copavin, Cap. (Lilly).
W/Cocaine phosphate, aluminum aspirin, phenylephrine HCl, chlorpheniramine maleate, Vit. C. See Calrex Compound, Caps. (Rowell)
W/Cocaine sulfate, aloin & Sod. salicylate. See: Copavin Compound, Elixir (Lilly).
W/Morphine sulfate. See: Asymarine Injection, Amp. (Brean).
- * Generic name recognized by N.F., U.S.P. or A.M.A.**
- W/Pantopon & atrolin.** See: Spasmolgin, Tab. & Amp. (Hoffmann-La Roche)
- PAPINE**, (Battie) Morphine HCl 1 gr., and chloral hydrate $3\frac{1}{2}$ gr./oz. Bot. 12 oz.
 Use: Narcotic analgesic
- *PARA-AMINOBENZOIC ACID, N.F., N.N.R. 1957;** Tabs, N.F. See: p-Aminobenzoic Acid (Various Mfr.)
- *PARA-AMINONIPPURIC ACID, N.F., N.N.E. 1950.** See: p-Aminohippuric Acid
- PARA-AMINOSALICYLIC ACID, U.S.P., N.N.D. 1962.** See: p-Aminosalicylic Acid, U.S.P.; p-Aminosalicylate, Calcium, U.S.P.; p-Aminosalicylate, Sodium, U.S.P.
- PARABROM.** See: Pyrabrom.
- PARAROMDYLAMINE.** * Brompheniramine 1-(p-bromophenyl)-1-(2-pyridyl)-3-dimethylamino propane maleate. See: Dimetane (Robins)
- PARACAIN, A.D.E.** See: Procaine Hydrochloride (Various Mfr.)
- PARACARINOXAMINE MALEATE, Carbinoxamine.** See Cistin Maleate, Elixir & Tab (McNeil).
- PARACETALDEHYDE.** See: Paraldehyde, U.S.P.
- PARACETAMOL.** See: N-Acetyl-p-Aminophenol
- PARACHLORAMINE HYDROCHLORIDE.** See: Sonamine, Tab. (Pfizer).
- PARACHLOROMETA XYLENOL** (Ottawa Chem. Co.)
 See: Dillof (R. T. French)
 Linal (Lincoln Re)
 Ottosept (Ottawa Chem.)
 Use: Phenolic antiseptic.
- *PARACHLOROPHENOL, N.F., A.D.R. p-Chlorophenol**
W/Comphor. See: Comphorated Parachlorophenol, N.F.
- PARACODIN.** See: Dihydrocodeine bitartrate. (Knoll)
- PARA CONESTRO.** (Para) Conjugated estrogens 0.625 mg. or 1.25 mg./Tab. Bot. 100s.
 Use: Menopausal syndrome
- PARACORT, N.N.D. 1962.** (Parke, Davis) Prednisone 2.5 and 5 mg./Tab. Bot. 30s, 100s.
 Use: Numerous conditions of inflammation, allergic states and collagen diseases
- PARACORTOL, N.N.D. 1961.** (Parke, Davis) Prednisolone 2.5 mg., or 5 mg./Tab. Bot. 30s, 100s.
 Use: Treatment of rheumatoid arthritis and related diseases
- PARADIONE, N.N.D. 1961.** (Abbott) Paramethadione 3.5 Dimethyl-5-ethylhexozolidine-2,4-dione Cap. D 15 Gm., O 3 Gm., Bot. 100s, 1000s. Sol. 0.3 Gm./cc. Bot. 50

- Use Petit mal, myoclonic & akinetic epilepsy
- PARAFLEX**, NND 1962 (McNeil) Chlorzoxazone [5 chlorobenxoxazolinone] Tab 250 mg Bot 50s 100s
Use Skeletal muscle relaxant
- PARAFOH**, (McNeil) Paraflex 125 mg plus Tylenol 300 mg Tabs Bot 24s 50s
Use Skeletal muscle relaxant
- PARAFOH W/CODEINE** (McNeil) Paraflex 125 mg, Tylenol 300 mg codeine phosphate 15 mg / Tab Bot 24s
Use Relief of severely painful muscular skeletal disorders
- PARAFON FORTE** (McNeil) Chlorzoxazone 250 mg, tylenol (acetaminaphen) 300 mg / Tab Bot 50s
Use Skeletal muscle relaxant
- PARAFON W/PREDHISOLONE** (McNeil) Paraflex 125 mg, tylenol 300 mg prednisolone 10 mg / Tab Bot 36s
Use: Relief of pain stiffness and limitation of motion
- PARAFORM**, ADR See Paraformaldehyde (Merck Sharp & Dahme)
- PARAFORMALDEHYDE**, NNR 1939, ADR (Merck Sharp & Dahme) Paraform Bot ¼ lb
See Aldeform Disks (Paul J Mandaback)
Formaldehyde, Pastils (Various Mfr)
Marvason Cream (Toblox)
Trioxymethylene (an incorrect term for paraformaldehyde)
Use: Essentially the same as formaldehyde
W/ Stearic acid, Pol carb & Triethanolamine
See: Linton Cream (McIntosh Linton)
Formil Tablets (Fatch)
- PARAGLYCYLARSANILIC ACID**, N Carb amylmethyl p aminobenzenearsanic acid the free acid of trypanamide See: Risorphen, Amp (Endo)
- PARA-HIST**, (Brain Kay & McQuarrie) Propenpyndamine maleate 10 mg thenyl pyrimide HCl 10 mg, pyrimidine maleate 10 mg / Tab Bot 100s
Use: Allergic disorders
- PARA HYCODAN**, (Endo) Dihydrocodeinone bitartrate 4.5 mg and terephthalate D38 mg homatrapine terephthalate 0.38 mg pentylene tetrazol 50 mg / 5 cc or Tab Bot 100s & 1000s Syr 1 pt & 1 gal Tab 100s & 1000s
Use: Antitussive
- PARAL** (Fellows) Paraldehyde 1 gm / Cap Bot 60s
Use: CNS depressant
- PARALDEHYDE**, USP Paracetaldehyde Bot ¼ lb, 1 lb — Amp See: Paraf, Cap (Fellows Testagar)
- Use IM, IV or Orally; Hypnotic & sedative
- PARALDEHYDE**, USP (Buffington) Amp 2 cc, Box 12s, 100s; 5 cc, Box 6s 25s, 100s; TD cc Box 6s, 25s, 100s
- PARALDEHYDE FELSULES** (Fellows) Paraldehyde 15 gr, hermetically sealed in soft gel Cap, Bot 60s
Use Sedative hypnotic
- PARAMAL**, (Columbus Pharm) Pyrimidine maleate 25 mg / tab Bot 100s
Use Antihistamine
- PARAMEPHRIN** See Epinephrine (Various Mfr)
- *PARAMETHADIONE**, HF NND 1962 (J 5 Dimethyl 5 ethyloxazoline 2,4 dione Caps & Soln NF See Paraldione Cap & Sol (Abbott)
- *PARAMETHASONE ACETATE** 16 α methyl 6 α fluoroprednisolone 21 acetate See Haldrone (Eli Lilly)
- PARAMINYL MALEATE**, NND 1962 (Buffington) Pyrimidine maleate 50 mg / tab Bot 100s 1000s
Use: Antihistamine
- *PARAMOMYCIN**
See Humatin (Parke Davis)
- *PARA-NITROSULFATHIAZOLE**, NF Susp NF NND 1962 See Hissulfazole NND 1958
- *PARAQUINYL HYDROCHLORIDE** (Merrell) α fluoren 9 ylidene p taluamidine hydrochloride
Use antiinflammatory agent
- PARA-PAS**, NND 1962 (Gold Leaf; Panray) p Aminosalicylic acid Tab D5 Gm Bot 100s, 500s, 1000s
Use Tuberculosis
- PARA-PAS SODIUM**, NND 1962 (Gold Leaf) Sod p aminosalicylate Tab D5 Gm, Bot 100s 500s 1000s; Effer, 2 Gm, Bot 100s Syr 4.5 Gm / fl oz Bot 16 oz
Use: Tuberculosis
- PARASAL**, NND 1962 (Panray) p Aminosalicylic acid
Pow: ¼, ½, 1 lb
Tab: Plain or E.C. (0.5 Gm) Bot 100s, 250s 500s; Tab SA (1 Gm) Bot 100s 250s, 500s; Tabs Buffered (0.5 Gm) Bot 100s 250s, 500s; Effer. Tabs. (2 Gm) Bot 20s 100s
Granules E.C., Bot
Use: Tuberculosis
- PARASAL CALCIUM**, (Panray) p Aminosalicylate Cal Tab D.5 Gm, Bot 100s, 250s 500s, 1000s
Use: Tuberculosis
- PARASAL-INH**, (Panray) p Aminosalicylic

- acid 0.5 Gm., Isoniazid 12.5 mg./Tab. buffered. Bot. 100s, 250s, 500s, 1000s.
 Use: Tubercularis.
- PARASAL-INH-20, BUFFERED.** (Panray)
 Aminosalicilic acid buffered 500 mg.,
 Isoniazid 20 mg./tab. Bot. 100s, 250s, 500s,
 1000s.
 Use: Tubercularis.
- PARASAL POTASSIUM, N N D. 1961.** (Panray)
 Pow. Bot. $\frac{1}{4}$, $\frac{1}{2}$ & 1 lb.
 Tab. (0.5 Gm.) Bot. 100s, 250s, 500s, 1000s
 Use: Tubercularis
- PARASAL SODIUM, N N D. 1962.** (Panray)
 p-Aminosalicylate Sod.
 Cap. (0.5 Gm.) 100s, 250s, 500s
 Pow. Bot. $\frac{1}{4}$, $\frac{1}{2}$ & 1 lb.
 Syr. (0.5 Gm. PAS/5 cc.) Bot. 1 pt.
 Granules: Enteric coated. Bot. 1 lb.
 Tabs.: Sugar coated (0.5 Gm., 0.69 Gm. &
 1 Gm.) Bot. 100s, 250s, 500s
 Tabs.: Enteric coated (0.5 Gm., 0.69 Gm. &
 1 Gm.) Bot. 100s, 250s, 500s
 Packets: (4.18 Gm.) Box 100s
 Vial: (5 Gm) 50 cc. (15 Gm) 100 cc.
 Use: Tubercularis.
- PARASED.** (Central) Camphorated opium
 tincture 0.3 ml., opium U.S.P. 1.2 mg., ben-
 zoic acid 1.2 mg./Palatab. Bot. 100s, 1000s
 Use: Control of diarrhea
- PARASYLLIUM.** (Abbott) Mineral oil 80%,
 psyllium seed 20%. Bot. 7 pt. Also avail.
 w/phenolphthalein. Bot. 1 pt.
 Use: Laxative
- PARASYMPATHOLYTIC AGENTS, Cholinergic**
 Blocking agents.
 See: Anticholinergic Agents.
 Antispasmodics.
 Mydriatics.
 Parkinsonism.
- PARASYMPATHOMIMETIC AGENTS, N N D.**
 1959. See: Cholinergic Agents.
- PARA TESTROGEN.** (Para) Ethinyl estradiol
 0.2 mg., methyl testosterone 5 mg./Tab
 Bot. 100s.
 Use: Geriatric
- PARATHANE.** (Graff) Chlorpheniramine ma-
 leate 1.25 mg., phenylpropanolamine HCl
 10 mg., phenylephrine HCl 5 mg., glyceryl
 guaiacolate 25 mg./5 cc. Bot. 1 pt.
 Use: Upper respiratory decongestants
- PARA-THOR-MONE, N N R. 1933.** See:
 Parathyroid Injection. (Lilly)
- PARATHYROID A ANTIGEN.** (Lederle)
 Febrile antigen diagnostic. Vial 5 cc.
 Use: Diagnostic reagent for macroscopic
 rapid slide agglutination.
- PARATHYROID B ANTIGEN.** (Lederle)
 Febrile antigen diagnostic. Vial 5 cc.
 * Generic name recognized by N F, U S P. or A.M.A.
- Use: Diagnostic reagent for macroscopic
 rapid slide agglutination method.**
- * PARATHYROID INJECTION, U S P. N N R.**
 1950. (Lilly) Aq. sol. of parathyroid hor-
 mone, 100 U.S.P. u./cc. Sol. Amp. 1 cc.,
 Box 6r, 100s; Amp. 5 cc. See: Paradrin,
 Vial (Parke, Davis).
 Use S.C. & I.M., parathyroid therapy.
- PARATHYROID THERAPY.**
 See: Dihydrotachysterol, U.S.P. Tachysterol.
 (Various Mfr.)
 Hytakerol, Sol. (Winthrop Labs.)
 Parathyroid Injection. (Various Mfr.)
 Paradrin, Vial. (Parke, Davis)
- PARAZINE, N N D. 1962.** (Tulag) Piperazine
 citrate 500 mg./5 cc. Bot. 4 oz., 1 pt., 1
 gal.
 Use: Vermifugal agent.
- PARAZONE.** See: p-Acetylamino benzalde-
 hyde thiosemicarbazone Thibione (Panray).
- PARREXIN, (C. D. Smith) Thiamine HCl 100**
 mg., niacinamide 100 mg., riboflavin 1
 mg., pyridoxine HCl 1 mg., Cal. pantho-
 thenate 5 mg., chlorobutanol 0.5%/cc.
 Vial 15 cc.
 Use: Vit. B complex.
- PARBEXIN-S. (C. D. Smith) Thiamine HCl**
 100 mg., niacinamide 150 mg., riboflavin
 0.5 mg., pyridoxine HCl 50 mg., panthenol
 5 mg., benzyl alcohol 2%/cc. Amp. 1 cc.,
 5r, 25r, 100r. Vial 10 cc., 15 cc.
 Use: Vit. B complex
- PARBOCYL.** (Elder) p-Aminobenzole acid
 0.25 Gm., Sod. salicylate 0.25 mg., Sod.
 ascorbate 20 mg./E.C. Tab. Bot. 100s,
 1000s.
 Use: (1-2 t.i.d.) Analgesic for rheumatic
 fever.
- PARROM.** (Central) Pyrebrom 50 mg. Tab.
 Bot. 100s, 500s, 1000s.
 Use: Premenstrual symptoms.
- * PARMUTOXATE.** β -dimethylaminoethyl 3-
 amino-4-butoxybenzoate
- PARDESS.** (Merck Sharp & Dohme) Thiamine
 HCl 25 mg., riboflavin 10 mg., pyridoxine
 HCl 5 mg., cal. pantothenate 50 mg., ni-
 acinamide 250 mg./6 cc. with or without
 diluent. Ros 25r, 50s, 100r.
 Use: R complex therapy.
- PARIDRIE HYDROBROMIDE, N N D 1962.**
 (Smith, Kline & French) Hydroxyampheta-
 mine HBr. Ophth Sol. 1% w/boric acid.
 Bot. $\frac{1}{2}$ oz. Tab. 20 mg, Bot 25r
 Use: Postural hypotension, heart block &
 nasal decongestant.
- W/Boric acid.** (Smith, Kline & French)
 Paradrin HBr 1%, boric acid 2%, Sod.

ethyl mercuri thiosalicylate 1.50 000 Bot 1/2 oz
 Use Ophthalmic solution, vasoconstrictor
 W/Hydrocortisone alcohol, phenylephrine
 HCl & thimerosal See Vasocort, Sol &
 Spraypok (Smith, Kline & French)
 W/Hydrocortisone, gramicidin, neomycin sul-
 fate, polymyxin B sulfate & phenylephrine
 See: Trisocort, Spraypok (Smith, Kline &
 French)

PAREDRINE-SULFATHIAZOLE. (Smith, Kline
 & French) Sulfathiazole 5%, paredrine
 HBr 1% O hydroxyphenyl mercurochloride
 1:20,000 Susp, Bot 1 fl oz, 12 fl oz
 Use: Nasal & sinus infections

PAREDBINOL. See: p (2 Methylaminopropyl)
 phenol (No Mfs currently lists)

PAREGORIC, U S P Tabs (5 min) Bot 500s
 1000s (Pilmon Moore)
 W/Aluminum hydroxide gel & bismuth
 subgallate See: Alpegara, Susp (Smith
 Dorsey)

W/Hyoscyamus ext, Ipecac syr, licorice ext
 See Crupic, Tab (Jenkins Labs)

W/Milk of bismuth, kaolin & pectin See
 Mul Sed, Susp (Pilmon Moore)

W/Pectalin & kaolin See: Parepectalin,
 Susp (Rorer)

W/Pepsin, bismuth subsalicylate, zinc sulfo-
 carbolate, salol & methyl salicylate See:
 Parepeptol, liq (Chicago Pharm)

W/Tartar emetic, ethyl nitrite, etc See:
 Brown Mixture, liq (Merrell)

PAREMYCIN. (O F Harvey) Neomycin sul-
 fate 300 mg, apium lactose 0.2 mg /oz
 Bot 3, 6 oz
 Use: Diarrheas

PARENAMINE, NND 1962 (Winthrop
 Labs) Protein hydrolysate Sol 6%, flask
 1000 cc 15%, Vial 100 cc
 Use: IV, IS & SCR dietary supplement

PARENEX. (Smith, Miller & Patch) Thiamine
 HCl 10 mg, niacinamide 50 mg, riboflavin
 1 mg, pyridoxine HCl 1 mg, Cal panto-
 thenate 5 mg, chlorobutanol 0.5%/cc Vial
 15 cc
 Use: Vit B complex therapy

W/Liver (Smith, Miller & Patch) Thiamine
 HCl 10 mg, niacinamide 20 mg, riboflavin
 1 mg, pyridoxine HCl 1 mg, Cal panto-
 thenate 5 mg, liver intl mcg, B 12 5
 mcg/cc Vial 15 cc
 Use: Anemias

PARENEX-C. (Smith, Miller & Patch) Thio-
 mine HCl 50 mg, niacinamide 50 mg,
 ascorbic acid 50 mg, riboflavin 3 mg
 pyridoxine HCl 5 mg, panthenol 5 mg
 gentiic acid ethanalamide 2 1/2%/cc Vial
 10 cc
 Use: Vit B complex & Vit C therapy

PARENEX MODIFIED. (Smith, Miller &
 Patch) Vit B 1 5 mg, B 2 2 mg, B 6 3 mg,
 niacinamide 50 mg, Vit C 15 mg, pan-
 thenol 3 mg, gentiic acid ethanalamide
 2 1/2%/cc Vial 10 cc
 Use: Vit B complex

PARENICILLIN, NND 1962 (Merrell)
 Aq Susp: Procaine penicillin G 300 000
 u/dose, Vial 10 cc

PARENOGEN, NND 1962 (Cutter) Fibrin
 ogen, human far IV inj Vial 1 Gm pow
 Use (2.6 Gm) Hypofibrinogenemia often
 associated w/premature separation of pla-
 centa

PARENS-400 (Hiss) Penicillin 400 000 u/
 Cap Bot 100s 500s
 Use: Infections due to penicillin sensitive
 organisms

PARENOSOL-B (Squibb) Vit B 1 10 mg,
 B 2 4 mg, niacinamide 200 mg/cc Amp
 1 cc, Box 5s Vial 10 cc
 Use: Pellagra

PARENTRACIN (Reed & Carnrick) Not listed
 Bacitracin 60 000 u Vial 1 dose
 Use: For hospital use only

PARENZYME, NND 1962 (National Drug)
 Trypsin Vial 5 mg/cc Cryst trypsin in
 sesame oil Vial 5 ml Pkg 1s Aq 25 mg
 w/5 ml diluent Pkg 1s Single dose pkg
 1 vial 12,500 NF u (5 mg) lyophilized
 trypsin
 Use: IM; reduces acute local inflammation
 (phlebitis, thrombophlebitis, traumatic
 wounds etc)

PARENZYME B (National Drug) Trypsin 5
 mg in polyethylene glycol 4000 base/
 Buccol Tab Vial 24s
 Use: Inflammation, sprains and edema

PARENZYME OINTMENT. (National Drug)
 Crystalline trypsin 2 mg, crystalline chymo-
 trypsin 6 mg, 9 aminocridine HCl 2 mg
 in water soluble base/Gm Tube 1 oz
 Use: Debridement and healing of external
 lesions

PAREPECTOLIN. (Rorer) Paregoric 37 cc
 pectin 2.5 gr, kaolin 85 gr/fl oz Susp
 Bot 4 & 8 fl oz
 Use: (1 2 tsp, tid) Nonspecific diarrheal

PAREPEPTOL. (Chicago Pharm) Paregoric
 60 min, pepsin 4 gr, bismuth subsalicylate
 4 gr, zinc sulfocarbolate 49 mg, solol 100
 mg, methyl salicylate 8 min/oz Bot 1 pl
 Use: Diarrhea

PARETHOXYCAINE HCl, NND 1959 See
 tetracaine HCl (Squibb)

PAR-FLUORO-VITE (Para Pharmacol) So-
 dium fluoride 221 mg, Vit A 2500 u,
 D 250 u, B 1 1 mg, ascorbic acid 30 mg,

- B-6 0.1 mg., B-12 3 mcg., and trace minerals/tab.
 Use: Control of dental caries.
- PARGAN CAPSULES.** (Squibb) Vit. A 5000 u., D 400 u., B-1 2 mg., B-2 2 mg., B-6 0.5 mg., C 50 mg., niacin 20 mg., pantothenic acid 5 mg., Vit. B-12 2 mg., folic acid 0.25 mg., menadione 2 mg./Cap. Bot. 30s, 100s, 1000s.
 Use: Multivitamin therapy.
- PARGAN-M.** (Squibb) Vit. A 5000 u., D 500 u., B-1 3 mg., B-2 3 mg., B-6 0.5 mg., C 50 mg., niacin 25 mg., pantothenic acid 5 mg., Vit. E 5 mg., B-12 2 mcg., folic acid 0.1 mg., menadione 2 mg. w/Col., Phos., Iodine, Iron, Pot., copper, Mang., Mag., cobalt, zinc & molybdenum/Cap. Bot. 100s, 250s.
 Use: Multivitamin therapy.
- * **PARGYLIN HYDROCHLORIDE.** (Abbott Lab) N-methyl-N-(2 propenyl)benzylamine hydrochloride See Etanyl (Abbott Labs)
- PARID.** (Burlington) Piperazine hexahydrate 500 mg./5 ml. Bot. 1 pt.
 Use: Treatment of pinworm and round-worm infections.
- PARISAMINE.** (Gold Leaf) Pyrilamine molecule 10 mg., aspirin 3½ gr., acetophenetidin 2½ gr., caffeine anhy. ¼ gr., p-aminobenzoic acid ½ gr./Tab. Bot. 100s, 1000s.
 Use: Head colds & hay fever.
- PARKINSONISM** (Parasympatholytic agents)
 See Atropa (Lederle).
 Carbamiphen HCl (Geigy).
 Cogentin, Tab. (Merck Sharp & Dohme)
 Diprival. (No Mfr. currently lists)
 Disipal (Riker)
 Kamadrin, Tab. (Burroughs Wellcome)
 Paglone HCl. (Lilly)
 Panparnit (Geigy).
 Parsidal (Warner-Chilcott).
 Phenothiazine HCl (Wyeth).
 Pipanol. (Winthrop Labs.)
 Prantal. (Schering)
 Trihexyphenidyl HCl (Various Mfr.)
- PARLITE.** (Lederle) Vitamins B-1 10 mg., B-2 10 mg., B-6 5 mg., B-12 25 mcg., C 500 mg., nicotinamide 150 mg., and sod pantothenate 10 mg./5 cc. Vial (dry) 5 cc., 25s.
 Use: IV, or I.M. Vitamin deficiencies
- PARLUGUAN.** (Panroy) Whole milk solids fortified w/dextrins & maltose 24 Gm., Mag. trisilicate 0.25 Gm., Mag. oxide 0.12 Gm., Cal. carb. 0.12 Gm., Mag. carb. D03 Gm./Trache. Bot. 25s.
 Use: 1-4 traches between meals; antacid in ulcer management.
- * Generic name recognized by N.F., U.S.P. or A.M.A.
- PARMAL.** (Central) Not listed Pyrilamine maleate Caps. Tamed disintegration (75 mg) Bot. 100s, 500s.
 Use: (1 cap. morning or evening) Anti-histaminic.
- PARNATE.** N.N.D. 1962. (Smith Kline & French) Tranylcypromine sulfate. 10 mg./Eob Bot. 50s, 500s.
 Use: Antidepressant; MAO inhibitor.
- PARODYNE.** See: Antipyrine (Various Mfr.)
- PARODIDIN.** N.N.R. 1950. (Parke, Davis) Parathyroid est., 100 U.S.P. u./cc. Vial 5 cc. Use: S.C. & I.M.; parathyroid therapy.
- PAROLEINE.** See: Petrolatum liquid (Various Mfr.)
- * **PAROMOMYCIN.** An antibiotic substance obtained from cultures of certain *Streptomyces* species, one of which is *Streptomyces elmoatus*.
 See: Humatin, Kapseal (Parke Davis) W/Chloramphenicol, hydrocortisone acetate, in a liquid petrolatum base. See: Humocort Ointment, (Parke Davis).
- PAROXYL.** See: Acetarsone, Tab. (Abbott).
- PARPANIT.** See: Carbamiphen Hydrochloride (Panparnit, Tab. Geigy)
- PARRISINE.** N.N.R. 1952. (Abbott) Paraffin paraffin. Parassed loce mesh surgical dressing, various sizes.
 Use: Protective surgical dressing
- PARSIDOL HCl.** N.N.D. 1962 (Warner-Chilcott) Ethopropazine, N.N.D. 1959 Propenamine HCl, lysivone, 10-(Diethylamino-1-propyl) Phenothiazine HCl. 10b (10 mg., 50 mg) Bot. 100s, 500s; (100 mg) Bot. 100s, 500s.
 Use: Parkinsonism.
- PARSLEY CONCENTRATE W/Garlic conc.** See: Allimin, Tab. (Von Poffen)
- PARZOCAL.** (Central) Zinc oxide 8%, colamine 8%, propolis 1%, benzyl alcohol 1%, pyrilamine maleate 1%, p-chloro-m-xylenol 0.8%, menthol 0.075%, camphor 0.05%. Bot 4 fl. oz., 1 pt., gal
 Use: Dermatitis.
- PARZONE.** (Mullinckrodt) Brand of dihydrocodeine bitartrate
- PAS.** N.N.D. 1962. (Lilly) p-Aminosalicylic acid, Tab. 7½ gr. Enseals. Bot. 500s, 1000s, 5000s
 Use: Tuberculosis.
- PASALATE.** N.N.D. 1962. (Dumas-Wilson) Sodium para-aminosalicylate Tab Bot. 100s, 1000s.
- PASANOL.** (Tilden) Digitalin 1/350 gr., straphanthin 1/25 gr., strychnine sulfate 1/25 gr., nitroglycerin 2/25 gr., cactus grandiflorus ½ gr./fl. oz. Bot. Pts Gals.
 Use: Cardiac and vasomotor stimulant

- ethyl mercuri thiosalicylate 1:50,000 Bot ½ oz
 Use: Ophthalmic solution; vasoconstrictor
 W/Hydrocortisone alcohol, phenylephrine HCl & thimerosal See: Vasocort, Sol & Spraypak (Smith, Kline & French)
 W/Hydrocortisone gramicidin, neomycin sulfate, polymyxin B sulfate & phenylephrine See: Trisocort, Spraypak (Smith, Kline & French)
- PAREDRINE-SULFATHIAZOLE.** (Smith, Kline & French) Sulfathiazole 5%, paredrine HBr 1% O hydroxyphenyl mercuric chloride 1:20,000, Susp, Bot 1 fl oz, 12 fl oz
 Use: Nasal & sinus infections
- PAREDROL** See: p (2 Methylaminopropyl) phenol (No Mfr currently lists)
- PAREGORIC, USP** Tabs (5 min) Bot 500s 1000s (Pitman Moore)
 W/Aluminum hydroxide gel & bismuth subgallate See: Alpegara, Susp (Smith Dorsey)
 W/Hyascyamus ext, Ipecac syr, Iscorice ext See: Crupic, Tab (Jenkins Labs)
 W/Milk of bismuth, kaolin & pectin See: Mul Sed, Susp (Pitman Moore)
 W/Pectolin & kaolin See: Parepectolin, Susp (Rorer)
 W/Pepsin, bismuth subsalicylate, zinc sulfocarbamate, salol & methyl salicylate See: Parepeptol, liq (Chicago Pharm)
 W/Tartar emetic, ethyl nitrite, etc See: Brown Mixture, liq (Merrell)
- PAREMYCIN.** (G F Harvey) Neomycin sulfate 300 mg, opium tincture 0.2 mg/oz Bot 3, 6 oz
 Use: Diarrheas
- PARENAMINE, NND 1962** (Wentrop Labs) Protein hydrolysate Sol 6%, Plask 1000 cc, 15%, Vial 100 cc
 Use: IV, I.S & S.C. dietary supplement
- PARENEX.** (Smith, Miller & Patch) Thiamine HCl 10 mg, niacinamide 50 mg, riboflavin 1 mg, pyridoxine HCl 1 mg, Cal pantothenate 5 mg, chlorbutanol 0.5%/cc Vial 15 cc
 Use: Vit B complex therapy
 W/Liver (Smith, Miller & Patch) Thiamine HCl 10 mg, niacinamide 20 mg, riboflavin 1 mg, pyridoxine HCl 1 mg, Cal pantothenate 5 mg, liver inj mcg, 8 12 5 mcg/cc. Vial 15 cc
 Use Anemias
- PARENEX-C.** (Smith, Miller & Patch) Thiamine HCl 50 mg, niacinamide 50 mg, ascorbic acid 50 mg, riboflavin 3 mg, pyridoxine HCl 5 mg, panthanol 5 mg, gentisic acid ethanalamide 2½%/cc Vial 10 cc.
 Use: Vit B complex & Vit C therapy
- PARENEX MODIFIED.** (Smith, Miller & Patch) Vit B 1 5 mg, B 2 2 mg, B 6 3 mg, niacinamide 50 mg, Vit C 15 mg, panthanol 3 mg, gentisic acid ethanalamide 2½%/cc Vial 10 cc
 Use: Vit B complex
- PARENCILLIN, NND 1962** (Merrell)
 Aq Susp: Procaine penicillin G 300,000 u/dose, Vial 10 cc.
- PARENOGEN, NND 1962** (Cutter) Fibrinogen, human for IV, int Vial 1 Gm pow
 Use (2.6 Gm) Hypofibrinogenemia, often associated w/premature separation of placenta
- PARENS-4DO** (Hiss) Penicillin 400,000 u/ Cap Bot 100s 500s
 Use Infections due to penicillin sensitive organisms
- PARENTOSOL-B.** (Squibb) Vit B 1 10 mg, B 2 4 mg, niacinamide 200 mg/cc Amp 1 cc, Box 5s Vial 10 cc
 Use Pellagra
- PARENTRACIN** (Reed & Cornrick) Not listed
 Bacitracin 60,000 u Vial 1 dose
 Use For hospital use only
- PARENZYME, NND 1962** (National Drug) Trypsin Vial 5 mg/cc Cryst trypsin in sesame oil Vial 5 ml Pkg 1s Aq: 25 mg w/5 ml diluent Pkg 1s Single dose pkg: 1 vial 12,500 NP u (5 mg) lyophilized trypsin
 Use: IM; reduces acute, local inflammation (phlebitis, thrombophlebitis, traumatic wounds, etc).
- PARENZYME B** (National Drug) Trypsin 5 mg in polyethylene glycol 4000 base/Buccal Tab Vial 24s
 Use: Inflammation, sprains and edema
- PARENZYME OINTMENT** (National Drug) Crystalline trypsin 2 mg, crystalline chymotrypsin 6 mg, 9 aminoacridine HCl 2 mg in water soluble base/Gm Tube 1 oz
 Use: Debridement and healing of external lesions
- PARPECTOLIN.** (Rorer) Paregoric 3.7 cc-pectin 2.5 gr, kaolin 85 gr/fl oz Susp-Bot 4 & 8 fl oz
 Use: (1-2 tsp, 11 d) Nonspecific diarrheas
- PARPEPTOL.** (Chicago Pharm.) Paregoric 60 min, pepsin 4 gr, bismuth subsalicylate 4 gr, zinc sulfocarbamate 49 mg, salol 100 mg, methyl salicylate 8 min./oz Bot 1 pt
 Uses Diarrhea
- PARITHOXYCAINE NCI, NND 1959** See Intracaine HCl (Squibb)
- PAR-FLUORO-VITE.** (Para Pharmacol) Sodium fluoride 2.21 mg, Vit A 2500 u, D 250 u, B-1 1 mg, ascorbic acid 30 mg.

- B-6 0.1 mg., B-12 3 mcg., and trace minerals/tab.
 Use: Control of dental caries.
- PARGRAN CAPSULES.** (Squibb) Vit. A 5000 u., D 400 u., B-1 2 mg., B-2 2 mg., B-6 0.5 mg., C 50 mg., niacin 20 mg., pantothenic acid 5 mg., Vit. B-12 2 mg., folic acid D25 mg., menadione 2 mg./Cap. Bot. 30s, 100s, 1000s.
 Use: Multivitamin therapy.
- PARGRAN-M.** (Squibb) Vit. A 5000 u., D 500 u., B-1 3 mg., B-2 3 mg., B-6 0.5 mg., C 50 mg., niacin 25 mg., pantothenic acid 5 mg., Vit. E 5 mg., B-12 2 mcg., folic acid 0.1 mg., menadione 2 mg., w/Cal., Phos., Iodine, Iron, Pot., copper, Mang, Mag., cobalt, zinc & molybdenum/Cap. Bot. 100s, 250s.
 Use: Multivitamin therapy.
- *PARGYLIN HYDROCHLORIDE.** (Abbott Lab) N-methyl-N-(2-propynyl)benzylamine hydrochloride. See Eulanyl (Abbott Labs.)
- PARID.** (Burlington) Piperazine hexahydrate 500 mg / 5 ml. Bot. 1 pt.
 Use: Treatment of pinworm and round-worm infections.
- PARISAMINE.** (Gold Leaf) Pyrilamine maleate 10 mg., aspirin 3½ gr., acetophenetidin 2½ gr., caffeine anhy. ¼ gr., p-aminobenzoic acid ½ gr./Tab. Bot. 100s, 1000s.
 Use: Head colds & hay fever.
- PARKINSONISM** (Parosympatholytic agents)
 See: Artone (Lederle).
 Coramiphen HCl (Geigy).
 Cogentin, Tab. (Merck Sharp & Dohme)
 Diparcol. (No Mfr. currently lists.)
 Disipal (Riker)
 Kemadrin, Tab. (Burroughs Wellcome)
 Pogitone HCl. (Lilly)
 Panpoint (Geigy).
 Parsidal (Warner-Chilcott).
 Phenothiazine HCl (Wyeth).
 Pipanol. (Winthrop Labs)
 Prantal. (Scheingl)
 Trihexyphenidyl HCl (Various Mfr.)
- PARLITE.** (Lederle) Vitamins B-1 10 mg., B-2 10 mg., B-6 5 mg., B-12 25 mcg., C 300 mg., niacinamide 150 mg., and sod. pantothenate 10 mg / 5 cc. Vial (dry) 5 cc., 25s.
 Use: I.V. or I.M. Vitamin deficiencies
- PARLUGUAN.** (Panray) Whole milk solids fortified w/dextrins & maltose 2.4 Gm., Mag trisilicate 0.25 Gm., Mag. oxide D.12 Gm., Cal. carb. 0.12 Gm., Mag. carb. 0.03 Gm./Trache Bot. 25s.
 Use: 1-4 troches between meals; antacid in ulcer management.
- *Generic name recognized by N.F.**
- PARMAL.** (Central) Not listed Pyrilamine maleate Caps Timed disintegration (75 mg) Bot. 100s, 500s
 Use: (1 cap. morning or evening) Antihistaminic.
- PARNATE.** N.N.D. 1962. (Smith Kline & French) Tranylcypromine sulfate. 10 mg / Tab Bot. 30s, 500s
 Use: Antidepressant; MO inhibitor
- PARODYNE.** See: Antipyrene (Various Mfr.)
- PARODIN.** N.N.R. 1950. (Parke, Davis) Parathyroid ext., 100 U.S.P. u./cc. Vial 5 cc.
 Use: S.C. & I.M.; parathyroid therapy.
- PAROLEINE.** See: Petrolatum Liquid (Various Mfr.)
- *PAROMOMYCIN.** An antibiotic substance obtained from cultures of certain *Streptomyces* species, one of which is *Streptomyces rimosus*.
 See: Humatin, Kapseal (Parke Davis)
 W/Chloramphenicol, hydrocortisone acetate, in a liquid petrolatum base See: Humacort Ointment, (Parke Davis).
- PAROXYL.** See: Acetartane, Tab. (Abbott).
- PARPANIT.** See: Coramiphen Hydrochloride (Panpanit, Tab. Geigy)
- PARESINE.** N.N.R. 1952. (Abbott) Plialle paraffin. Paraffined face mesh surgical dressing, various sizes.
 Use: Protective surgical dressing.
- PARSIDOL HCl.** N.N.D. 1962. (Warner-Chilcott) Ethopropazine, N.N.D. 1959. Prolomine HCl, lysivone, 10-(Diethylamino-1-propyl) Phenothiazine HCl, Tab (10 mg., 50 mg) Bot. 100s, 500s; (100 mg.) Bot 100s, 500s
 Use: Parkinsonism.
- PARSLEY CONCENTRATE W/Garlic conc.**
 See: Allimin, Tab. (Van Patten).
- PARZOCAL.** (Central) Zinc oxide 8%, colamine 8%, propesin 1%, benzyl alcohol 1%, pyrilamine maleate 1%, p-chloro-m-xyleneol 0.8%, menthol 0.075%, comphos 0.05%. Bot. 4 fl oz., 1 pt., gal
 Use: Dermatitis.
- PARZONE.** (Mallinckrodt) Brand of dihydrocodeine bitartrate
- PAS.** N.N.D. 1962. (Lilly) p-Aminosalicylic acid. Tab. 7½ gr. Enseals. Bot 500s, 1000s, 5000s
 Use: Tuberculosis.
- PASALATE.** N.N.D. 1962. (Dumas-Wilson) Sodium para-aminosalicylate. Tab. Bot. 100s, 1000s.
- PASANOL.** (Tilden) Digoxin 1/350 gr., strophanthin 1/25 gr., strychnine sulfate 1/25 gr., nitroglycerin 2/25 gr., cactus grandiflorus ½ gr./fl. oz. Bot. Pts Gals
 Use: Cardiac and vasomotor stimulant.

PASARA CALCIUM, NNO 1962 (Smith Darsey) Cal p aminosaliclylate Cap 0.5 Gm Bot 100s Pow Bot 1 oz, 25 lb Tab 0.5 Gm, Bot 100s

PASARA CALCIUM GRANULATE. (Smith Darsey) Calcium aminosaliclylate anhydrous 0.85 Gm equiv to para aminosaliclylic acid 0.75 Gm / Gm Bot 1 lb Packets 5 Gm / Packet Box 1b

PASARA SODIUM, NND 1962 (Smith Darsey) Sod p aminosaliclylate Cap 0.5 Gm, Bot 100s, 1000s Pow, Bot 1 oz, 1 lb Tab 0.5 Gm, Bot 100s 1000s

PASATON (Lanett) Sod p aminosaliclylate 0.5 Gm / tab EC chocolate Bot 100s 500s, 1000s

PASCA PACK GRANULES. (Barnes Hind Pasna) Calcium aminosaliclylate 5 Gm / Pack Granules are not enteric coated Boxes 90 packs

Use Calcium aminosaliclylate therapy

PASEM SODIUM, NND 1962 (Massengill) Sod p-aminosalicylate Cap 0.5 Gm Bot 500s Tab 0.5 Gm, Bot 500s

Use Tuberculosis

PASKALIUM, NNO 1962 (Glenwood) Pot p aminosaliclylate Tab (500 mg) Bot 100s 1000s Pow Bot 100 Gm, 1 lb Envules 3 Gm, sealed pkg 100

Use Antituberculosis agent

PASKATE, NNR 1957 (Lilly) Pot p amina salicylic acid 7 1/2 gr / poly Bot 250s, 1000s

Use: Indicated as an adjunct to other anti tuberculous drugs isoniazid, streptomycin & dihydrostreptomycin

PASMEO SODIUM, NNO 1962 (Inter-medico) Sod p aminosaliclylate Tab 0.5 Gm Bot 100s

PAS BUFFERED. (Lanett) p Aminosaliclylic acid 7 1/2 gr, sod biphosphate 3 gr / cap Bot 100s, 500s 1000s

Use: Tuberculosis

PASNA PACK GRANULES (Barnes Hind Pasna) Sodium aminosaliclylate 625 Gm / pack Granules are not enteric coated Boxes 90 packs

Use Sodium aminosaliclylate therapy

PASNA TRI-PACK GRANULES. (Barnes Hind Pasna) Tri Pack 300 / Isonazid 300 mg, sodium aminosaliclylate 15 Gm pyridoxine HCl 30 mg / 3 packs Tri Pack 600 Isonazid 600 mg, sodium aminosaliclylate 15 Gm, pyridoxine HCl 30 mg / 3 packs Boxes 90 packs

Use Sodium aminosaliclylate INH and Vit B 6 therapy

PAS RESIN See: Rexipos (Squibb)

PASSI-BARB. (C D Smith) Phenobarbital 1/4 gr, hyoscyamus ext 1/8 gr, Sod bicarb 1/2 gr, passiflora tr 1 min, chamomilla tr 1

min, pulsatilla tr, 2 min / Tab Bot 100s 1000s

Use: Antispasmodic & sedative

PASSIBARBITAL. (C D Smith) Same formula as Passi Barb Tab Bot 1000s

PASSIFLORA. (Penick) Dried flowering & fruiting tops of Passiflora incarnata

W/Phenobarbital Sod See: Barbiflora Compound, Sol (First Texas)

W/Stronium bromide See Bromoflora, Liq (Warren Teed)

PASSIPHEN (McNeil) Phenobarbital 1/4 gr, hyoscyamus ext 1/8 gr, passiflora ext 1 gr, valerian ext 1 gr / Tab Bot 100s 1000s

Use: 1 2 q 3 h; sedative & antispasmodic

PASTA DURET. (Doak) Sulfur, coal tar, camphor, chaulmoogra oil, zinc oxide paste Jar 4 oz, 1 lb, 5 lb

Use Pruritus

PASTURELLA-SALMONELLA-CHOLERA-SUIS (Cutter) Aluminum hydroxide adsorbed Vial, 10 dose, 50 cc; 50 dose, 250 cc

PASTURELLA TULARENSIS ANTIGEN 10 000 million organisms/cc

Lederle—Vial 5 cc 20 cc

Merck Sharp & Dohme—Vial 5 cc

Use Diagnostic use for tularemia

PATHEBA (Broyten) Vit D 300 u, 8 1 2 mg, C 25 mg, dicalcium phosphate 136 mg, ferrous lactate 50 mg, bone flour 400 mg, Vit 8 6 0.5 mg / Tab Bot 100s

Use: Calcium, vitamin & iron deficiencies

PATHIBAMATE-400 (Lederle) Meprobamate 400 mg, trihexethyl chloride 25 mg / Tab Bot 100s 1000s

Use Treatment of GI tract disorders associated with anxiety and tension

PATIBAMATE 200. (Lederle) Meprobamate 200 mg trihexethyl chloride 25 mg / Tab Bot 100s & 1000s

PATILON CHLORIDE, NNO 1962 (Lederle) Trihexethyl chloride 3 Diethylamino-1 cyclohexyl 1 phenyl 1 propanol ethachloride Tab 25 mg Bot 100s 1000s Amp (10 mg / cc) 1 cc Box 6s 25s

Use: 25 50 mg 3-4 times daily; anticholinergic in the treatment of peptic ulcer

W/Meprobamate See Pathibamate (Lederle)

W/Phenobarbital Pathilon 25 mg, phenobarbital 15 mg / tab Bot 100s, 1000s (Lederle)

Use Anticholinergic with sedative in the treatment of peptic ulcer

PATILON IODIDE (Lederle Lab) See Trihexethyl iodide

PATILON PEDIATRIC DROPS (Lederle) Trihexethyl chloride 5 mg, propylparaben

- 0.2 mg., methylparaben 0.8 mg / 5 cc. Bot. 15 cc.
 Use: Infant colic.
- PATHILON SEQUELS.** (Lederle) Tridihexethyl chloride 75 mg / Cop Bot 30s
 Use: Anticholinergic
- PATHILON WITH PHENOBARBITAL SEQUELS.** (Lederle) Tridihexethyl chloride 75 mg., phenobarb., phenobarbital 45 mg / Cop. See: Pathilon Sequels, Cops. (Lederle)
- PAUSTABS.** (Frankroy labs) Bot. 30s, 100s.
 Use: Treatment of menopause.
- PAVATRINE.** (Searle) Corbofluorene amine-ester. 2-diethylaminoethyl-9-fluorene-carboxylate HCl, Tab. 125 mg., Bot. 100s, 1000s.
 Use: [1-2 t.i.d.] Antispasmodic.
- W/Phenobarbital.** (Searle) Pavatrine HCl 125 mg., phenobarbital 15 mg./Tab. Bot. 100s, 1000s.
 Use: [1-2 t.i.d.] Antispasmodic & sedative.
- PAVERIL & AMYTAL.** (Lilly) Dioxylone phosphate 200 mg., amytal 16 mg./Tab. Bot. 100s, 1000s.
 Use: [1 t.i.d.] Antispasmodic & sedative.
- PAVERIL PHOSPHATE.** (Lilly) 6,7-Dimethoxy-1-(4'-ethoxy-3'-methoxybenzyl)-3-methylisoquinoline. Dioxylone phosphate. Tab. 1½ gr., 3 gr., Bot. 100s, 500s, 1000s.
 Use: [1-2 dolly] Relax smooth-muscle spasm.
- *PAXAMATE.** (Upjohn) 4-biphenyl methyl-carbamate
 Use: anti-inflammatory agent
- PAXITAL.** (Warner-Chilcott) See: Meperazine
- PAZILLIN, A.D.R.** (Merck Sharp & Dohme) Procaine penicillin G 300,000 u. w/2% aluminum monostearate/cc. In sesame oil. Cartridge 1 cc., w/disposable syringe. Vial, multiple dose, 10 cc.
 Use: I.M.; penicillin therapy.
- P-B-SAL-C.** (Ulmer) Sod. salicylate 250 mg., p-aminobenzoic acid 250 mg., vit. C 20 mg / tab. E.C. Bot. 100s, 500s, 1000s.
 Use: [1-2 q.i.d.] Gouty rheumatism.
- W/Calcifene.** (Ulmer) P-B-Sal-C formula with calcifene 0.25 mg / tab. Bot. 100s, 500s, 1000s.
 Use: (2 q. br.) Gout.
- W/Esoprine.** (Ulmer) Pot. salicylate 280 mg., p-aminobenzoic acid 250 mg., vit. C 20 mg., physostigmine salicylate 0.25 mg., homatropine methylbromide 0.50 gr./tab. Bot. 100s, 500s, 1000s.
 Use: 1-2 q.i.d.] rheumatic therapy.
- P-B-SAL-C SODIUM FREE.** (Ulmer) Ammonium salicylate 250 mg., p-aminobenzoic acid 250 mg., ascorbic acid 20 mg./tab. Bot. 100s, 500s, 1000s
 Use: Analgesic & antirheumatic.
- * Generic name recognized by N.F., U.S.P. or A.M.A.
- PBSC COMPOUND-BIOTIC.** (Biotic Drug Co) Pot. penicillin G 200M u., dihydrostreptomycin base 0.2 Gm., bacitracin 2M u., Sod. caprylate 0.2 Gm./Gm. Oint., Tube 1 oz.
 Use: Dental paste for sterilization of root canals.
- PBSC ENDODONTIC THERAPULES.** (Novocol Chem. Co.) Pot. penicillin G 200M u., bacitracin 2M u., streptomycin sulf. 0.2 Gm and sod. caprylate 0.2 Gm./Therapules. Bot. 100s
 Use: Systemic infections
- PBSC POLYANTIBIOTIC PASTE.** (Praco Sol Chem. Co.) Pot. penicillin G 200M u., dihydrostreptomycin base 0.2 Gm., bacitracin 2M u., Sod. caprylate 0.2 Gm./Gm. Oint., Tube 1 oz.
 Use: Dental paste for sterilization of root canals.
- PBSC THERAPULE.** (Novocol) Penicillin G pot. 200M u., bacitracin 2M u., streptomycin sulf. 0.2 Gm., sod. caprylate 0.2 Gm. in silicone/cc.
 Use: Root canal sterilization
- P/C BGAM.** (Ingram) Lemac bioflavonoid complex 150 mg., ascorbic acid 150 mg., rutin 30 mg./cap. Bot. 100s.
 Use: Capillary fragility.
- PCMX.** See Para Chloro Meta Xylenol.
- P-D-N DENTAL CONIS.** (Novocol) Sod. penicillin G 15,000 u., dihydrostreptomycin sulfate 35 mg., neopaine HCl 5 mg./Cone. Pkg. 12s.
 Use: Vincent's infections & other infections of the oral cavity.
- PEACOCK'S BROMIDES.** (Peacock) Liq., Pot. bromide 6 gr., Sod. bromide 6 gr., ammonium bromide 3 gr./dr. Bot. 8 oz. Tab., Pot. bromide 3 gr., Sod. bromide 3 gr., ammonium bromide 1½ gr./Tab. Bot. 100s.
 Use: Sedative.
- PEANUT OIL W/Dextrose.** See: lipomul Oral (Upjohn).
- PEBZICOLE.** (Cole) Pepsin 6 gr., lactic acid 1 min., glycerin 38 min., benzoic acid 0.9 gr./fl. oz. Bot. 1 pt., 1 gal.
 Use: Digestant.
- PECEGRAM.** (Ingram) Herperidin chalcane sod 50 mg., ascorbic acid 50 mg /Cap. Bot. 60s, 300s
 Use: Orally; hemorrhagic conditions.
- PECO.** (Superior Pharm.) Acetophenetidin 2 gr., camphor ¼ gr., quinine sulfate ¼ gr., opiate root pow. 1/2000 gr., oleo 1/16 gr., podophyllin 1/40 gr., capsicum pow ¼ gr./Tab. Bot. 700s.
 Use: Common cold.

PECTALUM (C D Smith) Colloidal kaolin 90 gr, aluminum hydroxide 12 gr, pectin 2 gr, aromatics q s / fl oz. Bot 4 fl oz, 1 pt, 1 gal Also avail as Tab
Use Gastric antacid

***PECTIN**, NF (Various Mfr) See: Appela Pow (Winthrop Labs)

W/Aluminum hydroxide, hamotropine methyl bromide Mag trisilicate See Alucen, Tab (Central)

W/Aluminum hydroxide gel & Mag carb See Alupecc, Liq (Pitman Moore)

W/Dihydrostreptomycin, kaolin See Diamycin Susp (Maurry)

W/Dihydrostreptomycin sulf, aluminum hydroxide gel, polymyxin B sulf, atropulgite See Polymagma (Wyeth)

W/Furazolidone kaolin See Furoxone Liquid (Eaton)

W/Hamotropine methylbromide, Mag trisilicate aluminum hydroxide gel dried & Mag hydroxide
See: Gel A Nulrin Tab (Walker)

W/Kaolin

See Kaopectate Liquid (Upjohn)

Keatin, Liq (Blue Line)

Pacatin Liq & Tab (Massengill)

Pecta Kalin, Liq (Paul Maney)

Pectoloid, Liq (G F Harvey)

Pektomalt, Liq (Warren Teed)

W/Kaolin & aluminum hydroxide See: Kalpec, Liq (Wyeth)

W/Kaolin & bismuth subcarb

See: K B P, Liq (Cole)

Kaamuth, Liq (Elder)

W/Kaolin hyoscyamine sulf atropine sulf hyoscine HBr phenab See Donnagel products (Robins)

W/Kaolin & milk of bismuth See Kaolin Bismuth Pectin Compound (Pitman Moore)

W/Kaolin, milk of bismuth & paregoric See Mul Sed, Liq (Pitman Moore)

W/Kaolin neomycin sulfate See Neopeccate, Syr (Tyler)

W/Kaolin apium paw hyoscyamine sulfate atropine sulfate hyoscine HBr, & phenobarbital See Donnagel PG Susp (A H Robins)

W/Kaolin & sulfathiazole See Pecta thiazole Susp (Paul Maney)

W/Kaolin & zinc phenalsulfonate See Pectacel, Liq (Lilly)

W/Malic acid apple & carn syr See Pama ca Syr (Brean)

W/Neomycin, kaolin solal, zinc sulfacarbo late See: Neo Kalefan (Burrington)

W/Phthalylsulfacetamide & albumen tannate See Mucamide, Cap (Warren Teed)

* Generic name recognized by NF USP or AMA

W/Phthalylsulfacetamide & kaolin See: Thalin, Susp (Abbott)

W/Sulfaguanidine & kaolin See: Pamalin, Susp (Brean)

PECTIN-AGAR IN DEXTRI-MALTOSE (Mead Johnson) Pectin & agar are blended with Dextrin Maltose Cons 1 lb

Use: To modify milk for infants and children suffering from diarrhea

PECTO-BISMUTH, (Horton & Converse) Bismuth subcarbonate 40 gr, pectin 10 gr, suspended with bentonite 3% / fl oz Bot 4 oz, 1 pt, 1 gal

Use: Gastric and intestinal distress

PECTOCEL, (Lilly) Kaolin 90 gr, pectin 4.5 gr, zinc phenalsulfonate 1½ gr./oz Bot 1 pt, 1 gal,

Use: Intestinal distress

PECTOGUANIDINE (Corrtane) Sulfaguanidine 20 Gm Ir apium 008 cc, pectin (0.225 Gm, kaolin 30 Gm / 15 cc Bot pt

Use: Diarrheas

PECTO-KALIN (Paul Maney) Kaolin 92 gr, pectin 14 gr / oz Bot 1 pt, 1 gal Also available W/Neomycin Sulf 300 mg / oz Bot 4 oz and 1 pt

Use: Intestinal distress

PECTOL, (Roerig) Mineral oil 60% w/pectin Emulsion, Bot 14 oz

Use: Laxative

PEDAMETH, (Durst) di Methionine 0.2 Gm / Cap Bot 30s, 1000s

Use: 1 Cap added to contents of baby's formula To control diaper rash

PEDIACOF SYRUP, (Winthrop) Codeine phos 50 mg, phenylephrine HCl 2.5 mg, chlorpheniramine mol 0.75 mg, potassium iodide 750 mg / tsp Bot 16 fl oz

Use: Antitussive

PEDIAGESIC, (Columbus) Acataminaphen 60 mg, salicylamide 60 mg / 5 cc Bot 3 oz

Use: Analgesic

PEDIAHIST, (Columbus) Pyrilamine maleate 5 mg, methapyrilene HCl 5 mg, prophen pyridamine maleate 5 mg / 5 cc Bot 1 pt 1 gal

Use: Antihistaminic

PEDIAMINTIC (Columbus) Piperazine hexahydrate 100 mg / cc Elix Bot 1 pt, 1 gal

Use: Treatment of roundworm and pinworm infestations

PEDIAQUIL, (Columbus) Dihydrocodeinone bitartrate 1/12 gr, Pot guaiaac sulfonate 6 gr, squill 1½ gr, acetic acid dilute 27 min / fl oz Bot 3 oz, 1 pt, 1 gal

Use: Cough preparation

PEDIATRON, (Paul Maney) Vit D 10,000 u, A 50,000 u, sad ascorbate 100 mg / 5 cc

Use: Minor respiratory diseases

PEDICOLE-PEDIATRIC. (Lemmon) d-Methorphan HBr 7.5 mg., glyceryl guaiacolate 25 mg., ammonium chloride 50 mg., sod. citrate 50 mg./5 cc. Bot. 3 oz., pts.

Use: Antitussive.

PEDICOO. (Parker) Lotion. Bot. 6 oz.

PEDICRAN. (First Texas Chemical) Vit. B-12 150 mcg., B-1 18 mg., B-6 6 mg., nicotinamide HCl 60 mg., cal. pantothenate 12 mg./fl. oz. Bot. 1 pt.

Use: Secondary anemias.

PEDICULICIDES.

See: Benzene Hexachloride, N.N.R. 1953 (Various Mfr.).

Benzyl Benzoate, N.N.R. 1953 (Various Mfr.).

Bornate (Wyeth).

Chlorophenothane, N.N.B. 1953 (Various Mfr.).

Cuprex (Merck).

DDT (Various Mfr.).

Gexane (Strassenburgh).

Hexacide (Central).

Kwell. (Reed & Carnrick)

PEDIDROPS. (Tilden) Ephedrine sulfate 1% in an aqueous buffered isotonic solution. Bot. 1/2 oz. w/dropper, pts.

Use: Nasal congestion.

PEDINOL NYDRISINOL CREME. (Pedinol) Sulfonated hydrogenated castor oil. Creme, Tube 1 oz.

Use: Skin irritations.

PEDISAL-PEDIATRIC. (Lemmon) Salicylamide 65 mg., chlorpheniramine maleate 0.75 mg./5 cc. Bot. 1 oz.

Use: Antihistaminic-antipyretic-analgesic.

PEDITUSS. (Sutliff & Cose) N-Acetyl-p-aminophenol 64.5 mg., Sod. salicylate 129 mg., phenylephrine 2 mg., glyceryl guaiacolate 20 mg./5 cc. Bot. 4 oz., 1 pt., 1 gal.

Use: Analgesic, expectorant.

PEDIVITE-C. (Crystal) Vitamin D 10M u., A 50M u., sodium ascorbate 100 mg./5 cc. Bot. 4 oz., pt.

Use: Vitamin deficiency.

PEDO-SOL. (Warren Pharmacol) Butabarbital 30 mg., chlorpheniramine maleate 1.5 mg., hyoscine hydrobromide 0.1 mg./5 cc. Elixir. Bot. 16 oz.

Use: Sedative.

PEDRIN. (Braemmel) Salicylamide 1 gr., ascorbic acid 10 mg./5 cc. Syr. Bot. 4 oz., 1 pt.

Use: (1/2-1 dr. q. 3-4 hr.) Pediatric, analgesic.

PEGANONE, N.N.D. 1962. (Abbott) Ethotain. 3-Ethyl-5-phenylhydantoin. Tab. 250 mg. or 500 mg. Bot. 100s, 1000s.

Uses Orally, anticonvulsant in the control of grand mal seizures.

PEGA PALO FORTIDOM Tr. (Quimica Dominicana) Hydro-alcoholic extract of the woody vine of *Elychnasia pyramidalis* (*Dalicholus pyramidalis*) containing 15% of the drug. Bot. 1 oz.

Use: Fatigue and as an aphrodisiac in aged men.

PEKTAMALT. (Warren-Teed) Koolin 100 gr., pectin 10 gr./oz. Bot. 4 oz., 1 pt.

Use: (1-2 tsp. p.r.n.) Intestinal distress.

PELARGON. (Nestle) Dried milk, glucose, sucrose, starch, iron & Vit. A 1725 u., O 460 u., C 230 mg., B-1 2.8 mg., B-2 0.6 mg./Qt. of sol. Pkg. 16 oz.

Use: Infant dietary feeding.

PELARGONIC ACID w/zinc pelargonate, sulfur. See: Pellar, Ointment and Powder (Crookes-Borne)

PELENTAN. See: Ethyl Biscoumacetate. (No Mfr. currently lists.)

PELLAR. (Crookes-Borne) Pelargonic acid 4%, zinc pelargonate 10%, sulfur 2%. Oint., Pelargonic acid 2 1/2%, zinc pelargonate 20%, zinc sulfide 3%. Pow.

Use: Treatment of fungus infections.

PELLBASS. (Kirkman) Pentobarbital 60 mg., phenobarbital 30 mg., d-amphetamine sulf./cap. timed disint. Bot. 100s, 500s, 1000s.

Use: To control emotional states.

PELLCAPS. (Kirkman) d-Amphetamine sulf. 10 mg. & 15 mg./cap. timed disintegration. Bot. 100s, 500s, 1000s.

Use: Central nervous system stimulant.

PELIDONNA. (Kirkman) Mephobarbital 65 mg., homatropine methylbromide 5 mg., atropine methyl nitrate 1 mg., hyoscine HBr 0.02 mg./cap. Bot. 100s, 500s, 1000s.

Use: Depressant-antispasmodic.

PELLITOL. (Pitman-Moore) Bismuth 5%, bismuth subgallate 1%, bismuth subnitrate 9%, zinc oxide 17%, calamine 10%, zinc-nacea 3%, code oil 1%, pyraligneous acid 0.4%. Oint., Tube 1 oz., Jar 1 1/2 oz., 1 lb.

Use: Skin antiseptic.

PELVICINS. (Schenley) Pot. penicillin G 200,000 u. Vag. Supp., Jar 6s, 12s, 50s.

Use: Penicillin therapy.

PIMBRIN, A.D.R. (Novaco) Aspirin 5 gr., pentobarbital Sod. 3/4 gr. & 1 1/2 gr./Cap. Bot. 100s, 500s

Use: (1-2 l.i.d.) Analgesic w/sedative.

PIMBULE. (Novaco) Pentobarbital sod. Cap. 3/4 gr., 1 1/2 gr., Bot. 100s, 500s

Use: Sedative

- *PEMOLINE.** (Norden Pharm Labs, Inc)
2 Imino 5 phenyl-4 oxazolidinone
Use Central nervous system stimulant
- PENZOLES** (Pitman Moore) Tyrothricin 1 mg, benzocaine 5 mg, phenyl mercuric nitrate 0.5 mg/Loz Bot 100s, 500s.
Use: Infections of throat & mouth
- PENAGE LIQUID** (Manne Labs) Pentylene tetrazol (Penage) 100 mg Vit B 1 10 mg; B 6 10 mg; B 12 mcg; alcohol 15% per tsp /5 cc Bot Pints Gals
Use Geniatric
- PENALEV, NND 1962** (Merck Sharp & Dahme) Tab 50 000, 100 000, 200 000, 250 000, 500 000 & 1 000 000 u of pot penicillin G Vial 12s 100s
Use: Soluble penicillin tab
- PENAMIN,** (Preme) Procaine penicillin 100,000 u, methoprylene hydrochloride 30 mg, salicylamide 150 mg, acetophenone hidin 120 mg, caffeine 30 mg /Tab
Use: Relief of symptoms due to colds
- PEN AQUA.** (Bristol) Procaine penicillin G 300,000 u, Pot penicillin G 100 000 u /dose Vial 1 dose 5 dose, 10 dose
Use: I.M.; penicillin therapy
- PENAZINE,** (Elder) Cryst Pot penicillin G 100,000 u, sulfadiazine D 17 Gm, sulfa merazine 0.17 Gm, sulfamethazine 0.17 Gm /Tab Bot 24s
Use: Penicillin w/triple sulfonamide therapy
- PENCARD** (Cole) Pentoerythritol tetranitrate 10 mg /Tab Bot 100s, 500s, 1000s Also avail with Phenobarbital ¼ gr
Use: Angina pectoris
- PENCETYLON,** (Elder) Acetylsalicylic acid 5 gr, pentobarbital ¼ gr /Tab Bot 100s, 500s
Use: Analgesic and sedative
- PEN DROPS.** (Pfizer) Pot penicillin G 100 000 u buffered/cc liq, Bot w/dropper, 10 cc.
Use: Orally; penicillin therapy
- PEN-DUOSTREP 6 ½ (Dry).** (Merck Sharp & Dahme) Penicillin G, procaine 400 000 u, penicillin G, pot 200 000 u, dihydrostreptomycin sulfate 0.25 Gm, streptomycin sulfate 0.25 Gm /2 cc dose Vial 1 dose, 5 dose
Use I.M., treatment of most moderate infections caused by susceptible pathogens
- PEN-DUOSTREP 6-1 (Dry).** (Merck Sharp & Dahme) Penicillin G, procaine 400,000 u, penicillin G, pot 200 000 u, dihydrostreptomycin sulfate 0.5 Gm, streptomycin sulfate 0.5 Gm /3 cc dose Vial 1 dose, 5 dose
Use: I.M., treatment of most moderate infections caused by susceptible pathogens
- PENESTHETTES,** (Schenley) Pot penicillin G 5000 u, benzocaine 15 mg /trache Bot 20s
Use: Infections of throat & mouth
- *PENETHAMATE HI.** See Benzylpenicillin 2 diethylaminoethyl ester HI
- PENFONYLIN.** See: Penid Sulfas, 1ab [Squibb]
- PEN-G-CAP.** (Upjohn) Cryst procaine penicillin G 200,000 u, in peanut oil containing 2% w/v aluminum monostearate/Cap Bot 12s, 100s, 500s
Use: Oral penicillin
- *PENICILLAMINE,** (Merck, Sharp & Dahme) D J mercaptavoline
Use: Lenticular degeneration
- PENICILLIN, NND 1960** Unless clarified, it means an antibiotic substance or substances produced by growth of the molds *Penicillium notatum* or *P. chrysogenum*
- PENICILLIN-152 POTASSIUM** See: Phen ethicillin Potassium
- PENICILLIN, ALPHA-PHENOXYETHYL.** See: Phenethicillin Pot
- *PENICILLIN, ALUMINUM, NND 1962** ADR (Hynson, Westcott & Dunning) 1ab 50M u, Vial 12s 1955
- PENICILLIN, CALCIUM,**
Squibb—Trache 20M u, Vial 12s
Upjohn—Oint 1M u/Gm, Tube 1 oz;
Ophth oint 1M u/Gm, Tube 1 dr
Winthrop—Trache 1M or 10M u, Vial 12s
Buffered, U S P XII
Parke, Davis—Tab Buffered w/Sod cit rate, 50M u
Merck Sharp & Dahme—Tab Buffered w/Cal carb, 50M or 100M u
In oil (Sesame oil) & wax (parenteral) 1M Sterone—N R 1952 Cortan (300M u) 1 cc Vial (300M u /cc) 10 cc
- PENICILLIN, DIMETHOXY-PHENYL**
See Dimocillin Vial (Squibb)
Staphicillin, Vial (Bristol)
- PENICILLIN-DIHYDROSTREPTOMYCIN, N F**
for Susp Sterile See Table Procaine Pen G Comb
- PENICILLIN, EORTIEED**
Sees Penicillin G, Procaine W/Penicillin G Potassium (Various Mfr)
Penicillin G, Procaine W/Penicillin G Sodium (Various Mfr)
W/Dihydrostreptomycin or streptomycin
See Penicillin G Procaine W/Penicillin G Potassium & Dihydrostreptomycin
- *PENICILLIN G, BENZATHINE, U S P**
NND 1962 ADR Steril Susp U S P N

*Generic name recognized by NF USP or AMA

N-Dibenzylethylene diamine dipenicillin G
See: Benzathine Penicillin G.

Bicillin. (Wyeth)
Neolin. (Lilly)
Permopen. (Pfizer)

PENICILLIN O, DIBENZYLETHYLENEDI-AMINE. See: Benzathine Penicillin G.

PENICILLIN G, DIETHYLAMINOETHYL BSTER HYDRIODIDE SALT. Procaine Penicillin G. See: Neo Penil. Vial (Smith, Kline & French).

PENICILLIN G 1-PHENAMINE. (Penicillin G salt of N-Methyl-1,2-diphenyl-2-hydroxyethylamine.) See: Campenamine, Vial (Commercial Solvents).

PENICILLIN G NYDBABAMINE, N.N.D. 195B. See: Compoicillin, Susp. (Abbott)

PENICILLIN, PHENOXYMETHYL, U.S.P. See: Penicillin V, Preps.

PENICILLIN G, POTASSIUM, U.S.P., N.N.D. 1962, A.D.R. (Potassium Penicillin, Benzyl Penicillin Pot.)

Inhalation, N.P. X

Cutter—Cartridge 100M u.
Schenley—Cartridge 100M u.
Upjohn—Cap. 100M u.
Abbott—Cartridge 100M u. Use in aerosol sol. Tab. 50M or 100M u.
Prema—Aerosol
Squibb—Dispulator
Lederle—Peniator

In oil, oral susp.

See: K-Cillin, Liq. (Bio-Roma).
Readicillin, Susp. (Upjohn).

Oral Liq., Buffered w/Sod. citrate.

See: Cilligran, Oral Sol. (Columbus)
L-Cillin (Leyden)
Cilforal. (Bristol)
Dram-Cillin (White).
Drop-Cillin (White).
Eskocillin 50 (Smith, Kline & French).
Eskocillin 100 (Smith, Kline & French).
Eskocillin 250 (See under Penicillin G, Procaine).
Flonacillin (Philadelphia Amp. Co)
Honey Cillin 400, Liq. (Physicians Products)
Liquopen (Pfizer).
Orapen (Schenley).
Penalev (Merck Sharp & Dohme)
Pen-Draps (Pfizer).
Penisem, Sol. (Mossangill)
Sharcilin Oral. (Merck Sharp & Dohme)
Sugracillin (Upjohn).

Oral Tab., Buffered w/Cal. carb.

Abbott—Dulcel 50M, 100M u. Tab. 50M, 100M, 250M, 500M u.
A.P.C.—Tab. 100M u.
Blue line—Tab. 250M u.

* Generic name recognized by N.F., U.S.P. or A.M.A.

Commercial Solvents—Tab. 50M, 100M u. K.P.C.-400, Tabs. (Physicians Products)
Buffered w/glyceride & Sod. salts of fatty acids.

Cutter—Tab. 50M, 100M u.
Eilly—100M u.
Parke, Davis—Tab. 50M, 100M u.
Prema—Tab. 50M, 100M u.
Rexall—Tab. 50M u.
Schering—Tab. 50M, 100M, 250M, 1000M u.
Winthrop—Tab. 50M, 100M, 250M u.
See: Bupenna, Tab. (Burrough Bros.)
Cilloral. (Bristol)
Coafels (Schenley).
Orapen (Schenley).
Pentads (Squibb).
Penioral (Wyeth)

Oral Tab., Soluble, buffered.

Abbott—50M, 100M u.
Wyeth—50M, 100M, 250M u.
See: Pen-Eff, Tab. (Smith, Kline & French).

Oral Tab., Soluble, not buffered.

Amer. Pharm.—100M u.
Schenley—50M, 100M u.
Schering—50M, 100M u.
Squibb—50M, 100M u.
Upjohn—50M, 100M u.
Wyeth—50M, 100M u.
See: Hyosorb Penicillin, Tab. (Key Corp.)
Jacillin, Tab. (Jamieson Pharm.)
Nebulabs, Tab. (Prema)
Penalev, Tab. (Merck Sharp & Dohme)
PSI Pot. Penicillin G, Tab. (Pacific States Lab.)
E. D. Dis Pen-Tabs. (Rexall)
E. D. Pentabs. (Rexall)
Soltabs, Tab. (Commercial Solvents).
Til-Pen, Tab. (Tilden)

Parenteral.

Abbott—Vial 100M, 200M, 500M, 1000M, 5000M u.
Bio-Roma—Vial 100M, 200M, 500M, 1500M u.
Bristol—Vial 100M, 500M, 100M, 200M, 5000M u.
Commercial Solvents—Vial 100M, 200M, 500M u.
Cutter—Vial 100M, 200M, 500M, 1000M u.
Dwight, B. E.—Vial 100M, 200M, 500M, 1000M u.
Lilly—Vial 100M, 200M, 500M, 1000M u.
Parke, Davis—Vial 100M, 200M, 500M, 1000M, 5000M u.
Pfizer—Vial 100M, 200M, 500M, 1000M u.
Prema—Vial 100M, 200M, 500M, 1000M u.
Schenley—Vial 100M, 200M, 500M, 1000M, 2000M, 5000M u.
Upjohn—Vial 100M, 200M, 500M, 1000M u.

Penicillin G Potassium & Sodium Combinations

| Penicillin G | Others | Manufacturer | Brand |
|--------------|---|---------------------|---------------------------------------|
| pot. 100M v. | Acet-Dia-Mer | Whala | Dram-Cillin W/Triple Sulfas/3 cc. |
| pot. 100M u. | Acet-Dia-Mer | Schenley | Pencambisul, Tab. |
| pot. 100M v. | Acet-Dia-Mer | Merrell | Pansulfas W/Penicillin, Tab. |
| pot. 250M u. | Acet-Dia-Mer | White | Dram cillin 250 W/Triple Sulfas/5 cc. |
| pot. 250M v. | Albomycin sod. 250 mg. | Upjohn | Alba Penicillin, Cap. |
| pot. 20M v. | Bacitracin 50 u. | Chas. Chem. | Chesillets, Troche |
| pot. 20M u. | Bacitracin 50 u. | Bristol | Clilorets, Troche |
| pot. 20M v. | Bacitracin 50 u. | Wyeth | Pendets, Troche |
| pot. 25M v. | Bacitracin 50 u. | Abbott | Bacillets, Troche |
| pot. 100M v. | Benamid 250 mg. | Preme | Candicillin, Troche |
| pot. 100M u. | Benamid 250 mg. | Merck Sharp & Dohme | Remanden-100, Tab. |
| pot. 250M u. | Benamid 250 mg. | Merck Sharp & Dohme | Remanden-100, Gran. |
| pot. 250M v. | Benamid 250 mg. | Merck Sharp & Dohme | Remanden 250, Tab. |
| pot. 100M v. | Bensciline pen. G 200M u. | Merck Sharp & Dohme | Remanden 250, Gran. |
| pot. 3M u. | Benscoline 15 mg. | Bristol | Dibensipen, Vial/5 cc. |
| pot. 100M u. | d Desoxyephedrine HCl 1/4 % | Schenley | Panasthetin, Troche |
| pot. 150M u. | Dia-Mer-Thia Sulfas | Elder | Cafed Nose Drops/7.5 cc. |
| pot. 200M u. | Dihydrostreptomycin 0.2 Gm., boctraclo 2M u., sod. ceprylate 0.2 Gm., coprylate 0.2 Gm. | Esall | Pen-Taba W/Triple Sulfas, Tab. |
| pot. 200M v. | Dihydrostreptomycin 0.2 Gm., boctraclo 2M u., sod. coprylate 0.2 Gm. | Boyle Drug Co. | PBSC Compound Biallic, Oint./Gm. |
| pot. 200M v. | Dihydrostreptomycin sulf. 200 mg. | Proca-Sol Chem. Co. | PBSC Polyantibiotic Paste/Gm. |
| sod. 15M v. | Dihydrostreptomycin sulf. 33 mg., Neopoline HCl 5 mg. | Schenley | Syncrobin, Vag. Supp. |
| pot. 200M v. | Dihydrostreptomycin sulf. 1 Gm., procaine pen G 300M v. | Noracot | P.-D.-N Dental Cones |
| pot. 50M u. | Meth-Dia-Mer Sulfas | Schenley | Syncrobin, Vial/dose |
| pot. 50M v. | Meth-Dia-Mer Sulfas | Ellly | Neolisine W/Penicillin-G, Savorels |
| pot. 50M u. | Meth-Dia-Mer Sulfas | Abbott | Truo-Cillin-50, Tab. |
| pot. 100M u. | Meth-Dia-Mer Sulfas | Frasst | Trufalin-50, Tab. |
| pot. 100M v. | Meth-Dia-Mer Sulfas | Frasst | Trufalin-100, Tab. |
| pot. 100M u. | Meth-Dia-Mer Sulfas | National Drug | Sulfosaf W/Penicillin/5 cc. |
| pot. 100M v. | Meth-Dia-Mer Sulfas | Schenley | Teinacillin, Tab. |
| pot. 100M u. | Meth-Dio Mer Sulfas | Eron | Tricillin No. 1, Tab. |
| pot. 100M v. | Meth-Dio Mer Sulfas | Vampelt & Brown | Tritocillin/5 cc |

| Penicillin G | Others | Manufacturer | Brand |
|--------------|---------------------|-----------------------|--|
| pot 100M u | Meth Dia Mer Sulfas | Success Chem. Co. | Sulfacilin, Tab |
| pot 100M u | Meth Dia Mer Sulfas | Upjohn | Sulfa Sugracilin/5 cc. |
| pot 100M u | Meth Dia Mer Sulfas | Tilden | Pentrizine, Tab |
| pot 100M u | Meth Dia Mer Sulfas | Bio Rama | Ramasul 100, Tab. |
| pot 100M u | Meth Dia Mer Sulfas | Lilly | Savareis, Tab. |
| pot 100M u | Meth Dia Mer Sulfas | Upjohn | Biasulfa 100M, Tab |
| pot 100M u | Meth Dia Mer Sulfas | Smith Darsey | Darsalfin W/Triple Sulfa, Tab. |
| pot 100M u | Meth Dia Mer Sulfas | Caemir Funk | Pentrisul, Tab. |
| pot 100M u | Meth Dia Mer Sulfas | Merck Sharp & Dahme | Pentresamide, Tab. |
| pot 100M u | Meth Dia Mer Sulfas | Merck Sharp & Dahme | Pentresamide Pediatric Gron./5 cc. |
| pot 100M u | Meth Dia Mer Sulfas | Elder | Phenazine, Tab |
| pot 100M u | Meth Dia Mer Sulfas | Cutter | P53, Tab |
| pot 100M u | Meth Dia Mer Sulfas | Smith, Kline & French | Estacilin Sulfas/5 cc. |
| pot 100M u | Meth Dia Mer Sulfas | Wassengill | Trisem Pen/5 cc. or Tab |
| pot 100M u | Meth Dia Mer Sulfas | Central Pharm | Trisulfazene W/Penicillin, Tab. |
| pot 100M u | Meth Dia Mer Sulfas | Elder | Penazine, Tab |
| pot 150M u | Meth Dia Mer Sulfas | Lilly | Neopenzine 150, Tab. |
| pot 150M u | Meth Dia Mer Sulfas | Wyeth | Sulfabiatic, Tab |
| pot 150M u | Meth Dia Mer Sulfas | Abbott | True-Cilin 150, Tab. |
| pot 200M u | Meth Dia Mer Sulfas | Lilly | Neopenzine Tab |
| pot 200M u | Meth Dia Mer Sulfas | Bio-Rama | Ramasul 200, Tab |
| pot 250M u | Meth Dia Mer Sulfas | Squibb | Penid Sulfas, Tab |
| pot 250M u | Meth Dia Mer Sulfas | Upjohn | Biasulfa 250M, Tab. |
| pot 250M u | Meth Dia Mer Sulfas | Armour | Deltamide W/Penicillin, Tab. or 5 cc. |
| pot 250M u | Meth Dia Mer Sulfas | Merck Sharp & Dahme | Pentresamide 250, Tab. |
| pot 250M u | Meth Dia Mer Sulfas | Cutter | P53 Tab |
| pot 250M u | Meth Dia Mer Sulfas | Bio Rama | Ramasul 250, Tab |
| pot 250M u | Meth Dia Mer Sulfas | Success Chem Co | Sulfacilin, Tab |
| pot 250M u | Meth Dia Mer Sulfas | Upjohn | Sulfa Sugracilin Fortified/5 cc. |
| pot 250M u | Meth Dia Mer Sulfas | Evron | Tricilin Na 2, Tab |
| pot 300M u | Meth Dia Mer Sulfas | Winthrop Labs. | Citamide, Susp /5 cc. |
| pot 300M u | Meth Dia Mer Sulfas | Bristol | Calloral Susp W/Triple sulfonamides/ 5 cc |
| pot 300M u | Meth Dia Mer Sulfas | Lilly | Neopenzine 300, Tab. |
| pot 300M u. | Meth Dia Mer Sulfas | Upjohn | Sulfa Redicilin Susp /5 cc |

Penicillin G Potassium & Sodium Combinations

| Penicillin G | Others | Manufacturer | Brand |
|--------------|---|----------------|------------------------------------|
| pot. 300M u. | Meth-Dio-Mer Sulfos | Abbott | Truo-Cillin-300, Tab. |
| sod. 350M u. | Neo-Synephrina HCl 1/4 % | Winthrop Labs. | Neo-Synephrina W/Penicillin/15 cc. |
| pot. 200M u. | Novahistine (Phenylephrine HCl 10 mg., prophenpyridamine molecule 12.5 mg) | Pitman-Moore | Novahistina W/Penicillin, Cap. |
| pot. 250M u. | Phenyltolazamine dihyd. cit. 25 mg., salicylamids 100 mg., phenacetin 100 mg. | Patch | Elpainet, Tab. |
| pot. or sod. | Procaina pen. G See: Tablets of Procaina Pen. G Combinations | | |
| pot. & sod. | Procaina pen. G and Benzothina Pen. G. See: Tablets of Procaina Pen. G Combinations | | |
| pot. 2M u. | Streptomycin sulf. 10 mg. | Schanley | Synacrin, Oint./Gm. |
| pot. 5M u. | Sulfanilamids 0.25 Gm. | Abbott | Powdulator, Tab. |

Penicillin (Merrell).

Penicillin G Procaine Combinations on pp. 552 and 553.

*PENICILLIN G, SODIUM, U.S.P., N.N.D. 1962, A.D.R.; Unbuffered Tab., U.S.P.

Inhalation, N.F.

Abbott—Pow., Cartridge 100M u.

Squibb—Pow., Dispulator 100M u.

Winthrop Labs.—Micronized, Cartridge 100M u.

Ointment, Topically. See: Ades-Illin. (Polaris)

Parenteral.

Bristol—Vial 100M, 200M, 500M, 1000M, 2000M, 5000M u.

Dwight—Vial 100M, 200M, 500M, 1000M u.

Merck—Vial (100M, 200M, 500M u.) 5 cc; (1000M u.) 20 cc.

Pfizer—Vial 100M, 200M, 500M, 1000M u.

Promo—Vial (100M, 200M, 500M, 1000M u.) 20 cc.

Upjohn—Vial 100M, 200M, 1000M u.

See: Peni-Crystin (Merrell).

Parenteral, buffered.

Bio-Roma—Vial 200M, 500M, 1000M u.

*PENICILLIN HYDRABAMINE PHENOXYMETHYL, N.N.D. 1962

PENICILLIN FOR INHALATION, N.F., N.N.D. 1962.

See: Penicillin Aerosol. (Abbott)

Penicillin Aerosol. (Promo)

Penicillin Dispulator. (Squibb)

Penicillin G, Potassium.

Penicillin O, Procaine.

Penicillin O, Sodium.

Penicillin Inhaler. (Upjohn)

Penlator. (Lederle)

*PENICILLIN O, CHLOROPROCAINE, N.N.D. 1962, A.D.R. See: Depo-Cer-O-Cillin Chloroprocaine (Wyeth)

PENICILLIN O, POTASSIUM, N.N.D. 1959.

*PENICILLIN O, SODIUM, N.N.D. 1962.

Allylmercaptomethyl penicillin See: Cer-O-Cillin (Upjohn)

W/Chloroprocaine penicillin O, dihydrostreptomycin sulfate, streptomycin sulfate. See Cer-O Streps, Vial (Upjohn)

PENICILLIN OINTMENT, N.F.

See: Penicillin, Calcium.

Penicillin, Potassium.

Penicillin G, Sodium.

Penicillin, phenoxyethyl. See: Penicillin Y.

*PENICILLIN PHENOXYMETHYL, N.N.D. 1962 Penicillin Y.

See: Compoicillin Y (Abbott)

Pen-Yee (Wyeth)

Y-Cillin (Lilly)

* Generic name recognized by N.F., U.S.P. or A.M.A.

| Procaine Penicillin G | | Others | | Manufacturer | | Brand | |
|-----------------------|--|--------|--|----------------------|--|-------|-------------------------------------|
| 300M u | Benzath ne pen G 300M u | | | Pfizer | | | Permapen Fortified In Aq Susp /cart |
| 300M u | Benzath ne pen G 600M u, pot pen G 300M u | | | Wyeth | | | Bicilin All Purpose Vial/2 cc. |
| 300M u | Benzathine pen G 600M u, pot pen G 300M u | | | Bristol | | | Panbioatic, Vial/dose |
| 300M u | Benzath ne pen G 200M u, pot pen G 100M u | | | Pfizer | | | Pronapen Plus, Vial/cc. |
| | Benzath ne pen G 600M u, pot pen G 300M u, dihydrosireptomycin 250 mg, streptomycin 250 mg | | | Wyeth | | | B cell mycin All Purpose, Vial |
| 100M u | Br stam n d hyd cit 25 mg, A.P.C. | | | Bristol | | | Bristapen, Tab |
| 150M u | Chlor Tr metan maleate 2 mg, A.P.C. | | | Schering | | | Corlicid n W/Proc. pen, Tab. |
| 400M u | Streptomycin 0.5 Gm as sulf | | | Lilly | | | Durycid A S, Vial/cc. |
| 250M u | Meth D a Mer Sulfas | | | Smith Kline & French | | | Estocillin 250 Sulfas/5 cc. |
| 100M u | Phenyltolaxam ne dihyd cit 25 mg, A.P.C. | | | Wyeth | | | A P Cillin, Tab. |
| 100M u | Phenyltolaxamine d hyd cit 25 mg, A.P.C. | | | Bristol | | | Bristapen, Tab |
| 200M u | Phenyltolaxam ne d hyd cit 25 mg, A.P.C. | | | Bristol | | | Bristapen 200, Tab |
| 100M u | Phenyltolaxamine dihyd cit 25 mg, A.P.C. | | | Wyeth | | | A P Cillin, Tab. |
| 600M u | Pat pen G 200M u | | | Abbott | | | Abboxillin 600M, Vial/cc. |
| 300M u | Pat pen G 300M u | | | Monsieg II | | | Aquapropen, Vial/cc. |
| 300M u | Pat pen G 100M u | | | Comm Solvents | | | Bi Pen, Vial/cc. |
| 300M u | Pat pen G 100M u | | | Lilly | | | Durac II n Fort Red Vial |
| 300M u | Pat pen G 100M u | | | Squibb | | | Cryst for 400, Vial/cc. |
| 600M u | Pat pen G 100M u | | | Squibb | | | Cryst for 800, Vial/cc. |
| 900M u | Pat pen G 200M u | | | Squibb | | | Cryst for 1200, Vial/cc. |
| 300M u | Pat pen G 100M u | | | Upjohn | | | Dijurnal Fort Red 400M, Vial/cc. |
| 300M u | Pat pen G 100M u | | | Smith-Dorsey | | | Donsall n 400 Vial/cc. |
| 300M u | Pat pen G 100M u | | | Elder | | | Eldi Pen Vial/cc. |
| 300M u | Pat pen G 100M u | | | Bristol | | | Flo-Cillin "98" Fort Red Vial/cc. |
| 300M u | Pat pen G 100M u | | | Premo | | | Forticillin, V al/cc. |
| 300M u | Pat pen G 100M u | | | Roche | | | Hamoxillin Vial/cc. |
| 300M u | Pat pen G 100M u | | | Lederle | | | Lederril in Parenteral, Vial/cc. |
| 300M u | Pat pen G 100M u | | | Standard Pharm | | | Magnar II n Vial/cc. |
| 300M u | Pat pen G 100M u | | | Mayer | | | Myropen Vial/cc. |
| 300M u | Pat pen G 100M u | | | Merrell | | | Parent illin Fort Red, Vial/cc. |
| 300M u | Pat pen G 100M u | | | Bristol | | | Pen Aqua Fortified Vial/cc. |
| 300M u | Pat pen G 100M u | | | March Sharp & Dohme | | | Sharcilin Fort Red, Vial/cc. |
| 300M u | Pat pen G 100M u | | | Wyeth | | | Wycillin Fortified Vial/cc. |
| 400M u | Streptomycin base 0.5 Gm | | | Squibb | | | Strept. Districillin A S |

Penicillin G Procaine Combinations

| Procaine Penicillin G | Others | Manufacturer | Brand |
|-----------------------|--|--------------|--------------------------------|
| 300M u. | Sod. pen. G 100M u. | Pfizer | Pranapen, Vial/cc. |
| 100M u. | Sod. pen. G 100M u. | Parke, Davis | Penicillin S-R, Vial/cc. |
| 300M u. | Sod. pen. G 100M u. | Pfizer | Pronapen, Vial/cc. |
| 300M u. | Sod. pen. G 100M u., streptomycin sulf. 0.5 Gm. | Pfizer | Strep. Combiotic aq. Vial/cc. |
| 300M u. | Sod. pen. G 100M u., streptomycin sulf. 1 Gm. | Pfizer | Strep. Combiotic P S, Vial/cc. |
| 300M u. | Sod. pen. G 100M u., streptomycin sulf. 0.5 Gm. | Lilly | Durycin F.A., Vial/cc. |
| 300M u. | Sod. pen. G 100M u., streptomycin sulfate 10 Gm. | Parke, Davis | Penicillin S-R-S, Vial/cc. |

*PENICILLIN POTASSIUM PHENOXYMETHYL, N.N.D. 1962. Potassium Penicillin V.

PENICILLIN, POTASSIUM, U.S.P., A.D.R. 1954.

Winthrop Labs.—Oint. 1000 u./Gm. Tube 1 oz.; Ophth. aint. 1000 u./Gm. Tube ½ oz. See: Gelu-Clillin, Tab. & Trache (Warner).

*PENICILLIN G, SODIUM, U.S.P., A.D.R. W/Sulfacetamide. See: P-S Dental Cones. (Novaco)

PENICILLIN PRODUROL. (Merck Sharp & Dohme) Procaine penicillin G and Buf. Cryst., Penicillin G for Susp. Sterile, N.F.

PENICILLIN S-R. (Parke, Davis) Procaine penicillin G & Sod. penicillin G. Vial, 400M u., 1 dose; 400M u., 1 dose w/diluent; 2000M u., 5 dose; 4000M u., 10 dose; 1000M u., 1 dose; 10,000M u., 10 dose.

*PENICILLIN STREPTOMYCIN FOR SUSP. STERILE, N.F.

See: Durycin F.A. (Lilly)
Strep-Combiotic (Pfizer)
Strep-Dicryllicin (Squibb)

PENICILLIN TROCHES, N.P.

See: Penicillin, Calcium,
Penicillin G, Potassium,
Penicillin G, Sodium.

PENICILLIN, PHENOXYETHYL. See: Phenethicillin Penicillin sol. 152.

*PENICILLIN, PHENOXYMETHYL, POTASSIUM, U.S.P., N.N.D. 1962. Penicillin V, Pot. Tab. U.S.P.

PENICILLIN-STREPTOMYCIN, READMIXED, STERILE AQUEOUS SUSPENSION. (Upjohn) Procaine Penicillin G 400,000 u., Streptomycin sulfate 0.5 Gm., Procaine HCl 2%/2 cc.

PENICILLIN V, N.N.D. 1962. Phenoxymethyl penicillin. A biosynthetic penicillin formed by fermentation, with suitable precursors, of Penicillin notatum

See: Compoillin V. (Abbott)
Pen-Vee-Oral, Preps. (Wyeth)
V-Clillin, Preps. (Lilly)

W/Benzathine Penicillin G. See: Bicillin-Vee, Tab. (Wyeth)

W/Dia-Mer-Sulfonamides. See: Pen-Vee Sulfos, Tab. (Wyeth)

W/Meth-Dia-Mer Sulfonamides. See: V-Chlin-Sulfa, Tab. (Lilly)

W/Salicylamide, promethazine HCl, phenacetin, mephentermine sulf. See: Pen-Vee-Cidin, Cap. (Wyeth)

PENICILLIN V HYDRABAMINE, N.N.D. 1962. Hydrabamine phenoxymethyl penicillin. See: Compoillin (Parke Davis)

* Generic name recognized by N.P., U.S.P. or A.M.A.

PENICILLIN V POT., NND 1961

Phenoxyethyl penicillin Pot
See Compicillin VK, Tob (Abbott)
Pen Vee K (Wyeth)
V Cillin K, Tob (Lilly)
W/Pyrrolbutamine compound, aspirin, acetophenelinin, caffeine See V Kar, Tob (Lilly)
W/Sulta meth dio mer See V Cillin K Sulta, Tob (Lilly)
W/Sulladiazine, sulfamerazine, sullamethazine See Compicillin V K Sulfos, Film Tob (Abbott)

***PENICILLINASE, NND 1962** An enzyme that hydrolyzes penicillin, obtained from several strains of bacteria (*B cereus*) See Neutropen, Inj (Schenley Labs)

PENICOMBISUL TABLETS, (Schering) Pot. penicillin G 100,000 u with 0.166 Gm each of sulfacetamide, sulfadiazine & sulfamerazine/Tob Bot 24s, 100s
Use: Penicillin & sulfonamide susceptible infections

PENICRYSTIN, (Merrell) Sod penicillin G Vial 1 million u
Use: IM; penicillin therapy

PENIDURAL, See Benzathine Penicillin G

PENIMYCIN, (Liberty Vitamins Corp) Procaine penicillin G 300M u, Pot penicillin G 100M u, dihydrostreptomycin sulfate 0.5 or 1.0 Gm /dose Vial 1 dose
Use: Penicillin & streptomycin therapy

PENIORAL (Wyeth) Pot penicillin G w/col carb 50,000 100,000 u /tab Bot 12s
Use: Orally; penicillin therapy

PENISEM (Mossingill) Pot penicillin G 100M u /5 cc Bot 2 oz
Use: Orally; penicillin therapy

PENITE (Carnrick) Nitroglycerin 0.9 mg, pentaerythritol tetranitrate 30 mg reserpine 0.12 mg /Cap Bot 50s
Use: Coronary vasodilation

PENITRACIN, (Penick) Bacitracin, bulk only
Use: Manufacturing Purposes

PEN-M (Pfizer) Oleandomycin salt of penicillin Under study

PENMYCIN-A S, (Philadelphia Ampoule) Procaine penicillin G 400,000 u, dihydrostreptomycin 0.5 Gm /2 cc Vial 10 cc
Use: Antibiotic

PENOTRANE, (Mead Johnson & Co) See Hydrargaphen

PEN-PHETAMINE (Clark & Clark) Profetamine phosphate 3 mg phenobarbital 1/4 gr, aspirin 5 gr /Tab Bot 100s, 250s, 1000s
Use: Orally (1 tid) analgesic & relaxant

PEN-TABS W/TRIPLE SULFONAMIDES, (Rexall) Pot penicillin G 150M u, sulfadiazine sulfathiazole & sulfamethoxazole of each 167 mg /Tab Bot 100s
eneric name recognized by NF USP or AMA

Uses: Orally; penicillin w/sulfonamide therapy

***PENTABAMATE, (Eiker) 3 methyl 2,4 pentacetal dicarbomate**
Use: Tranquilizer

PENTABROMIDES, (Merrell) Sod bromide 476 Gm, ammonium bromide 130 Gm, Pot bromide 130 Gm, Cal bromide 0.78 Gm, lithium bromide 0.25 Gm /30 cc Bot 1 pt, 1 gal
Use: Sedative

PENTACHLOROPHENATE SODIUM, See Podoson, Pow (Doak)

PENTACINE, (Walker Corp) Pentobarbital 1/2 gr, scopolamine HBr 1/400 gr /Tab Bot 100s
Use: Sedative

PENTACOLS, (Endo) Guaiaccol carb 1 gr, terpin hydrate 2 gr, hyoscyamus est 1/10 gr, phenobarbital 1/2 gr tricalcium phosphate 3 gr /Tab Bot 100s, 1000s
Use: (1 tsp tid, & hs) Sedative & expectorant

PENTACORT, (Dalin) Hydrocortisone alcohol 1/2% coal tar soln 2%, iodochlorhydray quinoline 3% dipiperon HCl 1/4% Vit A Vit D in Dalicarin base Tube 5 Gm 15 Gm

Use: Treatment of dermatologic disorders
PENTAERYTHRITOL CHLORAL, NND 1962
See: Periclor, Cap (Ives Cameron)

***PENTAERYTHRITOL TETRANITRATE, NND 1962**
See: Angicap, Timed Cap (Pro Actel)

Angitel, Tob (H G Horvey)
Circulin (Fleming)
Duotrate, Cap (Marion)

Lowtrate and Lowtrate CTR (Lowe)
Martrate-45 Caps (Marsh Emory Labs)
Metranil, Caps (Meyer Labs)

Myotrate 10, Tabs (Sig Inc)
Neo Corvas, Cap (Grant)
Nitratals Tab (Paul Maney)

Pescard, Tab (Cale)
Pentestee 80, Cap (Standex)
Pentelrate Unicelles (Hiss)

Pentrate Stronger Caps (Testagar)
Pentritol Tempules (Evron)
Pentryate, Cap (Testagar)

Pentryl, Cap (Bernhall)
Pentrate, Tab (Warner Chilcott)
PET Tob (Barre)

Rylthral Tab (Melra Lab)
Tanipent, Tab (Superior)
Tranite D Lay Caps (Lloyd Dabney Westerfield)

Vasodiatol, Tab (Rawell)
Vasa 80 Unicelles TD Caps (Hiss)

- W/Alseroxylan. See: Nitroax, Tab. (Smith-Dorsey)
- W/Aminaphylline. See: Amitrate, Tab (Kremers-Urban)
- W/Aminaphylline, aluminum hydrazide. See Peritrate w/Aminaphylline, Tab. (Warner-Chilcott)
- W/Amobarbital. See: Metranil-AM Deracap, Caps. (Meyer)
- W/Barbital sodium. See P.E.T.-R., Caps (Barre Drug)
- W/Butobarbital.
See: Pentrital-B, Tempules. (Evrae)
Caralel-T.D., Caps. (Crystal)
- W/Butobarbital Sod. See: Perbuzem, Tab. (Zommer)
- W/Butisal sod., reserpine. See: Pentrasine, Tab. (McNeil)
- W/Carbromal. See: Perosed, Tab. (Mollard)
- W/Dihydroxypropyl theophylline. See: Niphylline, Tab. (Paul Money)
- W/Ethaverine HCl
See: Papavatal, Tab. (Kenwood Labs)
Peritol, Tab (Ulmer)
- W/Ethylpapaverine HCl. See Papavatal L. A., Tab. (Kenwood)
- W/Hydrochlorothiazide. See: Perithiazide SA, Tabs. (Warner-Chilcott)
- W/Hydroxyzine See: Cartrax, Tab. (Roerig)
- W/Mephobarbital, phenobarbital. See: Rith-A-Tab, Tab. (Walker Corp.)
- W/Meprabamate.
See: Equanilate, Tab. (Wyeth)
Miltate, Tab. (Wallace Labs)
- W/Neathylline. See: Niphylline, Tab. (Money)
- W/Nitroglycerin.
See: Nitrate, Tabs (Arco)
Dilcaron, Tab. (Winthrop Lab)
Nitral, Tab. (Kremers-Urban)
Peritrate W/Nitroglycerin, Tab (Warner-Chilcott)
Vasolate, Caps. No. 1 (Miss)
Bulo-Penite, Tabs, Caps., (Cornick)
- W/Nitroglycerin, reserpine. See: Penite, Tab (Cornick)
- W/Nitroglycerin, secobarbital sod.
See: Anti-Anginal, Tabs. (Lemman)
Nitrosted, Tabs (Lemman)
- W/Nitroglycerin, Vit. B-1. See: Pentafort, Cap. (Sutliff & Case)
- W/Phenobarbital. See: Martrate-45 with Phenobarbital, Caps (Marsh-Emary Labs)
See: Nitratolans w/Phenobarbital, Tab (Paul Money)
Peribar, Tab. (Whittier)
Peritrate w/Phenobarbital S. A. (Warner-Chilcott)
Tanipent, Tab (Superior Pharm)
- W/Phenobarbital, ethylpapaverine HCl. See: Papavatal L.A. w/Phenobarbital, Caps (Keewood)
- W/Reserpine
See: Reserpatrate, Tab. (Vitamax)
Respet, Tab (Lloyd, Dabney & Westerfield)
- W/Reserpine, Butisal sod.
See: Pentaserpine, Tab. (Nysco)
Pentrasine, Tab. (McNeil)
- W/Reserpine, rutin, ascorbic acid. See Ruserpatol, Tab. (Lemman)
- W/Secobarbital. See: Caravas, Tymcaps. (Gross Chemical)
- PENTAFIN**, N.N.D. 1962. (Tutag)
- PENTAFORT**. (Sutliff & Case) Nitroglycerin 1/150 gr., pentaerythritol tetranitrate 15 mg., Vit. B-1 5 mg./Cap. Bot. 100s, 1000s
Use: Coronary vasodilator
- PENTAGISIC**. (Kremers-Urban) Acetylsalicylic acid 300 mg; salicylamide 100 mg; salicylsalicylic acid 100 mg; guaiacol glyceryl ether 200 mg; phenyltolazamine citrate 12.5 mg/Tab
Use: Analgesic/relaxant.
- PENTAL**. (Vanpell & Brown) pentobarbital Sod. Cap. 3/4 gr., 1 1/2 gr. Bot. 100s, 500s, 1000s.
Use: (1-3 gr. daily) Sedative A hypnotic.
- PENTAM RECTAL SUPP.** (Alcon) Aminaphylline 3 3/4 gr., benzocaine 1/2 gr./Supp. Box 12s
Use: Anesthesia
- PENTAMETHYLENETETRAZOL**, U.S.P., N.N.R. 1950.
See: Pentylenetetrazol U.S.P. (Various Mfr.)
- * **PENTAMOXANE HCl**. (Eli Lilly) 2-Isosmethylaminomethyl-1,4-benzodioxane HCl.
Use: Tranquilizing agent.
- PENTAPHEDRINE**. (Fint, Eaton) Pentobarbital Sod. 1/2 gr., ephedrine HCl 1/2 gr., Cal iodide 2 gr./fl. oz. Bot. 1 pt., 1 gal.
Use: Sedative antispasmodic.
- * **PENTAPIPERIDE METHYLSULFATE** (formerly Valpamate Methylsulfate), 4'-(1'-methylpiperidyl) 2-phenyl-3-methylvalerate dimethylsulfate. See: Quilene (Ayerst Labs.)
- PENTAPHONATE**. See: Claraphos (Bischoff)
- * **PENTAPIPERIDE METHYLSULFATE**. (Ayerst) 4'-(1'-methylpiperidyl) 2-phenyl-3-methylvalerate dimethylsulfate. See: Quilene.
- PENTAPLEX**. See: Eskay's Pentaplex, Liq. (Smith, Kiene & French).
- PENTAQUINE**, U.S.P., N.N.R. 1952. (Abbott) Pentaquine phosphate, 6-Methoxy-8-(5-isopropylaminoamylamino) quinaline, Tab. 13.3 mg Bot. 50s, 1000s
Use: Antimalarial.

* Generic name recognized by N.F., U.S.P. or A.M.A.

PENTASERPINE. (Nysco) Reserpine 01 mg, pentoerythritol tetranitrate 10 mg /Tab Bot. 100s, 1000s

User Angina pectoris

PENTASPAN. (Straits Pharmacal) Caps T D 30 mg, 100s

PENTASYN. (G F Harvey) Thiamine NCI 1000 mg, riboflavin 10 mg, pyridoxine HCl 20 mg, Sod pantothenate 100 mg, nicotinamide 500 mg /Vial Vial 10 cc

User Vit B complex therapy

PENTAVALENT GAS GANGRENE. Antitoxin, NNR 1952

PENTAZETS. (Merck Sharp & Dohme) Homarylamine HCl (N Methyl 3,4 methyl enedioxyphenethylamide HCl) 20 mg, zinc bacitracin 50 u, tyrothricin 1 mg, neomycin sulf 5 mg, benzocaine 5 mg /Troche Box 12s

User Symptomatic relief of sore throat

PENTAZYME (Ulmer) Pancreatic enzyme concentrate to supply amylase lipase & protease 100 mg /Tab Bot 50s

User Treatment of indigestion

PENTERGOT. (Benlex Pharm) Ergolamine tartrate 2 mg, caffeine 100 mg, hyoscyamine sulfate 0.25 mg, pentobarbital 60 mg /insert Box 12s

User: Rectal; analgesic & relief of tension headaches

PENTESTAN-80 (Stondex) Pentoerythritol tetranitrate 80 mg /Cop Bot 100s

User Angina

PENTETRATE UNICELLES. (Hiss) Pentoerythritol tetranitrate 30 mg /cop Bot 30s, 100s
No 1 W/Nitroglycerine 4/500 gr
No 2 W/Nitroglycerine 4/500 gr, amobarbital 50 mg

User Angina pectoris

***PENTHENATE BROMIDE, N F, NND 1962,** Tabs N F 2 diethylaminoethyl 2 cyclo pentyl 2 (2 threnyl) hydroxyacetate metha bromide See: Monardal Bromide (Winthrop Labs)

W/Mephobarbital Seer Monardal Meboral, Tab (Winthrop)

PENTHRANE. (Abbott) Methoxyflurane, 1,1 dichloro 2,2 difluoro 2 methoxyethane

User: Anesthetic for surgery

W/D 01% butylated hydroxytoluene as an antioxidant Bot 125 cc

PENTIOS, NND 1962 ADR (Squibb) Pot penicillin G 200M u buffered w/Cal carb or sol w/Sod citrate/Tab Pkg 12s 100s Cop (unbuffered) 200M u Bot 24s, 100s Syr Dissolve contents of Pkg W/35 cc of water Solutron provides 200M u /5 cc Bot

* Generic name recognized by N F, U S P, or A M A

2 oz Tab. "400" Bot 12s, 50s, 100s Syr "400," Bot 60 cc

User Orally; penicillin therapy

PENTID-SULPAS. (Squibb) Pot penicillin G 200000 u w/sulfamethazine, sulfadiazine & sulfamerazine 0.167 Gm. of each/Tab Bot 30s, 100s, 500s or 5 cc Syr. in 2 dose Bot

User: Orally; penicillin therapy w/triple sulfas

PENTIOS-SULPA for SYRUP. (Squibb) Penicillin G potassium (Pentids) 200,000 u, triple sulfanamides 0.5 Gm, buffered with sodium phosphate/5 cc Bot 60 cc

User Penicillin with sulfanamide therapy

PENTOBARBITAL, Elixir, N F X NND 1962 ADR (Various Mfr) 5 Ethyl 5-(1 methylbutyl) barbituric acid Seer Nembutal, Elix (Abbott)

Pentobarbital Combinations on pp. 557 to 559.

***PENTOBARBITAL CALCIUM, NND 1962,** ADR Seer Nembutal Calcium (Abbott)

W/Aminophylline See Aminophylline W/Pentobarbital Sod (Various Mfr)

W/Chlor Trimefon Seer Chlor Trimefon W/Sod Pentobarbital, Tab (Schering)

W/Deserpidine Seer Harmony-N, Tab (Abbott)

***PENTOBARBITAL SODIUM, USP NNR 1952 ADR,** Cop & Elixir USP; (nj), USP; Sterile; Tab N P (Embutal) Sod 5 Ethyl 5 (1 methylbutyl) borbiturate See Isobarb, Tab (US Standard)

Napental, Prep (Massengill)

Nembutal, Prep (Abbott)

Pembule Cop (Navco)

Pental Cop (Vanpell & Brown)

PENTOBARBITAL, SOLUBLE See Pentobarbital Sodium USP

PENTO-DEL. (Boyle) Pentobarbital Sod 1 1/2 gr, pentobarbital 7 gr/Cop Bot 100s 1000s

User Sedative

***PENTOLINIUM TARTRATE, NND 1962** Pentamethylene-1,5 bis(1-methylpyrrolidinum bitartrate) See: Ansalyzen, Tab, Vial (Wyeth)

W/Meprobamate See: Equolyzen, Tab (Wyeth)

PENTO-PHENO-BABB. (G F Harvey) Pento barbital sod 1/2 gr, phenobarbital 1/4 gr /Tab Bot 100s, 1000s

User Sedative and hypnotic

PENTOBLOCK. (Crystal) Pentylene-tetroxol 100 mg, nicotinic acid 50 mg, dihydroxypropyl theophylline 100 mg /Tab Bot 100s 1000s

User: Treatment of mental confusion, anorexia, fatigue

Pentobarbital Combinations

Pentobarbital or
Sod. Salt

Others

| Pentobarbital or Sod. Salt | Manufacturers | Brand |
|-------------------------------|---|---|
| 0.1 Gm. | Aminophylline 0.5 Gm., benzocaine 0.06 Gm. | |
| 0.05 Gm. | Aminophyllin 0.5 Gm., benzocaine 0.1 Gm., phenobarbital Sod. 0.05 Gm. | Bischoff Fellows |
| 20 mg. | d Amphetamine sulf., 5 mg. | |
| 1½ gr. | Amphetamine sulfate 5 mg., phenobarbital ½ gr. | Metan Pharm. |
| ½ gr. | d-Amphetamine sulf., 2.5 mg., ectophenentidin 2½ gr., espirin 3½ gr. | Columbus Arner-Stone |
| ¾ to 1½ gr. | Aspirin 5 gr. | |
| ½ gr. | Aspirin 5 gr. | Novocel |
| ¼ gr. | Aspirin 5 gr. | Massengill |
| ¼ gr. | Aspirin 5 gr. | Elder |
| 15 mg. | Aspirin 5 gr. | Parklins Abbott |
| 18-125 mg./isp. | Aspirin 3½ gr., phenacetin 2½ gr., caffeine ½ gr. | |
| 3/16 gr. | Atropine sulfate 0.02425 mg./isp., hyosine HBr 0.00472 mg./isp., hyoscyamine sulfate 0.1296 mg./isp., Vit. B-1 12.5 mg./isp. | First Toses |
| 60 mg. | Belladonna ext. ½ gr., phenobarbital 3/16 gr. | Kirkman |
| 15 mg. | Benzocaine 100 mg., vit. B-6 25 mg., nicotinamide 25 mg., me- thionine 100 mg. | Sandoz Elder |
| 1½ gr. | Butalbarbital ½ gr. | |
| 60 mg. | Butalbarbital Sod., 30 mg., d-desoxyephedrine HCl 5 mg. | |
| ½ gr. | Butalbarbital 8 mg., phenobarbital 8 mg. | |
| 1/8 gr. | Butalbarbital sod., 1/6 gr., phenobarbital ½ gr./3 cc. | |
| 6.075 mg. | Butalbarbital sod., 2.025 mg., mncobarbital sod., 6.075 mg., pheno- barbital 2.03 mg., belladonna pow., ext., 4.1 mg., phenyl- tetrazamine 20 mg., atthaverine HCl 12.5 mg. | Atcher Central Durst Barze Armour Labs. |
| 15 mg. | Dehydrocholic acid 200 mg. & methamphetamine HCl 5 mg. | |
| 30 mg. | Deserpidine D 25 mg. | |
| 30 mg. | d-Desoxyephedrine HCl 5 mg. | Melitin Abbott |
| ½ gr. | d-Desoxyephedrine HCl 2.5 mg., homatropine methylbromide 1/40 gr., methylcellulose 5 gr. | Abbott Central |
| ½ gr. | d Desoxyephedrine HCl 5 mg. & mephobarbital ½ gr. | |
| 30 mg. | Dianastrol D 5 mg., methylflinstosterone 2.5 mg., d-amphetamine sulf., 5 mg. | Walker Kremerr-Urbau |
| | Amphetamine without Cocaine, Tab. | |
| | Hydral. Liq. | |
| | Duodon, Tab. | |
| | Cofargel PB, Supp. | |
| | Naudal, Tab. | |
| | Twln-Borb Improved Tab. | |
| | Somnodax, Tab. | |
| | Quines, Tab. | |
| | Pyramid, Liq. | |
| | Tri-Syneer Plus, Tab. | |
| | Sifron, Tab. | |
| | Harmony-N. FilmTab | |
| | Darbutoj, Cop. | |
| | Endascal, Tab. | |
| | Mepha-d, Tab. | |
| | Amperon, Tab. | |

| Pentobarbital or Sod Salt | Others | Manufacturer | Brand |
|---------------------------|---|-----------------------|--------------------------------|
| 7.5 mg | 2 Diethylaminoethyl diphenylacetate HCl 250 mg activated 90 mg, aluminum hydroxide 75 mg | Palmedica | Spasmasarb Tab |
| 48.6 mg | Dihydrocadesonone bitartrate 5 mg & phenobarbital 16.2 mg | Brean | Bredative DHC, Tab |
| 1/2 gr /oz | Ephedrine HCl 1/2 gr./oz. Cal iod 8 gr./oz | Flint, Eaton | Pentaphedrine, Liq |
| 1/2 gr | Hexestrol 1 mg, scopolamine HBr 1/1200 gr | Walker Carp | Tal Estamine Tab |
| 8 mg | Homatropine methylbromide 2.5 mg, dehydrocholic acid 60 mg ox bile ext 150 mg | Lemman | Homachol, Tab |
| 1/4 gr | Homatropine methylbromide 1/25 gr, Mag trisilicate 5 gr | Lemman | Homapent, Tab |
| 1/4 gr | Homatropine methylbromide 1.2 mg, Mag trisilicate 2 gr, gastric mucin 1 gr, aluminum hydroxide 4 gr | Lemman | Antosed Tab |
| 7.5 mg | Homatropine methylbromide 2.5 mg, phenobarbital 7.5 mg | Meyer | Almesed Elixir |
| 8 mg | Homatropine methylbromide 2.5 mg, phenobarbital 8 mg | Rarer | Precibus Cap |
| 1/16 gr | Homatropine methylbromide 1/480 gr, phenobarbital 1/8 gr | Walker Carp | Sedadreps, Liq |
| 1/16 gr | Homatropine methylbromide 1/480 gr, phenobarbital 1/8 gr | Walker Carp | Sedaplex |
| 1/16 gr | Homatropine methylbromide 1/16 gr, phenobarbital 1/6 gr | Harrower | Selhyli W/Barb, Tab |
| 1/16 gr | Homatropine methylbromide 1/480 gr, phenobarbital 1/6 gr | Jackson Mitchell | Saised Pediatric Liq. |
| 48.6 mg | Hyoscyamine sulf 0.311 mg, atropine sulf 0.582 mg, hyoscyne HBr 0.0195 mg | Calymbus | Belakoids TT, Tab |
| 1/4, 1/2 or 1/2 gr | Mephnesin 0.5 Gm | Invenex | Talaphen Pentobarb., Tab |
| 90 mg | Mephnesin 425 mg | Smith Darsey | Nebralin, Tab |
| 30 mg | Methacopalamine bromide 5 mg | Marrin | Nutrop W/Pentobarbital, Cap |
| 8 mg | Methacopalamine nitrate 2 mg, pepsin 150 mg, glutamic acid HCl 200 mg, cellulose 9.14 mg, ox bile ext 100 mg, pancreatin 500 mg | Darsey | Kanumadic, Tab |
| 100 mg | Niacinamide | | |
| 1 gr | Pentobarbital 1/2 gr | Mason Labs | Niapent, Cap |
| 16 mg | Pepsin 32 mg, charcoal, activated 200 mg | Valentine | Valtwabarb, Tab, |
| 3/4 gr. | Phenobarbital 3/4 gr | Hoack Labs | Carget, Tab |
| 3/4 gr | Phenobarbital 1/4 gr | Flint, Eaton | Barfurate, Cap |
| 1 1/2 gr. | Phenobarbital 1/2 gr | Brean | Bredative, Tab |
| 1 1/2 gr. | Phenobarbital 1 gr | Superior | Duenite, Empule |
| 1 1/2 gr. | Phenobarbital 1/2 gr | Boyle | Penta Del, Cap |
| 1 gr | Phenobarbital sod 1 gr | Ascher | Twin Barb, Tab |
| | | Smith, Muller & Patch | Mill A Sed, Cap |

Pentobarbital Combinations

| Pentobarbital or Sed. Salt | Others | Manufacturer | Brand |
|----------------------------|--|------------------------|----------------------|
| 60 mg. | Phenobarbital 30 mg., d-amphetamine salt, 5 mg. | Mysco Labs. | Sedes. Tab. T.D. |
| ¼ gr. | Phenobarbital Sod., ½ gr., Cal. lactate 5 gr. | Rowell | Vio-Lactal, Cop. |
| 1/6 gr. | Phenobarbital 1/6 gr., epoxyminoxypropine Iropate HBr 0.25 mg., aminoxypyropine Iropate HCl 0.25 mg., procaine HCl 150 mg., Vi. B-6 10 mg., aluminum hydroxide gel dried 250 mg. | Harrower | Prametic, Tab. |
| 15 mg. | Phenobarbital Sod. 15 mg., barbital Sod. 15 mg. | Smith, Miller & Polich | Quadra-Sed. Sol. |
| 30 mg. | Phenobarbital 30 mg., secobarbital Sod. 30 mg. | Taylor | Barbclin, Tab. |
| 100 mg. | Pentylenetetrazol 300 mg. | Strong Cobb | Tetrapentobarb. Cop. |
| 27.5 mg. | Racephadrine HCl, 20 mg., choline theophyllinate 200 mg. | Napera | Cholorace, Tab. |
| ¼ gr. | Sallylamide 3/4 gr., phenacetin 2 ½ gr., caffeine ¼ gr. | Smith-Dorsey | Angeral, Tab. |
| ¼ gr. | Scopolamine HBr 1/1200 gr., hexestrol 1 mg. | Walker Corp. | Tab-Enamine, Tab. |
| ¼ gr. | Scopolamine HBr 1/400 gr. | Walker Corp. | Pentacine, Tab. |

PENTOTHAL SODIUM, N.N.D. 1959, A.D.R. (Sterile Thiopental Sod., U.S.P.) (Abbott) 5-ethyl-5-(1-methylbutyl) thiobarbiturate. Amp. 0.5 Gm., 1s, 6s, 25s, also w/diluent, 1 Gm., 1s, 6s, 25s, also w/diluent, 5 Gm., 25s; 10 Gm., 25s. Vial 2 Gm., for rectal application.

Sterile Pow. Amp. 6.25 mg, 12.5 mg
Use: Ultra-short acting anesthetic.

PENTOTHAL SOD. Rectal Susp. Thiopental sod. 400 mg/Gm. rectal susp. with sod. carb. 24 mg. in water miscible mineral oil susp. Box 25s & 100s

Use: Rectal. Basal anesthesia.

PENTOXYLON TABLETS, (Biker) Alseroxylon fraction of Rauwolfia 0.5 mg., pentaerythritol tetranitrate 10 mg/Tab. Bot. 100s, 1000s

Use: Angina pectoris.

PENTRALINE, (McNeil) Formerly Pentrosine Pentaerythritol tetranitrate 10 mg, buliso sod. 10 mg., reserpine 0.05 mg/tab. Bot 100s, 1000s.

Use: Prevention of angina attacks.

PENTRASINE, (McNeil) Pentaerythritol tetranitrate 10 mg, buliso sod. 10 mg., reserpine 0.05 mg/tab. Bot. 100s, 1000s

Use: Prophylactic management of angina pectoris

PENTRAZOL, (Kirkman) Pentylenetetrazol 100 mg/tab. Bot. 100s, 1000s, 5000s.

Use: Analeptic.

PENTRESAMIDE-250, (Merck Sharp & Dohme) Not listed. Pot. penicillin G 250M u., sulfadiazine 0.2 Gm., sulfamerazine 0.1 Gm., sulfamethazine 0.2 Gm./5 cc. Bot. 60 cc.

Use: Orally; penicillin & triple sulfonamide therapy.

PENTRESAMIDE PEDIATRIC GRANULES, (Merck Sharp & Dohme) Sulfamerazine 0.05 Gm., sulfadiazine 0.1 Gm., sulfamethazine 0.1 Gm., pot. penicillin G 100,000 u./5 cc. Bot. 60 cc.

Use: Orally; penicillin & triple sulfonamide therapy.

PENTRESAMIDE TABLETS, (Merck Sharp & Dohme) Sulfamerazine 0.1 Gm., sulfadiazine 0.2 Gm., sulfamethazine 0.2 Gm., pot. penicillin G 100,000 u. & 250,000 u./tab. Bot. 60s, 250s.

Use: Penicillin w/triple sulfonamide.

PENTRISUL, (Cosmir Funk) Pot. penicillin G 100M u., sulfadiazine 0.165 Gm., sulfamerazine 0.165 Gm., sulfamethazine 0.165 Gm./Tab. Bot. 30s, 100s, 500s.

Use: Orally; penicillin w/triple sulfonamide therapy.

PENTRITOL-B TEMPULES (Armour) Pento-

- erythritol tetranitrate 30 mg, butobarbital 50 mg/cap Bot 60s 250s
 User 1 2 daily coronary vasodilator with sedative
- PENTRITOL TEMPULES, NND 1962**
 (Armour) Pentaerythritol tetranitrate Cap 30 mg, Bot 60s 250s
 Use Coronary vasodilator
- PENTRIZINE** (Triden) Pot penicillin G 100M u, sulfadiazine 0.17 Gm, sulfamerazine 0.17 Gm, sulfamethazine 0.17 Gm/Tab Bot 24s
 Use Orally; penicillin w/triple sulfonamide therapy
- PEN-TROCHES, ADR 1953** (Cutter) Pot penicillin G 5000 u/Trache Vial 25s; Bot 500s
 Use Orally; throat & mouth infections
- PENTRYATE, (Testogor)** Pentaerythritol tetranitrate 30 mg/Timed Cap Bot 100s, 1000s
 Also available with Phenobarbital 15 mg/Tab
 Use: Angina pectoris
- PENTRYATE STRONGER, (Testogor)** Pentaerythritol tetranitrate 80 mg/Cap (timed release) Bot 30s 100s, 1000s
 Use Angina pectoris
- PENTRYL** (Bernhaft) Pentaerythritol tetranitrate 45 mg/S R cap
 Use: Treatment of anginal attacks
- *PENYLENETETRAZOL, NF NNR 1950**
 ADR; Inj, NF Pentamethylenetetrazol, Lepazol
 See Cardiazol (Knoll)
 Metrazol (Brhuber Knoll)
 Pentrazol Tab (Kirkman)
 Petrolone, Inj (Vitamax)
- W/d-amphetamine sulfate, caffeine anhydrous See Tempolrid, Liq or Tabs, (Smith, Miller & Patch)
- W/d Amphetamine sulf, vit B complex Seer Dexazyme, Tab (Gray Pharm)
- W/Dihydrocodeinone bitartrate dihydrocodeinone terephthalate, homatrapine terephthalate See Para Mycadon (Endo)
- W/Dihydrohydroxycodone, dihydrohydroxycodone terephthalate & homatrapine terephthalate Seer Nucodon, Tab (Endo)
- W/Methyltertartrate, ethinyl estradiol, nicotinic acid, and dicalcium glutamate See Senatric, Tab (Dumar Wilson)
- W/Methyltestosterone, ethinyl estradiol, strontium salicylate See Analeptone Anabolic (Reed & Carnrick)
- W/Nicotinic acid
 See Analeptone, Elix (Reed & Carnrick)
 Arterline Tab or Elix (Confield)
 Benzal, Tab, Elix, (Bentex)
- * Generic name recognized by N.F., U.S.P. or A.M.A.
- Geronozol, Tab (Columbus Pharmacol)
 Menrc, Tab (Geriatric)
 Metalex, Liq (Storck)
 Nralift (B F Ascher & Co)
 Napent (Ferndale)
 Nico Met, Tab & Elix (Knoll)
 Nrcozal Caps Elixir, Tabs (Hort Laboratories)
 Semalex, Tab (Durst)
 Tosec, Tab (Hoack)
 Vasa tabs, Tab (Paul Money)
- W/Nicotinic acid, ascorbic acid, bioflavonoids See Niatric, Tab (Ascher)
- W/Nicotinic acid, ascorbic acid, l-glutamic acid, niacinamide, riboflavin, d pantothenyl alcohol, alcohol See Lrp Nrcin PTZ (Brown Pharmaceutical)
- W/Nicotinic Acid, dienestrol methyltestosterone See Lucidon, Tab (Lloyd, Dabney & Westerfield)
- W/Nicotinic acid methyl testosterone ethinyl estradiol thiamine, riboflavin pyridoxine HCl Vit B 12, folic acid panthoal choline bitartrate inositol, l lysine monohydrochloride Vit E, iron (as ferric pyrophosphate) magnesium manganese zinc, alcohol See Nicozal Complex (Hart Laboratories)
- W/Nicotinic acid dihydroxypropyltheaphyl Ine See Pentarack Tab (Crystal)
- W/Nicotinic acid, hesperidin complex, Vit C See Nivalex, Tab (Mallard)
- W/Nicotinic acid, methyltestosterone, ethinyl estradiol, thyroid, Vit C See Nicotricline Tab (Ascher)
- W/Nicotinic acid neothylline See Neo Pent, Tab (Paul Money)
- W/Nicotinic acid, tetrahydro tyranine, methyl testosterone, ethinyl estradiol See: Tetra Forte, Tab (E S Miller)
- W/Nicotinic acid, thiamine HCl See Zalger, Tab (Superior Pharm)
- W/Nicotinic acid, Vit B 1, B 2, B 6 calcium pantothenate, Vit C See Nialift Tabs (Ascher)
- W/Thiamine HCl niacinamide cyanocobalamin, alcohol See Cenalene Liq (Central)
- W/Pentobarbital Sod Seer Tetropentobarb Cap (Strang Cobb)
- W/Phenobarbital Sod See: Tetraphenobarb Cap (Strang Cobb)
- W/Reserpine, niacin See: Nrcozal with reserpine, Tab (Drug Spectralter)
- W/Secobarbital Sod See: Tetrasecobarb Cap (Strang Cobb).
- W/Thiamine HCl, niacinamide, Vit B 12 alcohol See: Centalex, Liq (Central)
- PIN VEE, NND 1962** (Wyeth) Seer Phenoxymethyl Penicillin

PEN-VEE-CIDIN. (Wyeth) Penicillin V 100,000 u., salicylamide 194 mg., promethazine HCl 6.25 mg., phenacetin 130 mg., mephentermine sulf. 3 mg./Cap. Bot. 36c
Use: Treatment of the common cold

PEN-VEE-K, N.N.D. 1962. (Wyeth) Penicillin V Potassium 125 mg. 250 mg. Bot. 36a.
Use: Penicillin susceptible organisms.

PEN-VEE K LIQUID. (Wyeth) Penicillin V potassium 125 mg. or 250 mg /5 cc. Vial 40 cc.
Use: Antibiotic.

PEN-VEE-L.A. (Wyeth) Phenoxymethyl penicillin 250 mg. (400,000 u.)/Tab. Vial 24.
Use: Penicillin therapy.

PEN-VEE-ORAL, N.N.D. 1958. (Wyeth) Penicillin-V, Phenoxymethyl penicillin 125 mg. (200,000 u.) Tab., Bot. 36s. Tab. 300 mg. (500,000 u.) Bot. 12s. Susp. (300,000 u./5 cc.) Bot. 8D cc. (150,000 u./5 cc.) Bot. 2 oz. Drops (150,000 u./cc.) Bot. 10 cc.
Use: (1-2 Tab. q.i.d.) penicillin therapy.

PEN-VEE PEDIATRIC. (Wyeth) Benzathine penicillin V issued as dry powder which, when reconstituted with water, contains in each 5 cc., 250 mg. (400,000 u.). Bot. 2 fl. oz.
Use: Penicillin therapy.

PEN-VEE BULFAS, N.N.B. 1957. (Wyeth) Penicillin V 150,000 u., sulfadiazine 250 mg., sulfamerazine 250 mg./Tab. or 5 cc. Susp. Bot. 2 oz. Bot. 36s.
Use: Sulfonamide therapy w/penicillin.

PEPCATE. (Sutliff & Case) Pepsin 32.5 mg., pepsin 32.5 mg., sodium bicarbonate 160 mg., charcoal 65 mg., aromatics/tab. Bot 100s
Use: Digestant, adsorbant, antacid.

PEPSENCIA. (Winthrop Labs.) Hog stomach linings & calf rennets. Liq., Bot. 8 oz., 1 pt.
Use: Orally (3-4 dr. p.c.): aid digestion.

* **PEPSIN, N.F.** Comp. Elixir, N.F. Pow. or Granule. See: lactated Pepsin (Merck Sharp & Dohme)
Use: Digestive aid.

W/Atropine sulf., hyoscyne HBr, hyoscyamine sulf., activated charcoal

See Beldase, Cap. (Finley)
Donnazyme (Robins)

W/Betaine HCl. See: Acidol-Pepsin, Cap. (Winthrop Labs.)

W/Betaine HCl & methylcellulose See Stuart Narmacid, Tab (Stuart)

W/Cellulose, glutamic acid, pancreatin, ox bile ex. See: Kanulase, Tab (Smith-Dorsey)

W/Dehydracholic acid, ox bile ext., pancreatin. See: Neabile, Tab. (Haug Drug Co)

W/Diastase, rennin, hydrochloric acid &

* Generic name recognized by N.F., U.S.P. or A.M.A.

lactic acid. See: Diastanol, Liq. (Smith-Dorsey).

W/Enzyspons, lurocholic acid. See: Enzymet, Tabs. (Lloyd, Dabney & Westfield)

W/Glutamic Acid. See: Glutasin, Cap. (McNeil)

W/Lactic acid, caffeine & celery seed. See: Pan-Peptic, Elixir. (Merck Sharp & Dohme)

W/Pancreatin. See: Entozyme (Robins)

W/Pancreatin, ox bile, dehydracholic acid. See: Ro-Bile, Tab. (Rowell)

Entozyme (Robins)

W/Pancreatin, bile ext. and papain. See: Plurizyme, Tab. (Harrower)

W/Pancreatin and diastase. See: Peptenzyme Digestant, Tab. (Reed & Carnrick)

W/Pancreatin, mycozyme diastase, hog bile desiccated, ext. nux vomica. See: Enzobile, Tab. (Mallard)

W/Pancreatin, laurocholic acid. See: Enzymet, Tabs. (Lloyd, Dabney & Westfield)

W/Paregoric, bismuth subsalicylate, zinc sulfocarbamate, salol & methyl salicylate. See: Parepeptol, liq. (Chicago Pharm.)

W/Pentaborbital sod., charcoal activated. See: Carget, Tab. (Hoack Labs.)

W/Bennin. See: Pepsin Essence, N.F.

W/Taka-diastase & Col. carb. See: Celltase, Tab. (Parke, Davis).

PEPSIN ESSENCE, Pepsin & Rennin Elixir, N.F.

PEPSIN LACTATED, ELIXIR. Comp. Pepsin Elixir, N.F.

PEPEZYME DIGESTANT. (Reed & Carnrick) Pepsin 65 mg., pancreatin 32 mg., diastase 50 mg./Tab. Bot. 25s, 100s, 500s Elixir Bot. 8 oz., 1 pt., 1 gal.

Uses: Digestant.

PEPTO-FERRIN. (Ulmer) Iron peptonized 60 mg./cc. Vial 1 cc., 10 cc.

Use: Anemias.

PEPTOLIN. (Walker Labs.) Pipradol HCl 2 mg., vitamins, lipotropics, hesperidin methyl chalcone 30 mg./45 cc. w/iron and other essential minerals. Bot. pts.

Use: Tonic & mood elevator.

PEPTONIDS & CREOSOTE. (Arlington) Beechwood creosote, guaiacol, amino acids & polypeptides. Liq. Bot. 6 oz., 1 pt., 1 gal.

Use: Cough preparation.

PEPTRON. (Savage) Peptonized Iron 100 mg./2 cc. Vials. 10 cc. Amp. 2 cc. 6s.

Use: Iron deficiency anemia.

PEPULCIN. (Ives-Cameron) Scopolamine methylsulfate 2 mg., aluminum hydroxide 2.5 gr., magnesium hydroxide 0.8 gr., ascorbic acid 10 mg./Tab. Bot. 100s

- Use: Antacid in the treatment of peptic ulcer
- PERAEMON.** See: Vit B 12
- PERALGA.** (Warner Chilcott) Barbitol $1\frac{1}{2}$ gr, aspirin $3\frac{1}{2}$ gr, acetophenetidin 2 gr /Tab Bot 12s, 50s, 100s
Use: Analgesic & sedative
- PERANDREN, NND 1962** (Ciba) Testosterone propionate in oil Amp (25 mg) 1 cc, Carfan 6s, 50s Vial (25, 50, 100 mg/cc) 10 cc, Carton 1; Box 6s Buccalet 5 mg, Bot 30s, 100s
Use: I.M. & Orally; androgen therapy
- PERANDREN PHENYLACETATE.** (Ciba) Testosterone phenylacetate Vial (50 mg/cc) 10 cc Carton 1s, 6s
- PERASED** (Mallard) Pentaerythritol tetranitrate 10 mg, Corbromol 250 mg /Tab Bot 100s 100's
- Use: Coronary vasodilator for prophylaxis in angina pectoris
- PERASMAL.** (Preme) Theophylline 2 gr, ephedrine HCl phenobarbital $\frac{1}{2}$ gr /Tab Bot 24s
Use: Relief in bronchial asthma and hay fever
- PERAZIL HYDROCHLORIDE, NND 1962** (Burroughs Wellcome) (Chlorcyclizine HCl) N methyl N (4-chlorobenzhydryl) piperazine HCl Cream 1%, Tube 1 oz; Jar 1 lb Tab 25 mg, Bot 100s, 1000s 50 mg Bot 100s 500s 1000s
Use: Antihistaminic
- W/Cocaine phosphate sudafed acetophenetidin aspirin caffeine See Emprozil C, Tabs (Burroughs Wellcome)
- PERBUZEM** (Zemmer) Pentaerythritol tetranitrate 10 mg, butobarbital sodium 15 mg /Tab Bot 100s
Use: Angina pectoris, hypertension
- PERCAINE.** See: Dibucaine Hydrochloride (Nupercaine HCl Ciba)
- PERCHLORETHYLENE.** See: Tetrachlorethylene USP
- PERCHLORPEBAZINE.** See: Compazine Suppositories (Smith, Kline & French)
- PERCOBABB.** (Endo) Dihydrohydroxycodone HCl 4.5 mg, dihydrohydroxycodone terephthalate 0.38 mg, hexobarbital 100 mg, homatropine terephthalate 0.38 mg, aspirin 224 mg, phenacetin 160 mg, caffeine 32 mg /Cap Bot 100s, 500s
Use: Analgesic and sedative
- PERCOBABB-DEMI** (Endo) Dihydrohydroxycodone HCl 2.25 mg, dihydrohydroxycodone terephthalate 0.19 mg, hexobarbital 50 mg, homatropine terephthalate 0.19 mg, aspirin 224 mg, phenacetin 160 mg, caffeine 32 mg /Cap or Tab Bot 100s, 500s, 1000s
Use: Analgesic
- PERCODAN.** (Endo) Dihydrohydroxycodone HCl 4.50 mg, dihydrohydroxycodone terephthalate 0.38 mg, homatropine terephthalate 0.38 mg, acetylsalicylic acid 224 mg, acetophenetidin 160 mg & caffeine 32 mg /Tab Bot 100s, 500s, 1000s
Use: Analgesic
- W/Hexobarbital See: Percobarb, Cap (Endo)
- PERCOMORPH LIVER OIL, NNB, 1955, A.D.B. 1955.** (May be blended with 50% other fish liver oils; each Gm contains 60,000 u Vit A & 8500 u Vit D)
See: A.S. Percomorph Liver Oil Oileum Percomorphum
- PERCORTEN, NND 1962** (Ciba) Desoxy corticosterone acetate in oil Amp (5 mg) 1 cc, Carton 6s, 50s linguet 2 mg, 5 mg, Bot 30s, 100s Pellet 125 mg, 1s Vial (5 mg/cc) 10 cc, Carton 1s, 6s
Use: I.M., implantation & orally; adrenal cortex replacement therapy
- PERCORTEN TRIMETHYLACETATE.** (Ciba) Desoxycorticosterone trimethylacetate Vial (25 mg/cc) 4 cc
Use: I.M.; adrenocortical insufficiency
- PERDIPIGEN-FLUID.** (Lilly) Diphtheria toxoid with 50,000 million killed H pertussis organisms/cc Vial 1.5 cc, 7.5 cc Alum prec (Lilly) Diphtheria toxoid with 75,000 million killed H pertussis organisms/1.5 cc Vial 1.5 cc, 7.5 cc
Use: Immunization
- PERFECTION LOTION** (Suliff & Cose) Hexachlorophene 0.25%, silicone 4.0% Lot 2 oz 8 oz
Use: Siliconized protective lotion
- PERFECTOCHOL.** (Lafayette) Iodoaliphane acid Tab 0.5 Gm Envelope 6s; Box 3s
Use: Diagnostic agent in cholecystography
- PERFOLIN.** (Lederle) Vit A 25,000 u, D 1000 u, B 110 mg, B 2.5 mg, niacinamide 150 mg, Vit B 12 5 mcg, C 150 mg, folic acid 5 mg /Cap Bot 30s, 100s, 1000s
Use: (1 daily) Multivitamin therapy
- PERGANTENE** (Pergantene) Sodium fluoride 2.2 mg /Tab Bot 120s, 1000s Fluoride ion 1 mg /Tab and 5 drops liquid Plastic dropper bot 30 ml Topical, 2%, plastic bot 125 ml
Use: Prevention of dental caries
- PERGANTENE-V LIQUID & LOZ.** (Pergantene) Sodium fluoride 1.1 mg (0.6 cc liq and 2.2 mg loz) Vit A 5000 USP u, C 50 mg, D 1000 USP u, B-6 0.3 mg (0.6 cc liq or loz) /Liq, Bot 30 cc, 60 cc; Loz, 100s
Use: Prevention of dental caries

- PERHYDROL.** See: Hydrogen Peroxide 30%.
- PERIACTIN.** (Merck, Sharp & Dohme)
*Ciproheptadine HCl 4 mg./Tab. Bot. 100s.
Use: Antipruritic.
- PERIACIN SYRUP.** (Merck Sharp & Dohme)
Ciproheptadine HCl 2 mg./5 cc. Bot. 1 pt.
Use: Allergic disorders and pruritic dermatoses.
- PERIBAR.** (Whittier) Pentoerythritol tetranitrate 10 mg., phenobarbital 16.2 mg./Tab. Long acting Tabs. contain 3X. Bot. 100s.
Use: Treatment of angina.
- PERICLOR.** N.N.D. 1962. (Ives-Cameron)
Pentoerythritol chloral 5 gr./Cap. Bot. 36s, 100s.
Use: [2 h.s.] Sedative.
- PERI-COLACE.** (Mead-Johnson) Dioctyl sod. sulfosuccinate 100 mg., peristim 30 mg./Cap. Bot. 30s, 60s, 5yr. (Dioctyl sod. sulfosuccinate 60 mg., peristim 30 mg./dr.) Bot. R oz.
Uses: Fecal softener and laxative
- PERIDEX L-A.** (Whittier) Pentoerythritol tetranitrate 60 mg./Tab Bot. 100s.
Use: Angina pectoris.
- PERIDIAL.** (Cutter)
Use: Renal stappage, barbiturate poisoning, edema, hepatic coma, chronic uremia.
- PERIDINET.** (Beutlich) Ascorbic acid 100 mg., hesperidin complex 100 mg./Chewable Tab. Bot. 100s
Uses: Orally, infants and children for capillary fragility.
- PERIHEMIM.** (Lederle) Caps Ferrous fumarate 168 mg., Vit. B-12 with Astring 3/5 U.S.P. Oral Unit, Iron 55 mg., C 50 mg., folic acid 0.67 mg./Cap Bot. 100s, 500s, 1000s.
Liqs Ferrous gluconate 197.33 mg., folic acid 0.34 mg., Vit. B-12 4 mcgm, powdered stomach 40 mg., sol. liver fraction 140 mg./5 cc. Bot. 4 & 16 fl. oz., 1 gal. Jr. size approx. 1/4 the above formulas.
Use: Orally, anemias.
- PERIMYCIN** (formerly Aminomycin), (Warner-Lambert Res. Inst) An antibiotic of the heptane type elaborated by *Streptomyces coelicolor* var. *aminomycinus*
Use: Antifungal antibiotic.
- PERIN.** N.N.D. 1962. (Endo) Piperazine-calcium edothamil equiv. to 500 mg of piperazine hexahydrate/5 cc. Syr. Bot. 4 oz., 1 pt., 1 gal. Also as water (500 mg) Rox 30s.
Uses Anthelmintic (Pinworms & roundworms)
- PERISTALTINE.** (Ciba) Cascara glycosides. Tab. 1 1/2 gr., Rot. 15s, 100s.
Use: Orally [1 t.l.d.] laxative.
- PERISTIM.** (Mead Johnson) Cosanthronol.
Use: Stimulant laxative.
- PERISTIM.** An anthraquinone extracted from caccara. W/Dioctyl sod. sulfosuccinate. See: Peri-Colace, Cap., Syr. (Mead-Johnson)
- PERITRATE.** N.N.D. 1962. (Warner-Chilcott) Pentoerythritol tetranitrate. Tab. 10 mg., Bot. 100s, 500s, 5000c. Tab. (20 mg) Bot. 100s, 500s.
Use: [1-2 t.l.d.] Coronary vasodilator used in angina pectoris.
- PERITHAZIDE SA.** (Warner-Chilcott) Pentoerythritol tetranitrate 20 mg., hydrochlorothiazide 25 mg., in the immediate release layer, 60 mg pentoerythritol tetranitrate in the sustained-release layer/Tab, Bot, 100s.
Use: Cardiac edema
- PERITRATE W/AMINOPHYLLINE.** (Warner-Chilcott) Pentoerythritol tetranitrate 10 mg., aminophylline 100 mg., aluminum hydroxide 150 mg./Tab. Bot. 100s, 500s,
Uses: Treatment of angina pectoris
- PERITRATE W/NITROGLYCERIN.** (Warner-Chilcott) Pentoerythritol tetranitrate 10 mg., nitroglycerin 1/200 gr./Tab. Bot. 50s.
Uses: Angina pectoris.
- PERITRATE - PHENOBARRITAL.** (Warner-Chilcott) Peritrate 10 mg., phenobarbital 15 mg./Tab. Bot. 100s, 500s. Also available: Peritrate 20 mg., phenobarbital 15 mg./Tab. Bot. 100s, 500s.
Uses: Control of angina pectoris.
- PERITRATE W/PHENOBARRITAL.** (Warner-Chilcott) Pentoerythritol tetranitrate 20 mg., phenobarbital 15 mg./Tab Remaining 60 mg pentoerythritol tetranitrate and 30 mg. phenobarbital gradually released over a 12-hour period Bot. 100s.
Use: Anxious coronary patients.
- PERITRATE SUSTAINED ACTION.** (Warner-Chilcott) Pentoerythritol tetranitrate 20 mg. in plate base and 60 mg. in a wax base released over a period of 12 hr./Tab. Bot. 100s, 500s
- PERITRATE TABLETS E.C.** (Warner-Chilcott) Tab. (10 mg) Bot. 100s, 500s.
Use: 1 or 2 at bedtime w/peritrate plate; angina pectoris.
- PERIVAS.** (Amtra-Grant) Acetylcathomal 200 mg. mephenesin 200 mg., pentoerythritol tetranitrate 15 mg./Tube. Box 50s, 100s.
Use: Orally, relief and prevention of angina pectoris.
- PERLATAN.** See: Estrone (Various Mfr.).
- PERMAPEN AQUEOUS SUSPENSION,** N.N.D. 1960, A.D.R. (Pfizer) Benzethacil Dibenzylethylenediamine d'penicilline G

* Generic name recognized by N.F., U.S.P. or A.M.A.

600,000 u Cartridge, single dose for use w/Steraject syringe
Use: Prolonged blood levels of penicillin therapy

PERMAPEN FORTIFIED IN AQ. SUSP. (Pfizer) Dibenzylethylenediamine dipenicillin G 300M u, procaine penicillin G 300M u / Cart

Use: Penicillin sensitive organisms

PERMAPEN ORAL SUSPENSION, NNR 1955 (Pfizer) Dibenzylethylenediamine ds penicillin G 300,000 u / 5 cc Susp Bot 2 oz

Use: Oral penicillin therapy

PERMICORT (Person & Covey) Hydrocortisone alcohol 1/4% or 1/2% Cream 1/4%/1 oz, 1/2%/1/2 oz or 1/2%/1 oz
Use: Dermatoses

PERMICORT-TQ CREME. (Person & Covey) Hydrocortisone alcohol as 1/4%, 1/2%, or 1% with cool tar 5% iodochlorhydraxy quin 3% Tubes 1/4% 1 oz, 4 oz; 1/2% 1/2 oz, 1 oz, 4 oz; 1% 1/2 oz
Use: Dermatitis

PERMITIL, NND 1962 (White) Fluphenazine di HCl 0.25 mg / Tab Bot 50s 500s
Chronolab 1 mg / Tab Bot 30s
Use: Anti anxiety agent

PERNAEMON (PL) (Organon) Liver from pregnant cows & Vit B 12 20 mcg / cc Vial 10 cc, 6s
Use: Hematinic

PERNAJEK. (Elder) Vit B 12 300 mcg / cc. Vial 10 cc.

PERNOX, NNR 1954 (Pretested) Ricinoleic acid, propylene glycol, propylene glycol monostearate, wool fat & tragocanth/vaginol cap Bot 6s
Use: Intervaginal, contraceptive

PEROAGENT. (Merrell) Zinc peroxide 10% Cal peroxide 1%, ceepryn chloride 1.500 Oint, Jar 4 oz, 1 lb
Use: Antiseptic.

PEROXIDES.
See: Hydrogen Peroxide
Urea Peroxide
Zinc Peroxide Medicinal

PERPARIN. See: Ethaverine (Chema Puro)
PERPHENAZINE, NND 1962 (2 by droxyethyl) 4 [3 (2-chloro 10 phenothiazyl) propyl] piperazine See: Trilafan, Tab (Schering)

PERSANTIN (Geigy) Dipyridamole 25 mg / Tab; dipyridamole 5 mg / cc Tabs 25 mg / 100s; 10 mg amp / 10s, 10 mg amp / 50s
Use: Coronary or myocardial insufficiency

PER-SED. (Columbus) Dihydracadeinane bitartrate 1.67 mg, citric acid, ammonium chloride, chloroform & alcohol / 5 cc Bot 3 oz, 1 pt, 1 gal
Use: Cough preparation

PERSISTIN. (Sherman Labs) Aspirin 2 1/2 gr. salicylic acid 7 1/2 gr / tab Bot 90s
Use: Analgesic

PERSONAMINE. (Person & Covey) Chlor pheniramine maleate 2 mg, thenylpyramine fumarate 12 mg, pyrilamine maleate 12 mg / Tab Bot 100s, 1000s
Use: Antihistamine therapy

PERSONAPHEN (Person & Covey) Phenol 0.95% sodium chloride in special, readily modified w/a lotion Bot 4 oz
Use: Psoriasis

PERSONATINIC. (Person & Covey) Ferrus fumarate 200 mg diocyl sodium sulfo succinate 50 mg / Tab Bot 100s
Use: Anemia

PERSONATINIC FORTE. (Person & Covey) Ferrus fumarate 303 mg Vit B 12 w / intrinsic factor 1/2 orol u, B 12 10 mcg; C 100 mg, P 30 mg, B 1 10 mg, B 2 5 mg B 6 2 mg niacinamide 25 mg, liver desic cated 150 mg lysine monohydrochloride 100 mg / Tab Bot 100s
Use: Anemias

PERSONATIC STRESS TARS. (Person & Covey) Fe fum 400 mg, B 12 w / intrinsic factor 2 U S P orol u, C 150 mg, hesperidin 50 mg folic acid 2 mg liver stomach substance 100 mg / Tab Bot 50s
Use: Oral hematinic

PERTENAL. (Crookes Barnes) Veratrum viride 100 mg, hamatropine methylbromide 2.5 mg, mannitol hexonitrate 30 mg, pheno barbital 15 mg / Tab Bot 50s 100s, 500s
Uses: (1 q 4 h) Hypertension

PERTEXOL. (McKesson & Robbins) Tyrothricin as esed in Yadara
PERTOL (Ulmer) Pentaerythritol tetranitrate 10 mg ethverine HCl 30 mg / Tab Bot 100s
Use: Coronary vasod later

PETOLAC BACTERIAL VACCINE (US Standard) Hemophilus pertussis 4000 million staphylococcus aureus 200 million, streptococcus hemolyticus 100 million, M. catarrhalis 200 million, Br influenza 100 million / cc Vial 10 cc, 30 cc
Uses: (1/4-1/2 cc S C) Whooping cough

PETUSSIN. (Seeck & Kade) Thyme extract saccharated Bot 4 oz, 8 oz
Use: Orally (1 2 pr n); Cough

PETUSSIN ANTIGEN. (Lilly) H. pertussis undenatured antigen Vial 5 cc, 20 cc
Uses: Prevention & treatment of whooping cough

* Generic name recognized by N F, U S P or A M A

PETUSSIN VAPORIZER. (Chesebrough-P) Medicated spray. Aerosol. 5 oz.

Use: Sprayed into sickroom for nasal congestion.

PETUSSIS BACTERIN, N.N.R. 1930. (Merck Sharp & Dohme) *H. pertussis* 50,000 million/cc., whole culture. Vial 1.5 cc. (1 3-dose immunization). Vial 7.5 cc. (5 3-dose immunizations).

Use: Immunization against whooping cough.

PETUSSIS IMMUNE GLOBULIN, N.N.D. 1960 (Hyland Labs) A 16.5% solution of the gamma globulin fraction of blood from healthy, adult human donors hyperimmunized with pertussis vaccine. Vial 1.5 cc.

Use: Passive prevention and treatment of whooping cough.

***PETUSSIS IMMUNE HUMAN SERUM, U.S.P., N.N.D. 1962.**

Cutter—Antipertussis Serum (Hypertussis) Human, Vial 2.5 cc. 1 initial dose.

Hyland—Pertussis Immune Serum Human. Vial 20 cc. of dried serum with 20 cc. diluent 1.5 cc.

League Blood Center—Pertussis Immune Serum Human. Vial 10 cc., 20 cc.

Serum Exchange—Pertussis Immune Serum Human Despak. Vial 20 cc.

Use: I.V. or I.M.; whooping cough therapy.

PETUSSIS IMMUNOGEN, N.N.R. 1930. (Parke, Davis) *H. pertussis* 6000 million, *Staphylococci albus* & *ovirus* 1000 million, *Streptococci hemolyticus* & *nonhemolyticus* 1000 million/cc. Vial 10 cc.

Use: Immunization against whooping cough.

PETUSSIS "SEROBACTERIN" VACCINE MIXED, (Merck Sharp & Dohme) 5400 million mixed organisms/cc. Vial 5 cc., 20 cc.

Use: Treatment of secondary infections associated w/whooping cough.

PETUSSIS-TOPAGEN. (Merck Sharp & Dohme) Antigenic fraction of 20 billion *H. pertussis* organisms in sterile saline sol./5 cc. Vial w/dropper 5 cc.

Use: Intranasal (5 drops daily); for respiratory infections.

***PETUSSIS VACCINE, U.S.P., N.N.D. 1962** (Various Mfr.)

Use: Immunization against whooping cough.

***PETUSSIS VACCINE, ALUMINUM NITROXIDE ADSORBED, U.S.P., N.N.D. 1962.** (Cutter) Vial 7.5 cc.

***PETUSSIS VACCINE, ALUM PRECIPITATED, U.S.P., N.N.D. 1962.**

Lilly—Vial 1.5 & 7.5 cc.

National Drug—Vial 1.5 & 7.5 cc.

Pitman-Moore—Vial 5 cc.

PETUSSIS VACCINE & DIPHTHERIA TOX-

***Generic name recognized by N.F., U.S.P. or A.M.A.**

OID COMBINED. See: Diph-Pertussis, Vial (Parke, Davis).

W/Diphtheria toxoid & aluminum hydroxide adsorbed. See: Dipertussis Alhydrax, Vial (Cutter).

P-E-T. (Maurry Labs.) Progesterone 25 mg., estrone 6 mg., testosterone 25 mg./cc. Vial 10 cc.

Use: I.M.; triple hormone therapy.

P.E.T. (Barre) Pentaerythritol tetranitrate. Tab. 10 mg., 20 mg. Bot. 100s, 1000s.

Use: Long-lasting oral vasodilator.

P.E.T.-B. (Barre Drug) Pentaerythritol tetranitrate 30 mg., barbital sodium 45 mg./Cap. Bot. 30s, 100s.

Use: Vasodilator and sedative.

PETHIDINE HYDROCHLORIDE, N.N.D. 1962. U.S.P., A.D.R. See: Meperidine Hydrochloride, U.S.P.

PETN. See: Pentaerythritol tetranitrate.

***PETRICHLORAL, N.N.D. 1962.** See: Perchlor, Cap. (Ives-Cameron)

PETROGALAR, N.N.R. 1949. (Wyeth) Plain, liq. petrolatum in agar gel. Bot. 1 pt. Alkaline, liq. petrolatum w/milk of Mag Bot. 1 pt. Cascara, liq. petrolatum w/cascara B. ext. Bot. 1 pt. Phenolphthalein, liq. petrolatum w/phenolphthalein. Bot. 1 pt.

Use: Laxative.

PETROLATUM CRUDE. See: Nafolan (Rawell)

***PETROLATUM, LIQUID, U.S.P., N.N.R. 1949, A.D.R.** Emulsion, N.F.; Gauze, U.S.P. Adepsine Oil, Glymal, liquid Paraffin, Parolein, White Mineral Oil, Heavy Liquid Petrolatum.

See: Alkal. (Am. Pharm. Co.)

Albolene. (McKesson & Robbins)

Mineral Oil (Various Mfr.)

Nujol (Plough).

Saxol. (Chester Kent).

Petronol. (Lilly)

Use: Laxative.

PETROLATUM, LIQUID, EMULSION, N.F.

See: Milkol. (Kremers-Urban)

W/Agar-Gel.

See: Agarol Plain. (Warner-Chilcott)

Cremogol (Cremogol).

Nivea Oil & Cream. (Duke)

Petrogalar, Prep. (Wyeth).

W/Cascara.

See: Cremogol w/Cascara, Emul. (Cremogol)

Petrogalar w/Cascara, Emul. (Wyeth)

W/Irish Moss Emulsion. Also w/phenolphthalein or cascara ext. See: Kandremul, liq. (Patch).

W/Kaolin, ethyl ricinoleate & Mag. hydroxide. See: Kericin, Susp. (Merrell)

PETROLONE

566

W/Milk of magnesia

See: Haley's M O, Liq (Phillips)

Magoleum, Liq (Blue Line)

W/Pectin See Pectal, Emulsion (Roerig)

W/Phenolphthalein

See: Agaral (Warner Chilcott)

Cremagol w/Phenalph (Cremagol)

Petragal w/Phenalph (Wyeth)

Phenolphthalein in Liquid Petrolatum Emulsion NF

W/Psyllium seed See: Parasyllium (Abbott)

W/Thiamine See Geol B, Emulsion (C D Smith)

W/Vit A See Emulseral, Emulsion (Upjohn)

PETROLONE (Vitamix Corp) Pentylene-tetra-zol 100 mg, Sod Phos 25%/cc Vial 10 cc

Use: (IM) Central nervous system stimulant

PETRONE (Kremers Urban) Progesterone 25 mg, testosterone 25 mg, estrone 6 mg/cc Aq Susp Vial 10 cc

Use: IM; functional uterine bleeding

PETRONOL (Lilly) Liquid Petrolatum, USP Bot 1 gal

PETROSYLLIUM (Whitehall) Mineral oil & psyllium seed Also avail w/phenolphthalein Bot 1 pt

Use: Orally; laxative

PEV-GRAM (Ingram) Hesperidin choleone Sod Cap 50 mg Bot 60s, 300s

Use Orally capillary disease therapy

PGA, See Folic Acid, USP

PHAC, (Cole) Phenobarbital ½ gr, acetyl salicylic acid 5 gr, caffeine alkaloid ½ gr gelsemium ext 3/40 gr/Tab or Cap Bot 100s

Use Analgesic & sedative

PHAGENE POWDER (Dumas Wilson) Citric acid, lactase, papain, methylsalicylate menthol, eucalyptol & chlorothymol Bot 7 oz

Use: Vaginal antiseptic & cleanser

PHAN, (Sandaz) Aluminum hydroxide glycine 450 mg, mag oxide 60 mg/Tab Bot 100s

Use: Antacid

PHAHODORN, N NR 1951 (Winthrop Labs) Cyclaborbital cal, cyclahexenylethylborbitalate cal Tab 3 gr Bot 100s, 500s

Use: Orally; sedative & hypnotic

W/Epival See: Evidarn, Cap & Tab (Winthrop)

PHANOLAR, (Winthrop) Plaquenil 60 mg, aspirin 300 mg/Tab

Use: Treatment of rheumatoid arthritis

PHANTOS, (Cooper Tinsley Lab) Delayed action tab Immediate release: amphetamine sulf 5 mg, thyroid ½ gr, atropine 1/360 gr, alain ¼ gr Intermediate release:

amphetamine sulf 5 mg, thyroid ½ gr, atropine 1/360 gr, Final release: amphetamine sulf 5 mg, thyroid ½ gr, phenobarbital ¼ gr Tabs Bot 30s, 250s 500s, 1000s

Use Anti obesity

PHARAGESIC, (Marian) Outer shell benzocaine 5 mg; inner tablet chlorpheniramine maleate 4 mg, phenylpropanolamine HCl 25 mg, acetaminophen 300 mg

Use Pharyngitis

PHARMAGEL A & B (Atlantic Gelatin) Gelatin

PHARMALAX SUPPOSITORIES, (Pharmacia Labs) Sod bicarb 0.6 Gm Pot bitartrate 0.9 Gm in cocoa butter base Supp Box 4s, 20s, 60s

Use Laxative

PHARYCIDIN CONCENTRATE, (Purdue Frederick) Choline salicylate 870 mg, cetyl dimethyl benzyl ammonium chloride 15 mg/5 cc Liq Bot 4 oz 8 oz

Use Analgesic w/antiseptic

PHAZYME (Reed & Carnrick) Pepsin 100 mg, diastase 25 mg activated dimethyl polysiloxane 60 mg pancreatin 240 mg/Tab Bot 50s 100s

Use Relief of flatulence

PHAZYME W/PHENOBARBITAL (Reed & Carnrick) Outer layer—pepsin NF 100 mg, diastase 25 mg; activated dimethyl polysiloxane 20 mg; phenobarbital 15 mg Inner core—pancreatin NF 240 mg, activated dimethyl polysiloxane 40 mg/Tab (two phase) Bot 50s

Use Antiflatulent, enzyme combination

PHEBEN, (Sutliff & Case) Phenolphthalein 325 mg, alain 16 mg, belladonna ex 54 mg strychnine 0.8 mg, ipecac pow 43 mg/Pill Bot 1000s

Use Laxative

PHEBORINE, (Buffington) Diallylbarbituric acid ¼ gr, stramonium ext ½ gr, ephedrine HCl ½ gr/Pill Bot 100s, 1000s

Use Sedative

PHEDOXE, (Elder) dl Methamphetamine HCl 5 mg/Tab Bot 100s, 500s

Use CNS stimulant

PHEDRACOL (Mallard) Morphine sulfate ¼ gr, ephedrine HCl ½ gr, labela Infata 4 gr, Sod citrate 20 gr, Pot guaiaccol sulfonate, methyl & aromatics/A oz Syr. Bot 3 oz, 1 pt, 1 gal

Use (1 tsp q 3-4 h) Coughs

PHEDRAMIST, (Caldwell & Bicar) Acetaphen etidin 120 mg, salicylamide 250 mg, Dovers Pwd 24 mg, racephedrine HCl 5 mg, pyrilamine maleate 12 mg, Vit C 50 mg/Cap Bot 100s, 1000s

Use: Common cold

- PHEDRETAL.** (Rorer) Ephedrine HCl $\frac{3}{8}$ gr., phenobarbital $\frac{1}{4}$ gr./Tab. Bot. 100s, 500s, 1000s.
Use: (1-2 t.i.d.) Bronchial antispasmodic w/ sedative.
- PHEDRISOX.** (Ascher) Methamphetamine HCl (2.5 & 5 mg) Tab. Bot. 1000s. Vials (2 mg./cc.) 10 cc.
Use: Central nervous system stimulant.
- PHELANTIN KAPSEALS.** (Parke, Davis) D-lantoin 100 mg., phenobarbital 30 mg., desoxyephedrine HCl 2.5 mg./Kopseal. Bot. 100s.
Use: 1 t.i.d. epilepsy.
- PHEMERIDE.** See Benzethonium Chloride.
- PHEN-MER-NITE.** (Massengill) Phenylmercuric nitrate.
Cap.s ($\frac{1}{8}$ gr.) Bot. 100s, 500s, 1000s.
Oint., (1:1500) Tube 1 oz., Jar 1 lb.
Ophth. aint., (1:300,000) Tube $\frac{1}{8}$ oz. w/ phenacaine, Tube $\frac{1}{8}$ oz.
Sol., (1:1000) Bot. 1 pt., 1 gal.
Vaginal supp: Box 12s.
Tab. w/benzacaine: Pkg. 1 pt., 1 gal.
Tr. (1:3000) Bot. 1 pt., 1 gal.
Vaginal pow., Vial (5 Gm.) Jar 1 lb.
Use: Antiseptic.
- PHEMEROL CHLORIDE, N.N.D. 1962, A.D.R.** (Parke, Davis) Benzethonium chloride.
Ophth. sol.: Uncolored, 1:5000 w/boric acid 2%, Bot. 1 pt., 1 gal.
Sol.: Uncolored, 1:1000, Bot. 1 pt., 1 gal.; colored, 3%, Bot. w/applicator 1 fl.oz., Bot. 1 pt., 1 gal.
Tr.: Red colored, alcohol 65%, acetone 10%, Bot. 1 pt., 1 gal.
Use: Antiseptic & germicide.
W/Solleylanilide. See Sonicyl Cream (Parke, Davis).
- PHEMITHYN.** (Flint, Eaton) Benzethonium chloride monohydrate 3.17% Bot. W/ measuring cap. 4 oz.
Use: 1 cap full/1 quart warm water, vaginal douche.
- PHENAC.** (Sutliff & Case) Acetophenetidin B mg., strontium salicylate 32.5 mg., phenolphthalein 6.5 mg., camphar-monobromate 32.5 mg., eucalypt 16 mg., atropine sulfate 0.03 mg./Tab. Bot. 1000s.
Use: Analgesic & laxative.
- *PHENACAINE HYDROCHLORIDE, N.F., N.N.R. 1950.** (Upjohn) Ethanyl diethoxydiphenylamide HCl. Bot. 5 Gm., 1 oz. See: Holocaine HCl, Prep. (Winthrop Lab, Abbott).
Use: Local anesthetic.
W/Atropine, phenol, nut gall & zinc oxide.
See: Tonicaine, Oint. & Supp. (Upjohn).
- W/Bacitracin.** See: Topitracin Oint. W/ Phenacaine. (Commercial Solvents Corp.)
W/Ephedrine. (Upjohn) Phenacaine HCl 1%, epinephrine 1:25,000. Ophth. oint. Tube 1 dr.
W/Mercarbolid. (Upjohn) Holocaine HCl 2% & mercarbolid 1:3000. Ophth. oint., Tube w/applicator tip, 1 dr.
W/Sulfanitamide & cod liver oil. See: Morumide & Phenacaine Ophthalmic Ointment (Massengill).
- *PHENACEMIDE, N.N.D. 1962.** Phenylacetylurea. See: Phenurone, Tab. (Abbott)
- PHENACETIN, U.S.P., N.N.R. 1941, A.D.R.** (Various Mfr.). See: Acetophenetidin, Prep.
- PHENACETIN COMPOUND.** (Flint, Eaton) Dover's pow. $\frac{1}{4}$ gr., phenacetin $1\frac{1}{2}$ gr., acetylsalicylic acid 2 gr., camphor monobromate $\frac{1}{4}$ gr., caffeine $\frac{1}{2}$ gr./Cap. Bot. 100s, 500s, 1000s.
Use: (1 q. 3 h.) Analgesic & antipyretic.
W/Atropine. (Flint, Eaton) Dover's pow. $\frac{1}{4}$ gr., phenacetin $1\frac{1}{2}$ gr., aspirin 2 gr., camphor monobromate $\frac{1}{4}$ gr., caffeine $\frac{1}{2}$ gr., atropine sulfate 1/500 gr./Cap. Bot. 100s, 500s, 1000s. See: Datin, Cap. & Tab. (Massengill).
Use: (1 q. 3 h.) Analgesic & antipyretic.
- PHENACETYL.** (Smith-Dorsey) Acetylsalicylic acid 3 gr., acetophenetidin 2 gr., caffeine citrate $\frac{1}{2}$ gr., galium 1/10 gr./Cap. Bot. 100s, 1000s.
W/Codaine sulfate $\frac{1}{2}$ gr., Bot. 100s, 500s, 1000s.
Use: Analgesic & antipyretic.
- PHENACETYLCARBAMIDE.** See: Phenurone, Tab. (Abbott).
- PHENACETYLUREA.** See: Phenurone, Tab. (Abbott).
- PHENACRIDANE.** [p-Hexyloxyphenyl]-10-methylacridinium chloride. See: Micridium. (Johnson & Johnson)
- PHENAGESIC.** (Dahn) Phenylephrine HCl 10 mg., propylpropanolamine HCl 50 mg., pyrilamine maleate 25 mg., pheniramine maleate 25 mg., acetyl-p-aminopheno 300 mg./Tab. or 10 cc. Bot. Tab 50. Bot. Syrup 6 oz.
Use: Treatment for the common cold.
- *PHENAGLYCODOL, N.N.D. 1962.** See: Ultrac, Pulv. (Lilly)
- W/Acetylsalicylic acid, dextro-propoxyphene.** See: Darvo-Tran, Pulvule. (Lilly)
- W/Propoxyphene, acetylsalicylic acid.** See: Vondar, Caps (Lilly)
- PHENALCO.** (Chicago Pharm.) Phenylmercuric nitrate 1:3000 in alcohol-acetone base. Bot. 1 pt., 1 gal.
Use: Skin disinfectant.

* Generic name recognized by N.F., U.S.P. or A.M.A.

PHENALZINE DIHYDROGEN SULFATE. See: Nardil (Warner Labs)

PHENAMPROMID, (Lederle) Under study
Use: Analgesic

PHENANTOIN Mephenytoin N Methyl 5 5 phenyl ethylhydantoin See Mesantoin Tab (Sandoz)

PHENAPHEN (Robins) Phenacetin 3 gr as
pirin 2½ gr, hyascyamine sulfate 0031
mg, phenobarbital ¼ gr /Cap or Tab
Bot 100s, 500s, 1000s Also avail w/
codeine phosphate ¼ gr & ½ gr Bot
Caps 100s, 500s

Use: (1 2 prn) Analgesic

PHENAPHEN No. 4 (Robins) Aspirin 162
mg, phenacetin 194 mg, phenobarbital
162 mg, hyascyamine sulfate 0031 mg
codeine phosphate 64 E mg /Cap Bot
100s, 500s

Use: Analgesic

PHENAPHEN PLUS (Robins) Phenacetin 194
mg, aspirin 162 mg, phenobarbital 162
mg, hyascyamine sulf 0031 mg, phe
niramine maleate 125 mg, phenylephrine
HCl 10 mg /Tab Bot 100s

Use: Treatment of the common cold and
upper respiratory infections

PHENAPHTHAZINE Sod d nitro phenyl
azo naphthal d sulfonate See: Nitrazine
paper (Squibb)

PHENARSONE SULFOXYLATE, N N D 1959
See: Aldersone (Abbott)

PHENASPIRIN COMPOUND. (Lannett) Pho
nobarbital ¼ gr aspirin 3½ gr, phen
acetin 1½ gr /cap Bot 1000s, 5000s

PHENAZINE (Elder) Pot penicillin G 100,
000 u w/sulfamethoxazine, sulfadiazine &
sulfamerazine 0.167 Gm each/Tab Bot
24s

Use: Orally; penicillin w/triple sulfonamide
therapy

PHENAZOCINE HBr, N N D 1962 1,2,3,4
5,6-hexahydro E hydroxy 6 11 d methyl 3
phenethyl 2 6 methano-3 benzoxazine HBr
See: Prinadol, Tab, Sol (Smith, Kline &
French)

PHENAZODINE. (Lannett) Phenylazodiamine
pyridine HCl 0.1 Gm /Tab Bot 100s
Use: Urinary antiseptic.

PHENAZONE, N N R. 1940. See: Antipyrone
(Various Mfr.)

PHENAZOPYRIDINE HCl 2,6-Diamino-3
phenylazopyridine HCl
See: Phenylazo d amino pyridine hydro
chloride

PHENCAREAMIDE (Rier) 5 2-d ethyl
aminoethyl d phenylthiocarbamate
Use: Spasmolytic agent

PHENCYCLIDINE See: Sernyl (Parke Davis)

* Generic name recognized by N F U S P or A.M.A.

PHENOMETRAZINE BITARTRATE Colorie
content 0.3/Tab See Plegine (Ayerst Lab)

PHENEDRINE. (MacAllister) Amphetamine
sulfate Bot 1 oz, 1 pt, 1 gal

Use: Central nervous system stimulant

**PHENEEN GERMICIDAL SOLUTION & TINC
TURE.** (Uimer) Benzalkonium Chloride
USP 1% Bot 1 qt & 1 gal

* **PHENELZINE SULFATE, N N D 1962** Mono
amine oxidase inhibitor, beta phenylethyl
hydrazine sulfate See Nardil (Warner
Chilcalf)

PHENERGAN NCl, N N D 1962 (Wyeth)
(Promethazine) N (2'-O-methylamino 2
methyl ethyl)phenothiazine HCl

Cream 2%, Tube 1 12 oz Lot, Phenergan
2%, neocalamine 15% Bot 4 oz
Supp (25 mg) Box 12s

Use: Antihistaminic

PHENERGAN EXPECTORANT TROCHES
(Wyeth) Phenergan 15 mg, Ipecac pow
ext 23 mg, Pot guaiaac sulfonate 162
mg /Troche Jar 36s

Use: Expectorant

**PHENERGAN EXPECTORANT TROCHES W/
CODEINE** (Wyeth) Phenergan 15 mg,
codeine 81 mg, Ipecac pow ext 23 mg,
Pot guaiaac sulfonate 162 mg /Troche
Jar 36s

Use: Expectorant

PHENERGAN HYDROCHLORIDE, N N D
1960 (Wyeth) Expectorant Phenergan
HCl 5 mg, Ipecac fldxt 0.17 min, pot
guaiaac sulfonate 44 mg, chloroform 0.25
mg, citric acid 60 mg, Sod citrate 197
mg /5 cc Bot 1 pt Also avail w/codeine
phosphate 1/6 gr/dr Supp 25 mg Box
12s Syrup: Phenergan HCl 6.25 mg /5 cc.
Also Forte 25 mg /5 cc Bot 1 pt Tab:
Phenergan HCl 12.5 mg & 50 mg /Tab
Bot 100s

PHENERDAN INJECTION. (Wyeth) Prametha
zine HCl Vial (25 mg/cc) 10 cc, (50
mg/cc) 10 cc

Use: IM, IV, antihistaminic

PHENERDAN PEDIATRIC. (Wyeth) Dextra
methorphan HBr 7.5 mg, promethazine
HCl 5.0 mg, fluidextract Ipecac 0.17 min,
pot guaiaac sulfonate 44 mg, min chloro
form 0.25 min citric acid 60 mg, sod
citrate 197 mg /5 cc Bot pt

Use: Ant tussive

* **PHENERIDINE** 1 (b phenyl b ethyl)-4 car
bethoxy 4 phenylpiper dine

* **PHENETHICILLIN POEASSIUM, N N D**
1962 Alpha phenoxyethyl penicillin pot
Penicillin 152 pot

See: Alpen Tabs (Schering)

Brazil (Beecham Res)

Chemipen, Tab. (Squibb)

Dorsil (Wyeth)

Dramcillin-s (White)

Maxipen, Tab. (Roerig)

Ro-Cillin, Tabs. (Roell)

Semopen, Tab, Pow. (Massengill)

Synallin, Tab. (Bristol)

PHENETRON INJECTABLE. (Lanett) Chlorpheniramine maleate 100 mg./cc. Vial 3 cc.

PHENETSAL, N.N.R. 1948. See: Salophen, Tab. (Winthrop Labs.)

* **PHENFORMIN HCl, N.N.D. 1962.** N¹-phenethylbiguanide HCl. See: D.B.I. (U. S. Vitamin)

PHENICHTHOL (Parke, Davis) Phenol 2%, ammonium ichthosulfonate, alum & lead plaster. Oint., Jar 1 lb., 5 lb.

Use: Antiseptic.

* **PHENINDAMINE TARTRATE, U.S.P.** Syr. or Tab, U.S.P. 2-Methyl-9-phenyl-2,3,4,9-tetrahydro-1-pyridindene tartrate. See: Thepharin, Prep. (Roche)

W/Alumium aspirin, chlorpheniramine maleate, phenylephrine HCl. See: Dolco, Tabs. (Assher)

W/Chlorpheniramine maleate, phenylpropanolamine HCl. See: Nolamine, Tab & Elix. (G. W. Carrick)

W/Dextromethorphan, chlorpheniramine maleate, phenylephrine HCl, menthol See: Meladon, Syr. (Mayrand)

W/Dextromethorphan HBr, phenylephrine. See: Cerose, Pediatric (Ives-Cameron)

W/Phenylephrine hydrochloride, carbamphenethionedisulfonate. See: Dandril (Whitehall Labs.)

W/Phenylephrine HCl, chloroform, Ipecac fluid, glycerol, pot. guaiacal sulfonate, sod. citrate, citric acid. See: Cerose, Liq. (Ives-Cameron)

W/Phenylephrine maleate, chlorpheniramine maleate, phenylephrine. See: Carhist Elixir (Carrione)

W/Sulfadimethoxine, N-acetyl-p-aminophenol, caffeine. See: Modricide, Cap. (Roche)

* **PHENINDIONE, N.N.D. 1962** (2-Phenylindane-1,3-Dione; 2-Phenyl-1,3-Indandione; Phenylindanedione, Dindevan)

See: Danilone, Tab. (Schleffelin)

Hedulin, Tab. (Hed Pharm.)

Indan, Tab. (Parke, Davis)

PHENIODOL. See: Iodoalphonic Acid

* **PHENIPRAZINE HCl.** See: Cotran, Tab. (Lakeside)

PHENIRAMINE, 1-Phenyl-1-(2-pyridyl)-2-dimethylaminopropane. (No pharmaceutical form available.)

W/Codaine phosphate. See: Antosen, Syr. (Squibb)

* Generic name recognized by N.F., U.S.P. or A.M.A.

W/Scopolamine HBr. See: Nio-Piracene, Tab. (Nilon)

* **PHENIRAMINE MALEATE, N.F., N.N.D. 1962;** Tab, N.F.† Ophth. Sol, N.F. 1-Phenyl-1-(2-pyridyl)-2-dimethylamino-propane maleate. Prophenpyridamine.

See: Ichistae (Union).

Trimelae (Stering).

W/Acetophenetidin, aspirin, phenobarbital, hyoscyamine sulfate, phenylephrine HCl. See: Phenaphen Plus, Tab. (Robins)

W/Ammonium chloride, Sod. citrate & chloroform. See: Trimetose, Syr. (Stering)

W/d-Amphetamine sulfate, chloroform, methyl & alcohol. See: Tussate, Syr. (Pitman-Moore)

W/Dihydrocodeinone bitartrate, Pot. guaiacal sulfonate, Sod. citrate, citric acid & chloroform. See: Tussar, Liq. (Armoer)

W/Dihydrocodeinone bitartrate, Pyrilamine maleate pot. citrate. See: Citraforte. (Boyle)

W/Dihydrocodeinone bitartrate, phenylpropanolamine HCl, pyrilamine maleate, glyceryl guaiacolate, chloroform. See: Tus-saminic Expectoran, Sy. (Smith-Dorsey)

W/Dihydrocodeinone bitartrate, pyrilamine maleate, phenylephrine HCl, pot. guaiacal-sulfonate, cherry flavor. See: Nova-Tussa, Liq. (Drug Specialties-Prane)

W/Doxylamine succinate, pyrilamine maleate. See: Tridecamine, Tab. (National)

W/Ephedrine sulfate & ammonium chloride. See: Elixase, Syr. (Squibb)

W/Glyceryl guaiacolate, desoxyephedrine HCl & codeine phosphate. See: Robitussin A-C, Syr. (Robins)

W/Hesperidin, ascorbic acid, thienpyramine fumarate, pyrilamine maleate, salicylamide, caffeine. See: Tripac, Cap. (Person & Covey)

W/Hesperidin, phenylephrine HCl, methapyrilene HCl, pyrilamine maleate, vit. C, salicylamide, acetophenetidin, caffeine. See: Boy's Citra, Caps. (Boyle)

W/Narcolee, phenylephrine HCl, chloroform, menthol. See: Conar, Liq. (Massengill)

W/Phenylephrine HCl. See: Ichist, Ophth. Sol. (Broemme)

W/Phenylephrine HCl. See: Phenilan, Tab. (Chicago Pharm)

W/Phenylephrine HCl, acetyl-p-aminophenol, atropine sulfate. See: TPC, Tab. (Tennessee Pharm.)

W/Phenylephrine HCl, methapyrilene HCl, pyrilamine maleate, aspirin, acetophenetidin, caffeine, Vit. C, hesperidin. See: Darbacin, Cap. (Crestmed)

W/Phenylephrine HCl, pyrilamine maleate, ascorbic acid See: Carizahit, Tab. (Mason)

W/Phenylephrine HCl, pyrilamine maleate, methapyrilene HCl See: Citra H P, Cap (Boyle)

W/Phenylephrine HCl & APC See Nova histine W/A P C, Cap (Pitman Moore)

W/Phenylephrine HCl, chlorafarm & meethal See: Novohistine, Elixir (Pitman Moore)

W/Phenylpropranolamine HCl, glyceryl guaiacolate See Darathon, Sol (Groll)

W/Phenylpropranolamine HCl, pyrilamine maleate See: Triaminic, Preps (Smith Dorsey)

W/Phenylpropranolamine HCl phenylephrine HCl, pyrilamine maleate See: Synergie Elix (Lemmon)

W/Phenylpropranolamine HCl phenylephrine HCl, pyrilamine maleate acetyl p amina phena See Phenogestic (Dalin)

W/Pyrilamine maleate See: Triaminic, Tab (Smith Dorsey)

W/Pyrilamine maleate diaxylamine succinate See: Tridecamin, Tab (Merrill)

W/Pyrilamine maleate & methapyrilene HCl See: Incorporist, Tab (Blue Line)
Ped olist, Syr (Calumbus)
Pyma Timed, Cap (Testagar)

W/Pyrilamine maleate phenylpropranolamine HCl See: Triolier, Tab (Lemar)

W/Pyrilamine Maleate, phenylpropranolamine HCl, trisulfapyrimidines See: Trisulfaminic, Prep (Smith Dorsey)

W/Pyrilamine maleate & phenyltoloxamine dihydrogen citrate See: Mullihist, Cap & Syr (Smith Dorsey)

W/Salicylamide, acetophenetidin, caffeine ascorbic acid, hesperidin purified See: Coryban, Cap (Koenig)

W/Scopolamine hydrabramide olum num hydroxide gel, ethyl aminobenzoate See: Daptren, Tab (Davis & Sly)

W/Thienopyram ne HCl pyrilamine maleate phenylpropranolamine See: Histene, Tab (Paul Maney)

W/Tytrathridn, neamydn, benzocaine See: Neo-T-Calin, Lozenges (Chicago Pharmacal)

W/Vit C, esprin & alkalinizing base See: Cehitra Tab (Organon)

*PHENISONONE HBr 3,4-d hydroxy-alpha isopropylaminopropiophenone HBr See: Dopanone HBr (Merck, Sharp & Dohme)

PHENISTAN (Chicago Pharm) Phenylephrine HCl 10 mg, prophenpyridamine maleate 12.5 mg / tab Bot 50s, 100s 1000s
Uses: (1) (1-d) Treatment of nasal congestion sinusitis & the common cold

PHENISTIX BEAGENT STRIPS (Ames) Ferric end magnesium salts with cyclohexyl

*Generic name recognized by N.F., U.S.P. or A.M.A.

selfamic acid *Dip ond test for phenyl ketoneena Bot 50s

Use Test urine for phenylketone

PHENITOL (Chicago Pharm) Phenylmercuric citrate Oint 1:1500, Jar 1 oz, 1/2 lb, 5 lb Sol 1:1500, Bot, 1 pt, 1 gal

Use Fungicide

PHENMERZYL NITRATE (Pitman Moore) Phenylmercuric citrate Sol 1:1500 Bot 1 pt, 1 gal Tr 1:100D, Bot 1 pt, 1 gal

Uses: Topically; germicide & fungicide

*PHENMETRAZINE HCl, N M D 1962 See Preledin, Tab (Galay)

PHENOBANE. (Smith, Miller & Potch) Hyoscyamus ext 1/2 gr, phenobarbital 1/2 gr / Tab Bot 1000s

Uses: Analgesic & sedative

*PHENOBARBITAL, USP N N R 1949 A D R; Elixir, U.S.P., Tab, USP (Various Mfr) (Phenylethylbarbituric Acid, Phenyl ethylmalonylureo, Barbenyl, Dorminal, Durnoryl, Neurabarb, Numal, Phenonyl, Somonal)

See Amvicol X10 (Stuart)

Amylalfene Tab (Texas First)

Barbipil, Pill (Chicago Pharm)

Barbita, Pill (Chicago Pharm)

Barbevis Tab (Chicago Pharm)

Blu phen Tab (Lemmon)

Eskabarb Spansule (Smith Kline & French)

Gardenal, Tab (Wallau)

Liquatal, Elixir (Merck Sharp & Dohme)

Lixaphen, Elixir (Massengill)

Luminal, Prep (Winthrop Labs)

Mosobarb #1 & 2 Tabs (Mason Pharm)

Peribar LA & PI (Whittier)

Phenofuric, Liq (Ethical Pharm)

Phob, Gel Et Tabs (Brewer)

Sental, Extentabs (Robins)

Teolaxin Cap (Paul Maney)

PHENOBARBITAL W/AMINOPHYLLINE

See: Aminophylline

PHENOBARBITAL W/ATROPINE SULFATE

See: Atropine Sulfate

PHENOBARBITAL W/BELLADONNA. See: Belladonna

PHENOBARBITAL & BROMIDE, EFFERVESCENT. (Burroughs Wellcome) Phenobarbital Sod 1/4 gr, Pot bromide 3 gr, Sod bromide 3 gr, ammonium bromide 1 1/2 gr- effe salts q s / Tab Tube 25s
Uses: Sedative & hypnotic.

Phenobarbital with Central Nervous System Stimulants see pp 571 to 573

PHENOBARBITAL COMBINATIONS

See: Tables of.

Acetophenetidin

Aminophylline W/Phenobarbital

Aspirin Barbiturate

Phenobarbital with Central Nervous System Stimulants

| Phenobarbital | Others | Manufacturer | Brand |
|-----------------|---|----------------------|---------------------------|
| ¼ gr. | Aminophylline 1½ gr., ephedrine ¾ gr. | Kremer Urban | Enofan, Tab. |
| ¼ gr. | Amphetamine phosphate 3 mg., aspirin 5 gr. | Clark & Clark Pharm. | Pen-Phetamine, Tab. |
| ½ or ¼ gr. | Amphetamine HCl 5, 10 or 15 mg., thyroid ½ gr. | Distributors | Laktobie, Tab. |
| ¼ gr. | Amphetamine sulfate 10 mg. | G. F. Harvey | Ampharb, Tab. |
| ¼ gr. | Amphetamine sulfate 5 mg. | Gold Leaf | Amfebarbs, Tab. |
| 20 mg. | d-Amphetamine sulf., 15 mg., omnobarbital 60 mg. W/7 minerals, & 9 vitamins | Stuart | Amvicei-X, Tab. |
| ¼ gr. | d-Amphetamine sulf., 2.5 mg., nicotinic acid 25 mg., vit. B-1 5 mg., C 50 mg., toladiastase 300 mg. | Parks, Davis | Amphedase W/Phenob., Tab. |
| ½ gr. | Amphetamine sulfate 5 mg., pentobarbital Sod. 1½ gr. | Columbus | Somtrol, Tab. |
| ¼ gr., or ½ gr. | d-Amphetamine sulfate 15 or 10 mg. | Wynn Pharmacal | Desatal, Tab. No. 1 & 2 |
| ¼ gr. | di-Amphetamine sulf., 15 mg., atropine sulf., 2/360 gr., thyroid 3 gr., elain ¼ gr. | Taylor | Obetrlo, T.R. Tab. |
| ¼ gr. | d-Amphetamine 15 mg., thyroid 3 gr., olein ¼ gr., atropine sulfate 1/160 gr. | Lowe | Lowebses CTR. T.R., Cap. |
| 30 mg. | d-Amphetamine sulf., 5 mg., pentobarbital 60 mg. | Nysco Labs. | Sedex, Tab. T.D. |
| ½ gr. | d-Amphetamine 5 mg., mellose 2 gr., thyroid ½ gr., atropine sulfate 1/300 gr. | Dumas-Wilson | Anlibese, Tab. |
| ¼ gr. | d-Amphetamine 5 mg., choline bitartrate 400 mg., di-methylene 150 mg. & vitamins | Lakeside | Obalip, Cap. |
| ½ gr. | d-Amphetamine HCl 5 mg., cellulose 2½ gr. | Dumas-Wilson | Celladox, Tab. |
| ¼ gr. | d-Amphetamine HCl 5 mg., methylcellulose 200 mg., vitamins | Stuart | Amvical, Cap. |
| ¼ gr. | d-Amphetamine HCl 5 mg., methylcellulose 200 mg., vitamins, Mag. trisulfate 200 mg. | Smith-Dorsey | Nexarin, Cap. |
| ¼ gr. | d-Amphetamine phosphate 1.5 mg., saicylemide 3½ gr. | Irwin, Neister | Analdine, Tab. |
| ¼ gr. | d-Amphetamine phosphate 5 mg., thiamine 5 mg. | Patch | Blubon, Tab. |
| ¼ gr. | Amphetamine sulfate 15 mg., thyroid 3 gr., atropine sulfate 1/160 gr., elain ¼ gr. | Paul Maney | Nika, Cap. |
| ¼ gr. | d-Amphetamine sulf., 5 mg., thyroid 1 gr., methylcellulose 5 gr., atropine sulf., 1/360 gr./Tab. | Bell Pharmacal | Celudex, Tab. |
| 1/6 gr. | d-Amphetamine sulfate 2 mg., mephesisin 200 mg., hematropine methylbromide 2 mg. | Adams | Relaxamine, Tab. |
| ¼ gr. | Amphetamine HCl 10 mg., thyroid 1 gr. | Dumas-Wilson | Amroid, Tab. |
| 23 mg. | d-Amphetamine HCl 15 mg., thyroid 50 mg. | Dumas-Wilson | Obesity, Cap |

| Phenobarbital | Others | Manufacturer | Brand |
|---------------|--|-----------------------|-------------------------|
| 1/4 gr | benzedrine sulfate 5 mg | Smith, Kline & French | Benzebor, Tab. |
| 1/4 gr | d Desoxyephedrine HCl 2.5 mg, dl alpha tocopheryl acetate 5 mg | Brean | Transibarb Cap |
| 1/4 gr | dl Desoxyephedrine HCl 5 mg | Flint Eaton | Barbidex, Tab |
| 30 mg | dl Desoxyephedrine HCl 2.5 mg, diltantol 100 mg | Parke Davis | Phelantin, Caps |
| 16 mg | dl Desoxyephedrine HCl 5 mg, Mag glucaate 7 1/2 gr | Kendall | Magalin Tab |
| 1/4 gr | dl Desoxyephedrine HCl 5 mg, thyroid 1 gr | Meyer | Myrab e Na J, Tab. |
| 1/6 gr | Ephedrine 3/8 gr, theophylline 2 gr | Chicago | Barbephine, Tab |
| 1/4 gr | Ephedrine HCl 3/8 gr | Mallard | Ephabarb Cap |
| 1/4 gr | Ephedrine HCl 3/8 gr | Rarer | Phedrelal, Tab |
| 1/4 gr | Ephedrine HCl 3/8 gr, acetophenctidin 3 gr | Mallard | Phenadrin, Cap |
| 1/6 gr | Ephedrine HCl 1/12 gr., aminophylline 1 gr, ethyl aminobenzoate 1/2 gr, aluminum hydroxide 1/2 gr | Irwin Neisler | Dainite Pediatric, Tab. |
| 1/6 gr | Ephedrine sulif 3/8 gr, prednisolone 0.5 mg theophylline 2 gr | Vitamax | Predisophine, Tab |
| 1/4 gr | Ephedrine HCl 3/8 gr, Sad iodide 1 gr, theophylline 1 1/4 gr | Testagar | Tepadrine, Tab. |
| 1/6 gr | Ephedrine HCl 3/8 gr, theopropinol 1 1/2 gr | National | Theopropedrine, Tab. |
| 1/4 gr | Ephedrine HCl 1/4 gr, theophylline 1 gr | Medicane | Meditab, Tab. |
| 1/6 gr | Ephedrine HCl 3/8 gr theophylline 1 1/2 gr | Endo | Teprin, Tab |
| 3/16 gr | Ephedrine HCl 3/16 gr theophylline calcium sal cyclate 1 gr, pot iod de 2 1/2 gr | Knoll | Quadrinal, Susp /5 cc. |
| 8 mg | Ephedrine sulif 24 mg theophylline 100 mg, thenyldiamine HCl 10 mg, glyceryl guaiacolate 100 mg | Breac | Brankatab, Tab |
| 8 mg | Ephedrine HCl 11 mg theophylline 65 mg belladonna leaves 5 mg pyrilamine maleate 12.5 mg, pot iodide 65 mg, m-ogecium tr sil cate 130 mg | Suffin & Case | Theofed Pediatric, Tab. |
| 1/4 gr | Ephedrine sulfate 3/8 gr | Shermae | Ephedrital, Tab. |
| 1/4 gr | Ephedrine sulfate 3/8 gr | Suffin & Case | Ephedratol, Cap |
| 1/2 gr | Ephedrine sulfate 1/2 gr amseophylline 1/2 gr | Warree Teed | Eu Phed Amine |
| 8 mg | Ephedrine sulfate 24 mg, aminophylline 65 mg, pyrilamine maleate 12.5 mg | Roymex | Barbuhist, Tab |
| 10 mg | Ephedrine sulfate 10 mg, aspirin 3 gr, Cal gluconate 100 mg, hyoscyamus ext 10 mg | Flessner | Hyefolac, Tab |
| 1/4 gr | Ephedrine sulfate 1/4 gr belladonna pow 1/10 gr, atromonium 1 gr | Chemica | Acopnald, Tab |
| 1/2 gr | Ephedrine sulfate 1/2 gr belladonna tr 0.5 cc | Maskell | Belphe'dri'barb, Tab. |
| 1/2 gr | Ephedrine sulfate 1/2 gr, euphorbia ext 1 1/2 gr. | Warren Teed | Eu Phed Ital, Tab. |

Phenobarbital with Central Nervous System Stimulants

Phenobarbital

Others

Manufacturer

Brand

| | | | |
|----------------|--|----------------------------|---------------------------------------|
| 10 mg. | Ephedrine sulfate 10 mg., Hyoscyamus est. 10 mg., aspirin 3 gr., Cal. gluconate 100 mg. | Plesner | Hyafolin, Tab. |
| ½ gr. | Ephedrine sulfate ¼ gr., isopropyl arterenol 10 mg., theophylline 2 gr. | Leeming | Naphenalin, Tab. |
| ¼ gr. | Ephedrine sulfate ¼ gr., organidin 10 mg. | Wampala Professional Drugs | Sadarzyl, Liq. |
| ¼ gr. | Ephedrine sulfate ¼ gr., Pot. iodide 2½ gr., Cal. lactate 2½ gr. | Professional Drugs | Navalene, Tab. |
| (Sod.) ¼ gr. | Ephedrine sulfate ¼ gr., Pot. iodide 2½ gr., Cal. lactate 2½ gr. | Professional Drugs | Hista-Navalene, Tab. |
| (Sod.) ½ gr. | Ephedrine sulfate ½ gr., theophylline 2 gr. | Cal. Chem. Co. Brewer | Asmilyn, Tab. |
| (Sod.) ½ gr. | Ephedrine sulfate ½ gr., theophylline-Sod. acetate 3 gr. | Blue Line | Luamin, Tab. |
| 7.5 mg. | Ergamine tartrate 0.5 mg., caffeine 50 mg. | Winthrop | Ercafital, Tab. |
| 8 mg. | Isoproterenol 10 or 5 mg., benzylophedrine 32 mg., theophylline 130 mg. | Crystal Xabins | Isuprel-Franol |
| 8 mg. | Isoproterenol 10 mg., acephedrine HCl 16 mg., neohyline 100 mg. | Massengill | Asaphylline, Tab. |
| ½ gr. | Methamphetamine HCl 3.33 mg. | Star | Ambar, Tab. |
| 15 mg. | Methamphetamine HCl 3 mg., secobarbital Sod. 30 mg. | Chestmed | Hypnarm, Pill |
| ½ gr. | Methamphetamine HCl 10 mg., thyroid 1 gr. | Winthrop Labs. | Amphabese, No. 9, Tab. |
| 16 mg. | Methamphetamine HCl 5 mg., Vit. C 100 mg., B-1 mg., niacin 10 mg., methylcellulose 150 mg. | Rarer | Obestrol, Cap. |
| ¼ gr. | Neosynephrine HCl ½ gr., theophylline 2½ gr. | Superior Pharm. | Adnaphrin, Cap. |
| ¼ gr. | Nux vomica ext. ¼ gr., carbenzylamine 6 gr. | Warner-Chilcott | Nurakardiac, Tab. |
| 16 mg. | Pentamethyl tetranitrate 10 mg. | Strong Cobb | Tanipent, Tab. |
| 45 mg. | Pentamethyl tetranitrate 80 mg. | Searle | Peritrate W/Phenobarbital S. A., Tab. |
| (Sod.) 100 mg. | Pentylenetetrazol 150 mg. | Central | TetraPhenobarb, Cap. |
| ½ gr. | Racophedrine ¼ gr., aminophylline 1½ gr. | Sullin & Case | Amadrine, Tab. |
| ¼ gr. | Racophedrine ¼ gr., theophylline Sod. glycinate 5 gr. | Sullin & Case | Synaphedal, Tab. |
| 32.5 mg. | Theobromine 325 mg. | Sullin & Case | Theophal, Tab. |
| 16 mg. | Theobromine 325 mg., pot. iodide 160 mg. | Sullin & Case | Theophal W/KI, Tab. |
| 8 mg. | Theophylline 130 mg., ephedrine HCl 24 mg., belladonna pow. ext. 5 mg., Pyrimamine maleate, 25 mg., KI 65 mg., magnesium trisilicate 130 mg. | Zimmer | Theofed, Tab. |
| ½ gr. | Thyroid ¼ gr., d-desoxyephedrine HCl 5 mg. & methylcellulose 2½ gr. | | |

Hysabel, Tab.

- Atropine Hyoscyamine Hyoscine Combina-
tions
- Atropine Sulfate w/Phenobarbital
- Belladonna Ext Combinations
- Belladonna Products w/Phenobarbital
- Cocaine Combinations
- Ephedrine Product Combinations
- Homatropine Methylbromide & Phenobarbital Combinations
- Hyoscine Hydrobromide Combinations
- Hyoscyamus Products & Phenobarbital Combinations
- Mannitol Hexanitrate w/Phenobarbital
- Mephensin & Barbiturates
- Phenobarbital W/Central Nervous System Stimulants
- Solicylamide Phenobarbital Combinations
- Secobarbital Combinations
- Sodium Nitrite Combinations
- Theobromine W/Phenobarbital Combinations
- Theophyllin W/Phenobarbital Combinations
- Veratrum Viride W/Phenobarbital Combinations
- PHENOBARBITAL W/HOMATROPINE METHYLBROMIDE** See Homatropine Methylbromide
- PHENOBARBITAL W/HYOSCYAMUS.** See Hyoscyamus
- PHENOBARBITAL W/MANNITOL HEXANITRATE** See Mannitol Hexanitrate
- PHENOBARBITAL & THEOBROMIDE COMBINATIONS** See Theobromine
- PHENOBARBITAL W/THEOPHYLLINE** See Theophylline
- PHENOBARBITAL W/VERATRUM VIRIDE** See Veratrum Viride
- ***PHENOBARBITAL SODIUM, USP, NNR** 1949; ADR; Inj, NF; Sterile, USP; Tab, NP (Sod Phenylethylborbiturate Soluble Phenobarbital) (Various Mfr) See Luminal Sod (Winthrop Labs)
- PHENOBARBITAL SODIUM IN PROPYLENE GLYCOL.**
Endo—Amp (2 gr, 5 gr) 2 cc, Box 12s, 100s
Vial—Amp (0.12 Gm) 1 cc, Box 12s, 25s, 100s
- PHENOBARBITAL THEOCALCIN** (Knoll) Phenobarbital ¼ gr, theocalcin 7½ gr, Tab Bot 100s, 500s
Uses (1 q 4 h) Coronary vasodilator sedative, diuretic
- PHENOBELL TABS A SPANCAPS** (Dome) Hyoscyamine sulfate 0.1037 mg atropine sulfate 0.0194 mg hyoscine HBr 0.0065 mg phenobarbital 76.2 mg/Tab Nyasymamine sulfate 0.3111 mg, atropine sulfate
- *Generic name recognized by NF USP or AMA
- 0.0582 mg, hyoscine HBr 0.0195 mg phenobarbital 48.6 mg/Cap Bot 100s
Uses: Antispasmodic, Antisecretory, Sedative
- PHENO-BEPADOL** (Ives Cameron) Liq 1 Phenobarbital ¼ gr, thiamine HCl 1 mg, riboflavin 0.5 mg, niacinamide 5 mg, Col pantothenate 0.3 mg, pyridoxine HCl 0.15 mg, rice bran ext, yeast ext & soluble liver fraction/dr Bot 1 pt, 1 gal Tab 1 Phenobarbital ¼ gr, thiamine HCl 3 mg, riboflavin 2 mg, niacinamide 10 mg, Col pantothenate 1 mg, pyridoxine HCl 0.5 mg, secondary liver fraction & dried yeast qs/Tab Bot 100s 1000s
- PHENOBROMIDE** (Central) Phenobarbital sod 2 gr, sodium bromide 80 gr/fl oz Bot pt, gal
Uses: Sedative
- PHENOBROMIDINE ELIXIR.** (Tilden) Phenobarbital 2 gr pot bromide 30 gr, sod bromide 30 gr, fluid extract passion flower 16 min/fl oz Bot pts, gals
Uses: Tranquillizer and sedative
- ***PHENOBUTIDIL 2** (2,4,6-trimethoxy)butyric acid See Vesipaque (Warner Lambert Res Inst)
- PHENOCYAMUS.** (Cole) Hyoscyamus ext ¼ gr, phenobarbital ¼ gr/Tab Bot 100s
Uses (1 2 11 d) Antispasmodic w/sedative
- PHENO-DONNA.** (Meyers Kasser) Nat listed Sod phenobarbital ¼ gr, belladonna paw ext 1/6 gr/tab Bot 50s, 100s, 500s 1000s
Uses (1 2 11 d) Antispasmodic w/sedative
- PHENODRIN** (Mallard) Phenobarbital ¼ gr, ephedrine HCl ¾ gr, acetophenetidin 3 gr/Cap Bot 100s, 500s 1000s
Uses (1 2 q 4 h) Antispasmodic & sedative
- PHENODYNE.** (Blue Line) Aspirin 3½ gr, acetophenetidin 2½ gr, caffeine ½ gr phenolphthalein ¼ gr/Cap or Tab Bot 100s; Also avail w/codeine ½ gr & ½ gr, Bot 100s
Uses: Orally (1 2 11 d); Analgesic & anti-pyretic w/laxative
- ***PHENOL, USP, NNR** 1953 ADR; Liq, USP; Oint, NP (Various Mfr) Carbolic acid
Uses: 0.5% to 3% conc in pruritus
W/Benzocaine Ichthammol balsam peru elem exsic, oil cade oil eucalyptus See Alucaine Oint (Jenkins Labs)
W/Benzocaine clove oil, menthol See Otolgesic Liq (Blue Line)
W/Hexachlorophene resorcinol oil of tur rect Red zinc oxide See Retolan (Vanpelt & Brown)
W/Mineral oil See: P & S Liquid (C A Baker Lab)

- oxyethyl)hydrazine Drazine hydrogen maleate
- ***PHENPROCOUMON**, 3 (o ethylbenzyl)-4 hydroxycoumarin See Liquamar, Tab (Oranon)
- PHENSAL**, (Pitman Moore) Acetophenetidin 2 gr, aspirin 4 gr, galxemium tr 3 msn, caffeine 1/4 gr /Tab Also w/cocaine sulfate 1/4 gr & 1/2 gr Bat 100s, 1000s
Use: Orally (1 lid); analgesic & antipyretic
- ***PHENSUXIMIDE**, NND 1962 See: Milan tin Prep (Parke, Davis)
- ***PHENTERMINE**, NND 1962 Alpha, alpha dimethylphenethylamine Phenyl tertiary butylamine HCl
See lanamin (Strassenburgh)
Wilpo (Dorsey Lab)
- PHENTETIOTHALEIN SODIUM**, NNR 1937 (No Mfr currently lists) Iso tadeikon
Use: Radiopaque medium for cholecystography & for liver function test
- ***PHENTOLAMINE HYDROCHLORIDE**, NF, NND 1962; Tab, NF 2 (m Hydroxy N p talylaminomethyl) 2 imidazole HCl See Regitine Hydrochloride, Tab (Ciba)
- ***PHENTOLAMINE METHANESULFONATE**, USP, NND 1962; Inj, USP See Egitine Methanesulfonate, lyophilized, Amp (Ciba)
- PHENURONE**, NND 1962 (Abbott) (Phenocemide, Phenocetylcarbamide Phenocetyl urea) Tab 0.5 Gm, Bat 100s, 1000s
Use: Psychomotor epilepsy, grand mal & petit mal epilepsies
- PHENURONE ENTERAB**, NND 1958 (Abbott) Tab 0.5 Gm, Bat 100s, 1000s
Use: Anticonvulsant
- N-PHENYLACETAMIDE** See: Acetamidid
- PHENYLACETYLUREA**, NND 1962 See: Phenurane Tab (Abbott)
- PHENYLALANINE**. For phenylalanine—free products See: Phenylketanuria
- PHENYLAZO TABLETS (APC)** Phenylazodiamino-pyridine hydrochloride 1 1/2 gr / Tab Bat 100s, 1000s
Use: Urinary analgesic and antiseptic
- ***PHENYL p-AMINOSALICYLATE**. See: Phenyl PAS Tebamin, Tab., Paw (Purdue Frederick)
- PHENYLAZO-DIAMINO-PYRIDINE W/Sulfadiazine, sulfacetamide** See: Azadettes Tab (Plesner)
- W/Sulfacetamide atropine sulfate hyoscyne hydrobromide, hyoscyamine sulfate See: Urclin, Tab (Richard & Williams)
- W/Sulfamethoxy-pyridazine HCl See: Azo Kynex Tab (Lederle)
- ***PHENYLAZO-DIAMINO-PYRIDINE HCl** or HBr. Phenazopyridine HCl or HBr
See Diridone (Premo)
Maliophene, Disc, Paw, Tab (Mallinckrodt)
Phenazodine, Tab (Lanett)
Phenylazo, Tab (APC)
Pirid, Tab (Miserer)
Pyridium, Pills (Warner Lambert)
Tridium, Tab (Amco Drug)
Urslad Tab (Crystal)
Urodine, Tab (Nysco)
- W/Gantanol See Azo Gantanol, Tabs (Roche)
- W/Gantrisin See Azo Gantrisin, Tab (Roche)
- W/Hyoscyamine HBr, atropine sulfate, scopolamine HBr See: Urogesic, Tab (Edwards)
- W/Methenamine mandelate See: Azo-Mandelamine Tabs (Warner Chilcott)
- W/Methenamine mandelate, dl methionine See Azofate, Tabs (Amfre Grant)
- W/Methenamine, calcium mandelate, sodium phosphate monobasic See: Urinal, Cop (Central)
- W/Sulfacetamide
See Azonamide, Tab (G F. Harvey)
Sulfidum, Tab (Columbus)
- W/Sulfacetamide, atropine methyl nitrate See Cystizem, Tab (Zemmer)
- W/Sulfacetamide, atropine methyl nitrate See Tridium, Tab (Amco)
- W/Sulfacetamide atropine sulfate, hyoscyne HBr, hyoscyamine sulfate See: Pylasetamide, Tab (Vitamiz)
- W/Sulfacetamide, methenamine mandelate acid sodium phosphate See: Uramon W/Azo Tab (Superior Pharm)
- W/Sulfacetamide, sulfamethazine, buffered pine sulf, hyoscyne HBr See: Sulfid B A Tab (Columbus)
- W/Sulfacetamide, hyoscyamine sulf, atropine See: Sufazo, Cop (Sutliff & Case)
- W/Sulfadiazine, sulfamerazine, sulfamethazine See: Pyridium Tri Sulfaz, Tab (Warner Chilcott)
- W/Sulfamethizole See Thiazolidin A Forte Tabs (Ayerst)
- W/Sulfamethizole, tetracycline phosphate See: Azotrex, Cop (Bristol)
- W/Sulfamethoxy-pyridazine See: Azo Kynex, Tab (Lederle)
- W/Sulfamethylthiadiazole
See: Thiazolidin A, Tab (Ayerst)
Urosulfon, Tab (Nepera Chem)
- W/Sulfamethylthiadiazole & sulfadiazine See: Suladyne, Tab (Stuart)
- ***Generic name recognized by NF USP or A.M.A.**

W/Sulfamethylthiadiazole, trisulfapyrimidines. See: Ro-Sulla, Tabs. (Rawell)

W/Tetracycline and sulfamethizole. See: Azotex, Cap. (Bristol)

*PHENYLBUTAZONE, N.N.D. 1952. 4-n-Rutyl-1,2-diphenylpyrazolidine-3,5-dione. See: Butazolidin, Tab. (Geigy)

W/Aluminum hydroxide, mag. trisilicate, homatropine methylbromide. See: Butazolidin Alka, Cap. (Geigy)

W/Prednisone, aluminum hydroxide gel, magnesium trisilicate, homatropine methylbromide. See: Sterasolidin, Cap. (Geigy)

PHENYL-1-RUTYL-NORSUPRIFEN. See: Arlidin, Tab. (Arlington-Funk)

PHENYLCARRINOL. See: Benzyl Alcohol, N.F.

PHENYLCINCNONIC ACID, N.N.R. 1947. See: Aphon, Tab. & Pow. (Warner). Clinchphen, Prep. (Various Mfr.)

PHENYL-DRANE. (Mahan) Phenylephrine NCl 0.5% in buffered isotonic soln. pH 5.0 to 5.5 1 oz. in spec. squeeze bot. Use. Sinus congestion, hayfever, head colds. See: Sinus congestion, hayfever, head colds.

*PHENYTERPBRINE HYDROCHLORIDE, U.S.P., N.N.R. 1954, A.D.R.; l-m-Hydroxyphenyl-2-methylaminoethanol (Neophryn). inj., U.S.P. Sol., U.S.P. (Various Mfr.) Crookes-Barnes, 2.5% and 10% Ophthalmic Drops./Vials 7.5 cc. and Drop-perettes. See: Almetrin Sol. (Meyer). Censdrin (Du Mont) Derizene Sol. (Hollister-Stier) Efricef—10% (Professional Pharm.) Isophrin HCl (Broemmel) Neo-Synephrine HCl, N.N.R. (Winthrop Labs.) Nose Drops—Spray (Johnson & Johnson) Phenyl-Drane, Soln (Mahan) Sucraphen Sol. (Smith-Dorsey) Synasol (Texas Pharm.) Use: Sympathomimetic agent, vasoconstrictor & mydriatic.

PHENYLEPHRINE TANNATE W/Ephedrine tannate, carbetapentane tannate, chlorpheniramine tannate, pyrilamine tannate. See: Rynaton Expectarant, liq. (Irwin, Neisler)

W/Carbetapentane tannate, chlorpheniramine tannate, ephedrine tannate. See: Rynatuss, Tabule (Irwin, Neisler)

W/Prophepyridamine tannate, pyrilamine tannate. See: Rynaton Tabulet, Tab. (Irwin, Neisler)

W/Pyrilamine tannate, chlorpheniramine tannate, caffeine alkaloid. See: Rynalert, Caps (Neisler)

*Generic name recognized by N.F., U.S.P. or A.M.A.

PHENYL ESTER OF p-AMINOSALICYLIC ACID. See: Pheny-Pas-Tabamin. (Frederick)

*PHENYLETHYL ALCOHOL, N.F. Beta-phenylethanol. Benzylcarbinal. (Various Mfr.) Use: Antibacterial; preservative (ophthalmic). beta-PHENYL-ETHYL-HYDRAZINE, (phenazine dihydrogen sulfate). See: Nardil. (Warner-Lambert)

PHENYLETHYLMALONYLUREA, U.S.P., N.N.R. 1949, A.D.R. See: Phenobarbital (Various Mfr.)

PHENYLIC ACID. See: Phenol, U.S.P.

PHENYLISATIN. See: Diacetylhydroxyphenylisatin

W/Prone concentrate, thiamine mononitrate. See: Prunisatin, Tab. (Sutliff & Case)

*PHENYLISOPROPYLHYDRAZINE, BETA-PHENYLISOPROPYL HYDROZINE HCl, See: Catron (Lakeside)

PHENYLKETONURIA THERAPY, See: Ketanil, Pow. (Merck Sharp & Dohme) Lafenolac (Mead Johnson) Phenistix Reagent Strips (Ames)

PHENYLMERCURIC ACETATE. (Various Mfr.). Bot. 1 lb., 5 lb., 10 lb. See: Contra Creme. (Contra)

W/Aluminum caprylate, urea, citric acid, boric acid, lactose & dextrose. See: Mand-Serts, Tab. (Wintersmith).

W/9-Aminoacridine HCl, methylbenzethanol chloride, succinic acid, allantoin, lactose. See: Vagilob, Vag. Tab. (Ascher)

W/9-Aminoacridine HCl, methylbenzethanol chloride, tyrothricin, urea, succinic acid, lactose. See: Trisert Vaginal, Tab., Pow. (Ulmer)

W/9-Aminoacridine HCl, sulfacetamide, benzalkonium chloride. See: Kenhex, Vaginal Preps. (Kendall)

W/9-Aminoacridine HCl, tyrothricin, urea, lactose & sod. lauryl sulf. See: Trinalis, Vag. Supp. (Webster)

W/Copper undecylenate, zinc undecylenate & undecylenic acid. See: Di-Spor, Oint (Wintersmith).

W/Hydrocortisone acetate, tyrothricin, 9-aminoacridine HCl, polysorbate-80, urea, lactose. See: Aquacarl, Vag. Supp. (Webster)

W/Methylbenzethanol chloride, 9-aminoacridine NCl, succinic acid, allantoin & lactose. See: Vagilobs, Tab. (Ascher).

W/Naphazoline hydrochloride See: Clasol, Nasal Spray (Vitamax Corp.)

PHENYLMERCURIC BORATE, N.N.R. 1952; A.D.R. (F. W. Berk) Pkg. Custom packed

W/Benzyl alcohol, benzocaine & butyl p-

aminobenzoate See: Dermathyn, Orn
(Flint Eaton)
W/Sad pentachlorophenolate See: Mertricone,
Liq (Rorer).

PHENYLMERCURIC LACTATE. (No Mer cur
rently lists)

W/Ephedrine lactate, dextrose, tyrothricin
& cetyl trimethyl ammonium See Gluco
Thricil, Sol (Farke, Davis)

***PHENYLMERCURIC NITRATE, N F, N N D**
1961; ADR (Merphenyl Nitrate Phen
merayl Nitrate)

A P L.—Oint 1:1500, 1 oz, 4 oz, 1 lb.
Chicago Pharm.—Laz w/benzoate Bot
100s, 1000s Ophth oint, 13000, Tube
1/2 oz Sol, 1:20,000, Bot 1 pt, 1 gal
Vaginal supp., 1:5000, Rox 12s

See: Merphene (Barry Labs)
Merphenyl nitrate (Hamilton)
Phe Mer Nile, Prep (Massengill)
Phenolca, Sol (Chicago Pharm)
Phenital, Sol & Oint (Chicago Pharm)
Phenmerzyl Nitrate (Pilman Moore)

W/Undecylenic acid See: Merlenate, Oint
& Paw (Chicago Pharmacal)

PHENYLMERCURIC OLEATE. (No Mer cur
rently lists)

W/Undecylenic acid See: Mercene, Oint
(Maltbie)

PHENYLMERCURIC PICRATE. (Metal salts)
Pig Custan packed,
Urei Germicide

W/Picric acid, N N R 1952 See Merphenyl
Picrate W/Picric Acid, Tr (Hamilton)

a-PHENYLPHENOL
See: Ampul (Lehn & Fink)
Phenolor (Squibb)
Lysal (Lehn & Fink)
O Syl (Lehn & Fink)

See: Sol'n of (Darsey)
a-PHENYLPHENOL MERCURIC NITRATE.
See: Hydrophen, Oint (Amire).

**PHENYLPROPANOLAMINE HYDROCHLO-
RIDE, N N R 1957.** 2 Amino T phenyl-T-
propanol (Mydrilaine)

See: Makrion (Haack)
Panamine (Metro Lab)
Propadrine Hydrochloride (Merck Sharp &
Dohme)

PHENYL PROPYL ETHYLAMINE.
See: Antispasmine (Raymer)
Pentrapine (C D Smith)
Pralenil (C D Smith)

***PHENYLPROPYLMETHYLAMINE HYDRO-
CHLORIDE, N N D 1962** See: Yonedrine
HCl, (Merrell)

***PHENYL SALICYLATE, N F, Tab, N F** See:
Solal (Various Mfr.)

PHENYRAMIDOLNCl 2(B hydroxyphenethyl
*Generic name recognized by N F, U S P or A.M.A.

amra)-pyridine HCl See Analexin (Irwin
Nersler)

PHENYL-TERT-BUTYLAMINE. See Phenter-
mine

***PHENYLOXAMINE** o-Benzylphenyl 2
dimethylaminoethyl ether

**PHENYLOXAMINE W/Belladonna pow
der, ethaverine HCl, phenobarbital, buta
barbital sod, pentobarbital sod, secobarbi
tal sod** See: Tri-Synar Plus, Tab (Armour
Labs)

W/Narcotine, NH₄Cl in a thixotropic sur
pensan See Narcol, Liq (Barre)

**PHENYLOXAMINE DIHYDROGEN CIT-
RATE.** See Bristamin, Prep (Bristol Labs)

W/Acetophenetidin, salicylamide, caffeine
See: Acheore, Tab (Philadelphia Ampoule)

W/Acetyl p aminophenol, salicylamide, dl
amphetamine phosphate, metoprine See:
Acogesc, Cap (Strassenburgh)

W/Acetyl p aminophenol, salicylamide, Vit
C, citrus bioflavonoids, caffeine, racemic
amphetamine sulfate See Euphenex, Tab
(Lloyd, Dabney & Westerfield)

W/Acetaminophen, phenylpropanolamine
HCl, pyrriamine maleate methapyrilene
HCl, phenylephrine HCl, chlorpheniramine
maleate See: Leyahist, Caps (Leyden Labs)

W/Ammonium chloride, lpecac fl ext, Sod
citrate & menthal
See Bristalin, Liq (Bristol)

Dorcal, Cough Prep (Smith Darsey)

W/A P C See: Talaxidyne Tab (G F Harvey)

W/A P C & procaine penicillin G
See: A P Citlin, Tab (White)
Bristopen, Tab (Bristol)

W/Belladonna, Ethaverine HCl See: Tri
Synar, Tab (Armour)

W/Calamine & zinc oxide See: CZO w/
Antheslamine, Lat (Massengill)

W/Chlorpheniramine, phenylephrine HCl
phenylpropanolamine HCl See: Naldecob
(Bristol)

W/Chlorpheniramine maleate, d-ampheta
mine sulfate See: An Du Hist TT, Tabr
(Phillips Roxane)

W/Chlorpheniramine maleate, pyrriamine
maleate, phenylephrine HCl, phenylpro-
panolamine HCl See: Trislocamp, Tab, Liq
(Tallby Navon)

W/Chlorpheniramine maleate, rocephedrine
HCl See: Algic Tab (Spencer)

W/Codaine, chlorpheniramine, ephedrine
guaicol carbonate See: Omnil Tuss (Stras-
seburgh)

W/Dexamethasone See: Taldex, Tabs
(Pitman Moore)

W/Dihydrocodeinone bilartrate, racemic am
phetamine sulf, sod citrate pot guaicol

- sulfonate. See: Neotuss-PT, Liq. (Lloyd, Dabney & Westerfield)
- W/Dihydrostreptomycin sulf. See: Dihydrostreptomycin sulf. W/Bristamin, liq. (Bristol)
- W/Glyceryl guaiacolate. See: Guaiatussin, Syr. (White)
- W/Ipecac R. ext., ammonium chloride, Sod. citrate & menthol.
See: Bristalin, Syr. (Bristol Labs)
Dorcol. (Smith-Dorsey)
- W/Methampyrone, *n*-acetyl-*p*-aminophenol, adiphenine HCl. See: Anfac, Caps. (Pharmalac)
- W/Methapyrilene HCl & pyrilamine maleate. See: Trihistin, Tab. (Aicher)
- W/Neomycin sulf., tyrothricin, bithionol, zirconium oxide, dipiperdon HCl. See: Allergesic, Lot (Bryant)
- W/Penicillin potassium alpha-phenoxethyl, phenylephrine HCl, phenylpropanolamine HCl, chlorpheniramine maleate, acetyl-*p*-aminophenol. See: Syndecac, Tab. Sol. (Bristol)
- W/Penicillin G potassium, acetophenetidin, salicylamide & buffered with calcium carbonate. See: Elpafec, Tab. (Paich)
- W/Phenylpropanolamine HCl. See: Conex-DA, Dual Action Tab.
- W/Phenylpropanolamine HCl, glyceryl guaiacolate, pyrilamine maleate, salicylamide. See: Resprol, Syr. (Tilden)
- W/Prednisone, chlorpheniramine maleate, ephedrine sulfate. See: Esteril, Tab (Enley-Scott)
- W/Pyrilamine maleate, methapyrilene fumarate, aspirin, acetophenetidin, caffeine. See: Trish-A-lin, Tab. (Walker Corp)
- W/Pyrilamine maleate, meth-dia-mer sulfonamide, glyceryl guaiacolate, ephedrine sulfate. See: Sullusin, Susp. (Tilden)
- W/Pyrilamine maleate & propenpyridamine maleate. See: Maltihist, Cap. (Smith-Dorsey)
- W/Pyrilamine maleate, sulfa meth-dia-mer, glyceryl guaiacolate, ephedrine sulfate. See: Sultussin, Tab. (Tilden)
- W/Pyrilamine maleate, phenylpyramine HCl. See: Triamine, Tab. (Lemman)
- W/Pseudoephedrine HCl. See: Ephamine (Spencer Labs)
- W/Salicylamide, acetophenetidin & Pot. penicillin G. See: Elpafec, Tab. (Paich)
- W/Salicylamide, phenacetin, caffeine. See: Medache, Tab. (Organon)
- W/Salicylamide, phenylephrine HCl, *N*-acetyl-*p*-aminophenol. See: Absinol, Tab. (Lemman)
- W/Scopolamine HBr, niacinamide, benzocaine. See: Hyloxine, Tab. (Tutag)
- W/Secobarbital. See: Hytran. (Wampole)
- * Generic name recognized by N.F., U.S.P. or A.M.A.
- W/Tetracycline base, *n*-acetyl-*para*-aminophenol. See: Tetrex AP (Bristol)
- W/Theophylline, *d*-isoprenaline HCl. See: Astec, Tabs. (Arnar-Stone)
- W/Thiamine HCl, adiphenine HCl, pyridoxine HCl. See: Tapp, Caps. (Pharmalac)
- W/Zirconium oxide. See: Zircon, Lat. (Bristol)
- W/Zirconium oxide, colamine, camphor, phenyl, benzocaine. See: Histanium, Sol. (Columbus)
- W/Zirconium oxide, oxyquinaline sulfate. See: Ivy-Ess, liq. (Lloyd, Dabney & Westerfield)
- PHENYLOLOXAMINE NBr.** W/Aspirin, phenacetin, caffeine, tetracycline phosphate. See: Tetrex APC W/Bristamin, Cap. (Bristol)
- PHENYLOLOXAMINE RESIN.** See: Mixturex, Cap. (Strassenburgh)
- W/Dihydrocodeine resin complex. See: Tussionex, Prep. (Strassenburgh)
- PHENYLELIN.** (Crookes-Barnes) Zinc sulfate 1/2% and phenylephrine HCl 0.08%/Lacrivial 15 cc.
Use: Astringent decongestion.
- PHENY-PAS-TIBAMIN.** (Purdue Frederick) Phenyl *p*-methylsalicylate 0.5 Gm./Tab, or tablespoonful of powder, Tab. Sol. 500s, 5000s, Pow. Pkg. 1, 5, 25 lbs.
Use: Treatment of tuberculosis.
- * **PHENYRAMIDOL HCl.** *o*-(2-pyridylaminoethyl) benzyl alcohol. See: Analgin AF, Tab. (Irwin, Nelsler)
- * **PHENYRAMIDOL SALICYLATE.** See: Analgin Syrup (Nelsler)
- PHENYRAMIDOL W/Aluminum Aspirin.** See: Analgin AF, Tab. (Irwin, Nelsler)
- * **PHENYTHILONE.** 2-ethyl-2-phenyl-3,5-dihloromorpholideane. See: Thimarlone (Merck Sharp & Dahme)
- PHENYTOIN, SOLUBLE, U.S.P., N.N.D. 1959.** See: Diphenylhydantoin Sod., Prep. (Various Mfr.)
- PNITHARBITAL.** 5,5-diethyl-1-phenylbarbituric acid.
- PHETOGNIC.** (Myers-Carter) Estrogenic sub 20M w/w, *d*-amphetamine sulf. 20 mg./cc Vial 10 cc.
- PHISOAC.** (Winthrop) Fresh-colored cream containing colloidal sulfur 6%, resorcinol 1.5%, hexachlorophene 0.3%, *O*-phenylphenol 0.3%, alcohol 10% in a stable cream base.
Use: Treatment of acne.
- PHISOAC.** (Winthrop) Colloidal sulfur 6%, resorcinol 1.5%, hexachlorophene 0.3%, orthophenylphenol 0.3%, alcohol 10% (w/w) in fresh-colored, greaseless base Tabs 1 1/2 oz.
Use: Acne.

- PHISODERM.** (Winthrop Labs) Sod. octylphenoxethoxyethyl ether sulfonate Emulsion regular, oily or dry type Bot 5 fl oz, 12 fl oz, 1 gal Wall dispenser 10 fl oz
Use: Topically, detergent
- PHISOHEX, H N D 1962** (Winthrop Labs) Sod octylphenoxethoxyethyl ether sulfonate & hexachlorophene 3% Emulsion, Bot 16 fl oz, 1 gal; hand dispensers 3 fl oz; squeeze bot 5 fl oz; wall dispensers 10 fl oz; pedal dispensers 30 fl oz
Use: Antiseptic detergent
- PHISOHEX-PHISOAC COMBINATION.** (Winthrop) PhisoHex 5 fl oz, PhisoAc 1 1/2 oz / Plastic Bot 5 fl oz; Tube 1 1/2 oz
Use: Acne
- P-H-H ISOPTO.** (Alcon) Polymyxin 16 250 u, hydrocortisone 1.5%, neomycin 5 mg, and methylcellulose 0.5%/cc Drop tainer 5 cc.
Use: Ophthalmic therapy
- PHOAM CLEANSE PAC.** (Doak) Detergent cake W/bithional, sulfur and salicylic acid Box 9 cakes
Use: Skin cleanser, acne
- PHOB.** (Brewer) Phenobarbital Gel Et Tabs 1/2 gr or 1/4 gr Bot 100s, 500s, 1000s
Use: Sedative
- PHOBEX, H N D 1962** (Lloyd, Dabney & Westerfield) Benactyzine HCl 1 mg or 5 mg / Tab Bot 100s
Use: Treatment of phobias, anxiety and depression
- *PHOLCOLINE** The narcotic antitussive compound 3 (2 morpholinoethyl) ether of morphine
See Ethnine (Lab for Pharmaceutical Development, Inc)
Simplex (Purdue Frederick)
- PHORSIX.** (Texas) Oil in water emulsions of the nonionic type adjusted to a pH of 4.16 Cream, 4 oz and 1 lb jars Lotion, 8 oz, 1 pt btl
Use: For skin lubrication
- *PHOSCOLIC ACID** (Lederle Labs) 2,2-phosphinocadiacetic acid
Use: Adjuvant
- PHOS-FE-CAC.** (Sherman) Iron cocodylate 1/4 gr, Sod cocodylate 1/4 gr, strychnine cocodylate 1/125 gr, Sod glycerophosphate 2 gr, chlorobutanol 0.5%, procaine HCl 1%/cc Amp 1 cc, Box 10s, 100
- PHOSFO-HEPRIN** (Schiffelin) Epinephrine phosphate Sol 1,000, Bot 1 fl oz
Use: Sympathomimetic agent
- PHOSPHAL-D.** (Lanett) Dicalcium phosphate anhyd 330 mg, vit D 333 u/cap Bot 500s, 1000s
- *PHOSPHALJEL, N H D 1962** (Wyeth) Aluminum phosphate gel Bot 12 oz
Use: Orally (2-4 dr m p); antacid
- PHOSPHATABS TEST KIT.** (Warner-Chilcott) Phospholase tabs
Use: Diagnosis of cancer and jaundice
- PHOSPHATASE.** See: Phosphotabs Test Kit (Warner Chilcott)
- *PHOSPHENTASIDE.** See: Adenosine-5-monophosphate
- PHOSPHO-LECITHIN W/Strychnine** (Wampole) Bot 1 pt, 1 gal
Use: Fatty fat metabolism
- PHOSPHOLINE-IODIDE, N N D 1962** (Campbell Pharm) Echothiophate iodide Diethoxy phosphorothiocholine iodide 6.25 mg dry pow bot with 5 cc diluent Also avail to hospitals D 125 Gm bot
Use: (Anticholinesterase) Eye drops of D 125% or 0.25% for glaucoma
- PHOSPHOLIPIDS SOY.** See: Granulestin, Pow (Associated Concentrates)
- PHOSPHORATED CARBOHYDRATE SOLUTION.** See: Emelrol, Liq (Kinney)
- PHOSPHO-SODA, N H R 1950** (C B Fleet) Sod biphosphate 48 Gm, Sod phosphate 13 Gm / 100 cc Bot 2 1/2, 6, 16 fl oz
Use: A mild saline laxative
- PHOTOPOPE, N H D 1962**
- *PHTHALYLSULFACETAMIDE, N F, Tab, N F**
See Enterosulfon, Tab (Consol Midland)
Talsigel, Susp (Squibb)
Talsutin, Tab (Squibb)
Thalajen, Tab (Jenkins Labs)
Thalomyd, Tab (Schering)
Thalisl, Tab (Massengill)
W/Koolin & pectin
See: Thalajen, Prep (Jenkins Labs)
Thalocin, Susp (Abbott)
Thalita, Susp (Abbott)
W/Neomycin See: Neothalidine, Tab (Merck Sharp & Dahme)
W/Pectin & albumen linnate See: Mucamide, Cap (Warren Teed)
W/Polymyxin B sulfate & polyamine methylene resins See: Resion PMS, Liq (Notional Drug)
W/Sod Carboxymethylcellulose, Inorganic Salts & Glucose See: Kolacet, Susp (Danley-Evans)
- *PHTHALYLSULFATHIAZOLE, U S P, N N D 1962 A D R 1951; Tab U S P** Thalozone
See: Crema Thalidine, Susp (Merck Sharp & Dahme)
Sulfathalidine, Tab (Merck Sharp & Dahme)
W/Neomycin sulfate See: Sulfathalidine
W/Neomycin, Tab. (Merck Sharp & Dahme)

*Generic name recognized by N F U S P or A M A

PHYATROMINE-H. (Kremers Urban) Physostigmine salicylate 0.6 mg., 1-hyoscyamine HBr 0.3 mg./cc. Amp. 1 cc., 25s, 100s Vial 10 & 30 cc.

Use: Skeletal muscle spasm & associated pain.

PHYATROMINE-H PANTODREN-100.

(Kremers Urban) Phytotromine-H, Pantodren-100. Vial 30 cc., Box. 3.

PHYATROSUL. (New Mexico Pharm) Physostigmine salicylate 1 mg., atropine sulfate 0.65 mg., sodium bisulfite 0.1%, benzyl alcohol 2%, benzoic acid 0.35%/cc. Vial 30 cc.

Use: Control of muscle spasm in arthritis.

PHYLCARDIN. See: Aminophylline (Various Mfr.).

PHYLIPHEN. (Kremers-Urban) Aminophylline $1\frac{1}{2}$ gr., phenobarbital $\frac{1}{4}$ gr., Pot. Iodidn 1 gr./Tab. Bot. 100s, 500s, 1000s. Also Phyliphen-3 contains aminophyllin 3 gr., Pot. Iodidn 2 gr.

Use: Coronary vasodilator, diuretic & antispasmodic.

PHYLICIN. (Knoll) Theophylline-Cal. salicylate. Pow., Bot. $\frac{1}{2}$ oz., 4 oz. Tab. 4 gr., Tube 20s, Bot. 100s, 500s.

Use: (3-4 daily) Diuretic & myocardial stimulant.

PHYLLINDON. See: Aminophylline (Various Mfr.).

PHYLLOQUINONE.

See: Phytonadione, U.S.P.

Vit. K-1. (Various Mfr.)

PHYSEPTONE. See: Methedone

PHYLLOQUINONE. 2-Methyl-3-phytyl-1,4-naphthoquinone, Vit. K.

PHYLTONE. (Texaphyl Prod.) Pol. hydrogenophytochlorin.

Bulk pow.s Bot. 2 oz.

Lot. (3% in water) Bot. 1, 2 & 8 oz.

Oint. (3% in water-sol. base) Jar $\frac{1}{2}$, 1 & 6 oz.

Cap. ($\frac{1}{4}$ Gm) Bot. 100s.

Use: Decholesterolizing agent (arteriosclerosis, arthritis, allergies).

PHYSOSTIGMINE, ALKALOID. (Eserine)

Merck—Pow., Tube 1 gr. Bot. 5 gr., 15 gr., $\frac{1}{2}$ oz.

Use: Parasympathomimetic agent.

* **PHYSOSTIGMINE SALICYLATE, U.S.P.**

Eserine salicylate.

Burroughs Wellcome—Tabloid, hypodermic, 1/100 gr., Tube 20s.

Durst—Vial (2 mg./cc.) 10 cc.

Merck Sharp & Dohme—Pow., Tube 1 gr., 5 gr., 15 gr.

Parke, Davis—Tab. hypodermic, 1/100 gr., 1/50 gr., Tube 20s, Bot. 100s, 500s.

* Generic name recognized by N.F., U.S.P. or A.M.A.

See: Esromiofin (Crocker-Barnes)

Isopht-Eserine, Ophth. (Alcon)

Mioal (Prof. Pharm.)

Use: Parasympathomimetic agent

W/Aspirin, para-aminobenzoic acid, ascorbic acid, homatropine methylbromide. See: Mobisyl, Tab. (B. F. Ascher)

W/Atropine sulfate.

See: Atrostigmine, Vial. (G. F. Harvey)

Banmiolin (Iso-Sol)

Phyalromine-H, Amp. (Kremers-Urban)

Physotropin, Tab. & Vial (Durrig)

Tromine (Testagar)

W/Atropine sulfate, sodium bisulfite, benzyl alcohol, benzoic acid. See: Phytrosul, Inj. (New Mexico Pharm.)

W/Atropine sulfate, Sod. bisulfite & dextrose. See: Almetstigmine, Vial (Meyer).

W/Amobarbital, mephensin, homatropine methylbromide & salicylamide. See: Myomaphetone, Tab (Testagar)

W/Homatropine methylbromide.

See: Esoprine, Tab. (Ulmer)

Physal, Tab. & Vial. (Central)

W/Homatropine methylbromide, Vit. C, p-aminobenzoic acid & Sod. salicylate. See: P-B-SAL-C W/Esoprine, Tab. (Ulmer)

W/Homatropine methylbromide & Vit. B-12. See: Artha-B-12, Vial (Columbus).

W/Homatropine methylbromide & alpha-tocopherol acetate. See: Artha-E, Vial (Columbus).

W/Homatropine methylbromide, Sod. salicylate, p-aminobenzoic acid & ascorbic acid. See: Paliline, Tab. (Ascher).

W/Hyoscyamine HBr. See: Phyalromine-H, Amp. (Kremers-Urban)

W/Mephensin, homatropine methylbromide. See: Trimitis, Tab. (Bell Pharmacol)

W/Pilocarpin, methylcellulose. See: hopto P-E-S, Soln. (Alcon)

PHYSOTROPIN. (Durrig) Sol., Physostigmine salicylate 1 mg., atropine sulfate 0.6 mg./cc. Vial 30 cc. Tab., Physostigmine salicylate 0.5 mg., atropine sulfate 0.15 mg./Tab. Bot. 100s.

Use: Antispasmodic.

* **PHYTATE SODIUM.** (Squibb) Sod. salt of inositolhexaphosphoric acid.

PHYTIC ACID. Inositol hexaphosphoric acid.

PHYTIN. (Ciba) Cal.-Mag. salt of inositol hexaphosphoric acid, containing Phos. 22%, Cal. 12%, Mag. 1.5% in organic combination. Tab., Bot. 40s.

Use: Orally, mineral supplement.

PHYTOCO. (Drug Products) Thyroid 1/3 gr.,

phytalacca fructus ext. 2 gr., apocynum

- paw 1 gr, pituitary whole desic 1/10 gr / Cap Bot 100s, 1000s
 Use: Obesity
- W/Homatropine methylsalicylate, mephensin, salicylamide & amobarbital See Myamephetane Tab (Testagar)
- PHYTOMEHADIONE**. See Phylloeadione USP
- ***PHYTOHADIODIOL SODIUM DIPHOSPHATE**. Disodium salt of 2 methyl 3 phytol 1 4 naphthohydroquinone O¹, O⁴ diphosphoric acid See AquinNephyton (Merck Sharp & Dahme)
- ***PHYTOHADIONE**, USP Emulsion Sterile USP, Tabs USP NND 1962 Vit K 1 2 methyl 3 phytol 1 4 naphthoquinone See Kanakion Cap Amp (Bache) Mephyton, Amp (Merck Sharp & Dahme) Mono Kay Caps (Abbott)
- PICLOXYDIHE** Name Approved by BPC NN D1 [p Chlorophenylguanidinotormimic dyl]piperazine
- PICRAGOL**, NNR (Wyeth) Silver picrate Cryst, Bot 2 Gm Pow, Bot 5 Gm Vag Supp 0.065 Gm 0.130 Gm Box 12s
 Use: Silver antiseptic trichomonocide
- PICRIC ACID, TRINITROPHENOL**, NF See Silver salt
 W/Butesin See: Butesin Picrate (Abbott)
 W/Phenol & benzyl alc See Picro Benzyl Oint (Ulmer)
- PICRO-BENZYL OINTMENT**. (Ulmer) Picric acid 0.3%, phenol 0.5%, benzyl alcohol 1% Jar 1 lb
 Use: Burn therapy
- ***PICROTOXIN**, NF, NND 1961, ADR Inj, NF Cocculin
 Abbott—Vial 0.3%, 20 cc
 Lilly—Amp (3 mg), 1 cc, 6s, 100s Vial (3 mg / cc) 20 cc
 Use: Respiratory stimulant
- P-I-D** (Lincoln Labs) Ferrous sulfate 33.3 mg sorbitol laurate 150 mg / D 6 cc 30 cc Bot w/calibrated dropper/6s
 Use: Treatment of iron deficiency anemia
- PIG BILE EXTRACT** (Burroughs Wellcome) Tablet 4 gr, Bot 100s
 Use: Orally chologogue
- PIGMENT CEMOGEN SOL**. (Dermatologic Pre Labs) Buffered dihydroxyacetone in quick drying vehicle Bot 1 oz
 Use: Masking areas of vitiligo
- PILO B-1/2, -1, -2, -3, -4** (Broemmel) Pilocarpine nitrate 1/2, 1, 2, 3 and 4% solutions, methylcellulose 1/2%, boric acid sodium citrate buffer system preserved with D 1% chlorbutanol & stabilized with D 1% sod bisulfite Tubes 15 cc
 Use: Treatment of glaucoma
- ***Generic name recognized by NF USP or AMA**
- PILOCAR**. (Smith, Miller & Patch) Pilocarpine HCl D 5%, 1%, 2%, 3%, 4%, 6%, in a methylcellulose vehicle Vial 15 cc
 Use: Therapy of glaucoma
- ***PILOCARPINE HYDROCHLORIDE**, USP, ADR (Merck Sharp & Dahme; Mallinckrodt) Pilocarpine muriate
 Use: 0.5 to 2.0% sol topically as a miotic
- ***PILOCARPINE HYDROCHLORIDE**, USP, ADR (Lilly) Tab hypodermic 1/4 gr, Tube 20s; Bot 100s, 500s, 1000s, 5000s
 Use: Parasympathomimetic agent
 See Isapta Carpine, Ophth (Alcon)
 Pilocar (Smith, Miller & Patch)
 Pilacel 1% & 2% Sol (Professional Pharm)
 Pilavisc (Crookes Barnes)
- W/Physostigmine Salicylate See: Bonmiotin (Crookes Barnes)
- W/Eserine salicylate, methylcellulose See Isopto P ES Soln (Alcon)
- ***PILOCARPINE HIATRATE**, USP, ADR (Merck Sharp & Dahme) Pow, Bot 5 gr-15 gr 1/2 oz 1 oz
 Wyeth—Tab 5 mg Bot 100s
 Use: Parasympathomimetic agent
 W/Methylcellulose See Pilo No 1 & 2, Sol (Broemmel)
- PILOCEL** (Professional Pharm) Pilocarpine HCl 1/2, 1, 2, 3, & 4% buffered in methylcellulose base w/Zephiran Bot w/dropper 15 cc
 Use: Ophthalmic therapy, miotic
- PILOFRIN OPHTHALMIC DROPS** (Allergan) Pilocarpine nitrate buffered sol Sol (1% or 2%), Bot 1/2 oz
 Use: Miotic
- PILOID SPECIAL** (Suliff & Case) Tannis acid 10%, phenol 2%, benzocaine 1%, camphor 2%, stramonium ext 2.5%, petrolatum laealen base qs Oint Tubes 1 oz Jar 1 lb
 Use: Antiseptic w/local anesthetic
- PILOIN** (Tilden) Extract stramonium, benzocaine, boric acid, tannic acid, extract hydnosis extract witch hazel, pine oil iodine Applicator tube 1/2 oz
 Use: Hemorrhoids and rectal irritation
- PILOMIOTIN** (Crookes Barnes) Pilocarpine HCl stabilized solutions D 5 to 6% in Vials 15 cc and Dropperettes
 Use: Miotic
- PILOLONOL** (Seydel) Benzyl alcohol 79 parts benzocaine 20 parts, azyquinoline benzoate 1 part Sol Bot 25 cc
 Use: Cysts & pilonidal cysts
- PILOVISC**. (Crookes Barnes) Pilocarpine HCl 0.5% to 4% W/methylcellulose/Lacrival 15 cc
 Use: Miotic

- PIMARICIN.** See: Myprozina. (Lederle)
- ***PIMETINE HYDROCHLORIDE.** (Irwin Meisler) 4-benzyl-1-(2-dimethylaminoethyl)piperidine hydrochloride.
Use: Cardiovascular drug.
- PIMINODINE.** See: 1-[3-phenylaminopropyl]-4-phenylpiperidine-4-carboxylic acid ethyl ester.
- PIMINODINE ETHANESULFONATE.** Ethyl-4-phenyl-1-[3-(phenylamino) propyl] piperidine-4-carboxylate ethanesulfonate. See: Alvadine (Winthrop Lab)
Alvadine (Winthrop Lab)
P-1-N. (Lonnell) Isoinoxid 50 mg., Vit. B-6 1 mg./Tab. Bot. 100s, 1000s. Forte—[isoxiazid 100 mg., Vit. B-6 5 mg./Tab]
Use: Treatment of tuberculosis
- PINCETS.** (Walker Labs.) Piperazine phosphate 250 mg./tab. Bot. 50s.
Use: Treatment of pinworm or roundworm infections.
- PINEX ANTI-NISTAMINE, (Pinex)** Pyranisamine molecule. Tab. 25 mg., Bot. 32s.
Use: (I s.l.d., p.c. & h.s.) Common cold & hay fever.
- PIPROU, (Carrtane)** Piperazine phosp. 250 mg./Tab. Bot. 56s, 500s
Use: Pinworm, roundworm.
- PINSIRUP, (Walker)** deleted Piperazine phosphate equiv. to piperazine hexahydrate 250 mg./dr. Syr. Bot. 8 oz.
Use: Treatment of pinworm or roundworm infections
- PINUS ALBA, EUCALYPTOL & HONEY COMPOUND, (Merck Sharp & Dohme)** Chloroform, white pine bark, wild cherry bark, squill, senega, ipecac, sanguinaria, eucalyptol & honey. Bot. 1 gal.
Use: Cough.
- ***PIPAMAZINE, N N D 1962.** 10-[(3-(4 Carbamoylpiperidino) propyl)-2-chlorophenothiazine. See: Mornidine, Tab. & Amp. (Searle)
- PIPANOL, (Winthrop Labs.)** (Trihexyphenidyl HCl) alpha-phenyl-1-cyclohexyl-3-piperidyl-1-propanol HCl. Elixir 2 mg./4 cc. Bot. 1 pt., 1 gal. Tab. 2 mg., 5 mg., Bot. 100s, 1000s.
Use: Orally; antispasmodic in parkinsonism
- PIPAZETHATE.** See: Theratuss, Tabs. (Squibb)
- PIPENZOLATE METHYL BROMIDE, (Lakeside Labs)** Changed to Pipenzolate bromide.
- ***PIPENZOLATE BROMIDE, N N D. 1962.** 1-Ethyl-3-piperidyl benzilate methylbromide. See Pipal (Lakeside)
W/Acetaminaphen, phenobarbital. See: Pediatric Pipal Antipyretic (Lakeside)
W/Antacids. See: Promyl. (Crookes-Barnes)
- ***PIPERACETAZINE.** 2-acetyl-10-[3-[4-(b-hydroxyethyl)-piperidino]propyl]phenothiazine See Guide (Pitman-Moore)
- * Generic name recognized by N F, U S P or A M A
- PIPERAT, N.N.D. 1961.** (Lincoln) Piperazine tartrate 100 mg./cc. Bot. 1 pt. Tab. 250 mg. Bot. 200s, 1000s.
Use: Pin worm & roundworm treatment in children & adults.
- PIPERAZATE, (Leeming)** Piperazine phosphate 500 mg./water. Bot. 28s.
Use: Treatment of pinworm or roundworm infections.
- ***PIPERAZINE CALCIUM EDATHAMIL, N N D. 1962.** See: Perin, Syr. (Endo)
- ***PIPERAZINE CITRATE, U.S.P.** Piperazine Citrate Telsa, Syr. & Tabs. U.S.P., N N D. 1962.
See: Antepar, Syr. (Burroughs Wellcome)
Anthelcol, Tab. (Cale)
Multifuge Citrate, Prep. (Blue Line)
Ozicide, Syr. (Breon)
Parasac, Liq. (Tulog)
Pipizan Citrate, Syr. (Merck Sharp & Dohme)
Vermogo, Syr. (Lloyd, Dabney & Westfield)
- ***PIPERAZINE ESTRONE SULFATE, N.P., N N D 1962;** Tabs, N.F. See: Sulestrex Piperazine, Tab. (Abbott)
- PIPERAZINE GLUCONATE.** See: Vermizene, Syr. (Chicago Pharm)
- PIPERAZINE NEXAHYDRATE,** See: Antholazine, Syr. (Bowman)
Parid, Liq. (Buffington)
Pediamic, Elix. (Columbus)
Bandelim, Sol. (Tilden)
- PIPERAZINEPENTANOL, (Parke, Davis)** 4-[o-Propylthiophenyl]-1-piperazinepentanol. Under study.
Use: Ataractic treatment of anxiety
- ***PIPERAZINE PHOSPHATE,** See: Perin, Water & Syr. (Endo)
Pincels, Tab. (Walker Labs.)
Pinow, Tab. (Carrtane)
Pinsirup, Syr. (Walker)
Piperazate, Water. (Leeming)
- ***PIPERAZINE TARTRATE, N N D. 1962** See Piperate, Sol. (Lincoln)
- ***PIPERAZINE TRICHLORPHATE, (Parke, Davis)** Piperazine di[2,4,5-trichlorophenoxyde].
Use: Anthelmintic (nematodes).
- ***PIPERIDINE PHOSPHATE, (Abbott)** Piperidine phosphate.
Use: Psychiatric drug
- PIPERIDINOETHYL BENZILATE NCl.** See Sycatol, Tab. (Reed & Carnrick)
W/Scopolamine methylbromide, magnesium hydrazide, aluminum hydroxide See: Madutrol, Tab (Reed & Carnrick)
- PIPERIDONE.** See: Sedulan, Syr. (Roche)

- * **PIPERIDOLATE HCl, N N D 1962** See: Dartil (Lakeside)
- W/Ascorbic acid, hesperidin complex See: Doctil OB Tab (Lakeside)
- W/Cellulolytic enzyme amylolytic enzyme proteolytic enzyme, pancreatin, tauracholic acid See Dartilase, Tabs (Lakeside)
- * **PIPEROCAINE HYDROCHLORIDE, USP** N N R 1954, A D R: 1nj, Ophth Oint, USP 3 Benzoxyl-1-(2 methylpiperidino) propane HCl See: Metyroine, Prep (Lilly)
- * **PIPEROPNYLLIN, (M. H. Smith) Theophylline** 200 mg w/piperazine/Tab Bot 100s, 1000s
Use 1 q i d; theophylline therapy.
- * **PIPEROXAN HYDROCHLORIDE, N N D 1959** 2 (1 Piperidylmethyl) 1,4 benzodioxan HCl Faurneau 933 Benzodioxone See: Benadoine Hydrochloride, Amp (Merck Sharp & Dahme)
- * **PIPERPENIDOL HCl, 5 methyl-4 phenyl 1 (1 piperidyl) 3 hexanol HCl** See: Reline HCl (Merck Sharp & Dahme)
- * **PIPETHANATE HYDROCHLORIDE, N N D 1962** 2 (1 piperidino)ethyl benzilole hydrochloride See Sytolol (Reed & Cornick)
- * **PIPIZAN CITRATE SYRUP, N N D 1962** (Merck Sharp & Dahme) Piperazine citrate equiv to piperazine hexahydrate 0.5 Gm / 5 cc Bot 4 oz, 1 pl, 1 gal
Use: Treatment for pinworms & roundworms
- * **PIPRADROL HYDROCHLORIDE, NP, N N D 1962** Tabs NP α Diphenyl 2 piperidine methanol HCl α (2 Piperidyl) benzhydral HCl See Merotran Tab (Merrell)
- W/Hesperidin methyl chalcone, vitamins, Ioprotropics, iron, essential minerals See: Peptalin, Liq (Walker Labs)
- * **PIPTAL, N N D 1962** (Lakeside Labs) Pipenzolate methobromide N ethyl 3 piperidylbenzilata methobromide Tab 5 mg, Bot 100s
Use: 1 t i d, o c & 1 h s; treatment of peptic ulcer
- W/Doctil See: Trifol, Tab (Lakeside)
- * **PIPTAL PEDIATRIC ANTIPYRETIC (Lakeside)** Aretaminophen 60 mg pipenzolate methyl bromide 2 mg phenobarbital 3 mg / 6 cc Dropper bulbs of 3D cc w/droppers calibrated to deliver D 6 cc
Use: Analgesic & antipyretic
- * **PIPTAL PEDIATRIC, (Lakeside) Piptal** 4 mg, phenobarbital 6 mg / cc Bot w/calibrated dropper 3D cc
Use: Relief of colic, pyloric spasm, regurgitation & other functional gastrointestinal disorders
- * Generic name recognized by N F USP or A.M.A.
- PIRAMINE, (Dumas Wilson) Pyrimine** maleate 25 mg, as Iron 3.5 gr, caffeine 0.5 gr, acetophenetidin 2.5 gr / Cop Bot 100s, 1000s
Use: Orally (1 t i d); analgesic w/anti histamine
- * **PIRAYON, (Walker Labs) Pipradrol HCl** 2 mg, Vit. A 5000 u, D 500 u, B 1 5 mg, B 2 3 mg, Col pantothenate 2 mg, C 25 mg, niacin 5 mg, B-6 5 mg, B-12 2 mgr, folic acid 0.5 mg / Sust release Cop Bot 50s, 1000s
Use: Stimulant w/Vitamin supplement
- * **PIRIO, (Misemer) Phenylazo-diaminopyridine HCl** 1 1/2 gr / Tab Bot 100s, 1000s
Use: Urinary analgesic
- * **PIRIDAZOL, See: Sulfopyridine (Various Mfr)**
- * **PIRISOCAINE HCl, Beta-(2 Piperidyl) ethyl α -aminobenzoate** See: Lucoline HCl
- * **PIRITON, See: Chlorpheniramine**
- * **PIRITRAMIDE (Registered in Great Britain)** 4 (4 Corbamoyl-4 piperidinopiperidino-2,2 diphenylbutyranitrile
- * **PIROCAINE, (Sutliff & Cose) Benzocaine** benzoate 1%, picric acid 0.5%, petrolatum compound Jar 1 lb, 5 lbs Ointment
Use: Antiseptic and local anesthetic ointment
- * **PIROMEN (Plant Eaton) Sterile nonprotein & nonophyloctogenic bacterial prepared in stable colloidal dispersion** Vial (4 gamma per cc) 10 cc, (10 gamma per cc) 10 cc
Use: Allergic disorders
- * **PIRSEAL (Testogor) Aspirin** 5 gr / Enteric coated Tab Bot 100s
Use: Analgesic
- * **PITAYINE, See: Ouhnidine (Various Mfr)**
- * **PITOCIN, N N R 1952 (Parke, Davis)** Oxytocin Amp (5 u) 0.5 cc (10 u) 1 cc, Box 10s
Use: Stimulate uterine contraction
- * **PITRESSIN, N N R 1950 (Parke, Davis)** Vasopressin Methahypophamine Amp D 5 cc (10 pressor u) & 1 cc (20 pressor u) Box 10s
Use: Prevention & control of abdominal distention
- * **PITRESSIN TANNATE IN OIL, N N D 1962 (Parke, Davis)** Vasopressin Tannate Beta hypophamine tannate Amp (5 pressor u) 1 cc, Box 10s
Use: Where long continued action of Pitressin is desired
- * **PITUITARY ANTERIOR GONADOTROPINS** See: Andutrops, Vial (Rowell)
Biotropin, Vial (Biophysic)
Chimadin, Vial (Myers Carter)
F S L, Vial (Myers Carter)

Pituitary Gonadotropin, Amp. (Squibb)

Tropinote, Vial. (Mallord)

PITUITARY, ANTERIOR. The anterior lobe of the pituitary gland supplies protein hormones classified under following headings.

See preparations of:

Antuitrin (Parke, Davis)

Corticotropin, U.S.P.

Gonadotropin.

Growth Hormone.

Lactogenic Factor.

Luteotrophin (Parke, Davis)

Thyrotropic Factor.

W/Thyroid, adrenal cortex fresh gland, ovarian substance fresh gland, posterior pituitary. See: Mestrocina, Vial. (Mastro)

PITUITARY, ANTERIOR, & OVARY COMPOUND.

Breon—Amp. 2 cc., Box 12s, 25s. Vial 30 cc.

Lincoln—Vial 30 cc.

PITUITARY GONADOTROPIN. (Squibb)

Pituitary gonadotropin prepared from horse anterior pituitary gland. Vial 125 rat u., w/3 cc. amp. of sterile isotonic sol. as diluent.

Use: S.C. & I.M.; gonadotropin therapy.

PITUITARY GONADOTROPINS.

See: Biotropin, Vial (Biophysical)

Tropinate, Vial (Mallord).

PITUITARY, POSTERIOR, INF., U.S.P.

Pituitary Extract, Posterior

See: Infundin, Amp. (Burroughs Wellcome)

Pituitin, Obstetrical (Parke, Davis)

Pituitrin, Surgical (Parke, Davis)

PITUITARY, POSTERIOR, HORMONES.

(a) Vasopressin. Pressor principle, β -hypophamine, postlabn-Y.

See: Pitressin, Amp. (Parke, Davis)

Pitressin Tannate (Parke, Davis)

(b) Oxytocin. Oxytocic principle. α -hypophamine, postlabn-O.

See: Oxytocin, Inf., U.S.P., Pitocin, Amp. (Parke, Davis)

Syntocinon, Amp. (Sandoz)

PITUITRIN, N.N.R. 1950. (Parke, Davis)

Pituitary, posterior ext. Amp., obstetrical (5 u.) 0.5 cc.; (10 u.) 1 cc.; surgical (20 u.) 1 cc., Box 10s.

Use: Obstetrical, to stimulate uterine contraction. Surgical, prevention & control of abdominal distention.

* **PIVAZIDE.** N-Benzyl-N'-pivaloylhydroxamine.

Tersavid.

PIX ALBA. (Mallord) Purified colorless coal tar. Oint. 1 oz., 1 lb.

Use: Dermatology.

PIX CARBONIS. See: Coal Tar, U.S.P.

PIX-OEL. (Pitman-Moore) 5% Crude coal

* Generic name recognized by N.F., U.S.P. or A.M.A.

tar & 10% zinc oxide in vanishing base.

Jar 2 oz., 1 lb. Tube 1 oz.

Use: Topically; antipruritic.

PIX JUNIPERI. See: Juniper Tar, U.S.P.

PIXONAT OINT. (Dermatologicals, Inc.)

Micronized whole crude coal tar 3%, with

zinc oxide and corn starch in homogenized

emollient base. Jars 2 oz., 1 lb.

Use: Chronic dermatoses.

PIX-UNO IMPROVED. (Sulliff A Case) Zinc

oxide 10%, crude coal tar distillate 10%,

benzocaine 1%, washable type base q.s.

Oint. 1 lb.

PLACENTA. See: Gonadotropins, Chorionic.

PLACENTAL EXTRACT. See: Cantalone

(Paschall)

PLACIDYL, N.N.D. 1962. (Abbott) Ethyl-

chlorvynol. Ethylchlorvynol. Betachlorovinyl

ethyl ethynyl carbinal. Cap. 100 mg, 200

mg., 500 mg., Bot. 100s, A 1000s.

Use: (I h.s.) Nonbarbiturate hypnotic

* **PLAGUE VACCINE, U.S.P., N.N.R. 1950.**

(Cutter) 2000 million killed Pasteurella

pestis/cc. Vial 2 cc., 20 cc.

Use: S.C. (0.5 cc. followed by 1 cc. a week

later); vaccination.

PLANADALIN. See: Corbama (Various Mfr.).

PLANCELO. (American Farmer) Methylcal-

lufosa 75% w/plantago/tab. Bot. 50s, 500s,

1000s

Use: Laxative.

PLANOCAINE. See: Procaine Hydrochloride

(Various Mfr.).

PLANOCHROME. See: Merbromin, N.F.

(Various Mfr.)

PLANOLAR. (Winthrop Labs) Aspirin 300

mg., placebo 60 mg./Tab. Bot. 100s.

Use: Arthritis.

* **PLANTAGO, OVATA COATING, N.N.D.**

1962.

See: Betojel Powder (Floratos).

Konsyl, Pow. (Burton Parsons).

L.A. Formula (Burton Parsons).

Melomvel, Pow. N.N.D. (Searle)

Mucilosa, Flakes. (Winthrop Labs.)

W/Mineral oil. See: Parasyllium, liq. (Ab-

bott)

W/Sterculia, pectin & lactose. See: Kosilac,

Pow. (United Labs.).

W/Vit. B-1.

See: Siblin, Pow. (Parke, Davis).

Testarr, Pow. (Testagar).

PLANTAGO SEED, N.F. (Psyllium Seed, Plan-

tain Seed).

PLANTAMUCIN. (Elder) deated 70% Vege-

table meal, Mag carb. A sucrose, Cal.

carb. & Sod bicarb. Also avail. w/brewer's

yeast. Pkg. 4 oz., 1 lb., 5 lb.

Use: Bulk laxative.

PLAQUENIL SULFATE, NND 1962 (Winthrop Labs) Hydroxychloroquine sulf 200 mg/Tab Bot 100s

Use: Light sensitive diseases such as leprosy erythematous or antimalarial

W/Aralen phosphate, atabrine hydrochloride See Triquam, Tab (Winthrop)

W/Aspirin See Planalar, Tab (Winthrop Labs)

PLASMA

See: Hyland Liquid Plasma (Hyland)

Normal Human Plasma (Cutter)

Normal Human Plasma (Merck Sharp & Dahme)

* **PLASMA, ANTINEMOPHILIC ORIED, U S P** NND 1962 (Hyland) Freshly drawn blood plasma, treated w/ultraviolet radiation & processed to retain antihemaphilic activity Bot 50 cc w/diluent

Use: (V) reduce clotting time of hemaphilic blood

PLASMACILLIN AQUEOUS, (Rexall) Procaine penicillin G 600M u/cc Vial 1 dose

Use: Penicillin therapy

PLASMACILLIN IN OIL, (Rexall) Procaine penicillin G 300M u/cc Vial 1 dose

Use: Long acting penicillin therapy

PLASMA EXPANDERS OR SUBSTITUTES

See: Expondex (Commercial Solvents)

Gentran (Baxter)

Plavalex Dextran (Wyeth)

Plazmaid Infusion Solution (Upjohn)

Vinisil, Sol (Abbott)

PLASMA HYDROLYSATE, NNR 1955 See: Travamin, Sol (Baxter)

* **PLASMA, NORMAL HUMAN, U S P, NND 1961** (Cutter) Dried Vial 250 cc Merck Sharp & Dahme—(Lyovac) Vial 2.5 cc, 50 cc, 250 cc, 500 cc

PLASMA, NORMAL HUMAN OPHTHALMIC (Merck Sharp & Dahme) (Lyovac) Vial 2.5 cc. (dry)

Use: (In combination with thrombin) Physiologic suture in eye surgery or skin grafting

* **PLASMA PROTEIN FRACTION (HUMAN), NND 1962** For the plasma protein preparation obtained from human plasma using the Cohn fractionation technic See Plasmanate (Cutter Lab)

PLASMANATE, NND 1962 (Cutter Lab)

Human plasma proteins 5% in saline solution 0.67% pH 6.75-7.25 Bot 250 cc

Use: Shock Hypoproteinemia

PLASMIN (Registered in Great Britain) The proteolytic enzyme derived from the activation of plasminogen Actose Is Plasmin (Human).

PLASMOCHIN NAPHTHOATE, (Winthrop

* Generic name recognized by NF, USP or A.M.A.

Labs) Pamaquine naphthaate Tab 1/2 gr 3/4 gr Bot 25s, 100s, 500s

Use: Malaria

W/Quinine sulfata (Winthrop Labs) Plasmochin HCl 1/2 gr, quinine sulfate 2 gr/Tab Bot 500s

Use: Malaria

PLASMOQUINE NAPHTHOATE See: Pamaquine Naphthaate (Plasmochin Naphthaate) (Winthrop Labs)

PLASTIBASE, (Squibb) Formerly Squibb Base composed of heavy liquid petralatum 95% & high molecular weight polyethylene Jar 1 lb

Use: Hydrophobic ointment base

PLASTIBASE HYDROPHILIC, (Squibb) Formerly Squibb Base W.A., similar to Plastibase but w/hydrophilic ingredients Jar 1 lb

Use: Hydrophilic ointment base

PLAZMOID, NND 1960 (Upjohn) Gelatin 5 Gm, Sod chloride 0.9 Gm/100 cc Bot 500 cc, 1000 cc

Use: Blood plasma substitute.

PLEBEX, (Wyeth) Thiamine HCl 10 mg, riboflavin 2 mg, niacinamide 100 mg, pyridoxine HCl 5 mg, Cal pantothenate 5 mg/cc Vial 10 & 30 cc

Use: Vit B complex therapy.

PLEBILIN, (Plessner) Bile salts 3 gr/Tab Bot 100s, 500s, 1000s

PLEBILIN PLUS, (Plessner) Desoxycholic acid 1/2 gr, dehydrocholic acid 3/4 gr, malt diastase 3/4 gr, bile salts 1 1/2 gr, pancreatin 3 gr/E C Tab Bot 100s, 1000s

Use: Functional indigestion

PLECYAMIN, (Plessner) Vit B 12 Vial (100 mcg/cc) 10 cc; (1000 mcg/cc) 10 cc

Use: Pernicious anemia

PLEGICIL, (Chn Byla Lab France and Ayerst) Acepromazine

Use: Tranquilizer

PEGINE, (Ayerst) Phendimetrazine bitartrate 35 mg/Tab Bot 100s, 1000s

Use: Appetite suppressant

PLEINABEX, (Plessner) Vit B 1 15 mg, B 2 2 mg, niacinamide 10 mg, Vit B 6 0.1 mg, pantothenic acid 1 mg w/liver & yeast/Tab Bot 100s, 1000s

Use: Vit B complex

PLENAVITA, (Massengill) Vit A 5000 u, Vit C 1000 u, Vit B 1 3 mg, Vit B 2 3 mg, nicotinamide 50 mg, Vit B 6 0.5 mg, cal pantothenate 2 mg, Vit C 50 mg, Vit E 1 u/Peri Bot 100s, 500s, 1000s

Use: Dietary supplement

PLESTRAN, (Warner Chilcott) Ethinyl estradiol 0.005 mg, methyltestosterone 2.5 mg, proloval 15 mg/Tab Bot 100s, 500s

- Use: Orally, treatment of the geriatric patient.
- PLESTRIN**, N.N.B. 1950. (Harrower) Natural estrogenic substances. Aq. susp. [10,000 u./cc. & 20,000 u./cc.], 10 cc. in oil [10,000 u./cc. & 20,000 u./cc.], 10 cc., 30 cc.
- Use: Menopausal syndrome.
- PLETOCOL** (Plessner) d-Alpha tocopheryl acetate equiv. to Vit. E 100 u./Cap. Bot. 100s, 500s, 1000s
- PLEXCINE**, (Barry) Thiamine HCl 100 mg., riboflavin 1 mg., niacinamide 100 mg., pyridoxine HCl 10 mg., panthoel 10 mg., benzyl alcohol/cc. Vial 10 cc.
- Use: Vit. B complex therapy.
- PLEXCINE-12**, (Barry) Cryst. Vit. B-12 10 mcg., thiamine HCl 5 mg., riboflavin 2 mg., niacinamide 75 mg., pyridoxine HCl 5 mg./cc. Vial 10 cc.
- Use: Vit. B complex therapy.
- PLEXODERM**, (Crookes-Barnes) Styraz 6%, allantoin 1%, salicylic acid 1%. Cream, Pkg 2 oz., 4 oz.
- Use: Treatment of psoriasis.
- PLEXONAL**, (Suedaz) Barbitol Sod. 45 mg., phenobarbital Sod. 15 mg., sodium 25 mg., scopalamine HBr .08 mg., dihydroergotamine methanesulfonate 0.16 mg. Tab. Bot. 100s, 500s, 1000s.
- Use: [1-3 i.i.d.] Sedative-hypnotic.
- PLEXOPAR**, (Estra) Thiamine HCl 20 mg., riboflavin 1 mg., pyridoxine HCl 2 mg., niacinamide 100 mg., Cal. pantothenate 5 mg., benzyl alcohol 1% /cc. Amp. 1 cc., Box 6s, 12s, 25s, 100s. Vial 10 cc., 30 cc.
- Use: Vit. B complex therapy.
- PLEXOR**, (Taylor) Pancreatin, pepsin, betaine HCl, oxbile ext., thiamine HCl, riboflavin, pyridoxine HCl, ascorbic acid, niacinamide, calcium pantothenate, Vit. A, D, calcium, phosphorus, iron, magnesium, zinc/tab. Bot. 100s.
- Use: Digestive.
- PLEXTONE W/PBOCAINE**, (Nien) Vit. B-1 100 mg., B-2 0.5 mg., B-6 1 mg., nicotinamide 50 mg., iron cacodylate 10 mg., liver inj. 0.2 cc., procaine HCl 1% /cc. Vial 10 cc.
- Use: I.M., treatment of anemias.
- PLIMASIN**, (Ciba) Pyribenzamine HCl 25 mg., Ritalin 5 mg./Tab. Bot. 100s.
- Use: [1 q.i.d.] Antihistaminic w/stimulant.
- PLUBABRIN**, (Approved Pharm.) Vit. A 12,000 u., D 1200 u., B-1 4 mg., B-2 2 mg., B-6 0.5 mg., C 75 mg., niacinamide 30 mg., Cal. pantothenate 3 mg., tocopherol 5 mg., Vit. B-12 1 mcg./Cap. Bot. 100s, 1000s.
- Use: Vitamin supplement.
- PLUBAVIT DROP**, (Winthrop Labs.) Vit. A 5000 u., B-1 1 mg., B-2 0.4 mg., B-6 1 mg., nicotinamide 5 mg., pantothenic acid 2 mg., Vit. C 50 mg., D-2 1000 u., B-12 1 mcg./0.6 cc. Bot. 15 cc. w/dropper.
- Use: Nutritive for infants.
- PLURAVIT PELLETS**, (Winthrop Labs.) Vit. A 5000 u., B-1 2 mg., B-2 2 mg., B-6 0.5 mg., B-12 1 mcg., nicotinamide 20 mg., cal. pantothenate 1 mg., vit. C 50 mg., D-2 1000 u., folic acid 0.25 mg./pellet. Bot. 30s, 100s, 1000s.
- Use: Multiple vitamin deficiencies.
- PLURAXIN**, (Winthrop Labs.) Vit. A 10,000 u., D 400 u., B-1 10 mg., B-2 10 mg., B-6 2 mg., B-12 4 mcg., folic acid 1.5 mg., Cal. pantothenate 20 mg., nicotinamide 100 mg., Vit. C 300 mg., menadione 2 mg./Cap. Bot. 30s, 100s, 1000s.
- Use: Multivitamin therapy.
- PLUBICAPS-M**, (Corrione) Vit. A 25,000 u., D 1000 u., B-1 10 mg., B-2 5 mg., B-6 1 mg., B-12 5 mcg., C 150 mg., nicotinamide 100 mg., calcium 103.6 mg., phosphorus 80.2 mg., iron 10 mg., magnesium 5.5 mg., manganese 1.0 mg., potassium 5.0 mg., zinc 1.4 mg./Cap. Bot. 50s, 100s
- Use: Nutritional supplement.
- PLURIGLAND**, (Myers-Carter) Water-soluble ext. of adrenal cortex, thyroid, thymus, ant. pit., post. pit., lymphatics and ovaries. Vial 30 cc.
- PLUBISULFA**, (Kirk) Sulfadiazine 2½ gr., sulfamethazine 2½ gr., sulfamerazine 2½ gr./Tab. Bot. 100s, 1000s.
- Use: Orally; sulfonamide therapy.
- PLUBIZYME**, (Nelson-Boyer) Pancreatin 1½ gr., bile ext. 2 gr., pepsin 1 gr., papain ½ gr./Tab. Bot. 100s, 1000s.
- Use: Indigestion.
- PLUS-FACTOR**, (Sutliff & Case) Calcium carbonate 1 mg., hesperidic complex 100 mg., dioctyl sod. sulfosuccinate 120 mg., Vit. K 0.5 mg., C 100 mg., A 5000 u., D 1000 u., B-1 2 mg., B-2 1.2 mg., B-6 1 mg., B-12 2 mcg., calcium pantothenate 2 mg., nicotinamide 10 mg., folic acid 0.675 mg., iron 30 mg., cobalt 0.1 mg., manganese 5 mg., magnesium 5 mg., copper 3 mg., zinc 2 mg./2 Tab. Bot. 100s, 1000s, 50 disp-paks of 60s.
- Use: Phosphorus-free prenatal supplement.
- PMB "200"**, (Ayerst) Conjugated equine estrogens 0.4 mg., meprobamate 200 mg./Tab. Bot. 60s, 500s.
- "400" contains meprobamate 400 mg.
- Use: Relief from anxiety and tension.
- P-M-HISTINE**, (Pitman-Moore) Phenylephrine hydrochloride 25 mg., chlorpheniridine maleate 4 mg./lang-acting tab.
- Use: Antihistaminic and vasoconstrictor.

PMN OINTMENT. (Poythress) Phenylmercuric nitrate 1:1500 oint Tube 1 oz., Jar 1 lb

Use: Topically; fungicide

PNEUMOCOCCUS ANTIGEN. (Lilly) Susp of partially autolyzed pneumococci, 20,000 million/cc Types 1, 2, 3 & 4 Vial 5 cc

Use: Pneumococcus vaccines

PNEUMOCOCCUS TYPING SERA. (Wyeth) Mix "A," Type 1, 2 & 7; "B," Type 3, 4, 5 & 8; "C," Type 9, 12, 14, 15, 17 & 33; "D," Type 10, 11, 13, 20, 22, 24 & 34; "E," Type 16, 18, 19, 21 & 2R; "F," Type 23, 25, 27, 29, 31 & 32

PNEUMO-STREPTO VACCINE. (National Drug) Polyvalent 5500 million/cc Vial 30 cc

Use: SC or IM; prophylaxis & treatment of common cold

PNS, SUPPOSITORIES. (Winthrop Labs) Pontocaine HCl 10 mg, neo-synephrine HCl 5 mg, sulfamylam HCl 3 gr, bismuth subgallate 1½ gr, balsam of peru ¾ gr / Supp Rax 12s

Use: Rectal inflammatory conditions

POCHLORIN. (Texaphyl) Prophyrinic and chlorophyllic compound

Use: Anti hypercholesteremic agent

PODIACORT. (PodiatRex) Podiacort (hydrocortisone diethylaminoacetate HCl) 5 mg, neomycin sulfate 3.5 mg, undecylenic acid 50 mg, in petroleum base/Gm 5 Gm tube

Use: Control of inflammation and infection

PODIPEX. (PodiatRex) Multivitamins and minerals with intrinsic factor for podiatrists/chiropractors Cops Bot 60s

PODIVITE. (PodiatRex) Vitamins & minerals Cops Bot 60s

Use: Nutritional supplement

PODOBEN. (Maurry) Resin podophyllum 25%, zinc benzoin comp 50% v/v, iso-propyl alcohol liq Bot 5 cc

Use: Aches

***PODOPHYLLIN, USP** (Various Mfr) Resin podophyllin Pkg 1 oz., ¼ lb, 1 lb

PODOSAN POWDER (Ooak) Sod penta chlorophenate 01%, boric acid 5%, zinc peroxide 30% & boric acid 5% in kieselguhr Pkg 2 oz

Use: Fungicide

POISON IVL. (Cutter) Tax ivl Ext of Rhus toxin Bot w/dropper 13 cc

Use: See Poisonak (Cutter)

POISON IVY EXTRACT, NND 1962

Abbott—Amp 1 cc, 2s

Ledeile—Vial 1 cc, 2s

Parke, Davis—Amp 1 cc, 10s

Pitman-Moore—Pkg 1s for treatment

Merck Sharp & Dahme—Bot 1 oz

* Generic name recognized by NF, USP or AMA

See: Aqua Ivy, Viol. (Strong Cobb)

Aqua Ivy AP, Bot (Allergens)

Birhutox, Amp (Central)

Poisonivl (Cutter)

Poisonak, Ext, Bot (Cutter)

POISON OAK EXTRACT, NND 1952

Parke, Davis—Amp, Rhus toxicodendron antigen, 1 cc, Box 10s

Pitman Moore—Amp (Dried leaf ext 1 mg/cc) 1 cc

POISON OAK-IVY EXTRACT, NND 1952

Hollister-Stier—Poison Ivy ext & poison oak, Vial 5 doses

Pitman Moore—Poison Ivy ext & poison oak, 0.5 mg each/cc Vial 1 cc

See: Birhutox (Central)

Ivaka (Pitman Moore)

Ivyoak (Sherman)

Ivyol (Merck Sharp & Dahme)

* **POISON IVY EXTRACT, ALUM PRECIPITATED, NND 1962** An aqueous suspension of a pyridine extract of poison ivy which is precipitated with alum and adjusted to standard concentrations See: Aqua Ivy, AP (Synlex Chem Co, Inc)

POISON OAK-IVY-SUMAC EXTRACT (Wyeth) Poison ivy ext, oak & sumac Tube 1 cc, 5s See: Taxak, Amp (Cutter)

POISONOK. (Cutter) Taxok Alcoholic ext of Rhus toxin Bot w/dropper 13 cc

Use: Orally; in ascending dosage, in treatment of poison ivy & poison oak dermatitis (1 gtt daily in water, increased 1 gtt each day to 10, then finish bot with 10 gtt daily)

* **POLAMETNENE RESIN CAPRYLATE.** The physico chemical complex of the acid bind mg ion exchange resin, polyamine methyl ene resin and caprylic acid See: Kaprylex (R J Strassenburgh Co)

POLANIL. (Schering) Dexamethasone 0.25 mg, d chlorpheniramine maleate 2 mg; Vit C 75 mg / Tab Bot 50

Use: Seasonal allergies

POLARAMINE MALEATE, NND 1962 Dextro chlorpheniramine maleate (d isomer of Chlor Trimeton) Repetab 6 & 4 mg Bot 100s 1000s Tab 2 mg Bot 100s, 1000s

Syr [2 mg/dr] Bot 1 pt

Use: Antihistaminic

* **POLDINE METHYLSULFATE.** 2 benziloyl oxymethyl 1 1 dimethylpyrrolidinium methyl sulfate See: Nactan (McNeil)

POLIO EMMUNE GLOBULIN, HUMAN (Cutter) Antibody equivalent 40 cc of normal immune serum (165 mg gamma globulin/cc) Vial 2 cc, 10 cc

Use: Passive immunization

- * **POLIOMYELITIS IMMUNE GLOBULIN (HUMAN)**, N.N.D. 1962. Prep. of gamma globulin primarily assayed for content of poliomyelitis antibodies in accordance with standard procedures licensed by National Institutes of Health.
- POLIOMYELITIS IMMUNE GLOBULIN, HUMAN**, (Hyland) Antibody-bearing gamma globulin fraction of normal human plasma. Vial 2 & 10 cc.
Uses: M.M., measles & poliomyelitis.
- POLIOMYELITIS IMMUNE GLOBULIN, HUMAN**, (Merck Sharp & Dohme) gamma-Globulin. Vial (globulin 16.5 Gm./100 cc.) 2 & 10 cc.
- * **POLIOMYELITIS VACCINE, U.S.P., N.N.D. 1962.**
Cutter—Vial 3 cc.
Lilly—Vial 3 cc., 9 cc.
Parke, Davis—Vial 3 cc., 9 cc.
Pittman-Moore—Vial 1 cc., Pkg. 3s, Vial 9 cc.
Merck Sharp & Dohme—Vial 3 cc., 9 cc.
Wyeth—Vial 1 cc. Pkg. 3s, 9 cc.
- POLIOMYELITIS VACCINE**, (Parke, Davis) Poliomyelitis vaccine prepared by adsorbing poliomyelitis vaccine on aluminum phosphate suspension with the aid of a small amount of protamine sulfate. Vial 4.5 cc., enough for 9 doses.
- POLIO VACCINE, ORAL** (Pfizer) From attenuated Type 1 and Type 2 polioviruses which were dev. by Dr. A. Sabin.
Uses: Orally, Poliomyelitis.
- POLIVEX**, (G. F. Horvay) Vit. B-12 15 mcg. plus Vit. B complex present in crude liver sol./cc. Vial 30 cc.
Uses: Pernicious anemia
- POLLEN-X**, (S-K Research Labs) Pollen antigens, 6 different types according to 6 pollen zones in U.S. Bot. 1/2 & 1 cc.
Uses: Treatment of hay fever.
- POLONIL**, (Schering) Dexamethasone 0.25 mg., dexchlorpheniramine maleate 2 mg., ascorbic acid 75 mg./Tab. Bot. 50s, 500s.
Uses: Treatment of allergic and inflammatory disorders.
- * **POLOXALKOL**, N.N.D. 1962. Polyoxyethylene polyoxypropylene polymer. See: Polykal Drops (Upjohn)
W/Casanthrol See: Cosakol, Cap., & Syr. (Upjohn)
- POLYACRYLIC CATIONIC RESIN**.
See: Kolnil (National Drug)
Kayexalate (Winthrop)
Resins, Sodium Removing
Resins, Potassium Removing
Voids, Carborestin
- * Generic name recognized by N.F., U.S.P. or J
- * **POLYAMINE-METHYLENE RESIN, N.N.D. 1962.**
See: Basex. (Columbus)
Carbo-Resin (Lilly)
Esorbin (Ayers)
Resiant, Prep., N.N.R. (National Drug)
Resion, Liq. (National Drug)
Resipax (Squibb)
W/Gastric mucln. See: Resmican, Tab. (Whittier)
W/Polymyxin-B sulfate, Sod. aluminum silicate, Mag aluminum silicate & phthalyl-sulfacetamide. See: Resion-PMS, Susp. (National Drug)
- POLYAMINE RESIN**.
See: Polyamine-Methylene Resin, N.N.D. 1958.
W/Polyacrylic collicae resin Carbo-resin (Lilly). See: Vacld.
- POLYANHYDROGLUCOSE**, See: Dextran
- POLY-B**, (U.S. Vitamin) Vit. B-12 1 mcg., choline 20 mg., inositol 10 mg., thiamine NCI 10 mg., riboflavin 5 mg., pyridoxine NCI 1 mg., niacinamide 50 mg., Cal. pantothenate 5 mg., liver 100 mg., yeast 50 mg./Cap. or 5 cc. of Syr. Bot. 50s, 100s, 200s Bot. 4 & 8 cc., 1 pt., 1 gal.
Uses: Vitamin B complex therapy.
- POLY-B W/C**, (U.S. Vitamin) Same formula as Poly-B w/ascorbic acid 150 mg./Cap. or 1 cc Bot. 50s, 100s, 200s. Vial 10 cc.
- POLY-B FORTIS**, (U.S. Vitamin) Vit. B-1 50 mg., B-2 0.5 mg., B-6 3 mg., niacinamide 50 mg., Cal. pantothenate 5 mg./cc. Vial 10 cc., Box 1s, 6s.
- POLY-B SPECIAL**, (U.S. Vitamin) Liver crude 1 cc., Vit. B-1 20 mg., B-2 1 mg., B-6 3 mg., niacinamide 20 mg., Vit. B-12 activity 2 mcg./cc. Vial 10 cc., 1s, 6s Vial 30 cc.
- POLYBENZARSOL**, See: Benzarsol, (Pittman-Moore)
- POLYBRENE**, N.N.D. 1962, (Abbott) Hexadimethrine. Amp (10 mg/cc) 10 cc. Cart. 5 & 25s.
Uses: Neutralize action of heparin (1 mg for each 100 u heparin)
- * **POLYCARBOPHIL A** synthetic, loosely cross linked, hydrophilic resin of the polycarboxylic type.
W/Thiexinal methylbromide. See: Sorbaquel, Tab (White Labs)
- POLYCORT**, (Lemmon) Hydrocortisone 0.02%, neomycin 0.6 mg., polymyxin B sulf. 2,000 u., chlorpheniramine maleate 0.2%, phenylephrine HCl 0.5% 3/4 oz. Squeeze spray bot. Bot. 1 pt. & 3/4 oz. spray.
Uses: Decongestant, anti-inflammatory, anti-allergic nasal therapy.

POLYCYCLINE AQUEOUS 125, NND
1962 (Bristol) Cal tetracycline equiv to
tetracycline HCl 125 mg / 5 cc Bot 2 oz
Use: Orally; antibiotic therapy

POLYCYCLINE AQUEOUS 250, NND
1962 (Bristol) Cal tetracycline equiv to 250
mg of tetracycline HCl / 5 cc Bot 1 fl oz
Use: Orally; antibiotic therapy

POLYCYCLINE HYDROCHLORIDE, NND
1961 (Bristol) Tetracycline HCl
Cap: 100 mg, Bot 25s, 100s; 250 mg, Bot
16s, 100s

Intramuscular: 100 mg / vial
Oint: Tetracycline HCl 30 mg, xylocaine
2% / Gm Tube 1/2 oz

Ophth. Oint: Tetracycline 10 mg, xylo-
caine 2% / Gm Tube 1/4 oz
Susp, liq: 250 mg / dr, Bot 30 cc
Use: Antibiotic therapy

POLYCYCLINE INTRAVENOUS, (Bristol)
Tetracycline HCl Vial (100 mg, 250 mg
300 mg)

POLYCYCLINE PEDIATRIC DROPS, (Bristol)
Tetracycline 100 mg / cc Bot w/calibrated
dropper, 10 cc
Use: Tetracycline therapy

POLYCYCLINE SUSPENSION-250, (Bristol)
Tetracycline 1.5 Gm, dry pow when sus-
pended 250 mg / 5 cc (Crushed fruit flavor)
Bot 30 cc

Use: 1 tsp qid; anti infective

POLYCYCLINE T/S, (Bristol) Tetracycline
HCl 125 mg sulfadiazine, sulfamerazine
sulfamethazine of each 167 mg / 5 cc Susp
Bot 2 fl oz

Use: Antibiotic w/triple sulfonamide therapy

***POLYESTRADIOL PHOSPHATE, See** Estro-
durin, Amp (Ayerst)

***POLYETHYLENE GLYCOL 1540, NF, &**
4000, USP, NMR 1955 See Carbowax
(Carbide & Carbon)

***POLYFEROSE, An iron carbohydrate chelate**
containing approximately 45% iron in
which the metallic (Fe) ion is sequestered
within a polymerized carbohydrate derived
from sucrose See Jetrone (Pitman Moore)

POLYGALACTURONIC ACID W/Sennasides
A and B, Guar Gum See Gentlas Gran-
ules (Purdue Frederick)

POLYGESIC, (Arnar Stone) d Amphetamine
sulfate 25 mg, acetophenetidin 2 1/2 gr
pentobarbital cal 1/4 gr salicylamide 230
mg / tab Bot 100s, 1000s
Use: (1 2 q 4 h) Analgesic w/sedative

POLYKOL, NND 1962 (Upjohn) Drops
Polaxalkol 200 mg / cc Bot 30 cc Caps
250 mg Bot 16s, 100s
Use: Fecal normalizer for infants

* Generic name recognized by NF, USP or AMA

POLYMACMA, (Wyeth) Activated attapul-
gite 3 Gm, dihydrostreptomycin 300 mg,
polymyxin B sulf. 120M u, pectin 270 mg /
fl oz Bot 8 oz.

Use: Treatment of diarrhea
POLYMACMA PLAIN, (Wyeth) New name
Diamagma Attapulgitte activated 3 Gm,
pectin 270 mg and alumina gel/fl oz Bot
12 fl oz See: Spensin (Ives)

Use: Treatment and control of diarrhea
POLYMACMA TABLETS, (Wyeth) Polymyxin
B sulfate 25,000 u, dihydrostreptomycin
base (as sulfate) 75 mg, activated attapul-
gite 350 mg, pectin 45 mg, hydrated
alumina powder 70 mg / Tab Bot 48s

Use: Treatment of diarrhea
POLYMETNINE BLUE DYE, (3,3' diethylthio
dicarbocyanine iodide) See: Delvex, Tab-
(Lilly)

**POLY[METHYLENE-4-HYDROXYBENZENE-
ARSONIC ACID], See** Benzorsol (Pitman
Moore)

POLYMONINE, A formaldehyde polymer of
N methylmonoamylamine

POLYMUL, (Pharmaceutical Inds) Unsolu-
ble fatty acid glycerides 950 mg (pro-
viding linoleic and linolenic acids 600 mg,
lauric, penta, hexanoic acids 100 mg,
oleic and saturated acids 150 mg) mixed
tocopherols 5 mg polysorbate 80, 50
mg / cc

Use: Dietary supplement
POLYMYXIN, NND 1962 Generic term
for antibiotics obtained from fermentations
of various media by strains of *Bacillus*
polymyxa

Use: Polymyxins A, B & D active against
susceptible gram negative bacteria

POLYMYXIN B, NND 1962 (Various Mfr)
(No Pharmaceutical Form Available) Anti-
microbial substances produced by *Bacillus*
polymyxa

W/*Bacitracin* See: Candettes Trache (Pfizer)
W/*Bacitracin* & streptomycin Cal chloride
complex See: Multibiotic Gauze Pads (Bio-
rama Drug)

W/*Bacitracin* zinc, neomycin sulf benzal-
konium chloride See: Biotres, Oint (Central)
W/*Neomycin* sulfate zinc bacitracin, benzal-
konium chloride See: Biotres, Oint (Central
Pharmaceutical)

W/*Terramycin* See: Terramycin oint (Pfizer)

***POLYMYXIN B SULFATE, USP, NND**
1962; Tab USP Oint USP (Pfizer) Ophth
oint 20 000 u / Gm Tube 1/2 oz Oint
20 000 u / Gm Tube 1/2 oz Paw, Vial (200
000 u) 20 cc Tab 250 000 u Bot 24s Vial
(500 000 u) 10 cc

- See: Aerosporin, Prep. (Burroughs Wellcome).
- W/Bacitracin.
- See: Polycin, Prep. (Pitman-Moore).
- Polysporin, Oint. & Ophth. Oint. (Burroughs Wellcome).
- W/Bacitracin, isobutyl p-aminobenzoate
- See: Poly-Tracin, Troches. (Hance)
- W/Bacitracin, Neomycin.
- See: Neo-Polycin Oint. & Ophth. Oint. (Pitman-Moore)
- Neo-Polycin Ophth. Oint. (Pitman-Moore)
- Neosporin, Oint. & Ophth. Oint. (Burroughs Wellcome)
- Trimixin, Oint. (Hance)
- POLYOXYETHYLENE (20) (SORBITAN MONO-OLEATE)**, A.D.R. 1955. See Polysorbate 80, U.S.P.
- POLYOXYETHYLENE AD MONOSTEARATE**, See: Polyoxyl 40 Stearate
- POLYNOXYLIN**, Poly[methylenedi(hydroxymethyl)urea], Anaesthetic.
- POLYOXYETHYLENEOXYLPHENOL** W/AI-kyldimethylbenzylammonium chloride, methylrosaniline chloride, polyethylene glycol tert-didecylthioether. See: Hyva, Tab. (Holland-Rantas Co.)
- POLYOXYETHYLENE GLUCITAN MONO-LAURATE**, W/Ferrous gluconate. See: Simron, Cap. (Merrell)
- POLYOXYETHYLENE LAURYL ETHER** W/Hexachlorophene, sulfur. See: Fostril, Lot. (Westwood)
- W/Hydrocortisone, hexachlorophene, sulfur. See: Fostril HC (Westwood)
- ***POLYOXYL AD STEARATE**, U.S.P. Polyoxethylene 40 monostearate. See: Myrl 52. (Atlas Pow.)
- Use: Hydrophilic Oint.
- POLYPEP**, (Liberty Labs) Solution: Polysaccharid complex iron 75 mg., Vit. B-1 6 mg., B-2 6 mg., niacinamide 30 mg., panthenol 1.5 mg., B-6 1.5 mg., B-12 15 mcg./oz. Bot. 12 # ae. Tablets: Polysaccharid complex iron 25 mg., Vit. B-1 2 mg., B-2 2 mg., B-6 0.5 mg., panthenol 0.5 mg., niacinamide 10 mg./Tab. Bot. 60s.
- Use: Dietary supplement with iron and vitamins.
- POLYPROTE**, (Kirk) Sulfur colloidal, non-specific protein. Amp. 2 cc.
- POLYSAL**, (Cutter) Sod. chloride 0.496 Gm., Sod. acetate 0.64 Gm., Pot. chloride 0.0746 Gm., Cal. chloride 0.0368 Gm., Mag. chloride 0.0305 Gm., Sod. citrate 0.0785 Gm./100 cc. Saltflash 250 cc., 1000 cc.
- Use: I.V., to restore the electrolyte balance.
- POLYSAL ELIXIR**, (Cutter) Sod. & pot. chlorides 5.55 Gm., cal. & sod. tetrates 5.55 Gm., mag. gluconate 0.5 Gm./100 cc. Liq. Bot. 6 oz.
- Use: Orally, electrolyte therapy.
- POLYSORB**, (Fougera) Hydrophilic aint base of sorbitan sesquioleate in petrolatum base, 100 Gm., 1 lb., 5 lb.
- Use: Dermatological vehicle.
- POLYSORB HYDRATE**, (Fougera) A hydrodispersant emollient. Tube 50 Gm. Jar 425 Gm.
- Use: Treatment of dermatitis
- ***POLYSORBATE 80**, U.S.P., A.D.R. (Polyoxyethylene 20, Sorbitan Mono-Oleate).
- See: Monilan, liq. (Ives-Cameron).
- Sorbate, liq. (Abbott)
- Tween 80 (Atlas Pow. Co.)
- W/Choline & inositol, See: Monichol, liq. (Ives-Cameron)
- POLYSPORIN ANTIBIOTIC OINTMENT**, (Burroughs Wellcome) Polymyxin B sulfate 10,000 u., bacitracin 500 u. in special white petrolatum base/Gm. Tube 1/2 & 1 oz. Ophth. 1/2 oz.
- Use: Antibiotic active against gram-negative & gram-positive organisms.
- POLYSULFIDES**.
- See: Polythionates (Upsher-Smith)
- Vlem-Dome (Dome)
- POLYTAR BATH**, A blend of four different vegetable and mineral tars with emollient oils/Pkg. 10s.
- Use: Psoriasis.
- POLYTAR BATH, CAP.** (Silelef Lab) See: Cool Tar.
- POLYTAXIN**, (Winthrop Labs) Vit. A 5000 u., B-1 2 mg., B-2 3 mg., C 75 mg., nicotinamide 20 mg., Vit. D-2 400 u./cap. Bot. 100s, 1000s.
- Use: Prophylaxis against multiple vitamin deficiency.
- ***POLYTHIAZIDE**, 2-methyl-3,4-dihydro-3-(2,2,2-trifluoroethylthiomethyl)-6-chloro-7-sulfomyl-1,2,4-benzothiadiazine-1,1-dioxide. See: Renesa Tab. (Pfizer)
- POLYTHIONATE**, (Upsher-Smith) Sulfur 1.5%, polythionates of Sod. & zinc oxide lot. 2 oz., 4 oz.
- Use: Topically: antibacterial.
- POLY-TRACIN**, (Hance) Bacitracin 50 u., polymyxin B sulf. 1000 u. & isobutyl p-aminobenzoate 5 mg./trache. Vial 16s.
- Use: Treatment of throat irritations.
- ***POLYURETHANE FOAM**, N.N.D. 1962.
- Prepared by mixing a prepolymer (consisting of a trihydroxy resin reacted with an excess of an aramalic diisocyanate) with a tertiary amine catalyst. When the catalyst and prepolymer are mixed in 1:4 ratio, polymerization and liberation of carbon
- * Generic name recognized by N.F., U.S.P. or A.M.A.

- dioxide result finally in a rigid foam of approximately twice the volume of the original components. See Ostamer (Merrell)
- POLYVIDONE** See Polyvinylpyrrolidone
- POLY-VI-FLOR** (Mead Johnson) Fluvone (sod) 1 mg, Vit A 4000 u, D 400 u, C 75 mg, B 1 12 mg, B 2 15 mg, nicotinamide 15 mg/Tab Bot 50s
Use: Dental caries
- POLYVIMS** (E S Miller) Vit A 10,000 u, D 1000 u, B 1 6 mg, B 2 8 mg, B 6 1 mg, B 12 2 mcg, C 100 mg, E 0.5 mg, niacinamide 20 mg, Cal d pantothenate 2 mg, folic acid 1 mg, choline bitartrate 40 mg, inositol 10 mg, w/Phos, Cal, Iron, copper, Mang, molybdenum, iodine, Pol, zinc, Mag, cobalt/tab Bot 100s, 500s, 1000s
Use: Dietary supplement
- POLYVINYLPIRROLIDONE-IODINE COMPLEX. POVIDONE-IODINE COMPLEX.**
See: Betadine (Carbin & Conwell)
Iodine (Isodine Co)
- POLYVINYLPIRROLIDONE, POLYVIDONE, POVIDONE.**
See PVP Macrose, Sol (Schenley)
Vinisol, Sol (Abbott)
- W/Acetrisoate Sod** See Solpax, Vial (Ortho)
- POLY-VI-SOL** (Mead Johnson) Vit A 5000 u, D 1000 u, ascorbic acid 60 mg, thiamine 12 mg, riboflavin 15 mg, niacinamide 15 mg/5 cc Bot 4, 8, 12 oz Tob Bot 24s, 100s
Use: Diet supplement
- POLY-VITAMIN** (Mead Johnson) Vit A 5000 u, D 1000 u, thiamine 1 mg, riboflavin 0.8 mg, niacinamide 5 mg, ascorbic acid 30 mg/1sp Susp Bot 3 oz
Use: Dietary supplement
- POLY-VITAMIN** (E S Miller) Vit A 5000 u, B 1 6 mg, B 2 8 mg, B 6 1 mg, C 100 mg, E 0.5 mg, D 1000 u, niacinamide 20 mg, Cal pantothenate 2 mg, w/iron, iodine Mag trisilicate 200 mg/Tab Bot 100s, 500s, 1000s
Use: Dietary supplement.
- POLYVITAMINS** See Vitamin preparations
- POMALIN** (Winthrop Labs) Sulfaquinoxidine 2 Gm, pectin 225 mg, kaolin 3 Gm, opium tr 008 cc/15 cc Bot 1 pt
Use: Antidiarrhea
- PONDETS** (Wyeth) Pot penicillin G 20,000 u, bacitracin 50 u/Troche Con 48
Use: Topical; antibiotic preparation for dental procedures
- PONTOCAINE BASE, NNR 1953, ADR** (Winthrop Labs) (Tetracaine Dimethylamino ethyl p N butylaminobenzoate)
Creams: (1%) Tube 1 oz; Jar 1 lb
* Generic name recognized by N F U S P or A M A
- Ophth oint.: (0.5%) Tube ½ oz**
Use: Anesthetic.
- PONTOCAINE COMPOUND OINTMENT** (Winthrop Labs) 0.5% pontocaine & 0.5% menthol in white petrolatum, anhydrous lanolin & beeswax. Tube 1 oz Jar 1 lb, 5 lb
Use: Topical anesthetic
- PONTOCAINE DEXTROSE SOLUTION** (Winthrop Labs) Pontocaine 0.2%, dextrose 6% Amp 2 cc, Box 10s Pontocaine 0.3%, dextrose 6%. Amp 5 cc, Box 10s
Use: Spinal anesthetic
- PONTOCAINE HYDROCHLORIDE, NNR 1957** (Winthrop Labs) Anetholne, Tetracaine NCI
Ophth. Oint.: (0.5% of base) Tube ½ oz. Amp: Niponoid (10, 15 or 20 mg pow) for spinal anesthetic
Sol: (0.15% in isotonic solution of three chlorides) Bot 100 cc; (1.5 mg/cc) Bot 100 cc. for caudal anesthetic; (0.2% with 6% dextrose) 2 cc amp for saddle block anesthetic; (0.3% with 6% dextrose) 5 cc amp for spinal anesthetic; (0.5%) Bot 15 cc, 60 cc; (1%) "Niponoid" ampuls 250 mg Amp 2 cc; (2%) Bot 30 cc; 120 cc tinted w/methylene blue to prevent occlusion injection
Tab: (0.1 Gm) for use in sol for surface anesthetic, not for inj]
- POPPY-SEED OIL**, the ethyl ester of the fatty acids w/iodine See: Ethiodol, Amp (Fougero)
- POPPY-SEED OIL, 10%, IODIZED, NNR 1952**
- POQUIL** (Parke Davis) See: Pyvinium Chloride
- PORFIROMYCIN** (Upjohn) Antibiotic from *Streptomyces ardens*
Use: Inhibits gram negative and gram positive bacteria
- PORLYN** (Parke Davis) Benzilium bromide
Use: Anticholinergic
- POSTA-COLE** (Cole) Hyoscyamus ext ½ gr, arbutin ¼ gr, chromium sulfate ½ gr, tritium ext ½ gr, echinocoe ext ½ gr/Tab Bot 100s
Use: Urinary antiseptic
- POSTAFENE**. See: Bonamine, Tab (Pfizer)
- *POSTERIOR PITUITARY, U S P, NNR 1949; Inf, U S P**
See: Pituitary, Posterior (Various Mfr)
- POSTLOBIN-O**. See: Pituitary, Post., Hormone (b)
- POSTLOBIN-V**. See: Pituitary, Post., Hormone (a)

- POTABA.** (Glenwood) Pot. p-aminobenzoate. Paw., Bot. 100 Gm., 1 lb. Tab. 0.5 Gm., Bot. 100s.
Use: Rheumatoid arthritis.
- POTABA ± 6.** (Glenwood) Potassium p-aminobenzoate 0.5 mg., Vit. B-6 1 mg./Tab. or Cap. Caps. Bot. 240s, 1000s. Tab. Bot. 120s, 1000s.
Use: Antifibrinolytic therapy.
- POTASSIC SALINE (DARROW), N.N.R. 1955.** (Baxter) Vaculiter bot. containing 500 cc. w/Pot. chloride 2.7 mg., Sod. chloride 3 mg., Sod. lactate 6 mg./cc.
Use: Parenterally for dehydration and acidosis.
- POTASSIO-MERCURIC IODIDE DISCS, N.N.R. 1950, A.D.R. 1955.** (Parke, Davis) Pot. mercuric iodide. Disc, small & large, Bot. 100s, 500s.
Use: Antiseptic.
- POTASSIUM ACETATE.** (Cultnr) Pot. acetate soln. 49%. Ambats 10 cc.
Use: To avoid chloride when high concentration of pot. is needed
- POTASSIUM ADSORBING RESIN.** Sani Kallit, Pow. (Nallanal Drug)
- POTASSIUM p-AMINOZOATE.** See: Potaba, Pow. & Tab. (Glenwood).
W/Amm. salicylate, ascorbic acid, methamphetamine HCl. See: Neocydex, Tab. (Central Pharm. Co.)
W/Ascorbic acid, pot. salicylate, prednisolone. See: Neocylene, Tab. (Central)
W/Hydrocortisone alcohol, pot. salicylate, ascorbic acid. See: Pabolate-HC, Tab. (Robins)
W/Mephensin, salicylamide, ascorbic acid. See: Maxsal, Tab. (Haack)
W/Pot. salicylate, ascorbic acid. See: Pabolate-Sodium Fren (Robins)
W/Pyridoxine See: Potaba ± 6, Caps. or Tabs. (Glenwood Labs)
W/Salicylamide, potassium salicylate, dicalcium glutamate, ascorbic acid, prednisone. See: Glutalate W/Prednisone, Cap. (Dumas-Wilson)
- ***POTASSIUM p-AMINOSALICYLATE, N.N.O. 1962.** See: Paskatin, Fav. (Lilly)
- POTASSIUM p-AMINOSALICYLATE.** See: Inebacin Kalium (C.M.C.)
- ***POTASSIUM ASPARTATE.** W/Magnesium aspartate. See: Sportase, Tabs. (Wyeth)
- ***POTASSIUM BROMIDE, N.F., A.D.E.** (Merck Sharp & Dohme; Mallinckrodt) Granule, Bot. 3 oz., Paw., Pkg 1 lb., 5 lb.
W/Amnopyrin. See: Gynalgos, Liq. (Wyeth)
W/Hyoscyamus nlt. & chloral hydrate. See: Hypna-Bromic, Sol. (Wampoln).
- ***Generic name recognized by N.F., U.S.P. or A.M.A.**
- ***POTASSIUM CHLORIDE, U.S.P. Inj., U.S.P. Tab., U.S.P.** (Various Mfr.) Ensoals 5 gr., 15 gr. (Lilly) Bot. 100s, 500s, 1000s. (Parke, Davis) Amp. (20 mEq.) 10 cc.; (40 mEq.) 20 cc. (Abbott) Amp. (20 mEq.) 10 cc.; (40 mEq.) 20 cc.
Use: Potassium deficiency.
W/Prednisolone. See: K Predna Oame, Tab. (Dome Chemicals)
- POTASSIUM GLUCONATE.** See: Kaon, Elixir. (Warren-Teed)
- ***POTASSIUM GLUTAMATE.** The monopotassium salt of L-glutamic acid. See: Glutavenn K (Gray Pharm. Co.)
- POTASSIUM G PENICILLIN, N.N.O. 1962.** U.S.P., N.N.R. See: Penicillin G, Potassium, Prep. (Various Mfr.)
- POTASSIUM GUAIACOLSULFONATE, N.F.** Sullagualcol. Used in many cough preps.
W/Codaine. See: Thiatn (Polch)
W/Dihydrocodeinone bitartrate, phenylphrine HCl, chlorpheniramine maleate, ammonium chloride, sodium citrate. See: Efricon Expectorant. (Lonnert)
W/Phenylephrine HCl, theonyldamine HCl, dihydrocodeinone bitartrate. See: Dicodeine Syrup (Breon)
W/Pyrilamine maleate, N-acetyl-p-aminophenol, phenylephrine, d-methaphan HBr, alcohol. See: Tharmal, Liq. (Central)
- ***POTASSIUM IODIDE, U.S.P.** See: KI-N, Tabs. (Irwin-Nelsler)
W/Chloral hydrate, hyoscyamus ext. See: Bromide & Chloral Comp. (Merck Sharp & Dohme)
Bramidia, Liq. (Bottle)
Hypna-Bromic, Sol. (Wampole)
W/Phenobarb., theobromine. See: Brodital, Tab. (U.S. Standard)
W/Phenobarbital & theobromine sod. acetate. See: Theosophen Prep. (Starr)
W/Phenobarbital, theophylline anhydrous, ephedrine, p.e. belladonna leaves, pyrilamine maleate, magnesium trisilicate. See: Theolad Pediatric, Tab. (Suliff & Coe)
- POTASSIUM IODIDE ENTERIC COATED TABLETS.** (Lilly; Parke, Davis) See: Enkide, Tab. (Brewer)
W/Niacinamide. See: Iodo-Niacin, Tab. (Cole Chem. Co.)
- ***POTASSIUM ISOPROPICILLIN.** (Lilly) Potassium 6-(α -phenoxylisobutyramidal) penicillanate.
Use: Antibiotic.
- ***POTASSIUM LEVOPROPYLCILLIN.** (Eli Lilly and Company) Potassium 6-(1- α -phenoxylisobutyramidal) penicillanate.
- POTASSIUM NITRATE, N.F.**
Use in: Crataganita, Tab. (C. D. Smith)

POTASSIUM PENICILLIN V, N N D 1962
See: Phenoxymethyl Penicillin Pot

***POTASSIUM PENICILLIN O.** See: Penicillin O Pot

***POTASSIUM PERMANGANATE** (Paul B. Elder) See: Insta Perm

POTASSIUM PHENOXYETHYL PENICILLIN.
See: Phenethicillin Pot

***POTASSIUM PHENOXYMETHYL PENICILLIN, U S P, N N D 1962**
See: Penicillin Phenoxymethyl pot

Penicillin V Potassium

***POTASSIUM PHENETHICILLIN, N N D 1962** See: Phenethicillin Pot

***POTASSIUM PROPYLICILLIN.** (Eli Lilly) Potassium 6 (a phenoxycarbutylamido)penicillanate

Use: Antibiotic

POTASSIUM-REMOVING RESINS.
See: Kalini, Paw (Nolional Drug)

Kayexolate (Winthrop Lab)
Special Resin No. 648 (Smith, Kline & French)

POTASSIUM RHODANATE. See: Pot Thio cyanate

POTASSIUM SALICYLATE
W/Dicalcium glutamate, ascorbic acid, salicylamide, prednisone, potassium p-aminobenzoate See: Glutalate W/Prednisone, Cap (Dumas Wilson)

W/Hydrocortisone alcohol, pot p aminobenzoate ascorbic acid See: Fobalate HC, Tab (Robins)

W/Pot aminosalicylate, ascorbic acid See: Fobalate Sodium Free (Robins)

W/Prednisolone, ascorbic acid, pot p aminobenzoate See: Neocytone, Tab (Central)

W/Prednisone, cal pentathenate See: Salfort Delta, Tab (Massengill)

W/Salicylamide, p aminobenzoate pot, dicalcium glutamate, and Vit C See: Glutalate Cap (Dumas Wilson)

POTASSIUM SORBATE See: Sorbistat K Pow (Pfizer)

POTASSIUM SULFOCYANATE (Par Rhodanate) See: Pot Thiocyanate

POTASSIUM THERAPY.
See: Kaon, Elixir (Warren Teed)
Potassium Chloride (Viosius Mfr)

POTASSIUM THIOCYANATE, H F X (Pot sulfocyanate, Pot Rhodanate)
See: Artesiacyn, Elixir (Flint, Eaton)
Kocyan (McNeil)

W/Cascaramonic See: Thio Cara (Suliff & Case)

W/Phenobarbital See: Hyper Sed, Tab (Suliff & Case) Sybolan, Tab (Massengill)

* Generic name recognized by N.F., U.S.P. or A.M.A.

W/Rutin See: Hyper Ru, Tab (Suliff & Case)

POTASSIUM THIOCYANATE. (Lilly) Timed disintegrating tab (1 or 3 gr) Bot 100s, 500s, 1000s

Use: Hypertension

***POTASSIUM THIPHENCILLIN.** (Lilly) Potassium 6 (phenylmercaptoacetamido)penicillanate

Use: Antibiotic

POTENSORS. (Tailby Nasan) Not listed
Veratrum viride tr 2 min, acanite root tr 2 min, Sod nitrite 1 gr, bile salts 1/6 gr, cascara sagrada ext 1/2 gr /Tab blue E.C., Bot 100s

Use: Hypertension

W/Phenobarbital 1/4 gr./Tab pink E.C., Bot 100s

W/Rutin 20 mg, phenobarbital 1/4 gr /Tab yellow E.C., Bot 100s

POTENSORS FORTE. (Tailby Nasan) Not listed
Veratrum viride whole 1 1/2 gr, Sod nitrite 1 gr, bile salts 1/2 gr, cascara sagrada ext 1/2 gr /Tab orange E.C., Bot 100s

Use: Hypertension

POTENSOR FORTE, IMPROVED (Tailby Nasan) Rauwolfia serpentina whole root 50 mg, veratrum viride alkaloids 1 mg./Tab. Bot 100s

Use: 1 to 3 days hypertension

POVAN, N N D 1962 (Porke, Davis) Pyrvinium base 50 mg /Tab Bot 25s

Use: Treating pinworm infections

POVAN SUSPENSION. (Porke, Davis) Pyrvinium pamoate 10 mg/cc Bot 2 fl oz

Use: Control of pinworm infections

POVIDONE. See: Polyvinylpyrrolidone (Various Mfr)

***POVIDONE-IODINE COMPLEX, N N D 1962**
See: Iodine (Iodine Ca)
Betadine (Corbin & Canwell)

POWDALATOR, A D R 1954 (Abbott) Pot penicillin G 5000 u sulfonilamide 0.25 Gm /Tab Tube, Pkg 20s, 100s

Use: Topically, insufflation

POWDALATOR-ES (Abbott) Erythromycin ethyl succinate 10 mg erythromycin activity, sterile sulfonilamide 250 mg /tube Tubes Box 20s 100s

Use: Treatment of infections

POYALIVER STRONGER. (Testagar) Liver mg (eq 10 mcg B 12) Vit B 12 60 mcg, folic acid 5 mg /cc Vial 10 cc

POYAMIH (Testagar) Vit B 12 Cap 5 mcg. Bot 100s Liq 1 mcg /drop Bot w/ dropper 12 cc, 30 cc Vial (30 mcg /cc

- 50 mcg./cc.) 10 cc., 30 cc., (100 mcg./cc.) 10 cc.
 Use: Hematopoietic.
- P.P.D. TUBERCULIN.** See: Tuberculin, Purified Protein Derivative, U.S.P. (Various Mfr)
- P.P. FACTOR (PELLAGRA PREVENTIVE FACTOR).** See: Nicotinic Acid (Various Mfr.).
- PRADASE.** (E. S. Miller) Carminative oils 0.1 min., bile salts 150 mg., ketochalonic acids 60 mg., pancreatin 300 mg., betaine HCl 90 mg./Tab. Bot. 90s, 100s.
 Use: (1-2 h.s.) Cholagogue.
- PRAEQUINE NAPHTHOATE.** See: Pamaquine Naphthoate (Plasmochin Naphthoate. Winthrop-Stearns).
- PRAGMASUL.** (Smith, Kline & French) Sulfathiazole 5%, thimerosal 1:20,000. Oint. 1 oz.
 Use: Topically; sulfanamide therapy.
- PRAGMATAR.** (Smith, Kline & French) Cetyl alcohol-coal tar distillate 4%, near-colloidal sulfur 3%, salicylic acid 3% in oil-in-water type emulsion base. Jar 1½ oz.
 Use: Skin disorders.
- * **PRALIDOXIME CHLORIDE.** (Campbell Pharmaceuticals) 2-formyl-1-methylpyridinium chloride axime. See: Protropam Chloride.
- * **PRALIDOXIME IODIDE.** 2-pyridine aldoxime methiodide. See: Protropam (Campbell Pharm., Inc)
- * **PRALIDOXIME METHIODIDE.** See: Prothoxime iodide (Campbell Pharm., Inc)
- PRAMYL.** (Allied Prod.) Ammonium salicylate 5 gr., p-aminobenzoic acid, Pot. salt 3 gr./Tab. E.C. Bot. 100s, 500s.
 Use: Orally (1-2 t.i.d.) analgesic & antipyretic.
- PRAMEY.** (Abbott) Ferrous sulfate dried 200 mg., Vit. B-6 5 mg., C 100 mg., B-12 3 mcg., calcium carbonate 625 mg., Vit A 1.2 mg., D 10 mcg., B-1 3 mg., B-2 2 mg., nicotinamide 10 mg., calcium pantothenate 1 mg., iodine 1 mg., copper 0.15 mg./Tab. Bot. 100s.
 Use: Nutritional supplement.
- PRAMEY FILMTAB.** (Abbott) Ferrous sulfate 200 mg., pyridoxine HCl 5 mg., ascorbic acid 100 mg., cobalamin 3 mcg., calcium carbonate 625 mg., Vit. A 1.2 mg., Vit. D 10 mcg., thiamine mononitrate 3 mg., riboflavin 2 mg., nicotinamide 10 mg., calcium pantothenate 1 mg., iodine 0.1 mg., copper 0.15 mg./Tab. Bot. 100s.
 Use: Maternal supplement.
- PRAMILETS.** (Abbott) Phosphorus-free calcium, iron as ferrous fumarate plus Vit. A 4000 u., D 400 u., B-1 3 mg., B-2 2 mg., nicotinamide 10 mg., C 50 mg., B-6 1 mg., intrinsic factor 5 mg., B-12 2 mcg., folic acid 1 mg./Filmtab Bot. 100s, 1000s
 Use: Vitamin-mineral support for pregnancy and lactation.
- PRAMILETS-F FILMTAB.** (Abbott) Vit. A 4000 u., D 400 u., thiamine mononitrate 3 mg., riboflavin 2 mg., nicotinamide 10 mg., ascorbic acid 60 mg., pyridoxine HCl 3 mg., cobalamin 3 mcg., calcium pantothenate 1 mg., folic acid 1 mg., calcium carbonate 625 mg., ferrous fumarate 120 mg., magnesium 0.15 mg., zinc 0.025 mg., molybdenum 0.2 mg., iodine 0.1 mg., copper 0.15 mg./Tab. Bot. 100s, 1000s.
 Use: Dietary supplementation.
- * **PRAMOXINE HYDROCHLORIDE, N.F., N.N.D. 1962.** Cr., Jelly, Soln., N.F. gamma-Morpholinopropyl-4-N-butazyphenyl ether HCl. See: Tranothane HCl (Abbott)
- W/Diperodon HCl. See: Nascuta, Creme (Calumbeus)
- W/Hydrocortisone, polymyxin B sulfate, neomycin, propylene glycol, glycerin. See: Aurocart, Drops (Calumbeus)
- PRANONE.** (Schering) Ethisterone Tab 5 mg., 10 mg., 25 mg., Bot. 20s, 40s, 100s, 250s.
 Use: Orally; corpus luteum therapy.
- PRANTAL CREAM 2%.** N.N.D. 1962. (Schering) Diphenanthyl methylsulfate 20 mg./Oint. water-washable base Tube 50 Gm.
 Use: Dermatitis of many varieties.
- PRANTAL METHYLSULFATE, N.N.D. 1962.** (Schering) Diphenanthyl methylsulfate. Tab. 100 mg, Bot. 100s. Vial 25 mg./cc. 1s, 6s.
 Use: Anticholinergic.
- PRANTAL METHYLSULFATE RETABS, N.N.D. 1962.** (Schering) Diphenanthyl Retabs (50 mg. in core & 50 mg. outer coat) Bot. 100s, 1000s.
 Use: (1 q 8 h) Anticholinergic.
- W/Phenobarbital. (Schering) Prantal 100 mg., phenobarbital ¼ gr./Tab. Bot. 100s
 Use: (1 q.i.d.) Anticholinergic w/sedative.
- PRANTURON.** (Schering) Cholinic Ganodotrophin urine. Vial 5000 I.U., 10 cc.
 Use: I.M.; cryptorchidism.
- PRECALCIN.** (Walker) Calcitriol phosphate 0.5 Gm., bone phosphate 0.15 Gm., Vit. A acetate 2000 u., D 400 u., thiamine HCl 3 mg., riboflavin 2 mg., nicotinamide 10 mg., ascorbic acid 30 mg., cal. pantothenate 2.5 mg., pyridoxine HCl 3.33 mg., folic acid 0.33 mg., B-12 1 mcg., flavinol 33.3 mg., ferrous gluconate 100 mg., 8 trace elements/Cop. Bot. 100s, 500s, 1000s
 Use: Pregnancy & lactation.
- PRECALCIN-D.** (Walker) Cal. lactate 1 Gm., hesperidin complex 20.0 mg., Vit. A 5,000

* Generic name recognized by N.F., U.S.P. or A.M.A.

v, D 2 500 u, E 20 1U, K 0.5 mg, B 1 30 mg, B 2 30 mg nicotinamide 20 D mg, cal pentothenate 5 D mg, Vit B 6 2 5 mg, folic acid 0.5 mg, Vit B 12 20 mcg, C 750 mg, ferrous fumarate 1000 mg copper, mang, cobalt, zinc, mag, molybdenum, iodine pot, fluorine/Cop Rot 60s 300s

Use: Prenatal nutrition

PRECALCIN LACTATE (Walker) Calcium lactate 400 mg, Vit A 2000 u, D 400 u, B 1 3 mg, B 2 2 mg, C 30 mg, B 6 3 33 mg, B 12 1 mg, nicotinamide 10 mg, calcium pentothenate 25 mg, folic acid 0.33 mg, ferrous gluconate 100 mg, flavinol 100 mg, and B trace elements/Cop orange yellow Bot 100s, 500s, 1000s

Use: Phosphorus free prenatal therapy

PRECEPTIN (Ortho) Vaginal gel containing active ingredients ricinoleic acid & p diiso butylphenoxy polyethoxyethanol in base of synthetic gel forming agents Tube with or without applicator 3 oz

Use Effective gel to be used without dia phragm for control of conception

PRECIRUS, (Rorer) Homatropine methylbromide 2.5 mg, pentobarbital Sod 8 mg, phenobarbital 8 mg/Cop Bot 40s, 250s 1000s

Use: Orally (1 q 3-4 h); antispasmodic & sedative

PRE-CREATINE, (Mercury) Glycocyamine 100 mg, betaine 400 mg/Cop Bot 50s, 100s

Use: Dietary supplement for intermediary metabolism of body

PREDATEN, (Smith Miller & Patch) Ammonium chloride 3 gr, aspirin 2 gr, atropine sulfate 1/300 gr, caffeine 1/4 gr/Tab Bot 50r, 1000s

Use: Orally (1-2 tid pc) analgesic & antipyretic

PRE-DAYLIN DROPS, (Abbott) Vit A 5,000 u, D 1,000 u, C 50 mg/0.6 cc Bot w/ dropper 15 cc, 50 cc.

Use: Infant vitamin supplement

PREDEMA (Brayten) 2 Amino-2 methyl 1 propanol 8 bromo theophyllinate (Pembrom) Use: Pre-menstrual edema and tension

PREDICAINE, (Rhodes Pharm) Tyrothricin 0.5 mg, benzocaine 5%, p chlor m xylenol 1%, methylbenzethonium chloride 0.6%/Gm Oint, Tube 1/4 oz

Use: Topically skin infections.

PREDICORT-C, (Crystal) Ascorbic acid 50 mg, acetylsalicylic acid 5 gr, Prednisone 0.75 mg/Tab Rot 100r, 1000s

Use: Rheumatic arthritis, osteoarthritis, Intestinal gout

PREDICTARS (Drug Specialties) Anhydro-

progesterone 50 mg, ethinyl estradiol 0.03 mg/Tab Bot 24s

Use Pregnancy test amenorrhea, infertile, habitual abortion

PREDILOAMIDE (Delta) Prednisolone 0.005 Gm, salicylamide 0.25 Gm, Vit B 1 0.005 Gm/Tab

Use Treatment of arthritis

PRED-KAM, (First Texas) Prednisolone 5 mg dihydroxyaluminum aminoacetate 300 mg/Tab Bot 30s, 100s

Use: Antiarthritic

PREDLON, (Drug Specialties) Prednisolone 5 mg/Cop, released over 6 to 10 hour period Bot 30r, 100s

Use Timed disintegration steroid

PREDMYCIN, (Allergan) Prednisolone 0.2%, neomycin sulf 0.5%, phenylephrine HCl 0.12% Bot dropper 5 cc

Use: Infections of the eye

PREDMYCIN OPHTHALMIC OINTMENT (Allergan) Prednisolone alcohol 0.2%, neomycin sulfate 0.5%/Gm Applicator Tube 3.5 Gm

Use: Optic

PREDNAMIN (Dome) Prednisone 2.5 mg, chlorpheniradomine maleate 20 mg, ascorbic acid 250 mg/Tab Bot 30s 100s

Use: Treatment of bronchial asthma

PREDNE-DOME, (Dome) Prednisolone Tab 1 mg, 5 mg Bot 30r, 100s

Use: Allergic or inflammatory dermatoses

PREDNE-DOME & NEO-PREDNE-DOME AEROSOL SPRAY, (Dome) Predne Dome contains purified prednisolone, 20 mg/60 Gm Neo Predne Dome has neomycin sulfate, 20 mg/60 Gm added 60 Gm dispensers

Use: Eczematous and pruritic skin eruptions

PREDNEFRIN (Allergan) Prednisolone acetate 0.12%, phenylephrine HCl 0.12% Sol Bot 5 cc, 10 cc

Use Treatment of ophthalmic inflammation

PREDNEFRIN FORTE OPHTH. SUSP, (Allergan) Prednisolone acetate 1.0%, phenylephrine HCl 0.12%, in sterile, aqueous non irritating suspension Plar Bot 5 cc

PREDNEFRIN-S, (Allergan) Prednisone 0.2%, phenylephrine HCl 0.12% Sol Bot 5 cc

Use: Inflammatory ocular disorders

PREDNICHLOR, (Vitamix) Prednisone 0.75 mg, chlorpheniradomine maleate 4 mg/Tab Bot 100s

Use: Treatment of asthma and allergies

PREDNICIDIN, (Crocker Borne) Prednisone 0.2%, neomycin sulfate 0.5%, phenylephrine HCl 0.08%, gramicidin 0.005%/Lactival 7.5 cc

- Use: Allergic, inflammatory, infectious conjunctivitis.
- PREDNIS**, (Arlington-Funk) Prednisolone 2.5 mg. & 5 mg./Tab. Buf. 100s, 1000s.
Use: Treatment of rheumatic disorders
- PREDNIS-CVP**, (Arlington-Funk) Prednisolone 4 mg., bioflavonoid complex, ascorbic acid, aluminum hydroxide, magnesium oxide—each 100 mg./Cap. Buf. 30s, 100s
Use: Antiarthritic.
- ***PREDNISOLONE**, U.S.P., N.M.D. 1962; Oint., Tabs., U.S.P. Δ 1,2-Pregnadione-3,20-dione-11,17 α, 21-triol. Delta-1-hydrocortisone. 1-Dehydro-hydrocortisone. Metacort-ondralone.
- See: Cortalone, Tab. (Truxton)
Delta-Cortef, (Upjohn)
Fernisolone, Tab. (Fernole)
Hydeltra. (Merck Sharp & Dohme)
Meti-Derm, (Schering)
Meticortelone, Tab. & Susp. (Schering)
Paracort, Tab. (Parke, Davis)
Predne-Dame, Tab. (Dome)
Prednis, Tab. (U. S. Vitamin)
Stecane, Tab. (Pfizer)
Steralone, Tab. (Rowell)
Ulocort (Testagar)
- W/Ascorbic acid, mephenesin, salicylamide.
See: Salimeph/Prednisolone, Tab. (Kremers Urban)
- W/Aspirin, See: Cordex, Tab. (Upjohn)
- W/Aspirin, sod. ascorbate, aluminum hydroxide gel. See: Tempagen Forte, Tab. (Merck Sharp & Dohme)
- W/Atropine sulfate, methylcellulose. See: Isopta Hydrapred, Sus. (Alcon)
- W/Carisapradal. See: Somacort, Tab. (Wallace)
- W/Cytirus bioflavonoids, Vit. C, K. See: Sarlnoid-3, Caps. (Dumas-Wilson)
- W/Dihydroxy aluminum aminoacetate. See: Pred-Kom, Tab. (First Texas)
- W/Hexochlorophene. See: Lipalone, Cream (Spiri)
- W/Hydrocortisone acetate, prednisone, neomycin sulfate. See: Pansteroid, Oint. (Kenwood)
- W/Hydroxyquine HCl. See: Atoraxoid, Tab. (Pfizer)
- W/Iodochlorohydroxyquin. See: Iodo-Lipalone, Cream (Spiri)
- W/Magnesium trisilicate, aluminum hydroxide gel. See: Co-Hydeltra, Tab. (Merck Sharp & Dohme)
- W/Mepramate, dried aluminum hydroxide gel. See: Mepralane, Tab. (Merck Sharp & Dohme)
- W/Naphasaline HCl, chlorpheniramine maleate. See: Vasopred (Smith, Miller & Patch)
- W/Neomycin, sodium propionate. See: Otabione. (White's)
- W/Neomycin sulfate.
See: Hydeltrafin Topical Lotion. (Merck Sharp & Dohme)
Meti-Derm w/Neomycin, Oint. (Schering)
Predmycin Ophthalmic Ointment (Allergon)
- W/Neomycin sulf., phenylephrine HCl. See: Predmycin, Sol. (Allergon)
- W/Neomycin sulfate, phenylephrine HCl, gramicidin. See: Prednicidin, Oph. Soln. (Crookes Barne)
- W/Paraflex, tylenol. See: Parafon, Tab. (McNeil)
- W/Phenylephrine HCl.
See: Isopta Prednisolone (Alcon)
Prednetrin S, Liq. (Allergon)
- W/Phenylephrine HCl, methylcellulose. See: Isopta Prednisolone, Susp. (Alcon)
- W/Pot. chloride. See: K-Predne Dome, Tab. (Dome Chemicals)
- W/Pot. salicylate, ascorbic acid, pot. p-aminobenzoate. See: Neocylone, Tab. (Central)
- W/Salicylamide, ascorbic acid, mephenesin. See: Salimeph Forte with Prednisolone, Tab. (Kremers-Urban)
- W/Salicylamide, Vit. B-1. See: Predlamide, Tab. (Delta)
- W/Sulfacetamide sodium, methylcellulose. See: Isopta Celopred, Susp. (Alcon)
- W/Sodium sulfacetamide, phenylephrine HCl. See: Sultopred (Professional Pharmacol)
- W/Tetrahydrozalone HCl.
See: Cor-Tyzanol (Pfizer)
Cor-Tyzine, Drops (Pfizer)
- W/Tetrahydrozalone HCl, neomycin Sulf. See: Neacor-Tyzine, nasal drops (Pfizer)
- W/Theophylline, ephedrine sulfate, phenobarbital. See: Prednisophine (Vitamix)
- W/White's Vitamin A & D Ointment. See: White's Vitamin A & D Ointment w/Prednisolone (White Labs)
- ***PREDNISOLONE ACETATE**, U.S.P., N.M.D. 1962; Susp., U.S.P.
- W/Neomycin sulfate. See: Neo-Delta-Cortef Acetate Oint. (Upjohn)
- W/Neomycin sulfate & phenylephrine HCl. See: Neo-Delta-Cortef Nasal Spray. (Upjohn)
- W/Neomycin, Sod. propionate. See: Otabione (White)
- W/Neomycin sulfate, sulfacetamide sod. See: Metimyd, Oint. W/Neomycin. (Schering)
- W/Phenylephrine HCl. See: Prednetrin, Sol. (Allergon)
- W/Sodium sulfacetamide, phenylephrine

HCl See Blephamide Ophthalmic Liquifilm (Allergon)
 W/Sodium sulfacetamide phenylephrine NCl, propylene glycol, sodium phosphate mono basic sodium phosphate dibasic sodium thiosulfate in a methylcellulose base See: BufOpta Sulfapred (Professional Pharmacol)
 W/Sulfacetamide See Metimyd, Ophth Susp (Schering)
 ***PREDNISOLONE BUTYLACETATE, NND 1962** See Hydextra T B A (Merck Sharp & Dohme)
 ***PREDNISOLONE CYCLOPENTYLPROPIONATE** d1,4 pregnadiene 3 20-dione 11,17a diol 21 cyclopentylpropionate(prednisolone 21 cyclopentylpropionate)
 ***PREDNISOLONE 21-PHOSPHATE SOD.** NND 1962 See: Hydextra Sol, Prp (Merck Sharp & Dohme)
 W/Neomycin sulf See Neo Hydextra Sol, Viat (Merck Sharp & Dohme)
PREDNISOLONE SODIUM SUCCINATE, USP A hemisuccinate See Meticortelone Soluble (Schering)
PREDNISOLONE TERTIARY-BUTYLACETATE See Hydextra T B A, Viat (Merck Sharp & Dohme)
 ***PREDNISONE, USP NND 1962, Tabs,** USP Δ1,4 Pregnadiene 17 α, 21 diol 3 11 20 trione 1 Dehydro cortisone Metacort androcin
 See Colone Tab (Truxton)
 Deltasone, Tab (Upjohn)
 Delta, Tab (Merck Sharp & Dohme)
 Listacort (Testogar)
 Metasone, Cap (Wilfred)
 Meticorten, Tab (Schering)
 Paracort, Tab (Parke, Davis)
 Sterapred Tab (Mayrand)
 W/Aluminum hydroxide and mag trisilicate See: Co-Delta, Tab (Merck Sharp & Dohme)
 W/Ascorbic acid dicalcium glutamate potassium salicylate salicylamide potassium p-aminobenzoate See: Glutalate w/Prednisone Cap (Dumas Wilson)
 W/Ascorbic acid, salicylamide n acetyl para-aminophenol See Arthralgen PR, Tabs (Whittier)
 W/Aspirin, ascorbic acid, aluminum hydroxide See: Sigmagen, Tab (Schering)
 W/Cal pantothenate, dried aluminum hydroxide gel, mag trisilicate See: Predsem, Tab (Massengill)
 W/Chlorpheniramine maleate See: Prednichlor (Vitamax)
 W/Chlorphenopyridamine maleate ascorbic acid See: Prednamin, Tab (Danel)
 W/Chlor Trimeton See: Metretan (Schering)
 *Generic name recognized by NF USP or A.M.A.

W/Hydrocortisone acetate, prednisolone, neomycin sulfata See: Pansteraid, Qint (Kenwood)
 W/Magnesium trisilicate, aluminum hydroxide gel See: Co Delta, Tab (Merck Sharp & Dohme)
 W/Methylglantenedial, magnesium trisilicate, aluminum hydroxide gel, calcium carbonate, magnesium carbonate, guar gum, oat binder (Egraine) See: Stenisane, Tab (Organon)
 W/Phenylbutazone, dried aluminum hydroxide gel, magnesium trisilicate, homatropine methylbromide See: Sterasolidin, Cap (Geny)
 W/Phenylephrine HCl See: Prednefrin S, Sol (Allergon)
 W/Phenyltolaxamine dihydrogen citrate chlorpheniramine maleate, ephedrine sulfate See Esteril Tab (Ensley Scoll)
 W/Pot salicylate, cal pantothenate See: Salfcort Delta, Tab (Massengill)
 W/Salicylamide ascorbic acid See: Aspred C, Tab (Money)
 W/Salicylamide, Vit C, aluminum hydroxide See Loupred Tablets (Louisans)
 W/Salicylamide hesperidin conc, ascorbic acid Vit K, alum hydrox gel See Holv cort Tabs (Moller)
 W/Theophylline, ephedrine HCl, phenobarbital See: Deltasmyl (Roussel)
 W/Triafon See Steroinl, Tab (Schering)
PREDNISONE COMBINATION, (Massengill)
 Prednisone 50 mg, cal pantothenate 10 mg, dried aluminum hydroxide gel 0.2 Gm, mag trisilicate 0.1 Gm/Tab Bat 30s 100s
 Use Acute and subacute phases of various rheumatic conditions
PREDNISOPNINE, (Vitamax) Prednisolone 0.5 mg, theophylline 2 gr, ephedrine sulfate 3/4 gr phenobarbital 1/4 gr/Tab
 Use Asthmatic and antihistaminic
PREDNITABS, (Leeds) Prednisone 0.75 mg mephensan 250 mg, mephobarbital 10 mg/Tab Bat 50s
 Use: Antiarthritic
PREDNYL, (Arlington Funk) Prednisolone 1 mg, salicylamide 5 gr, bioflavonoid compound 33.3 mg, ascorbic acid 33.3 mg, aluminum hydroxide 50 mg/Tab Bat 100s
 Use: Anti inflammatory
PREDSEM, (Massengill) Prednisone 50 mg, cal pantothenate 10 mg, dried aluminum hydroxide gel 0.2 Gm, mag trisilicate 0.1 Gm/Tab Bat 30s, 100s, 1000s
 Use Acute phase of arthritis
PRIGENAL TABLETS, (Utter) Vit A 1500 u, D 200 u, B 1 1 mg, B 2 2 mg, B 6 2.5

- mg., C 30 mg., B-12 1 mcg., niacinamide 5 mg., menadione 0.25 mg., folic acid 0.34 mg., Vit. E 2 mg., cal. pyrophosphate 7.5 gr., ferrous sulf. 3 gr./Tab. Bot. 100s, 500s, 1000s.
 Use: Prenatal vitamin therapy.
- PRIGENT.** (Beutlich) Calcium carbonate 300 mg., Vit D 300 I.U., ferrous sulfate, asiccated 130 mg., ascorbic acid 100 mg., hesperidin complex 10 mg./Tab. Bot. 100s
 Use: Nutritional supplement.
- PREGNANCY TEST.** See: C.P.T. (C.P.T. Lab)
- PREGNANCY TEST TABLETS.** (Westward) Ethisterone 50 mg., ethinyl estradiol 0.03 mg./Tab. Bot. 24s
 Use: Treatment of amenorrhea, infertility and in prevention of habitual abortion.
- PREGNICAL.** (Uimer) Vits. A 6000 u., D 400 u., C 100 mg., R-1 1.3 mg., R-2 2 mg., iron 15 mg., calcium 150 mg./Tab Bot. 60s.
 Use: Nutritional supplement.
- PREGNENINOLONE.** See: Ethisterone, U.S.P. (Various Mfr.)
- *PREGNENOLONE.** Synthetic steroid hormone, 3-hydroxy-20-keto-pregene-5
 See: Enelone, Prep. (Funk)
 Natalone, Prep. (Notlon Orug).
 Prenolon, Prep. (Schering).
 Shormone, Prep. (Merck Sharp & Dohme)
- PREGNENOLONE ACETATE.**
 See: Enelone, Prep. (Funk)
 Enescorb, Tab. (Arlington-Funk)
 Prenolon, Tab. (Schering).
 Egnorone, (Winthrop Labs.)
 Shormone, Tab. (Merck Sharp & Dohme)
 W/Ascorbic acid. See: Enescorb, Tab. (Funk)
 W/Beta-estradiol, methyl testosterone & monobasic amphetamine phosphate. See: Longevan, Cap. (Northwest Lab.)
- *PREGNENOLONE SUCCINATE.** 3-hydroxy-5-pregnen-20-one, hydrogen succinate. See: Panzalone (Doak)
- PREGNYL.** N.N.D. 1960. (Organon) Human chorionic gonadotrophin. Vial 10,000 I.U., dry pow., 10 cc.
 Use: I.M.; cryptorchidism.
- PREGULAR.** (Bentley) Anhydrohydroxyprogesterone 50 mg., ethinyl estradiol 0.03 mg./Tab. Bot. 48s.
 Use: Pregnancy test; menstrual regulator.
- PRELAC.** (Dumas-Wilson) Ferrus gluconate 3 gr., cal. lactate 350 mg., Vit. A 1000 u., D 400 u., B-1 2 mg., B-2 2 mg., niacinamide 5 mg., C 20 mg., w/trace elements/Cap Bot 100s, 1000s.
 Use: (3 daily) A prenatal dietary supplement
- PRELACTON IMPROVED.** (Table Rock Labs) Vit. A 6000 u., D 400 u., B-1 1.5 mg., B-2 2 mg., B-12 2 mcg., B 6 1 mg., C 50 mg., niacinamide 15 mg., cal. pantothenate 3 mg., ferrous fumarate 15 mg., cal. carbonate 375 mg., hesperidin complex 50 mg., dioctyl sod sulfosuccinate 50 mg./Cap. Bot. 100s, 500s.
 Use: Stool softener w/vitamin and mineral supplement
- PRELANS.** (Lanett) Vit. A 4000 U.S.P. units, B-1 0.5 mg., B-2 2 mg., niacin 10 mg., R-12 2 mcg., D 400 units, C 50 mg., Co 200 mg., Fe 30 mg./Tab Bot 100s
 Use: Prenatal Vitamin-mineral supplementation.
- PRELUDIN.** N.N.D. 1962 (Geigy) 2-Phenyl-3-methyl tetrahydro-1,4 oxazine HCl, Phenmetrazine HCl, Tab. 25 mg., Bot. 100s. Endurets (75 mg) Bot 100s.
 Use: Orally (1 t.i.d., a.c.), steady progressive weight loss in overweight patients.
- PRELU VITE.** (Geigy Pharm) Preludin (phenmetrazine HCl) 25 mg., Vit A 2000 U.S.P. units, Vit D 200 U.S.P. units, thiamine mononitrate U.S.P. 20 mg., riboflavin U.S.P. 20 mg., niacinamide 200 mg., calcium pantothenate U.S.P. 30 mg., pyridoxine HCl U.S.P. 10 mg., Vit. B 12 0.5 mcg., ascorbic acid 37.5 mg., iron 50 mg., calcium 1400 mg., phosphorus 1080 mg., iodine 0.1 mg., copper 10 mg./Cap Bot. 100s.
 Use: Nutritional and anorectic therapy.
- PREMARIN.** N.N.D. 1962. (Ayerst) Estrogenic substance conjugated Water-soluble conjugated estrogens derived from pregnant mares' urine Tab. 2.5 mg., 1.25 mg., Bot. 20s, 100s; 0.625 mg., 0.3 mg., Bot. 100s, 1000s liq. 0.625 mg/4 cc. Bot. 120 cc.
 Use: Menopausal syndrome
- PREMARIN N. C.** (Ayerst) Premarin 0.625 mg/Gm., hydrocortisone 0.1%, in a non-liquefying base. Tube w/Applicator 1 oz the Vaginitis, vulva dermatoses.
- PREMARIN VAGINAL CREAM.** N.N.D. 1962. (Ayerst) Conjugated estrogens. Regular, 0.625 mg/Gm., 125 mg/Gm., Jar 1 oz, 2 oz Tube 1 1/2 oz. Vaginal, 0.625 mg/Gm. Tube with or without applicator. 1 1/2 oz.
- PREMARIN INTRAVENOUS.** N.N.D. 1962. (Ayerst) Conjugated estrogens from pregnant mares' urine Vial (20 mg) 5 cc w/diluent.
 Use: Hemostasis in functional uterine bleeding.
- PREMARIN LOTION.** N.N.D. 1962. (Ayerst) Conjugated estrogens lat. (1 mg/cc) Bot 60 cc.

* Generic name recognized by N.F., U.S.P. or A.P.A.

Use: Sebarrheic alopecia, seborrhea, acne vulgaris

PREMARIN W/MEPROBAMATE. (Ayerst)

Premarin 0.4 mg, meprobamate 400 mg / Tab Bot 60s, 500s See: PMB 200 (Ayerst)

Use: Control of the menopausal syndrome

PREMARIN W/METHYLTESTOSTERONE

(Ayerst) Conjugated estrogens equine (Premarin) 1.25 mg, methyltestosterone 10 mg / Yellow Tab

Conjugated estrogens equine (Premarin) 0.625 mg, methyltestosterone 5 mg / Red Tab Bot 100s, 1000s

Use: Estrogen androgen therapy

W/Methyltestosterone, d desoxyephedrine

HCl & vitamins See: Mediatric, liq (Ayerst)

PREMARIN W/PHENOBARBITAL. (Ayerst)

Conjugated estrogens equine (Premarin) 0.625 mg, phenobarbital 1/2 gr / Tab Bot 100s, 1000s

Use: Climacteric & its nervous manifestations

PREMATINIC. (Ingram) Ferrous fumarate 180 mg, Vit C 100 mg, B 6 35 mg. Cal carbonate 500 mg, Vit D 400 u phos phorus and sodium free / Tab Bot 100s

Use: Anemia

PRE-MENS (Purdue Frederick) Ammonium chloride 5 gr, homatropine methylbromide 0.5 mg, caffeine 33 mg, Vit B 1 2 mg, B 2 1 mg, B 6 0.5 mg, Cal pantothenate 1 mg, niacinamide 5 mg / Tab Bot 60s

Use: (2 1 1 d) Premenstrual tension

W/d Amphetamine (Purdue Frederick) d

Amphetamine sulfate 25 mg, ammonium chloride 330 mg, homatropine methylbromide 1 mg, Vit B 1 2 mg, B 2 1 mg, B 6 0.5 mg, Cal pantothenate 1 mg, niacinamide 5 mg / Tab Bot 60s

Use: 2 1 1 d, p c; premenstrual tension

PREMENSTRUAL TENSION

See: M. Minus 5, Tab (Whittier)

Matian Sickness

Neo Bramth Tab (Broyten)

Neoparabram Tab (Central)

Neo pyrapar, Tab (Ulmer)

Pamabrom (Various Mfr)

Pambramat, Tab (Whittier)

Pre Mens Tab (Purdue Frederick)

Pyranisamine Bromotheaphyllinote (Various Mfr)

Sanbrom Tab (Smith Darsey)

PREMESED. (Columbus) Vit B-6 10 mg

belome HCl 400 mg, levulose 112 Gm dextrose 112 Gm / fl oz Bot 3 oz, 1 pt 1 gal

Use: Control of functional nausea and vomiting

PREMOCEL (Prema) Methylcellulose Tab

0.5 Gm Bot 100s, 1000s

Use: Constipation

PREMOCILLIN. (Prema) Procaine penicillin G

300M u/cc Aq susp, Vial 1 dose Powder 2 oz bottles

Use: Penicillin therapy

PREMOCONES. (Prema) Tyrothricin 1 mg w/beezocaine, bismuth subgallate, balsam of Peru & zinc oxide / Supp Box 12s.

Use: Internal & external hemorrhoids

PREMODRIN. (Prema) dl Desoxyephedrine HCl Tab 10 mg, Bot 100s, 1000s

Use: Appetite depression

PREMOTRACIN. (Prema) See: Bacitracin USP

PREMOGEN. (Prema) Prednisone 0.75 mg, aspirin 325 mg, Vit C 20 mg, aluminum hydroxide 75 mg / Tab Bot 100s, 1000s

Use: Treatment of rheumatism

PRENALAC. (Lilly) Thiamine mononitrate 1.5 mg, riboflavin 3 mg, Vit B 12 7.5 mg, ascorbic acid 100 mg, nicotinamide 15 mg, folic acid 0.4 mg, Vit A 6000 u, D 400 u, iron 15 mg, Ferrous sulfate 75 mg, B 6 6 mg, Cal pantothenate 6 mg, dicalcium phosphate equivalent to Cal 1.5 Gm, Phos 14 Gm / 6 pulvules Bot 100s, 500s, 1000s

Use: Nutritional supplementation

PRENALIN-O (S J Tutag & Co) Ferrous fumarate 150 mg, rutin 10 mg, folic acid 0.25 mg, B 12 2 mcg, Vit K 0.25 mg, iodine 0.15 mg, Vit A 4000 USP u, Vit D 400 USP u, Vit B 1 2 mg, Vit B 2 2 mg, Vit B 6 0.8 mg, niacinamide 10 mg, Vit C 50 mg, calcium 600 mg / Cop Bot 100s, 1000s Also available in nursing bottles of 100 and plastic boxes of 50

Use: Treatment of Pre and Post Natal Nutritional Stress

PRENAMEL. (Marsh Emary Labs) Vit A 5000 u, D 400 u, B 1 2 mg, B 2 2.5 mg, niacin 12 mg, Vit B 6 10 mg, C 75 mg, calcium carbonate 600 mg, ferrous fumarate 160 mg / Tab Bot 100s

Use: Nutritional supplement

PRENAPHOS. (Chicago Pharm) Ferrous sulfate 90 mg, dicalcium phosphate 250 mg, Vit A 1000 u, D 100 u, B 1 0.25 mg, B 2 0.334 mg, R 6 1 mg, B 12 w/intrinsic factor 1/12 u, C 35 mg, K 0.25 mg, nicotinamide 3 mg, alpha tocopherol acet 0.5 mg, folic acid 0.167 mg, calcium pantothenate 0.5 mg, menaquinone 25 mg / Tab Bot 100s, 1000s

Use: Prenatal supplement

PRENASUP (Chicago Pharmacal) Complete vitamin formula w/minerals Tab, Bot 100s, 200s

Use: Dietary supplement

PRENAT (Minnesota Pharmaceutical) New name for Preacal

PRE-NATAL. (Horton & Converse) Vit A

2000 u., D 200 u., C 35 mg, B-1 2 mg, B-2 2 mg, B-6 3.3 mg, niacinamide 8 mg, cal. pantothenate 2 mg., hesperidin complex 30 mg, folic acid 0.1 mg., K 0.33 mg, B-12 1 mcg./Tab. Bot. 100s, 500s

Use: Dietary supplement in pregnancy & lactation

PRE-NATAL (West-Ward) Vit. A 2000 u., D 400 u., B-1 2 mg., B-2 2 mg., B-6 0.35 mg., B-12 1.5 mcg., niacinamide 10 mg., Cal. pantothenate 2 mg., folic acid 1 mg., Vit. C 35 mg., E 3 mg., dicalcium phosphate anhy. 900 mg., iron 6 mg., Mang. 0.2 mg., iodine 0.1 mg., copper 0.5 mg./Cap. Bot. 100s, 500s.

Use: Dietary supplement.

PRENATAL (Lederle) Vit. A 2000 u., D 400 u., thiamine HCl 2 mg., riboflavin 2 mg., ascorbic acid 35 mg., niacinamide 7 mg., Vit. B-12 1 mcg., folic acid 1 mg., Cal., Phos., iron, Mang./Cap. Bot. 100s, 1000s.

Use: Dietary supplement.

PRENATAL (Boyle) Vit. A 2000 u., D 200 u., B-1 1.66 mg., B-2 0.80 mg., niacinamide 16.60 mg., Vit. C 33.3 mg., ferrous gluconate, Cal. 0.25 Gm./Tab Bot. 100s.

Use: (1 t i d) For pregnant patients.

W/Vit. B-12 (Boyle) Cal. 0.75 Gm., ferrous gluconate 195 mg., Vit. A 6000 u., D 600 u., B-1 5 mg., B-2 2.5 mg., niacinamide 50 mg., Vit. C 100 mg., Vit. B-12 18 mcg., folic acid 0.75 mg., iodine 0.15 mg., Mang. 4.5 mg., copper 1 mg., zinc 2.25 mg., cobalt 0.15 mg., molybdenum 0.15 mg., Mag. 15 mg., nickel 0.15 mg./3 Tab. Bot. 100s.

Use: (1 t i d) For pregnant patients.

PRENATAL (Stuart) See: Stuart Prenatal, Tab. (Stuart)

PRENATAL DBI-KAPS. (Lederle) Vit. A 2000 u., D 400 u., B-1 2 mg., B-2 2 mg., niacinamide 7 mg., B-12 1 mcg., K 0.5 mg., C 35 mg., folic acid 1 mg., calcium 250 mg., iron 6 mg., ferrous sulf. assic. 20 mg., mang. 0.12 mg., phosphorus 190 mg./Cap. Bot. 100s.

Use: Dietary supplement in pregnancy and lactation

PRE-NATAL SUPPLEMENT. (West-Ward)⁴ Vit. A 2,500 u., D 250 u., B-1 2 mg., B-2 2 mg., niacinamide 10 mg., B-6 2 mg., cal. pantothenate 2 mg., folic acid 0.25 mg., B-12 2 mcg., C 37.5 mg., hesperidin 35 mg., B-12 W/intrinsic factor conc. 1/15 u., plus¹ minerals/Cap. Bot. 100s, 1000s.

Use: Dietary supplement in pregnancy and lactation.

PRENATUM TABLETS. (Ascher) Vit. A 6000 u., D 400 u., C 100 mg., B-1 3 mg., B-2 2 mg., B-6 5 mg., B-12 2 mcg., niacinamide

10 mg., menadione 1 mg., ferrous fumarate 150 mg., calcium 200 mg./Tab Bot. 100s

Use: Pregnancy and lactation.

PRE-NATUM-CAL. (Ascher) Same formula as Prenatum but with no Phos. & Cal. 375 mg as citrate & carb. Bot. 100s, 500s, 1000s.

Use: Prenatal care.

PRENATUM B-P. (New Mexico Pharm) Vit. A 1500 u., D 250 u., B-1 1 mg., B-2 1.5 mg., niacinamide 10 mg., B-6 0.1 mg., C 25 mg., folic acid 0.1 mg., calcium lactate 500 mg., ferrous gluconate 45 mg., E 1/4 u., menadione DDS mg., cobalt 0.1 mg., copper 0.1 mg., iodine 0.1 mg., manganese 1 mg., magnesium 5 mg., molybdenum 0.1 mg., potassium 5 mg., zinc 0.1 mg., cyanocobalamin 0.5 mg., methionine 5 mg., inositol 5 mg., choline bitartrate 5 mg./Cap. Bot. 100s.

PRENAUSEN. (Walker) Vit. B 6 10 mg., B-1 10 mg., potassium phosphate 100 mg., potassium citrate 100 mg., carbohydrates 750 mg./trache. Box 24s

Use: Relief of nausea and vomiting of pregnancy.

PSE-NAY. (Ortho) lanolin, lecithin, cholesterol cream. Jar 3 oz.

Use: Treatment of dry skin.

PRENDIBOL. (Squibb) Not listed Diethyl propanediol. Tab. 0.5 Gm., Bot 100s, 1000s.

Use: (1-2 q i d) Muscle relaxant.

PRENOLDN. (Schering) Pregnanolone. Vial, aq. susp. (100 mg/cc) 10 cc., Box 1s.

Use: I.M.; rheumatoid arthritis.

PRENOLON ACETATE. (Schering) Pregnanolone acetate. Tab. 100 mg., Bot. 24s.

Use: Orally; rheumatoid arthritis.

***PRENYLAMINE.** (Lloyd Bros) N-(3,3-Diphenylpropyl)-alpha-methylphenethylamine See: Segontin.

PREOBIN. (Person & Covey) Calcium 250 mg., B-6 10 mg., ferrous fumarate 120 mg., Vit. C 100 mg., hesperidin 30 mg., lysine HCl 50 mg., diacyl sod. selfosuccinate 15 mg., Vit. A 6000 u., D 400 u., B-12 5 mcg., K 0.5 mg., B-1 3 mg., B-2 3 mg., calcium pantothenate 5 mg., niacinamide 15 mg./3 Tabs. Bot. 100s, 1000s.

Use: Dietary supplement during pregnancy & lactation.

PREOBIN FORTE. (Person & Covey) Calcium 250 mg., Vit. B-6 10 mg., ferrous fumarate 120 mg., Vit. C 100 mg., hesperidin 30 mg., lysine HCl 50 mg., diacyl sod. selfosuccinate 15 mg., Vit. A 6000 u., D 400 u., B-12 5 mcg., K 0.5 mg., B-1 3 mg., B-2 3 mg., calcium pantothenate 5 mg., niacinamide 15 mg./Cap. Bot 100s, 250s

Use: Vitamin & mineral supplement.

⁴ Generic name recognized by N.F., U.S.P. or A.M.A.

- PRESID.** (Merck & Co., Inc.) See Hydroxindolol HCl
- PRESSOR AGENTS**
See Ephedrine Related Compounds
Epinephrine Related Compounds
- PRESTOGEN MYOSOL.** (Ulmer) Estrone 2.5 mg, progesterone 25 mg/cc
Use: Treatment of amenorrhea, menorrhagia and habitual abortion
- PRETAB** (U S Standard Products) Calcium lactate 500 mg, ferrous gluconate 125 mg, ascorbic acid 30 mg and Vit D 400 u/Tab Bot 100s
Use: Prenatal supplement
- PRETHCAMIDE** Mix of crocthamide & cetraramide See: Mitaren (Geigy)
- PREVIM** (Sig Labs) Vitamin A 4000 USP units, D 600 USP units B 1 2 mg, B 2 2 mg B 6 1 mg C 90 mg, niacinamide 10 mg B 12 1 mcg, rutin 8 mg, ferrous fumarate 100 mg, oyster shell 480 mg, Mn 10 mg inositol 5 mg, choline 5 mg/2 Tab Bot 100s 1000s
Use: Prenatal vitamin mineral amino acid combination
- PRI-CORT.** (K & M Labs) Hydrocortisone alcohol 1/2%; dexamethasone HCl 1/4%; iodochlorohydroxyquin 3%; solution of coal tar 2%; Vit A & D—twice the amount as contained in Cod Liver Oil/Gm
Use: Eczematous eruptions
- PRICULEX.** (Elder) Caprylic acid 2%, copper caprylate 1%, salicylanilide 3%, p chlor meta xylene 2% Oint 2 oz Pow 1/2 oz
Use: Topically; fungicide
- PRI-HIST.** (K & M Labs) Phenylephrine HCl 5 mg, phenylpropanolamine HCl 20 mg, chlorpheniramine maleate 2 mg, acetaminophen 150 mg/Tab Bot 50s
Use: Antihistamine
- PRIMACAINE.** (Navacal) 2' Diethylaminoethyl 2 butyl-3-aminobenzoate HCl 1.5%/2 cc Anestubes tins 50s.
Use: Dental local anesthesia
- PRIMAQUINE BASE W/Camaquin base** See: Camaprim, Tab (Parke, Davis)
- *PRIMAQUINE PHOSPHATE, USP, NND 1962;** Tab, USP (Winthrop Labs) 8-4 amino 1 methylbutylamino 6-methoxyquinoline diphosphate Tab 26.5 mg, Bot 14s, 100s
Use: Prevents relapses in nearly all cases of vivax malaria
- PRIMASE.** (Bauer) Amylase 36 mg, pepsin 250 mg, pancreatin 100 mg, ketocholanic acids 20 mg/Tab Bot 50s
Use: Digestion
- PRIME-AVITE** (Carrtone) Vit D 2 400 u,
* Generic name recognized by NF, USP or A.M.A., ascorbic acid 30 mg/0.6 cc Drop Bot 1 oz
Use: Vitamin supplement
- PRIMESERP.** (Prime) Reserpine alkaloids 0.25 mg/Tab Bot 100s
Use: Antihypertensive and calming agent
- PRIMETRATE 20 Mg. Tabs** (Prime) Pentaerythritol tetranitrate, 20 mg/Tab Bot 100s
Use: Vasodilator coronary
- PRIMIC** (Prime) Pheniramine maleate 25 mg, pyrimidine maleate 25 mg, phenylpropanolamine HCl 50 mg/timed release tab Bot 100s
Use: Common cold and allergies
- PRIMICOL.** (Sherman) Colostrum obtained from primiparas during 28th week of pregnancy Set 1s, 3s & 6s test
Use: ID; diagnostic aid as a test for pregnancy
- *PRIMIDONE, USP, NND 1962;** Tabs, USP 5 Ethylhexahydro-5-phenylpyrimidine 4,6-dione See Mysoline Tab (Ayerst)
- PRIMOSTRUM.** A prep of primiparous colostrum See: Que Test (General Pharm Co)
- PRINADOL, NND 1962** (Smith, Kline & French) Phenazocine (2 mg/cc) Amp 1 cc Box 10s, 100s Vial 10 cc, Box 20s
Use: Narcotic analgesic
- PRIODAX, NND 1962** (Schering) (Iodoaliphonic Acid) Tab 0.5 Gm, Envelopes of 6 (1 dose); Box 1s, 5s, 25s, 100s; Pkg 100s, in 4 rolls of 250 tab
Use: Orally; cholecystography
- PRI-RON** (K & M Labs) Iron ammonium citrate brown 18 grn, liver fraction #1 3 grn, Vit B 1 10 mg, B 2 4 mg, B 12 20 mcg, alcohol 8%/oz Bot 12 oz; ferrous sulfate 104 mg, dexticated liver 60 mg, w/Vit B 1, B 2, niacinamide and Vit B 12 4 mcg/Tab Bot 60s
Use: Anemia
- PRISCOL** (Ciba) Old name for Priscoline
- PRISCOLINE, NND 1962** (Ciba) Talazoline Hydrochloride, USP; Tabs: USP Priscol (Benzosaline HCl) 2 Benzyl imidazoline HCl Elixirs 25 mg/4 cc Bot 1 pt Tab 25 mg-tantabs (80 mg) Bot 100s, 1000s Vial (25 mg/cc) 10 cc
Use: Relieves pain, promotes healing & increases blood supply to extremities.
- PRISILIDENE HYDROCHLORIDE.** See Alpha-prodine HCl
- PRIVADORN.** See: Bromisovalum, NP.
- PRIVINE HYDROCHLORIDE, NND 1962** (Ciba) Naphazoline HCl Jelly: 0.05%, Tube 20 Gm Ophth. sol.: 0.1%, 15 cc-

- Bot. w/dropper. Sal. 0.05%, Bot. 1 R. oz., 16 fl. oz., D. 1%, Bot. 4 fl. oz.
Use: Vasodilator.
- PBIVINE NEBULIZER.** (Ciba) 0.05% Plastic squeeze bot. 15 cc.
- PBIZOLE HCl.** (DuMont) Naphazoline HCl 0.05% and 1% Bot. 1 pf.
Use: Nasal congestion
- PRN.** (Bristol) Phenyltolazamine Citrate Tabs. (100 & 200 mg.) Bot. 50s
Use: Ataractic agent
- P.R.N.** (Palmedica) d-*amphetamin*e phosphate dibasic (N.F.) 5 mg., phenacetin 162 mg., aspirin 227 mg., phenobarbital 24 mg., amobarbital 24 mg./Tab. Bot. 100s, 1000s
Use: Relief of pain.
- PRO-ACET DOUCHE CONCENTRATE.** (Pro-Acet) Lactic, citric, & acetic acids, sod. lauryl sulfate, lactose, dextrose & sod. acetate. Pkg. polyethylene envelope 10 cc. Contents of 1 envelope to be diluted with 2 quarts of water.
Use: Douche.
- PRO-ACTASE.** (Ortho) Pro fibrinolysin (human) 10,000 u./Amp. with Amp. of diffusin 150 u. (hyaluronidase).
Use: Topical. Wound debridement.
- PROANTHELINE BROMIDE.** See: Probanthine
- PROBANA.** (Mead Johnson) Milk protein, banana pow. & dextrose. Pow. 1 lb.
Use: Infant food.
- PRO-BANTHINE, N N D.** 1962. (Searle) (Propantheline Bromide) beta-Diisopropylaminoethyl xanthene 9-carboxylate methobromide. Amp. 30 mg., Box 6s, 25s, 100s; Tab. 7½ & 15 mg., Bot. 100s, 1000s.
Use: (I t.i.d. & h.s.) Anticholinergic.
- W/Phenobarbital** ¼ gr. Tab., Bot. 100s, 1000s.
- PRO-BANTHINE W/DARTAL.** (Searle) Probanthine 15 mg., dartial 5 mg./Tab. Bot. 100s, 500s.
Use: Functional gastro-intestinal complaints.
- PRO-BANTHINE P.A.** (Searle) b-diisopropylaminoethyl xanthene-9-carboxylate methobromide 30 mg. (sustained-release beads)/Tab. Bot. 100s.
Use: Anticholinergic action.
- PROBABBITAL CALCIUM, N N R.** 1951. See: Iprat Calcium, Tab. (Squibb).
- PROBARBITAL SODIUM, N F. X, N N B** 1955; Tab., N F. X, 5-Ethyl-5 Isopropylbarbiturate and See: Iprat Sodium, Elixir & Tab. (Squibb)
- W/Aspirin.** See: Iprat-Aspirin, Tab. (Squibb).
- PROBEC.** See: Stuart Probec, Tab. (Stuart)
- *PROBENECID, U S P., N N D.** 1962; Tabs. U S P. p-(Di-n-propylthioamyl) benzoic acid
* Generic name recognized by N F., U S P. or A.M.A.
- See: Benemid, Tab. (Merck Sharp & Dohme)
W/Colchicine. See: Colbenemid, Tab. (Merck Sharp & Dohme)
- W/Pol. penicillin G.** See: Remanden, Prep. (Merck Sharp & Dohme)
- *PROBENZAMIDE.** o-propoxybenzamide. (Warner-Lambert)
- PROBILAGOL.** (Purdue Frederick) d-Glucitol-(d-sorbitol) 4.5 Gm., homotropine methylbromide 1 mg./5 ml. Bot. 12, 6 oz.
Use: Cholagogue cholecystokinetic agent for normalized digestion.
- PROBITAL.** (Searle) Propantheline bromide 7.5 mg., phenobarbital 15 mg./Tab. Bot. 100s
Use: Orally, antispasmodic.
- PBC-BLEM.** (Hoyt) Hexachlorophane 0.1%, isopropyl alcohol 70%, cetyl alcohol 0.5%. Plastic Bot. squeeze 4 oz.
Use: Treatment of acne.
- PRORUTYLIN.** (Eorar) Procaine butyrate. Cap. 300 mg., Bot. 50s. Elixir 10%, Bot. 180 cc.
Use: Antinausea.
- PROCA-H.** (National Vitamin Prods.) Controlled pH of 3.5-4 containing procaine hydrochloride 2%, sodium chloride 0.2%, sodium bisulfite 0.19%, chlorobutanol 0.5%.
* **PROCAINAMIDE HYDROCHLORIDE, U S P., N N D.** 1962; Cap., U S P., Inj., U S P. p-Amino-N-(2 diethylaminoethyl) benzamide HCl. See: Pronetyl Hydrochloride, Cap. Vial (Squibb)
Use: Cardiac depressant in arrhythmias.
- PROCAINE ADENYLATE.** (Physiological Chemicals) 50 mg. procaine adenylylate equiv. to 30 mg. Adenosine-3-monophosphate, 20 mg. (2%) procaine base/cc. Amp. 1 cc.
Use: Arthritis & associated diseases.
- PROCAINE ASCORBATE.** See: Scorbacaine, Amp. (Physiological Chemicals).
- PROCAINE BASE.** (No Mfr. currently lists) **W/Benzocaine,** butyl aminobenzoate & benzyl alcohol in sesame oil. See: Analgacain, Vial (Mitra)
W/Benzyl alcohol & propyl-p-aminobenzoate. See: Bactocaine, Amp., Oint. & Supp. (Kirk)
W/Butesin & benzyl alcohol in oil. See: Zylcaine, Amp. (Abbott)
W/Butyl-p-aminobenzoate, benzyl alcohol, in sweet almond oil. See: Anucaine, Amp. (Calvin Chem.)
W/Isocaine & benzyl alcohol. See: Proctocaine, Amp. (Columbus)
W/Procaine HCl, butyl p-aminobenzoate & polyethylene glycol-300. See: Efoacaine, Vial (Fougeron)

W/Propylamin benzoate See: Propalcaline, Amp. (Chicago Pharm).

PROCAINE BORATE. See: Barocaine A & D (Halperin)

W/Theophylline, mersalyl See: Mercuprocyll, Inj (Barrows)

PROCAINE BUTYRATE. (Rorer) p Aminobenzoic diethylaminoethanol butyrate

Oint: (10%) w/Ephedrine 0.5%, cresol 0.5%. Tube 1 oz, Jar 4 oz, 1 lb

Ophth Oint: (2%) w/Epinephrine 1:25,000 Tube 1/8 oz

Sol: (2%) Amp 2 cc, Box 12s, 25s, 100s; Vial 30 cc, 100 cc; (2%) w/Epinephrine 1:50,000, Amp 2 cc, Box 12s, 25s, 100s

Amp 5 cc, Box 7s, 12s, 25s, 100s; Vial 30 cc, 100 cc

Supp: (10%) w/epinephrine 0.5%, cresol 0.5% Box 6s, 12s

Topical anesthetic: Procaine butyrate 50%, benzyl alcohol 41%, ephedrine 1%, menthol 5%, phenol 0.25% Sol 15 Gm, 25 Gm

Use: Anesthetic.

See: Prabutylin, Cap & Elixir (Rorer)

W/Ephedrine, tannic acid, phenol & ammonium alum See: Butritannic, Oint (Rorer)

W/Mag sulfate, aromatics & glycerin See: Auralol, Improved, Liq (Rorer)

PROCAINE C. See: Scorbacaine

***PROCAINE HYDROCHLORIDE, U.S.P.,** N.N.R. 1949 A.O.R. Inj, U.S.P.; Sterile, U.S.P.; Tab, N.P. X Diethylaminoethyl p-aminobenzoate HCl (Allocaïne, Bernocaine, Chlorocaine, Ethocaine, Irocaïne, Keracaine, Syncaïne)

See: Geritacaine H 3 (Kirk)

Kirkcaine, Vial (Kirk)

Lactocaine, Vial (United Lab)

Noucaïne, Sol & Tab (Taylor Lab)

Neocaine, Amp (Fougere)

Navocain (Winthrop Labs)

Scurocaine (Wallau)

W/Aminaxytropine tropate HCl, pentobarbital Sol, phenobarbital, epaxymoxoxy tropine tropate HBr, Vit B 6 & aluminum hydroxide gel dried See: Prometic, Tab (Harrower)

W/Ascorbic acid See: Ascarbocaine, Tab (Testagar)

W/Ascorbic acid, methapyriene HCl See: Vio Caine, Cap (Rawell)

W/Ascorbic acid, and Vit B 6 See: Cevicaine, Cap (Calumbus)

W/Charlone gonadotropin, Vit B 1, glutamic acid See: Glukor, Vial (Research Supplies)

*Generic name recognized by NF, U.S.P. or A.M.A.

W/Ephedrine HCl, U.S.P. See: Ephedrine-Novocain, Amp (Winthrop Labs)

W/N lactic acid See: Lactocaine, Vial (United Labs)

W/Pentobarbital, phenobarbital See: Barbicaine, Liq (Cutter)

W/Phenylephrine HCl See: Iso-Synocaine, Vial (Warren Teed)

W/Testosterone, thiamine HCl, glutamic acid See: Glutest, Vial (Research Supplies)

***PROCAINE PENICILLIN G, U.S.P., N.N.O.** 1962 See: Penicillin G, Procaine (Various Mfr)

PROCARDIUM. (Grant) Digitoxin 0.05 mg, strophanthin 0.05 mg, strychnine sulfate 0.10 mg/Tab Bot 100s, 500s

Use: Orally cardiac diseases

PROCARMIN. (Lantern) Hydrolyzate of primary yeast, nonfat milk solids & defatted whole liver w/added iron. Pow 12 oz, 5 lb, 25 lb; Tab 0.5 Gm, Bot 250s, 500s, 1000s

Use: Hypoproteinemia

PROCEBRIN. (Billy) Thiamin chloride 1 mg, riboflavin 0.5 mg, pyridoxine HCl 0.5 mg, nicotinamide 8 mg, ascorbic acid 60 mg, Vit A 3000 u, 0 800 u/0.3 cc, Bot 15 cc

Use: Vitamin deficiencies

PROCELATE SOLUTION. (Crookes Barnes) Edathamil Disod in two strengths 0.37 and 1.85% w/sod bicarb 0.1% Vials 100 cc

Use: Chelating agent for calcium cpds accidentally placed in eye

***PROCHLORPERAZINE MALEATE, U.S.P.,** N.N.D. 1962; Tabs U.S.P. 2 Chloral D [3-(1-methyl-4-piperazinyl)propyl] phenothiazine maleate U.S.P. Also ethanedithiolate, U.S.P., N.N.O. 1962 for inj U.S.P. See: Compazine preps (Smith, Kline & French)

W/Dextroamphetamine sulfate See: Exkatrol Span (Smith, Kline & French)

PROCHOLON (Squibb) Not listed Dehydrochloric acid Tab 0.25 Gm, Bot 100s, 500s

Use: Nonobstructive biliary stasis

PROCOLLAGENASE. (Children's Hospital Research Foundation Wash, D.C.) Enzyme

PROCO-SOL PENICILLIN DENTAL CONES (Proco-Sol Chem Co) Pat penicillin G 15M u/Cone Pkg 12s

Use: Prophylaxis of dry socket

PROCTALME (Meyer) Benzocaine, chlorbutanol neomycin, thymol iodide, ointment w/rectal applicator Tube 1 oz

Use: Hemorrhoids

PROCTOCAINE. (Columbus) Benzyl alcohol 4.16%, procaine base 1.25%, Isocaine 4.16% Amp 5 cc, Box 25s, 100s

Use: Local anesthetic

- PROCTOCAINE OINTMENT.** (Rowell) Rnzo-caine 5%, 10% & 25% with burbot liver oil oint. base. Tube w/pipn 1 oz.
Use: Anesthetic rectal use.
- PROCTODON.** (Rowell) Dipreron HCl 1%. In petrolatum and fish liver oil base. Oint. Tube 1 oz.
Use: Treatment of rectal irritations.
- PROCTOFORM.** (Testogor) Isobutyl amino-benzoate, bismuth subiodide, zinc oxide, bismuth subcarb. & boracic acid. Snp. Box 12s.
Use: Rectal supp. for local infections.
- PROCTOPLEX.** (Rowell) Tonic acid 5%, dipreron HCl 1%, zinc oxide 5%. Oint. Tube 1 oz.
Use: Rectal conditions.
- PROCTUROL.** (Hart) Ephedrine HCl 0.75%, chlorbutanol 3%, benzocaine 1%, w/phenol, menthol, lavender oil, camphor & Sod. chloride, in water-soluble jelly base. Tube 1 oz.
Use: Anesthetic, antipruritic, vasoconstrictor & antiseptic.
- ***PROCYCLIDINE HCl, N.N.D. 1962.** (1-Cyclohexyl-1-phenyl-3-pyrrolidinopropionol). See: Kemadrin, Tab. (Burroughs Wellcome)
- PROCYCLO.** (Metropolitan Lab.) alpha-Estradiol 0.267 mg., estrone 0.267 mg., progesterone 10 mg., in sesame oil, aq. susp./cc. Vial 10 cc.
Use: (I.M./cc) Estrogen-progesterone therapy.
- PROCYSTAMINE.** (Lakeside) 0.1 Gm. foreign protein, 0.01 mg cysteine & 30 mg thiamine/2 cc. Vial. 30 cc. Box 3s, 6s, 25s
Use: I.M. non-specific protein therapy.
- PRODECADRON, RESPIHALER.** (Merck Sharp & Dahme) Dexamethasone 21 phosphate 18 mg, isoproterenol sulfate 18 mg. In total weight of 12.6 gms. Aerosol—170 sprays.
Use: Branchial asthma.
- PRO-DERMA.** (Westwood) Nat listed. Silicone 52.5%. Oint. 1 oz.
Use: Skin protective.
- ***PRODILIDINE HCl.** (Mead Johnson & Co.) (1,2-dimethyl-3-phenyl-3-pyrrolidyl propionate HCl. See: Cogesic.
Use: Analgesic.
- PRODOX, N.N.D. 1960.** (Upjohn) Hydroxyprogesterone Acetate 25 and 50 mg./Tab. Bot. 25s.
Use: Progesterone activity.
- PRODOXAN.** (Smith-Dorsey) Ethisterone. Tab. 10 mg, 25s, 100s.
Use: Carpus luteum therapy.
- * Generic name recognized by N.F., U.S.P. or A.M.A.
- PRO-DUOSTERONE.** (Roussel) Anhydrohydroxyprogesterone 50 mg, activated by ethinyl estradiol 0.03 mg./Tab. Bot. 50 mg./0.03 mg./24s.
Use: Narmane for oral pregnancy test.
- PRODYN.** (Elder) Vit. A 5000 u., D 800 u., B-1 1.5 mg., B-2 2 mg., niacinamide 5 mg., cal. pantothenol 1 mg., Vit. C 30 mg., E 1 u./Cap. Bot. 100s, 500s, 1000s
Use: Multivitamin therapy
- PRO-ESTROBEV E.** (Key Corp.) Progesterone 12.5 mg., estradiol benzoate 2.5 mg. In sesame oil/cc. Vial 10 cc.
- PROFAMINA.** See: Amphetamine (Various Mfr.)
- PROFENAMINE HCl.** Ethoprapazine HCl, N.N.D. 1959. See: Parsidol, Tab (Warner-Chilcott)
- PROFENIL.** (Smith, Miller & Patch) Sestronyl bis-gamma-Phenyl-propylethylamine. Citrate salt for oral; HCl salt for parenteral use. Amp. (45 mg) 1 cc., Pkg. 5s, 25s. Tab. (120 mg) Bot. 100s, 1000s, w/phenobarbital 1/4 gr. Bot. 100s, 1000s
Use: Antispasmodic.
- See: Antispasmin, Tab. (Roemer)
Fentropine (Smith, Miller & Patch)
- PROFENIL CITRATE W/Homatropine methylbromide.** See: Fentropine, Tab. (C. D. Smith)
- W/Homatropine methylbromide, butobarbital sod** See: Fentropine w/Butobarbital, Tab (Smith, Miller & Patch)
- PROFIRON.** (Central) Oral hematinic with protein hydrolysate and vitamin B complex. Bot. pt, gal.
- PROFERRIN.** (Merck Sharp & Dahme) Saccharated iron oxide. Sterile soln. cont. 20 mg. of available iron/cc. Vial 20 cc.
Use: Hypochromic microcytic anemia.
- PROFETAMINE PHOSPHATE.** (Clark & Clark) dl-Amphetamine phosphate. Amp. (5 mg./cc.) 1 cc., 25s, 100s. Chewing gum 10 mg, Box 21s. Pow. Bot. 1 oz. Tab. 5 mg, 10 mg., Bot. 100s.
Use: Central nervous system stimulant.
W/Phenobarbital & aspirin. See: Pen-Phetamine, Tab. (Clark & Clark).
- PROFLAVINE, N.F. IX, N.N.R. 1950.** (Mol linekrodil) Pow., Bot. 5 Gm., 25 Gm.
Use: Topically; antiseptic.
- PROGELAN.** (Lannett) Progesterone 25 mg./cc. In oil. Vials 10 cc.
- PROGELAN AQUEOUS.** (Lannett) Progesterone 25 mg. aq. susp./cc. Vials 10 cc.
- PROGEST.** (Hart) Progesterone 10 or 25 mg./cc. Vial 10 cc.
- PROGESTAB.** (Massengill) Ethisterone Tab. 10 mg, Bot. 100s.

Use: Orally; threatened abortion, premenstrual tension & dysmenorrhea

PROGESTAO. (New Mexico Pharm) Progesterone 50 mg, procaine HCl 1%, chlorobutanol 0.5%, polyoxyethylene sorbitan 0.24% Vial 10 cc

Use: Progesterone therapy

PROGESTEROID. (Sherman) Progesterone In aq susp Vial (25 mg / cc, 50 mg / cc) 5 cc

Use: Corpus luteum therapy

PROGESTEROL. (Smith Miller & Patch) Progesterone sol in peanut oil or susp in Sod chloride sol
In oil, Vial (10 & 25 mg / cc) 15 cc; (50 mg / cc) 10 cc

Susp: Vial (25 mg & 50 mg / cc) 10 cc

Use: I.M.; corpus luteum replacement therapy

* **PROGESTERONE, U.S.P., NND 1962;** Inj., Tabs U.S.P.; Susp NF Corpus luteum hormone (Flavalutan Luteagon, Luteosan, Lutren) See: Progestin

Aq susp, U.S.P. (Various mfr)

See: Calprosterone, Vaginal Tab (Ayerst)

Gesterol, Vial (Testagar)

Lipa-Lutin (Parke, Davis)

Lucarlem, Amp (Merck Sharp & Dohme)

Macrogestin, Vial (Kirk)

Myosal Progesterone (Ulmer)

Pragelan Aq, Vial (Lanett)

Progesterol, Vial (Sherman)

Progesterol, Vial (C. D. Smith)

Progesteronaq Vial (Central)

Syngesterone, Vial (Pfizer)

In oil, U.S.P. (Various mfr)

See: Carlutone, Sol in oil (Gold Leaf)

Carpomane, Vial (Preme)

Gesterol, Vial (Testagar)

Glanestin, Amp & Vial (Inglalaid)

Lipa-Lutin, Amp (Parke, Davis)

Lutocycin, Amp & Vial (Ciba)

Lutramane, Amp (Endo)

Lutrone, Amp (Bunting)

Nalutran, Vial (Winthrop Labs)

Pragelan, Vials (Lanett)

Progestin, Vial (Harveys; Kirk; Organon)

Progesterone, Vial (Carnrick)

Progestranol, Vial (Central)

Pralutan Vial (Schering)

Syngesterone, Vial (Pfizer)

In propylene glycol See: Migesterone Vial (Durst)

Tab, sublingual, U.S.P. (Various mfr)

See: Linguarbs (Ayerst)

Lutocycin, Linguets (Ciba)

Membrettes (Wyeth)

Pralutan, Tab (Schering)

Syngestrals, Tab (Pfizer)

W/Estradiol (beta)

* Generic name recognized by N.F., U.S.P. or A.M.A.

See Aquaprazol, Vial (Estra)

Corpa Gen, Vial (Barry)

Di Pro, Vial (Organon)

Estratone, Vial (Chicago Pharm)

Estratone, W/Progesterone, Vial (Lakeside)

Myosal Tergesten (Ulmer)

Neo Femotrone Vial (Testagar)

Progestradial, Vial (Ciba)

Promestran, Vial (Schering)

Prostradiol, Vial (Kirk)

W/Estradiol benzoate

See Fasigyn in Oil, Vial (Pfizer)

Pro Estrabev E, Vial (Key Corp)

Progestradial, Amp (Ciba)

W/Beta estradiol & estrone

See: Alimestrane, Vial (Meyer)

Procydo, Vial (Metro)

W/Estradiol, methyltestosterone See: In

hormonal Tab (Key Corp)

W/Estradiol & testosterone

See: Tergesten, Vial (Ulmer)

Tregonaq Vial (Central)

Trihorm, Vial (Ascher)

Trihormogen, Vial (Mallard)

W/Estradiol & testosterone propionate

See: Espronate, Vial (Chicago Pharm)

Tregestradiol, Vial (Kirk)

W/Estradiol, testosterone, procaine HCl

procaine base See: Horma Triad, Vial (Bell

Pharmaceutical)

W/Estrogonic substance See Estrogenic Sub

stance W/Progesterone (Various Mfr)

W/Estrogonic substance

See: Harvestrin and progesterone, Vial (C. F

Harvey)

Lipa-Mistrane, Vial (E. S. Miller)

W/Estrogonic substance & testosterone See

Tremilone, Vial (Miller)

W/Estrogyne See: Progestagyne (Estra)

W/Estrone

See: Crinogen, Vial (Merrell)

Korep, Vial (Karmel Co)

Myosal Prestagen (Ulmer)

W/Estrone & alpha estradiol

See: Alimestrane, Vial (Meyer)

Procydo, Vial (Metropolitan Labs)

W/Estrone, estradiol, benzyl alcohol in pe

nut oil See: Ovagest, Vial (C. D. Smith)

W/Estrone & Testosterone

See: Kar Tep (Karmel)

P E T, Vial (Maury Labs)

Patrane, Vial (Kremers Urban)

Pro-Mal O Fem, Aq Susp (Testagar)

Termone, Vial (Metropolitan Labs)

Tristerone, Tube (Wyeth).

W/Estrone, testosterone, procaine HCl See

Kar Tep, Aq Susp (Karmel Co)

W/Testosterone propionate estradial bet

- zoate in oil. See: Trihormonal, Vial (Key Corp)
- PROGESTIN**, N.N.D. 1962. Progesterone. (Various mfr.)
- PROGESTOGENE**. (Estra) Estrogyne 50,000 I.U., progesterone 10 mg./cc. Vial. 3 cc., 10 cc.
- Use: Primary & secondary amenorrheas.
- PROGESTONE**. (Carrick) Progesterone. Hypoglossal 10 mg., 25 mg. Bot. 50s, 100s, 1000s. Vial. In oil (10 mg./cc., 25 mg./cc.) 10 cc. Box 1s, 5s, 10s, 25s.
- PROGESTORAL**, (Organon) Ethisterone. Tab. 5 mg., Box 20s, 50s, 100s, 250s; 10 mg., Box 10s, 20s, 40s, 100s, 250s; 25 mg., Bot. 20s, 100s. Also buccal 10 mg., Bot. 30s, 100s.
- Use: Corpus luteum replacement therapy.
- W/Phenobarbital & thiamine HCl. See: Surdan, Tab. (Organon).
- PROGESTRADIOL** (Ciba) Progesterone 12.5 mg., estradiol benzoate 2.5 mg., in oil/cc., Amp. 1 cc., 2s, 6s.
- Use: I.M. (1 cc. daily for 2 days); estrogen & androgen therapy.
- PROGESTROID**, (Brown Pharmaceuticals) Anhydroxyprogesterone 10 mg., ethinyl estradiol 0.01 mg., thyroid extract 10 mg., Resperdin 100 mg., Vit. C 100 mg., glutamic acid 50 mg., thiamine hydrochloride 10 mg./Tab. Bot. 100s, 500s.
- Use: Treatment of habitual abortion.
- PROGESTROLAN**, (Lanett) Progesterone 25 mg., estrogenic sub. natural 25,000 u. in oil/cc. Vial 10 cc.
- PROGESTRONAO**, (Central) Progesterone. Vial (25 & 50 mg./cc.) 10 cc.
- Use: Corpus luteum therapy.
- PROGESTRONOL**, (Central) Progesterone, in sesame oil 25 mg./cc. Vial 10 ml.
- PROGONASYL**, (Progonasyl Co) O-Iodebenzoic acid 1%, oleic acid 35%, mineral oil 32%, veg. oil 22%, triethanolamine 11%. Bot. 55 & 110 Gm.
- Use: Hemorrhoids.
- PROGUANIL HYDROCHLORIDE**.
- See: Chloroguanide Hydrochloride. (Lilly)
- Paludrine, Tab (Ayerst)
- PROGYNON**, N.N.D. 1962. (Schering) Estradiol, dihydroxyestrie. Buccal Tab. 0.125 mg., 0.25 mg., Bot. 30s, 100s. Pellets 25 mg., Vial 1s, 3s. Tab. 0.1 mg., 0.2 mg., 0.5 mg., Bot. 30s, 60s, 250s. Vial, Sosp. (0.25 mg., 1.0 mg./cc.) 10 cc.
- PROGYNON-B**, N.N.D. 1961. (Schering) Estradiol benzoate. Amp. 1000 u., 2000 u., 6000 u., 10,000 u./cc., 1 cc., 6s, 50s, 100s; 1 cc., 1s, 6s.
- PROHALONE**, (Ayerst Lab) Halaprogesterone.
- Use: Gestational agent.
- PRONEPIN-V**, (Merrell) Liver ext. 10 u., thiamine HCl 50 mg./cc. Vial 10 cc., 30 cc.
- Use: Pernicious anemia.
- PROHYDRION**, (Upjohn) Cal. mandelate 5 gr., methenamine $1\frac{1}{2}$ gr., ammonium acid phosphate $2\frac{1}{2}$ gr./Tab. Bot. 100s, 1000s.
- Use: Orally; urinary antiseptic.
- PROKETAZINE MALEATE**, (Wyeth) Carphenazine maleate.
- Use: Management of schizophrenia.
- PROKLAR**, (Lloyd, Dabney & Westerfield) Sulfacetamide 250 mg., methenamine mandelate 250 mg./cap. Bot. 100s.
- Use: Urinary infections.
- PROLACTIN**. See: luteotrophic, Vial (Squibb).
- PROLADYL**, (Lilly) Pyrrobutamine, 1-pyrrolydyl-3-phenyl-4-(p-chlorophenyl)-2-butene phosphate 0.015 Gm./Tab.
- Use: Antihistaminic.
- PROLAGEN**, (Drug Products) Estrogenic substances in sesame oil. Vial, eq. susp. (2000 I.U., 10,000 I.U., 20,000 I.U./cc.) 10 cc. Vial, in oil (10,000 I.U., 20,000 I.U./cc.) 30 cc.
- Use: Estrogen therapy.
- PROLAIRE**, (Stuart) Butobarbital 32 mg., D-amphetamine sulfate 5 mg., acetylsalicylic acid 160 mg., salicylamide 160 mg., acetaminophen 160 mg./Tab. Bot. 100s.
- Use: Analgesic.
- PROLAIRE-R**, (Stuart) Butobarbital 32 mg., d-amphetamine sulfate 5 mg., aspirin 160 mg., salicylamide 160 mg., N-acetyl-p-aminophenol 160 mg., hyoscyamine sulf. 0.1384 mg., atropine sulf. 0.0258 mg., hyoscyne HBr 0.0086 mg./Tab. Bot. 100s.
- Use: Analgesic & sedative.
- PROLASE 300**. Proteolytic enzyme from corica papaya. See: Papain.
- W/Mylose, cellulase, calcium carbonate, magnesium glycinate. See: Zylase, Tabs. (Bryant)
- W/Mylose-100, dehydracholic acid, hyoscyamine HBr, hyoscyne HBr, atropine sulfate See: Almezyme, Cap. (Meyer)
- PROLAX**, N.N.D. 1962. (Cala) Mephenesin. Tab. 0.5 Gm. Bot. 100s, 500s, 1000s.
- Use: (1 Gm. I.D.) Muscle relaxant.
- W/Atomatropine methyl bromide. (Cala) Mephenesin 0.5 Gm., hamatropine methyl bromide 1.25 mg./Tab. Bot. 100s.
- Use: 2 I.D.; muscle relaxant for certain spastic & neurologic disorders.
- PROLIFEROL "T" INJECTION**, (Ulmer) Same as "B" plus phenol 6.25%. Vials 60 cc.
- PROLIFEROL** (Ulmer) B, Est. al catechu, wild brier, rhatany & huckleberry, phlobaphenes, thymol & alcohol. Vial 60 cc. T,

contains thuja & phenol plus above constituents Vial 60 cc

Use: Varicose veins

PROLIFEROL "R" INJECTION. (Ulmer) Phlobatannins 0.5%, thymol 0.5%, alcohol 80% Vials 60 cc

Use: Hernia Injection

PROLIXIN. (Squibb) Fluphenazine 4 (3 [2 trifluoromethyl 10 phenothiazyl] propyl) 1 piperazine ethanol dihydrochloride Tabs 1 mg, 2.5 mg, 5 mg Bot 50s Elixir (2.5 mg/ml) Bot w/dropper 60 ml, 1 pt Vial (2.5 mg/cc) 10 cc

Use: Treatment of schizophrenia, chronic brain syndrome, psychoneurosis, etc

PROLOID. (Warner Chilcott) Thyroid globulin ext Tab ¼ gr, ½ gr, 1 gr, 1½ gr, 3 gr, 5 gr, Bot 100s, 500s, 1000s

Use: Myxedema, cretinism & hypothyroid states

W/Ethinyl estradiol, methyltestosterone See: Plestron Tab (Warner Chilcott)

PROLOXIN, NND 1962 (Lanett) Me phenesin 0.25 Gm/tab Bot 250s, 1000s Also 0.5 Gm Bot 250s, 500s, 1000s

Use: Skeletal muscle relaxant

PROLUTON, NND 1962 (Schering) Progesterone Succol Tab 10 mg, Bot 30s, 100s Vial, in oil (1 mg, 2 mg, 5 mg, 10 mg) 1 cc, 6s, 30s; (10 mg, 25 mg, 50 mg/1 cc) 10 cc, 6s

Use: I.M. & Orally (10 mg, daily) corpus luteum therapy

PROMACETIN. (Parke, Davis) Acetosulfone. Tab. (0.5 Gm) Bot 500s

Use: Treatment of leprosy (Hansen's disease)

PROMAGETIN. (Parke, Davis) Acetosulfone Tab 0.5 Gm, Bot 500s

Use: (1-3 daily) Hansen's disease

PRO-MAL-O-FEM. (Testogon) Aq. Susp.—Testosterone 25 mg, estrone 6 mg, progesterone 25 mg/cc Vial 10 cc

In Oil—Testosterone propionate 25 mg, progesterone 25 mg, estradiol benzoate 1 mg/cc Vial 10 cc

Use: I.M., estrogen and androgen therapy

PROMANULARE. (Lakeside) Mang butyrate & foreign protein Vial 60 cc, 6s, 12s, 25s

Use: Nonspecific protein therapy

***PROMAZINE HCl, HND 1962** 10 (gamma Dimethylamino-n Propyl)-phenothiazine HCl See: Sparine, Tab, Vial (Wyeth) W/Meprobamate See: Proxine, Cap (Wyeth)

PRO-MEPERDAN. (Barr) Meperidine HCl 50 mg/cc Amps 2 cc Vial 30 cc

Use: Analgesic

***PROMETHAZINE HYDROCHLORIDE, U S P** NND 1962; Tab, Syr, Inj, U S P (D)

* Generic name recognized by H.F., U.S.P. or A.M.A.

methylamino-2 methyl-2 ethyl) N dibenzoparathiazine See: Phenergan, Prep (Wyeth)

W/Aspirin, phenacetin, mepentermine sulf See: Synalgos, Cap (Ives Cameron)

W/Meperidine HCl See: Mepergan, Amp (Wyeth)

W/Penicillin V, Salicylamide, phenacetic, mephestermine sulf. See: Pen Vee Cidin, Cap (Wyeth)

***PROMETNESTROL.** 3,4 Di (4 hydroxy-3 methylphenyl)hexone. See: Meprone, Tab (Reed & Carnrick)

***PROMETNESTROL DIPROPIONATE, NND 1962** See: Meprano Dipropionate Tab (Reed & Carnrick)

Use: Estrogen deficiency

W/Phenobarbital See: Meprone Phenobarbital, Tab (Reed & Carnrick)

PROMETIC W/AMPHETAMINE. (Warner Chilcott) Phenobarbital 10.8 mg, Sod pentobarbital 10.8 mg, amphetamine 1 mg, epoxyaminoxytropine tropate HBr 0.25 mg, aminoxytropine tropate HBr 0.25 mg, procaine HCl 150 mg, Vit B 6 HCl 10 mg, aluminum hydroxide 250 mg/Tab Bot 50s

Use: Anti emetic

PROMETRON. (Schering) Estradiol benzoate 2.5 mg, progesterone 12.5 mg/cc Amp 1 cc, 2s Vial 10 cc, 1s, 6s

Use: I.M., estrogen & corpus luteum therapy

PROMIN SODIUM, NND 1962 (Parke, Davis) Glucosulfone sodium Amp (2 Gm) 5 cc, 6s, 25s 100s, (5 Gm), 12½ cc, 25s, 100s Jelly Tube 1¼ oz

Use: Leprosy & tuberculosis

PROMINAL. See: Mebaral, Tab (Winthrop Stearns)

PROMINATE. (Schleffelin) Protein hydrolysate Bot 8 oz

Use: Dietary supplement

PROMIZOLE, NND 1960 (Parke, Davis) 2 Amia 5 sulfamizythiazole (Thiazolsulfone) 0.5 Gm/Tab Bot 500s

Use: Orally; adjunct streptomycin therapy

PROMOMS. (Kenwood) Calcium 300 mg, Vit C 100 mg, A 4000 u, D 400 u, B 12 2 mg, B 2 2 mg, B 12 1 mcg, iron 45 mg, sodium diacetyl succinate 50 mg, Vit B 6 2 mg, potassium 15 mg, zinc 5 mg, manganese 1 mg/Tab Bot 30s

Use: Vitamin and mineral supplement

***PROMOXALANE, NND 1962** 2,2 Diisopropyl 1,3 dioxolane 4 methanol See: Di methylene, Cap (National Drug)

PROMOXIHE HYDROCHLORIDE. gamma Morpholinopropyl-4 N butoxyphenyl ether NCl See: Tranothane HCl (Abbott).

- PROMULOEN.** (Robinson Wagner) A versatile nonionic emulsifying ointment base. Jars 1 lb.
- PROMULSE.** (E. S. Miller) Progesterone in emulsifiable oil 100 mg./cc. Vial 10 cc.
- PROMYL.** (Crookes-Barnes) Pipenzolate methylbromide 5 mg., aluminum mag. hydroxycarbonate 300 mg., aluminum proclinate 200 mg./Tab. Rat. 100s.
Uses: Anticholinergic and antacid.
- PRONAC.** (Fougera) Stabilized sulfated potash 0.60 Gm. w/equimolecular amount of zinc sulfate 0.40 Gm., & inert binders Pow. pack, 12s.
Uses: Topically for acne.
- PRONAPEN, A.D.E.** (Pfizer) Procaine penicillin G 300,000 u., Sod. penicillin G 100,000 u./cc. Vial 1 cc., 5 cc., 10 cc.
Uses: I.M.; penicillin therapy.
- PRONAPEN PLUS.** (Pfizer) Benzethacil 200,000 u., Pat. penicillin O 100,000 u., procaine penicillin G 300,000 u./dose. Vial, single dose.
Uses: I.M.; penicillin therapy
- PRO-NASYL.** (Progenasyl Co) o-Iodobenzoic acid 1%, triethanolamine 11%, dibucaine base 1%, in a neutral hydrophilic base.
Uses: Treatment of sinusitis.
- PRONE.** (Barre) Ethisterone. Tab. 5 mg., 10 mg., 25 mg.
Uses: Corpus luteum replacement therapy.
- PRONEMIA.** (Ledarie) Vit. B-12 30 mcg., ferrous sulfate 400 mg., Vit. C 150 mg., stomach pow. 300 mg., folic acid 4 mg., intrinsic factor conc. 10 mg./Cap. Rat. 30s.
Uses: (1 daily) Hematinic.
- PRONESTYL HYDROCHLORIDE, N.N.D.** 1962. (Squibb) Procaine amide HCl. Cap. D 25 Gm., Bot. 100s., 1000s. Sol. 100 mg./cc., Vial 10 cc.
Uses: Treatment of ventricular arrhythmias & related cardiac conditions.
- PRO-OTASYL OTIC DROPS.** (Prognasyl) Triethanolamine 5 1/2%, o-Iodobenzoic acid 1/2%, neutral hydrophilic base/Bot. 15 cc., 60 cc.
Uses: Otolologic.
- PROPADRINE HYDROCHLORIDE, N.N.E.** 1937. (Merck Sharp & Dohme) Phenylpropanolamine HCl. Cap. 1/4 gr., 1/2 gr., Bot. 25s., 100s., 1000s. Pow., Bot. 1/2 oz. Effair, D.12 Gm./fl. oz., Rat. 1 pt., 1 gal.
Uses: Nasal decongestant & bronchodilator.
W/Hydrocortone, neomycin sulfate. See: Hydrocortone, Nasal Suspension. (Merck Sharp & Dohme)
- W/Phenylephrine See: Neo-Vadrin, Sol. (First Texas)
- * Generic name recognized by N.P., U.S.P. or A.M.A.
- W/Theophylline. See: Prophyllin Tablet. (Merck Sharp & Dohme)
- W/Tyrosine. See: Tyrosine Liquid. (Merck Sharp & Dohme)
- W/Phenacetin & Aspirin. See: Riona, Cap. (Merck Sharp & Dohme)
- W/Thyroid. See: Altepose, Tab. (Merck Sharp & Dohme)
- PROPAGESIN.** See: Propyl p-Aminobenzoate. (Various Mfr.)
- W/Benzyl alcohol, urea & p-chloro-m-xylene. See: Cenathesin, Oint. (Central).
Dunaguent, Oint. (Ascher).
- PROPALCAIN.** (Chicago Pharm.) Procaine base 0.075 Gm., propylamine-benzoate 0.30 Gm., benzyl alcohol 0.25 Gm., vegetable oil as vehicle/5 cc. Amp. 5 cc., 25s., 100s.
Uses: Local anesthesia.
- PROPAMINE.** (Finley) Phenylpropanolamine HCl 12.5 mg., chlorpheniramine maleate 2.0 mg./Cap. or 5 cc. syrup Cap. 50s., 250s. Syrup pts.
Uses: Decongestant and antihistaminic.
- PROPANEDIOL DIACETATE, 1,2.** See: VoSol (Wampole).
- * **PROPANTHINE BROMIDE, U.S.P., N.N.D.** 1962; Tab. U.S.P. See: Pro-Santhine, Prep. (Searle)
- W/Phenobarbital. See: Probital, Tab. (Searle)
- PROPORACAIN HCl, N.N.D.** 1962. 2-Diethylaminoethyl 3-amino-4-propoxybenzoate HCl. See: Ophthaine.
- PROPASA, N.N.D.** 1959 (Merck Sharp & Dohme) p-Amino-salicylic acid. Tab. eff. 1 Gm., Vial 20s.
Uses: Tuberculosis.
- * **PROPYL NITRATE.** 2-ethyl-2-hydroxy-methyl-1,3-propanediol trinitrate. See: Etrynit (Bison)
- * **PROPZOLAMIDE.** 2-propionylamino-1,3,4-thiadiazole-5-sulfonamide See: lanaze (Eli Lilly)
- * **PROPENZOLATE HYDROCHLORIDE.** (National Drug Company) 1-methyl-3-piperidyl o-phenylcyclo-hexaneglycolate hydrochloride. See: Delinal (National Drug)
Uses: Parasympatholytic agent.
- PROPETHYLENEOXIDES.** See: Magnocyl, Cap. (Elder)
- PROPHEPYRIDAMINE.** See: Pheniramine, N.N.D. 1939.
- PROPHEPYRIDAMINE MALEATE, N.F., N.N.D.** 1962; Tab. N.F., Ophth. Sol., N.F. See: Pheniramine Maleate.
- PROPHEPYRIDAMINE TANNATE W/Phenylephrine tannate, pyrilamine tannate.** See: Rynatan Tablets, Tab. (Irwin, Nelster)
- PROPHYLLIN.** (Ryston) Sod. propanate &

- chlorophyll "a" paw Baz 12 packets Bot 4 oz, 16 oz Oint 1½ oz, 4 oz Tube
Use: Antiseptic.
- PROPICILLIN.** (Registered in Great Britain) 6 (α Phenoxybutyramido)penicillanic acid (1 Phenoxypropyl)penicillin Brocillin is the potassium salt; Ultrapen is the potassium salt
- PROPIODAL.** See: Entadan, Amp (Winthrop Labs)
- ***PROPIOMAZINE.** (Wyeth) 1 [1D (2 di methylaminopropyl) phenothiazine 2 yl] 1 propanone See: Larga
- PROPIONATE SALTS.**
See: Copper
Potassium
Sodium
Zinc
- PROPIONATE-CAPRYLATE MIXTURES.** NNR 1955 See: Sopronal, Prep, NNR (Wyeth)
- PROPIONATE COMPOUND.** NNR 1955 See: Propion Gel, NNR (Wyeth)
- PROPION GEL.** NNR 1955 (Wyeth) Cal propionate 100%, Sod propionate 100%, propionic acid 1% in water miscible base Tube 95 Gm with or without applicator
Use: Treatment of mycotic vulvovaginitis (moniliasis)
- PROPIONIC ACID.**
See: Propionate Caprylate Mixtures, NNR. Prozin (Calumbus)
- PROPION OPTHALMIC.** (Wyeth) Sod propionate 5% in neutral sol Bot 5 fl. dr w/ dropper
Use: Control of superficial infections of conjunctiva, cornea & eyelid
- PROPIONYL ESTER OF ERYTHROMYCIN** W/Meth dia mer sulfas See: Ilosone Sulfate Tab (Lilly)
- PROPIONYL ERYTHROMYCIN LAURYL SULFATE** See: Erythramycin Propionate Lauryl Sulfate
- PROPISAMINE.** See: Amphetamine (Various Mfr)
- PRO-PLESTRIN** (Harrower) Natural estrogenic substances equiv to estrone 10000 IU, progesterone 10 mg, in sesame oil/cc Vial 5 cc
Use: Estrogen & corpus luteum therapy
- PROPOQUIN.** (Parke, Davis) See: Amopyraquin HCl
- ***PROPOXAMIDE.** α propoxybenzamide (Warner Lambert) Name is withdrawn
- ***PROPOXYCAINE HYDROCHLORIDE, N.F., ADR 1955; Inj NF 2** Diethylaminoethyl 4 amino-2 propoxybenzoate HCl
See: Blackal, Sol (Breon)
Ravocaine Hydrochloride (Cooke Waite)
- * Generic name recognized by NF USP or A.M.A.
- ***PROPOXYCHLORINOL.** See: Toloxychlorinol (Warner Chilcott)
- PROPOXYPHENE HCl, U.S.P., NND 1961; Caps USP 1,2-diphenyl 3 methyl-4-di methylamine 2 propoxybutane** See: Darvon, Pulv (Lilly)
W/Acetylsalicylic acid, acetophenetidin See: Darvon Compound—65, Caps (Lilly)
W/Phenalglycolal, acetylsalicylic acid See: Vondor, Caps (Lilly)
- PROPYL p-AMINORENZOATE.** [Various Mfr] Propaestin
Use: Local anesthetic
W/Anilpyrine, sulfathiazole Sod See: Sulf Aurette, liq (Raymer)
W/Benzyl alcohol, urea, p chlor m xylenol See: Dunaguent Oint (Ascher)
W/Rismuth subnitrate, peru balsam, zinc oxide, salicylic acid, calamel, resorcin phenol, calamine & code oil See: Raymer Derm, Oint (Raymer)
W/Gramicidin, neomycin sulfate See: Orbiotic Gum Tab (White)
W/Neomycin sulfate & bacitracin See: Du Biotic, Troche (White)
W/Procaine base & benzyl alcohol See: Propolcain, Amp (Chicago Pharm). Tactocaine, Oint & Supp (Kirk). Trithesia, Amp (Breon)
W/Zinc oxide, calamine, benzoin See: Raymer Derm, Oint (Raymer)
- ***PROPYLHEREDRINE, U.S.P., NND 1962;** 1 Cyclohexyl 2 methylaminopropane; Inhalant USP See: Benzedrex, Inhalant (Smith, Kline & French)
- ***PROPYLIODONE, U.S.P.** Sterile Susp USP N propyl 3,5 diiodo 4 pyridone N acetate Sterile oil susp (peanut oil) Sterile water susp
- n-PROPYLNICOTINATE** W/Methyl salicylate menthol, thymol, nicotinylsalicylic acid See: Nisolin, Oint (National Drug)
- Is-PROPYLNORADRENALINE.** See: Isoproterenol
- ***PROPYLPARABEN, U.S.P., ADR** [Various Mfr] Propyl p hydroxybenzoate Propyl Chemosept (Chema Pura)
- ***PROPYLTIOURACIL, U.S.P., NND 1962;** Tab USP 2 Mercapto-4 hydroxy 6 n propylpyrimidine 25 mg, 50 mg Bot 100, 1000s
- PRORENATA ¼ & ½ TAB** (Palmedical) Phenacetin 162 mg, d amphetamine phosphate 5 mg, salicylamide 227 mg, phenobarbital 24 mg, codeine phosphate ¼ & ½ gr/tab Bot 100s
Use: Analgesic

- PROSCOMIDE**, (Smith, Miller & Patch) Methscopolamine bromide 7.5 mg./cap. Bot. 30s, 100s.
Use: Anticholinergic in treatment of peptic ulcer.
- PROSECA**, (Westwood) Polyoxyethylene lauryl ether 3%, sulfur 4%, hexachlorophene 1%, in a special base. Bot. 4½ oz.
Use: Acne.
- PROSOL**, (Torch) Mercury oleate 0.25%, coal tar crude 0.6%, salicylic acid 1.5%, phenol 0.25%, p-nitrophenol 0.25%. Emulsion, Bot. 2 oz.
Use: Antiseptic, antipruritic & fungicide.
- PROSTALL**, (Metabolic Products Corp.) Glycine, alanine glutamic acid mixture 6% gr./Cap. Bot. 100s, 250s.
Use: Benign prostatic hypertrophy.
- PROSTAPHLIN**, (Bristol) Sodium oxacillin 250 mg. and 500 mg./Tab. Bot. 48s.
Use: Staphylococcal infections.
- PROSTIDMIN BROMIDE**, N.N.O., 1955. (Roche) Neostigmine bromide, Ophth. sol. 5%, Bot. 7½ cc. Tab. 15 mg., Bot. 20s, 100s, 250s, 1000s.
Use: Parasympathetic stimulant.
- W/N-Acetyl-p-amino-phenol, hyescine hydrobromide**. See: Dorth (Oenob)
- PROSTIGMIN METHYLSULFATE**, N.N.O. 1955 (Soche). Neostigmine methylsulfate, U.S.P., Amp. (1:2000) 1 cc., Rox 12s, 50s, (1:4000) 1 cc., 12s, 100s. Vial (1:1000, 1:2000) 10 cc.
Use: Parasympathetic stimulant.
- W/Hyoscyamine HBr & benzyl alcohol**. See: Kelogen, Vial (Paul-Money)
- PROSTRADIOL**, (Kirk) beta-Estradiol 50,000 u., progesterone 10 mg./cc. Vial 10 cc.
Use: I.M., estrogen & corpus luteum therapy.
- PROSTRUGEN**, (Kremers-Urban) Estrogens 2.5 Gm., progesterone 12.5 mg./cc. In oil, Vial 10 cc. Also in aq. susp., Vial 10 cc.
Use: I.M., estrogen w/corpus luteum therapy.
- PROTALRA**, N.N.O. 1962 (Pitman-Moore) Protoveratrine A. (0.2 mg.) Tab. Bot. 100s.
Use: Hypertension.
- PROTALRA-R**, (Pitman-Moore) Protoveratrine A. 0.2 mg., reserpine 0.05 mg./Tab. Bot. 100s.
Use: Hypertension.
- PROTAMIDE**, (Sherman) Sterile, colloidal sol. of processed & denatured proteolytic enzymes. Amp. 1.3 cc. Pkg. 10s.
Use: (I.M., 1.3 cc. twice weekly) Tabes dorsalis, herpes zoster, nerve root pain, chicken pox & neuritis.
- PROTAMINAL**, (Brewer) Autolysed primary yeast 0.5 Gm., desiccated liver 0.1 Gm.,
* Generic name recognized by N.F., U.S.P. or:
- dicalcium phosphate 60 mg., ferrous gluconate 3 mg., fat-free milk solids/Tab. Bot. 500s.
- *PROTAMINE SULFATE INJ.**, U.S.P., N.N.D. 1962.
Celta—Pow.
Lilly—Amp. 1%, 5 cc., 6s, 25s.
Upjohn—Vial (10 mg./cc) 5 cc., 1s, 5s, 25s.
Use: I.V., heparin overdosage.
- PROTAMINE ZINC INSULIN**, U.S.P., N.N.S. See: Insulin, Protamine Zinc. (Various Mfr.)
- PROTAPE**, (Triflor) Simult oxyiodide, benzoin, aloes, staraz, Tolu balsam, alcohol 65% Bot. 1 oz.
Use: Increases adhesion and decreases sensitivity to adhesive tapes.
- PROTARGIN MILD**. See: Silver Protein Mild (Various Mfr.)
- PROTARDOL**, N.N.R. 1949. (Winthrop Labs.) Strong silver protein. Pow., Bot. 1 oz.
Use: Topically silver antiseptic.
- PROTEASE**, W/Amylase, lipase. See: Trulose, Tabs. (Smith, Mills & Patch)
- PROTEF**, (Upjohn) Rectal supp. Hydrocortisone 15 mg., neomycin sulf. 15 mg. supp. Rox. 12s.
- PROTEGEL**, (Wyeth) Alumina & kolin gel. Tube, 1 oz., 4 oz.
Use: Pruritus ani (moist type).
- *PROTEIN HYDROLYSATES INJ.**, U.S.P., N.N.D. 1962.
See: Amoc, Pow. (Blue Line).
Amigen, Sol. (Mead, Johnson)
Aminogen, Amp. (Christina).
Aminonut, Pow. (National)
Aminosol, Sol. (Abbott)
Hyprologen, Sol. (Don Baxter)
Lacotain, Vial (Christina)
Porenamine, Sol. (Winthrop Labs.)
Travamin, Sol. (Baxter Lab.)
W/Vitamins & minerals. See: Vitamin, Prep.
- *PROTEIN HYDROLYSATES ORAL**, N.N.D. 1962.
See: Amateen, Pow. (Dumas-Wilson)
Amino Acids (Stuart)
Amino-concamin (Marrell)
Caminoids, Pow. (Arlington)
Co-Amino (Physiological)
Ketoad, Pow. (Merck Sharp & Dohme)
Lefenamalc (Mead, Johnson)
- PROTEIN, NONSPECIFIC THERAPY**.
See: Alan, Amp. (Duke)
Clesilac, Vial. (Pitman-Moore)
Estratol, Amp (Estron)
Kirkavin, Vial (Kirk)
Kirkolac (Kirk)
Lacotain (Christina)
Mangolac, Vial (Endo)

Mucosal, Vial (Kremers Urban)
 Niadalin, Vial (Lincoln)
 Omnadin
 Polyprate, Amp (Kirk)
 Promanulake, Amp (Lakeside)
 W/Vit B Complex See: Betathalin, Amp (Lincoln)
PROTENUM. (Mead-Johnson) Milk solids, Cal caseinate, minerals & fat Pkg 1 lb, 4 lb.

Use: Dietary supplement
PROTEINAMYL. (Strassenburgh) Ceylon papain 16 gr w/malt ext./fl. oz Sol 1 pt, 1 gal
 Use: Digestant
PROTEOLAC. (Searle) Hydrolyzed proteins of defatted milk Amp, Serum Type, S cc, 6s, 25s, 100s
 Use: I M; nonspecific protein therapy

PROTEOLYTIC ENZYMES
 See: Annonax Tabs (Rorer)
 Papase (Warner Chilcott)
 W/Amylolytic enzyme d sorbitol See: Kuzyme, Cap (Kremers Urban)
 W/Cellulolytic enzyme See Gustazyme Tabs (Geriatric Pharmaceutical)
 W/Amylolytic, cellulolytic See: Trienzyme Tabs (Testagar)
 W/Amylolytic enzyme, cellulolytic enzyme, aluminum hydroxide gel See: Molozyme, Tabs (Halco)

W/Amylolytic enzyme, hamotropine methyl-bramide d sorbitol (Papain) See: Converzime Liq (Ascher)
 W/Amylolytic enzyme lipolytic enzyme See: Farmulase, Tabs (Kremers Urban)
 W/Neomycin palmitate hydrocortamate HCl; water miscible base See Chymar (Armaur Pharmaceutical)
 W/Piperidolate hydrochloride, cellulolytic enzyme, amylolytic enzyme, pancreatin 3X, tauracholic acid See: Dactilose, Tabs (Lakeside)
 W/Pyridoxine HCl, pynilamine maleate, amylolytic enzyme See: Nemisis Tabs (Beutlich)

PROTERNOL. (Key) 3D mg Isoproterenol HCl/Tab Bot 30s, 500s
 Use: Heart block

PROTEUS OX 19. (Lederle) Febrile antigen Vial 5 cc.
 Use: Diagnostic aid

PROTINIALATE (New Mexico Pharm) Sodium thiosalicylate In ampul water 50 mg, benzyl alcohol 2%/cc Vial 30 cc
 Use: Relief of arthritic and rheumatic pain

PROTINIOLIN. (New Mexico Pharm) Foreign protein 50 mg, methionine 5 mg, thiamine HCl 15 mg, niacinamide 50 mg, riboflavin

* Generic name recognized by N.F., U.S.P. or A.M.A.

1 mg, calcium pantothenate 2 mg, pyridoxine HCl 1 mg, benzyl alcohol 2%/cc Vial 30 cc
 Use: Anti arthritic

***PROTHIPENOYL HCl.** [(4 dimethyl amino propyl pyrida (3 2 B) benzothiazine)] HCl monohydrate See Timovan (Ayerst)

PROTHRICIN. (Merck Sharp & Dahme) Tyrothricin 0.02%, propadrine HCl 15%. Bot 1 oz w/dropper
 Use: Decongestant w/antibiotic

PROTHROMASIN, NNO 1962 (G F Harvey) Warfarin sod, 3 (α-acetyltolbenzyl) 4 hydroxycoumarin 5 mg, 10 mg or 25 mg/Tab Bot 25s, 100s Also called Warcoumin
 Use: Anticoagulant

PROTHROMBIN. (No Pharmaceutical firms listed)

W/Thrombekinase & antithrombin See Neo-Hemoplastin, Vial (Parke, Davis)

PROTHYRIN. See: Thyractin (Winthrop)

PROTIDE. (Paul Maney) Sod iodide 12 gr iodinated protein 6 gr/2 cc Vial 30 cc
 Use: Liquefaction of mucous secretions

PROTINAL. (National Drug) Carbohydrate 30%, protein 61.25% Pow, Bot, ½, 1, 5 & 25 lb
 Use: Protein therapy.

PROTINIC. (Dumas-Wilson) Liver intrinsic factor complex 250 mg, ferrous sulf 325 mg, Vit C 60 mg, Vit B 12 15 mcg, folic acid 1 mg, w/trace elements/Cap Bot 100s, 1000s
 Use: Hematinic

PROTORAN. (UVC) Protein 12 Gm, carbohydrate 100 Cal, Iron 20 mg/oz, Pow, Bot 1 lb.
 Use: Orally; nutritional therapy

PROTOGON. (Maare Kirk) Chazianic ganadotropin 250 IU, pyridoxine HCl 25 mg, protein hydrolysate 60 mCg, w/pracaine HCl 1%/cc Vial 10 cc
 Use: Hypogonadism

***PROTOKYLOL HCl, NND 1962 a** [(α methyl 3,4-methylenedioxyphenethylamino) methyl]protocatechuyl alcohol HCl See: Caytine (Lakeside Labs)

PROTOLYSATE, NNR. 1954 (Mead Johnson) Dried enzymic digest of casein consisting of amino acids & small peptides. Pkg 1 lb, 5 lb
 Use: Orally; dietary supplement

PROTONUCLEIN. (Reed & Carnrick) Thyroid ¼ gr, spleen ½ gr, pancreas ½ gr, seprerant ¼ gr, thymus ¼ gr, brain ¼ gr, lymphatic ¼ gr./Tab Bot 100s, 500s 1000s Pow 1 oz, R oz
 Use: Orally; thyroid therapy

- PROTOPAM.** (Campbell Pharm., Inc.) See: Prolidaxime iodide and chloride.
Use: Anticholinergic.
- PROTOPLEX.** (Walker) Casein, lactalbumin, yeast, liver desic. & carbohydrate. Pow., Can 1 lb. Tab. 200s.
Use: Protein therapy.
- PROTOS.** (Heun) Protein, lactalbumin, minerals & fat 0.4%, Sod. 0.05%. Can 1 lb., 25 lb.
Use: Protein therapy.
- PROTOSIL.** (Bryant) Tyrothricin 0.01%, hexachlorophene 0.5%, benzocaine 2%, W/Vit. A, D, silicone & zinc oxide. Tube 1 oz.
Use: Protective agent.
- *PROTOVERATRINE A, N.N.D. 1962.** See: Protalba, Tab. (Pitman-Moore)
W/Hydroflumethiazide, reserpine. See: Saltesin, Tab. (Bristol)
- W/Beserpine.** See: Protalba-B, Tab. (Pitman-Moore)
- *PROTOVERATRINES A & R MALEATE, N.N.D. 1962.** See: Pavell Maleate, Tab. (Lilly).
Veralba, Vial, N.N.R. (Pitman-Moore).
W/Rauwolfia serpentina. See: Baupiate, Tab. (Vale)
W/Rauwolfia serpentina whole root & phenoxanzamine HCl. See: Mio-Presin, Cap. (Smith, Kline & French)
W/Reserpine.
See: Veralba-R, Tab. (Pitman-Moore)
Veropene, Tab. (Wampole)
- PROVATOL.** (Wyeth) Vit. A 6820 u., D 1360 u./cc. Sol. w/dropper 50 cc.
Use: Orally; Vit. A & D therapy.
- PROVELL MALEATE, N.N.D. 1962.** (Lilly)
Prolaveratrines A & B. Tab. 0.2 & 0.5 mg., Bot. 100s, 1000s.
Use: Vasodilator.
- PROVERA, N.N.D. 1962.** (Upjohn) Medroxyprogesterone acetate 25 mg., or 10 mg./Tab. Bot. 25s, 100s.
Use: Threatened and habitual abortion.
- PROVERA-TESTPAC.** (Upjohn) Medroxyprogesterone acetate 10 mg./tab Box 5s.
Use: Determination of pregnancy.
- PROVIDONE.** See: Vinsil, Sol. (Abbott)
- PROVIGEN.** (Mead Johnson) Non-fat milk solids, calcium caseinate, sucrose, vitamins and minerals. Pow.
Use: Dietary supplement.
- PRO-VITAMIN A, N.N.B. 1954.** See: Coralene.
- PROVITE "B."** (Ives-Cameron) Vit. B-1 25 mg., B-2 12.5 mg., niacinamide 130 mg., Cal. pantothenate 5 mg., Vit. B-6 1.5 mg., choline dihydrogen citrate 100 mg., Inositol 50 mg., liver fraction 40 mg., yeast dried 20 mg./Cap. Bot. 100s, 1000s.
Use: Orally (1 daily); Vit. B complex therapy.
- W/Vit. C.** (Ives-Cameron) Vit. B-1 25 mg., B-2 12.5 mg., niacinamide 100 mg., Cal. pantothenate 5 mg., Vit. B-6 1.5 mg., B-12 2 mcg., choline dihydrogen citrate 100 mg., ascorbic acid 100 mg., inositol 50 mg., liver fraction 60 mg./Cap. Bot. 100s.
Use: Orally (1 daily); Vit. B complex w/Vit. C.
- PROXYDRIN.** (Professional Pharm. Co.) di-Desoxyephedrine HCl 1/8%, benzalkonium chloride and chlorobutanol. Bot. 1 oz., 1 pt
Use: Nasal spray
- PROZINE.** (Wyeth) Meprobamate 200 mg., promazine HCl 25 mg./Cap. Bot. 50s.
Use: Control of emotional disturbances.
- PROZINE HALF-STRENGTH CAPS.** (Wyeth) Meprobamate 100 mg., promazine HCl 12.5 mg./Cap. Bot. 50s.
Use: Control of emotional disturbances.
- PRURILAX.** (Ball Pharmacal) Diacetyldihydroxyphenylisatin 5 Gm., prune conc. 5 gr./Tab. Bot. 100s, 1000s.
Use: (1-2 h.s.) Laxative.
- PRULET.** (Mission Pharmacal) Diacetyldihydroxyphenylisatin. Tab. 5 mg., Bot. 60s.
Use: (1 morning & night) Laxative.
- PRULOSE COMPLEX.** (Harrowes) Tab., Prune conc. 2 gr., methylcellulose 6 gr., diacetyldihydroxyphenylisatin 1/65 gr./Tab. Bot. 100s, 500s. liq., Prune conc. (70% solids) 52 gr., carbosymethylcellulose Sod. 11.5 gr., diacetyldihydroxyphenylisatin 1/32 gr./15 cc. Bot. 12 oz.
Use: Functional constipation.
- PRUNE POWDER CONCENTRATED DEHYDRATED.** (Chemo Pure) Pkg. 35 lb., 103 lb. See also: Diacetyldihydroxyphenylisatin.
W/Carboxymethylcellulose Sod. & diacetyldihydroxyphenylisatin. See: Prulose Complex. liq. & Tab. (Harrower).
W/Cascara fl. ext. aromatic & psyllium husk pow. See: Cassylum, Pow. (Upjohn).
W/Diacetyldihydroxyphenylisatin. See: Prulilax, Tab. (Ball Pharmacal)
W/Diacetyldihydroxyphenylisatin, thiamine mononitrate. See: Prunisatin, Tab. (Sutliff & Case)
W/Methylcellulose. See: Prulose, Tab. (Harrower).
W/Methylcellulose & diacetyldihydroxyphenylisatin. See: Hydrelase Fortified, Syr. (Upjohn)
W/Mg(OH)₂, dextran. See: Danthro-lex, Tab. (Mallard)

* Generic name recognized by N.F., U.S.P. or A.M.A.

PRUNE PREPS.

- See: Casyllium, Liq (Upjohn)
 Prulose Complex (Harrower)
- PRUNICODEINE.** (Lilly) Cadeine phosphate
 $\frac{7}{8}$ gr, terpin hydrate, prunus virginiana,
 pinus strabus & sanguinaria Bot 1 pt
 1 gal
 Use Orally (1 2 dr q 3 h); Cough.
- PRUNISATIN.** (Sulfin & Case) Diacetyl
 hydroxyphenylisatin 5 mg, prune concen-
 trate 30 mg, thiamine mononitrate 1 mg /
 Tab Bot 100s, 1000s
 Use Constipation
- PRURINE OINTMENT.** (Rand) 1% Oamine
 N (n dodecyl) N methyl N N dimethyl ethyl-
 ene diamine Tube 1 oz
 Use Prolonged soothing anesthetic
- PRURISAT.** (Dale) Phenol, menthol, chlor-
 butolol, hydracortisone 0.3% Tube/5 Gm,
 $\frac{1}{2}$ oz
 Use Dermatologic
- PRYDONNAL SPANSULE.** (Smith, Kline &
 French) Belladonna alkaloids 0.4 mg, phe-
 nobarbitol 1 gr/Spansule Bot 30s
 Use Antispasmodic w/sedative
- PRYDON SPANSULE.** (Smith, Kline & French)
 Belladonna alkaloids Spansules (0.4 & 0.8
 mg) Bot 30s
 Use: Anticholinergic & antispasmodic
- PRYMEDON.** (Prime) Hyoscyamine sulfate
 0.1037 mg atropine sulfate 0.0194 mg,
 hyoscine HBr 0.0065 mg, phenobarbitol
 16.2 mg / Tab Bot 100s
 Use Antispasmodic antisecretory sedative
- PRYMEDON SPASCAPS** (Prime) Hyoscy-
 amine sulfate 0.311 mg, atropine sulfate
 0.0582 mg, hyoscine HBr 0.0195 mg
 phenobarbitol 48.6 mg / timed disintegra-
 tion cap Bot 100s
 Use Antispasmodic antisecretory sedative
- P33.** (Cutter) Sulfamerazine 0.167 Gm sal-
 fadiazine 0.167 Gm, sulfamethazine 0.167
 Gm / Tab W/Pat penicillin G 100M u or
 250M u Tab 100 (100M u penicillin) Vial
 20s; 250 (250M u penicillin) Vial 20s
 Use: Orally; penicillin w/triple sulfonamide
 therapy
- P-3 DENTAL CONES.** (Navacol) Penicillin
 Sad 15,000 u. w/Sad sulfacetamide 0.46
 Gm & sulfacetamide 0.04 Gm / Cone Bot
 50s, 100s
 Use: Dry socket & dental infections
- * **PSEUDOEPHEDRINE HYDROCHLORIDE**
 (Burroughs Wellcome) Isaeophedrine HCl
 See: Sudafed, Tab (Burroughs Wellcome)
 Use: Orally ($\frac{1}{4}$ - $\frac{1}{2}$ gr b i d); sympathom-
 etic agent for asthmatic attacks
- W/Chlarcyclizine HCl See: Fedrazil, Tab
 (Burroughs Wellcome)
- W/Chlarcyclizine HCl, acclaphenetidin, as-
 pirin, caffeine See: Emprazil, Tab (Bur-
 roughs Wellcome)
- W/Chlarpheniramine maleate See: Isodor,
 Tab (Haskell)
- W/Tripalidine HCl See: Actifed, Tab & Syr
 (Burroughs Wellcome)
- W/Tripalidine HCl, cadeine phosphate
 glyceryl guaiacolate See: Actifed C, Syr
 (Burroughs Wellcome)
- PSICOFURANINE.** (Upjohn) 6-amino 9 D
 psicofuranasyipurine
 Use: Antibacterial against experimental
 Staph aureus, etc
- PSL DIHYDROSTREPTOMYCIN SULFATE**
 (Pacific States Lab) Dihydrostreptomycin sul-
 fate 10 Gm / Vial 1 dose
 Use: Gram-negative organism infections
- PSL PENICILLIN G POTASSIUM TABS.,
 BUFFERED.** (Pacific States Lab) Pot. peni-
 cillin G 50M u / Tab Bot 50s, 100s
 Use: Penicillin therapy
- PSORIACIDE.** (Daak) Salicylic acid 10%,
 anhralin 0.25% or 0.50%. Stick 30 Gm
 Use Treatment of psoriasis
- PSYCHOTHERAPEUTIC DRUGS.** See: Alar-
 zic Agents
- PSYLLIATE SODIUM, NND 1958** See
 Synasol, Amp (Searle)
- PSYLLIUM NUSK POWDER W/Cascara fl**
 ext aromatic & prune paw See: Casyllium,
 Pow (Upjohn)
- PSYLLIUM HYDROCOLLOID.** See: Effersyl-
 lum (Stuart)
- PSYLLIUM HYDROPHILIC MUCILLOID.**
 NND X 1962 See Plantaga Mucilloid
 Products NND (Various Mfr)
- W/Oi (acetylhydroxyphenyl) Isalin See: Ef-
 fersyl, Gran (Stuart)
- PSYLLIUM SEED, N F W/Mineral Oil** See:
 Parosyllium (Abbott)
- PSYLLIUM SEED GEL.** See: Betajel Powder
 (Floralose Lab)
- PTEROIC ACID.** The compound formed by
 the linkage of carbon 6 of 2 amine 4
 hydroxypteridine by means of a methylene
 group with the nitrogen of paminobenzoic
 acid
- PTEROYLGLUTAMIC ACID, U.S.P. NND**
 1962 See: Folic Acid, Prep
- PTEROYLMONOGLUTAMIC ACID.** Pteroyl
 glutamic acid See: Folic acid
- PTEROYLTRIGLUTAMATE SOD** See: Terap-
 terin, Amp (Lederle)
- PTHALOX.** (Sulfin & Case) Cal ricinateate
 65 mg, phenalpthalein 32.5 mg, sad
 glycocholate 32.5 mg, sad tauracholate 16

* Generic name recognized by NP USP or AMA

- mg., cocora socrado ext. 8 mg./Tab. Bot. 1000s.
 Use: Laxative.
- PURICIOE.** (Warren-Teed) Benzyl benzoate 12½%, DDT 1%, benzocaine 2%. Liq. Bot. 4 oz., 1 pt.
 Use: Topically; scabicide & pediculicide.
- PULMO-PAK.** (Abbott) Disposable heart-lung apparatus.
 Use: Isolate perfusion and dry-field surgery.
- PULMOSEROL.** (El Paso Pharm.) Liquid. Bot. 1 pt.
- PURALIV.** (Philadelphio Ampoule) Liver Injection containing B-12 activity of 20 mcg./cc. Viol 10 cc.
 Use: Anemias
- PURALIVER.** (Myers-Carter) Liver ext. (B-12 20 mcg./cc.) Viol 10 cc.
- PURIFIED CORTICOTROPIN-GEL.** N.N.R. 1955. (Wilson Lab.) Purified corticotropin. Viol 5 cc., avail. in 20, 40, 80 or 100 U.S.P. u./cc.
 Use: Adrenal cortex stimulant.
- PURIFIED OXGALL.** See: Rile Ox Extract (Various Mfr.).
- PURIFIED PROTEIN DERIVATIVE OF TUBERCULIN.** U.S.P., N.N.R. 1955. (Parke, Davis) See: Tuberculin, Old, U.S.P.
 Use: Mantoux TR test.
- PURINETHOL.** N.N.D. 1962. (Rurrroughs Wellcome) 6-Mercaptopurine. Tab. (50 mg) Bot. 25s, 250s.
 Use: (2-4 daily) Acute leukemia & chronic myelogenous leukemia
- PURITAL.** (A. J. Parker) Theobromine Cal. equiv. to 3.25 gr. of theobromine/tab. Bot. 100s, 1000s. Pow. Bot. 1½ oz.
 Use: Cardiovascular therapy.
- PURITAL R.S.** (A. J. Parker) Theobromine Cal. equiv. to 3.25 gr. of theobromine, & terpine 0.25 mg./tab. Bot. 100s.
 Use: Cardiovascular therapy w/hypotensive & tranquilizing agent.
- PURITON.** (Eaton) See: Nihydrozone
- PURIVAX VACCINE.** (Merck Sharp & Dohme) Types 1, 2, and 3 poliomyelitis virus grown on monkey kidney and treated with formaldehyde. 2 cc. (4 dose).
 Use: Poliomyelitis vaccine.
- PYROBABB.** (A. J. Parker) Purital w/phenobarbital 1/6 gr. Bot. 100s, 1000s.
 Use: Cardiovascular therapy w/mild sedation.
- W/Rutin** 30 mg. Bot. 100s, 1000s. (A. J. Parker)
 Use: Cardiovascular therapy capillary fragility protection.
- PURDIGIN.** N.N.R. 1950. (Wyeth) (Digitoxin, Crystalline)
 Amp. (0.2 mg./cc.) 1 cc., 6s, 25s.
- Tab. (0.05 mg.) Bot. 100s. (0.1, 0.15, 0.2 mg) Bot. 100s, 500s, 0.2 mg. Viol 30s
 Use: Cardiac decompensation & congestive heart failure.
- PUROGEL.** (Bruce) Emulsion zirconium carb. 2%, ethyl aminobenzoate 3%, Isopropyl alcohol 8%, liq. pot. 20%, pH 6-7. Bot. plastic 4 oz.
 Use: Burns and wounds
- PURPASAL-C.** (Meyers-Rosser) Sod. salicylate 5 gr., ascorbic acid 10 mg, p-aminobenzoic acid 5 gr./Tab. Bot. 100s, 500s.
 Use: (2-4 q. 3 gr.) Analgesic & antipyretic.
- PURPLE FOXGLOVE.** See: Digitalis (Various Mfr.).
- PYORONE.** (Brean) Dipyrone. Amp. (500 mg./2 cc.) 2 cc., Box 25s,
 Use: S.C., I.M. (1 or 2 cc.); analgesic & antirheumatic.
- PYGMAI.** (Astier) Tartaric acid, boric acid, Rurow's sol., glycerin, starch, talc & ben-tonite. Oint., Tube 1½ oz.
 Use: Antiseptic skin preparation.
- PYLORIC SUBSTANCE W/Liver,** vitamins & Iron. See: Stoffin, Tab. (Orgonon).
- PYLOSITAMIDE.** (Vitomix) Phenylazodiaminopyridine HCl 50 mg, sulfacetamide 0.5 Gm., atropine sulfate 0.0195 mg, hyoscine HBr 0.0065 mg, hyoscyamine sulfate 0.1040 mg./tab.
 Use: Urinary analgesic and anti-infective
- PYMADEX.** (Testagor) Pyramisamine maleate 25 mg, dextroamphetamine HCl 2 mg/cc. or Tab. Viol 10 cc., Bot. 100s, 1000s.
- PYMADEX TIMED, CAPSULES.** (Testagor) Pyramine maleate 25 mg., methoprylene HCl 25 mg, prophenpyridamine maleate 10 mg, d-amphetamine HCl 6 mg/cap. Bot. 100s.
 Use: Timed disintegrating antihistaminic for daytime use only.
- PYMAFED.** (Lloyd) Pyrilamine maleate. Tab., Bot. 100s, 500s, 1000s.
- PYMA TIMED CAPSULE.** (Testagor) Pyrilamine maleate 25 mg., methoprylene HCl 25 mg., prophenpyridamine maleate 10 mg/cap. Bot. 100s.
 Use: Timed disintegrating antihistaminic.
- PYOKTANIN.** (Merck Sharp & Dohme) Gan-han violet, N.N.R. Pow., Bot. 1 oz.
 Use: Topically, instillation & orally; antiseptic & anthelmintic.
- PYRABROM.** (Pyrilamine R bromotheophyllinate).
 See: Bramth, Tab. (Brayten).
 Glybrom, Tab. (Patch).
 Mensalin, Tab. (Walker).
 W/Acetopheneridin. See: M-Mrus-4, Tab. (Whittier).

PYRACORT. (Lemmon) Hydrocortisone alcohol 0.02%, tyrothricin 0.02%, phenylephrine HCl 0.25%, thienylpyramine HCl 0.2%
 Bot ¼ oz plastic spray, 1 pt
 Use: Treatment of acute and chronic allergic rhinitis

PYRADONE. See: Aminopyrine (Various Mfr)

PYRANIST. (Brewer) Pyrilamine maleate 5/12 gr., aspirin 3½ gr., phenacetin 2½ gr., caffeine ½ gr., Vit C 2½ gr./Cap Bot 100s, 500s, 1000s

Use: (1 q 3 h) Antihistamine w/analgesic

PYRALDINE (VanPelt & Brown) Dextromethorphan HBr 15 mg., pyrilamine maleate 12½ mg., ammonium chloride & citric acid/5 cc Bot 1 pt, 1 gal

Use Orally (1 dr q 3 h) cough

PYRALDINE No 2 (VanPelt & Brown) Dextromethorphan HBr 90 mg pyrilamine maleate 75 mg phenylephrine HCl 30 mg., ammonium chloride 6 gr., w/citric acid/oz Syr Bot 1 pt 1 gal

Use: Antitussive agent

PYRALDINE PEDIATRIC. (VanPelt & Brown) d methorphan HBr, 45 mg; pyrilamine maleate, 75 mg; phenylephrine HCl, 30 mg; ammonium Cl, 6 gr; citric acid, 5 gr/oz Syr Bot Pt

Use: Relief of cough and congestion

PYRALGIN PEDIATRIC DROPS. (Savage) Methampyrone sodium (Dipyrone) 250 mg/cc Bot 1 oz

PYRA-MALEATE, NND 1962 (VanPelt & Brown) Pyrilamine maleate

PYRAMAL MALEATE, NND 1962 (Columbus) Pyrilamine maleate Tab 50 mg, Bot 100s, 1000s

PYRAMAL UHICELLES. (Hiss Pharm) Pyrilamine maleate 75 mg/cap Bot 100s

Use: Delayed action, antihistaminic

PYRAMID (Barre) Pentobarbital sod 1/6 gr., butobarbital sod 1/6 gr., phenobarbital ½ gr/5 cc Bot 16 oz, gal

Use: In anxiety states

PYRAMIDE. (Columbus) Salicylamide 5 gr pyrilamine maleate 25 mg, amphetamine sulfate 2.5 mg/Tab Bot 100s, 500s, 1000s

Use: (1 tid) Premenstrual tension & primary dysmenorrhea

PYRAMIDON, NNR 1949 (Winthrop Labs) [Aminopyrine, NF]

Elixir: (160 mg per 4 ml) Bot 16 fl oz

Pow: Bot 1 oz, 1 lb

Tab: (5 gr) Bot 100s

Use: Analgesic

PYRAMINAL (Winthrop Labs) Aminopyrine

* Generic name recognized by N.F., U.S.P., or A.M.A.

2½ gr. luminal ¼ gr/5 cc Elixir, Bot 12 fl. oz., 1 gal

Use: Sedative & antispasmodic

PYRAMINE CREAM. (Flint, Eaton) Pyrilamine HCl 2%, colamine 15%, aluminum hydroxide 5%, phenol 1%. Tube 1½ oz

Use: Local anesthetic & antihistaminic.

PYRAMINYL. See: Pyrilamine Maleate

PYRANEURIN INJ. (Nutrition Control Prod) Vitamin B 1 100 mg & B 6 100 mg/cc

Vials: 1D cc

Use: B Vitamin Inj

PYRANILAMINE MALEATE. See: Pyrilamine Maleate, NND 1959

PYRAHISAMINE BROMOTNEOPYLLI-NATE. See Pyrabrom.

PYRANISAMINE MALEATE, NNR 1950

See: Pyrilamine Maleate, NND 1959

PYRAPNED. (Lemmon) Phenylephrine HCl ½%, thienylpyramine HCl 0.2%, & tyrothricin 0.02% Bot plastic spray ¼ oz

Use: Nasal congestion

PYRA-SED. (Columbus) Dihydrocodeinone bitartrate 18 mg, pyrilamine 15 mg, desoxyephedrine HCl 1 mg, w/ammonium chloride citric acid, chloralform & alcohol/5 cc Bot 3 oz, 1 pt, 1 gal

Use Orally (1 dr q 3 h); cough

PYR-ASPRO. (VanPelt & Brown) Salicyl amide 3½ gr., acetophenetidin 2½ gr., caffeine ½ gr., pyranisamine 125 mg., belladonna alkaloids 0.07 mg/Tab Bot 100s, 500s

Use: Orally (1 2 q 3 h), analgesic & antihistaminic

* **PYRATIAZINE HYDROCHLORIDE, N.F., NND 1962** Tab NF [N(2 Fyralidine ethyl)-phenothiazine See: Pyralazate, PreP (Upjohn)]

W/Orthoxine HCl, A.P.C. Also with Codeine

See: Pyrazate, Cap (Upjohn)

PYRATNYN. (Flint, Eaton) Methapyrilene HCl Tab 25 mg, 50 mg, Bot. 100s, 500s, 1000s

Use: (50 mg 2 3 h) Antihistaminic

W/Aluminum hydroxide & phenol See: Pyramine Cream (Flint, Eaton).

W/A.P.C. See: Capathyn, Cap (Flint, Eaton)

* **PYRAZINAMIDE, NND 1962** (Merck Sharp & Dohme) 2 Carbamyl pyrazine Aldinamide Tab (500 mg) Bot 500s

Use: Tuberculosis

PYRAZINOIC ACID AMIDE. See: Pyrazinamide, Tab (Merck Sharp & Dohme)

PYRAZOLINE. See: Antipyrine (Various Mfr)

PYRETHRUM, N.F., NNR 1955 [Upsher Smith] Oint 27%, Jar 100 Gm.

Use: Externally; scabicide

PYBETS. (Bernhoff) Chlorprophenpyridamine maleate 4 mg., N-acetyl-p-aminophenol 300 mg., phenylephrine HCl 10 mg., phenylpropanolamine HCl 10 mg., ephedrine sulfate 10 mg./tab.

Use: Relief of common cold.

PYBENZAMINE, N.N.D. 1962. (Ciba) (Tripeleannamine Citrate as HCl Salt) N-Benzyl-N',N'-dimethyl-N-2-pyrrolidylethylammonium HCl.

Amp.s (25 mg./cc.) 1 cc., 5s. Vials 7D cc.

Creams (2%) Tube 50 Gm., Jar 1 lb.

Elizir Citrate salt (7.5 mg./cc.) Bot. 1 pt., 1 gal.

Lantabs (33 mg. for immediate action and 67 mg. for delayed action/Tab.) Bot. 100s

Olet.s (2%) Tube 50 Gm., Jar 1 lb.

Sol.s Nasal (0.5%) bot, w/dropper 15 cc.; nasal spray (0.5%) nebulizer 15 cc.

Tab.s Regular (50 mg.) Bot. 100s, 1000s; delayed action (50 mg.) Bot. 100s, 1000s.

Use: Antihistamine.

W/Antistina. See: Dibistina, Tab. (Ciba).

W/Penicillin-PBZ & Pot. penicillin G. See: Penicillin-PBZ, Tab. (Ciba).

W/Ephedrine. (Ciba) Tripeleannamine HCl 25 mg., ephedrine sulfate 12 mg./Tab. Bot. 100s.

Use: Orally; antihistamine.

W/Nupercaline base & combination. See: Tricalinal, Oint. & Supp. (Ciba)

W/Privine. (Ciba) Tripeleannamine HCl 0.25%, naphazoline HCl 0.25% in aqueous solution. Plastic nebulizer 15 ml.

Use: Relieve nasal congestion.

W/Ritalin (methyl-phenidyl-acetate). See: Plitmasin, Tab. (Ciba)

PYBENZAMINE ANESTHETIC. (Ciba) Tripeleannamine HCl 2%, chlorbutanol 1/2% Jelly, Tube 1 oz. Sol. Bot. 30 cc.

PYBENZAMINE ANESTHETIC. (Ciba) Tab.s 600 mg. Boxes 12s and 100s; Sol.s 2% Bot. 30 ml.; Jellys 2% Tube 1 oz.

Use: Most procedures requiring a local anesthetic

PYBENZAMINE EXPECTORANT W/Codine and Ephedrine. (Ciba) Pyribenzamine citrate 7 1/2 gr., codeine phosphate 2 mg., ephedrine sulfate 2 1/2 gr., amm. chloride 20 mg./cc. Bot. 1 pt., 1 gal.

Use: Antitussive agents for control of severe cough.

PYBENZAMINE EXPECTORANT W/Ephedrine. (Ciba) Pyribenzamine 7 1/2 gr., ephedrine sulfate 2 1/2 gr., amm. chloride 20 mg./cc. Bot. 1 pt., 1 gal.

Use: Relief of cough.

* Generic name recognized by N.F., U.S.P. or A.M.A.

PYBENZAMINE HYDROCHLORIDE W/Zirconium azide. See: Antivy, Lot. (Ciba)

PYRIBEX "50." (Nian) Vit. B-1 50 mg., Vit. B-6 50 mg./cc. Vial 10 cc.

Use: (I.V. or I.M./cc.) Twice weekly.

PYBICARDYL. See: Nikethamide (Various Mfrs.)

PYRICIDIN. (Nepera) (Isanioxid, Isonicotinic Acid Hydroxide) Tab. 50 mg., Bot. 100s, 1000s.

Use: Orally; antitubercular therapy.

PYRIDEX. (Lantest) Pyridoxine 25 mg., thiamine HCl 3 mg./Tab. Bot. 10s, 50s, 100s.

Use: Orally; Vit. B-6 therapy.

2-PYRIDINE ALDOXIME METHIODIDE. See: PAM. (Borden Co.)

PYRIDINE-3-CARBOXYLIC ACID. See: Nicotinic Acid

PYRIDINE-BETA-CARBOXYLIC ACID DIETHYL AMIDE, N.N.R. 1944. See: Nikethamide (Various Mfrs.)

3-PYRIDINEMETHANOL. See: Ronlocol, Prep. (Roche)

PYRIDIUM (Warner-Chilcott) (Phenylazopyridine HCl) 2,6-Diamino-3-phenylazopyridine HCl. Tab (1 1/2 gr.) Tube 12s, Bot. 50s, 500s, 1000s. Sol. (1%) Bot. 100 cc.

Use: Urinary antiseptic and anesthetic

PYRIDIUM TBI-SULFA. (Warner-Chilcott) Phenylazodiamino-pyridine HCl (Pyridium) 150 mg. W/167.0 mg. each of sulfadiazine, sulfamerazine and sulfomethazine/Tab. Bot. 30s.

Use: Acute urinary tract infections.

* **PYRIDOSTIGMINE BROMIDE, N.N.O.** 1962, U.S.P. Tabs. U.S.P. Dimethyl carbamic ester of 3-hydroxy-1-methylpyridinium bromide. See: Mestinon, Tab (Roche)

PYRIDOXAL. Part of pyridoxine, Vit. B-6. 3-Hydroxy-4-formyl-5-hydroxymethyl-2-methylpyridine.

PYRIDOXAMINE. Part of pyridoxine, Vit. B-6. 3-Hydroxy-4-aminomethyl-5-hydroxymethyl-2-methylpyridine.

* **PYRIDOXINE HYDROCHLORIDE, U.S.P., N.N.D.** 1961, A.D.R. Vit. B-6. Mixture of pyridoxal, pyridoxal and pyridoxamine 1s., Tabs. U.S.P.

See: Box-dox (Barry-Martin)

Beadox, Tab. (Promo)

Bedoxine, Amp. (Massengill)

Beesix, Tab. (Tetagar)

HexaSetalin, Amp., Tab., Vial (Lilly)

Hexavitex, Amp., Tab., Vial. (Parke, Davis)

Use: Motion sickness, nausea & vomiting.

W/Bentyl HCl, decapryl succinate. See: Bendin, Tab. (Marrell)

W/Bonamina See: Bonadoxin, Tab (Pfizer)
 W/Isoniazid. See: P I N, Tab (Lanett)
 W/Linoleic Acid See: Arcofac, Emul (Armour)
 W/Linoleic acid, Vit. E See: Elnodazine Emul (Pfizer)
 W/l Lysine HCl, & milk powder defatted See: Lysidox, Pow (Gray)
 W/Marezine See: Moredox, Tab (Burroughs Wellcome)
 W/Mecizine HCl See: Bonadoxin, Tab (Roerig)
 W/Pantothenylol See: Pyripan, Oint (Danley Evans)
 W/Pantothenyl alcohol, hexachlorophene, & menthal See: Vi Cuta, Skin Cream (Columbu)
 W/Phenobarbital See: Daxilane, Tab (Haack)
 W/Pyrimidine maleate See Naudox, Cap (Dumas Wilson)
 W/Salicylamide methenamine mandelate See Lithrol w/Pyridoxine Tab (Columbus)
 W/Vit B 1 See Thiamine HCl, Prep (Various Mfr)
 W/Vit B 1, Vit B 12 See Orazin, Tab (Stuart)
 W/B 1 potassium phosphate potassium citrate carbohydrates See Prenausen Trache (Walker)
 W/Vit B 12, Intrinsic factor, vit C, folic acid choline See: Covitrol, Cap (Flint, Eaton)
 W/Vitamins See: Vitamin Prep
PYRIDOXOL, Part of pyridoxine, Vit B 6 3 Hydroxy 4,5-dihydroxymethyl 2-methylpyridine
 ***PYRIDYL CARRINOL** See Ronicalol Et & Tab (Roche)
PYRINHIST COUGH SYRUP. (Evron) Dihydrocadinane bitartrate 10 mg, pyrimidine maleate 50 mg, Pot guaiacalsulfonate 0.518 Gm, ammonium chloride 0.518 Gm, tartar emetic 5 mg / 8 oz Bot 4 oz, 1 gal Use: Sedative expectorant
PYRINHIST JR. (Evron) Same formula as Pyrihist Cough Syrup but cherry flavor Bot 4 fl oz
PYRILAMINE PROMOTHEOPHYLLINATE. See: Pyrabrom
 W/2-amino-2-methyl-1-propanol See: Bromaleate
 ***PYRILAMINE MALEATE, USP, NND 1962**; Tab, USP 2 [(2-Dimethylaminoethyl) (p-methoxybenzyl)amino] pyridine maleate Pyranisamine, NNR 1950, Pyranitamine Pyramyl, Antipyridamine Available: Cap sustained action, Cream, Syrup, Tab lets

* Generic name recognized by NF USP or A.M.A.

Uses: (25 to 50 mg. i i d) Antihistamine
 See: Alfa Hist, Tab (A.P.C.)
 A N, Tab (Boyle)
 Antamine, Tab (Grave)
 Anihist, Tab (Lipton)
 Antopic, Cream (Smith Dorsey)
 Capsamine, Tab (Durist)
 Diamide, Tab (Merck Sharp & Dohme)
 Dorantamin, Tab (Smith Dorsey)
 Enurmay, Tab (Tilden)
 Nstaborf, Tab (Lincoln)
 Nstacop, T D, Tab (Ethical Pharm)
 Nstacin, Tab (Walker)
 Nstalan, Tab (Hance)
 Nstlan, Tab (Sherman)
 Nstapyran, Prep (Hartan & Converse)
 Kriptin Tab (Whitehall)
 Maranhist, Tab (Morla)
 Minihist, Tab (Ives Cameron)
 Neo Antergon, Tab, NNR (Merck Sharp & Dohme)
 Nisaval, Tab (Vale)
 Paramol Tab (Columbus)
 Paraminyl, Tab, NNR (Buffington)
 Parmol, T D Cap (Central Pharmacol)
 Pymafed Tab (Lloyd)
 Pyramol, Tab (Columbus)
 Pyramaleate Tab (Vanpelt & Brown)
 Renstamin Tab (Strassenburgh)
 Stangen, Tab, NNR (Physicians Drug)
 Stalamin, Tab (Bowman Bras)
 Thylogen, Tab, NNR (Rorer)
 W/B Bromotheophylline See Pyrabrom
 W/Pamabrom See: Pamabrom W/Pyrimidine
PYRILAMINE TANNATE W/Chlorpheniramine tannate, d-amphetomine tannate See Nalertian, Tab (Neisler)
 W/Chlorpheniramine tannate, phenylephrine tannate, ephedrine tannate, carbapentane tannate See Rynatan Expectorant Liq (Irwin, Neisler)
 ***PYRIMETNAMINE, USP NND 1962**; Tabs USP 24 Diamina 5 p chlorophenyl 6 ethyl pyrimidine See: Daraprim Tab (Burroughs Wellcome)
PYRIPAN OINTMENT. (Danley Evans) Pyridoxine & pantothenol in water abarbing base Jar 1 oz
 Use Topically; dermal disorders
PYRISAL (Klug) Pyrilamine maleate 15 mg, ephedrine sulfate 5 mg salicylamide 140 mg caffeine 20 mg Vit C 30 mg / Cap Rot 100s
 Use Cold therapy
PYRISTAN (Arcum) Phenylephrine HCl 8 mg, phenylpropanolamine HCl 15 mg, chlorpheniramine maleate 3 mg, pyrilamine maleate 10 mg / Cap Bot 50s
 Use: Decongestant and antihistaminic

- PYRITHEN.** See: Chlorothen Citrate, U.S.P.
- PYRITHIAD.** (Lakeside) Vit. R-6 20 mg., B-1 2 mg./tab. Bot. 100s. Parenteral: lyophilized. Vial 10 cc.
Use: Antinausea.
- PYRI-THIME INJ.** (Ortega) Thiamine HCl 100 mg., pyridoxine HCl 100 mg./ml. Vial 10 cc.
Use: Nausea of pregnancy, seborrheic dermatitis, thiouracil-induced granulopenia, hyperemesis gravidarum.
- PYRIZIDIN, N.N.D. 1959.** (Nepera) Isoniazid Tab. 50 mg., 100 mg., Bot. 100s, 1000s.
- PYRODOL.** (Central) Salol, camphor, eucalyptol in an emollient base. Jar 1 lb.
Use: Soothing dressing for minor burns and scalds.
- PYROGALLOL, N.F. X.** Pyrogallol acid
- PYROGESIC PEDIATRIC DROPS.** (Hydrex) Methampyrone sodium 250 mg./cc. Bot. 1 oz.
Use: Analgesic-Antipyretic.
- PYROGESIC INJ. & LIQUID.** (Hydrex) Liquid: Methampyrone sodium 500 mg./sp. Injectable: Methampyrone sodium 500 mg./ml. Bot. 1 pt., 1 gal. Vial 10 cc.
Use: Severe pain.
- PYRONIL, N.N.D. 1962.** (Lilly) Pyrobutamine phosphate. Tab. 15 mg. Bot. 100s.
Use: Antihistaminic.
- W/Histadyl, clopane.** See: Co-Pyranil, Cop. (Lilly)
- *PYROPHENINOANE, 1-(1-methyl-3-pyrrolidylmethyl)-3-phenylindane.** (Mead Johnson)
- PYROTHIAMIN.** (New Mexico Pharm.) Pyridoxine HCl 100 mg., thiamine HCl 100 mg., chlorbutanol 0.4%/cc. Vial 10 cc., 30 cc.
Use: Pernicious vomiting of pregnancy, adolescent acne, etc.
- *PYROXAMINE MALEATE.** (Robins) 3-(p-chlorobenzhydryloxy)-1-methyl-pyrrolidone maleate
Use: Antihistamine.
- *PYROXYLINE, U.S.P.** Saluble guncotton. Cellulose nitrate.
- PYRRALAN.** (Lanett) Methopyrilene HCl 25 mg. & 50 mg./tab. Bot. 100s, 500s, 1000s. Also with A.P.C. as Pyrralan Compound. Also Pyrralan Expectoant & Pyrralan Expectoant DH (dihydrocadinane).
Use: Antihistamine.
- *PYRROBUTAMINE PHOSPHATE, N.N.D. 1962.** 1-p-chlorophenyl-4-pyrrolidino-2-butene. See: Pyranil, Tob. (Lilly)
- W/Clopane HCl, Histadyl.** See: Co-Pyranil (Lilly)
- W/Penicillin V potassium, aspirin, acetophenetidin, caffeine.** See: V-Kor, Tab. (Lilly)
- *Generic name recognized by N.F., U.S.P. or A.M.A.**
- W/Reserpine.** See: Sandril W/Pyranil, Tob. (Lilly)
- *PYRROCAINE.** (Endo) 1-pyrrolidinoacetamide See: Endocaine.
- PYRROLAZOTE, N.N.D. 1962.** (Upjohn) Pyrrhiazole. Elixir, as citrate, 2.5 mg./cc. Bot. 3 pt., 1 gal. Tab., as HCl, coated white 25 mg., 100 mg. Bot. 100s, 1000s, 5000s; laminated (HCl) coated pink, prolonged action, contains 2.25 mg. layers, Bot. 100s
Use: Antihistamine.
- W/Orthoxine HCl, acetophenetidin, orpirin & caffeine.** See: Pyrrazote, Cop. (Upjohn)
- N-(PYRROLIDINOMETHYL)TETRACYCLINE.** Pyrralidinomethyl.
See: Raltetracycline.
Sytetrin (Rialol)
Vefacycline (Squibb)
- PYRROXATE.** (Upjohn) Chlorpheniramine maleate 2 mg., orthoxine HCl 25 mg., acetophenetidin 2½ gr., aspirin 3½ gr., caffeine ½ gr./Cap. Bot. 24s, 500r. Disp. pk 600s.
Use: Treatment of common cold & allergic rhinitis.
- W/Codeine Phosphate.** (Upjohn) Chlorpheniramine maleate 2 mg., orthoxine HCl 25 mg., acetophenetidin 2½ gr., aspirin 3½ gr., caffeine ½ gr., codeine phosphate ¼ gr./cap. Bot. 100s, 500s.
Use: Treatment of common cold
- *PYRVINUM PAMOATE, N.N.D. 1962.** 6-Dimethylamino-2-[2-(2,5-dimethyl-1-phenyl-3-pyrryl)vinyl]-1-methylquinolinium Pamoate. Poquill. Vanquin. See: Pavan (Parke, Davis)
- PYTOSIN LIQUID.** (Blue Line) Benzoic acid 52 gr., salicylic acid 26 gr., benzocaine 8 gr./oz. Bot. 1 oz., 1 pt., 1 gal.
Use: Topical; fungicide & desquamative.
- PZA.** See: Pyrazinamide, Tob. (Merck Sharp & Dohme)
- Q. A. E. (Hallinghead Corp.) 2-Hydroxypropylamine nitrile.** Paw. 1 lb.
Use: In solution, 0.05 & 0.1% as rust inhibitor.
- QUADA, CREAM.** (Sutcliffe & Care) Tyrrhricin 0.1%, benzocaine 5%, pyrilamine 1%, cetyl dimethyl ethyl ammonium bromide 2%. Cream, Jar 1 lb.
- QUADAMINE.** (Tutog) d-Amphetamine sulf. 15 mg., amobarbital 45 mg., Vit. A 6600 u., D 500 u., R-1 1.6 mg., B-2 2.5 mg., niacinamide 155 mg., C 30 mg., ferrous sulf. 20 mg., cobalt sulf. 0.49 mg., copper sulf. 2.8 mg., sodium molybdate 0.45 mg., zinc sulf. 3.9 mg., pot. iodide 0.13 mg./Timed Cap. Bot. 100s, 500s, 1000s
Use: Treatment of the obese patient

QUADETT DIPLETT. (Elder) Sulfadiazine 13 mg, sulfamerazine 37 mg, sulfamethazine 62 mg, sulfaguanidine 13 mg /Tab White & red mottled Bot 100s, 1000s

QUADETTs, TABS. (Elder) Sulfadiazine 50 mg, sulfamerazine 150 mg, sulfamethazine 250 mg, sulfaguanidine 50 mg /Tab Yellow Bot 100s, 1000s

QUADRACINE. (Millard) Scopolamine MB 0.93 mg, aluminum hydroxide 200 mg, magnesium hydroxide 200 mg, ascorbic acid 7.5 mg, milk protein 100 mg /Tab Bot 100s

Use: Ulcer treatment

QUAD-RAMCID, SUSP. (Elder) Sulfamethazine 250 mg, sulfamerazine 150 mg, sulfadiazine 50 mg, sulfaguanidine 50 mg, alcohol 2%/5 cc Bot 2 oz, 1 pt, 1 gal

Use: Sulfonamide therapy

QUADRA-SED LIQUID. (C D Smith) Pentobarbital Sod 15 mg, phenobarbital Sod 15 mg, butobarbital Sod 15 mg, Secobarbital Sod 15 mg /5 cc Bot 1 pt

Use: Sedative

QUADRATES (Cutter) Sodium chloride—(4 molar) [4 mEq/cc] Vial 30 cc, Box 12s
Sodium lactate—(4 molar) [4 mEq/cc] Vial 30 cc, Box 12s
Ammonium chloride—(4 molar) [4 mEq/cc] Vial 30 cc, Box 12s
Potassium acetate—(4 molar) [4 mEq/cc] Vial 30 cc, Box 12s

Use: Electrolyte therapy

QUADRIGEN (Parke, Davis) Immunization of children against diphtheria tetanus, pertussis and poliomyelitis Vial 4.5 cc

Use: IM (1 cc) immunization

QUADRINAL. (Knoll) formerly Quadrodide
Ephedrine HCl 24 mg, phenobarbital 24 mg, phyllin 2 gr Pot iodide 5 gr /Tab Bot 100s 500s 1000s

Use: Asthma & related allergic respiratory conditions

QUADRINAL SUSPENSION. (Knoll) Ephedrine HCl 3/16 gr, phenobarbital 3/16 gr, theophylline calcium solicylate 1 gr, potassium iodide 2 1/2 gr /5 cc Bot pts

Use: Bronchospasm and wheezing

QUADRIGEN VACCINE. (Parke Davis) (Diphtheria, tetanus pertussis poliomyelitis multiple) Vials 4.5 cc

Use: IM, or SC immunization

QUADRQDIDE See: Quadrinal (Knoll)

QUADRUPLE SULFONAMIDES. See: Sulfonamides

QUADSUL TARS AND QUADSUL JR. WAFERS (Vale) Tabs, Sulfadiazine 100 mg, sulfamerazine 100 mg, sulfamethazine 100 mg, sulfacetamide 200 mg /tab

* Generic name recognized by N.F., U.S.P. or A.M.A.

Wafers: Contain 1/4 of the above amt / wafer
Tabs: Bot 100s, 500s; Wafers: 100s, 500s, 1000s

Use: Bacterial infections

QUAD-SULFA. (Zemmer) Sulfadiazine 2 1/2 gr, sulfamerazine 2 1/2 gr, sulfamethazine 1 1/2 gr, Sod sulfacetamide 1 gr /tab Bot 100s.

Use: Sulfonamide therapy

QUALATUM. (Almay) Oily type oint for 4 oz, 1 lb, 5 lb

Use: Ointment base.

QUANTI-VITE-F. (Hoyt) Vit A 3000 USP u, D 400 USP u, C 60 mg, B-1 1 mg, B-2 12 mg, B-6 1 mg, niacinamide 10 mg, fluoride 0.5 mg /D 6 cc Pediatric Drops Bot 50 cc

Use: Dental preparation, fluoride/vitamin preparation

QUANTI-VITE (F) IQZI-TARS. (Hoyt) Vit A 4000 u, D 400 u, C 75 mg, B 12 mg, B-6 2 mg, niacinamide 15 mg, fluoride 1 mg /orange Bot 60s

Use: Vitamin and sodium fluoride combination

QUANTRIL. (Roerig) Benzquinamide

Use: CNS depressant

QUARZAN BROMIDE. (Hoffmann LaRoche) Clidinium Bromide

Use: Anticholinergic agent

QUATRASAL. (Dome) Salicylanilide 5% B hyamine 1% (T.C.A.R.) liq, Bot 1 oz, 4 oz, 1 pt, also 2 oz spray bot

Use: Topically, antibacterial & fungicidal

***QUAZOLINE.** (Mead Johnson) 4 [(1-methyl-3-pyrrolidiny)methyl] aminoquinoline

Use: Bronchodilator, anti-inflammatory agent

QUELICIN CHLORIDE, NND 1962 (Abbott) Succinylcholine chloride Amp (50 mg/cc) 10 cc Amp (100 mg/cc) 10 cc Vial (20 mg/cc) 10 cc Abbo Vials (25 mg/ml) 20 ml & 40 ml Box 1s, 10s

Use: Short-acting muscle relaxant

QUELIDRINE SYRUP. (Abbott) Oestromethorphan hydrobromide 10 mg, chlorpheniramine maleate 2 mg, ephedrine HCl 5 mg, phenylephrine HCl 5 mg, ammonium Cl 40 mg, specac fluidextract 0.005 cc, ethyl alcohol 2%/5 cc Bot 4 oz, 1 pt

Use: Allergic disorders

***QUERCETIN.** Active constituent of rutin See: Quercin, Tab (Abbott)

W/Aminophylline, ascorbic acid, reserpine See: Rutaminol RO, Tab (Schenley)

W/Reserpine & Vit C. See: Quer Serpin Tab (Haskell)

QUER-SERPIN. (Haskell) Reserpine 0.25 mg., quercetin 10 mg., Vit. C 50 mg./Tab. Bot. 100s.

Use: Hypertension.

QUERTINE. (Abbott) Quercetin. 3,3',4',5,7-Pentahydroxyflavone. Tab. (10 mg., 20 mg) Bot. 100s, 1000s.

Use: (10 to 20 mg. I.I.d.) For treatment of capillary fragility.

W/Ascorbic acid. (Abbott) Quertina 10 mg., ascorbic acid 50 mg./Tab. Bot. 100s, 1000s. Use: (1 I.I.d.) Treatment of capillary fragility.

QUIACTIN, N N O. 1962. (Merrell) Oxamide, [2-ethyl-3-propylglycidamide]. Tab 400 mg. Bot. 50s.

Use: Tranquillizer. See: Oxamide (Merrell)

QUIRRON. (Mead Johnson Lab.) Theophylline 150 mg., glyceryl guaiacolate 90 mg./Cap. Bot. 100s.

Use: Asthmatic disorders.

QUIRRON ELIXIR. (Mead Johnson) Theophylline 150 mg., glyceryl guaiacolate 90 mg., in a hydro-alcoholic vehicle/15 ml. Bot. R. N. oz.

Use: Treatment of asthma.

QUIDE. (Plitman-Moore) Piperacetazine, 2-acetyl-10-(3-[4-(β-hydroxyethyl)-piperidino]propyl)phenothiazine.

Use: Treatment of psychoses.

QUISS. (Durst) Pentobarbital ¼ gr., butobarbital 8 mg., phenobarbital 8 mg./Tab. Bot. 100s, 500s, 1000s.

Use: (1-2 m.p.) Sedative.

QUILINE. (Ayerst) Pentapiperide Methylsulfate (formerly Volpipamate Methylsulfate).

Use: Anticholinergic and antisecretory agent.

QUILLAJA EXT. W/Coal tar ext., nitromersol, benzoin, Isopropyl alcohol. See: Roricide, Sol. (Kormet Co.)

QUIMQTRASE. (Smith, Miller & Patch) Alpha-chymotrypsin 750 Perry units, sterile isotonic solution 10 cc./Pkg. Vial (w/ diluent) 10 cc.

Use: Chymotrypsin preparation, colorost surgery aid.

QUINACRINE HYDROCHLORIDE, U.S.P., N.N.D. 1961; Tab., U.S.P. Mepacrine HCl, 3-Chloro-7-methoxy-9-(1-methyl-4-diethylaminoethylamino) acridine dihydrochloride. See: Atobrine HCl. (Winthrop Labs.)

W/Chloroquine, hydroxychloroquine. See: Triquin, Tab. (Winthrop)

QUINAGLUTE. (Wyne) Quinidine gluconate 60 mg., monoethyglycerol 1%, phenol

* Generic name recognized by N.F., U.S.P. or A.M.A.

0.25%/cc. Vial 10 cc. Tab. (5 gr./5 M. Tab) Bot. 30s, 100s, 250s.

Use: I.M., I.V., cardiac arrhythmias.

QUINALBARRITONE. See: Seconal Sod.
QUINAMM. (Walker) Quinine sulf. 250 mg., amleophylline 200 mg./Tab. Bot. 50s.

Use: Treatment and relief of muscle cramps.

QUINETAZONE. 7-chloro-2-ethyl-6-sulfamoyl-1,2,3,4-tetrahydro-4-quinazolinone. (Lederle Labs.)

QUINETTES. (Starck) di-tetrahydroxyquinoline 1.5 gr., sulfadiazine 7.5 gr., hyamine 4 mg., stilbestrol 0.1 mg., sarbic acid 4 mg./Insert. Pkg. 24s.

Use: Trichomonas, vaginitis.

QUINGESTRONE. (Warner-Lambert) 3-(cyclopentylloxy)pregno-3,5-dien-2-one
Use: Progestational agent.

QUINICARDINE. (Feugero) Quinidine sulfate 0.2 Gm./Tab. Bot. 20s.

Use: Orally; quinidine therapy.

QUINIDATE. (M. R. Thompson) Quinidine sulfate, Amp. (3 gr.) 1 cc., Rox 6s, 25s, 100s. Cap. (3 gr.) Bot. 40s, 100s, 1000s.

Use: Auricular fibrillation & paroxysmal tachycardia.

QUINIDEX I.A. (Whittier) Quinidine sulfate 300 mg./T.D. Tab. Bot. 30s.

Use: Auricular fibrillation.

QUINIDINE GLUCONATE, N.F. Inj., N.P. (Ellyl Amp. (80 mg./cc.) 10 cc. (Wynn) Tabs. 5 gr. Bot. 30s, 100s.

See: Quinaglute, Vial (Wynn)

Use: Quinidine therapy.

QUINIDINE HYDROCHLORIDE.

Brewer—Amp. (0.6 Gm.) 5 cc.

Davis, Rose—Tab. (0.125 Gm.) Bot. 50s, 500s

QUINIDINE POLYGALACTURONATE. See: Cardioquin (Purdue Frederick Co)

QUINIDINE SULFATE, U.S.P., N.N.R. 1949; Cap., N.F. Tab., U.S.P.

See: Azarum Compound, Sol. (First Texas) Quinacardine, Tab. (Varick)

Quinidate, Amp. & Cap. (M. R. Thompson) Quinidex I.A., T.D. Tabs (Whittier)

W/Aminophylline. See: Quinamm. Tab (Walker)

QUININE ARSENATE W/Strychnine arsenate, iron arsenate, nucleic acid. See: Arsenoin, Tab. (Elder)

QUININE RISULFATE, N.F.

Use: Orally; analgesic & antipyretic & anti-malarial.

QUININE CARRACRYLIC RESIN, N.N.D. 1959. See: Diagnex. (Squibb)

QUININE DIHYDROCHLORIDE, N.F., Inj., N.F.

Use: I.M., I.V.; antimalarial therapy.

QUININE ETHYLCARBONATE, N N R 1949

See: Equinine, Paw (Various Mfr)
 * **QUININE HYDROCHLORIDE, N F**
 W/Nupercaine Benzyl Alcohol See Anesacaine (New Mexico Pharm)
 W/Urethan (Horlan & Converse) Quinine HCl 0.25 Gm, urethan 0.12 Gm / 2 cc Amp 2 cc, Box 12s, 25s, 100s See Sclerosing agent
 W/Urethan (Lilly) Quinine HCl 2 parts w/urethan 1 part Amp (quinine HCl 0.266 Gm, urethane 0.133 Gm) 2 cc, Box 12s 100s

Use: Sclerosing agent
QUININE GLUCONATE, (Lilly) Amp (0.68 Gm/cc) 10 cc

Use: (0.4 Gm IM) For those who cannot take medication orally

W/Cal glucanagalacta gluconate See: Cal gluquine, Amp (Sandoz)

* **QUININE SULFATE, N F, N N R 1953;**
 Cap Tab N F See Coco Quinine Liq (Lilly)

W/Acetonid, podophyllin, aloin, aconite root, capsicum, atropine sulfate & camphor See: CCO, Tab (Noyes)

W/Acetophenetidin, camphor, podophyllin, aloin, atropine sulfate See Special Formula No 2 Tab or Cap (Upjohn)

W/Aminophylline See Quinamm, Tab (Walker)

W/Ammanium chloride camphor, belladonna fl ext, aconite fl ext See: Carylzo, Tab. (Merck Sharp & Dahme)

W/Ferric chloride, magnesium sulf See: A I A, Sol (Woods)

W/Niacin 3 alpha locopheryl acid succinate See Myadyne Tabs (Canfield)

W/Pyrimine maleate solicylamide, phenacolin, caffeine atropine sulf ascorbic acid See: Histaquin Cap (Taylor)

W/Urea HCl N F; Inj, N F
 Endo—Amp (1½ gr) 2 cc, Box 12s, 25s, 100s

Kirk—Amp (1% & 5%) 2 cc, Box 12s, 100s

Parke, Davis—Amp (1%) 5 cc, Box 6s 25s; (5%) 2 cc, Box 12s 100s Tab, hypodermic (2 gr) Bot 100s, 500s; (5 gr) Bot 100s

Use: SC, IM, IV; antimalarial & sclerosing agent

QUIN-O-CREME, (Walker) Iodachlorhydroxyquin 3% in a washable base Jar ½ lb, 1 lb

Use: Treatment of skin infections

QUINOFORM, (C & M Pharmacal) Iodachlorhydroxyquin 3% Oint, Tube 2 oz, Jar 1 lb

Use: Topically, eczemas & dermatitis

* Generic name recognized by N.F. U.S.P. or A.M.A.

QUINOLINE SULFATE W/Hydroxychloroquine sulfate See: Plaquenil, Tab (Winthrop Labs)

QUINOLOR COMPOUND OINTMENT, (Squibb) Equal parts of white petrolatum & deodorized anhy lanolin containing 10% benzoyl peroxide & 0.5% chlorhydroxyquinolin Tube 1 oz, Jar 1 lb

Use: Impetigo & dressing for infections
 W/Coal Tar See: Tarquinor, Cr (Squibb)

QUINOPHAN, See: Cinchaphen (Various Mfr)

QUINOPLEX, (Gray) Anthraquinolidal glycosides Sennaside A and B 10 mg / Tab Bot 30s

Use: Treatment of chronic constipation

QUINTESS (Lilly) Activated attapulgit 3 Gm, activated attapulgit, colloidal 0.9 Gm citrus pectin 0.3 Gm / 30 cc Plas Bal 6 oz, Glass Bot pint

Use: Control of diarrhea

QUINTEX, (O F Harvey) Vit A 5000 u D 1000 u, B 1 1.5 mg, B 2 2 mg, B-6 0.1 mg, B 12 1 mcg, E 1 U, Cal pantothenate 1 mg, C 37.5 mg, niacinamide 20 mg / Cap Bot 100s, 1000s

Use: Multiple avitaminosis

QUINTRATE, N N D 1962 (Glenn A Beard) Pentaerythritol Tetranitrate

QUINTRONS (Vitamin Corp) Iron & ammonium citrates 3 gr, liver conc 5 gr stomach pow substance 100 mg, Vit B 1 1 mg, B 2 0.5 mg, B 6 0.15 mg, Cal pantothenate 0.3 mg, niacinamide 10 mg, folic acid 0.1 mg, Vit B 12 1 mcg / Cap Bot 100s, 250s, 1000s

Use: Dietary supplement

QUINPENYL NAPHTHOATE, See Pamaquine

Naphthoate (Plasmachin naphthoate, Winthrop Labs)

QUISYRUP, (Elder) Pyrimine maleate 100 mg, ipecac fl ext ammanium chloride / oz Syr 2 oz, 1 pt, 1 gal

Use: Antihistaminic expectorant

QUISYRUP PLUS (Elder) Pyrimine maleate 100 mg, dihydrocodeinone bitartrate 10 mg, ammanium chloride 8 gr, ipecac fl ext 1 min / fl oz Bot 2 oz, 1 pt, 1 gal

Use: Antihistaminic & antitussive

QUOTANE HCl, N N D 1962 (Smith, Kline & French) Dimethisoquin HCl 1 (β Dimethyl aminoethoxy) 3 N butylisoquinoline HCl Oint (½%) Tube 1 oz w/applicator lot (½%) Bot 2 fl oz

Use: Dermatologic conditions

W/Gramicidin, polymixin B sulfate & cetyl dimethylbenzylammonium chloride See: Drixalax, Lax (Smith, Kline & French)

QUOTULES. (Central) Vit. A 5000 u., D 500 u., B-1 3 mg., B-2 2.5 mg., B-6 0.5 mg., C 40 mg., niacinamide 20 mg., pantothenic acid 5 mg., Vit. B-12 2 mcg., folic acid 0.25 mg./Cap. Bot. 100s, 500s, 1000s
Use: Multivitamin therapy.

RABELLON. (Merck Sharp & Dohme) Hyoscyamine HBr 0.4507 mg., atropine sulfate 0.0372 mg., scopolamine HBr 0.0119 mg./tab. Bot. 100s, 1000s.
Use: (1 q. 3 h.) Relief in postencephalitic parkinsonism.

RABIES IRADOGEN. (Parke, Davis) Pkg. 7 doses.

Use: Preventing rabies after exposure.

***RABIES VACCINE, U.S.P., N.N.R. 1950.**

Lederle—Vial 0.5 cc., 7s, 14s.

National—Vial 0.5 cc., 7s, 14s.

Parke, Davis—Vial 7 doses, 14 doses.

Pitman-Moore—Vial 1 cc., 7 doses, 14 doses.

Merck Sharp & Dohme—Vial 7 doses.

Wyeth—Vial 7 doses.

***RABIES VACCINE, N.N.D. 1962.** (Lilly)

Dried killed virus (Duck embryo) developed by the Powell-Culbertson method but has 1 to 4000 addition of beta-propiolactone and buffers. Each dose in individual vial (dry) with amp. of diluent. 7 doses/pkg.

Use: Treatment 1 subcutaneous dose daily for 14 days (free of paralytic factor)

RABUDIX-O.R. (Lynn) d-Amphetamine sulf.

15 mg., dl-amphetamine sulf. 15 mg., butabarbital sod. 45 mg./Cap. Bot. 100s, 500s, 1000s

Use: Treatment of obesity

RACEMETHORPHAN, N.N.R. 1953. See: Dramoran.

RACEMIC CALCIUM PANTOTHENATE, U.S.P.

See: Calcium Pantothenate, Racemic, U.S.P.

RACEMIC DESOXY-NOR-EPHEDRINE,

N.N.R. 1934. See: Amphetamine (Various Mir.).

RACEMIC EPHEDRINE HCl, N.N.R. See:

Racephedrine HCl. (Upjohn)

RACEMIC PANTOTHENIC ACID, U.S.P. See:

Vitamin, Preparations.

RACEMISTAT. (Pascal Co.) dl-Epinephrine

HCl B to 100 and zinc chloride 2 to 100.

Bot. 1/2 oz., 1 oz.

Use: Dental topical anticoagulant

***RACEMORPHAN, N.N.R. 1953** See: Dramoran, Amp. & Tab. (Roche).

***RACEPHEDRINE HYDROCHLORIDE, N.F.,**

N.N.D. 1959; Cap., N.F., Sol., N.F. (Upjohn)

Synthetic racemic ephedrine HCl. Cap. (1% gr.) Bot. 40s, 250s, 1000s. Sol. (1%) Bot. 1

oz.

* Generic name recognized by N.F., U.S.P. or A.M.A.

fl. oz., 1 pt., 1 gal. See: Ephetanin (Merck Sharp & Dohme)

Use: Vasoconstrictor & nasal decongestant. See also Ephedrine Combinations.

W/Acetophenetidin, aspirin, pyrilamine maleate. See: Nicapfen, Tab. (Ulmer)

W/Acetophenetidin, salicylamide, pyrilamine maleate, ascorbic acid, Dover's powder. See: Cenamal, Cap. (Central)

W/Aminophyllin & diphenhydramine.

See: Hyadrin, Tab. (Searle).

Hydrillin W/Racephedrine Hydrochloride, Tab. (Searle).

W/Aminophylline & phenobarbital. See: Amadrine, Tab (Searle).

W/Chlorpheniramine maleate, phenyltoloxamine dihydrogen citrate See: Algic, Tab (Spencer)

W/Isoproterenol, phenobarbital, neohydrillin. See: Asophylline, Tab. (Crystal)

W/Pentobarbital, choline theophyllinate. See: Chalarace, Tab. (Nepera)

W/Phenyltoloxamine dihydrogen citrate. See: Ephoxamine (Spencer Labs)

W/Secobarbital, synaphylate, potassium iodide, niacinamide. See: Synate-M, Tabs. (Central)

W/Synaphylate, secobarbital, potassium iodide, niacinamide See: Synole, Tab (Central)

W/Theophylline Sod. glycinate & phenobarbital.

See: Glythionate W/Phenobarbital & Racephedrine, Tab. (Palk).

Synaphedal, Tab. (Central).

RACNOMATE, N.N.D. 1962. (Abbott) Radio-Chromate (Cr⁵¹), Sod.

Use: Radioactive Isotopes.

RA-CO-LIN. (Paul Money) Heparin sod. 25 mg., choline chloride 100 mg., Vit. B-12 50 mcg., folic acid 1.5 mg., niacinamide 50 mg., liver injection 2 u./cc. Amp. 10 cc

Use: Anticoagulant and hematinic.

RADIOACTIVE IODINATED BLOOD ALBUMIN. See: Risa, Vial (Abbott)

RADIO-ACTIVE ISOTOPES.

See: Alphanon (Radioactive Products)

Aurcoloid

Rochamate

Radio-Gold Colloid

Radio-Iodinated Serum Albumin (Human)

Risa, (Abbott)

Sodium Radio-Chromate

Sodium Radio-Iodide

Sodium Radio Phosphate

RADIOCAPS, N.N.D. 1962. (Abbott) See:

Radio-Iodide, Sod

RADIO-CHROMATE (Cr⁵¹), SODIUM, NND 1962 See Rachomate (Abbott)

***RADIOCYANOCOBALAMIN SOLUTION**, USP, See Racobalamin Solution (Abbott)

***RADIO-GOLD** (Au¹⁹⁹) COLLOID, USP NND 1962 See Aurcolaid (Abbott)

Use Irradiation therapy
RADIO-HIPPURAN (Abbott) Radioactive hippuran (I¹³¹) Vial 0.1, 0.25, 0.5, and 1 to 10 millicuries

Use: Diagnostic agent for study of kidney function

RADIO-IODIDE (I¹³¹), SODIUM, NND 1962 (Abbott)

Use Radioactive isotopes

***RADIO-IODINATED** (I¹³¹) **SERUM ALBUMIN** (Human), USP NND 1962 See Risa Vial (Abbott)

RADIO-PHOSPHATE (P³²), SODIUM, NND 1962

Use Radioactive isotopes

RADIOL GERMICIDAL SOLUTION, ADR (Novacol Chem) Cetyl dimethyl ethyl ammonium bromide 0.25%, Sod bicarbonate 0.08%, Sod carbonate 0.17%, Sod nitrite 0.25%, propylene glycol 2%, color & water Regular, Bot 1 qt Concentrate w/ anti rust tab, Bot 1 qt

Use Bactericidal agent

RADON, NNR 1934 See: Alphatron (Radioactive Prod)

RAMETIN, NND 1962 (Bio Rama) vit B 12 Tab (5 mcg, 10 mcg, 30 mcg) Bot 100s Vial (10 mcg, 15 mcg, 30 mcg/cc) 10 cc

Use: Anemias

RAMOCILLIN, (Bio Rama Drug Co) Procaine penicillin G 300M u/dose Aq Susp, Vial 1, 5, 10 doses

Use: Penicillin therapy

RAMOSUL-100 (Bio Rama Drug Co) Pot penicillin G 100M u w/sulfadiazine, sulfamerazine & sulfamethazine 167 mg of each/Tab Bot 100s

Use: Penicillin w/triple sulfonamide therapy

RAMOSUL-200 (Bio Rama Drug Co) Pot penicillin G 200M u, w/sulfadiazine, sulfamerazine & sulfamethazine 167 mg of each/Tab Bot 100s

Use: Orally; penicillin & triple sulfonamide therapy

RAMOSUL-250 (Bio Rama Drug Co) Pot penicillin G 250M u, w/sulfadiazine, sulfamerazine & sulfamethazine of each 167 mg/Tab Bot 100s

Use: Penicillin & triple sulfonamide therapy

RAMSES JELLY, NNR 1935 (Schmid) Dodecethylene glycol monolaurate 5%, boric acid 1%, glycerin, fragaanth carboxy

*Generic name recognized by NF, USP or A.M.A

methylcellulose & preservative Tube 5 oz. Use: Contraceptive

RAMYCIL-AS, (Bio Rama Drug Co) Procaine penicillin G 400M u, dihydrostreptomycin sulfate 0.5 Gm/dose, Vial 1 dose

Use: Penicillin & streptomycin therapy

RANDAMIN "B" (Rand) Vit B 1 25 mg, B 2 12 mg, B 6 15 mg, nicotinamide 125 mg, Cal pantothenate 5 mg, choline 100 mg, Inositol 50 mg, liver, yeast/Cap Bot 50s, 100s, 1000s

Use: (3 daily) Vitamin B complex therapy

RANDAPLEX, (Rand) Vit B 1 3 mg, B 2 3 mg, B 6 0.25 mg, Cal pantothenate 5 mg, niacinamide 20 mg, liver conc/Cap Bot 100s, 1000s

Use Dietary supplement

RANOCINE, (Cook-White Lab) Propoxy caine HCl ADR

RAOLEIN, (Abbott) Iodine¹³¹ labeled triolein

Use: Diagnostic aid

RAOVIN (Abbott) Tolpovidone I¹³¹

Use Differential diagnosis of source of hypotubulinemia

RAPACODIN BIARTRATE (Knoll) Para codin Drocade Dihydrocodeine bitartrate Amp 30 mg, 1 cc Box 6s 100s Vial, multiple dose (30 mg/cc) 20 cc 1abr (10 mg, 20 mg) Bot 100s

Use Analgesic and antitussive

RAPADYNE (Knoll) Dihydrocodeinone bitartrate 20 mg, acetylsalicylic acid 200 mg, phenacetin 150 mg, caffeine 50 mg/Tab

Use Analgesic

RAPAX INSERTS, (Elder) Sod bicarbonate 0.7 Gm, sod biphosphate 0.8 Gm In a water soluble base/Insert Rot 12s

Use Rectally, laxative

RAPHARIL, (Sandoz) Raphanus sativus niger principle (black radish) 5 Gm, dehydrocholic acid 0.15 Gm/Tab Rot 40s, 200s

Use: Choleragogue & cathartic

RAPHETAMINE PHOSPHATE, NND 1962 (Strassenburgh) di Amphetamine phosphate Elixir (5 mg/4 cc) Bot 1 pt, 1 gal Sol (10 mg/cc) Vial 10 cc, 6s Tab (5 mg) Bot 100s, 1000s

Use: Sympathomimetic agent

RARICAL, (Ortho) Iron 25 mg, Cal 85 mg as ferrous Cal citrate & tricalcium citrate/Tab Rot 100s, 1000s Also with Vitamins Rot 100s

Use: (2 t i d w/meals) Calcium & Iron therapy

RASBERIN, (E S Miller) Salicylamide 1 gr/D 6 cc Bot w/dropper 30 cc, 1 pt

Use: 1.5 gr q 4 hr; analgesic & antipyretic

RASHENE, (Squibb) Protective creme. Tube 1½ oz., 3 oz.
Use: Prevent or relieve diaper rash.

RASTINON. See: Tolbutamide

RA-TEST, (Hyland Labs.) Lotex-globulin, control serum, glycine-saline buffer. Testz 100s.
Use: Screening test for rheumatoid arthritis.

RATTLESNAKE RITE THERAPY. See: Antivenin, Snake Polyvalent.

RAUCAP, (Ethical Pharm.) Reserpine alkaloid 0.25 mg., 50 mg./Tab. timed disintegration. Bot. 30s
Use: Sedative and tranquilizing agent

RAUCYLATE, (Flint, Eaton) Rauwolfia serpentina whole root 25 mg., aspirin 300 mg., aluminum hydroxide gel dried 65 mg., ascorbic acid 25 mg./Cap. Bot. 100s.
Use: (1-2 t.i.d.) Antacid w/sedative and tranquilizing agent.

RAUDIXIN, N.N.D. 1962. (Squibb) Rauwolfia whole root, Tab. (50 mg. & 100 mg.) Bot. 100s, 1000s.
Use: 1-2 q.i.d., hypotensive A tranquilizing agent.

W/Benzhydroflumethiazide, potassium chloride. See: Rautrax-N, Tab. (Squibb)

W/Flumethiazide, Pot. chloride. See: Rautrax, Tab. (Squibb)

RAUDOLFIN, (Preme) Rauwolfia serpentina 50 mg., 100 mg./Tab. Bot. 100s, 1000s.
Use: Hypotensive agent

RAUDONNA, (Vanpelt & Brown) Reserpine 0.1 mg., hyoscyamine sulfate 0.1286 mg., atropine sulfate 0.0250 mg., hyoscyne HBr 0.0074 mg./Tab. Bot. 100s, 1000s.
Use: (1 q.i.d.) Antispasmodic A tranquilizing agent.

RAUFIA ENCOTE, (Paul Money) Mannitol hexanitrate 30 mg., veratrum viride 100 mg./Tab. Bot. 100s, 1000s.
Use: Essential hypertension.

RAULEN, (Paul Money) Reserpine 0.25 mg., 1.0 mg. Bot. 100s.
Use: Hypertension.

RAULOR, (Taylor Labs.) Rauwolfia serpentina alkaloids 1 mg./Tab. Bot. 100s, 1000s.
Use: Sedative & tranquilizing agent.

RAULODYN, N.N.D. 1962 (Tutog) Reserpine.

RAUNESCINE, (S. B. Penick) An alkaloid of Rauwolfia serpentina. Under study.
Use: Tranquilizing agent.

RAUNORMINE, (S. B. Penick) 11-Desmethoxy reserpine.

RAUPENA, (Crookes-Barnes) Rauwolfia serpentina whole root 50 mg. Tab., Bot. 100s A 1000s.
Use: 4 daily; hypotensive A tranquilizing agent.

RAU-PERTINAL (Crookes-Barnes) Rauwolfia serpentina 50 mg., veratrum viride ext. 75 mg., mannitol hexanitrate 30 mg., homatropine methylbromide 2.5 mg./Tab. Bot. 50s, 100s, 500s
Use: 1 t.i.d.; hypertension.

RAUPROTE, (Vale) Rauwolfia serpentina 50 mg., protoveratrine A B 0.2 mg./Tab. Bot. 100s.
Use: (1 t.i.d.) hypertension.

RAURESCIN, (Columbus) Rescinnamine 0.25 mg./Tab. Bot. 100s, 500s, 1000s.
Use: Antihypertensive and tranquilizing agent.

RAURINE, N.N.D. 1962. (Lloyd, Dobney & Westerfield) Reserpine. Tab. (0.1 mg. A 0.25 mg.) Bot. 100s, 0.5 mg. Bot. 100s.
Use: Orally, sedative A tranquilizing agent.

W/Pentaerythritol tetranitrate, See: Respal, Tab. (Lloyd, Dobney & Westerfield)

RAU-SED, N.N.D. 1962. (Squibb) Reserpine. Tab. (0.1 & 0.25 mg.) Bot. 100s, 1000s; (0.5 mg.) Bot. 100s, 500s; (1 mg.) Bot. 30s, 100s, 500s. Tab. 4 mg., Bot. 100s. Inf. (2.5, 5 A 10 mg./cc.) Amp 2 cc. Box 5s.
Use: 0.5-2 mg. daily; hypotensive A tranquilizing agent.

RAUSERIA, (New Eng. Phr. Co.) Rauwolfia serpentina 50 mg./Tab. Bot. 100s

RAUSERPA, N.N.D. 1962. (Tutog) Rauwolfia serpentina.

RAUSINGLE, (Columbus) Reserpine 25 mg./Tab. Bot. 100s, 1000s
Use: Tranquilizing agent

RAU-TAB, (National Drug) Alseraxylon fraction of Rauwolfia serpentina 2 mg./Tab. Bot. 60s, 500s.
Use: 1-2 daily; hypotensive A tranquilizing agent.

RAUTENSIN, (Smith-Dorsey) Alkaloid ext. from rauwolfia serpentina. Tab. (2 mg.) Bot. 100s, 500s, 1000s.
Use: 1-2 daily; hypotensive A tranquilizing agent.

RAUTINA, (Testagar) Rauwolfia serpentina whole root 50 A 100 mg./Tab. Bot. 100s.
Use: Hypotensive agent.

RAUTOTAL, (Columbus) Rauwolfia serpentina, whole root 50 mg., 100 mg./Tab. Bot. 100s, 1000s
Use: Treatment of hypertension

RAUTRAX, (Squibb) Raudixin 50 mg., flumethiazide 400 mg., Pot. chloride 400 mg./Tab. Bot. 100s.
Use: Hypertension.

RAUTRAX IMPROVED, (Squibb) Flumethiazide 50 mg., raudixin 50 mg., potassium chloride 625 mg./Tab
Use: Antihypertensive and diuretic.

RAUTRAX-N (Squibb) Standardized whole root Rauwolfia serpentina 50 mg, benzydromethiazide 4 mg, polorasium chloride 400 mg /Cap shaped Tab Bot 100s
Use: Hypertension

RAUVAL, NND 1962 (Vale Chem) Rauwolfia serpentina whole root Tab 50 mg 100 mg Bot 100s 1000s
Use: Hypotensive & tranquilizing agent

RAUVERA (Smith Dorsey) Rauwolfia serpentina alkaloid 1 mg, veratrum viride alkaloid 3 mg /Tab Bot 100s 500s, 1000s
Use: (1 1/2, p c) Hypertension

RAUWICON (Consol Midland) Rauwolfia serpentina whole root 50 mg, or 100 mg Tab Bot 100s, 200s 1000s
Use: Tranquilizing agent

RAUWIDRINE, (Riker) Alkaloidal ext from rauwolfia serpentina 1 mg, amphetamine sulfate 5 mg /Tab Bot 100r, 1000s
Use: (1 2 p c) Neuroses, alcoholism & obesity

RAUWILOID, NND 1962 (Riker) Al seroxyton Rauwolfia alkaloids Tab (2 mg) Bot 60s 1000s

W/Hexamethonium (Riker) Rauwolfia alkaloids 1 mg hexamethonium 250 mg /Tab Bot 100s, 1000s
Use: Hypotensive & tranquilizing agent

W/Veriloid (Riker) Rauwolfia alkaloids 1 mg, alhaverir 3 mg /Tab Bot 100s, 1000s
Use: (1 1/2) Hypertension

RAUWISTAN-MRT (M. R. Thompson) Contains both alkaloidal & nonalkaloidal components of Rauwolfia serpentina Tab (1 MRT u or 100 mg of Rauwolfia serpentina pow) Bot 100r 1000s 50 mg, Bot 100s 1000s
Use: 1 3 daily; hypotensive & tranquilizing agent

RAUWOLDIN (G F Harvey) Rauwolfia serpentina whole root Tab (50 mg, 100 mg) Bot 100s 1000s
Use: 4 R daily; hypotensive & tranquilizing agent

RAUWOLFIA CANESCENS ALKALOID
See: Harmony, Tab (Abbott)
Rauwolficine

RAUWOLFIA SERPENTINA ACTIVE PRINCIPLES (ALKALOIDS), Deserpidine Res cinnamine NF
See: Rererpine

RAUWOLFIA SERPENTINA ALKALOIDAL EXTRACT (ALSEROXYLON), NND 1962
See: Abten, Tab (Paul Maney)
Kogluoid, Tab (Panray)
Raudalfin, Tabs (Premo)
Rau Tab (National Drug)
Raulor, Tab (Taylor Labs)

* Generic name recognized by NF U.S.P. or A.M.A.

Raulensin, Tab (Smith Dorsey)
Rauwford, Tab (Riker)
Via Serpine Tab (Rawell)
W/Amphetamine sulfate See: Rauwidrine Tab (Riker)

W/Hexamethonium chloride See: Rauwloid
W/Nexamethonium, Tab (Riker)

W/Luminal & theabramine See: Theominal R S, Tab (Winthrop Labs)

W/Mannitol hexanitrate See: Nitranitol R S, Tab (Merrill)

W/Nealhylline & rutin See: Laten Encate (Paul Maney)

W/Pentaerythritol tetranitrate See: Nitralox, Tab (Smith Dorsey)

W/Proloveratrine A & B See: Rauprole, Tab (Vale)

W/Rutin, veratrum viride & mannitol hexanitrate See: Raufia Encate (Paul Maney)

W/Theominal & luminal See: Theominal R S, Tab (Winthrop Labs)

W/Veratrum viride
See: Rauvera, Tab (Smith Dorsey)
Rau Vertin Tab (National Drug)
Rauwloid W/Veriloid, Tab (Riker)

* **RAUWOLFIA SERPENTINA, NF (WHOLE ROOT)**, NND 1962
See: Ekans, Tab (Haag)

Hyperloid Tab (Perron & Covey)
Raudixin Tab (Squibb)

Raupena, Tab (Croches Barnes)
Rauserfia Tab (New England Phr)

Raulina (Tertagar)
Rautotal Tab (Columbur)

Rauval, Tab (Vale Chemical)
Rauwicon, Tab (Consol Midland)

Rauwistan Tab (Thompson)
Rauwoldin Tab (G F Harvey)
Verubar, Tab (Ulmer)

Walfina, Tab (Lloyd & Dabney)

W/Aspirin aluminum hydroxide gel dried ascorbic acid See: Rauylate, Cap (Flint Rotan)

W/Benzhydroflumethazide See: Rautrax N Tab (Squibb)

W/Flumethiazide pot chlor de See: Rautrax Tab (Squibb)

W/Mannitol hexanitrate, rutin See: Maxitate W/Rauwolfia, Tab (Strorenburgh)

W/Methyltestosterone See: Andro Walfia (Ortega)
W/Proloveratrine A & B & phenoxybenzamine HCl (Dibenzylina) See: Mio-Pressin Cap (Smith, Kline & French)
W/Veratrum viride alkaloids
See: Patensor Forte, Improved Tab (Tailby Nasan)
Vertina, Tab (Elder)
W/Veratrum viride ext, mannitol hea-

- nitrate & homotropine methylbromide. See: Roe-Pariental, Tab. (Crookes-Bornes)
- W/Veratrum viride whole root. See: Verwolfia, Tab. (Thompson)
- RAUWOLFSCINE.** (S. B. Penick) An alkaloid of *Rauwolfia canescens*. Under study.
Use: Tranquilizing agent.
- RAYOCAINE HYDROCHLORIDE,** A.D.R. (Winthrop Labs.) Propoxycaïne HCl
- RAWL-VITE.** (Rawl) Vit. A 10,000 u., D 500 u., B-1 10 mg., B-2 5 mg., C 125 mg., B-6 1 mg., nicotinamide 50 mg., Vit. B-6 1 mg., Cal. pantothenate 5 mg., Vit. E 3.4 mg./Tab. Bot. 30s, 100s.
Use: Orally; multivitamin therapy.
- RAWL WHOLE LIVER VITAMIN B COMPLEX.** (Rawl) Whole liver 500 mg. and amino acids found in the whole liver, B-1 1 mg., B-2 2 mg., niacinamide 5 mg., choline chloride 12 mg., B-6 0.2 mg., cal. pantothenate 0.2 mg., inositol 5 mg., biotin 0.6 mcg., B-12 0.3 mcg./Cap. Bot. 100s, 500s
Use: Deficiency of Vitamin B complex
- RAY-AURETONE.** (Raymer) Antipyrine 5% in glycerin, liq., Bot. w/dropper ¼ R. oz., Box 12s, 1 pl.
Use: Topically; analgesic ear drops.
- RAYBANOL** (Raymer) Sod. monobenzylsuccinate,
W/Aminophylline, ephedrine sulfate & phenobarbital. See: Rayphyllin, Cap. (Raymer).
W/Aminopyrine & caffeine. See: Ray-Pyrine, Cap. (Raymer).
- RAY-BURN OINT.** (Cosmos Drug) Aloe Vera 55% in a cholesterolized petrolatum-lanolin base, Tube 1 oz.
Use: Burns of radiation, chemical, flash, etc.
- RAY-D.** (Neoca) Vits. D 1500 u., B-1 1 mg., B-2 2 mg., iodine 0.1 mg., calcium 375 mg., phosphorus 300 mg./6 tabs. Bot. 100s, 500s.
Use: Dietary supplement.
- RAY-FORMOSIL** (Raymer) Formic acid 10 mg., colloidal silicic acid 4.5 mg./2 cc. Amp. (2 cc.) Box 25s, 50s, 100s.
Use: (I.M. 2 cc. twice weekly) Arthritis & rheumatism.
- RAY-GLUCIRON.** (Raymer) Ferrous gluconate 5 gr./tab. Bot. 100s, 500s, 1000s.
Use: (1 to 2 i.d.) Hematinic.
- RAY-MANCOL** (Raymer) Manganic hydroxide colloidal (0.25% mg.), procaine HCl 0.2%/amp. 1 cc. Box 12s, 25s, 100s.
Use: (I.M. 1 cc. q. 5 to 7 days) Furunculosis, acne, psoriasis, sinusitis & impetigo.
- RAY-MER-DERM.** (Raymer) Zinc oxide, calamine, propyl p-aminobenzoate, benzoin, water-miscible base. Tube 1 oz., Jar 1 lb.
Use: Antipruritic analgesic.
- RAY-MER-LAX.** (Raymer) Phenolphthalein, podophyllin, cascara, aloin, sod. monobenzylsuccinate/tab. Bot. 100s, 500s, 1000s.
Use: (1 daily) Constipation.
- RAYMER OINTMENT.** (Raymer) Propyl p-aminobenzoate, bismuth subnitrate, Peru balsam, zinc oxide, salicylic acid, calomel, resorcin, phenol, calamine, cade oil, Jar 1 lb.
Use: Topical anesthetic dressing.
- RAYMINOTOL NO. 2.** (Raymer) Aminophylline 2½ gr., Pot. iodide 2 gr., phenobarbital ¼ gr./Tab. Bot. 100s, 500s, 1000s.
Use: (1 q. 3 h.) Treatment of congestive heart failure.
- RAYPHYLLIN.** (Raymer) Aminophylline 1½ gr., phenobarbital ½ gr., ephedrine sulfate ½ gr., raybanol 1 gr./cap. Bot. 100s, 1000s.
Use: (1 i.d.) Asthma.
- RAY-PYRINE.** (Raymer) Aminopyrine 2½ gr., caffeine ¼ gr., raybanol 2 gr./Cap. Bot. 100s, 1000s.
Use: (1 m.p.) Analgesic & antipyretic.
- RAY-SAL W/Gentisate sod.** See: Genlorth, Tab. (Raymer)
- RAY-SAL W/Succinate.** (Raymer) Salicylic acid 45%, Iodine buffered 3%, succinic acid 2 gr./Tab. Bot. 100s, 1000s.
Use: Orally (1-2 q.i.d.) analgesic & antipyretic.
- RAY-TIC LAP PACKS.** (Johnson & Johnson) Barium sulfate impregnated gauze packs. Boxes 100s.
- RAYTHESIN.** (Raymer) See: Propyl p-Aminobenzoate.
- RAYTOL.** (Raymer) Ox bile ext. 1½ gr., phenolphthalein ½ gr., Sod. salicylate ½ gr., cascara sagrada ext., menthol ½ gr./Tab. Bot. 100s, 500s, 1000s.
Use: Choleric, laxative.
- RAY-TRI-MIDIS, N.N.R. 1955.** (Raymer) Sulfadiazine 167 mg., sulfamerazine 167 mg., sulfamethazine 167 mg./Tab. or 5 cc. Tab. Bot. 100s, 500s, 1000s. Susp., Bot. 1 pt., 1 gal.
Use: Triple sulfonamide therapy.
- RAY-TROIE IMPROVED.** (Raymer) Veratrum viride 65 mg., nitroglycerin 0.25 mg., Sod. nitrite 30 mg., phenobarbital 15 mg./Cap. Bot. 100s, 1000s.
Use: (1 q. 3-4 h) Hypertension.
- W/Rutin. (Raymer) Rutin 20 mg./Tab. In addition to the preceding formula. Bot. 100s, 500s, 1000s.
- RAY-X SPONGES.** (Parke, Davis) Barium sulfate surgical sponges. Size 3 x 3 (12 ply), 4 x 4 (6 ply), 4 x 8 (12 ply). Box 100s; 4 x 4 (8 ply), Box 200s.
Use: To eliminate exploratory reoperations.

- R. D. DIS-PEN-TABS.** (Rexall) Pot penicillin G 50M u/Tab Bot 100s
Use: Orally; penicillin therapy
- R. D. PENTABS.** (Rexall) Pot penicillin G Tab (100M & 250M u/Tab) Bot 100s
Use: Penicillin therapy.
- REACTROL.** (Tailby Nason) 1 p chloro benzyl 2 pyridylidymethyl benzimidazole HCl, Clemizole HCl 20 mg/Tab Bot 60s
Use: Antiallergenic therapy
- READICILLIN, NND 1959** (Upjohn) Penicillin G Pot In veg oil 300 000 u/5 cc Bot 60 cc
Use: Orally; penicillin therapy
- REALDINE NO. 3** (Massengill) Sod salicylate 120 gr, Cimicifuga 15 gr, gelsemium 8 gr, Pot Iodide 4 gr, methyl salicylate q s/R or Bot 1 pl, 1 gal
Use: Analgesic antipyretic
- REALPHENE.** See Acetarsone, Tab (Abbott)
- REBAC.** (Reid) Bacitracin 500 u/Gm Ointment Tube ½ oz
Use: Treatment of minor burns, cuts and wounds
- RECANESCINE** (Riker Labs) See Deserpidine
- RECORTEX.** (Testogor) Adrenal cortex eq equiv to 100 mcg of hydrocortisone/cc Vial 10 cc, 50 cc
Use: Treatment of adrenal insufficiency
- RECTALAD ENEMA.** (Wampole) Glycerin, Sodium Stearate, Dioctyl Sodium Sulfate succinate and water/5 cc Disp plastic container Also pediatric, 2 cc
Use: Enema
- RECTALAD-AMINOPHYLLINE.** (Wampole) Aminophylline 100 mg/cc Bot 3 cc, 45 cc
Use: Diuretic myocardial stimulant
- RECTALAD-MIGRAINE** (Wampole) Ergotamine tartrate 2 mg, caffeine 25 mg, scopolamine aminoxide 0.4 mg, chloral hydrate 200 mg/2 cc Bot 2 cc units
Use: Aborting migraine attacks
- RECTALAD MINIATURE ENEMA.** (Wampole) Glycerine, potassium stearate, and sodium dioctyl sulfosuccinate Small plastic applicator with rectal sp. 2 cc for children 5 cc for adults
Use: Constipation
- RECTAL MEDICINES.** (Medicone Co) Anesthetin 2 gr, ephedrine HCl 1/16 gr, balsam Peru, oxyquinoline sulfate, bismuth subgallate/Supp Box 12s, 24s
Use: Antiseptic & local anesthetic.
- RECTAL MEDICONE-HC Supp.** (Medicone) Anesthetin, 2 gr; ephedrine HCl 1/16 gr; balsam Peru, oxyquinoline sulfate, bismuth subgallate hydrocortisone acetate, 10 mg/Supp Box 12s.
Use: Severe anorectal inflammation
- RECTAL MEDICONE UNGUENT.** (Medicone) Benzocaine 2%, oxyquinoline sulf 0.5%, ephedrine HCl 1/8%, zinc oxide 10%, peru balsam Oint Tube 1 ½ oz w/applicator
Use: Treatment of hemorrhoids
- RECTAL, NO. 15A.** (Merck Sharp & Dohme) Opium pow ½ gr, belladonna ext ¼ gr/supp Box 20s
Use: Antispasmodic
- RECTAL, NO 16A.** (Merck Sharp & Dohme) Opium pow 1 gr, belladonna ext ¼ gr/supp Box 20s
- RECTAL, NO 35A** (Merck Sharp & Dohme) Stramonium ext 30 mg, lead carb 60 mg, Pot alum 60 mg, creosote 30 mg, sal lead subacetate 0.12 cc/supp Box 20s
Use: Antiseptic & astringent
- RECTALGAN.** (Doh Chem) Benzocaine 45%, phenol 175%, menthol 0.5% ephedrine alk 0.125% in oil Sol, Bot 2 oz, 1 pl
Use: Relief of perineal, rectal & hemorrhoidal pain
- RECTALYT HC.** (Mallin) Hydrocortisone alcohol, 10 mg; sulfauridin (sodium dimethyl acrylyl sulfonilamide) 105 mg; menthol 175 mg; benzocaine, 175 mg in water miscible polymer base to 3 Gm/Rectisert 12 u each containing 3 Gm
Use: Antiseptic and local anesthetic
- RECTOCAINE.** (Kirk) Phenol 1%, Propyl P-aminobenzoate 7%, benzyl alcohol 7% procaine 0.5%, Amp 5 cc, 6s, 25s, 100s Oint water soluble 1 oz, 1 lb Supp, Bot 12s
Use: Rectally; local anesthetic
- RECTODYNE OINTMENT.** (Massengill) Opium pow 2 gr, tannic acid 20 gr, phenol 5 gr, stramonium ext 5 gr, Hamamelis leaf ext 20 gr, balsam Peru 15 gr, benzocaine 3% gr, thymal & ricinoinol q s/oz Tube 1 oz, Jar 1 lb
Use: Astringent & antipruritic.
- RECTODYNE SUPP.** (Massengill) Boric acid 8%, ammonium alum 4%, belladonna ext 1%, opium pow ¾%, benecoline 5%, cocoa butter q s/Supp Box 12s
Use: Astringent & antipruritic
- RECTOID.** (Massengill) Atropine sulfate 0.028%, ephedrine sulfate 1%, benzocaine 3%, phenol 0.37%, camphor 0.68%, menthol 1%, zinc oxide oint q s Tube 1 oz, Jar 1 lb.
Use: Astringent & antipruritic.
- RECTULES.** (Fellows) Chloral hydrate 20 gr/Supp in water soluble base, Pkg 12s
Use: Sedative, hypnotic.

- RED BONE MARROW EXTRACT.** See: Carborol, Vial. (Hermelin)
- REDEXCEL.** (Central) d-Desoxyephedrine 3.5 mg., pentobarbital Sod. $\frac{1}{2}$ gr., methylcellulose 5 gr., hamatrapine methylbromide 1/40 gr./Tab. Bot. 100s, 500s.
Use: (1-2 i.l.d.) Central nervous system stimulant w/sedative.
- REDEXCEL IMPROVED.** (Central) Pentobarbital sodium 8 mg., methylcellulose 0.3 Gm, methamphetamine HCl 5 mg., hamatrapine methylbromide 3 mg./Tab. Bot. 100s, 500s, 1000s
Use: C.N.S. stimulant and sedative
- REDICYTE CAPSULES.** (Merck Sharp & Dohme) Thiamine HCl 2 mg., riboflavin 2 mg., pyridoxine HCl 1 mg., vit. B-12 10 mcg., folic acid 1 mg., ascorbic acid 50 mg., ferrous sulfate 0.3 Gm., liver fraction 2.75 mg., stomach desic. [nonofficial] 75 mg./Cap. Bot. 100s, 1000s.
Use: (1 i.l.d.) Hematinic.
- REDIFACT.** (Merck Sharp & Dohme) Vit. B-12 25 mcg., intrinsic factor concentrate 5 mg./Tab. Vial 36s, 100s.
Use: Nutritional supplement.
- REDIPACT FORTE.** (Merck Sharp & Dohme) Vit. B-12 50 mcg., intrinsic factor concentrate 10 mg./tab. Vit. 36s, 100s. Elisir, Bot. 8 oz.
Use: Nutritional supplement.
- REDILEV SOLUBLE TABLETS.** (Merck Sharp & Dohme) Vit. A 3000 u., D 1000 u., thiamine HCl 3 mg., riboflavin 1.5 mg., pyridoxine HCl 3 mg., niacinamide 15 mg., ascorbic acid 50 mg., vit. B-12 5 mcg./tab. Vial 30s.
Use: (1-2 daily) Vitamin deficiency states.
- REDI-MYCIN.** (Elder) Procaine penicillin G 1.5 million u., Pot. penicillin G 500M u., dihydrostreptomycin sulfate 2.5 Gm./5 cc. Vial 3 dose.
Use: L.M.; gram-positive & gram-negative infections.
- REDI-PEN.** (Elder) Penicillin G Procaine. Vial (300,000 u./cc.) 10 cc.
- REDIPLETE PEDIATRIC SYRUP.** (Merck Sharp & Dohme) Vit. A 3000 u., Vit. D 1000 u., C 50 mg, B₆ 1.0 mg., B-2 1.5 mg, B-1 1.5 mg., Cyanocobalamin 5 mcg., Nicotinamide 10 mg./5 cc. Aerasal cans of 237 cc.
Use: Nutritional Supplement.
- REDISOL.** N.N.D. 1962 (Merck Sharp & Dohme) Vit. B-12. Elisir (5 mcg/5 cc.) Bot. 1 pt., 1 gal. Inj. (30 & 100 mcg./cc.) 10 cc., (1000 mcg./cc.) 5 cc. Tab: (25 mcg., 50 mcg) Bot. 36s, 100s. (100 mcg) Bot. 36s, (250 mcg) Vial 12s.
Use: Orally & parenterally; Vitamin B-12 therapy.
- REDITRIN.** (Merck Sharp & Dohme) Vit. B-12 w/intrinsic factor $\frac{1}{2}$ u., B-12 crystalline 10 mcg., ferrous sulfate 300 mg., Vit. C 50 mg., folic acid 1 mg., Vit. B-1 2 mg., B-2 2 mg., B-6 1 mg., niacinamide 10 mg., liver fraction #2 50 mg, desic. stomach 50 mg./Cap. Bot. 60s, 250s
Use. Nematonic w/intrinsic factor.
- REDITRIN-T.** (Merck Sharp & Dohme) Vit. B-12 w/intrinsic factor concentrate 1 oral v., ferrous sulf. 600 mg., B-12 10 mcg., Vit. C 100 mg., folic acid 2 mg., Vit. B-6 1 mg., molybd. 1.5 mg., copper 1 mg., manganese 2 mg., zinc 1.5 mg., cobalt 0.5 mg./Cap. Bot. 30s, 100s
Use: High potency hematinic with intrinsic factor and minerals
- RED MERCURIC IODIDE.**
Used In: Auralcaine, Liq. (Ethical Pharm)
- REDÖDERLIN.** (Testagar) Döderlein's bacillus (*Lactobacillus vaginalis* and *Lactobacillus crassus*) 200,000/Vials (lyophilized) 5 cc Pkg. 6s.
Use: Vaginitis therapy
- REDONA.** (Lemman) Beterpine 0.1 mg., hyoscyamine sulfate 0.1 mg., atropine sulfate /5 cc. or Tab. Bot. 3 fl. oz., 1 pt., 1 gal., Bot. 100s
Use: Antispasmodic—sedative
- REGITINE.** N.N.D. 1962. (Ciba) Phentolamine methanesulfonate & HCl. Nhydrchloridol Tab. (50 mg) Bot. 100s. Methanesulfonate: Vial (5 mg.) 1 cc. Box 1s, 6s.
Use: Diagnostic test for pheochromocytoma.
- REGNOSONE.** (Winthrop Labs.) Pregnenolone acetate. Tab. (100 mg) Bot. 24s.
Use: Rheumatoid arthritis.
- REGONOL.** A vegetable mucin. Cyamopsis tetragonoloba gum.
W/Aluminum hydroxide, Mag. trisilicate, Cal. carb., Mag. carb. & egraine. See: Trenidal, Tab. (Organon).
- REGUTOL.** (Pharmeca) Diacetyl sod. sulfasuccinate 100 mg/Tab. Bot. 30s, 100s.
Use: Stool softener.
- RENTOM.** (Superior Pharm.) Vit. A acetate 2000 u., D 250 u., K 0.5 mg., B-12 2 mcg., folic acid 1 mg., thiamine HCl 2 mg., riboflavin 1.5 mg., niacinamide 10 mg., pyridoxine HCl 0.1 mg., ascorbic acid 25 mg., calcium pantothenate 1 mg., E acetate $\frac{1}{2}$ LU., oyster shell-calcium carbonate, ferrous sulfate anhydrous, cobalt, copper, iodine, manganese, magnesium, molybdenum, potassium, zinc, methionine inositol.

choline bitartrate/Empule Bat 100s, 500s
 Use Prenatal nutritional supplement
RELA. (Schering) NND 1962 Carisapro
 del 350 mg /Tab Bat 30s, 100s
 Use Muscle relaxant and analgesic
RELAGEN (Paul Maney) Neostigmine meth
 ylsulfate 0.25 mg, hyoscyamine HBr 0.25
 mg, benzyl alcohol 3% /cc Vial 10 & 30 cc
 Use: To increase muscle tone
RELANE (Cutter) See Dexoxadrol HCl
RELAXAMINE A-P. (Adams) Solicylamide
 2½ gr, phenacelin 2½ gr, mephenesin
 375 mg, phenobarbital 16 mg, hamatro-
 pine methylbromide 1.5 mg /cap Bat 30s
 100s
 Use: Analgesic & sedative
RELAXANT. (Hart) Mephenesin 7.7 gr /Tab
 Bat 100s, 1000s
 Use: Muscle relaxant
RELAXERPINE. (Adams) Reserpine 0.1 mg,
 mephenesin 400 mg /Tab Bat 30s, 100s
 Use: (1 1 d) Sedative & tranquilizing agent
***RELAXIN,** NND 1962 A purified ovarian
 hormone of pregnancy (obtained from sows)
 responsible for pubic relaxation or separa-
 tion of the symphysis pubis in mammals
 See: Cervixin (Nall)
 Lutrexin (H W D)
 Relaxin (Warner Chilcott)
RELAXIN, PURIFIED. See Cervixin, Vial
 (National Drug)
RELEASIN, NND 1961 (Warner Chilcott)
 Relaxin Amp 1 cc (20 mg /cc) 3000 G P U
 Use: Halt premature labor
RELETUSS. (Scott Alston) D Methorphan HBr
 15 mg, chlorpheniramine maleate 1 mg,
 phenylephrine HCl 5 mg, phenylpropanol
 amine HCl 5 mg, n Acetyl p aminophenol
 120 mg, glyceryl guaiacolate 100 mg /5 cc
 Syrup Bat 4 oz, 1 pl
 Use: Treatment of coughs due to colds
REMANDEN-1DD, TABLETS. (Merck Sharp &
 Dahme) Penicillin G Pot 100,000 u &
 (Benemid) probenecid 0.25 Gm /Tab Vial
 12s, 100s
 Use: Oral "repository" penicillin therapy
REMANDEN-25D, TABLETS. (Merck Sharp &
 Dahme) Penicillin G Pot 250,000 u, Bene-
 mid 0.25 Gm /Tab Bat 12s
 Use: Oral "repository" penicillin therapy
REMANDEN-10D, SUSPENSION. (Merck
 Sharp & Dahme) Pot penicillin G 100,000
 u, probenecid 250 mg, aluminum hydrox-
 ide 250 mg, in peanut oil/5 cc Bat 60 cc
 Use: Treatment of wide range of infections
 due to penicillin susceptible organisms
REMANDEN-25D, SUSPENSION (Merck
 Sharp & Dahme) Pot penicillin G 250,000

u, probenecid 250 mg, aluminum hydrox-
 ide 250 mg, in peanut oil/5 cc Bat 60 cc
 Use See Remanden-10D, Susp (Merck Sharp
 & Dahme)
RENACIDIN. (Guardian Chem) Antilithic
 agent composed of lactones anhydrides
 and acid salts of edible, multivalent or
 organic acids selected from the group: glu-
 conic, citric and malic (Hemiacidrin) Bat
 150 Gm; 300 Gm
 Use Bladder irrigation to aid removal of
 stones composed of calcium and magnesium
 phosphate
RENAL (Squibb) Sodium phytate 3 Gm /
 4.5 Gm Pkg 4.5 Gm packet/90s
 Use Hypercalciuria
RENESE TABLETS, (Pfizer) Polythiazide 1, 2
 or 4 mg /Tab Bat 30s
 Use: Diuretic hypotensive
***RENANOLONE,** 3a Hydroxypregnane 11,
 20 dione
RENESE. (Pfizer) Polythiazide 1 mg 2 mg
 4 mg Bat 100s
 Use Diuretic
RENIR. (Massengill) Reserpine 0.25 mg,
 ephedrine 8 mg /tab Bat 100s
 Use (1 1 d) Hypertension
***RENNIN,** N F, N N R 1935 Milk curdling
 enzyme obtained from the fourth stomach
 of the calf
RENOFORM. See: Epinephrine (Various
 Mfr)
RENOGRAFIN INJECTION, NND 1962
 (Squibb) Methylglucamine diacetylamino-
 iodobenzoate Amp 20 ml w/amp 1 ml of
 the solution to test for sensitivity
 Use: IV, diagnostic agent
RENOLON. Cal acetylsalicylate See Zuma-
 rid, Tab (Stanley)
RENOVIST INJ. (Squibb) Diatrizoate methyl
 glucamine 34.3%, sodium 35%, equivalent
 to 59.7% base which contains 62% iodine
 sol contains 37% firmly bound iodine /
 Vial 25 cc
 Use Radiopaque medium
RENTAMIN (Strasenburg) Pyrimilamine male-
 ate Tab, Bat 100s
 Use: Antihistaminic
REPENDO. (Standex) Hesperidin methyl
 chalcone 100 mg, ascorbic acid 100 mg
 menadione D 1 mg /Cap Bat 100s
 Use: Treatment of capillary wall perme-
 ability
REPLAVIT. (Travenol Lab) Vit B 1 25 mg-
 B 2 10 mg, B 6 20 mg, C 50 mg, B-12
 25 mcg, niacinamide 100 mg, sod panto-
 thenate 20 mg /Vial add 2 to 5 cc water
 Use: IM or IV Infusion

* Generic name recognized by N F U S P or A J A

- REPOSANS.** (Talby-Nason) Not listed.
Phenobarbital $\frac{1}{2}$ gr., hyoscyamus est. $\frac{1}{2}$ gr., valerian, peppermint flower, sumbul R comphar/Tab. Bot. 100s.
Use: Orally (1 m p); sedative.
- RIPOSO-E.** (Canfield) Estradiol valerate 10 mg. in sesame oil/cc. Vial 5 cc.
Use: Ergogenic therapy
- RIPOSO-TE.** (Canfield) Testosterone enanthate 90 mg., estradiol valerate 4 mg. in sesame oil/cc. Vial 5 cc.
Use: Menopausal syndrome
- RIPOSO-TMD.** (Canfield) Testosterone enanthate 100 mg. in sesame oil/cc. Vial 10 cc.
- *RISCIINAMINE.** N.F., N.M.D. 1967.
See: Maderil, Tab. (Pfizer)
Rourescin, Tab. (Columbus)
Tenamine, Tab. (Miss)
- W/Vitamins. See: Neary, Tab. (Blaine Drug)
- RESERBUTAL.** (Vitamins) Butobarbital sodium $\frac{1}{4}$ gr., reserpine 0.1 mg./Tab. Bot. 100s
Use: Hypotensive.
- RESERCEN.** (Central) Reserpine 0.15 mg./tab. Bot. 100s, 500s, 1000s.
Use: Reserpine therapy.
- *RESERPINE, U.S.P., N.M.D. 1967.** In: Tab U.S.P. Pure alkaloid from Rauwolfia serpentina.
See: Anquil, Tabs. (Dymat Wilson)
Banasil, Tab. (Ulmer)
Crystaserpine, Tab. (Smith-Dorsey)
Elserpine, Tab. (Canright)
Ersarerp, Cap. (Smith, Kline & French)
Key Serpine (Key Chr)
Kiline, Tab. (Superior Pharm.)
Lemlerp, Ells., Tab. (Lemman)
Lowserp (Lowe)
Roucap, Caps. (Ethical Pharm.)
Roulen (Paul Money)
Rouline, Tab. (Lloyd Dabney Co.)
Rou-Sed, Tab. (Squibb)
Roulingie, Tab. (Columbus)
Rouwoleaf, Tab. (Gold Leaf)
Rerercen, Tab. (Central)
Reterior, Tab. (Taylor Lab)
Rereroid, N.N.R., Tab. (Upjohn)
Resine, Tab. (Kirkman)
Resipal, Tabs., Elisir (Prama)
Rertran, Tab. (Cole)
Roxinoid, Tab. (Merck Sharp & Dahme)
Sandril, N.N.R. Tab. (Lilly)
Serfn, Tab. (Parke, Davis)
Seralko, Tab. (Ascher)
Serpoid (Canfield)
Serpolan, Tab. (Lanett)
Serporeay, Tab. (Panray)
Serparyl, N.N.R., Tab. (Ciba)
Serpate, Tab. (Vale)
Serpeno, Tab. (Hoag)
- Serpican, Tab. (Conrad, Midland)
Serpiloid, N.N.R. Tab. (Riker)
Serpinte (Vitamins)
Sertabs (Tablet Rock)
Sertina, Tab. (Terogar)
Sessidol, Tab. (Taylor Drugs)
Via-Serpine, Tab. (Rowell)
W/Aminophylline, ephedrine sulfate. See: Serphedrine, Cap. (Norway).
W/Amesium chloride, homatropine methylbromide, caffeine. See: Amiens, Tab (Hoag)
W/Rutabarbital sodium. See: Rererbutal, Tab. (Vitamins)
W/Buical Sod. See: Rutiserpine, Tab. R. Elia. (McNeil)
W/Rutabarbital, Hydrochlorothiazide. See: Rutiserpogide Prertab (McNeil)
W/Choline bitartrate, anhydrous theophylline. See: Theracor, Tab. (C. D. Smith)
W/Cryptenamine tartrate. See: Unitensin-R, Tab. (Irwin, Nelsler)
W/Cryptenamine, methylothiazide. See: Diutensin-R, Tabs. (Irwin, Nelsler)
W/Diethylstilbestrol R methyltestosterone. See: Tylandril, Tab. (Lilly)
W/Druril. See: Dilupres, Tab. (Merck Sharp & Dahme)
W/Ephedrine. See: Renir, Tab. (Messengill)
W/Estrogenic substance & glutamic acid. See: Glutrone, Vial. (Jeffery)
W/Ethylal. See: Ethrosed, Tab. (Chicago Pharmacal)
Estra-Serp. (Omega)
Sergynol, Tab. (Ascher)
W/Hesperidin, ascorbic acid. See: Flavorerp, Cap. (Blue Line)
W/Hexamethonium chloride. See: Hexalina, Tab. (Terogar)
W/Hydroxazine HCl. See: Serparil-Apresoline, Tab. (Ciba).
W/Hydroxazine HCl, Hydrochlorothiazide. See: Ser-Ap-Es, Tab. (Ciba)
W/Hydrochlorothiazide. See: Hydropres, Tab. (Merck Sharp & Dahme) Serparil-Eridrix, Tab. (Ciba)
W/Hydrochlorothiazide, potassium chloride. See: Hydropres-KA-25 or Hydropres-KA-50, Tabs. (Merck Sharp & Dahme)
W/Hydroflumethiazide, protervatrine A. See: Solutensin, Tab. (Bristol)
W/Hyoscyamine sulfate, atropine sulfate R hyoscyne HBr. See: Roudonna, Tab. (Vanpel & Brown)
Redona, Ells., Tab. (Lemman)
Serpodon, Tab. (Walker)
W/Mannitol hexanitrate, theophylline, rutin & ascorbic acid. See: Neo-Semhyten, Cap. (Messengill)

* Generic name recognized by N.F., U.S.P. or A.M.A.

- W/Mannitol hexanitrate, rutin, ascorbic acid
See: Ruhexolaf W/Reserpine, Tab (Lemmon)
- W/Mebaral, Tab
See: Reserpine W/Mebaral (Winthrop Labs)
- W/Mephenesin See: Relaxerpine, Tab (Adams)
- W/Mephenesin & acetylcarbamol See: Amri, Tab (Leeming)
- W/Meratron HCl See: Meratron W/Reserpine, Tab (Merrell)
- W/Methamphetamine HCl
See: Duoria Tab (Ascher)
Resydess, Tab (Chicago Pharm)
Sealress, Tab (Wampale)
- W/Methium See: Methium W/Reserpine, Tab (Warner Chilcott)
- W/Methyl testosterone, dienestrol See: Teserene Tab (Lloyd, Dabney and Westerfield)
- W/Nembutal Calcium See: Nambu Serpin Filmtab (Abbott)
- W/Nidar See: Nitensar, Tab (Armour)
- W/Nitroglycerin, pentoerythritol tetranitrate
See: Penite, Tab (Carrick)
- W/Pentoerythritol tetranitrate
See: Pentaserpine, Tab (Nysco)
- W/Pentoerythritol tetranitrate See: Reserpitate Tab (Vitamax)
Respet, Tab (Lloyd Dabney & Westerfield)
- W/Pentoerythritol tetranitrate, butisal sod
See: Pentrasine, Tab (McNeil)
- W/Pentoerythritol tetranitrate rutin ascorbic acid See: Ruserpatol, Tab (Lemmon)
- W/Pentylenetetrazol, niacin See: Nicoxol W/Reserpine Tab (Drug Specialties)
- W/Phenobarbital, centrine hydrogen sulfate
See: Neuro Centrine, Tab (Bristol)
- W/Phenobarbital, glytheonate, dihydroxy aluminum aminoacetate See: Glytheonate w/Phenobarbital & Reserpine Tab (Patch)
- W/Phenobarbital, Vit B 1
See: Besertal, Tab (Central)
Neo-Slawien, Tab (Patch)
- W/Protaveratrine
See: Veralba R, Tab (Pitman Moore)
Verapene, Tab (Wampale)
- W/Protaveratrine A See: Protalba R, Tab (Pitman-Maare)
- W/Pyrrabutamine See: Sandril W/Pyranil, Tab (Lilly)
- W/Quercetin aminaphylline ascorbic acid
See: Rutaminat RQ Tab (Schenley)
- W/Quercetin & Vit C See: Quer Serpin, Tab (Haskell)
- W/Ritalin See: Serpatin, Tab (Ciba)
- W/Salicylamide See: Salserp, Tab (G W Carrick)
- W/Theobromine Cal See: Purital R S, Tab (A. J. Parker)
- W/Theobromine, phenobarbital See: Theobarb R, Tab (Vanpell & Brown)
- W/Trichlarmethiazide See: Naquival (Schering)
- W/Vit B 12, B 1. See: Samatavite, Tab (Purdue Frederick)
- W/Vitamin B complex See: Rozel, Elix (Merck Sharp & Dohme)
- RESERPINE ELIXIR.** (Bryant) Reserpine 0.25 mg / 5 cc Bot 1 pt
Uses: Sedative & tranquilizing agent
- RESERPINE W/MEBARAL.** (Winthrop Labs) Reserpine 0.15 mg., mebaral 30 mg / tab Bot 100s
Uses: (1) (1 d) Management of anxiety & tension states
- RESERPOID, NND 1962** (Upjohn) Reserpine Tab (0.1 mg., 0.25 mg.) Bot 100s & 500s; (1 mg.) Bot 100s (4 mg.) Bot 100s Elix 0.25 mg / 5 cc Bot 1 pt
Uses: 0.75 to 1 mg daily; hypotensive & tranquilizing agent
- RESERPOTRATE.** (Vitamax) Reserpine 0.1 mg., pentoerythritol tetranitrate 10 mg / Tab
Uses: Angina pectoris
- RESIDERM** (Rowell tab) Quaternary amine type anion exchange resin (base form) 10%, triethanolamine 2.5%, sorbitol, propylene glycol Bot 2 & 6 oz
Uses: Topical Poison Ivy
- RESINAT, NND 1962** (National Drug) Polyamine methylene resin Cap (0.25 Gm) Bot 50s, 100s, 300s, 1000s Tab plain (0.5 Gm) Bot 36s, 100s, 1000s
Uses: Orally; anion exchanger, for gastric antacid & pepsin inactivator
- RESINAT H-M-R.** (National Drug) Polyamine methylene resin 0.5 Gm, hamatrapide methylbromide 1 mg / Tab Bot 36s, 100s
Uses: (1 q 2 3 h) Peptic ulcer
- RESIN CAPRYLATE.** See: Kaprylex, Cap (Strassenburgh)
- RESINE.** (Kirkman) Reserpine 0.1 mg., 0.25 mg., 0.5 mg & 1 mg / tab
Use: Hypotensive & tranquilizing agent
- RESINOL.** (Resinol Chem) Resarcin, cade oil, calamine, zinc oxide, bismuth subnitrate, boric acid Oint 1 1/4 oz., 3 1/2 oz
Uses: Ointment; antiseptic
- RISINS, ANTACID.** See: Polyamine methylene Resins
- RESINS, ION EXCHANGE, NND 1960**
- RESINS, POTASSIUM REMOVING** See: Kalnat, Paw (National Drug)
- RESINS, SODIUM REMOVING.**
See: Carbo Resin, Paw (Lilly)
Kafonsum Paw (Winthrop Labs)
Nafnatil, Paw (National Drug)

- Resodex, Pow.** (Smith, Kline & French)
Vaccid
- RESION.** (National Drug) Polyamine methyl-
one resin 10%, Sod. aluminum silicate
10%, Mag. aluminum silicate 1½%. Bot
4 oz., 12 oz.
- Use: 12 tsp. q. h. for 4 doses, then 2 tsp
q 3-4 h.) Gastro-intestinal cond. rans
- RESION-PM3.** (National Drug) Polyamine-
methylene resin 10%, Sod. aluminum
silicate 10%, Mag. aluminum silicate
1.25%, polymethyl sulfate 125,000 g.,
phthalylacetamide 1 Gm. w/parabens/
15 cc. Bot. 4 oz.
- Use: Diarrheas.
- RESISTAB.** (Bristol-Myers) Theozylamine
HCl. Tab., Bot. 12, 36.
- Use: Antihistamine.
- RESISTOPIN.** (Squibb) Sodium oxacillin 250
mg/cap Bot 4oz
- Use: Staphylococcus infections.
- RESMICON.** (Whittier) Not listed Action ex-
change polyamine resin 500 mg, gastric
mucin 170 mg/Tab Bot 84.
- Use: Antacid.
- RESOCHIN.** See: Chloroquine Phosphate,
U.S.P.
- RESODIC.** (Smith, Kline & French) Polycar-
boxylic cation exchange resin. Pow., 15
Gm., Ptg. 21.
- Use: Sodium-removing resin.
- RES-O-GEL.** (Sanford) Aluminum hydroxide
magnesium carbonate, 375 mg/Tab. Bot.
100s, 1000s.
- Use: Antacid.
- RES-O-PHYLLIN.** (Sanford) Glycerol guai-
acolate 100 mg, methamphetamine hydro-
chloride 1 mg, ephedrine 195 mg,
phenylephrine hydrochloride 12 mg, buta-
barbital sodium 16 mg, and aluminum
hydroxide magnesium carbonate 172 mg./
Cap Bot 100s.
- Use: Treatment of bronchial asthma.
- RESORCIN.** See: Resorcinol.
- *RESORCINOL, U.S.P.; Compound Oint.,
N.F.** (Various Mfrs)
- Use: External; for skin diseases.
- W/Atropine** activated, zinc oxide, col-
loidal sulfur, Isopropylal. See: Sebaborb
lotion. (Summer Lab)
- W/Baric acid,** phenol, fuchsian, acetone &
alcohol.
- See: Corfasin, Eliq. (Rorer).
- Fusinol, Eliq. (Kendall).
- W/Cool** tea distillate & salicylic acid. See:
Tasmel, Oint. (Smith, Kline & French).
- W/Hesachlorophene,** phenol, oil of tar cetyl-
fied, zinc oxide. See: Betafen (Yanpelt &
Brown)
- * Generic name recognized by N.F., U.S.P. or A.M.A.
- W/beta Naphthol,** peru balsam, cellus. See:
A.V.G. Oint. (First Texas)
- W/Pyrimamine** mainate, colomine, zinc oxide,
eucalyptol, phenol, baric acid. See: Solva-
list, Oint. (Columbus).
- W/Sulfur.**
- See: Acromel, Oint. (Smith, Kline & French)
Appliderm #4. (White)
- W/Sulfur,** zinc oxide, bentonite, iron oxide,
talc & propylene glycol. See: Resulin, Oint.
& Lot (Almay).
- W/Sulfur,** zinc oxide & sepetah "5." See:
Resaltah, Cream (Tallby-Nason).
- *RESORCINOL MONOACETATE, N.F.,
M.N.R. 1850** (Various Mfrs) Resorcin acetate.
See: Euresol, Liq. (Knoll).
- Euresol Pro Capitis, Eliq. (Knoll)
- Uses Resorcinol therapy.
- W/Panthenol & sulfur,** See: S. S. Pantho-
derm Cream (U.S. Vitamin).
- W/Salicylic acid.** See: Resorcitate, Sol. (Al-
may).
- W/Sulfur.** See: Sulforcin, Lot. & Oint. (First
Texas).
- W/Salicylic acid.** See: Resorcitate, Sol. (Al-
may).
- W/Sulfur.** See: Sulforcin, Lot. & Oint. (First
Texas).
- RESORCINOLPHTHALIN SODIUM.** See:
Fluorescein Sodium, U.S.P.
- RESORCITATE.** (Almay) Resorcinol mono-
acetate 1.5%, salicylic acid 1.5% in oil or
non-oily base. Bot. 8 fl. oz.
- Use: Antiseptic skin detergent.
- RESPET.** (Lloyd, Dabney & Westfield)
Pentaerythritol tetranitrate 10 mg., cecarpine
0.1 mg./Tab. Bot 100s.
- Use: Angina pectoris.
- RESPIHALES DICADROM.** (Mersch Shosp &
Dohme) Dexamethasone 21-phosphate, sor-
biton trioleate, fluorochlorohydrocarbon 18
mg, alcohol 2%/12.6 Gm in aerosol con-
tainers.
- Use: Bronchial asthma
- RESPIRATORY VACCINE.** See: Cold Vac-
cines
- RESPIRAZONE.** (Tilden) Sodium bromide 3½
gr., potassium iodide 7 gr., motherwort 45
gr., lobelia 10 gr., Ipecac 6 gr./fl. oz. Bot.
pts., gals.
- Use: Sedative and expectorant.
- RESPI-SED.** (Chicago Pharm.) Codeine sul-
fate 0.5 gr., chloralform 0.75 min., Sod. cit-
rate 24 gr., ammonium chloride 8 gr., men-
thol 0.1 gr., grindella 0.25 gr., cenega 0.25
gr./fl. oz. Bot. 1 oz., 4 oz., 1 pt., 1 gal.
- Use: 1 tsp. q. 3 h.; sedative, expectorant.
- RESPITAL.** (Pisano) Resepine 0.1 mg., 0.25
mg., 1.0 mg., 4.0 mg./Tab. Elisis 0.25 mg./

- Tsp Tab Bot 100s, 1000s Elixir Bot pts
Use Hypertension
- RESPREL**. (Tilden) Phenylpropanolamine HCl 10 mg, glyceryl guaiacolate 50 mg, phenyltoloxamine dihydrogen citrate 6.25 mg, pyriminone maleate 6.25 mg, salts cylamide 25 mg / 5 cs Bot pt, gel
Use: Decongestant, antispasmodic, expectorant antihistaminic action
- RESROGEN** (Parke, Davis) Combination of adenovirus vaccine types 3, 4 and 7, and influenza virus vaccine, absorbed on aluminum phosphate One dose (1 cc) contains the equivalent of 1 cc of each of the two vaccines contained in the formula Vial 5 cc
Use: Immunization of adults against adenovirus, types 3 and 4 and 7, and endemic strains of epidemic influenza
- RESTOPHEN**. (Noyes) Phenobarbital $\frac{1}{2}$ gr Hyoscyamus ext $\frac{1}{2}$ gr, cypripedium, valerian & camphor/Tab Bot 100s, 1000s Also available in half strength.
Use: Orally (1 m p); sedative
- RESTRAN**. (Cale) Reserpine 0.25 mg, 1 mg / Tab Bot 100s, 500s, 1000s
Use: Tranquilizer
- RESTROL**, NND 1962 (Central) Dienestrol Susp (5 mg / rs) Vial 10 sc Tab (0.1, 0.5) Bot 100s, 500s, 1000s
Use: I.M.; estrogen therapy.
- RESULIN**. (Schieffelin) Sulfur, resorcinol, zinc oxide, talcum, iron oxide, glycerin, bentonite, alcohol Oint. Tube $1\frac{1}{2}$ oz, Bot 4 oz Soapt (sulfur, resorcinol & salicylic acid).
Use: Antiseptic skin preparation
- RESULIN-P**. (Schieffelin) Resorcinol monoacetate 1.5%, sulfur 2%, hydrocarbons 0.5%/cream Tube 15 Gm
Use: Acne therapy
- RESULTAH**. (Tollby-Nason) Resorcinol 2%, sulfur 6%, zinc oxide 3%, superfath "5" 5%. Cream, Tube $1\frac{1}{2}$ oz.
Use: Topically; acne vulgaris
- RESYDESS**. (Chicago Pharm) Reserpine 0.1 mg, di-desoxyephedrine 8 mg / Tab Bot 100s.
Use: (1 1 d) Obesity
- RITAFEN**. (VonPelt & Brown) Liq (hexachlorophene, phenol, resorcinol) Bot 2 oz Oint (hexachlorophene 1%, phenol 0.9%, resorcinol 1%, tar oil 0.5%, zinc oxide 4%) Tube 1 oz
Use: Antiseptic.
- RITICULEX**. (Lilly) Vit. B-12 10 mg, C 50 mg., folic acid 0.67 mg., ferrous sulfate 200 mg., liver stomach 400 mg./Pulv. Bot 100s, 500s. Also available without iron.
Use: Orally (2 1, 1 d h) anemias.
- RETICULOGEN**. (Lilly) Vit. B 1 2.5 mg, B 12 10 mcg / 0.5 cc Amp 0.5 cc, 3s, 50s, 100s; 5 cc, 25s Fortified: (Vit B 1 5 mg, B 12 20 mcg, liver ext 20 mcg / cc) Amp 0.5 cc, 3s, 50s, 100s; 5 cc, 25s
Use: I.M.; anemias
- RETICULOSE**. (Chemico) Protein, lipoids, nucleic acid Amp. 2 cc, 25s, 100s.
Use: I.M.; nonspecific protein therapy
- RETROGRAFIN**. (Squibb) Sodium and methylglucamine diatrizoates (Benografin) 30%, neomycin (as sulfate) $2\frac{1}{2}\%$. Vial 50 cc
Use: Contrast medium
- RETROPAQUE**. (Winthrop) Skiodon sodium (brand of methiodal sodium), 20% (weight/volume) with neomycin sulfate 4.2% (equivalent to 2.5% neomycin base, weight/volume) Skiodon contains 52% iodine firmly bound in organic form Vials of 50 cc
Use: Retrograde Pyelography
- REVAC SUPPRETTES**. (Webster Co) Diacetyl sod sulfosuccinate 100 mg in neocera base/Supp Jar 12s
Use: Constipation
- REVAMODEX, 18 AND 6D**. (Rand) d Amphetamine sulf 15 mg, amobarbital 60 mg / Cop Bot 100s, 1000s
Use: Sedative w/CNS stimulant
- REVICAPS**. (Lederle) d Amphetamine sulfate 5 mg, methylcellulose 200 mg, Vit A 1670 u, D 167 u, B 1 1 mg, B 2 1 mg, niacin 20 mg, pantothenic acid 0.34 mg, Vit B 6 0.34 mg, C 20 mg, B 12 $\frac{1}{2}$ mcg, folic acid $\frac{1}{2}$ mg, w/Cof, Iron, Phos, Iodine, fluorine, copper, Pot, Mang, zinc boron, Mg / Cop Bot 100s, 500s
Use: (1 1 d) Obesity
- REVIDEX**. (Rand) d Amphetamine sulf 15 mg / Cop Bot 100s, 1000s
Use: CNS stimulant
- REVOBESSE, 15**. (Rand) d Amphetamine sulf 15 mg, thyroid $1\frac{1}{2}$ gr, atropine sulf 1/180 gr, acetphenolisatin 0.5 gr, ascorbic acid 50 mg, phenobarbital $\frac{1}{4}$ gr / Cop Bot 100s, 1000s
- REVOBEX, 15**. (Rand) d Amphetamine sulf 15 mg, or 30 mg, thyroid 3 gr, atropine sulf 1/180 gr, aloin $\frac{1}{4}$ gr, phenobarbital $\frac{1}{4}$ gr / Cop Bot 100s, 1000s
- REXALL SUPER PLENAMINS** (Bexall) Vit A 8000 u, D 1000 u, B-1 2.5 mg, B-2 2.5 mg, C 50 mg, niacinamide 20 mg, B-6 0.03 mg, B-12 3 mcg, folic acid 0.2 mg, E 1 mg, panthanol 3 mg., liver conc. 100 mg., w/minerals/Tab Bot 100s
Use: Multivitamin therapy
- REXAURUM**. (Kohlenberg) Colloidal gold

- 0.1% sol. Bot. 4 fl. oz. Amp. (2 cc.), Box 12s.
Use: I.V., I.M., gold therapy.
- REXIN.** (Dumas-Wilson) Tab. Bot. 100s, 1000s.
Use: Dysmenorrhea.
- REZAMID.** (Dermik) N-Sulfanilylacetonamide 5%, sulfur colloid 3%, salicylic acid 1% / Susp. plas. Bot. 4 oz.
Use: Acne.
- REZAMID CREAM.** (Dermik) N-Sulfanilylacetonamide 8.5%, salicylic acid 1%, sulfur colloid 3%, 2,2' methylenebis 0.5% in a greasable vegetable oil vehicle/cream. Tube: 1 oz.
Use: Dermatologic.
- REZIFILM.** (Squibb) Methacrylate resin w/ tetramethylthiuram disulfide in ethyl acetate and propellants. Dispenser aerosol can 2 oz., 6 oz.
Use: Surgical spray dressing.
- REZIPAS.** N.N.D. 1962. (Squibb) PAS resin, p-aminosalicylic acid 30% adsorbed on an anionic exchange resin. Pkg. 1 lb; 8 Gm. pk. 100s.
Use: (24 to 30 Gm. daily) Tuberculosis.
- "RQ" LECITHIN WAFERS.** (Glidden Co.) Choline 145 mg., inositol 145 mg., 17% (130 mg) of the minimum adult daily requirement of phosphorus/3 Wafers. Box 60 wafers.
Use: Fat metabolism.
- "RQ" SOYA LECITHIN.** (Glidden Co.) lecithin 29.5%, cephalin 29.5% & plant products. Yellow granules, Bot. 8 oz.
Use: Lipid metabolism & supply phospholipids.
- R-GENE.** (Cutter) L-Argine HCl 5% Solution. Flask 500 cc. w/inj. equipment.
Use: Hepatic coma.
- RHAMNOTIN.** (Strassenburgh) Ascorbic acid 75 mg., rutin 60 mg./Tab. Bot. 100s. Also available w/ascorbic acid 300 mg./Tab. Bot. 100s.
Use: (1 i.i.d.) Capillary permeability.
- RHEASTAT.** (VonPelt & Brown) Phenobarbital 16.2 mg., hyoscyamine sulf. 0.1286 mg., atropine sulf. 0.025 mg., hyoscin HBr 0.0074 mg., kaolin 6 Gm., pectin 320 mg., sod. chloride 6 mEq., pot. chloride 4 mEq. w/aluminum hydroxide gel/fl. oz. Susp. Bot. 1 pt.
Use: Treatment and management of diarrhea
- RHEUMATISM PHYLLACOGEN.** (Parke, Davis) Strep. 75%, antigen isolated from rheumatic infections & Staph. albus & Aureus 25%/cc. Vial 10 cc.
Use: S.C. (1 cc.); arthritis.
- RHEUMATOID ARTHRITIS VACCINE.** (Bed-
wile) Hemolytic streptococci 100 or 1000 million/cc. Vial 5 cc.
Use: I.M. (50 million), I.V. (5 million); arthritis.
- RHINALGAN HC.** (Doho) Hydrocortisone alcohol 0.02%, di-desoxyephedrine hydrochloride 0.22%, pyrilamine maleate 0.01%, anilpyrine 0.28%, gentian violet.
Use: Nasal decongestive.
- RHINAZINE.** (Lederle) Sulladiazine 1%, sulfathiazole 0.1%, desoxyephedrine, racemic, 0.1%. Bot. 1 oz., 1 pt., 1 gal.
Use: Nasal vasoconstrictor w/sulfa therapy.
- RHINIDEIM PALATAB.** (Central) Solicylamide 60 mg., belladonna tr. 0.015 ml, r-ephedrine HCl 5 mg /Palatob. Bot. 100s, 500s, 1000s
Use: Rhinitis in children
- RHINIDRIN TABLET.** (Central) Solicylamide 5 gr., belladonna alkaloids equivalent to 0.06 ml. belladonna tincture, r-ephedrine HCl 20 mg./Tab. Bot. 100s, 500s, 1000s.
- RHINITIS FULL STRENGTH C.C.T.** (Merck Sharp & Dahme) Belladonna leaf fl. ext. 0.015 cc, quinine sulfate 30 mg., camphor 30 mg./tab. Bot. 100s, 1000s.
- RHINITIS HALF STRENGTH TABLETS.** (Parke, Davis) Camphor 1/4 gr., quinine sulfate 1/4 gr., fl. ext. belladonna root 1/2 min./Tab. Bot. 100s.
- RHINORAL.** (Rhinopto) Phenylephrine HCl 3.5 mg., phenylpropanolamine HCl 6.5 mg., chlorpheniramine maleate 1.25 mg., oral-aminophen 120 mg /5 cc. Bot. pts.
Use: Nasal congestion associated with colds.
- RHITIC.** (Lamson) Chlorphenamine 1.25 mg., phenylephrine HCl 2.5 mg., neomycin 3.5 mg., polymyxin B sulfate 2000 u., benzocaine 5 mg./tox. Bot. 100s.
Use: Upper respiratory congestion, post-nasal drip, sore throat.
- RHODANATE.** See: Potassium Thiocyanate.
- RHODANIDE.** More commonly rhodanate, same as thiocyanate (CNS). See: Potassium thiocyanate.
- RHOPEX.** (Vitarine) Vit. B-12 29.3 mcg., B-1 20 mg., B-2 3 mg., B-6 1 mg., pantothenic acid 3 mg., niacinamide 100 mg., choline chloride 20 mg., inositol 10 mg., liver inj. equiv. to Vit. B-12 30 mcg., procaine HCl 1%/cc. Vial 5 cc., 10 cc.
Use: (1 cc. daily) Hematopoietic.
- RHUEARB ALKALINE EXTRACT.** See: Neutralizing (Merck Sharp & Dahme)
- RHUBARB & PANCREATIN COMPOUND, ALKALINE.** (Merck Sharp & Dahme) Ehubarb 1.2 Gm., pancreatin 0.25 Gm., hydrastis 0.53 Gm., pot. bicarb. 1.2 Gm., alcohol 18%/30 cc. Bot. 1 gal.

RHUBARB & SODA MIXTURE, NF. (Merck Sharp & Dohme) Rhubarb fl ext 15 cc, ipecac fl ext 0.3 cc, sod bicarb 3.5 Gm spirit peppermint 3.5 cc, alcohol 3%/100 cc) Bot 1 pt, 1 gal

RHUCEN. (Central) Zirconium oxide 4%, pyrilamine moleate 1%, p chloro m xylenol 0.8% benzalkonium chloride 0.01% Oint Tube 1 oz
Use: Rhus dermatitis

RHULICREAM. (Lederle) Benzocaine 1%, zirconium oxide 1%, menthol 0.5%, camphor 0.3%, w/paraben & Isopropyl alcohol Oint 1 oz
Use Topically; local anesthetic

RHULIFOAM (Lederle) Zirconium Oxide 0.90% Colamine 2.8% Benzocaine 0.94% Menthol 0.11%, Camphor 0.11% 2 oz pressure pack can
Use Relieve minor itching and pain due to ivy and oak poisoning as well as hives, insect bites, and other minor skin irritations.

RHULI-LOTION (Lederle) Benzocaine 1%, zirconium oxide 1%, menthol 0.1%, calamine 3%, camphor 0.1%, methylparaben and propylparaben as preservatives Bot 4 oz
Use: Relief of itching of ivy and oak poisoning

RHULISPRAY. (Lederle) Zirconium oxide 1%, benzocaine, menthol, camphor, calamine & Isopropyl alcohol Aerosol 3 oz
Use: Topically analgesic, anesthetic for treatment of minor skin irritations

RHULITOL (Lederle) Tannic acid 3%, phenol 0.4%, chlorobutanol 1%, camphor, ammonium alum Isopropyl alcohol Bot 4 oz
Use: Poison ivy oak & sumac

RHUS-ALL ANTIGIH (Barry) Ivy, oak & sumac ext In oil Vial 1 cc, 4s
Use: Treatment & desensitization in Rhus dermatitis

RHUS IN OIL SOLUTION. See: Ivyoak, Vial (Sherman).

RHUS TOX ANTIDEN. (National Drug) A hydro-alcoholic solution containing the antigenic principles of rhus toxicodendron [poison ivy, poison oak] Vial 1 ml
Use: I.M., treatment of poison ivy or poison oak

RHUS TOXIN TINCTURE. (Mullford) Poison ivy ext in alcohol Bot 1 oz.
Use: Orally; Rhus dermatitis

RHUS TOX TINCTURE. (National Drug) Bot 1 fl oz
Use: Oral use in the prophylaxis of rhus dermatitis

RHUTOX LOTION (Blue Line) lead acetate
* Gener c name recognized by NF, USP or A.M.A.

grindelia, echinacea in alcohol Bot 4 oz, 1 pt, 1 gal

Use Antiseptic skin preparation
RNYTYMOL. (Roche) 2,6 bis(1 piperidyl methyl 4 (alpha di methylbenzyl)phenol d hy drabromide

Use Treatment of cardiac arrhythmias
RIAMINE. (Kirk) Vit B 1 50 mg, pyridoxin 50 mg/cc Vial 10 cc

RIASOL (Shield) An oily emulsified liq containing 0.45% mercury chemically combined w/soaps, 0.5% phenol & 0.75% cresol Bot 4 & 8 fl oz
Use Topically; psoriasis

RIATRIN (Ascher) Estrane 0.5 mg, testosterone 10 mg, B 1 50 mg B 2 2 mg nicotinamide 100 mg, B 6 5 mg, panthanol 10 mg, cyanocobalamin 100 mcg, liver In jection equiv in B 12 activity to cyanocobalamin 2 mcg, choline chloride 25 mg, inositol 25 mg, dl Methionine 25 mg, procaine HCl 1%, benzyl alcohol 1.5%/ml Susp Vials 10 ml
Use: Gynecitics

RIBEX. (Walker Vitamin) Thiamine HCl 1.5 mg, riboflavin 2 mg, pyridoxine 0.3 mg, Cal pantothenate 1 mg, niacinamide 10 mg/Tab Bot 30s, 100s
Use: Dietary supplement

RIBODERM. (Meyer Chem) Riboflavin 10 mg/cc Vial 5 cc 10 cc
Use: Psoriasis

***RIBOFLAVIN, USP** NND 1962, ADRJ Inj. USP; Tab, USP (Vit B 2, Vit G yellow enzyme, lactoflavin)
See: Flomatide, amp (Philadelphia Amp) Flavoxin Amp & Tab (Winthrop Lab) Hyflavin, Amp (Endo) Hyrye, Tab & Vial (Durst) Riboderm, Vial (Meyer) W/Nicotinamide

See: Niro, Vial (Barry) Ribotinic, Amp (Abbott)

W/Nicotinamide (Lilly) Riboflavin 5 mg; nicotinamide 200 mg/cc or Tab Amp 1 cc, Box 6s 25s 100s Tab Bot 100s, 500s
Use: I.M., I.V.; Vitamin B therapy

W/Vitamins, See: Vitamin Preparations
RIBOTLAVIN-5-MONOPHOSPHATE SODIUM See: Dermaject (Columbus Pharm)

RIBOFLAVIN, METHYLOL NND 1962
See: Hyflavin Amp, Vial (Endo)

RIBOFLAVIN-5-PHOSPHATE SODIUM
See: Flomatide, Amp (Philadelphia Amp) Hyrye, Tab & Vial (Durst)

RIBO-PLEX. (Warren Teed) Vit B 1 10 mg- B 2 3 mg- B-6 5 mg, & nicotinamide 20 mg/10 cc Vial 10 cc

- RIBOTHIRON ELIXIR.** (Merck Sharp & Dahme) Ferrous sulfate 1.2 Gm., thiamin HCl 2 mg., riboflavin 2 mg./30 cc. Bot. 1 pt.
Use: Hypochromic anemia.
- RIBOTHIRON TABLETS.** (Merck Sharp & Dahme) Ferrous sulfate exsicc. 0.2 Gm., thiamine HCl 1 mg., riboflavin 1 mg./tab. Bot. 100s, 1000s.
Use: Hypochromic anemia.
- RIBOTHYN.** (Flint, Eaton) Vit. B-1 10 mg., B-2 5 mg., B-6 5 mg., Cal. pantothenate 10 mg., nicotinamide 100 mg./cc. Vial 10 cc.
Use: I.V., I.M., Vitamin B complex.
- RIBOTINIC SOLUTION.** (Abbott) Riboflavin 5 mg., nicotinamide 200 mg., in water for inj./cc. Amp. 1 cc., 6s, 100s.
- RIBRANEX.** (Wyeth) Vit. B-1 0.6 mg., B-2 1.3 mg., B-6 0.3 mg., niacinamide 9.4 mg., pantothenic acid 0.4 mg. & rice/5 cc. Elixir, Bot. B oz.
Use: Vitamin B complex therapy.
- RICE BRAN EXT. CONCENTRATE.** (Flint, Eaton) Thiamine HCl 1.5 mg., riboflavin 1 mg., nicotinic acid 10 mg., pyridoxine HCl 0.75 mg., pantothenic acid 1 mg., w/ choline, inositol & biotin/dr. Bot. 4 fl. oz., 1 pt., 1 gal.
Use: Vitamin B complex therapy.
- RICINOLEATE, CALCIUM.** See: Pihatoz. (Suttlin & Case)
- RICINOLEATE, ETHYL W/Mineral oil, mag hydrox., koalin.** See: Karidin, Liq. (Merrell)
- RICINOLEATE SODIUM, N.N.R. 1954.**
See: Sarlein, Prep. (Merrell)
Ricinoleic acid prep. (See: Aci-Jel. (Ortho))
- RICINOLEIC ACID W/Bile Salts.** See: Byferic. (Merrell)
- RID-OBESSE T.D.** (American Pharm. Co) Amphetamine sulf. 15 mg., thyroid 195 mg., atropine sulf. 0.34 mg., aloin 16 mg & phenobarb. 16 mg/Cap. Bot. 1000s
Use: (8 hr. duration) Control of overweight
- RIDUPOIS.** (Elder) Thyroid 1/5 gr., dl-methamphetamine HCl 2.5 mg., Vit. B-1 1 mg., B-2 0.5 mg., sod. carboxymethyl cellulose 4 gr./Tab. Bot. 100s, 500s
Use: Anobesity
- RIFOMYCIN.** (Registered in Great Britain) Antibiotics isolated from a strain of *Streptomyces mediterranei* (specific substances are designated by a terminal letter thus, Rifomycin B).
RIFOMYCIN B. Under study.
Use: Antibiotic.
- RIMIFON, N.N.D. 1962.** (Roche) Isoniazid. Tab. [50 mg] Pkg. 100s, 1000s (100 mg) Bot. 100s, 1000s.
Use: Orally; tuberculosis.
* Generic name recognized by N.F., U.S.P. or A.M.A.
- RINOFEDS.** (Broemmel) Cap. 24s, 100s, 500s; Cap. 1/2 strength 24s, 100s; Syr. 4 oz., pt., gal.
- RINOHIST.** (Broemmel) Cap. 24s, 100s, 500s; Syr. 4 oz., pt., gal; Cap. 1/2 strength 24s, 100s
- RIODINE.** (Gallia) Iodinated castor oil. Cap. (0.2 Gm. equiv. to 1/2 gr) Iodine, Bot. 50s, 300s
Use: Orally; iodine therapy.
- RIOGON, N.N.R. 1953.** (Brean) Chorionic gonadotropin, Vial (5000 u. & 10,000 u.) 10 cc.
Use: Cryptorchism.
- RIONA.** (Merck Sharp & Dahme) Propadrine HCl 50 mg., phenacetin 0.12 Gm., aspirin 0.2 Gm./cap. Bot. 30s, 100s, 1000s.
Use: (1-3 daily) Central nervous system stimulant w/analgesic.
- RIOPAN.** (Ayers) Monalium Hydrate. Hydrated magnesium aluminate 400 mg/Tab. or 5 cc. Tab. Bot. 60s, 500s Susp. 12 oz.
Use: Antacid & G.I. disturbances.
- RI-PLEX.** (Warren-Teed) Thiamine HCl 18 mg., riboflavin 6 mg., pantothenic acid 4.5 mg., Vit. B-6 18 mg., nicotinamide 60 mg. & rice bran/fl. oz. Elixir, Bot. 1 pt., 1 gal. Also Vials 10 cc.
Use: Vitamin B complex therapy.
- RISA, N.N.D. 1962.** (Abbott) Radioactive iodinated blood albumin. Vial 1 millicurie.
Use: I.V., diagnostic aid.
- *RISTOCETIN, N.N.D. 1962.** An antibiotic from species of actinomycetes, *Nocardia lurida*. See: Spontin, Vial (Abbott)
- RITALIN HYDROCHLORIDE, N.N.D. 1962.** (Ciba) Methylphenidylacetate HCl (Methylphenidate) Tab. 5 mg., 10 mg., Bot. 100s, 500s, 1000s. Tab. 20 mg., Sol. 100s, 1000s (inj) 100 mg) Vial 10 ml.
Use: Psychomotor stimulant.
W/Methyltestosterone, ethinyl estradiol, B-1, B-2, B-6, B-12, nicotinamide, dicalcium phosphate. See: Ritonic, Cap. (Ciba)
W/Pyribenzamine HCl. See: Plimastin, Tab. (Ciba)
W/Reserpine. See: Serpatilin, Tab. (Ciba)
- RITONIC.** (Ciba) Ritalin 5 mg., methyltestosterone 1.25 mg., ethinyl estradiol 5 mcg w/ Vit. B-1, B-2, B-6, B-12 nicotinamide and dical. phos. Cap. Bot. 100s.
Use: Stimulant and tonic.
- RITH-A-TAL.** (Walker Corp) Pentaerythritol tetranitrate 10 mg., mephobarbital 1/2 gr., phenobarbital 1/2 gr./Tab. Bot. 500s, 1000s
Use: Treatment of angina pectoris
- R-M COLE.** (Cole) Salicylamide 5 gr., col-

- choline 1/500 gr, cimetidine 1/2 gr, physalocin 1/4 gr/Cap Bot 100s
 Use: Gastric therapy
- ROBALATE**, N N O 1962 (Robins) Dihydroxy aluminum aminoacetate 7 1/2 gr/Tab Bot 100s, 500s Liq (7 1/2 gr/5 cc) Bot 12 oz
 Use: (1 2 p c & h s) Hyperacidity
- ROBANUL**, See Robinul
- ROBARIN-750** (Robins) Methocarbamol 750 mg Tabs Bot 100s 500s
 Use: Skeletal muscle relaxant
- ROBAXIN**, N N O 1962 (Robins) Methocarbamol 3 (o Methoxyphenoxy) 2 hydroxypropyl carbamate (Glyceryl guaicolate carbamate) Tab. 500 mg, Bot 50s, 250s In[—] 1 Gm/10 cc of a 50% aqueous solution of polyethylene glycol 300 Amp 10 cc
 Use: Skeletal muscle relaxant
- ROBAXIN-750** (Robins) Methocarbamol 750 mg/Tab Bot 50s 500s
 Use: Acute skeletal muscle hyperactivity
- ROBAXISAL** (Robins) Methocarbamol (Robaxin) 400 mg, Acetylsalicylic acid 325 mg/Tab Bot 100s, 500s
 Use: Muscle Relaxant/Analgesic
- ROBAXISAL-PH**, (Robins) Methocarbamol 400 mg, phenacetin 97 mg, acetylsalicylic acid 81 mg, hyoscyamine sulfate 0.016 mg, phenobarbital 81 mg/Tab Bot 100s, 500s
 Use: Analgesic
- RO-BILE** (Rowell) Pepsin 260 mg, pancreatin 260 mg, ox bile 100 mg, dehydrocholic acid 30 mg/Tab Bot 50s
 Use: Digestant
- ROBINUL**, (Robins) *Glycopyrrolate 1 mg/Tab Bot 100s 500s
 Use: Anticholinergic
- ROBINUL-PN** (Robins) Glycopyrrolate 1 mg phenobarbital 1/4 gr/Tab Bot 100s 500s
 Use: Duodenal and gastric ulcer
- ROBITUSSIN**, (Robins) Glyceryl guaicolate 100 mg/5 cc Syr, Bot 4 oz, 1 pt, 1 gal
 Use: Orally, antitussive & expectorant
- ROBITUSSIN-AC**, (Robins) Glyceryl guaicolate 100 mg, pheniramine maleate 7.5 mg codeine phosphate 10 mg/5 cc Bot 4 oz, 1 pt, 1 gal
 Use: Expectorant
- ROBUDEN** (Crookes Baines) a tissue extract Amps & Tabs 200 mg
 Use: Gastric and duodenal ulcer
- ROCCAL** (Winthrop Labs) Benzalkonium chloride technical Sol 10%, Bot 1 pt, 1 qt, 1 gal
 Use: Cationic germicide
- RO-CILLIN**, (Rowell) Potassium Phenethicillin
 * Generic name recognized by N F, U S P or A.M.A.
- 250 mg/Tab and powder for oral solution providing 125 mg./5 cc Bot 24s, 100s
 Oral solution in 60 cc Bot
 Use: Treatment of various infections caused by penicillin sensitive organisms
- ROCKY MOUNTAIN SPOTTED FEVER ANTIGEN**, (Lederle) Vial 5 cc., 15 tests
- ROCKY MOUNTAIN SPOTTED FEVER VACCINE**, N N R 1932
 Lederle—Vial 3 cc, 20 cc
 Merck Sharp & Dohme—Vial 15 cc
- ROCYTE TABS. AND INJ**, (Sig) Tabs: Ferric fumarate 5 gr, Vit C 100 mg, copper sulfate 2 mg, desiccated liver 15 mg, cobalt chloride 10 mg, vit B 12 25 mcg/tab Bot 100s, 1000s Inj: Vit B 12 200 mcg, beef liver NF 10 u, ferric iron combined w/enzymically digested protein 15 mg, cobalt gluconate 5 mg, sodium citrate 10 mg, benzyl alcohol/cc 10 cc vials
 Use: Anemias
- RO-DERM** (Lemmon) Neomycin sulfate 0.35%, dipradon HCl 1%, allantoin 0.30% Sol roll on applicators 5 cc
 Use: Skin irritations
- RO-DEX**, (Rowell) Vit A 5000 u, D 1200 u, B 1 3 mg, B 2 3 mg, B 6 1 mg, C 100 mg, niacinamide 20 mg, cal pantothenate 2 mg, E 1 mg, d amphetamine HCl 2.5 mg phenobarbital 1/4 gr/Cap Bot 100s, 500s
 Use: Treatment of obese children
- ROENTEN**, (Bayer) Cassia angustifolia senna 1 oz can
 Use: Evacuant of bowels before x ray
- ROENTENOGRAPHY**, See: Iodine Products Diagnostic
- ROETINIC**, (Roerig) Vit B 12 w/intrinsic factor 1 u, folic acid 2 mg, Vit C 100 mg ferrous sulfate exsic 400 mg, molybdenum copper, cobalt, Mang, zinc, each D 5 mg/Cap Bot 30s, 100s.
 Use: Hematinic agent
- ***ROFLURANE** (Dow Chem Co) 2 bromo 1,1 2 trifluoroethyl methyl ether
- ROGITINE**, See: Phentalamine
- ROLAIDS**, (American Chicle Co) Dihydroxy aluminum Sodium carb Tab, Roll 10s
 Use: Antacid
- ROLICTON**, N N D 1961 (Searle) Amlinetradine Aminoisometradine 1 Methylallyl 3 methyl 6 aminotetrahydropyrimidinedione Tab 400 mg Bot 100s, 500s
 Use: Diuretic
- ROLIEF**, (Merrell) Zirconium oxide diethane ceepryn and menthal Lot Bot 1 oz
 Use: Poison ivy, Insect bites and skin irritations
- ROLITETRACYCLINE**, N [pyrrolidino methyl tetracycline]

See: Synthes (Bristol Lab)
 Yelocryline (Sqs/hb)

ROMEX TROCHES & LIQUID. [American Pharmaceutical Co] Polymyxin B sulfate 1000 u., benzocaine 5 mg., tetracycline chloride 2.5 mg., gramicidin 100 mcg., chlorpheniramine maleate 0.5 mg., pyrazinole 2 mg./Troche, 2 LIQUID Glyceryl glycoacetate 200 mg., dextromethorphan hydrobromide 60 mg., chlorpheniramine maleate 12 mg., phenylephrine hydrochloride 30 mg./oz
 Use: Minor throat and mouth irritations.

ROMICIL. See: Olanedemycin

ROMILAR, NHO 1962 (Roche) d-Methorphan N-methylmorphine HBr, d-Methorphan HBr, Tab. [15 mg] Sol. 20%, 100%, 500%. Syr. [15 mg/5 cc] Sol. d. oz., 16 oz. 8 gal.
 Use: Cough therapy.
 W/Homotropic methylbramide, pyrilamine maleate, ammonium Cl, sodium citrate. See: Endostin 104, Syr. [Tada]

ROMILAB-CP. (Roche) d-Methorphan HBr 15 mg., chlorpheniramine maleate 1.25 mg., phenylephrine HCl 5 mg., N-acetyl-*p*-aminophenol 120 mg/5 cc. Syr., Sol. 3 oz., 1 pt., 1 gal.
 Use: Treatment of the common cold

ROMILAR EXPECTORANT. (Roche) d-Methorphan HBr 15 mg., Amm. chloride 90 mg w/sod. benzoate/5 cc. Sol. 8 pt., 8 gal.
 Use: [1-2 dr. 11 d] Expectorant.

RONCOVITE. (Lloyd) Cobalt chloride 15 mg. & ferrous sulfate 0.28 Gm./Tab. or 5 cc. Drops [15 cc.] Sol. w/dropper. Sol. 1 pt., 1 gal. Tab., Sol. 100%, 500%, 1000%.
 Use: Hypochromic anemia.

W/Vit. B-12. (Lloyd) Sol. 100%, 500%, 1000%.

RONCOVITE-MP. (Lloyd Bros) Cobalt chloride 15 mg., ferrous sulf. sat. 100 mg./Tab. Sol. 100%
 Use: Treatment of anemias

RONCOVITE-OB, TAB. (Lloyd Bros) Cobalt chloride 15 mg., ferrous sulfate 0.2 Gm., col. lactate 0.9 Gm., Vit. O 250 u/Tab. Sol. 100%
 Use: 1 cc., h; anemia of pregnancy.

RONIACOL (Roche) beta Pyridyl carbinol. (Picolinyl alcohol) Ellals (50 mg/5 cc.) Sol. 1 pt., 1 gal. Tab. (50 mg) Sol. 100%, 500%, 1000%. Timespan 150 mg. Sol. 50%
 Use: (1 tsp. or Tab. 11 d) Vasodilator.

W/Aminophylline (Roche) Banlacol 50 mg., aminophylline 100 mg. 8 Mag. trisilicate/Tab. Sol. 100%, 500%
 Use: Peripheral vasodilator.

W/Aminophylline 8 Mag. trisilicate. See: Banlacol W/Aminophylline, Tab. (Roche).

W/Tigon. See: Tigacol, Caps. (Roche)

RONIACOL TIMESPAN TABS. (Roche) Beta-pyridyl carbinol, 150 mg/Tab. Sol. 50%.
 Use: Impaired peripheral circulation.

RONONE. (Abbott) Ronone, C.P. 1.96% w/w chloroform 4.19% w/w, aqueous mucilage of Irish moss & quince seed 10.85%. Sol. 4 fl. oz., 1 pt.
 Use: Siccicide.

BONUVEX. (Kromers-Urban) Ferrous fumarate 200 mg., copper sulfate 3 mg., thiamine HCl 2 mg., riboflavin 2 mg., niacinamide 10 mg., folic acid 0.25 mg., Vit. B 12 O 50 mcg., liver fraction 2, 100 mg/Tab. Sol. 100%, 500%, 1000%
 Use: Anemias.

BONUVIN. (Kromers Urban) The daily dose of 3 tablets contains ferrous fumarate 400 mg., copper sulfate 15 mg., Vit. B-12 w/ folic acid conc. 1/3 U.S.P. Unit. (oral) Vit. B-12 (activity equivalent) from streptomycin fermentation) 2.5 mcg., folic acid 2 mg., ascorbic acid 100 mg., thiamine mononitrate 5 mg., riboflavin 5 mg., niacinamide 25 mg., molybdenum (from sodium molybdate) 1.5 mg., cobalt (from cobalt sulfate) 0.5 mg., manganese (from manganese sulfate) 0.5 mg., zinc (from zinc sulfate dried) 0.5 mg/Tab. Sol. 100%, 500%, 1000%
 Use: Anemias

RONYNIC. (Robinson) Nisotinic acid 500 mg/Tab.
 Use: Sedation of high cholesterol levels

BOPAD. (Priman-Moore) Dexamethorone 0.075 mg., acetylsalicylic acid 325 mg., aluminum hydroxide 75 mg., ascorbic acid 20 mg/Tab. Sol. 100%
 Use: Analgesic in the treatment of rheumatic and arthritic conditions.

ROSA GALLICA T. See: Estlin, Sol (Schleffelin)

ROSANTINE DYES, MNR, 1954, A D B
 See: Fuchsln.
 Methylrosaniline Chloride. [Various Mfr.]

RO-SULFA. (Rowell) Sulfamethylthiadiazole 250 mg., triethylpyrimidine 250 mg., phenylazo-diaminopyridine HCl 75 mg/Tab. Sol. 100%
 Use: Urinary tract infections.

B-D-SULFA PLAIN. (Rowell) Sulfamethizole 250 mg., Tris Sulfa 250 mg/Tab. Sol. 100%
 Use: Urinary tract infections.

RO-TABS. (Rowell) Vit A 5000 u., D 1000 u., B-12 5 mcg., C 50 mg., thiamine 2 mg., niacinamide 10 mg., d-calcium pantothenate 2 mg., pyridoxine HCl 0.2 mg., riboflavin 2 mg/Chewable Tab. Sol. 100%, 1000%, 5000%
 Use: Dietary supplement.

ROTENONE.

- See: Ronane, Lot (Abbott)
 Rotacide, Lot (Blue line)
- * **ROTHETAMINE** (Smith Kline & French)
 o, o-dimethylphenethylamine
 Use: Anorexic compound
- * **ROTOXAMINE TARTRATE** +{(-)-2 [p-chloro a[2 dimethylaminoethoxy]benzyl] pyridine D tartrate} See Twiston tartrate (McNe I)
- ROVICOTE** (Hoffmann LaRoche) B vitamins (Thiamine riboflavin pyridoxine, niacinamide) coated by an inert edible, water resistant material
 Use: Chewable tablets vitamin supplement
- ROXEL** (Merck Sharp & Dahme) Reserpine 0.1 mg, vit B 1.5 mg, B 6 5 mg, B 12 25 mcg niacinamide 10 mg / 15 cc Bot 1 pt
 Use (1 tablespoonful i d) Treatment of emotional instability, neuroses & mild labile hypertension
- ROXINOID, NND 1959** (Merck Sharp & Dahme) Reserpine Tabs 0.1 mg, 0.25 mg 0.5 mg 1 mg Bot 100s, 1000s
 Use 0.1 to 0.5 mg daily; hypotensive & tranquilizing agent
- R S PANTHODERM CREAM, (U S Vitamin)**
 Panthenol 2% resorcinol monoacetate 2% sulfur 5% Cream Jar 2 oz, 1 lb
 Use: Acne dandruff & dermatitis
- R-5-THESODATE, (Brewer)** Theobromine sodium acetate 7 1/2 gr, Rauwolfia serpentina 50 mg / E C Tab Bot 100s 500s
 Use (1 t d and h s) Treatment of essential hypertension
- RU-AN CEE** (Sull ff & Case) Vitamin C 50 mg rut n 20 mg / Tab Bot 100s, 1000s
 Use: O tory supplement
- RUBAPLEX-C** (Lanett) Vit B 12 100 or 1000 mcg Vit B 1 50 mg Vit B 2 5 mg niacinamide 75 mg, pyridoxine HCl 5 mg panthenol 6 mg / cc Vial 20 cc
- RUBECEN** (Central) Vitamin B 12 activity 1000 mcg / ml Vial 10 cc
- RUBICIN** (Sull ff & Case) Shubarb root 0.78 Gm sodium bicarbonate 0.39 Gm cinna man 0.39 Gm, berberis aquifolium 0.39 Gm glycerin 0.39 Gm al peppermint 1.50 cc / fl oz Bot pts gals
 Use: Mild laxative and antacid
- RUBIFAX** (Merck Sharp & Dahme) Capsicum oleo resin, mustard oil syn, turpentine oil cedarwood oil, methyl salicylate & camphor in a suitable ointment base Tube 1 oz
 Use: Rubefacient and counterirritant
- RUBIGUENT** (Ives Cameron) Histamine dihydrochloride 0.1% methyl nicotinate 1%

* Generic name recognized by NF USP or AMA

- glycol monosalicylate 10%, capsicum oleo-resin 0.1% Cream, Tube 1 oz
 Use: Rubefacient & counterirritant
- RUBIPHEN.** (Tilden) Phenobarbital 1/12 gr, Ipecac lr 24 min, Sod citrate 8 gr, ammonium chloride 1 gr & chloroform 1/2 min / oz Bot 4 oz, 1 pt, 1 gal
 Use: Sedative for cough
- RUBRAFERATE, (Squibb)** Vit B 12 4 17 mcg, folic acid 0.28 mg, ferrous sulfate exsic 130 mg, ascorbic acid 50 mg / Cop Bot 100s, 1000s
 Use: Anemia
- RUBRAFOLIN** (Squibb) Cap (Vit B 12 25 mcg, folic acid 1.67 mg) Bot 100s, 1000s Vial (Vit B 12 15 mcg, folic acid 5 mg / cc) 5 cc
 Use: Anemias
- RUBRAGRAN** (Squibb) Vit B 12 25 mcg Vit B-12 intrinsic factor 1/2 u, ferrous sulfate exsic 340 mg, folic acid 1 mg, ascorbic acid 100 mg, liver 100 mg / Cop Bot 60s, 500s
 Use: 1 b l d; onemias
- RUBRAMIN, NND 1962** (Squibb) Vit B 12 Amp (15 & 30 mcg) 1 cc, Box 5s Vial (15 30, 50, 100, 1000 mcg / cc) Sol; Amp (15 mcg) 1 cc; Vial (30 mcg / cc) 10 cc Cop (25 mcg) Bot 100s, 500s 1000s
 Use: Hematopoietic
- RUBRAMIN PC.** (Squibb) Vit B 12 USP (cyanocobalamin) 30 50 100 1000 mcg / cc (30, 50 100, 1000 mcg / cc) Vial 10 cc (1000 mcg / cc) Amp 1 cc
 Use: Vitamin for various anemias
- RUBRAPLEX.** (Lanett) Vitamins B 1 215 mg, B 2 2 mg, B-6 0.5 mg, B 12 5 mcg, cal pantothenate 1 mg, niacinamide 20 mg / cap Bot 100s, 1000s, 5000s
- RUBRATON.** (Squibb) Iron 38 mg, Vit B-12 4 mcg, B 1 1 mg, B 2 1 mg, B 6 0.5 mg niacinamide 5 mg, pantothenic acid 1.5 mg / 5 cc Bot B fl oz, 1 pt
 Use: Anemia
- RUCEEKAY.** (Kremers Urban) But n 10 mg Vit C 30 mg, menadiene 1 mg / Tab Bot 100s, 1000s
 Use: Orally (1-3 daily) capillary protection w/Vitamin K therapy
- RUCEMIN** (Kinney) Rutin 20 mg, ascorbic acid 60 mg / Tab Bot 100s
 Use: Orally; capillary fragility
- RUCEVITIN** (Kremers Urban) Butin 20 mg-Vit C 150 mg / Tab Bot 100s, 500s 1000s Rucevita-60 contains rutin 60 mg, Bot 25s 100s, 1000s
 Use: Orally (1 t i d); capillary protect on
- RUCON** (Parke, Davis) Butin 100 mg, as

- corbic acid 100 mg. & dicalcium phosphate anhy. 400 mg./Cap. Bot. 100s, 1000s.
 Use: (1 l.i.d.) Correction of increased capillary fragility and permeability.
- RUCYN CAPTAB.** (Invenex) Vit. C 400 mg., niacinamide 250 mg., rutin 10 mg., B-12 5 mcg./Tab. Bot. 100s.
 Use: Capillary fragility.
- RUDIONATE-C.** (Ascher) Meeadione 0.5 mg., ascorbic acid 25 mg., rutin 10 mg., CaL lactate 7.5 gr./Tab. Bot. 100s, 500s.
 Use: (1-2 q.i.d.) Hemorrhagic conditions.
- RUFOLBEX.** (Central) Cyanacabalamie 50 mcg., folic acid 5 mg., methamphetamine HCl 2.5 mg., niacinamide 100 mg./ml. Vials 10 ml.
 Use: Anemias.
- RUFOLCEN.** (Central) Cyanacabalamie 50 mcg., folic acid 5 mg., methamphetamine 2.5 mg., niacinamide 100 mg./ml. Vial 10 cc.
- RUFOLFX.** (Lanett) Thiamine mononitrate 1.5 mg., riboflavin 1.5 mg., pyridoxine HCl 1 mg., Vit. B-12 5 mcg., ascorbic acid 50 mg., niacinamide 10 mg., ferrous fumarate 200 mg., d-sorbitol 200 mg., folic acid 0.25 mg./Cap. Bot. 100s.
 Use: Iron & vitamin deficiencies.
- RUGAR.** (McKesson & Robbins Lab.) A colloidal dispersed barium sulfate 72.73%. Jars 10 oz.
 Use: Radiopaque medium
- RUNEXATAL.** (Lemmon) Phenobarbital 15 mg., mannitol hexanitrate 30 mg., rutin 20 mg., ascorbic acid 10 mg./Tab. Bot. 100s
 Use: Treatment of hypertension and angina pectoris
- RUNEXATAL W/RESERPINE.** (Lemmon) Reserpine 0.1 mg., mannitol hexanitrate 30 mg., rutin 20 mg., ascorbic acid 10 mg./Tab. Bot. 100s
 Use: Treatment of hypertension
- RULIVAN.** (Squibb) Vit. B-12 20 mcg. as found in liver lnj. & Vit. B-12 30 mcg. from other sources/1 cc. Vial 10 cc.
 Use: Anemias.
- RUMA.** (El Paso Pharm.) Tab. Bot. 100s.
- RUMARID.** (Stanley Drug Prods.) Renofon. Cal. acetylsalicylate. Tab., Bot. 100s.
 Use: Analgesic.
- RU-NITRAL.** (Plessner) Mannitol hexanitrate 32 mg., rutin 20 mg., phenobarbital 16 mg./Tab. Bot. 100s, 1000s.
 Use: (1 q.i.d.) Vasodilator.
- W/Theophylline.** (Plessner) Mannitol hexanitrate 32 mg., theophylline 1.5 gr., rutin 20 mg., phenobarbital 1/4 gr./tab. Bot. 100s, 1000s.
 Use: (1 q.i.d.) Vasodilator.
- * Generic name recognized by N.F., U.S.P. or A.M.A.
- RUPHENAMIN.** (Lanett) Amieaphylline 100 mg., phenobarbital 15 mg., rutin 20 mg./cap. Bot. 500s, 1000s.
 Use: (1 l.i.d.) Cardiovascular disorders.
- RUPHON.** (Massengill) Sod. salicylate 5 gr., Pot. iodide 1 gr., Gelsemium ext. 1/2 gr., thiamine 1 mg., Cimicifuga ext. 1/2 gr., riboflavin 0.5 mg./Tab. Bot. 100s, 500s, 1000s.
 Use: (1-2 t.l.d.) Analgesic, antipyretic & relief of rheumatoid arthritis.
- RUPHYLLIN.** (Searle) Aminaphylline 100 mg., rutin 20 mg., phenobarbital 1/4 gr./Tab. Bot. 100s, 1000s.
 Use: (1 l.i.d.) Cardiovascular disorders.
- RUSERPATOL.** (Lemmon) Reserpine 0.2 mg., pentaerythritol tetranitrate 20 mg., rutin 40 mg., ascorbic acid 20 mg./Tab.
 Use: Treat hypertension and prevent angina pectoris.
- RUSON CHLORIDE.** (Northwest Labs.) alpha-Pyridinium acetic acid ester of beta-coprylamide ethanol Cl. N-coprycalamine formylmethyl pyridinium chloride. Sterilized chloride. Bot. 1 pt., 1 gal.
 Use: Disinfection, fungicidal & bactericidal action.
- RUSSELL VIPER VENOM.** See: Styree (Burroughs Wellcome)
- RUST INHIBITOR.**
 See: Anti-Rust, Tab. (Winthrop Labs.)
 Q. A. I., Pow. (Hollingshead Corp.)
 Sodium Nitrite
- RUTAMINAL.** (Schenley) Rutin 60 mg., ascorbic acid 25 mg., aminaphylline 100 mg., phenobarbital 15 mg./Tab. Bot. 100s, 1000s.
 Use: (1-2 t.l.d., p.c.) Cardiovascular disorder.
- RUTAMINAL-RQ.** (Schenley) Aminaphylline 100 mg., quercetin 15 mg., ascorbic acid 25 mg., reserpine 50 mcg./tab. Bot. 100s.
 Use: (1 p.c. & h.s.) Treatment of cardiovascular diseases.
- RUTASCORB.** (U.S. Vitamin) Ascorbic acid 50 mg., rutin 20 mg./Tab. Bot. 100s. Also available in triple strength.
 Use: (2 t.l.d.) Capillary permeability.
- RUTGERS 612.**
 See: Ethohexadiol, U.S.P.
- * **RUTIN.** N.F., A.D.R. 1955; Tab. N.F. (Various Mfr.) 3-rhamnoglucoside of 5,7,3',4'-tetrahydroxyflavone. Eldrin, globularictin, myricitorin, oxyritin, phytomelin, rutoside, rapharin.
 Lloyd Bros.—Sol. (20 mg./5 cc.)
 Various Mfr.—Tab. (20, 50, 60 & 100 mg.)
 Use: Vascular disorders.
 Rutin Combinations on pp. 642 to 644.

Rutin Combinations

| Rutin | Others | Manufacturer | Brand |
|------------------|--|---------------------|-------------------------------|
| 20 mg | Aminophylline 50 mg, mannitol hexanitrate 16 mg, phenobarbital 16 mg | Kendall | Hexarutan Tab |
| 30 mg | Am naphylline 97 mg phenobarbital 16 mg | Suliff & Case | Amo-Ru Tab |
| 20 mg | Aminophylline 100 mg, phenobarbital $\frac{1}{4}$ gr | Searle | Ruphylin Tab |
| 60 mg | Aminophylline 100 mg, phenobarbital $\frac{1}{4}$ gr, ascorbic acid 25 mg | Schenley | Rutaminol, Tab |
| 20 mg | Aminophylline 100 mg, phenobarbital 15 mg, veratrum viride 95 mg | Adams | Verutamin, Tab |
| 20 mg | Ascorb c acid 100 mg | A P C | Ascor Rutin, Tab, No 1 |
| 60 mg | Ascorbic acid 300 mg | A P C | Ascor Rutin, Tab, No 2 |
| 100 mg | Ascorbic acid 100 mg | Schenley | Capardin, Tab. |
| 100 mg | Ascorb c acid 100 mg | Parke Davis | Rutin with C, Tab |
| 20 mg | Vit C 50 mg | Suliff & Case | Ru An Cee, Tab |
| 50 mg | Ascorbic acid 50 mg | Maltbie | Clavone, Tab |
| 50 mg | Ascorb c acid 50 mg | Parke, Davis | Rutin with C, Tab |
| 20 mg | Ascorbic acid 60 mg | Strassenburgh | Rhamnitolin Tab |
| 20 mg | Ascorbic acid 50 mg | Kinney | Rucemin, Tab |
| 20 mg | Ascorbic acid 100 mg | U S Vitamin | Rulascarb, Tab |
| 20 mg | Ascorbic acid 100 mg | Squibb | Rularbin, Tab |
| 60 mg | Ascorb c acid 25 mg, aminophylline 100 mg, phenobarbital $\frac{1}{4}$ gr | Lannett | Lanoscarbin Tab |
| 30 mg | Ascorbic acid 40 mg, dicumarol 50 mg | Schenley | Rutaminol Tab. |
| 25 mg | Ascorb c acid 100 mg, hesperidin 100 mg | Strassenburgh | Dicumeral W/Rhamnitolin, Tab. |
| 15 mg | Ascorb c acid 25 mg, mannitol hexanitrate $\frac{1}{2}$ gr | Arnar Stone | Ascarulin 8 o, Tab |
| 50 mg | Ascorbic acid 50 mg, mannitol hexanitrate 30 mg, phenobarbital $\frac{1}{4}$ gr | Cale Chem Co | Mannucee Tab |
| | | Chicago Pharm Co | Manncor, Tab |
| 10 mg | Ascorbic acid 15 mg, mannitol hexanitrate $\frac{1}{2}$ gr, phenobarbital $\frac{1}{4}$ gr, theophylline $1\frac{1}{2}$ gr | Mastengill | Semhylen, Cap |
| 15 mg | Ascorb c acid 30 mg, mannitol hexanitrate $\frac{1}{2}$ gr, Vit B 12, phenobarbital $\frac{1}{4}$ gr | Strassenburgh | Mazitate W/Rhamo-8-12, Tab. |
| 10 mg | Ascorbic acid 30 mg, menadione 1 mg | Kremert Urban | Rucekay, Tab |
| 20 mg | Ascorbic acid 50 mg, theobromine 5 gr, phenobarbital $\frac{1}{2}$ gr | Haskell | Ruzan, Tab |
| 20 mg | Ascorbic acid 30 mg, veratrum viride 1 gr, aspirin 5 gr | Mallard | Aceto-Ver, Tab |
| 100 mg | Citrus b flavanoids complex 100 mg, ascorbic acid 100 mg | Plesner | Cap Jan, Tab |
| $\frac{1}{2}$ gr | Crotalaria 1/6 gr, viscum album 25 gr | Lloyd | Hyper Viscra W/Rutin, Tab |
| 20 mg | Crotalaria tr 2 min, d oilyl barbituric acid $\frac{1}{4}$ gr, nitroglycerin 0 26 mg Sed n tr te 1 gr | Buffington | Vasorutin, Pulls |

| Rutin | Others | Manufacturer | Brand |
|-------|---|------------------|---------------------------------|
| 20 mg | Rauwolfia serpentina alkaloids 0.75 mg, neohyline 100 mg | Paul Maney | Loten Encofe |
| 20 mg | Reserpine 0.1 mg, mannitol hexanitrate 30 mg, ascorbic acid 10 mg | Lemmon | Ruhexatal W/Reserpine, Tob. |
| 10 mg | Reserpine 0.1 mg, mannitol hexanitrate 30 mg, theophylline 100 mg, ascorbic acid 15 mg | Mossengill | Neo Semhyfen, Cap |
| ½ gr | Sod nitrite 1 gr, veratrum viride ¾ gr | Superior | Verutinium Tab |
| ½ gr | Sod n. trite 1 gr, phenobarbital ¼ gr, veratrum viride ¾ gr | Superior | Verutinium W/Phenobarbital, Tab |
| ½ gr | Theobromine 5 gr, phenobarbital ½ gr | Van Pelt & Brown | Theobarb W/Rutin, Tab |
| 20 mg | Theobromine 5 gr, phenobarbital ½ gr, ascorbic acid 50 mg | Haskell | Ruzen, Tab |
| 10 mg | Theophylline 1 ½ gr, phenobarbital ¼ gr, ascorbic acid 15 mg, mannitol hexanitrate ½ gr | Mossengill | Semhyfen Cap |
| 20 mg | Veratrum viride 100 mg, manitol hexanitrate 30 mg, rauwolfia ser pent mg 0.75 mg | Paul Maney | Raufo Encofe |
| 20 mg | Veratrum viride 65 u, phenobarbital 15 mg | Irwin, Neister | Veroflex, Tab |
| 20 mg | Veratrum viride 95 mg, phenobarbital ¼ gr aminophylline 100 mg | Adams | Verutamin, Tab |
| 10 mg | Veratrum viride 100 mg, phenobarbital ¼ gr, mannitol hexanitrate 30 mg | Rand | Verutal, Tab |
| 20 mg | Veratrum viride 32 mg, phenobarbital ¼ gr, mannitol hexanitrate 16 mg, choline methyl tartrate 100 mg | Hallard | Amvero, Tab |
| 20 mg | Veratrum viride 1.5 gr, phenobarbital 15 gr, neohyline 1.5 gr | Paul Maney | Vir I Phyl, Tab |
| ½ gr | Veratrum viride ¾ gr, phenobarbital ¼ gr, Sod nitrite 1 gr | Superior | Verutinium W/Phenobarbital, Tab |
| ½ gr | Veratrum viride ¾ gr, Sod nitrite 1 gr | Superior | Verutinium Tab |

RUTINSOL. (Meer Corp) Water soluble rutie.

RUTOL. (Pitman-Moore) Mannitol hexanitrate $\frac{1}{2}$ gr., rutie $\frac{1}{6}$ gr., phenobarbital $\frac{1}{8}$ gr./Tab. Bot. 100s, 500s, 1000s.
Use: (1-2 t.i.d.) Vasodilator.

RUTORRIN. (Squibb) Ascorbic acid 100 mg., rutie 20 mg./Tab. Bot. 100s, 1000s. Also available in triple strength (300 mg. & 60 mg.) Rot. 50s.
Use: Orally (1-2 t.i.d.) capillary permeability.

RUTOSIDE. See: Rutin, N.F.

RUVAMIN. (New Mexico Pharm.) R-21 25 mcg./Cap. Bot. 100s, 1000s.

RUZAN. (Haskell) Ascorbic acid 50 mg., theobromine 5 gr., rutin 20 mg., phenobarbital $\frac{1}{2}$ gr./Tab. Rot. 100s, 500s.
Use: (1-2 t.i.d.) Vasodilator.

RYBUTOL. (Vitamins Corp) Vit. R-1 15 mg., R-2 6 mg., C 30 mg., nicotinamide 10 mg., Cal, pantotheate 3 mg., Vit. B-6 0.5 mg., Vit. E 5 mg., whole dried liver 100 mg., dried debittered brewer's yeast 100 mg., ferrous glucoate 30 mg., choline dihydrogen citrate 20 mg., inositol 20 mg., dimethylacetate 20 mg., folic acid 0.1 mg., Vit. B-12 3 mcg., dicalcium phosphate anhy. 200 mg., Pot. iodide 0.15 mg., Mag. sulfate dried 7.2 mg., copper sulfate dried 5 mg., Mang. sulfate 3.4 mg., cobalt sulfate 0.2 mg., Pot. chloride 1.3 mg./Cap. Bot. 25s, 50s, 100s, 250s, 1000s.
Use: Dietary supplement.

RYDAR. (Cutler Lab) See: Dioxodrol HCl.

RYDIAMIN. (Kremers-Urban) Pyridoxine HCl 100 mg., thiamine HCl 100 mg., di-methionine 100 mg., benzyl alcohol 40 mg./2 cc. Vial 20 cc. Pyridoxine HCl 20 mg., thiamine HCl 5 mg., di-methionine 300 mg./tab. Tab., Bot. 100s.
Use: Nutritional supplement.

RYNALERT. (Irwin, Neisler & Co) Phenylephrine tannate 37.5 mg., pyrilamine tannate 25.0 mg., chlorpheniramine tannate 50 mg., caffeine alkaloid 200.0 mg./cap Bot 30s.
Use: For the common cold.

RYNATAN TABLETS. (Irwin, Neisler) Phenylephrine tannate 25.0 mg., prophepyridamine tannate 37.5 mg., pyrilamine tannate 37.5 mg./Tab. Bot. 30s, 100s, 500s.
Use: Upper respiratory congestion resulting from colds, sinusitis, allergic rhinitis.

RYNATAN EXPECTORANT. (Irwin, Neisler) Carbetapentane tannate 30 mg., ephedrine tannate 5 mg., phenylephrine tannate 5 mg., chlorpheniramine tannate 2 mg., pyrilamine tannate 12.5 mg./5 cc. Bot. pt.
Use: Cough due to colds.

RYNATUSS. (Irwin, Neisler) Carbetapentane tannate 60 mg., chlorpheniramine tannate 5 mg., ephedrine tannate 10 mg., phenylephrine tannate 10 mg./Tablet. Bot 30s, 500s.

Use: Treatment of Common Cold.

RYNATUSS SUSPENSION. (Irwin, Neisler) Carbetapentane tannate 30 mg., chlorpheniramine tannate 4 mg., ephedrine tannate 5 mg., phenylephrine tannate 5 mg./5 cc. Susp. Bot. 8 oz.

Use: Antitussive.

RYNATUSS TABLETS. (Irwin, Neisler) Carbetapentane tannate 60 mg., chlorpheniramine tannate 5 mg., ephedrine tannate 10 mg., phenylephrine tannate 10 mg./Tab. Tabs. Bot. 30s.

Use: Antitussive.

RYNODYNE. (Canfield) Dexamethorphan HBr 60 mg., chlorpheniramine maleate 7.5 mg., phenylephrine HCl 30 mg., ammonium chloride 500 mg./oz. Bot. 3 or 4 fl. oz., 1 pt., 1 gal.

Use: Control of cough.

RYPROGEN. (Organon) Typhlized Reller proteic antigen for accurate serodiagnosis of syphilis. Ampul 10 cc.
Use: Diagnostic aid.

RYSTAL. (Grain) N-acetyl-p-aminophenol 120 mg., glyceryl gualcolate 50 mg./5 cc. Byr. Bot. 1 pt.

Use: Analgesic for the relief of minor muscular aches and pains.

RYTHRITOL. (Metropolitan Lab) Pentaerythritol tetranitrate 10 mg./Tab. (green) Bot. 100s, 1000s

Use: Vasodilator

RYTHROGEN. (O. P. Harvey) Liver w/cryotalline Vit. B-12 36 mcg. & thiamine HCl 4.5 mg./cc. Vial, 10 cc.

Use: Anemias.

S AND A. (Tilden) Cascara sagrada, gentia, taraxacum, senna, trifolium. Bot. pts, gals.

Use: Laxative, stomachic.

SABABIN. (Haag) Salicylamide 250 mg., YABA 125 mg., mephenesin 125 mg./Tab. Bot. 100s.

Use: 2 q. 3-4 h. analgesic, relaxant.

SABIN ORAL POLIO VACCINE. (PRx) Live poliomyelitis virus, type 1./vial. Vials 10 dose.

Use: Immunization.

SACAGEN. See: Simran, Cap (Merrell)

SACASU. (Smith, Miller & Patch) Sulfur 2%, salicylic acid 2%, camphor 15%. Amp. 1 cc., 5s, 25s, 100s.

Use: Nonspecific antirheumatics.

SACCHARATED IRON OXIDE, STERILE SOLUTION. See: Proferrin, Vial (Merck Sharp & Dohme)

***SACCHARIN, USP, ADR** (Benzosulfimide, Gluside)

Mallinckrodt—Pkg 1 oz, ¼ lb, 1 lb

Merck—Pkg 1 oz, ¼ lb, 1 lb

See Saxon, Tab (Burroughs Wellcome)

Use: Sweetening agent when sugar is contraindicated

W/Aspirin See Theryl, Tab (Church)

SACCHARIN 12. (Armour) Saccharin Sod

¼ gr, Vit. B 12 1 mcg/Tab Bot 100s

Use: Sweetening agent

***SACCHARIN CALCIUM, USP**

***SACCHARIN SODIUM, USP, ADR, Tab, USP** (Mallinckrodt) (Benzosulfimide Sod,

Soluble Gluside, Soluble Saccharine) Pow,

Bot 1 oz, ¼ lb, 1 lb Tab usual sizes

(Various Mfr) See Sweete, Sal (Squibb)

Use: Sweetening agent & test for circula-

tion time of blood

SACCHARIN SOLUBLE. See: Saccharin Sodium USP

SAFENON. (Massengill) Iron peptonized 0.25

Gm/Tab Sol 100s, 1000s

SAFF, NND 1962 (Abbott) Safflower oil

65%, linoleic acid 46% Oil emulsion Bot

1 pl

Use: Management of atherosclerosis

SAFFLOR CAPSULES, NND 1962 (Haug

Drug) 750 mg safflower oil/Cap Sol 100s

Use: Lowering cholesterol blood level

SAFFLOWER CAPS. (American Pharm Co)

Safflower oil 750 mg/Cap Sol 100s, 1000s

Use: Cholesterol lowering agent

***SAFFLOWER OIL, NND 1962** See: Del

benol Cap (Brawn Pharm) Safflor Capsules

(Haug Drug) Safflorin, Cap (Superior)

W/Linoleic Acid See: Emul (Abbott)

W/Pyridoxine, mixed tocopherols See:

Salplex, Emulsion (Lloyd, Dabney &

Westerfield)

W/Vit B-6 Vit E See: Alestral, Cap

(Testogor)

SAFFOLA. (Saffola Lab) Refined safflower

oil

Use: Cruffy, to decrease cholesterol blood

level

SAFFOLIN. (Tutag) Safflower oil 500 mg,

Vit E 2.5 IU pyridoxine 5 mg, niacin 20

mg/Cap

Use: Anticholesterol

SAFFLEX EMULSION (Lloyd, Dabney &

Westerfield) linoleic acid 6.5 Gm, pyridox-

ine HCl 5 mg, mixed tocopherols 10 mg/

15 cc Bot 1 pl, 1 gal

Use: Lower blood cholesterol level

SAJODIN, NNB 1950 (Winthrop Labs)

* Generic name recognized by N.F., U.S.P. or A.M.A.

Cal iodobenohene Paw 1 oz Tob 1 gr,
Bot. 24s; Tob 8 gr, Bot. 20s

Uses: Iodine therapy

SALACETIN. See: Acetylsalicylic Acid

SALADIN. (Sirasenburgh) Acetophenetidin

2 gr, salaf 2 gr, acetylsalicylic acid 2 gr/

Tab Bot 100s, 500s, 1000s

Use (1-2 q 3 h) Antipyretic & analgesic

SALAMIDE, NND 1962 (Columbus) Soli-

cyamide Tab 5 gr, Bot 100s, also avail

w/colchicine 1/250 gr/Tab Bot 100s

Use (1-3 q 3 h) Analgesic & antipyretic.

SAL-A-SUL. (Dane) (Su-A Sa) Vit A 100,000

u, sulfur 2%, salicylic acid 2% Cream 1

oz, 2 oz, 4 oz, 1 lb

Use: Antiseptic

SALATAR CREAM. (Lannell) Coal tar solu-

tion 5%, salicylic acid 3% Jar 4 oz, 1 lb

SALSEX. (Wintersmith) Aminobenzoic acid

275 gr, ascorbic acid 10 mg/Tab Bot

100s, 1000s

Use: (1-2 q 3 h) Analgesic & antipyretic

SALCYDE OX. (Massengill) Sod salicylate 5

gr, aluminum hydroxide gel, dried 2 gr,

Cal ascorbate 1 gr, Cal carb 1 gr/Tab

Bot 100s, 500s

Use: (3-4 q 4 h) Rheumatic & arthritic con-

ditions

SALCIPAB. (Kirkman) Sod salicylate 5 gr,

Sod aminobenzoate 5 gr/Tab Also avail

w/Vit C 50 mg/Tab Sol 100s, 500s

1000s

Use: Rheumatic conditions

SALCOCE. (Cale Chem Co) Sodium salicyl-

ate 0.5 Gm, calcichine 0.25 mg, ascorbic

acid 20 mg/Tab Bot 100s

Use: Antirheumatic & gout

SALCOBBINE. (Chicago Pharm) Sod salicyl-

ate 5 gr, ascorbic acid 100 mg, calcichine

1/150 gr/Tab Bot 100s, 500s, 1000s

Use: Arthritic gout

SALCOBE. (Massengill) Cortisone acetate 2.5

mg, Sod salicylate 0.3 Gm, aluminum hy-

droxide 0.12 Gm, Cal ascorbate 60 mg-

Cal carb 60 mg/Tab Bot 100s

Use: 12-16 initially, then 6-8 daily; rheu-

matism

SALCOBT-DELTA. (Massengill) Prednisone 1

mg, pot salicylate 300 mg, w/col. pen-

tathenate and buffers/Tab Bot 100s

Use: Treatment of rheumatic disorders

SALCOBT SODIUM FREE. (Massengill) Corti-

sone acetate 2.5 mg, pot salicylate 0.3

Gm, alum hydroxide gel 120 mg, cal

ascorbate 60 mg, cal carbonate 60 mg/

tab Bot 100s

Use: Sodium free, treatment of rheumatic &

arthritic conditions

- SAL-ESTER-SAL.** (Cole) Salicylic ester of salicylic acid. Tab. 1 gr., Bot. 100s.
Use: Analgesic & antipyretic.
- SAL ETHYL CARBONATE, N.N.S. 1948.** (Parke, Davis) Carbonic acid ester of ethyl salicylate. Tab. 5 gr., Bot. 100s, 500s, 1000s. Pow. Bot. 4 oz., 1 lb.
Also avail. w/amineopyrine 1½ gr. & sal ethyl carb. 3½ gr. Sol. same.
Use: Analgesics, antipyretics & antirheumatics.
- SALÉTIN.** See: Acetylsalicylic Acid.
- SALITO.** (Mallard) Salicylamide, acetophenetidin, caffeine, camphor monobromated, & aspirin. Tabs. Sol. 12s, 30s, 100s.
Use: Analgesic.
- SAL-EZE.** (Cole) Salicylamide 3 gr., Mag salicylate 2 gr., Vit. C 20 mg., p-aminobenzoic acid 2 gr./Tab. Bot. 100s.
Use: 2 t.i.d. arthritides.
W/Cochicine. Sal-Eze formula plus colchicine 1/250 gr./Tab. Bot. 100s.
Use: Gout.
- SALGILATE.** (Walker) Salicylamide 4 gr., caffeine ½ gr., Hyoscyamus ext. ¼ gr./Tab. Bot. 100s, 500s, 1000s.
Use: Symptomatic treatment of rheumatic disorders.
- SALHABT.** (Mort) Sod. salicylate 4 gr., p-aminobenzoic acid 4 gr., ascorbic acid ½ gr., physostigmine salicylate 1/250 gr., homatropine methylbromide 1/120 gr./Tab. Bot. 200s, 500s.
Use: Analgesic for rheumatic conditions.
- SALISAR Jr.** (Jenkins Labs.) Aspirin 2 gr., phenobarbital ½ gr./tab. Sol. 100s, 1000s.
Use: Analgesic and sedative.
- SALICLM.** (Irwin, Neisler) Salicylamide. Tab. 5 gr., Bot. 1000s.
Use: (1-3 q. 3 h.) Analgesic & antipyretic.
- SALICLOHYL.** (Upjohn) Sod. salicylate 10 gr., Sod. bicarb. 20 mg./tsp. Granulated eff. pow. 4 oz., 8 oz.
Use: Antacid.
- SALICITUM.** (Smith, Miller & Patch) Sod. salicylate 5 gr., colchicine 1/1000 gr., Pot. iodide ½ gr./Tab. Bot. 100s, 1000s.
Use: Analgesic & antirheumatic.
- SALICLUM.** (Taylor) Sod. salicylate 4 gr., Sod. p-aminobenzoic acid 4 gr., ascorbic acid ½ gr./Tab. Bot. 100s, 1000s.
Use: Rheumatic Fever.
- SALICOL.** (Taylor) Sod. salicylate 4 gr., p-aminobenzoic acid 4 gr., ascorbic acid ½ gr., colchicine 1/250 gr./Tab. Sol. 100s, 1000s.
Use: Gout therapy.
- SALICOLCHIN.** (Gold Leaf) Sod. salicylate
- 3 gr., colchicine 1/1000 gr., Pot. iodide ½ gr./Tab. Bot. 100s, 1000s.
Use: (1-2 q. 3 h.) Analgesic.
- SALICOBATE.** (Cole) Sod. salicylate 10 gr., ascorbic acid 32 mg./Tab. Bot. 1000s. Also avail. in half strength
Use: (2-3 q. 3 h.) Antirheumatic.
- SALICESIN.** (Upjohn) Benzoic acid 2%, salicylic acid 2%, sec. amylicresols 0.1%, o-hydroxyphenyl mercuric chloride 0.1%. Cream 2 oz. Liq. 1 oz., 4 oz., 16 oz., 1 gal.
Use: Fungus infections of skin.
- SALICYL ALCOHOL.** (Delta) Saligenin. Bot. 100 Gm.
Use: 2 to 8% solution for anesthesia.
- SALICYLAMIDE, N.F., N.N.D. 1962.** Tab. N.F. & Hydroxybenzamide
Use: Analgesic.
See: Atka-Sal, Tab. (A.P.C.)
Amid-Sal, Tab. (Glenwood)
Dropsprin, Liq. (M. H. Thompson)
Liquiprin (Johnson & Johnson)
O.H.S., Tab. (Baulich)
Raspberin, Sol. (Smith, Miller & Patch)
Salomide, Tab. (Columbus)
salicem, Tab. (Irwin, Neisler)
Salrin, Tab. (Warren-Teed).
- Salicylamide-Artenanthenetidin Combinations on pp. 648 and 649.
Salicylamide-Mephenesin Combinations on p. 649.
Salicylamide-Phenobarbital Combinations on p. 650.
Salicylamide-Vit. C Combinations on pp. 650 and 651.
- SALICYLANILIDE, N.F., N.N.D. 1962.** N-Phenyl salicylamide.
Use: Antifungal agent
See: Anadol, Oint., N.N.D. 1958. (Rorer)
Safinidal, Oint., N.N.D. 1958. (Dank).
- W/3-Chloro-salicylanilide, 5,3'- & 5,4'-Dichlorosalicylanilides; undecylenic acid; and zinc undecylenate. See: Salundek New. Oint. (Wallace & Tiernan)
W/Hyamine. See: Quatrol, Oint. (Doma).
W/Octraphenate. See: Sylamin, Cream (Abbott).
W/Phameral chloride. See: Sanicyl, Cream (Parke, Davis).
- SALICLATED SILE EXTRACT.** See: Cholegastin, Tab. (Strang)
- SALICYLAZOSULFAPYRIDINE, N.N.D., 1962.** See: Azulfidina, Tab., N.N.D. 1958. (Pharmacia)
- SALICYLIC ACID, U.S.P., N.N.R. 1949.** Col-lodion, U.S.P. Plaster, U.S.P. Orthohydroxybenzoic acid. Cryst., Pkg. 1 oz., ¼ lb., 1 lb. Pow., Pkg. ¼ lb., 1 lb.
Use: Keratolytic agent.

* Generic name recognized by N.F., U.S.P. or A.M.A.

Salicylamide-Acetophenetidin Combinations

| Salicylamide | Acetophenetidin | Others | Manufacturer | Brand |
|--------------|-----------------|---|------------------------|----------------------------|
| 150 mg | 120 mg | Achromycin HCl 125 mg, caffeine 30 mg, chlorothen citrate 25 mg | Lederle | Achroctidin, Tab |
| 150 mg. | 120 mg | Achromycin 125 mg, Vit C 25 mg, pyrilamine maleate 15 mg | Lederle | Achroctidin Syr /5 cc. |
| 200 mg | 120 mg | Amobarbital 15 mg, d amphetamine sulf 2 mg, hyoscyamine sulf 0.05 mg | Finley | Sedemasal, Cap |
| 1 1/4 gr. | 1 1/4 gr. | Caffeine 1/2 gr | Paul Maney | Anodyn, Tab. |
| 2 gr. | 2 1/2 gr. | Caffeine 1 gr | Bristol Myers | Excedrin, Tab |
| 3 1/2 gr. | 2 1/2 gr. | Caffeine 1/2 gr, phenobarbital 1/4 gr | Smith Darsey | Angelal, Tab. |
| 3 1/2 gr. | 2 1/2 gr | Caffeine 1/2 gr | Rowell | Salprin, Tab also w/Codene |
| 1 gr | 1 gr | Caffeine 1/2 gr, aspirin 2 gr, atropine sulf 1/1000 gr, phenylephrine HCl 5 mg | Clapp | Emagrin Forte, Tab |
| 2 gr. | 2 gr | Caffeine 1/20 gr, phenobarbital 1/8 gr, hyoscyamus ext 1/8 gr | Lemmon | Aspifedrine, Tab & Cap |
| 190 mg. | 130 mg | Chlor trimeton maleate 4 mg, vit C 50 mg, methamphetamine HCl 1.25 mg, caffeine 30 mg | Schering | Concoidin Forte, Cap. |
| 2 1/2 gr. | 2 1/2 gr | dl Desoxyephedrine HCl 2 mg | Wampole | Tryada, Tab. |
| 250 mg. | 120 mg | Dover's powder 24 mg, acephedrine HCl 5 mg, pyrilamine maleate 12 mg, vit C 50 mg | Central | Cenamal, Cap |
| 200 mg | 120 mg | Hesperidin 100 mg, vit C 50 mg, pyrilamine maleate 8.33 mg, methopylone HCl 8.33 mg, phenylephrine HCl 5 mg, prophepyridamine maleate 6.25 mg, caffeine 30 mg | Boyle | Boyle Citra, Cap. |
| 230 mg. | 120 mg | Hesperidin purified 20 mg, vit C 20 mg, prophen pyridamine maleate 10 mg, caffeine 30 mg | Koerig | Caryban, Cap |
| 3.5 gr. | 2.5 gr. | Hesperidin 25 mg, caffeine 0.5 gr, ascorbic acid 50 mg, pyrilamine maleate 25 mg | Bryant Pharmacal Corp. | Buacal, Cap. |
| 194 mg | 130 mg | Penicillin V 100,000 u, promethazine HCl 6.25 mg, mepentermine sulf 3 mg | Wyeth | Pen Vee Cadin, Cap. |
| 100 mg. | 100 mg | Pet penicillin G 250,000 u, phenyltolaxamice di hydrogen citrate 25 mg | Patch | Elpafec, Tab |
| 3 gr. | 2 gr. | Pyrilamine maleate 25 mg, caffeine 1/2 gr, atropine sulf 1/600 gr, quinine sulf 1/2 gr, Vit C 25 mg | Taylor | Histaquin, Cap. - |

Salicylamide-Acetophenesisdin Combinations

| Salicylamide | Acetophenesisdin | Others | Manufacturer | Brand |
|--------------|------------------|---|------------------|-------------------|
| 3½ gr. | 2½ gr. | Pyrilamine molecule 12.5 mg., caffeine ½ gr., hyoscine sulf. 0.0702 mg., atropine sulf. 0.0140 mg., hyoscyne HBr 0.0042 mg. | Van Pelt & Brown | Pyro-Astro, Tab. |
| 150 mg. | 120 mg. | Tetracycline HCl 125 mg., Secizine HCl 15 mg., caffeine 30 mg. | Pfizer | Tetracyclin, Tab. |

Salicylamide-Mephesisin Combinations

| Salicylamide | Mephesisin | Others | Manufacturer | Brand |
|--------------|------------|--|----------------|---------------------------------------|
| 2½ gr. | 3½ gr. | | Columbus | Mephamide, Tab. |
| 300 mg. | 300 mg. | | Miller | Talagestic, Tab. |
| 200 mg. | 300 mg. | | Barre | Cerulet A, Tab. |
| 195 mg. | 195 mg. | Butabarbital Sod. 10 mg. | Kendall | Benesin, Tab. |
| Pres. | Pres. | Butabarbital & Vit. C | Ethical | MeBuSal-C, Cap. |
| 300 mg. | 150 mg. | Col. p-aminobenzoate 50 mg., Vit. C 50 mg. | Grant | Causalin, Tab. |
| 100 mg. | 250 mg. | O-Benzyl succinate | M. H. Smith | Esposmus, Tab. |
| 125 mg. | 125 mg. | Dipyrone 50 mg., Vit. C 50 mg. | Barre | Barizol, Cap. |
| 250 mg. | 250 mg. | Homatropine methylbromide 1 mg. | Mallard | Mephotine, Tab. |
| 150 mg. | 200 mg. | Homatropine methylbromide 2 mg. | Iemman | Hamesol, Tab. |
| 130 mg. | 400 mg. | Homatropine methylbromide 0.6 mg., omobarbital 8 mg., physostigmine 0.25 mg. | Testagar | Myamephetone, Tab. |
| 2½ gr. | 375 mg. | Homatropine methylbromide 1.5 mg., phenobarbital 16 mg., acetophetidin 2½ gr. | Adams | Relaxamine A-P, Tab. |
| 250 mg. | 125 mg. | Hydrocarbons acetate, 5 mg., 200 mg. | Walker | Articon, Tab. |
| 195 mg. | 195 mg. | PABA 125 mg. | Hoag | Sobasin, Tab. |
| 4 gr. | 2 gr. | PABA 195 mg., Vit. C 10 mg. | Lloyd & Dabney | Mybasol, Tab. |
| 250 mg. | 250 mg. | Pentobarbital ½ gr., phenobarbital ¼ gr., acetophenesisdin 2 gr., caffeine ¼ gr. | Thompson | Sopano-MAT, Tab. |
| 500 mg. | 250 mg. | Vit. C 15 mg. | Kremers-Urban | Solimph-C, Tab. |
| 162 mg. | 125 mg. | Vit. C 20 mg., prednisolone 0.75 mg. | Kremers-Urban | Solimph Forte with Prednisolone, Tab. |
| | | Vit. D 50,000 u. | Whittier | Ertron S-M, Cap |

Salicylamide-Phenobarbital Combinations

| Salicylamide | Phenobarbital | Others | Manufacturer | Brand |
|--------------|---------------|---|---------------|-------------------------------|
| 3½ gr | ¼ gr | N Acetyl p-aminophenol 2½ gr, caffeine ½ gr | Chicago Pharm | Cylamide W/Phenobarbital, Tab |
| 3½ gr | ¼ gr | d Amphetamine phosphate 1.5 mg | Irwin Neisler | Analdine Tab |
| 3½ gr | ¼ gr | Caffeine ¼ gr, phenacetic 2½ gr | Smith Darvey | Angelal, Tab |
| 3 gr | ¼ gr | dI Desoxyephedrine HCl 1/20 gr, acetophenetidin 2 gr, hyoscyamus ext ¼ gr | Lemman | Aspifedrine, Tab |

Salicylamide-Vit C Combinations

| Salicylamide | Vit C | Others | Manufacturer | Brand |
|--------------|--------|--|----------------|----------------------|
| 300 mg | 15 mg | | Lemman | Asal Tab |
| 62 mg | 10 mg | | Boermmel | Pedrin, Syr |
| 190 mg | 50 mg | Acetophenetidin 130 mg, chlor trimeton maleate 4 mg, methamphetamine HCl 125 mg, caffeine 30 mg | Schering | Carcedin Forte, Cap |
| 250 mg | 50 mg | Acetophenetidin 120 mg, Dovers powder 24 mg, rocephedrine HCl 5 mg, pyrilamine maleate 12 mg | Central Pharm | Cenamal, Cap |
| 230 mg | 20 mg | Acetophenetidin 120 mg, hesperidin purified 20 mg, prophenpyridamine maleate 10 mg, caffeine 30 mg | Koerlig | Caryban, Cap |
| 200 mg | 50 mg | Acetophenetidin 120 mg, hesperidin 100 mg, pyrilamine maleate 8.33 mg, methapyrilene HCl 8.33 mg, phenylephrine HCl 5 mg, prophepyridamine maleate 6.25 mg, caffeine 30 mg | Boyle | Boyle Citra, Cap |
| 300 mg | 50 mg | p-Aminobenzoate Cal 50 mg, mephensine 150 mg | Grant | Cavallin, Tab |
| 225 mg | 20 mg | p-Aminobenzoate pot 225 mg, salicylate pot 165 mg, dicalcium glucomate 75 mg | Dumas Wilson | Glutalate, Cap |
| 5 gr | 85 mg | Bioflavonoid complex 50 mg | Chicago Pharm | Citrisan, Tab |
| 250 mg | 25 mg | Cal succinate 200 mg | Hastell | Slimdate Tab |
| 300 mg | 15 mg | Calcitric 0.25 mg | Lemman | Asalcine, Tab |
| 125 mg | 50 mg | Dipyron 125 mg, N-acetyl p-aminophenol 125 mg | Arnor Stone | Dipraion Forte, Tab. |
| 1 gr | 100 mg | Hesperidin 100 mg, prophenpyridamine maleate 6 mg, phenylpyramine fumarate 6 mg, pyrilamine maleate 6 mg, caffeine ¼ gr | Person & Cavey | Tripac, Cap |

Salicylamide-Vit. C Combinations

| Salicylamide | Vit. C | Others | Manufacturer | Brand |
|--------------|--------|---|----------------|------------------------|
| 3.5 gr. | 50 mg. | Hesperidin 25 mg., phenacetin 2.5 gr., caffeine 0.5 gr., pyrilamine maleate 25 mg. | Bryant Pharm. | Bical, Cap. |
| 250 mg. | 15 mg. | Mephensin 250 mg. | Kremers-Urban | Salimeph-C, Tab. |
| 150 mg. | 25 mg. | Phenacetin 120 mg., ethromycin 125 mg., pyrilamine maleate 15 mg. | Lederle | Achrocidin, Syr./5 cc. |
| 3 gr. | 50 mg. | Prednisone 0.75 mg. | Mosey | Aspred-C, Tab. |
| 3 gr. | 25 mg. | Pyrilamine maleate 25 mg., phenacetin 2 gr., caffeine 1/2 gr., atropine salt, 1/600 gr., quinine sulf., 1/2 gr. | Taylor | Histaqin, Cap. |
| 150 mg. | 10 mg. | Sod. genitrate 150 mg., d-desoxyephedrine 0.5 mg. | Sulliff & Case | Casacomb, Tab. |
| 1/2 gr. | 30 mg. | Sod. salicylate 5 gr., Sod. PAB 3 gr., dihydrogen aluminum orthonoacetate 1/2 gr. | Patch | Elpagen, Tab. |
| 250 mg. | 40 mg. | Saccharic acid 250 mg. | Danley-Evans | Decosal, Tab. & El. |

W/Borotannic Complex, ethyl alcohol, See Onycho-Phytex.

W/Gelsamium comp. (Merck Sharp & Dahme) Sod. salicylate 2.6 Gm., cimicifuga 1 Gm./30 cc. Bot. 1 gal.

Use: Analgesic.

W/Sulfur, Sod. diacetyl sulfasuccinate, Sod. alkyl polyether sulfonate, Sod. sulfacetate, Kerohydric, hexachlorophene. See: Sebulex, Liq. (Westwood)

Salicylic Acid Combinations on p. 652.

SALICYLIC COMPOUND ELIXIR. (Warner-Chilcott) Sod. salicylate 6 gr., Cimicifuga 1.72 gr., Sod. bicarb. 0.32 gr./fl. dr. Bot. 12 oz., 1 gal.

Use: Analgesic, antipyretic & antirheumatic.

SALICYLIC ETHYL ESTER CARBONATE W/ Acetophenetidin, sod. salicylate, colchicum root pow., ext. See: Tetragen, Tab. (Endo)

SALICYLINE, (Kendall) Sod. salicylate 6 gr., w/Pot. Iodide, colchicum arm ext., Cimicifuga ext. & gelsamium ext./Tab. Bot. 100s, 500s.

Use: (1-2 q. 3 h.) Analgesic, antipyretic & antirheumatic.

SALICYLSALICYLIC ACID, N.N.R. 1948. See: Salytal, Tab. (Rora-Golen).

W/Aspirin. See: Persistin, Tab. (Sherman)

SALIGENIN. (Delta) Salicyl alcohol, Bot. 100 Gm.

W/Acetylcholine HCl. See: Acacoline, Amp. (Anglo-French).

W/Sulfathiazole & propylene glycol. See: Olozole, Liq. (Hort).

W/Limodin & Thio-dia-mer-sulfonamide. See: Trisulfacoline, Loz. (Central Pharm.)

SALINEXIN. (Abbott) (Hexamethylenamine, Acetaminosalicylic Acid Sod } Amp. (1 Gm.), 10 cc., 6s, 23s, 100s, Cap. 2 1/2 gr., Bot. 100s, 1000s; 5 gr., Bot. 100s.

Use: (5-10 gr. daily) Urinary antiseptic.

SALIMEPH-C. (Kremers-Urban) Salicylamide 250 mg., mephensin 250 mg., ascorbic acid 15 mg./tab. Bot. 100s, 500s, 1000s. Also available w/codaine phosphate 1/2 gr./tab. Bot. 100s. Also available w/calchicine 0.25 mg./tab. Bot. 100s.

Use: (1-2 t i.d.) Muscle relaxant w/analgesic.

SALIMEPH FOETE. (Kremers-Urban) Salicylamide 500 mg., mephensin 333 mg., ascorbic acid 50 mg./Tab. Bot. 100s, 500s, 1000s. Use: Analgesic.

SALIMEPH/PREDNISOLONE. (Kremers-Urban) Salicylamide 500 mg., mephensin 333 mg., ascorbic acid 50 mg., prednisolone 0.75 mg./Tab. Bot. 100s.

Use: Anti-arthritis.

SALINIDOL, N.N.D. 1962. (Doak) Salicylanil-

| Salicylic acid | Others | Manufacturer | Brand |
|----------------|--|----------------------|--------------------------------|
| 2% | Allantoin 0.2%, bith onol 1% | Dome | Domarine Shampoo |
| 2% | Benzoic acid 2%, amyli tritresols 0.1%, o hydroxyphenyl mercuro chlor 0.1% | Upjohn | Salicresin, Liq |
| 6% | Benzoic acid 6%, chloromercuri covacrol 1% | Smith Dorsey | Monarden, Oint & Pow |
| 26 gr./oz. | Benzoic acid 52 gr./oz., benzocaine 5 gr./oz. | Blue Line | Pylosin, Liq |
| 1% | Benzalc acid 2%, benzocaine 3/4%, chlorothymol 1/30% | P. Imou Moore | Ben Sal Coine, Oint. |
| 2 1/2% | Benzoic acid 2 1/2%, bor e acid thymol isopal | Dook | Mycofan, Oint |
| 6% | Benzoic acid 12%, chlorothymol 1/4% | Upjohn | Benzo Salicylic Com-pound Oint |
| 2% | Coal tar 10%, calamine 20% | Manhattan Eye Salve | Distaral Colamine, Oint |
| 3% | Coal tar sol 5%, sulfur 2.5%, zinc oxide 5% | Merck Sharp & Dohme | Topotar, Cream |
| 1.5% | Coal tar crude 0.6%, mercury oleate 0.25%, phenol 0.25%, p antrophenol 0.25% | Torch | Prosol, Emulsion |
| 3% | Coal tar crude 1.2%, mercury oleate 0.45%, phenol 0.5% | Torch | Esteral, Emulsion |
| 3% | Coal tar 5%, sulfur 3% | Ulmer | Sal Su Tar, Oint |
| 3% | Coal tar distillate 4%, sulfur 3% | Smith Kline & French | Pragmatar, Oint |
| 3% | Coal tar sol 5% | Lannett | Salatar, Cream |
| 1/2 gr | Magnesium salicylate 2 gr., strontium salicylate 2 gr., sod bicarbonate 3 gr., acetophenetidin 1/4 gr., w/methyl salicylate pancreatin, diastase | Jenkins Lobs | Foursalco, Tab. |
| 1.5% | Resorcinol monoacetate 1.5% | Almay White | Resorcitate, Sol. |
| 3% | Sulfur 3% | | Appliderm No 6, Oint |
| 2% | Sulfur 2%, Yt A 100 000 u/oz | Dome | Su A So |

- ide 5%, in carbowax 25%. Oint., Jar 4 oz., 1 lb., 5 lb.
- Use: Fungus infections of skin.
- SALIPHEN.** (Smith-Dorsey) Acetophenetidol 2 gr., aspirin 4 gr., Gelsemium 2/10 gr., caffeine 1/4 gr./Tab. Bot. 100s, 1000s.
- Use: (1-2 q. 3 h.) Analgesic & antipyretic.
- SALIPYRIN.** N.N.R. 1934. (Biedel) Antipyrine 58%, salicylic acid 42%. Pow. 1 oz., 3 oz., 1 lb. Tab. 7 1/2 gr., Bot. 20s, 50s, 500s.
- Use: (1-2 t.i.d.) Analgesic & antipyretic.
- SALISIL.** (Elder) Aspirin 5 gr., Mag. trisilicate 2 gr./Tab. Bot. 100s, 1000s.
- Use: (1-2 t.i.d.) Analgesic & sedative.
- W/Phenobarbital.** (Elder) Phenobarbital 1/2 gr., aspirin 5 gr., Mag. trisilicate 2 gr./Tab. Bot. 100s, 1000s.
- Use: (1-2 t.i.d.) Analgesic & sedative.
- SALI-ZEM No. 2.** (Zemmer) Colchicine 1/400 gr., Sod. salicylate 2 1/2 gr., Sod. p-aminobenzoate 2 1/2 gr., Vit. B-1 1 mg., B-2 1 mg., C 10 mg./Tab. Bot. 100s, 500s, 1000s.
- Use: 1-2 t.i.d.; rheumatic conditions.
- SALMONELLA DIAGNOSTIC SERA.** (Lederle) Sera to identify 98% of Salmonella infections in the U. S. Vial 1 cc. w/control.
- SALOL.** N.N.R. 1936. (Mallinckrodt) Phenyl salicylate, Cryst. Granular, Pow., Pkg. 1/4, 1 lb.
- Use: Analgesic & antipyretic.
- W/Acetanilid, acetophenetidin & caffeine.**
- See: Ferbrilal, Liq. (Tilden).
- W/Methanamine, atropine sulfate, hyoscyamine, benzal acid & methylene blue.**
- See: Meth-Atra-Mine, Tab. (Breon).
- W/Neomycin, kaolin, pectin, zinc sulfocarbamate.** See: Neo-Kaleton, (Burrington)
- SAL-NISOLONE.** (Ortego) Prednisolone 2 mg., potassium salicylate 0.28 Gm., potassium paraminobenzoate 0.20 Gm., ascorbic acid 20 mg./Tab. Bot. 50s, 100s.
- Use: Rheumatic fever, rheumatoid arthritis, gout.
- SALOPHEN.** N.N.R. 1948. (Winthrop Labs.) p-Acetamidophenyl salicylate; Phenacetol. Pow., Bot. 1 oz. Tab. 5 gr., Sat. 100s.
- Use: (1 t.i.d.) Analgesic.
- SALOSTAN.** (Patch) Tin oxide & lin. Tab. Bot. 75.
- Use: Tin therapy in treatment of boils & acne.
- SAL PABA.** (Evron) p-Aminobenzoate Sod. 324 mg., Sod. salicylate 324 mg., ascorbic acid 50 mg., thiamine 1 mg./E.C. Tab. Bot. 100s, 500s, 1000s.
- Use: 2 q.i.d.; antiarthritic.
- SAL PABA No. 2.** (Evron) Same, except Sod. salts replaced w/ammonium & Pot.
- SALPACINS.** (Vale) Sod. salicylate 5 gr., Sod. p-aminobenzoate 3 gr., calcichine 1/350 gr., Vit. C 15 mg./Tab. Bot. 100s, 500s, 1000s.
- Use: Analgesic in rheumatic disorders.
- SAL-PNENESIN.** (Marion Labs) Mephenesin 100 mg., ammonium salicylate 600 mg., p-aminobenzoic acid 2 gr./Tab. Bot. 100s.
- Use: (1-2 q. 2-3 h.) Arthritic conditions.
- SALPIX.** (Ortho) Sod. acetrizoate 530 mg., polyvinylpyrrolidone 230 mg./sc. Vial 6 & 10 cc.
- Use: Diagnostic agent.
- SALPRIN.** (Ewell) Salicylamide 3 1/2 gr., phenacetol 2 1/2 gr., caffeine 1/2 gr./Tab. Bot. 30s, 100s, 1000s, 5000s. Also avail. w/codaine.
- Use: (1 q. 3 h., m.p.) Analgesic.
- SALRAMIN.** (Normand) Sod. salicylate 5 gr., p-aminobenzoic acid 5 gr./Tab. Bot. 500s, 1000s.
- Use: (1-3 t.i.d.) Combating pulmonary tuberculosis.
- SALRIN.** (Warren-Teed) Salicylamide. Tab. 0.3 Gm., 0.6 Gm., Bot. 100s, 1000s.
- Use: (1-3 t.i.d.) Analgesic, antipyretic & antirheumatic.
- SALSET.** (Direct Labs.) Tabs. Bot. 100s
- Use: Muscle relaxant and sedative.
- SALSO-COL.** (Breon) Sod. salicylate 2.07 Gm., Sod. iodide 0.32 Gm., colchicine 0.65 mg., Sod. cocodylate 0.22 Gm./20 cc. Amp. 20 cc., Box 25s.
- Use: (5-10 cc.) Analgesic & antirheumatic.
- SALSUNATE.** (Drug Service) Aspirin 8 gr., ascorbic acid 20 mg., Cal. succinate 2 gr., succinic acid 3 gr./Tab. Bot. 100s, 1000s.
- Use: (1-2 q. 3 h.) Analgesic.
- SAL-SU-TAR.** (Ulmer) Salicylic acid 2%, sulfur 3%, coal tar ext. 5%, menthol. Oint. Jar. 1 oz., 1 lb.
- Use: Fungicide, bacteriostatic.
- SALTIPO.** (Sunif & Case) Sodium salicylate 325 mg., potassium iodide 65 mg., thiamine chloride 0.26 mg., ascorbic acid 11 mg., gelsemium 8 mg., simicifuga 8 mg./Tab. Bot. 100s, 1000s.
- Use: Analgesic and antirheumatic.
- SALT SUBSTITUTES.**
- See: Adolph's Salt Sub. (Adolph's)
- Co-Salt (Funk)
- Diasal (Faugera)
- Gustamate (Arlington)
- Neocortasal (Winthrop Labs.)
- SALUNDEK.** N.N.B. 1955. (Wallace & Tiernan) Not listed. 2% Undecylenic acid, 25% zinc undecylenate, 5% salicylanilide in carbowax base Oint., Tube 1 oz., Jar 1 lb.
- Use: Tinea capitis (ringworm) & athlete's foot.
- SALUNDEK, NEW.** (Maltbia) Zinchlorundesal.

- Salicylanilide 30 mg 5 chloro salicylanilide 20 mg, 5 3 & 5 4 dichlorosalicylanilides of each 10 mg undecylenic acid 2D mg zinc undecylenate 100 mg /Gm Oint Tube 2 oz; Jar 454 Gm
 Use: Tinea capitis
- SALUNDEK SOLUTION** (Maltbie) Salicylanilide 3% mono and dichloro salicylanilides 2%, undecylenic acid 2%, alcohol 55% Sol 3 oz
 Use: Treatment of ringworm of the scalp
- SALURON** (Bristol) Hydroflumethiazide 50 mg /Tab or 5 cc Tab Sol 50s Syr Bot 8 oz
 Use: Diuretic
- SALUTENSIN** (Bristol) Hydroflumethiazide 50 mg reserpine 0.125 mg, prochlorperazine A 0.2 mg /Tab Bot 60s
 Use: Antihypertensive
- SALUVIMS** (Pioneer) 12 vitamins, including B 12 activity 5 mcg, 12 minerals Tab Bot 80s
 Use: Comprehensive dietary supplement
- SALVAHIST**, (Columbus) 3% Pyranisamiso maleate, 10% calamine, 2% zinc oxide 2% resorcinol, eucalyptal phenol, boric acid & corn starch Oint, Tube 1 oz; Jar 1 lb
 Use: Antihistamine & antiseptic cream
- SALVARSAN**, N.N.R. 1947 See: Arspenamine (Various Mfr.)
- SALVITAL**, (Columbus) Resorcinol, phenol boric acid zinc oxide, eucalyptal, amyllum, balsam of peru Oint, Tube 1 1/2 oz; Jar 16 oz
 Use: Antipruritic.
- SALYDOL**, (Eastern) Sod aspirin, Sod salicylate acetic acid Sod acetate & glyceric Elixir 6 oz
 Use: Analgesic
- SALYPHEC**, (Rare Galen) Salysal 2.5 gr caffeine 1/2 gr, acetophenetidin 2 1/2 gr /Tab Bot 25s, 100s, 1000s
 Use: Analgesic.
- SALYROAN** See: Mersaly (Various Mfr.)
- SALYROAN SUPP.** (W Nitrop Labs) Salysyrgan 0.4 Gm./Supp Box 12s
 Use: Diuretic
- SALYSAL**, N.N.B. 1948 (Rare Galen) Salicylic ester of salicylic acid Tab 5 gr, Bot 50s 250s 1000s Pow Bot 1 oz, 1 lb
 Use: Analgesic.
- W/Caffeine & acetophenetidin See: Salyphec, Tab (Rare-Galen).
- SANDALWOOD OIL, ACTIVE PRINCIPLE**, N.N.R. 1941
 See: Arheol, Cap (Astler)
 Ganesan, Cap (Riedel)
- SANDOPTAL**, N.N.R. 1952 [Sandoz] Not listed Allylbarbituric acid, isobutylallyl barbituric acid
- W/Aminopyrine caffeine See: Optalidan Tab (Sandoz)
- W/Caffeine, aspirin & acetophenetidin See: Fiorinol Tab (Sandoz)
- W/Dihydroergotamine methanesulfonate scopolamine, barbiturates See: Plaxonal, Tab (Sandoz)
- SANDOPTAL SODIUM** W/Barbital Sod., phenobarbital Sod., scopolamine HBr, dihydroergotamine methanesulfonate (DHE 45) See: Plaxonal, Tab (Sandoz)
- SANDOSTENE**, (Sandoz) (Withdrawn) Methyl amino phenyl phenyl piperidine tartrate Thinalidine Tab (25 mg) Bot 100s Space-labs 75 mg Bot 100s
 Use: Antipruritic & antihistaminic therapy
- SANDOSTENE** W/Calcium (Sandoz) (Withdrawn) Amp (50 mg w/10% cal gluconate) 10 cc, Box 5s, 20s
 Use: I.V., Anti-allergic and antipruritic
- SANDRIL**, N.N.D. 1962 (Lilly) Reserpine 0.1 mg 0.25 mg, 1 mg Bot 100s, 1000s Tab 5 mg Bot 30s, 500s Elixir (0.25 mg / 5 cc) Bot 1 pt 1 gal Amp (2.5 mg /cc) (5 mg /cc) 10 cc Dropt (2 mg /cc) Bot w/dropper 30 cc
 Use: Hypotensive & tranquilizing agent
- W/Pyroell (Lilly) Reserpine 0.25 mg, pyrambutamine 7.5 mg /Tab Bot 100s 1000s
 Use: To relieve nasal stuffiness in reserpine (hypertension) therapy
- SANICYL CREAM** (Parke, Davis) Salicylanilide 5 Gm, phenol chloride 1.25 Gm / 110 Gm Tube 2 oz Jar 1 lb, 5 lb
 Use: Ringworm
- SANLUOL**. See: Arspenamine (Various Mfr.)
- SANMETTO**, (OD Peacock) Saw palmetto sandalwood, corn silk & alcohol Bot 8 & 8 oz
 Use: (1 2 tsp q 4 h) Urinary sedative
- SAN-NAUS** (Lemmon) Hyoscine HBr 0.4 mg, Phenyltolaxamine d hydrogen citrate 25 mg, Pyridoxine HCl 10 mg, Methamphetamine HCl 1.25 mg, /Tab Bot 100
 Use: Prevention nausea and vomiting of pregnancy
- SANSIRT**, (Sandoz) Methysergide Maleate 2 mg /Tabs Bot 50s
 Use: Vascular headaches
- SANTA** (Santa) Vit A 5000 u, D 1000 u, B-1 1 mg, B-2 1 mg, B-6 1 mg, C 50 mg, alcatnamide 10 mg, panthenol 3 mg / D 6 cc Bot 3D cc, 50 cc.
 Use: Vitamin supplement for infants and some adults
- SANTALOL ALCOHOL**. See: Ganesan, Cap (Riedel)

- SANTALOL SALICYLATE.** See: Santyl (Knoll)
- SANTAMINIC SYRUP.** (Santo) Pyrilamine maleate 50 mg., di-desoxyephedrine HCl 2 mg., potassium guaiacol sulfonate 8 gr., ammonium Cl 8 gr., tartar emetic 1/12 gr., chloroform 2 min./fl. oz. Bot. pt., gal.
Use: Expectorant for coughs.
- SANTA-THIRA.** (Santo) Vitamin A 30,000 u., D 1000 u., K 2 mg., B-1 12 mg., B-2 12 mg., B-6 0.5 mg., B-12 10 mcg., folic acid 0.1 mg., niacinamide 100 mg., d-calcium pantothenate 5 mg., C 150 mg., E 5 mg., Ca 21 mg., P 16 mg., I 0.15 mg., Fe 15 mg., K-sulfate 5 mg., Co-sulfate 0.1 mg., Cu-sulfate 1 mg., Mn-sulfate 1 mg., Mg-sulfate 6 mg., Mo 0.2 mg., Zn-sulfate 1.5 mg./Cap. Bot. 100s.
Use: Vitamin deficiency in postoperative, postpartum and after serious illness.
- SANTOL OIL.** (Merck Sharp & Dohme) Cap. 5 min., 10 min., Bot. 100s.
Use: Treatment of dysuria.
- SANTONIN, N.F.** Tab., N.F. (Mal-linckrodt) Cryst. & Pow. Bot. 1/8 oz., 1/4 oz., 1 oz.
Use: Anthelmintic.
- SANTONIN & CALOMEL.** (Merck Sharp & Dohme) Santonin 30 mg., calomel 30 mg./tab. Bot. 100s.
- W/Cascara sagrada.** (Winthrop Labs.) Santonin 1 gr., cascara sagrada 20 gr./fl. oz. Bot. 2 oz.
Use: 1/4 to 1 tsp. b.i.d. for 3 days) Ascari-cide w/laxative.
- SANTONIN & MILD MERCUROUS CHLO-RIDE TABLETS, N.F.** (Various Mfr.) See: Santonin & Calomel. (Merck Sharp & Dohme)
- SANTYL, N.N.R. 1941.** (Knoll) Santalol salicylate. Cap. 6 min., Box 30s. Vial 15 cc.
Use: Urinary antiseptic.
- SAPONATED CRESOL SOLUTION, N.F.** See: Cresol Compound (Lilly)
- SAPONINS, WATER SOLUBLE.** See: Sas-par, Tab. (Bischoff)
- SABATOGA OINTMENT.** (G. F. Harvey) Boric acid, zinc oxide, eucalyptol, prepared w/et & white petrolatum. Tube 1, 2 & 4 oz. Can 1 & 5 lb.
Use: Skin irritations & protective.
- SARCOPEN LIQUID.** (Dome) Bot. 4 oz.
Use: Treatment for psoriasis of the scalp.
- SARINOID-3.** (Dumas-Wilson) Citrus bio-flavonoids 100 mg., Vit. C 100 mg., K 0.75 mg., prednisolone 5 mg./Cap. Bot. 50s.
Use: Antiinflammatory.
- S-A-VITE.** (Ayerst) Vit. A 25,000 u., D 1000 u., B-1 25 mg., B-2 12.5 mg., Nicotinamide 50 mg., Vit. B-6 3 mg., calcium pantothenate 10 mg., C 250 mg., B-12 5 mcg., intrinsic factor concentrate 15 mg / sustained action tab. Bot. 60s, 500s.
Use: Therapeutic multivitamin.
- SAVOBETS, Penicillin W/Sulfas.** (Lilly) Pot. penicillin G. 50M u. or 100M u. w/sulfa-diazole, sulfamerazine & sulfamethazine 83 mg. each/Tab. Bot. 24s, 100s.
Use: Orally; penicillin & triple sulfanamide therapy.
- SAWRILCO.** (Sulliff & Cose) Saw palmetto extract 65 mg., rhus aromatica 65 mg., aquilegium 65 mg., corn silk extract 32.5 mg., hyoscyamine 0.03 mg., belladonna extract 1.3 mg./Tab. Bot. 1000s.
Use: Antispasmodic and diuretic.
- SAXIN.** (Burroughs Wellcome) Saccharin 1/4 gr./Tab. Bot. 200s, 1000s.
Use: Sweetening agent.
- SAXOL.** See: Petrolatum Liquid (Various Mfr.)
- SAYOLIN.** (Elder) Phenobarbital 1/4 gr., hyoscyamus ext. 1/2 gr., camphor mono-bromated 1 gr., passiflora 2 gr./Tab. Bot. 100s, 500s
- SCABINZATE.** (Hort) Benzyl benzoate 30% w/soap and alc. Bot. 4 oz., 1 pt., 1 gal.
Use: Scabies and pediculosis
- SCABICIDES, N.N.R. 1953.**
See: Benzyl Benzoate (Various Mfr.), Eurax Cream (Geigy), Pyrethrum. (Upsher-Smith) Rotenone. (Various Mfr.) Sulfur (Various Mfr.)
- SCARLET FEVER STREPTOCOCCUS TOXIN, N.N.R. 1950.**
Lederle—Dick Test, Vial 2 cc., 5 cc.; Im-munization, Vial 1 cc.
National Drug—Disk Test, Vial 2 cc., 11 cc.
Parke, Davis—Dick Test, Vial 2 cc., 10 cc.; Im-munization, Vial 0.1 cc., 1 cc.
Wyeth—Dick Test, Vial 5 & 50 doses; Im-munization, 1 & 10 doses.
- SCARLET FEVER CONVULSANT SERUM, N.N.D. 1958.** See: Scarlet Fever Immune Serum, Hyman (Milwaukee Blood Center)
- SCARLET FEVER IMMUNE SERUM, HU-MAN, N.N.D. 1960**
Jr. League Blood Center, Milwaukee—Vial 10 cc., 20 cc.
Michael Beese Research Found.—Vial 10 cc., 20 cc.
- SCARLET RED, N.F.** Oint., N.F. o-Tolyl-azo-o-tolyl-azo-beta-naphthal.
Merck Sharp & Dohme—Bot. 1 oz., 1/4 lb., 1 lb.
Parke, Davis—Oint. 5%, Tube 1 oz., Jar 1 lb., 5 lb.
Use: To stimulate epithelial cell growth.

- SCHAMBERG'S LOTION.** (C & M Pharm) Menthol D 15%, phenol 1 D%, zinc oxide, peanut oil, lime water and dispersant Bot. pt
Use Pruritic type eczema
- SCHEDULIN.** (Louisiana Ethical Pharm) Cas anthranol 25 mgm, diethyl sodium sulfosuccinate 50 mgm / Cap Bot 25s, 100s, 500s.
Use Laxative
- SCHICK TEST.** See: Diphtheria Toxin, Diagnostic, Inactiv, U S P
- SCHICK TEST CONTROL.** (Lilly) Diphtheria Toxin, Diagnostic
- SCHICK TEST TOXIN.** See: Diphtheria Toxin, Diagnostic, U S P
- SCHLESINGER'S SOLUTION.** See Morphine HCl & Scopolamine Compound (Merck Sharp & Dohme)
- SCILLAREN.** N N R 1952 (Sandoz) Mix of glycoside scillaren A & B Amp (0.5 mg / cc) 1 cc, Box 6s, 12s, 50s Liq 0.5 mg / cc, Bot 20 cc, 200 cc Supp 1 mg, Box 10s, 30s Tab DS mg, Bot 100s, 250s, 1000s
Use See Digitalis
- SCINOCORT.** (Testogor) Hydrocortisone acetate 50 mg, neomycin sulfate 5 mg / cc
Use Anti-inflammatory
- SCLEROSINO AGENTS.**
See: Etalate (Forka, Davis)
Etholote (Metropolon)
Invert Sugar (Various Mfr)
Menolate (Abbott)
Proliferol B (Ulmer)
Proliferol T (Ulmer)
Quinine & Urethol Injection (Lilly)
Quinine & Urea HCl (Various Mfr)
Sodium Morrhuate N N R (Various Mfr)
Soricin (Merrell)
Solradecal, (Wallace & Tiernan)
Sylnasol (Searle)
Varisol (Abbott)
- SCOLAMINE.** (Nyca Labs) Genoscolamine HBr Hyoscine aminoxide HBr 1 mg / Tab TD Bot 100s
Use: Antispasmodic
- SCOLATE.** (Central) Butabarbital 15 mg, methscopolamine nitrate 1 mg, atropine methyl nitrate 1 mg / Tab Bot 100s
Use: Antispasmodic
- SCOLEN UHICELLES.** (Hiss Pharm) Hyoscine aminoxide HBr 1 mg (Scopolamine Aminoxide HBr) Cap Eat 100s
Use: Delayed action, motion sickness
- SCOLINE** (Kendall) Methscopolamine bromide 2.5 mg / tab Bot 100s
Use: Antispasmodic
- SCOLINE-AMOBARBITAL.** (Kendall) Methscopolamine bromide 2.5 mg, amobarbital 16.2 mg / tab Bot 100s
Use: Antispasmodic w/sedative
- SCOPAMAL.** (E. S. Miller) Scopolamine methylbromide 1.5 mg, phenobarbital 8 mg, Cal. carb 200 mg, Mag trisilicate 200 mg, Mag oxide 200 mg / Tab. Bot 100s, 1000s
Use: Anticholinergic & antacid
- SCOPATRATE.** (Deseret) Scopolamine methyl nitrate 5 mg, phenobarbital 32 mg / Tab. Bot 100s
Use: Antispasmodic w/sedative
- SCOPAN.** (Dumas Wilson) Tab Bot 100s, 1000s
Use: Psychosomatic relief
- SCOPHENESIN, TABLET.** (Nelson Bayer) Mephobarbital 1/4 gr, mephenesin 3/4 gr, genoscolamine 1/640 gr, betaine HCl 1 gr Bot 100s
Use: (2 11 d. p c) Antispasmodic
- SCOPODEX.** (Fougero) Scopolamine aminoxide HBr genoscolamine HBr Pallet 0.5 mg, Box 20s; Bot 100s
Use: (2 p r n) Motion sickness
- SCOPOLAMINE.** (Hyoscine; Scopolamine) Epoxytropine Iropote
Use: As scopolamine hydrobromide
- SCOPOLAMINE AMINOXIDE.** See Genoscolamine
- SCOPOLAMINE AMINOXIDE HBr W/Methapyrilene HCl.** See Nite Rest, Cop (APC)
- SCOPOLAMINE CAMPHORATE W/Hyoscine amine Camphorate.** See Vosono Tab (Schering)
- *SCOPOLAMINE HYDROBROMIDE, U.S.P.** N N R 1949, In] U S P Tab, N F (Hyoscine HBr) 1 Scopolamine HBr
Burroughs Wellcome—Amp (1/200 gr) 1 cc, (1/150 gr) 0.5 cc, (1/130 gr) 1 cc, (1/100 gr) 1 cc Box 100s
Merck Sharp & Dohme—Cryst or Pow
Viol 5 gr, 15 gr, Bot 1/2 oz Tab 1/150 gr, 1/100 gr, Bot. 100s, 1000s
Use: Hypnotic & sedative.
- Scopolamine Hydrobromide Combination.** See Hyoscine HBr
- SCOPOLAMINE METHOBROMIDE.** See Methscopolamine Bromide, Preps
- SCOPOLAMINE METHYLBROMIDE.** See Methscopolamine Bromide, Preps
- SCOPOLAMINE METHYL NITRATE.** See Methscopolamine Nitrate
- SCOPOLAMINE SALTS.**
See Hyoscine salts
- SCOPOMAL.** (E. S. Miller) Scopolamine methylbromide 1.5 mg, cal carb 200 mg

- mg. oxide 200 mg., mag. trisilicate 200 mg., phenobarb. 8 mg./Tab. Bot. 100s
 Uses: (3 to 9 w/meal) Antispasmodic
- SCORBACAINI.** (Physiological Chemicals)
 Procaine ascorbate 1% sol. Amp. 1%, 5 cc., Box 10s, 25s, 100s.
 Uses: Local anesthetic.
- SCRIPAC.** (Armour) Thyroid. Tab. 1 gr., Pkg. 30s.
 Uses: Thyroid therapy.
- SCUBENALINE.** See: Epinephrine. (Various Mfr.).
- SCUROFORME.** See: Butyl Aminobenzoate (Various Mfr.).
- SEBACTIN.** (Summers) Sulfur 3%, salicylic acid 2%, hexachlorophene 1%. Cream, Jars 4 oz. Liq. Plastic Bot. 4 R. oz.
 Uses: Treatment of seborrhea.
- SEBA-NIL.** (Texas Pharm.) Cleansing mixture of alcohol, acetone, sorbitan monofaurate & water. Bot. 8 oz., 1 pt., 1 gal.
- SEBAQUIN SHAMPOO.** (Summers Labs.) Diiodohydroxyquin, U.S.P. 3%, in a neutral, soapless detergent, lathering base with hypallergenic lanolin derivatives./Liq. Bot. 4 oz. plastic.
 Uses: Anti-seborrheic.
- SEBABORB LOTION.** (Summer Labs.) Activated colloidal attapulgit 10%, zinc oxide 5%, colloidal sulfur 5%, resorcinol 2%, hexachlorophene 1%, isopropanol 20%. Bot. 2 oz.
 Uses: Acne.
- SEBELLA.** (Wyeth) Phenobarbital ½ gr., helladonna ext. ½ gr., aluminum hydroxide 5 gr. Tab. Bot. 100s, 1000s.
 Uses: (1-2 t.i.d.) Antacid w/antispasmodic & sedative.
- SEBICAL CREAM.** (Reed & Cornick) Allantoin 2%, Coal tar extract 2%, Hexachlorophene 1% in a greasless vanishing base. Tubes 2 oz.
 Uses: Scalp Dermatoses.
- SEBICAL SHAMPOO.** (Reed & Cornick) Allantoin 0.2%, Hexachlorophene 1%, Special coal tar extract (Tarbanis) in a detergent cream base Tubes 2 oz.
 Uses: Relief of itching and scaling.
- SEBIZON LOTION.** (Schering) Sulfacetamide Sod. 100 mg., methylparaben 1 mg./Gm. Bot. 4 oz., plastic squeeze.
 Uses: Scaling dermatoses
- SEBULEX LIQUID.** (Westwood) Bot. 4 oz.
 Uses: Treatment of itchy scaly scalp.
- SEBULEX SHAMPOO.** (Westwood) Sod. sulfocacetate, Sod. alkyl aryl polyether sulfonate, Sod. dioctyl sulfosuccinate, Keratin.
- hydic, sulfur 2%, salicylic acid 2%, hexachlorophene 1%. Bot. 4 oz.
 Uses: Dermatologic shampoo.
- BECADERM.** (Thompson) Benzocaine 2%, colamine 5%, pyrilamine moleate 0.5%, phenol 0.5%, menthol 0.3%. Oint. Tube 1 ½ oz., Jar 2 oz.
 Uses: Topically, antiseptic & antihistaminic.
- *SECOBABBITAL.** U.S.P., N.N.D. 1961, A.D.B. (Various Mfr) Elixir U.S.P. 5-allyl-5-(1-methylbutyl)-barbituric acid. Paw., Bot. ½ oz., 1 oz. See: Seconal, N.N.R. (Lilly)
- Secobarbital Combinations** on pp. 658 and 659.
- SECOBARBITAL ELIXIR.** See: Seconal Elixir (Lilly)
- *SECOBARBITAL SODIUM.** U.S.P., N.N.D. 1961, A.D.E.; Cap. U.S.P., N.N.O. 1958. Sod. 5-allyl-5-(1-methylbutyl)-barbiturate.
 Hance—Cap. 1 ½ gr., Bot. 100s, 1000s.
 Keith-Victor—Cap. 0.1 Gm. Bot. 100s.
 Lannett—Cap. ¾ gr., 1 ½ gr., Bot. 100s, 500s.
 Massengill—Cap. 1 ½ gr., Bot. 100s, 5000s.
 Raymer—Cap. ¾ gr., 1 ½ gr., Bot. 500s, 1000s.
 See: Barbosec, Cap. (Rowell).
 Evronal Sodium, Cap., N.N.D. (Evron).
 Seconal Sodium, N.N.O. (Lilly)
 Synate (Central)
- SECOBARBITAL SOL.** N.N.D. 1961. See: Secobarbital sodium.
- BECODEX.** (Nyco Labs.) d-Amphetamine sulf. 15 mg., secobarbital 60 mg./Tab. Bot. 100s.
 Uses: C.N.S. stimulant w/sedative.
- SECOBIN.** (Prama) Secobarbital Sod. 50 mg., d-methamphetamine HCl 5 mg./Tab. Bot. 100s, 1000s.
 Uses: (1 p.c. t.i.d.) Control of obesity.
- SECOBIN ENCORE.** (Prama) Secobarbital 60 mg., methamphetamine HCl 10 mg./Tab. Bot. 30s, 100s
 Uses: Delayed action, treatment of mental distress
- SECONAL.** N.N.D. 1961. (Lilly) Secobarbital Elixir 2 gr./oz., Bot. 1 pt., 1 gal.
 Uses: Sedative.
- SECONAL SODIUM.** N.N.D. 1961. (Lilly) Secobarbital Sod.
 Ensat: (½ & 1 ½ gr.) Bot. 100s, 500s.
 Pow.r Bot. ½ oz.
 Pulvule: (½, ¾ & 1 ½ gr.) Bot. 100s, 500s
 Supp.s (½, 1, 2 & 3 gr.) Bot. 12s, 144s
 Vial: (0.25 Gm.) in 50% polyethylene glycol/5 cc., 6s, 25s, 100s (50 mg/cc) 20 cc.
 Uses: Hypnotic & sedative.
 W/Amlytal Sod. See: Tuinal, Cap. (Lilly).
 W/Ephedrine sulfate. See: Ephedrine & Seconal Sodium, Cap. (Lilly).
- * Generic name recognized by N.F., U.S.P. or A.M.A.

Secobarbital Combinations

| Secobarbital | Others | Manufacturer | Brand |
|---|---|----------------|------------------------------------|
| (Sod) 3/4 gr 60 mg | Amobarbital Sod 3/4 gr | G F Harvey | Dusajal, Cap |
| 35 mg | d Amphetamine sulf 15 mg | Nysco Labs | Secodax, Tab TD |
| 45 mg | d Amphetamine tartrate 17.5 mg | Nessler | Seco Synotan, Tab |
| (Sod) 3/4 gr | d Amphetamine sulf 15 mg | Webster | Sedadex, Cap |
| (Sod) 1/2, 3/4, 5/8, 3/2 gr | Aspi " | Webster | Aspirin Secobarb, Supp |
| 50 mg | Aspirin 1, 3, 5 gr | Webster | Aspirin Secobarbital, Supp |
| (Sod) 8 mg | Aspirin, Vit B-1 | Patch | Ambral, Tab |
| 1 1/2 gr | Aspirin 300 mg, phenacetin 150 mg, caffeine 30 mg, amobarbital 8 mg, buffered w/aluminum & magnesium hydroxides | Lemmon | Buffodyne W/Barbita rates, Tabs |
| (Sod) 1/4 gr 50 mg | Butobarbital 1/2 gr | Hiss Pharm | Butasec Unicefles, Tab. |
| (Sod) 15 mg | Butobarbital Sod, 1/4 gr, phenobarbital 1/4 gr | Bowman | Buiseco, S CT, Tab |
| (Sod) 1 1/2 gr, 1/2 gr | Butobarbital 30 mg, phenobarbital 50 mg | Lincoln | Duroform Tab. |
| (Sod) 1/2 gr 15 mg | Butobarbital Sod 15 mg, phenobarbital Sod 15 mg, pentobarbital Sod 15 mg | C D Smith | Quadra Sed, Sol |
| 25 mg | Diethylbarbituric acid 3/4 gr | Ulmer | Barlwo, Tab. |
| 30 mg | Ephedrine 3/4 gr, theophylline 2 gr, isoproterenol 5 mg | Hart | Alured Tab |
| (Sod) 1/2 gr 15 mg | Hametrapine methylbromide, Cal carb, charcoal | Rowell | Via Aika Phen, Cap |
| 25 mg | Hydrocyanic sulfate 0.25 mg | Kremer Urban | Levsin sulfate W/Seco-barbital Tab |
| 30 mg | Mannitol hexanitrate 30 mg, aminophyllin 90 mg, Pot Iodide 60 mg, phenobarbital 8 mg, rutin 10 mg, Vit C 25 mg, aluminum hydroxide 60 mg | Genetric | Manophyllin, Tab. |
| (Sod) 1/2 gr (Sod) 1/2 gr 15 mg | Mephenezin 400 mg | Crookes Barnes | Secanetin, Tab |
| (Sod) 50 mg (Sod) 15 mg (Sod) 100 mg 50 mg 6.075 mg | Mephenezin 6 gr | Hoack | Haxsen, Tab |
| | Mephenezin 5 gr., homatropine methylbromide 1/20 gr | Rowell | Via Phen, Cap. |
| | Mephenezin 180 mg, phenobarbital 8 mg, solicylic Sod 300 mg, Vit C 50 mg, aluminum hydroxide 60 mg | Genetric | Gerisaf, Tab |
| | d Methamphetamine HCl 5 mg | Prema | Secodrin, Tab |
| | N traglycerin 0.4 mg, penterythritol tetranitrate 15 mg | Lemmon | Nitrosed, Tabs |
| | Pentylene tetrazol 300 mg | Strong Cobb | Tetrasecobarbital, Cap. |
| | Pentacerythritol tetranitrate 30 mg | Grant Chem | Caravas, Tymcaps |
| | Pentobarbital sod 6.075 mg, butobarbital sod 2.025 mg, phenobarbital 2.03 mg, belladonna pow ext 41 mg, phenyltoloxamine 20 mg., ethoverine HCl 12.5 mg | Armaur Labs | Tri Synar Plus, Tab |

Secobarbital Combinations

| Secobarbital | Others | Manufacturers | Brand |
|---------------|--|---------------|-----------------|
| (Sod.) ½ gr. | Phenobarbital ½ gr. | Thompson | Delonal, Tab. |
| 50 mg. | Phenobarbital 50 mg., aspirin 3 gr., Vit. B-1 5 mg. | Patch | Ambrol, Tab. |
| (Sod.) 50 mg. | Phenobarbital 50 mg., butobarbital Sod. 30 mg. | Wyeth | Ethabral, Cap. |
| (Sod.) ½ gr. | Pentobarbital Sod. | Armour | Nidar, Tab. |
| (Sod.) 30 mg. | Phenobarbital 30 mg., pentobarbital Sod. 30 mg. | Taylor | Sorbiclin, Tab. |
| (Sod.) 30 mg. | Phenobarbital 15 mg., methamphetamine HCl 5 mg. | Massengill | Hypnorm, Pill |
| 60 mg. | Phenyltolazamine citrate 25 mg., eucalyptol 75 mg., inner core | Wampole | Hypiron, Tab. |
| 20 mg. | Synaphylate 360 mg., racephedrine HCl 30 mg., Pot. iodide 300 mg., niacinamide 40 mg. | Central | Synata, Tab. |

SECONESIN. (Crookes-Barnes) Mephensin 400 mg., secobarbital 30 mg./Tab. Tab., Bot. 100s.

Uses: (1-2 t.i.d.) Relaxant & sedative.

SECOBATE. (Kirk) Vit. C. Amp. (100 mg./cc.) 2 cc., (500 mg./cc.) 5 cc., 6s, 25s, 100s; (1000 mg./cc.) 10 cc., 6s, 25s, 100s.

Use: Vitamin C deficiencies.

SECO-SYNATAN. (Neisler) d-Amphetamine tartrate 17.5 mg., secobarbital 35 mg./Tab. Bot. 100s.

Use: Treatment of nervous tension and mental distress.

SECOTRESS. (Wampole) Not listed. Eserpina 0.1 mg., methamphetamine HCl 2.5 mg./tab Bot. 100s.

Uses: (1-2 b.i.d.) Management of psychological & psychosomatic reactions to stress.

SECBAN CAP. (First Texas) Vit. B-12 10 mcg., ferrous sulfate 195 mg., folic acid 0.25 mg., dicalcium phosphate 227 mg., Vit. C 15 mg., B-6 1 mg., B-1 3 mg., nicotinamide 10 mg./Cap. Bot. 100s, 500s, 1000s.

Uses: (3-6 daily) Anemias.

SECRAN LIQ. (First Texas) Vit. B-12 10 mcg., B-1 3 mg., nicotinamide 10 mg., Cal. pantothenate 2 mg., Vit. B-6 1 mg./5 cc. Bot. 1 pt., 1 gal.

Use: (2-4 tsp. daily) Anemias.

SECREMOL. (Patch) Ammonium chloride 15 gr., lobelia fl. ext. 6 min., Sod. citrate, citric acid, tartaric acid, aerosol O.T./oz. Bot. 1 pt., 1 gal.

Use: (1-2 tsp. p.r.n.) Expectorant.

SECTIN. (Teitagar) Soybean lecithin, choline, inositol & other phosphatides, especially cephalin/Cap. Bot. 100s, 250s, 1000s.

Use: Dietary supplement.

SID. (Leyden Labs.) Butobarbital sodium 15 mg., pentobarbital sodium 15 mg./Cap. Bot. 100s.

Use: Sedative.

SEDACO. (Strassenbergh) Ethylmorphine HCl ½ gr., chloroform 2 gr., ammonium chloride 4 gr. w/terpin hydrate, talu syr., wild cherry syr./fl. oz. Bot. 4 fl. oz., 1 pt., 1 gal.

Use: (1 tsp. q. 3 h.) For coughs.

SEDACOF. See Semcap, Prep (Massengill).

SEDAEIN. (Smith-Dorsey) Codeine phosphate 1 gr., Sod. citrate, Pot. gealacal sulfonate, citric acid, senega fl. ext. 1½ min., menthol/fl. oz. Syr. Bot. 1 pt., 1 gal.

Use: (1 tsp. q.i.d.) Sedative expectorant.

SEDADEX. (Novasol) Secobarbital ½ gr. & d-amphetamine sulf. 3 mg./Cap. Bot. 50s

Use: Premedication in dentistry

SEADROPS. (Walker) Phenobarbital ½ gr., pentobarbital Sod. 1/16 gr., homo-

atropine methylbromide 1/480 gr /10 drops
 Sol 15 cc w/dropper, 30 cc, 1 pt
 Use Sedative & hypnotic

SEDAFORM See: Chlorobutanol (Various Mfr)

SEDALCO. (Mallard) Phenobarbital ¼ gr,
 Sod bromide 4 gr, Pot bromide 3½ gr,
 Hyoscyamus Ir 10 min /Tab Sol 100s
 500s 1000s
 Use: Sedative

SEDALOID (Horton & Converse) Phenobarbital ¼ gr, hyoscyamine sulfate 0.103 mg
 atropine sulfate 0.02 mg hyoscyine HBr
 0.006 mg /Tab Bot 100s 500s 1000s
 Use Sedative & antispasmodic

SEDAMASAL (Finley) Amobarbital 15 mg,
 salicylamide 200 mg phenocetin 120 mg
 d amphetamine sulfate 2 mg hyoscyamine
 sulfate 0.05 mg /Cap Bot 100s
 Use Analgesic

SEDAMYL. (Schenley) Acetylcarbromal 4 gr
 /Tab Tube 20s Bot 100s, 1000s
 Use: Sedation without hypnosis

SED-A-NAT (National Drug) Phenobarbital
 ¼ gr, hyoscyamus ext ½ gr, passiflora
 ext ¼ gr, valerian ext ¼ gr /Tab Bot
 100s, 1000s
 Use Hypnotic and sedative

SEDANS (Parke Davis) Black How 60 gr
 hydrostine represent ng 30 min fl ext
 hydrastis, jamaica dogwood 30 gr, W/
 aromatics and alcohol 28% Sol 1 pt
 Use Relief in painful menstruation

SEDAPHEN (C D Smith) Phenobarbital Sod
 20 mg, Sod bromide 300 mg, Pot bromide
 200 mg, Cal bromide 100 mg /5 cc Bot
 4 oz, 1 pt, 1 gal
 Use Sedative

SED-A-PINE. (Smith Darsey) Codeine sulfate
 3/16 gr, squil Ir 20 min, ipecac Ir B
 min, acid phosphoric, diluted 16 min /R
 oz Syr, Bot 1 pt, 1 gal
 Use: (1 tsp q 3 h) Expectorant & sedative

SEDAPLEX ELIXIR (Walker) Phenobarbital
 ½ gr, pentobarbital Sod 1/16 gr, homo
 atropine methylbromide 1/480 gr, alcohol
 18% v/v/4 cc. Bot 4 oz, 1 pt
 Use Daytime sedation & simple insomnia

SEDAPLEX (TABLETS), (Walker) Phenobarbital ½ gr, pentobarbital Sod 1/16 gr,
 homatropine methylbromide 1/480 gr /Tab
 Bot 100s
 Use: Daytime sedation

SEDASPASM (Metropolitan) Phenobarbital
 ¼ gr, hyoscyamine sulfate 0.1040 mg
 atropine sulfate 0.0195 mg, hyoscyine HBr
 0.0065 mg /Tab Tab white grooved, Bot
 100s, 1000s 5000s
 Use: Antispasmodic.

SEDATAMIN, (C D Smith) Pentobarbital
 Sod 50 mg, phenobarbital sod 15 mg
 thiamine HCl 10 mg /Cop Bot 100s, 1000s
 Use: Sedative

SEDATIVES

See Bromides

Barbiturates

Non Barbiturates

Carbamide (Urea) Compounds

Chloral Hydrate Preparations

Chlorobutanol

Chlartan (Wampoles)

Doriden (Ciba)

Dormisan (Schering)

Naludor (Roche)

Paraldehyde

Pentaerythritol Chloral, Caps (Ives
 Cameron)

Persicor (Ives Cameron)

Placidyl (Abbott)

Sominat (National Drug)

Ultron (Lilly)

Valmed (Lilly)

Viadril (Pfizer)

SEDATOLE (Merck Sharp & Dohme) Codeine
 sulfate 30 mg /30 cc w/wild cherry bark
 sanguinaria squill balm of gilead buds
 phosphoric acid diluted & menthol Bot
 4 oz, 1 pt, 1 gal
 Use Sedative expectorant

SEDATUSSIN (Lilly) Cephaeline HCl 1/30
 gr, Sod benzoate 4 gr, sanguinaria 3¼
 gr, squill syr 48 min aromatics/R oz
 Bot 1 pt, 1 gal
 Use (1 tsp q 3 h) Coughs

SEDAYSON. (Tailby Nason) Acetylcarbromal
 150 mg bromisovalum 100 mg /tab Bot
 100s 1000s
 Use (2 ts d) Sedative & hypnotic

SEDEVAL See Barbitol (Various Mfr)

SEDEX (Nysco Labs) Pentobarbital 60 mg
 phenobarbital 30 mg, d-amphetamine sulf
 5 mg /tab T D Bot 100s
 Use CNS stimulant w/sedative

SEDINYL (C D Smith) Ethinyl estradiol 0.03
 mg, phenobarbital Sod 15 mg /5 cc liq
 Bot 4 oz, 1 pt, 1 gal
 Use (1 tsp once or twice daily) Estrogen
 therapy

SEDLIN UNODAY Ph. 754 (Paul Maney)
 Phenobarbital ¾ gr, hyoscyamine HBr 0.54
 mg, homatropine methylbromide 0.0447
 mg, ecopalamine HBr 0.0144 mg thiamine
 HCl 3 mg /cop Bot 100s
 Use: Sedative spasmolytic agent

SEDOBARS (Whitt er) Phenobarbital ¼ gr,
 pentobarbital Sod ½ gr /Tab Bot 100s
 1000s
 Use: (1 pr n) Sedative

SEDOBARS No 2 (Whitt er) Phenobarbital

- $\frac{1}{2}$ gr., pentobarbital sod. 1 gr./Tab. Bot. 100s, 1000s.
Use: Sedative.
- SEDOCRINE.** (Mallard) Thyroid $\frac{1}{4}$ gr., phenobarbital $\frac{1}{4}$ gr., Hyoscyamus tr. 10 min./Tab. Bot. 100s, 500s, 1000s.
Use: Thyroid therapy w/sedative.
- SED-O-ET.** (Drug Products) Phenobarbital 15 mg., Hyoscyamus pow. ext. 8 mg., pasciflora 60 mg., valerian 60 mg./Cap. Bot. 100s, 1000s.
Use: (1 q. 3 h.) Sedative.
- SEDORMID.** (Roche) Allyl-isopropyl-acetyl-carbamide. Pow., Bot. 1 oz. Tab. 4 gr., Box 10s; Bot. 100s.
Use: Sedative.
- SEDORZYL.** (Wampole) Ephedrine sulfate $\frac{1}{4}$ gr., phenobarbital $\frac{1}{8}$ gr., organidin 10 mg./dr. Bot. 1 pt.
Use: Cough.
- SED-YENS.** (Lemmon) Amobarbital 50 mg., homatropine methylbromide 7.5 mg./Tab. Bot. 100s, 500s, 1000s.
Use: Sedative with antispasmodic agent.
- SEDULON.** (Roche) Dihyprylene, 3,3-diethyl-2,4-dioxo-piperidine. (Piperidone). Syr. 60 mg./5 cc. Bot. 4 oz., 16 oz., 1 gal.
Use: (1 tsp. q. 2 h.) Sedative.
- SEEK.** (Tracy) Tab. Bot. 12s, 25s, 50s.
- SEEF.** (See! Lab.) Tab. Bot. 100s.
- SEGO LIQUID.** (Pet Milk Co) Protein 77 Gm., Carbohydrate 103 Gm., Fat 20 Gm., Vit. A 5000 U.S.P. u., D 400 U.S.P. u., C 100 mg., B-1 2 mg., B-2 3 mg., niacinamide 15 mg., B-6 2 mg., Copanthenate 10 mg., I 150 mcg., K 3.5 Gm., Na 0.9 Gm., Mn 2 mg., Zn 5 mg./4 cont. Can 10 oz.
Use: Liquid diet food.
- SEGONTIN.** (Lloyd Bros) Prenylamine.
- SELENIUM SULFIDE, U.S.P., N.N.D. 1962.**
Deterg. Susp. U.S.P.
See Selsun, Susp., N.N.B. 1959. (Abbott)
- SELSUN CREAM.** (Abbott) Selenium sulfide 0.5%. Tube $\frac{1}{4}$ oz.
Use: Seborrhea of the glabrous skin & biopharitis marginalis.
- SELSUN SUSPENSION, N.N.D. 1961.** (Abbott) Selenium sulfide 2½%. Bot. 4 fl. oz.
Use: Seborrheic dermatitis of scalp.
- SELSUNEF, OINT.** (Abbott) Selenium disulfide 0.5%, hydrocortisone acetate 0.5% Tube 5 Gm.
Use: Antiseborrheic
- SEMALCYNE.** (Massengill) Phenobarb. $\frac{1}{4}$ gr., hyoscyamus ext. 1/6 gr., aspirin 2½ gr., phenacetin 1 gr./Cop. Bot. 100s.
Use: Analgesic.
- SEMAPIRIN.** (Strafenburgh) Acetophenetidin 2½ gr., acetylsalicylic acid 3½ gr., caffeine $\frac{1}{4}$ gr., Gelsemium tr. 2 min./Tab. Bot. 100s, 500s, 1000s. W/caffeine $\frac{1}{4}$ gr./Tab. Bot. 100s, 500s, 1000s.
Use: Antipyretic & analgesic.
- SEMBOX.** (Massengill) Liver inj. (eq 8-12 10 mcg) Vit. B-12 20 mcg., folic acid 10 mg./cc. Vial 10 cc.
- SEMCOF.** (Massengill) Codeine phosph. 10 mg., sod citrate 0.2 Gm., ammonium chloride 0.1 Gm., ephedrine HCl 5 mg., tartar emetic 1 mg./5 cc. Bot. 4 oz., 1 pt., 1 gal.
Use: Treatment of coughs
- SEMCOF-AH.** (Massengill) Methapyrilene HCl $\frac{1}{4}$ gr., Sod. citrate, ammonium chloride, ephedrine HCl, antimony Pot. tartrate/dr. Syr., Bot. 1 pt., 1 gal.
Use: (1 tsp. q. 4 h) Coughs, asthma & bronchitis.
- SIMCOF-CP.** (Massengill) Codeine phosphate 1/6 gr., w/Sod. citrate, ammonium chloride, ephedrine HCl, antimony paw. tartrate, aromatics/dr. Syr., Bot. 1 pt., 1 gal.
Use: (1-2 tsp. p.r.n) Cough.
- SIMCOF-M.** (Massengill) Morphine sulf. 1/24 gr., w/Sod. citrate, ammonium chloride, ephedrine HCl, antimony Pot. tartrate, aromatics/dr. Syr., Bot. 1 pt., 1 gal.
Use: (1-2 tsp. q. 4 h) Cough.
- SEMESTIBIN.** (Massengill) Natural estrogenic substances. Amp. In oil (10,000 u., 20,000 u./cc.) 1 cc., 10 cc.; eq. susp. 20,000 u., 50,000 u./cc.) 1 cc. Cop. 10,000 u., Bot. 100s, Tab. 4000 u., Bot. 100s
Use: Estrogen therapy.
- SEMETS.** (Massengill) Tyrothricin 1 mg., hyamine 2 mg., benzocaine 3 mg./toz. Bot. 12s & 500s.
Use: Infections of throat & mouth.
- SEMHYTEN.** (Massengill) Phenobarbital $\frac{1}{4}$ gr., maanitol hexanitrate $\frac{1}{2}$ gr., theophylline 1½ gr., rutin 10 mg., ascorbic acid 15 mg./Cop. Bot. 100s, 500s, 1000s.
Use: (1 q. 4d) Essential hypertension.
- SEMI-LENTE ILETIN.** (Lilly) A complex substance resulting from the interaction of zinc chloride and insulin, to produce a small particle size. It has a duration of action of 12-16 hrs. Vial (40 u./cc.) U-40 10 cc.; (80 u./cc.) U 50 10 cc.
Use: To shorten duration of lente insulin
- SEMIKON HYDROCHLORIDE, N.N.D. 1962.** (Massengill) Methapyrilene HCl Cream 2%, Tube 1 oz.; Jar 1 lb. Tab. 25 mg., 50 mg., 100 mg., Bot. 100s, 500s.
Use: Antihistaminic.
- W/Atropine sulf., A.P.C. See: Dasikon, Cop. (Massengill)
- W/Semoxyrine HCl. See: Hisdrin, Tab. (Massengill).
- Generic name recognized by N.F., U.S.P. or A.M.A.

SEMOPEN, NND 1962 (Massengill)
Pheneticillin alpha phenoxyethyl penicillin
Pot 125 mg or 250 mg/Tab Bot 24s
Powder for oral soln 125 mg/5 cc Bot 2
oz

Use Penicillin therapy

**SEMOXYDRINE HYDROCHLORIDE, NND
1962 (Massengill)** Methamphetamine HCl
Tab 2.5 mg, 5 mg, 7.5 mg, Bot 100s,
500s, 1000s Elix (5 mg/5 cc) Bot 1 pt,
1 gal

Use Central nervous system stimulant

SEMPERL. (Massengill) Vit A 10,000 u,
D 1000 u, thiamine HCl 5 mg, riboflavin
5 mg, ascorbic acid 75 mg, niacinamide
75 mg, pyridoxine HCl 0.5 mg, Cal panto-
thenate 5 mg, folic acid 1 mg, liver conc
(1:20) 56 mg, E 2 IU, ferrous sulfate
78 2, B 12 10 mcg/Perle Bot 100s

Use Dietary supplement

SEMPYLEX, (Massengill) Pyridoxine HCl 35
mg, thiamine HCl 10 mg, riboflavin 10
mg, nicotinamide 25 mg, glucose 20 Gm,
aromatics q s /ft oz Bot 1 pt, 1 gal

Use: Nausea & vomiting of pregnancy

SEMVIMIN, (Massengill) Vit A 5000 u,
D 1000 u, B 1 3 mg, B 2 2.5 mg, nia-
cinamide 20 mg, pantothenic acid 5 mg,
Vit B 6 0.75 mg, C 50 mg, E 3 mg, B 12
1.5 mcg, folic acid 0.33 mg, w/Cal, iron,
Phos, Pot iodide Mong, zinc, copper/
Perle Bot 100s, 1000s

Use Multivitamin therapy

SEMVITA, (Massengill) Vit A 12,500 u,
D 1000 u, thiamine HCl 10 mg, riboflavin
10 mg, pyridoxine HCl 0.5 mg, ascorbic
acid 75 mg, Cal pantothenate 2 mg,
nicotinamide 75 mg, d-alpha tocopheryl
acetate 2 IU, B 12 2 mcg/Perle Bot 100s,
500s, 1000s

Use: Dietary supplement

SEMVITA DROPS, (Massengill) Vit A 5000
u, D 1000 u, thiamine HCl 5 mg, ribo-
flavin 1 mg, nicotinamide 10 mg, pyri-
doxine HCl 1 mg, ascorbic acid 100 mg/
0.6 cc Bot w/dropper 30 cc

Use: Dietary supplement

SEMVITA PEBLE, (Massengill) Also avail
w/Vit B 12 2 mcg

SENAGEST, (Tablet Rock) Pentylene-tetraol
100 mg, ethinyl estradiol 0.0033 mg,
methyltestosterone 0.833 mg, mylose 30
mg, prolase 6 mg, cellulose 2 mg, choline
bitartrate 50 mg, inositol 25 mg, rutin 5
mg, Vit A acetate 5000 u, Vit E 0.5 mg,
Vit B-12 1 mcg, thiamine HCl 125 mg,
riboflavin 125 mg, niacin 10 mg, niacin-
amide 10 mg, pyridoxine HCl 0.25 mg,
calcium pantothenate 125 mg, ascorbic

acid 25 mg, iron 10 mg, lysine HCl 150
mg/Cap Bot 100s, 500s

Use Geriatric therapy, anabolic, hormone/
enzyme preparation

SENATRIC, (Dumas Wilson) Ethinyl estradiol
3.3 mcg, methyltestosterone 66 mg, pen-
tylene-tetraol 100 mg, nicotinic acid 100
mg, dicolcium glutamate 65 mg/Tab Bot
100s, 1000s

Use: A balanced androgen estrogen therapy
with respiratory and circulatory stimulants

SENAZOL, (Tablet Rock) Ethinyl estradiol 0.01
mg, methyltestosterone 2.5 mg, metrazol
100 mg, Vit A 10,000 u, D 500 u, E
1 mg, B 12 1 mcg, B 1 2.5 mg, B-2 2.5
mg, ascorc 20 mg, niacinamide 20 mg,
B 6 0.5 mg, cal pantothenate 2.5 mg,
C 50 mg, folic acid 1 mg, choline bitar-
trate 100 mg, inositol 50 mg, rutin 10
mg, w/iron, cal, phosp, iodine, copper,
cobalt, fluorine, mang, pot, zinc, mag
nesium/Cap Bot 30s, 100s, 500s

Use Orally, treatment of the geriatric pa-
tient

SENILAVITE, (Finley) Vit A 5000 u, C 100
mg, thiamine 2.5 mg, riboflavin 2 mg,
nicotinamide 10 mg, pyridoxine HCl 1 mg,
calcium pantothenate 5 mg, B 12 with In-
trinsic factor concentrate 0.133 u, ferrous
fumarate 150 mg, glutamic acid HCl 150
mg, dioctyl sodium sulfosuccinate 50 mg/
Cap Bot 100s

Use: Nutritional supplement

SENILEX, (Durst) Pantylene-tetraol 100 mg,
nicotinic acid 50 mg/Tab Bot 96s

Use: Psychotonic in geriatrics

SENNA, Alexandrian

See Aperans, Tab (Brayton)

Senokal Pow & Tab (Purdue Frederick)

**SENNA POD EXTRACT, W/Dehydrocholic
Acid** See: Senobile (Purdue Frederick)

**SENNA POWDER COMPOUND, NF
(Perick)** Camp Licarlice pow Bot 1/4 lb, 1 lb.

Use: Mild cathartic

**W/Cascara pow & methylcellulose. See:
Cellusena, Tab (Babst)**

SENNOSIDES A & B.

See: Glyxennid, Tab. (Sandaz)

Ovinoplax, Tab (Groy)

Senotab, Tab (Purdue Frederick)

**W/Guar gum, polygalacturonic acid See:
Gentils Granules (Purdue Frederick)**

SENOBILE (Purdue Frederick) Dehydrocholic
acid 125 mg active principles of Cassia
acutifolia pods 300 mg/Tab Bot 30s, 60s.

Use: Bileous constipation

SENODIN SYRUP, (Squibb) Codeine sulfate
3/8 gr, squill, senego, sanguinaria, licorice,
menthol, alcohol/8 oz Bot 1 pt, 1 gal

Use: (1 tsp q 3-4 h) Sedative expectorant

- SENOKAP.** (Purdue Frederick) Diocyl sod. sulfosuccinate 50 mg., *Cassia acutifolia* active principles 225 mg./Cap. Bot. 30s, 60s
Use: Stool softener in the treatment of functional constipation
- SENOKOT.** (Purdue Frederick) Aleandrian senna 600 mg./Tsp. Granules. Bot. 4 & 8 oz. Tabs.—Two tablets are equiv. to one teaspoonful Senokot granules. Bot. 16s, 30s, 100s.
Use: Laxative.
- SENOKOT W/PSYLLIUM.** (Purdue Frederick) Senna pod principles with psyllium. Pkg. 5.5 oz.
Use: Management of constipation
- SENOTAB.** (Purdue Frederick) Sennalids A & B 10 mg. (*Cassia acutifolia* pods 300 mg) Tab. Bot. 50s.
Use: Treatment of constipation.
- SENSODYNE.** (Block Drug) Strontium Cl, 10%; deionized water, 36.8%; polyols, 24%; water soluble gums, 1.8%; anionic detergent, 1.7%; non-ionic detergent, 0.5%; diatomaceous silico, 23%; flavoring, saccharin, inert agents and preservatives 2.2%/Cream. Tube: 2.2 oz.
Use: Dental.
- SENTI-PHEN.** (Vitarline) Adiphenine HCl 50 mg., phenobarbital 20 mg./Tab. Bot. 100s.
Use: Sedative w/antispasmodic agent.
- SENTIV.** (Vitarline Co.) Adinaphine HCl 75 mg./Tab. Bot. 100s, 500s, 1000s.
Use: Spasmolytic, sedative and anesthetic.
- SENTOR.** (Dunbor) Sticks.
Use: Dermatalogic.
- SEPTAMIDE.** (Lemman) Methenamine mandelate 250 mg., sulfacetamide 250 mg., homalrapine methylbromide 2.5 mg./Tab. Bot. 100s.
Use: G-u tract infections.
- SEPTIDERM.** (Fougero) 1% chlorocresol, a substituted phenol (3,5-dimethyl-4-chlorophenol), in water-washable base. Cream 1 oz. tubes and 1 lb. jars. HC Cream ½%, ¼ oz. and 1 oz. tubes.
Use: Infected dermatoses.
- SEPTISOL.** N.N.R. 1953, A.O.R. (Vnsta) Soap containing 2% hexachlorophen. Poil 3.7R L, 113.5 L; Drum 208.1 L
Use: Local anti-infective soap.
- SEQUESTRENE A.** (Alrose Chem.) Edathamil, N.N.R., Ethylenediaminetetraacetic acid. Pkg. according to demand
Use: Sequestering agent.
- SER-AP-ES.** (Ciba) Reserpine 0.1 mg., Hydralazine HCl 25 mg., Hydrochlorothiazide 15 mg./Tab. Bot. 100s.
Use: Hypertension
- SERC.** (Spencer) Methohistone Hydrochloride. Use: Diamine oxidase inhibitor.
- SERENIUM.** (Squibb) Ethazocene; Diominoethoxy-azobenzene HCl. Tab. D.I Gm., Bot. 25s, 50s, 500s.
Use: Infections of urinary tract
- SERFIN.** N.N.D. 1962. (Parke, Davis) Reserpine. Tab. (0.25 & 1 mg) Bot 100s, 500s.
Use: Hypotensive & tranquilizing agent.
- SERFOLIA.** (Mallard) Rauwolfia serpentina 2 mg./Tab. Bot. 100s, 1000s.
Use: Hypotensive agent.
- SERGINOL.** (Ascher) Reserpine 0.167 mg., ethinyl estradiol 0.02 mg./tab. Box 60s, Bot. 100s.
Use: Tranquilizing agent w/estrogen therapy.
- SERICINASE.** A proteolytic enzyme.
- SERNYL.** (Parke, Davis) Phencyclidine, 1-(1-Phenylcyclohexyl) piperidine HCl. Under study.
Use: Anesthetic.
- SEROLIA.** (Ascher) Reserpine. Tab. 0.5 mg Bot. 60s.
Use: Hypertension.
- SEROMYCIN.** N.N.D. 1962. (Lilly) Antibiotic produced by a strain of *Streptomyces orchidaceus* Cyclasterine (D-4-amino-3-isoxazolidinone), 250 mg./Pel. Bot. 40s & 500s.
Use: (1 twice daily) Pulmonary tuberculosis.
- SEROMYCIN W/ISONIAZID.** (Lilly) Seromycin 250 mg., isoniazid 150 mg./Pulv. Bot. 40s, 500s
Use: Treatment of tuberculosis
- SERPANRAY.** (Panray) Reserpine. Tab. (0.25 & 0.1 mg) Bot. 100s, 500s, 1000s. Tab. (0.5 mg. & 1.0 mg) Bot. 100s.
Use: Hypotensive & tranquilizing agent.
- SERPASIL.** N.N.D. 1962. (Ciba) Reserpine Tabs (0.1, 0.25 & 1 mg) Bot. 100s, 500s, 1000s, 5000s. Tab. (2 mg. & 4 mg) Bot 100s.
Amp. (2.5 mg./cc.) 5s.
Vial (2.5 mg./ml.) 10 ml.
Elixir (0.2 mg. or 1.0 mg./4 ml.) Bot. 1 pt., 1 gal.
Use: Hypotensive & tranquilizing agent.
For neuropsychiatric use only. Parenteral sol. Amp. 2 cc. (each cc. containing 2.5 mg Serpasil). Tab. 2 & 4 mg.
- SERPASIL—APRESOLINE.** (Ciba) Reserpine 0.2 mg., hydralazine HCl 50 mg./Tab. Bot. 100s, 500s, 1000s. Also avail. in half strength.
Use: Hypertensive disorders.
- SERPASIL-ESIDRIX.** (Ciba) #1 Serpasil 0.1 mg., Esidrix 25 mg./Tab. #2 Serpasil 0.1 mg., Esidrix 50 mg./Tab. Bot. 100s.
Use: Mild to moderate hypertension

- SERPATE, NND 1962** (Vale) Reserpine Tab 0.1 & 0.25 mg Bot 100s, 500s, 1000s
Use Hypotensive & tranquilizing agent
- SERPATILIN.** (Ciba) Reserpine 0.1 mg, ritalin 10 mg /tab Bot 100s
Use Psychomotor stimulant w/tranquilizing agent
- SERPATRATE.** (Amfro Grant) Phenobarbital 8 mg, monoflavin hexanitrate 30 mg, rauwolfia serpentina 50 mg, hesperidin complex 50 mg, ascorbic acid 50 mg /Tab Bot 100s 1000s
Use Treatment of hypertension
- SERPEDON.** (Walker) Reserpine 0.1 mg, hyoscyamine sulfate 0.105 mg, atropine sulfate 0.02 mg, hyoscyne HBr 0.007 mg /tab or 5 cc Bot 100s, 1000s Elix Bot 1 pt
Use Hypertension with antispasmodic
- SERPEN** (Hoag) Reserpine Tab (0.25 mg) Bot 100s, 1000s
Use: Hypotensive and tranquilizing agent
- SERPHEDRINE,** (Harvey) Reserpine 0.1 mg, aminophylline 100 mg, ephedrine sulfate 16 mg /tab
Use: Treatment of bronchial asthma
- SERPHYLLINE,** (Blair) Tab Bot 100s
Use Angina pectoris
- SERPICON,** (Consol Midland) Reserpine 0.1 mg or 0.25 mg /tab Bot 50s, 100s, 1000s
Use: Tranquilizing agent
- SERPILOID, NND 1959** (Riker) Reserpine Tab 0.25 mg, Bot 100s 500s, Tab 1, 2, 3, 4 & 5 mg Bot 100s, Amp (2.5 or 5 mg /cc) 2 cc
Use 1 lid; hypotensive and tranquilizing agent
- SERTRIM-R A.** (Arnell) Dextro amphetamine 15 mg, amobarbital 90 mg, atropine 3/500 gr /sustained tab Bot 50s
Use Management of obesity
- SERTINA.** (Testagar) Reserpine 0.25 mg /Tab Bot 100s 1000s
Use Hypotensive and ataraxic
- SERTS.** (Farrest) Bismuth subgallate 1 gr, balsam peru 1 gr, cetyl alcohol/Supp Box 12s Also avail w/ephedrine 1/16 gr, benzocaine oleate 1/2 gr /Supp
Use: Hemorrhoids
- SERUM, ALBUMIN, NORMAL HUMAN** See: Albumin Normal Serum, Human, USP, NND 1960
- SERUM, ALBUMIN, RADIOIODINATED,** NND 1960
- SERUM, ANTIHEMOPHILUS INFLUENZAE TYPE B (RABBIT), NND 1959** See: Anti hemophilus Influenzae Type B Serum (Rabbit), NND 1958
- SERUM, GLOBULIN (HUMAN), IMMUNE.** See Globulin Immune Serum (Human) USP, NND 1958 (Cutter; Lederle; Merck Sharp & Dohme; Pitman Moore)
- SERUM, MEASLES IMMUNE, HUMAN** See Measles Immune Human Serum, HF, NND 1959
- SERUM, NORMAL HUMAN, NND 1962** (Michael Reese Research Found) Bot 20 cc, 250 cc
- SERUM PERTUSSIS IMMUNE, HUMAN,** USP See: Pertussis Immune Serum, Human, NND 1962
- SERUM, SCARLET FEVER IMMUNE.** See Scarlet Fever Immune Serum, NND 1960
- SISPRON.** (Mossengill) Natural estrogens 20 000 IU 2 mg, progesterone in sesame oil 20 mg /cc, 10 cc
Use Estrogen & corpus luteum therapy
- SESTRAMIN.** (Polch) Compounded in 2 strengths, conjugated estrogens 0.625 mg, 1.25 mg w/vitamin content of each tableth brewers' yeast 100 mg, thiomine HCl 3 mg, riboflavin 2 mg, niacinamide 10 mg, pyridoxine HCl 1 mg, Col pantothenate 5 mg, ascorbic acid 25 mg, Vit B 500 IU /Tab Bot 100s, 500s
Use: Estrogens w/vitamin supplement
- SESTRON, N-ETHYL-3,3'-DIPHENYL-DI-PROPYLAMINE.**
See Antisiposmin (Royster)
Fentopine (C O Smith)
Prolend (C O Smith)
- W/Phenobarbital** See Antisiposmin w/phenob (Royster)
- SETRA-V.** (Worren) Vit B 1 10 mg, B 2 10 mg, B 6 5 mg, niacinamide 50 mg, calcium pantothenate 10 mg, Vit C 100 mg, bioflavonoid complex 100 mg /Tab Bot 100s
Use Nutritional supplement
- SEVETOL.** (Ives Cameron) Vit A 25,000 u; D 1000 u, B 1 20 mg, B 2 10 mg, niacinamide 75 mg, Vit B 6 3 mg, C 200 mg, B 12 5 mcg /cap Bot 100s
Use Multivitamin therapy
- SHADOCOL** (Davies, Rose) Sod Tetraiodophenolphthalein, iodophthalein Sod Bot 20 Gm
Use Cholecystography
- SNARCILLIN AQUEOUS SUSPENSION.** (Merck Sharp & Dohme) Procaine penicillin G Vial (300M u/cc) 10 cc; (600M u) 1 dose
Use: Penicillin therapy
- SNARCILLIN FORTIFIED** (Merck Sharp & Dohme) Procaine penicillin G 300 000 u, pot penicillin G 100 000 u /dose Vial 1 dose, 10 dose
- SNARCILLIN FORTIFIED DI** (Merck Sharp & Dohme) Procaine penicillin G 300M u, pot penicillin G 100M u, dihydrostrepto-

- mycin sulfate 0.5 or 1 Gm./dose. Vial 1 dose.
 Use: I.M., penicillin w/streptomycin therapy.
- SHARCILLIN ORAL**, (Merck Sharp & Dohme) Pot. penicillin G 300M u./tsp. Bot. 60 cc.
 Use: Orally; penicillin therapy.
- SHARMONE**, (Merck Sharp & Dohme) Pregnenolone acetate. Tab. 0.1 Gm., Bot. 24s.
 Use: Rheumatoid arthritis.
- SHEBDEX**, (Sherman) Thiamin HCl 100 mg., riboflavin 1 mg., pyridoxine HCl 5 mg., nicotinamide 100 mg., Col. pantothenate 1 mg., benzyl alcohol 100 mg., dextrose 0.5 Gm./5 cc. Amp. 5 cc., Sox 5s, 50s, 100s.
 Use: Dietary supplement.
- SHEBESTOL**, (Sherman) Estrogenic substance (2 mg) 20,000 I.U., testosterone 10 mg./cc. Aq. susp., Vial 10 cc.
 Use: Estrogen & androgen therapy.
- SHERMAPLEX**, (Sherman) Thiamine HCl 100 mg., riboflavin 1 mg., nicotinamide 100 mg., pyridoxine HCl 5 mg., Col. pantothenate 1 mg./cc. Vial 20 cc.
 Use: Vitamin B deficiencies.
- SHEVAC**, (Sherman) 60 billion killed bacteria of common upper respiratory tract/Tab. Bot. 20s, 60s, 100s.
 Use: Prophylaxis for respiratory infections.
- SHIGELLA GROUPINO BERA**, (Lederle) Five shigella species agglutinins with control. (1 cc. in 3 cc. Bot.)
 Use: Diagnostic.
- SI-AMINO-BETAINE CAPS & POWDER**, (Siler) Glycine 100 mg., betaine anhydrous 400 mg., l-arginine HCl 5 mg., yeast 100 mg./Cap. or Tbs. or pwd Bot. 100s, 8 oz jars.
 Use: Food supplement.
- SIBEL**, (Plough) Methylpolysiloxane 25 mg., belladonna extract 16 mg., butobarbital sodium 16 mg./Tab. Bot. 100s.
 Use: Antispasmodic.
- SIBLIN**, (Parke, Davis) Water-absorbent material from plantago combined with 2 mg. Vit. B-1/Tsp. Box 4 oz., 1 lb. Tab., Bot. 100s, 500s.
 Use: Chronic constipation & diarrhea.
- SIGMAGEN**, (Schering) Prednisone 0.75 mg., aspirin 325 mg., ascorbic acid 20 mg., aluminum hydroxide 75 mg./tab. Bot. 100s, 1000s.
 Use: Rheumatoid arthritis.
- SIGNAMYCIN**, (Pfizer) See: Signemycin (Pfizer)
- SIGMOL ENEMA**, (Pharmaseal) Sorbitol solution N.F. 43 Gm., pot. dioctyl sebacinate 0.12 Gm./120 cc. Plastic bag & tip
 Use: Enema
- SIGNATAL**, (Sig. Labs) Vit. A 4000 U.S.P.
 * Generic name recognized by N.F., U.S.P. or A.M.A.
- u., D 400 U.S.P. u., B-1 5 mg., B-2 5 mg., B-6 4 mg., B-12 2 mcg., C 50 mg., K 1 mg., niacinamide 10 mg., ferrous fumarate 90 mg., l 01 mg., potassium .85 mg., Mn .05 mg., calcium carbonate 500 mg., choline 10 mg., inositol 25 mg./Cap. Bot. 100s, 1000s.
 Use: Prenatal vitamin-mineral-amino acid combination.
- SIGNEMYCIN**, (Pfizer) Formerly named Coso-Signemycin Caps.
 Caps. Oleandomycin Phos. 83 mg. & tetracycline HCl 167 mg. (total 250 mg./Cap. Bot. 16s, 100s. Oleandomycin 33 mg. & tetracycline 67 mg. (total 100 mg./Cap. Bot. 25s, 100s.
 Susp., Oral: Oleandomycin 42 mg. & tetracycline 53 mg. (total 125 mg.)/5 cc. Bot. 60 cc. Mint flavored, 1 pt. Syr.; Oleandomycin 41.7 mg. as triacetate and tetracycline 83.3 mg. as HCl/5 cc. Bot. 2 oz., 1 pt. Intravenous: Oleandomycin 166 mg. & tetracycline 334 mg. (total 500 mg./Vial Oleandomycin 83 mg. & tetracycline 167 mg. (total 250 mg.) buffered w/ascorbic acid/Vial
 Use: (1 to 2 cap. q.t.d.) Infections of staph, strep, & pneumo; also gram-positive bacteria, rickettsiae, viruses & protozoa.
- SIGNEMYCIN V**, (Pfizer) A mixture of oleandomycin 83 mg. & tetracycline 167 mg. (total 250 mg.) phosphate buffered/Cap. Bot 16s, 100s
- SIGTAS**, (Upjohn) Vit. A 5000 u., D 500 u., B-1 10 mg., B-2 10 mg., C 333 mg., B-6 2 mg., S-12 4 mcg., folic acid 0.4 mg., nicotinamide 100 mg., cal. pantothenate 20 mg./Tab. Bot. 30s, 90s
 Use: 1 daily vitamins.
- SILAIN TABLETS**, (Plough) Dimethicone. Methylpolysiloxane 50 mg./Tab. Bot. 100s.
 Use: Abdominal gas distention.
- SILAIN-GEL**, (Plough) Dimethicone. Methylpolysiloxane 25 mg., magnesium hydroxide 85 mg., codried magnesium carbonate and aluminum hydroxide 282 mg./Tab. Bot 100s.
 Use: Antacid
- SILALOID**, (Van Pelt & Brown) Atropine sulfate 1/5000 gr., hyoscyamine sulfate 1/1250 gr., phenobarbital 1/8 gr., Mag. trisilicate 7 gr./Tab. Bot. 100s, 500s, 1000s.
 Use: (1 q. 3-4 h) Antispasmodic & sedative.
- SILCONEX**, (Hygiene Research) Silicone 77% in an inert base. Oint. Tube 1 oz., 3.2 oz., 5 oz., Jar 5 lb.
 Use: Protective agent.
- * **SILICONE**, U.S.P. (Dow Chemicals) Dimethicone. Liq., Bot 1 oz. Bulk Pkg Oint. U.S.P.

W/Cod liver oil, zinc oxide, hexachlorophene See Hollandex Silicone, Oint (Holland Rantos)

W/Hexachlorophene See Perfection, Lot (Sutliff & Case)

W/Magnesium hydroxide See Casil (Reed and Carnick)

W/Nitro cellulose & castor oil See Covicone, Cream (Abbott)

W/Triethylene glycol & mineral oil See Allergex, Liq (Hallister Sher)

W/Vitamin A & D, tyrosine, hexachlorophene See Protasil, Oint (Bryant)

SILICONE LOTION. See: Aquasil (Lentheric)

SILICONE OINTMENT, USP DIMETHICONE, NND 1962 Dimethyl polysiloxane See: Be Ba, Cr (Crookes)

Cavicone Cream (Abbott)

Damicone, Cream (Dane)

Pro-Derma (Westwood)

Siliconex, Oint (Hygiene Research)

Silicote (Arnar Stone)

SILICOTE, NND 1962 (Arnar Stone) 30% silicone oils Oint, Tube 1 oz; Jar 1 lb, Spray Disp 3 oz

Use: Skin protectant

SILICOTE PROTECTIVE CREAM (Arnar Stone) Silicones (Dimethicones) 30%, titanium oxide 1%; in a petrolatum water repellent base/Cream Tubes 1 oz
Use: Dermatologic

SILMACOL (Amfre) Mag trisilicate 50%, kaolin 25%, aluminum hydroxide 25%
Cap & Tab 8 gr, Bot 60s, 120s, 1000s
Pow, Bot 6 oz
Use: Antacid

SILMAGEL (Lannett) Aluminum hydroxide 25 gr, magnesium trisilicate 385 gr/tab
Bot 1000s, 5000s

Use: Protective antacid

SILMIDATE (Haskell) Salicylamide 250 mg
Cal succinate 200 mg, ascorbic acid 25 mg/tab
Bot 100s, 500s

Use: Analgesic

SILMIDATE-M. (Haskell) Mephenesin 500 mg, salicylamide 200 mg, ascorbic acid 25 mg/Tab
Bot 100s Also avail w/ codeine sulfate 8 mg/Tab
Use: 2-6 tid; various types of muscle spasm

SIMILAC W/IRON (Ross) Fat 3.4%, lactose 6.7%, protein 1.7%, minerals 0.5%, Vit A 2500 u, D 400 u, C 50 mg, B-1 65 mg, B-2 1 mg, niacin 7 mg, equiv elements B 6 155 mcg B 12 and folic acid in breast milk quantities, ferrous iron 12 mg / OI Liq Can 13 oz Case 24s Pow case 12s
Use: Iron supplementation in infancy

*Generic name recognized by N.F., U.S.P. or A.M.A.

SILVER CHLORIDE, COLLOIDAL, N.F. IX, NNR 1950 See Lunosol (Hille)

SILVER COMPOUNDS.

See Lunosol (Hille)

Neo Silvol (Parke, Davis)

Silver Chloride, Colloidal (Hille)

Silver Iodide (Parke, Davis)

Silver Nitrate (Various Mfr.)

Silver Picrate (Wyeth)

Silver Protein, Mild (Various Mfr.)

Silver Protein, Strong (Various Mfr.)

Silvagon (Bischoff)

SILVER IODIDE COLLOIDAL, N.F., NNR 1950 See: Neo Silvol (Parke, Davis)

SILVER, METALLIC COLLOIDAL. See: Col sargen, Liq (Crookes Barnes)

SILVER, METALLIC COLLOIDAL W/d Desoxyphedrine See: Colsedron, Liq (Crookes Barnes)

***SILVER NITRATE, USP, NNR 1949, ADR** (Various Mfr.) Ophth Sol USP Cryst & Granular Bot 1 oz Applicator, Pkg 100s See: Arzol Appl (Arzol Chem Co)

Use: Astringent, caustic & antiseptic

SILVER NITRATE OPHTHALMIC SOLUTION USP (Wax Amp)

Lilly—Amp 1%, 24s, 100s

Parke Davis—Cap 1%, 2s, 100s

Merck Sharp & Dohme—Amp 1%, 12s, 100s

Wyeth—Amp 1%, 2s

***SILVER NITRATE, TOUGHENED, N.F.** Silver nitrate plus 4% silver chloride See Arzol appl (Arzol Chem)

SILVER PICRATE, NNR 1957 (Delta) Bot 10 Gm See: Picrogol, NNR 1954 (Wyeth)
Use: Antiseptic

W/Kaolin See: Picragol Vaginal Supp. & Pow, NNR 1954 (Wyeth)

***SILVER PROTEIN, MILD, N.F., NNR 1949** (Various Mfr.) (Argentum Vitellinum; Cergentos, Nucleinate Mild Protargin Mild) See: Argyn NNR (Abbott)

Argyrol (Barnes)

Lunargen (Lilly)

Silvol NNR (Parke, Davis)

Selargentum Oint (Gilmore)

W/Kaolin, lactose See: Argypulvis Vag Cop (Crookes Barnes)

SILVER PROTEIN, STRONG, N.F., NNR 1949 See: Protargol (Winthrop Labs)

SILVOL, NNR 1949 (Parke, Davis) Mild silver protein. Cap (0.389 Gm) Pow, Bot 1 oz, ¼ lb

Use: Topically, silver antiseptic

- *SIMETHICONE.** (Dow Corning) A mixture of liquid dimethyl polysiloxanes.
See: Antifoam A Compound
Mylicon
Silain
- SIMILAC.** (M. & R. Dietetic Lab.) Modified milk & vitamins & minerals. Liq., Can 13 oz. Pow. 1 lb.
Use: Infant food.
- SIMILAC W/IRDN.** (M. & R. Dietetic Lab.) Similac formula w/ferrous iron 12 mg. per quart. Liq. Case 24s. Pow. case 12s.
Use: Iron supplementation in infancy.
- SIMPLASTIN.** (Warner-Chilcott) Defeted. Thromboplastin ext. w/Cal. & Sod. chloride. Vial 20 determinations
Use: Diagnostic aid.
- SIMRON.** (Merrell) Elemental iron as Ferrous gluconate 10 mg., Sacogen (polyoxyethylene glycol monolaurate) 400 mg./Cap Bot. 100s, 1000s.
Use: Anemia.
- SIMRON PLUS.** (Merrell) Iron (as ferrous gluconate) 10 mg., Sacogen, brand of polyoxyethylene glycol monolaurate 400 mg., oxyethylene glycol monolaurate 400 mg., Vit. B-12 with intrinsic factor 1/3 μ , ascorbic acid 50 mg., pyridoxine HCl 1 mg., folic acid 0.5 mg./Cap. Bot. 100s, 1000s
Use: Iron deficiency anemias.
- SINAN, N.N.D. 1962.** (Warren-Teed) Mephenesin. Tab. 0.5 Gm., Bot. 100s, 1000s.
Use: Spastic & neuromuscular disorders.
- SINAXAN, N.N.D. 1962.** (Armour) Styromate (2-phenyl, 2-hydroxy ethyl carbamate) 200 mg./Tab. Bot. 50s.
Use: Muscle relaxant.
- SINCO.** (Layden) Aluminum aspirin 5 gr., Chlorpheniramine maleate 2 mg., Phenylephrine HCl, 5 mg./Tab. Bot. 100s.
Use: Common cold.
- SINGD SERP, N.N.D. 1962.** (Ciba) Syrosingopine, 1 mg./Tab. Bot. 100s.
Use: Antihypertensive agent.
- SINGD SERP-ESIDRIX.** (Ciba) Tabs. No. 1 Singoerp 0.5 mg., Eisdrix 25 mg./Tab. Bot. 100s. Tabs. No. 2 Singoerp 1 mg., Eisdrix 25 mg./Tab. Bot. 100s.
Use: Moderate hypertension.
- SINOGRAFIN.** (Squibb) Dialrisozalm and iodipamide methylglucamines. Vial 10 cc.
Use: Diagnostic.
- SINTRDM, N.N.D. 1962.** (Geigy) Acenocoumarol. 3-(α -Acetyl-4-nitro-benzyl)-4-hydroxycoumarin. Tab. 4 mg., Bot. 50s.
Use: Drally, anticoagulant therapy
- SINUTAB.** (Warner-Chilcott) Acetyl-p-aminophenol 150 mg., phenacetin 150 mg., phenylpropranolamine HCl 25 mg., phenyltolaxamine dihydrogen citrate 22 mg./Tab. Bot. 30s.
Use: Decongestant & analgesic.
- SINUTAB WITH CODEINE.** (Warner-Chilcott) Codeine phosphate 15 mg., acetaminophen 150 mg., acetophenetidin 150 mg., phenylpropranolamine HCl 25 mg., phenyltolaxamine citrate 22 mg./Tab. Bot. 24s.
Use: Decongestant and analgesic.
- SIDMINE, N.N.B. 1952.** (Pilman-Moore) Iodoformine. Methanamine tetraiodide. Cap. 1 gr., 2 gr., 3 gr., Bot. 100s, 500s, 1000s.
Use: Alterative & urinary antiseptic.
- W/Sod. salicylate.** (Pilman-Moore) Methanamine tetraiodide $\frac{1}{2}$ gr., Sod. salicylate 7.5 gr./Cap. Bot. 100s.
Use: Anodyne.
- SIPPYLAC.** (Purdue Frederick) A brand of defatted dry milk protein solids.
W/Aluminum hydroxide, magnesium trisilicate, vit. B-1, B-2, B-6, C, col. pantothenate, niacinamide, menadione. See: Sippyplex, Pow. (Purdue Frederick)
- SIPPYPLEX.** (Purdue Frederick) Sippyloc 32.77 Gm., aluminum hydroxide 1 Gm., magnesium trisilicate 2 Gm., vit. B-1 20 mg., B-2 4 mg., B-6 2 mg., col. pantothenate 2 mg., niacinamide 20 mg., vit. C 140 mg., menadione 1 mg./8 tsp. Dry pow. Pkg. 8 oz.
Use: Comprehensive therapy for peptic ulcer.
- SIPREMEX.** (Ayerst) Vial, Estrone (200,000 I.U.) 20 mg., conjugated estrogens equine (Premarin), 10 mg. 10 cc.
Use: Estrogen therapy.
- SIRNDSITOL.** (Reed & Cornick) Choline gluconate 7.41 Gm., inositol 0.75 Gm./15 cc. Bot. 1 pt.
Use: Lipotropic.
- SIRDIL.** (Siroff) Mercuric oleate, creosol, vegetable & mineral oil. Emulsion, Bot. 8 oz.
Use: Antiseptic skin therapy.
- SISTOMYCOSIN.** (Porke, Davis) Antibiotic from *Streptomyces viridosporus*
- SITAB.** (Canright) Lobeline Sulf. 2.5 mg., benzocaine 4 mg., methapyrflane HCl 10 mg./Tab. Bot. 100s.
Use: Discourage smoking.
- SITABS KIT.** (Canright) Tab. A: pancreatic 10% gr., lysocyanine sulfate 0.0047 mg., pentobarbital sodium 108 mg./Tab. B d-amphetamine sulfate 12 mg., pentobarbital sodium 65 mg./Tab. C: lobeline sulfate 1.5 mg., aluminum hydroxide magnesium carbonate co-dried gel 150 mg., methapyrflane HCl 2 mg., benzocaine 2 mg./Tab. Kit contains 10 A 5 B and 51 C.
Use: Smoking inhibitor.
- * Genetic name recognized by N.F., U.S.P. or A.M.A.

***SITOSTEROLS**, N.F., NND 1962 A mixture of 80 to 90 per cent beta sitosterol and 10 to 20 per cent dihydro beta sitosterol See Cytellin, Susp (Lilly)

SITRIMIN, (Hart) Mag trisilicate 20 gr, colloidal kaolin 10 gr, with a trace of oil of peppermint and a trace of soluble saccharin/Tsp Mag trisilicate 7 gr, colloidal kaolin 3.5 gr with a trace of oil of peppermint and a trace of soluble saccharin/Tob Pow 1 oz, 4 oz, 1 lb Tab 100s, 500s

SIX B VITAMINS, (Raymer) Vit B 12 1 mcg, thiamine HCl 2 mg, riboflavin 1 mg, pyridoxine HCl 0.1 mg, Cal pantothenate 0.25 mg, niacin 10 mg, brewer's yeast 325 mg/Cap Bot 100s, 500s, 1000s
Use: Dietary supplement

SIX GLYCEROPHOSPHATES (No Sugar) ELIXIR (Merck Sharp & Dahme) Col glycerophosphate 8 gr, sod glycerophosphate 16 gr, iron glycerophosphate 1½ gr, man glycerophosphate 1 gr, quinine glycerophosphate ½ gr, strychnine glycerophosphate 1/16 gr/A oz Bot 1 pt, 1 gal
Use: (1 2 tsp t i d) Tonic

SKAYLOS LOTION, (Lee Colbert) Allantoin 2%, coal tar extract 5%, squalone 2%/cc Bot 8 oz
Use: Psoriasis

SKELAXIN (A. H. Robins) Metaxalone 400 mg/Tob Bot 50s, 500s
Use: Skeletal muscle relaxant

SKELETAL MUSCLE RELAXANTS,
See Adrenergic blocking agents

Ataraxic agents
Curare Preparations N N B
Dilyl (Elder)
Ganglionic blocking agents
Glyceryl Guaracolate
Mephenesin, N N R
Meprobamate
Penderol (Squibb)
Robaxin (Robins)
Sinaxar (Armour)
Succinylcholine Chloride, N N R
Tharozine (Smith, Kline & French)

SKIABARYT, N N R 1949 (Merck Sharp & Dahme) Barium sulfate
Oral: Mix of barium sulfate 80-85%, sugar, tragacanth, vanillin, cinnamon & cocoa Can 1 lb

Rectal: Mix of barium sulfate 95%, sugar & tragacanth Can 1 lb
Use: X-ray diagnosis

SKIN FILM, (Dome) Water soluble protective coating Cream, Jar 95 Gm, 390 Gm, 2200 Gm

Use: Skin protective

* Generic name recognized by N F U S P or A M A

SKIN LOTION,

See: Labana (Ulmer)

Lotocreme (Abbott)

SKIODAN, N N R 1952 (Winthrop Labs) (Methiodal Sod U S P; Sod Iodamethanesulfonate) Pow, Bot 20 Gm Sal 20%, Bot 50 cc, 40%, Bot 50 cc, 100 cc, Tab 1 Gm, Bot 100s, 1000s

Use: Radiopaque contrast medium

SKIODAN ACACIA SOLUTION, (Winthrop Labs) Sod iodamethanesulfonate Amp 40%, 10 cc

SKOPLATE NITRATE, N N D 1962 (Strassenburgh) Methscopolamine nitrate Tab (2 mg) Bot 100s, 1000s

Use: Treatment of peptic ulcers

SKOPOLATE D.A. (Strassenburgh) Methscopolamine nitrate Tab (2 mg), delayed action Bot 100s, 1000s

Use: Long term management of peptic ulcers

SKOPOLATE INJECTION, (Strassenburgh) Methscopolamine nitrate N N D 1959 Amp (0.5 mg/cc) 10 cc

Use: Treatment of peptic ulcers

SKOPOLATE P A M, N N D 1959 (Strassenburgh) Methscopolamine nitrate 1 mg, aluminum hydroxide 125 mg, Mag carb 125 mg Sod dodecyl sulfate 125 mg/Cap Bot 100s, 500s

Use: Management of peptic ulcers

SKOPYL DROPS, N N D 1962 (Pharmacia Labs) Methscopolamine nitrate 0.6 mg/Drop Bot 5 ml

Use: Spasmodic action

SLEEPWELL, (Amerit) Methapyrilene HCl 25 mg, W/Vit B complex/Cap Bot 24s
Use: Insomnia

SLO-JAYDEX, (Mid Atlantic) Amobarbital 1 gr, dextroamphetamine sulfate 15 mg/tab

Use: Appetite depressant

SLOWTEN, (Patch) Phenobarbital ¼ gr, thiamine HCl 5 mg/Tab or dr. Elixir, Bot 1 pt, 1 gal Tab, Bot 100s, 1000s
Use: Sedative

S M A. CONCENTRATED LIQUID, (Wyeth) Protease 1.5%, lactase 7%, fat 3.5% plus vitamins Can 13 oz.

Use: Infant feeding

S M A. POWDER (Wyeth) Detatted milk cocoa nut oil, beef fat, soybean oil, corn oil, Vit A, carotene, Vit B 1, C, niacinamide, lactase, Pot chloride, ferrous sulfate Can 1 lb

Use: Infant feeding

S M A. PROTEIN, (Wyeth) 2¼ times as much protein, ½ less fat, ⅓ less lactase and 3 times as much ash as human milk

- Contains vit. A 9000 u., D 600 u., B-1 0.67 mg., C 50 mg./Quart. Con B oz.
 Use: Infant feeding.
- S.M.A. NEW FORMULA 3-26.** (Wyeth Laboratories) The 40-60 casein: whey protein ratio is identical to human milk. Liquid, 13 fl. oz. cans, Pow. 1 lb. cans.
 Use: For infant feeding.
- *SMALLPOX VACCINE, N.N.D. 1962; U.S.P., N.N.B. 1949.**
 Cutter—Tube 1s, 5s, 10s, vaccinations.
 Lederle—Tube 1s, 5s, 10s, vaccinations.
 Lilly—Tube 1s, 5s, vaccinations
 Merck Sharp & Dohme—Tube 5s, 10s
 National Drug—Tube 1s, 5s, 10s, vaccinations.
 Parke, Davis—Tube 1s, 5s, vaccinations.
 Wyeth—Tube 1s, 5s, 10s, vaccinations.
- SMOOTHETS.** (Heinrich-Rickerstadt) Tob. Bot. 100s.
- SN. 13,272.** See: *Primoquine Phos.*, U.S.P.
- SNAKE VENOM.** See: *Antivenin*. (Wyeth)
 Use: S.C., I.M. & orally; trypanosomiasis.
- SOAPS GERMICIDAL.**
 See: *Cleasil* (Eastco)
Dial (Armour)
Fastex (Westward)
Lifebuoy (Lever Bros.)
pHisoex (Winthrop Labs.)
Thylax
Vestol (Vestal)
Woodbury (Jergens)
- SOAPS, MODIFIED.** See: *Neutrogena*. (Nalane)
- SOAP SUBSTITUTES.**
 See: *Acidolate* (White)
Acne-Dome, *Cleanser*. (Dome)
Breck Cleaner (Breck)
Dermolate (White)
Demerina, *Shampoo*. (Dome)
Lowila (Westwood)
pHisaderm (Winthrop Labs.)
Tersus (Daak)
- SOBEI, N.N.B. 1932.** (Mead Johnson) Soybean & precooked food. Pkg 15½ oz.
 Use: Infant food.
- SOCIN.** (Sun Lab) Tob. 100s.
- SODA ALUM.** See: *Aluminum Sodium Sulfate*.
- SODA MINT.** (Merck Sharp & Dohme) Tab. 5 gr. Bot. 100s, 1000s.
- SODASCORBATE.** (Van Patten) Sod. ascorbate. Amp. (100 mg./cc) 2 cc., 5 cc., 10 cc.; (500 mg./cc) 2 cc., 5 cc.; (1000 mg./cc) 10 cc. Tob. 120 mg. equiv. 100 mg. Vit. C. Bot. 40s, 100s, 1000s. Tab. 300 mg. Bot. 50s, 100s, 1000s.
 Use: Vitamin C therapy.
- SODASULF.** (G. F. Harvey) *Sulfadiazine*
 *Generic name recognized by N.F., U.S.P. or A.M.A.
- 0.083 Gm., *sulfamerazine* 0.083 Gm., *sulfamethazine* 0.083 Gm., Sod. bicarb. 0.250 Gm./Tab. Bot. 100s, 500s, 1000s.
 Use: Triple sulfonamide therapy.
- SODESTRATE.** (Maltbie) Conjugated estrogens. Tob. 1.25 mg., Bot. 50s, 100s, 500s.
 Use: Estrogen therapy.
- SODESTRIN.** (Tulog) *Potassium estrone sulfate* 1 mg., *estrone* 2 mg./cc. Vial 10 cc.
 Use: Menopausal; postmenopausal symptoms.
- SODIO-PHOS.** (Merck Sharp & Dohme) Sod. phosphate 39 Gm./100 cc. Bot. 1 gal
- SODIUM ACETATE & THEOPHYLLINE, N.N.R. 1950.**
 ***SODIUM ACETAZOLAMIDE, N.N.D. 1962.**
 See: *Diamox*, Prep (Lederle)
- ***SODIUM ACETRIZOATE INJECTION, U.S.P., N.N.D. 1962.** Sod. 3-acetamido-2,4,6-triiodobenzoate, Diagonal. See: *Urokin Sodium*, N.N.B. (Mallinckrodt), *Solpax*. (Ortho)
- SODIUM ACTD PHOSPHATE.** See: *Sodium Biphosphate*. (Various Mfr.)
- ***SODIUM ALGINATE, N.F., A.D.B.**
 See: *Algin*.
Kelgin (Kelco)
Kelpin. (Wallace)
 W/Col. alginic acid complex. See: *Kelpin*, Fow. (Wallace).
- SOD. ALKYL SULFOACETATE, A.D.R.** See: *Lathonal LAL*, Fow. (National Aniline Corp.)
- SODIUM ALUM.** See: *Aluminum Sodium Sulfate*.
- SODIUM ALUMINUM SILICATE.** See: *Ludoxan*, Tob. & Pow. (Schering).
- ***SODIUM AMINOHIPPURATE INJECTION, N.F., N.N.D. 1962** (Merck Sharp & Dohme) Sol. 20%, Bot. 10 & 30 cc.
 Use: I.V., to determine kidney tubular excretion function
- ***SODIUM AMINOPTERIN, U.S.P., N.N.D. 1962.** Tabs. U.S.P. See: *Aminopterin Sod.* (Lederle)
- ***SODIUM AMINOSALICYLATE, U.S.P., N.N.D. 1962** Tabs. U.S.P. See: *Aminosalicylate Sodium*. (Various Mfr.)
- SODIUM AMYTAL, N.N.R. 1947.** (Lilly) (Amobarbital Sod; Sod. Isoamylethylbarbiturate)
 Amps. (0.065 & 0.125 Gm.) Box 6s, 100s; (0.25, 0.5, 0.75 & 1 Gm.) Box 6s, 25s, 100s; (1 Gm.) w/Amp. 10 cc. water. Box 1.
 Enzals: (0.2 Gm.) Bot. 500s.
 Tablets: (1 & 3 gr.) Bot. 100s, 500s.
 Supps. (3 gr.) Bot. 12s, 100s.
 Use: I.M., S.C. & I.V., sedative.
- ***SODIUM ANAZOLENE.** (Ayerst) B-(4-anilino-5-sulfa-1-naphthylazo)-1-naphthol-3,6-disulfonic acid trisodium salt. See: *Coomassie Blue*.

- ***SODIUM ASCORBATE**, N N D 1962
See: Ascorbin, Amp (Lakeside)
Natrocarb (U S Vit)
Natri C, Tab (Physiological Chem)
Sodascorbate Amp & Tab (Von Patten)
- SODIUM AUROTHIOMALATE**. See: Gold
Sodium Thiosulfate N F
- SODIUM BENZOATE SOLUTION** (Hynson
Wescott & Dunning) Amp 1.77 Gm, 20 cc
Use: Liver function test
- ***SODIUM BIPHOSPHATE**, U S P N N R
1943 Monosodium Phosphate Sod Acid
Phosphate, Sod Dihydrogen Phosphate
See: Batophos Pow (Alphaden)
W/Sodium bicarbonate See: Ropox Inserts
(Elder)
W/Sod phosphate
See: Clyseral, Sol (Clyseral)
Phospho Soda Liq (Fleet)
W/Methenamine (Various Mfr)
- SODIUM BISMUTH CEVITAMATE** See
Bismuth Cevitamate Amp (C D Smith)
- SODIUM BISMUTH TARTRATE**, N N R 1952
See: Binazol Vial (Rorer)
Bismuth Sod Tartrate Preps
- ***SODIUM BITHIONOLATE** (Vonderbill)
Disod um 2,2 thiobis (4,6 dichlorophen
oxide) See: Yoncide BN
- ***SODIUM BROMIDE**, N F A D R
W/Antipyrine sodium salicylate caffeine
alkaloid See: Bromoso Liq (Tilden)
W/Citrate carb See: Bromo Vess, Tab
(Ames)
W/Phenobarbital See: Bromobarb Elixir
(Hort)
- ***SODIUM CACODYLATE**, N F W/Iodine
See: Cacodyne Iodine Amp (Res Medica
ions Inc)
W/Mercury binoxide Sodium iodide Phos
phorus See: Endoarsen (Endo)
- SODIUM CALCIUMDEATE** See: Calcium
Disodium Versenate (Eker)
- ***SODIUM CAPRYLATE**, Inj N N D 1962
Chemo Pura—Pow Bulk
Ingram—Amp 33%, 1 cc Pkg 12s, 25s,
100s
Strassenburgh—Sol 0.2 Gm/cc, Bot 60
cc, 480 cc, 383 L
See: Caprylate Sodium
Use: Fungus infections of skin
W/Zinc caprylate
See: Caprylic Compound, N N R 1954
Naprylate, N N R 1954 (Strassenburgh)
- ***SODIUM CARBOXYMETHYLCELLULOSE**,
U S P N N D 1962 A D R C M C Cellulose
Gum Tabs U S P See: Carboxymethylcel
lulose Sod (Various Mfr)
- SODIUM CELLULOSE GLYCOLATE** See:
Carboxymethylcellulose Sod (Various Mfr)
*Generic name recognized by N F U S P or A M A
- ***SODIUM CEPHALOTHIN**, (Eli Lilly) 7-(2
thienylacetamido)cephalosporanic acid so
dium salt
Uses: Antibiotic
- SODIUM CHLORIDE SUBSTITUTES**. See: Salt
substutes
- SODIUM CHLORIDE THERAPY**.
See: Thermolene, Tab (Lannett)
Thermotabs (Merck Sharp & Dohme)
- SODIUM CHLOROTHIAZIDE**, N N D 1960
See: Diuril, Prep (Merck Sharp & Dohme)
- SODIUM CITRATE** W/Sod tartrate See:
Citra Soda Pow (Abbott)
- SODIUM COLISTIN METHANESULPHATE**
Colistimethate Sod The sodium methane-
sulfonate salt of an antibiotic substance
elaborated by *Aerobocillus colistinus* See:
Coly Mycin (Warner Chilcott)
- ***SODIUM CYCLAMATE**, N F, N N D 1962
Sod Cyclohexylsulfamate Cpd Saln, Cpd
Tab N F See: Sucaryl Salts
- ***SODIUM CYCLONEXYLSULPAMATE**. See:
Sucaryl salts
- ***SODIUM DEHYDROCHOLATE**, Inj N F,
N N R 1953
Paul Money—Amp 20%, 5 cc, 6s, 25s
Breon—Amp 20%, 5 cc, 6s, 25s
Endo—Amp 20%, 3 cc, 5 cc, 10 cc, 6s,
25s
G F Harvey—Amp 20%, 5 cc, 6s, 12s,
25s, 100s
Strassenburgh—Amp 20%, 5 cc, 6s, 25s
Use: Choleraic & diagnostic agent
See: Decholin Sodium (Ames)
Dilabil Sodium (Breon)
- ***SODIUM DEXTROXYNIXINE** Sodium
D 3,3,5 Tetraiodothyronine See: Cholaxin
(Flint Eaton)
- ***SODIUM DIATRIZOATE**, U S P N N D
1962, Inj. U S P See: Hypoaque (Winthrop
Labs)
W/Methylglucamine diatrizoate sodium
citrate disodium ethylenediamine tetra
acetate dihydrate methylparaben, propyl
paraben See: Renovist (Squibb)
- SODIUM DICROMATE** W/Boric acid See:
Di O Chroma, Liq (Warren Teed)
- SODIUM DIMETHOXYPHENYL PENICILLIN**
See: Methicillin
- ***SODIUM DIPNICILLIN** (Smith Kline &
French) Sod um 2 biphenylpenicillin See:
Ancillin
- ***SODIUM DIPROTRIZOATE**, U S P N N D
1962 See: Makin Inj (Mallinckrodt)
- ***SODIUM ETNASULFATE** (Abbott) Sodium
2 ethyl 1 hexanoate sulfate
Use: Detergent

- SODIUM ETHYL-MERCURI-THIO-SALICYLATE**, N.N.R. 1949, A.D.E.
See: Thimerosal, N.F.
Merthiostat (Lilly).
- * **SODIUM FLUORIDE**, U.S.P., A.D.R.
See: Flura Drops (Kirkman Pharm.)
Flural, Liq. (Novocall)
Karidium, Tab. (Lorvic Corp)
Perganlene, Soln. (Fergontene)
T-Fluoride (Tennessee Pharm.)
W/Vit. A, D, C. See: Tri-Vi-Flar Drops.
(Mead Johnson)
- * **SODIUM FOLATE**, N.N.D. 1962: See:
Folvite Sodium (Lederle)
- SODIUM POLYVITE**, N.N.D. 1962.
- SODIUM FORMALDEHYDE SULFOXYLATE**.
(Various Mfr.) $H_2C(OH)SO_2Na$
Use: Eeducing agent
- SODIUM-FREE SALT**.
See: Co-Salt (Funk).
Diosol (Fougera).
Gustomate (Arlington).
Neocurtosol. (Winthrop Labs.)
- SODIUM GENTISATE**. See: Gentisate, Sod.
- * **SODIUM GLUCOSULFONE INJ.**, N.N.D.
1962. See: Promin Sod. Inj. (Parke, Davis)
- * **SODIUM GLUTAMATE**, N.N.D. 1962 See:
Glutamate.
- SODIUM GLYCOCHOLATE**, A BILE SALT.
See: Sile Salts, N.N.R. 1949
W/Sod. taurocholate, Sod. salicylate, pan-
creatlin & Sod. bicarb. See: Tablogestin,
Tab. (Sirang)
- * **SODIUM HEXACYCLONATE**. Sodium 3-
hydroxymethylcyclohexanecarboxylate. (Warner-
Chilcott)
- SODIUM HYDROGEN CITRATE**. See: Ci-
tralka, Liq. (Parke, Davis)
- SODIUM HYPOCHLORITE SOLUTION**, N.F.,
A.D.E.; Diluted, N.F.; N.N.E. 195D
See: Antiflarmin
Cloroxin (Guardian)
Dakin's Soln.
Fexeral (Fesler Co.)
Hyclorite, N.N.R. 1950
Millon Antib. Soln. (Walker)
Zonite (Zonite Prod)
- SODIUM HYPOSULFITE**. See: Sodium Thio-
sulfite
- * **SODIUM INDIGOTINDISULFONATE**,
U.S.P. Inj., U.S.P. See: Indigo Carmine.
- * **SODIUM IODIDE**, U.S.P., N.N.E. 1947, Inj.,
N.F.
W/Caseln.
See: Coside, Vial (Endo)
Iodotain, Vial (Myers-Carter)
W/Iodinated peptane.
See: Iodaloke-S, Amp. & Vial. (Lakeside)
Prolide, Vial. (Poul Maney)
- * Generic name recognized by N.F., U.S.P. or A.M.A.
- * **SODIUM IODIPAMIDE**, N.N.D. 1962. See:
Chalografra Sod., Amp. (Squibb)
- SODIUM IODONYDROXYQUINOINE-
SULFONATE** W/Purified Casein. See:
Activin, Amp. (Bischoff).
- SODIUM ortho-iodohippurate**, N.N.E.
1954.
See: Hippuran, Amp. (Mallinckrodt)
Medapaque, Vial (Bell Craig)
- * **SODIUM IODOMETHAMATE**, N.F., N.N.E.
1954; Inj., N.F. (Iodoxyl, Pyelectan, Uropac)
See: Neolapox, Amp. (Schering)
- SODIUM IODOMETHANESULFONATE**. See:
Methiodal Sodium, U.S.P.
- SODIUM IODOPROPANOL SULFONATE**.
See: Iyxanthine (Gallico)
Neo-Eiadin, Amp. (Asliier)
- SODIUM IOTHALAMATE**. (Mallinckrodt)
See: Angia-Conray.
- SODIUM IOTHIOURACIL**, N.N.D. 1962. Sod.
5-iodo-2-thiouracil. See: Itrumil, Tab. (Ciba)
- * **SODIUM IPODATE**, Sodium 3-(dimethyl-
aminomethyl)benzoino)-2,4,6-triiodohydro-
cinnamate. See: Oraglofin Sodium (Squibb)
- SODIUM ISOAMYLETHYLBARBITURATE**,
N.N.R. 1949. See: Sodium Amytal (Lilly).
- SODIUM LACTATE** W/Chlorides of pat. &
sod. See: Darrow's Solution, U.S.P.
W/Pat. citrate. See: Citra Lactate, Soln.
(Donley-Evans)
- SODIUM LACTATE SOLUTION**. (Don Baxter)
One-sixth molar. Vials 150 cc., 250 cc.,
1000 cc.
Use: I.V., S.C., I.S., I.P., treatment of
acidosis
- * **SODIUM LAURYL SULFATE**, U.S.P., A.D.E.
See: Amtracin. (Amlra)
Duponal.
Neutrozyme, Supp. (Smith-Dorsey)
W/Dihydroxyaluminum aminoacetate, meth-
scopolamine bromide & magnesium hydrox-
ide.
See: Aiu-Scop, Cap, Susp. (Lloyd, Dabney
& Westerfield)
Kolantyl. (Marrell)
- SOD. LAURYL SULFOACETATE**. See: Latho-
nal LAI, Pow. (National Aniline Corp)
- * **SODIUM LEVOTHYRONE**, U.S.P., N.N.D.
1962; Tabs, U.S.P. See: Synthroid Sod.,
Tab. (Travenol)
- * **SODIUM LIOTHYRONINE**, U.S.P., N.N.D.
1962; Tabs., U.S.P.
See: Cytomet, Tab (Smith, Kline & French)
Liothyronine Sodium.
Trianine, Tab. (Roche)
- SODIUM MALONYLUREA**. See: Barbitol
Sod. (Various Mfr.)
- * **SODIUM MERALEIN**. (Hynsan, Wittcott &
Dunning) mono-sodium salt of 2,7-diiodo-

4 hydroxy mercuriresorcinulfonphthalein
See: Merodicein

***SODIUM METHICILLIN** Sod 2,6-dimethoxy
phenyl penicillin monohydrate
See: Dimocillin (Squibb)
Staphycillin (Bristol)

SODIUM METHOXYCELLULOSE. (Mixture of
methylcellulose and sod CMC) W/Meth
amphetamine HCl See: Obetamine, Tab
(Sly)

SODIUM MORGHUATE, Inj, USP, NNR
1954 (Various Mfr) Amp 5%, 2 cc., 5 cc.,
30 cc., 60 cc
Use: Sclerosing agent

W/Sulfadiazine Sod See: Morusul, Vial
(Endo)

SODIUM NICOTINATE. Sod pyridine-3-
carboxylate 24 mg/cc.

Mint Eaton—Amp 5 cc., Box 6s; 15 cc.,
Box 1s

Upjohn—Amp 10 cc., Box 6s; 25s; Vial
(140 mg/cc) 10 cc

Use: IV, nicotinic acid therapy

SODIUM NITRATE. (Various Mfr) Granules,
Pkg 1 lb

W/Nitroglycerin, labeia & pilocarpine See:
Hiperline, Tab (Westerfield)

W/Sod nitrite, ox bile & echinococ See:
Hyper Zem, Tab 2 (Zemmer)

***SODIUM NITRITE**, USP, Tabs NF P (Var-
ious Mfr) Gran Bot ¼ lb 1 lb

See: Anti Rust Tabs
Pimerine, Soln (Novocof)

Use: Vasodilator

Sodium Nitrite Combinations on p 673

***SODIUM OXACILLIN** 5 Methyl 3 phenyl
4 isoxazolyl penicillin

See: Prostophlin (Bristol)
Resistopen (Squibb)

SODIUM PARA-AMINO BENZOATE, NNR
1949 See: p Aminobenzoate Sod (Var-
ious Mfr)

SODIUM PARA-AMINO HIPPURATE, Inj,
NF NND 1960 (Merck Sharp & Dohme)
Sol 20%, Bot 50 cc

Use: IV to determine kidney function

SODIUM PARA-AMINOSALICYLATE, USP
NND 1962 See: p Aminosalicylate Sod
(Various Mfr)

SODIUM PANTOTHENATE (Parke, Davis)
Cap 10 mg Bot 100s

Use: Orally dietary supplement

***SODIUM PENICILLIN G**, USP NND
1962 ADR See: Penicillin G Sodium

***SODIUM PENICILLIN O** See: Cer O Cillin
Sod (Upjohn)

**SODIUM 3,3-PENTAMETHYLENE-4-HY-
DROXYBUTYRATE** See: Hexacycloneate so-
dium Under study

*Generic name recognized by NF USP or A.M.A.

SODIUM PENTOBARRITAL, NNR 1952.
See: Pentobarbital, Sod (Various Mfr).

SODIUM PERRORATE, NF, ADR. Sod
Peroxyborate Sod peroxyhydrate

See: Amoran (Knox)
Vince, New (Standard Lab)

SODIUM PEROXYBORATE See: Sod Per-
borate, NF

SODIUM PEROXYHYDRATE. See: Sodium
Perborate

SODIUM PHOSPHATE, NF, NNR. 1930,
Effer, NF; Essic, NF; Sol, NF; Disodium
phosphate

W/Sod biphosphate

See: Clystral solution (Clystral Lab)
Phospho-Soda, Liq (Fieels)

***SODIUM PHYTATE** (Squibb) Sodium salt
of inositolhexaphosphoric acid See: Renal
(sodium phytate granules)

SODIUM POLYSTYRENE SULFONATE, See:
Kayaxalate, Pkg (Winthrop Labs)

***SODIUM PROPIONATE**, NF, NNR 1955
5% Soln Eye drops (Crookes Barnes)
Lacrival 15 cc

See: Propion Gel (Wyeth).
Sopronal (Wyeth)

W/Neomycin sulfate See: Otiobiotic, Ear
Drops (White)

W/Phenylephrine HCl See: Neo-Propisal
(Ophthalmal)

W/Prednisolone acetate, neomycin See:
Otiobione (Whites)

W/Sod caprylate, propionic acid & un-
decylenic acid See: Verdefam (Texas
Pharm)

***SODIUM PSYLLATE**, Inj, NF NND
1962 See: Slynosal (Searle)

Use: Sclerosing agent

SODIUM PTEROYLGLUTAMATE, NND
1962 See: Sodium Folate (Lederle)

SODIUM PTEROYLTRIGLUTAMATE, NNR
1957 See: Teropterin, Amp, NNR (Led-
erle)

***SODIUM RADIO-CHROMATE** Inj, USP.
NND 1962 See: Rad-a-Chromate (Cr⁵¹)
Sod

**SODIUM RADIO IODIDE (I¹³¹) SOLU-
TION, DIAGNOSTIC** (Abbott) Sod um
Radio Iod de (I¹³¹) 25 microcuries/cc Vials
10 cc

Use: Liver function thyroid function & pro-
tein bound iodine

***SODIUM RADIO IODIDE SOLUTION**,
USP, NND 1962 (Abbott) Sodium Radio
Iod de (I¹³¹) 5 15 15-40 & 40 100 mth
curies respectively Vials 10 20 30 cc

See: Iodatope (Squibb)
Onod de (Abbott)

Sodium Nitrite Combinations

| Sodium Nitrite | Others | Manufacturers | Brand |
|-------------------|--|-----------------------|---|
| 130 mg. 1 gr. | Barbital Sod. 65 mg. Diethylbarbital $\frac{1}{4}$ gr., nitroglycerin 1/250 gr., crotaegus tr. 2 min. | Chicago Buffington | Barbiodite, Pills Diatroegus, Pill |
| 1 gr. | Diethylbarbital $\frac{1}{4}$ gr., nitroglycerin 0.26 mg., crotaegus tr. 2 min., rutin 20 mg. | Burlington | Vasorulin, Pill |
| 0.13 Gm. | Glycolate Sod. 8.1 mg., chionanthin 16 mg., phenobarbital 16 mg., irisin 6.3 mg. | Cafe Chem. Co. | Norb. Tab. |
| $\frac{1}{2}$ gr. | Mannitol hexanitrate $\frac{1}{4}$ gr., erythryl tetranitrate $\frac{1}{8}$ gr., ammonium hippurate 1 gr. | Burroughs Wellcome | Vasodilator Comp., Tab. |
| 1 gr. | Ox bile, Sod. nitrate 2 gr., echinacea | Zemmer | Hyper-Zem, Tab. 2 |
| 1 gr. | Phenobarbital $\frac{1}{4}$ gr., veratrum $\frac{1}{4}$ gr. | Metropolitan | Verital, Tab. |
| 1 gr. | Phenobarbital $\frac{1}{4}$ gr., veratrum 3 Crow u. | Normand | VJ-Rumal, Tab. |
| 1 gr. | Phenobarbital $\frac{1}{4}$ gr., veratrum $\frac{1}{4}$ gr., rutin $\frac{1}{8}$ gr. | Superior | Verulinum, Empule |
| 65 mg. | Phenobarbital 16 mg., veratrum viride 40 u. | Irwin-Nelson | Veratril, Tab. |
| 2 gr. | Phenobarbital $\frac{1}{4}$ gr., veratrum viride tr. 4 min. | Lehman | Nirital, Tab. |
| 1 gr. | Phenobarbital $\frac{1}{4}$ gr., veratrum viride tr. 2 min., aconite tr. 2 min., cascara $\frac{1}{8}$ gr., bile salts 1/6 gr. | Tailby-Nason | Polensars, Tab., also w/Phenobarbital $\frac{1}{4}$ gr. & Rutin 2 mg. Manverin, Cap. |
| 50 mg. | Phenobarbital 10 mg., veratrum viride 50 mg., mannitol hexanitrate 30 mg., rutin 20 mg. | Sullivan & Case | Nyamin, Tab. |
| $\frac{1}{2}$ gr. | Veratrum viride 2/5 gr., aconite 2/5 gr., nitroglycerin 1/250 gr., Pot. nitrate 2/5 gr. | Elder | Polensars Forte Cratoganite, Tab. |
| 1 gr. | Veratrum viride 1 1/2 gr., cascara $\frac{1}{2}$ gr., bile salts $\frac{1}{8}$ gr. | Tailby-Nason | |
| 1 gr. | Veratrum viride 1 gr., nitroglycerin 1/100 gr., Pot. nitrate 1 gr., crotaegus tr. 1 min., opuncium tr. 1 min. | C. D. Smith | |

- Radiocaps (Abbott)
Theriodide (Abbott)
Uses: Thyroid tumors, hyperthyroidism & cardiac dysfunction
- ***SODIUM RADIO-PHOSPHATE, U.S.P.**, NND 1962 See: Radio Phosphate [p³²], Sod
- SODIUM REMOVING RESINS.** See: Resins
- SODIUM RHODANATE.** See: Sodium Thiocyanate
- SODIUM RHODANIDE.** See: Sodium Thiocyanate
- SODIUM RICINOLEATE, NNR 1954** See: Sorican (Merrell)
- ***SODIUM SALICYLATE, U.S.P.**, NND 1962; Tab, U.S.P.; Inj., NF [Various Mfr]
See: Nadisal Liquid (Lilly)
W/Acetophenetidin salicylic ethyl ester carbonate, colchicum root pow ext See: Tetragen, Tab (Endo)
W/Iodide (Merck Sharp & Dohme) Sod salicylate 1 Gm, sod iodide 1 Gm/20 cc. Amp Rox 25s
Uses: Intravenous injection
W/Iodide & calcichine (Merck Sharp & Dohme) Sod salicylate 1 Gm, sod iodide 1 Gm, calcichine 0.6 mg/20 cc Amp, Box 25s
Use: IV, gaul
W/Vegetable oil See: Lipa Sol, Liq (Donley Evans)
- SODIUM SALICYLATE, NATURAL.** See: Aly sine Elixir (Merrell)
- Sodium Salicylate Combinations on pp. 675 and 676
- SODIUM SALICYLATE & IODIDE, NF** (Merck Sharp & Dohme) Sod salicylate 15 gr, sod iodide 15 gr/20 cc Amp 20 cc, Box 25s
- SODIUM SALICYLATE & IODIDE W/COLCHICINE, NF** (Merck Sharp & Dohme, Parke, Davis) Sod salicylate 15 gr, sod iodide 15 gr, calcichine 0.6 mg/20 cc Amp 20 cc, Box 25s
- SODIUM SOTRADECOL, NND 1962** (Wallace & Tiernan) Sol of Sod tetradecyl sulfate w/benzyl alcohol 2% Vial (1%, 3%, & 5%) 2D cc
Use: Sclerosing agent
- SODIUM SUCCINATE.**
Horton & Converse—Tab 5 gr, Bot 100s, 1000s
Lilly—Tab 5 gr, Bot 100s, 1000s
Use: Alkalinize urine & awaken patients following barbiturate anesthesia
See: Saduxin Amp (Brewer)
- SODIUM SULAMYD, NND 1962** See: Sulamyl Sodium, Sod & Ophth Oint, NNR (Schering)
- ***SODIUM SULFACETAMIDE, NND 1962** ADR Sol, Bot (10%) 15 cc (Crookes-Barnes)
See: Oc U Mid (Blue Line)
Sulamyl Sodium, NNR (Schering)
Uses: Ophthalmic infections
W/Neomycin sulfate See: Sulfacidin, Oph Soln (Crookes-Barnes)
W/Phenylephrine HCl See: Sulfostatol, Oph Soln (Crookes-Barnes)
W/Prednisolone acetate, phenylephrine HCl See: Riephamide Ophthalmic Liquifilm (Allergan)
Sulfapred (Professional Pharmacol)
- SODIUM SULFADIAZINE, NND 1961** See: Sulfadiazine Sodium (Various Mfr)
- SODIUM SULFAMERAZINE, NNR 1945.** See: Sulfamerazine Sodium (Various Mfr)
- SODIUM SULFAPYRIDINE, NNR 1947.** See: Sulfapyridine Sodium (Various Mfr)
- SODIUM SULFATHIAZOLE, NNR 1948.** See: Sulfathiazole Sodium (Various Mfr)
- SODIUM SULFOACETATE, W/Sod alkyl aryl polyether sulfonate, Sod dioctyl sulfosuccinate, Kerahydric, sulfur, salicylic acid, hexachlorophene** See: Sebulex, Liq (Westwood)
- SODIUM SULFOCARBOLATE, (Mallinckrodt)** Sod phenolsulfonate Granules, Rot 1 lb Pow, Bot 1 lb
See: Sodaxilin, Liq (Abbott)
Uses: Internal antiseptic
- SODIUM SULFOCYANATE.** See: Sodium Thiocyanate, NF
- ***SODIUM SULFOXONE, U.S.P.**, NND 1962, Tabs, U.S.P. See: Diasone Sodium, Tab (Abbott)
- SODIUM TARTRATE** W/Sod citrate See: Citro Soda, Pow (Abbott)
- SODIUM TAUROCHOLATE, A BILE SALT** See: Bile Salts, NNR 1949
- ***SODIUM TETRADECYL SULFATE, NND 1962** Tergital See: Sodium Sotradecol Vial (Wallace & Tiernan)
- SODIUM TETRAIODOPHENOLPHTHALEIN, NNR 1950** See: Iodophthalein Sod, USP
- SODIUM THIACTYPHENARSAMIDE.** The trivalent organic arsenical p [bis (carboxymethylmercapto)arsina] benzamide See: Ceparsolate Sodium (Abbott)
- ***SODIUM THIMERFONATE** (Lilly) Sodium p ethyl mercuri thiophenylsulfonate See: Sulfa Merthiolate
- ***SODIUM THIOCYANATE, NF X.** Sodium Sulfocyanate Sod Rhodanide
See: Mimosed, Liq (Tilden)
Nitro Cyanates, El (First Texas)

*Generic name recognized by NF, U.S.P. or A.M.A.

Sodium Salicylate Combinations

| Sodium Salicylate | Others | Manufacturer | Brand |
|-------------------|---|----------------------|------------------------|
| 25 gr. | Anilpyrine 24 gr., sodium bromide 14 gr., caffeine alkaloid 4 gr. | Tilden | Bramoto, liq./a. oz. |
| 5 gr. | Cal. ascorbate 1 gr., aluminum hydroxide 2 gr., Cal. carb. 1 gr. | Mossengill | Solcedraz, Tab. |
| 15 gr. | Cal. carb., sod. bicarb. | Tablet Rock | Bufasal, Gran./dr. |
| 80 mg. | Chlorophenpyridamine maleate 0.75 mg., glycine 25 mg. | Schering | Carlin, liq./cc. |
| 5 gr. | Calcichine 1/250 gr., ascorbic acid 15 mg. | McNeil | Cinbisol, Tab. |
| 5 gr. | Calcichine 1/150 gr., Pot. iodide 1 gr. | Chicago | Caltalide, Tab. |
| 300 mg. | Carlisone acetate 2.5 mg., aluminum hydroxide gel dried 120 mg., Cal. ascorbate 60 mg., Cal. carb. 60 mg. | Pharm Co. Mossengill | Saltort, Tab. |
| 5 gr. | Menadione 0.125 mg. | | |
| 5 gr. | Mephensin 3 gr., saccharin 1/4 gr., phenobarbital 1/8 gr., Vit. C 5/6 gr., aluminum hydroxide 1 gr. | Fellows | Erqa, Tab. & liq. |
| 1/4 gr. | PABA 1/4 gr. | Geriatric | Garisol, Tab. |
| 1/4 gr. | PABA 1/4 gr., Vit. C 20 mg. | | |
| 5 gr. | PABA 5 gr., Vit. C 10 mg. | Ulmer | P-B-Sol, Tab. |
| 5 gr. | PABA 5 gr., Vit. C 10 mg. | Ulmer | P-B-Sol-C, Tab. |
| 5 gr. | PABA 5 gr., Vit. C 10 mg. | Warren-Teed | Hyslyte, Tab. |
| 5 gr. | PABA 5 gr., Vit. C 10 mg. | Kendall | Nucarsol, Tab. |
| 4 gr. | PABA 4 gr., Vit. C 20 mg. | Meyer-Besser | Purpasol, Tab. |
| 250 mg. | PABA 250 mg., Vit. C 20 mg. | Taylor | Salicin, Tab. |
| 5 gr. | PABA Sod. 5 gr., Vit. C 10 mg. | Sly | Cavipaba, Tab. |
| 5 gr. | PABA Sod. 5 gr., Vit. C 50 mg. | Ives-Cameron | Pabozyl, Tab. |
| 5 gr. | PABA Sod. 5 gr., vit. C 50 mg. | Armour | Armyl, Tab. |
| 1/4 gr. | PABA 1/4 gr., Vit. C 20 mg., calcichine 20 mg. | Robins | Pabalate, Tab. |
| 4 gr. | PABA 4 gr., Vit. C 20 mg., calcichine 1/250 gr. | Ulmer | P-B-Sol W/Calcichine |
| 250 mg. | PABA 250 mg., Vit. C 20 mg., physostigmine salicylate 0.25 mg., homatropine methylbromide 0.5 mg. | Taylor | Salical, Tab. |
| 1 gr. | Phenalthalain 1/2 gr., cascara 1/2 gr. | Siler Medical Corp. | Salisigmine, Tab. |
| 2.5 Gm. | Pot. bicarb. 3 Gm., sod. elb. 1.1 Gm. | McNeil | Biatco, Tab. |
| 5 gr. | Salicylamide 2 1/2 gr., sod. p-aminobenzoate 5 gr., Vit. C 50 mg., butobarbital sod. 1/2 gr. | Merrell | Alytine/oz. |
| 10 gr. | Sod. bicarb. 20 mg. | Allison | Bisolate, Tab. |
| 28 gr. | Sod. bromide 4 gr./oz., onitpyrine 24 gr./oz., caffeine 4 gr./oz. | Upjohn | Salicylonyl, Paw. |
| 5 gr. | Sod. lactate 1.5 Gm. | Tilden | Bramoto, Sol./oz. |
| 6 gr. | Vit. C, calcium, PAB 3 gr., menadione 0.5 mg. | Thompson | Lacta-Salicylate, Syr. |
| | | Paul Maney | Pabacee, Tab. |

- $\frac{1}{2}$ gr., calcichine 1/250 gr./Tab. Bot. 200s, 500s.
Use: Acute gout & arthritis.
- SOLACTHYL INJECTION.** (Squibb) Corticotrapin Inj., U.S.P. Vial (40 u./cc.) 5 cc.
Use: ACTH therapy.
- SOLAGEST.** (Canright Corp.) Pancreatin 10 $\frac{1}{2}$ gr., hyoscyamine sulf. 0.072 mg., atropine sulf. 0.013 mg., hyoscyne HBr 0.0047 mg., pentobarb. sad. 10.8 mg./Tab. Bot. 100s.
Use: Dyspepsia.
- SOLAHIST.** (Vitamax Corp.) Pyrilamine maleate 0.5%, ephedrine hydrochloride 0.75%, chlorbutanol 0.5%. Nasal spray.
Use: Vasocostrictor and antihistaminic.
- SOLANITAL ELIXIR.** (Smith-Dorsey) Phenobarbital 2 gr., Hyoscyamus ext. $\frac{1}{2}$ gr., belladonna ext. $\frac{1}{2}$ gr., hyoscyamine sulfate 1/200 gr., atropine sulfate 1/130 gr., scopolamine HBr 1/400 gr./fl. oz. Bot. 1 pt., 3 gal.
Use: Antispasmodic & sedative.
- SOLANITAL B-C CAPSULES.** (Smith-Dorsey) In addition to ingredients in tablets, Vit. B complex w/Vit. C has been added. Cap. Bot. 100s, 500s, 1000s.
Use: Antispasmodic w/vitamins.
- SOLANITAL TABLET.** (Smith-Dorsey) Phenobarbital $\frac{1}{2}$ gr., Hyoscyamus 1/12 gr., belladonna ext. 1/20 gr., hyoscyamine sulfate 1/1200 gr., atropine sulfate 1/300 gr., scopolamine HBr 1/2400 gr./Tab. Bot. 100s, 1000s.
Use: Antispasmodic & sedative.
- SOLARGENTUM, N.N.R. 1947.** See: Mild silver protein (Various Mfr.).
- SOL DE ORO.** (El Paso Pharm) Cap. Bot. 50s. Eye Drops Bot. $\frac{1}{2}$ oz.
- SOLESTRO.** (Merck Sharp & Dohme) Estradiol benzoate in oil. Vial (0.33 mg./cc.) 10 cc. (1 mg./cc.) 10 cc.
Use: Estrogen therapy.
- SOL-FERR.** (Jackson-Mitchell) Ferrous sulfate 125 mg., citric acid 56.5 mg./cc. Bot. 3 & 2 fl.oz.
Use: Hematinic.
- SOLFO-SERPINE.** (Paythress) Phenobarbital $\frac{1}{4}$ gr., colloidal sulfur $\frac{1}{2}$ gr., dispersed on colloidal bentonite w/reserpine 0.06 mg./Tab. Bot. 100s, 500s.
Use: Hypertension.
- SOLFOTON.** (Paythress) Phenobarbital $\frac{1}{4}$ gr., colloidal sulfur $\frac{1}{2}$ gr., dispersed on colloidal bentonite./tab. & cap. Bot. 100s, 500s.
Use: Orally, mild sedation.
- SOLONAL, N.N.D. 1952.** (Schering) Aureothiolglucate, N.F. Amp. 10 mg., 25 mg., 50 mg./T.5 cc., Sox 1s, 12s, Vial 10 cc., 10 mg., 50 mg., 100 mg./cc. Box 1s
Uses: I.M., gold therapy, antiarthritic.
- SOLGUATONE.** (Pitman-Moore) Strychnine hypophosphate 1/32 gr., Pat. guaiacol sulfonate T6 gr., creasote sulfonate 16 gr., hypophosphites of Cal., Iron, Mang. & Pot 5 gr./dr. Bot. 1 pt., 1 gal.
Use: Expectarant & tonic.
- SOLINIMENT.** (Breon) Methyl salicylate 12.5%, menthol 2.5%, eucalyptol 2.5%, camphor 5.5%, in axycholesterinated aini. base. Tube 1 oz. Jar 1 lb., 5 lb.
Use: Counterirritant.
- SOLMOIES.** (Merck Sharp & Dohme) 5 gr. each ammonium, sod. & pot. bromide/tab. Also avail. in $\frac{1}{2}$ amount. Sat. 25s.
Uses: Sedative.
- SOLSEO.** (Jackson-Mitchell) Pentobarbital 1/16 gr., phenobarbital $\frac{1}{2}$ gr., hamotrapine methylbromide 1/480 gr./0.6 cc. Bot. 15 cc.
Use: Pediatric sedative.
- SOLSTIGMINE.** (Siler Medicine) Sod. salicylate 0.25 Gm., p-aminobenzoic acid 0.25 Gm., ascorbic acid 20 mg., physostigmine salicylate 0.25 mg., hamotrapine methylbromide 0.50 mg./Tab.
Uses: (2 q.i.d., o.c., h.s.) Antirheumatic.
- SOLTABS.** (Commercial Solvents) Cryst. Pot. penicillin G. Tab. 50,000 u., 100,000 u., 250,000 u. Pkg. 24s, 100s.
Use: Antibiotic therapy.
- SOLU-A-50.** (New Mexico Pharm. Co) Vit. A 50,000 u./Cap. Bot. 100s, 1000s.
Use: Acute vitamin A deficiency.
- SOLU-B.** (Upjohn) Thiamine HCl 10 mg., riboflavin 10 mg., pyridoxine HCl 5 mg., Cal. pantothenate 50 mg., nicotinamide 250 mg., B-12 25 mcg./Vial. Vial 10 cc., w/5 cc. amp. of sterile distilled water or Mix-a-Vial 2 cc. Also avail. w/ascorbic acid 500 mg./Vial 10 cc. or 5 cc. Mix-a-Vial.
Use: Vitamin B complex therapy.
- SOLU-B 5X.** (Upjohn) Thiamine HCl 50 mg., riboflavin 50 mg., pyridoxine HCl 25 mg., Cal. pantothenate 250 mg., nicotinamide 1250 mg., B-12 125 mcg./Vial. Vial 30 cc. Pkg. 5s, 25s
Use: Vitamin B complex therapy.
- SOLU-B-PORTE.** (Upjohn) Thiamine HCl 250 mg., riboflavin 50 mg., Vit. C 1000 mg., B-6 50 mg., nicotinamide 1250 mg., sod. pantothenate 500 mg./10 cc. Mix-a-Vial.
Use: I.M. or I.V. Vitamin therapy.
- SOLU-CORTEF, N.N.D. 1961.** (Upjohn) Hydrocortisone sod. succinate buffered w/sod. biphosphate & sod. phosphate. Vial (100 mg) 2 cc. with or without amp. of water. Mix-O-Vial (100 mg., 250 mg)

Use: (V) Treatment of acute adrenal cortical insufficiency

SOLULEXIN. (Upjohn) Thiamine HCl 10 mg, riboflavin 10 mg, pyridoxine HCl 5 mg, Cal pantothenate 50 mg, nicotinamide 250 mg, liver conc 250 mg / 5 cc Vial 10 cc, w/5 cc amp sterile water Pkg 5s, 25s
Use: Vitamin B complex therapy

SOLULEXIN FOLIC ACID. (Upjohn) Folic acid 5 mg, thiamine HCl 10 mg, riboflavin 10 mg, pyridoxine HCl 5 mg, Sod pantothenate 50 mg, nicotinamide 250 mg / 5 cc Vial 10 cc, with 5 cc amp sterile water Box 5s 25s 50s 100s
Use: Vitamin B complex w/folic acid therapy

SOLU-MEDROL, NND 1962 (Upjohn) Methylprednisolone Sod succinate Mix O Vial 40 mg with 1 cc of water
Use: Treatment of emergency hyperseesalivity

SOLUTHRICIN, NND 1962 (Merck Sharp & Dohme) Tyrothricin 0.5 mg / cc Bot 240 cc
Use: Topically gram positive & gram negative organism infections

SOLUTHRICIN, NND 1961 (Merck Sharp & Dohme) Tyrothricin 25 mg / cc Vial 10 cc, 20 cc

SOLUTION G, SUBY'S See: Serologic Soln O (Abbott)

SOLU-TROL (Schieffelin) lyophilized plasma to make 0.5 ml Tubes 5s with one 3 ml tube of diluent 10s with two 3 ml tubes of diluent
Use: Determination of prothrombin time

SOLU-ZYME (Upjohn) Vit B 12 25 mcg folic acid 5 mg thiamine HCl 10 mg, riboflavin 10 mg pyridoxine HCl 5 mg, Sod pantothenate 50 mg, nicotinamide 250 mg / Vial Vial 10 cc Also with 5 cc amp sterile water also avail w/ascorbic acid 500 mg / Vial Also 2 cc Mix a Vial
Use: Vitamin B complex therapy

SOLVETS. (Lilly) Bacitracin 2500 u / Tab Bot 24s, 500s
Use: Preparation of topical solutions for aphthoidic or intranasal therapy

SOLVILITH (E S Miller) Salicylamide 600 mg, vit A 10 000 u / tab Bot 100s 1000s
Use: (I t d) Phosphocalcemia

SOLVIFLEX-STRONG (Rorer) Th amine HCl 10 mg riboflavin 2 mg niacinamide 50 mg, Cal pantothenate 1 mg pyridoxine HCl 1 mg w/urea 100 mg / cc Vial 10 cc
Use: Vitamin B complex therapy

* **SOLYPERTINE TARTRATE** (Winthrop Labs) 7 {2 [4 (a methoxyphenyl) 1 piperaziny] ethyl} 5H 1,3 diaxato [4,5 f] indole tartrate
* Generic name recognized by N F U S P or A M A

SOMA, NND 1962 (Wallace) Isomepramate, Carisoprodol 350 mg / Tab Bot 50s Caps 250 mg / Cap 50s

Use: Treatment of inflammatory conditions of muscles and joints & muscle relaxant

SOMA COMPOUND. (Wallace) Carisoprodol 200 mg, acetophenetidin 160 mg, caffeine 32 mg / Tab Bot 50s

Use: Analgesic with muscle relaxant

SOMA COMPOUND W/CODEINE. (Wallace) Carisoprodol 200 mg, acetophenetidin 160 mg, caffeine 32 mg, codeine phosphate 16 mg / Tab Bot 50s

Use: Analgesic

SOMACORT. (Wallace) Carisoprodol 350 mg, prednisolone 2 mg / Tab Bot 50s

Use: Muscle Relaxant, Steroid

SOMAGEN. (Upjohn) Milk proteins, yeast B liver conc w/thiamine HCl 5 mg, riboflavin 5 mg, pyridoxine HCl 2 mg, Cal pantothenate 10 mg, nicotinamide 30 mg, folic acid 2 mg / 100 Gm w/minerals Bot 1 lb, 3 lb

Use: Dietary supplement

SOMATAL. (Person & Covey) Chloral hydrate 200 mg, d-amphetamine sulfate 5 mg, mephenesin 200 mg / cap Bot 100s, 1000s

Use: I t t i d; sedative with muscle relaxant

SOMATOTROPIN. Growth hormone derived from the anterior pituitary gland

SOMATOVITE (Purdue Frederick) Reserpine 0.05 mg, vit B 12 10 mcg, B 1 10 mg / tab Bot 100s

Use: I t t i d; appetite stimulation

SOMATOVITE LIQUID (Purdue Frederick) Thiamine mononitrate 10 mg, Vit B 12 10 mcg, reserpine 0.05 mg / 5 ml Bot 12 fl oz

Use: Promote weight gain

SOMBROL (Columbus) Pentobarbital Sod. $\frac{3}{8}$ gr, phenobarbital Cal $\frac{3}{8}$ gr / Cap Bot 100s, 500s, 1000s

Use: (I 2 p r n) Sedative

SOMBULEX. (Schenley) Hexobarbital N methyl cyclohexenyl methylbarbituric acid 4 gr / tab Bot 100s

Use: (I 2 h s) Sedative

SOMINAT (National Drug) Dichloralphenazine Dichloralantipyrene 600 mg equiv to chloral hydrate 382 mg antipyrene 218 mg / Tab Bot 100s

Use: Relief of insomnia

SOMNALERT. (Warren Teed) 5 (1 Cyclohexenyl) 1,5 d methylbarbituric acid 250 mg / Cap Bot 100s 5000s

Use: Sedative

SOMHESIN See: Darmlon (Schering)

- SOMNICAPS.** (Am. Pharm. Co.) Methapyri-
lene HCl 25 mg./Cap. Vial 36s.
Use: Sedative.
- SOMNOS.** (Merck Sharp & Dohme) Chloral
hydrate. Cap. (3¼ & 7½ gr.) Rot. 100s.
Elixir 25 gr./fl. oz., Bot. 1 pt., 1 gal.
Use: Sedative.
- SOMONAL.** See Phenobarbital (Various
Mfr.).
- SOMORSED.** (Columbus) Chloral hydrate 3¼
gr., mephenesin 2 gr., scopolamine HBr
0.065 mg./Cap. Bot. 100s, 1000s.
Use: (1-2 m.p.) Sedative.
- SOMTROL.** (Columbus) Pentobarbital Sod.
1½ gr., phenobarbital ½ gr., amphetamine
sulfate 5 mg./Tab. Rot. 100s, 1000s.
Use: (1 h.s.) Sedative.
- SONAZAR.** (Tutag) Mephenesin 250 mg.,
amobarbital 25 mg., 3-(o methoxyphenoxy)
propionediol-1,2. 50 mg./Tab. Bot. 100s,
1000s.
Use: Relief of tension and anxiety.
- SONERYL.** See Butethal, N.F.
- SONILYN.** (Mollinckrodt) Sulfachloropyrida-
zine 0.5 Gm./Tabs. Bot. 100s.
Use: Sulfonamide for G.U. infections.
- SOOTHENE.** (Paul Money) Tyrothricin 5 mg.,
benzocaine 3%, In vit. A and O aint. Oint.
Tube 1 oz.
Use: Topically, antiseptic & protective.
- ROPENA-MRT.** (Thompson) Salicylamide 4
gr., acetophenetidin 2 gr., pentobarbital
¼ gr., phenobarbital ¼ gr., mephenesin
2 gr., caffeine ¼ gr./Tab. Bot. 50s, 100s,
1000s. Also available w/phenobarbital ½
gr.
Use: Analgesic.
- SOPRONOL OINTMENT.** (Wyeth) Sod. propi-
onate 12.3%, propionic acid 2.7%, Sod.
caprylate 10%, zinc caprylate 5%, dioctyl
Sod. sulfosuccinate 0.1%, inert ingredients
69.9% Including n-propyl alcohol 10%.
Tube 1 oz., 4 oz.
Use: Dermatophytosis (Athlete's foot).
- SOPRONOL POWDER.** (Wyeth) Sod. propi-
onate 5%, zinc propionate 5%, sod. capry-
late 10%, inert ingredients 75%. Con 2
oz., 5 oz.
Use: Prophylactic fungicide.
- SOPRONOL SOLUTION.** (Wyeth) Sod. propi-
onate 12.3%, Sod. caprylate 10%, propi-
onic acid 2.7%, dioctyl Sod. sulfosuccin-
ate 0.1%, inert ingredients 74.9% includ-
ing n-propyl alcohol, 12.5%. Bot. 2 fl. oz.
Use: Dermatophytosis.
- SOR-B-12.** (Bentex) Ferrus sulfate 195 mg.,
thiamine HCl 10 mg., pyridoxine HCl 2 mg.,
folic acid, 0.5 mg., 5-12 15 mcg., sorbitol
gr./Tab. Sol. 100s.
Use: Vit. B deficiencies.
- SORREFAVIN OINTMENT.** (Foster-Dock) Cal-
tan seed oil, stearine, petrolatum, camphor,
carbolic acid, zinc oxide. 1 lb. jar, 5 lb.
tin.
Use: Soothing, analgesic, emollient in skin
conditions.
- *SORBIDE NITRATE.** 1,4:3,6-Dianhydro-
sorbitol dinitrate.
- SORBITAT-K.** (Pfizer) Potassium sorbate
Pkg. bulk.
- RORBITAN MONO-OLEATE POLYOXYETH-
YLENE DERIVATIVES.** See Polysorbate 80,
U.S.P.
- SORRITANS.**
See Polysorbate 80, U.S.P.
Topisol. (Texas Pharm. Co)
- *SORBITOL SOLN., U.S.P., A.D.R.** (Abbott)
Amp. 50%, 50 cc., 6s, 25s, 100s.
See: Sigmal Enema.
Sorbo (Atlas Paw)
Use: Diuretic; dehydrating agent.
- W/Phenylpropanolamine HCl, phenindamine
tartrate, chlorpropenpyridamine moleale**
See: Naloxone Elixir, Liq. (Carrick)
- SORBO.** (Atlas Paw.) Sorbitol, N.F.
- SORBOOUL.** (White Lab.) Polycarbophil 0.5
Gm., Thiexinal Methylbromide 15 mg./
Tab. Bot. 50s, 250s
Use: Diarrhea.
- W/Neomycin Tabs, Polycarbophil 0.4 Gm.;
Ithexinal methylbromide 15 mg.; neo-
mycin (from sulfate) 150 mg./tab. Bot 15s.**
Use: Diarrheal disorders.
- SORBUS.** (Dalin) d-methorphan HBr 10
mg., glycerol guaiacolate 100 mg., potas-
sium citrate 85 mg., citric acid 35 mg.,
ipeacac fluid extract 0.05 min /5 cc. Bot. 4
oz., 16 oz.
Use: For coughs.
- SORETHYAN (20) MONO-OLEATE.** See
Polysorbate 80, U.S.P.
- RORICIN, N.N.R. 1954.** (Merrell) Sod. ricin-
oleate. Tab. 5 gr., Bot. 100s. Sol. 2%, Vial
20 cc.
W/bite salts: Sod. ricinoleate 5 gr., bite
salts ½ gr./Tab. Bot. 100s.
Use: Intestinal distress.
- SORIBEX.** (Kahlenberg) Mercury bialeate
2%, chloral hydrate 2.2%, zinc oleate, and
olein stearin base. Tube ½ oz
Use: Psoriasis
- SORLA-BILEIN.** (Abbott) Sorlate or polysor-
bate 80. 0.4 Gm. bilein (dried & purified
ox bile) 30 mg., dehydrocholic acid 30
mg./Cap. Bot. 100s, 1000s.
Use: (2 q Ld) Fat emulsifying properties.

SORLATE. (Abbott) Not listed Polysorbate 80, U.S.P., Sorbitan mono oleate polyoxyethylene derivative Cap 0.5 Gm Bot 100s, 1000s liq, Bot w/dropper 50 cc.

Use: (4-6 Gm, daily) Fat emulsification & absorption

W/Bilein & dehydrocholic acid See Sorla Bilein, Cap (Abbott)

SOROPON. (Purdue Frederick) Triethanolamine polyepoxide cococate condensate 12%, tythrucin 0.1% in propylene glycol Bot 4 oz

Use: Cradle cap Seborrheic infections

SORPARIN. (McNeil) Sorbus aucuparia ext 3 gr/Tab Bot 100s, 1000s

Use: (2 lid) Cholerae.

SORSIS-ALPHA. (Ar Ex) Ammoniated mercury, salicylic acid, tar phenol in non lipidal base Oint Tube 1 oz

Use: Softening cream to aid in removal of scales

SORSIS-BETA. (Ar Ex) Ammoniated mercury, Ichthomol, tar, boric acid in nonlipodal base Oint 1 oz

Use: Aid healing of lesions

SOTRADECOL SODIUM, NND 1960 (Wallace & Tiernan) Sod tetradecyl sulfate Sol 1%, 3%, 5%, Vial 20 cc

Use: Varicose veins

SOYA LECITHIN. (Glidden) Soybean extract Pow Pkg B oz.

Use: Phosphorus therapy

SOYALOID. (Dome) Soyaloid (50% colloidal protein) W/irritation reducing action of PBP (polyvinylpyrrolidone) 2% Box 5, Pkg 3 oz

Use: Acute and subacute dermatitis

SOYA PHOSPHATIDES (Central Soya Co) Lecithin 29.5%, chemical cephalin 29.5%, inositol phosphohde 31.6%, carbohydrates, soybean oil Polyunsaturated fatty acids 2.75 Gm, monounsaturated acids 1.25 Gm, saturated acids 0.60 Gm, choline 250 mg, inositol 250 mg/teaspoonful Bot 8 oz, 1 lb

Use: Cholesterol reduction in treatment of cardiovascular diseases

SOYBEAN LECITHIN W/Choline, inositol & other phosphatides See: Sectin, Cap (Tes tagar)

SOYBORO. (Dome) Aluminum sulf, cal acetate, boric acid, soyaloid (colloidal soybean complex) each pow packet added to one pint of ordinary tap water instantly provides the equivalent of 120 Burrows solution

SOY-DOME CREME. (Dome) Colloidal fraction of soy bean, pH 5 Tube 2 oz, Jar 1 lb

Use: Acid restoring skin cleaner

* Generic name recognized by N.F., U.S.P. or A.M.A.

SOY-DOME LOTION. (Dome) Hexachlorophene 3%, polyvinylpyrrolidone 1%, colloidal soy bean complex, in a blond lotion base/cc Squeeze Bot w/appli 6 oz

Use: Cleansing of irritated skin.

SOY-SITZ. (Dome) Colloidal soy bean complex, polyvinylpyrrolidone 1%, aluminum sulfate, calcium acetate, sodium sesqui carbonate, tetrasodium pyrophosphate, lauryl sulfoacetate Packets/5s 2 oz

Use: Dermatologic

SPANOREX D.L.A. (Cooper, Tinsley) Vit B 1 15 mg, B-6 6 mg, B 12 30 mcg, L-lysine, 50 mg/Tab Bot 30s

Use: Appetite Stimulant

SPARINE HCl, NND 1962 (Wyeth) Promazine HCl 10 (gamma Dimethylamino n propyl)phenothiazine HCl Tab. 10 mg, Bot 50s Tab. 25 mg, 50 mg, 100 mg, Bot 50s 500s Tab 200 mg, Bot 50s, 500s Amp (50 mg/cc) 1 & 2 cc, 10 cc Amp (25 mg 1 cc) Vials 10 cc Syr. 10 mg/5 cc, Bot 4 oz, Tubex (50 mg/cc) 2cc; Vial 10 cc Oral liq. 100 mg/cc. and 30 mg/cc. Bot 4 oz

Use: Anxiolytic

W/Meprobamate See Prozine (Wyeth)

* **SPAROSMYCIN.** (Upjohn) Agent derived from cultures of Streptomyces sparsogenes

Use: Anti neoplastic agent

SPARTASE. (Wyeth) Potassium aspartate 250 mg, magnesium aspartate 250 mg/Tab Bot 100s

Use: Fatigue syndrome

* **SPARTEINE SULFATE, NF VIII** (March Sharp & Dahme) Pow, Bot. ½ oz, 1 oz.

See Tocamine sulfate (Trent)

Use: Cardiac arrhythmias

W/Camphor See: Camphoral Sparteine Amp (Torigan)

SPARTOLOXIN. (Spartan) Mephesisin 250 mg & 500 mg/Tab 100s, 1000s

SPASCAPS. (Nyco Labs) Mephobarbital 65 mg, homatropine methyl bromide 5 mg, atropine methyl nitrate 1 mg, hyoscine HBr 0.02 mg/Tab 1 D Bot 100s

Use: Antispasmodic

SPASLOIDS. (G F Harvey) Hyoscyamine sulfate 0.140 mg, atropine sulfate 0.0195 mg, hyoscine HBr 0.0065 mg, phenobarbital ¼ gr/tab Bot 1000s, 5000s

Use: Antispasmodic & sedative

SPASMABAR. (Medco) Atropine sulf 1/66 gr, phenobarbital ¼ gr/Tab Bot 30s 100s

Use: Sedative w antispasmodic action

SPASMALGIN. (Roche) Papaverine HCl 3 gr, pantopon 1/6 gr, atrolol (atropine sul

- lots) 1/60 gr. Tab. or cc. Amp. 1 cc., 6s.
Tab. Tube 20s; Sol. 100s.
Use: Antispasmodic & sedative.
- SPASMANOL.** (Bullington) Aprabarbitol
(allyl-isopropyl barbituric acid) 50 mg.,
homatropine methylbromide 2 mg., hyo-
scine HBr 0.0065 mg./Tab. Bot. 100s, 1000s.
Use: Antispasmodic & sedative.
- SPASMASORB.** (Palmedico, Inc.) Adiphenine
hydrochloride .025 gm., pentobarbital
sodium .0075 gm., pharמושorb .090 gm.,
aluminum hydroxide .075 gm./Tab. Bot.
100s.
Use: Antacid-Antispasmodic.
- SPASMOLYD.** (Columbus) Aminopentamide
hydrogen sulfate 0.5 mg., butobarbital sod.,
15 mg./Tab. Bot. 100s, 1000s
Use: Hypertonic and spastic states
- SPASMOLYN, N N D.** 1962. (Heun) Mephene-
rin, Tab. 0.5 Gm., Bot. 100s, 1000s.
Use: (1-2 t.i.d.) Muscle relaxant.
- SPASMOLYTIC AGENTS.** See: Antispas-
modics.
- SPASMOPHEN.** (Lannett) Phenobarbital 15
mg., hyoscyamine sulf. 0.1037 mg., atropine
sulf. 0.0194 mg., hyoscyne HBr 0.0065 mg./
Tab. Bot. 500s, 1000s, 5000s.
Use: Antispasmodic-sedative.
- SPASNIL.** (Cala) Adiphenine HCl 75 mg./
Tab. Bot. 100s, 500s, 100s. Also oval. with
phenobarbital 20 mg. and Adiphenine HCl
50 mg./Tab.
Use: Antispasmodic
- SPASTICEN.** (Metropolitan) Phenobarbital ¼
gr., belladonna ext. ¼ gr., hyoscyamus
ext. ¼ gr./Tab. Tab. Gray, Sol. 100s,
1000s, 5000s.
Use: Antispasmodic.
- SPASTICOL P. R.** (Key Corp.) Homatropine
methylbromide 2.5 mg., phenobarbital ¼
gr./Tab. Bot. 100s, 1000s
Use: Antispasmodic
- SPASTICOL S.A.** (Key Corp.) Homatropine
methylbromide 5 mg., phenobarbital ¼ gr.
/Tab. Bot. 50s, 100s, 1000s
Use: Antispasmodic & sedative.
- SPASTILOWE CTR.** (Lawa) Hyoscyamine sul-
fate 0.3 mg., atropine sulfate 0.06 mg.,
hyoscyne HBr 0.02 mg., phenobarbital 50
mg./Cap. Bot. 100s, 250s.
Use: Smooth muscle spasms.
- SPASTOLATE.** (Nysco Labs) Phenobarbital
16.2 mg., hyoscyamine sulf. 0.1037 mg.,
atropine sulf. 0.0194 mg., hyoscyne HBr.
0.0065 mg./cap. Bot. 100s, 1000s.
Use: Antispasmodic w/sedative.
- SPASTOSED.** (Chicago Pharm.) Phenobar-
bital 1.8 gr., atropine sulfate 1/500 gr.
- Col. carb. 3½ gr., Mag. carb. 2½ gr./
Tab. Bot. 500s, 1000s, 6000s, 13,000s.
Use: (1-2 t.i.d.) Antispasmodic & antacid.
- SPECIAL FORMULA NO. 1.** (Upjohn) Ace-
tophenetidin 1¼ gr., acetylsalicylic acid 1¼
gr., camphor (monobromated) ½ gr., cof-
feine ¼ gr., phenolphthalein 1/5 gr., atro-
pine sulfate 1/1000 gr./Tab. Bot. 3000s,
10,000s.
Use: Symptomatic treatment of the acute
common cold.
- SPECIAL FORMULA NO. 2.** (Upjohn) Ace-
tophenetidin 2 gr., quinine sulfate ¼ gr.,
camphor ¼ gr., podophyllin 1/40 gr.,
alain 1/16 gr., atropine sulfate 1/2000 gr./
Tab., or Cap. Tab. Bot. 100s, 1000s, Cap.
1000s,
Use: Symptomatic treatment of the acute
common cold.
- SPECIAL RESIN NO. 645.** (Smith, Kline &
French) A cotton resin to control hyper-
kalemia. Under study. Bot. 1 lb.
- SPECIAL VITAMIN FORMULA.** (Squibb)
Vit. A 5000 u., D 800 u., thiamine HCl 2
mg., riboflavin 3 mg., niacinamide 20 mg.,
ascorbic acid 75 mg./Cap. Bot. 100s, 1000s
Use: (1 daily) Supplement to diet.
- SPECTINOMYCIN.** (Upjohn) An antibiotic
isolated from broth cultures of *Streptomyces*
spectabilis. See: Tetracycline.
- SPECTRO-RIOTIC.** (American Pharmaceutical
Co.) Bacitracin 500 u., neomycin sulfate 5
mg., polymyxin B sulfate 5000 u./Gm. Oint.
Use: Infection in cuts, wound, etc.
- SPECTROCIIN.** (Squibb) Oint. Neomycin 2.5
mg., gramicidin 0.25 mg. Tube 15 Gm.
Ophth. oint. Each Gm. contains 2.5 mg.
neomycin & 0.25 mg. gramicidin. Tube 3.6
Gm. Spectrocin-Ts Each raspberry-flavored
troche contains 2.5 mg. neomycin, 0.25 mg.
gramicidin, 10 mg. benzocaine as topical
anesthetic. Vial 12s. Spectrocin-Ns: Each
cc. aq. susp. contains 5 mg. neomycin, 5
mcg gramicidin, 2.5 mg. (¼%) phenyl-
ephine HCl as vasoconstrictor. Bot. 15 cc.
w/dropper.
Use: Effective against gram-positive &
gram-negative organisms.
- W/Hydrocortisone acetate. See: Florinet-S,
Preps. (Squibb)
- SPENSIN.** (Ives-Cameron) Activated attapul-
gite 3 Gm., pectin in special alumina gel
270 mg./oz. Bot. 6 fl. oz.
Use: Antidiarrheal.
- SPENSIN-PS. SUSP.** (Ives-Cameron) Dihydro-
streptomycin sulf. 300 mg., polymyxin B
sulf. 120 M u., attapulgite activated 3 Gm.,

- pectin 270 mg w/alumina gel/8 oz Bot 4 oz
 Use Antidiarrheal
- SPENSIN-PS TABLETS** (Ives Cameron) Dihydrostreptomycin sulfate equivalent to 75 mg of the base, polymyxin B sulfate 25,000 u, Altopulgit 350 mg, pectin 45 mg, hydrated alumina powder 70 mg /Tab Bot 48s
 Use: Treatment of diarrhea
- SPERMINE** Diaminopropyltetromethylene See Latane, Tab (Westerfeld)
- ***SPIDER-BITE ANTIVENIN, USP, NND 1962** See Antivenin Black Widow Spider
- SPINOCAINE.** (Winthrop Labs) Viscous 10% sol of Novocain w/specific gravity lower than that of spinal fluid, Amp 3 cc, Box 10
 Use Spinal anesthetic
- SPIRA-DINE.** (Warren Teed) Aspirin soluble 66 gr /tsp 1 pt & 1 gal
- ***SPIRAMYCIN.** (Ives Cameron Co) Antibiotic substance from cultures of *Streptomyces ombifaciens*
 Use Antibiotic
- SPIRAZINE HCl.** (Burroughs Wellcome)
 * Spiratrizine HCl
 Use Anthelmintic
- SPIRODON.** (Cutter Lab) See Tetratin
- ***SPIRONOLACTONE, NND 1962 3-43** oxo 7 α acetylthio 17 β hydroxy 4 androsten 17 α yl) propionic acid gamma lactone See Aldolone A, Tab (Searle)
- W/Hydrochlorothiazide** See Aldactazide A Tabs (Searle)
- ***SPIROTRIAZINE HCl.** (Burroughs Wellcome) 2,4 diamina 5(p-chlorophenyl) 9 methyl 1,3,5 triazaspiro[5.5]undeca 1,3-diene HCl
 Use Anthelmintic
- SPONTIN, NND 1962** (Abbott) Ristocelin (An antibiotic) Vial 500 mg lyoph pow
 Use: (IV) Infections
 Gram-positive infections
- SPOBOSTACIN (Ortho) (Chlordantain).** 3 trichloramethylthio 5 (1 ethyl)amylthydon tain 1% Cream Tube 95 Gm with or without applicator
 Use Treatment of vaginal moniliasis
- SPOBOSTACIN LOTION (Ortho)** Chlordantain 1%, benzalkonium chloride 0.5% compounded with glyceryl monostearate, phosphoric acid lanolin, cetyl alcohol 1%, cholesterol polyethylene, glycol mono stearate, propylparaben, catanac methyl paraben, water /Bot 60 cc
 Use Dermatologic antifungal agent
- SPOBOSTACIN SOLUTION.** (Ortho) Chlordantain 1%, benzalkonium chloride 0.5%
 * Generic name recognized by NF USP or AMA
 compounded with Isopropyl alcohol 93%, water / Applicator bot 30 cc
 Use Dermatologic, antifungal agent
- SPADERM** (Westwood Pharmacol) Menthol 0.25%, phenol 0.5%, alcohol 30%, talc 15%, zinc oxide 15%, distilled water q-s Pkg aerosol 6 oz
 Use: Antipruritic and protective spray
- SPREADING FACTOR.** See: Hyaluronidase (Various Mfr)
- SQUALENE.** W/Allantoin, coal tar extract
 See Skaytas Lotion (Lee Colbert)
- SQUIBB RASE.** See: Plastibase, Oint (Squibb)
- SQUILL, GLYCOSIDAL MIXTURE.** See: Scillaren, Amp (Sandoz)
- S-R PENICILLIN.** (Parke, Davis) Procaine penicillin G 300,000 u, Sod penicillin G 100,000 u Vial 1 dose
- S-R-S PENICILLIN.** (Parke, Davis) Procaine penicillin G 300,000 u, Sod penicillin G 100,000 u, w/streptomycin sulfate 0.5 Gm Vial 1 dose
 Use Penicillin & streptomycin therapy
- S.T. 37.** (Merck Sharp & Dahme) Hexyl resorcinol 1,100 Bot 5 oz, 12 oz
 Use Antiseptic solution
- STABIL A** (Dumas Wilson) Vit A acetate Cap 25,000, 50,000, 100,000 u Bot 100s 1000s
 Use: Vitamin A deficiencies
- STABIL-A-PLUS** (Dumas Wilson) Vit A acetate 20,000 u Vit C 50 mg, niacin 18 mg, Vit B 2 4 mg, Vit B 12 2 mcg, non inhibitory intrinsic factor 1/15 unit/Cap Bot 100s
- STABIL-ACEE.** (Dumas Wilson) Vit A 100,000 u, ascorbic acid 100 mg /Cap Bot 100s 1000s
 Use: Vit A and C therapy
- STABILISOL.** (Squibb) Each 1.5 cc of susp. contains 0.195 Gm bismuth subsalicylate w/3% chlorbutanol & 0.03% mercurated chlorxylenol as antiseptic Bot 30 cc, 60 cc
 Use I G₂ therapy of syphilis w/penicillin where indicated
- STAMINE, NND 1961** (Tutag) Pyrimamine Maleate
- STAMYL.** (Winthrop Labs) Pancreas ext con taining trypsin, amylase, steapsin (lipase) w/hemicellulase & ax bite Tab, Bot 50s 500s
 Use Digestant
- STAN-A-DOME** (Dane) Dihydrotestosterone 0.3 Gm, Vit A 500,000 u /oz Tube 1/2 oz 1 oz
 Use Senile skin atrophy and ichthyosis senile palmar and plantar hyperkeratosis

- STANGEN MALEATE, N.N.D.** 1962. (Physicians Drug) Pyrilamine maleate. Tab. 25 mg., 50 mg., Bot. 100s, 1000s.
Use: Antihistaminic.
- STANNINE.** (Blue line) Tin proteinate. Tab. 22 mg., Bot. 100s.
Use: Tin therapy.
- STANNOXYL** (Anglo-French) Metallic tin & tin oxide. Amp. 2 cc., Bz. Oint. 1 oz. Tab. 80s, 500s, 1000s.
- STANNO-YEAST.** (Staddard) Tin proteinate. Tab. 65 mg., 100s, 500s, 1000s.
- *STANOLONE, N.N.D.** 1962. Androstano-17 β -ol-3-one.
See: Androlone, Vial. (National Drug)
Nedrol, Vial. (Pfizer)
- *STANOZOL.** (Winthrop Labs) 17 β hydroxy-17 α -methylandrosterone [3 2 c] pyrazole.
See: Winstrol (Winthrop Lab)
Use: Anabolic agent
- STAPHICILLIN.** (Bristol) Sodium Methicillin Sodium dimethoxyphenyl penicillin 1 Gm., equiv to 900 mg dimethoxyphenyl penicillin activity/Vial.
Use: Penicillin therapy.
- *STAPHICIDIN.** (Mead Johnson) An antibiotic derived from *Staphylococcus staphylolyticus*
Use: Antibiotic
- STAPHYLOCOCCUS AMBOTOXOID.**
(Squibb) Toxoid. Vial 5 cc.
Use: I.D., S.C. & I.M., staphylococcal immunization.
- STAPHYLOCOCCUS TOXOID, N.N.R.** 1951.
Lederle—Vial detoxified (100 u./cc., 1000 u./cc.) 5 cc. Vial, digest modified (10,000 u./cc.) 5 cc.
Parke, Davis—Vial, 1st strength (3 u./cc.) 5 cc.; 2nd strength (5 u./cc.) 5 cc.
Merck Sharp & Dohme—Vial, No. 1 (100 S.N.D./cc.) 5 cc.; No. 2 (1000 S.N.D./cc.) 5 cc.
Use: Localized staphylococcal infections.
- STAPHYLOCOCCUS TOXOID-VACCINE.**
(National Drug) *Staphylococcus vaccine* combined with a standardized staphylococcus toxin detoxified with formaldehyde to form a toxoid. Vial 6 ml.
- STAPHYLOCOCCUS VACCINE.**
Kirk—Vial 2000 million killed *S. aureus*/cc. 5 & 20 cc.
Lilly—Vial 2000 million killed each *S. aureus* & *S. albus*/cc. 5 cc.
- STARCH DERIVATIVE DUSTING POWDER,**
N.N.D. 1962. See: Bio-Sorb (Ethicon)
- STARLAC.** (Borden) Vit. A 60 I.U., thiamine HCl 0.24 mg., riboflavin 0.93 mg., niacin
- 1.06 mg., protein 36.5 Gm., lactose 52 mg., Cal. 0.3 Gm., Phos. 1 Gm., minerals (total) 8 Gm., milk fat 1 Gm., moisture 2.5 Gm./100 Gm. Pkg., 1 lb.
Use: Dietary supplement.
- *STATOLON.** (Lilly) Antiviral agent derived from *Penicillium stoloniferum*
Use: Antiviral agent.
- STATOMIN MALEATE TABLETS, N.N.D.** 1962. (Bowman) Pyrilamine maleate Tab. 25 mg., Bot. 100s.
Use: Antihistaminic.
- STEAP SIN W/Betaine & Thyroid.** See: Chylipose, Tab. (Columbus Pharm.)
W/Oxidized bile acids, ox bile, phenobarbital & homotropine methylbromide.
See: Oxochofin, Tab. (Columbus).
- W/Thyroid & betaine.** See: Chylipose, Tab. (Columbus)
- STEARODINE, N.N.R.** 1944. (Parke, Davis) Cal. iodostearate. Tab. 1/6 gr. Iodine, Bot. 100s, 500s.
Use: Simple gatter.
- STEARYL ALCOHOL, W/Patrolatum, lanolin.**
See: Eiderol (Elder)
- STEC LIN HYDROCHLORIDE, N.N.D.** 1962. (Squibb) Tetracycline HCl. Cap. (50 & 100 mg) Bot. 25s, 100s; (250 mg) Bot. 16s, 100s.
Use: Broad spectrum antibiotic.
- STEC LIN I.M.** (Squibb) Tetracycline HCl 100 mg. dry pow./vial. Reconstitute w/2 cc. of water.
- STEC LIN I.V.** (Squibb) Tetracycline HCl. 500 mg./vial.
- STEC LIN ORAL SUSP., N.N.D.** 1962. (Squibb) Tetracycline 1.5 Gm dry pow./bot. Susp. (250 mg/5 cc when mixed w/water) Bot. 1 oz.
- STEC LIN PEDIATRIC DROPS, N.N.D.** 1962. (Squibb) Cal. tetracycline 100 mg/cc. Bot. 10 cc.
- STEC LIN SYRUP, N.N.D.** 1962. (Squibb) Tetracycline HCl 125 mg/5 cc. Bot. 2 oz.
- STEINBERG'S SOLUTION.** See: Keromin Inj (Campbell)
- STELAZINE HCl, N.N.D.** 1962. (Smith, Kline & French) Trifluoperazine 10-[2-(1-methyl-4-piperazinyl)propyl]-2-trifluoromethylphenothiazine dihydrochloride. Tab. 1 mg., 2 mg., 5 mg., 10 mg., Bot. 50s, 500s.
Use: Tranquilizing agent.
- STELAZINE PARENTERAL, N.N.D.** 1962 (Smith, Kline & French) Trifluoperazine, 2 mg/cc Vials 10 cc. & Amps 1 cc. Oral Liq. (10 mg/cc) Bot. 4 oz.
Use: Rapid control of nausea & vomiting.
- STEMETIL.** See: Prochlorperazine.
- *Generic name recognized by N.F., U.S.P. or A.M.A.

- STEMUTROLIN.** (Lincoln) Chorionic gonadotropin 500 IU, Vit. B 1 25 mg, L glutamic acid 52.5 mcg/cc Vial 10 cc.
Use: Male senility
- STENEDIOL.** (Organon) Methylandrostenedial
Vial (25 & 50 mg/cc) 10 cc, 6s
Tab: (10 mg) Bot 30s, 100s; (25 mg) Bot 15s, 100s Sublingual (10 mg) Bot 30s, 100s; (25 mg) Bot 15s, 100s
Use (25 mg 2 to 5 times a week) Tissue building
W/Vit B complex, C, A, D & minerals. See: Covisten, Tab (Organon)
- STENIBELL.** (Bell Pharmacol) Methylandrostenedial Vial (25 mg, 50 mg/cc) 10 cc Tab 25 mg, Bot 100s
Use: Anabolic
- STENIFORM STRONGER.** (New Mexico Pharmacol Co) Methyl androstenedial 50 mg, ocaio 1%, procaio HCl 0.5%, thimiasol 1 10 000/cc Vial 10 cc
Use: Endocrine deficiency
- STENISONE.** (Organon) Prednisone 5 mg, methylandrostenedial 20 mg, magnesium trisilicate 100 mg, aluminum hydroxide gel 60 mg, calcium carbonate 70 mg, magnesium carbonate 40 mg, guar gum 66.6 Gm ass binder (Egionine) 30 mg /Tab Bot 30s, 100s
Use: Hormone & anabolic
- STENOGEN.** (Chemica) Cal glycerophosphate 3 gr, Sod glycerophosphate 2 gr, strychnine glycerophosphate 1/100 gr, gentian paw 1/2 gr /Tab Bot 100s, 500s, 1000s
- STENTAL EXTENTARS.** (Robins) Phenobarbital 3/4 gr (1/4 gr released for immediate action, 1/2 gr. released gradually) Tab Bot 100s, 500s
Use: Delayed action sedative
- STERAMINE-OTIC** (Mayrand) Hydrocortisone 5 mg, neomycin sulfate 6 mg, chlorpheniramine maleate 1 mg, cetalkansum chloride 0.05%, diperodon HCl 1%/cc drops
Use: Treatment of acute and chronic otitis media and externa
- STERAMINE-OPHTHALMIC.** (Mayrand) Polymyxin B sulfate 16,500 u, hydrocortisone 5 mg, neomycin sulfate 5 mg/cc Drop Bot 5 cc
Use: Inflammation
- STERANE.** NND 1962 (Pfizer) Prednisolone Tab (1 mg) Bot 100s; Tab 5 mg, Bot 20s, 100s; Inj (25 mg/cc) 5 cc
Use: Rheumatoid arthritis, allergies & various inflammatory skin conditions.
- STERAQ.** (Ascher) Desoxycorticosterone acetate Aqueous susp 5 mg/cc Vial 10 cc
Use: I.M. adrenal cortex therapy
- STERASAL-K.** (Mayrand) Prednisone 2.5 mg, salicylamide 486 mg, potassium chloride 25 mg, Vit C 50 mg /Tab Bot 100s, 500s
Use: Treatment of painful disorders involving muscles
- STRAZOLIDIN.** (Geigy) Phenylbutazone 50 mg, prednisone 1.25 mg, dried aluminum hydroxide gel 100 mg, magnesium trisilicate 150 mg, hamatropine methylbromide 1.25 mg /Cap Bot 100s
Use: Treatment of rheumatoid arthritis
- STERCULIA GUM, N.P., ADR. 1955** See: Karaya Gum (Penick)
W/Mag trisilicate See: Bossaron Plain (Merrell)
W/Mag trisilicate, cascara See: Bossaron with Cascara (Merrell)
- STERICIDE CHLORIDE.** See: Ruson Chloride (Northwest)
- STERI-DEX MEDICATED.** (Lehn & Fink) Sulfonated alkyl benzenes, Salicylic acid Citric acid Hexachlorophene, Alcohol 30% volume Throwaway pads 42s
Use: Acne
- STERI-EZE** (Buffington) Diadol 1/4 gr, mephesisin 5 grs, N-acetyl-p-aminophenol 5 grs, hamatropine methyl bromide, 1.2 mg /Tab Bot 100s
Use: Musculo-skeletal pain
- STERILE TROPICAL SODIUM, U.S.P.** ADR 1955, See: Pentothal Sodium (Abbott)
- STERISIL, NND 1962** (Warner Chilcott) Hexetidine 5 Amino 1,3 bis(beta ethylhexyl)-5 methylhexahydropyrimidine Vaginal gel 0.1%, Tube 1 1/2 oz & 6 appl
Use: Vaginal infections of bacteria, trichomonas & monilia
- STERISOL** (Lactona) Hexetidine solution Bot 7 oz
Use: Antiseptic
- STEROGYL.** (Eoussel) Calciferol, Vit. D 2D Amp (600,000 u) 1.5 cc, Box 2s, 6s
Use: I.M.; Vitamin D deficiencies
- STEROID DETERMINATION.** See: Ketodase, Sol (Warner Chilcott)
- STEROLONE, NND 1962** (Ewell) Prednisolone 1 mg, or 5 mg /Tab Bot 60s, 250s.
Use: Rheumatoid arthritis
- STEOSAN, NND 1962** (Geigy) (Chloroquinolone) 5,7 dichloro 8 hydroxyquinolone Cream 3%, Tube 30 Gm Oint 3%, Tube 3D Gm; Jar 1 lb
Use: Topical; superficial pyogenic & mycotic skin disorders
- STEOSAN-HYDROCORTISONE** (Geigy) 5,7 Dichloro-8 hydroxyquinolone 3%, hydrocortisone 1% Oint Tube 5 Gm, 20 Gm
Use: Treatment of skin disorders

- STEBOTATE MYOSOL.** (Ulmer) Testosterone 25 mg./cc.
Use: Inhibit lactation.
- STEBOTRIL.** (Schering) Prednisone 2.5 mg., Trilofan 2 mg./Tab. Bot. 30s, 100s.
Use: Corticoid-tranquilizes therapy.
- STETRIC.** (Smith-Dorsey) Dioctyl sod. sulfosuccinate 50 mg., hesperidin complex 50 mg., Vit. C 50 mg., A 2000 u., D 133 u., K 0.5 mg., B-1 1 mg., B-2 1 mg., B-6 2 mg., nicotinamide 10 mg., cal. pantothenate 2.5 mg., folic acid 0.5 mg., B-12 1 mcg., cal. carbonate 500 mg., ferrous sulf. 20 mg., w/trace minerals/Tab. Bot. 100s, 500s
Use: Prenatal vitamin & mineral supplement
- STIBAMINE GLUCOSIDE, N.N.D. 1951.** See: Neostom, Paw. (Burlington Wellcome)
- STIBOPHEN, U.S.P. Inj., U.S.P., N.N.D. 1958.** See: Fudin, Amp., N.N.E. (Winthrop Labs)
Use: Antiprotazoan & antischistosomal.
- SEIG, (Hart) Physostigmine salicylate 1 mg., homotriprine methylbromide 2 mg./cc. Vial 10 cc.**
- STIGMANOL.** (Gold Leaf) Neostigmine methylsulfate, Amp. (1:2000) 1 cc., 6s, 25s, 100s. Vial (1:1000, 1:2000) 10 cc.
Use: Parasympathetic stimulant.
- STIGMONENE BROMIDE, N.N.D. 1962.** (Warner-Chilcott) Benzpyrimium bromide. Amp. 1:500 sol., 1 cc.; 1:2000 sol., 1 cc.; 1:1000 sol., 5 cc.; 1:2000 sol., 10 cc.
Use: Parasympathetic stimulant.
- STILBAL.** (Sreon) Diethylstilbestrol 0.2 mg., phenobarbital 1/4 gr./Tab. Bot. 100s, 500s, 1000s.
Use: Menopausal disorders.
- STILBAMIDINE ISETHIONATE, N.N.E. (Merrill) 4,4'-stilbenedicarboxamide isethionate.** Amp. 150 mg. paw. See also: Hydroxystilbamidine. (Merrill) See also: Hydroxystilbamidine isethionate.
- Use: T.V. Blastomycosis & leishmaniasis.
- STILBESTROL, N.N.D. 1962.** See: Diethylstilbestrol, U.S.P. (Various Mfr.)
- STILBESTRONATE.** See: Diethylstilbestrol Dipropionate (Various Mfr.)
- STILBETIN, N.N.D. 1962.** (Squibb) Diethylstilbestrol, Tab., E.C. & plain, 0.1 mg., 0.25 mg., 0.5 mg., 1 mg., 5 mg., Bot. 100s, 1000s; Plain, 25 mg., 100 mg., Bot. 50s.
Use: Estrogen therapy.
- STILBINOL.** (Merck Sharp & Dohme) Diethylstilbestrol 4 mg./fl. oz. Elixir, Bot. 1 pt., 1 gal.
Use: Estrogen therapy.
- STILBOESTROL, N.N.E. 1959.** See: Diethylstilbestrol (Various Mfr.)
- STILBOESTROL DP.** See: Diethylstilbestrol Dipropionate (Various Mfr.).
- STILBOPHEN, N.N.D. 1959.** Stibophen, U.S.P. See: Fudin. (Winthrop Labs)
- STILESTRATE.** (Upjohn) Diethylstilbestrol 0.5 mg., phenobarbital 32.4 mg./Cap. Bot. 100s, 1000s.
Use: (I.I.D.) Estrogenic & sedative prep
- STILPHOSTIBOL.** (Ames) Diethylstilbestrol diposphate. Amp. (250 mg./5 cc.) 5 cc. Box 20s. Eob. 50 mg., Bot. 50s
Use: I.V. & oral; estrogen therapy
- STIL-ROL DISPELLETS.** (Ballard) Diethylstilbestrol 100 mg./Dispellet. Bot. 60s.
Use: Sustained action, estrogen therapy.
- STILRONATE.** See: Diethylstilbestrol Dipropionate (Various Mfr.).
- STILES.** (Key Corp.) Diethylstilbestrol 25 mg./Tab. Bot. 100s
- STIM INJECTION.** (Foy) d-Pantothene alcohol 500 mg./2 ml. Ampules 2 ml., In 6s, 12s or 100s.
Use: Peristaltic stimulant.
- STIMAVITE TASTITABS.** (Roerig) Vit B-1 10 mg., B-6 3 mg., B-12 20 mcg., sod. ascorbate 25 mg., L-lysine 15 mg./Tastitab. Sol 30s, 100s
Use: Dietary supplement
- STIMDEX.** (Ulmer) Methamphetamine HCl 5 mg./tab. Bot. 100s, 1000s.
Use: (1/2 to 2 daily) Central nervous system stimulant.
- STIMENSIL.** (Wollau) Ovarian gland ext. 0.30 Gm./cc. Bot. 8 cc., 15 cc.
Use: Hormone therapy.
- STIMOGEN.** (Lemmon) Methyltestosterone 1.25 mg., ethinyl estradiol 5 mcg., d-amphetamine Sulf. 1.25 mg., ascorbic acid 50 mg., ascorbic acid 25 mg./Tab. Bot. 100s.
Use: Anabolic stimulation.
- STIMULA.** (Smith Miller & Patch) Camphor, capsicum oleoresin, menthol, methyl salicylate & turpentine oil Tube 1 1/2 oz., 1 lb
Use: Analgesic.
- STOLIC.** (Merck Sharp & Dohme) Mannitol hexonitrate 15 mg., rutin 20 mg., delvalin 30 mg./Tab. Bot. 100s, 1000s.
Use: (1-2 q. 4-6 h) Vasodilator.
- STOLIC FORTE TABLETS.** (Merck Sharp & Dohme) Mannitol hexonitrate 30 mg., rutin 20 mg., delvalin (vinbarbital) 30 mg. Bot. 100s, 1000s.
Use: Vasodilator.
- STOLIMIN.** (Organon) Liver desic. 345 mg., Vit. B-12 1 mcg., natural ptyloric substance 115 mg., ferrous gluconate 170 mg./Tab. Bot. 100s, 1000s.
Use: Nutritional anemias & iron deficiency

STOMACH MUCOSA. See: Kormocaral Tab (Karmel)

STOMACH MUCOSA, HOG EXTRACT W/ Vit B 12 See: Gastran W/Vit B 12 (Winthrop Labs)

STOMACH, POWDERED, N.F., N.N.R. 1952
Dried stomach

STOMACH SUBSTANCE DESICCATED W/ Vit B 12, cobalt gluconate & ferrous gluconate See: Cobedace Cap (Lincoln)

STOMEZE (Richlyn) Mag trisilicate, aluminum hydroxide & chlorophyll Tab, Bot 100s

Use: Antacid

STORAXOL OINTMENT. (Parke Davis) Contains storax, menthol resorcin, camphor sulfur, phenol (1.8%) Tube 1 oz
Use: Skin infections

STORCAVITE (Roerig) Vit A 2,000 u, D 200 u, E 1 IU B 1 1 mg, B 2 1 mg, B 6 0.5 mg, niacinamide 5 mg, calcium pantothenate 2.5 mg, C 33.4 mg, folic acid 0.05 mg, S 12.1 mcg, calcium 67 mg, iron 10 mg, copper 0.3 mg, cobalt 0.03 mg, manganese 0.3 mg, molybdenum 0.03 mg, magnesium 10 mg, zinc 0.2 mg, potassium 1 mg/tab Bot 100s

Use: Pregnancy and lactation

STOVAINE (Wallace) Amylocaine HCl Pow, Bot 5 Gm, 10 Gm
Use: Local anesthetic

STOVARSOL, N.N.R. 1949 See: Acetorsone Tab (Abbott)

Use: Trichomonas vaginalis, vaginitis, amebiasis & Vincent's angina

STOXIL. (Smith Kline & French) Idoxuridine

Use: Antiviral agent for herpetic keratitis

STRAMONIUM, N.F.; Cap, N.F.; Yr, N.F. (Penick) Pow, Sat, 1/4 lb, 1 lb

Use: Sedative & antispasmodic

W/Belladonna, phenobarbital & ephedrine sulfate See: Acapnold Tablet

W/Diethylbarbituric acid & ephedrine HCl See: Phedarine Pill (Burlington)

STRASCOSIC. (Straßenburgh) Acetyl p

aminophenol 300 mg, salicylamide 200 mg, raphatamine phosphate 2 mg, metopine (methylnatrapine nitrate) 0.5 mg/Tab Bot 100s, 1000s

Use: Analgesic.

STRASCO SPECIAL VITAMIN. (Straßenburgh)

Vit A 5000 u, D 500 u, thiamine HCl 2 mg, riboflavin 2 mg, ascorbic acid 30 mg/Tab Bot 100s, 500s, 1000s

Use: Dietary supplement

STRASCO-THIAZIN. (Straßenburgh) Sulfathiazole 100 mg, sulfadiazine 30 mg/

Troche Bot 100s, 500s, 1000s
Use: Orally for sulfonamide therapy

STREP-COMBIOTIC, AQUEOUS (Pfizer) Procaine penicillin G 400 mu streptomycin 0.5 Gm as sulfate/1 dose Sterajets or 5 dose Vials

Use: Anti infective

STREP-COMBIOTICS PS. (Pfizer) Procaine penicillin G 300 mu pot penicillin G 100 mu streptomycin 0.5 Gm as sulf/Vial dose or 5 dose vial

Use: Anti infective

STREP-CRYSDIMYCIN (Squibb) Aqueous suspension containing 600 000 u procaine penicillin G and 1 D Gm streptomycin per 3 cc dose Vials 3 cc, 15 cc

Use: Penicillin Streptomycin Therapy

STREP-DICRYSTIN (Squibb) Procaine penicillin 300 000 u penicillin G 100 000 u streptomycin sulfate 0.5 Gm/Vial 1 dose (2.5 cc) Vials; 2.5 Gm/5 dose Vial

Use: Prophylaxis in surgery in or near infection

STREP-DISTRYLLIN AS (Squibb) Procaine penicillin G 400 mu, streptomycin 0.5 Gm as sulf/vial 1 and 5 dose vials

Use: Anti infective

STREPTIMIDONE. (Parke Davis) Streptomyces rimosus var paramamycinus

Use: Antifungal

STREPTOCOCCI, STAPHYLOCOCCI, PNEUMOCOCCI, & E COLI CULTURES. In cream base

See: Dermo Vee Oint (Sherman)

STREPTOCOCCUS W/Staphylococcus (Lilly) Undenatured antigens of Staph albus & aureus to 0.025 mg nitrogen, streptococcus to 0.025 mg nitrogen/cc Vial 5 cc, 20 cc

STREPTOCOCCUS IMMUNOGEN (Parke Davis) Streptococci hemolytic & non hemolytic 2000 million/cc Vial 10 cc

Use: Treatment of certain types of infections

STREPTOCOCCUS TOXIN, SCARLET FEVER. N.N.R. 1950 See: Scarlet Fever Streptococcus Toxin

STREPTOCOCCUS VACCINE. (Lilly) Arthritis Strain, Vial (20 or 30 million/cc) 20 cc

I.V., Vial (100 or 400 million/cc) 20 cc

STREPTO-COMBINED VACCINE. (National Drug) Streptococcus hemolyticus 1,000 million, streptococcus anhemolyticus 1,000 million, streptococcus viridans 500 million/ml Vial 30 ml

Use: S.C., I.M., as an aid in the treatment of various forms of rheumatism and related diseases

STREPTOCONNIN. (Coniat Midland) Streptomycin Isoniazid sulfate Vial 1.4 Gm in dry powder

Use: Tuberculosis

- * **STREPTODORNASE.** See: Streptokinase-Streptodornase.
- * **STREPTODUOCIN.** Par. Inj., U.S.P., N.N.B. 1955. A mix. of equal parts of streptomycin & dihydrostreptomycin sulfates.
See: Bi-Strept. (Bio-Bamo)
Combistrep, Vial. (Pfizer)
Dihydrostreptomycin Sulfate, Amp. (Lilly)
Districin (Squibb)
Duo-Strep. (Merck Sharp & Dohme)
Mutamycin (Bristol).
- * **STREPTOHYDRAZID.** (Pfizer) Streptomycin-diene Isonicotinyl hydrazine sulfate. * Streptonicoxid. Streptomycin Isoniazid. Vial (equiv. to streptomycin base 1 Gm., isoniazid 236 mg.)
Ure; Tuberculosis.
- * **STREPTOKINASE-STREPTODORNASE.** NND 1962. See Varidare (Lederle)
- * **STREPTOMAGMA.** (Wyeth) Dihydrostreptomycin eulfate 0.3 Gm., koolin 45 gr., pectin 4 gr./fl.oz., Bot. 3 oz., 1 pt.
Use: Injections; bacterial diarrheas.
- * **STREPTOMYCIN.** w/Penicillin G. See: Streptomycin Combiotic (Pfizer) Streptomycin-Districin (Squibb)
- * **STREPTOMYCIN CALCIUM CHLORIDE.** NND. 1962. Streptomycin Calcium Chloride Complex.
Bio-Bamo—Vial 1 Gm., 20 cc; 5 Gm., 50 cc.
Merrell—Vial 1 Gm., 5 cc.; 5 Gm.
Premo Pharm.—Vial 1 Gm.
Merck Sharp & Dohme, N.N.R.—Vial 1 Gm./3 cc., 5 Gm./20 cc.
Winthrop Labs.—Vial 1 Gm., 5 Gm.
- W/Bacitracin & polymyxin B. See: Multibiotic Gauze Pads (Bio-Bamo).
- * **STREPTOMYCIN ISONIAZID.**
See: Streptoconin, Vial. (Consol. Midland)
Streptohydrozid, Vial. (Pfizer)
- * **STREPTOMYCIN SULFATE.** U.S.P., N.N.D. 1962. A.O.R.; Inj., U.S.P. See: Strycin, Syr. (Squibb)
W/Bacitracin & polymyxin-B sulfate. See: Bio-Band Adhesive Stripr. (Multibiotic Corp. of America)
- W/Bicillin, Pot. penicillin G, procaine penicillin G & dihydrostreptomycin sulfates. See: Bicillimycin All-Purpose, Vial (Wyeth)
- W/Dihydrostreptomycin sulfate, N.N.D. 1960
See: Streptoduocin. (Votaur Mfr.)
- W/Dihydrostreptomycin eulfate, chlorprocaine penicillin O, penicillin O sod. See: Cer-O-Strept. Vial. (Upjohn)
- W/Dihydrostreptomycin sulfate & procaine penicillin G. See: Crysidimycin A.S., Vial (Squibb).
- W/Dihydrostreptomycin eulfate & procaine penicillin G & Pot. penicillin G.
- * Generic name recognized by N.F., U.S.P. or A.M.A.
- See: Chlilmycin, Improved (Wyeth).
Dicrysticin Improved, Vial (Squibb).
Pen-Duoastrep, Vial. (Merck Sharp & Dohme)
- W/Gluconolactone. See: Gluco-Mycin, Tab (Commercial Solventr).
- W/Mafenide HCL. See: Sulfamylon, Lq. (Winthrop Labs.)
- W/Neomycin sulfate. See: Intramycin, Pow. (Pitman-Moore)
- W/Penicillin G, Procaine. See: Streptomycin-Crysidimycin A.S. Aqueous surp (Squibb)
- W/Penicillin G, procaine, crytollin sodium penicillin G. See: Strep-Dicrysticin Fortir, Powder (Squibb)
- W/Procaine Penicillin. See: Palfilin. (Phila. Ampoule) Strep-Dicrysticin (Squibb)
- * **STREPTOMYCYLIDINE ISONICOTINYL HYDRAZINE SULFATE.** See: Streptohydrozid, Vial (Pfizer)
- * **STREPTONICOZID.** (Pfizer) Streptomycin-diene Isonicotinyl hydrazine sulfate. See: Streptohydrozid (Pfizer)
- * **STREPTONIGBIN.** (Pfizer) Antibiotic isolated from both filtrates of Streptomyces flocculus.
Use: Antineoplastic agent
- * **STREPTONIVICIN.** See: Novobioctin. (Varioux Mfr.)
- * **STREPTO-STAPHYLO VATOX.** (National Drug) Vaccine from 1000 million each, Str. hemolyticus, onhemolyticus viridans & 2000 million Staph. aureus/cc. Vial 10 cc.
- * **STREPTOVARICIN.** An antibiotic composed of several related components derived from cultures of Streptomyces variabilis See: Dalacin (Upjohn)
- * **STREPTOZOTOCIN.** (Upjohn) Under study.
Use: Antibiotic.
- * **STRESSCAPS.** (Ledaile) Vit. B-1 10 mg., B-2 10 mg., B-6 2 mg., C 300 mg., B-12 4 mcg., niacinamide 100 mg., Cal. pantothenate 20 mg., menadione 2 mg., folic acid 1.5 mg./Cap. Bot. 30s, 100r, 5000s.
Use: Vitamin B complex therapy.
- * **STRESS FORMULA VITAMINS.** (Amer. Pharm. Co) Vit. B-1 10 mg., B-2 10 mg., B-6 2 mg., B-12 4 mcg., C 300 mg., K 2 mg., niacinamide 100 mg., folic acid 1.5 mg., cal. pantothenate 20 mg./tab. Bot. 30s 100s.
Use: Vitamin-stress therapy.
- * **STIBEXATE.** (Marrengitt) Styromate 200 mg, salicylamide 210 mg., acetophenetidin 150 mg., caffeine 30 mg./Tab. Bot. 50r
Use: Muscle relaxant
- * **STRIATRAN.** (Merck Sharp & Dohme) Emylcomate 200 mg./Tab. Bot. 100s.
Use: Tranquillizer

- STRI-EZE.** (Buffington's) Diethylborbituric acid ¼ gr, mephenesin 5 gr, N acetyl p amino phenol 5 gr hamatropine methylbromide 1/48 gr/Tab Bot 100s
Use Treatment of musculoskeletal conditions
- STRON-CAL.** (Vitamix) Col gluconate 0.5 Gm, strontium bromide 0.5 Gm/10 cc
Vial 10 cc
Use IM relief, itching skin
- STRONCYLATE.** (Upjohn) Strontium salicylate 5 gr, aromatics qs/Pill Bot 100s, 500s 1000s
Use Analgesic
- STRONCYLATE (Estra Chem Co)** Strontium bromide 0.5 Gm, calcium gluconate 0.5 Gm, calcium saccharate 45 mg/10 cc
Amps 25s, 100s
- STRONG SILVER PROTEIN, U.S.P., N.N.R.** 1949 See Protargol (Winthrop Labs)
- STRONICAPS.** (Elder) Not listed Strontium salicylate 5 gr, calcichine 1/200 gr/w methyl salicylate/Cap Bot 100s, 1000s
Use Analgesic
- STRONTIUM BROMIDE, N.F.** Cryst or Granule Bot ¼ lb, 1 lb Amp 1 Gm/10 cc
Use Sedative & antiepileptic.
See Ekzebral, Amp (Tesse)
W/Col gluconate & dextrose
See Bromcalate Amp (Endo)
Coldebrom, Amp (G P Horvey)
Glucalbram, Amp (Kirk)
Stron Col, Vial (Vitamix)
W/Passiflora See: Bromo-Flora, Liq (Warren Teed)
- STRONTIUM CHOLATE** W/Strontium salicylate & phenolphthalein diacetate See: Agachalan, Tab (Knoll)
- STRONTIUM LACTATE TRIHYDRATE** See: Strontolac, Cap (Wyeth)
- STRONTIUM SALICYLATE** See: Stroncyolate (Upjohn)
W/Pentylentetrazol methyltestosterone, ethinyl estradiol See Analeptone Anabolic (Reed & Cornrick)
W/Strontium cholate & phenolphthalein diacetate See Agachalan, Tab (Bilhuber-Keoff)
- STRONTOLAC.** (Wyeth) Strontium lactate trihydrate Cap 11 gr, Bot 100s
Use: (3 w/water ac) Management of postmenopausal, senile or idiopathic osteoporosis
W/Passiflora See: Bromo-Flora, Liq (Warren Teed)
- STROPNANTHIN, N.F., N.N.R.** 1947, 1e]. N.F. K strophanthin
Burroughs Wellcome—Tabloid H.T. (1/500 gr) Tube 50s
- W/Digitoxin & strychnine sulfate See: Procardium, Tab (Grant)
- W/Digitoxin, strychnine sulfate nitroglycerin, cactus grandiflorus See: Pasunal, Liq (Tilden)
- STRYCHNINE, N.F. X** (Mollinckradt, Merck Sharp & Dohme) Pow, Bot ¼ oz, 1 oz
W/Alain, belladonna lvs ext & cascara sagrada See A B S & C, Tab. (Horton & Converse)
- W/Cascara sagrada extract, alain, podophyllin belladonna extract, oleoresin ginger See: Alocas No 2 (Sutliff & Case)
- W/Digitalis tr crataegus tr & strophanthus tr See Cardaf, Tab (C D Smith)
- W/Phenolphthalein, alain, belladonna ext ipecac paw See Pheben Pill (Sutliff & Case)
- STRYCHNINE ARSENATE** W/Quinine arsenate, iron arsenate, nucleic acid See: Arsenoin, Tab (Elder)
- STRYCHNINE CACODYLATE** (Various Mfr on request)
W/Sod glycerophosphate, iron cacodylate & procaine HCl See Ampophos, Amp (Buffington)
W/Iron cacodylate Sod glycerophosphate & cacodylic acid See: Ferruginous Compound Amp (Mallard)
- STRYCHNINE GLYCEROPHOSPHATE, (N.Y. Quinine)** Pow, Bot ¼ oz, 1 oz
W/Sod glycerophosphate & Col glycerophosphate See: Glycano, Elixir (Schiefelin)
- STRYCHNINE HYPOPHOSPHITE, (N.Y. Quinine)** Pow, Bot ¼ oz, 1 oz
W/Guaiacol sulfonate, creatate sulfonate hypophosphites of Col, iron, Mang & Pol See: Solguotone, Liq (Pitman Moore)
- STRYCHNINE NITRATE, N.F. X**
- STRYCHNINE PHOSPHATE, N.F.** See: Anort (Chicago)
- STRYCHNINE SULFATE, N.F.** Tab, N.F. (Various Mfr) Cryst, Bot ¼ oz, 1 oz Pow, Bot ¼ oz, 1 oz, 5 lb
W/Acetaphenetidie, acetylsalicylic acid, caffeine & camphor monobromate gelsemium See: Narelic, Tab (Elder)
- W/Belladonna lvs ext, phenolphthalein, alain, podophyllin, ipecac & capsicum oleoresin See: Alipco, Cap (Drug Products)
- W/Digitoxin, nitroglycerin See: Unifloxin, Liq (Tilden)
- W/Phenolphthalein, alain, belladonna lvs ext & ipecac paw See: Alothal, Tab (Mallard)
- W/Iron cacodylate & Sod glycerophosphate. See: Ferrous Compound
W/Strophanthin & digitoxin See: Procardium, Tab (Grant).

W/Straphanthin, digitoxin, nitroglycerin, cactus grandiflorus. See: Posanol, Liq (Tilden)

W/Thyroid, yohimbine. See: Andre-Mudicone, Tab. (Medicone)

W/Vitamin B complex. See: Antecibus, Liq. (Rand)

STRYGIN. (Squibb) Streptomycin sulfate. Syr. 50 mg./cc., Bol. 4 oz.

Use: Streptomycin therapy.

STUART AMINO ACIDS. (Stuart) Total casein acid hydrolysis with 1% L-tryptophane. W/vit. B-12. Pow., Bol. 6 oz. Tab., Bol. 100s, 500s.

Use: Dietary supplement.

STUART BUCLADIN. (Stuart) Buclizine HCl 50 mg., Vit. B-6 10 mg., scopolamine HBr 0.2 mg., atropine sul. 0.05 mg., hyoscyamine sul. 0.05 mg./Tab. Bol. 100s

Use: Treatment of nausea

STUART CMC. (Stuart) Sod. carboxymethylcellulose. Cap. 0.65 Gm. Bol. 100s, 1000s.

Use: Nonirritating bulk laxative.

STUART DIALOSE. (Stuart) Diethyl sod. sulfosuccinate 100 mg., sod. carboxymethylcellulose 400 mg./Cap. Bol. 100s, 1000s

Use: Treatment and prevention of constipation

STUART FORMULA. (Stuart)

Liq. Vit. A 5000 u., D 800 u., B-1 4 mg., B-2 4 mg., P-P 30 mg., B-6 0.2 mg., panthothenal 4.3 mg., Vit. E 0.2 mg., iron, Mang sulfate, iodine/isp. Bol. 1 pt., 1 gal. Tab. Vit. A 10,000 u., D 1000 u., C 100 mg., E 0.3 IU., B-1 5 mg., B-2 5 mg., B-6 0.2 mg., B-12 1 mcg., niacin and niacinamide 30 mg., panthothenal 4.3 mg., w/iron, iodine, calcium, ca. cop., mang., zinc, mag., pot./2 Tab. Bol. 100s, 250s, 500s.

Use: Dietary supplement.

STUART HEMATINIC. (Stuart)

Liq. Ferrous gluconate 3 gr., thiamine chloride 1.7 mg., riboflavin 1.7 mg., niacinamide 10 mg., panthothenal 1.43 mg., pyridoxine HCl 0.5 mg., liver fraction #1 derived from 0.055 Gm. whole liver/5 cc. Bol. 1 pt.

Tab. Ferrous gluconate 3 gr., copper sulf. 2.5 mg., liver desic. 100 mg., B-1 1.7 mg., B-2 1.7 mg., niacinamide 25 mg., cal. panthothenal 1.7 mg., B-6 0.34 mg., B-12 w/noninhibitory intrinsic factor conc. 0.167 oral u., C 25 mg./tab. Bol. 100s, 1000s.

Use: Hematinic.

W/Folic acid & vit. B-12. (Stuart) Ferrous gluconate 3 gr., copper sulf. 2.5 mg., liver desic. 100 mg., vit. B-1 1.7 mg., B-2 1.7 mg., niacinamide 25 mg., cal. panthothenal 1.7 mg., B-6 0.34 mg., folic acid 0.5 mg., B-12

w/intrinsic factor conc. 0.167 oral u., C 25 mg./tab. Bol. 100s.

STUART HEMATINIC FORTIFIED. (Stuart)

Ferrous gluconate 3 gr., copper sulf. 2.5 mg., liver desic. 100 mg., vit. B-1 1.7 mg., B-2 1.7 mg., niacinamide 25 mg., cal. panthothenal 1.7 mg., B-6 0.34 mg., folic acid 0.5 mg., B-12 7.5 mcg., B-12 w/noninhibitory intrinsic factor conc. 0.167 oral u., C 50 mg./tab. Bol. 100s.

Use: Hematinic w/dietary supplement.

STUART LIPOTAINE. (Stuart)

Liq. Betaine 3 Gm., choline 210 mg., liver fraction 210 mg., vit. B-12 12 mcg., B-6 3 mg./15 cc. Bol. 1 pt., 1 gal.

Cap. Betaine 333 mg., choline 35 mg., liver desic. 35 mg., vit. B-12 2 mcg., B-6 0.33 mg./cap. Bol. 100s, 1000s.

Use: Lipotropic therapy.

STUART MULTIVITAMIN DBOPS. (Stuart)

Vit. A 5000 u., D 1000 u., C 60 mg., B-1 1 mg., B-2 1 mg., B-6 0.5 mg., B-12 5 mcg., panthothenal 2.6 mg., niacinamide 10 mg./0.6 cc. Bol. w/dropper 18 cc.

Use: Multivitamin therapy.

STUART MULVIDREN. (Stuart) Vit. A 5000

u., D 1000 u., C 60 mg., B-1 2 mg., B-2 2 mg., B-6 1 mg., B-12 5 mcg., niacinamide 10 mg., cal. panthothenal 3 mg./tab. Bol. 50s, 100s.

Use: Multivitamin therapy.

STUART NEOTINIC. (Stuart) Vit. B-12 w/non-

inhibitory intrinsic factor conc. 1/2 oral unit, ferrous gluconate 2.5 gr., ferrous sulf. 2.5 gr., copper sulf. 5 mg., liver desic. 100 mg., vit. B-1 3.34 mg., B-2 3.34 mg., niacinamide 33.4 mg., cal. panthothenal 6.67 mg., B-6 1 mg., folic acid 0.5 mg., B-12 1.34 mcg., ascorbic acid 100 mg./tab. Bol. 100s, 1000s.

Use: (1 & 2) Anemias.

STUART NOBMACID. (Stuart) Betaine HCl

440 mg., pepsin [1:10,000] 32.4 mg., methylcellulose 110 mg./tab. Bol. 100s, 1000s

Use: (1 a.c. & 1 p.c.) Hydrochloric acid deficiencies & chronic digestive disturbances

STUART OBEXIN. (Stuart) Vit. B-12 25 mcg.,

pyridoxine HCl 5 mg., thiamine mononitrate 10 mg./Tab. Bol. 30s, 100s, 500s.

Use: Nutritional supplement.

STUART PRENATAL. (Stuart) Vit. A 2000 u.,

D 200 u., B-1 1 mg., B-2 1 mg., niacinamide 20 mg., Vit. B-6 3.3 mg., Cal. panthothenal 3.3 mg., Vit. C 66.7 mg., B-12 2 mg., folic acid 0.33 mg., ferrous gluconate 3 gr., K 0.5 mg., w/Cal., Phos., copper, zinc, Mang., Mag., fvarine/tab. Bol. 100s, 1000s.

Use: Vitamin & mineral supplement in pregnancy & lactation.

STUART PRENATAL-F. (Stuart) Vit. A 5000

u D 400 u C 100 mg, B 1 3 mg, B 2 3 mg, B 6 3 mg, B 12 4 mcg niacinamide 20 mg, cal pantothenate 33 mg, 7 minerals bioflavonoids sod fluoride 11 mg / Tab Bot 100s, 500s

STUART PROBEC. (Stuart) Vit C 150 mg, B 1 10 mg, B 2 10 mg, B 6 5 mg, B 12 5 mcg, niacinamide 150 mg, Cal pantothenate 10 mg, plus liver fraction #2 / Tab Bot 50s, 100s, 500s

STUART PROLAR. (Stuart) Salicylamide 2.5 gr, acetaminophen 2.5 gr, acetylsalicylic acid 2.5 gr butobarbital 0.5 gr, d amphetamine sulf 5 mg / Tab Bot 100s
Use Analgesic

STUART PROLAR-R (Stuart) Butobarbital 0.5 gr, d amphetamine sulf 5 mg, acetyl salicylic acid 2.5 gr, salicylamide 2.5 gr N acetyl paminophenol 2.5 gr, hyoscyamine sulf 0.1384 mg, atropine sulf 0.0258 mg, hyoscine HBr 0.0086 mg / Tab Bot 100s

Use Analgesic w/belladonna alkaloids

STUART THERAPEUTIC MULTIVITAMIN (Stuart) Vit A 25,000 u, D 1000 u, C 150 mg, S 1 10 mg, B 2 10 mg, B 12 5 mcg, niacinamide 150 mg / Cap Sol 50s, 100s, 500s

STUART THERON. (Stuart)
Liq: Vit A 25,000 u, D 1000 u, B 1 10 mg, B 2 10 mg, B 6 5 mg, C 300 mg, B 12 5 mcg niacinamide 100 mg, panthenol 17.2 mg / 5 cc Bot 4 oz

Use: Therapeutic multivitamin

Tab: Vit B 1 10 mg, B 2 10 mg, niacinamide 100 mg, Cal pantothenate 20 mg, Vit B 6 2 mg, ascorbic acid 300 mg, Vit B 12 4 mcg, K 2 mg, A 25,000 u, D 1000 u, w/Cal, Phos, Iron, copper, iodine, Mang, Mag, Pot, zinc, cobalt / Tab Bot 100s, 1000s

Use: 1-2 daily; multivitamin with mineral therapy

STUARTINIC (Stuart) Iron 100 mg, copper 2.5 mg, desiccated liver 100 mg, Vit B 1 2 mg, Vit B 2 2 mg, Vit B 6 0.5 mg, Vit B 12 2 mcg, niacinamide 10 mg, calcium D pantothenate 2 mg, Vit C 50 mg, Vit B 12 absorption enhancing complex / Tab Bot 100s, 500s

Use Hematinic

STUART V-C-M. (Stuart) Vit A 5000 u, D 400 u, C 50 mg, B-1 3 mg, B 2 3 mg, B 6 0.5 mg, E D 85 mg, niacinamide 15 mg, panthenol 4.31 mg, folic acid D 5 mg, Vit B 12 1 mcg, Cal, Phos, Iron, copper, iodine, cobalt, zinc, Mang, Mag, Pot / cap Bot 100s, 1000s

Use Dietary supplement

* Generic name recognized by NF USP or A.M.A.

STYPTICIN. (Merck Sharp & Dohme) Calcium chloride Bot 1/8 oz, 1/4 oz, 1 oz
Use Hemostatic

STYPTIRENAL. See Epinephrine (Various Mfr)

STYPTOL. (Knoll) Cotarnine biphthalate Pow 1/8 oz, Bot 100s, Tab 3/4 gr, Bot 100s
Use Hemostatic

***STYRAMATE.** NND 1962 2 Phenyl 2 hydroxy ethyl carbamate
See Myospaz (Chicago Pharm)
Sinaxar (Armour)

W/Salicylamide acetaphenetidin, caffeine
See Strexate (Massengill)

STYRONATE RESINS. Mix of ammonium polystyrenesulfonate 75%, Pot polystyrene sulfonate 25%. See Katonium (Winthrop Labs)

SU-A-SA. (Dome) New name for Sal A Sul Cream (Dome)

SUAVITIL. NND 1962 (Merck Sharp & Dohme) Benzydizine hydrochloride 1 mg / Tab Bot 100s, 1000s
Use Anxiety, tension and depression

SURENON (Seydel) Cal salt of benzoic acid & benzyl ester of succinic acid Tab 5 gr Sol 100s, 500s, 1000s
Use (1-2 q d) Analgesic

SUBY'S SOLUTION G. Citric acid 3.24 Gm, sod carb 0.43 Gm, mag oxide 0.35 Gm / 1000 cc See: Urologic Soln G (Abbott)

SUCARYL CALCIUM. NND 1961, ADR (Abbott) Cyclamate Cal Sod 6%, saccharin cal 0.6% Bot 4 oz 1 pt Tab (Cyclamate cal 50 mg & saccharin cal 5 mg / Tab) Bot 100s, 1000s

Use Noncaloric sweetening agent

SUCARYL CALCIUM. NND 1962 (Abbott) Cal cyclamate 10% & saccharin cal 1% Paw Pkg w/sifter tap 2 oz
Use: Sweetening agent

SUCARYL SODIUM. NND 1961, ADR (Abbott) Sodium Cyclohexylsulfamate Cycloamate Sod Sol 6% & saccharin 0.6%, Bot 4 oz 1 pt, Tab (Cyclamate sod 50 mg & saccharin 5 mg) 2 gr, Bot 100s, 1000s

Use Noncaloric sweetening agent

SUCCINATE, DISODIUM HEXAHYDRATE See: Soduvin Amp (Brewer)

SUCCINATES.

See Calcium succinate

Sodium succinate

Succinic acid

SUCCINIC ACID W/p Aminosalicylate Sod

See: Tubercate Tab (Heun)

W/Aspirin, vit C, calcium succinate See

Arthatabs, Tabs (Columbus)

W/Salicylamide, vit C See: Decasal, Elial

& Tab (Donley Evans)

- W/Sod. salicylate, cal. succinate, vit. C. See: Arterabs No. 2 (Columbus)
- ***SUCCINYLSALICYLIC ACID, U.S.P.**, NND 1967, A.D.R.; Inj., U.S.P.
See: Anectine Chloride, Amp., N.N.R. (Burroughs Wellcome).
Ovelcin Chloride, Amp., N.N.R. (Abbott).
Succostrin Chloride, Amp., N.N.R. (Squibb).
Saxinyl, Vial. (Feigera)
- ***SUCCINYLSULFATHIAZOLE**, NND 1962, A.D.R. 1951; Tab., N.P. See: Sulfasezidine (Merck Sharp & Dohme)
W/Phthalylsulfathiazole. See: Intestal (First Texas)
- SUCCUS CINERARIA MARITIMA**. (Walker Pharm) Bot., 2 dram.
Use: Early stages of cataract.
- SUCOSTRIN CHLORIDE**, NND 1962. (Squibb) Succinylcholine chloride Vial (20 mg/cc) 10 cc; (50 mg/cc) 10 cc.
Use: Muscle relaxant.
- SUCRAPHEN**. (Smith-Dorsey) Phenylphrine HCl. Sol. ¼%, 1%. Bot. 1 oz., 1 pt.
Use: Vasoconstrictor.
- SUCRASE**. See: Invertase (Various Mfrs).
- SUCRITS**. (Merck Sharp & Dohme) Neostigmine 1:1000. (oz., Box 24).
Use: Minor throat irritations.
- ***SUCROSE**, U.S.P. (Saccharose, Sugar) Don Basler—Sol., 50%, Vial 30 cc. Box 1, 6, 25, 100.
Lilly—Sol., 50%, Amp. 50 cc. Box 1, 6, 25.
Use: I.V. diuretic & dehydrating agent.
- SUDAFED**. (Burroughs Wellcome) Pseudoephedrine HCl ½ or 1 gr./Tab. Bot. 100s. Syr. (30 mg/5 cc) Bot. 1 pt.
Use: Decongestant of respiratory tract mucosa.
- W/Chlorcyclizine HCl. See: Fedrazil, Tab. (Burroughs Wellcome)
- W/Cocaine phosphate, Perazil, acetophenanthidin, aspirin, caffeine. See: Emprazil C, Tabs. (Burroughs Wellcome)
- SUGRACILLIN-125**. (Upjohn) Pot. penicillin G 25,000 u./cc. when dissolved in enough water to make 60 cc. sol. Bot. 60 cc. size.
Use: Orally; penicillin therapy.
- SUGRACILLIN-250**. (Upjohn) Pot. penicillin G 50,000 u./cc. when dissolved in enough water to make 60 cc. of sol. Bot. 60 cc. size (3 million u.).
Use: Orally; penicillin therapy.
- SULTRATE**. (Patch) Not listed Sod. phenobarbital 15 mg., methscopolamine bromide 25 mg., alcohol 15%/cc. Bot. 4 oz., pt.
Use: G.I. disorders.
- *Generic name recognized by N.F., U.S.P. or A.M.A.
- SULADYNE**. (Stuart) Sulfamethylthiadiazole 125 mg., sulfadiazine 125 mg., phenylazodimnapyridine HCl 75 mg./tab. Bot. 100s.
Use: Urinary tract infections.
- SULAMYD**, NND 1962. (Schering) Sulfacetamide, Pow., Bot. 5 Gm. Tab. 0.5 Gm., Bot. 100s, 1000s.
Use: Ocular sulfonamide therapy.
- SULAMYD SODIUM**, NND 1962. (Schering) Sulfacetamide Sod. Ophth. Sol. 30%, Bot. sterile w/eye dropper 5 cc., 15 cc.; Nasal, Bot. w/dropper 15 cc. Ophth. sol. 10%, Tube ½ oz.; Box 12x. Ophth. sol. 10%, w/methylcellulose sterile, Bot. w/dropper 15 cc.
Use: Topically; sulfonamide therapy.
- W/Cortagen (Cortisone acetate). See: Cortomyd Ophthalmic Suspension (Schering).
- SULAMYD SODIUM**, NND 1962 Ophthalmic Solution. (Schering) Sod. sulfacetamide 10%, w/methylcellulose 0.5%, Bot. 15 cc.
Use: Antibacterial & anti-inflammatory.
- SUL-DETOX**. (Columbus) Sulfadiazine 1.5 Gm., sulfamerazine 1.5 Gm., ascorbic acid 120 mg., cystine 260 mg., choline chloride 520 mg., alcohol 3%/fl. oz. Bot. 1½ oz., 1 pt., 1 gal.
Use: Sulfonamide susceptible organisms.
- SULESTREX-MT NO. 2**. (Abbott) Sulestrex piperazine 1.5 mg., methyltestosterone 5 mg./S.T. Bot. 25, 100s, 1000s.
- SULESTREX PIPERAZINE**, NND 1961. (Abbott) Piperazine estrone sulfate. Elizin (1.25 mg./3 cc) Bot. 4 oz., 1 pt. Tab. (0.75 mg) Bot. 100s, 1000s; (1.5 mg) Bot. 25s, 100s, 1000s; (2 mg) Bot. 25s, 100s, 1000s.
Use: (1.5 mg. daily) Menopausal symptoms.
- SULIASENZAMIDE**. Sulfanilylbenzamide, N'-benzoylsulfanilamide.
See: Isotogen (Brenntag)
Sultrin, Vag. Tab. (Ortho)
Triple Sulta Cr. (Ortho)
- SULFABID**. (Physicians Prod Co) Sulfaphenazole 0.5 Gm./Tab. Bot. 50s, 250s, 1000s Susp. 500 mg/5 cc Bot 4 oz., 1 pt
Use: (2 tabs. daily) Ur genital tract infections.
- SULFABTIC**. (Wyeth) Sulfadiazine 0.167 Gm., sulfamerazine 0.167 Gm., sulfamethazine 0.167 Gm., Pot. penicillin G 150,000 u./Tab. Bot. 24s, 100s, 250s.
Use: Orally; triple sulfonamide & penicillin therapy.
- SULFA-BON**. (Banar) Sulfacetamide 0.166 Gm., sulfadiazine 0.166 Gm., sodium citrate 0.5 Gm./5 cc. Syr Bot 1 pt.
Use: Treatment of infections.

- SULFACINE CREAME.** (Brewer) Sulfamerazine 2½%, sulfadiazine 2½%, benzocaine 4% Jar 1 oz, 1 lb
Use: Topically; sulfonamide therapy
- SULFACINE TABLETS** (Brewer) Sulfamerazine 19 gr, sulfadiazine 19 gr, orthoform ½ gr / Tab Bat 100s, 500s, 1000s
Use: Sulfa therapy w/local anesthetic, dissolve on tongue
- SULFA-CEEPRYN (CREAM).** (Merrell) Sulfathiazole 10%, sulfanilamide 10%, Ceepryn (brand of cetylpyridinium chloride) 1.500 Tube 1 oz Jar 1 lb
Use: Topically; bacteriostatic
- SULFACEL-15, BUFOPTO.** (Professional Pharm Co) Sod sulfacetamide 5% and methylcellulose ¼%
Use: Common eye pothogens
- SULFACET,** See: Sulfacetamide
- * **SULPACETAMIDE, N.F., A.D.R.** Tab NND 1962 (A.P.C.) N acetylsulfanilamide Tab. 77 gr Bat 100s 1000s
See Bufopto Sulfacel 15 (Professional Pharm)
Isopto Cetamide Ophth (Alcon)
Sulamyd, Pow & Tab, NNR (Schering)
Urosulfon, Tab (Consol Midland)
- W/2,6-Diamino-3-phenylazopyridine HCl See: Azonamide, Tab (G.F. Harvey)
- W/Hexoestrol sulfometh diemer See Yagi Flex, Cream (Rowell)
- W/Mandelic acid & methenamine See: Fene Sul, Tab (Kendall)
- W/Meth dio mer sulfonamides See Sulfa Plex Vaginal Cream (Rowell)
- W/Methenamine mandelate See Proktor, Tabs (Lloyd Dobney & Westerfield)
- W/Methenamine mandelate homatropine methylbromide See Septomide, Tabs (Lemmon)
- W/Methenamine mandelate phenylazodiaminopyridine HCl, acid sodium phosphate See: Uraman w/Azo Tab (Superior Pharm)
- W/Methenamine mandelate, acid sodium phosphate See Uraman Tab (Superior Pharm)
- W/Phenylazo-diamino pyridine HCl See: Azonamide, Tab (G.F. Harvey)
Sulfidum Tab (Columbus Pharm)
- W/Phenylazo diamino pyridine HCl, hyoscyamine sulf, atropine sulf, hyoscyne HBr See: Pylosetamide, Tab (Vitaminix)
Sulfid B-A Tab (Columbus)
- W/Phenylazo diamino pyridine HCl, atropine methylnitrate See Cyslizem, Tab (Zemmer)
Tridium, Tab (Amco)
- W/Phenylmercuric acetate, 9-aminoacridine
- * Generic name recognized by N.F., U.S.P. or A.M.A.
- HCl, benzalkonium chloride See: Kenhez Vaginal Preps (Kendall)
- W/Phenylazo-diamino pyridine HCl See Sulfidum, Tab (Columbus)
- W/Phenylazodiaminopyridine, atropine sulfate, hyoscyne hydrobromide, hyoscyamine sulfate See Uricin, Tab (Richard & Williams)
- W/Prednisolone See Cetapred (Alcon)
- W/Sulfadiazine, sulfamerazine & Sod citrate See: Acet Dio Mer Sulfonamides, NNR
- W/Sulfadiazine, sulfamerazine See: Sulfa Bon (Bonar)
Sulfard (Deseret)
- W/Sulfadiazine, phenylazo diamino pyridine See: Azadelta, Tab (Flessner)
- W/Sulfamethazine, phenazopyridine HCl, citrate citric acid buffered base. See Bufazo, Cap (Sutliff & Case)
- W/Sulfamethylthiodiazole & hyoscyamus ext See: Uronamide, Tab (Flint, Eaton)
- W/Sulfathiazole, N¹-benzoylsulfanilamide, urea See Sulftrin Tab (Ortho)
- W/Sulfur colloid, salicylic acid See Rezo mid, Susp (Dermik)
- * **SULFACETAMIDE SODIUM, USP NND 1962;** Ophth Oint USP; Ophth Sol USP
See Sebizon Lotion (Schering)
Sulamyd Sodium (Schering)
- W/Cortisone acetate See: Cortomyd, Ophth Susp (Schering)
- W/d-Desoxyephedrine HCl See Sulfedexon Liq (Abbott)
- W/Methylcellulose See Bufopto Sulfacel 15 Sol (Professional Pharmacol)
- W/Penicillin Sod See P.S. Dental Cones (Novocoll)
- W/Prednisolone methylcellulose See: Isopto Cetapred Susp (Alcon)
- SULFACETAMIDE, SULFADIAZINE, & SULFAMERAZINE SUSPENSION, N.F.;** Tab; N.F. See Acet Dio Mer Sulfonamides, NNR 1955
- * **SULFACHLOROPYRIDAZINE, N¹** (6-chloro-3-pyridazinyl)sulfonilamide
See Consulid (Ciba)
Sonilyn (Mallinckrodt)
- SULFACIDIN** (Craikes Borneis) Sodium sulfacetamide 10% neomycin sulfate 0.5% / Lacrival 10 cc and Dropperettes
Use: Infections of the eye
- SULFACILLIN** (Success Chemical Co) Pot penicillin G 100M or 250M u, w/sulfadiazine sulfamerazine sulfamethazine 167 mg of each/tab Bat 100s
Use: Penicillin & triple sulfonamide therapy
- SULFACOMP ORALETs.** (C.D. Smith) Sulfa

- thiazole D.25 Gm., benzocaine 6.5 mg., urea 0.0648 Gm., menthol 0.8 mg./Tab. Bot. 500s.
 Use: Antibacterial.
- SULFACTOL** (Winthrop Labs.) Sodium thio-sulfate crystals in amp. of 1 Gm. Box 10s, 100s. Also Tabs. 0.5 Gm. Bot. 100s.
 Use: (I.V. 1 Gm./10 cc.) Antidote for metallic poisoning (Hg, As, Bi & Pb). Tabs. 1 or 2 i.i.d.
- * **SULFADIAZINE**, U.S.P., A.D.R.; Tab., U.S.P., N.N.D. 1961. 2-Sulfonamidopyridine.
 See: *Coco-Diazine Suspension*. (Lilly)
Codiazine Suspension (Massengill).
Cremodiazine Suspension. (Merck Sharp & Dohme)
Cremodiazine-A. (Merck Sharp & Dohme)
Diazine-Citrate, Tab. (Massengill).
Di-Azo-Mul, Susp. (First Texas).
Eskadiazine, Susp. (Smith, Kline & French).
Gluco-Sulfadiazine, Susp. (Donley-Evans).
Lipo-Diazine, Susp. (Donley-Evans)
Lipo-Levazine. (Hiss)
Liquadiazine, Susp. (EliLecol Pharm.).
Mogmeld Sulfadiazine (Pitman-Moore).
Microsulfon, Tab. (Consolid. Mid.)
Fecta-Diazine, Susp. (Funk).
Pyrimol, Tab. (Valtex).
Yano-Diazine, liq. (Vilomix)
 Use: Antibacterial.
- Sulfadiazine Combinations* on p. 694.
- SULFADIAZINE AND SULFAMERAZINE**, Tab., N.F. See: *Dio-Mer-Sulfonamides*, N.N.R. 1955.
- * **SULFADIAZINE SODIUM**, U.S.P., N.N.D. 1962, Inj., U.S.P.; Sterile.
 Use: See *Sulfadiazine*.
- W/Sod. bicarb.
 Massengill—Tab. 7 gr., Bot. 100s, 500s, 1000s.
 Pitman-Moore—Tab. 5 gr., Bot. 100s, 500s, 1000s; 2½ gr., Bot. 100s, 500s, 1000s.
 Upjohn—Tab. 3.85 gr., Bot. 500s, 1000s.
 W/Sod. tartrate. See: *Morusol*, Vial. (Endol)
 See: *Rhinazine*, Nasal Sol. (Lederle).
 W/Tyrosine & butylaminobenzoate. See: *Suthagen*, Traches. (Thompson)
- SULFADIAZINE, SULFAMERAZINE & SULFACETAMIDE SUSPENSION**, N.F.; Tab., N.F. See: *Acet-Dio-Mer-Sulfonamides*, N.N.R.
- SULFADIL LOTION**, (Lilly) Histadyl, sulfacaine, titanium dioxide. Bot. plastic 75 cc
- SULFADIMER**, N.N.R. 1955. (Pitman-Moore)
 Dio-Mer-Sulfonamides, N.N.R. Mogmeld susp. w/Sod. lactate 50 mg. each of sulfadiazine & sulfamerazine/cc. Bot. 1 pt., 1 gal.
- * **SULFADIMETHOXINE**, N.N.D. 1962. See: *Madibon*, Tab. and Susp. (Roche)
- * Generic name recognized by N.F., U.S.P. or A.M.A.
- W/Phenadamine tartrate, N acetyl-p-aminophenol, caffeine. See: *Modricidin*, Cap. (Roche)
- SULFADIMETINE**, N¹-(2-6-Di-methyl-4-pyrimidyl) sulfanilamide. Sulfosomidine, N.N.D. 1959 See: *Elkosin*, Prep (Ciba)
- SULFADIMIDINE**. See: *Sulfamethazine*.
- SULFADINE**.
 See: *Selfadimidine*.
Sulfamethazine.
Sulfapyridine (Various Mfr.)
- * **SULFAETHIDOLE**, N.N.D. 1962. Sulfoethylthiadiazole. N¹-(5-ethyl-1,3,4-thiadiazole-2-yl) sulfanilamide. Sulfoethylthiadiazole.
 See: *Sul-Spanolam* (Smith, Kline & French)
Sul-Spantab (Smith, Kline & French)
- SULFAETHYLTHIADIAZOLE**. See: *Sulfoethidole*, N.N.D. 1959.
- SULFATURAZOLE**. See: *Sulfisoxazole*
- * **SULFAGUANIDINE**, N.F., N.N.R. 1949, A.D.R. 1955, Tab., N.F. (N-Guanylsulfonamide); *Abiguanil*; *Ganidons*; *Guanidol*
 Use: G.I. tract infections.
 W/Koofin & pectin.
 See: *Poguan Susp.* (Massengill).
Pomalin, Susp. (Braun).
 W/Sulfamethazine, sulfamerazine & sulfadiazine.
 See: *Quadetz*, Tab. (Elder).
Quadromoid, Susp. (Elder)
- SULF-ALKA-LAC**. See: *Mogmeld Sulf-Alkaloc*, Sosp. (Pitman-Moore).
- SULFAJEN Jr**, No. 2. (Jenkins Labs) Sulfadiazine 42 mg, sulfamerazine 42 mg., sulfomethazine 42 mg./Tab. Bot. 100s, 1000s.
 Use: Triple sulfonamide therapy.
- SULFALOID TABLETS**, N.N.B. 1955. (Lloyd & Dohney) Meth-Dio-Mer-Sulfonamides, N.N.R. 1954. Tabs. 0.25 Gm. (containing 83 mg each of sulfadiazine, sulfamerazine & sulfomethazine); 0.5 Gm. (0.167 Gm. each). Bot. 100s, 1000s.
- SULFALOXINGE**. (Rorer) Sulfathiazole 0.1 Gm., sulfadiazine 0.1 Gm./Loz. Bot. 100s, 500s.
 Use: Topically; susceptible organisms of throat & mouth.
- * **SULFAMERAZINE**, U.S.P., A.D.R.; Tab., N.F., N.N.D. 1962. N¹-(4-methyl-2-pyrimidyl) sulfanilamide. (Sumedine) See: *Cremamerazine*, Sosp. (Merck Sharp & Dohme)
 Use: Infections caused by susceptible bacteria.
- Sulfamerazine Combinations* on p. 695.
- SULFAMERAZINE SODIUM**, N.N.R. 1951; Inj., Tabs., N.F.
 Use: I.V.; 5% solution.
- SULFAMERAZINE & SULFADIAZINE**, N.F. See: *Dio-Mer-Sulfonamides*, N.N.R. 1955.

Sulfadiazine Combinations*

| Sulfadiazine | Others | Manufacturer | Brand |
|------------------|---|-----------------|--|
| 0.15 Gm | Allantoin 7.5 mg, Iopacote 1.5 Gm | National Drug | Sulfanamets Loz |
| 5 gr | Aspirin 3 gr | Squibb | Diazprin, Wafer |
| 6.48 Gm./100 cc. | Glucose Sod lactate | Danley Evans | Glucio Sulfadiazine, Susp |
| 1.67 mg | Ipratone 125 mg, sulfamerazine 1.67 mg, sulfamethiazole 1.67 mg | Lilly | Ilatone Sulfa Tab |
| 1.67 mg, | Penicillin V Pot 125 mg, sulfamerazine 1.67 mg, sulfamethiazole 1.67 mg | Abbott | Compacilin V W/Sulfas, Filmtab |
| 125 mg | Sulfacetamide 125 mg, phenylazo diamine pyridine 50 mg | Pflesner | Azadettes Tab |
| 0.25 Gm | Sulfamerazine 0.25 Gm | Sulfliff & Case | Chera Sulfabs |
| | Sulfamerazine (See Tablets N F) | Various Mfr | Dia Mer Sulfanamides N R R, 1954 |
| 0.12 Gm | Sulfamerazine 0.12 Gm, sod citrate 0.25 Gm | Messeng Il | Aldiazol M Tab |
| 0.75 Gm | Sulfamerazine 0.75 Gm, sod citrate 3 Gm | Messengill | Ald azal M/15 cc |
| 50 mg/cc. | Sulfamerazine 50 mg/cc, Sod lactate sulfamerazine sulfathiazole | Pitman Moore | Sulfadimer N R R 1954 |
| | Sulfamerazine Sulfacetamide | Various Mfr | Acet Dia Mer Sulfanamides, N R R 1954 (See Susp N F, Tab, N P) |
| | Sulfamerazine Sulfamethiazole | Various Mfr | Meth Dia Mer Sulfanamides, N R R, 1954 |
| 0.167 Gm | Sulfamerazine 0.167 Gm, sulfamethiazole 0.56 Gm, sulfacetamide 0.111 Gm | Chicago | Kairatul, Tab & Susp |
| 0.167 Gm | Sulfamerazine 0.167 Gm, sulfamethiazole 0.056 Gm, sulfacetamide 0.111 Gm | Armour | Deltamide, Tab & Susp (Also W/Pot Penicillin G) |
| 50 mg | Sulfamerazine 150 mg, sulfamethiazole 250 mg, sulfa guanine 50 mg | Elder | Quadetts, Tab |
| 50 mg / 5 cc. | Sulfamerazine 150 mg / 5 cc, sulfamethiazole 250 mg / 5 cc Sulfaguandine 50 mg / 5 cc | Elder | Quad Ramoid, Susp. |
| 125 mg | Sulfamerazine, Sulfathiazole | Various Mfr | Dia Mer Thia Sulfanamide |
| 250 mg | Sulfamethiazole 125 mg, oxadiazole 75 mg | Stuart | Suladyne, Tab |
| 4 gr | Sulfathiazole 250 mg | Schering | Combustol TD, Tab |
| 30 mg | Sulfathiazole 4 gr | Kremer Urban | Disulfadine, Tab |
| 100 mg | Sulfathiazole 100 mg | Strassenburgh | Strasco Thiazin, Tr. |
| 0.12 Gm | Sulfathiazole 0.12 Gm, sod citrate 0.25 Gm | Rarer | Sulfalazone |
| 0.75 Gm | Sulfathiazole 0.75 Gm, sod citrate 23 gr, sod lactate 27 1/2 gr | Messengill | Aldiazol S, Tab |
| | | Messengill | Aldiazol S/15 cc |

* See Sulfamerazine Table

Sulfamerazine Combinations*

| Sulfamerazine | Others | Manufacturer | Brand |
|-------------------------|--|------------------------|---|
| 6.48/100 cc. 167 mg. | Glucose, Sod. lactate Penicillin V Pot., sulfadiazine 167 mg., sulfamethazine 167 mg. | Donley-Evans Abbott | Gluco-Sulfamerazine, Susp. Compeccillin-V W/Sulfas, Filmtab |
| 125 mg. | Sulfacetamide 125 mg., phenazopyridine HCl 65 mg., citrate-citric acid buffer 150 mg. | Suliff & Case | Bulfaza, Cap. |
| 0.25 Gm. | Sulfadiazine 0.25 Gm. | Suliff & Case | Chera Sullabs |
| 0.25 Gm./5 cc. | Sulfadiazine 0.25 Gm./5 cc. | Funk | Duo-Sulfanyl, Syr. |
| 1 Gm./30 cc. | Sulfadiazine 1 Gm./30 cc. | National Drug | Sulfonol, Susp. |
| 20 gr./30 cc. | Sulfadiazine 20 gr./30 cc. | Firf Texas | So-Di-Zola, Susp. |
| 0.22 Gm. | Sulfadiazine 0.22 Gm., Sod. bico.b. 5 gr. | Miller | Sul-Dij-Mill, Tab. |
| 22½ gr./oz. | Sulfadiazine 22½ gr./oz., Sod. citrate 45 gr./oz. | Tilden | Til-Zine, Susp. |
| 2½ % | Sulfadiazine 2½ % benzocaine 4 % | Brewer | Sulfacaine, Cream |
| 1.9 gr. | Sulfadiazine 1.9 gr., orthoform ½ gr. | Brewer | Sulfacaine, Tab. |
| 15 gr./oz. | Sulfadiazine 0.5 gr./oz., sulfacetamide 15 gr./oz., Sod. citrate 45 gr./oz. | Tilden | Tilamide, Liq. |
| 0.167 Gm. | Sulfadiazine 0.167 Gm., sulfacetamide 0.111 Gm., sulfamethazine 0.167 Gm. | Armour | Delamide, Tab. & Susp. Also W/Pot. Penicillin G |
| 0.167 Gm. | Sulfadiazine 0.167 Gm., sulfacetamide 0.111 Gm., sulfamethazine 0.167 Gm. | Chicago Pharm. Co. | Kotrasul, Tab. & Susp. |
| 50 mg. | Sulfadiazine 150 mg., sulfamethazine 250 mg., sulfaguanidine 50 mg. | Elder | Quodeth, Tab. |
| 5 mg./cc. | Sulfadiazine 0.167 Gm., sulfacetamide 0.111 Gm., sulfamethazine 50 mg./cc. | Elder | Quad-Ramold, Susp. |

* See Sulfadiazine Table.

SULFAMERAZINE, SULFADIAZINE & SULFAMETHAZINE 696

- SULFAMERAZINE, SULFADIAZINE & SULFAMETHAZINE.** See: Meth Dia Mer Sulfonamides, N N R 1955
- SULFAMERAZINE, SULFADIAZINE & SULFATHIAZOLE, N N R. 1950** See: Dia Mer Thia Sulfonamides
- ***SULFAMETHAZINE, U S F, N N D 1962** A D R (Merck Sharp & Dohme) 4.6 dimethyl 2 sulfanilamidopyrimidin Tab 7½ gr, Bot 100s, 1000s See: Cremamethazine (Merck Sharp & Dohme)
- W/Penicillin V Pot, sulfadiazine, sulfamerazine See Compoellin-V W/Sulfas, Filmtab (Abbott)
- W/Sulfadiazine, sulfamerazine & sulfacetamide See: Deltamidn, Tab & Susp (Armour) Katrasul, Tab & Susp (Chicago Pharm. Co)
- W/Sulfadiazine, sulfamerazine & Sod sulfacetamide See: Duod Sulf, Tab (Zemmer)
- W/Sulfadiazine, sulfamerazine, sulfacetamide & Pot penicillin G See: Deltamide w/Fenclillin, Tab & Liq (Armour)
- W/Sulfadiazine, sulfamerazine & sulfaguanidine See: Duodetts, Tab (Elder) Duod Ramold, Liq (Elder)
- W/Sulfamerazine sulfadiazine, Ilosone See: Ilosone Sulf, Tab (Lilly)
- ***SULFAMETHIZOLE, N F, N N D 1962,** Susp Tabs, N F 2 Sulfanilamido-5 methyl 1,3,4 thiazolain Sulfamethylthiazazole Urolucastl See: Thiasulfil Susp & Tab (Ayerst)
- W/Methenamine mandelate See: Mesulfiln, Tabs (Ayrst)
- W/Oxytetracycline w/glucosamine, phenylazo-diaminopyrimidin HCl See: Uro-biotic, Cap (Pfizer)
- W/Phenylazo diamino pyridone HCl See: Thiasulfil A Forte, Tabs (Ayerst)
- W/Phenylazodiamino-pyridone HCl See: Thiasulfil A, Tab (Ayrst) Uraulfiln, Tab (Nepira Chem)
- W/Sulfacetamidn & hyoscyamus ext See: Ureamidn Tab (Flinr, Eaton)
- W/Sulfadiazine & esodyn See: Suladyne, Tab (Stuart)
- W/Tetracycline end phenylazo-diamino-pyridone HCl See: Azatraz, Cap (Bristol)
- W/Trisulfapyrimidinns, phenylazo diamino pyridone HCl, See: Es-Sulfo, Tabs (Rownt)
- ***SULFAMITHOMIDINE, (Warner Lambert)** N-(6-methoxy 2 methyl 4-pyrimidinyl)sulfanilamidn monohydrate Use Ant bacterial sulfonamide
- ***SULFAMITHOXAZOLE (Hoffmann LaRoche)** Methyl sulfanilamidolixazole 5 methyl 3
- ***Generic name recognized by N.F., U.S.P. or A.M.A.**
- sulfanilomido-Isaxozain N-(5 methyl 3 isaxozaly)sulfanilomide See Gantonal (Roche Lab)
- ***SULFAMETNOXYPYRIDAZINE, U S P, N N D 1962,** Syr, U S P 3 Sulfanilamido 6 methoxy pyridazine See Kynex, Tab (Lederle) Medical, Tab (Parke, Davis)
- ***SULFAMETHOXYPYRIDAZINE ACETYL, N N D 1962** See Medical Acetyl Suspension, Liq (Parke, Davis)
- SULFAMETNOXYPYRIDAZINE HCl** W/Phenylazodiaminopyridine HCl See: Aza Kynex, Tab (Lederle)
- SULFAMETNYLTNIADIAZOLE.** See: Sulfamethazole, N N D 1959
- SULFAMEZANTINENE.** See: Sulfamethazine
- ***SULFAMONOMETNOXINE, (Lederle Labs)** N¹ (6 methoxy 4 pyrimidinyl)sulfanilamide Use Antibacterial sulfonamide
- ***SULFAMOXOLE, N¹ (4,5 dimethyl 2-methoxy)sulfanilomide** See Nuprin (Upjohn Co)
- p-**SULFAMOYL BENZYLAMINE HCl,** See: Sulfbenzamide
- SULFAMYLON HYDROCHLORIDE, (Winthrop Labs)** (Melenide HCl) p aminomethyl phenylsulfanamide 4 hamosulfanilamide HCl Sol: (5%) Bot 1 & 8 oz Viscous: (5%) w/methylcellulose 1%, Bot 30 cc Vial: (5%) w/streptomycin sulfate 20,000 u, 100 cc See Sulfbenzamine
- W/Benzocaine in propylene glycol See: Olamylon, Ear Drops (Winthrop Labs)
- M/Pantocaine, Nno Synephrine See: FNS Supp (Winthrop Labs)
- SULFA-NEOLIN, (Lilly)** Not listed Dibnzyli ethylendiaminn dipenicillin G 300,000 w sulfadiazine 0.167 Gm, sulfamerazine 0.167 Gm, sulfamethoxine D 167 Gm / 5 cc Sol 2 oz of finished product Use: Penicillin w/triple sulfonamide therapy
- ***SULFANILAMIDE, N F, N N R 1949,** A D R 1955; Tab, N F p Aminobenzene sulfonamidn
- Sulfanilamide Combinations on p. 697.
- SULFANILIC ACID OF ZINC.** See: Nizix (Barroughs Wellcome)
- N-**SULFANILYLACETAMIDE.** See: Sulfacetamide, Prep (Various Mfr)
- SULFANILYLBENZANIDE.** See: Sulfbenzamide
- SULPAPAC, (Abbott)** Sulfathiazole 60%, benzyl alcohol 3%, metaphen 1,500 lb paste of acetic & glycerin, Jar 1 oz

Sulfanilamide Combinations*

| Sulfanilamide | Others | Manufacturer | Brand |
|---------------|--|-------------------|------------------------------------|
| 10% | Allantoin 2% | National Drug Co. | Allantamide, Oint. |
| 10% | Allantoin 2%, aminoacridine HCl 1.500 | National Drug Co. | ACR-Allantamide, Oint. |
| 15% | Allantoin 2%, aminoacridine 0.2% | National Drug Co. | AVC Improved, Cr. & Supp. |
| 5% | Allantoin 2%, sulfathiazole 5% | National Drug Co. | Allantamide w/Sulfathiazole |
| 2% | Allantoin 0.4%, sulfathiazole 1%, ephedrine sulfate 0.5% | National Drug Co. | Allantamide, Liq. w/Ephedrine |
| 10% | Cod liver oil | Massengill | Marumide, Oint. |
| 5% | Cod liver oil, phenacaine HCl 1% | Massengill | Marumide & Phenacaine Ophth, Oint. |
| 320 mg. | Lipiodol 40% | Fougere | Visiodol, Susp./cc. |
| 0.25 Gm. | Pot. penicillin G 5000 u. | Abbott | Powdotalar, Tube |
| 0.063 Gm. | Sod. lactate 12.8%, Pot. citrate 4.9% | Denley-Evans | Glucio-Sulfanilamide, Susp. |
| 12.5% | Sulfathiazole 12.5% | Haskell | Duesulfan, Oint. |
| 6.5% | Sulfathiazole 6.5% | Haskell | Duesulfan, Susp. |
| 6.5% | Sulfathiazole 6.5%, ephedrine sulfate 1% | Haskell | Duesulfan, Susp. w/Ephedrine, Liq. |
| 50% | Sulfathiazole 50% | Flint, Eaton | Sulfanamide Comp. Pow. |
| 3.7% | Sulfathiazole 1.25% | Pott | Thiasulfamilx, Cream |
| 2% | Sulfathiazole 2%, urea 10% | Drug Products | Sulfarea, Oint. |
| 8.8% | Urea 8.8%, benzyl alcohol 3%, propylene glycol | Strasburgh | Surbyl, Liq. |
| 5% | Urea 10%, chlorbutanol 5% | Warren-Teed | Tympanide |
| 5% | Urea 10%, chlorbutanol 2% | White | Otomide, Liq. |
| 2% | Urea 5%, saligenin 1% | Prof. Pharm. | Sulfaural, Liq. |

* See Sulfathiazole Table.

- *SULFAPHENAZOLE N (1 phenyl 5 pyrazolyl) sulfanilamide
See Orisul (Ciba)
Sulfabid (Physicians Prod Co)
- SULFA-PLEX VAGINAL CREAM** (Rowell)
Sulfacetamide sulfadiazine sulfamerazine & sulfamethazine 2.5% of each Tube w/ or without applicator
Use Treatment of vaginal infections
- SULFAPRED** (Professional Pharmacal) Fred
nisalane acetate 0.25% sodium sulfacetamide 10% phenylephrine HCl D 12% in a sterile buffered Methulose suspension
Plas Bot 5 cc
Use Allergic and inflammatory diseases of the eye
- SULFAPYRIDINE, USF, ADR; Tab USF**
2 Sulfanilamidopyridine
- SULFAPYRIDINE SODIUM, Sterile NF X**
(Merck) Pow, Bot 1 oz, ¼ lb, 1 lb
- SULFAQUIN** (Nysco) 0.1 Gm each of sulfathiazole, -diazine -acetamide, merazine and methazine/Tab Bot 1000s
- SULFAREA** (Drug Products) Sulfanilamide Sulfathiazole each 2% urea 10% Oint, Tube ¾ oz Ophth oint Tube w/ tip ¾ oz
- SULFARSPHENAMINE BISMUTH, NNR**
1951 See: Bismarsen Amp (Abbott)
- SULFASED** (Chicago) Succinylsulfathiazole 3 Gm koalin 3 Gm pectin 0.45 Gm homotropine methylbromide 3.5 mg/oz Susp Bot pl
Use Diarrheas
- SULFASOMIDINE** See: Eikosin
- ***SULFASOMIZOLE** (Ives Cameron Co) N
(3 methylsulfathiazolyl) sulfanilamide
Use: Ant bacterial sulfonamide
- SULFASTATIN** (Crookes Barnes) Sodium sulfacetamide 10% phenylephrine HCl D 12%/Lactin al 7.5 cc
Use: Prophylactic ocular ant septic
- SULFA-SUGRACILLIN** (Upjohn) Pot penicillin G buffered 1,500,000 u sulfadiazine 1 Gm sulfamerazine 1 Gm, sulfamethazine 1 Gm, w/sugar & flavoring/60 cc Bot 60 cc.
Use Chemabiotic therapy
- SULFA-SUGRACILLIN FORTIFIED** (Upjohn)
Pot penicillin G 250,000 u, sulfamethazine 0.167 Gm sulfadiazine 0.167 Gm, sulfamerazine 0.167 Gm/3 cc Fow mixed os directed Bot 1 oz, 2 oz
Use Penicillin w/triple sulfonamide therapy
- SULFASUXIDINE, NND 1962** (Merck Sharp & Dohme) Succinylsulfathiazole Fow, Bot ½ oz, 1 lb Tab 7½ gr, Bot 100s 500s
Use: Bacteriostatic for GI tract infections
- *Generic name recognized by NF, USF or A.M.A.
- W/Koalin, pectin, homotropine methylbromide See Sulfased Susp (Chicago)
W/Neomycin, pectin & koalin See: Crema mycin, Susp (Merck Sharp & Dohme)
W/Pectin & koalin See: Cremosuxidine Susp (Merck Sharp & Dohme)
- ***SULFASYMAZINE** (Lederle Lab) N¹ (4,6-drethyls triazin 2-yl)sulfanilamide
Use Antibacterial sulfonamide
- SULFA-TER TABLETS** (American Pharm Co)
Trisulfapyrimidines, USP Bot 1000s
- SULFATNALIDINE, NND 1962** (Merck Sharp & Dohme) Fthalylsulfathiazole Tab 7½ gr, Bot 100s, 500s, 1000s See Crema thalidine, Susp (Merck Sharp & Dohme)
W/Hexylresorcinol See Thalexyl, Cap (Merck Sharp & Dohme)
W/Neomycin sulfate See Neothalidine, Liq (Merck Sharp & Dohme)
W/Neomycin (Merck Sharp & Dohme) Phthalylsulfathiazole 300 mg, neomycin sulfate 200 mg /tab Viol 40s
Use: Orally (5 tab q 4 h for 3 days) control of bacteria in the intestinal tract
- ***SULFATHIAZOLE, NF, NND 1961**
ADR; Tab NF 2 Sulfanilamidothiazole See Cerazole Susp (Massengill)
Coco Thiazole Susp (Lilly)
Gluco Sulfathiazole, Susp (Danley Evans)
Magmoid Sulfathiazole Susp (Pitman Moore)
Pecto Thiazole Susp (Paul Maney)
W/Chlorophyllin See: Thiaphyll Cr (Lonnert)
W/Coal tar, crude See: Sul Torbonis, Oint (Torbonis)
W/Meth dia mer sulfonamides & Sod citrate See: Til For, Susp (Tilden)
W/Paredrine HBr See: Paredrine Sulfathiazole Liq (Smith, Kline & French)
W/Tyrosalricin, 2 chloro 4 phenylphenol & methyl p hydroxybenzoate See: Alginea Cream (Pitman Moore)
- Sulfathiazole Combinations on p 699
- SULFATHIAZOLE CARBAMIDE** See: Otomosan Liq (Daha)
- SULFATHIAZOLE GUM** (White) Tab 3¼ gr-Flg 24s
Use: Typically effective in oral infections.
- ***SULFATHIAZOLE SODIUM, NF, IN, NF, J**
Sterile, NF See: Sulthigel, Oint (Brean)
W/Anthipyrine & propyl p amino-benzoate. See: Sulf Auretane Liq (Raymer)
W/di Desoxyephedrine HCl
See: Sulfedex, Liq (Abbott)
Sulf Opto Sol (Abbott)
Thizodrin, Sol (Lilly)
W/Sulfadiazine Sod & di desoxyephedrine. See: Rhinzazine, Nasal Sol (Lederle)

Sulfathiazole Combinations*

| Sulfathiazole | Others | Manufacturer | Brand |
|-------------------------|--------|-----------------------|------------------------------------|
| 5% (Carbamide) 20.5% | | | Allulion Sulfathiazole |
| 2% | | Wyeth | Otasmose, Liq. |
| 0.25 Gm. | | Daho | Myrinal, Ear Drops |
| | | Smith-Dorsey | Sultacomp., Tablets & Tab. |
| | | C. D. Smith | |
| 5% | | Zorer | Carbathiazole, Oint. |
| 5 gr. | | | Tricobela, Vaginal Tab. |
| 2% | | Sly | Setabar, Vaginal Cap. & Tab. |
| 5% | | Tarboeis | Sub-Tarbanis, Oint. |
| 5% | | Uimar | Ultrazole, Oint. |
| 6 gr./oz. | | Paul Money | Pacto-Thiazole, Liq. |
| 3% | | Hart | Otazole, Liq. |
| 6.48 Gm./100 cc. | | Dorley-Evans | Glucio-Sulfathiazole, Liq. |
| 171 mg. | | Ortbo | Sultrin, Tab. |
| 4 gr. | | Kremers-Urban | Disulfadine, Tab. |
| 0.13% | | Massegill | Aldiazal-M, Tab. Also Susp. |
| 12.5% | | Hoskell | Duetifon, Oint. & Susp. |
| 1.25% | | Patch | Thiasulfamiz, Cream |
| 5% | | National Drug Co. | Allantamide w/Sulfathiazole, Oint. |
| 1% | | National Drug Co. | Allantamide Liq. w/Ephedrine |
| 6.25% | | Hoskell | Duosulfoe Susp. w/Ephedrine |
| 10% | | Merrell | Sulfa Caspryn, Cream |
| 2% | | Drug Products | Sulfarea, Oint. |
| 20% | | Smith, Kline & French | Sulfathiazole, Susp. Microform |
| 5% | | Smith, Kline & French | Primasul, Oint. |
| 10% | | Pitman-Moore | Algeniz, Oint. |
| 10% | | Daho | Otasmose, Liq. |
| 10% | | Drug Products | Sulfarea, Oint. |
| 2% | | | |

* See Sulfanilamide Table

SULFA-TNREE IMPROVED. (Buffington) Sulfadiazine, sulfamerazine, sulfamethazine, each 0.15 Gm / tab Bot 100s, 1000s

SULPATHRIN. See: Algenix, Oint (Pitmoore Moore)

SULPATRIAD (Smith, Miller & Patch) Sulfadiazine 37.5%, sulfamethazine 37.5%, sulfamerazine 25% / tab Bot 100s, 1000s

Use Triple sulfonamide therapy
SULPA-TRI-AZINE, NNR 1955 (M. R. Thompson) Meth dia-mer-sulfonamides, NNR 1954 Sulfadiazine 0.166 Gm, sulfamerazine 0.166 Gm, sulfamethazine 0.166 Gm, Sod lactate 1.5 Gm / 5 cc Susp, Bot 16 fl oz, 1 gal

Use: Sulfonamide susceptible organisms

SULPA-TRIO, (Rezell) Sulfadiazine 167 mg, sulfamerazine 167 mg, sulfamethazine 167 mg / Tab Bot 100s

Use: Triple sulfonamide therapy
SULFAURAL, (Professional Pharm) Sulfonilamide 3%, urea 5% and saligenin 1% in propylene glycol Sol dropper 15 cc

Use: Ear infections
SULP-AUBETONE, (Raymer) Antipyrine 5%, sulfathiazole Sod 5%, propyl p-aminobenzoate 1% in glycerin vehicle Sol w/drop per 1/4 fl oz Bot 1 pt

Use: Analgesic earache drops
SULPAVITIN, (Chicago Pharm) Sulfathiazole w/Vit A from marine liver & Vit D Oint, Tube 1 oz, 2 oz; Jar 1 lb, 5 lb

Use: Second & third degree burns & infected wounds
SULPREZAMINE HYDROCHLORIDE, (Fint, Eaton) Mafenide HCl p Methylaminophenyl sulfonamide HCl Pow, Bulk

Use: Bacteriostatic
Otiomyon Ear Drops (Winthrop Lab)
PNS, Supp (Winthrop Lab)

See: Sulfamylon HCl (Winthrop)
W/di Desoxyephedrine & alkyl dimethyl benzylammonium chloride See: Mordin, Liq (Fint, Eaton)

SULPEDEX, (Abbott) Sod sulfathiazole 2.5%, di-desoxyephedrine 0.125%, Sod sulfate 2% Sol w/dropper 1 fl oz, 1 pt, 1 gal

See: Sulf-opta (Abbott)
Use: Bactericidal nasal solution
SULPEDEXAN, (Abbott) Sod sulfacetamide 2.5%, Sod thiosulf, di-desoxyephedrine HCl 0.25% Sol 1 fl oz, w/dropper, 1 pt

Use: Bactericidal, nose drops
SULFEBROUS, (Chicago Pharm) Ferrous sulfate esic 3 gr / Pill Sol 100s, 500s, 1000s

Use: (1 2 1 f d) Hematinic
SULPHDRYL ION. See: Hydrasulphesol (Lientz)

* Generic name recognized by N.P., U.S.P. or A.M.A.

SULPID. (Calumbus) Phenylazodiamino-pyridine HCl 50 mg, sulfacetamide 250 mg / Tab or 1/2 dr Tab Bot 100s, 500s, 1000s
Susp Bot 4 oz, 1 pt

Use Urinary tract infection
SULPID B-A. (Calumbus) Phenylazodiamino-pyridine HCl 50 mg, sulfacetamide 250 mg, hyoscyamine sulf 0.05185 mg, atropine sulf 0.0097 mg, hyoscine HBr 0.00325 mg / tab Bot 100s, 1000s Double strength Bot 30s, 500s

Use Antispasmodic w/sulfonamide therapy at urinary tract infections
SULPID PEDIATRIC (Calumbus) Phenylazodiamino-pyridine HCl 25 mg, sulfacetamide 125 mg / 5 cc Susp Bot 4 oz

Use Orally, sulfonamide therapy w/urinary antiseptic
SULPIDIUM, (Calumbus) Phenylazodiamino-pyridine HCl 50 mg, sulfacetamide 250 mg / tab Bot 100s, 500s, 1000s

Use: 2 q i d, p c; anti infective
* **SULFINPYRAZONE, NND 1962** 1,2 di-phenyl 4 {2' phenyl sulfanyl-ethyl} 3,5 pyrazolidine dione

See: Anturan, Tab (Geigy)
* **SULPISOMIDINE, NND 1962** Sulfadimidine See: Elikasin, Prep (Ciba)

* **SULFISOXAZOLE, USP, ADR, Tab, USP, NND 1962** 5 p Abinobenzenesulfonyl amide 3,4 dimethylisoxazole See: Gantrisin, Prep NNR (Roche)

* **SULFISOXAZOLE, ACETYL, USP, NND 1962** Oral Susp, USP See: Gantrisin Acetyl (Roche)

* **SULFISOXAZOLE DIETHANOLAMINE, USP, NND 1962;** Inj, Ophth Oint, Oint Soln, USP See: Gantrisin Diethanolamine (Roche)

* **SULTOBBOMOPHTHALEIN SODIUM, USP;** Inj, USP, NNR 1952 (Gotham) Amp (150 mg) 3 cc, 10s, 25s, 100s See: Bromsulphalein, Amp, NNR 1952 (Hynson, Wescott & Dunning)

Use: Liver function test.
SULFOCARBOLATES. Salts of Phenolsulfonic Acid, N.F. Usually Ca, Na, K, Cu, Zn

See: Sodaxilin, Liq (Abbott)
Sulphadine, Tab (Strassenburgh)
Tymcide (Tailley-Nason)

Zinc Salt
SULFOCYANATE. See: Potassium Thiocyanate

SULFOGUAIACOL. See: Pot Guaiacolsulfonate
SULFO-MERTHIOATE POWDER, (Lilly) (Sodium Thimerfonate) 1:1000 Sod p Ethylmercuri thiophenyl sulfonate Bot 1 1/2 oz

Use: Antiseptic

SULFONAMETS W/TOPICAINE. (National Drug) Sulfadiazine 0.15 Gm., Iopicaine 1.5 Gm., allantoin 7.5 mg./Loz. Bot. 25s, 100s.
Use: Local anti-infective.

SULFONAMIDE COMPOUND, N.N.D. 1960. A.D.R. 1955. (Flint-Eaton) Sulfanilamide & sultathiazole 50% each. Tube, insufflator, 1/2 oz.

Use: Topically; antiseptic pow.

SULFONAMIDE, DOUBLES. See: Dia-Mes-Sulfonamides.

SULFONAMIDE DUPLEX, (Lilly) Sulfadiazine & sulfamerazine 250 mg./tab. or 5 cc. Tob. Bot. 24s, 100s, Susp. Bot. 1 pt.

Use: Di-sulfonamide therapy.

SULFONAMIDE PREPS.

See: Acet-Dia-Mer, N.N.R. 1955 (Various Mfr.)

Dia-Mer, N.N.R. 1955 (Various Mfr.)

Dia-Mer-Thia (Various Mfr.)

Meth-Dia-Mer, N.N.R. 1955 (Various Mfr.)

SULFONAMIDES, QUADRUPLE.

See: Delmets, Tab. (Durst)

Deltamide, Tab. & Susp. (Armour)

Kotrasul, Tab. & Susp. (Chicago Pharm.)

Quadett Diplet, Tab. (Elder)

Quadetts, Tab. (Elder)

Quad-Ramold, Susp. (Elder)

Quad-Sulfa, Tab. (Zimmer)

SULFONAMIDES, TRIPLE.

See: Acet-Dia-Mer Sulfonamides.

Acet-Meth-Mer.

Dia-Mer-Thia Sulfonamides.

Meth-Dia-Mer Sulfonamides.

SULFONAMIDES TRIPLEX, (Lilly) Sultathiazole 0.167 Gm., sulfadiazine 0.167 Gm., sulfamerazine 0.167 Gm./Tab. Bot. 100s, 1000s.

Use: Triple sulfonamide therapy.

SULFONATED HYDROGENATED CASTOR OIL. See: Pedinal Hydrisinal, Creme (Pedinal Labs.).

SULFONES, N.N.D. 1960.

See: Clatanone, Tab. & Susp. (Roche Lab)

Diazone (Abbott).

Glucosulfone Sodium (Parke, Davis).

Promacellin (Parke, Davis).

Promin (Parke, Davis).

Promizole (Parke, Davis).

Sulfaxone Sodium (Abbott).

Thiazolsulfone (Parke, Davis)

SULFONIC EXCHANGE RESIN. See: Katanium, Pow. (Winthrop Labs)

SULFONPHTHAL. See: Phenolsulfonphthalein, Prep. (Various Mfr.).

SULFONSOL. (National Drug) Sulfadiazine 0.667 Gm., sulfamerazine 0.667 Gm., sulfa-

methazine 0.667 mg./30 cc. Susp., Bot. 2 fl. oz., 4 fl. oz., 1 pt.

Use: Sulfonamide therapy.

W/Penicillin. (National Drug) Pot. penicillin G 300,000 u, w/sulfadiazine, sulfamerazine & sulfamethazine 0.167 Gm. each/5 cc. when diluted as instructed. Bot. 2 oz.

Use: Penicillin & triple sulfonamide therapy.

SULFONSOL ADULT. (National Drug) Sulfadiazine 1 Gm., sulfamerazine 1 Gm., sulfamethazine 1 Gm./30 cc. Susp., Bot. 4 oz., 1 pt.

SULF-OPTO. (Abbott). Not listed. Sod. sultathiazole 1%, di-desoxyephedrine 0.1% in aq sol. Bot. 1/2 fl. oz., 1 pt. See: Sultedex, (Abbott)

Use: Locally; antibacterial & vasoconstrictor.

SULFURCIN. (Texas Pharm. Co.) Base, sulfur 8%, resorcinol monoacetate 3%. Jar 1/2 oz. Also available in 1/2 strength. Cream, Sulfur 5%, resorcinol monoacetate 3%. Jar 1 lb. Lot., some conc. as cream, Bot. 4 oz., 1 pt.

Use: Topically; antiseptic skin preparations.

SULFORD. (Deseret) Sulfacelamide 2.5 gr., sulfadiazine 2.5 gr., sulfamerazine 2.5 gr./Tab. or 5 cc. of susp. that contain Sod. Citrate.

Use: Treatment of infections.

SULFOREX. (Mottard) 0.083 gr. each of sulfamerazine, sulfadiazine, sulfacelamide. (3.85 gr. total sulfonamide per tab)/Tab. Bot. 100s, 1000s.

Use: Infections responsive to sulfonamides.

SULFOSALICYLIC ACID TABLETS. See: Bumintest, Reagent Tab. (Ames).

SULFOSALICYLIC ACID, W/Aq. Menthol

See: Albumin Test Solution. (Endo)

SULFOSE. (Wyeth) Sulfadiazine 0.167 Gm., sulfamerazine 0.167 Gm., sulfamethazine 0.167 Gm./5 cc. or Tab. Susp., Bot. 1 pt. Tab., Bot. 100s, 1000s.

Use: Triple sulfonamide therapy.

* **SULFOXONE SODIUM, U.S.P.** Tab., U.S.P., N.N.D. 1962. See: Diazone Sed, N.N.D. 1958. (Abbott)

SULFUR. (Various Mfr.) Precipitated, Pkg 1 lb. Sublimed (flowers of sulfur), Pkg. 1 lb. Washed, Pkg. 1 lb. See: Bensulfold, Pow. (Paythress)

Sulfur Combinations on p. 702.

SULFUR, ANTIARTHRITIC.

See: Endogen-A (Endo).

Intra-Sul (Crookes-Barnes)

Polyprate, Amp. (Kirk)

Sacosin (C. D. Smith).

Sul-Gen (Vitamins)

* Generic name recognized by N.F., U.S.P. or A.M.A.

Sulfur Combinations

[702]

| Sulfur | Others | Manufacturer | Brand |
|--------------|--|-----------------------|---------------------------------|
| 1/2 gr | Altopine sulfate 1/200 gr., phenobarbital 1/2 gr. | Paythress | Antracol, Tab |
| 1/2 gr | Belladonna alkaloids 1/600 gr., phenobarbital 1/2 gr | Hart | Taludon, Tab |
| 10 mg /2 cc. | Glucose | Dook | Sulfur Diasporal, Amp. |
| 2% | Pelargonic acid 4%, zinc pelargonate 10% | Crookes Barnes | Pellor, Oint |
| 13.3% | Peru balsam, taya bean oil | Blue Line | Perusal, Oint. |
| 2% | Polyoxyethylene lauryl ether 6%, hydrocortisone 1/4%, hexachlorophene 1% | Westwood | Fastril HC, Oint |
| 2% | Resorcinol 0.5% | White | Appliderm No. 4, Oint, |
| 5% | Resorcinol acetate 2%, panthenol 2% | U S Vitamin | R.S. Panthoderm, Cream |
| 20% | Resorcinol 2 1/2%, beta naphthol 1/2%, peru balsam 7 1/2% | First Texas | A V G, Oint |
| 6% | Resorcinol 2%, zinc oxide 5% | Taibby Mason | Resulch, Cream |
| 3.05% | Salicylic acid 3% | White | Appliderm No. 5, Oint, |
| 2.5% | Salicylic acid 3%, coal tar sol 5%, zinc oxide 5% | Merck Sharp & Dahme | Topotar, Cream |
| 2% | Salicylic acid 2%, Vit A 100,000 u /oz | Dome | Su A Sa |
| 2% | Salicylic acid 2%, camphor 15% | Smith, Muller & Patch | Socasu, Amp |
| 3% | Salicylic acid 3%, coal tar distillate 4% | Smith, Kline & French | Progmatel, Oint, |
| 3% | Salicylic acid 2%, isopropyl alcohol | Dook | Sulfur Salicyl Diasporal, Cream |
| 1.2% | Silican dioxide 15%, aluminum hydroxide 10% | Dook | Aluifa, Cream & Lot. |
| 0.35% | Surfactant | Wallace Lab. | Intraderm Sulfur |
| 5% | Zinc oxide 10%, ellagitigite 10%, resorcinol 2%, iso- propenal 20% | Summer Labs. | Sebasorb, Lot |
| 0.8% | Zinc oxide 22.5%, camphor 1 2%, chaulmoogro oil 0.3%, coal tar 1.5% | Dook | Pasta Durel |
| 5% | Zinc sulfate 3% | Tannett | Acnederm, Uq |

- Sulphocol (National Drug).
Sulfur Diasporal (Doak).
Thiocyl, Amp. (Torigion)
- SULFUR DIASPORAL** (Doak) Sulfur, colloidal in dextrose sol. Amp. (10 mg.) 2 cc.; (30 mg.) 5 cc.
Use: I.V. (10-100 mg. weekly); arthritis.
- SULFUR, SALICYL DIASPORAL** (Doak) Sulfur 3%, salicylic acid 2%, isopropyl alcohol in diasporal base. Cream, Jar 4 oz., 1 lb., 5 lb.
Use: Topically; antiseptic.
- SULFUR, TOPICAL**
See: Alsulfo, Lot. (Doak)
Colla-Sul, Lot. (Crookes-Barnes)
Dalla-Sul, Oint. (Crookes-Barnes)
Polysulfides, Oint. (Upjohn).
Polythionate, Lot. (Upsher-Smith)
Thylox (Shulton, Inc.)
- SUL-GEN** (Vitamis) Coseln 5%, sulfur 10 mg./2 cc. Vial 10 cc.
User I.M., arthritis
- SULKOWITZ REAGENT, IMPROVED** (Winthrop Labs.) Oxalic acid, amm. oxalate, pot. biphthalate, pot. phosphate, pot. bicarbonate, chlorphenol red. Sol. Bot. 2 oz.
User Diagnostic aid, determination of calcium in the urine
- SULMEFRIN**, (Squibb) Not listed. Sulfathiazole Sod. 1.25%, sulfadiazine Sod. 1.25%, with 0.125% di-desoxyephedrine in stabilized, aq. isotonic sol. Bot. w/dropper 1 oz., 1 pt.
Use: Chronic & acute sinusitis, & in nasopharyngeal infections.
- SULMEFRIN OPHTHALMIC**, (Squibb) Sulfathiazole Sod. 0.5%, sulfadiazine Sod. 0.5%, with 0.1% di-desoxyephedrine in stabilized, aq. isotonic sol. Bot. w/dropper 1 oz.
Use: Indicated in keratoconjunctivitis & in chronic & acute conjunctivitis.
- SULMERADINE**, N.N.R. 1955. (Physicians Drug) Di-mer-sulfonamides, N.N.R. Tab. 0.5 Gm., contains 0.25 Gm. each sulfadiazine & sulfamerazine. Bot. 100s, 1000s.
- SULCABBILATE**. 2-hydroxyethyl p-sulfamylcarbonylate.
- SULPHASENZIDE**. See: Sulfabensamide
- SULPHAMETNOXYDIAZINE**. Name Approved by B.P.C. 2-p-Aminobenzene-sulphonamido-5-methoxy-pyrimidine, 5-Methoxy-2-sulphonilamidopyrimidine Durenate.
- SULPHOCOL** (National Drug) Sulfur, colloidal in nonspecific protein colloid. Cap. 5 gr., Bot. 100s, 1000s. Amp. 2 cc., 25 cc.
Use: Rheumatism.
- SULPHODINE** (Strassenburgh) Bismuth subgallate 2 gr., Col. sulfocarbalate 1 gr., Sod. sulfocarbalate 2 gr., copper sulfocarbalate 1/2 gr., coloxidine (iodised Col.) 1/2 gr./Tab. Bot. 500s, 1000s.
Use: Gastro-intestinal astringent.
- SUL-PONDETS**. (Wyeth) Cryst. Pot. penicillin G 20,000 u., bacitracin 50 u., sulfadiazine 2 gr., benzocaine 3 mg., in hard candy base. Jar 36s.
Use: Antibiotic chemotherapeutic agent for oral cavity infections.
- SULOVA IMPROVED BUFFERED SUSPENSION**. (Suliff & Case) Sulfadiazine 125 mg., sulfamerazine 125 mg., sulfacetamide 185 mg., sulfamethazine 62 mg., sodium citrate 454 mg./5 cc. Bot. 2 oz., pts., gals.
Use: Multi-sulfonamide therapy.
- SULOVA IMPROVED TABLETS**. (Suliff & Case) Sulfadiazine 125 mg., sulfamerazine 125 mg., sulfacetamide 185 mg., sulfamethazine 62 mg., sodium citrate 65 mg./Tab. Bot. 100s, 1000s.
Use: multi-sulfonamide therapy.
- SULQUAPAP SUSPENSION**. (Suliff & Case) Sulfadiazine 125 mg., sulfamerazine 125 mg., sulfacetamide 185 mg., sulfamethazine 62 mg., N-ethyl-p-aminophenol 65 mg., sodium citrate 454 mg./5 cc. Bot. 2 oz., pts., gals.
Use: Buffered multi-sulfonamide therapy.
- SUL-SPANSION**, N.N.D. 1962. (Smith, Kline & French) Sulfoethidole 10 gr./5 cc. Susp. & R. oz.
Use: Sustained action broad-spectrum antibacteriol
- SUL-SPANTAB**, N.N.D. 1962. (Smith, Kline & French) Sulfoethylthiadiazole 0.65 Gm./Tab. Bot. 50s. Liq. (650 mg./dr) Bot. 5 oz.
Use: Respiratory and genito-urinary infections.
- SULTACOF**. (Tablet Rack) Sulfadiazine 166 mg., sulfamethazine 166 mg., sulfamerazine 166 mg., pyrimamine maleate 6.25 mg., chlorpheniramine maleate 2 mg., potassium guaicolulfonate 100 mg., phenylephrine HCl 5 mg./5 cc. Susp. Bot. 4 oz., 1 pt. Also tabs. double strength. Bos 50s, 250s.
Use: Upper respiratory disorders.
- SUL-TARBONIS**. (Tarbonis) Sulfathiazole 5%, liquor carbonis detergens 5%, lanolin, menthol. Oint., Jr. 2 1/2 oz., 1 lb.
Use: Topically; antiseptic skin preparation.
- SULTHIAME** (Riker) p-(tetrahydro-2H-1,2-thiazin-2-yl)-benzenesulfonamide, S,S-dioxide See: Tolane.
Use: Anticonvulsant.
- SULTHIGEL** (Breon) Sulfathiazole Sod. 5% in base of propylene glycol 15% & methyl cellulose. Jar, 1 lb., 5 lb.
Use: Sulfonamide treatment.

* Generic name recognized by N.F., U.S.P. or A.M.A.

SULTRI. (Cala Chem Co) Sulfathiazole 0.162 Gm, sulfadiazine 0.162 Gm, sulfamerazine 0.162 Gm /Tab Bot 100s, 500s, 1000s

Use: Triple sulfanamide therapy

SULTRIN (Ortho) Sulfathiazole 171 mg, sulfacetamide 185 mg, N¹ benzoylsulfanilamide 185 mg, urea 13 mg /Tab Bot 20s w/appl

Use: Gynecologic conditions

SULTRIN CREAM. (Ortho) Sulfathiazole 3.42%, sulfacetamide 2.86%, N¹ benzoylsulfanilamide 3.7%, urea 0.64% Tube 78 Gm with or without measured dose applicator

Use: Vaginal infections

SULTRIM (Crystal) Tab Bot 1000s

SULTUSSIN. (Tilden) Sulfadiazine 0.166 Gm, sulfamerazine 0.166 Gm, sulfamethiazole 0.166 Gm, pyrimidine maleate 6.25 mg, phenyltoloxamine dihydrogen citrate 6.25 mg, glyceryl guaiacolate 50 mg, ephedrine sulfate 3 mg/cc Bot 4 oz

Use: (2 dr q 3 h) Treatment of the common cold and related upper respiratory tract infections

SULTUSSIN TABLETS (Tilden) Sulfamethiazole 0.083 Gm, pyrimidine maleate 3.125 mg, phenyltoloxamine dihydrogen citrate 3.125 mg, glyceryl guaiacolate 25 mg, ephedrine sulfate 25 mg /Tab Bot 100s

Use: Anti-infective for upper respiratory tract disorders

SULZADESTRIN. (Brean) Sulfathiazole 5%, fish liver oil in oxycholesterinated base Oint, Jar 1 lb

Use: Topically; sulfonamide therapy

SUMYCIN, NND 1962 (Squibb) Tetracycline Phosp equiv to HCl salt 250 mg /Cap Bot 16s 100s Susp (125 mg/5 cc) Bot 60 cc Pediatric Drops (100 mg/cc) Bot 10 cc Inj, I.M., Vial (100 mg & 250 mg) Syr. (125 mg/5 cc) Bot 60 cc 1 pt Aqueous, Tetracycline phosphate complex equiv to 100 mg tetracycline HCl/cc Pres sulcs/1 cc/12s

Use: Tetracycline phosphate complex

SUNDARE LOTION. (Texas Pharmacal) 2 ethoxyethyl p-methoxycinnamate 1.5%, ethyl alcohol 51.8% Plastic Bot 4 oz

Use: Prevention of sunburn

SUPAC-B. (Mission) N-acetyl p-aminophenol 160 mg, aspirin 230 mg, caffeine 33 mg, aluminum hydroxide 33 mg, calcium gluconate 60 mg /Tab Bot 36s, 100s

Use: Analgesic

SUPER A VITAMIN. (Upjohn) Vit A Cap 25,000, 50,000, 100,000 u, Bot 100s

* Generic name recognized by N.F., U.S.P. or A.M.A.

1000s liq 50,000 u/Gm, Bot 15 cc, 50 cc

Use: Vitamin A deficiencies

SUPERBEV. (Key Corp) Vit B 1 100 mg, B 2 1 mg, B 6 2 mg, niacinamide 50 mg, cal pantothenate 10 mg/cc Vial 10 cc, 30 cc

Use: I.V., I.M., Vitamin B complex

SUPER BEVPLEX FORTIFIED. (Lincoln) High potency B complex Vial 30 cc

Use: Vitamin supplement

SUPER O (Upjohn) Drop, Fish liver oil concentrate Vit A 85,000 u, D 8,500 u/Gm Bot 5 cc, 10 cc, 30 cc w/dropper Perle, Vit A 14,450 u, D 1445 u Bot 30s, 100s 250s

Use: Vitamin A & D therapy

SUPER OASS. (Ferdale) Brewers yeast 210 mg, dioctyl sodium sulfosuccinate 120 mg /Tab Bot 100s

Use: Constipation

* **SUPERINONE.** Tylaxapal

See Alevaire (Winthrop Labs)

Trilon WR 1339

SUPEROXOL. (Merck Sharp & Dohme) Hydrogen peroxide 30%. Bot ¼ & 1 lb

SUPERTAN. (Tailby-Nason) Coal tar fraction 1.25% combined with zinc oxide and starch in proper proportions Tube 1½ oz

Use: Neurodermatitis

SUPERTAN-S (Tailby-Nason) Crude coal tar fraction equal to 5% in a special base Tube 2 oz Jar 2 oz, 1 lb Also avail w/sulfur & salicylic acid

Use: Eczema & allied skin diseases

W/Hydrocortisone See: Neo-Supertan S Cream (Tailby-Nason)

SUPERTAN-10. (Tailby-Nason) Crude coal tar fraction equivalent to 10% tar in a special base Tube 2 oz Jar 2 oz, 1 lb

Use: Eczema & allied skin diseases

SUPERTAN H-C. (Tailby-Nason) Hydrocortisone 1%, with a coal tar fraction 1.25% in a washable base Tube ½ oz 0.590, Tube ½ oz

Use: Dermatoses

SUPERTAN S/S. (Tailby-Nason) Coal tar fraction 1.25% combined with sulfur 4%, and salicylic acid 3%. Tube 1½ oz.

Use: Seborrheic dermatitis

SUPERTROPIC. (Superior Pharm) Choline Bitartrate 35 mg, methaerine 12.5 mg, betaine HCl 25 mg, thiamine HCl 10 mg, riboflavin 2 mg, niacinamide 25 mg, Vit B 12 activity 3 mcg, liver desiccated N.F. 1 gr /Tab Bot 100s, 500s

Use: Liver dysfunction

SUPER ULEX. (Ulmer) Caps.—Vit. B 1 10 mg, B 2 5 mg, B 6 1 mg, cal. pantothenate

25 mg., nicotinamide 50 mg., liver desic. 50 mg., vit. C 200 mg./cap. Bot. 100s, 500s, Inf.—Vit. B-1 100 mg., B-2 3 mg., B-6 50 mg., pantothenic acid 5 mg., nicotinamide 200 mg./cc. Vial 10 cc.

Use: Vitamin B complex therapy.

SUPLEX. (New Mexico Pharm.) Thiamin HCl 150 mg., calcium pantothenate 10 mg., niacinamide 150 mg., inositol 20 mg., riboflavin 2 mg., pyridoxine HCl 10 mg., choline Chloride 20 mg., benzyl alcohol 1.5%/cc. Vial 30 cc.

Use: Super B complex vitamin.

SUPLEX-C. (Bauer) Ascorbic acid 100 mg., choline bitartrate 250 mg., methionine 33 mg., inositol 50 mg., betaine HCl 130 mg., Vit. A 1000 u., riboflavin 1 mg., thiamine mononitrate 1 mg., pyridoxine HCl 1 mg., cobalamin concentrate (B-12) 1 mCg./Tab. Bot. 100s, 1000s

Use: Vitamin supplement.

SUPLIGOL. (Americas Ferment) Ketocholalic acid 1 gr., whale bile, desic. 4 gr./Tab. Bot. 50s, 500s.

Use: (1-2 t.i.d.) Stimulate excretion of bile.

SUPLAMIN FERROUS. (Ayerst) Vit. A 3200 u., carotene 800 u., Vit. D 400 u., C 30 mg., ferrous sulfate 194 mg., Cal. phosphate, dibasic 860 mg./Cap. Bot. 100s, 1000s.

Use: Dietary supplement.

SUPLAVITE. (Ayerst) Cap. (Ferrus sulfate 66 mg., dicalcium phosphate 100 mg., thiamine HCl 1 mg., riboflavin 1 mg., nicotinamide 5 mg., pyridoxine HCl 0.5 mg., Cal. pantothenate 1 mg., Vit. C 25 mg., A 2000 u., D 200 u./cap.) Bot. 100s, 1000s.

SUPPLECAP. (Coral) Calcium 400 mg., Iron 100 mg., purified hesperidin 20 mg., Vit. A 2000 u., Vit. D-2 400 u., Vit. B-1 3 mg., Vit. B-2 2 mg., niacinamide 10 mg., Vit. C 30 mg., calcium pantothenate 2.5 mg., pyridoxine hydrochloride 3.33 mg., folic acid 0.33 mg., Vit. B-12 1 mcg., copper 0.2 mg., manganese 0.2 mg., cobalt 0.04 mg., zinc 0.2 mg., magnesium 0.2 mg., molybdenum 0.04 mg., iodine 0.02 mg./Cap. Bot. 100s, 1000s.

Use: Vitamine-mineral supplement.

SUPPLECAP-G. (Coral) Vit. A 12,500 u., Vit. D 1000 u., Vit. B-1 5 mg., Vit. B-2 2.5 mg., niacinamide 40 mg., Vit. B-6 0.5 mg., calcium pantothenate 4 mg., folic acid 0.5 mg., Vit. B-12 1 mcg., Vit. C 75 mg., Vit. E 2 International u., choline bitartrate 31.4 mg., DL-methionine 10 mg., inositol 15 mg., calcium 75 mg., phosphorus 58 mg., Iron 30 mg., cobalt 0.04 mg., copper 0.45 mg., iodine 0.075 mg., magnesium 3 mg., molyb-

denum 0.1 mg., potassium 2 mg., zinc 0.5 mg./Cap. Bot. 60s.

SUPPLECAP-Y. (Coral) Vit. A 25,000 u., Vit. D 1000 u., Vit. B-1 10 mg., Vit. B-2 5 mg., niacinamide 100 mg., Vit. B-6 1 mg., Vit. B-12 5 mcg., Vit. C 150 mg., calcium 103.6 mg., phosphorus 80.2 mg., Iron 10 mg., cobalt 0.1 mg., copper 1 mg., iodine 0.098 mg., magnesium 5.56 mg., manganese 0.954 mg., molybdenum 0.17 mg., potassium 4.93 mg., zinc 1.4 mg./Cap. Bot. 60s

Use: Vitamins & mineral supplement.

SUPPLIFORY ELIXIR. (White) Vit. B-1 3.3 mg., B-2 3.3 mg., B-6 0.75 mg., niacin 33 mg., pantothenic acid 3.3 mg., B-12 1.67 mcg., lysine HCl 263 mg., methionine 50 mg., choline 100 mg., inositol 100 mg., w/Co, I, Fe, Co, Pot., Mg, Mn, Mo, Zn/15 cc. Bot. 1 fl., 1 gal.

Use: Dietary supplement in the treatment of underweight

SUPRADIN. (Roche) Vit. A 25,000 u., D 1000 u., B-1 10 mg., B-2 5 mg., B-6 3 mg., C 150 mg., niacinamide 50 mg., pantothenic acid salt 10 mg., Vit. E 3 mg./Cap. Bot. 30s, 100s, 1000s

Use: (1-3 daily) Therapeutic vitamin.

SUPRANEPHRIN SOLUTION. (Roche) Epinephrine HCl. Amp. (1:1000) 1 cc., Box 12s, 25s, 100s. Vial (1:1000) 1 fl. oz. (1:100) 6 cc. (For Inhalation).

Uses: S.C., I.M. & Topically; central nervous system stimulant.

SUPRARENAL. N.F. VIII. Is the dried, partially defatted and powdered adrenal gland of cattle, sheep or swine.

SUPRARENIN. N.N.R. 1949, A.D.S. (Winthrop Labs.) Synthetic epinephrine marketed in form of bitartrate salt. Amp. (0.05 Gm.) 1s; (in oil 1:500) 1 cc., 10s. Sol. (1:1000) Bot. 1 oz.

Use: Vasoconstrictor & hemostatic.

SUPREL. (Winthrop Labs) Epinephrine HCl. Sol. 1:100, Bot. 7.5 cc. w/nebulator.

Use: Vasoconstrictor.

• **SURAMIN SODIUM.** U.S.P., N.N.R. 1954. Hexa-Sod bis-(m-aminobenzoyl-m-amino-p-methylbenzoyl-1-naphthylamine-4,6,8-trisulfonate)carbamide. (Bayer 205; Germanin) See: Naphuride Sod., N.N.R. (Winthrop Labs) Deleted.

SURBEX. (Abbott)

Tab. Vit. B-1 6 mg., B-2 6 mg., B-6 1 mg., B-12 2 mg., niacinamide 30 mg., pantothenic acid salt 10 mg., liver 300 mg., yeast 150 mg./filmabot 100s, 500s, 1000s.

Syr. Vit. B-1 6 mg., B-2 6 mg., B-6 1 mg.

* Generic name recognized by N.F., U.S.P. or A.M.A.

- B 12 2 mg, niacinamide 30 mg, yeast 200 mg /5 cc Bot 1 pt, 1 gal.
 Use: (1 dr or 1 Tab daily) Vitamin B complex therapy
- W/ Vit C. (Abbott) Same as Surbex Tab w/Vit C 150 mg/tab Bot 100s, 500s, 1000s Filmtabs Bot 60s
- SURBEX-Y** (Abbott) Vit B 1 15 mg, B 2 10 mg, nicotinamide 100 mg, Vit B 6 5 mg, B 12 4 mcg, calcium pantothenate 20 mg, Vit C 500 mg, desiccated liver 75 mg. liver fraction 275 mg /Filmtab, Bot 100s
 Use: Vitamin deficiencies
- SURBYL**. (Strassenburgh) Sulfanilamide 8 8%, urea 8 8%, benzyl alcohol 3%, propylene glycol liq, Bot 1 pt, 1 gal
 Use: Topically; antibacterial skin therapy
- SURDON**. (Organon) Ethisterone 5 mg, phenobarbital 15 mg, thiamine HCl 15 mg /Tab Bot 30s
 Use: (1 2 daily) Corpus luteum therapy
- SURFACAIN**, NND 1962 (Lilly) (Cyclo methacaine, Topocaine) Cream 0.5%, 1 oz, 12s; 4 oz, 12s; 1 lb, 5 lb Jelly 0.75%, Tube 1/2 oz, 1 oz, 5 oz Oint, 1%. Tube 1 oz 1 lb Supp, 10 gr, Box 12s, 100s Aerosol (0 15 Gm) 2 oz
 W/Sulfadiazine, Box 12s
 W/Zinc oxide, Box 12s
 Use: Local anesthetic
- W/Histadyl See Histadyl & Surfacaine (Lilly)
- W/Histadyl & titanium dioxide See Surfadil, Lal (Lilly)
- SURFACAIN COMPOUND CREAM**, (Lilly) Surfocaine 1/2%, calomine 5%, zinc oxide 5%, methiolate 13000 Tube 1 oz
 Use: Local anesthetic & protective cream
- SURFACAIN & PENICILLIN**, (Lilly) Surfocaine 1%, Pot penicillin G 10,000 u/Gm. Oint, Tube 1 oz.
 Use: Topically; local anesthetic w/penicillin therapy
- SURFACE-ACTIVE AGENTS**, NND 1960 See Antiseptics Surface Active
- SURFADIL** (Lilly) Cream: Histadyl 2%, surfocaine 1/2% Tube 1 oz, Jar 1 lb, 5 lb.
 Lat: Histadyl 2 Gm, surfocaine D 5 Gm, titanium dioxide 5 Gm./100 cc. Plastic bot 75 cc, 1 pt
 Use: Dermatitis
- SURFAK**. (Lloyd Bros) Calcium bis (dioctyl sulfosuccinate) 50 mg or 240 mg /Cap 50 mg Bot 30s, 100s; 240 mg Bot 15s, 100s
 Use: Fecal softener
- SURFRIDER**. (Rawland) Ointment 1 oz
- SURGAIRE**. (Cyanamid) A topical deodorant in aerosol can for use on wound dressings.

- SURGAMYCIN TOPICAL SPRAY OINTMENT**. (Cyanamid) Tetracycline 15 mg, neomycin sulfate 15 mg /Gm 300 Gm /100 Gm aerosol
 Use: Wounds, abrasions, ulcers and in burns
- SURGI-CEN**, NND 1962 (Central) Liq Soap 1% hexachlorophene Bot 375 L (4 larger sizes)
- SURGIDINE**. (Continental Labs) Nonylphenoxypoly (oxyethylene) ethanolic iodine complex (provides 8% ppm available iodine) triethanolamine dodecyl benzene sulfonate and essential oils 24%, water 76%/1 US gallon
 Use: Surgical soap
- SURIN**. (McKesson & Robbins) Methacholine chloride 0.25% w/camphor, menthol & methyl salicylate Oint Jar 1 1/2 oz, 6 oz, 1 lb
 Use: Topically; relief of superficial aches pains & local congestion
- SURITAL SODIUM**, NND 1962 (Parke Davis) Thiamylal Sod Amp: (0.5, 1 & 5 Gm) 1s, 25s, 100s Vials: (1 Gm) 1s, 25s, 100s
 Rectal instillation. Vial (1.5 & 3 Gm) Box 10s A green dye is added to distinguish this from the sterile I.V. form
 Use: I.V.; short acting anesthetic
- SURPLEX FERROUS**. (Abbott) Vit B 1 1 mg, B 2 1 mg, niacinamide 6 mg, ferrous sulfate 200 mg /5 cc Bot 12 oz, 1 gal
 Use: Hematinic (1 tsp 5 times daily)
- SUROFENE**, NND 1962 Hexachlorophene
- SUROMATE**. (Patch) Sulfadiazine 100 mg, sulfamerazine 100 mg, sulfacetamide 100 mg, hyoscyamine ext 575 mg, pot citrate 200 mg /Tab Bot 100s, 500s
 Use: Urinary tract infections
- SURTENOL**. (Guardian Chem) Phenoxynol 1 gal
 Use: Antiseptic
- SUS-PHRINE**. (Brewer) Epinephrine HCl 1:200 Vial 0.5 cc, 12s 2 cc, 5s; 5 cc, 1s
 Use: Relief of frequent asthmatic attacks
- SUSTAGEN**. (Mead Johnson) Vit A 5000 u, D 500 u, ascorbic acid 300 mg, Vit B-1 10 mg, B 2 10 mg, niacinamide 100 mg, Cal pantothenate 40 mg, Vit B 6 5 mg, choline bitartrate 500 mg, folic acid 2.5 mg, Vit B 12 4 mcg, Iron, Cal, Phos Sod, Pot & Protein 210 Gm, fat 30 Gm, carbohydrate 600 Gm, calories 3500/900 Gm Pow, Can 2 1/2 & 5 lb
 Use: Dietary supplement
- SUSTASED**. (Infermedica) Hyoscyamine sulfate D 36 mg, atropine sulfate 0.0585 mg

- hyoscine HBr 0.021 mg., phenobarbital $\frac{3}{4}$ gr./Tab. Bot. 50s, 250s.
 Use: 1-2 on arising; antispasmodic.
- SUSTINEX.** (McNeil) Thiomine HCl 36 mg., riboflavin 12 mg., niacinamide 180 mg., panthenol 6 mg., pyridoxine HCl 6 mg., Vit. B-12 30 mcg./30 cc. Syr., Bot. 8 oz., 1 pt.
 Use: Nutritional disorders.
- SUTHOGEN.** (M. R. Thompson) Tyrothricin 2 mg., sulfadiazine 2 mg., butylaminobenzoate 2 mg./troche. Pkg. 12s, 42s.
 Use: Infections of throat & mouth.
- SUTICIDE.** (Suthiff & Cose) Cetyl dimethyl ethyl ammonium bromide 10%, isopropyl alcohol 19%, inert ingredients 71%, Bot. 45 cc.
 Use: Antiseptic sol.
- SUVREN.** (Ayers) Captoprilomine HCl Bot. 50 mg., 100 mg., 100s, 1000s.
 Use: Antispasmodic.
- SUXAMETHONIUM CHLORIDE, N.N.D. 1962.**
 See: Succinylcholine Chloride.
- SUXINYL CHLORIDE.** (Fougere) Succinylcholine chloride. Vial (20, 50 & 100 mg./cc.) 10 cc.
 Use: I.V. short-acting muscle relaxant.
- SUXSED.** (Roemer) Carbromal 150 mg., scopalamine HBr 0.1 mg., ray-succinate 125 mg./Tab. Bot. 100s, 1000s.
 Use: (1-2 q. 4 h.) Antispasmodic & sedative.
- SWEETA.** (Squibb) Saccharin. Sol. $\frac{1}{4}$ gr./2 drops, Bot. 24 cc., 4 oz.
 Use: Sweetening agent.
- SWEETENING AGENTS, A.D.R. 1955.**
 See: Saccharin (Various Mfr.).
 Sucaryl (Abbott).
 Cyclomate (Abbott).
 Sweeto (Squibb).
- SYLOTAN.** (Mossengill) Pot. sulfocyanate $1\frac{1}{2}$ gr., phenobarbital $\frac{1}{4}$ gr./Tab. Bot. 100s, 1000s.
 Use: Vasodilator & sedative.
- SYCYTROL, N.N.D. 1962.** (Reed & Corrick) Pipethanate HCl, piperidinoethyl benzilate HCl 3 mg./Tab. Bot. 100s.
 Use: Antiphobic.
- SYLAPAR.** (Buffington) Salicylamide 0.25 Gm., p-aminobenzoate Sod. 0.25 Gm., ascorbic acid 10 mg./E.C. Tab. Bot. 100s, 1000s.
 Use: Antirheumatic & antiarthritic.
- SYLAPAR-HC.** (Buffington) Hydrocortisone 1.25 mg., salicylamide 0.25 Gm., sod. p-aminobenzoate 0.25 Gm., ascorbic acid 20 mg./Tab. Bot. 100s, 500s.
 Use: Arthritis, rheumatism.
- SYLLAMALT.** (Borchardt) Diastatic malt extract powder 15%, malt soup extract powder 35%, hemicellulose of the husk of blond psyllium seed 30%. Powder. Jars $\frac{1}{2}$ lb.
 Use: Treatment of constipation.
- SYNASOL, N.N.D. 1962.** (Searle) Sol. of Sod. psyllate, N.F. 5% Amp., serum type, 5 cc., Box 6s, 25s. Vial, serum type, 60 cc.
 Use: Sclerosing agent.
- *SYMETINE HCl.** (Lilly) 4,4'-(ethyleneedioxy)-bis[N-hexyl-N-methylbenzylamine] dihydrochloride.
 Use: Antiepileptic compound.
- SYMPHIST CTR.** (Lowe) Pseudoephedrine HCl 15 mg., pyrilamine maleate 45 mg./Cap. Bot. 100s, 250s.
 Use: Common cold and allergies.
- SYMPATHOLYTIC AGENTS, N.N.R. 1953.**
 See: Adrenergic-Blocking Agents.
- SYMPATHOMIMETIC (PRESSOR) AGENTS.**
 See: Ephedrine-Related Compounds,
 Epinephrine-Related Compounds.
- SYMPATOL.** (Winthrop Labs) Synephrine Tartrate, N.N.E. 1934 (p-Methylaminoethanolphenol Tartrate, Synephrine Tartrate, N.N.R. 1934)
 Sol. (6%) Amp. 1 cc., 6s; (10%) Bot. 1 oz. Tab. (100 mg) Bot. 50s
 Use: Peripheral circulatory stimulant which produces pressor effect.
- SYMPOCAINE.** (Winthrop Labs.)
 Use: Spinal anesthetic.
- SYNA-CLEAR.** (Pruvo) Decangestant plus Vit. C 25 mg./Tab. Bot. 30s.
 Use: Decangestant.
- SYNACOL.** (Reed & Corrick) Scopalamine methylbromide 1 mg., guar gum 420 mg., diethyl sod. sulfosuccinate 20 mg./Cap. Bot. 50s
 Use: Treatment of functional disturbances of lower bowel.
- SYNALAR.** (Syntex Chem.) Fluocinolone acetonide, 6a,9a-difluoro-16a,17a-isopropylidene-dioxy- Δ^1 -hydrocortisone 0.025%. Tube 15 Gm. and 0.01%. Tube 45 Gm.
 Use: Treatment of dermatoses.
- SYNALGESIC.** (Central) Phenobarbital 15 mg., acetylsalicylic acid 0.15 Gm., acetamidol 0.10 Gm., hyoscyamus ext. 10 mg./Tab. Bot. 100c, 1000s
 Use: Analgesic, antipyretic
- SYNALGOS.** (Ives-Cameron) Promethazine HCl 6.25 mg., aspirin 3 gr., phenacetin 2.5 gr., methamphetamine sulfate 7.5 mg./cap. Bot. 48s.
 Use: Treatment of mild to moderate pain.
- SYNALGOS-DC.** (Ives-Cameron) Dihydrocodeine bitartrate 16 mg., promethazine 6.25 mg., aspirin 3 gr., phenacetin 2.5 gr.

- mephentermine sulf. 7.5 mg /Tab. Bot 48s
Use Analgesic relaxant
- SYNANDRETS.** (Pfizer) Methyltestosterone Tab (S1) 10 mg, 25 mg Bot 100s.
Use: Sublingual; testosterone therapy
- SYNANDROL, NND 1960** (Pfizer) Testosterone propionate Vial (25 mg/cc) 1 cc, 5 cc; (50 mg/cc) 1 cc, 5 cc; (100 mg/cc) 1 cc, 5 cc
Use: Androgen therapy
- SYNANDROL P, NND 1960** (Pfizer) Testosterone oil susp Vial (25 mg/cc) 1D cc; (50 mg/cc) 10 cc; (100 mg/cc) 1D cc
Use: Androgen therapy
- SYNANDROTABS.** (Pfizer) Methyltestosterone Tab 10 mg, 25 mg, Bot 100s
Use: Orally; androgen therapy,
- SYNAPOIDIN.** (Parke, Davis) Comb of chorionic gonadotropin & follicle stimulating hormone from anterior pituitary Contains 15 synergy RU/cc Vial 10 cc
Use: IM; chorionic & pituitary gonadotropin therapy.
- SYNASAL.** (Texas Pharm Co) Phenylephrine HCl 1/4%, Spray bot 1 oz
- SYNATAN.** (Irwin, Neisler) Tonphelamin (dextroamphetamine tartrate) 17.5 mg tab Bot 50s
Use: Delayed action stimulant
- SYNATAN FORTE.** (Irwin Neisler) Tonphelamine 26.25 mg/Tab Bot 50s
Use: Obesity
- SYNATE** (Central) Secobarbital 2D mg, synophylate 360 mg, rocephedrine HCl 30 mg potassium iodide 300 mg, niacinamide 40 mg/Tab Bot 100s 500s, 1000s
Use Bronchial asthma
- SYNATE JR** (Central) Secobarbital 1D mg synophylate 180 mg rocephedrine HCl 5 mg, potassium iodide 150 mg niacinamide 20 mg/Tab Bot 100s 500s
Use Bronchial spasm
- SYNATE-M.** (Central) Secobarbital 45 mg Synophylate 330 mg, rocephedrine HCl 25 mg, potassium iodide 300 mg niacinamide 40 mg/Tab Bot 100s, 500s
Use Bronchial asthma
- SYNCAINE.** See: Procaine Hydrochloride (Various Mfr)
- SYNCELOSE, NND 1962** (Elue Line) Methyl cellulose Tab 0.5 Gm, Bot 100s 1000s
Use: Hydrophilic colloid laxative
- SYNCILLIN, NND 1962** (Bristol) Phenethylamine pot Alpha phenoxyethyl penicillin, potassium 125 mg or 250 mg Tab Bot 25s 100s 125 mg/5 ml as powder for oral solution for reconstitution, fruit flavor Pediatric Drops; 125 mg/0.6 cc Bot 72 cc w/calibrated dropper
Use: Penicillin therapy with broader spectrum
- SYNCORT.** See: Desoxycorticosterone Acetate (Various Mfr)
- SYNCORTYL.** See: Desoxycorticosterone Acetate (Various Mfr)
- SYNCROBIN.** (Schenley)
- SYNCURINE, NND 1962** (Burroughs Wellcome) (Decamethanum Bromide) Deco methylene 1,10 bistrimethyl-ammonium di bromide Vial (1 mg/cc) 10 cc
Use: Skeletal muscle relaxant
- SYNDECON** (Bristol) Pot alpha phenoxy ethyl penicillin 62.5 mg, pheylephrine HCl 2.5 mg, Phenylpropanolamine HCl 10 mg, Phenyltoloxamine citrate 3.75 mg, Chlorpheniramine maleate 1.25 mg, Acetyl p omeaophenol 120 D mg/Tab or 5 cc Tab Bot 25s As powder for reconstitution to 60 ml
Use: Relief of the common cold
- SYNERFLEX INJ.** (Ries) Estradiol 1 mg testosterone 10 mg, Vit C 100 mg, B1 25 mg, B2 5 mg, B6 5 mg, niacinamide 25 mg, pantothenol 10 mg/cc Vial (multiple dose) 20 cc
Use Androgen estrogen
- SYNDIOL.** (Central Pharm Co) Estradiol Vial (0.5 mg/cc) 10 cc Vial (0.2 mg/cc) 30 cc
Use IM (0.2 mg m p); estrogen therapy
- SYNDROX HYDROCHLORIDE, NND 1962** (McNeil) Methamphetamine HCl Elixir 30 mg/oz, Bot 1 pt, 1 gal Tab 5 mg, Bot 100s, 1000s
Use: Sympathomimetic.
- SYNEPHRICOL.** (Winthrop Labs) Neo synephrine HCl 5 mg, codeine phosphate 7.7 mg pot guaicol sulfonate 7D mg, ammonium chloride 7D mg, menthol 1 mg, chloroform 0.0166 cc, 8% alcohol/4 cc Bot 1 pt, 1 gal
Use: (12 tsp q 2-4 h) Coughs due to colds
- SYNEPHRICOL YHENFACIL** (Winthrop Labs) Neo synephrine HCl 5 mg, theladil HCl 4 mg, dhydrocodeinone bitartrate 1.33 mg, pot guaicol sulfonate 7D mg, am
Use: Antispasmodic
- W/Aluminum hydroxide, Col carb & Mag** Bot 1 pt, 1 gal
Use: (12 tsp q 2-4 h) Coughs due to colds
- SYNEPHRINE {TARTRATE}, NNR 1934** See: Oxedrine (Bawmoe)
Sympafol, Amp (Winthrop Labs)
Synhenate Tartrate, Amp (Breco)
- SYNERGEN.** (Lemmon) Phenylpropanolamine HCl 12.5 mg, phenylephrine HCl 1.25 mg.

- pheniramine maleate 6.25 mg., pyrilamine maleate 6.25 mg./5 cc. Elixir Bot. 3 oz.
Use: Antihistaminic for colds.
- SYNERGISTIN.** (Consolidated Midland Corp) Procaine penicillin G 300M u., Pot. penicillin G 100M u., dihydrostreptomycin sulfate 0.5 or 1 Gm./dose. Vial 1 dose
Use: I.M., penicillin & streptomycin therapy
- SYNBONE.** (Pitman-Moore) Testosterone propionate in sesame oil, Vial 25 mg./cc. 10 cc.
Use: I.M. (25-50 mg. q. 2-3 days); androgen therapy.
- SYNSETBOL.** N.N.D. 1962. (White) Diene-sol. Tab. 0.1 mg., 0.5 mg., 10 mg., Bot. 100s, 1000s.
Use: Synthetic estrogen therapy.
- SYNETRIN.** (Bristol) Vials. I.M. 150 mg. 350 mg IV. 700 mg.
- SYNGESTERONE.** N.N.D. 1960. (Pfizer) Progesterone in aq. susp. Vial (25 mg./cc.) 10 cc.; (50 mg./cc.) 10 cc.
Use: Corpus luteum therapy
- SYNGESTRETS.** (Pfizer) Progesterone. Tab. (sublingual) 10, 20, 50 mg., Bot. 100s.
Use: Sublingual; androgen therapy
- SYNIODIN.** (Blue line) Iodine 1 gr. combined w/protein/Tab. Bot. 100s
Use: (1 fl.d) Iodine therapy
- SYNIRIN.** (Paythress) Aspirin 5 gr., pentobarbital 1/2 gr./Tab. Bot. 100s.
Use: Analgesic w/sedative.
- SYNIST.** (Luka) Chlorpheniramine maleate 1 mg., pyrilamine maleate 12.5 mg., phenylpropranolamine hydrochloride 25 mg., phenylephrine hydrochloride 2.5 mg./Cap. S.-SA contains in addition: acetylsalicylic acid 300 mg./Cap. Bot. 50s.
Use: Treatment of common cold.
- SYNKAMIN.** (Parke, Davis) 4-Amino-2-methyl-1-naphthol HCl. (Vit. K analogue). Amp. (1 mg.) 1 cc., 6s, 25s, 100s. Capsule 4 mg., Bot. 100s, 1000s.
Use: Prophylaxis & correction of hypoprotrombinemia.
- SYNKAYVITE.** N.N.D. 1961. (Roche) [Menadiol Sod. Diphosphate] 2-Methyl-1,4-naphthalenediol disphosphoric acid ester, tetrasodium salt. Amp. (1 mg., 2.5 mg.) 0.5 cc. Box 12s (5 mg., 10 mg.) 1 cc., Box 6s, 25s, 100s. Amp. (37.5 mg./2 cc.) Box 6s, 25s, (75 mg.) 2 cc., Box 6s, 25s. Tab. 1 mg., 2.5 mg., 5 mg., Bot. 100s, 500s, 1000s.
Use: Synthetic Vitamin K for hemorrhagic tendency.
- SYNKAYVITE-C.** (Roche) Water-soluble Vit. K comp. w/Vit. C. Synkayvite 5 mg., ascorbic acid 200 mg. Loz., Pkg. 20s, 100s, 500s. Drop. Bot. w/dropper 30 cc.
Use: Prevent hypoprotrombinemia & promote wound healing.
- SYNKAYVITE-C.B.** (Roche) Synkayvite 5 mg., ascorbic acid 200 mg., Vit. B-1 5 mg., B-2 5 mg., niacinamide 25 mg., panthenol 5 mg./Cap. or Loz. Cap., Bot. 20s, 100s, 500s. Loz., Bot. 20s, 100s.
Use: Vitamin K therapy w/vitamin supplement.
- SYNKONIN.** See: Dihydrocortisone (Various Mfr.).
- SYNODAL.** (Winthrop Labs.) Comb. of lipoproteins w/emeline, each 6 cc. containing 1 gr. emeline HCl. Amp. 6 cc., Box 6s, 25s.
Use: I.V. & I.M., for gastric & duodenal ulcer.
- SYNOPHEDAL.** (Central) Phenobarbital 1/4 gr., acephedrine 3/8 gr., theophylline Sod. glycinate 5 gr./Tab. Bot. 100s, 500s, 1000s
Use: (1-2 fl.d.) Bronchodilation in treatment of asthma.
- SYNOPHYLATE.** N.N.D. 1962. (Central) Theophylline Sod. glycinate.
Amp. (0.4 Gm.) 10 cc., Box 6s, 25s. Elix.: (0.33 Gm./15 ml.) Sol. 1 pt.
Supp. (12 gr.) Box 12s.
Tab.: (2 1/2 & 5 gr.) Bot. 100s, 500s, 1000s.
Use: Diuretic, antispasmodic & coronary vasodilation.
W/Phenobarbital. (Central) Theophylline Sod. glycinate 5 gr., phenobarbital 1/4 gr./Tab. Bot. 100s, 500s, 1000s.
Use: Diuretic w/sedative.
W/Secobarbital, acephedrine HCl, Pot. Iodide, niacinamide. See: Synote, Tab. (Central)
- SYNOPHYLATE JR.** (Central) Synophylate 3 gr., pentobarbital sod. 1/2 gr./Pediatric Sepp. Box 12s.
Use: Bronchial asthma.
- SYNOREX.** (Schilling) di-amphetamine 6 mg., d-amphetamine 6 mg., butabarbital 30 mg./Cap.
Use: Obesity control.
- SYNSALYL-C.** (Tilden) Salicylamide 4 gr., Sod. p-aminobenzoate 4 gr., vit. C, Sod. 25 mg./Tab. Bot. 100s, 500s.
Use: Salicylate therapy.
- SYNTA.** (Dumas-Wilson) Synthetic coal tar equivalent to 13% natural product, in a lipophilic base of petrolatum, zinc oxide and starch Synta w/hydrocortisone contains in addition hydrocortisone alcohol 0.5%. Plain Tube 1 1/2 oz., 1 lb., w/MC Tube 1/2 oz.
Use: Dermatologic.
- SYNETRIN.** (Bristol) N-(pyrrolidinomethyl) tetracycline. 150 mg., 350 mg r ascorbic

- acid, lidocaine—1 M Inj 700 mg, plus ascorbic acid—IV Inj Vials
 Use: Tetracycline sensitive infections
- SYNTHALOIDS.** (Buffington) Benzocaine, eucalyptol, Col iodised, hexamethylene amine & menthol/Trache Box 15; Bot 100s, 1000s
 Use: Throat & mouth infections
- SYNTHENATE TARTRATE.** (Breon) Not listed
 Synephrine tartrate Amp 10% 1 cc, 25s
 Use: SC or IM, peripheral circulatory stimulant
- SYNTHOESTRIN.** See: Diethylstilbestrol (Various Mfr.)
- SYNTROID SODIUM, NND 1962** (Travenol Lab) Sod levo thyroxine Tab (0.05 0.1 & 0.2 mg) Bot 100s
 Use: Thyroid deficiencies
- SYNTIL** (McNeil) Syndrax HCl 2.5 mg, butisol Sod 15 mg/Tab Bot 100s, 1000s
 Use: Sedative & central nervous system stimulant
- SYNTOCINON** (Sandoz) A sterile aqueous sol of synthetic oxytocin Amp (10 u/cc) 1 cc, Sax 12s 100s Amp (5 u/0.5 cc) 0.5 cc Box 12s 100s Bot (40 u/cc) 5 cc
 Use: Induction and/or labor and for prevention and control of postpartum hemorrhage
- SYNTOCINON NASAL SPRAY.** (Sandoz) Syn oxytocin 1 cc of aqueous solution of 40 USP u 2 cc, 5 cc squeeze bottle
 Use: Lactation
- SYNTOPHEROL ACETATE** (Abbott) dl alpha tocopherol acetate Tab 25 mg, 50 mg, Bot 25s, 100s, 250s, 1000s
 Use: Vitamin E therapy
- SYNTOPHEROL ACETATE IN OIL.** (Abbott) dl-alpha tocopherol acetate Vial (200 mg/cc) 10 cc.
 Use: I.M., vitamin therapy
- SYNTRATE.** (Central) Phenobarbital 1/4 gr, mannitol hexanitrate 1/2 gr, theophylline Sod glycinate 2 1/2 gr/Tab Bot 100s, 500s, 1000s
 Use: (1 2 tid) Vased later
- SYNTROGEL.** (Roche) Aluminum hydroxide 2 1/4 gr, syntropan 3/40 gr Cel carb 1 1/2 gr, Mag perox 1 1/2 gr/Tab Bot 50s, 100s 250s 1000s
 Use: (1 2 qid) Antacid & antispasmodic.
- SYNTRONAL.** (Roche) Syntropan 50 mg, phenobarbital 15 mg/Tab Bot 100s, 1000s
 Use: 1 2 q 3-4 h; relaxes spastic smooth muscles
- SYNTROPAN, NNB 1950** (Roche) (Amproropine Phosphate) dl tropic acid ester of 3-diethylamino 2,2 dimethyl 1 propanol phosphate Tab 50 mg, 100 mg, Bot 1000s Pow, Vial 5 Gm, 25 Gm
 Use: Antispasmodic
 W/Aluminum hydroxide Cel Carb & Mag peroxide See: Syntrogel, Tab (Roche)
 W/Phenobarbital See: Syntrol, Tab (Roche)
- SYNTUSSIN.** (Ives Cameron) Chlorpheniramine maleate 2 mg, phenylephrine HCl 7.5 mg, d methorphan HBr 10 mg terpin hydrate 64.8 mg Tab Bot 48s
 Use: Nasal congestion, cough
- SYNTUSSIN COMPOUND.** (Ives Cameron) Chlorpheniramine maleate 2 mg, phenylephrine HCl 7.5 mg dextromethorphan HBr 10 mg terpin hydrate 64.8 mg, acetaminophen 194 mg ascorbic acid 25 mg / Cap Bot 48s
 Use: Symptomatic relief in colds and acute upper respiratory infections
- SYRSULFAS SYRUP.** (Upjohn) Sulfadiazine 0.167 Gm, sulfamerazine 0.167 Gm, sulfamethazine 0.167 Gm / 5 cc Bot 1 pt
 Use: Triple sulfonamide therapy
- SYRATINIC** (Dumas Wilson) Folic acid 3 mg Vit B 6 12 mg, B 12 50 mcg, Iron 72 mg / oz Bot 1 pt, 1 gal
 Use: Treatment of anemias
- SYRCODATE** (Blue Line) Codeine phosphate 1 gr, ammonium chloride 8 gr Sod benzoate 4 gr, aromatics flavors in sugar syr Bot 1 pt, 1 gal
 Use: (1 2 tsp q 3-4 h) Sedative expectorant
- *SYROSINGOPINE, NND 1962** 4 ethoxy carbonyl 3,5 dimethoxybenzoic acid ester of methyl reserpate See: Singoserp Tab W/Hydrochlorothiazide See: Singoserp Elixir (Ciba)
- SYRPALTA.** (Emerson) Syr containing comb of fruit flavors Bot 1 pt, 1 gal
 Use: Vehicle for masking drug taste
- SYSTIN** (Sulfik & Case) Systin 325 mg / Tab Bot 1000s
 Use: Urinary antiseptic
- SYSTIN WITH SODIUM ACID PHOSPHATE** (Sulfik & Case) Systin 325 mg, sodium acid phosphate 325 mg / Tab Bot 1000s
 Use: Urinary antiseptic
- SYTOREX, NND 1962** (Parke Davis) Vit B 12 (New name for Cytobex). Vial (15 mcg/cc) 10 cc; (30 mcg/cc) 10 cc; (100 mcg/cc) 10 cc; (1000 mcg/cc) 10 cc. & 1 cc. See 10s
 Uses: Pernicious anemia

TAB. (Tilden) Methyl nicotinate, camphor, menthol, methyl salicylate, Bot. 3 oz., pts., gals.

Use: Analgesic Balm.

TABAMINS. (APC) Vitamins A 5000 u., D 1000 u., B-1 2 mg., B-2 2 mg., C 50 mg., B-6 1 mg., B-12 5 mcg., niacinamide 20 mg., folic acid 0.25 mg., calcium pantothenate 5 mg./Tab. Bot. 60s.

Use: Vitamin.

TABFERR. (Jackson-Mitchell) Ferrous sulfate 4 gr., citric acid 2 gr./Tab. Bot. 100s.

Use: (1-2 t.i.d.) Iron therapy.

TABLOGESTIN. (Strong) Bile salts 2 gr., Sod. salicylate 2½ gr., pancreatin 5 gr., Sod. bicarb. 5 gr./3 Tab. Bot. 72. See: Chologestin (3 Tab./15 cc.) (Strong).

Use: (1 to 3 t.i.d.) Choleric & chologogue.

TABOREA. (Tablet Rock) Bromisovalerylurea 2 gr., bromadethylocetylurea 3 gr./Tab. Bot. 100s, 500s.

Use: Daytime sedative.

TAC-33. (Norwich) Aspirin, phenacetin & caffeine/Tabs. Bot. 100s.

TACARYL. N.N.D. 1961. (Mead Johnson) Methdilazine HCl 10-(1-methyl-3-pyrroldidyl methyl) phenothiazine hydrochloride Tab. 8 mg. Bot. 100s Syrup 4 mg per 5 cc. Bot. 16 oz.

Use: Antiallergic Therapy

TACARYL CHEWABLE TAB. (Mead Johnson) Methdilazine, 3.6 mg /Tab. Bot. 100s.

Use: Antiallergic-antipruritic action

TACE. N.N.D. 1962. (Merrell) Chlorotriani- tene, Tri-p-anisylchloroethylene Cap. (12 mg. in corn oil) Bot. 70s, 350s, (25 mg) Bot. 12s, 60s.

Use: Estrogen therapy.

TACE W/ANDROGEN. (Merrell) Tace 6 mg., methyltestosterone 2.5 mg /Cap. Bot. 100s

Use: Balanced estrogen-androgen therapy

TACE W/ERGONOVIN. (Merrell) Chlorotriani- tene 25 mg., ergonovine maleate 0.1 mg /Cap. Bot. 12s, 60s.

Use: Prevention of postpartum hemorrhage

TACOL. (Mossengill) Dexamethorphan hydrobromide 15 mg., chlorpheniramine maleate 2 mg., pyrilamine maleate 12 mg., phenylephrine hydrochloride 10 mg., acetaminophen 325 mg., caffeine anhydrous 20 mg., ascorbic acid 100 mg /Tab. Bot. 50s

Use: Relief of common cold symptoms.

TACOSAL. (Tablet Rock) Hydrocortisone 2.5 mg., Vit. C 50 mg., Pot. salicylate 0.3 Gm., Al. hydrox. 120 mg., Pot. PABA 0.3 Gm /Tab. Bot. 100s

Use: Antirheumatic

TACT. (Hart) Boric acid 63%, sod borate 69%, ammonium alum 97%, glucono delta

lactone 4.9%, lactose 10%, carboric acid 0.86% w/camphor, menthol, eucalyptol, methyl salicylate and aerosol O.T. Jar 3 oz., 6 oz., 1 lb.

Use: Douche.

TADONA. (Tablet Rock Labs.) Bromisovaleryl- urea 300 mg., homatropine methylbromide 0.125 mg., hyoscyamine HBr 0.125 mg., scopolamine HBr 0.0065 mg./Tab. Bot. 100s, 500s.

Use: 1 t.i.d.; antispasmodic.

TAGATHEN. N.N.D. 1962. (Lederle) Chloro- then citrate, U.S.P., N,N-dimethyl-N'-(2-pyr- idyl)-N-[5-chloro-2-phenyl]-ethylenediamine citrate. Tab. (25 mg) Bot. 100s, 1000s.

Use: (1 q.i.d.) Antihistamine.

TAIN. (Smith-Dorsey) Triacetyloleandomycin equiv. to oleandomycin 125 mg., phenyl- propanolamine HCl 12.5 mg., pheniramine maleate 6.25 mg., pyrilamine maleate 6.25 mg., calcium acetosalicylate carbamide equiv to aspirin 300 mg /Inlay-Tab. Bot. 50s

Use: Upper respiratory infections

TAIN ORAL SUSP. (Dorsey) Triacetyloleando- mycin equiv to oleandomycin 125 mg., phenylpropanolamine HCl 12.5 mg., phenir- amine mol. 6.25 mg., pyrilamine mal. 6.25 mg., acetaminophen 150 mg /isp. Bot. 8 oz. Use. Common cold, secondary complications due to susceptible organisms

TAKA-COMBEX ELIXIR. (Parke, Davis) Taka Diastase 2½ gr., Vit. B-1 2 mg., B-2 1 mg., B-6 0.5 mg., nicotinamide 5 mg., panto- thenic acid 2 mg./4 cc. Bot. 1 pt.

Use: Vitamin B complex deficiencies.

TAKA-COMBEX KAPSEALS. (Parke, Davis) Taka-Diastase 2½ gr., Vit. B-1 10 mg., B-2 10 mg., B-6 0.5 mg., B-12 1 mg., panto- thenic acid 3 mg., nicotinamide 10 mg., Vit. C 30 mg., liver conc. 0.17 Gm., & liver fraction # 2 0.17 Gm./cap. Bot. 100s, 1000s. Use: Vitamin B complex and Vitamin C deficiencies.

TAKA-DIASTASE. (Parke, Davis) Aspergillus oryzae enzyme, a starch-converting fer- ment. Cap. (2½ gr.) Bot. 100s, 500s, liq. (20 gr./oz) Bot. 1 pt., 1 gal. Pow., Bot. 1 oz., ¼ lb., 1 lb. Tab. (2½ gr) Bot. 100s, 1000s.

Use: Symptoms of inadequate starch diges- tion.

W/Pepsin. Cal. carb. See: Calbase, Tab (Parke, Davis)

W/Phenobarbital. d-amphetamine sulfate & vitamins. See: Amphetasid-Phenobarbital, Tab. (Parke, Davis)

W/Phenobarbital & belladonna ext. See: Eardase, Tab. (Parke, Davis).

TAKA-DIASTASE COMPOUND. (Parke,

- Davis) Taka Diastase 1 gr, pepsin 2 gr, nux vomica ext 1/12 gr, capsicum 1/8 gr, papain 1 gr/Cap Bot 500s, 1000s
- TAKA-DIASTASE & PANCREATIN** (Parke, Davis) Taka Diastase 2 gr, pancreatin 3 gr/Tab Bot 100s
- TAKA-DIASTASE, PEPSIN & PANCREATIN**, (Parke, Davis) Taka Diastase 2 gr, pepsin 1 gr, pancreatin 2 gr/Tab Bot 100s, 1000s
- TAKAZYME** (Parke, Davis) Taka Diastase w/Cal carb, Mag carb, & bismuth subcarb Pow, Bot 2 oz, 5 oz
Use: Relief of gastric hyperacidity
- TA-LASE** (Tablet Rock) Dehydrocholic acid 130 mg, deoxycholic acid 65 mg, mylase 30 mg, prolose 6 mg, cellose 2 mg, atropine sulfate 0.0194 mg, hyoscyamine hydrobromide 0.1037 mg, hyosine hydrobromide 0.0065 mg/Cap Bot 100s 500s
Use Digestant, enzyme preparation
- TALATROL** (Abbott) See Tramethamine
Use Treatment of acidemia
- ***TALBUTAL**, NND 1962 5 allyl 5 sec butylbarb acid See Latusate, Tab (Winthrop Stearns)
- TAL-ESTAMINE** (Walker Corp) Pentobarbital 1/4 gr, scopolamine HBr 1/1200 gr, hexestral 1 mg/Tab Bot 100s.
Use: Estrogenic substance w/sedatives
- TALMIN** (Jackson Mitchell)
Tab: (Dihydroxy aluminum aminoacetate 5 gr, Mag carb 2 gr/tab) Bot 100s, 500s
Comp: (Dihydroxy aluminum aminoacetate 5 gr, belladonna ext 1/8 gr, Mag carb 2 gr, phenobarbital 1/4 gr/tab) Bot 100s, 500s
Use (1 2 1 d) Antacid
- TALON** (Warren Teed) Methylbenzethonium chloride 1:1000 phenol 0.5% w/acetonilid zinc oxide, calomine, eucalyptol in a water repellent base Oint, Tube 1 oz; Jar 1 & 5 lb
Use: Skin protective & antiseptic
- TALSIGEL** (Squibb) Phthalylsulfacetamide 1 Gm, galin 1 Gm, pectin 0.03 Gm, belladonna 0.3 cc/5 cc Susp, Bot 1 pl
Use: Diarrheas of nonspecific origin
- TALUDON**, (Hort) Belladonna alkaloids 1/600 gr, phenobarbital 1/4 gr, sulfur 1/2 gr/Tab Bot 100s 500s 1000s
Use (1 2 qid) Antispasmodic & sedative
- TAMA IRRIGATING SOLUTION**, (Tama) Isotonic solution
Use Flush ng agent for rinsing the eyes
- TAMA LACHRYMAL** (Tama) Simulated tear lubricant Solution
Use Tear lubricant for contact lens wearers
- *Generic name recognized by NF USP or AMA
- TAMA SOAKING SOLUTION** (Tama) Solution sterile formula Bot 4 oz
Use: Soaking contact lenses overnight
- TAMA WETTING SOLUTION** (Tama) Solution for contact lens wearers Plus Bot 2 oz
Use: Contact lens wearers—cleaning activity
- TANACAR**, (Enda) Tannic acid 2%, phenol 1%, otcahal 2% liq Bot 4 oz, 8 oz, 1 gal
Use: Antiseptic mouth spray or gargle
- TANBISMUTIN**, See: Bismuth Tannate
- TANCID**, (Superior) Tab Bot 500s, 1000s
- TANDEABIL** (Geigy) Oxyphenbutazone 100 mg/Tab Bot 100s
Use Rheumatoid arthritis
- TANICAINE**, (Upjohn) Oint (Atropine 1/4 gr, phenacaine 5 gr, w/phenol, nut gall, camphor, menthol & zinc oxide) 1 oz Supp. (Atropine 1/60 gr, phenacaine 1/2 gr w/phenol, nut gall & zinc oxide) Box ds 12s
Use: Local anesthetic & antiseptic
- TANIPENT** (Superior) Na 1—Pentaerythritol Tetranitrate 10 mg Na 2—Pentaerythritol Itranitrate 20 mg Na 3—Pentaerythritol Tetranitrate 10 mg phenobarbital 16 mg/Tab Bot 100s, 500s
Use Vasodilator
- TANNALBIN**, (Knoll) Tannin albuminate ex sec Pow, Bot 1 oz, 8 oz Tab (5 gr) Bot 100s, 500s, 1000s
Use Intestinal astringent & for diarrhea
- TANNIC ACID**, NF, NNR, 1941, A.D. 2; Gallotannic acid Glycerite, NF Pkg 1/4 lb, 1 lb
Use: 1 to 20% solution as an astringent
W/Benzocaine, quinine & urea HCl See Tanquen Oint (Sherman)
W/Benzyl alc p tertiary amyl phenol, alc See Burntanik, Sol (Horton & Converse)
W/Formaldehyde See Tannafarm, Pow (Meyer Drug)
W/Hexylresorcinol See: Hexyltan Jelly (Merck Sharp & Dahme)
W/Merthiolate See: Amerlan, Jelly (Lilly)
W/Phenol, Alcohol See: Tanacar, Liq (Enda)
W/Phenol & benzocaine See: Tanurot, Oint (Brean)
W/Procaine butyrate, ephedrine phenol alum See Butritannic, Oint (Rorer)
W/Secondary amyltricalcals See: Tanipaste (Upjohn)
- TANNIC ACID COMPOUND JELLY**, See Amerlan (Lilly)
- TANNIGEN**, NNR 1941 (Winthrop Labs) Acetyltannic acid Pow, Bot 1 oz
Use Astringent

- TANNOFORM.** (Mayer Drug) Helgofan. Tannin-formoldehyde. Bot. 1 oz.
Use: Astringent.
- TANPHENAMIN.** (Oxtramphetamine tannale). See: Synotan, Tab. (Irwin, Meisler)
- TANQUIN OINTMENT.** (Sherman) Benzocaine 1½%, tannic acid 3%, quinine & urea HCl Tube ¾ oz.
Use: Topically; local anesthetic & antiseptic.
- TANUROL.** (Breon) Tannic acid 3%, benzocaine 1% & phenol. Oint. 1 oz. Tube 1 oz.
Use: Topically; local anesthetic & antiseptic.
- TAO, NND 1962.** (Eaerig) Triethyl oleandomycin w/gleucosamine Cap. 250 mg., 125 mg Bot. 60s. Inj. (500 mg) Vial. Susp. 1.5 Gm., (125 mg /Tsp) Bot. 2 oz.
Use: Antibiotic therapy.
- TAO-AC.** (Roerig) Tao 125 mg, phenacetin 120 mg, caffeine 30 mg, salicylamide 150 mg., Buclizine HCl 15 mg./Cap. Bot. 36s.
- TAO PEDIATRIC OROPS.** (Eaerig) Oleandomycin base 100 Gm./cc. Vial 10 cc.
Use: Staphylococci infections in infants and children.
- TAOMID.** (Roerig) Trilocyloleandomycin 75 mg, sulfa-meth-dio-mer 113 mg. each/Tab. Bot. 60s.
Use: G.U. tract infections & upper respiratory diseases.
- TAOMID FOR ORAL SUSPENSION.** (Roerig) Trilocyloleandomycin 125 mg., sulfa meth-dio-mer 167 mg. each/tsp. Bot. 60 cc.
Use: G.U. tract infections & upper respiratory diseases.
- TAPAZOLE, NND. 1962.** (Lilly) Methimazole. 1-Methyl-2-Mercaptimidazole. Tab. (5 mg. & 10 mg.) Bot. 100s, 1000s.
Use: (20 mg. daily) Hyperthyroidism.
- TAPP.** (Pharmaloc) Thiamine HCl 25 mg, adiphenine HCl 20 mg, pyridoxine HCl 25 mg, phenyltoloxamine citrate 25 mg/Cop Bot 100s
Use: Antihistaminic w/vitamin supplement
- TAR.** See: Coal Tar.
- TARACTAN.** (Roche) Chlorprothixene Thi-oxanthene derivative 10, 25, 50 or 100 mg / Tab Bot. 50s/10 mg; 500s/10 mg; 50s/25 mg; 500s/25 mg; 50s/50 mg; 500s/50 mg; 500s/100 mg; 25 mg/2 cc., 100s.
Use: For emotional disorders.
- TARALBA.** (Upjohn) Formula B (Cool tar distillate 15%) Tube 1 oz. & 2 oz.
Use: Topically; antiseptic skin preparation.
- TAR BATH.** (Almay Div.) Juniper tar (code oil) 35 Gm., sulfonated oil, fatty acid esters & water/100 Gm. Bot. 8 oz., 1 pt., 1 gal.
Use: Antiparasitic and antipruritic.
- TARBONIS.** (Eed & Carrick) Tar ext. 5%, lanolin & menthol in a special vanishing type cream base. Tube 2¼ oz., 8 oz.
Use: Eczemas, psoriasis, seborrheic dermatitis, etc.
- W/Hydrocortisone.** See: Tarcartin Lotion. (Eed & Carrick)
- TARCAINE.** (Corbiselphoil) Cool tar 5%, benzocaine 1%, lanolin. Oint. 2 oz., 1 lb.
Use: Antipruritic.
- TARCORTIN.** (Eed & Carrick) Hydrocortisone 0.5% in tarbonis oint. base. Tube 7 Gm. & 1 oz. Aerosol 95 Gm.
Use: Topically, dermatitis
- TARCORTIN LOTION.** (Eed & Carrick) Tarbonis w/hydrocortisone 0.5%. Plastic squeeze bot. 15 cc
Use: Dermatology.
- TARCREME.** (Starenburgh) Cool tar, sol. 5% in oint. base. Oint., Tube ¾ oz., Jar 14 oz.
Use: Dermatitis.
- TAROLIN.** (Smith-Dorsey) Naphthalene, anthracene, phenanthrene, carbazole, phenol, cresol, benzene, alpha picoline, pyridine, quinoline & benzyl benzoate. Oint. 1 oz., 1 & 5 lb.
Use: Topically; antiseptic skin preparation
- TAROXIDE, MILD (NO. 1) OINTMENT.** (Abbott) Special cool tar distillate 1.5%, zinc oxide 15%, starch 24.5%, petrolatum 58%. Tube 1 oz., Jar 1 lb.
Use: Eczema.
- TARPASTE.** (Dook) Cool tar, distilled 5% in zinc paste. Jar 4 oz., 5 lb.
Use: Dermatitis.
- TARQUINOR.** (Squibb) Crude coal tar 1%, quinolor. 0.2% in hydrophilic cream, or Oint. in Plastibase. Jar 1 oz.
Use: Cutaneous disorders.
- TARTAR EMETIC.** See: Antimony Potassium Tartrate, U.S.P.
- TASNAN CREAM.** (Roche) Vit. A 10,000 u., d-pantothene 50 mg, vit. D-2 1000 u., E 5 mg/Gm. Tube 1 oz.
Use: Topically, symptomatic relief of burning & itching due to many skin irritations.
- TA-TEST.** (Hyland) Tert reagent combines purified thyroglobulin with polystyrene latex particles. Speedily detects thyroglobulin autprecipitin in serum specimens. Kits 20s
Use: Diagnostic agent for Hashimoto's disease and primary myxedema.
- TAURGLYPAN.** (Sherman) Bile salts 1 gr., nux vomica ext. 1/6 gr., papain 2 gr., cascara ext. ½ gr., phenolphthalein ½ gr., Capsicum oleoresin 1/40 gr./tab. Bot. 100s, 500s, 1000s.
Use: (1-2 h.) Laxative.
- TAUROCHOLIC ACID.** W/Pancreatin, pepsin. See: Enzymet. Tabs (Lloyd, Dobney & Westerfield)
- TAUROPHEN.** (Drug Prods.) Sod. succinate. Sod. salicylate. Sod. glycocholate, taurocho-

- late, glycodeoxycholate, taurodeoxycholate & bile salts present in ox gall cascara ext, phenolphthalein menthol Pulvoids, Bot 100s, 1000s
 Use: Chologogue, laxative
- TA-VERM** (Table Rock) Piperazine citrate 100 mg/cc Syr Bot 1 pt
 Use: Anthelmintic
- TAVILEN PLUS** (Table Rock Labs) liver Solution 1 Gm, Ferric Pyrophosphate Soluble 500 mg, B 12 24 mcg, B 1 6 mg, B 2 72 mg, B 6 3 mg, Panthenol 3 mg, Niacinamide 60 mg, Lysine HCl 300 mg, 5% alcohol, in a d sorbitol base/30 cc Bot 16 oz 1 gal
 Use: Hypochromic anemias
- TAXOL** (Lobico Debrulle) Bile salts 1 gr, aloes 2/5 gr, Hyoscyamus ext 1/13 gr, w/pancreas, duodenum/tab Bot 50s, 500s
 Use: (1 2 11 d) Laxative
- TAYVIMINS** (Taylor Drug) Vit A 25000 u, D 1000 u, B 1 10 mg, B 2 5 mg, B 6 1 mg, B 12 5 mcg, C 150 mg niacinamide 100 mg, 10 minerals/Cap Bot 100s, 500s, 1000s
 Use: Vitamins minerals therapy
- TCA P.** (Wallace) Cream: Trimethyl cetyl cilynic acid 1% Cream, Tube 1 oz Shampoo Trimethyl cetyl ammonium pentochlorophenol 1%, Bot 180 cc
 Use: Pungous infections of the skin
- TCM** (Wallace & Tiernan) Trichloroethylene Use: Germicidal
- T.C.S. TABLETS.** (Poythress) Theobromine salicylate 6 gr, Cal salicylate 1 gr & phenobarbital 1/4 gr Tab Bot 50s, 250s
 Use: Hypertension, angina pectoris, myocardial insufficiency
- TEAB-EFRIN** (Crookes Barnes) Methylcellulose 0.45% phenylephrine HCl 0.12%/Lochvial 15 cc
 Use: Lubricant decongestant
- TEARISOL** (Crookes Barnes) Methylcellulose 1/2% in sterile isotonic sol
 Use: Artificial tears in many eye conditions
- TECAL.** (Premo) Theobromine Cal salicylate, Tab, NF Paw, Bot 1 oz, 1/2 lb, 1 lb Tab (7 1/2 gr) Bot 100s, 1000s
 Use: Diuretic & myocardial stimulant
- ***TECLOZAN** (Winthrop) NN-(p phenylene dimethylene) bis[2,2 dichloro N (2 ethoxy ethyl)acetamide]
 Use: Amebicide
- TEDBAL** (Warner Chilcott) Theophylline 2 gr, ephedrine HCl 3/8 gr phenobarbital 1/8 gr/tab, supp or 4 cc of susp Also available in half strength Bot 8 oz Tab EC, Box 24s 120s 1000s Supp Box 12s
 Use: Symptomatic relief in asthma
 * Generic name recognized by NF USP or AMA
- TEDRAL ANTI-H.** (Warner Chilcott) Theophylline 130 mg, ephedrine HCl 25 mg, phenobarbital 8 mg, chlorpheniramine maleate 2 mg/Tab Bot 50s
 Use: Asthma
- TEORAL-SA.** (Warner Chilcott) Theophylline 180 mg, ephedrine HCl 48 mg, phenobarbital 25 mg/Tab Bot 100s
 Use: Prevention of asthmatic attacks
- TEEBACIN.** (Cansol Midland) Sod p-aminosalicylate 0.5 Gm & 0.69 Gm/tabs Bot 500s, 1000s, 5000s EC Tabs 0.5 Gm. & 1 Gm Bot 500s 1000s, 5000s Pow. Bot 1 lb Also special water soluble tablet (iterpack) equal to 300 Gm of free acid
 Use: (10 to 16 Gm daily) Tuberculosis.
- TEEBACIN ACID BUFFERED.** (Cansol Midland) p Aminosalicilic acid 0.5 Gm, with magnesium glycinate & carbonate/tab Bot 500s, 1000s, 5000s EC. Tab Bot 500s, 1000s, 5000s
 Use: (10 to 16 Gm, daily) Tuberculosis
- TEEBACIN KALIUM** (C.M.C.) Potassium p-aminosalicylate, 1 Gm/Tab Bot 500s, 1000s
 Use: Tuberculosis
- TEEBACONIN.** (Cansol Midland) Isoniazid 50 & 100 mg/tab Bot 30s, 100s, 250s 1000s
 Use: (1 or 2 tabs daily) Tuberculosis
- TEEBAZONE.** (Cansol Midland) Tibione, 4 acetyl amino benzaldehyde thiosemicarbazone 25 mg/tab Under study
 Use: Antituberculous
- TEEDS.** (Warren Teed) Tyrothricin 1 mg benzocaine 10 mg, w/chlorophyll/Tab Bot 100s, 1000s
 Use: Mouth & throat infection
- TEEK** (Norwich) Codeine sulfate 1/9 gr, chloroform, squill ext, ipecac ext, san guinaria ext & cherry syr./5 cc Bot. 3 oz Use: (1 dr. q 3 h) Cough preparation w/ narcotic
- TEENAC.** (Elder) Red mercuric sulfide 0.5%, colloidal sulfur 1.5% and urea
 Use: Treatment of acne
- ***TEFLURANE.** (Dow Chem Co) 2 bromo 1 1 1 2 tetrafluoroethane
 Use: General inhalation anesthetic
- TEENAMINS** (Rowell) Vit A 15000 u, D 500 u, B 1 5 mg, B 2 6 mg, B 6 2 mg C 75 mg, B 12 3 mcg, Calcium pantothenate 5 mg, Niacinamide 20 mg/Tab Bot 100s 250s and 1000s
 Use: Vitamin supplement
- TEGACID.** (Goldschmidt Chem Corp) See: Glyceril monostearate
- TEGA-BRON** (Ortega) Theophylline sodium glycinate 360 mg racephedrine HCl 30

- mg., potassium iodide 300 mg., niacinamide, 40 mg., secobarbital 20 mg./Tab. Bot. 100s, 500s.
 Use: Bronchial asthma, chronic asthmatics
- TIGA-NOLE.** (Ortega) Gomenol D1 Gm., gvaliacol 0.1 Gm., eucalyptol D05 Gm., iodoform 0.02 Gm., camphor D05 Gm., benzyl alcohol 2% / 2 cc. Vial 30 cc.
 Use: Bronchodilator.
- TIGA-OTIC.** (Ortega) Neomycin sulfate 4 mg., hydrocortisone 5 mg., pyrilamine maleate 0.6%, cetylaluminum chloride 0.5%, dipropion HCl 1%, urea 5%. Bot. 5 cc.
 Use: Treatment of otitis media and otitis externa.
- TIGA-SPIBITAL** (Ortega) Hesperidin (purified) 50 mg., ascorbic acid 50 mg., theobromine 5 gr., butobarbital sodium $\frac{1}{4}$ gr./Cap. Bot. 100s, 500s.
 Use: Cardiovascular diseases, capillary fragility.
- TIGUMIN CREME.** (Warren-Teed) Thienylpyrimine HCl 2%, calamine 8%, benzocaine 3%, hexachlorophene 3%. Cream, Tube 1 oz.
 Use: Topically; antiseptic & local anesthetic skin preparation.
- TILDIN SPANSULES, N.N.D. 1962.** (Smith, Kline & French) Chlorpheniramine maleate. Spansules (8 & 12 mg) Bot. 30s, 250s.
 Use: (1) In the morning & 1 in the evening) Antihistamine.
- TILOPAQUE, N.N.D. 1962.** (Winthrop Labs) Iodopanoic acid, 3-(3-amino-2,4,6-triiodophenyl)-2-ethylpropanoic acid. Tab. (0.5 Gm.) Envelope 6s, Box 5 & 25 envelopes
 Use: Orally; cholecystographic medfem.
- TELES SUSPENSION.** (Torch Labs.) Tellurium dioxide 2.5%, Bot. 4 oz.
 Use: Seboreheic dermatitis.
- TELLURIUM DIOXIDE.** See: Teles, Susp. (Torch Labs.)
- TEM.** See: Triethylene Metamine (Lederle).
- TEMABIL, N.N.D. 1962.** (Smith, Kline & French) Trimeprazine tartrate 2.5 mg./Tab. dt. 10-(3-dimethylamino-2-methylpropyl)-phenothiazine. Bot. 50s & 500s. Syr. (2.5 mg / 5 cc) Bot. 4 oz. Spansule 5 mg. Bot. 30s.
 Use: Antipruritic.
- TEMPOGEN.** (Merck Sharp & Dahme) Prednisolone 1 mg., acetylsalicylic acid 0.3 Gm., sod. ascorbate 60 mg., aluminum hydroxide gel. 0.2 Gm./Tab. Bot. 100s
 Use: Treatment of rheumatoid arthritis
- TEMPOGEN FORTI.** (Merck Sharp & Dahme) Prednisolone 2 mg., acetylsalicylic acid 0.3 Gm., sod. ascorbate 60 mg., aluminum hydroxide gel. 0.2 Gm./Tab. Bot. 100s
- Use: Treatment of rheumatoid arthritis and related diseases
- TEMPOSIL.** (Lederle) Under study
 Use: Antialcoholic.
- TEMPOTRIAD.** (Smith, Miller & Patch) Pentyl-enetetrol 100 mg., d-amphetamine sulf. 2.5 mg., caffeine anhydrous 100 mg./Tab. or 5 cc. Bot. 50s; Liq Bot. 4 oz
 Use: Stimulant—vasodilator.
- TEMPRA.** (Mead-Johnson) N-Acetyl-p-aminophenol. Bed draps 1 gr./0.6 cc. Bot. 15 cc. Green syrup 2 gr./5 cc. Bot. 3 cc.
 Use: Relief of fever and pain
- TEMPTEE-TABS.** (Bryant) Vit. A 5000 u., Vit. D 1000 u., Vit. C 60 mg., Vit B-1 2 mg., Vit. B-2 2 mg., Vit B-6 1 mg., Calcium pantothenate 3 mg., niacinamide 15 mg., Vit B-12 5 mcg / tab Chewable. Containers 40s. Apothecary jars 100s, 250s.
 Use: Nutrition supplement.
- TENACHE.** (Amfre-Grant) Butobarbital 15 mg., acetaminophen 200 mg., phenocellin 120 mg., caffeine 30 mg./Tab. Bot 100s, 1000s.
 Use: Analgesic and sedative for the relief of tension headache.
- TENAMINE.** (Hiss) Rescinnamine, 0.25 mg. Tab. Bot. 30s, 100s & 250s.
 Use: Hypotensive.
- TENICIDE.** (Tailby-Nason) Zinc sulfocarbolate, salicylic acid, iodine, phenol, formaldehyde, thymol, camphor, Set. 1, 2, 6 & 8 R. oz.
 Use: Antiseptic.
- TENNECETIN.** (Messengill) Tetraene antibiolic.
 Use: Antifungal.
- TENOC.** (Hiss) Isometheptene bitartrate. Vial 10 cc.
- TENSILON, N.N.D. 1962.** (Roche) Edrophonium chloride. Vial (10 mg./cc.) 10 cc., 1s.
 Use: Curare antagonist.
- TENSODIN.** (Knoll) Theophylline-Cal. salicylate 3 gr., ethavrine HCl $\frac{1}{2}$ gr., phenobarbital $\frac{1}{4}$ gr./Tab. Bot. 100s, 500s, 1000s.
 Use: Coronary and peripheral vasodilator, and sedative
- TENSOR.** (Crystal) Mannitol hexanitrate 30 mg., veratrum viride 100 mg., rutin 20 mg., Rauwolfia serpentina alkaloids 0.75 mg./Tab. Bot 100s.
 Use: Hypertension.
- TENTONE, N.N.D. 1962** (Lederle) Withdrawn. Methoxypramazine maleate 10, 25 & 50 mg / Tab. Bot 100s, 1000s
 Use: Tranquillizer.
- TENUATE, N.N.D 1962** (Marrell) Diethylpropion (1-phenyl-2-diethylaminopropane-

- 1 HCl Tab 25 mg, Bat 100s, 1000s Also Darpon Tabr 75 mg Bat 100s
 Use: Treatment and control of obesity
- TEOLAXIN.** (Poul Maney) Phenobarbital 100 mg/C D Cap Bat 100s, 1000r
 Use: Controlled distintegration, long-acting redative
- TEOTINE.** (Smith Darrey) Mannitol hexantrate $\frac{1}{4}$ gr, phenobarbital $\frac{1}{8}$ gr, theobromine $2\frac{1}{2}$ gr/Tab Bat 100s, 1000s
 1000r
 Use: (1 2 1 d) Vasodilator.
- TEOTINE R.C.** (Smith Darrey) Mannitol hexantrate $\frac{1}{4}$ gr, theobromine $2\frac{1}{2}$ gr, phenobarbital $\frac{1}{8}$ gr, rutin $\frac{1}{4}$ gr, Vit C 20 mg/Tab Bat 100s
 Use (1 2 1 d) Vasodilator
- TEPADRINE.** (Tertagor) Ephedrine NCl $\frac{3}{8}$ gr, phenobarbital $\frac{1}{4}$ gr, Sod Iodide 1 gr, theophylline $1\frac{1}{4}$ gr/Tab Bat 100r, 1000s
 Use: (1 1 d) Asthma
- TEPANIL.** NND 1962 (National) Diethyl propion Tab 25 mg Bat 100s
 Use Treatment and control of obesity
- TEPANIL TEN-TAB.** (National Drug) Diethyl propion 75 mg/Tab Bat 30s
 Use: Treatment of obesity
- TEPRIN.** (Endo) Not listed Ephedrine NCl $\frac{3}{8}$ gr, phenobarbital $\frac{1}{8}$ gr, theophylline $1\frac{1}{2}$ gr/Tab Bat 40s 100s 500r 1000s
 Use: (1 2 1 d) Asthma
- TEPRIN.** (Endo) Clonazepam
 Use Antihistaminic
- TERALASE.** W/Pancreatin, dehydrocholic acid, polysorbate 80 See: Digolase, Cap (Boyle)
- TERFONYL.** NNR 1955 (Squibb) Meth Dia Mer Sulfonamide, sulfadiazine 0.167 Gm sulfamerazine 0.167 Gm/Tab or 5 cc of Susp, Bat 1 pt Tab, Bat 100s 1000s See: Meth Dia Mer Sulfonamides, NNR
 Use: Triple sulfonamide therapy
- TERGEMIST.** (Abbott) 2 Ethylhexyl sulfate sod 0.125%, pot iodide 0.1% Abba Vial 40 cc, Bat 250, 500 cc
 Use: A mucolytic detergent To thin thick, tenacious sputum by nebulization of 3 to 5 cc q 1 d for 30 min
- TERGESTEN.** (Ulmer) Estradiol 1.2 mg, progesterone 25 mg, testosterone 25 mg/cc Aq Susp., Vial 10 cc
 Use: I.M., triple hormone therapy
- TERNYDROL.** (Irwin, Nelrier) Terpin hydrate 16 gr, chloroform 4 min, menthol, eucalyptol flavar q. /fl oz Bat 1 pt, 1 gal Also available w/codaine
 Use: (1 2 1 r q 2 3 h) Cough
- TERIDAX.** NND 1962 (Schering) fophenac acid, alpha ethyl beta 2,4,6 triado-3-hydroxyphenyl propionic acid Tab (0.75 Gm) Bat 30s, 100s, 500s
 Uses Diagnostic agent, cholecystography
- TERJINE PNOs.** (Merck Sharp & Dohme) Antleridine Phosphate
- TERJOLATE.** (White) Household detergent Bat 8 oz, 16 oz & 1 gal
 Use: Detergent nonirritating to skin
- TERMINAL.** (Benton) Vit A 6000 u, D 400 u B 1 5 mg, B 2 5 mg, B 6 2 mg, B 12 activity 2 mcg, C 100 mg, folic acid 0.25 mg niacinamide 20 mg, calcium pantothenate 5 mg, iron 58 mg, calcium 150 mg, diacetyl sodium sulfocuccinate 100 mg/Cap Units 100s
 Use Nutritional supplement
- TERMONE.** (Metropolitan Lab) Estrone 6 mg, testosterone 25 mg, progesterone 25 mg/cc Vial 5 & 10 cc
 Use Uterine hemorrhage
- TEROPIERIN SOLUTION.** (Lederle) Not listed Sod pteroyl triglutamate (1D mg/cc) Vial 1D cc
 Use IM, IV; palliation of symptoms in malignancy
- TERPANTONE.** (Strassenburgh) Terpin hydrate 4 gr, ethylmorphine HCl $\frac{1}{8}$ gr, Cal glycerophosphate $\frac{1}{4}$ gr, Sod glycerophosphate $1\frac{1}{2}$ gr, creosote carb 1/6 gr/fl oz Bat 1 pt, 1 gal
 Use: (1 1 r q 3 4 h) Cough; expectorant
- TERPIDEINE.** (Central) Codeine sulfate 1 gr terpin hydrate 20 mg/R oz Bat 1 pt, 1 gal
 Use: (1 1 s p q 3 h) Expectorant and cough sedative
- *TERPIN HYDRATE.** NF; Elixir, NF; Codeine Elixir, NF (Various Mfr) Opipentone Glycol Cryst & Pow, Pkg $\frac{1}{4}$ lb 1 lb
 Use Expectorant for chronic cough
 W/Caffeine, ammonium salicylate, ammonium gluconate, glycine, benzo'kenium chloride See: Martussin Special, Liq (M R Thompson)
 W/Carbetapantane citrate & chloroform See: Talcose Expectorant Compound (Pfizer)
 W/Chlorpheniramine maleate, phenylphthaline NCl, d methorphan HBr See: Syntussis Tab (Ives Cameron)
 W/Codeine phosphate, ammonium hypophosphite, Pot guaiaicolulfonate See: Bronch Tussin No 2, Sol (First Texas)
 W/Codeine phosphate, chloroform, menthol & glycine See: Coturus, Syr (National)
 W/Codeine phosphate & pectones See: Peptamoids W/Terpin Hydrate, Liq (Arlington)

- W/Cocaine sulfate. See: Terplidine, Liq. (Central).
- W/Creosote. See: Creo-Terpin, Liq. (Wampole).
- W/Dextromethorphan hydrabromide. See: T. H. & M., Syr. (Upjohn)
- W/Dextromethorphan, phenylephrine HCl, chlorpheniramine maleate, acetaminophen, ascorbic acid. See: Syntussin Compound, Cap. (Ives-Cameron)
- W/Dihydrocodonine bitartrate. See: Dro-daine, Syr. (Pitman-Maore).
- W/Dormethan, traminic. See: Tussaminic, Tab. (Smith-Darsey)
- W/Ethylmorphine HCl. See: Diaterpin, Liq. (Schleffelin).
- W/Ethylmorphine HCl, Cal. glycerophosphate, Sod. glycerophosphate, creosote carb. See: Terpatone, Liq. (Strazensburgh).
- W/Ethylmorphine HCl, creosote, eucalyptol. See: Kreo-Dionin-Terpin Compound, Liq. (Tilden).
- W/Guaiacol, glycerophosphates, ammonium chloride, tartar emelle, creosote carb., hyoscyamus fr. See: Neodyn, Syr. (National Drug)
- W/Morphine acetate, pot. guaiacol sulfonate, amm. hypophosphite. See: Broncho-Tussin No. 1. (First Texas)
- W/Norcollin, doxylamine succinate. See: Concol, Susp. (Merrell)
- W/Noscapine, sodium citrate, pyrilamine maleate, alcohol. See: Nektussin (Bruce)
- W/Tytrathricin & benzocaine. See: Bio-Kets, Traches. (Rexall)
- TERRABOH.** (Pfizer) Terramycin (as calcium di-terramycin) 125 mg./5 cc. homogenized mixture. Bot. 2 fl. oz., 1 pt.
Drops: Oxytetracycline 100 mg. in a specially homogenized fruit flavored mixture/cc. Bot. 10 cc.
Use: Orally, broad-spectrum antibiotic.
- TERRA-CORTIL** (Pfizer)
Topical Oint.: (Hydrocortisone 1%, terramycin 3%) Tube 1/6 & 1/2 oz.
Ophth. Susp.: (Hydrocortisone 15 mg terramycin HCl 5 mg./cc.) Bot. w/dropper 5 cc.
Topical Aerosol Spray. (Pfizer) Terramycin 300-mg., hydrocortisone 100 mg./85 Gm. Aerosol can 85 Gm.
Use: Anti-inflammatory.
- TERRAMYCIN, N.N.D. 1962, A.D.R.** Oxytetracycline, U.S.P. (terramycin amphoteric) Pediatric Drops: 1 Gm. dry pow. vial add water, 100 mg./cc. Bot. w/dropper 10 cc.
Oral Susp.: 1.5 Gm. dry pow. vial add water, 50 mg./cc. Bot. 1 oz.
Tab.: (50 mg.) Bot. 25s, 100s, (100 mg.) Bot. 25s, 100s, (250 mg.) Bot. 16s, 100s.
- Traches: (15 mg. ea.) Box 24 wrapped in foil.
See: Terraban (Pfizer)
- W/Hydrocortisone. See: Terra-Cortril Topical Aerosol Spray (Pfizer)
- TERRAMYCIN HYDROCHLORIDE, N.N.D. 1962, A.D.R.** (Pfizer) Oxytetracycline HCl, U.S.P.
Aerosol: (0.5 Gm.) 10 cc. (75% propylene glycol).
Cap.: (50 & 100 mg.) Bot. 25s, 100s, (250 mg.) Bot. 16s or 100s.
Conee, dental: (5 mg./cone) Vial 25s, Bot. 100s.
Elixir: (250 mg./fl.oz.) Bot. 1 fl. oz.
Drops, oral: (200 mg./cc.) when mixed w/diluent, Bot. 10 cc.
Intramuscular: Vial (100 mg.) 1s.
Intravenous: Vial (250 or 500 mg.)
Olet.: Tube 1/2 or 1 oz. (30 mg./Gm.)
Ophth. Oint.: (5 mg./Gm.) Tube 1/2 oz.
W/polymyxin B sulfate 1 mg./Gm.
Paste, dental: (30 mg./Gm.) Jar 5 Gm.
Otic: Dry pow. 5 mg./cc. W/polymyxin B sulfate 1 mg./cc. & benzocaine 5%. Comb. Pkg. Vial 25 mg. & Vial 5 cc. solvent 90% propylene glycol.
Paw., topicality: (30 mg./Gm.) W/polymyxin B sulfate 1 mg./Gm., Bot. 1 oz.
Sol. nasal, droppers: Vial 5 cc. (5 mg./cc.)
W/dextroxyphedrine HCl 0.25%.
Sol. ophtha (25 mg./5 cc.) buffered drop-per-vial 5 cc.
Sopp. vaginal: (100 mg./Supp.) W/polymyxin B sulfate 10 mg./Supp.
Tab. soluble: (50 mg./tab.) Bot. 24s.
Trocher 15 mg., Box 24s
- W/Neomycin sulfate. See: Enterobiotic, Tab. (Pfizer)
- W/Hydrocortisone acetate.
See: Cortril Acetate W/Terramycin Hydrochloride, Ophth. Susp. (Pfizer)
Terra-Cortril (Pfizer)
- TERRAMYCIN L.M. SOLUTION.** (Pfizer) Oxytetracycline 100 mg. or 250 mg w/dicaine/2 cc. Amp. 2 cc., Pkg. 5s, 100s.
- TERRAMYCIN NASAL W/Dextroxyphedrine HCl** (Pfizer) Terramycin HCl 5 mg., dextroxyphedrine hydrochloride 0.25%/cc. Sol. Bot. w/dropper 5 cc.
Use: Topically, antibiotic w/vasoconstrictor.
- TERRAMYCIN OTIC W/POLYMYXIN.**
(Pfizer) Terramycin 5 mg., polymyxin B 10,000 u., benzocaine 5%/cc. Bot. w/dropper 5 cc.
- TERRAMYCIN-SF.** (Pfizer) Oxytetracycline 250 mg., ascorbic acid 75 mg., thiamine mononitrate 2.5 mg., riboflavin 2.5 mg., niacinamide 25 mg., pyridoxine hydrochloride

- ride 0.5 mg, Cal pentathenate 5 mg, Vit B 12 1 mcg, folic acid 0.375 mg, menadione 0.5 mg/cap or 5 cc Cap Bot 16s, 100s Susp Bot 1 oz
Use: 4 daily; terramycin & eutritional therapy
- TERRAMYCIN TOPICAL** (Pfizer) Terramycin 30 mg, polymyxin 1000 u/Gm Pow Vial shaker top 1 oz
- TERRAMYCIN-VITAMIN** (Pfizer) Terramycin 250 mg, thiamine mononitrate 2.5 mg, riboflavin 2.5 mg, nicotinamide 25 mg, ascorbic acid 75 mg menadione 0.5 mg/cap
Use: Treatment of infections
- TERRASTATIN** (Pfizer) Oxytetracycline 25 mg gluco amine 250 mg eystate 250,000 u/Cap or Tsp Cap Bot 50s Susp Bot 2 oz
Use Antibiotic
- TERSASEPTIC** (Dook) A liquid cleanser, pH 6.8 w/hexochlorophene 1% Bot 4 oz, 1 pt, 1 gal
Use Liquid detergent
- TERSAVID** (Roche) N¹ pivaloyl N² benzyl hydrax ne
Use Monoamine oxidase inhibitor
- TERSULFAS, TAB, NNR, 1953** (Boyle) Sulfadiazine, sulfamerazine & sulfamethazine 0.167 Gm ea Tab (0.5 Gm) Bot 100s, 1000s See: Meth D a Mer Sulfonamides NNR 1954
Use: Tri sulfa therapy
- TERSUS** (Dook) Detergent Bot 4 oz, 1 pt, 1 gal
Use: Skin cleanser
- TERTIARY AMYL ALCOHOL, NNR, 1941**. See: Amylene Hydrate, USP (Various Mfr)
- TERTIASUL** (Rorer) Sulfamerazine 0.166 Gm, sulfadiazine 0.166 Gm, sulfathiazole 0.166 Gm/Tab Bot 100s, 1000s
Use: Triple sulfonamide therapy
- TESERENE** (Lloyd, Dabney and Westerfield) Dienestril 0.25 mg, methyl testosterone 5 mg, reserpine 0.1 mg/Tab Bot 100s
Use: Management of senility
- TESSALON, NND 1962** (C bo) w/methoxy poly (ethyleneoxy) ethyl p-butylam nabenzonate Benzonatate 100 mg/Cap Bot 100s Also pedatric perles 50 mg Bot 100s Amps. 15 mg/cc. 1 cc Cart 5s
Use: Control of coughs
- TESTACIDS IMPROVED**. (Reed & Carnrick) Methyltestosterone w/three factors at the vitamin B complex Tab Bot 100s, 500s
Use: Androgen w/vitamin therapy
- TESTADIOL, DEPO-** (Upjohn) Testosterone cyclopentylpropionate 50 mg., estradiol cyclopentylpropionate 2 mg./cc Vials 1 cc., 10 cc
Use I.M. hormone therapy
- TESTALDIOL** (Meyer) Estradiol 0.5 mg & testosterone 10 mg./cc. Vial 5 cc. Buccal Tabs (Estradiol 0.125 mg & methyltestosterone 5 mg./tab) Bot 100s.
Use Estragen-androgen therapy
- TESTANDRONE, NND 1962** (Carrick) Testosterone cryst Hypoglossals (10 mg.) Bot 100s, 1000s Vial, Aq Susp (25 & 100 mg./cc) 10 cc., 5s, 10s
Use I.M. (25 mg., 50 mg weekly); androgen therapy
- TES-TAPE**. (Lilly) Diagnostic test for glucose in urine Pkg 100s
Use Diagnostic agent
- TESTAPLEX**. (Horton & Converse) Methyl testosterone Tab 10 mg., Bot 30s, 100s; 25 mg., Bot 15s, 100s Sublingual Tab 5 mg., Bot 100s In oil Testosterone propionate 25 mg./cc Vial 10 cc. Aq. Inj. Testosterone susp 25 mg./cc Vial 10 cc.
Use Androgen therapy
- TESTARR**. (Testagor) Plantago seed mucilage. Also available w/thiamine Pkg 180 Gm., 390 Gm
Use: (2 tsp) Laxative
- TESTASCORBIC**. (Testagor) Ascorbic acid. Cap (250 mg) Bot 100s, 1000s.
Use Vitamin C therapy
- TESTAVOL** (Testagor) Vit A 100,000 u./Cap or cc Amp (1 cc) 10s Cap., Bot 100s.
Use I.M. & oral; Vitamin A therapy
- TESTAVOL-S**. (Testagor) Vitamin A palmitate Cap 25,000 u., 50,000 u., Bot 100s, 1000s
Use: Vitamin A deficiencies
- TESTEED**. (Warren Teed) Methyltestosterone 5 mg., (ethinyl estradiol) 0.025 mg./tab. Bot 30s, 100s
Use: Female & male climacteric.
- TESTODET**. (Merck Sharp & Dohme) Testosterone propionate in oil Vial (25 mg./cc.) 5 cc.
- TESTOFEROL**. (Ascher) di-alpha Tocopheryl acetate 100 mg., testosterone propionate 25 mg./cc Vial 10 cc.
Use: I.M., 1 cc. twice weekly, menopausal disorders
- TESTORA**. (Chicago Pharm.) Methyltestosterone Tab (25 mg.) Bot 30s, 100s.
Use: Androgen therapy
- TESTOSTEROID, NND 1962** (Shermae) Testosterone crystalline aq susp Vial (25 mg./cc.) 10 cc., (50 mg./cc.) 10 cc.
Use: I.M. (25 to 50 mg weekly); androgen therapy

- ***TESTOSTERONE, N.F., N.N.D. 1962;** Inj; Pellets, N.F.
Aqueous Susp.
 See: Androlin, Aq. Susp. & In oil, Vial. (Lincoln)
 Andronaq, Aq. Susp., Vial. (Central Pharm. Co.)
 Andrusol, Aq. Susp., Vial (C. D. Smith).
 Aqua-Testosterone, Aq. Susp., Vial (Endocrine)
 Diphasal Testosterone, Aq. Susp., Vial (Kremers-Urban).
 Harvogen, Vial (G. F. Harvey)
 Malastrone, Aq. Susp., Vial (Kirk).
 Martestote, Aq. Susp., Vial (Breon).
 Sterotote (Ulmer)
 Neo-Hombreal-F, Aq. Susp., Vial (Organon).
 Oretan-F, Aq. Susp. & Pellets (Schering).
 Synandrol-F, Aq. Susp., Vial (Pfizer).
 Testandrone, Aq. Susp., Vial (Cornrick).
 Testobase, Aq. Susp., Vial. (Merck Sharp & Dohme)
 Testosteroid, Aq. Susp., Vial. (Sherman)
 Testrone, Aq. Susp., Vial (Miller).
 Testryl, Aq. Susp., Vial (Squibb).
- W/Estradiol.**
 See: Aquodiol w/Testosterone, Vial. (National Drug)
 Cambandrets, Tab. (Pfizer).
W/Estradiol & chlorobutanol in cottonseed oil. See: Depo-Testadiol, Vial. (Upjohn)
W/Estradiol, procaine HCl. See: Testrogyn, Vial (Ascher).
W/Estradiol & progesterone.
 See: Tergesten, Vial. (Ulmer)
 Trigonaq, Vial (Central)
 Trihorn, Vial. (Ascher)
 Tri-Hormogen, Vial. (Mallard)
- W/Estradiol, progesterone, procaine HCl, procaine base.** See: Hormo-Triad, Vial. (Bell Pharmacal)
W/Estrogenic substance.
 See: Andreitol, Vial (Central).
 Kestos, Vial (Burlington).
 Sherestos, Vial (Sherman).
 Testrogen, Vial (Rowell).
- W/Estrogenic substance & thiamina.**
 See: Duohorm, Vial (Elder).
 Thelal, Vial. (Durril)
- W/Estrone.** (Ayers) Each cc. provides 20 mg. of testosterone & 2 mg. (20,000 I.U.) of estrone when completely suspended. Vial 10 cc.
 See: And-Est, Vial. (Ulmer)
 Andasterone, Vial. (Lincoln)
 Androvarin, Vial (Warren-Teed).
 Anestra, Vial (Mallard).
 Angen, Vial. (Eaton)
 Di-Steroid, Vial (Kremers-Urban).
- Generic name recognized by N.F., U.S.P. or A.M.A.**
- Duogen, Vial (C. D. Smith).
 Hormestrin-T, Vial (Miller).
 Mal-O-Fem, Aq. (Testagor)
W/Estrone & progesterone.
 See: P-E-T, Vial (Maury).
 Patrone, Vial (Kremers-Urban).
 Pro-Mal-O-Fem, Vial. (Testagor)
 Termone, Vial. (Metrol)
W/Estrone, progesterone & procaine HCl. See: Kar-Tep, Aq. Susp. (Kormel Co.)
W/Progesterone, estrogenic substance. See: Trimitann, Vial. (Miller)
W/Progesterone, estradiol. See: Tergesten Myasal, Inj. (Ulmer)
W/Thiamine HCl, glutamic acid, procaine HCl. See: Glutest, Vial. (Research Supplies)
- ***TESTOSTERONE CYCLOPENTYLPROPIONATE, U.S.P., N.N.D. 1962;** Inj., U.S.P.
 See: Depo-Testosterone Cyclopentylpropionate, in Oil (Upjohn)
 Testadiol, Depo. (Upjohn)
W/Estradiol 17-cyclopentylpropionate. See: Depo-Testadiol, (Upjohn)
- ***TESTOSTERONE ENANTHATE, U.S.P., N.N.D. 1962;** Inj., U.S.P.
 See: Delatestyl Solution, Vial (Squibb)
 Reposo-TMD, Vial (Confield)
- W/Estradiol valerate.**
 See: Deladumone, Vial (Squibb)
 Estomone (Key)
 Reposo-TE, Vial (Confield)
- TESTOSTERONE HEPTANOATE, N.N.D. 1962.**
 See: Testosterone enanthate.
- TESTOSTERONE PHENYLACETATE.** See: Perandree Phenylacetate, Vial. (Ciba)
- ***TESTOSTERONE PROPIONATE, U.S.P., N.N.D. 1962;** Inj., U.S.P.
 See: Aq. Susp.:
 Androlin, Vial. (Lincoln)
 Andro-Hart, Vial (Hart).
 Andrusol, Vial (C. D. Smith).
 Oretan, Aqueous, Vial (Schering)
 Synandrol, Vial (Pfizer).
- In Oils**
 Androlin, Vial. (Lincoln)
 Andro-Hart, Vial (Hart).
 Androlan, Vial. (Lanett)
 Androlin, Oil Susp. (Lincoln)
 Andrusol-P. (C. D. Smith)
 Hormafe, Vial (Key Corp)
 Masenate, Vial (Schleffelin).
 Neo-Hombreal, Vial (Organon).
 Orchic-ol, Vial. (Cole Chem. Co.)
 Oretan, Vial (Schering)
 Perandran, Vial (Ciba).
 Synerann, Vial (Pitman-Moore).
 Testodet, Vial. (Merck Sharp & Dohme)
 Testrone-P, Vial. (Miller)
- W/Estradiol benzoate.**

- See: Bi Ger Mone, Vial (Kirk)
 Combandrin, Vial (Pfizer)
 Di Mel BP, Vial (Organon)
 Extramal Improved, Vial (Key Corp.)
 Gynelone, Vial (Schering)
 Mal O Fem, Vial (Testagar)
 Testrial, Vial (G F Harvey)
- W/Estragenic substances (Ayerst) Each cc provides 10 mg of testosterone propionate & activity equiv to 10,000 IU (1 mg) of estrane or estragenic substances Vial 10 cc.
- W/Progesterone & estradiol
 See: Espranote, Vial (Chicago Pharm.)
 Trigestandrial, Vial (Kirk)
 Triharmonal, Vial (Key Corp.)
 Trimilane, Vial (Miller)
- W/Progesterone & estradiol benzoate See: Pro Mal O Fem, Vial (Testagar)
- W/dl alpha Tocopherol acetate
 See Andropheral, Vial (Central)
 Testoferal, Vial (Ascher)
- W/Alpha Tocopherol Chlorobutanol See: Andratocopherol (New Mexico Pharm)
- TESTRADIOL** (Metropolitan Eob) Estradiol benzoate 1 mg, testosterone propionate 20 mg/cc Aq susp & in oil, Vial 10 cc
 Dose: 0.5 to 1.5 cc
 Use: Menopausal syndrome
- TESTRAMONE** (G F Harvey) Testosterone 12.5 mg, vit B1 10 mg, B2 2 mg, B6 5 mg, niacinamide 40 mg, inositol 50 mg, choline chloride 10 mg, methionine 10 mg/cc Amp 10 cc
 Use: Androgen therapy with Vitamin B complex
- TETRA-CORT OINT** (Federal Pharm) Hydrocortisone free alcohol 5 mg neomycin base 3 mg polymixin B sulfate 4000 u tyrothricin 0.05%, gramicidin 0.01%/Gm Tube 5 Gm 15 Gm
 Use: Dermatitis
- TESTRAQ.** (New Mexico Pharm) Testosterone 100 mg, sodium chloride D9% thimerosal 1:10,000 distilled water/cc Vial 10 cc
 Use: Mammary carcinoma
- TESTRIOL** (G F Harvey) Estradiol 1 mg, testosterone propionate 10 mg in sesame oil/cc Vial 10 cc
 Use: Menopausal complaints
- TESTROGEN.** (Rowell) Testosterone 25 mg & natural estrogen 2 mg/cc Vial 10 cc
 Use: I.M.; estrogen & androgen therapy
- TESTROGYN.** (Ascher) Beta estradiol 0.5 mg, testosterone 10 mg, procaine HCl 2%/cc Vial 10 cc.
 Use: Male & female hormones
- TESTROLIX.** (Chicago Pharm) Ethynyl estradiol 0.02 mg, methyltestosterone 10 mg/4 cc Elx Bot 4 oz, 1 pt, 1 gal
 Uses: Male & female hormone therapy
- TESTRONE, NND 1962** (Smith, Miller & Patch) Testosterone in aq susp Vial (25 & 100 mg/cc) 10 cc Box 6s, 25s (50 mg/100 cc)
 Use: I.M.; androgen therapy
- TESTRONE-P.** (E S Miller Labs) Testosterone propionate in oil Vial 25 mg/cc. 10 cc.; 50 mg/cc. 10 cc.; 100 mg/cc. 10 cc. 1s, 6s 25s
- TESTRYL, NND 1962** (Squibb) Testosterone Vial (50 mg/cc.) 10 cc.
 Use: Androgen therapy.
- TETANUS.** Diphtheria Combinations. See: Diphtheria and Tetanus Toxoids, Combined
- ***TETANUS ANTITOXIN, U S P**
 Cutter—Vial 1500, 5000, 10,000 & 20,000 u
 Lederle—Vial 1500 & 3000 u
 Lilly—Vial 1500, 5000, 10,000 & 20,000 u
 National Drug—Vial 1500, 3000, 5000, 10,000 & 20,000 u
 Parke, Davis—Vial 1500, 5000, 10,000 & 20,000 u
 Merck Sharp & Dahme—Vial 1500 u, 10,000 u
 Wyeth—Vial 1500, 20,000 & 40,000 u
- TETANUS-DIPHTHERIA TOXOIDS COMBINED, ALUMINUM PHOSPHATE ADSORBED, ULTRAFINED** (Wyeth) Vial 5 cc
 Use: I.M., Primary Immunization or maintenance in persons over 10 years old
- TETANUS-GAS GANGRENE ANTITOXIN NP, NNR 1952**
 Lederle—Vial globulin modified 1 dose
 Parke, Davis—Vial 1500 u
 Wyeth—Vial, polyvalent
- TETANUS IMMUNE GLOBULIN, HUMAN**
 See: Hyper Tet Injection (Cutter)
- ***TETANUS TOXOID, U S P, NND 1961**
 Cutter—Vial 1.5 & 15 cc
 Lederle—Vial 1.5 & 7.5 cc
 Lilly—Vial 1.5 & 7.5 cc
 Parke, Davis—Vial 1.5 & 7.5 cc
 Merck Sharp & Dahme—Vial 1.5 & 15 cc
 U S Standard—Vial 1.5, 7.5 & 22.5 cc
 Wyeth—Vial 7.5 cc
- ***TETANUS TOXOID, ALHYDROX, NND 1962** (Cutter) Tetanus toxoid, aluminum hydroxide adsorbed, U S P Vial 1.5 & 7.5 cc
- ***TETANUS TOXOID, ALUMINUM HYDROXIDE ADSORBED, U S P NND 1962** (Cutter) Vial 1 & 5 cc
- ***TETANUS TOXOID, ALUMINUM PHOSPHATE ADSORBED, U S P**
 See: Tetanus Toxoid, alhydrox Cutter—Vial 1.5 & 7.5 cc

*Generic name recognized by N F U S P or A M A

- ***TETANUS TOXOID, ALUM PRECIPITATED, U.S.P., N.N.R. 1957.**
 Cutter—Vial 1 & 5 cc.
 Lilly—Vial 1 & 5 cc.
 National—Vial 0.5, 2 & 5 cc.
 Parke, Davis—Vial 1 & 5 cc.
 Pitman-Moore—Vial 5 cc.
 Merck Sharp & Dohme—Vial 1 & 5 cc.
 Squibb—Vial 2 & 10 cc.
 U. S. Standard—Vial 1 & 5 cc.
 Wyeth—Vial 5 & 10 cc.
- TETIOTHALEIN SODIUM, N.N.R. 1944. See: Iodophthalin Sodium, U.S.P.**
- TETRAREE. (Endo) Thiamine HCl 100 mg., Nicotinamide 100 mg., riboflavin 1 mg., pyridoxine HCl 1 mg./cc. Vial 10 cc. Urr: Pellagra.**
- TETRARIOTIC. (Bryant) Polymixin B sulf. 4 M g., neomycin sulf. 3 mg., tyrothricin 0.5 mg., gramicidin 0.125 mg., benzocaine 10 mg./Gm. Oint. Tube ½ oz.**
 Use: Topically, antibiotic treatment.
- TETRACON V SYRUP. (Pfizer) Tetracycline phosphate complex equivalent to 125 mg. of the HCl/5 cc. Bot. 2 oz., 1 pt.**
- ***TETRACAIN HYDROCHLORIDE, U.S.P., A.D.B., Inj., U.S.P.**
 See: Pentacaine Hydrochloride, N.N.E. Winthrop Labs.)
 Tetracel, Bulepto (Prof. Pharm.)
 W/Isocaline & benzalkonium chloride. See: Isotraline, Oint. (Columbus).
 W/Isocaline, tyrothricin & menthol. See: Tyrisotraline, Lox. (Columbus)
- TETRACAIN HYDROCHLORIDE 0.5%.**
 Dropper has .5%/1 cc. (Iso-Sol) Droptainer, Ophth., 15 cc. (Alcan)
- ***TETRACHLOROETHYLENE, U.S.P., N.N.R. 1949. Cap., U.S.P. Perchloroethylene, tetrachloroethylene, U.S.P. (Various Mfr.)**
- TETRACEL, BUFOPTO. (Professional Pharm.)**
 Tetracaine HCl ½% in methylcellulose soln. Bot. dropper 15 cc.
 Use: Anesthetic in ophthalmology
- TETRACHLOROETHYLENE, U.S.P., N.N.R. 1949. See: Tetrachloroethylene, U.S.P.**
- TETRACHLORSALICYLANILIDE. See: Impregon (Fleming)**
- TETRACILLIN. (Schenley) Penicillin G Pot. 100,000 u., sulfadiazine 0.167 Gm., sulfamerazine 0.167 Gm., sulfamethazine 0.167 Gm./Tab. or 5 cc. Tab., Rat. 10s, 50s, 250s. Liq., Bot. 60 cc.**
 Use: Triple sulfonamide with penicillin therapy.
- ***TETRACYCLINE, U.S.P., N.N.D. 1967. Oral Susp. U.S.P.**
 W/N acetyl-para-amino-phenol, phenyltoloxamine citrate See: Tetrax Ap (Bristol)
- ***TETRACYCLINE HYDROCHLORIDE, U.S.P., N.N.D. 1962, A.D.R., Cap., U.S.P.; Inj., U.S.P.; Ophth. Sol., U.S.P.; Orol Surp., U.S.P.; Tab., N.F.**
 See: Achromycin (Lederle)
 Panmycin, Prop. (Upjohn)
 Polycycline (Bristol)
 Stactin, Cap. (Squibb)
 Tetrabon, Susp. (Pfizer)
 Tetracycl (Roerig; Pfizer)
 W/Hydrocortisone. See: Achromycin W/Hydrocortisone, Oint. & Ophth. Oint. (Lederle)
 W/Hydrocortisone acetate & phenylephrine NCL. See: Achromycin Noral Suspension, (Lederle)
 W/Meth-dia-mer sulfenamides. See: Polycycline W/Triple Sulfenamides, Susp (Bristol)
 W/Citric Acid. See: Achromycin V (Lederle)
 W-Neomycin sulfate. See: Surgamycin Topical Spray Ointment, Aeroral (Cyanamid)
 W/Novobiocin calcium, potassium metaphosphate. See: Penalba KM Drops (Upjohn)
 W/Nystatin.
 See: Achrostatin V, Cap. (Lederle)
 Comycin, Cap. (Upjohn)
 Tetrostatin, Cap. & Susp. (Pfizer)
 Mystecilin, Cap. (Squibb)
 W/Oleandomycin.
 See: Signomycin (Pfizer)
 W/Phenacolin, caffeine, salicylamide, buclizine HCl. See: Tetrocydin, Tab. (Pfizer)
 W/Stress formula vitamins. See: Tetrabon SF, Susp. (Pfizer)
 W/Vitamins & stress formula.
 See: Achromycin SF, Caps. (Lederle)
 Tetracycl-SF, Cap. & Susp. (Pfizer)
 W/Xylocaine. See: Polycycline, Ophth. Oint. (Bristol)
- ***TETRACYCLINE PHOSPHATE COMPLEX, N.N.D. 1962.**
 See: Panmycin Phosphate, Cap. (Upjohn)
 Sumycin (Squibb)
 Tetrax (Bristol)
- W/Amphotericin R. See: Mystecilin F (Squibb)
 W/Aspirin, phenacolin, caffeine, phenyltoloxamine HCl. See: Tetrax APC W/Bristolmin, Cap. (Bristol)
- W/Mycostatin. See: Mystecilin-V, Cap. (Squibb)
- W/Nystatin
 See: Achrostatin-V, Cap. (Lederle)
 Comycin, Cap. (Upjohn)
- W/Sulfamethizole, phenylazodiaminopyridine HCl. See: Azotrex, Cap. (Bristol)
- TETRACYCLINE, SUSPENSION, ORAL, U.S.P. See: Brand names under the hydrochlorides.**
- ***Generic name recognized by N.F., U.S.P. or A.M.A.**

TETRACYDIN. (Pfizer) Tetracycline HCl 125 mg, phenacetin 120 mg, caffeine 30 mg, salicylamide 150 mg, buclizine HCl 15 mg / Tab Bot 24s, 100s

Use: Analgesic-antihistaminic

TETRACYN, N N O 1962 (Roerig) Old name Cosa Tetracycln Tetracyclina amphoteric Susp oral chocolate paw 250 mg / 5 cc Bot 1 oz

TETRACYN HYDROCHLORIDE, N N D 1962 (Roerig) Tetracycline HCl with glucosamine Cap (125 mg) 25s & 100s, (250 mg) 16s, 100s

I M. Vial 100 mg & 250 mg

I V. (250 and 500 mg w/procaine HCl Mag chloride Vit C/2 cc) Vial

Syr 125 mg N acetylglucosamine eq to 125 mg of gluc HCl/5 cc Bot 2 oz, 1 pl Pediatric Drops (100 mg/cc) Bot 10 cc Tabs - 250 mg Bot 100s

TETRADECYL SULFATE, SODIUM, N N D 1962 See Sodium Sotradecol

TETRADENE. (Smith Miller & Patch) Pentyl ene tetrazol 100 mg, nicotinic acid 50 mg, tetraiodothyranine 10 mcg, methyl testolone 1 mg, ethinyl estradiol 0.002 mg / Tab or 5 cc Bot 100s, 1000s Elix Bot 1 pt Use Geriatrics

TETRAETHYLAMMONIUM BROMIDE

{T*ARI

Use: Diagnostic & therapeutic agent in peripheral vascular disorders Diagnostic in hypertension

***TETRAETHYLAMMONIUM CHLORIDE, N N D 1962** (McNeil) Vial (0.1 Gm/cc) 20 cc See Etamyl Chloride, Vial (Parke, Davis)

Use I V, ganglionic blocking agent

TETRAETHYLTHIURAM DISULFIDE. See: Disulfiram, N N D 1960

TETRAGLYCINE HYDROPERIODIDE, See: Globaline, Tab (Wallace and Tiernan)

Brusoline (Burnham Soluble Iodine)

TETRAHYDROPHENOBARBITAL See: Cyclo barbitol, Prep (Winthrop Labs)

TETRAHYDRO-PTEROYLGLUTAMIC ACID See: Leucovorin Vial (Lederle)

***TETRAHYDROZOLOINE HYDROCHLORIDE, N F, N N D 1962** 2 (Tetrahydrozoline HCl 1%)imidazoline HCl

See: Cor Tyzine Nasal Sol n (Pfizer)

Cor Tyzine Pediatric Nasal Drops (Pfizer)

Tyzine, Sol (Pfizer)

Visine (Pfizer)

/Prednisolone See: Cortyzomal (Pfizer)

Cor Tyzine, drops (Pfizer)

/Prednisolone, neomycin sulf See: Neo-

cor Tyzine, Nasal drops (Pfizer)

TETRAIODOPHENOLPHTHALEIN SODIUM,

Generic name recognized by N F, U S P or A M A

N N R 1950 See: Iodophthalein Sodium, U S P

TETRAIODOPHTHALEIN SODIUM, N N R 1950 See: Iodophthalein Sodium, U S P

I-3,5,3',5'-TETRAIODOTHYRONINE SODIUM Levothyroxine Sod, N N D 1958

See: Synthroid Tab (Troveal Labs)

W/Pentylene tetrazol, nicotinic acid, methyl testosterone, ethinyl estradiol See: Tetra Forte, Tab (E S Miller)

TETRAMETHYLENEDIMETHANESULFONATE See: Busulfan, U S P

TETRAMETHYLTHIURAM DISULFIDE. Thiram See: Rezifilm (Squibb)

Use Antibacterial & antifungal agent

***TETRANTOIN, 7,8 benzo 1,3 diazospiro (4,5)decane 2,4 diane** See: Spirodon (Culter Lab)

TETRASTATIN (Pfizer) Tetracycline HCl 250 mg glucosamine HCl 250 mg, nystatin 250 000 u / Cap Bot 16s 100s

Use Antimicrobial and anti infective

TETRASTATIN ORAL SUSP. (Pfizer) Tetracycline 125 mg glucosamine HCl 125 mg, nystatin 125 000 u / 5 cc Bot 2 oz

Use Antimicrobial and anti infective

TETRAYAX. (Merck Sharp & Dahme) Multiple vaccine 4 in 1, Vial 9 cc

Use: To provide protection against diphtheria, whooping cough, tetanus and polio myelitis

TETRAZETS (Merck Sharp & Dahme) Zinc bacitracin 50 u Tyrothricin 1 mg, neomycin sulfate 5 mg, benzocaine 5 mg / troche Vial 12s

Use Sore throat due to either gram positive or gram negative bacteria

TETREX, N N D 1962 (Bristol) Tetracycline phosphate complex (quantities as hydrochloride equiv)

Cap: (250 mg) Bot 16s, 100s (100 mg) Bot 25s, 100s

Pediatric Drops: (100 mg/cc) Bot 10 cc Syr: (125 mg/5 cc) Bot 60 cc

Intramuscular w/Xylacaine: (100 mg or 250 mg/Vial)

W/Sulfamethizole and phenylazo diamino pyridine HCl See: Azalrex, Cap (Bristol)

TETREX AP SYRUP. (Bristol Lab) Tetracycline base 125 mg, N acetyl para amino phenol 120 mg, Phenyltoloxamine citrate 12.5 mg / 5 cc Bot 2 oz

Use Respiratory infections

TETREX APC W/Bristamin (Bristol) Tetracycline phosphate 125 mg, aspirin 150 mg, phenacetin 125 mg, caffeine 30 mg

phenyltoloxamine HBr 25 mg / Cap Bot 24s

Use: Treatment of the common cold

- TETREX W/TRIPLE SULPAS.** (Bristol) Tetracycline HCl 125 mg., sulfadiazine, sulfamerazine, sulfamethoxazole 167 mg. ea./5 cc. Bot. 2 oz.
Use: Severe bacterial infections.
- TETROGEN.** (Endo) Acetophenetidin 65 mg., sod. salicylate 320 mg., salicylic ethyl ester carbonate 32 mg., calcium root pow. ext. 1.6 mg./Tab. Bot. 100s, 1000s
Use: Analgesic in the treatment of rheumatic pains
- TETROPENTOBARBITAL.** (Strong Cobb) Pentobarbital Sod. 100 mg., pentylene-tetrazol 300 mg./Cap. Bot. 24s, 100s, 500s.
Use: Sedative.
- TETROPHENOBARBITAL.** (Strong Cobb) Phenobarbital Sod. 100 mg., pentylene-tetrazol 150 mg./Cap. Bot. 24s, 100s, 500s.
Use: Sedative.
- TETROSECOBARBITAL.** (Strong Cobb) Secobarbital Sod. 100 mg., pentylene-tetrazol 300 mg./Cap. Bot. 24s, 100s, 500s.
Use: Sedative.
- TEXACORT.** (Texas Pharm.) Texacort-25-hydrocortisone alcohol 0.25% 1 oz. tubes & dexamethasone in a non-ionic lubricating base 4 oz. jars in a non-ionic lubricating base pH 4.6, texacort-50-hydrocortisone alcohol 0.5% ½ oz. & 2 oz. tubes; texacort-100-hydrocortisone alcohol 1.0% 10 Gm. tubes.
Use: Sedative with diuretic.
- TEXACORT LOTION 25**—2 fl. oz., 4 fl. oz. & pt. bottles.
Use: Dermatitis.
- TEXACORT LOTION 50**—2 fl. oz. bot
Use: Dermatitis.
- T-FLUORIDE.** (Tennessee Pharm.) Sodium fluoride 2.31 mg./Tab. Bot. 100s, 1000s.
Use: Dental caries and prophylactic.
- THALAJEN.** (Jenkins Labs.) Phthalylsulfacetamide w/kaolin & pectin, Tab. 500 mg., Bot. 100s, Tab. Jr. (250 mg.) Bot. 100s. Susp. (1.0 Gm./5 cc.) Bot. 1 pt.
Use: Sulfanamide therapy.
- THALAMYD.** (Schering) Phthalylsulfacetamide, Tab. (0.5 Gm.) Bot. 100s, 1000s.
Use: Sulfanamide-susceptible organisms of the intestinal tract.
- THALREXYL CAPSULES.** (March Sharp & Ohme) Phthalylsulfathiazole 0.5 Gm., hexylresorcinol (in oil) 0.1 Gm./Cap. Bot. 100s.
Use: Acute & chronic urinary-tract infections.
- THALFED.** (Massengill) Aminophylline 1½ gr., ephedrine ½ gr., phenobarbital ½ gr./5 cc. Elixir, Bot. 1 pt., 1 gal. Yab., Bot. 100s, 500s, 1000s.
Use: (1-2 q. 3-4 h.) Asthma.
- THALIDOMIDE.** α[N-phthalimido]glutaramide See: Kevadan (Merrell Co.)
- THALINOL.** (M. R. Thompson) Not listed.
- ^a Generic name recognized by N.F., U.S.P. or A.M.A.
- Phenolphthalein. Cap. (2 gr. in oil) Bot. 20s, 100s, [5 gr. in oil] Bot. 30s, 100s
Use: Laxative.
- THALOCTIN.** (Abbott) Phthalylsulfacetamide 1 Gm., kaolin 1 Gm., pectin 80 mg. in a flavored aq. vehicle. Contains no sugar/5 cc. Susp., Bot. 1 pt.
Use: Gastro-intestinal tract infections.
- THALOSEN.** (Abbott) Phenolphthalein 1½ gr., w/senna, sulfur & aromatics/Tab. Bot. 100s, 1000s.
Use: Laxative.
- THALTIN.** (Abbott) Phthalylsulfacetamide 15 gr., kaolin 15 gr., pectin 1 gr./5 cc. Susp., Bot. 1 pt.
Use: (1-3 dr. daily) Sulfanamide therapy for intestinal infections.
- THANTIS LOZENGES.** (Hynson, Westcott & Dunning) Mercapocain ½ gr., soligenin 1 gr./toz. Vial 12s.
Use: Antiseptic lozenge.
- THARB.** (Cole) Phenobarbital Sod. ¼ gr., theobromine & Sod. salicylate 5 gr./Tab. Bot. 100s, 500s, 1000s
Use: Angina pectoris.
- THASMATAL.** (Miller) Phenobarbital ¼ gr., theophylline 1½ gr., ephedrine HCl ½ gr./tab. Bot. 100s, 500s, 1000s.
Use: Sedative with diuretic.
- THAVERINE.** (Pitman-Moore) Eltheverine ½ gr. & theophylline Sod. glycinate 3 gr./Tab. Bot. 100s.
Use: Coronary insufficiency.
- THEACITIN.** (Rorer) Theophylline & Sod. acetate, Tab. (2½ gr.) plain or E.C., Bot. 100s.
Use: (1-2 q.i.d.) Diuretic.
- W/Phenobarbital.** (Rorer) Theophylline & Sod. acetate 2½ gr., phenobarbital ½ gr./plain or E.C. Tab. Bot. 100s.
Use: (1-2 i.i.d.) Diuretic with sedative.
- THEAMIN.** Monoethanolamine salt of theophylline. See: Monotheamin, Prep. (Lilly)
- W/Amytal.** See: Monotheamin & Amytal, Cap. (Lilly).
- THEBA-PHEN.** (Meyer & Rosser) Theobromine 5 gr., phenobarbital ½ gr./Yab. Bot. 100s
Use: (1 i.i.d.) Diuretic.
- THELANDROL ORAL.** (Parke, Davis) Menagen (estrogens) 10,000 I.U., methyltestosterone 10 mg./cap. Bot. 100s. Also available in half strength.
Use: Climacteric symptoms in either sex.
- THELIN, U.S.P., N.N.O. 195B.** (Parke, Davis) Estrone, U.S.P. Each mg. of Theelin contains 10,000 I.U. of estrone
Aq. Susp. Amp. 1 cc. (1 mg.) 10,000 u.
(2 mg.) 20,000 u. (5 mg.) 50,000 u. Vial

10 cc; (2 mg) 20 000 u/cc Boxes 10s;
Steril Vial 5 cc; (5 mg) 50 000 u/cc
Amp In oil 1 cc; (2 mg) 2000 u, (0.5
mg) 5000 u, (1 mg) 10 000 u Boxes 10s;
Steril Vial 10 cc; (1 mg) 10 000 u
Supp. (vaginal); (0.2 mg) 2000 u glycerol
gelatin) Box 6s & 50s

Use: Estrogen deficiency
W/Testosterone propionate See: Theolandrol
Vial (Parke Davis)

THELIN R-P. (Parke Davis) Theelin, natu-
rally-occurring water soluble & water in-
soluble Vial (Theelin 2 mg, Pot theelin
sulfate 1 mg/cc) 10 cc

Use: Rapid & prolonged estrogen therapy
THEIOL, NMD 1958 (Parke Davis) Estrsol
Capsules (0.24 mg or 2400 I.U. estradiol) Bot
20s; 100s; 250s See: Estrisol, Cap (Abbott)

Use: Estrogen therapy
THIETHYPERAZINE MALEATE (Sandoz)

See: Tarenan
THEKOBARB (Lloyd) Atropine methylurate
2 mg, phenobarbital 1/4 gr, thiomine HCl
5 mg/Tab Bot 100s; 500s; 1000s

Use: (1 lid) Autonomic dysfunction
THELESTRIN, NND 1958 (Cornrick) Es-
trone, USP Aq Susp (10 000, 20 000 &
50 000 u/cc) Vial 1 cc, 10 cc in sesame
oil (10 000 u/cc) Amp 1 cc; Vial 10 cc
25 cc, Hypoglossals (10 000 u) Bot 50s
100s, 1000s

Use: I.M., orally; estrogen therapy
THELESTRONE (Cornrick) Estrogens mixed
Hypoglossals (10 000 u) Bot 50s; 100s
1000s

Use: Estrogen therapy
***THIMALIDINE** See: Sondostene (Sandoz)

THEMFADIL-A P C. (Winthrop Labs) Then-
fadil HCl 15 mg, acetylsalicylic acid D3
Gm, phenacetin 0.2 Gm, caffeine 15 mg/
tab Bot 50s

Use: Common cold
THEMFADIL HYDROCHLORIDE. (Winthrop
Labs) Thenyldiamine HCl, USP, Diethylan-
diamine 2 [(2-dimethylaminoethyl) 3 phenyl-
amino] pyridine Tab (15 mg) Bot 100s

Use: (1-6 daily) Antihistaminic, ant allergic
agent

W/Neo-synephrine HCl, Zephiran Chlor
See: NT2 Nasal Soln (Winthrop Labs)

THEMFADIL-SA. (Winthrop Labs) Two layers
containing Thefadil HCl 1/4 gr each Bot
100s, 1000s

Use: Antihistaminic, delayed
THEMFADIL-S P C. (Winthrop Labs) Then-
fadil 15 mg., salicylamide 300 mg., phen-
acetin 200 mg., caffeine 15 mg./tab Bot
50s.

Use: 1 lid's colds.

* Generic name recognized by H.F., U.S.P. or A.M.A.

***THEMIUM CLOSYLATE** (Burroughs Well-
come) N,N dimethyl H 2 phenoxy ethyl N 2
thenylammonium p chlorobenzene sulfonate
Use: Canine hook worm

See: Bancaris (Burroughs Wellcome)

THENYLDIAMINE CHLORIDE See: Thentadil
HCl (Winthrop Labs)

***THENYLDIAMINE HYDROCHLORIDE, N F**
Tab N F 2 [(2 dimethylaminoethyl) 3 phenyl-
amino] pyridine HCl thenyldiamine chlo-
ride Diethylandiamine See: Thentadil (Win-
throp Labs)

W/Neo-synephrine HCl, N-acetyl p amino
phenol, caffeine See: Neo-Synephrine Cam-
pound Cold Tablets (Winthrop Labs)

W/Phenylephrine HCl isoetharine methane
sulfonate freon propellant See: Branko
meter (Brean)

W/Phenylephrine HCl hydrocortisone, neo-
mycin sulf., polymyxin B sulf See: Biocly-
nephrene (Winthrop Labs)

THENYLENE-A P C CAPSULES (Abbott)
Thenylene HCl 25 mg, acetylsalicylic acid
230 mg, phenacetin 150 mg, caffeine 30
mg/Cap Bot 100s, 500s; 1000s

Use: Antihistaminic A P C

THENYLENE & DESOXYN TABLETS (Ab-
bott) Thenylene HCl 50 mg, desoxyne HCl
2.5 mg./Tab Bot 100s; 500s; 1000s

Use: To provide sympathomimetic action
with antihistamine

THENYLENE EXPECTORANT (Abbott) Ce-
daine phosphate 1 gr thenylene fumarate
1 1/2 gr, Sod citrate 25 gr, Ipecac syr 30
min, menthol & aromatics qs in a ben-
zoated syr/fl oz Bot 1 pl, 1 gal

Use: (1 tsp q 3-4 h) Coughs
THENYLENE HCl, NND 1960 (Abbott)
Methapyrilene HCl Cream (2%) Tube 1 oz
Jar 1 lb, 5 lb, Tab (25 mg, 50 mg, 100
mg) Bot 100s, 500s; 1000s

Use: Antihistaminic
W/Ephedrine Hydrochloride See: Thenyfred
Tab (Abbott)

THENYLFRED (Abbott) Thenylene HCl 50
mg, ephedrine HCl 25 mg/Tab Bot 100s
1000s

Use: Antihistamine with central nervous sys-
tem stimulant

THENYPYRAMINE, NND 1962 See
Methapyrilene Hydrochloride

THEO-BARB (Haack) Theobramine 5 gr-
phenobarbital 1/2 gr, 1/4 gr or 1/2 gr/Tab.
Bot 100s, 1000s

Use: Diuretic & sedative
W/Pot iodide (Haack) Theobramine 4 gr-
phenobarbital 1/4 gr, Pot iodide 2 1/2 gr/
tab Bot 100s; 1000s

Use: Diuretic & sedative

- THEOBARB.** (VanPelt & Brown) Maenitol hexonitrate $\frac{1}{2}$ gr., theobromine 5 gr., phenobarbital $\frac{1}{2}$ gr./tab. Also available w/phenobarbital $\frac{1}{4}$ gr./tab. Bot. 50s, 500s. Use: (1 q. 3-4 h.) Vasodilator with sedative & diuretic.
- W/Ephedrine. (VanPelt & Brown) Ephedrine sulfate, theobromine & phenobarbital. Tab., Bot. 50s, 500s.
- W/Rutin. (VanPelt & Brown) Theobromine 5 gr., phenobarbital $\frac{1}{2}$ gr., rutin $\frac{1}{2}$ gr./tab. Bot. 50s.
- Use: (1-2 i.l.d.) Vasodilator.
- THEOBARB-R.** (VanPelt & Brown) Theobromine 324 mg., phenobarbital 10 mg., reserpine 0.1 mg./tab. Bot. 50s, 500s. Use: Tranquilizing agent w/sedative.
- THEOBARRIDE.** (Cole) Phenobarbital $\frac{1}{4}$ gr., theobromine 5 gr., Pot. Iodide $1\frac{1}{2}$ gr./Tab. Bot. 100s. Use: Hypertension.
- ***THEOBROMINE, N.F. 3,7-Dimethylxanthine.** Tab. (5 gr.).
- W/Carbromol & Cal-ureo-Iodide. See: Krozonine, Tab. (Blue Line)
- W/Ephedrine sulf., phenylpyramine HCl. See: Asmi-Hisl, Tab. (Cele)
- W/Ethylenediamine 2 dihydroiodide. See: Xanlamne, Tab. (Pilman-Moore)
- W/Luminal & rauwolfia serpentina. See: Theominal R.S., Tab. (Winthrop Labs.)
- W/Mebarol, Cal. Iodide, triethanolamine. See: Jacopral, Tab. (Winthrop Labs.)
- W/Phenylpropanolamine, d-Amphetamine sulfate. See: Apsed, Tab. (First Texas Chem.)
- Theobromine with Phenobarbital Combination on p. 726.
- THEOBROMINE CALCIUM.** See: Purital, Tab. (A. J. Parker)
- W/Mersalyl Acid. See: Mercurital, Tab. (A. J. Parker)
- W/Reterpine. See: Purital R.S., Tab. (A. J. Parker)
- THEOBROMINE CALCIUM GLUCONATE.** Bites—Tab., Bot. 100s, 1000s. Also available w/phenobarbital. Grant—Tab., Bot. 100s, 500s, 1000s. See: Calpurate, Prep. (Mallinck)
- W/Phenobarbital. See: Calpurate w/Phenobarbital, Tab. (Mallinck)
- Glarb, Tab. (Cole Chem. Co.)
- ***THEOBROMINE CALCIUM SALICYLATE,** N.F., Tab., N.F., N.N.R. 1948. Bites—Tab., Bot. 100s, 1000s. Normand—Tab., Bot. 100s, 1000s. See: Theocalcin, Tab. N.N.R. (Knoll)
- ***THEOBROMINE MAGNESIUM OLEATE.** See: Athemal, Tab. (Meyer)
- ***THEOBROMINE SODIUM ACETATE, N.F., N.N.R. 1948,** Cap., N.F., Tab., N.F., Agurin, N.N.R. 1928. See: Thesodate, Tab. (Brewer) W/Dicalcium phosphate & phenobarbital. See: Barbltheo, Cap. & Tab. (C. D. Smith)
- W/Ephedrine ethylenediamine, caffeine, and acetophenetidin. See: Kradon, Tab. (Field Labs.)
- W/Phenobarbital. See: Thesodate w/Phenobarbital, Tab. (Brewer).
- Theosophen w/Phenobarbital, Tab. (Starr).
- ***THEOBROMINE SODIUM SALICYLATE,** N.F. See: Diareks (Knoll) Neathylline, Tab. (Paul Money)
- W/Col. lactate. See: Diurbitol, Tab. (Grant)
- W/Phenobarbital Sod. See: Tharb, Tab. (Cole Chem. Co.)
- THEOBROMITAL,** (Warren-Teed) Theobromine D.3 Gm., phenobarbital 30 mg./Tab. Bot. 100s. Use: Diuretic and myocardial stimulant
- THEOCALBITAL.** (Forrest) Cal. lactate 1 gr., phenobarbital Sod. $\frac{1}{4}$ gr., theobromine $3\frac{1}{2}$ gr./Tab. Bot. 35s, 100s, 500s. Use: (1-2 i.l.d.) Diuretic with sedative.
- THEOCALCIN,** N.N.R. 1948. (Knoll) Theobromine-Cal. salicylate basic. Pow., Bot. 1 oz., 8 oz. Tab. ($7\frac{1}{2}$ gr.) Bot. 100s, 500s, 1000s. Also available w/phenobarbital. Also available w/Pot. Iodide. Use: Myocardial stimulant & diuretic.
- THEOCARDONE.** (Hoack) Theobromine 5 gr., phenobarbital $\frac{1}{4}$ gr./Tab. Bot. 100s, 1000s. Use: (1 i.l.d.) Diuretic with sedative.
- THEOCIN SOLUBLE,** N.N.R. 1950. (Winthrop Labs.) Theophylline & sod. acetate. Tab. ($1\frac{1}{2}$ gr. & $2\frac{1}{2}$ gr.) Bot. 25s. Use: Coronary vasodilator.
- THEODIGITAL** (Drug Prods) Phenobarbital $\frac{1}{2}$ gr., theobromine $1\frac{1}{2}$ gr., Cal. salicylate $1\frac{1}{2}$ gr., digitalis leaf pow. $\frac{1}{4}$ gr./Cap Fulvicoids, plain or E.C., Bot. 100s, 1000s. Use: Vasodilator.
- THEOFED.** (Suliff & Case) Phenobarbital 8 mg., theophylline 130 mg., ephedrine HCl 24 mg., powdered extract belladonna 5 mg., pyrimine maleate 25 mg., potassium iodide 65 mg., magnesium trisilicate 130 mg./Tab. Bot. 100s, 1000s. Use: Diuretic, antispasmodic, sedative, bronchodilator and antihistamine.
- THEOFED-PEDIATRIC.** (Suliff & Case) Phenobarbital R mg., theophylline 65 mg., ephedrine HCl 11 mg., belladonna leaves 5 mg., pyrimine maleate 12.5 mg., potas-

| Theobromine | Phenobarbital | Others | Manufacturers | Brand |
|---------------------------|----------------|--|-----------------------|--|
| 5 gr | 1/8 gr | | Hoack | Theo Barb, Tab |
| 5 gr | 1/4 gr | | Hoack | Theocardone Tab, |
| 5 gr. | 1/4 gr | | Winthrop Labs | Theorinal, Tab |
| 5 gr | 1/4 gr | | Robins | Theorate Tab |
| 5 gr | 1/2 gr | | Smith Dorsey | Theotal, Tab |
| 5 gr | 1/2 gr | | Meyer & Rossen | Theba Phen., Tab |
| 5 gr | 1/2 gr | | Haskell | Zanthinol, Tab. |
| 325 mg | 32.5 mg | | Sulchiff & Case | Theophol, Tab |
| (Sod acetate), 3 1/2 gr. | 1/4 gr. | | Starr | Theophen w/Phenobarbital, , Tab |
| 3 1/2 gr. | 1/4 gr | Col gluconate 1 1/2 gr. | Cole Chem Co | Globb, Tab |
| 3 1/2 gr | (Sod), 1/4 gr | Col lactate 1 gr. | Forrest | Theocarbitol, Tab |
| (Sod acetate), 3 gr | 1/4 gr | Droctrum phosphate 2 1/2 gr | Smith Miller & Peitch | Borbtheo, Cap & Tab. |
| (Col salicylate), 3 gr. | 1/4 gr | Digitalis lvs 1/4 gr | Drug Products | Theadigital, Cap |
| 5 gr | 1/2 gr | Ethyl diiodo-brassidate 1 gr | Haskell | Iozanthol, Tab |
| 2 1/2 gr. | 1/4 gr | Ethylenediamine dihydroiodide 1/2 gr | Pitman Moore | Xantophen, Tab. |
| (Cal salicylate), 5 gr. | 1/4 gr | Ethylenediamine dihydroiodide 1/2 gr | Pitman Moore | Xancylo's, Tab |
| 5 gr. | 1/4 or 1/2 gr | Mannitol hexonitrate 1/2 gr | Van Pelt & Brown | Theoborb, Tab Also w/Ephed- rine alto w/Kutin |
| 2 1/2 gr | 1/2 gr. | Mannitol hexonitrate 1/4 gr | Smith Dorsey | Teatine, Tab |
| 2 1/2 gr. | 1/2 gr | Mannitol hexonitrate 1/2 gr., rutin 1/2 gr., Vit C 1/2 gr | Smith Dorsey | Teatine R C, Tab |
| 2 1/2 gr | 1/4 gr. | Pot iodide 2 1/2 gr | U S Standard | Brodital, Tab |
| 4 gr | 1/4 gr | Pot iodide 2 1/2 gr | Hoack | Theo Barb w/Pot Iodido, Tab |
| 5 gr | 1/4 gr. | Pot iodide 1 1/2 gr | Cole Chem Co | Theobarbide Tab |
| 325 mg | 16 mg | Pot iodide 160 mg | Sutcliff & Case | Theophal w/Pot Iodide, Tab. |
| (Sod acetate), 5 gr. | 1/4 gr. | Pot iodide 2 gr | Starr | Theasaphen w/Pot iodide |
| 150 mg | 15 mg. | Pot iodide 120 mg., crotaegus tr. 0.3 mg | Central | Theonatal, Tab |
| 324 mg | 10 mg | Reserpine 0.1 mg. | Van Pelt & Brown | Theoborb R, Tab |
| (Cal. salicylate), 160 mg | (Sod), 1/4 gr. | Sod nitrate 64 mg | Drug Products | Theoviscum, Tab |
| 4 gr. | 1/4 gr | Watermelon seed ext 2 gr. | Table Rock | Carhol, Cap |

- sium iodide 65 mg., magnesium trisilicate 130 mg./Tab. Bot. 100s, 1000s
 Use: Diuretic, antispasmodic, bronchodilator, sedative and antihistaminic
- THEOFENEDRINE.** (Premo) Phenobarbital $\frac{1}{2}$ gr., theophylline 2 gr., ephedrine HCl $\frac{3}{4}$ gr./Tab. Box 30s, 120s, 1000s.
 Use: Paroxysms of bronchial asthma.
- THEOGEN.** (Sig-Tabs) Conjugated estrogens 125 mg./Tab. Bot. 50s
 Use: Menopausal syndrome.
- THEOGLYCINATE, N N D.** 1962. (Brayten-Theophylline-Sod. glycinate.
 Sol. Aero; 3 Gm./30 cc. Sol. 6 fl. oz.
 Tab. (0.325 Gm.) Bot 100s
 Tab. (0.325 Gm. theophylline-Sod. glycinate 16 mg., phenobarbital) Bot. 100s.
 Tab. (0.325 Gm. theophylline-Sod. glycinate 16 mg., phenobarbital 25 mg., rafephedrine HCl) Bot. 100s.
 Supp. (0.78 Gm.) Box 12s.
 Syr. (0.13 Gm./5 cc.) Bot. 8 oz.
 Use: Theophylline therapy.
- THEOHIST.** (Massengill) Methapyrilene HCl $\frac{3}{4}$ gr., ephedrine sulfate 0.3 gr., theophylline $\frac{1}{2}$ gr./Tab. Bot. 1000s
 Use: (1 l.l.d.) Antihistamine with sympathomimetic action.
- THEOLAPHIN.** (Massengill) Theobramine & Sod. salicylate 3 gr., Cal. lactate $1\frac{1}{2}$ gr., phenobarbital $\frac{1}{4}$ gr./Tab. Bot. 100s, 500s, 1000s.
 Use: 1 l.l.d.; vasoconstrictor, sedation.
- THEOLAXIN Ph. UNODAY 748.** (Paul Maney) Phenobarbital 100 mg./delayed action cap. Bot. 100s.
 Use: 1 daily; sedative.
- THEOLIPO.** (Babs) Theophylline Sod. glycinate 120 mg., choline bicarbonate 143 mg., dl-methionine 130 mg., inositol 65 mg., phenobarbital 15 mg./Tab. Bot. 36s, 100s.
 Use: Various degenerative cardiovascular disorders.
- THEOMERSYL.** (Central Pharm.) Theophylline Sod. glycinate.
 Use: Diuretic.
- THEOMINAL.** (Winthrop Labs.) Theobramine 5 gr., & luminal $\frac{1}{2}$ gr./tab. Bot. 100s, 500s.
 Use: Coronary disease.
- THEOMINAL M.** (Winthrop Labs.) Theobramine 5 gr., luminal $\frac{1}{4}$ gr./tab. Bot. 100s, 500s.
- THEOMINAL R.S.** (Winthrop Labs.) Theobramine 5 gr., luminal $1\frac{1}{6}$ gr., efheraxylan fraction 1.5 mg./tab. Bot. 100s.
 Use: Hypertension.
- THEOMINE.** (Smith-Dorsey) Theobramine 5 gr., Pot. iodide $2\frac{1}{2}$ gr., phenobarbital $\frac{1}{4}$ gr./Tab. Bot. 100s, 1000s.
 Use: Cardiac edema.
- THEONATAL.** (Central) Phenobarbital 15 mg., theobramine 0.15 Gm., potassium iodide 0.12 Gm., calceogus fr. 0.3 ml./Tab. Bot. 100s, 500s, 1000s
 Use: Afferative, diuretic, sedative
- THEO-NITRAL.** (Babs) Theophylline Sod. glycinate 130 mg., mannitol hexanitrate 15 mg., phenobarbital 15 mg./Tab. Bot. 100s.
 Use: Hypertension.
- THEONYLDIAMINE HCl** W/Phenylephrine HCl, pot. guaiacol sulfonate, dihydrocodeinone bitartrate. See: Dicodrine Syrup, Syr. (Bran)
- THEO-ORGANIDIN.** (Wampole) Theophylline 120 mg., iodopropylidene glycerol 30 mg./15 cc. Sol. 16 oz.
 Use: Treatment of asthma
- THEOPHAL.** (Suliff & Case) Theobramine 325 mg., phenobarbital 32.5 mg./Tab. Bot. 100s, 1000s
 Use: Bronchodilator and sedative
- THEOPHAL WITH POTASSIUM IODIDE.** (Suliff & Case) Theobramine 325 mg., potassium iodide 160 mg., phenobarbital 16 mg./Tab. Bot. 100s, 1000s
 Use: Bronchodilator and sedative.
- *THEOPHYLLINE, N F, N.N.R. 1949;** Tab. N.F. 1,3-Dimethylxanthine.
 See: Elizaphyllin, Sol. (Sherman)
 Theocin, Tab. (Winthrop Labs.)
 Use: Coronary vasodilator & diuretic.
 W/Amobarbital, ephedrine, alcohol See: Asmafield, Liq. (Canfield)
 W/Ephedrine sulfate & methapyrilene HCl See: Theohist, Tab. (Massengill).
 W/Ephedrine HCl, butobarbital See: Numa Dura-Tabs. (Wynn)
 W/Ephedrine HCl, phenobarbital See: Parosmal, Tabs. (Premo)
 Tedrol SA, Tabs. (Werner-Chilcatt)
 W/Ephedrine hydrochloride, phenobarbital, glyceryl guaiacolate. See: Thephecac, Tab. (Marsh-Emory Labs.)
 W/Ethaverine HCl, Isopropylarterenal, butobarbital. See: Benedrin, Tab. (Bene)
 W/Glyceril guaiacolate. See: Oubran (Mead Johnson)
 W/Hydroxyzine HCl, ephedrine sulfate. See: Marax. (Roerig)
 W/d-isoephedrine HCl, phenyltalaxamine. See: Azeac, Tabs. (Arnold-Stone)
 W/Isoprotrenol, benzylephedrine HCl, phenobarbital. See: Isuprel-Franol. (Winthrop)
 W/Isoprotrenol, ephedrine, & secobarbital. See: Aluted, Tab. (Hart)
 W/Luminal, ephedrine sulfate, Isuprel, po-

- lassium iodide alcohol See: Isuprel Com
pound (Winthrop)
- W/Mercury in organic form.
- See: Cumetilin, Amp (Endo)
- Diuremin, Amp (C D Smith)
- Diursal, Amp (Harvey)
- Mercuhydrin, Amp (Lakeside)
- Mercuzanthin, Amp (Campbell)
- Mersolyn, Amp (Kirk)
- Merthyline, Amp (Drug Prods)
- Neamersyl, Vial (Central)
- Salyrgan w/Theoph (Winthrop Labs)
- W/Mersalyl See Merthiphyl (New Mexico
Pharm Co)
- W/Mersalyl, procaine barate See: Mercu
pracyl, Inj (Barrows)
- W/Organidin, alcohol See: Theo Organidin
(Wampale)
- W/Prednisone, Ephedrine HCl, phenobarbi-
tal See: Deltasmyl (Roussel)
- W/Reserpine, choline bitartrate See: Thera-
cor, Tab (C D Smith)
- W/Reserpine mannitol hexanitrate, rutin &
ascorbic acid See: Neo Semhyten, Cap
(Massengill)
- Theophylline with Phenobarbital Com-
binations on pp 729 and 730
- THEOPHYLLINE AMINOISOBUTANOL** (Bu-
topyllamine) Theophylline w/2 amino 2
methyl-1-propanol See: Butophyllamine,
Butoid, Supp (Merrell)
- W/Phenobarbital Nelthamine HCl See:
Nelthaphyl, Cap (Merrell)
- THEOPHYLLINE BETAINE** W/Mersalyl See:
Neamersyl, Vial (Central)
- THEOPHYLLINE, 8-BROMO, AMINOISO-
BUTANOL** See Pomabrom
- THEOPHYLLINE, 8-BROMO, PYRILAMINE.**
See: Pyroabram
- THEOPHYLLINE CALCIUM SALICYLATE.**
See: Phyllicin, Tab (Knall)
- THEOPHYLLINE, 8-CHLORO, DIPHENY-
DRAMINE.** Dimenhydrinate, U.S.P. See
Dramamine (Searle)
- THEOPHYLLINE CHOLINE SALT** See: Chole-
dyl, Tab (Nepera)
- THEOPHYLLINE ETHYLENEDIAMINE, N N R**
1954 See: Aminaphylline, USP (Various
Mfr)
- THEOPHYLLINE MAGNESIUM.** See: Magne-
phylline Tab (Sherman)
- THEOPHYLLINE AND MERSALYL SODIUM,**
N N D 1959
- ***THEOPHYLLINE-METHYLGLUCAMINE,**
N N D 1962 W/Nembutal, Supp See Glu-
cophylline & Nembutal (Abbott)
- THEOPHYLLINE MONOETHANOLAMINE**
See Clysmathone Rectal Unit
Monatheamin (Lilly)
- *Generic name recognized by N F, U S P or A M A
- THEOPHYLLINE PIPERAZINE SALT, See:**
Piperophyllin, Tab (M. H. Smith)
- ***THEOPHYLLINE SODIUM ACETATE, N F,**
N N R 1950; Tab, N F.
See: Theacilin, Tab (Rorer)
- Theacin Safuble, Tab (Winthrop Labs)
- W/Ephedrine sulfate & Sod phenobarbital
See: Luosmin, Cap (Brewer)
- W/Phenobarbital See: Theacilin w/Pheno-
barbital, Plain or E.C. Tab (Rorer)
- ***THEOPHYLLINE SODIUM GLYCINATE,**
N F, N N D 1962; Tabs, N F.
See: Cleophyl, Tab, N N R (Ascher)
- Darsaphyllin, Elixir, Supp, Syr, Tab-
N N R (Smith Darsay)
- Glylene, Tab (Flinn, Eaton)
- Glynozox, Syr, Tab, N N R (First Texas)
- Glytheonate, Supp, Syr, Tab, N N R
(Patch).
- Pemophyllin, Tab (Pitman Moore)
- Synophylate, Aerosol, Syr, Tab, N N R
(Central)
- Theoglycinate, Syr, Supp, Tab, N N R
(Brayten)
- W/Choline bitartrate, dl methionine, Inasi-
tol, phenobarbital See: Theo Lipa, Tab
(Bobst)
- W/Cocaine phosphate, pyranisamine male-
ate & ephedrine HCl See: Gyralene, Syr
(Patch)
- W/Ephedrine sulfate, phenobarbital See
Glynozox EP, Tabs (First Texas)
- W/Ethoverine See: Thaverine, Tab (Pitman
Moore)
- W/Mannitol hexanitrate, phenobarbital See:
Thno Nitrol, Tab (Bobst)
- W/Phenobarbital
See Glytheonate w/Phenobarbital, Tab
(Patch)
- Synophylate w/Phenobarbital, Tab (Cen-
tral)
- W/Phenobarbital & mannitol hexanitrate
See: Syntrate, Tab (Central)
- Nitrophyl, Tab (Ascher)
- W/Phenobarbital & racephedrine HCl See
Glytheonate w/Phenobarbital and Rac-
ephedrine, Tab (Patch)
- W/Phenobarbital, reserpine, & dihydroxy
aluminum aminoacetate See: Glytheonate
- W/Phenobarbital & Reserpine Tab (Patch)
- W/Phenobarbital & Rutin See: Glytheonate
w/Phenobarbital & Rutin, Tab (Patch)
- W/Racephedrine & phenobarbital See
Synphedal, Tab (Central)
- THEORATE.** (Robins) Theobramine 5 gr &
phenobarbital ¼ gr/Tab Bat 50s, 500s
Use: Mild sedation & diuretic
- THEOSOPHEN** (Starr) Theobramine Sod

Theophylline with Phenobarbital Combinations

| Theophylline | Phenobarbital | Others | Manufacturer | Brand |
|----------------------------|---------------|---|-----------------------|---|
| (Isopropanolamine), 1½ gr. | ¼ gr. | | National Drug | Theopropanal w/Phenobarbital, Tab. |
| (Sod. acetate), 2½ gr. | ¼ gr. | | Rorer | Theocite w/Phenobarbital, Tab. plate & E.C. |
| (Sod. glycinate), 5 gr. | ¼ gr. | | Central Pharm. Co. | Synaphylate w/Phenobarbital, Tab. |
| 1½ gr. | ¼ gr. | Ammonium chloride 5 gr. | Mallard | Borphyllin, Tab. |
| 2 gr. | 2 gr. | Bephedin ½ gr. | Winthrop Labs. | Fractal, Tab. |
| (Sod. glycinate), 120 mg. | ¼ gr. | Choline tartrate, 143 mg., di-meth- ioctee 120 mg., inositol 65 mg. | Babst | Theo-Lipa, Tab. |
| 2 gr. | ½ gr. | Ephedrine ¾ gr. | Chicago Pharm. Co. | Barbaphine, Tab. |
| 1 gr. | ½ gr. | Ephedrine HCl ¼ gr. | Medicane | MediTab, Tab. |
| 1½ gr. | ¾ gr. | Ephedrine HCl ¾ gr. | Enda | Teprin, Tab. |
| 1½ gr. | ¾ gr. | Ephedrine HCl ¾ gr. | Mittler | Thasmatol, Tab. |
| 120 mg. | 6 mg. | Ephedrine HCl 24 mg., pow. ext. belladonna 5 mg., pyrilamine maleate 25 mg., pot. iodide 65 mg., mag. trisulfate 120 mg. | Sutliff & Case | Theated, Tab. |
| 65 mg. | 8 mg. | Ephedrine HCl 11 mg., belladonna leaves 5 mg., pyrilamine maleate 12.5 mg., pot. iodide 65 mg., mag- nesium trisulfate 120 mg. | Sutliff & Case | Theated-Pediatric, Tab. |
| 15 mg. | 4 mg. | Ephedrine sulfate 12 mg., chlorphe- iramine maleate 1 mg., glyceryl sebacolate 50 mg. | Breco | Bronkotab/5 cc. Eliz. |
| 1¼ gr. | ¼ gr. | Ephedrine HCl ¾ gr., Sod. iodide 1 gr. | Testagar | Tepadrine, Tab. |
| 2 gr. | (Sod.) ½ gr. | Ephedrine sulfate ½ gr. | Cole | Amnyl, Tab. |
| 2 gr. | (Sod.) ½ gr. | Ephedrine sulfate ½ gr. | Cole Chem. Co. | Asmynl, Tab. |
| (Sod. glycinate), 3 gr. | (Sod.) ½ gr. | Ephedrine sulfate ½ gr. | Brewer | Luasmin, Cap. |
| 1½ gr. | ½ gr. | Ephedrine sulfate 3/16 gr., alludrine 5 mg. | Leeming | Nephenafin, Pediatric |
| 2 gr. | ½ gr. | Ephedrine sulfate ¾ gr., Isopropyl arsenol 10 mg. | Leeming | Nephenate, Tab. |
| (Sod. glycinate), 162 mg. | ¼ gr. | Mannitol hexeacetate 32.4 mg. | Ascher | Nitraphyl, Tab. |

Theophylline with Phenobarbital Combinations

| Theophylline | Phenobarbital | Others | Manufacturer | Brand |
|---------------------------|---------------|--|--------------------|-------------------------------|
| (Sed Glycinate) 2 1/2 gr. | 1/4 gr. | Monsel hexestrate 1/2 gr. | Central Pharm. Co | Syntrate, Tab. |
| (Sed Glycinate) 130 mg | 1/4 gr. | Monsel hexestrate 15 mg. | Babst | Theo Nitral, Tab |
| 1.5 gr | 1/4 gr. | Monsel hexestrate 37 mg. rufin 20 mg | Plassner | Ku Nitral w/Theophylline, Tab |
| 2 1/2 gr. | 1/4 gr. | Neo-synephrine HCl 1/2 gr. | Winthrop Labs | Adnephrine, Cap |
| 2 gr | 1/2 gr | Prednicolose 0.5 mg., ephedrine sulf 1/2 gr | Vitamin | Prednisophine, Tab |
| (Sed Glycinate), 5 gr. | 1/4 gr | Racephedrine 3/8 gr | Central Pharm. Co. | Synophedal, Tab |

acetate Tab (3 3/4 gr & 7 1/2 gr) Bot 100s, 500s, 1000s

Use Diuretic

W/Phenobarbital (Starr) Theobromine Sod acetate 3 3/4 gr, phenobarbital 1/4 gr or 1/2 gr/Tab Bot 100s, 1000s

Use (1 2 q 3 h) Diuretic & sedative

W/Pot iodide & phenobarbital (Starr) Theobromine Sod acetate 5 gr, phenobarbital 1/4 gr, Pot iodide 2 gr/Tab Bot 100s, 500s, 1000s

Use (1 2 1 s d) Diuretic with sedative

THEOTAL Na 1. (Smith Dorsey) Theobromine 5 gr, phenobarbital 1/2 gr/Tab Bot 100s, 1000s

Use (1 2 1 s d) Diuretic & sedative

THEOVISCUM. (Drug Prods) Sod phenobarbital 16 mg, theobromine 80 mg, Col salicylate 80 mg, Sod nitrite 64 mg, Yescum album 16 mg, Crataegus oxyacantha 16 mg/Cap E C Bot, 100s, 1000s

Use: (1 q 3 4 h) Vasodilator, diuretic & myocardial stimulant

THEPHECON. (Morch Emory Labs) Theophylline 4 gr, ephedrine hydrochloride 1/4 gr, phenobarbital 1/2 gr, glyceryl guaiacolate 200 mg/Tab Bot 100s

Use: Bronchial dilatation

THEPHORIN (Roche) Phenindamine tartrate, 2 methyl 9 phenyl tetrahydro 1-pyridindene hydrogen tartrate

1st: (5%) Bot 4 oz, 1 pt; (10%) Bot, 2 oz, 1 pt

Oint: (5%) Tube 1 1/2 oz, Jar 1 lb

Pow: (5 & 25 Gm) Vials

Syr: (10 mg /4 cc) Bot 1 pt, 1 gal

Tab: (10 & 25 mg) Bot 100s, 500s, 1000s

Use: (25 mg 1 d) Antihistaminic

W/Madribon, N acetyl p aminophenol, col feine See: Madricidin, Cap (Roche)

THEPHORIN-AC. (Roche) Thephorin 10 mg., aspirin 2 1/2 gr, acetaphenetidin 2 1/2 gr, caffeine 1/4 gr/Tab Bot 12s, 25s, 100s, 1000s

Use: (1 2 1 d) Common cold

THEPHORIN EXPECTORANT. (Roche) Thephorin 10 mg, codeine phosphate 4 mg, papaverine 4 mg, w/ammonium chloride & chloroform/4 cc. Bot 1 pt, 1 gal

Use: (1 2 tsp 1 d) Coughs

THEPTINE. (Smith, Kline & French) Dexamphetamine sulfate 2.5 mg, thiamine HCl 5 mg, riboflavin 0.45 mg, niacin 6.7 mg/dr (Liale, Bot 12 fl oz

Use: (1 tsp 1 d, p.c) Obesity

THERAZEN. (Bentax Pharm) Vit A 25,000 u. D 1000 u., B-1 10 mg, B 2 5 mg., C 150 mg., nicotinamide 150 mg/cap Bot 100s, 1000s

Use: Vitamin therapy

- THERABEX**, (Bryant) Vit. A 33,000 u., D 1500 u., B-1 20 mg., B-2 10 mg., B-6 1 mg., B-12 10 mcg., C 150 mg., niacinamide 150 mg., folic acid 1 mg./Cap. Bot. 100s.
Use: Multivitamin therapy
- THERABILE**, (Ascher) Ferrated ox bile 200 mg., desoxycholic acid 25 mg., dl-methionine 100 mg., pancreatin 250 mg., homarine 100 mg., methylbromide 2.5 mg., aleoresin 100 mg., tocopherols 25 mg./Cap. Bot. 100s, 300s.
Use: Gall bladder disease.
- THERACEBIN**, (Lilly) Vit. A 25,000 u., D 1500 u., B-12 10 mcg., thiamine chloride 15 mg., riboflavin 10 mg., pyridoxine 3 mg., niacinamide 150 mg., folic acid 0.3 mg., ascorbic acid 150 mg., Cal. pantothenate 20 mg., tocopherols 25 mg./Cap. Gelseals, Bot. 30s, 100s, 300s.
Use: Multiple vitamin deficiencies.
- THERA-COMBEX**, (Parke, Davis) Vit. B-1 25 mg., B-2 15 mg., C 150 mg., B-6 1 mg., niacinamide 100 mg., pantothenic acid 10 mg., folic acid 2.5 mg./Kapsak. Bot. 100s, 1000s.
Use: Vitamin B complex therapy.
- THERACOMBIN**, (Bryant Pharm.) Vit. A 6,000 u., D 1,200 u., B-1 5 mg., B-2 3 mg., B-6 1 mg., B-12 5 mcg., pantothenol 1 mg., niacinamide 25 mg., calcium 10 mg., phosphorus B mg., cobalt gluconate 0.2 mg./3 cc. Bot. B oz., 1 pt.
Use: Treatment of acute multiple avitaminosis.
- THERA-CONCEMIN**, (Merrell) Vit. A palmitate 25,000 u., D 1000 u., thiamine nitrate 5 mg., riboflavin 5 mg., niacinamide 150 mg., ascorbic acid 150 mg./Cap. Bot. 30s, 100s, 300s, 1000s.
- THERACOR**, (Smith, Miller & Patch) Anhydrous theophylline 100 mg., reserpine 0.05 mg., choline bitartrate 209 mg./Tab. E C. Bot. 100s.
Use: Hypertensive-cardiovascular ill.
- THERACORT**, (C & M Pharm) Hydracortisone 0.25%, colloidal sulfur 4%, salicylic acid 1.75%, resorcinol 1.5%. Lot. Bot. 1 oz.
Use: Treatment of acne
- THERA-DEIMAL**, (Boyle) Vit. A acetate 25,000 u., D 1500 u., C 150 mg., B-1 10 mg., B-2 5 mg., B-6 0.5 mg., B-12 5 mcg., K 2 mg., tocopherols 5 mg., niacinamide 100 mg., Cal. pantothenate 5 mg., folic acid 0.1 mg., ferrous sulfate 130 mg., dical. phos. 250 mg., iodine 15 mg., copper 1 mg., cobalt 0.15 mg., Mang. 3 mg., Mag. 10 mg., Pot. 5 mg., molybdenum 2 mg., zinc 1.5 mg., nickel 0.1 mg./cap. shaped tab. Bot. 30s, 100s.
Use: Therapeutic vitamin & mineral therapy.
- THERAFORMS**, (Horton & Converse) Vit. A 25,000 u., D 1000 u., C 150 mg., B-1 10 mg., B-2 10 mg., B-6 1 mg., B-12 5 mcg., folic acid 100 mcg., niacinamide 100 mg., B 2 mg., col. pantothenate 5 mg., Vit. E 5 IU. w/minerals/Tab. Bot. 30s, 100s, 250s, 500s, 1000s
Use: Vitamin-mineral deficiencies
- THERAFORTE**, (Hiss) Vit. C 300 mg., B-1 10 mg., B-2 10 mg., niacinamide 100 mg., B-6 2 mg., col. pantothenate 20 mg., B-12 4 mcg., folic acid 1.5 mg./Cap. Bot. 30s, 100s, 500s.
Use: Therapeutic stress formula.
- THERAFORTE SOLUTION**, (Hiss) Thiamine HCl 50 mg., riboflavin 5 mg., niacinamide 20 mg., pantothenic acid 5 mg., pyridoxine HCl 5 mg., ascorbic acid 50 mg./cc. 10 cc. Multiple dose vial, 2 & 5 cc. ompts.
- THERAFORTIS**, (United Lab) Vit. A 12,500 u., D 1000 u., B-1 5 mg., B 2 5 mg., B-6 1 mg., B-12 3 mcg., niacinamide 50 mg., pantothenic acid salt 10 mg., Vit. C 150 mg., folic acid 0.5 mg./Cap. Bot. 100s, 500s.
Use: (1 daily) Multivitamin therapy.
- THERAGBAN**, (Squibb) Vit. A 25,000 u., D 1000 u., thiamine mononitrate 10 mg., riboflavin 10 mg., niacinamide 100 mg., ascorbic acid 200 mg., B-6 5 mg., Cal. pantothenate 20 mg., B-12 5 mcg./Cap. Bot. 180s. Also available as liquid; 1 teaspoonful of liquid is equiv. to 1 capsule. Bot. 4 oz. Cap. Bot. 30s, 60s, 100s.
Use: Multiple avitaminosis.
- THERAGBAN JUNIOR**, (Squibb) Vit. A 5000 u., D 1000 u., B 1 5 mg., B 2 5 mg., niacinamide 30 mg., C 100 mg., B-6 2 mg., d-cal. pantothenate 3 mg., Vit. B-12 10 mcg./Cap. Bot. 30s, 100s.
Use: Vitamin.
- THERAGBAN-M**, (Squibb) Vit. A 25,000 u., D 1000 u., C 150 mg., B-1 10 mg., B-2 10 mg., niacinamide 100 mg., Cal. pantothenate 5 mg., Vit. B 6 0.5 mg., E 5 mg., B-12 5 mcg., folic acid 0.1 mg., menadione 2 mg., w/Cal., Phos., iodine, Iron, Pot., copper, Mang., Mag., cobalt & zinc/cap. Bot. 180s.
Use: Multivitamin therapy.
- THERALOWE**, (Lowe) Vit. A 25,000 U.S.P. u., D 1000 U.S.P. u., B-1 10 mg., B 2 10 mg., B 6 0.5 mg., B-12 5 mcg., folic acid 0.1 mg., niacinamide 100 mg., d calcium pantothenate 5 mg., C 150 mg., E 5 I.U., K 2 mg., Ca 105 mg., I 0.15 mg., Fe 15 mg., K 5 mg., Cu 1 mg., Mn 1 mg., Mg 6 mg., Zn 1.5 mg./Tab. Bot. 100s
Use: Vitamin and mineral deficiencies.

THERA-MINERAL. (Rowell) Vit A 25,000 u, D 1200 u, B 1 25 mg, B 2 10 mg, B-6 1 mg, B 12 6 mg, niacinamide 100 mg, Vit E 5 mg, C 150 mg, cobalt sulfate, copper sulfate, ferrous gluconate, Mang sulfate, Pot iodide Pot sulfate, Sod molybdate zinc sulfate, bone phosphates/Tab Bot 100s 300s

Use Nutritional deficiency

THERANATES, ESKAY'S (Smith, Kline & French) Strychnine glycerophosphate anhy 1 mg, Sod glycerophosphate 0.13 Gm, Cal glycerophosphate 0.13 Gm, phosphoric acid 75% 0.102 cc, Vit B 1 0.75 mg / 2 tsp Bot 12 R oz

Use Tonic

THERAPADS (Euller Pharm) Salicylic acid 1 1/2%, ethyl alcohol 50%, distilled water w/pads Jar 40s

Use Acne vulgaris

THERAPAS (Barnes Hind Pasno Co) Calcium salt of 4 benzamidosalicylic acid (B PAS)/Jars 1 lb

Use Treatment of tuberculosis

THERAPEUTIC COMBINATION (West Ward)

Vit A 25,000 u, D 1000 u, thiamine mononitrate 10 mg, Vit B 2 10 mg B-6 3 mg, B-12 5 mcg C 150 mg, niacinamide 100 mg, cal pantothenate 5 mg, folic acid 0.34 mg, Vit E 5 Int u, rutin 10 mg, Vit K 0.15 mg, viatin 10 mcg, Vit B-12 with intrinsic fact conc 2/15 u, feman bioflavono d complex 15 mg, choline bitartrate 17.3 mg inositol B 3 mg, methionine 4.4 mg, llysine HCl 100 mg, plus 11 minerals and trace-elements/Cap Bot 30s 100s, 1000s

Use: Therapeutic vitamin

THERAPEUTIC FORMULA (Milwell Pharma)

cal) Vit A 25,000 u, D 1000 u, B 1 10 mg, B-2 5 mg, niacinamide 150 mg, C 150 mg /Cap Bot 100s

Use: Multiple vitamin therapy

THERAPEUTIC FORMULA W/Vit E (Ives-Cameron) Vit A 25,000 u, D 1000 u,

ascorbic acid 150 mg, thiamine HCl 10 mg, riboflavin 10 mg, niacinamide 150 mg, d-alpha tocopherol acetate 3.4 mg / Cap Bot 100s, 1000s

Use: Multiple vitamin deficiencies

THERAPEUTIC FORMULA W/R-12 (West Ward) Vit A 25,000 u, D 1000 u, thiamine

10 mg, riboflavin 10 mg, ascorbic acid 150 mg, niacinamide 150 mg, crystalline Vit B-12 5 mcg /Cap Bot 100s, 1000s

Use: Therapeutic vitamin

THERAPEUTIC NEMATINIC. (Boyle) Vit B-1

6 mg, B-2 3 mg, C 150 mg, B-12 37.5 mcg, niacin 54 mg, folic acid 1.5 mg, duodenum desic. 100 mg, liver desic. 100

mg, copper 1 mg, cobalt 0.05 mg, ferrous sulfate 505 mg /tab Bot 100s

Use (1 daily) Hematinic & dietary supplement

THERAPEUTIC MULTIVITAMINS. See Vitamins, Multitherapeutic (Various Mfr)

THERAPEUTIC VITAMIN (Ulmer) Vit A 25,000 u, D 1,000 u, B 1 10 mg, B-2 5 mg, C 150 mg, niacinamide 150 mg, E 10 mg /cap Bot 100s

Use: Dietary supplement

THERAPEUTIC VITAMIN FORMULA.

(Brewer) Vit A 25,000 u, B 1 10 mg, B-2 10 mg, B-6 2 mg, cal pantothenate 5 mg niacinamide 150 mg, C 150 mg, D 1,000 u, alpha tocopherol 10 mg, folic acid 1 mg /cap Bot 10s, 30s, 100s

Use Therapeutic vitamins

THERAPEUTIN. (Walker Labs) Vit A 25,000 u, D 1000 u, B 1 10 mg, B 2 5 mg, niacinamide 150 mg, ascorbic acid 150 mg, folic acid 0.67 mg, Vit B 12 5 mcg w/ minerals/Cap Bot 100s

Use Multivitamin therapy

THERASULEA. (Mollard) Meth Dia Mer-Sulfonamides, sulfamerazine, sulfamethazine sulfadiazine 0.166 mg each/Tab Bot 100s, 500s, 1000s

Use: Triple sulfonamide therapy

THERATUSS (Squibb) Pipazethate 10 mg and 20 mg /Tab Bot 100s

Use Coughs

THERA-VI-MIN (Endocrine) Contains 14 vitamins and 11 minerals/Cap Bot 50s, 100s

THERA-VITA (Warner) Vit A 12,500 u, D 1250 u, R 1 10 mg, R 2 10 mg, niacinamide 100 mg, Vit B-6 1 mg, panthanol 10 mg, Vit C 150 mg /Cap Bot 25s, 100s, 1000s

Use: Vitamin deficiencies

THERA-ZYMACAP. (Upjohn) Vit A 25,000 u, D 1000 u, B 12 2.5 mcg, B 1 10 mg, R 2 10 mg, B-6 2 mg, Cal pantothenate 10 mg, ascorbinamide 150 mg, folic acid 2.5 mg, Vit C 150 mg /Cap Bot 30s, 100s

Use: Multiple vitamin deficiencies

THERIDOL (Parke, Davis) Vit A 25,000 u, D 1000 u, B-1 10 mg, B-2 5 mg, B-12 2 mcg, niacinamide 150 mg, Vit C 150 mg /Cap Bot 30s, 100s, 1000s

Use: Vitamin therapy

THERIODIDE, NND 1962 (Abbott) See Sodium Radio iod de

THERMOLENE (Lanett) Sodium chloride 7 gr, dextrose 3 gr, vit B-1 1 mg /tab Bot 1000s, 5000s

THERMOTARS. (Merck Sharp & Dohme) Sodium chloride 0.45 mg, pot chloride 30 mg, cal

- carb. 18 mg., dextrose 0.2 Gm./Tab. Bot. 100s, 1000s.
 Use: Heat foliæ.
- THIBON.** See: Stuart's Thibon, Tab. & Eq. (Sivart)
- THIBUHISTIN.** (Ayerst) Isotripesdyt Hydrochloride (N-dimethylamino-isopropyl-thiophenyl-pyridylamine) Tab. 4 mg. Bot. 100s, 1000s. Syr. 4 mg./5 cc. Bot. 1 pt.
 Use: (1 tab. or 2 tsp. q.l.d.) Antihistamine
- THIBUHISTIN-SA FOSTE.** (Ayerst) Isotripesdyt HCl 24 mg./Tab. Sustained Action. Bot. 100s, 1000s.
 Use: Long-acting antihistamine.
- THI-VI-MIN.** (New Mexico Pharm) Vit. A 25,000 u., D 1000 u., thiamine HCl 10 mg., riboflavin 5 mg., ascorbic acid 150 mg., niacin amide 150 mg./Cap. Bot. 100s, 1000s.
 Use: Therapeutic formula vitamin therapy
- THIBYL.** (Church) Saccharinated aspirin 5 gr., w/benzocaine. Tab., Bot. 40s, 100s, 500s.
 Use: 5 L. (1-2 q. 3 h.); analgesic & antipyretic.
- THIODATE.** N.R. 1948. (Brewer) Thiobromine & Sod. acetate, N.F., Cap. (3½ gr. & 7½ gr.) Bot. 100s, Tab. E.C. (3½ & 7½ gr.) Bot. 100s, 500s, 1000s. Also available w/phenobarbital. Also available w/Pot. Iodide.
 Use: (1 cc. & h.c.) Vasodilator & cardiac stimulant.
- THIVASIN.** (Uncoln) Glyceril guaiacolate 50 mg., atroverline HCl 25 mg., aspirin 165 mg./Tab. Bot. 100s, 1000s.
 Use: 1-2 q. 3 h. analgesic, sedative & expectorant.
- THEX, CAP.** (Ingram) Thiamine HCl 10 mg., riboflavin 10 mg., pyridoxine HCl 2 mg., Col. pantothenate 5 mg., niacinamide 50 mg., ascorbic acid 150 mg., Vit. B-12 4 mcg./Cap. Bot. 100s, 500s.
 Use: Vitamin B complex deficiencies.
- THEX-FOSTE, PABENTERAL.** (Ingram) Double salt agents except choline & inositol & niacinamide/cc. Vit. (1 cc.) Box 12s, 25s, 100s Vit. 15 cc.
 Use: I.V. or I.M.
- THEX, PABENTERAL.** (Ingram) Vit. B-1 50 mg., B-2 1 mg., B-6 1 mg., pantothenic acid 1 mg., niacinamide 75 mg., choline 10 mg., inositol 6 mg./cc. Vit. (1 cc.) Box 12s, 25s, 100s; Vit. 15 cc.
 Use: I.V. or I.M.
- THEX W/C.** (Ingram) Vit. B-1 100 mg., B-2 10 mg., B-6 10 mg., Pantothenol 12 mg., niacinamide 250 mg., ascorbic acid 100 mg./2 cc. Gentisic acid Ethanolamide 2.5%.
- THI (5 cc)** lyophilized, Box 1s, 6s, 25s
 Use: I.V. or I.M.
- *THIABENDAZOLE.** 2-(4'-thiazolyl)-benzimidazole See: Thiabendazole (Merck & Co., Inc)
- THIAEV.** (Key Corp) Vit. B-1 100 mg./cc. Vit. 20 cc.
- THIACITAZONE.** See: p-Acetylamino benzaldehyde thiosemicarbazone
- THIACTIN.** (Bristol) Amphoteric antibiotic containing nitrogen and sulfur from *Streptomyces hawaiiensis*
 Use: Inhibits gram positive organisms.
- THIA-DIA-MES-SULFONAMIDES.** Sulfadiazine w/sulfamerazine & sulfathiazole.
 See: Coca-Sulfonamides Triplex (Lilly)
 Combazine, Tab. (Ives-Cameron)
 Combiul, Susp. & Tab. (Schering)
 Otiemerazole, Susp. (Parke)
 Magmoid Sulco, Susp. (Pitman-Moore)
 Thi-Di-Mes (Pitman-Moore)
 Titul, Susp. (Tilden)
 Trimerazole, Tab. (Mallord)
 Trionamide, Seep. (Pflint, Eaton)
 Trisulfo, Susp. (Central Pharm. Co.)
 Tri-Selfanyl, Susp. (Fenik).
- W/Soligenin & Iimodin.** See: Trisulfacetone, (Central Pharm. Co.)
- W/Urethan.**
 See: Tridermol, Cream. (Central Pharm. Co.)
 Trigynol, Jelly. (Central Pharm. Co.)
- THIA-DIA-METH W/Pot.** penicillin G. See: Pen-Tabs W/Triple Sulfonamides. (Rexall)
- THIADOXINE.** (Estro) Thiamine HCl 50 mg., pyridoxine HCl 50 mg./cc. Vit. 10 cc.
 Use: Nausea & vomiting of pregnancy.
- THEA-FER.** (Pitman-Moore) Ferrous sulfate 3 gr., thiamine HCl 0.375 mg./Tab. or dr. Eq. Bot. 1 pt., 1 gal. Tab., Bot. 500s, 1000s.
 Use: Hematinic.
- THIA-FERRIN.** (Mayer & Rosses) Not listed. Ferrous gluconate 5 gr., vit. B-1 5 mg./tab. Bot. 100s.
 Use: (1 to 2 t.i.d.) Hematinic.
- THIAHEP.** (Lannett) Liver conc. 2½ gr., ferrous sulf. 2 1/10 gr., vit. B-1 1 mg./cap. Bot. 100s, 500s, 1000s.
- THIALBARBITAL.** See: Kamithal
- THIAMALT B COMPLEX.** (Borchardt)
 Syr. (Vit. B-1 12 mg., B-2 12 mg., B-6 1-5 mg., niacinamide 120 mg., pantothenol 30 mg., Vit. B-12 12 mcg., Mong. sulfate 30 mg., w/brewer's yeast conc. & ext. of malt barley/h.oz.) Bot. 8 fl.oz., 1 gal.
 Cap. (Vit. B-1 2 mg., B-2 2 mg., niacinamide 20 mg., Vit. B-6 0.25 mg., Col. pantothenate 5 mg., Mong. sulfate 5 mg., diastatic malt ext. pow. 165 mg., w/brewer's

- yeast conc & ext. of malt barley/cap)
Bot 50s, 100s, 250s.
Use: Dietary supplement
- THIA-MANNITOL** W/Phenobarbital (Walker)
Mannitol hexanitrate ½ gr, Thiamine HCl
3 mg, phenobarbital ¼ gr / Cap Bot 100s,
1000s
Use: (T 2 1 1 d) Hypertension
- *THIAMINE HYDROCHLORIDE, USP,**
NND 1961, ADR Elixir, USP; Inj,
USP, Tab USP Aneurine HCl, thiamine
chloride Vit B 1 HCl
See Apatate Drops (Kenwood Lab)
B Amin, Tab (Buffington)
Beatine (Prema)
Betalin S Amp, Elixir Tab (Lilly)
Betaxin, Amp, Elixir, Syr, Tab (Winthrop
Labs)
Bethiamin, Cap, Elixir, Vial (Masiengil)
Bewon Elixir (Wyeth)
Eikaphen, Elixir, Tab (Smith, Kline &
French)
Uxa Beta Elixir (Warner)
Slowten, Elixir, Tab (Patch)
Thiamal, Elixir (Strassenburgh)
Trophite, Elixir (Smith, Kline & French)
Yinathiam, Elixir (Merck Sharp & Dohme)
W/Ascorbic acid See: B Scorbit, Tab (War-
ner Chilcott)
W/Aspirin See Thi Prin Tab (Covendish)
W/Cal gluconate & ferrus sulf See: Iro-
theran B, Tab (Tilden)
W/Charionic ganadotropin, procaine HCl &
glutamic acid See: Glukor, Vial (Research
Supplies)
W/Petrus sulfate
See: Betaron, Syr (Warren Teed)
Bironex, Elix & Tab (Robins)
Terra Betalin, Tab (Lilly)
W/Gentian fl ext, taraxacum fl ext, phos-
phoric acid See: B Jen, Elix (Fellows)
W/Methylcellulose See Bevell, Tab (Irwin
Neisler)
W/Mineral oil See Geal B, Emulsion
(Smith, Miller & Patch)
W/Neostigmine methylsulfate, atropine sul-
fate See Thiastigmine, Vial (Kirk)
W/Pentylentetrazol nacinamide, cyana-
cobalamin, alcohol See Cenolene, liq
(Central)
W/Phenobarbital
See Benthiamine, Tab (Bentex Pharm)
Vanaphen Prep (VanPell & Brown)
W/Phenobarbital, reserpine See: Nea slow-
ten, Tab (Patch)
W/Scopolamine HBr homatropine methyl-
bromide, hyoscyamine HBr, and phenobar-
bital See Sedlin Unaday, Cap (Paul
Maney)
- W/Testosterone, glutamic acid, procaine HCl**
See: Glutest, Vial (Research Supplies)
W/Thyroid See: Berald, Cap (Walker)
W/Thyroid substance, riboflavin & choline
dihydrogen citrate See: Methyroid, Tab
(Lakeside)
**W/Vit B 2, cal, sad, pot & mang glyce-
ro-phosphate** See: B G-Phas Elix (Merck Sharp
& Dohme)
W/Vit B 6 (Various Mfr)
See B Dax (Lannett)
Grabexine, Amp. (United)
Grandax, Sol. & Tab (Lederle)
Neadax, Amp (Central Pharm. Co.)
Nudax, Vial (Ulmer)
Pyridex, Tab (Lanteed)
Pyrilthad, Tab, Amp (Lakeside)
Thiadaxine, Vial (Estra)
Triandaxine, Vial, (Philadelphia Ampoule
Labs)
W/Vit B 12
See: Apatate, Drops (Kenwood Labs)
Appetum, Tab, chewing gum (Phila Am
poule Labs)
Thicobin, Tab (Bryant)
- *THIAMINE MONONITRATE, USP (Var-**
ious Mfr) Thiamine nitrate Pow, Bot ¼,
½ & 1 lb
W/Corbromal, bromisovalum See: Kartryl,
Tab (Direct Lab)
W/Methylcellulose See: Bevell, Tab (Irwin,
Neisler)
W/Methylcellulose & d amphetamine sulfate
See Celladrine, Tab (Sulliff & Case)
W/Reserpine & vit B 12 See Samalevite
Tab (Purdue Frederick)
- THIAMINTOL.** (Rarer) Thiamine HCl 6 mg /
fl oz Elixir, Bot 14 fl oz, 1 gal
Use: Vitamin B 1 therapy
- THIAMOL.** (Strassenburgh) Thiamine HCl
Elixir (1 mg./5 cc) Bot 1 pt, 1 gal
Use: Vitamin B 1 therapy
- *THIAMYLAL SODIUM,** Inj USP, NND
1962 ADR See Suvital Sodium, NNR
(Parke, Davis)
- THIANTOIN,** NNR 1952 (Lilly) Phethanyl-
ate Susp, oral (24 mg./cc) Bot 1 pt
Use: Anticonvulsant
- THIANTOIN SODIUM,** NNR 1952 (Lilly)
Phethenylate Sod, NNR 1952 Pulvule
(2 gr & 4 gr) Bot 100s, 1000s
Use: Anticonvulsant
- THIAPHYLL CREAM** (Lannett) Sulfathiazole
5%, chlorophyllin 1%. Jar 4 oz & 1 lb.
Use Topically to promote healing
- THIASTIGMINE.** (Kirk) Vit B 1 100 mg-
neostigmine methylsulfate 0.5 mg., atropine
sulf 1/300 gr/cc Amp 1 cc, 10 cc

- Use: Treatment of herpes zoster, neuralgias & auritis.
- THIASULFAMIX.** (Patch) Sulfanilamide 3.75%, sulfathiazole 1.25%. Cream, Tube 1 1/5 oz., 1 lb.
Use: Antibacterial.
- THIATE WITH CODEINE.** (Merrill) Codeine phosphate 10 mg. Pot. guaiacolsulfonate (Thiate) 0.43 Gm./5 cc. Bot. 1 pt., 1 gal.
Use: (1 tsp. q. 4 h.) Coughs.
- THIA-TON.** (Banor) Vitamin B-1 10 mg., B-12 25 mcg./5 cc.
Use: Appetite and growth stimulant.
- THIAZINC.** (Hart) Sulfathiazole 10%, prepared colomine 10% in an aqueous, non-fatty base. Jar 1 oz., 1 lb
Use: Impetigo.
- THIAZOINT.** (Hart) Sulfathiazole 10% in an emulsion base Oint. Jar. 1 oz., 1 lb., 5 lb
Use: Supplement to the accepted surgical procedures of incision and drainage.
- THIAZOLSULFONE, N N D 1960.** See: Promizole, Tab. (Parke, Davis)
- THIBENZOLE.** (Merck & Co., Inc) Thiazobenzazole.
Use: Anthelmintic. Trichinosis.
- THI-COBAL IMPROVED ELIXIR.** (Amco) Vit. B-1 10 mg., B-12 25 mcg./5 cc. Bot. 1 pt.
Use: Dietary supplement at vitamins B-1 & B-12.
- THI-COBIN.** (Bryant) Thiomine HCl 10 mg., vit. B-12 25 mcg./tab. Bot. 100s, 1000s.
Use: Orally, dietary supplement.
- THI-DI-MER.** (Pitman Moore) Sulfathiazole 0.166 Gm., sulfadiazine 0.166 Gm., sulfamerazine 0.166 Gm./Tab. Bot. 100s, 500s, 1000s.
Use: Sulfanamide therapy.
- *THIETHYLPEBAZINE.** (Sandoz) Torecon, 2-ethylmercaptio-10-3 (1-methyl-4 piperazinyl)-propyl phenothiazine dimaleate See: Torecon (Sandoz)
Use: Antiemetic.
- THI-FER-HEPTUM.** (Cavendish) Amp. (Iron peptonate 30 mg., Vit. B-1 10 mg., liver—pres/sc. 1 cc., Box 12s, 25s, 100s. Cap. (Ferrous sulfate 130 mg., liver conc. 486 mg., Vit. B-1 0.4 mg., B-2 0.6 mg., nicotinamide 5 mg./Cap. Bot. 50s, 100s.
Use: Hematinic with vitamin supplement.
- *THIHEXINOL METHYLBROMIDE.** alpha-di-thienyl-(4-dimethylamino-cystahexyl)-carbinal-methylbromide.
W/Polycarbaphil. See Sarbaquel, Tab. (White Labs) See: Entaqueil, Syp. (White)
- THI-LL-FEB.** (Kirk) Liver inj. 5 mcg., vit. B-1 3 mg., iron 2 mg./cc. Amp. 1 cc., 10 cc.
* Generic name recognized by N.F., U.S.P. or A.M.A.
- Also available as Syrup, Bot. 8 oz., & Caps. Bot. 100s, 1000s.
Use: Hematinic.
- THI-LL-FER POSTE.** (C. F. Kirk) Vit. B-12 30 mcg., liver inj. 0.25 cc., iron (ferric citrate) 28 mg./cc. Vial 10 cc.
Use: I.M. Hematinic.
- THILIRON.** (Elder) Liver 3 gr., ferrous sulfate 2 gr., Mang. citrate 4 mg., & Vit. B-1 1 mg./Tab. Bot. 100s, 1000s.
Use: Hematinic.
- THIMECIL, N N D. 1962.** (Physicians Drug) Methylthiourasil Tab. (30 mg.) Bot. 100s, 500s, 1000s
- *THIMEBOSAL, N.F., N.N.R. 1950, A.D.R. 1955,** Oint., N.F.J. Sol., N.F.J. Tr., N.F.J. Sod. Ethylmercurithiosalicylate.
See: Merthiolate (Lilly)
W/Sulfathiazole.
See: Microform Sulfathiazole, Susp. (Smith, Kline & French)
Progmutl, Susp. (Smith, Kline & French).
- THIMORLONE.** (Merck Sharp & Dahme) See Phenylthilane.
- THIOACETAZONE.** See: Tibione
- THIOBAN.** (Parke, Davis) Under study.
Use: Tuberculosis.
- THIO-BISMOL, N.N.R. 1952,** (Parke, Davis) Sed. bismuth thioglycollate. Amp. [0.2 Gm] Box 10s; Amp (2 Gm) Box 10s.
Use: I.M., antisyphilitic bismuth preparation
- 2,2-THIOBIS(4,6-dichlorophenol).** See: Bithional, U.S.P.
- THIO-CARA.** (Suliff & Case) Pot. thiocyanate 2 gr., coscaromatic 6.7 min/5 cc. Bot 1 pt., 1 gal.
Use: (1 dr. t.i.d.) Vasodilator.
- THIOCARBANIDIN.** Under study.
Use: Tuberculosis.
- THIOCARBASONNE.** (Lilly) 4-Carbomida-phenyl di-(carboxymethylthia) arsenite. Envents (25 mg., 50 mg) Bot. 100s. Vial [0.5 Gm] Pkg 6s.
Use: Amebicide.
- THIOCOL.** (Rashe) Pot. guaiacalsulfonate Syr. (8 gr./5 cc) Bot. 6 oz., 1 pt., 1 gal.
Use: (1 tsp. q. 3-4 h.) Cough.
- THIOCYAN-TARS.** (Suliff & Case) Calcium thiocyanate 45 mg/Tab Bot 100s, 1000s
Use: Hypotensive.
- THIOCYL.** (Farigian) Sulfur & Sod. salicylates organically combined. Amp. (2 cc.) Box 25s, 100s.
Use: I.M. (2 cc. 3 times a week) Rheumatism.
- THIODIPHENYLAMINE.** See: Phenothiazine, N.F.

THREAZI. (Irwin, Neister) Aspirin 2½ gr, acetophenetidin 2½ gr, d amphetamine phosphate 1/25 gr/Tob Bot 100s, 500s, 1000s

Use: 12 q 3 h; analgesic & relaxant

THREE BROMIDES, NF (Merck Sharp & Dahme) Ammonium bromide 9 Gm, potassium bromide 8 Gm, sodium bromide 8 Gm, alcohol 4% / 100 cc Elixir, Bot 1 pt, 1 gal
Use: Sedative

THREE BROMIDES, EFFERVESCENT. (Burroughs Wellcome) Sodium bromide 6 gr, Potassium bromide 6 gr ammonium bromide 3 gr, effervescent salts q s / Tab Tube 25s
Use: Sedative

THREE SULFAS SYRUP (Horton & Converse) Sulfadiazine 1 Gm, sulfamethoxazole 1 Gm, sulfamerazine 1 Gm / oz Bot 4 oz, 16 oz
Use: Sulfonamide therapy

THREE SULFONAMIDES, (Upjohn) Sulfadiazine 0.162 Gm, sulfamerazine 0.162 Gm, sulfathiazole 0.162 Gm / Tob Bot 100s, 500s

Use: Triple sulfonamide therapy

THROCAINE-T (Smith Miller & Fatch) Tyrothricin 1 mg benzocaine 5 mg menthol, eucalyptal, anise oil / Tob Bot 100s, 1000s
Use: Infections caused by gram positive organisms

THROCAINE, TAB (Smith, Miller & Fatch) Merc iodide red 1/80 gr, belladonna 1/1000 gr, Phylloca 1/80 gr Hydrastis 1/500 gr, benzocaine 1/10 gr, benzoic acid 1/12 gr eucalyptal 1/22 gr menthol 1/80 gr anise oil 1/80 min / Tob Bot 1000s

Use: Stomach cramps, antispasmodic

***THROMBIN, USP NND 1962 ADR**
Thrombin, topical mammalian origin
Cutter—Vial (5000 u) w/diluent 10 cc
Parke Davis—Amp (1000 u) w/diluent 6 cc; (5000 u) w/diluent 6 cc
Upjohn—Vial (1000 u & 5000 u) 30 cc

Use: To control capillary bleeding

THROMBODEHT (Merck Sharp & Dahme)
Thrombin, human 1 u, tyrothricin 1 u Dental cases, Box 12s, 50s

Use: To accelerate clot formation

THROMBOKINASE W/Prothrombin anti-thrombin See Neo Hemoplastin, 1q (Parke Davis)

THROMBOLYSIN LYOVAC, NND 1961 (Merck Sharp & Dahme) Fibrinolytic Vial 50 000 u 100 cc

Use: Lysis of Thrombi

***THROPHYLLINE** (Mead Johnson Lab) See Quibron

THRY-BEHZO (Jamieson Pharm Co) Tyro

*Generic name recognized by NF USP or AMA

thricin 2 mg, benzocaine 10 mg / troche Pkg 12s

Use: Minor throat & mouth infections

THRYCIN. (Horton & Converse) Tyrothricin 2 mg, benzocaine 5 mg, cetyl dimethyl benzylammonium chloride 1 mg / Troche Bot 12s

Use: Throat and mouth infections

THYBADEN (Bell Pharmacal) Adenosine 5-mono-phosphoric acid 25 mg, vit B-12 60 mcg / cc Vial 10 cc

Use: Treatment of bursitis & osteoarthritis

THYCALSIN. (Pitman Moore) Cal iodized ½ gr, Anesthesin 1/12 gr, thyme 1 gr, & aromatis / oz Bot 100s, 1000s

Use: Mouth & throat infections

THYDRON. (Merrill) Tob (Ferrous sulfate 5 gr, thiamine HCl 1.5 mg / Tob) Bot 100s 1000s Syr (Ferrous sulfate 16 gr, thiamine HCl 4.5 mg / R oz) Bot 1 pt, 1 gal
Use: Anemias

THYLOGEN MALEATE, NND 1962 (Rorer)
Pyridamine maleate Tob (25 & 50 mg) Bot 100s, 500s, 1000s

Use: Antihistamine

THYLOKAY. (Squibb) Menadiolone Sodium phosphate Vit K analogue Tob (5 mg) Bot 100s Sol, Amp (5 mg or 10 mg / cc) 1 cc, Box 5s, 100s

Use: To check bleeding caused by prothrombin deficiency or vitamin K therapy

THYLOQUINONE. (Squibb) See Menadiolone USP

THYLOSE SODIUM, NND 1962 (Jackson Mitchell) Sodium carboxymethylcellulose Tob (5 Gm) Bot 100s

THYLOX. (Shulton Ins) Medicated shampoo contains absorbable sulfur

Use: Seborrheic dermatitis

THYME W/Peppercorn syr, Sod citrate, benzoic acid See Citra Thyme, Syr (Pitman Moore)

THYME, EXTRACT SACCHARATED See Pertussin, Syr (Seck & Kade)

THYMOLIDINE (Blue Line) Phenol 0.5% iodine, camphor, thymol in mineral oil Liq 1 pt, 1 gal Pow, Bot 2/5 oz Supp Box 12s

Use: Topically antiseptic

THYMOL IODIDE, NF X NNR 1949 ADR (Merck Sharp & Dahme) Aristal Pow (Winthrop Labs)

W/Bis oxydide boric acid zinc oxide bis subgallate See Aseptogill, Pow (Mast sengliff)

THYNAL (G F Harvey) Thyroid 1/20 gr, ferrous sulfate 1½ gr strychnine sulfate 1/100 gr zinc phosphide 1/50 gr, calcium gluconate ½ gr / Tob Bot 100s, 1000s

Use: Treatment of general debility

- THYBROM.** (Mills) Brominated Thyroid 1 gr., or 2 gr./tab.
Use: Endocrine obesity.
- THYPHOR INJ.** (Maney) Sod. thiasalicylate 50 mg/cc. Vial 30 cc.
Use: I.M. Antirheumatic.
- THYRACTIN.** (Winthrop Labs) Iodine protein substance isolated from thyroid gland Tab. (25 mg.) Bot. 25s, 100s, 500s.
Use: Hypothyroidism.
- THYRAR.** (Armour) Bovine thyroid preparation. Tab. (½ gr., 1 gr. & 2 gr.) Bot. 100s, 1000s.
Use: Hypothyroid states.
- THYROREX.** (McNeil) Thyroid duo-sayed 30 mg., liver desic. 250 mg., thiamine mononitrate 1 mg., riboflavin 1.2 mg., niacinamide 10 mg., vit. R-12 1 mcg./tab. Bot. 100s, 1000s.
Use: (1 to 6 daily o.c.) Thyroid deficiency conditions.
- THYRROROM.** (Amfre) Brominated Thyroid. Tab (½ gr., 1 gr. & 2 gr.) Bot. 100s, 500s, 1000s.
Use: (1 to 2 gr. daily) Thyroid therapy.
- THYROFED.** (Cole Chem. Co) dl-Desoxyephedrine HCl 10 mg., thyroid ½ gr./tab Bot. 100s, 500s, 1000s.
Use: Obesity.
- THYROGLOBULIN PREP.**
See: Endothyrin (Nelson-Royer)
Proloid (Warner-Chilcott)
Thyrodois (Reed & Cornick)
Thyroprotein (Parke, Davis)
Thytractin (Winthrop)
Thyrrar (Armour)
- THYROID, U.S.P., N.N.D. 1960.** Tab, U.S.P. Thyroid Combinations on pp. 740 to 742.
- THYROID BROMINATED.** See: Thyrobrom, Tab. (Amfre).
- THYROID HORMONES.**
See: l-thyronine Sod. N.N.D. 1959.
Synthroid (Troveval)
Thyroxin.
- THYROID PREPARATIONS.**
See: Endothyrin, Cap. (Warner-Chilcott)
Proloid, Tab. (Warner-Chilcott)
Scirpac, Tab. (Armour).
Thyrodois, Tab. (Reed & Cornick).
Thytractin, Tab. (Winthrop Labs.)
Thyrai, Tab. (Armour).
Thyro-Brom, Tab. (Amfre).
Thyroxin, Prep.
Thyroxin Fraction, Tab. (Squibb)
W/Cat. Iodobenolate, phenolphthalein & Vit. C. See: Cavalytin, Tab. (Cavendish).
W/Steapsin & betaine. See: Chylipase, Tab. (Columbus)
- THYROLOGEN.** (United Lab.) Glycerinated tr. of iodine. Iodine (5 mg./10 min) Bot. 15 cc.
Use: Iodine therapy.
- *THYROPROPIC ACID.** 3,5 diiodo-4-(4 hydroxy-3 iodophenoxy)phenylpropionic acid
See: Triopron (Warner-Chilcott Labs)
- THYROPROTEIN.** (Parke, Davis) Purified ext. of thyroid gland adjusted to contain 0.33% organic iodine. Amp. (1/50 gr.) 1 cc., Box 12s, 100s.
Use: Thyroid therapy.
- THYRO-THIAMINE.** (Parke, Davis) Thyroid strong ½ gr., thiamine HCl 2 mg./Tab. Bot. 100s, 1000s.
Use: Hypothyroidism.
- THYROTROPIC PRINCIPLE OF BOVINE ANTERIOR PITUITARY GLANDS.** See: Thyropar, Vial. (Armour Labs)
- THYROXIN, N.N.R. 1947.**
Orgonon—Sol. (2 mg./cc.) Bot. 15 cc.
Squibb—Cryst. Vial 100 mg Sol. (5 mg./2.5 cc) Bot. 15 cc. Tab. (0.2, 0.4, 0.8 & 2 mg) Bot. 100s.
Use: Thyroid deficiency.
W/Ethinyl estradiol, methyltestosterone, d-amphetamine sulf. & vitamins. See: Neobon, liq. (Roerig)
- THYROXIN FRACTION.** (Squibb) Disodium salt of thyroxin with protein. Tabs. 0.2, 0.4, 0.8, and 2.0 mg Bot 100s.
Use: Thyroid therapy.
- THYROXIN GLYCEROL SOLUTION.** (Squibb) Thyroxin 5 mg/2.5 cc. of glycerol sol. Vial 2.5 cc.
Use: To reduce tension in eye infections.
- I-THYROXINE, SODIUM.** See: Synthroid, Tab (Troveval)
W/Methyl testosterone, potassium estrone sulfate. See: Vi-Tetrin (Savage)
- THYROPAR.** (Armour) Thyrotropic principle of bovine anterior pituitary glands. Vial (10 U.S.P. u) 6 cc.
Use: Thyroid myxedema due to pituitary insufficiency.
- THYRIDOXINE.** (Philadelphia Ampoule Labs.) Pyridoxine 100 mg, thiamine 100 mg/cc. Vial 10 cc.
Use: Morning sickness due to pregnancy.
- TIBIONE.** (Schentley) p Acetylamino benzaldehyde semicarbazone Amithiosene. Thiocastozone. Contabon. Tab. (25 mg & 50 mg) Vial 100s. Bot. 1000s. Pow. Bot. 5 Gm.
Use: Certain forms of tuberculosis.
- TIDEX.** (Dan Hall) Desoropamphetamine sulfate, 5 mg/Tab Bot 100s, 1000s
Use: Obesity control
- TIGACOL.** (Roche) Nicotinic alcohol (Rhone

| Thyroid | Others | Manufacturer | Brand |
|---------|--|----------------------------|----------------------------------|
| 1/12 gr | Adrenal substance ¼ gr, spermine 1 gr, lecithin ¼ gr | Westerfield | Lofone, Tab |
| 16 mg | Amphetamine sulf 5 mg | Dumas Wilson | Nubese Tab |
| 1 gr | d Amphetamine HCl 10 mg, atropine ¼ gr, atropine sulfate 1/360 gr | Dumas Wilson | Amphetamins Combinations, Tab |
| 3 gr | d Amphetamine 15 mg, atropine ¼ gr, atropine sulfate 1/180 gr, phenobarbital ¼ gr | Low | Lawbase CTR, T R Cap |
| 3 gr | d Amphetamine sulf 15 mg, atropine sulf 2/360 gr, atropine ¼ gr, Taylor phenobarbital ¼ gr | Taylor | Obeltra, T R Tab |
| ¼ gr | d Amphetam ne HCl 5 mg, atropine sulfate 1/300 gr, mefloze 2 gr, phenobarbital ¼ gr | Dumas Wilson | Antibese |
| ½ gr | d Amphetamine HCl 5, 10 or 15 mg, phenobarbital ½ or ¼ gr, | Pharm Distributors | Lalobse, Cop |
| 1 gr | d Amphetamine HCl 5, 10 or 15 mg, phenobarbital ½ or ¼ gr | Pharm Distributors | Lalobse, Cop |
| 50 mg | d Amphetamine HCl 15 mg, phenobarbital 25 mg | Dumas Wilson | Obesity, Cop |
| 1 gr | Amphetam ne HCl 10 mg, phenobarbital ¼ gr | Dumas Wilson | Amroid, Tab |
| 10 mg | d Amphetamine phosphate 5 mg, vitamins | Adams | T V O Tab |
| 3 gr | d Amphetamine sulfate 15 mg, atropine sulfate 1/180 gr, atropine ¼ gr, phenobarbital ¼ gr | Paul Maney | Nika Caps Ph 749, Cop. |
| 1 gr | d Amphetamine sulf 5 mg, methylcellulose 5 gr, atropine sulf 1/360 gr, phenobarbital ¼ gr | Beff Pharmacia | Celudex, Tab |
| 1 gr | d Amphetamine 5 mg, atropine 50 mg, atropine sulfate 0.18 mg, vitamins | Normand | Ampholin, afternoon Tab, |
| 1 gr | d Amphetamine HCl 5 mg, atropine sulfate 0.18 mg, vitamins | Normand | Ampholin, morning Tab, |
| 1 gr | d Desoxyephedrine 1/12 gr, atropine sulfate 1/320 gr | Marton | O B E C, Tab. A |
| 1 gr | d Desoxyephedrine HCl 5 mg, atropine sulfate 1/360 gr | Meyer | Myrbie # 2, Tab |
| 1 gr | d Desoxyephedrine HCl 1/12 gr, atropine sulfate 1/360 gr, atropine ¼ gr | Hance | Amphroid, Tab |
| 1 gr | d Desoxyephedrine HCl 5 mg, atropine sulfate 1/360 gr, atropine ¼ gr | Meyer | Myrbie # 1, Tab. |
| ¼ gr. | d Desoxyephedrine HCl 5 mg, methylcellulose 2½ gr, phenobarbital ½ gr | Zimmer | Hyzabel, Tab |
| 1 gr | d Desoxyephedrine HCl 5 mg, phenobarbital ¼ gr | Meyer | Myrbie # 3, Tab. |
| ½ gr | d Desoxyephedrine HCl 3/40 gr, Vit B 1 3/40 gr, | Calumbus | Desolds, Tab |
| ½ gr | d Desoxyephedrine HCl 4 mg, Vit B complex w/20 mg Vit. C 4 mg | Kendall | Deegians, Tab |
| ¼ gr | D enevitol 1/640 gr Estrogens | Warner Chilcott Carrick | Manocrin Tab Hormotone T, Tab |

Thyroid Combinations

| Thyroid | Others | Manufacturer | Brand |
|---------|---|------------------|-------------------------------|
| 1/2 gr | Vit B 1 mg, Vit B 2 mg, niacinamide 10 mg, Vit B-6 0.1 mg, Cal pantothenate 2 mg | Haskell | Pantabeeroid, Tab. |
| 30 mg. | Vit B-1 mg, Vit B-2 mg, niacinamide 10 mg, Vit. B-12 1 mcg, liver desic. 250 mg | McNeil | Thyrabex, Tab. |
| 32 mg. | Vit B-1 3 mg, Vit B-2 1 mg, niacinamide 5 mg, Vit B-6 6 mg, Vit C 15 mg, Vit D 2000 u | Warren Teed | Vitaroid, Tab |
| 1/2 gr. | Vitamins | Kenwood | Infertab M, Tab. |
| 1/5 gr. | Vitamins multiple, minerals & metabolic substances Zinc phosphide 1/10 gr, iron pyrophosphate 2 gr | Raering Elder | Vj Thyro, Cap Nepylol, Tab |

col) 50 mg, trimethobenzamide HCl 100 mg/Cap Bot 50s

Use: Relieve dizziness and nausea

TIGADOXAL. (Rache) Trimethobenzamide and pyridoxine/Tabs

Use: Anti-nauseant in pregnancy

TIGAN HCl, N N D 1962 (Rache Labs) (Tri methobenzamide hydrochloride) 4 (2 di methylaminoethoxy) N (3,4,5 trimethoxy benzoyl) benzylamine hydrochloride Cap 100 mg Bot 100s 500s Cap 250 mg Bot 20s 50s Amp. (100 mg/cc) 2 cc Box 6 & 25s; Vial 20 cc Supp 200 mg Box 6s Tmespan Tab (300 mg) Bot 50s 500s

Use: Nausea and vomiting

W/Rational See: Tigacal Caps (Rache)

TILAMIDE. (Tildan) Sulfacetamide 15 gr, sulfadiazine 15 gr, sulfamerazine 15 gr, Sod citrate 45 gr/fl oz Bot 4 oz, 1 pt
Use: Triple sulfonamide therapy

TIL-FOR. (Tilden) Sulfanamides meth d a mer thiazole 0.125 Gm ea, Sod citrate 0.5 Gm, lemon color & flavor Bot 4 oz 1 pt

Use: Oral susp; sulfonamide therapy

TIL-ZINE (Tilden) Sulfadiazine 22 1/2 gr, sulfamerazine 22 1/2 gr, Sod citrate 45 gr/fl oz Susp, Bot 4 oz, 1 pt

Use: Sulfonamide therapy

TIMED AMODEX. (Testagar) Dextroamphetamine HCl 15 mg amobarbital 60 mg/Cop Bot 100s

Use: (1 daily) Obesity, nervous anxiety & depression

TIMED ARDA. (Testagar) d Amphetamine sulfate 15 mg amobarbital 45 mg, Vit A 6600 u, D 400 u B 1 16 mg, B 2 25 mg, niacinamide 155 mg, C 30 mg, ferrous sulfate 28 mg, cobalt sulfate 0.49 mg, copper sulfate 28 mg, sodium molybdate 0.45 mg, zinc sulfate 39 mg, potassium iodide 0.13 mg/Cop

Use: Appetite depressant and vitamin and mineral supplement

TIMED TRIDEX. (Testagar) Dextroamphetamine HCl 15 mg/Imed Cap Bot 100s

Use: (1 daily) Obesity

TIMED TRIDEX JUNIOR. (Testagar) d Amphetamine HCl 7.5 mg/Cop Bot 100s 1000s

Use: Anti-obesity

TIMOFAX (Burroughs Wellcome) Undecylenic acid Oint (10%) Tube 3/4 oz, Jar 1 lb
Pow (10%) 2 1/2 oz (shaker top)

Uses: Fungicidal

TIMOYAN (Ayerst) Prothipendyl HCl, 25 mg or 50 mg/Tab Bot 100s 1000s

Use: Relief of tension

TINAZOLR ELIXIR (Blaine Co) Pentylam-

- tetraol 200 mg., nicotinic acid 100 mg., alcohol 5%/Tsp. Bot. Pis. Gals.
Use Analeptic-respiratory stimulant.
- TINDAL.** (Schering) Acetophenazine dimaleate 20 mg./Tab. Bot. 100s.
Use: Anti-anxiety agent used in the treatment of anxiety, tension, hyperexcitement, fear, irritability, nervousness, and apprehension.
- TIN THERAPY.**
See Chondro-Stann (Christina).
Callosal Stannum (Crookes-Bornes)
Oxistan (Chicago).
Safistan (Patch).
Stannine (Blue Line).
Stannoxyd. (Anglo-French)
Stanno-Yeast (Stoddard).
Tin Tabs. (Cole Chem. Co)
- TIN-TABS.** (Cole Chem. Co) Tab., Bot. 100s.
Use: Tin therapy.
- TINUVIN P.** (Geigy) 2(2'-hydroxy 5'-methyl-phenyl)benzotriazole/Drum 310 lb
Use Ultraviolet absorber.
- TINY VITAPRIME DROPS.** (Vitaprime Co)
Bot. 60 cc.
Use Superficial lesions.
- TIPEX.** (Filmon-Moore) Tyrothricin 0.5%, pectin 6%, propylene glycol 20%. Tube 1 oz.
Use Superficial lesions.
- TISIN, N.N.D. 1962.** (U. S. Vitamin) Isoni- acid. Tab. (50 mg. & 100 mg.) Bot. 100s, 1000s.
Use: In streptomycin-resistant cases of pulmonary tuberculosis.
- TIS-U-SOL.** (Boaxter) Sod. chloride 800 mg., pot. chloride 40 mg., magnesium sulfate 20 mg., sod acid phosphate 875 mg., pot. acid phosphate 625 mg., dextrose 100 mg./100 ml. Bot 1000s Sterile Sol Bot. 1000 ml.
Use: Irrigation.
- TITANIUM DIOXIDE, U.S.P.**
- TITANIUM OXIDE.** (Texas) Titanium oxide 10%. Oint. 1 lb.
Use: Skin antiseptic.
- TITRALAC, N.N.B. 1955** (Schenley) Amino- acetic & Cal. carb., glycine (30%) & Cal. carb. (70%). Tab., Bot. 100s, 1000s, Box 40s Liq., Bot. 8 fl. oz. Pow., Jar 4 oz.
Use: Gastric antacid.
- T. K. O.** (Modern Products) Dimono-triethyl- amine salicylate w/benzocaine. 1½ oz. roll-on container
Use: Relief of pain in headache, neuritis, neuralgia, etc.
- T.L.C.** (Polychem) Silicone and hexachloro- phenol Bot. 8 oz.
Use: Skin care, body massage
- * Generic name recognized by N.F., U.S.P. or A.M.A.
- T-M LOZENGES.** (Drugmaster) Tyrothricin 2 mg., benzocaine 5 mg./trache. Pkg. 12s.
Use: Superficial infections of throat & mouth.
- TOBINE.** (Amfre) Thiourea, o-phenylphenol & benzocaine. Oint. 1 oz., 3 oz.
Use: Antiseptic with local anesthetic.
- TOBIE TABLETS.** (Testogor)
No. 1: Regular strength. Desoxedrine 4 mg., thyroid 1 gr., atropine sulfate 1/360 gr., atain ¼ gr./tab. Bot. 100s, 1000s. Also avail. in double strength.
No. 2: Regular strength. Desoxedrine 4 mg., thyroid 1 gr., atropine sulfate 1/360 gr./tab. Bot. 100s, 1000s. Also avail. in double strength.
No. 3: Regular strength. Desoxedrine 4 gr., thyroid 1 gr., phenobarbital ¼ gr./tab. Bot. 100s, 1000s. Also avail. in double strength.
Use: Antiobesity.
- TOCIRANA EXPECTORANT CONCEN- TRATE.** (Bruce) 1 part is diluted with 4 parts of syrup or honey; it contains per 30 cc: pyrilamine maleate 80 mg., ammonium chlor- ide 500 mg., sodium citrate 500 mg., saccharin 250 mg., menthol 5 mg., glycerin 20%, tolu sulfoxide 8%, cocillona ex- tract 2.5%.
Use: Antihistaminic and decongestant
- TOCLASE, N.N.D. 1962.** (Pfizer) Carbetapen- tone citrate Syrup 7.25 mg./5 cc. Bot. 1 pt.
Use: p.r.n. cough.
- TOCLASE EXPECTORANT COMPOUND.** (Pfizer) Carbetapentone citrate 7.25 mg., terpin hydrate 16.67 mg., chloroform 2.45 mg./5 cc. Bot. 1 pt.
Use: p.r.n. cough.
- TOCOPHEREX.** (Squibb) Vit. E. Cap. (25 & 100 u) Bot. 100s, 1000s.
Use: Vitamin E therapy.
- * **dl-alpha-TOCOPHEROL, NF 57.8** Tri- methyltolcol, alpha-Tocopherol, dl alpha- tocopherol, vit. E, wheat germ oil.
See
Cap. & Tab.:
Alfacol, Cap. (Asher)
Almetrol, Cap. (Meyer)
Aqualol, E. Cap. (30 mg., 100 mg) (U.S. Vit.)
Cavacol, Perles (3 min) wheat germ oil (Cavendish).
Denomane, Cap. (3 min) wheat germ oil (Vio-Bin).
Ecofrol, Cap. (30 mg., 100 mg) (Breon).
Eprolin 5, Gelseal (5 mg) (Lilly)

Epsilon M, Cap (30 mg, 100 mg) (Warren Teed)
 Etorb, Cap (50 mg) (Wyeth)
 E Toplex, Cap (30 mg, 100 mg, 150 mg) (U S Vitamin)
 E Valine, Cap (50 mg) (Smith Dorsey)
 Evipheral, Cap (50 mg) (Kirk)
 E Vitum, Cap (50 mg) (Ives Cameron)
 Myopano, Perles (120 mg) wheat germ oil (Drug Prods)
 Vasculals, Cap (30 mg, 100 mg) (U S Vitamin)
 Verral, Cap (50 u) (McNeil)
 Vi E, Cap (50 mg) (Blue Line)
Oint
 Myopano (55%) (Drug Prods)
Sal
 Aquasol E Sal (50 mg/cc) (Funk)
 Vasculals, Sal (100 mg/cc) (U S Vitamin)
Vial,
 Alfacol Vial (Asher)
 Epsilon, Amp (200 mg/cc) (Warren Teed)
 Epsilon Phosp, Amp (50 mg/cc) (Warren Teed)
 E Vitine, Vial (200 mg/cc) (Smith Dorsey)
 Evipheral, Vial (200 mg/cc) (Kirk)
 W/Ethisterone, hesperidin complex, ascorbic acid Vit K analogue See: Nugestlarol, Tab (Organon)
 W/Phenobarbital & d desoxyephedrine HCl See: Transibarb Cap (Brean)
 *dl-alpha-TOCOPHERYL ACETATE, NF
 See: E Hart, Vial (Hart)
 Ephynal Acetate (Roche)
 Syntapheral Acetate (Abbott)
 Tocaphrine (Barry)
 Vi Ea, Cap (50 mg, 100 mg) (E S Miller)
 W/Testosterone propionate See: Testoferol Vial (Ascher)
 *TOCOPHEROLS, MIXED, CONCENTRATE, NF (Natural)
 See: Epralin, Gelseals (Lilly)
 Germal E (Ayerst)
 Mixed Tocopherols (Parke, Davis)
 Natapheral, Cap (Abbott)
 Tolaxin, Cap (Winthrop Labs)
 Vitamin E Conc. Cap (Uplahn)
 Vitamin E, Cap (Walker)
 *d-alpha-TOCOPHERYL ACETATE, NF
 See: Cotapheral Vial (Endo)
 Ecotal, Cap (Brean)
 Econ, Cap (50 mg) (Cals)
 E Feral, Cap, Vial (Testagar)
 Mixed Tocopheryl Acetate (Parke Davis)
 Plesoral (Plesner)

Tocopherex, Cap (Squibb)
 Tocopherol, Cap (A P C)
 Tokols, Cap (Ulmer)
 Vi Dom "E" Pillettes (Dame)
 Vs E (Blec Line)
 W/Testosterone propionate, chlorobutanol See Androtocopheral (New Mexico Pharm)
 *d-alpha-TOCOPHERYL ACETATES, CONC, NF (Parke, Davis) d alpha Tocopheryl acetate 74 mg/cap Bot 100s 1000s
 Uses Treatment of habitual & threatened abortion
 *d-alpha-TOCOPHERYL ACID SUCCINATE, NF See E Feral Succinate (Testagar)
TOCOPHRINE, (Barry) dl alpha Tocopherol acetate Vial (200 mg/cc) 10 cc
 Uses Vitamin E therapy
TOCOSAMINE SULFATE (Trent) Sparteine sulfate 150 mg sod chl 4.5 mg/cc Amps 1 cc Box 12s
 Use 1 M uterine inert o
TOFAXIN, (Winthrop Labs) Mixed tocopherals Amp (200 mg/cc) 1 cc, 5s Cap (30 mg, 100 mg) Bot 100s Vial (200 mg/cc) 10 cc
 Uses Vitamin E therapy
TOFRANIL, NND 1961 (Geigy) Imipramine HCl 5 (3 Dimethylaminopropyl 10 11 d hydro 5H dibenz (b f) azepine HCl 25 mg/Tab Bot 100s & 1000s Amps 25 mg/2 cc) box 100s & 50s
 Use Antidepressant Thymoleptic
TOIN UNICELLES (Miss) Diphenylhydantoin base 250 mg/Tab Bot 30s 100s
 Use Epilepsy
TOKOLS, (Ulmer) d alpha Tocopheryl acetate Cap 30 mg Bot 100s 500s, 1000s
 Uses Vitamin E therapy
TOLAGESIC, (Smith Miller & Patch) Salicylamide 300 mg, mephenesin 300 mg/Tab Bot 100s, 500s, 1000s Elixir (1 teaspoonful is equiv to 1 Tolagesic Tab) Bot 1 pt, 1 gal
 Uses 1 2 q 3 h; muscle relaxant with analgesic
TOLAMIC, (Kremers Urban) Mephenesin 0.25 Gm, & glutamic acid HCl 0.25 Gm/Cap Bot 100s, 500s, 1000s
 Uses (2 3 q 3 h) Relief of pain and spasm in muscle injuries & arthritis
TOLAMIC IMPROVED (Kremers Urban) Mephenesin 0.25 Gm, Cal glutamate 0.25 Gm/Cap Bot 100s, 500s, 1000s
 Uses Muscle relaxant
TOLANATE, (Reed & Carnrick) Inositol hexantrate Tab (10 mg) Bot 100s, 1000s
 Uses (1 tid) Vasodilator in management of hypertension

* Generic name recognized by N F, U S F or A M A

- W/Phenobarbital. (Reed & Carnrick) Inositol hexanitrate 10 mg., phenobarbital $\frac{1}{4}$ gr./Tab. Bot. 100s, 1000s
 Use: (1 t.l.d.) Vasodilator with sedative.
- TOLANSIN**, N.N.D. 1962. (Garde Drug) Mephenesin, Tab. (0.25 & 0.5 Gm) Bot 100s.
- TOLAPHEN-PENTOBARBITAL**. (Invenex) Mephenesin 0.5 Gm. w/pentobarbital Sod. available as $\frac{1}{4}$ gr., $\frac{1}{2}$ gr. & $\frac{1}{2}$ gr./Tab. Bot. 100s, 1000s.
 Use: Anxiety & tension.
- TOLAX**, (Sutlif & Case) Mephenesin, N.F. liq. (0.5 Gm./5 cc.) Bot. 1 pt., 1 gal. Tab (0.5 Gm) Bot. 100s, 1000s.
 Use: (1 Gm. t.l.d.) Muscle relaxant.
- TOLAZAMIDE**, (Upjohn) 1-(hexahydro-1-azepinyl)-3-p-tolylsulfanyurea.
- TOLAZOLINE HYDROCHLORIDE**, U.S.P., N.N.D. 1962; Inj., Tab., U.S.P. Renexazoline HCl, 2-benzyl-2-imidazoline HCl. Vasimid, Vasodil.
 See: Arterodyl TT (Columbus) Friscoline. (Ciba)
 Tolpal, Vial (Philadelphia Ampoule)
- TOL-BLUE CHLORIDE**, (Abbott) See: Blutene, Tab. (Abbott).
- TOLBUTAMIDE**, U.S.P., N.N.D. 1962. N-butyl-N'-toluene-p-sulfanyurea. Bastnon. See: Orinase (Upjohn)
- TOLBUZEM**, (Zemmer) Mephenesin 0.5 mg., butabarbital 6 mg./Tab. Bot. 100s.
 Use: Relaxant to relieve tension and nervousness.
- TOLDEX**, (Pfliman-Moore) Dexamethasone 0.5 mg., phenylloxamine dihydrogen citrate 88 mg./Tab. Bot. 30s.
 Use: Inflammatory and allergic disorders.
- TOLERON**, N.N.D. 1962. (Mollinckrodt) Ferrus fumarate
- TOLFERAIN**, (Ascher) Ferrus fumarate 300 mg providing 100 mg. elemental iron/Tab. (sustained release) Bot. 100s, 1000s.
 Use: Iron deficiency anemia.
- TOLPIC**, (International Vitamin Div.) Vit. A 50,000 u., D 1000 u., thiamine HCl 10 mg., riboflavin 10 mg., ascorbic acid 150 mg., niacinamide 150 mg./Cop. Bot. 30s, 100s.
 Use: Dietary supplement.
- TOLFRINIC**, (B. F. Ascher) Ferrus fumarate 600 mg., Vit. B-12 w/intrinsic factor concentrate 1 U.S.P. u., Vit. C 100 mg., folic acid 2 mg./Tab. Bot. 50s.
 Use: Hematinic.
- TOLHART**, (Hart) Mephenesin, Tab. (0.5 Gm.) Bot. 100s, 1000s.
 Use: (1 Gm. t.l.d.) Muscle relaxant.
- TOLONIUM CHLORIDE**, N.N.D. 1962. 3-Amino-7-dimethylamino-2-methyl-phenazone
 * Generic name recognized by N.F., U.S.P. or A.M.A.
- thionem. See: Blutene, Tab., N.N.D. 1958. (Abbott)
- TOLOSAT**, N.N.D. 1962 (Brewer) Mephenesin, Tab. (0.25 & 0.5 Gm) Bot. 100s.
 Use: (1 Gm. t.l.d.) Muscular relaxation.
- TOLOXYDYNE**, (G. F. Harvey) Phenyltoloxamine dihydrogen citrate 25 mg., aspirin 4 gr., phenacetin 2 gr., caffeine $\frac{1}{2}$ gr./Tab Bot. 100s
 Use: Analgesic w/antihistaminic.
- * TOLOXYCHLORINAL** (Warner-Chilcott) 1,1'-(3-o-tolylxypropyleneoxy) bis (2,2,2-trichloroethanol). See: Propoxychlorinal
- TOLOXYN**, N.N.D. 1962 (Delta Chemical) Mephenesin, Bot. 100 Gm.
 Use: Muscle relaxant.
- 3-o-TOLOXY-1,2-PROPANEDIOL**. See: Mephenesin, Prep.
- TOLPAL** (Philadelphia Ampoule) Tolazoline HCl 25 mg./cc. Vial 10 cc.
 Use: Vasodilator
- * TOLPOVIDONE** ⁽¹²¹⁾, (Abbott) 1-(w-p-lodo-1131-benzylpolyethylene)-2-pyrrolidinone. See: Roovin.
- TOLPROPAMINE**, [Registered in Great Britain] NN-Dimethyl-3-phenyl-3-p-tolylpropylamine.
- * TOLPYRAMIDE**, (Burroughs Wellcome) N-p-tolylsulfanyl-1-pyrrolidincarboxamide.
 Use: Antidiabetic compound.
- TOLSERAM**, N.N.D. 1962. (Squibb) Mephenesin carbamate, Tab. (7 $\frac{1}{2}$ gr) Bot. 100s, 1000s. Susp. (1.0 Gm./5 cc) Bot. 1 pt., 1 gal.
 Use: Muscle relaxant.
- TOLSEROL**, N.N.D. 1962. (Squibb) Mephenesin Caps. (0.25 Gm.) Bot. 100s, 1000s. Elixirs (0.1 Gm./cc) Bot. 1 pt., 1 gal. Sol.: (20 mg./cc) Amp. 50 & 100 cc. Tab.: (0.25 & 0.5 Gm) Bot. 100s, 1000s.
 Use: (1 Gm. t.l.d.) Muscle relaxant.
 W/Codeine. (Squibb) Tolserol 0.5 gr., & codeine sulfate $\frac{1}{2}$ gr./Tab. Bot. 100s, 1000s.
 Use: Acute, painful rheumatic disorders.
- TOLTEL**, (Pharma-Seal) Mephenesin 500 mg./Tab. Bot. 100s, 1000s
- TOLUDINE SYRUP**, (G. F. Harvey) Dihydrocodeinone bitartrate $\frac{1}{6}$ gr., lobelia $7\frac{1}{2}$ gr., chloroform $2\frac{1}{2}$ min., tartar emetic $\frac{1}{2}$ gr., Roid tale 14 min./fl. oz. Bot. 4 oz. 1 pt., 1 gal.
 Use: Symptomatic relief of dry cough
- TOLUIDINE BLUE O CHLORIDE**. See: Blutene, Tab. (Abbott).
- * TOLULEXIN**. See: Tolulax (E S Miller Lab)
- TOLUOX**, N.N.D. 1961. (Smith, Miller & Patch) Mephenesin 500 mg/Tab. or $\frac{1}{2}$ tea-

- spoonful Tab Bot 100s 1000s Elixir Bot 8 oz 1 gal
Use: Muscle relaxant
- TOLYPHY.** (Chicago Pharm) Mephenesin D 5 Gm, physostigmine 0.35 mg, atropine D 2 mg / Tab Bot 100s 500s 1000s
Use (1 to 2 t i d) Skeletal muscle relaxant
- TOLYSIN** (Lederle) Neocinchophen
- TOLYSPAZ.** N N D 1962 (Chicago Pharm) Mephenesin Tab (0.5 Gm) Bot 100s 1000s 6000s
Use (1 Gm t i d) Muscle relaxant
- TOMETIN** (Ayerst) Nickel pectinate 150 mg, Sod chloride 50 mg, Pot nat less than 20 mg together with other factors naturally present in dried fresh tamala pulp 780 mg / Gm Bot 50 Gm
Use Diarrheas of nonspecific origin
- TONGA & SALICYLATE COMPOUND** (Merck Sharp & Dahme) Tonga bark 4 1/4 Gm Cimicifuga 1.04 Gm, sod salicylate 2.08 Gm, pilocarpine nitrate 5 mg, alcohol 16% / 30 cc Bot 1 gal
Use Antirheumatic
- TONICINE** (Reed & Carnrick) Strychnine sulfate 0.36 mg, iodoquinine sulfate 4 mg iron Sod maleate 11 mg, Sod biphosphate 115 mg, thiamine HCl 0.33 mg, riboflavin 0.33 mg / 4 cc Elixir, Bot 8 oz, 1 gal
Use: Tonic
- TONIGESTINE** (Blue Line) Papain, pepsin, amylapsin gentian salts & glycerin liq Bot 1 pt, 1 gal
Use: Digestant
- TONIH** (Central) Ferrous gluconate 65 mg, Vit B 1 D.5 mg, Vit D 400 units / Palatob Bot 100s 1000s
- TONOCO** (Sutliff & Case) Strychnine sulfate 0.8 mg, thyroid 4 mg, zinc phosphide 4 mg, iron 65 mg calcium gluconate 97 mg / Tab Bot 1000s
Use: Systemic stimulant and tonic
- TOOTHACHE THERAPY.** See: Dentalone liq (Parke, Davis)
- TOPAMINIC.** (Merck Sharp & Dahme) Metha pyrilene HCl 2 Gm, calamine prep B Gm, benzocaine 3 Gm, hexylated cresols DD5 Gm / 100 Gm Cream, Tube 1 cc
Use: Anesthetic analgesic & antihistaminic cream
- TOPASIL** (Cyanamid) Silicone preparation Aersal can
Use: Protectant
- TOPATAR CREAM** (Merck Sharp & Dahme) Coal tar, sal 5%, sulfur colloidal 2.5 Gm, salicylic acid 3 Gm zinc oxide 5 Gm / 100 Gm Cream, Jar 2 oz
Use: Dermatitis, eczema, neuradermatitis, psoriasis
- TOPNOSAN** (Cavendish) Cinchophen Tab, Bot 100s
Use Analgesic & antirheumatic
- TOPICORT OINTMENT.** (Rausser) Hydrocortisone 1% & 2.5% Tube 5 Gm
Use Topically, dermatitis & related infections
- TOPICORT SPRAY.** (Rausser) Hydrocortisone D.5% Vial, aerosol spray 15 cc
Use Topically, dermatitis & related infections
- TOPISOL** (Texas Pharm Co) A solvent composed of alcohol 15%, propylene glycol and sorbitan monolaurate Bot 1 pt
- TOPITRACIN.** (Ayerst) Bacitracin Vial (50 000 u) 50 cc
Oint (500 u / Gm) Tube 1/2 oz, 1 oz, 4 oz Ophth Oint (500 u / Gm) Tube 1/2 oz Ophth Oint W/Phenacaine (1000 u / Gm W/Phenacaine HCl 1/4%) Tube 1/8 oz
- TOPITRACIN** (Reed & Carnrick) Bacitracin 500 u / Gm Oint 1/2, 1, 4 oz Ophth Oint, Tube 1/8 ac Vial (50 000 u) 50 cc
Use Topically; skin infections
- TOPITRACIN OINT.** W/Phenacaine (Reed & Carnrick) Bacitracin 1M u, phenacaine HCl 2.5 mg / Gm Oint, Tube 1/8 oz
Use: Infections of the skin
- TOPOCAINE.** See: Surfacaine (Lilly)
W/Sulfadiazine allantoin See: Sulfanamets, Iox (National Drug)
- TOPOCIDE** (Lilly) Emulsion of benzyl benzoate, DDT, benzocaine polyoxyalkylene derivatives of sorbitan monooleate Bot 4 oz 1 pt, 1 gal
Use: Scabicide, pediculicide
- TORANTIL.** (Winthrop Labs) Hag kidney ext containing a histamine destroying enzyme histaminase in active & stable form Tab EC (10 u) Bot 100s 500s
Use: (3D u, t i d) Antiallergic
- TORICAN** (Sandoz) Thiethylperazine maleate 10 mg / Tab Bot 100s; 5 mg / cc Ampuls 2 cc Box 12s, 100s
Use: Antiemetic and antinauseant
- TOROCOL.** (Plessner) Bile salts 1 gr, cascara sagrada ext 1 gr, phenolphthalein 1 gr, Capsicum oleoresin 1/40 gr, pepper mint oil 1/200 min / Tab Bot 100s, 500s 1000s
Use: (1 t i d, a c) Choleric
- TOROCOL COMPOUND** (Plessner) Bile salts 1 gr, papain 2 gr, cascara sagrada ext 1/2 gr, phenolphthalein 1/2 gr, Capsicum oleoresin 1/40 min / Tab Bot 100s, 500s, 1000s
Use: Choleric
- TORULA YEAST, DRIED, U.S.P.** Obtained by

- growing *Candida* (toruopsis) utilis yeast on wood pulp wastes (Chartes Bowman & Co) Use (10 Gm. q.t.d.) Supplemental source of protein.
- TORULEXIN.** (Upjohn) Thiomine HCl 2.5 mg., riboflavin 1 mg., nicotinic acid amide 5 mg., Cal. pantothenate 0.25 mg., pyridoxine HCl 1 mg./Cap. Bot. 5Ds, 500s, 1000s. Use: Vitamin B complex.
- TOBULOX, N N R.** (Miller) Mephenesin, Tab (0.25 & 0.5 Gm) Bot. 100s, 500s, 1000s. Elixir, Bot. 8 oz., 1 gal. Use: (1 Gm. i.i.d.) Muscle relaxant.
- TOSANON.** (Organon) Mephenesin 75 mg., dihydrocadenone bitartrate 1.67 mg., Pot. citrate 400 mg., pyrrolamine maleate 7.5 mg./5 cc. Syr., Bot. 1 pl. Use: (1 tsp. q. 3-4 h.) Sedative & expectorant.
- TOSEC.** (Hoack) Pentylametetrazol 3 gr., niacin $1\frac{1}{2}$ gr./tab. Bot. 100s, 1000s. Use: Respiratory & circulatory stimulant.
- TOSEDRIIN.** (Amfre-Grant) Glyceryl guaiacolate 50 mg., chlorpheniramine maleate 2 mg., phenylpropranolamine HCl 10 mg., ammonium chloride 120 mg./5 cc. Liq Bot. 4 oz. Use: Treatment of coughs due to colds.
- TOTAMIN.** (Buffington) Vit A 5000 u., D 500 u., thiamine chloride 1.5 mg., riboflavin 2 mg., niacinamide 20 mg., ascorbic acid 37.5 mg., Vit. B-12 2 mcg., folic acid 0.25 mg., Cal. pantothenate 5 mg., pyridoxine HCl 0.5 mg./Tab. Bot. 100s, 1000s. Use: Dietary supplement.
- TOTAQUINE, N.P. X.** Alkaloids from Cinchona bark, 7 to 12% quinine anhyd, 70% to 80% total alkaloids (cinchonidine, cinchonine, quinidine & quinine) Mallinckrodt—Pow., Bot. 1 oz., Canister, 5 oz., 25 oz. Smith-Dorsey—Pow., Bot. 5 oz., 50 oz.
- TOTA-VI-CAPS.** (Elder) Vit. A 5000 u., D 1000 u., B-1 2.5 mg., B-2 2.5 mg., niacinamide 20 mg., Vit. B-6 D 5 mg., cal. pantothenate 5 mg., folic acid D 5 mg., Vit. B-12 1 mcg., C 40 mg., E 2 u./Cep. Bot. 100s, 500s, 1000s. Use: Vitamin supplement
- TOTAL.** (Key) Oyster shell 625 mg., Vit. A, D, C, B-6, B-12, ferrous fumarate 150 mg./Tab. Bot. 100i. Use: Prenatal therapy.
- TOVILAC.** (Chicago Pharmacol) Vit. A 700 u., D 250 u., B-1 D.167 mg., B 2 D.334 mg., C 25 mg., niacinamide 5 mg., E D.34 1 u., folic acid D.167 mg., B 12 w/intrinsic factor conc. 1/30 u., ferrous sulf. basic. $1\frac{1}{2}$ gr., cal. lactate 5 gr./tab. Bot. 100s. Use: Dietary supplement.
- TOXICOGENON OUBECIFOLIUM EXTRACT.** See: Anergex, Vial (Mullford Colloid Labs)
- TOXIVL.** (Cutter) Poison Ivy, oak & sumac alcoholic ext Amp 1 cc. Box 3s w/syringe & needles. Use: Shus dermatitis.
- TOXOID, DIPHThERIA ADSORBED, N N D 1962** See: *Diphtheria Toxoid*, Alhydrox (Cutter) *Diphtheria Toxoid*, Aluminum Phosphate, Adsorbed
- TOXOID, TETANUS ADSORBED, N N D. 1962** See: *Tetanus toxoid*, alhydrox (Cutter)
- TOXOID MIXTURE.** See: *Tri-Solgen* (Lilly)
- TOROK.** (Cutter) Poison Ivy, oak & sumac ext. Amp. 1 cc Vial 2D cc. Use: Shus dermatitis.
- TPC.** (Tennessee Pharmaceutical) Phenylephrine HCl 5 mg., acetyl-p-aminophenol 200 mg., atropine sulfate 0.2 mg., pheniramine maleate 2 mg./Tab. Bot. 100s. Use: Common cold
- TPN.** (Hoack) L-Tyrosine 400 mg., pyridoxine HCl 2.5 mg., niacinamide 10 mg./Tab Bot 100s, 500s. Liq. Bot. 8 oz. Use: Antiallergic agent.
- TPNH.** (Sigma Chemical) Triphosphopyridine nucleotide: co-enzyme II, co-dehydrogenase II. Vials 0.3 mg., 0.6 mg., 1.0 mg., 5.0 mg., 100 mg., cartons of 10.
- TRABAS.** (Smith-Dorsey) Mannitol hexanitrate $\frac{1}{2}$ gr., phenobarbital $\frac{1}{4}$ gr./Tab. Bot 100s, 1000s. Use: (1 to 2 i.i.d.) Vasodilator w/sedative.
- TRACERAL.** (G. F. Harvey) Multivitamin w/minerals. Tab., Bot. 100s, 1000s. Use: Nutritional supplement.
- TRACERVIAL.** See: *Sodium Badio-ladide*, N N D. 1961.
- TRACIN TABLETS.** (Tracy) Tab. Bot. 12s
- TRACINETS.** (Merck Sharp & Dohme) Bac-tracin 50 u., tytharicin 1 mg., benzocaine 5 mg./troche. Vial 12s. Use: (1 q. 3 h.) Secondary infection of mouth & throat.
- TRACNE.** (Tablet Rock) Colloidal sulfur, phenylmercuric borate, resorcinol monoacetate, N trichloromethylmercapto-4 cyclohexene-1,2 dicarboximide/cream. Tube 1 oz. Use: Treatment of acne vulgaris.
- TRACO-DISCS.** (Neoca) Mag. trisulfate 3 gr., Cal. carb. 7 gr./disc. Bot. 100s. Use: Antacid.
- TRAL, FILMTABS, N N D. 1962** (Abbott) Hexacyclium methylsulfate. Tab. 25 mg., Bot 100s, 1000s. Use: Anticholinergic agent
- W/Ethylurea.** See: *Tralcyon, Filmtab* (Abbott)
- W/Phenobarbital** 15 mg Bot. 100s

- TRAL, GRADUMET, NND 1962** (Abbott) Hexocyclium methylsulfate 50 mg or 75 mg / Gradumet Bot 50s, 500s
Use: Anticholinergic agent
- TRAL INJECTABLE.** (Abbott) Hexocyclium methylsulfate 5 mg / cc Amp 10 mg / 2 cc 5s, 25s, 100s
Use: IV or IM Glaucoma or pyloric stenosis (anticholinergic)
- TRAL-PHENOBARBITAL GRADUMET.** (Abbott) Hexocyclium methylsulfate 50 mg, phenobarbital 30 mg / Tab Bot 50s
- TRAL WITH PHENOBARBITAL DROPS.** (Abbott) Hexocyclium methylsulfate 25 mg, phenobarbital 15 mg / 0.6 ml Bot 5 cc 12s
Use: Depressant and anticholinergic
- TRALCYON** (Abbott) Tral 25 mg, eptylurea 300 mg / film tab Bot 50s, 500s
Use: Management of gastrointestinal dysfunction caused or aggravated by anxiety or tension
- TRAMCINOLONE DIACETATE** See Aristocort Acetanide Cream (Lederle)
- TRANCIN** (Schering) Fluphenazine 0.25 mg / Tab
Use: Treat anxiety and tension
- TRANCOPAL,** (Winthrop Lab) 2 (4 Chlorophenyl) 3 methyl 4 methiazonone 1 dioxide 100 mg or 200 mg / Cap Bot 100s, 1000s
Use: Central relaxant and tranquilizer
- TRANCOPRIN,** (Winthrop) Chlorazepate 50 mg, acetylsalicylic acid 300 mg / Tab Bot 100s, 1000s
Use: Analgesic w/ muscle relaxant
- TRAND,** (Burlington) Methamphetamine 5 mg, Vit B 1 2 mg, B 2 3 mg, niacin amide 10 mg, Vit C 25 mg / Tab Bot 100s
Use: (1 2 tid) Obesity
- TRANE,** (Ciba) Tyrothricin 2 mg & benzoic acid 5 mg / loz Bot 100s, 500s, 1000s
Use: (1 q 3 h) Relief of minor throat irritations
- TRANIDON ELIXIR.** (Merck Sharp & Dohme) Hyoscyamine HBr 0.225 mg, atropine sulfate 0.019 mg, scopolamine HBr 0.006 mg, phenobarbital 15 mg / 5 cc Bot 1 pt, 1 gal
Use: (1 tsp q 3 to 4 h) Antispasmodic agent
- TRANITE D-LAY.** (Lloyd Dobney & Westfield) Pentaerythritol tetranitrate 30 mg / Cap Bot 100s
Use: Angina pectoris
- TRANQUILIZING AGENTS.** See: Atracetic agents
- TRANSED.** (Burlington) Diethylbarbituric acid 15 mg, d desoxyephedrine HCl 5 mg, thiamine chloride 2 mg, riboflavin 3 mg.
* Generic name, recognized by NF, USP or A.M.A.
- niacinamide 10 mg, ascorbic acid 25 mg / Tab Bot 100s, 1000s
Use: (1 tid) Emotional & mental distress
- TRANSIBARB** (Brean) d Desoxyephedrine HCl 25 mg, phenobarbital 1/4 gr, dl alpha tocopherol acetate 5 mg / Cap Cop, Bot 100s, 500s, 1000s
Uses: (1 tid, oc) Central nervous system stimulant w/sedative
- TRANSPOSE TABLETS** (Whittier) Mephenoxalone 5 (o methoxyphenoxymethyl) 2 oxazolidinone 400 mg / Tab Bot 50s
Use: Tranquilizer
- * TRANYLYCPROMINE SULFATE, NND 1962** Transyl 2-phenylcyclopropylomine
See Parnate (Smith, Kline & French)
- TRASENTINE,** (Ciba) Adiphenine HCl, di-phenylacetyldiethylaminoethanol HCl Amp (50 mg) 1.5 cc, Corlon 5s, 20s Tab (75 mg) Bot 100s, 500s Supp (100 mg) Box 5s
Use: (1 2 tid) Antispasmodic
W/Phenobarbital (Ciba) Trasentine 50 mg, phenobarbital 20 mg / Tab Bot 100s, 500s
Use: (1 2 tid) Antispasmodic w/sedative
W/Sod carboxymethylcellulose & Mag oxide See: Carmethose Trasentine, Tab (Ciba)
- TRAVAD ENEMA** (Flint, Eaton) Sodium dihydrogen phosphate 12 Gm, sod citrate 10 Gm / 100 ml Ready to use disposable enema unit 4 1/2 fl oz
Use: Cleansing of lower bowel
- TRAVAMIN, NND 1962** (Baxter) Protein hydrolysate (plasma) Sol (5%) Bot 150 250 500 & 1000 cc Sal (5% w/dextrose 5%) Bot 150 250, 500 & 1000 cc
Use: IV; protein therapy
- TRAVERT, NNR 1955** (Baxter) Invert sugar Sol (5%) water or saline 50 mg / cc, Bot 150, 1000 cc Sol (10%) water or saline 100 mg / cc, Bot 150, 500, 1000 cc
- TRAVLEX.** (Louisiana) Scopolamine aminoxide HBr 1 mg / 5 R Cop
Use: Treatment of motion sickness
- TREATOR.** (Ives-Cramer) * Ethionamide 250 mg / Tab Bot 100s, 500s
Use: Anti-tubercular agent
- TRELA.** (Kendall) Pentylene tetrazol 100 mg, niacin 50 mg, glutamic acid 200 mg / Tab or 5 cc liq Bot 8 oz, Tab Bot 100s, 1000s
Use: Geriatric therapy
- TREPARSOL.** (Wollau) Formyl A acetyl amides of m amino p-oxyphenylsuccinic acid Tab 0.25 Gm, Bot 100s
Use: Treatment of amebiasis

- TREPIDONE.** (Lederle) Mephenoxalone. 400 mg./Tab. Bot. 50s.
Use: Alaractic and muscle relaxant.
- TRESAMIDE.** (Merck Sharp & Dohme) Sulfamerazine 0.1 Gm., sulfadiazine 0.2 Gm., sulfathiazole 0.2 Gm./Tab. Bot. 100s, 1000s.
Use: Triple sulfonamide therapy.
- TRESANOIDS.** (Merck Sharp & Dohme) Tyrosine 1 mg., propadrine HCl 20 mg., benzocaine 15 mg., bismuth subgallate 150 mg., zinc oxide 150 mg./supp. Box 12s.
Use: Proctologic conditions.
- TRETAMINE.** See: Triethylene Melamine (Lederle)
- TRETHECANIC ACID.** 3-Hydroxy-3,7,11-trimethyl-dodecanoic acid.
- TRETHYLENE.** (Davis, Base) Trichloroethylene. Liq. Bot. 1 fl. oz.
Use: Volatile analgesic & anesthetic.
- TREVIDAL.** (Organon) Col. carb. 0.105 Gm., Mag. carb. 0.060 Gm., Mag. trisilicate 0.150 Gm., aluminum hydroxide 0.090 Gm., pepsin 0.100 Gm., & egraine 0.045 Gm./Tab. Bot. 100s. Liq. 1 sp. ls equiv. to 1 tab. Bot. 12 oz.
Use: Antacid.
- TREXINEST.** (Hynson, Wescott & Dunning) Equine estrogens 1 mg., lutrexin 250 u./Tab. Bot. 50s
Use: Estrogen therapy
- TREYHIST TABS. & ELIXIR.** (Meril) Thienpyramine HCl 10 mg.; chlorpheniramine mol. 2 mg.; pyrilamine mol. 10 mg./tab. or sp. Bot. 100s, 1000s Tabs. Bot. Pts., gals. Elixir.
Use: Antihistamine Comb.
- TRIACETIN.** N.N.D. 1962, U.S.P., A.D.R. Glyceril triacetate.
See: Enzadin, Cream (Ayers)
Fungocelin, Preps. (G. F. Harvey)
- TRIACETYLOLEANDOMYCIN.** N.F., N.N.D. 1962, Caps., N.F.
See: Cyclamycin, Prep. (Wyeth)
Signemycin, Prep. (Pfizer)
Tao, Cap. (Roerig)
W/Glucosamine. See: Tao, (Roerig)
W/Phenylpropanolamine HCl, pheniramine maleate, pyrilamine maleate, & calcium acetylsalicylate carbamide. See: Tain, Tab. (Smith-Dorsey)
W/Sulfa meth-dia-mer. See: Toamid, Preps (Roerig)
W/Triaminic, acetaminaphen. See: Tain Oral Suspension (Dorsey)
- TRIACETYL PYROGALLOL.** N.N.R. 1949.
See: Lenigallol, Prep. (Bilhuber-Knofl).
- TRIALDIN.** (Rorer) Belladonna ext. 5 mg., Mag. trisilicate 0.3 Gm., aluminum hydroxide gel 0.15 Gm./Tab. Bot. 100s, 500s, 1000s.
Use: (1-2 q. 3-4 h.) Gastric hyperacidity
- TRIALLER.** (Lemor) Phenylpropanolamine HCl 50 mg., pyrilamine maleate 25 mg., pheniramine maleate 25 mg./tab. Bot. 100s, 500s.
Expect. 1 pt., 1 gal., Pediatric Drops 15 cc. 3yr. 1 pt.
Use: Antihistamine-decongestant
- *TRIAMCINOLONE.** N.N.D. 1962 9-Alpha fluoro-16 alpha hydroxyprednisolone.
See: Aristocort, Tab (Lederle)
Kenacort, Tab. (Squibb)
W/Salicylamide, aluminum hydroxide, ascorbic acid. See: Aristogestic, Cap. (Lederle)
W/Chlorpheniramine maleate & ascorbic acid. See: Aristamin, Cap (Lederle)
- *TRIAMCINOLONE ACETONIDE.** U.S.P., N.N.D. 1962; Lol, U.S.P. 9-Alpha fluoro-16-alpha 17 alpha isopropylidenedioxy Δ^1 hydrocortisone
See: Aristocort Acetonide, Cr. & Oint (Lederle)
Aristoderm Foam (Lederle)
Kenalog Inj. & Spray (Squibb)
W/Benzyl alcohol See: Vetalog, Susp. (Squibb)
W/Neomycin base, gramicidin See: Kenalog-5, Preps. (Squibb)
W/Neomycin sulfate, methylparaben, propylparaben. See: Neo-Aristoderm Foam, Aerosol (Lederle)
W/Neomycin sulfate, light liquid petrolatum, wool fat, white petrolatum. See: Neo Aristocort Acetonide Eye-Ear Ointment. (Lederle)
W/Neomycin, gramicidin, Nystatin. See: Mycolog, Preps (Squibb)
- *TRIAMCINOLONE DIACETATE.** 9-Alpha-fluoro-16-alpha hydroxy prednisolone diacetate
See: Aristocort Diacetate, Preps. (Lederle)
Aristocort Diacetate Forte (Lederle)
- TRJAMEL CREAM & OINT.** (Fuller) Hamamelis Water, 20%; with Vit. A and D pH is about 4.6. Tubes 2 oz.
Use: Dermatologic.
- TRJAMEL-HC.** (Fuller) Hydracortisone acetate 0.5%, hamamelis water 20.0%, Vit. A and D in a special water-washable bland base. Tube 10 Gm.
Use: Combination entipruritic, antiinflammatory.
- TRIAMINIC.** (Dorsey) Pyrilamine maleate 25 mg., prophepyridamine maleate 25 mg., phenylpropanolamine HCl 50 mg./Timed-release tab Bot. 50s, 250 cc.
Use: Antihistaminic
W/Calcium acetylsalicylate carbamide. See: Ursinus, Tab. (Dorsey)
- *Generic name recognized by N.F., U.S.P. or A.M.A.

W/Dormethan, terpin hydrate See Tussaminic Tab (Dorsey)

TRIAMINIC CONCENTRATE. (Dorsey) Phenylpropanolamine HCl 20 mg, pheniramine maleate 10 mg, pyrilamine maleate 10 mg/ml Dropper bot 15 cc

Use Decongestant and antihistaminic action

TRIAMINIC-HC. (Dorsey) Phenylpropanolamine HCl 25 mg pheniramine maleate 12.5 mg pyrilamine maleate 12.5 mg hydrocortisone 10 mg/Tab Bot 30s

Use Allergic disorders

TRIAMINIC JUVELETS (Dorsey) Phenylpropanolamine HCl 25 mg pyrilamine maleate 12.5 mg, pheniramine maleate 12.5 mg/Tab Bot 50s

Use: Common cold

TRIAMINIC SYRUP (Dorsey) Phenylpropanolamine HCl 12.5 mg pyrilamine maleate 6.25 mg pheniramine maleate 6.25 mg/5 cc Bot 1 pt 1 gal

Use Colds and allergies

TRIAMINICIN (Dorsey) Phenylpropanolamine HCl 25 mg pyrilamine maleate 12.5 mg pheniramine maleate 12.5 mg aspirin 225 mg phenacetin 150 mg caffeine 30 mg VI C 50 mg aluminum hydroxide gel dried 180 mg/Tab Bot 100s 500s

Use Treatment of the common cold

TRIAMINICIN NASAL SPRAY (Dorsey) Phenylpropanolamine HCl 0.75% phenylephrine HCl 0.25% pheniramine maleate 0.125% pyrilamine maleate 0.125% benzalkonium Cl 1:10000/35 cc Spray Bot 15 cc

Use Nasal decongestant

TRIAMINICOL (Dorsey) Phenylpropanolamine HCl 12.5 mg pyrilamine maleate 6.25 mg pheniramine maleate 6.25 mg Dormethan 15 mg ammonium chloride 90 mg/5 cc (cough syr) Bot 3 oz, 1 pt 1 gal

Use Cough depressant

TRIAMINILONE-16,17-ACETONIDE. See Triaminalone acetamide

***TRIAMTERENE** (Smith Kline & French) 2.47 triamino 6 phenylpteridine

Use Diuretic

TRIANATE See Aminatrate Phosphate

TRITROPHENE. (Lannett) Magnesium trisilicate 7½ gr phenobarbital ½ gr, atropine sulfate 1/1000 gr/Tab Bot 100s

Use: GI sedative

TRIAZIL. (Central) Susp: (Sulfadiazine 1 Gm sulfamerazine 1 Gm sulfacetamide 1 Gm Sod lactate 3 Gm/oz) Bot 1 pt

Palatabl: (Sulfadiazine 0.083 Gm, sulfamerazine 0.083 Gm, sulfacetamide 0.083 Gm/100s, 1000s)

Tab: (Sulfadiazine 0.166 Gm, sulfamerazine 0.166 Gm, sulfacetamide 0.166 Gm/Tab) Bot 100s, 500s

Use: Triple sulfanamide therapy.

TRIAZINE. (US Standard) Sulfadiazine 0.1 Gm, sulfamerazine 0.17 Gm, sulfamethazine 0.17 Gm/Tab Bot. 100s, 1000s

Use: Sulfanamide therapy

TRIAZOLINE DULCET TABLETS. (Abbott) Sulfadiazine 0.1 Gm, sulfamerazine 0.1 Gm, sulfathiazole 0.1 Gm/Tab Bot 100s

Use: Sulfanamide therapy

TRIAZOLINE, TABLETS. (Abbott) Sulfadiazine 0.167 Gm, sulfamerazine 0.167 Gm, sulfathiazole 0.167 Gm/Tab Bot 100s 1000s

Use: Sulfanamide therapy

TRI-AZO-MUL. (First Texas) Meth diamer sulfanamides, sulfadiazine 3.38 Gm, sulfamerazine 3.381 Gm, sulfamethazine 3.38 Gm/100 cc Susp, plain or citrated Bot 1 pt

Use Sulfanamide therapy

TRI-AZO-TABS (First Texas) Meth diamer sulfanamides sulfadiazine 2.57 gr, sulfamerazine 2.57 gr sulfamethazine 2.57 gr/Tab Bot 100s, 500s 1000s

Use: Sulfanamide therapy

TRIBACT (Oermatologicals) Polymyxin B sulfate 5000 u bacitracin 400 u, neomycin sulfate 5 mg/Gm Oint

Use: Dermatologic

***TRIBROMOETHANOL, US P,** Sol, USP NNR 1957 Tribromoethyl alcohol, USP See Avertin, Sol (Winthrop Labs)

TRIBROMOETHYL ALCONOL, NNR 1957 See: Tribromoethanol, USP

TRIBROMOMETNANE. See Bromofarm Prep (Various Mfr)

TRIBROMOPNEUL-BISMUTH W/Xeroform See: Xeroform Ophthalmic Ointment (Mecol)

TRIBURAN CHLORIDE, NND 1962 (Roche Lab) Tribromsanium chloride NN bis(1-methyl 3 (2,2,6-trimethylcyclohexyl)propyl) NN dimethyl 1,6 hexanediamine bis (methochloride) hemihydrate Triburan Ointment 0.1% Tube 1 oz, jar 1 lb, Vag Cream (0.1%) Tube 3 oz w/applicator Vag Supp Box 25s

Use: Primary and secondary skin and wound infections

TRIBURAN-NC. (Roche Lab) Oint: Triburan 0.1% hydrocortisone 0.5% Tube 5 and 20 Gm Cream: Triburan 0.1%, hydrocortisone 1% Tube 5 and 15 Gm

Use Topical microbicide

*Generic name recognized by NF USP or AMA

- TRIS VAGINAL SUPPOS. as CREAM.** (Roche) Triclobisaniol chloride 0.1% / Suppos. With reusable applicator. 24s. Cream; 3 os w/18 applic.
Use: Vaginal and cervical infection.
- TRICAINAL.** (Ciba)
Oint. (Pyrbenzamine HCl 0.5 Gm., Nupercaine base 0.125 Gm., w/zlec oxide, bismuth subgallate, lanolin, petrolatum, acetone Sod. bisulfite/100 Gm.) Tube 1 oz.
Supp. (Pyrbenzamine HCl 10 mg., Nupercaine base 2.5 mg., acetone Sod. bisulfite 1 mg., bismuth subgallate 100 mg., zlec oxide 250 mg./supp.) Box 12s.
Use: Antipruritic & anesthetic.
- TRI-CALSATE.** (Paxton) Tri-Cal. phosphate 35.5%, tri-Sod. phosphate 12.5%, Sod. citrate 52%. Aq. susp., Bot. 4½ oz., 18 oz.
Use: Antacid.
- TRI-CARB-CREAM.** See: Cremo-Carboeatin (Merck Sharp & Dohme)
- TRICARRS.** (Roemer) Mag. trisilicate, Mag. carb., Cal. carb. Tab., Bot. 100s, 500s, 1000s.
Use: Antacid.
- TRICETAMIDE.** (Riker) 3,4,5-trimethoxybenzoylglycine diethylamide.
Use: Sedative.
- TRICHINELLA EXTRACT, N.N.R. 1949.** (Led-arts) Vial 1 cc.
Use: Diagnosis of trichinosis.
- TRICHLORISONIUM.** See: Triburan, (Roche)
- TRICHLORAN.** See: Trichloroethylene.
- TRICHLORMETHIAZIDE.** 3-Ochloromethyl-6-chloro-7-sulfamyl-3,4-dihydro-1,2,4-benzothiazine-1,1-dioxide.
See: Flutra, Tabs. (Schering)
Methhydrin, Tab. (Lakeside)
Nagua, Tab. (Schering)
Naquival (Schering)
W/Reserpine. See: Naquival (Schering)
- TRICHLOROACETIC ACID OINTMENT.** See: Eikon. (Merck Sharp & Dohme)
- TRICHLORORUTYL ALCOHOL.** See: Chlorobutanol
- TRICHLOROCARBANILIOE.** (Texas) W/Salicylic acid, sulfur. Medet Shampoo/Cleaner.
- TRICHLOROETHYLENE, U.S.P., N.N.R. 1962.** A.D.R. Trichloroethene, trichloran, trichloroethylene, U.S.P.
See: Chlorlylen (Schering)
Triethylene. (Davis, Rose)
Trilene, N.N.R. (Ayers).
Trimar (Ohio Chemical).
- TRICHOJEL.** (Reckel) 9-Aminoacridin propionate 0.35%. Jelly, Tube 4 oz.
Use: Vaginal antiseptic.
- TRICHOLINE CITRATE.** See: Choline citrate.
- TRICHO RAD.** See: Trilheon (Ortho)
- TRICHOTINE.** (Fesler) Sod. lauryl sulf., sod. perborate, sod. borate, thymol, eucalyptol, menthol, methyl salicylate. Powder, Pkg. 5, 12, 20 oz.
Use: 2 tablespoonful dissolved in a quart of warm water, vaginal douche
- TRICHOIN.** (Boyle) Neomycin sulf. 3.5 mg., gramicidin 0.25 mg., polymixin B sulf. 2000 u., benzocaine 1%/Gm. Oint Tube ½ oz.
Use: Topically, antibiotic and anesthetic ointment
- TRICILLIN.** (Evran)
No. 1 Pow. Sulfadiazine, sulfamerazine, sulfacetamide, of each 167 mg. w/Pat penicillin G 100M u./5 cc. when water is added Rot. 60 sc.
No. 2 Pow. Same triple sulfa w/Pat. penicillin G 250M u./5 cc. Rot. 60 cc.
No. 1 Tab. Same as No. 1 Pow./tab. Bot 100s, 500s, 1000s.
No. 2 Tab. Same as No. 2 Pow./tab Rot 100s, 500s, 1000s.
- *TRICLOSINIUM CHLORIDE, NND 1962** See Triburan, oint. (Roche Labs)
W/Hydrocortisone See Triburan-HC Ointment (Roche Labs)
- *TRICLOFENOL PIPERAZATE.** (Parke, Davis) Piperazine di(2,4,5-trichlorophenoxy).
Use: Anthelmintic (nematodes).
- TRICLOFOS.** Name approved by R.P.C. 222. Trichlorosthyl dihydrogen phosphate trichloryl.
- TRICOBETA.** (Cale Chem. Co) Sulfathiazole 5 gr., beta lactose 15 gr./tab. Vaginal Tab., Bot. 100s.
Use: Antibacterial.
- TRICOFURON IMPROVED.** (Eaton) Powder Furoxone Powder 0.1% and Mirafur 0.5%. Pkg 15 Gm. in squeeze bot. w/3 tips Vag supp. Furoxone 0.25% and Micalor 0.375%. Pkg 12s w/applicator 12s, 24s
Use: Vaginal trichomoniasis, menitis, haemophilus vaginalis vaginitis
- TRICLOFID CHLORIDE, NND 1962.** (Burroughs Wellcome) Tricyclamal Tab. 50 mg. Rot 100s, 1000s.
Use: [I.I.d.] Management of peptic ulcers.
- W/Phenobarbital.** (Burroughs Wellcome) Phenobarbital 16 mg., tricyclamal chloride 50 mg./Tab Rot 100s, 1000s.
Use: Nervous indigestion & peptic ulcer.
- TRICOMER SULFONAMIDE** Sulfacetamide 0.166 Gm., sulfadiazine 0.166 Gm., sulfamerazine 0.166 Gm./Tab. or 4 cc. Liq., Bot. 16 oz. Tab., Bot. 100s, 1000s
Use: Triple sulfonamide therapy.
- TRI-COSTIVIN.** (Professional Labs) Guggula, karaya gum, lactose & Vit. B complex.

Granule Pkg 160 Gm, 280 Gm, 600 Gm

Use: Laxative

TRI-CREAMALATE. (Winthrop Labs) {Aluminum hydroxide gel 300 mg, Mag trisilicate 600 mg /S cc} {Aluminum hydroxide gel 324 mg, mag trisilicate 162 mg /tab} Liq; Bot 8 12 fl oz

Tab; Bot 100s, 500s

Use: Antacid

***TRICYCLAMOL CHLORIDE, N N D 1962 1** Cyclohexyl 1 phenyl 3 pyrrolidone 1 propanol methyl chloride Lergina

See: Elorine Pulverul, NNR (Lilly)

Tricoloid, Tab (Burroughs Wellcome)

W/Phenobarbital See: Tricoloid W/Phenobarbital, Tab (Burroughs Wellcome)

***TRICYCLATE NCI 1** methyl 3 pyrrolidyl methyl benzilate HCl

TRIDAL (Lakeside) Dactil 50 mg, pipital 5 mg /tab Bot 50s

Use: GI antispasmodic

TRIDECAMINE (Merrell) Doxylamine succinate 20 mg pyrilamine maleate 20 mg pheniramine maleate 20 mg /Dorpon Tab Bot 50s 100s

Use: Allergic disorders

TRIDENEX LIQUID (Spiri) Olive oil made dispersible with acylphenoxxyethanol Nos 3% hexochlorophane added Bot 8 oz, qts gals

Use: Itching and/or burning after bathing

TRIDENOL (Spiri) Olive oil bath additive made completely dispersible with 8 phenoxxyethanol Plastic Bat 8 oz

Use: To relieve dryness itching scaliness hyperkeratotic discomfort in atopic dermatitis, etc

TRIENZYME (Testagar) Amylolytic 30 mg, proteolytic 10 mg cellulolytic 3 mg /Tab Bot 30s 100s 1000s

Use: Relief of functional digestive disorders

TRIDERMOL (Central) Sulfadiazine 1.67%, sulfamerazine 1.67%, sulfathiazole 1.67%, urethane 2%, sod alkyl benzene sulfonate 0.1% Cream 1/4 lb, 1 lb

Use: Topically; anti-infective

TRIDESTRIN See: Estril, Cap

TRIDEX TAB, TIMED TRIDEX CAP, TIMED

TRIDEX JR. CAP (Testagar) Changed to Doro Tab, Doro Timed Cap, Doro Jr Timed Cap

TRI-DEX. (Testagar) Dextro-amphetamine HCl Tab 5 mg., Bot 100s, 1000s

Use: Central nervous system stimulant

***TRIDIHETNYL CHLORIDE, N F, N N D 1962;** Tabs NF 3 Diethylamino 1 phenyl 1 cyclohexyl 1 propanol ethochloride See: Pathlan chloride Tab (Lederle)

W/Mepramate

*Generic name recognized by N.F., U.S.P. or A.M.A.

See: Milpath, Tab (Wallace)

Pathibamate, Tab (Lederle)

W/Propylparaben, methylparaben See:

Pathlan Pediatric Drops (Lederle)

TRIDILATIN. (Columbus) Nitroglycerin 1/200 gr, Sod nitrite 1/2 gr, pentaerythritol tetra nitrate 10 mg /tab Bot 100s, 1000s

Use: 1 prn; vasodilator in hypertension

TRIDIONE, N N D 1962 (Abbott) Trimethadione (Traxidone) Cap 0.3 Gm, Bot 100s 1000s Dulcet Tab 0.15 Gm, Bot 100s Sol 0.15 Gm /fl oz, Bot 1 pt, 1 gal

Use: (0.3 Gm to 2 Gm daily) Epilepsy

TRIDIPIGEN, ALUM PRECIPITATED, N N R 1955 (Lilly) Diphtheria & tetanus toxoids & pertussis vaccine combined, alum precipitated Vial 1.5, 7.5 cc

Use: IM 0.5 cc every 4-6 wks for 4 doses; active immunization

TRIDIPIGEN, FLUID (Lilly) Diphtheria tetanus toxoids pertussis vaccine combined, fluid Vial 1.5, 7.5 cc

Use: IM 0.5 cc every 4-6 wks for 3 doses; active immunization

TRIDIUM (Amco) Phenylazo diamino pyridine HCl 50 mg, sulfacetamide 250 mg atropine methyl nitrate 1 mg /Tab Bot 100s

Use: Treatment of urinary infections

TRIDROXAL, (Warren Teed) Mag trisilicate 80 gr, aluminum hydroxide gel 196 gr /fl oz Bot 3 fl oz, 1 pt, 1 gal

Use: Antacid

TRIENZYME, (Testagar) Amylolytic enzyme 30 mg, proteolytic enzyme 10 mg, cellulolytic enzyme 3 mg /Tab Bot 100s

Use: Digestive aid

TRIETHANOLAMINE SALICYLATE. See: Myoflex, Cream (Warren Teed)

TRIETHANOLAMINE TRINITRATE BIPHOSPHATE Trinitrate Phosphate

See: Metamine, Tab (Leeming)

Nigina (Haag)

Nitrelamine Phosphate, Tab (Squibb)

W/Butobarbital See: Metamine W/Butobarbital, Tab (Leeming)

***TRIETHYLENEMELAMINE, U.S.P., N N D 1962** (Lederle) (Tretamine) (TEM) (2,4,6-tris ethylenimino) S triazine 5 mg /Tab Bot 12s

Use: (Initial dose 2.5 mg) Antileukemic

***TRIFLUOPERAZINE NCI, N N D 1962 2** Trifluoromethyl 10 3 (1" methylpiperazinyl) propyl phenothiazine dihydrochloride

See: Stelazine (Smith, Kline & French)

TRIFLUOROPYRAZINE DIHYDROCHLORIDE. See: Stelazine (Smith, Kline & French)

***TRIFLUPROMAZINE NCI, N N D 1962 10-**

{3-dimethylaminopropyl}-2-(trifluoromethyl) phenothiazine HCl. See: Vesprin, Prep. (Squibb)

*TRIFLUTAMINE HCl. (Smith, Kline & French) 1-(4'-trifluoromethylphenyl)-2-aminopropane HCl.

Use Anorectic.

TRIFOCILIN. (VanPelt & Brown) Pot. penicillin G 100,000 u., sulfadiazine 0.167 Gm., sulfamerazine 0.167 Gm., sulfamethazine 0.167 Gm./5 cc. when water is added. Bot. 2 oz.

Use: Penicillin w/triple sulfonamide therapy.

TRIFONACIL-250. (Physicians Products) Buffered penicillin G potassium 250,000 u., sulfa meth-dia-mer each 0.167 Gm./Tab. or 5 cc. Tabs. 50s.

Use Penicillin-sulfa therapy.

TRIFONAMIDE, N.N.R. 1955 (VanPelt & Brown) (Meth-Dia-Mer-Sulfonamide) Sulfadiazine 0.166 Gm., sulfamerazine 0.166 Gm., sulfamethazine 0.166 Gm./Tab. or 5 cc. or susp. Susp., Bot. 12 fl. oz. Tab., Bot. 100s, 500s, 1000s.

Use: Triple sulfonamide therapy.

TRIGELMA. (Buffington) Liq. Bot. 12 oz. (Aluminum hydroxide gel q.s. & Mag. trisilicate 4 Gm./oz.) Tab. Bot. 100s, 1000s. (Aluminum hydroxide 2½ gr., Mag. trisilicate 5 gr./ea)

Use Antacid.

W/Mesamine. (Buffington) Scopolamine methylbromide 2.5 mg., magnesium trisilicate 150 mg., aluminum hydroxide gel 300 mg./tab. Bot. 100s.

Use Anticholinergic w/antacid.

TRIGELMA, H.M. (Buffington) Liq. Bot. 12 oz. (Homatropine methylbromide 1/12 gr., Mag. trisilicate 60 gr., conc. aluminum hydroxide gel q.s./oz.) Tab. Bot. 100s. (Homatropine methylbromide 1/48 gr., Mag. trisilicate 2½ gr., aluminum hydroxide q.s./tab)

Use Adrenergic & antispasmodic.

TRIGELMA W/MESCOMINE. (Buffington) Mesamine (scopolamine methylbromide) 2.5 mg., magnesium trisilicate 150 mg., aluminum hydroxide gel 300 mg./tab. Bot. 100s.

Use: (1) 1 d., a.c.) Treatment of peptic ulcer.

TRIGESIC. (Squibb) Acetyl-p-aminophenol 2 gr., aspirin 3½ gr., caffeine ½ gr./Tab. Bot. 100s, 1000s.

Use: Analgesic.

TRIGESIC W/CODEINE. (Squibb) Same analgesic comp. as Trigesic, w/codeine phosphate ½ gr., ¼ gr., ½ gr., 1 gr. added/Tab. Bot. 100s, 1000s.

Use: Severe pain.

*Generic name recognized by N.F., U.S.P. or A.M.A.

TRIGESTRANOROL. (Kirk) Progesterone 25 mg., testosterone propionate 25 mg., estradiol 1 mg. in Oil or Aq. Susp./cc. Vial 10 cc.

Use: I.M., estrogen & corpus luteum therapy.

*TRIHEXTPHENILOYL HTOROCHLORIDE, U.S.P., N.N.D. 1962, Tabs. U.S.P. Alpha-phenyl-1-cyclohexyl-3-piperidyl-1-propanol HCl

See: Artane, Elixir & Tab., N.N.R. (Lederle)

Pipanol, Elixir & Tab. (Winthrop Labs)

TRIHISTA-COO. (Recsel) Dihydrocodeinone bitartrate 1.33 mg., tri-histin 10 mg., pot. guaicol sulfanate 60 mg., NH₄Cl 80 mg., sod. citrate 100 mg., chloroform 15 mg./4 cc. Bot. 4 oz.

Use: Bronchial irritation, excessive cough or difficult expectoration.

TRI-HISTIN. (Recsel) Chloroethenylpyramine citrate 4%, pyrimidine maleate 34%, thienylpyrimidine 26%. Expectoant, Bot. 1 pt., 1 gal. liq. 10 mg./dr., Bot. 1 pt., 1 gal. Tab. 25 mg., 50 mg., Bot. 100s, 500s, 1000s.

Use: Triple antihistamine therapy.

W/Benzyl alcohol, chlorobutanol & isopropyl alcohol.

See: Derma-Pax, liq. (Recsel)

W/Dihydrocodeinone, pot. guaicol sulfanate, NH₄Cl, sod. citrate, chloroform. See: Trihista-Cod, liq. (Recsel)

W/Ephedrine HCl, aminophyllin, mephobarbital. See: Asmanol, Tab (Recsel)

W/Kl, ephedrine HCl, mephobarbital, aminophyllin. See: Neo-Asmanol, Cap. (Recsel)

TRIHORM. (Ascher) Estradiol 1.2 mg., progesterone 25 mg., testosterone 25 mg./cc. Aq. Susp. Vial 10 cc.

Use: Functional uterine bleeding.

TRI-HORMOGEN. (Mollard) Estradiol 1.2 mg., progesterone 25 mg., testosterone 25 mg./cc. Vial 10 cc.

Use: I.M., estrogen & androgen therapy.

TRIHORMONAL. (Key Corp.) Estradiol 0.05 mg., progesterone 0.5 mg., methyltestosterone 0.5 mg./Tab. Bot. 100s. Inj.: Progesterone 25 mg., testosterone propionate 25 mg., estradiol benzoate 1 mg. In sesame oil/cc. Vial 10 cc.

TRIHIDROXYESTRADIENE. See: Estriol.

TRIHIDROXYESTRINE. Trihydroxyestrin

See: Estriol, Cap. (Abbott)

TRIHIDROXYETHYLAMINE. Triethanolamine.

TRI-IMMUNOL. (Lederle) Diphtheria & tetanus toxoids & pertussis vaccine combined, aluminum phosphate-adsorbed purigenated

Vial I.S., 7.5 cc.

Use: I.M., 0.5 cc. every 4-6 wks. for 3 doses; active immunization.

- TRIIODOETHIONIC ACID.** See: Teridax, Tab (Schering)
- TRIHODOMETHANE, A D B** See: Iodoform NF (Various Mfr)
- L-TRIIODOTHYRONINE SOD.**
See Cytamel, Tab (Smith, Kline & French)
Liothyronine Sod NND 1958
- di-TRIIODOTRYONINE SODIUM** See: Tri none, Tab (Roche)
- 3,7,12-TRIKETOCHOLANIC ACID** See De hydrocholic Acid (Various Mts)
- TRIKETOL.** (Endo) Dehydrocholic (Ketocholanic) acid obtained from bile Pow., Bot 1 oz., 4 oz Tab 4 gr Bot 100s 1000s
Use: (1 2 fl d, p c) Choleric & hydro choleric
- TRILAFON, NND 1962** (Schering) Per phenazine Tab 2 4 8 mg., & 16 mg for hospital use Bot 50s Inj (5 mg/cc) 1 cc, 10 cc Box 6s Syr (2 mg/5 cc) Bot 4 oz Repeats 8 mg Bot 30s, 100s Supp (2, 4 8 mg) Box 6s Concentrate 16 mg/5 cc Bot 4 oz w/drapper
Use: Tranquilizer
W/Prednisone See Steratril Tab (Schering)
- TRILAMINE,** (Lemmon) Pyrilamine maleate 15 mg, phenylpyramine HCl 15 mg, phenyl tolaxamine dihydrogen citrate 15 mg/Tab Bot 100s available in double strength
Use: Treatment of allergic states
- TRILAPHEN** (Lonnell) Magnesium trisilicate 7½ gr, phenobarbital ½ gr, atropine sulf 1/1000 gr/Tab Bot 1000s, 5000s
Use: Antacid sedative
- TRILCARBS,** (G F Harvey) Mag trisilicate 3 gr, mag carb 3 gr, cal carb 4 gr/Tab Bot 1000s, 5000s
Use: Antacid
- TRILENE, NND 1962 A D B** (Ayerst) Tri chloroethylene Tubes 15 cc Contains 300 cc
Use: Inhalation analgesia In obstetrics & surgery
- TRILOX.** (Blue Line) Susp, Mag trisilicate 75 gr, aluminum hydroxide gel 90 gr./8 oz Bot 1 pt, 1 gal Tab, Mag trisilicate 10 gr, aluminum hydroxide gel, dried 3 gr./Tab Bot 100s, 500s
Use: Gastric antacid
- TRILUTE.** (Paythress) Trocinate 100 mg, the ophyline 100 mg, pyrilamine maleate 25 mg/Cop Bot 25s, 100s
Use: Dysmenorrhea and premenstrual tension
- TRIMACILL.** (Massengill) Tartaric acid citric acid dextrose potassium bitartrate, potassium alum adhesive Pow., Bot 5 oz, Vag nal inserts Boxes 24s
Use Vag nal inserts
- * Generic name recognized by NF USP or A.M.A.
- TRIMAGNOL** (Patch) Mag trisilicate 7.5 gr ascorbic acid 65 mg/Tab Bot 100s 1000s
Use: Antacid
- TRIMAS, A D R** (Ohio Chemical) Trichlorethylene Bot 1 pt
Use Inhalation anesthetic & analgesic
- TRIMAX** (Winthrop Labs) Magnesium trisilicate pow Bot 1½ oz
Use (1 tsp) Hyperacidity
- * **TRIMAZINOL** (Irwin Neisler) o-[[[4,6 diamino s triazin 2 yl]amino] methyl]benzyl alcohol
Use Anti-inflammatory agent
- * **TRIMEPRAZINE TARTRATE, NND 1962** dl 10 (3-Dimethylamino 2 methylpropyl) phenothiazine See Temarit, Tab (Smith, Kline & French)
W/Acetaminophen, phenylpropranolamine HCl See: Coplexen Liquid (Smith Kline & French)
- TRIMERAZOLE,** (Maffard) 0.166 Gm each sulfamerazine, sulfathiazole & sulfadiazole/Tab Bot 100s 500s, 1000s
Use: Triple sulfanamide therapy
- TRIMESTER,** (Haug Drug Co) Cal carbonate 450 mg, hesperidin 50 mg, vit C 100 mg B 1 15 mg, B 2 2 mg, niacinamide 15 mg A 6000 u, D 400 u/3 caps Bot 100s, 500s
Use Prenatal therapy
- TRIMETAPHAN** See: Trimethaphan
- * **TRIMETHADIONE, USP, NND 1962;** Cop USP, Sol NF Tab NF 3,3,5 Tri methyl 2,4 oxazolinedione See: Tridione NNR (Abbott)
- * **TRIMETHAMIDE** (Lederle Labs) N(2 amino 6 methyl 3 pyridylmethyl) 3,4,5 trl methoxybenzamide
Use: Hypotensive agent
- * **TRIMETHAPHAN CAMPHORSULFONATE, USP, NND 1962;** Inj USP d 3,4-(1',3'-d benzyl 2' keto midazolido) 1,2 trimethyl enethiophanium d camphorsulfonate See Arfanod Camphorsulfonate Amp (Roche)
- TRIMETHAZINE,** (Narmand) Sulfadiazine D 167 Gm, sulfamerazine D 167 Gm, sulfa methazine D 167 Gm/Tab Bot 500s, 1000s
- * **TRIMETHIDIUM METHOSULFATE, NND 1962** d (N methyl N (gamma trimethyl am moniumpropyl)) 1-methyl B 8-di methyl-3-azabicyclo (3 2 1) octane dimethosulfate See: Ostensin (Wyeth)
- * **TRIMETHOBENZAMIDE HCl, NND 1962** 4 (2 D methylaminoethoxy)N (4,5 trimethoxybenzoyl) benzylamine HCl See: Tigon (Roche Labs)
W/Beta Pyridylcarbinol nicotiny alcohol (Banacol) See: Tigocal Caps (Roche)

- *TRIMETHOPRIM.** (Burroughs Wellcome) 2, 4 diamino-5 (3',4',5'-trimethoxybenzyl)pyrimidine.
Use: Antibacterial agent.
- TRIMETHYLENE, N.N.D. 1961.** See: Cyclopropane, U.S.P. (Mallinckrodt, Squibb)
- TRIMETHYL XANTHINE.** W/N-acetyl-p-aminophenol, acetyl salicylic acid, chlorpheniramine maleate, phenylephrine HCL. See: Thoresin Cold Capsules. (Gillette Labs)
- TRIMETON, N.N.R. 1953.** (Schering) Prophepridamine. Tab. 25 mg., Bot. 100s, 1000s.
Use: Antihistaminic.
- TRIMETON MALEATE, N.N.D. 1962.** (Schering) [Prophepridamine Maleate] 1-Phenyl-1-(2-pyridyl)-3-dimethyl-amino-propane maleate.
Elixir (7.5 mg / 4 cc.) Bot. 4 oz., 1 pt.
Tabl. (25 mg) Bot. 100s, 1000s.
Use: Antihistaminic.
- TRIMETOSE.** (Schering) Prophepridamine maleate 7.5 mg., ammonium chloride 104.8 mg, Sod. citrate 43 mg., chloroform 17.5 mg / dr. Bot. 1 pt., 1 gal.
Use: (i-2 tsp. i.i.d.) Coughs.
- TRIMETIS.** (Bell Pharmacol) Mephenesin 500 mg., physostigmine salicylate 0.5 mg., homatropine methylbromide 1.2 mg / tab. Bot. 100s, 1000s.
Use: Muscle relaxant.
- TRIMIXIN.** (Mance) Bacitracin 200 u., polymixin B sulfate 4000 u., neomycin sulfate 3 mg./Gm. Oint., Tube 1/2 oz.
Use: Topically, antibiatic therapy
- TRINALIS.** (Webster) 9-Aminoacridine HCl 2 mg., phenylmercuric acetate 3 mg., tyrothricin 1 mg., urea 20 mg., lactate 100 mg., sod lauryl sulf. 5 mg./vag. supp. Jars 14s.
Use: Treatment of vaginal infections.
- TRINAVAC, N.N.R. 1954.** (Merck Sharp & Dohme) Diphtheria & tetanus toxoids w/ pertussis vaccine combined, alum precipitated, N.N.R. H. pertussis 30,000 million/cc. w/diphtheria & tetanus toxoids, Vial 1.5 cc., 7.5 cc.
Use: Immunization against diphtheria, tetanus & whooping cough.
- TRINESIUM.** (Abbott) Mag. Trisilicate. Tab. 7 1/2 gr., Bot. 100s, 500s.
Use: Adaptive & acid-neutralizing.
- TRINFAGEN VACCINE.** (Pitman-Moore) Multiple antigen fluid vaccine containing diphtheria toxoid, tetanus toxoid and pertussis toxin vaccine. Vials 5 cc
Use: Recall injections in children over 5 yrs who have been basically immunized against D.P.T. & Polio.
- TRINIDEX.** (Don Baxter) Dextrose 5 or 10% w/nicotinamide 100 mg., riboflavin 20 mg., thiamine HCl 10 mg./L. Dispensing container 1000 cc.
Use: I.V., to correct tissue & circulating fluid loss
- TRINITRIN TABLETS.** See: Glyceryl Trinitrate Tablets, U.S.P.
- TRINITROPHENOL, 2,4,6-Trinitrophenol.** See: Picric Acid
- TRINOLYTE.** (Don Baxter) Dextrose 50 Gm., ethyl alcohol 25 cc., sod. lactate 5.3 Gm., pot. chloride 0.75 Gm., sod chloride 5 Gm., citric acid 0.34 Gm., cal. chloride 0.35 Gm., mag. chloride 0.31 Gm., nicotinamide 100 mg., Vit. B-1 10 mg., B-2 20 mg./1000 cc. Vascular 1000 cc.
Use: A balanced electrolyte solution with dextrose, vitamins, and alcohol
- TRINSICON.** (Lilly) Liver-stomach conc. 150 mg., Intrinsic factor 1/2 u., Vit. B-12 7.5 mcg., ferrous sulfate, anhy., 300 mg., ascorbic acid 75 mg., folic acid 1 mg./pulvule. Bot. 60s, 500s.
Use: Hemostatic in anemias.
- TRINSOCIN-M.** (Lilly) Identical in formulation to Trinsocin except for the elimination of folic acid Bot 60s
- TRIOCAPS TD.** (Evran) Amphetamine sulfate 15 mg., thyroid 194 mg., atropine sulfate 0.361 mg., alain 16 mg., phenobarbital 16 mg./cap. Bot. 100s, 250s, 1000s.
Use: 1 daily; obesity therapy.
- TRIOCIL.** See: Hexehidine
- TRIOGEN.** (Parke, Davis) Diphtheria-tetanus-pertussis, aluminum hydroxide adsorbed. Vial 1.5 cc.
Use: I.M., 0.5 cc. at intervals of 4-6 wks. for 3 doses; active immunization
- TRIOHISTIN.** (Aicher) Methopropylene HCl 10 mg., pyritamine maleate 10 mg., phenyl tolaxamine dihydrogen citrate 10 mg / Tab. Bot. 100s, 1000s.
Use: Triple antihistamine therapy.
- TRIOLEIN OXONIDE OF OLIVE OIL.**
See: Oilzo, liq (G. F. Harvey)
Ozettes, Susp. (G. F. Harvey)
- TRIONACIL.** (Flint, Eaton) Pot. penicillin G 100M u., gastrisin 0.5 Gm., sulfadiazine, sulfamerazine & sulfacetamide, of each 167 mg / tab. Bot. 24s, 100i.
Use: Orally; penicillin & triple sulfonamide therapy.
- TRIONAMIDE SUSPENSION, N.N.R. 1955**
(Flint, Eaton) (Meth Dio-Mer-Sulfonamide) Sulfamethazine 0.167 Gm., sulfadiazine 0.167 Gm., sulfamerazine 0.167 Gm., Sod. citrate 0.330 Gm./5 cc. Susp., Bot. 2 oz., 4 oz., 1 pt., 1 gal
Use: Triple sulfonamides
- *Generic name recognized by N.F., U.S.P. or A.M.A.

- TRIONINE.** (Roche) di Triiodothyronine Sod., di Liothyronine Sod Tab 5, 25, 50 mcg, Bot 100s, 1000s
 Use: Orally, hypothyroidism, with or without myxedema and cretinism
- TRIPRON** (Warner Chilcott) Thyroid hormone analog See Thyropropic Acid
 Use: Lowers blood cholesterol
- TRIOXANE** See Trioxymethylene
- TRIOXYMETHYLENE, N N R 1943, A D R**
 (Name is incorrectly used to denote para formaldehyde in some pharmaceuticals)
 See Paraformaldehyde (Merck Sharp & Dahme)
- W/Sod oleate, glycerin & stearic acid** See Veritas Vaginal Kreml N N R 1954 (Veritas)
- W/Sod oleate, triethanolamine, dioctyl Sod sulfosuccinate, stearic acid & aluminum silicate** See Cooper Creme N N R 1954 (Whittaker)
- TRIPAC.** (Person & Covey) Hesperidin 100 mg ascorbic acid 100 mg, propenpyridine maleate 6 mg, phenylpyrimine fumarate 6 mg, pyrilamine maleate 6 mg salicylamide 2 gr caffeine 1/4 gr/cap Vial 18s Syr Bot 4 oz
 Use: (2 f i d) Treatment of the common cold
- TRIPAC-DC.** (Person & Covey) Chlorpropen pyridamine maleate 1 mg phenylpyrimine fumarate 6 mg, pyrilamine maleate 6 mg hesperidin 100 mg Vit C 100 mg, phenyl propanolamine HCl 12.5 mg salicylamide 2 gr caffeine anhydrous 1/4 gr/Tab Bot 100s
 Use: Cold therapy
- *TRIPARANOL, N N D 1962 Withdrawn** 1(p (beta diethylaminoethoxy) phenyl) 1(p faly) 2 (p chlorophenyl) ethanol See Mer/29 (Merrell)
- TRIPAZINE** (Ortega) Sulfadiazine 0.167 Gm sulfamerazine 0.167 Gm sulfamethiazine 0.167 Gm/Tab Bot 100s
 Use: Sulfonamide
- *TRIPLENNAMINE CITRATE, USP N N D 1962 A D R 1955 Elixir USP 2** (Benzyl [2 dimethylaminoethyl] amino) pyridine citrate See Pyribenzamine Citrate (Ciba)
- *TRIPLENNAMINE HYDROCHLORIDE, USP, N N D 1962 A D R Tab, USP** See Pyribenzamine Hydrochloride (Ciba)
- TRIPHENYLMETNANE DYES, N N R 1953, A D R**
 See Fuchsin
 Methylrosaniline Chloride
- TRIPERAZINE DICTRITATE, NYDROUS**
 See Piperazine Citrate USP N N D 1962
- TRIPLE ANTIGEN** (Wyeth) *H pertussis* 30
 *Generic name recognized by N F USP or A M A
- billion/cc w/alum precipitated diphtheria & tetanus toxoids Vial 1.5 cc, 7.5 cc
 Use: I M & S C; protection against diphtheria tetanus & whooping cough
- TRIPLE BROMIDES, EFFERVESCENT TABLETS.** See: Salicides (Merck Sharp & Dahme)
- W/Phenobarbital** See: Palagren, liq (Kendall)
- TRIPLE HORMONE.** (Brean) Progesterone 25 mg, testosterone 125 mg & estradiol 12 mg/cc Vial 10 cc
 Use: Uterine bleeding, amenorrhea and threatened abortion
- TRIPLE SPRAY.** (Philadelphia Amp Co) Hydrocortisone, phenylephrine HCl, gramlicidin, neomycin sulf, methapyrilene HCl, & polymyxin B sulf in buffered sol Bot plastic 20 cc
 Use: Nasal allergies
- TRIPLE SULFA CREAM** (Ortho) Sulfathiazole 3.42%, N'-acetylsulfonilamide 2.86% N' benzoylsulfonilamide 3.7%, urea peroxide 1% Tube with or without applicator 3 oz
 Use: Bacteriostatic & bactericidal action
- TRIPLE SULFAS SUSPENSION & TABLETS** (Lederle) Sulfadiazine 0.167 Gm, sulfamethazine 0.167 Gm sulfamerazine 0.167 Gm /5 cc Susp or Tab Susp, Bot 1 pl Tab, Bot 100s, 1000s
- TRIPLE SULFOID.** (Vale) Trisulthopyrimidine 5 Gm /Tab Bot 100s
- TRIPLE SULFONAMIDE**
 See Acet Dio Mer Sulfonamides, N N R
 Acet Meth Mer
 Dia Mer Thio Sulfonamides
 Meth Dia Mer Sulfonamides, N N R
- W/Penicillin** See Meth Dia Mer W/Penicillin G Pat Prep (Various Mfr)
- W/Aureomycin** See Aureomycin Triple Sulfas Tab (Lederle)
- W/Gamtricin & penicillin G pat** See: Tri-nacid Tab (Flint, Eaton)
- TRIPOTON MALEATE** (Schering) Pheniramine Maleate, N F
- *TRIPROLDINE NCI, N N D 1962 Trans 1** (4 methylphenyl) 1 (2 pyridyl) 3 pyrrolidone propylene HCl See Actidol (Burroughs Wellcome)
- W/Codaine phosphate pseudoephedrine NCI glyceryl guaracolate** See Actifed C Syr (Burroughs Wellcome)
- W/Pseudoephedrine HCl** See Actifed Tab & Syr (Burroughs Wellcome)
- TRIPSUL M** (Carrione) Sulfadiazine 166.7 mg, sulfamerazine 166.7 mg, sulfamethazine 166.7 mg/Tab or 5 cc Bot 100s pint
 Use: Systemic infections

- TRIPTEDE.** (Schwarz) Sod. glutathione. Vial 2 Gm.
- TRIPTOLAC.** (Massengill) Milk proteins obtained by tryptic digestion of skim milk Vial 30 cc.
- Use: I.M. & I.O.; nonspecific protein therapy.
- TRIOUIN.** (Winthrop Labs) Quinacrine (Altabrine HCl) 25 mg., chloroquine (Aralen PO₄) 65 mg., hydroxychloroquine (Plaque-nin) 50 mg./Tab. Bot. 100s.
- Use: Dermatologic.
- TRISEM.** (Marsengill) (Meth-Dia-Mer-Sulfonamide) Sulfamerazine 0.17 Gm., sulfadiazine 0.17 Gm., sulfamethazine 0.17 Gm./5 cc. Susp. or Tab. Bot. 1 pl., 1 gal.
- Use: Triple sulfonamide therapy.
- TRISEM I.A.** (S. E. Marsengill Co) Sulfadiazine 0.167 Gm., sulfamerazine 0.167 Gm., sulfamethazine 0.167 Gm.; and vegetable oil, 2.5 cc./Tsp. Bot. 8 oz.
- Use: Systemic gastrointestinal or urinary.
- TRISEM-PEN.** (Marrengill) Pot. penicillin G 100M u., sulfadiazine 167 mg., sulfamerazine 167 mg., sulfamethazine 167 mg./5 cc. or Tab. Pkg. for dry pow. for sol. 60 cc. Tab., Bot. 50r.
- Use: Orally; penicillin & triple sulfonamide therapy.
- TRISERT VAGINAL.** (Ulmer) 9-Aminoacridine HCl 2 mg., phenylmercuric acetate 3 mg., methylbenzethonium chloride 1.8 mg., tyrothridin 0.6 mg., urea 100 mg., succinic acid 12.5 mg., lactose 0.65 Gm./Tab. Bot. 100s.
- Also available in powder form.
- Use: Vaginal antiseptic.
- TRISIDONNA.** (Rorer) Mag. trisulfate 7½ gr., activated charcoal 1 gr., belladonna ext. 1/12 gr./Tab. Bot. 100s, 500r, 1000s.
- Also avail. w/phenobarbital.
- Use: (1-2 tid) Gastric antacid & antispasmodic.
- TRISOCORT SPRAYPAK.** (Smith, Kline & French) Hydrocortisone alcohol 0.02%, gramicidin 0.005%, neomycin sulfate equiv. to neomycin base 0.6 mg., polymyxin-B sulfate 2000 u., phenylephrine HCl 0.125%, povidone HBr 0.5%, thimerosal 1:100,000/cc. Squeeze bot. 15 cc.
- Use: Nasal injections.
- TRISOFOR.** (Smith, Kline & French) Ferrus sulfate 6 gr., thiamine HCl 2.25 mg., riboflavin 2 mg., nicotinic acid 20 mg./6 fl. dr. Elixir, Bot. 12 fl. oz.
- Use: Hematinic.
- TRISOGEL.** (Lilly) liq. Mag. trisulfate 9 gr., aluminum hydroxide 23 gr./5 cc. Bot. 12 cc. Tab., Mag. trisulfate 4½ gr., aluminum hydroxide 1½ gr./Tab. Bot. 100s, 1000s
- Use: Antacid.
- TRI-SOLGIN.** (Lilly) Diphtheria and tetanus toxoids and pertussis vaccine combined Alum precipitated Vial 7.5 cc.
- TRISOMIN.** (Lilly) Mag. trisulfate. Pulvule 5 gr., Bot 100s, 1000r Tab 7½ gr., Bot 100s, 500s.
- Use: Antacid.
- TRI-SPAN.** (Walker Lab) Vit A 12,000 u., D 1200 u., E 3 mg., B₁ 10 mg., B₂ 5 mg., B₆ 3 mg., Calcium pantothenate 5 mg., Niacinamide 50 mg., B₁₂ 3 mcg., folic acid 0.25 mg., C 100 mg./Cap. Bot. 30s.
- Use: Space-release therapeutic multiple vitamin
- TRISTACOMP.** (Tailby Naran) Chlorpheniramine maleate 2.5 mg., phenyltoloxamine citrate 12.5 mg., pyrilamine maleate 25 mg., phenylephrine HCl 10 mg., phenylpropranolamine HCl 30 mg./Tab. or 4 tip Tab. Bot. 100r, liq. Bot. pt
- Use: Antihistaminic and decongestant.
- TRIST-A-LIN.** (Walker Corp) Pyrilamine maleate 8.3 mg., phenyltoloxamine dihydrogen citrate 8.3 mg., methapyrilene fumarate 8.3 mg., aspirin 3 gr., acetophenofidin 2 gr., caffeine ½ gr./Tab. Bot. 500s, 1000s
- Use: Antihistaminic
- TRISTAMIN.** (Physicians Prods) Chlorpheniramine maleate 12.5 mg., phenyltoloxamine citrate 6.25 mg., pyrilamine maleate 12.5 mg.
- Use: Antihistaminic.
- W/Phenylephrine HCl, salicylamide, acetyl-p-aminophenol, caffeine. See: Conitromal-CP, Cap (Physicians Prods)
- TRISTAMINE.** (Barre) Phenylpropranolamine HCl 50 mg., phenylephrine HCl 10 mg., chlorpheniramine maleate 4 mg./Tab. Bot. 100s
- Use: Nasal decongestant
- TRISTAMINE FORTI.** (Barre) Phenylpropranolamine HCl 25 mg., phenylephrine HCl 5 mg., chlorpheniramine maleate 2 mg., salicylamide 2.5 gr., acetylparaminophenol 2 gr., caffeine 0.5 gr., ascorbic acid 50 mg./Cap. Bot. 50s
- Use: Nasal decongestant & analgesic
- TRISTERONE.** (Wyeth) Estrone 6 mg., testosterone 25 mg., progesterone 25 mg./cc. Tube (I cc) Pkg. 3s; Vial 5 cc.
- Use: Control of uterine bleeding.
- TRISUL.** (Haug Drugs) Sulfadiazine 0.5 Gm., sulfamethazine 0.5 Gm., sulfamerazine 0.5 Gm., Sod. citrate 0.78 Gm./fl. oz. Susp. or Tab. Susp. 1 pt. Tab., Bot. 100r, 1000s.
- Use: Sulfonamide therapy.
- TRISULDONE.** (G. F. Harvey) (Meth-Dia-Mer Sulfonamide) Sulfamethazine, sulfadiazine

- sulfamerazine each 0.167 Gm / Tab Rot 100s 500s, 1000s
Use Triple sulfanamide therapy
- TRI-SULPA** (Central) 2½ gr each Thio Dia Mer Sulfanamide / Tab Tab Grooved Bot 100s 500s 1000s
- TRISULPAGEN**, (Central) (Meth Dia Mer Sulfanamide) Sulfadiazine 40 mg sulfamerazine 40 mg, sulfamethazine 40 mg, salicyl alcohol 15 mg / Lax Lax Bot 100s, 500s
Use Triple sulfanamide therapy
- ***TRISULPAPYRIMIDINES SUSPENSION**, Oral USP; Tab, USP See Meth Dia Mer Sulfanamides NNR 1955 (Varies Mfr)
- W/Pheniramine maleate pyrilamine maleate phenylpropanolamine HCl See Tri sulfaminic Prep (Smith Dorsey)
- W/Sulfamethylthiazole phenylazo diaminopyridine HCl See Ro Sulla Tabs (Rawell)
- TRISULFAMETH**, NNR 1955 (Arlington Funk) (Meth Dia Mer Sulfanamide) Sulfamethazine 0.162 Gm, sulfadiazine 0.162 Gm, sulfamerazine 0.162 Gm / 5 cc or Tab Syr Bot 4 oz, 1 pt Tab Bot 50s 100s 250s 500s, 1000s
Use Sulfanamide therapy
- TRISULFAMINIC** (Smith Dorsey) Triaminic 25 mg (phenylpropanolamine HCl 12.5 mg pyrilamine maleate 6.25 mg pheniramine maleate 6.25 mg) Trisulfapyrimidines 0.5 Gm / Tab or 5 cc Susp Pint Tab 50s
Use Upper respiratory disorders
- TRI-SULFANTYL** (Arlington Funk) Sulfathiazole 0.162 Gm, sulfadiazine 0.162 Gm, sulfamerazine 0.162 Gm / 5 cc or Tab Syr, Bot 4 oz, 16 oz Tab, Bot 50s 100s 250s 500s 1000s
- TRISULPAZINE SUSPENSION** (Central) Sulfadiazine, sulfamerazine, sulfamethazine of each 0.167 mg / 5 cc Bot 1 pt, 1 gal
Use: Triple sulfanamide therapy
- TRISULFAZINE TABLETS**, NNR 1955 (Central) (Meth Dia Mer Sulfanamide) Sulfadiazine 0.167 Gm sulfamerazine 0.167 Gm sulfamethazine 0.167 Gm / Tab or ½ above amounts for each Palatab Palatab, Rot 100s, 500s 1000s Tab Bot 100s 500s 1000s
Use: Triple sulfanamide therapy
- TRISULPIN** (Central) Sulfadiazine 0.083 Gm, sulfathiazole 0.083 Gm, sulfamerazine 0.083 Gm / Palatab Bot 100s 500s 1000s
- TRISULPYL**, (Approved Pharm) (Meth Dia Mer Sulfanamide) Sulfadiazine 167 mg Sulfamethazine 167 mg, sulfamerazine 167 mg / Tab Bot 100s 1000s
Use: Triple sulfanamide therapy
- *Generic name recognized by N.P., U.S.P. or A.M.A.
- TRISULPEN TABLETS**, (Prema) (Meth Dia Mer Sulfanamide) Sulfadiazine 0.166 Gm sulfamerazine 0.166 Gm, sulfamethazine 0.166 Gm, Cryst Pot Penicillin G 300 000 u / Tab Bot 24s, 100s, 1000s
Use Sulfanamide therapy w/penicillin
- TRISUREID** (Reid) Sulfadiazine 2½ gr, sulfamerazine 2½ gr, sulfamethazine 2½ gr / Tab Bot 100s
Use Infections
- TRI-SYNAR**, (Forrest Company) Belladonna ext 4.1 mg, phenyltolazamine 20 mg ethaverine HCl 20 mg / Tab Bot 100s
Use 1 f i d; antispasmodic
- TRI-SYNAR PLUS**, (Forrest Company) Belladonna pow ext 4.1 mg, phenyltolazamine 20 mg, ethaverine HCl 12.5 mg, phenobarbital 203 mg, butobarbital sod 2025 mg, pentobarbital sod 6075 mg, secobarbital sod 6075 mg / Tab Bot 100s
Use Sedatives w/antispasmodic
- TRITAL** (Testagar) Secobarbital 65 mg, amobarbital 40 mg phenobarbital 25 mg / Tab Bot 30s 100s, 1000s
Use Hypnotic
- TRITHEON**, (Orlha) Aminotriazole, 2 acetyl amino 5 nitratiazole Acinitriazole Tab Bot 30s 180s
Use: Treatment of Trichomoniasis
- TRITHESIA** (Breon) Propyl p aminobenzoate 4%, procaine base 1.5%, benzyl alcohol 5% Amp 5 cc Box 6s, 25s
Use Pruritus ani & anal fissure
- TRITIGEN** (Searle) Aq infusion of cough grass Bot 1 pt, 1 gal
Use Genita urinary sedative
- TRITON A-20** (Rahm & Haas) Oxyethylated tertiary acetyl phenol formaldehyde polymer
- TRITON WR-1339**, (Rahm & Haas) Oxyethylated tertiary acetyl phenol formaldehyde polymer
- W/Sod bicarb & glycerin See: Alevale Sol (Winthrop Labs)
- TRITONE**, (Myers Carter) Testosterone 25 mg progesterone 25 mg, estrone 6 mg / cc Vial 10 cc
- TRITRANQUIL** (Meyer) Mephensin 400 mg mephobarbital 10 mg, hyoscine HBr 0.25 mg / Tab Bot 50s, 500s
Use Relaxant
- TRIVA** (Boyle) Alkyl aryl sulfonate 35% Sod sulfate 53%, oxyquinoline sulfate 2% lactose 10%, and disodium ethylene bis iminodiacetate (versene) Packet 33 Gm 24s
Use: (b f d) Douche for trichomonal and gonorrheal infections
- TRIVA-COMBINATION**, (Boyle) Oxyquinoline benzoate and sulfate alkyl aryl

- sulfonate, disodium ethylenediamine-tetraacetate, 9-aminoacridine HCl, copper sulfate/3 Gm. Douche Packet 24s; Jel Tube with applicator 85 Gm.
Use: Therapeutic vaginal douche and jelly.
- TRIVALENT GAS GANGRENE ANTITOXIN, N.F., N.M.R. 1952.** Each pkg. contains 10,000 u. Clostridium perfringens & Clostridium septicum antitoxins & 3500 u. of Clostridium oedematiens (Navy) antitoxin. (National Drug Parke, Davis, Wyeth).
- TRI-VIDA.** (Daserel) Vit. A 5000 u., Vit. D 1000 u., C 50 mg./0.6 cc.
Use: Vitamin supplement.
- TRI-VI-FLOR.** (Mead Johnson) Fluoride (sod) 0.5 mg., Vit. A 3000 u., D 400 u., C 60 mg./0.6 cc. Bot. 30 cc.
Use: Dental caries in infants.
- TRI-VI-SOL.** (Mead Johnson) Vit. A 5000 u., D 1000 u., ascorbic acid 50 mg./0.6 cc. Bot. w/dropper 15, 30 & 50 cc. Tab., Sol. Bot. 24s, 100s.
Use: Orally; diet supplement.
- TRI-VI-SOL WITH IRON.** (Mead Johnson) Vit. A 3000 u., C 60 mg., D 400 u., iron 10 mg./0.6 cc. Bot. 30 cc.
Use: Dietary supplement.
- TRIVITRIMINE.** (Myers-Carter) Vit. B-1 100 mg., B-6 100 mg., B-12 100 mcg./cc. Vial 10 cc.
Use: Triple sulfonamide therapy.
- TRIZYL SUSPENSION.** (Central) Sulfacetamide 0.167 Gm., sulfadiazine 0.167 Gm., sulfamerazine 0.167 Gm./5 cc. Bot. 1 pt., 1 gal.
Use: Triple sulfonamide therapy.
- TRIZYL TABLETS.** (Central) Sulfadiazine, sulfamerazine, sulfacetamide of each 0.167 mg./Tab. Bot. 100s, 500s, 1000s
Also avail. at Patlabs, half strength.
Use: Triple sulfonamide therapy.
- TROEX.** (Tetagar) Methionine 25 mg., R-1 50 mg., niacinamide 100 mg., R-2 2 mg., choline 50 mg., B-6 2 mg., pantothenol 4 mg./cc. Vial 20 cc.
Use: I.M., I.V., vitamin deficiencies w/lipotropics.
- TROXICIN.** (Upjohn) Spectinomycin. Actinospectacin, antibiotic isolated from a strain of Streptomyces spectabilis.
Use: Gram positive and gram-negative infections.
- TROCASE.** (Sulliff & Case) Tyrothricin 2 mg., benzocaine 5 mg./Loz. Bot. 100s.
Use: Throat & mouth infections.
- TROCHONS.** (Ives-Cameron) Tyrothricin 2 mg., benzocaine 5 mg., sweetening & flavoring q.s./Troche. Vial 12s.
Use: Throat infections.
- TROCINATE.** (Paythress) Thiphenamil. beta-Gamic name recognized by N.F., U.S.P. or A.M.A.
- Diethylaminoethylidiphenylthioacetate HCl. Tab. 100 mg., Bot. 40s, 250s Also avail. w/phenobarbital 1/4 gr., 1 gr./Tab.
Use: 1-2 t.i.d., antispasmodic.
- W/Thiophylline, pyrilamine maleate. See Trilute, Cap. (Paythress)
- TROFONAMIDE.** (VanPelt & Brown) Triisulopyrimidine Tab., U.S.P.
- TROKELS.** (Humphrey) Tyrothricin 2 mg., benzocaine 5 mg./Loz. Bot. 12s.
Use: Throat & mouth infections.
- TROKETTES.** (Bryant) Polymyxin B, tyrothricin, gramicidin, benzocaine and celalcolium/Troche. Pkg. 10s.
Use: Throat and mouth infections.
- TROLONE.** (Riker) Sulthiame
Use: Anticonvulsant
- TROLAR.** (G. W. Carnick) Hematropine methylbromide 5 mg., butobarbital 15 mg./5 cc. or Tab. Elix., Bot. 12 R. or. Tab. Bot 100s, 1000s
Use: Antispasmodic.
- TROLNITRATE PHOSPHATE, N.M.D. 1962.** See. Triethanolamine Trinitrate Phosphate.
- TROMETHAMINE.** (Abbott) 2-Amino-2-(hydroxymethyl)-1,3-propanediol See Talatrol.
Use: Acidemia.
- TROMEXAN, N.M.D. 1962.** (Geigy) (Ethyl Biscoumarate) 3,3-carboxy-methane bis-(4-hydroxycoumarin)ethyl ester. Tab. 150 mg., Bot. 50s, 250s, 1000s. 300 mg., Bot. 25s, 50s, 250s, 1000s.
Use: Anticoagulant that depresses prothrombin activity.
- TROMINE.** (Taitagar) Phycostigmine sulfocylate 0.6 mg., atropine Sulf. 0.6 mg./cc. Vial 30 cc.
Use: Stimulate smooth muscle. Parasympathomimetic.
- TRONIC.** (Merck Sharp & Dahme) Protein hydrolyzate 6.8 Gm., thiamine HCl 4 mg., riboflavin 2 mg., pyridoxine HCl 1 mg., niacinamide 20 mg., cal. glycerophosphate 130 mg., Sod. glycerophosphate 260 mg., Pal. glycerophosphate 24 mg., Mang. glycerophosphate 16 mg., alcohol 17% / 1 1/2 fl. oz. Bot. 1 pt., 1 gal.
Use: Dietary supplement.
- TRONOLEN LOTION.** (Abbott) Tronothane HCl 1%, di-paralene HCl 2%, calamine, zinc oxide, menthol. Squeeze bot. 75 cc.
Use: Topically; local anesthetic w/antihistaminic.
- TRONOTHANE HYDROCHLORIDE, N.M.D. 1962.** (Abbott) (Promazine HCl) gamma-Morpholinopropyl 4-n-butylphenyl ether HCl Jelly (1%) Tube 7 oz. Cream (1%)

- Tube 1 oz Lot (1%) Bot 2 oz Sol (1%)
Bot 100 cc Aerosol (1%) 120 cc
Use: Local anesthetic
- TRON-OTO OTIC.** (Abbott) Trosathane 1%, erythrocine 5 mg, polymyxin B 10000 u, in glycols/1 cc Dropper Bot. 5 cc
Use (3 to 4 gtt) Simple earache
- TROPANE.** (Myers Carter) Methscopolamine bromide 1 mg/cc Vial 10 cc.
- TROPH-FRON** (Smith, Kline & French) Vit B 12 25 mcg, B 1 10 mg, ferric pyrophosphate 250 mg/5 cc or Tab Bot 50s Bot 4 fl oz
Use: 1 tsp daily to increase appetite & promote growth
- TROPHITE.** (Smith, Kline & French) Vitamin B 12 25 mcg, B 1 10 mg/5 cc Elixir, Bot 4 oz Tab, equiv to 5 cc of elix Bot 50s
Use: Appetite stimulating
- ***TROPICAMIDE** N ethyl 2 phenyl N (4 pyridylmethyl)hydrocrylamide Bis Tropane See Mydrinyl (Alcan Labs)
- TROPINAL.** (Rand) Atropine sulfate 1/200 gr, phenobarbital 1/4 gr, Vit B 2 0.50 mg, niacinamide 5 mg, Cal panlathenole 0.30 mg, Vit B 6 0.15 mg/4 cc Bot E os
Use: Antispasmodic & sedative
- TROPINATE** (Mollard) Pituitary gonadotropin 125 RU/5 cc Vial 5 cc
Use: Gonad stimulating hormone
- TROPINE BENZOATORTL ESTER METHANESULFONATE** (also named benztropine methanesulfonate) See Cogentin Methane sulfonate, Tab (Merck Sharp & Dohme)
- TROPOMIN TABLETS.** (Mollard) Choline bitartrate 100 mg, inositol 50 mg, dl methionine 50 mg, Vit B 1 5 mg, B 6 0.25 mg, C 30 mg B 12 2 mcg, folic acid 0.25 mg, poethenal 1 mg, rutin 10 mg, cabalt sulfate 1 mg/Tab Bot 100s 500s, 1000s
Use: 1 qid, o c; Ipatropic & metabolic stimulation
- TROSCINE UNICELLES.** (Miss Pharm) Mephobarbital 65 mg homatropine methylbromide 5 mg, atropine methyl nitrate 1 mg hyoscine HBr D02 mg/cap Bot 100s
Use: (1-2 daily) Delayed action, antispasmodic w/sedative
- TROSE.** (Cole) beta Lactose 15 gr, boric acid 5 gr, lactic acid 1/2 min, oxyquinoline sulfate 1/4 gr/vaginal tab Bot 100s
Use: Trichomonas vaginalis vaginitis
- TROSINONE** (Abbott) Ethisterone Tab 5 mg, 10 mg, Bot 20s, 100s
Use: Corpus luteumlike effect
- TROXIDONE.** See Tridione (Abbott)
- TROXONIUM TOSTLATE** (Registered in Great Britain) Triethyl 2 (3,4,5 trimethoxybenzoyloxy) ethylammonium tosylate
* Generic name recognized by N.F., U.S.P. or A.M.A.
- [Tosylic acid is the trivial name for p toluenesulphonic acid]
- TRYOPTEROLIUM TOXYLATE.** (Registered in Great Britain) N Ethyl N 2 (3,4,5 trimethoxybenzoyloxy) ethylpyrrolidinium tosylate (Tosylic acid is the trivial name for p toluenesulphonic acid)
- TRULASE.** (Smith, Miller & Patch) Amylolytic enzyme 30 mg, proteolytic enzyme 6 mg, lipolytic enzyme 25 mg/Tab Bot 30s, 100s
Use: Digestive aid
- TRUO-CILLIN-50.** (Abbott) Pot. penicillin G 50000 u, w/sulfamethazone 0.167 Gm, sulfadiazine 0.167 Gm, sulfamerazine 0.167 Gm/dulcet Tab Bot 25s, 100s
Use: Penicillin w/triple sulfonamide therapy
- TRUO-CILLIN-150** (Abbott) Pot penicillin G 150000 u, w/sulfamethazine 0.167 Gm, sulfadiazine 0.167 Gm, sulfamerazine 0.167 Gm/Tab Bot 25s, 100s, 1000s
- TRUO-CILLIN-300.** (Abbott) Pot penicillin G 300,000 u, w/sulfamethazine 0.167 Gm, sulfadiazine 0.167 Gm, sulfamerazine 0.167 Gm/Tab Bot 25s, 100s, 1000s
- TRUOZINE DULCET TABLETS, N.N.R. 1955** (Abbott) (Meth Dia Mer Sulfonamide) Sulfadiazine 0.1 Gm, sulfamerazine 0.1 Gm, sulfamethazine 0.1 Gm/Tab Eat 100s
- TRUOZINE SUSPENSION, N.N.R. 1955** (Abbott) (Meth Dia Mer Sulfonamide) Sulfadiazine 0.1 Gm, sulfamerazine 0.1 Gm, sulfamethazine 0.1 Gm, Sod citrate U.S.P., 1.5 Gm/5 cc Bot 1 pl, 1 gal
Use: Triple sulfonamide therapy
- TRUOZINE TABLETS, N.N.R. 1955** (Abbott) (Meth Dia Mer-Sulfonamide) Sulfadiazine 0.167 Gm, sulfamerazine 0.167 Gm, sulfamethazine 0.167 Gm/Tab Bot 100s 1000s
- TRYADS.** (Wampole) Not listed Salicylamide 3 1/2 gr, dl desoxyephedrine HCl 1/30 gr, acetophenetidin 2 1/2 gr/Tab Bot 100s, 500s
Use: Analgesic, antipyretic & central nervous system stimulant
- TRYCOGEN** (Alphodes) Sod thiosulfate thymol, oxyquinoline sulfate, wormwood oil, boric acid, starch Mag carb insert Bot 1Bs, 100s Pow, Vial 25 Gm
Use: Trichomonas vaginalis vaginitis
- TRYNAZIN OINTMENT.** (Abbott) Sulfathiazole 10 gr, aluminum sulfacarbolate 10 gr, butyn sulfate 7 1/2 gr, atropine sulfate 1/12 gr in hydrocarb-water emulsion base/oz Oint. Tube 1 oz
Use: Analgesic & antiseptic
- TRYPTAFILAVIN** (Delta) Acriflavine HCl Bot 10 Gm
Use: Antiseptic

- *TRYPASPAMIDE, U.S.P., N.N.R. 1949**
(Wollou) Sod. N-carbamylmethyl p-aminobenzenesulfonate, Paraglycylsulfanilic acid sod. salt. Amp. 1 Gm., 2 Gm., 3 Gm.
Use: Antisyphilitic.
- W/Bismuth tartrate. See: Bisolphen, Amp. (Endo)**
- TRYPP. (U. S. Vit) Trypsin 7500 u., Phenylephrine HCl 0.25% / 8 cc. B cc. drop. bot./ envelope of powd./ diluent.**
Use: Nasal congestion, sinusitis, and hay fever.
- TRYSIGEL CREME. (Bruce) Allantoin 0.25%, trypsin 1%, zirconium carbonate 2%, ethyl aminobenzoate 1%, benzalkonium chloride 0.1%, Amerchol-9 15%. For 1 lb., flip top plastic 3 oz.**
Use: Necrotic tissue and pyogenic membranes
- *TRYPsin CRYSTALLIZED, N.F., N.N.D. 1962.**
See: Orenzyme, Tab. (National Drug)
Patenzyme (National Drug)
Trypar, N.N.D. 1958. (Armour)
Trypsin (Testagor)
W/Chymotrypsin.
See: Chymaral (Armour)
Orenzyme, Tab. (National Drug)
W/Phenylephrine HCl. See: Trypp, Sol. (U. S. Vit)
- TRYPtiAR, N.N.D. 1962. (Armour) Purified crystalline trypsin. Vial containing 250,000 Armour u., 250 mg. tryptic activity w/30 cc. vial containing 25 cc. of diluent (Sorensen's Phosphate Buffer sol.) pH 7.3, plus plastic adapter for use w/pow. blower. Vial (50,000 u.) 10 cc. (125,000 u.) 20 cc.**
Use: Dissolves necrotic tissue and removes debris without injury to living tissue.
- TRYPtiAR ANTIBIOTIC OINTMENT. (Armour) Trypsin 5000 u., Chymotrypsin 5000 u., bacitracin 500 u., polymyxin B sulfate 5000 u./Gm. Oint. Tube 1/2 oz. & 2 oz.**
- TRYPtiEST INJ. (Testagor) Proteolytic activity from trypsin, 12,500 u., gelatin base 50 mg./cc. Vial 5 cc. multiple dose.**
Use: Soft tissue inflammations.
- T-10NIC ELIXIR. (Tennessee) Methamphetamine HCl 2 mg., niacin 50 mg., Vit. B-12 10 mcg., B-1 10 mg., alcohol 20%, w/ cyclamate sodium (sweetening agent)/tbsp. (15 cc.). Bot 1 pt.**
Use: Geriatric therapy.
- TUAMINE, N.N.D. 1961. (Lilly) Tuaminoheptane. Inhaler 325 mg.**
Use: Inhalant in rhinitis & allergic conditions.
- TUAMINE SULFATE, N.N.D. 1961. (Lilly) Tuaminoheptane sulfate; 1-Methylhexylam-
*Generic name recognized by N.F., U.S.P. or A.M.A.**
- mine Sulfate; 2-Aminoheptane Sulfate) Sol 1%. 1 fl. oz., 1 pt. 2%. Bot 2 fl. oz., 1 pt
Should be used only in office by physician
Use: Nasal congestion.
- *TUAMINOHEPTANE SULFATE, N.F., N.N.D. 1962 See: Tuamine. (Lilly)**
- TUAZOLE. (Strassenburgh Co) Methaqualone HCl**
Use: C.N.S. calmative agent and central muscle relaxant.
- TUAZOLE W/METHSCOPOLAMINE. See: Akalon-T-5, Akalon-T-10 (Strassenburgh)**
- TUBADIL, N.N.D. 1962. (Endo) d-Tubocurarine chloride pentahydrate 25 mg. In peanut oil, aeycholesterol derivatives & beeswax/cc. Vial 5 cc.**
Use: I.M. relaxation of voluntary muscle spasm.
- TURABINE, N.N.D. 1962. (Burroughs Wellcome) d-Tubocurarine chloride. Vial (3 mg./cc.) 10 cc.**
Use: Neuromuscular blocking agent; muscle relaxant.
- *TUBERCULIN, OLD, U.S.P., N.N.R. 1950 (Cenc. Tuberculin, Crude Tuberculin, Tuberculin-Koch).**
Lederle—Old Tuberculin Cenc. Vial 1 cc.
Patch test, 1s, 10s, 1000s.
Lilly—Old Tuberculin. Vial 1 cc. Mantoux test 10s.
Parke, Davis—Old Tuberculin. Vial 1 cc. Mantoux test 10s; Van Parquet test 3s, 10s.
Merck Sharp & Dohme—1st & 2nd test 10s, 100s; intermediate test 10s.
Wyeth—Old Tuberculin. Vial 1 cc., 10 cc.
Use: Mantoux test.
- *TUBERCULIN, PURIFIED PROTEIN DERIVATIVE, U.S.P., N.N.D. 1960 (Tuberculin P.F.D.) (Parke, Davis)**
Tab. 1st strength: Pkg. 2 vials (10 tests) w/ 1 cc. diluent; pkg. 10 tab. (100 tests) w/10 cc. diluent.
Tab. 2nd strength: Pkg. 2 vials (10 tests) w/ 1 cc. diluent; pkg. 10 tab. (100 tests) w/10 cc. diluent.
Tab. 1st & 2nd strength: Pkg. sufficient for 20 tests. In sets of 2 vials 1 tab. ea. w/ 5 cc. diluent.
- *TUBERCULIN TINE TEST. (Lederle Labs)**
Each disposable test unit consists of a stainless steel disc, with four lines (or prongs) 2 millimeters long, attached to a plastic handle. The lines have been dip-dried with Old Tuberculin. The entire unit is sterilized by ethylene oxide gas. The test has been standardized by comparative studies, showing 0.05 mg. U.S. Standard Old Tuberculin (5 International Units) or 0.0001 mg. U.S. Standard (5 International Units)

by the Mantoux technique The reliability appears to be comparable to the standard Mantoux

***TUBERCULOSIS VACCINE, HND 1962** BCG Vaccine (Bacillus Calmette Guerin vaccine)

TUBERLATE. (Heun) Sod p-aminosalicylate 12 gr, succinic acid 4 gr/Tab Bot 500s 1000s

Use Tuberculosis

***TUROCURARINE CHLORIDE, U.S.P.** HND 1962 Int., U.S.P. (d-Tubocurarine chloride) Abbott—(3 mg) 10 cc, 20 cc; (15 mg/cc) 1 cc 5 cc In oil & wax (30 mg/cc) Vial 5 cc

Burroughs Wellcome—Vial (3 mg/cc) 10 cc

Cutter—Vial (3 mg/cc) 10 cc

Endo—Vial (3 mg/cc) 10 cc

Lilly—Vial (3 mg/cc) 10 cc, 20 cc

Parke Davis—Amp (15 mg/cc) 1 cc Vial (3 mg/cc) 10 cc

Squibb—Vial (3 mg/cc) 10 cc, 20 cc

Amp (15 mg/cc) 1 cc Vial 10 cc, 20 cc

Upjohn—Vial (3 mg/cc) 10 cc

See Delacurarine, Vial (Squibb)

Tubadil, Vial NND (Endo)

Tubarine, Vial, NND (Burroughs Wellcome)

TUROCURARINE CHLORIDE IN OIL W/ Myricin See: Delacurarine, Vial (Squibb)

TUROCURARINE CHLORIDE PENTONATE. See: Tubadil, Vial (Endo)

***TUROCURARINE CHLORIDE, DIMETHYL,** HND 1962 Dimethyl ether of d-tubocurarine chloride See Mecostin Chloride (Squibb)

TUROCURARINE IODIDE, DIMETHYL, NND 1962 Dimethyl ether of d-tubocurarine iodide See: Metubine, Vial (Lilly)

TUBULIN (Incrato) Oeproteinized renal ext Amp 10 mg/cc/10s, oral, 15 cc of 1% sol/15 cc

Use: Chronic indolent ulcers associated with varicose veins, thrombophlebitis, etc., also with angina pectoris

TUCKS. (Feller Pharm) Disks of ointment flannel, medicated w/sol nl witch hazel & glycerin Disk (10 envelopes w/3 disks), 30s

Use: Proctologic & dermatologic disorders

TUHAL. (Lilly) Equal parts second Sod & amyl Sod Pulvule ¼ gr, Bot 100s, 1000s; 1½ gr, Bot 100s, 1000s; 3 gr, Bot 100s, 1000s

Use: Sedative & hypnotic

TUMS. (Howe Co) Mag trisilicate, cal carb & mag carb Tabs pk, 10s

Use: Antacid

TURASED. (Smith, Miller & Patch) Pentobarb

* Generic name recognized by N.F., U.S.P. or A.M.A.

bital Sod ¼ gr, Pot thiocyanate ¼ gr, Sod nitrite ½ gr, rutin 10 mg/Tab Bot 100s, 500s

Use (1-2 t i d) Hypertension

TURENOL. (Raymer) Ox bin ext 1½ gr, phenolphthalein ½ gr, Sod succinate 1½ gr, Sod salicylate ½ gr, cascara sagrada ext ½ gr, menthol qs/Tab Bot 100s, 1000s

Use Choleric, cholagogue

TURGASEPT. (Daha) Deodorizer aerosol can 16 oz

TURICUM (Whittier) Sod carboxymethylcellulose 0.36 Gm, Mag hydraxide 0.6 Gm/2 tsp Bot 1 pl

Use (2-4 tsp h s) Laxative

TURIOC. (Armour) Meturedepa

Use Anti neoplastic agent

T-URY. (Tennessee Pharmaceutical) Sulfacetamide 125 mg, phenylazo diamino pyridine HCl 25 mg, hyascyamide sulfate 0.5185 mg, atropine sulfate 0.097 mg, hyoscine HBr 0.0325 mg/Tab Bot 100s

Use Urinary tract infections

TUSILAN, HND 1959 Dextramethorphan HBr

TUSSABAR (Bruce) Acetaminophen 400 mg, salicylamide 500 mg, potassium guaiac sulfonate 120 mg, pyrilamine maleate 30 mg, ammonium chloride 500 mg, sodium citrate 500 mg, phenylephrine HCl 30 mg/oz Bot pls, gals

Use: Analgesic antipyretic, decongestant and expectorant

TUSSACAP (Ensley Scott) Capsules Bot 24s

TUSSADINE. (VanPell & Brown) Codeine phosphate 1 gr, ammonium chloride 6 gr, Pot guaiac sulfonate 16 gr, citric acid 0.325 Gm, chloroform 0.12 cc, menthol 5 mg/fl oz Bot 1 pl, 1 gal

Use: (1 tsp q 3 h) Sedative expectorant

TUSSAGESIC. (Smith Dorsey) Triaminic 50 mg (phenylpropranolamine HCl 25 mg, pheniramine maleate 12.5 mg, pyrilamine maleate 12.5 mg), dextramethorphan HBr 30 mg, terpin hydrate 180 mg, N-acetyl-p-aminophenol 325 mg/Tab Bot 50s & 250s

Use: Common cold

TUSSAGESIC SUSPENSION. (Smith Dorsey) Triaminic 25 mg (Phenylpropranolamine HCl 12.5 mg, Pheniramine maleate 6.25 mg, Pyrilamine maleate 6.25 mg), Dexamethorphan 15 mg, Terpin hydrate 90 mg, N-acetyl-p-aminophenol 120 mg/5 ml liq Bot pl

TUSSAHIST (Finley) Codeine phosphate 10 mg, phenylpropranolamine HCl 12.5 mg, chlorpheniramine maleate 2 mg, pyrilamine

- salsate 7.5 mg., glyceryl guaiacolate 100 mg./5 cc.
 Use: Treatment of coughs.
- TUSAMINIC.** (Smith-Dorsey) Triamfenic 100 mg., phenylpropranolamine HCl 50 mg., pyrilamine maleate 25 mg., d-methorphan HBr (Dormethan) 30 mg., terpin hydrate 100 mg./Tab. T.R. Bot. 50s.
 Use: Antitussive and expectorant action.
- TUSAMINIC EXPECTORANT.** (Dorsey) Dihydrocodeinone 1.67 mg., glyceryl guaiacolate 100 mg., Triaminic 25 mg., chloralam 13.5 mg./cc. Bot. 1 pt.
 Use: Treatment of cough.
- TUSTAFAP.** (Sullivan & Case) N-Acetyl-p-aminophenol 195 mg., aspirin 120 mg., phenylephrine HCl 5 mg., noscapine 5 mg., caffeine 32.5 mg., pyrilamine maleate 12.5 mg., methapyrilene fumarate 12.5 mg./Tab. Bot. 100s, 1000s.
 Use: Treatment of common cold.
- TUSTAR.** (Armour) d-Methorphan HBr 60 mg., w/Pot. guaiacol sulfonate, Sod. citrate, citric acid, prophepyridamine maleate 45 mg., phenylpropranolamine HCl 30 mg., chloralam & precervaline/4 oz. Bot. 1 pt.
 Use: Coughs due to colds.
- TUSTILOR.** (Taylor Labs) Dihydrocodeinone bitartrate 1/6 gr., amm. chloride 8 gr., pot. guaiacol sulf. 8 gr., chloralam 0.12 ml., tartar emetic 1/12 gr., pyrilamine maleate 50 mg., di-desoxyephedrine HCl 2 mg./oz. Syr. Bot. 1 pt.
 Use: Treatment of coughs.
- TUSSIN LIQUID.** (Dumas-Wilson) Dextromethorphan hydrobromide 15 mg., chloralam 22 mg., benzyl alcohol 40 mg., sodium citrate 42 mg., potassium guaiacol sulfonate 90 mg., ammonium chloride 90 mg./5 cc. Bot. 1 pt. Caps. Bot. 100s, 1000s.
 Use: Antitussive.
- TUSTIONEX.** (Strosenburgh) Dihydrocodeinone 5 mg., and phenyltoloxamine 10 mg. as the resin complex of each/Tab. or 5 cc. Bot. 100s, liq. Bot. 1 pt.
 Use: Antitussive
- TUSU-ORGANIDIN.** (Wampole) Codeine phosphate 10 mg.; Organidin 30 mg.; chlorpheniramine maleate 2 mg.; alcohol 15%/5 cc Bot 16 oz
 Use: Common colds, etc.
- TUSU-ORNADE LIQUID.** (Smith, Kline & French) Cocamiphen ethanedisulfonate 5 mg., chlorpheniramine maleate 2 mg., phenylpropranolamine HCl 15 mg., isopropamide 0.75 mg./5 cc. Bot. 1 pt
 Use: Common cold.
- TUSU-ORNADE.** (Smith, Kline & French) Cocamiphen ethanedisulfonate 20 mg., chlorpheniramine maleate 8 mg., phenylpropranolamine HCl 50 mg., isopropamide 2.5 mg./Cap Bot. 30s
 Use: Common cold
- T.V.D. FORMULA.** (Adams) Thyroid 10 mg., Vit. B-1 2 mg., B-2 2 mg., niacinamide 10 mg., Vit. C 20 mg., d-amphetamine phosphate 5 mg./Tab. Bot. 50s, 100s.
 Use: Obesity.
- T-V-P.** (Sullivan & Case) Potassium thiocyanate 97 mg., veratrum viride 97 mg., phytolacca 16 mg./Tab. Bot. 100s, 1000s.
 Use: Hypotensive compound.
- TWEEN RD.** See: Polysorbate 80. (Various Mfrs.)
- TWIL-RE INJECTION.** (Pitman-Moore) Vit. B-12. Vial (15 mcg./cc, 1000 mcg./cc) 10 cc.
 Use: Anemias.
- TWIL-RE-60.** (Pitman-Moore) (Vit. B-12 60 mcg./cc.) 10 cc.
- TWILTONE.** (Pitman-Moore) liver 10 mcg., folic acid 10 mg., cryst. Vit. B-12 50 mcg./cc. Vial 10 cc.
 Use: Anemia
- TWILY-8E-1000.** (Pitman-Moore) (Vit. B-12 1000 mcg./cc.) 10 cc.
- TWILYCO.** (Sherman) Vit. A 5000 u., D 1000 u., liver conc. 120 mg., Vit. B-1 3.4 mg., B-2 3.4 mg., inositol 50 mg., Vit. B-6 1 mg., C 150 mg., folic acid 0.5 mg., pantothenic acid 1 mg., Vit. B-12 5 mcg., ferric sulfate 130 mg., clovech subit. 70 mg./cap. Bot. 100s.
 Use: Hematinic w/vitamin supplement.
- TWINDIX PR.** (Allison) Dextroamphetamine sulfate 20 mg., phenobarbital 1 gr./Tab. Bot. 30s, 100s.
 Use: Obesity and mood stimulant.
- TWIN-BARR.** (Archer) Pentobarbital Sod. 1 1/2 gr., surrounding enteric-coated nucleus of phenobarbital 1/2 gr./Tab. Bot. 100s, 1000s.
 Use: Twin-action sedative.
- TWIN-BARR IMPROVED TAB.** (Archer) Pentobarbital Sod. 97 mg. in outer core, butobarbital 32 mg. in E.C. core/tab. Bot. 100s, 500s.
 Use: Prolonged sedation.
- TWISTON.** (McNeil) Rotaxamine 2 mg./Plain Tab. Bot. 100s R-A 4 mg./Tab. Bot. 50s
 Use: Antihistaminic, treatment of allergic disorders
- TWISTUSSIN SYRUP.** (McNeil) Rotaxamine (d-carbinoxamine as tartrate) 2 mg., phenylephrine HCl 5 mg., glyceryl guaiacolate 100 mg., dihydrocodeinone bitartrate 1.67 mg./cc. Bot. pts.
 Use: Cold therapy.
- TWIX.** (Cartone) Vit. A 5000 u., D 500 u.,

- C 50 mg, B 1 2 mg, B 2 2 mg nicotina-
mide 15 mg B 12 5 mcg /Tab Bot 50s
Use Dietary supplement
- TWIX LIQUID & DROPS.** (Carrosee) 8 Vit,
d biotin, inositol, L lysine HCl (liquid) Drops
contain 8 Vit, L lysine HCl inositol Liquid
8 oz Drops 1 oz
Use Multivitamin tissue building formula
- TWO STREPTOMYCINS.** (Upjohn) Dihydra-
streptomycin sulfate 0.25 Gm, streptomycin
sulfate 0.25 Gm/cc Vial 2 cc, 1D cc
- TY-B.** (Paul Maney) Adenine 5 phosphoric
acid 25 mg, Vit B 12 65 mcg/cc Amp
1 cc
Use Enzyme and vitamin therapy
- TYCIN** (Richard & Williams) Troches Bot
1000s
- TYCOPAN.** (Lilly) Vit A 20 000 u, D 1000
u, thiamine mononitrate 10 mg, riboflavin 8
mg, pyridoxine HCl 10 mg, Cal panto-
thenate 32 mg, nicotinamide 32 mg, Vit
B 12 15 mcg, ascorbic acid 100 mg,
alpha lipoic acid 20 mg biotin 0.16 mg,
folic acid 3.3 mg, p-aminobenzoic acid 33
mg, inositol 160 mg choline dihydrogen
citrate 160 mg/3 cap Bot 100s
Use Chronic alcoholism
- TYDEX.** (Tyler) d Amphetamine sulfate 10
mg & 15 mg/TR Cap Bot 100s 1000s
Use Control of obesity
- TYDEX-PLUS** (Tyler) d Amphetamine sul-
fate 15 mg, amobarbital 95 mg/TR Cap
Bot 100s 1000s
Use Control of obesity
- TYLAMIN** (Tyler) Vitamin A 5000 u, D
1000 u, B 1 2.5 mg, B 2 2.5 mg, B-6 0.5
mg, B 12 2 mcg, C 50 mg, niacinamide
20 mg, calcium pantothenate 5 mg, folic
acid 0.25 mg, calcium 20 mg, phosphorus
13 mg, iron 15 mg, iodine 0.15 mg, man-
ganese 1 mg, cobalt 0.1 mg, copper 1 mg,
magnesium 1 mg potassium 5 mg, molyb-
denum 0.2 mg zinc 1 mg /cap
Use Dietary supplement
- TYLAMIN JR** (Tyler) Vitamin A 3000 u, D
500 u, B 1 1.5 mg, B 2 2 mg, B 6 0.5 mg,
niacinamide 8 mg calcium pantothenate
0.5 mg, folic acid 0.1 mg B 12 2.5 mcg,
C 50 mg, E 1 u, calcium 20 mg, phos-
phorus 15 mg, iron 5 mg, cobalt 0.25 mg,
manganese 0.3 mg, copper 0.25 mg,
iodine 0.05 mg, zinc 0.3 mg /cap
Use Dietary supplement
- TYLAMIN SR.** (Tyler) Vitamin A 12,500 u
D 1000 u, B 1 5 mg, B 2 2.5 mg, B 6
1 mg, B 12 1 mcg, C 75 mg niacinamide
40 mg, calcium pantothenate 4 mg, E 2 u,
folic acid 0.5 mg, calcium 75 mg, phos-
phorus 58 mg, choline bitartrate 314 mg,
*Generic name recognized by NF USP or A.M.A.
- inositol 15 mg, dl methionine 10 mg, iron
30 mg, cobalt 0.04 mg, copper 0.45 mg,
manganese 0.5 mg, molybdenum 0.1 mg,
iodine 0.075 mg, zinc 0.5 mg, potassium
2 mg /Cap
Use: Dietary supplement
- TYLANDRIL** (Lilly) Diethylstilbestrol 0.25
mg, methyltestosterone 5 mg, reserpine
0.1 mg/tab Bot 100s
Use Orally, androgen estrogen therapy with
relief from emotional tension
- TYLENOL** (McNeil) Acetaminophen N Acetyl-
p-aminophenol 120 mg/5 cc Elixir, Bot 4
oz, 12 oz Drops (60 mg/0.6 cc) Bot w/
dropper 15 cc Tabs (300 mg) Bot 100s
Use (1 2 dr q 4 6 h) Analgesic
W/Paraflex, prednisolone See Parafon
Tabs (McNeil)
- TYLER OTIC** (Tyler) Neomycin sulfate 6 mg,
hydrocortisone 5 mg pyriminoleate
0.6% celastrolum chloride 0.05%, diper-
adon hydrochloride 1%, urea 5%/cc Bot
plastic dropper 5 cc
Use Treatment of otitis media and otitis
externa
- TYLOSIN** (Lilly) Under study
Use Antibiotic
- TYLOSTERONE.** (Lilly) Diethylstilbestrol 0.25
mg, methyl testosterone 5 mg/Tab Bot
30s 100s, 1000s
Use (1 daily) Combined androgen estrogen
therapy
- ***TYLOXAPOL** (Winthrop) Oxyethylated
tertiary octylphenol formaldehyde polymer
See Mucilose Super (Winthrop)
Suprinone (Winthrop)
- TYMAFAST.** (Mason Pharm) d Amphetamine
sulfate 5 mg, sodium pentobarbital 20 mg,
ascorbic acid 50 mg /Cap Bot 100s
Use Anorectic in control of obesity
- TYMAHIST.** (Mason Pharmaceuticals) Chlor-
pheniramine maleate 8 mg, 12 mg/
Cap Bot 100s
Use Antihistaminic
- TYMPAGESIC W-T** (Warren Teed) Phenyl-
ephine HCl 1/4%, antipyrine 5%, benzo-
caine 5%, Sod metabisulfite Liq. Bot w/
dropper 15 cc
Use Antihistaminic & analgesic ear drops
- TYMPANIDE** (Warren Teed) Sulfamylamide
5%, urea 10% chlorbutanol 3%, in
propylene glycol Bot w/dropper 1/2 fl oz
Use Antiseptic
- TYOCIN.** (Rand) Pyranisamine maleate 10
mg, Sod citrate 3 gr, citric acid 3/4 gr+
chloroform 8 mg/4 cc Syr, Bot 1 pt-
1 gal
Use Antihistaminic

- TYTROCIN.** (Merck Sharp & Dahme) Tyrothricin 0.05%, benzocaine 1.25%, antipyrine 5%, hezylresorcinal 0.1%, propylene glycol anhy. 30%, glycerin anhy. q.s. 100%. Sol., Bot. w/ dropper 15 cc.
Use: Ear drops for infections of external auditory canal.
- TYPHOID ANTIGEN N.**
Lederle—Vial 5 cc.
Lilly—Vial (100 million/cc.) 2 cc., (25 million/cc.) 5 cc.
- TYPHOID ANTIGEN O.** (Lederle) Vial 5 cc.
- TYPHOID-PARATYPHOID VACCINE, U.S.P., N.N.R. 1947.**
Cutter—Vials 7.5 & 15 cc. (immunization)
Lederle—Vial, immunization, 1.5 cc., 15 cc.; protein therapy (100 million/cc.) 5 cc.
Lilly—Vial, immunization, 1.5 cc., 7.5 cc., 15 cc.
Parke, Davis—Vial (100 million/cc.) 5 cc., (1500 million/cc.) 1.5 cc., 15 cc.
Squibb—Vial, immunization, 1.5 cc., 7.5 cc.
- TYPHOID VACCINE, U.S.P., N.N.R. 1950.**
Lilly—Typhoid bacilli 1500 million/cc.
National Drug—Typhoid bacilli 1000 million/cc., Amp. 5 cc., 20 cc.
Parke, Davis—Typhoid bacilli 1000 million/cc., Amp. 1.5 cc.
Sherman—Typhoid bacilli 1000 million/cc., Amp. 12.5 cc., 20 cc.
- TYPHUS VACCINE, U.S.P.**
Lederle—Vial, epidemic type, 1 cc., 20 cc.
Lilly—Vial, prophylaxis, 1 cc., 20 cc.; epidemic & murine type, 1 cc., 15 cc.
- TYPSOGEN TABLET.** (Carnrick) Principle derived from pancreas. Tab., plain or E.C., Bot. 100s, 1000s.
Use: (1-2 daily) Aid in digestion.
- TYBISOTRAINE.** (Columbus) Tyrothricin 1 mg., isocaine 4 mg., tetracaine HCl 0.75 mg., menthol 0.65 mg./loz. Bot. 100s, 500s, 1000s.
Use: 1 q. 3 h prophylactic against respiratory infections.
- TYROBENZ.** (Mallard) Tyrothricin 7 mg., benzocaine 5 mg., aromatics q.s./Troche. Bot. 100s, 1000s.
Use: Sore throat, mouth infections.
- TYROBENZO.** (G. F. Horvey) Tyrothricin 1.25 mg., benzocaine 5 mg./tab. Bot. 100s, 500s.
Use: Throat & mouth infections.
- TYROBENZOCAIN.** (New Mexico Pharm) Tyrothricin 2 mg., benzocaine 5 mg./loz. Bot. 100s, 1000s.
Use: Throat irritations.
- TYRO-CEDRIN.** (Reckei) Tyrothricin 0.015%, allantoin 0.2%, desoxyephedrine 0.25% Sol., Bot. 1 oz.
Use: Topically; antibiotic w/antiseptic
- TYROCIDIN W/Tyrothricin & benzocaine**
See: Nymore, Loz. (Durst).
- TYRODIRM.** (Merck Sharp & Dahme) Tyrothricin 0.5 mg./Gm. Cream, Tube 1 oz.; Jar 1 lb.
Use: Minor infections.
- TYRO-GRAM A OINT.** (Lauisani) Tyrothricin 0.005%, gramicidin 0.001%, sulphur 5%, resorcinal 2%, hexochlorophene 0.25%, bithional 1%, calamine, pyrilamine maleate in a non-staining water washable base Tube 1 1/4 oz.
Use: Acne.
- TYRO-GRAMINE OINT.** (Lauisani) Chlorpheniramine maleate 0.5 mg., pyrilamine maleate 5 mg., dipiperdon 0.25%, tyrothricin 0.01%, gramicidin .001%, benzalkonium chloride 2%, menthol and camphor in greaseless, washable base. Tube 1 oz.
Use: Minor skin irritations.
- TYROHIST.** (Columbia) Tyrothricin, pyrilamine maleate, desoxyephedrine HCl, cetalkonium chloride. Sol. plastic spray bot. 20 cc.
Use: Relief of nasal congestion.
- TYROLARIS.** (Merck Sharp & Dahme) Tyrothricin 0.02%, panthenol 0.02%. Sol., Bot. 8 oz.
Use: Antibiotic mouth wash.
- TYROPHILE TROCHES.** (U. S. Standard Prods) Tyrothricin 2 mg., benzocaine 7.5 mg., carbamide 7.5 mg., chlorophyllin 2 mg./troche. Bot. 12s, 100s, 500s.
Use: Prophylactic against throat & mouth infections.
- TYROSAN LOZINGES.** (Reckei) Tyrothricin 3 mg., Cal. ascorbate 100 mg./Loz. Bot. 20s.
Use: For throat & mouth infections.
- TYROSCABE.** (Merck Sharp & Dahme) Benzyl benzoate 36%, tyrothricin 0.053%, benzocaine 3.230%, inert 60.707%. Sol., Bot. 4 oz.
Use: Antibiotic scabicide solution.
- L-TYROSINE W/Pyridoxine HCl, niacinamide.** See: TPN, Liq. and Tab. (Haack)
- TYROSUM.** (Summers Labs) Tyrothricin 0.2 mg./cc. w/isopropanol, polyorbate 80 & acetone. Bot. 4 oz.
Use: Skin cleaner and treatment of secondary infections
- *TYROTHRICIN, N.F., N.N.D. 1962, A.D.F.**
Spray, N.F. Sol., N.F. Troches, N.F. As antibiotic from *Bacillus brevis*
See: Alfa-Thricin, Cream & Loz. (A.P.C.)
Bactratycin, Oint. (Wallace Labs.)
Intraderm Tyrothricin, Oint. (Wallace Labs.)
- ^cGeneric name recognized by N.F., U.S.P. or A.M.A.

Neo Yogisol, Supp Vog (Smith Darsey)
 Premecones, Supp (Pramo)
 Saluthricin, Sol. (Merck Sharp & Dohme)
 Tipex, Oint (Pitman Moore)
 Tyroderm, Oint (Merck Sharp & Dohme)
 Tyrolaris, Spray (Merck Sharp & Dohme)
 Tyrothricin Combinations on pp 767 to 769

TYROTHRINIC COMPOUND LOZENGES.
 (Raymer) Tyrothricin 1 mg, propyl p amino benzoate 7.5 mg Bot 100s, 500s, 1000s

TYROTRACB. (Merck Sharp & Dohme) Bacitracin 500 u, tyrothricin 500 mcg in white petrolatum base/Gm Oint, Tube 1/2 oz, 4 oz

Use: Skin infections

TYRO TROCHBS. (Evron) Tyrothricin 2 mg benzocaine 5 mg /troche Pkg 12s
 Use: Prophylactic against throat & mouth infections

TYROZAIN. (Jenkins Labs) Tyrothricin 2 mg, benzocaine 5 mg /troche Vial 25s Bot 100s, 1000s

Use: Alleviation of minor throat irritations

TYROZITS. (Merck Sharp & Dohme) Tyrothricin 1 mg, benzocaine 5 mg /loz Vial 12s
 Use: Antibiotic therapy of minor mouth & throat infections

TYSOBIX. (Flint, Boton) Not listed Tyrothricin 2 mg, benzocaine 10 mg /Wafer Vial 10s, Corten 12 vials
 Use: Bactericidal & anesthetic effect

TYVID, NND 1962 (Isoiazid (Merrell))
TYZINE HCl, NND 1962 (Pfizer) Tetrahydrozoline HCl A2 sol (0.1%) Rot w/ dropper 1 oz; squeeze bot 15 cc, 1 pt
 Use: 2-3 drops q 3 h; nasal decongestion
 W/Prednisolone See: Cor Tyzine (Pfizer)

TYZINE PEDIATRIC NASAL DROPS (Pfizer)
 Tetrahydrozoline HCl 0.05% sol Bot w/ dropper 1/2 oz
 Use: For infants & children with common cold

U D. PHYLOFED. (Rexall) Phenobarbital 1/2 gr, ephedrine sulf 1/2 gr & theophylline sod acetate 3 gr /Cap & Tab Also half strength Bot 100s
 Use: Bronchial asthma

ULACORT. (Testagar) Prednisolone 5 mg / Tab Bot 100s, 1000s
 Use: Rheumatoid arthritis

ULCER THERAPY.
 See: Antacids
 Anticholinergic Agents

ULCETROL. (Flint Eaton) Alenimem hydroxide dried gel 100 mg, calcium carbonate 90 mg, magnesium trisilicate 100 mg, magnesium oxide 90 mg, Vit A acetate

4000 U S P u, C 100 mg, B-1 1.5 mg, B-2 2.5 mg, niacinamide 15 mg, calcium pantothenate 5 mg, B-6 0.5 mg, B-12 activity concentrate 2 mcg /Tab Bot 100s

Use: Vitamin deficiencies in ulcer diets
ULCORT. (Ulmer) Hydrocortisone 5 mg / Gm, in a petrolatum base Tube 1/2 oz
 Use: Dermatitis

ULCORTAR CRBAM. (Ulmer) Coal tar 1%, hydrocortisone 1% Tube 1/2 oz
 Use: Resistant and chronic dermatoses

ULCYN. (Pitman Moore) Endobenzylamine Bromide
 Use: Anticholinergic agent

ULIX. (Ulmer) Cap.—Vit B 1 3 mg, B 2 3 mg, B 6 0.25 mg, cal pantothenate 5 mg nicotinamide 20 mg, plus yeast/cap Bot 100s, 1000s In.—Vit B 12 30 mcg, R 1 25 mg, B 2 2 mg, B 6 5 mg, nicotinamide 100 mg, pantothenol 5 mg/cc Vial 10 cc

Use: Vitamin B complex deficiencies
ULMETS. (Durst) Methylcellulose, Cal carb-elm bark & saccharin Tab, Pkg 48s
 Use: Gastric antacid; demulcent

ULO. (Riker) SL 501, Bayer B 186 Deligan Chlorphedianol HCl alpha (2 Dimethylamino ethyl) a chlorobenzhydrol HCl Syr (25 mg / 5 cc) Bot 12 fl oz

Use: Non narcotic, antitussive
ULOGSIC. (Riker) Chlorphedianol HCl 7.5 mg, diphenylpyrrolone HCl 0.5 mg, phenyl ephrine HCl 2.5 mg, glyceryl guaiacolate 25 mg, acetaminophen 162.5 mg /Tab Bot 100s

Use: Antitussive
ULOMINIC. (Eiker) Chlorphedianol HCl 15 mg, diphenylpyrrolone HCl 10 mg, phenyl ephrine HCl 50 mg, glyceryl guaiacolate 100 mg, alcohol 6%/5 cc Rot 1 pt

Use: Control of cough
ULTANDREN, NND 1962 (Ciba) Fluoxymesterone 2 mg & 5 mg /Tab Bot 40s
 Use: Testosterone-like androgenic agent

UL-TAR, OINT. (Ulmer) Coal tar eq to 5% liq carbonis detergens in vanishing cream base Jar 1 ox & 1 lb

ULTARZOLE, OINT. (Ulmer) Sulfathiazole 5%, liq Carbonis detergens 5% in vanishing cream Jar 1 ox, & 1 lb

ULTRAREX. (Approved Pharm) Vit B 12 5 mcg, B 1 15 mg, R 2 6 mg, B-6 0.5 mg, C 50 mg, ferrous gluconate 194.4 mg, niacinamide 50 mg, pantothenol 5 mg, liver fraction 150 mg, inositol 20 mg, dl methionine 20 mg, choline dihy citrate 35 mg, folic acid 0.3 mg /Cap Bot 50s, 100s, 1000s

Use: Dietary supplement
ULTRACAIN. (Herzog) Cod liver oil 20%

Tyrosine Combinations

Tyrosine

Others

Manufacturer

Brand

| | | | |
|--------------|--|------------------------|--|
| 0.015% | Allantoin 0.2%, desoxyephedrine 0.25% | Reccel | Tyro-Cedrin, Sol. |
| 1 mg. | Bacitracin 50 u., neomycin sulfate 5 mg., benzocaine 5 mg. | Merck Sharp & Dohme | Tetraxels, Traches |
| 1 mg. | Bacitracin 50 u., hydrocortisone 5 mg., neomycin sulf. 5 mg., benzocaine 5 mg. | Merck Sharp & Dohme | Hydraxels, Trache |
| 2 mg. | Benzocaine 10 mg. | McKesson & Robbins | Neo-Aqua-Drin, Laz. |
| 2 mg. | Benzocaine 5 mg. | Barragham Brat. | Bu-Thricin, Traches |
| 2 mg. | Benzocaine 5 mg. | Blackman | Beneirycin, Traches |
| 2 mg. | Benzocaine 5 mg. | Lannett | Lanaxels, Laz. |
| 2 mg. | Benzocaine 5 mg. | Drugmaster | T-M Laz. |
| 2 mg. | Benzocaine 5 mg. | Evron | Tyro, Traches |
| 2 mg. | Benzocaine 5 mg. | National Drugs | Oromels, Laz. |
| 2 mg. | Benzocaine 5 mg. | Cals Chem. Co. | Trane, Laz. |
| 2 mg. | Benzocaine 5 mg. | Sullin & Case | Trocasse, Laz. |
| 2 mg. | Benzocaine 5 mg. | Ives-Cameron | Trochans, Trache |
| 1.25 mg. | Benzocaine 5 mg. | Humphrey | Trobelli, Laz. |
| 1 mg. | Benzocaine 5 mg. | Harvey | Tyrobenzo, Tab. |
| 2 mg. | Benzocaine 5 mg. | C. D. Smith | Throcaline-T, Tab. |
| 1 mg. | Benzocaine 5 mg. | Jenkins Labs. | Tyrozalin, Trache |
| 2 mg. | Benzocaine 5 mg. | Merck Sharp & Dohme | Tyroxels, Laz. |
| 2 mg. | Benzocaine 10 mg. | Flint, Eaton | Tysoxer, Waters |
| 2 mg. | Benzocaine 10 mg. | Jamieson | Thy-Benzo, Traches |
| 0.03% | Benzocaine 1.25%, entipyrine 5%, hexylresorcinol 0.1%, propylene glycol 50%, glycerine | Merck Sharp & Dohme | Tyolactin, Sol. |
| 1 mg. | Benzocaine 5 mg., bacitracin 50 u. | Merck Sharp & Dohme | Tracinetz, Traches |
| 10 mg. | Benzocaine 200 mg., benzathonium chloride 10 mg., p-chloro-m-xylene 400 mg. | Lincoln | Hexathricin Otic (10 cc) |
| 40 mg. | Benzocaine 1.6 Gm., benzethonium chloride 30 mg., p-chloro-m-xylene 800 mg., hexakese 8 mg. | Lincoln | Herathricin Alropak (6 oz.) |
| 0.33% | Benzocaine 3.33%, benzyl benzoate 36% | Merck Sharp & Dohme | Tyrosabe, Sol. |
| 1 mg. | Benzocaine, bismuth subgallate, bismuth of Peru & zinc oxide | Primo | Premocanes, Supp. |
| 1 mg. | Benzocaine 5 mg., cetylaluminum chloride 5 mg. | Sullin & Case | Cetyben, Trache |
| 2 mg. | Benzocaine 3.5 mg., cetyl dimethyl benzyl ammonium chloride 5 mg. | McNeil | Amatrichin, Laz. |
| 0.5 mg / Gm. | Benzocaine 5 mg., cetyl dimethyl benzyl ammonium chloride 0.1% Benzocaine, p-chloro-m-xylene & methylbenzethonium chloride | Approved Pharm. Rhodes | Iso-Thricin, Laz. Predicaine, Oint. |

| Tyrothricin | Others | Manufacturer | Brand |
|-------------|---|---------------------|------------------------------|
| 2 mg | Benzocaine 7.5 mg, carbamide 7.5 mg & chlorophyllin 2 mg | U S Standard Prods | Tyrophyll, Traches |
| 1 mg | Benzocaine 8 mg, hexylresorcinol 2 mg | Rowell | Vio-Thricin, Loz |
| 1 mg | Benzocaine 3 mg, hyamine 2 mg | Massengill | Semels, Loz |
| 0.05% | Benzocaine 2.5%, ichthammol phenol 0.25%, zinc oxide | Lemman | Brevosal, Oint |
| 1 mg | Benzocaine 5 mg, phenylmercuric nitrate 0.5 mg | Pitman Moore | Pemzoles, Loz |
| 1 mg | Benzocaine 5 mg, tyrocidin ½ mg | Durst | Nymare, Loz |
| 3 mg | Cal ascorbate 100 mg | Reisel | Tyrosan, Loz |
| 0.02% | Desoxyephedrine HCl 0.125% | Maflbie | Malcodez, Sol |
| 0.0002% | Ephedrine lactate 1%, cetyl trimethyl ammonium acetate, dextrose | Parke, Davis | Gluco-Thricol, Sol |
| 0.02% | Ephedrine sulfate 1.3%, chlorobutanol 0.52%, menthol, alcohol | Sulliff & Case | Kerizol, Liq |
| 0.05% | Hexachlorophene 0.25%, sulfur 8%, resorcinol 2%, zinc oxide 10% | Pfizer | Eez Acnesore, Aerosol |
| 1 mg | Homorylamine HCl 20 mg, zinc bacitracin 50 u, neomycin sulf 5 mg, benzacaine 5 mg | Merck Sharp & Dohme | Penlagets, Troche |
| 0.02% | Hydrocarfisono alcohol 0.02%, phenylephrine HCl 0.25%, thenyl pyrimine HCl 0.2% | Lemman | Pyratort, Liq |
| 2.5 mg | Hydrocarfisono acetate 10 mg, phenylmercuric acetate 3 mg, 9 aminoacridine HCl 3 mg, polysorbate 80 10 mg urea 20 mg, lactose 50 mg | Webster | Aquocort Vag Supp. |
| 2 mg | Isobutyl p aminobenzoate 5 mg, benzethonium chloride 0.1% | Hance | Cyclo-Thyrcin, Loz. |
| 1 mg | Isobutyl p-aminobenzoate 4 mg & tetracaine 0.75 mg | Columbus | Tyrisothrine, Loz. |
| 1 mg | Neomycin sulf, 5 mg, benzocaine 5 mg | A.P.C. | Benlyron, Lozenge |
| 0.02% | Neomycin sulf 0.2%, phenyltoloxamine dihydrogen citrate 1%, bithional 0.5%, zircanium oxide 4%, diperodon HCl 0.25% | Bryant | Allergetic, Lot |
| 2 mg | Neomycin 3.5 mg, prophen pyridomine maleate 12.5 mg, benzo-caine 8 mg | Chicago Pharmacol | Neo-T Coin, Lozenge |
| 0.02% | Panthenol 0.02% | Merck Sharp & Dohme | Tyrolaris, Sol. |
| 0.05% | Pectin 6%, propylene glycol 20% | Pitman Moore | Tlplex, Oint |
| 0.03% | Phenylephrine HCl 0.25%, pyrrolomine maleate 0.2%, retiktonum chloride 0.05%, phenylmercuric acetate | Hance | Neotrol, Sol |
| 1 mg. | Phenylmercuric acetate 3 mg, 9-aminoacridine HCl 2 mg, urea 20 mg, lactose 100 mg, sod lauryl sulf 5 mg | Webster | Trinolix, Vag. Sepp. |
| 0.6 mg | Phenylmercuric acetate 3 mg, methylbenzethonium chloride 1.8 mg, 9 aminoacridine HCl 2 mg, urea 100 mg, succinic acid 12.5 mg, lactose 650 mg | Ulmer | Triserit Vaginal, Tab., Pow. |

Tyrosine Combinations

| Tyrosine | Others | Manufacturers | Brand |
|------------------|---|---------------------|---|
| 0.5 mg. | Phenylmercuric nitrate 3 mg., beta-lactose 845 mg., propain 25 mg., Sol. lauryl sulfate 3 mg., succinic acid 25 mg. | Smith-Dorsey | Vagisol, Supp. & Pow. |
| 2 mg. | Polymyzin B sulfate 1000 u., benzocaine 5 mg., cetylaluminum chloride 2.5 mg., gramicidin 100 mg., chlorpheniramine maleate 0.5 mg. | A.P.C. | Somez, Troches |
| 0.02% | Propadrine 1.5% | Merck Sharp & Dohme | Prothricin, Liq. |
| 1 mg. | Propadrine 20 mg., benzocaine 15 mg., bisnuth subgallate 150 mg., zinc oxide 150 mg. | Merck Sharp & Dohme | Tresanoids, Supp. |
| 2 mg. | Prepsin 2 mg. | White | Lozilles, Tab. |
| 1 mg. | Propyl p-aminobenzoate 7.5 mg. | Boyer | Tyrosine Comp., Sol. |
| 0.1% | Pyrimidine 1%, benzocaine 5%, cetyl dimethyl ethyl ammonium chloride 2% | Sutlin & Case | Quado, Cream |
| 0.02% | Pyrimidine maleate 0.25%, benzocaine chloride 0.05%, phenyl- ephedrine HCl 0.25% | Bryce | Deterjel Aromatic Spray |
| 0.5 mg. | Pyrimidine maleate 10 mg., benzocaine 10 mg. | Amer. Pharm. | Hyrocin, Oint. |
| 2 mg. | Sulfadiazine 2 mg., butylaminobenzoate 2 mg. | Thompson | Suthagen, Troches |
| 0.05% | Sulfathiazole 10%, 2-chloro-4-phenylphenol 0.1%, methoben Tetracaine HCl 0.5% & epinephrine 2% of a 1:1000 sol. | Pitman-Moore | Alginex, Cream |
| 0.5 mg./ Gm. | Tetracaine HCl 0.5% & epinephrine 2% of a 1:1000 sol. | Columbus | Ophthalocaine w/tyrosine, Ophth. Oint. |
| 0.25 mg./ Gm. | Thenylpyrimine HCl 2%, benzocaine 2% | Frano | Dermeze-Frano, Oint. |
| 1 mg. | Tyrosine 1/2 mg., benzocaine 3 mg. | Durst | Nymore, Sol. |
| 1 mg. | Tyrosine 0.5 mg., benzocaine 3 mg. | Durst | Nymore, Troches |
| 0.05% | Zinc caprylate 5%, Zinc propionate 5% | Rowell | Gyno-Flex, Cr. |

benzocaine 2%, amyl p aminobenzoate 0.7%. Oint, Tube 1 oz; Jar 1 lb

Use: Anesthetic & aid to granulation

ULTRACAPS. (Hart) Vit A 3000 u, D 1000 u, thiamine HCl 3 mg, riboflavin 3 mg, pyridoxine HCl 0.25 mg, ascorbic acid 75 mg, niacinamide 20 mg, Cal pantothenate 5 mg, mixed natural tocopherols 5.67 mg / Cap Bot 100s, 1000s

Use: Vitamin deficiencies

ULTCORTEN See Prednisone

ULTRADIAZOL. (Wirth) Tab Bot 1000s

ULTRA-LENTE ILETIN. (Lilly) A complex between zinc ions and insulin 40 u/cc 10 cc; 80 u/cc 10 cc

Use: Long acting insulin (36 hrs)

ULTRAN, NND 1962 (Lilly) Phenaglycodol 2 p Chlorophenyl 3 methyl-2,3 butanediol 300 mg / Pulv 200 mg / Tab Bot 100s

Use: Tranquilizing

W/Darvon A S A See Darvo Tran (Lilly)

ULTRAPAQUE-A, B AND C, (Bell Craig) A contains 100% barium sulfate, type B 98% barium sulfate, C 95% barium sulfate Containers 10 lbs, 50 lbs 100 lbs C is also available in individual dose packages of 4 1/2 oz 120 per box

Use: Alimentary tract radiography

ULTROGEN AQUEOUS INJECTABLE. (Ulmer)

Natural estrogens 2 mg / cc Vial 10 cc.

Use: I M, Estrogen therapy

ULTROGEN-D A, (Ulmer) Theelin 2 mg, pethelin sulfate 1 mg / cc Vial 10 cc

Use: Dual action estrogenic hormone therapy

ULTROGEN IN OIL. (Ulmer) Natural estrogens 2 mg in sesame oil / cc Vial 10 cc

Use: I M Estrogen therapy

ULTROGEN MYOSOL. (Ulmer) Natural estrogenic substances 20000 u / cc

Use: Estrogenic

ULTROGEN TABLETS. (Ulmer) Natural estrogens (water soluble) 1.25 mg / tab Bot 100s, 1000s

Use: Estrogen therapy

ULVICAL. (Ulmer) Vit A 1500 u, D 200 u, C 16.6 mg, B 1 3 mg, B 2 2 mg, E 2 mg, w/ferrous sulfate & Cal pyrophosphate / Tab Bot 100s, 1000s

Use: Vitamin & mineral supplement in pregnancy & lactation

UMBRATHOR. (Heyden Chem) Tharium dioxide 25% sol Bot 200 cc.

Use: Oral, rectal, urethral, x ray contrast media

UNACAINE (Navocal) Metabutethamsine HCl 2 Isobutylaminoethyl m-aminobenzoate HCl

* Generic name recognized by N F, U S P or A M A

Anestubes (38 mg / cc) 1, 1.8, 2.2 & 2.3 cc Tins 50s

Amps (38 mg / cc) 2 cc Tins 50s

Tab (76 mg to make 2 cc)

Use: Injectable local anesthesia

UNDECAP. (Cole) Undecylenic acid Cap

D 44 Gm, Bot 100s, 500s, 1000s

Use: Psoriasis & neurodermatitis

* **UNDECOYLUM CHLORIDE-IODINE,** NND 1962 See: Virac, Prep (Rusan Labs)

* **UNDECYLENIC ACID, N F,** NND 1962

(Loeuff) Cap D 44 Gm, Bot 100s, 500s 1000s

See Desenex Sol, NNR (Wallace & Tiernan)

Undecap, Cap (Cole Chem Co)

Undesol, Oint (Veltex)

W/Copper undecylenate See Kudelate

Oint (Ulmer)

W/Copper undecylenate & aerasol See

Decupyl, Prep (Croakes Barnes)

W/Copper undecylenate, zinc undecylenate

& phenylmercuric acetate

See: Di Spor, Oint (Wintersmith)

Undeslin, Oint (Cutter)

W/Diphenamyl methylsulfate, salicylic acid

See Advicin, Cream, Pow (Schering)

W/Hexachlorophene See Appliderm No 7,

Oint (White)

W/Hexachlorophene & zinc undecylenate

See Hexacide, Oint (Central Pharm)

Undex Improved (VanPelt & Brown)

W/Hydrocortisone diethylaminoacetate HCl

Neomycin See Podiacart Oint (PodialRx)

W/Phenylmercuric nitrate See Merlenate

Oint & Pow (Chicago Pharmacal)

W/Phenylmercuric oleate See Mercene

Oint (Maltbie)

W/Salicylanilides; 5 chloro salicylanilides; 5,3'

& 5,4' dichlorosalicylanilides; zinc un

decylenate See Salundek New, Oint (Wal

lace & Tiernan)

W/Sod propionate, Sod caprylate, pro

pramic acid, salicylic acid, copper un

decylenate & Sod dioctylsulfosuccinate

See Verdofam, Sol (Texas)

W/Triethanolamine & zinc stearate See

Trydecyl, Cream (Patch)

W/Triethyl cetyl ammonium pentachlor

phenate See: T C A P Cream (Wallace Lab)

W/Zinc undecylenate

See Desenex, Prep (Wallace & Tiernan)

Epiderma Dex, Oint (Sulliff & Case)

Undex, Pow (VanPelt & Brown)

W/Zinc undecylenate, hexachlorocyclohexene

isomer & benzyl ammonium chloride See

Hexacide, Oint (Central Pharm)

W/Zinc undecylenate, menthal & benzocaine

See: Decyl, Cream (Paul Maney)

UNDECYLENIC ACID POTASSIUM SALT.

See: Timalox (Burroughs Wellcome).

UNDECYLENIC ACID SALTS.

See: Calcium.

Copper.

Sodium.

Zinc.

UNDESILIN. (Cutter) Sod. undecylenate 10%,

copper undecylenate 1% & undecylenic

acid 5%. Oint. Jar 1 oz. Pew. s shaker,

Bot. 1 1/4 oz.

Use: Fungus infections.

UNDESOL. N.N.S. 1955. (Veltex) Zinc-

undecolate, N.N.R., undecylenic acid w/zinc

undecylenate.

Oint. (Zinc undecylenate 0.2 Gm., unde-

cylenic acid 50 mg./Gm.) Jar 30, 60, 454

& 2270 Gm.

Pows (Zinc undecylenic acid 2%) Container

90 Gm., Jar 454 Gm.

Use: Local infections & fungicide.

UNDEX IMPROVED. (VonPelt & Browe) Hex-

echlorophene 2%, Zinc undecylenate 20%,

Undecylenic acid 5%. Ointment 1/2 oz. Tube.

Use: Antibacterial—Antilungol.

UNDEX POWDER. (VonPelt & Brown) Zinc

undecylenate 20%, undecylenic acid 2%.

Pkg. 1 1/2 oz. Also avail. w/hexachloro-

phene 2%.

Use: Topically, for fungus infections.

UNDULANT FEVER DIAGNOSIS.

See: Brucella Abortus.

Brucella Abortus Antigen.

Brucellergen, Vial. (Marck Sharp & Dahme)

UNDULANT FEVER VACCINE, N.N.R. 1933.

See: Brucella Vaccine.

National Drug—2500 million each of Br.

abortus & Br. suis killed organisms/cc. Vial

30 cc.

Parke, Davis—1000 million each of Br.

abortus & Br. melitensis organisms/cc. Vial

30 cc.

Sherman—1000 million each of Br. abortus

& Br. melitensis organisms/cc. Vial 12.5 cc.

20 cc.

Use: S.C. & I.M.; diagnosis & treatment for

undulant fever.

UNIBASE. (Parke, Davis) Water-absorbing

oint. base. Jar 1 lb., 5 lb.

Use: Ointment base.

UNICAP. (Upjohn) Vit. A 5000 u., D 500 u.,

ascorbic acid 50 mg., thiamine mononit.

2.5 mg., riboflavin 2.5 mg., pyridoxine HCl

0.5 mg., Cal. pantothenate 5 mg., nicotinic

acid amine 20 mg. & Vit. B-12 2 mcg./

Cap. Bot. 24s, 100s, 250s, 1000s.

Use: Dietary supplement.

UNICAP M. (Upjohn) Vit. A 5000 u., D 500

u., C 50 mg., B-1 2.5 mg., B-2 2.5 mg.,

B-6 0.5 mg., cal. pantothenate 5 mg., nica-

tinamide 20 mg., Vit. B-12 2 mcg., cobalt

0.1 mg., copper 1 mg., iron 10 mg., manga-

nese 1 mg., magnesium 6 mg., molybdenum

0.2 mg., zinc 1.5 mg., pot. 5 mg., iodine

0.15 mg. & calcium/Cap. Bot 30s, 90s,

180s

Use: Multivitamin-mineral, supplement

UNICAP CNEWABEL. (Upjohn) Vit. A 4000

u., D 400 u., B-1 2 mg., B-2 2 mg., C 75

mg., nicotinamide 18 mg., B-6 1 mg.,

calcium pantothenate 5 mg., Vit. B-12 2

mcg./Tab. Bot. 24s, 100s.

Use: Dietary supplement.

UNICAP SENIOR. (Upjohn) Vit. A 5000 u.,

D 400 u., thiamine mononitrate 3 mg., ribo-

flavin 3 mg., ascorbic acid (sodium ascor-

bate) 90 mg., nicotinamide 30 mg., pyri-

doxine HCl 1.5 mg., calcium pantothenate

9 mg., Vit. B-12 3 mcg., iron (sulfate) 10

mg., iodine (potassium iodide) 0.15 mg.,

calcium (lactate and carbonate) 50 mg.,

magnesium (sulfate) 6 mg., copper (sulfate)

1 mg., manganese (sulfate) 1 mg., potassium

(sulfate) 5 mg./Tab Bot. 24s, 100s.

Use: Supplement to the often deficient diet.

UNICAP THERAPEUTIC TABLETS. (Upjohn)

Vit. A 5000 u., D 500 u., C 150 mg., B-1 10

mg., B-2 10 mg., B-6 2 mg., cal. panto-

thenate 20 mg., nicotinamide 100 mg., B-12

4 mcg., folic acid 1.5 mg., cobalt 0.1 mg.,

copper 1 mg., iron 10 mg., manganese 1

mg., molybdenum 0.2 mg., zinc 15 mg.,

magnesium 6 mg., pot. 5 mg., iodine 0.15

mg., calcium 60 mg./tab. Bot. 30s, 90s,

100s, 500s.

Use: Nutritional deficiencies.

UNIDIGIN, TABLETS. N.N.E. 1950. (Merrell)

Digitoxin, Tab. 0.1 mg., 0.2 mg., Bot. 100s.

UNIDOTE. (Panray) Activated charcoal 2

parts, magnesium oxide 1 part, tannic acid

1 part/pockette 15 Gm.

Use: Poison antidote

UNITENSEN ACETATES, N.N.O. 1962. (Irwin,

Neisler) Cryptanamine acetate. Amp. (2

mg./cc.) 5 cc.

Use: Vasodilator.

UNITENSEN-PHEN TABLETS. (Irwin, Neisler)

Cryptanamine tannate 1 mg., phenobarbital

15 mg./tab. Bot. 100s.

Use: 1 t.i.d.; hypertension.

UNITENSEN-R. (Irwin, Neisler) Cryptanamine

tannate 1 mg., reserpine 0.1 mg./tab. Bot.

50s, 100s.

Use: Hypertension.

UNITENSEN TANNATES, N.N.O. 1962.

(Irwin, Neisler) Cryptanamine tannate, alka-

loids from veratrum viride. Tab 2 mg., 50s,

100s.

Use: (2 daily morn. & night) Hypertension.

UNITROXIN. (Tilden) Digitoxin 0.6

- nitroglycerin 17 mg, strychnine zolf 85 mg / 100 cc Bot 1 pt
 Use: Treatment of cardiovascular diseases
- UNIVERSAL LINIMENT.** (Suliff & Case) Oil eucalyptus 4.68%, oil camphor 3.12%, oil cajepul 2.33%, camphor 2%, menthol 0.75% Bot pts, gals
 Use: Relief of muscular soreness.
- UNIVITE.** (Lannett) Vitamins A 5000 u, D 500 u, B 1 3 mg, B 2 215 mg, B 6 0.5 mg, folic acid 0.25 mg, B 12 1 mcg, C 40 mg, niacinamide 20 mg, cal pantothenate 5 mg / cap Bot 100s, 500s, 1000s
- UNNA'S ROOT.** See: Zinc Gelatin, U S P
- UNSATURATED ACIDS.** See: Fatty acids, an saturated, also check fats, unsaturated
- URACIL.** (Chicago Pharm) Sed salicylate 5 gr, Pot Iodide 1 gr, Cimicifuga ext 1/7 gr, Gelsemium 1/8 gr, thiamine HCl 1 mg / Tab Bot 100s, 500s, 1000s
 Use: Analgesic
- ***URACIL MUSTARD.** (Upjohn) 5 [bis(2 chloroethyl)amino]uracil 10 mg / Cap Bot 50s
 Use: Leukemias and lymphomas
- URADAL.** See: Carbramal (Various Mfr)
- URAMAN.** (Superior) Methenamine Mandelate 125 mg, sulfacetamide 125 mg, acid sodium phosphate 125 mg / Tab Bot 100s, 500s
 Use: Urinary tract infections
- URAMAN W/AZO.** (Superior Pharm) Methenamine Mandelate 125 mg, sulfacetamide 125 mg, phenylazodiaminopyridine NCI 50 mg, acid sodium phosphate 125 mg / Tab Bot 100s, 500s
 Use: Urinary antiseptic & analgesic.
- URANTOIN-VIOLET.** (Irwin, Nelsler) Atan-tain 0.6%, crystal violet 1:1000 urea 15% Oint, Jar 2 & 4 oz, 1 lb
 Use: Dressing in burns & irritations of skin
- UR-AT.** (Paul Money) Phenylazo diamino pyridine HCl 50 mg, sulfacetamide 250 mg, homatropine methylbromide 0.015 mg / Tab Bot 100s, 1000s
 Use: Urinary tract infections
- URAZIUM.** (Carrigan) Phenylazo diamino-pyridine HCl 0.1 Gm / Tab Bot 100s
 Use: Urinary antiseptic
- URRITRON.** (Kremers Urban) Dicalcium phosphate 7 1/2 gr, ferrous sulfate exsicc 1 1/2 gr, thiamine HCl 1 mg, riboflavin 1 mg, Vit O 400 u / Cap Bot 100s, 500s, 1000s
 Use: (2 1/2 d) Vitamin deficiencies
- ***UREA.** N.F., N.M.R. 1950, A.O.R. 1953 Carbamide (Various Mfr)
 See: Ureaphyl Inj (Abbott)
 U-evert (Travenol Lab)
- Use: Orally & topically; tissue stimulant & diuretic
- W/Allantoin & crystal violet See: Urantoin Violet, Oint (Irwin, Nelsler)
- W/Allylisobutylbarbituric acid, Methamphetamine sulfate See Ourebese Tabs (Biber)
- W/Antipyrine, benzocaine & glycerine See Aurazine, Liq (Smith Oarsey)
- W/Calcium chloride See: Afenil, Amp (Bil-luber Knoll)
- W/Chlorophyll & benzocaine
 See C B U, Oint (Warren Teed)
 Ureacaine Oint (Chicago Pharm)
- W/Propesin, benzyl alcohol & p chloro meta xylenol See Ounaguent, Oint (Ascher)
- W/Sulfacetamide, neocalamine, talc, glyc-erol, camphor & menthol See: Neolana-mide, Liq (McNeil)
- W/Sulfacetamide, sulfathiazole, N¹benzoyl sulfanilamide See: Sultrin, Tab (Ortho)
- W/Sulfanilamide, benzyl alcohol & propylene glycol See: Surbyl, Liq (Strasburgh)
- W/Sulfanilamide & sulfathiazole See: Sul-torea, Oint (Drug Products)
- W/Sulfathiazole, benzocaine & menthol See: Sulfocamp, Oralel & Tob (C D Smith)
- W/Tyrothricin, benzocaine, chlorophyllin See: Tyrophyl, Troches (U S Standard Prod)
- W/Tyrothricin, phenylmercuric acetate, β -aminoacridine NCI, lactose & sed louryl sulf See: Trinalls, Vag Supp (Webster)
- W/Tyrothricin, phenylmercuric acetate, meth-ytbenzethonium chloride, β -aminoacridine NCI succinic acid, lactose See: Trisert Vaginat, Tab, Pow (Ulmar)
- UREA PEROXIDE.** See: Glycerite of Hydro-gen Peroxide W/Carbamide (Intern Pharm Corp)
- UREACAINE.** (Chicago Pharm) Urea 33 1/2%, chlorophyll 1%, benzocaine 10%, Jar 1/2 lb, 1 lb
 Use: Anesthetic & local in stimulating granulation
- UREAPHIL.** (Abbott) Lyophilized urea 40 Gm / 250 cc Abba-liter Bot
 Use: Diuretic.
- UREASE.** (Hynson, Westcott & Dunning) Tab 25 mg, Bot, 100s, 500s, 1000s
 Use: Diagnosis of urea in urine & blood
- URECHOLINE CHLORIDE.** N.N.D. 1962 (Marck Sharp & Dahme) [Bethanechol Chloride] beta methylcholine carbamate chloride, Amp. (3 mg) 1 cc, 6c Tab 5 mg & 10 mg Bot 100s
 Use: Parasympathomimetic
- ***UREDEPA.** [Armour] Ethyl[bis(1-astidinyl)phosphoryl]carbamate See: Aviner
- p-UREIDORENZENEARSONIC ACID.** See: Carbarone, U S P

- URESE** (J. B. Boerig & Co) See: Benzthionide
- UBISTREIN**, (Upjohn) Estrogens mixed.
- UBISTREIN**, (Upjohn) Estrogens mixed.
 Cap.s (1000 u) Bot. 100s, (2000, 4000 & 10,000 u) Bot. 30s, 100s.
 Ellix (2000 u./4 cc.) Bot. 4 oz., 1 pt., 1 gal.
- Ural**, eq. asp.s (2 mg./cc.) 1 & 10 cc.s (5 mg./cc.) 5 cc.
- Ural**, in oil (2000 & 10,000 u./cc.) 1 cc., 10 cc.s (20,000 u./cc.) 10 cc.
- URESTIN & VITAMIN R**, (Upjohn) Estrogenic substances 30,000 u., thiamine HCl 30 mg., riboflavin 30 mg., nicotinamide 300 mg./fl. oz. Syr., Bot. 12 fl. oz.
- UBETHANE**, U.S.P. (Lilly) Tabs, U.S.P. (Ethyl Carbonate, Ethyl Urethane, Urethane) Enseal 5 gr., Bot. 100s, 500s, 1000s.
- Amp.s** LV., 4 Gm./30 cc. Pk. 6s
 Supp.s 1 Gm., Pk. 6s
 Use: leukemias.
- W/Prolant**, Homatropine methylbromide.
 See: Pantropine Inj. (Smith, Miles & Potch)
- W/Thio-Der-Mar** Sulfonamide. See: Toldermal, Cream, (Central Pharm.)
- W/Thio-Dia-Mes** Sulfonamide. See: Trigynol, Jelly, (Central Pharm.)
- URETHANE W/QUININE HYDROCHLORIDE**.
 See: Quinine Hydrochloride & Urethane, Amp. (Morton & Converse).
- URVERT FOR INJECTION**, (Baxter) 30% urea in invert sugar solution. Bot. 93 ml, 210 ml.
- URICOSURIC AGENTS**.
 See: Dexamethasone
 Benamid, Tab. (Mack Sharp & Dahme)
- URICIN**, (Richard & Williams) Phenylazodiaminopyridine 50 mg., sulfacetamide 0.5 Gm., atropine sulfate 0.0195 mg., hyoscine hydrobromide 0.0063 mg., hyoscyamine sulfate 0.104 mg./Tab. Bot. 50s.
 Use: Treatment of urinary tract infection and spasm.
- URGININ**, N.N.R. 1954. (Grisord) Water-insoluble glycosides, urginin A & B derived from dried bulb of squill, *Urginea maritima*. Tab. 0.5 mg., Bot. 100s, 1000s.
 Use: See Digitalis.
- URIVERT**, (Trovinal) Urea 90 Gm. lyophilized, to be mixed with 210 cc. of 10% invert sugar.
 Use: IV. 60 Drops/min. to reduce intracranial pressure.
- U.R.I. DILACAPS**, (Sig. Labs) Phenylephrine HCl 15 mg., phenylpropranolamine HCl 25 mg., chloropheniramine maleate 8 mg./Cap. Bot. 50s, 100s.
 Use: Antihistamine/decongestant.
- U.R.I. EIQUID**, (Sig. Labs) Phenylpropranolamine HCl, 10 mg.; phenylephrine HCl, 5 mg.; pyramine maleate, 5 mg.; chlorpheniramine maleate, 2 mg./Tsp. Bot. pl.
 Use: Antihistamine/decongestant
- URIDINAL**, (Crystal) Phenylazodiamine pyridine HCl 1.5 gr./Tab. Bot. 100s.
 Use: Urinary antiseptic
- URIDINAL-R**, (Crystal) Tab. Bot. 500s.
- URILENE**, (Crystal) Tab. Bot. 1000s.
- URINE RUGAB TEST**. See: Clinistix, (Amer)
- URINARY ACIDIFIER**. See: Chlor-Ethamine, Tab. (Pitman-Moore)
- URIPLEX**, (Floyd-Dobney) Sulfacetamide 250 mg., phenyl-azo-diamino pyridine & methscopolamine nitrate 1 mg./Tab. Bot. 100s
 Use: Urinary infections
- URIRID**, (Chicago Pharm) Atropine sulfate 1/2000 gr., hyoscyamine 1/2000 gr., w/ methenamine, methylene blue, benzoic acid, salol & Gelsemium/tab. Bot. 500s, 1000s.
 Use: (1-2 t.i.d.) Relief of pain in urinary infections.
- URISEPT**, (Graft) Calcium mandelate 0.2 Gm.; methenamine 0.2 Gm.; phenylazodiaminopyridine HCl 50 mg.; sodium phosphate, monobasic, 80 mg./Cap. Bot. 100s.
 Use: Urinary tract antiseptic.
- URISEPTIN**, (Blaine) Methenamine 2 gr., phenylsulficylate 1/2 gr., methylene blue 1/10 gr., benzoic acid 1/2 gr., atropine sulf. 1/1000 gr., hyoscyamine sulfate 1/2000 gr./Tab. Bot. 100s, 1000s.
 Use: Urinary antiseptic.
- URISTIX**, (Amer) Test strips, Pkg 125s.
 Use: Test for protein and glucose in urine
- URITONE**, (Farka, Davis) Methenamine. Amp. (2 Gm) 5 cc., 6s, 25s, 100s.
 Ellix: 40 gr./oz., Bot. 1 pt., 1 gal.
 Ellix, comp.s Methenamine 10 gr., sandalwood 30 gr., corn silk 120 gr., raw palm-ella berries 120 gr., alcohol 20%/oz., Bot. 1 pt., 1 gal.
 Tab.s 7 1/2 gr. Bot. 100s, 1000s; w/Sod. acid phosphate, 5 gr. each, Bot. 100s, 1000s.
 Use: Urinary antiseptic.
- URITRAL**, (Central) Calcium mandelate 3 gr., methenamine 3 gr., phenylazodiaminopyridine HCl 1/2 gr., sod. phosphate monobasic 1 1/2 gr./cap. Bot. 100s, 500s
 Use: (2 t.i.d.) Urigenital antiseptic and analgesic.
- UROANTHELONE**. See: Kutrol, Cap. (Farka, Davis)
- URICOSUBIC AGENTS**. See: Benamid, Tab (Mack Sharp & Dahme)
- * Generic name recognized by N.F., U.S.P. or A.M.A.

UROBIOTIC. (Pfizer) Oxytetracycline HCl W/glucosamine 125 mg, sulfamethizole 250 mg, phenylazo diaminopyridine HCl 50 mg /Cap Bot 50s

Use Urinary tract infections

URO-CHLOR. (Vanpelt & Brown) Methenamine 0.5 Gm, ammonium chloride 0.5 Gm /Tab Bot 100s, 500s

Use: Urinary antibacterial

URODINE. (Nysca) Phenylazo diamino pyridine HCl 100 mg /Tab Bot 1000s

Use Urinary analgesic and antiseptic

URODYNE NMB (Beach Pharm) Phenylazo diamino pyridine HCl 150 mg, homatropine methylbromide 10 mg /Tab Bot 50s

Use Urinary pain or spasm

***UROENTERONE.** See Kutral, Kapseal (Parke Davis)

UROGESIC. (Edwards) Phenylazodiaminopyridine HCl 100 mg, hyoscyamine HBr 0.12 mg atropine sulfate 0.08 mg, scopolamine HBr 0.003 mg /Tab Tabs Bot 50s, 100s

Use: Analgesic for urinary tract disturbances

UROGESIC SULFA (Edwards) Sulfacetamide 500 mg, phenylazo diamino-pyridine HCl 75 mg, hyoscyamine HBr 0.09 mg, atropine sulfate 0.006 mg, scopolamine HBr 0.003 mg /Tab Bot 50s 500s

Use: Sulf sensitive urinary tract infections

UROGRAPH (Warner Chilcott) Urea nitrogen chromatography papers

Use: Determination of urea nitrogen

UROGRAPHY AGENTS

See Diodrast (Winthrop Labs)

Iodhippurate Sodium (Mallinckrodt)

Iodopyracet (Winthrop Labs)

Iodopyracet Compound (Winthrop Labs)

Methiodal (Winthrop Labs)

Neo-Iopax (Schering)

Skidan (Winthrop Labs)

Sodium Acetozoate (Mallinckrodt)

Sodium Iodamethamate (Schering)

UROKON SODIUM, NND 1962 (Mallinckrodt) (Sod Acetozoate) Sod. 3-acetylaminobenzoate Amp 30%, 25 cc., Box 1s, 5s 20s, 50% 25 cc., Box 1s, 5s, 20s, 70% 25 cc., Box 1s, 5s, 20s Vial 70% rubber-diaphragm stoppered 50 cc

Use: I V; x-ray contrast medium for urinary tract visualization

UROLATE. (Borcherdt) Methenamine 2 gr, Cal mandelate 2 gr, atropine sulfate 1/1000 gr, hyscyamine alkaloid 1/1000 gr./Tab Bot 100s, 500s.

Use: Urinary antiseptic.

UROLITHIA. (Borcherdt) Methenamine 20 gr, lithium benzoate 5 gr., Sod benzoate 2 gr.,

* Generic name recognized by N.F., U.S.P. or A.M.A.

In soothing, demulcent menstruum of couch grass & corn silk /Tsp Bot. 10 oz.

Use: Urinary antiseptic

UROLOCAINE. (Arnar Stone) Benzocaine 10%, oxyquinoline benzoate 0.25% in bland, water soluble liq. vehicle Bot 4 oz

Use: Urethral topical anesthetic

UROLOGIC SOLUTION G. (Abbott) (Suby's Solution G) Citric acid monohydrate 32.40 Gm, Magnesium Oxide (anhydrous) 3.84 Gm, Sodium Carbonate (anhydrous) 4.28 Gm Abba Liters

Use: Urinary tract irrigation for calculi

UROLUCOSIL. See Thiasulfil (Ayerst)

UROMETN (Beach) di methianine 500 mg /Tab Bot 100s

Use Urinary tract antiseptic and acidifier

URONA SPECIAL. (Sutliff & Case) Systin 2.05 Gm, sodium acetate 2.6 Gm, juniper berries 2.6 Gm, corn silk 3.9 Gm, apocynum 0.26 Gm /fl oz Bot pts, gals

Use Urinary antiseptic and diuretic

URONAL. See Barbitol (Various Mfr)

URONAMIDE. (Flint, Eaton) Sulfamethythiazidazole 250 mg, sulfacetamide 250 mg, hyoscyamine ext 10 mg /tab Bot 50s, 100s Syr Bot 1 pt, 1 gal

Use (1 2 1sd) Antibacterial, spasmolytic & analgesic action in urinary tract infections

URO-PHOSPHATE. (Paythress) Methenamine 7 1/2 gr, acid Sod phosphate 10 gr /Tab Bot 100s

Use: Urinary antiseptic

UROQUID-AMIDE. (Beach) Methenamine mandelate 350 mg, di methianine 200 mg, solicylamide 250 mg /Tab Bot 100s

Use Treatment of urinary calculi

UROSULFIN (Nepera Chem) Sulfamethylthiadiazole 250 mg, phenylazodiaminopyridine HCl 50 mg /tab. Bot 50s

Use: Analgesic w/sulf therapy

UROSULFON. (Cansel Midland) Sulfacetamide 0.5 Gm /tab Bot 30s, 100s, 250s, 1000s

Use: Sulfanamide therapy

UROTROPIN, N.N.R. 1949 (Warner Chilcott) Methenamine, Tab 5 gr., Box 30s, Bot 100s, 500s; 7 1/2 gr., Box 20s, Bot 100s 500s

Use: Urinary antiseptic.

UROTROPIN NEW. See: Methenamine Anhydromethylenecitrate (Various Mfr)

UROTROPIN & SODIUM ACID PHOSPHATE. (Warner Chilcott) Tab. 5 gr., Bot 100s.

Use: Urinary antiseptic.

URSINUS. (Smith Dorsey) Calcium acetyl

- sallylate carbamide (Calurin) 300 mg., tri-
 omnic 50 mg./Tab. Bot. 100s.
 Use: Acute and chronic sinusitis.
- U.S.T.** (Miden) Pot. bromide 1D gr., Hy-
 scyamus flexi. 8 min./fl. oz. Bot. 1 pl., 1 gal.
 Use: [1 dr. i.l.d.] Sedative
- UTASED.** (Massengill) Viburnum opulus 4 gr.,
 wild yam 4 gr., scutellaria 4 gr., life root
 2½ gr., cinnamon 1½ gr., clove 1½ gr.,
 ginger 1/6 gr./5 cc. Elisir, Bot. 1 pl., 1 gal.
 Use: (1-2 t.p. i.l.d.) Antispasmodic, carmina-
 tive & uterine sedative.
- UTERINE RELAXING FACTOR.** See: Lutelin,
 Est., Tab. (Hynson, Wescott & Dunning)
- UTONIUM.** (Massengill) Viburnum prunif-
 lorum ext. 1 gr., viburnum opulus ext.
 1 gr., star grass ext. ½ gr., helonias ext.
 ½ gr., squaw vine ext. ½ gr., corylophyllum
 ext. ¼ gr./tab. Bot. 100s, 500s.
 Use: (1 q. 2 h. p.r.n.) Antispasmodic &
 sedative.
- UVALERAL.** See: Bromisovalum, N.P.
- U-VITA.** (Pücher Pharmacol.) Cabbage Juice
 conc. 9 gr./cap. Bot. 100s.
 Use: Peptic ulcer.
- VABROCID.** See: Nitrofurazone (Eaton).
- VACAGEN.** (Merck Sharp & Dahme) Anti-
 genic fraction of diplococcus pneumonia
 50,000 million, streptococcus 20,000 million,
 Haemophilus influenzae 10,000 million,
 Neisseria catarrhalis 5000 million, Klebsiella
 pneumoniae 5000 million, Staphylococcus
 aureus 10,000 million/tab. Bot. 20s, 60s,
 100s.
 Use: Immunization against virulent patho-
 genic organism infections of upper respira-
 tory tract.
- V/Methapyrilene HCl.** See: Neovacogen,
 Tab. (Merck Sharp & Dahme)
- VACCINE, RACILLUS CALMETTE-GUERIN,**
 N.N.D. 1960. See: Tuberculosis Vaccine.
- VACCINE, BCG, N.N.D. 1962.** See: Tubercu-
 losis Vaccine.
- VACCINE, INFLUENZA VIRUS, MONOVA-
 LENT TYPE A (ASIAN STRAIN), N.N.D.**
 1962. See: Influenza Virus Vaccine Mono-
 valent Type A Asian Strain.
- VACCINE, INFLUENZA VIRUS, POLYVA-
 LENT, N.N.D. 1962.** See: Influenza Virus
 Vaccine, Polyvalent. (Various Mfr.)
- VACCINE, MUMPS, N.N.O. 1962.** See:
 Mumps Vaccine. (Lederle)
- VACCINE, PERTUSSIS, N.N.D. 1962.** See:
 Pertussis Vaccine. (Various Mfr.)
- VACCINE, PERTUSSIS, ALUMINUM HY-
 DROXIDE ADSORBED, N.N.D. 1962.** See:
 Pertussis Vaccine, Aluminum Hydroxide Ad-
 sorbed. (Cutter)
- VACCINE, PERTUSSIS, ALUM PRECIPIT-
 ATED, N.N.D. 1962.** See: Pertussis Vaccine,
 Alum Precipitated. (Various Mfr.)
- VACCINE, POLIOMYELITIS, N.N.D. 1962**
 See: Poliomyelitis Vaccine. (Various Mfr.)
- VACCINE, RABIES (DUCK EMBRYO),
 N.N.D. 1962.** See: Rabies Vaccine. (Various
 Mfr.)
- VACCINE SMALLPOX, N.N.D. 1962.** See:
 Smallpox Vaccine
- VACCINE, TUBERCULOSIS, N.N.O. 1962.**
 See: Tuberculosis Vaccine (Various Mfr.)
- VACCINE, WHOOPING COUGH, N.N.O.**
 1962 See: Pertussis Vaccine. (Various Mfr.)
- VACID.** (Fester) Polyacrylic cationic exchange
 resin, Vog. Supp. Rox 6s
 Use: pH stabilizer and treatment of vaginal
 infections
- VACON.** (Ethical Pharm.) Phenylephrine HCl
 D.20% with benzalkonium chloride. Nasal
 Solution, Bot. 1 oz.
 Use: Vasconstrictor
- VACUET SUPPOS.** (Medco) Sodium bicar-
 bonate, sodium pyrophosphate and sodium
 biphosphate in blend of polyethylene gly-
 cols of high and low mol. weight/supposi-
 tory. Adult and pediatric sizes in jars of
 3 and 12.
- VACUOSA PELLOIDS.** (Ayerst) Hydrophilic
 hemicellulose obtained from carob seed
 Pelfoid, Bot. 4 oz., 8 oz., 16 oz.
 Use: All types of constipation.
- VACU-TOR.** (Beulich) 6.7 Gm. suppository
 contains sodium bi-phosphate and sodium
 bicarbonate. Suppositories—Jars 10s.
 Use: Relief of constipation.
- VAD.** (Cele) Vit. A 5000 u., D 500 u./Cap.
 Bot. 100s.
 Use: Sickets.
- VAD SOFCREAM.** (Walker) Vitamin A
 100,000 u., D 10,000 u., allantoin 0.1%/oz.
 Aerosol can 5 oz. Also available w/Neo-
 mycin suff.
 Use: Treatment of diaper rash, etc.
- VADCON DROPS.** (Walker Vitamin) Vit. A
 palmitate 10,000 u., D 2000 u., ascorbic
 acid 100 mg./cc. Bot. w/dropper 30 cc.
 Use: Dietary supplement.
- VADRIN FORTIS COMPOUND.** (First Texas
 Chem) Chlorpheniramine maleate 2 mg.,
 methapyrilene HCl 20 mg./Tab. Fortis
 Chlorpheniramine maleate 4 mg., metha-
 pyridene HCl 40 mg./Tab. Fortis Comp.
 Phenylpropanolamine HCl 40 mg., chlor-
 pheniramine maleate 4 mg., methapyrilene
 HCl 40 mg./Tab. Bot. 100s, 1000
 Use: Antihistaminic.
- VADRIN FORTIS COMPOUND.** (First Texas
 Chem) Chlorpheniramine maleate, metha-

pyrilene HCl, phenylpropranolamine HCl/
Tab Bot 100s, 1000s

Use: Antihistaminic

VADZOLE OINT. (Ulmer) Sulfathiazole 5%
with vit A & D Jas 1 oz & 1 lb

Use: Bacteriostatic & granulation stimulant

VAFLOL (Breen) Vit A Cap 25,000 u, Bot
100s, 500s, 1000s

Use: Vitamin A therapy

VAGAGILL. (Massengill) Phenylmercuric ni-
trate 15000 oxyquinoline sulfate, boric
acid, carbolic acid Bot 3 oz Also avail as

Vaginal Supp

Use: Antiseptic

VAGA-SPRAY. (Menlo Park Labs) Cetyl tri-
methyl ammonium bromide and etoxylated

lanolin alcohals Bot 4 oz aerosol

Use: Vaginal infections

VAG-ASTRING POWDES. (Sutliff & Case)
Phenol 2% eucalyptol 0.1%, menthol 0.1%,
thymol 0.1% berberine 0.01% Myamine

1622 0.01%, alum 10%, boric acid 43.7%,
dextrose 43.7% Jar 4 oz, 1 lb, 5 lbs

Use: Mild antiseptic and astringent for
feminine douche

VAGICIDE. (Minnesota Pharm) 9 Amino-
acridine undecylenate 1 mg n myristyl 3

hydroxybutylamine HCl 1 mg methyl
benzethonium chloride 18 mg succinic

acid 12.5 mg/Tab or Gm Bot 30s, 100s
w/inserter; Powder 4 oz; 4 Gm insufflator

cart 12s

Use: Gyneciatric

VAGINAL DOUCHE POWDES. (Sutliff &
Case) Boric acid 39.7%, zinc sulfate 29.8%,
alum ammonium 29.8% salicylic acid

0.124%, carbolic acid 0.124%, menthol
0.062%, hydrastine 0.006% eucalyptol,
thymol oil wintergreen Jar 3 oz

Use: Mild antiseptic and astringent powder
for feminine douche

VAGINOIDS. (Merck Shore & Dohme) Ich-
thammal, phenol iodine, zinc phenolsul-
fonate & boric acid/supp Box 12s

Use: Antiseptic & anesthetic

VAGI-PLEX. (Rowell) Hexestrol D05 mg
sulfacetamide -dazine, -merazine, -metha-
zine ea 1.25%/Gm Tube 3 oz

Use: Vaginal infections

VAGISOL POWDES. (Smith Dorsey) Phenyl-
mercuric acetate 3 mg, tyrothricin 0.5 mg,
w/succinic acid, papain, Sod lauryl sulfate

& lactose/Gm Bot 45 Gm

Use: Vaginal antiseptic.

VAGISOL SUPPOSITABS (Smith Dorsey)
Phenylmercuric acetate 3 mg, tyrothricin
0.5 mg, succinic acid 25 mg, Sod. lauryl

sulfate 3 mg, beta lactose 845 mg, papain

* Generic name recognized by N F U S P or A M A

25 mg/Vaginal Tab Bot 36s Vagisol
comb Pkg 36 supp, 1 applicator

Use: Trichomonas & manilia vaginitis

VAGITAB. (Ascher) 9 Aminoacridine HCl 2
mg, phenylmercuric acetate 3 mg, methyl

benzethonium chloride 18 mg, succinic
acid 12.5 mg, allantoin 20 mg, lactose

0.75 Gm/vag tab Bot 30s, 100s

Use: Vaginal infections

VAG-M. (KLR Lab) Gentian violet 0.05%
w/undecylenic acid, alum, benzoic acid/
pkg Box 5s

Use: Vaginal fungicide

VAG SOL. (Mixer) Phenylmercuric acetate
1500 (0.2%) in buffered solvent of 50%
alcohol menthol, with pH adjusted to 4.9

Bot pts

Use: Vaginal cleanser and antiseptic

VALACHOL. (Vale) Ox bile ext Cap, Bot
100s, 1000s

Use: Choleric

* **VALETHAMATE BROMIDE, NND 1962**
See Murel Amp, Tab (Ayerst)

W/Phenobarbital See: Murel W/Phenobar-
bital, Tab (Ayerst)

VALIOL (Novacal) Menthyl valerate Vial
10 cc

Use: Relief of nausea

VALIUM (Hoffmann La Roche) Diazepam

VALLERGINE. See: Promethazine HCl, N F

VALLESTRIL, NND 1962 (Searle) Melhal-
lenestril Tab 3 mg, Bot 100s 20 mg,
Bot 250s

Use: Estrogen therapy

VALMID, NND 1962 (Lilly) Ethinamate
Tab 7½ gr Bot 100s, 1000s

Use: 1-2 tab hs; nonbarbiturate sedative.

VALMIDATE. See: Ethinamate

VALMYCIN. (Moore) Neomycin sulf 60 mg,
sacalin 160 mg, pectin 30 mg/Tab Bot
100s

Use: Orally, treatment of diarrhea in chil-
dren

* **VALNOCTAMIDE.** 2 Ethyl 3 methylvaler-
amide See Axiquel (McNeil)

VALOCELL. (Valentine) Methylcellulose Tab
0.5 Gm, Bot 50s, 1000s.

Use: Laxative

VALOCTIN. (Knoll) Actin 1 gr, bromural 4
gr/Tab Bot 100s, 500s, 1000s

Use: (1 q 4-6 h) Sedative & antispasmodic.

VALPIN. (Enda Lob) See: Octatropine
Methylbromide

VALPIPAMATE METHYLSULFATE (new
Pentapipeside Methylsulfate). See:
Quifene (Ayerst Lab)

VALTOBABB. (Valentine) Sod pentobarbital

- 1 gr., pentobarbital 0.5 gr./tab. Bot. 100s, 1000s.
 Use: Daytime sedation.
- VANA-DIAZINE.** (Vitamilz) Sulfadiazine 40 gr., Vit. B-1 24 mg./fl. oz. Bot. 4 oz., 16 oz.
- VANAPHEN.** (VonPelt & Brown) Phenobarbital 1/4 gr., Thiamine hydrochloride 5 mg./fl. dram, or Tab. Elizir. Bot. 6 oz., gal. Tab. Bot. 100s, 500s, 1000s.
 Use: Sedative—hypnotic.
- VANAY.** (Ayerst) Glyceril triacetate 250 mg./Gm. Creams Tube 1 1/2 oz. w/opplcator.
 Use: Monilia vaginitis.
- VANCIDE BN.** (Vanderbilt) Sodium Bithionolate.
 Use: Bactericide and fungicide.
- VANCDIN, N.N.D. 1962.** (Lilly) Vancomycin 500 mg/10 cc. Rubber-stoppered Amp. Vial 10 cc.
 Use: Severe resistant infections.
- VANCOMYCIN NCl, N.N.D. 1962.** (Lilly) An antibiotic from *Streptomyces orientalis* See: Vancocin.
 Use: (I.V.) Gram-positive (Staph.) infection
- VAN-COTT COMBINED VACCINE.** (Kirk) Staph. albus 200 million, Staph. aureus 200 million, Strep. mixed 100 million, Pneumococcus 200 million, E. coli 200 million/cc. Vial 20 cc.
 Use: Mixed bacterial infections.
- VANESIA.** (Horton & Converse) Dried aluminum hydroxide gel 5 gr., magnesium trisilicate 2 1/2 gr., sod. carboxymethyl cellulose 1/2 gr./Tab. Bot. 20s, 100s, 500s. Also available as suspension. Bot. 8 oz., 32 oz. 1 gal.
 Use: Antacid
- VANILLA.** See: Ethyl Vanillin, N.F.
- VANILLIC DIETHYLAMIDE.** (U. S. Vitamin) A central nervous system stimulant. See: Emvon, Vial, Tabs. (U. S. Vitamin)
- VANIRDME.** See: Ethyl Vanillin, N.F.
- VANDPHEN W/Vit. B-1** (VonPelt & Brown) Phenobarbital 1/4 gr., thiamine HCl 5 mg./dr. Bot. 8 fl. oz., 1 gal.
 Use: Sedative.
- VANDTEM.** (VonPelt & Brown) Salicylamide 130 mg., pyrilamine maleate 6.25 mg/5 cc.
 Use: Treatment of hay fever.
- VANPLEX.** (VonPelt & Brown) Brewer's yeast dried 65 mg., rice bran conc. 46 mg., thiamine HCl 5 mg., riboflavin 5 mg., nicotinic acid 20 mg., pyridoxine 1 mg., Cal. pantothenate 10 mg./Tab. Bot. 100s, 500s, 1000s.
 Use: Multiple Vitamin B deficiencies.
- VANQUIN.** See: Pyvinium Pamote.
- VANUL.** (Vanguard) Vegetable mucilage and belladonna Tinct. 2 cc., and licorice 0.5 cc./fl.oz. Susp. Bot. 1 pt.
 Use: Gastric and duodenal ulcers.
- VANULETS.** (Bryant) Vit. A 3000 u., D 1000 u., B-1 3 mg., B-2 3 mg., B-6 0.5 mg., B-12 3 mcg., C 50 mg., cal. pantothenate 5 mg., folic acid 0.25 mg., niacinamide 20 mg./tab. Bot. 100s.
 Use: Dietary supplement.
- VANVITA IMPROVED.** (VonPelt & Brown) Vit. A 25,000 u., B-1 20 mg., B-2 5 mg., niacinamide 150 mg., Vit. C 150 mg., D 1000 u./Cap. Bot. 100s, 500s, 1000s.
 Use: Dietary supplement.
- VANZDATE, N.N.D. 1959.** (VanPelt & Brown) Benzyl benzoate 25%. Lot, Bot. 11E cc, 3.78 L.
 Use: Externally; scabicide.
- VAPBDLE.** (Burroughs Wellcome) Amyl nitrite. 3 min., Box 12s, 100s; 5 min., Box 12s, 100s.
 Use: Vasodilator & antispasmodic.
- VABAMEL.** (Baker) Vit. A 2500 u., D 800 u., B-1 0.6 mg., B-2 1.2 mg., niacin 5 mg., w/modified cow's milk/Con. Can 12 1/2 oz.
 Use: Infant food.
- VARIDASE, N.N.D. 1962.** (Lederle) Topically: Vit. Streptokinase 100,000 u., streptodornase 25,000 u. Jelly: Streptokinase 100,000 u., streptodornase 25,000 u. w/ carboxymethylcellulose, Jar 15 cc. Buccal Tabs. (12,500 u) Bot. 24s, 100s.
 Use: To dissolve accumulations of pus & blood.
- VARISOL.** (Abbott) Invert sugar 30%, Sod. chloride 10%, benzyl carbonyl 1%. Amp. 10 cc., Box 6, 25s.
 Use: Sclerosing agent.
- VASANO.** (Schering) Hyoscyamine camphorate 0.4 mg., scopolamine camphorate 0.1 mg./Tab. Tube 12s.
 Use: Treating seasickness, carsickness & airsickness.
- VASCUALS.** (Vitamin Corp) dl-alpha-Tocopherol, N.F. Cap. (30 & 100 mg) Bot. 100s, 1000s. Drop (100 mg./cc) Bot. 60 cc.
 Use: Vitamin E therapy.
- VASCUTOL.** (Schenley) Choline 3 Gm., inositol 500 mg., pyridoxine HCl 4 mg/15 cc. Bot. 1 pt.
 Use: Aid in metabolism at fat.
- VASCUTUM IMPROVED.** (Schenley) Choline bitartrate 349 mg., inositol 167 mg., d-methionine 84 mg., B-6 0.67 mg., B-12 2 mcg., quercetin 15 mg., ascorbic acid 12.5 mg./cap. Bot. 100s, 1000s.
 Use: Combating impaired fat metabolism.
- VASIMID.** See: Talazoline HCl, U.S.P.
- VAS I ZINC.** (Ophthalmics, Inc.) Zinc sulf

¼%, phenylephrine HCl ½% Stars Torner
15 cc

VASOCON-A OPHTHALMIC. (Smith, Miller & Patch) Naphazoline HCl 0.05%, azo-
zoline phosphate 0.5% boric acid 1.2%,
phenylmercuric acetate 0.002%, w/sod
chloride sod carbonate anhydrous/cc
Plastic squeeze bot w/dropper tip 15 cc
Use Ocular decongestant and aethista
mina

VASOCON OPHTHALMIC. (Smith Miller &
Patch) Naphazoline HCl 0.1%, boric acid
1.2%, phenylmercuric acetate 0.002%,
w/sodium chloride & sod carbonate an-
hydrous/cc Bot plastic squeeze w/dropper
tip 15 cc

Use Ocular decongestant

VASOCONSTRICTOR.

See: Epinephrine Prep

Kutapressin (Kremers Urban)

VASOCORT. (Smith, Kline & French) Hydro-
cortisone alcohol 0.02%, paredrine HBr
0.5%, phenylephrine HCl 0.125%, thimerol
sol 1:100,000 Sol, Bot w/dropper ½ oz
Spraypak, Bot w/spray ½ oz

Use: Scleritis & nasopharyngitis

VASODIATOL, NND 1962 (Rowell) Pento-
erythritol tetranitrate 10 mg or 20 mg/
Tab Bot 100s, 500s, 1000s Available
w/Phenobarbital 16 mg

Use: Angina pectoris

VASODIL. See: Telazolone HCl, USP

VASODILAN, NND 1962 (Mead Johnson)
Isosuprine HCl 10 mg/Tab Bot 100s.
Amp 2 cc (5 mg/cc) box of 6

Use: Arterial insufficiency Peripheral vas-
cular disease Cerebrovascular disorders

VASODILATORS.

See: Atkavervir

Amyl Nitrite (Various Mfr)

Arlanad (Roche)

Arlidin (Arlington Park)

Cornacton (Cavendish).

Cartueon (Cartunon)

Cryptenamine

Ethyl Nitrite (Various Mfr).

Erythryl Tetranitrate (Various Mfr)

Inositol Hexanitrate (Various Mfr)

Khellin (Various Mfr).

Kutapressin, Vial (Kremers Urban)

Mannitol Hexanitrate (Various Mfr).

Metamine (Leeming)

Myocardone (Chemico).

N tetramine, Tab (Squibb)

Ocyl Nitrite (Hynson).

Padutin (Winthrop Labs)

Papaverine (Various Mfr).

Pentylethryl Tetranitrate (Various Mfr)

Ranivac (Roche)

Sodium Nitrite (Various Mfr)

Veratrum Viride (Various Mfr)

VASODILATORS, CORONARY.

See: Glyceryl Trinitrate (Various Mfr)

Khellin (Various Mfr)

Papaverine (Various Mfr)

Pentylethryl Tetranitrate (Warner Chil-
cott)

Peritrate (Warner Chilcott)

Ranivac (Roche)

VASODRINE. (Premo) Epinephrine HCl

1:1000 in Isotonic Sod chloride sol Amp

1 cc, Rax 6s, 12s Vial 1 fl oz Bot 1 fl oz

Use I.M., S.C. & topically; sympathomimetic

agent.

VASOPRED. (Smith, Miller & Patch) Predni-
solone 0.25%, naphazoline HCl 0.1%,

chlorpheniramine maleate 0.3%, methyl

cellulose 0.25% 5 cc plas dropper tipped

squeeze vial

Use Inflammatory and allergic eye dis-

orders

***VASOPRESSIN INJECTION, USP, NND**

1930 beta Hypophomies Post pituitary

pressor hormone See: Pitressin Amp

(Parke, Davis)

***VASOPRESSIN TANNATE INJECTION,**

USP, NND 1962 beta Hypophomies

tannate See: Pitressin Tonnate (Parke

Davis)

VASORUTIN. (Buffington) Sod nitrite 1 gr-

crofaegus Ir 2 min, rutin 20 mg, nitro-

glycann 0.26 gr, diethyl barbituric acid

¼ gr/Pill Bot 100s 1000s

Use: (i q d) Vasodilator

VASO-TABS. (Paul Money) Pentylenetetrazol

100 mg, nicotinic acid 50 mg/Tab

100s, 1000s

Use: Cerebral stimulation

VASOTRATE. (Hiss) Pentoerythritol tetra-

nitrate 30 mg/unicelle No 1 contains

pentoerythritol tetranitrate 30 mg, & nitro-

glycerine 4/500 gr/Unicelle No 2 con-

tains pentoerythritol tetranitrate 30 mg,

ethylglycerine 4/500 gr, amobarbital 30

mg/Unicelle Bot 30s 100s

Use: Control of angina attacks

VASO-RD UNICELLES. (Hiss) Pentoerythritol

tetranitrate 80 mg/Timed release cap Bot

30s, 100s

Use: Angina pectoris

VASOXYL HCl, NND 1962 (Burroughs

Wellcome) (Methoxamine HCl) beta hydrox-

ybeta-(2,5 dimethoxyphenyl)-isopropylamine

HCl Inf. (10 or 20 mg/cc) 1 cc., 5cc

12s, 100s Nasal Sol 0.25%, Bot 1 pt., 1

gal Spray. Bot 15 cc. & Bot w/dropper

15 cc

Use: Pressor action.

* Generic name recognized by N.F., USP or A.M.A.

- W/Polymyxin B sulf., gramitidin. See: Aerodrin, Sol. (Burroughs Wellcome)
- VASOXYL IN PROCAINE SOLUTION.** (Burroughs Wellcome) Methoxyamine HCl. Paralerol (15 mg./cc. with 1% procaine HCl) 1 cc., Box 12s, 100s
Use: Maintain or restore blood pressure.
- VASTRAN AMP.** (Wompole Labs.) Vit. B-12 75 mcg., sod. nicotinate 20 mg., adenosine-5-monophosphoric acid 25 mg., benzyl alcohol 2%/cc. Vial 3 cc.
Use: I.M., osteoarthritis, bursitis & tendinitis.
- VASTRAN ELIXIR.** (Wompole) Nicotinic acid 50 mg., ascorbic acid 100 mg., riboflavin 5 mg., thiamine mononitrate 10 mg., pyridoxine HCl 1 mg., calcium pantothenate 5 mg./5 cc. Bot. pls.
Use: Vasodilator and metabolic stimulant.
- VASTRAN FORTE.** (Wompole) Vit. C 50 mg., B-1 5 mg., B-2 2.5 mg., R-12 1 mcg., col. pantothenate 2.5 mg., R-6 0.5 mg., nicotinic acid 375 mg./Cap. Bot. 100s.
Use: Treatment of high cholesterol levels
- VASTRAN TABLETS.** (Wompole Labs.) Nicotinic acid 50 mg., ascorbic acid 100 mg./Vial. B-1 10 mg., B-2 5 mg., B-6 1 mg., B-12 2 mcg., col. pantothenate 5 mg./Tab. Bot. 100s, 500s
Use: Orally, treatment of geriatric patients.
- VA-TAR "A" OINT.** (Ulmer) 1% selected crude coal tar with low (0.2%) phenol-cresol content. Jar 1 oz. & 1 lb.
- VA-TAR "R" OINT.** (Ulmer) 2% selected coal tar with low (0.4%) phenol-cresol content. Jar 1 oz. & 1 lb.
- VATINE W/Vit.** B-1 Sol. (Smith-Dorsey) Vit. B-1 100 mg., B-2 2 mg., B-6 3 mg., Cal. pantothenate 5 mg., niacinamide 100 mg./cc. Vial 10 cc., 30 cc.
Use: I.V., Vitamin B complex.
- VB.** (VanPelt & Brown) Lotion. Bot. 3 oz., Pf. gal.
Use: Emollient.
- V-CILLIN, N N D.** 1962. (Lilly) Phenoxymethyl penicillin 125 mg./cap. Bot. 50s; 250 mg./cap. Bot. 24s. V-Cillin Drops. Phenoxymethyl penicillin 1.5 Gm. Bot. with dropper.
Use: Orally, penicillin therapy.
- V-CILLIN-K, N N D.** 1962. (Lilly) Phenoxymethyl penicillin, Tab. 125 mg., Bot. 50s, 100s; 250 mg., Bot. 24s, 100s.
Use: Orally, penicillin therapy
- W/Pyranil, hitadyil, clopane, aspirin, acetaminophen, caffeine. See: V-Kor, Tab. (Lilly)
- V-CILLIN PEDIATRIC—125.** (Lilly) Phenoxymethyl penicillin 125 mg. (200,000 u) Tip. Susp. Bot. 40 cc., 80 cc., 150 cc.
Use: Orally (1 tip. q 4-6 hr.) Penicillin therapy
- V-CILLIN PEDIATRIC—250.** (Lilly) Phenoxymethyl penicillin 250 mg./5 cc. Bot. 60 cc.
- V-CILLIN-K PEDIATRIC.** (Lilly) Potassium Phenoxymethyl penicillin 125 mg./5 cc. Bot. 40 & 80 cc
Use: Orally, penicillin therapy.
- V-CILLIN K SULFA.** (Lilly) V-Cillin K (Penicillin V Pot.) 125 mg. (200,000 u.), sulfadiazine 167 mg., sulfamerazine 167 mg., sulfamethoxine 167 mg./Tab. Bot. 50s.
Use: Treatment of mixed infections.
- V-CILLIN-SULFA.** (Lilly) Penicillin V 125 mg. (200,000 u.), sulfamethoxine, sulfadiazine, sulfamerazine 0.167 Gm. each/tab. Bot. 50s.
Use: (1-2 q.i.d.) Antibiotic & sulfonamide therapy.
- V-CILLIN SULFA PEDIATRIC.** (Lilly) Phenoxymethyl penicillin with sulfonamides. Each 5 cc. equals 1 tab Bot. 60 cc.
- V-C-M.** (Stuart) Vit. A 5000 u., D 400 u., Y-1 3 mg., B-2 3 mg., niacin 15 mg., Vit. B-6 0.5 mg., pantothenic acid 5 mg., Vit. C 50 mg., B-12 1 mg., folic acid 0.5 mg./Cal., Phos., iron, copper, cobalt, iodine, zinc & Manganese. Bot. 100s, 1000s.
Use: Dietary supplement.
- V.D.R.L. ANTIDEN.** (Lederle) Cordialipin 0.03%, cholesterol 0.9%, lecithin. Vial 5 cc. antigen mixture, 60 cc., V.D.R.L. buffered saline diluent.
Use: Blood test in diagnosis of syphilis.
- VEIGUM.** (R. T. Vanderbilt Co.) Magnesium aluminum silicate
Use: Emulsifying and suspending agent
- VEGOLYSEN.** See: Hexamethonium Bromide
- VELACYCLINE.** (Squibb) Rolitetracycline.
- VELACYCLINE INTRAMUSCULAR W/XYLCAINE.** (Squibb) N-(pyrrolidinomethyl)tetracycline 150 mg., or 350 mg., Xylcaine 40 mg., ascorbic acid 300 mg./single-dose vial.
Use: Antibiotic.
- VELACYCLINE INTRAVENOUS.** (Squibb) N-(pyrrolidinomethyl)tetracycline 700 mg., ascorbic acid 300 mg./vial.
Use: Antibiotic.
- VELBAN.** (Lilly) Ext. from *Vinca rosea* Linn. Vinblastine sulfate, lyophilized 10 mg./Amp. Vial 10 mg./10 cc.
Use: Antineoplastic drug
- VELVACHOL.** (Texas) Hydrophilic oint. base. Jar 1 lb.
- VINESETIC.** See: Amobarbital Sodium (Various Mfr.)
- VINETHENE.** See: Vinethene. (Merck Sharp & Dohme)
- VENIBAR.** (Ulmer) *Rauwolfia serpentina* whole root 30 mg./tab. Bot. 100s.
Use: Sedative & tranquilizing agent.

VENTREX. (Parke, Davis) Stomach ext 5 gr, ferrous sulfate 2 gr, Vit B 1 0.5 mg, B 2 0.25 mg, folic acid 1 mg /Cap Bot 100s, 1000s

Use: (1 1 d) Antianemic therapy

VENTRICULIN, NNR 1952 (Parke, Davis) Antianemic substance derived from gastric tissue, each 40 gr represents 1 USP u Bot with or without Iron 100 Gm, 500 Gm Use: Orally; pernicious anemia.

VENTRILEX KAPSEALS. (Parke, Davis) Stomach ext 0.3 Gm folic acid 2.0 Gm, liver conc. 0.1 Gm, ferrous sulfate 0.2 Gm, Vit B 1 2.0 mg, ascorbic acid 50 mg /Cap Bot 100s, 1000s

Use: Anti-anemia therapy

VENZOATE, NNR 1952 (Veltax) Benzyl benzoate 25% Lot, Bot 450 cc, 38 L Use: Externally; scabicide.

VERACOLATE (Standard Lab) Bile salts 1 1/2 gr, phenolphthalein 1/2 gr, Capsicum oleoresin 0.05 min, cascara ext 1/2 gr /Tab Bot 50s, 100s, 1000s Also avail w/papsin & pancreatic

Use: (1 1 d) Laxative

VERAFLEX. (Irwin, Neisler) Whole pow veratrum viride 65 u, rutin 20 mg, phenobarbital 15 mg /Cap Bot 100s, 500s, 1000s

Use: Hypertension

VERALBA INJECTION, NNO 1962 (Pitman Moore) Protoveratrine A & B 0.2 mg /cc Vial 10 cc

Use: I.M. & I.V.; treatment of hypertension

VERALBA-R. (Pitman Moore) Protoveratrine A & B 0.4 mg, reserpine 0.08 mg /tab. Bot 100s, 1000s

Use: (4 daily) Hypertension

VERALBA TABLETS, NNR (Pitman Moore) Protoveratrine A & B Inj 0.2 mg /cc Tab (0.2 & 0.5 mg) Bot 100s

Use: Hypertension

VERANCEL OPHTH SOL'N (Professional Pharm) Prophepryridamine maleate, 0.5% phenylephrine HCl, 1/2% la buffered methylolose base Bot 15 cc

Use: Relieve ophthalmic symptoms in hay fever & other allergies

VERAPINE. (Wampole) Reserpine 0.1 mg, protoveratrine A & B 0.4 mg /tab Bot 100s & 500s

Use: 1 1 d, p.c.s hypertension

VERASTAN-MRT. (M. R. Thompson) Contains the total activity of the whole drug Veratrum viride Tab (1 MRT u or 100 mg at V.V. pow) Bot 40s, 100s, 1000s; Elq (1 MRT u./ip) Bot 8 as. 1 gal

Use: 1 2 Tab b.i.d.; essential hypertension

VERATRABAR TABLETS. (Smith Dorsey) Atheroverir 2 mg, mannitol hexanitrate 10

mg, phenobarbital 15 mg /Tab Bot 60s 100s, 500s, 1000s

Use: (1-2 11 d) Hypertension

VERATRITE. (Irwin, Neisler) Veratrum viride 40 u, Sod nitrite 1 gr, phenobarbital 1/2 gr /Tab. Bot. 100s, 500s, 1000s

Use: (1-2 1 d) Hypertension.

VERATRONE. (Parke, Davis) Veratrum viride 0.25 Gm of alkaloids/100 cc Bot 1 fl oz.

Use: Hypertension

VERATRUM ALBA.

See: Protoveratrine A & B (Various Mfr)

Pravell Maleate, Tab (Lilly)

Veralbo, Tab (Pitman-Moore)

Verolba, Inj (Pitman Moore)

VERATRUM VIRIDE, NF Approx 1 gr equals 4 raw units

See: Cryptenamine, NND 1959

Carlone, Vials (Eric, Kirk & Gary)

Unitensin, Tab & Vial (Irwin, Neisler)

Verastan, Liq & Tab (Thompson)

Veratrone, Liq (Parke, Davis)

Veratrol, Vial (Irwin, Neisler)

Vergitryl, Tab (Squibb)

Vergitryl, I.M. (Squibb)

Vergitryl, I.V. (Squibb)

Verloid, Tab (Riker)

Verloid Sol, I.M. (Riker)

Verloid Sol, I.V. (Riker)

W/Atheroxylon See: Rouvera, Tab (Smith-Dorsey)

W/Aconite, omniophylline & atropine sulfate See: Aminite, Tab (Mallard)

W/Monitol hexanitrate, rutin & rauwolfia serpentina See: Roubin, Encate (Poul Maney)

W/Rauwolfia serpentina, whole root See: Palensar Forte, Improved, Tab (Tailby-Nason)

Verlina, Tab (Elder)

Verwalifa, Tab (Thompson)

W/Rauwolfia serpentina, mannitol hexanitrate & homatropine methylbromide See: Rau Perlen, Tab (Crookes Barnes)

W/Rutin, aspirin & Vit C. See: Acelo-Ver, Tab (Mallard)

W/Sod nitrite, acanite, cascara & bile salts. See: Palensars, Tab (Tailby-Nason).

W/Sod nitrite, acanite nitroglycerin & Pot nitrate See: Nyamin, Tab (Elder).

W/Sod nitrite, nitroglycerin, Pot nitrate apocynum fr. & crotaegus fr See: Crotaegonite, Tab (C. O. Smith).

Veratrum Viride with Phenobarbital Combinations on p. 781.

VERATRUM VIRIDE UNIT. See: CSR Unit

VEBAZIPTOL (Newman) Chlorothymol, eucalyptol, menthol, phenol, boric acid & zinc sulfate Pow, Bot 3 oz, & oz Use: Vaginal douche

Veratrum Viride with Phenobarbital Combinations

Veratrum Viride Phenobarbital

Others

| | Manufacturer | Brand |
|--|----------------|---|
| 10 cc. | | |
| (Est.) 1.5 gr. | Irwin-Neisler | Vertavis-Phen, Tab. |
| (Est.) 95 gr. | Paul Maney | Vir-I-Phyl, Tab. |
| 100 mg. | Adams | Verufomin, Tab. |
| | Crookes-Barnes | Pertenal, Tab. |
| (Est.) 20 mg. | Stoddard | Vernille, Tab. |
| 100 mg. | G. F. Harvey | Manujal, Tab. |
| (Est.) 2/5 gr. | F. Trout | Vermojin, Tab. |
| 100 mg. | Rand | Verulol, Tab. |
| 100 mg. | Paul Maney | Vir-I-Tin, Tab. |
| 50 mg. | Sulliff & Case | Manverin, Cap. |
| (Est.) 32 mg. | Mallard | Amvera, Tab. |
| (Tr.) 4 min., 40 cc. | Lemmon | Nipitol, Tab. |
| 1/4 gr. | Irwin-Neisler | Veratril, Tab. |
| 1/4 gr. | Metropolitan | Verital, Tab. |
| 1/4 gr. | Normand | Vi-Rumal, Tab. |
| 1/4 gr. | Tailby-Norton | Polenfort. w/Phenobar- bital, Tab. |
| 1/4 gr. | Superior | Polenfort. w/Phenobar- bital & Rutin |
| 65 cc. | Irwin-Neisler | Verutinim, w/Pheno- barbital, Tab. |
| 1.5 gr. | Paul Maney | Veroflex, Cap. Vir-I-Phyl, Tab. |
| Aconite 20 mg., neohyline 1.5 gr. | | |
| Aminophylline 100 mg., rutin 20 mg. | | |
| Mannitol hexanitrate 30 mg., homatropine methylbromide 2.5 mg. | | |
| Mannitol hexanitrate 1/4 gr., homatropine methylbromide 1 mg. | | |
| Mannitol hexanitrate 1/2 gr., rutin 1/6 gr. | | |
| Mannitol hexanitrate 1/2 gr., rutin 1/2 gr. | | |
| Mannitol hexanitrate 30 mg., rutin 10 mg. | | |
| Mannitol hexanitrate 30 mg., rutin 20 mg. | | |
| Mannitol hexanitrate 30 mg., sod. nitrite 50 mg., eulin 20 mg. | | |
| Mannitol hexanitrate 1/4 gr., rutin 20 mg., choline methio- mine tartrate 100 mg. | | |
| Sod. nitrite 2 gr. | | |
| Sod. nitrite 1 gr. | | |
| Sod. nitrite 1 gr. | | |
| Sod. nitrite 1 gr. | | |
| Sod. nitrite 1 gr., acetate tr. 2 min., cocca 1/2 gr., bile salt 1/4 gr. | | |
| Same as above plus rutin 20 mg. | | |
| Sod. nitrite 1 gr., rutin 20 mg. | | |
| Rutin 20 mg. | | |
| Sod. nitrite 20 mg., neohyline 1.5 gr. | | |

VERCLYSYL INFUSION SOLUTION (Abbott)

Mix of equal parts of dextrose & levulose 5% & 10% w/thiamine HCl 10 mg, riboflavin 10 mg, nicotinamide 250 mg, pyridoxine HCl 5 mg, Vit B 12 3 mcg/1000 cc Bot 1000 cc

VERDEPAM (Texas Pharm Co) Sod propionate 2%, Sod caprylate 2%, propionic acid 3%, endecylenic acid 5%, salicylic acid 5%, copper undecylenate 1/2%, Sod dioctyl sulfosuccinate 0.1% Sol Bot 2 oz
Use: Fungus Infections

VERDEFAM CREAM (Texas Pharm) Sodium propionate 1% sodium caprylate 1%, propionic acid 3%, salicylic acid 3%, undecylenic acid 2%, copper undecylenate 1/2%, vanishing cream base 1 oz tube
Use: Treatment of various fungal infections

VERENTERAL (Irwin, Nelsler) Each cc. contains 100 CSR u of Veratrum viride Vial 20 cc CSR unit is amount necessary to abolish the pressor response of the carotid sinus reflex resulting from occlusion of the carotid arteries

Use: (I V) Severe pre eclampsia & eclampsia

VEREQUAD TABLETS, BUSPENSION (Knoll) Ephedrine HCl 24 mg phenobarbital 8 mg, glyceryl guaiacolate 100 mg theophylline calcium salicylate 130 mg/Tab or 10 ml Tab Bot 24s Susp 4 oz
Use Bronchodilator

VERGITRYL (Squibb) Veratrum viride 1 M. each cc contains 15 u of Squibb standard in aq sol w/acetate buffer, 0.5% chlorobutanol & 1% procaine HCl Vial 10 cc I V each cc contains 0.1 u of Squibb standard with 5% dextrose Inj Vial 5 cc Tab, each tab contains 1 u of Squibb standard Bot 100s
Use: Hypertension

VERGO CREAM (Daywell Labs) Cal pantothenate 3% ascorbic acid 2% Tabe 1/6 oz, 1/2 oz
Use Treatment of plantar warts

VERIBIL (Merck Sharp & Dohme) Dehydrocholic acid 60 mg ox bile ext 60 mg, whole desic pancreas 30 mg, choline ds hydrogen citrate 150 mg/tab Bot 100s 500s
Use Cholagogue

VERIDERM MEROL ACETATE (Upjohn) D 25 or 1% methylprednisolone acetate in synthetic lipid base Tubes 5 Gm
Use: Skin lesions when adrenocortical steroids are indicated

VERIDERM NEO MEDROL ACETATE (Upjohn) D 25 or 1% methylprednisolone acetate .5% neomycin sulfate Tubes 5 Gm

Use: Skin lesions when adrenocortical steroids and broad spectrum antibiotics are indicated

VERILOID INTRAMUSCULAR SOLUTION NND 1962 (Riker) 1 mg of Alkaverin in buffered isotonic aq sol w/procaine HCl 1% cc Amp 2 cc, 6s

Use: Severe hypertension

VERILOID INTRAVENOUS SOLUTION, NND 1962 (Riker) Alkaverin, 0.4 mg/cc Amp 5 cc., 6s

Use All hypertensive emergencies

VERILOID TABLETS, NND 1962 (Riker) Alkaverin Tab 1 mg, 2 mg, 3 mg, Bot 100s 1000s

Use: Treatment of oral forms of hypertension

VERILOID-VPM. (Riker) Veriloid 2 mg phenobarbital 15 mg, mannitol hexanitrate 10 mg/Tab Bot 100s, 1000s

Use: Hypertension

W/Rouviloid (Riker) Alseroxylon 1 mg-alkaverin 3 mg/Tab Bot 100s

Use (I I d) Hypertension

VERINA, VIAL (Testogor) Ethoverine 15 mg/cc Vials 10 cc

Use: Relaxes smooth muscle

VERITAL (Metropolitan) Phenobarbital 1/4 gr, Veratrum viride 3/4 gr Sod nitrite 1 gr/Tab Tab 5 C, orange, Bot 100s 1000s, 5000s

Use: Hypertension

VERITAS VAGINAL KREME, NNR. 1955 (Veritas) Perotormaldehyde 0.1%, oleate glycerine & stearic acid Tube 2 1/2 oz., 4 1/2 oz with or without nozzle

Use: Contraceptive

VERITHOL (Veritas) Oxyquinoline sulfate lactic acid, boric acid aminoacetate acid glycerin, tragacanth Jelly, Tube 3 oz, 5 1/2 oz with or without nozzle

Use: Contraceptive

VERMAGO (Lloyd, Dabney & Westerfield) Piperazine citrate equiv to 500 mg piperazine hexahydrate/5 cc Bot 1 pt

Use: Pinworm and roundworm Infestation

VERMANTIN (F Trout) Phenobarbital 1/4 gr Veratrum viride pow ext 2/5 gr, mannitol hexanitrate 1/2 gr, rutin 1/2 gr/Tab Bot 100s, 1000s

Use: Hypertension

VERMIZINE SYRUP. (Chicago Pharm) Piperazine gluconate equiv to piperazine hexahydrate 100 mg/cc. Bot 8 oz, 1 pt

Use: Treatment of pinworms & roundworms.

VERNACEL (Professional Pharmacol Co) Prophepryridamine moleate 1/2%, phenylephrine HCl 1/2% in a methylcellulose solution Plastic Bot 15 cc

Use: Ophthalmic allergic symptoms

- VERNITE.** (Stoddard) Phenobarbital 8 mg., Veratrum viride ext. 20 mg., monninal hexanitate 16 mg., homotropine methylbromide 1 mg./Tab. Bot. 100s, 500s, 1000s.
Use: (1-2 I.Ld.) Vasodilator w/sedative.
- VERNAL SODIUM.** See: Barbitol Sodium (Various Mfr.).
- VEROPHYLIN.** (Mallord) Aminophylline $\frac{1}{2}$ gr., veratrum viride ext. 1 gr., ocanite pow. 1/5 gr., atropine sulf. 1/1000 gr., phenobarbital $\frac{1}{4}$ gr./Tab. Bot. 100s, 1000s
Use: Treatment of hypertension
- VERSACAIN.** (Lederle) Chlorprocaine hydrochloride 2%, epinephrine 1:100,000
Use: Local anesthetic for use in dentistry.
- VERSENATE, CALCIUM DIBODIUM, N.N.D.** 1962. See: Calcium Disodium Versenate, Amp. (Riker)
- VERSENE.** (Various Mfr.) Tetrasodium salt of ethylenediaminetetraacetic acid. Pkg. according to demand. See: Edothomil Tetrasodium
Use: Sequestering agent.
- VERTAVIB.** (Irwin, Nelster) Veratrum viride. Tab. 10 Crow u., Bot. 100s, 500s, 1000s.
Use: Severe, resistant essential hypertension.
- VERTAVIS HALF-STRENGTH.** (Irwin, Nelster) Veratrum viride. Tab. 5 Crow u., Bot. 100s, 500s, 1000s.
Use: Hypertension.
- VERTAVIS-PHIN.** (Irwin, Nelster) Veratrum viride 10 crow u. & phenobarbital $\frac{1}{2}$ gr./Tab. Bot. 100s, 500s, 1000s.
Use: Essential hypertension.
- VERTINA.** (Elder) Rauwolfia serpentina whole root 50 mg., veratrum viride alkaloids 1.1 mg./tab. Bot. 100s.
Use: (i p.c. & h.s.) Treatment of hypertension.
- VIBUTAL.** (Rond) Phenobarbital $\frac{1}{4}$ gr., monninal hexanitate 30 mg., Veratrum viride 100 mg., rutin 10 mg./Tab. Bot. 100s, 1000s.
Use: (1-2 I.Ld.) Vasodilator w/sedative in treatment of hypertension.
- VERUTAMIN.** (Adams) Phenobarbital 15 mg., aminophylline 100 mg., Veratrum viride 95 mg., rutin 20 mg./Tab. Bot. 100s, 500s.
Use: (1 I.Ld.) Vasodilator w/sedative.
- VERUTINIUM.** (Superior Pharm.) Veratrum viride $\frac{3}{4}$ gr., Sod. nitrite 1 gr., rutin $\frac{1}{2}$ gr./Enpule. Bot. 90r.
Use: Hypertension.
- W/Phenobarbital.** (Superior) Phenobarbital $\frac{1}{4}$ gr., veratrum viride $\frac{1}{4}$ gr., Sod. nitrite 1 gr., rutin $\frac{1}{2}$ gr./Enpule. Bot. 90s.
Use: Hypertension.
- VERY ALERTNESS CAPSULES.** (APC) Cof-faine 200 mg./Cap. Vial 15.
- Use: Maintain physical and mental alertness.
- VERWOLFIA.** (Thompson) Rauwolfia serpentina whole root 50 mg., veratrum viride whole root 50 mg./tab. Bot. 100s
Use: Treatment of hypertension
- VESIPAQUE.** (Warner-Lombert Res. Inst) Phenobutadol.
Use: Bodily opaque diagnostic aid
- VESPIN, N N D 1962.** (Squibb) Trifluoromethyl-2-(3-Dimethylaminopropyl)-2-(trifluoromethyl) phenothiazine HCl Tab. 10, 25, 50 mg, Bot 50s, 500r Inj. (20 mg./cc) 1 cc, Box 5s. (10 mg./cc) 10 cc. Inj. dirp. unit (0.75 mg./cc) 1 cc. Emulsion (10 mg./cc) Bot 30 cc., 120 cc, Caps. 100 mg. Bot. 100s, Oral Liq. (10 mg./cc) Bot 30 & 125 cc. I.V. 3 mg./cc, Vials 10 cc.
Use: Tranquillizer
- VESPIN INJECTION UNIMATIC.** (Squibb) Trifluoromethyl-2-(3-Dimethylaminopropyl)-2-(trifluoromethyl) phenothiazine HCl 15 mg./0.75 cc. In new type disposable syringe. Parenteral 15 mg. disposable syringe
Use: Psychotropic.
- VITALOG.** (Squibb) Triamcinolone acetonide 6 mg., benzyl alcohol 0.9%/cc. Susp.
Use: Treatment of dermatoses.
- VIACAPB.** (Manno Labs) Vit. A (soluble) 50,000 units; C 500 mg./Caps. Bot. 60s, 120s, 250s, 500s, 1000s
Use: Acne therapy.
- VIACETS.** (Walker) Vit. A 5000 u., D 1000 u., B-1 1 mg., B-2 1 mg., B-6 1 mg., B-12 2 mcg., col. pontathene 1 mg., niacinamide 5 mg., folic acid 0.1 mg., E 1 I.U., C 50 mg./Candisphere. Bot. 120s
Use: Nutritional supplement
- VIADRIE, N N D 1962.** (Pfizer) Hydroxydl-ones sodium; 21-hydroxy-pregnandione sod. succinate. Vial 500 mg. (dry)
Use: Surgical anesthetic in 1% sol.
- VIAGE CAPS.** (Manno Labs) Methamphetamine HCl 5 mg; methylterterterone 1.25 mg; ethinyl estradiol 5 mcg; Vit. B-1 7.5 mg; B-2 2.4 mg; B-6 2 mg; B-12 3 mcg; niacinamide 30 mg.; L-Lysine monohydrochloride 25 mg; iron (as ferrous fumarate) 10 mg; choline 25 mg; methionine 100 mg /Caps. Bot. 60s, 120r, 250r, 500s, 1000s.
Use: Geriatric.
- VI-ALPHA.** (Lederle) Vit. A 50,000 u /Cap. Bot. 50s, 100r, 1000s.
Use: Vitamin A deficiency.
- W/Tocopheryl Acetate** (Lederle) Vit. A 50,000 u., d-alpha-tocopheryl acetate 368 mg /Cap. Bot. 100, 1000s.
Use: Vitamin A & E deficiency.
- VI-AQUA CAPSULES.** (U. S. Vitamin) Vit. A 5000 u., D 500 u., thiamine HCl 5 mg., riboflavin 5 mg., Vit. B-12 1 mcg., nic-

cinamide 20 mg, pyridoxine HCl 0.5 mg, Cal pantothenate 5 mg, ascorbic acid 50 mg, dl alpha tocopheryl acetate 1 mg / Cap Bot 50s, 100s, 500s 1000s

Use: Multiple vitamin deficiencies

VI-AQUA SYRUP. (U S Vitamin) Vit A 4000 u, D 800 u, B 1 1.5 mg, B 2 12 mg B 6 0.3 mg, C 50 mg, nicotinamide 10 mg, pantothenic acid 2 mg / 5 cc Bot 4 oz, 1 pt 1 gal

Use: (1-2 dr daily) Multivitamin therapy

VI-AQUA THERAPEUTIC CAPSULES. (U S Vitamin) Vit A 12,000 u, D 1000 u, C 150 mg, B 1 10 mg, B 2 5 mg, B 12 2 mcg, niacinamide 100 mg, Vit B 6 1 mg, Cal pantothenate 10 mg, alpha tocopheryl acetate 5 mg / Cap Bot 30s 100s 500s

Use (1-2 daily) Therapeutic multiple vitamin

VI-AQUAMIN. (U S Vitamin) Vit A 5000 u D 800 u, B 1 3 mg, B 2 3 mg, B 6 0.5 mg, C 50 mg, nicotinamide 25 mg, pantothenic acid 5 mg, Vit E 1 mg, B 12 1 mcg w/Cal Phos iron copper, iodine, Mang, Mag zinc, cobalt / Cap Bot 100s, 500s 1000s

Use: (1 daily) Multivitamin & mineral therapy

VI-AQUAMIN THERAPEUTIC (U S Vitamin) Vit A 25,000 u, D 1000 u, C 150 mg, B 1 10 mg niacinamide 100 mg, Vit B 2 5 mg B 6 1 mg, B 12 5 mcg Cal pantothenate 5 mg Vit E 5 mg, dicalcium phosphate 200 mg, w/iron, copper, iodine Mang, Mag, zinc, cobalt, molybdenum / Cap Bot 50s 100s 500s

Use: Multivitamin w/mineral therapy

VI-ASPIRIN (Amuro) Aspirin 325 mg ascorbic acid 30 mg / Tab Bot 25s 100s

Use: Colds

VIATRIC (Omas Wilcox) Ethinyl estradiol 0.01 mg methyltestosterone 2 mg, d Amphetamine sulf 1 mg, ferrous sulf 60 mg, l lysine 50 mg, Vit A 5000 u D 500 u, B 1 5 mg, B 2 5 mg, B-6 1 mg, nicotinic acid 15 mg, cal pantothenate 10 mg, folic acid 1 mg, B 12 1 mcg, intrinsic factor conc 0.7 mg, cal ascorbate 30 mg, menadione 1 mg, Vit E 5 u w/trace elements / Cap Bot 100s 1000s

Use: Androgen estrogen therapy with nutritive vitamin and mineral supplements

VIAZOLE (VanPelt & Brown) Sulfanilamide 5%, sulfathiazole 5%, urea 2%, Vit D 700 u / oz Tube 1 oz Jar 1 lb

Use: Sulfanamide susceptible skin infections

VI-4-B FORTIS (Drug Products) Thiamine HCl 100 mg, riboflavin 2 mg, nicotinamide

100 mg, pyridoxine HCl 2 mg / cc. Vial 10 cc, 30 cc.

Use: Vitamin B therapy

VIBALT, NND 1959 (Roerig) Vit B 12 200 & 1000 mcg / cc Vial 5 cc.

Use: Hematinic

VIBAZINE HYDROCHLORIDE. (Pfizer) Buzilzine HCl Tab (50 mg) Bot 100s

Use: Antihistaminic.

VIBEFOL. (Brewer) Vit B-12 25 mcg, folic acid 2 mg / Cap Bot 100s

Use: Pernicious anemia

***VIBESATE, NND 1962** Polivite 93% and malrasal 31% with propellants See Aeroplast, Spray (Aeroplast Corp)

VIBETA. (Columbus) Vit B 1 1.5 mg, Vit B 2 2 mg, niacinamide 10 mg, cal pantothenate 1 mg, Vit B 6 1 mg, dried yeast 100 mg, liver desiccated 70 mg / Cap Bot 100s, 1000s

Use: Dietary supplement

VIBETA No 3. (Columbus) Thiamine chloride 10 mg, niacinamide 200 mg, riboflavin 5 mg, pantothenic acid 2.5 mg pyridoxine HCl 2.5 mg, iron cacodylate 15 mg / cc Vial 30 cc

Use: Vitamin B therapy

VIBETA-C. (Columbus) Thiamine HCl 500 mg, riboflavin 50 mg, niacinamide 500 mg, pyridoxine HCl 100 mg, Sod pantothenate 100 mg, ascorbic acid 1000 mg / 10 cc Vial 10 cc

VIBEX. (Parke, Davis) Thiamine HCl 12 mg riboflavin 24 mg, pyridoxine HCl 6 mg, nicotinamide 120 mg, pantothenic acid 12 mg / 8 oz Bot 4 oz

Use: Vitamin B therapy

VI-BLOIDS, (Rexall) Vit A 25,000 u, D 1000 u, C 150 mg, B 1 10 mg, B 2 10 mg niacinamide 60 mg, Vit E 10 mg / Cap Bot 100s

Use: Multivitamin therapy

VIBRUOEN (Philadelphia Ampoule Labs) Adenosine 5 monophosphoric acid 50 mg, Vit B 12 100 mcg, nicotinic acid 20 mg, procaine HCl 1%, benzyl alcohol 1%, cc Vial 10 cc

Use: IM, treatment of bursitis & osteoarthritis

VIBUMIN (Philadelphia Ampoule Labs) Vit B 12 Vial (30 mcg, 60 mcg 100 mcg / cc) 10 cc 30 cc Vial (250 mcg / cc) 10 cc Vial (500 mcg / cc) 10 cc Vial (1000 mcg / cc) 5 cc, 10 cc Vial (2000 mcg / cc) 10 l

VICALIBON (Chicago Pharm) Ferrous sulfate 1 1/2 gr, Cal phosphate dibasic 3-gr, Vit D 250 u / Tab Bot 500s, 1000s

Use: Dietary supplement

* Generic name recognized by NP USP or AMA

VICALMIN. (C. D. Smith) Cal. carbonate 400 mg., ferrous sulf. essic. 150 mg., vit. A 2500 u., D 500 u., cal. ascorbate 60 mg., B-1 1 mg., B-2 1 mg., niacinamide 10 mg., B-12 1 mcg., K 1 mg., folic acid 0.25 mg., B-6 3 mg., w/iodine 0.05 mg., manganese 0.33 mg., cobalt 0.033 mg., pot. 1.66 mg., molybdenum 0.066 mg., copper 0.33 mg., zinc 0.40 mg., magnesium 2 mg./tab. Bot. 100s, 1000s

Use: Pregnancy & lactation.

VICAM. (Testagor) Vit. A 2500 u., D 250 u., B-1 1 mg., C 33 mg., B-6 0.1 mg., intrinsic factor w/B-12 1/64 u., Vit. B-12 3 mcg / Pushile Bot. 100s

Use: Multi-vitamin therapy

VICIN. (Biewer) Ascorbic acid, Amps (100 mg) 2 cc., Pkg. 12s, 100s, (100 mg/cc) 5 cc., Pkg. 25s, 100s, (500 mg) 5 cc., Pkg. 6s, 25s, 100s, (1000 mg.) 10 cc., Pkg. 6s, 25s, 100s.

Tab. (25, 50, 100, 250 & 500 mg.) Bot. 100s, 500s, 1000s.

Use: Vitamin C therapy.

VICKS TRI-SPAN. (Vick Chemical Co) Acetyl p aminophenol 3 gr., pyrilamine maleate, caffeine, ephedrine sulfate, phenylpropanolamine HCl/Time-care Tab. Bot. 16s.

Use: Analgesic with vasoconstrictors.

VICKS VAPOSTEAM. (Vick Chemical Co) Polyisobutylene dodecanol, menthol, camphor, oil of eucalyptus, fr. of benzoin, & alcohol, Bot. 3 fl. oz.

Use: Steam medication.

VICKS COUGH SYRUP. (Vick Chemical Co) d Methaphon HBs, cetylpyridinium chloride, sod. citrate, omm. chloride, eucalyptus, menthol, camphor & other aromatics. Bot 3 1/2 oz., 6 fl. oz.

Use: Control of coughs.

VICKS FORMULA 44. (Vick Chemical Co) d Methaphon HBs, dextylamine succinate, cetylpyridinium chloride, sod. citrate & chloroform Bot. 3 1/2 fl. oz., 6 fl. oz.

Use: Control of coughs.

VICON.M. (Meyer) Vit. A 25,000 u., D 1000 u., thiamine mononitrate 10 mg., riboflavin 5 mg., nicotinamide 100 mg., pyridoxine HCl 1 mg., C 150 mg., B-12 5 mcg., calcium 103.6 mg., phosphorus 80.2 mg., van 10 mg., cobalt 0.1 mg., copper 1 mg., iodine 0.1 mg., magnesium 5.56 mg., manganese 0.954 mg., zinc 1.4 mg., molybdenum 0.17 mg., potassium 4.93 mg./Tab. Bot. 100s, 1000s.

Use: Vitamin and mineral supplement.

VI-CUTA SKIN CREAM. (Columbus) Pantothenyl alcohol 1%, pyridoxine HCl 0.5%.

hexachlorophene 0.5%, menthol 0.2%. Tube 1 oz.

Use: Topically, dermatitis.

VIDAC. (Endo) Vit. A 5000 u., D 1200 u., C 60 mg., wheat germ oil/0.6 cc. Bot. w/dropper 15 cc., 30 cc.

Use: Dietary supplement

VI-DAYLIN. (Abbott) Vit. A 3000 u., D 1000 u., thiamine HCl 1.5 mg., ascorbic acid 50 mg., B-2 1.2 mg., B-6 0.5 mg., nicotinamide 10 mg., Vit B-12 3 mcg /0.6 cc New 12 fl. oz "Pressure Pak" 3 oz., B ac., 1 pt.

Use: Multiple vitamin therapy.

VI-DAYLIN AEROSOL. (Abbott) Vit A 3000 u., D 800 u., thiamine HCl 1.5 mg., riboflavin 1.2 mg., ascorbic acid 40 mg., nicotinamide 10 mg., pyridoxine HCl 1 mg., B-12 3 mcg./5 cc. Push button container 12 fl. oz Bot 3 fl. oz., B fl. oz., pts.

Use: Pediatric vitamin supplement.

VI-DAYLIN CHEWABLE DULCET. (Abbott) Vit. A 3000 u., D 400 u., thiamine mononitrate 1.5 mg., riboflavin 1.2 mg., nicotinamide 10 mg., cobalamin 3 mcg., pyridoxine HCl 1 mg., ascorbic acid 50 mg / Tab. Bot. 30s, 100s

Use: Vitamin supplement

VI-DAYLIN DULCET. (Abbott) Vit. A 3000 u., D 800 u., thiamine mononitrate 1.5 mg., riboflavin 1.2 mg., pyridoxine HCl 1 mg., B-12 3 mcg., ascorbic acid 40 mg., nicotinamide 10 mg / Tab. Bot. 30s, 100s

Use: Pediatric vitamin supplement.

VIDAYLIN.M. (Abbott) Vit. A 3000 u., Vit. D 800 u., B₁ 1.5 mg., B₂ 1.2 mg., C 40 mg., Nicotinamide 10 mg., B₆ 1 mg., B₁₂ 3 mcg., Panthelol 5 mg., iron 3 mg., iodine 75 mcg., Calcium 40 mg., phosphorus 20 mg., manganese 0.5 mg., potassium 25 mg., magnesium 3 mg., Zinc 0.5 mg., choline 5 mg., Inositol 5 mg., Cysteine HCl 5 mg / 5 cc. Bot. 3 fl. oz., B fl. oz., 1 pt.

Use: Nutrition supplement for children

VI-DAYLIN-V. (Abbott) Vit. A 5000 u., D 800 u., B-1 3 mg., B-2 3 mg., B-6 1 mg., nicotinamide 20 mg., ascorbic acid 75 mg., B-12 3 mcg./5 cc. Bot. 3 oz., B ac., 1 pt.

Use: For infants, dietary supplement

VI-DE-CAPS. (Dumas-Wilson) Vit. A 25,000 u., D 1000 u., B-1 15 mg., B-2 7.5 mg., B-6 2 mg., C 150 mg., nicotinamide 150 mg., cal. pantothenate 10 mg., B-12 5 mcg., folic acid 1 mg., E 20 mg., with trace elements/Cap. Bot. 100s, 1000s.

Use: Vitamin and mineral deficiency.

VI-DELTA EMULSION. (Ledette) Vit. A 2500 u., D 375 u./dr. Bot. B ac., 1 pt.

Use: (1 dr. t.i.d.) Vitamin A B D therapy.

- VI-DERM SOAP** (Arthrins) Bolan, an extract of *Amaryllidaceae* Pkg 1 cake
Use: Hypo allergenic and antipruritic
- VIDEROL** (Parke, Davis) Vit D 50,000 u, mixed tocopherol 50 mg, whole bits 150 mg /Cap Bot 100s 500s
Use: (1 2 11 d) Chronic arthritis & associated fibrositis
- VI-DEXEMIN** (Smith, Kline & French) Dextrose sulfate 5 mg Vit A 5000 u, D 500 u, B 1 3 mg B 2 2 mg B-6 0.5 mg, Cal pantothenate 1 mg Vit B 12 1 mcg folic acid 0.4 mg, Vit C 25 mg, nicotinamide 20 mg Cal phosphate 500 mg, iron 12 mg, iodine 0.1 mg /Tab Bot 100s
Use: (1 1 d, a c) Anti obesity
- VI-DIEM** (McNeil) Vit A 5000 u, thiamine HCl 1 mg, riboflavin 2 mg nicotinamide 10 mg, ascorbic acid 30 mg, Vit D 400 u /Tab Bot 100s, 500s, 1000s
Use: Dietary supplement
- VI-DI-PEN** (Vidarine) Procaine penicillin G 300M u, Pot penicillin G 100M u, dihydro streptomycin sulfate D.5 or 1 Gm /dose Vial 1 dose
Use: I.M. penicillin & streptomycin therapy
- VI-DOM-A CREME** (Dome) Vit A 100,000 or 500,000 u /3D Gm in greaseless base Tube 1, Jar 2, 4 oz & 1 lb
Use: Topically stimulate healing of skin conditions
- VI-DOM-A ORAL** (Dome) Vit A 150,000 u or 75,000 u /Tab Bot 25s 100s
Use: Vitamin A deficiencies
- VI-DOM-A PILLETTES** (Dome) 25,000 u Bot 100s
Use: Vitamin A therapy
- VI-DOM-A-C** (Dome) Vit A 25,000 u, C 125 mg /Tab Bot 100s
Use: (buccal) Acne
- VI-DOM-C** (Dome) Vit C 1,335 mg /Ruccal Tab Bot 25s, 60s 100s
Use: Treatment of scurvy
- VI-DOM-D-2 PILLETTES** (Dome) Not listed Vit A 5000 u, Calciferol 50,000 u /Pilette Bot 100s 500s 1000s
Use: (1 daily) Vit A & D deficiencies
- VI-DOM 'E'** (Dome) Syn d-alpha tocopheryl acetate 50 mg /p illette Bot 30s 100s 1000s
Use: p.r.n.t; neuromuscular disorders
- VI-DRAPE** (Aeroplast) One dozen rolls per box
Use: Plastic film
- VI EA** (Miller) alpha Tocopherol acetate Cap 50 u, 100 u, Bot 100s, 500s, 1000s Vial (100 mg /cc.) 15 cc., 30 cc. (200 mg /cc.) 10 cc.
Use: Vitamin E therapy
- VI-E CAPS.** (Blue Line) d alpha Tocopherol acetate Cap 50 mg, Bot 100s
Use: (1 daily) Vitamin E therapy
- VI-FERRIN CAPSULES** (Lederle) Ferrous sulfate dried 200 mg, folic acid 1 mg, Vit B 1 0.25 mg, B 2 0.19 mg, niacinamide 2.5 mg, Cal pantothenate 4 mg, choline 10 mg insoluble liver fraction 500 mg, Vit B 12 1 mcg /cap Bot 100s
Use: Hematinic & vitamin deficiencies
- VI-FERRIN TABLETS** (Lederle) Ferrous sulfate dried 200 mg, folic acid 1 mg, Vit B 1 0.25 mg, B 2 0.35 mg, niacinamide 2.5 mg, Cal pantothenate 1 mg, choline 4 mg, insoluble liver fraction 250 mg, Vit B 12 1 mcg /tab Bot 100s, 1000s
Use: Hematinic & vitamin deficiencies
- VIELAVA** (Calumbe) Citrus bioflavonoid 100 mg, ascorbic acid 100 mg, menadiolone 0.5 mg /Cap Bot 100s, 500s, 1000s
Use: (1 qid) In diseases associated with capillary fragility
- VIEO-IRON** (VonPell & Brown) Ferrous gluconate 100 mg, ferrous sulfate 100 mg, Vit B 1 1 mg, B 2 2 mg, B-6 0.5 mg, C 30 mg, niacin 5 mg, pantothenic acid 1 mg, folic acid 1 mg, liver conc (1:20) 50 mg /Tab Bot 100s, 500s
Use: (1 2 11 d) Hematinic
- VIFOLIRON** (VonPell & Brown) Vit B 1 1 mg, B 2 2 mg, B-6 0.5 mg, C 30 mg, folic acid 1 mg, nicotinamide 5 mg, ferrous sulfate 100 mg, ferrous gluconate 100 mg, liver conc 50 mg /Tab Bot 100s 500s
Use: (1 2 11 d) Hematinic A vitamin deficiencies
- VIEORT CAPSULES** (Endo) Vit A 5000 u, O 500 u, C 60 mg, B-1 3 mg, B 2 2 mg, niacinamide 20 mg, Vit B-6 0.3 mg, Cal pantothenate 5 mg, tocopherol 2 mg /Cap Bot 30s, 100s, 250s, 1000s
Use: Polyvitamin therapy
- VIFORT DROPS** (Endo) Vit A 5000 u, O 12000 u, C 60 mg B 1 1 B mg, B 2 0.6 mg, niacinamide 8 mg, Vit B-6 0.3 mg, Cal pantothenate 1.2 mg /0.6 cc Bot w/drop- per 15 cc, 30 cc., 60 cc.
Use: Polyvitamin therapy
- VIEORT NECTAR** (Endo) Vits A 5000 u, O 1000 u, C 50 mg, B 1 3 mg, riboflavin 3 mg, niacinamide 10 mg, pyridoxine 1 mg panthanol 10 mg, cyanocobalamin 5 mcg /Tsp Bot 4 oz., 1 pt
Use: Polyvitamin therapy
- VIGITARS** (Ascher) Phenylmercuric acetate 3 mg, methylbenzethonium chloride 1 B mg, 9-aminoacridine HCl 2 mg, succinic acid 12.5 mg, allantoin 20 mg, lactose 0.75 Gm. /Tab Bot 30s, 100s
Use: Yeagnat antiseptic.

VIGORAMIN. (Consolid. Midland) Vitamins A 5000 u., D 500 u., B-1 1.5 mg., B-2 1.5 mg., B-6 0.1 mg., C 37.5 mg., E 0.34 u., cal. pantothenate 1.5 mg., niacinamide 15 mg./spheres. Bot. 100s, 250s, 1000s, 5000s.

VIGORAMIN THERAPEUTIC. (Consolid. Midland) Vitamin A 25,000 u., D 1000 u., B-1 10 mg., B-2 5 mg., C 150 mg., niacinamide 150 mg./sphere. Bot. 30s, 100s, 250s, 1000s.

VIGRAN. (Squelbb) Vit. A 5000 u., D 500 u., B-1 2 mg., B-2 2 mg., pyridoxine HCl 0.5 mg., niacinamide 20 mg., Vit. C 45 mg., Cal. pantothenate 1 mg./Cap. Bot. 25s, 50s, 100s, 250s, 1000s. Liq. Bot. 8 oz.

Use: Multiple vitamin therapy.

VIGRAN CHEWABLES. (Squelbb) Provide adults and children with at least 125% of the M.D.R. of vitamins. Vit. A acetate 5000 u., D 500 u., thiamine mononitrate 3 mg., riboflavin 3 mg., pyridoxine HCl 1 mg., cyanocobalamin 2 mcg., niacinamide 12.5 mg., Vit. C 50 mg., calcium pantothenate 5 mg./Tab. Bot. 30s, 90s.

Use: Multivitamin.

VIGRAN M. (Squelbb) Same as Vigran plus minerals. Bot. 30s, 100s

VI-HESTIB. (Aeroplast) Pkg. 12 oz.
Use: A clear adhesive especially formulated for adhering plastic film or surgical gauze to the skin.

VIKALUM. (C. D. Smith) Kaolin colloidal 96 gr., aluminum hydroxide 12 gr., thiamine HCl 4 mg./fl. oz. Bot. 4 fl. oz., 1 pt., 1 gal.
Use: Gastric antacid.

VI-LITRON. (U. S. Vitamin) Syr., liver conc. (equiv. to whole liver 10 Gm.), iron pantothenate & iron ammonium citrate 5½ gr., thiamine HCl 2 mg., riboflavin 2 mg., niacinamide 10 mg., d-Cal. pantothenate 1 mg., pyridoxine HCl 0.25 mg., brewer's yeast ext. 1 Gm./dr. Bot. 4 oz., 8 oz., 1 pt. Cap., liver conc. (equiv. to whole liver 12 Gm.) 0.218 Gm., ferrous sulfate 3 gr., Vit. C 20 mg., thiamine HCl 2 mg., riboflavin 2 mg., niacinamide 10 mg., d-Cal. pantothenate 1 mg./Cap. Bot. 50s, 100s, 200s, 500s.

Use: Multiple vitamin therapy.

VI-LITRON THERAPEUTIC. (U. S. Vitamin) Vit. B-12, B-12b 10 mcg., folic acid 1 mg., liver conc. 300 mg., ferrous sulfate 185 mg., ascorbic acid 50 mg., thiamine HCl 2 mg., riboflavin 2 mg., niacinamide 10 mg., pyridoxine HCl 0.5 mg., d-Cal. pantothenate 1 mg./Cap. Bot. 50s, 100s, 200s, 500s.

Use: Anemias.

W/Folic acid. (U. S. Vitamin) Liver conc. 0.218 Gm., ferrous sulfate 3 gr., ascorbic acid 20 mg., folic acid 2 mg., thiamine

2 mg., riboflavin 2 mg., niacinamide 10 mg., d-Cal. pantothenate 1 mg., liver 12 Gm./Cap. Bot. 50s, 100s, 200s, 500s.

Use: Anemias.

W/Folic acid. (U. S. Vitamin) Liver conc. 0.218 Gm., ferrous sulfate 3 gr., ascorbic acid 20 mg., folic acid 2 mg., thiamine

2 mg., riboflavin 2 mg., niacinamide 10 mg., d-Cal. pantothenate 1 mg., liver 12 Gm./Cap. Bot. 50s, 100s, 200s, 500s.

Use: Anemias.

VIMACOL. (Brewer) Vit. A 3334 u., D 333 u., B-1 0.67 mg., B-2 1.4 mg., iron & ammonium citrate 12 gr., red bone marrow ext. glycerinated 10 min., alcohol 5%, malt ext. (non-diastatic) q.s./fl. oz. Bot. 1 pt., 1 gal.

Use: Dietary supplement.

VIMAGNA. (Lederle) Vit. A 5000 u., D 500 u., thiamine HCl 3 mg., riboflavin 3 mg., niacinamide 20 mg., Cal. pantothenate 1 mg., pyridoxine HCl 0.2 mg., ascorbic acid 75 mg., Folvite (folic acid) 1 mg., Vit. B-12, 1 mcg./Cap. or 5 cc. Syr. Cap. Bot. 100s, 250s, 1000s. Syr., Bot. 4 oz., 1 pt., 1 gal.

Use: Dietary supplement.

VIMINASCORBE. (Kirkman) Combines 11 vitamins & 11 minerals/cap. Bot. 100s, 1000s.

VIMINETS. (Richard & Williams) Vit. A acetate 5000 u., D-2 500 u., B-1 3 mg., B-2 2 mg., B-6 0.1 mg., B-12 1 mcg., C 30 mg., calcium pantothenate 1 mg., niacinamide 20 mg., folic acid 0.2 mg., iron 15 mg., copper 1 mg., manganese 0.5 mg., zinc 0.5 mg., calcium 50 mg., phosphorus 40 mg., magnesium 0.5 mg., iodine 0.1 mg./Tab. Bot. 100s.

Use: Vitamin and mineral supplement.

VI-MIX. (Lilly) Vit. A 5000 u., D 1000 u., C 75 mg., B-1 1 mg., B-2 1 mg., niacinamide 10 mg., pantothenic acid 3 mg., Vit. B-6 0.5 mg., B-12 3 mg./O 6 cc. Bot. 30, 60 cc.

Use: Multivitamin therapy.

VINACTANE SULFATE, N.N.D. 1962. (Ciba) Viomycin sulfate. Vial 1.5 Gm.
Use: Antituberculous antibiotic.

* **VINBABBITAL, N.F., N.N.D.** 1962. See: Delvinol (Merck Sharp & Dohme)

* **VINBABBITAL SODIUM, N.F., N.N.D.** 1962. Cap., N.F. Inj., N.F. 5-Ethyl-5(1-methylbut-1-enyl)-borbiturate sod. See: Delvinol (Merck Sharp & Dohme)

W/Aspirin. See: Delvinol-Aspirin Capsules (Merck Sharp & Dohme)

W/Belladonna alkaloids. See: Delvinol Tablets (Merck Sharp & Dohme)

W/Diethylstilbestrol. See: Delvinol-Diethylstilbestrol, Tab. (Merck Sharp & Dohme)

* **VINBLASTINE SULFATE.** Vincalcestoblastine. Alkaloid extracted from *Vinca rosea* Linn. See: Velban (Lilly)

VINCALEUKOBLASTINE. [Lilly] Alkaloid from *Vinca rosea* See: Vinblastine
Use: Treatment of various kinds of cancer.

- VINCE.** (Standard Labs) Sod barate peroxy hydrated 37%, Cal carb 33.8%, Sod aluminum sulfate anhy 24% Mag trisilicate 3% Cal phosphate tribasic 1% Pkg 2 oz 5 oz
 Use: Oral Infections
- ***VINCRIStINE** (Eli Lilly & Co) An alkaloid extracted from *Vinca rosea* Linn
 Use: Anticancer agent
- VINETHENE** (Merck Sharp & Dohme) Vinyl Ether U.S.P. Divinyl Oxide Bot 10 cc 25 cc, 50 cc 75 cc
 Use: Inhalation anesthetic
- VINISIL** (Abbott) Povidone; polyvinylpyrrolidone; PVP Sol (2.5% in Ringer's Sol or 3.5% in saline). Abbotlet bot 500, 1000 cc
 Use: 500-1000 cc IV; plasma extender
- ***VINEUROStINE** (Eli Lilly & Co) An alkaloid extracted from *Vinca rosea* Linn
 Use: Anticancer agent
- VINOTHIAM.** (Merck Sharp & Dohme) Thiamine HCl 8 mg./fl oz Bot 1 pl, 1 gal
 Use: Vitamin B-1 deficiency
- VI-NOVEM** (Corrigan) Vit A 5000 u, D 500 u, B 1 3 mg B 2 2.5 mg, C 50 mg, nicothamide 20 mg, B-6 1 mg, B 12 2 mcg pantothenic acid 1 mg/Cap Bot 100s
 Use: Dietary supplement
- ***VINROSIDINE** (Eli Lilly & Co) An alkaloid extracted from *Vinca rosea* Linn
 Use: Anticancer agent
- VINUFER.** (C. D. Smith) Ferrous gluconate 100 mg, vit B-12 10 mcg, B 1 2 mg, B 6 2 mg./5 cc. liq, Bot 1 pt
 Use: Treatment of anemias
- VI-NUTRO.** (Vitamin Corp) Vit A 10,000 u, D 2400 u, C 75 mg, B 1 4 mg, B-2 1.5 mg, niacinamide 10 mg., Vit B-6 D 6 mg., Cal pantothenate 2 mg, tacopherole mixed 5 mg./cc. Bot 15 cc., 30 cc.
 Use: Multivitamin drops
- ***VINYL ETHER, U.S.P., N.N.R., A.D.R.** See Vinethene (Merck Sharp & Dohme)
- VINYZENE** (Scientific Chemicals) Bromochlorenone
 Use: Fungicide and bactericide
- VIO-12.** (Rowell) Vit A 10,000 u., D 1200 u., B-1 10 mg., B-2 10 mg., B-6 1 mg., B-12 1 mcg., C 150 mg., niacinamide 40 mg., Cal pantothenate 10 mg., sholine 20 mg., inositol 20 mg., folic acid D.5 mg./Cap Bot 100s, 1000s
 Use: Dietary supplement
- VIO-ALKA-PHEN.** (Rowell) Homotropine methylbromide 1/20 gr., Sod secobarbital 1/8 gr., Cal carb. 3 1/2 gr., charcoal 2 1/2 gr./Cap Bot 100s, 500s, 1000s
 *Generic name recognized by N.F., U.S.P. or A.M.A.
- Use: [12 p.c. & h.s.] Antacid w/antispasmodic
- VIO-B** (Rowell) Vit B 1 5 mg., B 2 5 mg., Cal pantothenate 10 mg., Vit B 6 0.5 mg., niacinamide 20 mg., brewer's yeast ext 3 1/2 gr., liver, dehy 3 1/2 gr./Cap Bot 100s 1000s
 Use: Dietary supplement
- VIO-BEC.** (Rowell) Thiamine HCl 25 mg., riboflavin 25 mg., niacinamide 100 mg., Cal pantothenate 40 mg., pyridoxine 10 mg., ascorbic acid 250 mg./Cap Bot 100s 1000s
 Use: Vitamin & mineral supplement
- VIO-BIOTIC TROCHES** (Rowell) Neomycin sulfate 5 mg., tyrothricin 1 mg., polymyxin B sulfate 1000 u., benzocaine 5 mg./Troche Box 12s
 Use: Minor mouth and throat irritations
- VIO-BONE.** (Rowell) Vit D 500 u., C 50 mg., purified bone 7 1/2 gr., Cal 163 mg., Phos 77 mg., w/traces of Mag, fluoride/Cap Bot 100s 1000s
 Use: Supplement in pregnancy & lactation. W/Iron Cap (Rowell) Vit D 400 u., ascorbic acid 30 mg., thiamine HCl 2 mg., riboflavin 2 mg., niacinamide 5 mg., ferrous gluconate 2.5 gr., Vit B 12 1 mcg., folic acid 0.5 mg., purified pow bone 5 gr., Cal. 33.65%, Phos 15.77%, trace fluoride Mag/Cap Bot 100s, 1000s
 Use: Vitamin & mineral supplement
- VIO-C.** (Rowell) Ascorbic acid Tab 25 mg., 50 mg., 100 mg., 250 mg., Bot 100s, 1000s
 Use: Vitamin C therapy
- VIO-C W/B** (Rowell) Vit B 1 15 mg., B-2 15 mg., cal pantothenate 20 mg., niacinamide 50 mg., Vit B-6 5 mg., C 500 mg./Tab Bot 100s 250s
 Use: Vitamin supplement
- VIO-CAL** (Rowell) Dicalcium phosphate 10 gr., Cal gluconate 5 gr., Vit D 700 u./water or 2 Cap Cap, Bot 100s, 500s
 Water, Box 60s, 150s
 Use: Vitamin D w/calcium therapy
- W/Iron** (Rowell) Dicalcium phosphate 7 1/2 gr., ferric pyrophosphate 1 gr., Vit D 350 u./Cap Bot 100s 500s
- W/Iron & Iodine** (Rowell) Vio-Cal w/Pot Iodide D 12 mg./Cap Bot 100s, 500s.
- VIOMIC SULFATE, N.N.D. 1962** (Pfizer) Viomycin Sulf Vial 1 B 5 Gm
 Use: Against streptomycin resistant organisms
- VIO-CORT.** (Rowell) Hydrocortisone HCl 0.5%, neomycin D.5% Tube 1/6 oz., 1/2 oz.
 Use: Skin Inflammations
- VIOTIN.** (Pfizer) Vit A 5000 u., D 1000 u., thiamine HCl 1.5 mg., riboflavin 2 mg.

pyridoxine 0.1 mg., Col. pantothenate 1 mg., niacinamide 20 mg., ascorbic acid 37.5 mg./Cap. Bot. 100s, 500s, 1000s

Uses: Dietary supplement.

VIO-D. (Rowell) Vit. D 50,000 u./Cap. Bot. 100s, 1000s.

Uses: Rickets.

VIODENUM. (VioBin) Duodenum desic. Tnb. 10 gr., Bot. 100s, 1000s.

Use: (1-2 t.i.d.) Peptic ulcers.

VIO-DEX. (Rowell) Vit. A 5000 u., D 1200 u., B-1 3 mg., B-2 3 mg., B-6 1 mg., C 100 mg., niacinamide 20 mg., Col. pantothenate 2 mg., Vit. E 1 mg., d-phenetamine HCl 5 mg., phenobarbital $\frac{1}{4}$ gr./Cap. Bot. 100s, 1000s.

Uses: Obesity.

VIO-DEX TIMELETS. (Rowell) d-Amphetamine sul. 10 mg. or 15 mg. w/Vit. A 5000 u., D 1200 u., B-1 3 mg., B-2 3 mg., B-6 1 mg., C 100 mg., niacinamide 20 mg., col. pantothenate 2 mg., Vit. E 1 I.U., phenobarbital 32 mg./Sustained release Timelet. Bot. 100s, 250s, 1000s

Use: Treatment of obesity

VIO-DEXOSE. (Rowell) Dextroamphetamine phosphate 2.5 mg., mephobarbital 8 mg., dextrose 2.5 gm., Vit. A 1000 u., D 100 u., B-1 0.5 mg., B-2 0.5 mg., B-6 0.15 mg., C 15 mg., niacinamide 3 mg., col. d-pantothenate 0.3 mg./Tab. Boxes 100s.

Use: Treatment of obesity and control of appetite.

VIO-E. (Rowell) Mixed tocopherols equiv. to 30, 50 or 100 mg. alpha-tocopherol/cap. Bot. 100s, 1000s.

Use: Habitual abortion.

VIO-FERONATE. (Rowell) Liver dex. 4 gr., ferrous gluconate 5 gr., thiamin HCl 3 mg., riboflavin 2 mg., niacinamide 5 mg., Col. pantothenate 1 mg., pyridoxine 1 mg., Vit. C 30 mg., folic acid 1 mg., Vit. B-12 3 mcg./Tab. Bot. 100s, 1000s.

Uses: Anemia.

VIOFORM, N.N.R. 1952. (Ciba) Idochlorhydroxyquinoline.

Creams (3%) Tube 50 Gm., Jar 1 lb.

Tab. (250 mg) Bot. 20s, 100s, 500s

Tablets: 250 mg., Bot. 20s, 100s, 500s.

Lot: (3%) Bot. 80 ml.

Dist. (3%) Tube 50 Gm., Jar 1 lb.

Pow. (5 Gm) Bot. $\frac{1}{2}$ oz.

Invert: Vioform 25%, boric acid 10%, zinc stearate 20%, lactose 42.5%, lactic acid 2.5%. Bot. 1 & 8 oz.

Tablets: Vioform 250 mg., lactic acid 25 mg., boric acid 100 mg. Bot. 40s, 100s

Use: Trichomonas vaginitis.

VIOFORM-HYDROCORTISONE MILD CREAM. (Ciba) Vioform 3% & hydrocortisone 0.5% in a water-washable base. Tubes 1 oz. Also oint.

Uses: Acute & chronic dermatologic disorders.

VIOFORM-HYDROCORTISONE LOTION. (Ciba) Idochlorhydroxyquin 3% and hydrocortisone 1% with methylparaben 0.5% and propyl paraben 0.025%. Plastic squeeze bot. 15 ml.

VIO-GERIC. (Ewell Labs) Vit. A 10,000 u., D 1000 u., B-1 10 mg., B-2 5 mg., B-6 0.5 mg., B-12 5 mcg., C 100 mg., E 5 I.U., niacinamide 20 mg., col. pantothenate 5 mg., folic acid 0.25 mg., w/minerals Co 0.1 mg., Cu 0.5 mg., Mn 1.5 mg., Mg 3.0 mg., Pot. 4.4 mg., Zn 0.5 mg., I 0.1 mg., ferrous sulfate 130 mg., col. phosphate dibasic 325 mg., choline dihydrogen citrate 75 mg., inositol 25 mg., rutin 25 mg./Tab. Bot. 60s, 250s, 1000s. Also Vio-Geric-H estrogen-androgen added.

Uses: Dietary supplement.

VIO-GERIC-H. (Ewell) Conjugated estrogen (equine) 0.25 mg., methyltestosterone 2.5 mg., rutin 20 mg., inositol 25 mg., methionine 50 mg., plus 11 vitamins and 11 minerals./Tab. Bot. 60s.

Use: Multiple vitamin-mineral nutritional supplement.

VIO-JUNIOR. (Rowell) Vit. A 5000 u., D 500 u., thiamine HCl 2 mg., riboflavin 2 mg., pyridoxine 0.2 mg., Vit. C 30 mg., E 1 mg., folic acid 0.25 mg., Col. pantothenate 2 mg., niacinamide 10 mg., Vit. B-12 5 mcg./Tab. Bot. 100s

Use: Nutritional therapy.

VIO-JUNIOR DROPS. (Rowell) Vit. A 5000 u., D 1000 u., B-1 1 mg., B-2 1 mg., B-6 1 mg., B-12 5 mcg., C 60 mg., niacinamide 10 mg., pantothenol 3 mg./0.6 cc. Bot. 30 cc. w/plastic dropper.

Use: Pediatric nutritional supplement.

VIOKASE. (VioBin) Pancreas, whole. Tab. 5 gr., Bot. 100s, 500s, 1000s. Pow. 4 oz.

Use: Aid digestion.

VIO-LACTAL (Rowell) Cal. lactate 5 gr., Sod. pentobarbital $\frac{1}{4}$ gr., Sod. phenobarbital $\frac{1}{4}$ gr./Cap. Bot. 100s, 1000s.

Use: Sedative.

VIO-LAX. (Rowell) Diacetyldihydroxyphenylsulfate 5 mg., dantrolen 75 mg./Tab. Bot 100s.

Use: (1 daily) laxative.

VIO-LIV. (Rowell) Liver desic. 0.485 Gm., ferrous sulfate 3 gr., folic acid 1 mg., thiamine HCl 3 mg., riboflavin 1 mg., niacinamide 5 mg., Col. pantothenate 1 mg.,

pyridoxine 1 mg, Vit. B 12 1 mcg /Cap Bot 100s, 1000s

Use: Secondary anemias

***VIOMYCIN SULFATE**, NND 1962 (Parke, Davis) Viomycin sulfate Vial 1 & 5 Gm

Use Against streptomycin resistant organisms

VIOMYCIN SULFATE, NND 1962 Sulfate salt of antibiotic produced by selected strains of actinomycete to which *Streptomyces puniceus* & *Streptomyces floridus* have been applied Steri Vial 5 Gm (Parke, Davis)

See: Vinactane Sulfate (Ciba)

Viacin Sulfate (Pfizer)

VIO-NATAL (Rawell) Ferrous gluconate 15 gr, calcium lactate 15 gr., vit A 6000 u, D 1200 u, C 100 mg, K 1 mg, B 13 mg, B 2 3 mg, B 6 10 mg, B 12 6 mcg, folic acid 0.75 mg, niacinamide 15 mg /6 tab Bot 100s

Use: Dietary supplement

VIOHATR-L (Squibb) Vit A 40,600 u, D 3 5000 u, B 1 40 mg, B 2 46 mg, B 6 5 mg, B 12 17.5 mcg, E 13.8 u, niacinamide 100 mg, calcium pantothenate 25 mg, iodine D779 mg, iron 12.4 mg, cobalt 0.12 mg copper 5.92 mg, magnesium 11.8 mg, manganese 1.87 mg, calcium 1.814 mg, phosphorus 1.417 mg, salt 567 mg /oz pellets
Use: Vitamin mineral supplement for large animals

VIO-PHEN, (Rawell) Hamalopine methylbromide 1/20 gr, secobarbital Sod 1/8 gr, mephenesin 5 gr /Cap, Bot 100s, 500s, 1000s

Use: (1 2 o.c., & h.s.) Antispasmodic w/ muscle relaxant

VIOPLEX HIGH POTENCY, (Preme) Vit B 1 25 mg, B 2 2 mg, B-6 2 mg, niacinamide 150 mg /cc Vial 10 cc

Use: Vitamin B deficiency

VIO-PRENATE, (Rawell) Vit A 1500 u, D 200 u, C 10 mg, B-1 1 mg, B 2 1 mg, Cal lactate 4 gr, ferrous gluconate 5 gr, Vit B 12 1 mcg, folic acid 0.3 mg, Pot iodide 0.15 mg /Cap Bot 100s

Use: Dietary supplement

VIO-PRENATE WITH FLUORIDE, (Rawell) Addition of fluoride to previously released Vio Prenate formula

Use: Prenatal vitamin fluoride combination

VIO-SAL, (Rawell) Potassium salicylate 150 mg, strontium salicylate 150 mg, calcium salicylate 150 mg, Vit K 0.5 mg /Tab Bot 100s

Use: Analgesic antipyretic

VIO-SERPINE, NND 1962 (Rawell) Rauwolfia serpentina alkaloids Tab. (3) 1 mg,

* Generic name recognized by N.F., U.S.P. or A.M.A.

D 25 mg, 0.5 mg, 1.0 mg, or 4.0 mg) Bot 1000s

Use Sedative & tranquilizing agent.

VIOSTEROL, See: Calciferol, USP

VIOSTEROL W/HALIBUY LIVER OIL, NNR 1953, ADR Vit A 50,000 u, Vit. D 10,000 u /Gm

Abbott—Bot 5 cc, 20 cc, 50 cc Cap Vit A 5000 u, D 1000 u

Ives-Cameron—Cap Vit. A 5000 u, D 1700 u

Parke, Davis—Bot 5 cc, 50 cc Cap Vit A 5000 u, D 1000 u

VIOSTEROL IN OIL, USP, NNR 1952 ADR (Oleovitamin D, Synthetic USP; Vit D-2, Calciferol)

Abbott—Bot 20 cc, 50 cc
Horton & Converse—10,000 u /Gm, Bot 2 oz, 4 oz, 1 pt

Ives-Cameron—Bot 10 cc, 60 cc
Parke, Davis—Bot, 5 cc, 50 cc

Squibb—Vial w/dropper 20 cc Bot, 50 cc

Use: Vitamin D therapy

VIO-THENE, NND 1962 (Rawell) Oxyphencyclimine HCl 10 mg /Tab Bot 100s, 250s, 1000s

Use: Anticholinergic, treatment of peptic ulcers

VIO-THENE W/PHENOBARBITAL, (Rawell) Oxyphencyclimine HCl 10 mg, Phenobarbital 30 mg /Tab Bot 100s, 1000s

Use: Anticholinergic with sedative

VIO-THRICIN, (Rawell) Tyralthricin 1 mg, benzocaine 8 mg, hexylresorcinol 2 mg /Loz Vial 14s

Use: Infections of mouth & throat

VI-PENTA DROPS #1, (Roche) Synkayvite 1 mg, Vit E 5 mg, C 25 mg /0.6 cc Vial 5 cc

Use: Progressive vitamin therapy for specific age groups

VI-PENTA DROPS #2, (Roche) Vit A 5000 u, D 1000 u, C 50 mg, E 2 mg /0.6 cc Vial 15, 50 cc

Use: Progressive vitamin therapy for specific age groups

VI-PENTA DROPS #3, (Roche) Vit A 5000 u, B 1 1 mg, B 2 1 mg, B 6 1 mg, C 50 mg, D 1000 u, d panthenol 10 mg, niacinamide 10 mg /0.6 cc Vial 15, 30, 50 cc

Use: Progressive vitamin therapy for specific age groups

VI-PENTA PERLES PORTE, (Roche) Vit A 5000 u, D 1000 u, thiamine HCl 3 mg, riboflavin 3 mg, pyridoxine HCl 1 mg, niacinamide 20 mg, panthenol 3 mg, ascorbic acid 75 mg, Vit E acetate 1 mg

- 12 2 mcg./Cap. Vial 25s, 100s. Bot. 250s, 000s.
Use: (1-3 daily) Multivitamin deficiencies.
- VI-PENTA NOMOGENATE.** (Roche) Nine vitamins w/Vit. B-6 & B-12. Bot. 4 oz., 3 pt. Uses: Vitamin deficiencies.
- VI-PENTA ZESTABS.** (Roche) Vit. A 5000 U.S.P. u., O 1000 U.S.P. u., C 60 mg., B-1 1.2 mg., B-2 1.5 mg., B-6 1.2 mg., B-12 3 mcg., d-biotin 40 mcg., niacinamide 10 mg., E 2 I.U., d-panthanol 10 mg./chewable tab. 30s.
Use: Nutritional supplement.
- VIPEPTOLAC.** (Wyeth) Protein 50 Gm., carbohydrate 37 Gm., iron 10 mg., folic acid 2 mg./100 Gm. Plain, Can 1 lb. Chocolate flavored, Can 1 lb.
Use: Dietary supplement.
- VIPIBONE.** (Vitomix Corp.) Oipyrone 300 mg./cc. Vials 30 cc.
- VIPER VENOM.** See: Stypren, Vial. (Burrroughs Wellcome)
- VI-PROTALIN.** (National Drug) Protein 18.4 Gm., carbohydrate 7.8 Gm., choline dihydrogen citrate 300 mg., Vit. A 1500 u., O 150 u., B-1 3 mg., C 30 mg., B-2 3 mg., B-6 0.3 mg., Cal. pantothenate 1.5 mg., Vit. B-12 4.5 mcg., folic acid 0.6 mg., aluminum 30 mg., Phos. 140 mg., Col. 180 mg., iron 4.5 mg./30 Gm. Pow., pkg. 8 oz., 1 lb.
Use: Dietary supplement where increased protein, vitamins & minerals are required.
- VIPRYNIUM EMBONATE.** 6-Oimethylamino-2-[2-(2,3-dimethyl-1-phenyl-3-pyrrolidyl)vinyl]-1-methylquolinium embonate. [Embonic acid is adopted as the trivial name for 4,4'-methylenebis-(3-hydroxy-naphthalene-2-carboxylic acid)] (Vanquin)
- VIBAC, N.N.D. 1962.** (Rusan Labs) Undecoylim chloride-iodine. Iodine complexed with a cationic detergent. Rex Sol. Bot. Squeeze 4 oz., 1 pt., 1 gal. Surgical sol. Bot. 4 oz., 1 gal.
Use: Antiseptic.
- VI-BANDIN.** (Rand) Vit. A 5000 u., D 1000 u., B-1 3 mg., B-2 2 mg., B-6 0.25 mg., B-12 1 mcg., C 37.5 mg., Cal. pantothenate 5 mg., niacinamide 20 mg., mixed tocopherols 5 mg./cap. Bot. 100s, 1000s.
Use: Dietary supplement.
- VIB-I-PHYL.** (Poul Money) Veratrum viride 1.5 gr., rufin 20 mg., phenobarbital 15 mg., neohyline 1.5 gr./Tab. Bot. 100s, 1000s
Use: (1-2 t.i.d.) Hypertension.
- VIB-I-TIN.** (Poul Money) Mannitol hexanitrate 30 mg., Veratrum viride 100 mg., rufin 20 mg., phenobarbital 1/4 gr./Tab. Bot. 100s, 1000s.
- Use:** (1-2 t.i.d.) Vasodilator in hypertension.
- VISION-1.** (Lincoln) Ascorbic acid 2000 mg., thiamine HCl 50 mg., riboflavin 5 mg., pyridoxine HCl 10 mg., nicotinamide 100 mg., methylparaben 0.89 mg., propylparaben 0.22 mg., sodium bisulfite 10 mg., disodium calcium sequestrene 1 mg./cc. 10 cc. Vial 6 (10 cc) of Lyophilized Powder and 6 (10 cc) of Diluent
Use: Treat viral infections.
- VIBON-2.** (Lincoln) Ascorbic acid 2000 mg., sodium bisulfite 0.1%, disodium calcium sequestrene 0.01%/ 10 cc. Amp. (score-break) 10 cc. Box 25s.
Use: Treat viral infections.
- VIBOSTIBONE.** (Endo) Testosterone. Aq. Susp., Amp. (5 mg./cc., 10 mg./cc., 25 mg./cc., 100 mg./cc.) 1 cc., Box 6s, 25s, 100s.
Use: Male sex hormone therapy.
- VIROSTIBONE PROPIONATE.** (Endo) Testosterone propionate. In oil, Vial (25 mg./cc.) 10 cc.
- VIRUGON.** (Winthrop) Under study. Anhydro bis-(beta-hydroxyethyl) biguanide derivative.
Use: Treatment of influenza, mumps, measles, chicken pox, and shingles.
- VI-RUMAL.** (Normand) Veratrum viride 3 Crow u., Sod. nitrite 1 gr., phenobarbital 1/4 gr./Tab. Bot. 500s, 1000s.
Use: (1-2 t.i.d.) Vasodilator in hypertension.
- VISAMMIN.** (Physicians Drug) Kheilin, Tob. 20 mg., 50 mg., Bot. 100s, 500s.
Use: Angina pectoris.
- VISCIODOL.** (Fougera) Sulfanilamide 320 mg., Ipidol 40%/cc. Susp., Vial 15 cc.
Use: Bronchography, diagnostic agent.
- VISCO MODIFIED.** (Pitman-Moore) Sod. nitrite 1 gr., mistletoe [Viscum album] 1 gr./Tab. Bot. 100s, 1000s
Use: Hypertension.
- VISCOPE.** (Lynn) Scopolamine methyl nitrate 5 mg./Disintegrating cap. Bot. 60s, 250s.
Use: Peptic ulcers
- VISCUM ALBUM, EXTRACT.**
See: Visco Modified (Pitman-Moore)
Viscysate (Bischoff)
Visnic (Drug Products)
Use: Vasodilator.
- VISINE, N.N.D. 1962.** (Pfizer) Tetrahydrozoline HCl 0.5%. Bot. 1/2 oz.
Use: Eye irritations.
- VISNICO.** (Drug Products) Viscum album pow. ext. 1/4 gr., Sod. nitrite 1 gr., Pot. nitrate 2 gr., crataegus oxyacantha ext. 1/2 gr., phenobarbital 1/4 gr./cap. Bot. 100s, 1000s.
Use: Hypertension.
- VI-SORBIN.** (Smith, Kline & French) Crystal.

¹ Generic name recognized by N.F., U.S.P. or A.M.A.

line Vit B 12 25 mcg, B 6 6 mg, iron (as soluble ferric pyrophosphate) 300 mg, folic acid 1.5 mg, D sorbitol/15 cc Bot 8 fl oz
Use: Tonic for convalescent

VISTABOLIC (Organon) Hydrocortisone 1 mg, methylandrostenediol 10 mg, Vit B 12 w/intrinsic factor conc $\frac{1}{2}$ u/Tab Bot 30s 100s Also avail as Inj Vial 10 cc.
Use: Geriatrics

VISTARIL, NND 1962 (Pfizer) Hydroxyzine pamoate 25 mg, 50 mg 100 mg/Cap Bot 100s Hydroxyzine HCl Vial 25 mg/cc 10 cc Steroject 25 mg/2 cc w/needle Pkg 10s & 100s Susp (25 mg/5 cc) Bot 1 pt
Use: Psychotherapeutic agent

VI-SYNERAL ADULT. (U S Vitamin) Vit A 10,000 u, B 1 3 mg, B-2 2.5 mg, B-6 0.75 mg, niacinamide 15 mg, Cal pantothenate 3.5 mg, Vit C 50 mg, D 1000 u, E 3 mg., yeast 90 mg, Cal 50 mg, Phos., iron, copper, iodine, Mang, Mag, zinc, cobalt/Cap Box 30s 50s, 100s, 200s
Use: Dietary supplement

VI-SYNERAL CHEWABLE TABLET (U S Vitamin) Vit A 5,000 u, D 500 u C 75 mg B-1 1.5 mg B-2 0.5 mg B-6 2 mg B 12 3 mcg niacinamide 10 mg d calcium pantothenate 2 mg., Vit E 2 mg/Tab Bot 100s
Use: Vitamin supplement

VI-SYNERAL DRDPS. (U S Vitamin) Vit A 5000 u., D 1000 u, B-1 1 mg, B 2 0.4 mg, B-6 0.1 mg, niacinamide 5 mg, pantothenic acid 2 mg, Vit C 50 mg/0.6 cc. Bot w/dropper 15 cc., 30 cc.
Use: Dietary supplement

VI-SYNERAL DRDPS FORTIFIED. (U S Vitamin) Vit A 5000 u, D 1000 u, C 50 mg, B-12 2 mcg, choline 12.5 mg, betaine 12.5 mg., Inositol 15 mg, B-1 1 mg, B-2 0.6 mg., niacinamide 10 mg., B-6 1 mg, pantothenol 3 mg, E 1 IU/0.6 cc. Bot w/dropper 15 cc., 30 cc., 45 cc.
Use: Dietary supplement

VI-SYNERAL EXPECTANT & NURSING (U S Vitamin) Vit A 12,000 u., B-1 4 mg, B-2 3 mg, B-6 1 mg, niacinamide 18 mg, Cal. pantothenate 4 mg, Vit C 60 mg, D 1750 u, E 4 mg., yeast 70 mg, Cal 1 Gm., Phos., iron, copper, iodine, Mang, Mag, zinc/Cap Box 50s, 150s
Use: Dietary supplement

VI-SYNERAL DEVID. (U S Vitamin) Vit A 5000 u., D 500 u., B-1 2.5 mg, B-2 2.5 mg., B-6 0.5 mg., B-12 0.5 mcg, folic acid 0.5 mg., niacinamide 10 mg., choline 25 mg., pantothenol 5 mg., Vit C 75 mg, d-alpha-tocopherol 2.5 mg., yeast 200 mg, Cal 110 mg., iron 25 mg., Phos 100 mg, copper

D75 mg, iodine 0.05 mg, Mang 0.5 mg Mag 0.75 mg, zinc 0.5 mg, cobalt 0.1 mg/1 Vit (dark color) & 1 mineral (light color) cap Bot 50s, 100s, 200s

Use Dietary supplement for expectant & nursing mothers

VI-SYNERAL INJECTABLE (U S Vitamin) Vit A 10,000 u, D 1000 u, thiamine HCl 10 mg, riboflavin 1 mg, pyridoxine HCl 3 mg, niacinamide 20 mg, ascorbic acid 50 mg, alpha-tocopherol 2 mg/2 cc. Amp 2 cc, Box 6s, 25s, 100s
Use I.M. only; replacement of depleted vitamins

VI-SYNERAL MULTIVITAMIN INJECTION. (U S Vitamin) Ascorbic acid 500 mg, Vit A 10,000 u, D 1000 u, dl alpha-tocopherol acetate 5 mg, thiamine HCl 50 mg, riboflavin 10 mg, niacinamide 100 mg, pyridoxine HCl 15 mg pantothenol 25 mg/10 cc Vial Vial 10 cc Box 1s, 5s, 25s, 100s
Use: IV after dilution; vitamin therapy

VI-SYNERAL ONE-CAP5 (U S Vitamin) Vit A 6000 u, D 600 u, ascorbic acid 75 mg, Thiamine mononitrate 3 mg, riboflavin 3 mg, pyridoxine HCl 1 mg, niacinamide 20 mg, Vit B 12 2 mcg, d calcium pantothenate 5 mg, d-alpha-tocopherol acetate 1 ml e, dicalcium phosphate 170 mg, ferrous sulfate anhydrous 50 mg, copper 1 mg, iodine 0.1 mg, manganese 1 mg, magnesium 1 mg, zinc 1 mg/Cap Bot 28s 100s
Use: Vitamin and mineral supplement

VI-SYNERAL SPECIAL (U S Vitamin) Vit A 12,000 u, B 1 5 mg, B 2 3.5 mg., B 2 mg niacinamide 20 mg., Cal pantothenate 5 mg, Vit C 75 mg, D 1200 u E 4 mg, yeast 50 mg, Cal, Phos., iron, copper, iodine, Mang, Mag, zinc, cobalt/Cap Box 50s, 100s, 200s
Use: Dietary supplement

VI-SYNERAL THERAPEUTIC. (U S Vitamin) Dark colored cap, Vit A 25,000 u, D 1000 u, ascorbic acid 150 mg, Vit B 12 1 mcg., folic acid 1.76 mg, thiamine HCl 15 mg., niacinamide 150 mg., riboflavin 10 mg., pyridoxine HCl 5 mg, d-alpha-tocopherol 10 mg, desicc liver & liver fraction No 2 derived from 7.5 Gm liver light colored cap., Choline 20 mg., Inositol 10 mg., d-Cal. pantothenate 15 mg., Cal 160 mg., Phos. 132 mg, iron 15 mg., copper 1.5 mg., Mang 1 mg, Mag 1 mg., zinc 1 mg., iodine 0.1 mg, cobalt 0.1 mg See 30s, 50s, 100s
Use: Dietary supplement

VITA-12 FORTIFIED (Crystal) Ferrus gluconate 170 mg, Vit B 12 1.25 mcg, thiamine HCl 1.25 mg, riboflavin 0.67 mg,

niacinamide 6.67 mg., liver-stomach powder 333 mg./Tab. Bot. 100s.
 Use: Treatment of anemias.

VITABEV. (Key Corp.) Vit. B-12 29.3 mcg., B-1 20 mg., B-2 3 mg., B-6 1 mg., pantothenic acid 3 mg., niacinamide 100 mg., choline chloride 20 mg., inositol 10 mg., procaine HCl 1%, liver inj. crude 50% w/Vit. B-12 equiv. to 2 mg./cc. Vial 10 cc.

VITACAIN. (Parker) Lotion, Bot. 4 oz.

VITACEE. (Morse Lab.) Ascorbic acid. Tab. 25 mg., 50 mg., 100 mg., Bot. 100s.
 Use: Vitamin C therapy.

VITA-CNEW. (Broemmel) Vit. A 5000 u., D 1000 u., C 600 mg., B-1 2 D mg., B-2 2.0 mg., B-6 2 D mg., B-12 5.0 mcg., calcium pantothenate 5.0 mg., niacinamide 150 mg./Tab. Bot. 30s, 100s.
 Use: 1-3 daily, dietary supplement.

VITACOL. (Mulfard) Vit. B-1 1.5 mg., B-2 0.60 mg., niacinamide 5 mg., colloidal iron 3 gr., liver conc. 200 mg., Mang. 1.8 gr./Cop. Bot. 40s, 100s, 1000s.
 Use: Hematinic.

VITACOL IMPROVED. (Mulfard) Vit. B-1 6 mg., B-2 2.75 mg., B-6 0.15 mg., niacinamide 30 mg., iron, Mang., peptone/oz. Bot. 8 oz.
 Use: Anemias.

VITADEX-B. (Cutter) Dextrose 5%, 10%, thiamine HCl 5 mg., riboflavin 7.5 mg., niacinamide 50 mg., pyridoxine HCl 3 mg./L. Sulfaflox 1000 cc.
 Use: I.V.; replace illost & circulating fluid.

VITA-DRINK—INSZANT. (Sivart) Vit. A 5000 u., D 1000 u., C 75 mg., B₁ 2 mg., B₂ 2 mg., B₆ 1.2 mg., B₁₂ 3 mcg (as an absorption-enhancing complex), Niacinamide 10 mg., d-calcium pantothenate 3 mg./4.3 Gm. Powder in 4.5 oz. pack. 3 flavors—cherry, grape, orange.
 Use: Vitamin supplement.

VITAE. (Mourry) 14 vitamins & lipotropic factors, 5 essential minerals, 5 trace minerals./Tab. Jar 150s.

VIZAFER. (National Drug) Iron & ammonium citrate (green) equiv. to 40 mg. iron, Col. gluconate 4 gr., Mang. gluconate 3 gr., thiamine HCl 2000 u./fl. oz. Bot. 1 pt., 1 gal.
 Use: Hematinic.

VIZAGUENT. (Tailby-Nasan) Cod liver oil 40%, zinc borosalicilate. Oint., Tube 1 oz., 4 oz. Jar 1 lb.

W/Bensocaine. Tube 1 oz., 4 oz. Jar 1 lb.

W/Sulfathiazole 5%. Tube 4 oz. Jar 1 lb.
 Use: Burns.

VITAKAPS. (Abbott) Vit. A 5000 u., D 500 u., thiamine HCl 3 mg., riboflavin 2.5 mg., niacinamide 20 mg., pyridoxine HCl 0.5 mg., ascorbic acid 50 mg./Cap. Bot. 50s,

100s, 250s. Also avail. w/Vit. B-12 1 mcg.
 Use: Dietary supplement.

VITAKAPS-M. (Abbott) Same vitamins as Vitakaps plus 9 minerals. Bot. 50s, 100s, 250s

VITA-L. (Klug) Methylcellulose 7.5 gr., B-1 1.5 mg., B-2 2 mg., C 20 mg., phenylpropanolamine HCl 25 mg./2 Tabs. Bot. 100s.
 Use: Constipation, obesity.

VITALEAF AMPULS. (Gold Leaf) Vit. B-1 25 mg., B-2 0.5 mg., B-6 1 mg., niacinamide 50 mg./cc. Amp. 1 cc., 10 cc., 30 cc.
 Use: Vitamin B deficiencies.

VITALEAF DRDPS. (Gold Leaf) Vit. A 6000 u., D 1200 u., B-1 1.8 mg., B-2 0.6 mg., C 60 mg., B-6 0.3 mg., niacinamide 3 mg., Cal. pantothenate 1.2 mg./0.6 cc. Bot. w/dropper 15 cc., 30 cc.
 Use: Dietary supplement.

VITALEAF SYRUP. (Gold Leaf) Vit. B-1 3 mg., B-2 1 mg., C 30 mg., B-6 0.83 mg., niacinamide 15 mg., pantothenic acid 2.3 mg./3 Tsp. Bot. 4 oz., 8 oz., 1 pt.
 Use: Dietary supplement.

VITALEAF TABLETS. (Gold Leaf) Thiamine HCl 10 mg., riboflavin 5 mg., pyridoxine HCl 1 mg., pantothenic acid 5 mg., niacinamide 50 mg., ascorbic acid 75 mg./tab. Bot. 25s, 100s, 500s.
 Use: Dietary supplement.

VITALERT. (Drug Products) Vit. A 5000 u., D 1000 u., B-1 3 mg., B-2 3 mg., C 25 mg., niacinamide 20 mg., Cal. pantothenate 3 mg., Vit. B-6 0.1 mg./Cop. Box 30s, 100s, 500s.
 Use: Dietary supplement.

VITAMS. (Primo) An eleven vitamin formulation including B-12, folic acid and vitamin E/lozenges. Bot. 30s.
 Use: Vitamin supplement.

VITAMALS. (Vitamin Corp.) Vit. A 10,000 u., D 1000 u., B-1 6 mg., B-2 4 mg., B-6 0.5 mg., C 100 mg., Cal. pantothenate 5 mg., niacinamide 10 mg., Vit. E (wheat germ oil) 5 mg., folic acid 0.1 mg., Vit. B-12 2 mcg./Cop. Bot. 50s, 100s, 250s, 1000s.
 Use: Dietary supplement.

VITA-METRAZDL. (Knott) Metrazol 100 mg., Vit. B-1 1 mg., B-2 1 mg., niacinamide 10 mg., B-6 1 mg., pantholol 2 mg./5 cc. Bot. 1 pt. Tab. (w/Vit. C 25 mg.) Bot. 100s.
 Use: Normalize cerebral functions and provide supplemental vitamin medication.

VITAMEZDL. (Strosenburgh) Hydrolyzed meat proteins, amine acids, maltose, minerals, Vit. B-1, B-2. Bot. 1 pt., 1 gal.

W/Liver B Iron. Iron, ammonium chlorates, liver conc. Bot. 6 B. oz., 1 pt., 1 gal.
 Use: Tonic.

*VITAMIN A WATER MISCIBLE, OR SOLUBLE USP NND 1962

- See Acon Cap & Sol NNR (Endo)
 Aquacel Oleum A (Biopharmaceut cals)
 Aquosol Vit A Drops NNR (U S Vitamin)
 A Vitol (Hance Bros & White)
 Sol A Caps (Bryant)

VITAMIN A NND 1962 ADR

- See D spots Tob's (Person and Covey)

Ampuls

- A Vitam n (100,000 u/cc) 1 cc (Squibb)
 Anatola (50,000 u/cc) 1 cc (Parke Davis)
 Aquosol A (25,000 u 50,000 u) 2 cc (U S Vitamin)
 A Vita (25,000 u) 1 cc (Brewer)
 H A Vita (50,000 u) 1 cc (200,000 u) 2 cc (Brewer)
 Testaval (Testagar) (100,000 u) 1 cc.

Capsules

- A Vitamin 25,000 u 50,000 u (Squibb)
 Acon 25,000 u 50,000 u (Endo)
 Aloxin 10,000 u 25,000 u 50,000 u (Winthrop Labs)
 Alphol n 10,000 u 25,000 u 50,000 u (Lilly)
 Alphasteral Caps 25,000 u 50,000 u (Meyer)
 Anatola 25,000 u 50,000 u (Parke Davis)
 Aaral 25,000 u 50,000 u (Massengill)
 Apexol 50,000 u (Roerig)
 Aquosol A Caps 25,000 u 50,000 u (U S Vitamin)

W/V I C & V I D

See

- ADC Drops (Parke Davis) /0.6 cc.
 Adcats (Walker) /Vasphere
 Aquosol A C D Drops (U S Vitamin) /0.6 cc.
 Aquosperse Vitamin ACD Drops (White) /0.6 cc.
 Pre Dayl n Drops (Abbott) /0.6 cc
 Tr Vit Sol (Mead Johnson) /0.6 cc
 Vadcon (Walke) /1 cc.
 Y dac (Endo) /0.6 cc

W/V I D

See

Capsules or Tablets

- Adcaps (Blue Line)
 A & D Caps (Rowell)
 Adex (Squibb) /Tab
 Adleaf Cap (Gold Leaf)
 A D Percomorph (Abbott) /Cap
 Alpha Delta n # 1 (Lilly) /Gelseat
 Alpha Delta n # 2 (Lilly) /Gelseat
 Apolarthron (Roerig) /Cap
 A Vita D # 2 (Pitman Moore) /Cap
 Rurbot Liver Oil (Rowell) /Cap
 Car tol w/Vitamin D (Wyeth) /Cap

*Generic name recognized by N.P. U.S.P. or A.M.A.

Aquasynth 25,000 u 50,000 u (U S Vitamin)

Atav 50,000 u (Cale Chem Co)

A Vita 25,000 u (Brewer)

Av tum 50,000 u (Ives Cameron)

H A Vita 50,000 u (Brewer)

Nio-A Let 25,000 u 50,000 u (Non)

Stabil A 25,000 u 50,000 u 100,000 u (Dumas Wilson)

Super A Vitamin 25,000 u 50,000 u 100,000 u (Upjohn)

Testaval 100,000 u (Testagar)

Testaval S 25,000 u 50,000 u (Testagar)

Vafat 25,000 u (Breon).

VI Alpha 50,000 u (Lederle)

VI Dom A Pill 25,000 u 50,000 u (Dome)

Ointment

V Dom A Creme (Dome)

Solutions

Acon 25,000 u/cc Bat 30 cc (Endo)

Afax n 65,000 u/Gm Bat 30 cc (Winthrop Labs)

Super A Vitamin 50,000 u/cc Bat 15 cc 50 cc (Upjohn)

Aquosol A 50,000 u/cc Bat 15 cc 30 cc (U S Vitamin)

W/Estrogen See Agera Oint (Biopharmaceut cals)

W/Pantathenol See Aquosol A W/Pantathenol Oint (U S Vitamin)

W/Salicylamide See Salilith Tab (Mile)

W/Sulfur & salicylic acid See Sal A Sol Oint (Dome)

| | A | C | D |
|--|----------|--------|---------|
| | 5,000 u | 50 mg | 1,000 u |
| | 5,000 u | 50 mg | 1,000 u |
| | 5,000 u | 75 mg | 1,000 u |
| | 5,000 u | 50 mg | 1,000 u |
| | 5,000 u | 50 mg | 1,000 u |
| | 5,000 u | 30 mg | 1,000 u |
| | 10,000 u | 100 mg | 2,000 u |
| | 5,000 u | 60 mg | 1,200 u |

| | A | D |
|--|----------|----------|
| | 5,000 u | 500 u |
| | 5,000 u | 500 u |
| | 5,000 u | 800 u |
| | 5,000 u | 500 u |
| | 5,000 u | 1,870 u |
| | 5,000 u | 500 u |
| | 15,000 u | 1,500 u |
| | 30,000 u | 25,000 u |
| | 5,000 u | 500 u |
| | 2,215 u | 315 u |
| | 5,000 u | 1,000 u |

W/Vit. D.

| | A | D |
|---|-----------|----------|
| Cod Liver Concentrate (White) /Cap. | 5,000 u. | 500 u. |
| Cod Liver Concentrate (White) /Tab. | 3,120 u. | 312 u. |
| Deltalin Alpha No. 1 (Lilly) /Cap. | 5,000 u. | 500 u. |
| Deltalin Alpha No. 2 (Lilly) /Cap. | 15,000 u. | 1,500 u. |
| Drisdol w/Vitamin A (Winthrop Labs) /Cap. | 5,000 u. | 1,000 u. |
| Holiver Oil (Various Mfr.) /Cap. | 5,000 u. | 83 u. |
| Holiver Oil w/Viosterol (Various Mfr.) /Cap. | 5,000 u. | 1,000 u. |
| Hepicoleum (Lilly) /Globule | 5,000 u. | 1,000 u. |
| Hi-Vi-Tol (Patch) /Cap. | 5,000 u. | 500 u. |
| Natala (Parke, Davis) /Cap. | 5,000 u. | 1,000 u. |
| Navitol w/Viosterol (Squibb) /Cap. | 5,000 u. | 1,000 u. |
| Oleum Percamorphum w/Viosterol (Mead Johnson) /Cap. | 5,000 u. | 700 u. |
| Super D (Upjohn) /Perte | 14,450 u. | 1,445 u. |
| Vod (Cole Chem. Co.) /Cap. | 5,000 u. | 500 u. |
| Vi-Dom D-2 (Dome) /Pill | 50,000 u. | 5,000 u. |

Liquid

| | | |
|---|-----------|-----------|
| A-D Drops (Walker) /Gm. | 62,500 u. | 10,000 u. |
| A-D Percamorph Oil (Abbott) 10.1 cc. | 5,450 u. | 773 u. |
| A & D Vit. Conc. (Horton & Converse) | 500 u. | 1,700 u. |
| Adsorl (Abbott) /0.1 cc. | 5,000 u. | 1,000 u. |
| Ad-Vitum (Ives-Cameron) /0.1 cc. | 3,000 u. | 500 u. |
| Aquasol A & D (U. S. Vitamin) /0.1 cc. | 3,636 u. | 727 u. |
| Burbot Liver Oil (Rowell) /Gm. | 4,480 u. | 640 u. |
| Cod Liver Oil (Various Mfr.) /Gm. | 850 u. | 85 u. |
| Cod Liver Oil w/Molt Extract (Burroughs Wellcome) /oz | 6,450 u. | 645 u. |
| Cod Liver Oil w/Viosterol (Various Mfr.) /2 cc. | 3,600 u. | 800 u. |
| Drisdol w/Vitamin A (Winthrop Labs) /0.1 cc. | 5,000 u. | 1,000 u. |
| Holibut Oil (Various Mfr.) /Gm. | 60,000 u. | 600 u. |
| Holibut Oil w/Viosterol (Various Mfr.) /Gm. | 60,000 u. | 10,000 u. |
| Hepicoleum (Lilly) /Gm. | 55,000 u. | 11,000 u. |
| Natala (Parke, Davis) /0.1 cc. | 5,000 u. | 1,000 u. |
| Oladol (Abbott) /0.1 cc. | 5,000 u. | 500 u. |
| Oladol (Abbott) /0.1 cc. | 7,500 u. | 1,500 u. |
| Pravitol (Wyeth) /Gm. | 5,000 u. | 500 u. |
| Super D Drops (Upjohn) /0.1 cc. | 5,000 u. | 375 u. |
| Vi-Delta Emulsion (Lederle) /5 cc. | 2,500 u. | 3,000 u. |
| Hi-Vi-Tol (Patch) /Gm. | 30,000 u. | 3,000 u. |

Ointment

| |
|---|
| Allantoin Compound (Upjohn). Cod liver oil 1.82%, allantoin 2%, secondary amylicresol 1/2%. |
| Adestrin (Brean). Fish liver oil. |
| Burba-Caine (Rowell). Burbot liver oil 30%, benzocaine 5%. |
| Burba Liver Oil Ointment (Rowell). Burbot liver oil 50%. |
| Cod Liver Oil w/Metaphen (Abbott). Cod liver oil 70%, metaphen 1:2500. |
| Clearcam (Upjohn). Cod liver oil. |
| Desitin (Desitin) Cod liver oil, zinc oxide & talc. |
| Gadament (Patch). Cod liver oil 70%, phenol less than 1/2%. |
| Morogues (Masseguill). |
| Vitamin A & D (White). Fish liver oils. |

W/Aluminum hydroxide, benzocaine, vit. D, calcium, iron. See, Calcifac, Tab. (Brwle-Neiser)

* Generic name recognized by N.F., U.S.P. or A.M.A.

W/Vit. D & sulfathiazole. See: Sulfavitin, Oint. (Chicago Pharm.)

W/Vit. D, carotene & dibasic Cal. phosphata. See: Supplamin, Cap. (Ayerst).

W/Vit. E.

See: Corital, Cap. (Wyeth).

Vi-Alpha, Cap. (Lederle).

W/Vitamin, mult. See: Multivitamin Tablets.

VITAMIN Bc. See: Folic acid.

VITAMIN B-1, N.N.D. 1962, A.D.R. See: Thiamine Hydrochloride, U.S.P.

VITAMIN B-2, N.N.D. 1962, A.D.R. See: Riboflavin, U.S.P.

VITAMIN B-6, N.N.D. 1962, A.D.R. See: Pyridoxine Hydrochloride, U.S.P.

VITAMIN B-12, U.S.P., N.N.D. 1962, A.D.R. Cyanocobalamin, U.S.P. Cobalamine.

See:

Capsules & Tablets

Almerat, 25 mcg. Cap. (Mayer)

Berubigen, 25 mcg./Cap. (Upjohn).

- Bevidox Concentrate 25 mcg /Cap, 10 mcg /Oulcet (Abbott)
 Beva'ine 12 (Smith Dorsey)
 Bexii (Chicago)
 Bexitab Tab (Chicago Pharm)
 Biocres, 30 mcg /tab (U M A)
 B Twelv Ora 25 mcg /Cap (Sherman)
 Cotel 25 mcg /tab (Jeffery)
 Docibin 10, 25, 50 mcg /Tab (Walker)
 Dodecavite 25 mcg /Tab (U S Vitamin)
 Normocytin 30 mcg /Tab (Lederle)
 Payamin 5 mcg /Cap (Testagar)
 Rametin 5 10, 30 mcg /Tab (Bio-Ramo)
 Redisal 25, 50 mcg /tab (Merck Sharp & Dahme)
 Rubramin 25 mcg /Tab (Squibb)
- Salutlan**
- Docibin 1 mcg /Drap (Walker)
 Dodecavite 10, 30 mcg /cc (U S Vitamin)
 Hepcovite 16 mcg /cc (Endo)
 Payamin 1 mcg /Drop (Testagar)
 Redisal 5 mcg /cc, Elixir (Merck Sharp & Dahme)
 Rubramin 30 mcg /cc (Squibb)
- Vials & Ampuls**
- Almeret 30 100 1000 mcg /cc (Meyer)
 Redoxe 30, 60 200, 1000 mcg /cc (Lincoln)
 Re Dodec 15 mcg /cc (Schieffelin)
 Rubrigen 30 50 1000 mcg /cc (Upjohn)
 Betalin 12 15 30 100 1000 mcg /cc (Lilly)
 Bevidox 15, 30 150 1000 mcg /cc (Abbott)
 Bexii 15, 30 mcg /cc (Chicago Pharm)
 Bexiifort 100 mcg /cc (Chicago Pharm)
 Bexii M. 1000 mcg /cc (Chicago Pharm)
 B Twelv 30 50 mcg /cc (Sherman)
 B Twelve Amp (Asher)
 Cobavite 15, 30 100 1000 mcg /cc (Metra)
 Cobiane 15 mcg /cc (Merck Sharp & Dahme)
 Casma Rey 1000 mcg /cc (Jeffery)
 Crystamin 30 60, 120 2000 mcg /cc. (Armaur)
- Cytobex 15 mcg /cc (Parke, Davis)
 Docibin 30 mcg /cc (Walker)
 Dodecavite 15, 30, 50, 100, 1000 mcg /cc (U S Vitamin)
 Dodecabee 15, 30, 100, 1000 mcg /cc (Miller)
 Dadex 15, 30, 100, 1000 mcg /cc (Organon)
 Ducabee 15, 30, 100, 1000 mcg /cc (Breon)
 Hemomin 30, 60 120, 1000 mcg /cc (Kirk)
 Normocytin 30 60 mcg /cc (Lederle)
 Pernajek, 300 mcg /cc (Elder)
 Rametin 10, 15 30 mcg /cc (Bio-Ramo)
 Redisol 30, 100, 1000 mcg /cc (Merck Sharp & Dahme)
 Rubasal 30, 50, 120, 250, 1000 mcg /cc (Central Pharm)
 Rubramin 15 30, 1000 mcg /cc (Squibb)
 Sytobex 15, 30, 100, 1000 mcg /cc (Parke, Davis)
 Twel Be 15, 60, 1000 mcg /cc (Parke Davis)
 Vibalt Vial (Roerig)
 Vibrumin 30 60, 100, 250 500 1000 2000 mcg /cc (Philadelphia Ampoule Labs)
 Vitral 15 30, 60, 1000 mcg /cc (Fint Eaton)
 Vi Twel 30, 60, 120, 1000 mcg /cc (C D Smith)
- W/Adenosine 5-manaphosphate**
- See Adeno B 12 Vial (Kirk)
 Adenovite Vial (Metropolitan Labs)
 Cobaden Amp (Rand)
 Thybadein Vial (Bell Pharmacol)
 TY B Amp (Paul Maney)
- W/Adenosine 5 manaphosphate, sod nicotinate procaine** See: Bastran Amp (Wom pole)
- W/Chlorophyllin ferrous sulf** See Cuferyll Tab (Lakeside)
- W/Duodenal ext conc** See Bevidoral, Film tabs (Abbott)
- W/Duodenal whole desic, ferrous gluconate** See: Btrinsic Cap (Elder)

W/Folic acid

See:

- B Pol dox (Abbott) /Cap
 Dodecafol (U S Vitamin) /Tab
 Rubrafal n (Squibb) /Cap
 Rubrafolin (Squibb) /cc.

Vit B-12

- 25 mcg
 30 mcg
 25 mcg
 15 mcg

Folic acid

- 1.7 mg
 1.76 mg
 1.67 mg
 5 mg

W/Folic acid & liver

See:

- Anplex (Chicago Pharm)
 Anplexolt (Chicago Pharm) /cc
 Anplexfort (Chicago Pharm) /cc

Vit B 12

- 10 mcg
 25 mcg
 30 mcg

Folic acid

- 5 mcg
 5 mg
 10 mg

Liver

- 2 u
 20 mcR
 20 mcR

| W/Folic acid & liver. | Vit. B-12 | Folic acid | Liver |
|----------------------------------|-----------|------------|---------|
| Anplexix (Chicago Pharm.) /cc. | 60 mcg. | 10 mg. | 10 mcg. |
| Asplexul (Chicago Pharm.) /cc. | 10 mcg. | 5 mg. | 2 mcg. |
| Dolocyte (Miller) /cc. | 30 mcg. | 10 mg. | 10 mcg. |
| Folaxel (Elder) /cc. | 10 mcg. | 5 mg. | 10 mcg. |
| Hepteryl-12 (Kremers-Urban) /cc. | 20 mcg. | 5 mg. | 10 mcg. |
| Lifalbox (Central Pharm.) /cc. | 5 mcg. | 2.5 mg. | 5 mcg. |
| Lifalbox Strong (Central) /cc. | 50 mcg. | 10 mg. | 10 mcg. |
| Palivite (Sherman) /cc. | 10 mcg. | 5 mg. | 2 mcg. |
| Tweltone (Pitman-Moore) /cc. | 50 mcg. | 10 mg. | 10 mcg. |

W/Folic acid & other nutritional supplements.

| See: | Vit. B-12 | Folic acid | Others |
|---------------------------------|-----------|------------|---|
| Armatnic Special (Armour) /Cap. | 10 mcg. | 1 mg. | Vit. C 50 mg., Liver & Duodenum 350 mg. |
| Crystamin Forte (Armour) /Cap. | 100 mcg. | 1.75 mg. | Duodenum 75 mg. |

W/Homatropine methylbromide & physostigmine salicylate. See: Artha-B-12, Vial (Columbus)

W/Intrinsic factor concentrate, N.F. Cap. Tab., N.F., N.N.D. 1962.

See: Bevidoral, Tab. (Abbott)
 Bexltab, Tab. (Chicago)
 Bifactan, (Organon)
 Biogeman (Organon).
 Biopor (Armour).
 Glabotrin (Patch).
 Intrinsic, Tab. (Upjohn)

W/Liver & other nutritional supplements.

| See: | Vit. B-12 | Liver | Others |
|-----------------------------|-----------|---------|--|
| Heptabee, Amp. (Endo) /cc. | 10 mcg. | 0.2 cc. | Vit. B-1 10 mg., B-2 2 mg., Nicotin 15 mg. |
| Malplee, Vial (McNeil) /cc. | 5 mcg. | 2 mcg. | Vit. B-1 10 mg., B-2 2 mg., Nicotin 40 mg. |

W/Liver & other nutritional supplements.

| See: | Vit. B-12 | Liver | Others |
|---|-----------|---------|---|
| Iexavite, Vial (Lilly) /cc. | 8 mcg. | 2 mcg. | Vit. B-1 10 mg., B-2 5 mg., Nicotin 150 mg., Pantothenic Acid 5 mg., Vit. B-6 5 mg. |
| Bythragee, Vial (Harvey) /cc. | 30 mcg. | Present | Vit. B-1 4.5 mg. |
| Baticulgee-Fortified, Amp. (Lilly) /0.5 cc. | 20 mcg. | 20 mcg. | Vit. B-1 5 mg. |

W/Methylandrostanedial, hydrocortisone.

See: Vistabolic, Tab. (Organon)
W/Serpine & vit. B-1. See: Somatovite, Tab. (Purdue Frederick)

W/Pyridic substance. See: Dodex, Tab. (Organon). Vit. B-12 2 mcg., pyridic substance 50 mg.

W/Salicylamide, p-Aminobenzoic acid & di-pyrone. See: Cabasal, Tab. (Rand Pharm.)

W/Stomach substance. See: Harvite-12, Tab. (G. F. Harvey). Vit. B-12 25 mcg., stomach substance 50 mg.

W/Thiamine HCl.

Mal-Iron Panemic. (White Lab.)

Pronemia (Rederis)

Redfact, Tab. (Merck Sharp & Dohme)

Trinsicon (Lilly).

W/Iron. Vit. B-1. See: Troph-Iron, Tab. (Smith, Kline & French)

W/Liver.
 See: Cru-B-12 (Thompson) Liver equiv. 2 mcg., Vit. B-12 10 mcg./10 cc.
 Sulivon (Squibb) Liver equiv. 20 mcg., Vit. B-12 30 mcg./10 cc.

See: Apatele, Drops. (Kenwood Labs) Vit. B-12 25 mcg., B-1 15 mg./cc.

See: Appetum, Tab. chewing-gum. (Phila. Ampoule Labs.)

Cristivite, liq. & Vial. (Hiss)

Throcabin, Tab (Bryant)

Trophite, Elixir & Tab. (Smith, Kline & French) Vit. B-12 25 mcg., B-1 10 mg./dr. or tab.

VITAMIN B COMPLEX, N.N.D. 1962, A.D.B.
 Conc. ext. of dried brewer's yeast & ext. of corn processed w/Clostridium acetobutylicum.

M R Thompson—Syr, Vit B 1 0.3 mg & other B factors from 2 Gm dried yeast/ B 2 0.2 mg B 6 0.1 mg, niacin 1.4 mg cc Bot 6 oz, 1 pl & 1 gal

Vica Product Co.—Syr (Same as above)

VITAMIN B COMPLEX (Upjohn) Thiamin HCl 3 mg riboflavin 3 mg, pyridoxine HCl 0.25 mg nicotinamide 15 mg calcium pantothenate 5 mg /Cap or Tab Bot 100s 1000s

Use Vitamin B therapy

VITAMIN B COMPLEX TABLETS [E S Miller] Thiamine HCl 2 mg, riboflavin 0.5 mg pyridoxine HCl 0.1 mg, calcium pantothenate 0.5 mg, niacinamide 5 mg In a base of brewer's yeast 230 mg /Tab Bot 100r 500s, 1000s

VITAMIN B COMPLEX W/SEDATIVES

See: Belbarb B (Haskell)

Beplete (Wyeth)

Beplete Belladonna (Wyeth)

Fenabex (Strobenburgh)

Pheno Bepadol (Ives-Cameron)

VITAMIN C, USP, NND 1962, ADR

See: Ascorbic Acid, Prep

VITAMIN CANDIES, (West Ward) Vitamins A, D, B 1, B 2, C, B 12 E B 6 niacinamide folic acid in daily amounts needed/tab. Bot 30s, 1000s

Use: (3 daily) Nutrition

VITAMIN D, SYNTHETIC, NND 1962

See Activated 7 Dehydro cholesterol

Calciferol

VITAMIN D-2, NND 1962 ADR

See: Calciferol, Prep (Various Mfr.)

Vitosterol (Various Mfr)

Sterogyl (Roussel)

VITAMIN D-3, ADR (No specific compounds listed)

See: Activated 7 dehydrocholesterol

Calciferol Prep for related activity

VITAMIN D-3-CHOLESTEROL, Cpd of crystalline vit D 3 & cholesterol

VITAMIN D-4, See: Dihydrocholesterol Prep (Various Mfr)

VITAMIN D W/CALCIUM SALTS.

See: Calcicaps. (Nan)

Calcigranules (Walker)

Calc D (Irwin, Neisler)

Calcifortir D (United)

Calcium Gluconate W/Vit D (Uby)

Calciflofers (Nion)

Calcglyphos-D (Maltbie)

Caltrad (Winthrop)

Calnutrion (Upjohn)

Calcicarbonate (Cole Chem)

Devical. (Prama)

Dical-D (Abbott)

Diostate-D (Upjohn)

Dynacal (McNeill)

Fleoras D (National)

Or Cal (Marion)

Sherdical (Sherman)

Vitamin D Calcium Phosphate (C D Smith)

Via Cal (Rowell)

VITAMIN E, NND 1962, ADR

See Tocopherol, Prep (Various Mfr)

Wheat Germ Oil (Various Mfr)

VITAMIN F.

See Fats, Unsaturated

Fatty Acids, Unsaturated

VITAMIN G, ADR See: Riboflavin

VITAMIN H, See: Biotin, Prep (Various Mfr.)

VITAMIN K, NND 1962, ADR

See Menadiol, Sodium Diphosphate Prep (Various Mfr)

Menadiolone, Prep (Various Mfr)

Menadiolone Sod Bisulfite, Prep (Various Mfr)

VITAMIN K-1, USP, NND 1962

See Aquamephyton Inj (Meick Sharp & Dahme)

Konakion, Caps & Amps (Rache)

Mephyton, Amp & Tabs (Meick Sharp & Dahme)

Phytonadione

VITAMIN K₂, NND 1962 See: Menadiolone

VITAMIN K-5, See: Synkamin, Kapsel (Parke, Davis)

VITAMIN K OXIDE Not available, but usually K-1 is desired

VITAMIN M See: Folic Acid, Prep (Various Mfr)

VITAMIN P, ADR (Citrin)

See: Bio Flavonoid Cpd (Various Mfr.)

Hesperidin Preps (Various Mfr)

Quercetin (Various Mfr)

Rutin, Prep (Various Mfr)

VITAMIN U Present in cabbage juice See

U Vita, Cap (Pitcher)

VITAMINS W/ANTI-OBESITY AGENTS

See: Adjudgets (Wyeth)

Amphedase (Parke, Davis)

Amplus (Roerig)

Amvical (Stuart)

Bephoid (C D Smith)

Deeglan (Kendall)

Desaxet (Abbott)

Fetamin (Misslan)

Norexlin (Smith Dorsey)

Obedrin (Massengill)

Obocell (Irwin, Neisler)

Opidice (Boyle)

Raphetabex (Strobenburgh)

Revicaps (Lederle)

Trand (Bunnington)

VI Dexamin (Smith, Kline & French)

VITAMINS W/LIVER & LIPOTROPIC AGENTS.

- See: B Complex, Fort. (Walker)
 Bejectol W/Liver, (Abbott)
 Ebicol, (Thompson)
 Ferralip Plus. (Flint, Eaton)
 Forty Plex. (Philo, Amp)
 Geratose. (Patch)
 Hepanine. (Bell Pharm.)
 Hepluna. (Roerig)
 Hyanal. (Walker)
 Jaco-B-Co. (Pitman-Moore)
 Lederplex. (Lederle)
 Lipo-Heplex. (U. S. VII.)
 Lipomic. (Chicago)
 Mejaln. (Mead Johnson)
 Methhepanex. (Rawl)
 Methischol. (U. S. VII.)
 Omin-Beta. (Warner-Chilcote)
 Pantabee. (Haskell)
 Poly-B. (U. S. VII.)
 Poly-B W/C. (U. S. VII.)
 Pravite-B. (Ives-Cameron)
 Pravite-B W/C. (Ives-Cameron)
 Randamin-B. (Rand)
 Stuart Lipotaine. (Stuart)
 Burbex. (Abbott)
 Yi-Syneral Therapeutic. (U. S. Vit.)

VITAMINS W/LIVER, LIPOTROPIC AGENTS & MINERALS.

- See: Gerject, In. (Kirk)
 Hyetole. (Merck Sharp & Dahme)
 Hyetole-12. (Merck Sharp & Dahme)
 Jerial. (Brewer)
 Livipeptone. (Borchardt)
 Mejaln. (Mead Johnson)
 Pantabee. (Haskell)
 Yi-Ferrin. (Lederle)

VITAMIN, MAINTENANCE FORMULA.

- See: Octaplex, Cap. (Am. Pharm. Corp.)
 Pargran, Cap. (Squibb)
 Pluravit, Drops. (Winthrop Labs.)
 Pluravit, Pallet. (Winthrop Labs.)
 Pluraxin, Cap. (Winthrop Labs.)
 Surgimin, Cap. (Walker Lab.)
 Yi-Magna, Cap. (Lederle)

VITAMINS: STRESS FORMULA.

- See: A.S.F. (Roerig)
 Becemar. (Chicago)
 Becamin. (Ayerst)
 Cebefartis. (Upjohn)
 Convalets, Pflm. (Abbott)
 Falbesyn. (Lederle)
 National Research Council, Formula.
 Navogran. (Squibb)
 Nvlactan. (Organon)
 Panalins-T. (Mead Johnson)
 Pravite B W/C, Cap. & Tab. (Ives-Cameron)
 Solu-zyme C. (Upjohn)
 Stresscaps. (Lederle)

- Surgimin-T. (Walker)
 Thera-complex Kaps. (Parke, Davis)

VITAMINS: STRESS FORMULA W/ANTI-BIOTICS.

- See: Achromycin-SF. (Lederle)
 Aureomycin-SF. (Lederle)
 Terramycin-SF. (Pfizer)
 Tetracyc-SF. (Pfizer)

VITAMINETS. (Hoffman-La Roche) Vit. A 1667 u., B-1 0.667 mg., B-2 0.667 mg., D 167 u., E 1 mg., d-panthelol 5 mg., niacinamide 5 mg., iron, Cal., Phos., Mang. Mag./Tab. & ascorbic acid 16.67 mg. Bot 100s, 250s & 1000s.

Use: Dietary supplement.
VITAMIX A.T.A. (Vitamin Corp) Morning: dl-Amphetamine HCl 1/6 gr., thyroid 1 gr., atropine sulf. 1/360 gr., alein 1/4 gr./Tab Noon: dl-Amphetamine HCl 1/6 gr., thyroid 1 gr., atropine sulf. 1/360 gr./Tab. Afternoon: Phenobarbital 1/4 gr., dl-Amphetamine HCl 1/6 gr., thyroid 1 gr./Tab. Bot. 100s, 300s.

Use: Treatment of obesity.
VITAMNAL. (Prama) Vit. A 15,000 u., D 1500 u., B-1 5 mg., B-2 5 mg., B-6 1 mg., B-12 1 mcg., niacinamide 50 mg., C 120 mg., cal. pantothenate 10 mg., folic acid 0.25 mg., Vit. E 5 LU, biotin 50 mcg., inositol 20 mg./Cap. Bot. 100s.

Use: Vitamin deficiencies.
VITAMOIDS. (Walker) Vit. A 5000 u., D 500 u., B-1 3.0 mg., B-2 2.0 mg., B-6 0.1 mg., B-12 1.0 mcg., C 300 mg., cal. pantothenate 10 mg., niacinamide 200 mg., folic acid 0.2 mg., iron, copper, mang., zinc, cal., phos., mag., iodine/Cap. Bot 100s, 250s, 500s.

Use: Dietary.
VITANATE. (Chicago Pharm.) Ferrous sulfate 1 1/2 gr., Dicalcium phosphate 3 1/2 gr., Vit A 700 u., B-1 0.16 mg., B-2 0.334 mg., D 250 u., E 0.34 mg., niacinamide 5 mg./Tab. Bot. 100s, 500s, 1000s. Also avail-w/Vit. C 25 mg.

Use: Supplement for diet.
VITANATE-C. (Chicago Pharm.) Ferrous sulfate, excic., 1.5 mg., dicalcium phosphate 3.75 gr., Vit. A 700 u., B-1 0.167 mg., B-2 0.334 mg., C 0.25 mg., D 250 u., E 0.34 mg., nicotinamide 5 mg., folic acid 0.167 mg., Vit. B-12 w/intrinsic factor 1/30 u./Tab. Bot. 100s, 200s, 1000s.

Use: Dietary supplement.
VITANEX. (Estra) Thiamine HCl 100 mg., riboflavin 2 mg., pyridoxine 10 mg., Cal. pantothenate 1 mg., niacinamide 200 mg., benzyl alcohol 2%/cc. Amp. 1 cc. Vial 10 cc., 30 cc.

Use: Deficiencies of vitamin B complex.

- VITA-NINE** (Sherman) Vit A 5000 u, D 1000 u, thiamine HCl 15 mg, riboflavin 2 mg, ascorbic acid 37.5 mg, pyridoxine 0.1 mg, niacinamide 20 mg, Cal pantothenate 1 mg, mixed tocopherols 2 mg / Tab Bot 100s
Use: Dietary supplement
- VITAPETOSE W/Karaya** (Strassenburgh) Gum Karaya 66 gr, apple pectin 2 gr, Vit B 1 0.75 mg/dr Bot 3 1/2 oz, 1 lb
Use: Laxative
- VITAPLEX C.** (Torrigian) Thiamine HCl 10 mg, riboflavin 6 mg, pyridoxine HCl 6 mg, niacinamide 80 mg, ascorbic acid 100 mg / 2 cc Amp 2 cc, Box 12s, 25s, 100s
Use: Vitamin B deficiencies
- VITA-PLUS H LIQUID.** (Scot Tuzian) Vit B 1 10 mg; l-lysine monohydrochloride 100 mg; B 12 25 mcg; B 6 5 mg iron pyrophosphate soluble 100 mg / tsp Bot pts
Use: Hemaline
- VIT-AQUA.** (Taylor) Water miscible Vit A 25M u & 50M u / Cap Bot 100s, 1000s
- VITAPRIME** (Vitaprime Co) Tab Bot 100s, 270s, 540s
- VITA RESPIRAL.** (Premo) Reserpine 0.25 mg, d-amphetamine sulf 2.5 mg, Vit A 12,500 u, D 500 u, B 1 5 mg B 2 5 mg, B 6 2.5 mg, C 150 mg, niacinamide 50 mg, pantothenol 9 mg / 5 cc Bot 4 oz Also supplied as a sealal with 5 mcg Vit E 12 added
Use: Treatment of anxiety, tension, and depression
- VITAROID.** (Warren Teed) Thyroid 32 mg, Vit D 200 u, ascorbic acid 15 mg, riboflavin 1 mg, thiamine HCl 3 mg, nicotinamide 5 mg / Tab, Bot 100s, 1000s
Use: Thyroid replacement therapy
- VITA-SED.** (Klug) Phenobarbital 1/2 gr, Vit B 1 1 1/2 mg, B 2 2 mg, C 20 mg / Tab Bot 500s
Use: Sedative, vitamin combination
- VITASIRUP.** (Walker Vitamin) Thiamine HCl 2 mg, riboflavin 0.25 mg, niacinamide 6 mg, folic acid 0.10 mg, Sod hypophosphite 70 mg, Pot hypophosphite 70 mg, cobaltous chloride 2 mg, Mang hypophosphite 10 mg, ferrous sulfate 50 mg / dr Bot 4 oz
Use: Nutritional supplement
- VITASOL IMPROVED.** (Mallard) Vit B 1 12 mg, B-6 0.5 mg, pantothenol 1 mg, B 2 1.3 mg, B-12 5 mcg, nicotinic acid 5 mg, Iron & ammonium citrate 40 mg, zinc sulf 1 mg, amm chloride 1 mg, mang sulf 1 mg, mag chloride 1 mg, cobalt sulf 1 mg, pot chloride 1 mg, choline dihydrogen citrate 50 mg / 30 cc. Bot B 8 oz, 1 pt, 1 gal
Use: Nutritional deficiencies
- VITATHYRIN.** (National Drug) Thyroid 1 gr thiamine HCl 6 mg / Tab Bot 100s, 500s, 1000s, 5000s
Use: Thyroid deficiencies.
- VITA-TREAT.** (Kirkman) Vit A 5M u, D 1M u, C 100 mg / Tab Bot 100s
- VE-TEENS.** (Lanteen) Super potency, Vit A 5000 u, B 1 4 mg, B 2 3 mg, niacinamide 20 mg, pyridoxine 1 mg, Vit C 75 mg, D 500 u, mixed tocopherols 5 mg, B complex factors in brewer's yeast, w/ niacinamide 15 mg, iron 10 mg / Tab Pkg 30s 90s, 250s Homogenized vitamins Vit A 3000 u, B-1 1 mg, B 2 1.5 mg, C 40 mg, D 800 u, niacinamide 4 mg / dr Bot 2 fl oz, 4 fl oz, 8 fl oz
Use: Dietary supplement
- VITEBBA.** (Roerig) Cobalt 0.1 mg, copper 1 mg, iron 10 mg, iodine 0.15 mg, Cal 213 mg, Mang 1 mg, Mag 6 mg, molybdenum 0.2 mg, Phos 165 mg, Pot 5 mg, zinc 12 mg, Vit A 5000 u, D 500 u, thiamine HCl 3 mg, riboflavin 3 mg, pyridoxine HCl 0.5 mg, niacinamide 25 mg, Vit E 37 IU, ascorbic acid 50 mg, pantothenate 5 mg, B 12 1 mcg / Cap Bot 100s
Use: Mineral & vitamin supplement
- VITERRA TABTAR** (Roerig) Vit A 5000 u, D 1000 u, B 1 1 mg, B 2 2 mg, B 6 1 mg, R 12.2 mcg, C 50 mg, niacinamide 12 mg, cal pantothenate 2 mg, cobalt 0.014 mg, copper 0.07 mg, iodine 0.05 mg, iron 1 mg, pot 0.016 mg, Ma 0.01 mg, mang 0.028 mg, mag 0.108 mg, zinc 0.071 mg / Tab Bot 100s, 250s
Use (1 daily) Vitamin & mineral supplement
- VITERRA THERAPEUTIC.** (Roerig) Vit A 25,000 u, D 1000 u, R 1 10 mg, B 2 5 mg, B-6 0.5 mg, niacinamide 100 mg, Vit C 150 mg, B 12 5 mcg, Cal pantothenate 1.5 mg, w/Cal, cobalt, copper, iodine, iron, Mang, Mag, molybdenum, Phos., Pot & zinc/cap Bot 30s, 100s
Use: Nutritional deficiency
- VI-TESTRIN.** (Savage) Methyl testosterone 5 mg, potassium estrone sulfate 0.25 mg, 1 thyroxine sodium 0.025 mg, Vit A 5000 u, C 50 mg, D 500 u, B 1 2.0 mg, R 2 2.0 mg, B-6 0.5 mg, niacinamide 150 mg, choline 13.5 mg, Inositol 25 mg, dl methionine 25 mg, molybdenum 0.1 mg, iodine 0.15 mg, d pantothenol 5 mg, calcium 60 mg, phosphorus 45 mg / 45 cc Bot pint, gal
Use: Geriatric
- VITETBIN, N N B 1935, A D R.** (Squibb) Vit B 1 10 mg, B-2 5 mg, niacinamide 150 mg

ascorbic acid 150 mg./Tab. Bot. 30s, 100s, 250s, 1000s
 Use: Vitamin B complex w/Vitamin C.

VI-THYRO. (Roerig) Thyroid 60 mg., Vit. B-1 3 mg., B-2 2 mg., B-12 1 mcg., niacinamide 5 mg., cal. pantothenate 5 mg., Vit. C 25 mg., B-6 1 mg., A 3000 u., D 200 u., E 1 mg., cobalt 0.2 mg., iodine 0.1 mg., mag. 10 mg., mang. 2 mg., inositol 10 mg., choline bitartrate 50 mg., methionine 100 mg./cap. Bot. 100s.
 Use: Thyroid therapy w/vitamins, minerals & metabolic substances.

VITIKON SYRUP. (Upjohn) Thiamine HCl 2 mg., riboflavin 3 mg., ascorbic acid 100 mg., nicotinamide 30 mg./dr. Bot. 4 fl. oz., 12 fl. oz.
 Use: Dietary supplement.

VITO-LEXIN. (National Drug) Thiamine HCl 6 mg., iron glycerophosphate 1½ gr., Cal. glycerophosphate 4 gr., Mang. glycerophosphate 1 gr., strychnine glycerophosphate 1/60 gr., hyoscyamus alkaloids D.2 mg./dr. Bot. 8 fl. oz., 1 pt., 1 gal.
 Use: Hematinic.

VITON. (Dumas-Wilson) Vit. A 5000 u., O 500 u., B-1 3 mg., B-2 2 mg., B-6 D.25 mg., C 50 mg., niacinamide 20 mg., cal. pantothenate 1 mg., B-12 1 mcg., folic acid 0.25 mg., E 1 mg., with trace elements/Tab. Bot. 100s, 1000s.
 Use: (1 dolly) Dietary supplement

VITONASE. (Lekaside) Pancreatin (activity of pancreatin N.F. 300 mg) 100 mg., proteolytic enzyme 6 mg., amylolytic enzyme 10 mg., tauracholic acid 15 mg, Vits. B-1 1.5 mg., B-2 1 mg., nicotinic acid 10 mg., C 25 mg., Vit. A 2000 u., ferrous fumarate 50 mg./Tab. Bot. 60s.
 Use: Digestive agents.

VI TOT. (Mizemes) Vit. A 5000 U.S.P. u., D 500 U.S.P. u., B-1 2.5 mg., B-2 2.5 mg., C 50 mg., nicotinamide 20 mg., B-6 0.5 mg., sodium pantothenate 5 mg., B-12 2 mcg., folic acid 0.25 mg./Tab. Bot. 100s.
 Use: Vitamin supplement.

VITRAMONE. (O. F. Harvey) Natural estrogenic hormone 1 mg., w/vit. B-1 10 mg., B-2 2 mg., B-6 5 mg., niacinamide 40 mg., inositol 50 mg., choline chloride 10 mg., methionine 10 mg./cc. 10 cc.
 Use: Estrogen therapy w/Vitamin B complex.

VITRATE. (Upjohn) Iron & ammonium chlorates 4 gr., Vit. A 25,000 u., D 1200 u., malt & yeast ext./oz. Bot. 10 fl. oz., 24 fl. oz.
 Use: Hematinic.

VITREXIM-M. (Page) Tab Bot. 100s.
VITREXIN. (Page) Tab. Bot. 50s, 100s.
VITRIMIN TABSULES. (Poul Maney) Estrogenic substance conjugated 0.25 mg., meth-

folic acid 1.5 mg., B-1 2 mg., B-2 2 mg., nietyfloxsterone 2.5 mg., Vit. B-1 5.5 mg., C 55 mg., B-12 2 mcg., folic acid 2 mg., terrous sulfate dried 100 mg., brewer's yeast 200 mg., d-desoxyephedrine HCl 1 mg./Tab. Bot. 100s, 500s & 1000s.
 Use: Geriatric & postmenopausal patients.

VI-TRINS. (U. S. Vitamin) Vit. B-12 w/intrinsic factor conc. 0.5 u., liver des. 200 mg., ferrous sulfate exs. 340 mg., vit. C 100 mg., cinamide 15 mg., B-6 0.5 mg., d-cal. pantothenate 1.5 mg./cap. Bot. 50s, 100s.
 Use: (2 dolly) Anemias.

VITRON. (Smith, Miller & Patch) Ferrous sulfate 200 mg. thiamine HCl 1 mg., riboflavin 1 mg., Vit. B-12 1 mcg./Tab. Bot. 100s, 1000s.
 Use: Vitamin supplement.

VITRON-C. (Smith, Miller & Patch) Ferrous fumarate 200 mg., ascorbic acid 125 mg./Tab Bot. 100s, 1000s.
 Use: Iron Deficiency Anemia.

VITRON MODIFIED. (Smith, Miller & Patch) Ferrous gluconate 600 mg., Thiamine HCl 8 mg., fl. oz. liq Bot. 4 oz., pt., gal.

VITRONBEX. (Smith, Miller & Patch) Ferrous sulfate, exs. 3 gr., thiamine HCl 1.5 mg., riboflavin D.5 mg., niacinamide 5 mg. w/Vit. B complex/Tab. Bot. 100s, 1000s.
 Use: Hematinic w/Vitamin B supplement.

VITT. (Warren-Teed) Vit. A 17,800 u., D 890 u., iron & ammon. citrate 40 mg./oz Bot. 1 pt., 1 gal.
 Use: Hematinic

VITULES. (Wyeth) Vit. A 5000 u., D 1000 u., ascorbic acid 75 mg., riboflavin 3 mg., nicotinamide 20 mg., pyridoxine HCl 0.2 mg., Cal. pantothenate 20 mg., Vit. B-1 2 mg./Cap. Bot. 30s, 100s.
 Use: Dietary supplement.

VIT-U-PEPT. (Ovesseas) Extract of brassica capitate (anti-ulcer factor "U" from raw cabbage) 150 mg, bismuth carbonate 50 mg, sodium bicarbonate 120 mg., magnesium carbonate 200 mg./Tab. Bot. 24s, 72s, 144s.
 Use: Gastric and intestinal ulcers.

VI-TWEL. (Smith, Miller & Patch) Vit. B-12. Vial 30, 60, 120 mcg/cc. In 15 cc. vials; 1000 mcg/cc. In 10 cc. vials.

VITWILBEX. (Smith, Miller & Patch) Vit. B-12 20 mcg., thiamine HCl 10 mg., niacinamide 50 mg., riboflavin 3 mg., pyridoxine HO 5 mg., panthelal 5 mg., gentils acid ethanalamine 2½%/cc. Vial 10 cc.
 Use: Vitamin B complex therapy.

VI-TYKE. (Lederle) Vit. A 3000 u., D 800 u., B-1 1.5 mg., B-2 1.5 mg., B-6 1.0 mg., C 40 mg., B-12 3 mg., pantothenic acid 1

mg, niacinamide 10 mg, w/cherry flavor/
Tsp Push button 12 fl oz

Use: Dietary supplement

VI-TYKE DROPS. (Lederle) Vit A 5000 u,
D 1000 u B 11 mg, B 2 0.8 mg, B 6 1 mg,
C 50 mg B 12 1 mcg, niacinamide 10 mg
pantothenic acid 2 mg /0.6 cc Dropper
Bot 50 cc

Use: Nutritional supplements

VI-ZO-B. (Pitman Moore) Vit A 5000 u,
D 800 u, thiamine HCl 1.5 mg, riboflavin
2 mg, ascorbic acid 30 mg, niacinamide
5 mg, Cal pantothenate 1 mg, mixed
natural tocopherols 6.16 mg /Cap Bot
100s, 250s

Use: Dietary supplement

V-KOR. (Lilly) Y Cillin K (Penicillin V Pot)
62.5 mg (100 000 u) Pyronil 7.5 mg
histadyl 12.5 mg, clopene 6.25 mg, as-
pirin 114 mg, acetaphenetidin 80 mg, caf-
feine 16 mg /Tab Bot 50s

Use: Antibiotic-antihistaminic-analgesic ther-
apy

VLEM-DOMB (Dane) Vlemincx solution in
powder form Calcium polysulfide, calcium
thiosulfate, sulfur Pkg 12s, 100s, Jar 4 oz
Use: (1 pk in 1 pt) Skin conditions (acne).

VLEMINCCKX'S LOTION (Ulmer) Lime sulfur
rated sal Bot 1 pt

Use: Antiseptic

VLEMINCCKX SOLUTION. (Ulmer) Solution of
Sulfurated Lime NF Bot 1 pt See Vlem-
Domb Pow (Dane)

VONDAR (Lilly) Propoxyphene 32 mg,
phenaglycolol 150 mg, acetylsalicylic acid
325 mg /Cap

Use: Analgesic and relaxant

VONEDRINE HYDROCHLORIDE, NND
1962 Sal (Merrill) Vonedrine (phenylpro-
pylmethylamine) HCl 2.8%, ceepryn chlo-
ride 0.02% Bot 1 oz, 1 pt

Use: Nasal decongestant

VONTIL (Smith Kline & French) Thioperazine.

Use: Tranquilizer antiemetic

VoSol DROPS (Wampale) 1.2 propaned ol
diacetate 3%, acetic acid 2%, benzethon-
ium Cl 0.02%, in propylene glycol Plas
Bot 15 cc Also available w/hydrocortisone
1% Bot 7 1/2 cc

Use: Central of all his externa

VYCOM B PLUS (Bristol) Thiamine HCl
5 mg, riboflavin 5 mg, niacinamide 50
mg, pyridoxine HCl D.5 mg, Cal panto-
thenate 2 mg yeast 100 mg ascorbic acid
50 mg /Tab Bot 100s, 500s

Use: Nutritional supplements

VYONS (Vitamin Corp) Vit A 6000 u, B-1
6 mg, B 2 3 mg C 50 mg D 600 u
niacinamide 10 mg, Cal pantothenate 3

* Generic name recognized by NF USP or A.M.A.

mg, Vit B-6 0.5 mg, E 5 mg, whole dried
liver pow 10 mg, dried debittered brewer's
yeast 10 mg, choline dihydrogen citrate 20
mg, folic acid 0.1 mg, Vit B-12 2 mcg,
Iron, Cal, Phos, Iodine Mag, copper,
Mang, cobalt, Pot, zinc, molybdenum/
Cap Bot 24s, 50s, 100s, 250s, 1000s

Use: Dietary supplement

VYTONE CREAM. (Dermik) Hydrocort sane
alcohol 0.5%, diadaphoxyquinoline 1%;
in a greaseless vegetable oil vehicle/
cream Tube 1 oz

Use: Dermatologic

WALCO-NESIN. (Walker Corp) Mephenesin
Tab 5 gr Bot 100s, 1000s

Use: (1 Gm tid) Muscle relaxant

WARCOUMIN, NND 1959 (G F Harvey)
Warfarin sod Tab 5 mg, 10 mg, 25 mg
Bot 100s See also Prothromadin

Use: Anticoagulant

WARPARIN (Various Mfr) Comp 42.3
(alpha Acetonylbenzyl)-4 hydroxy coumarin
Use: Rodenticide

* **WARFARIN SODIUM, USP, NND 1962**
Inj, Tab USP 3-(alpha Acetonylbenzyl)-
4 hydroxycoumarin 8 Hs sod salt
See Coumadin Sodium (Endo)

Panwarfin Tab (Abbott)

Prothromadin, Tab (G F Harvey)

Warcoumin, Tab (G F Harvey)

WATERBURY'S NUTRITIVE. (Standard Lab)
Iron 12 mg, Vit B 1 0.6 mg, Mang Cal
Phos & Pot /oz Bot 14 oz

Use: Tonic

**WATERMELON SEED EXTRACT W/Pheno-
barbital & theobramine** See: Cital Cap
(Table Rock)

WATER-MISCIBLE VITAMIN A, USP
NNR 1955, NND 1959 See Vitamin A
Water Miscible

WELDERS EYE LOTION (Ashlan) Ethacaine
HCl, boric acid camphor glycenn & chlor
butanol Bot 1/2 oz, 1 oz, 2 oz
Use: Relief for arc burn

WESCODYNE. (West Disinfecting Co) Poly-
ethoxy polypropoxy ethanol iodine complex
7.75%, nanyphenylether al propylene gly-
col iodine complex 3.75% (total 1.6%
available iodine)

Use: Highly diluted as disinfectant

WESTHAZOLE VAGINAL (Westwood) 10%
Sulfathiazole 3% Lactic acid & 1% Acetic
acid in polyethylene glycol Jelly 1.5 oz wth
or without applicator, 3 oz

Use: Vaginitis & cervicitis

WET TONE B (Braemmel) Myristyl gamma

picolinium chloride 0.002%, chlorobutanol 0.1%. Soln. Bot. 2 oz.

Use: Wetting solution for contact lenses.

WHEAT GERM MEAL W/ Liver desic. & yeast autolysate. See: Aminothyn, Pow. (Filt-Eaton)

WHEAT GERM OIL (Various Mfr.) See: Tocopherol Prep. (Various Mfr.)

WHEAT GERM OIL (Hilly) Bot. 60 cc., 1 pt. Use: Vitamine E therapy.

WHITE LINIMENT. (Merck Sharp & Dohme) Mullard formula. Contains oil turpentine, ammonia water, oil arignon artificial, & water, emulsified to form permanent white product. Bot. 1 gal.

WHITE LOTION, U.S.P.
See: Latiobloc, (Arnor-Stee)
Fronac, Pow. (E. Faugero)

WHITE PRECIPITATE. See: Ammoniated Mercury, U.S.P.

WHITE'S OTOLITIC. (White) Neomycin sulfate equiv. to 3.5 mg. of base, Sod. propionate 50 mg./cc. Bot. 15 cc.
Use: Otitis externa.

WHITE'S VITAMIN A AND D OINTMENT. (White) Vitamins A and D in a lanolin-petrolatum base. Tube 1½ oz., 4 oz., Jar 1 lb., 5 lb.

Use: To promote healing and granulation

WHITE'S VITAMIN A & D OINTMENT W/ PREDNISOLONE. (White Labs) Prednisolone 0.5%. Oint. 10 Gm. Tube.
Use: Dermatologic.

WHITFIELD'S OINTMENT, U.S.P. (Merck Sharp & Dohme) Benzoin acid 12%, salicylic acid 6%. Tube 1 oz. Jar 1 lb.
Use: Fungicidal.

WHOOPIING COUGH VACCINE, N.M.D. 1962. See: Pertussis Vaccine, U.S.P.

WIGRAINE. (Organon) Ergotamine tartrate 1.0 mg., caffeine 100 mg., belladonna alkaloids 0.1 mg., acetophenetidol 130 mg./Tab. Box 20s. Bot. 100s. Also supp. Box 12s.
Use: 1 q.h. for 6 doses; migraine headache syndrome.

WILCORT. (Wilson) Corticotropin (ACTH) 40 U.S.F. v. Vii. B-12 1000 mcg./cc. Vial 5 cc.
Use: I.M., I.M. treatment & control of inflammation in neuritis & related disorders

WILPO. N.M.D. 1962. (Darsey Labs) Phenentermine HCl, 8 mg./Tab. Bot. 100s.
Use: Obesity.

WIN-CODIN. (Winthrop) Codeine phosphate 15 mg., phenylephrine HCl 10 mg., acetylsalicylic acid 300 mg., chlorpheniramine maleate 2 mg., ascorbic acid 50 mg./Tab. Bot. 100s.
Use: Analgesic.

WINSTROL. (Winthrop) Stanozolol 2 mg./Tab. Bot. 100s.

Use: Anabolic.

WINTETIL. (Winthrop Prod.) Chloromphenicol 250 mg./Cap.

WOLFINA. (Eloyd & Dobney) Rauwolfia serpentina whole root. Tab. (50, 100 mg.) Bot. 100s, 500s, 1000s.

Use: Hypertensive & tranquilizing agent.

WOLSERTINE. (Medco) Tab. 50 mg. Bot. 100s, 100 mg. Bot. 100s.

WOOD CREOSOTE, N.N.R. 1936, A.D.R. 1953. See: Creosote (Various Mfr.).

WOOL FAT, A.D.R. 1954. See: Lanolin Anhydrous (Various Mfr.).

WRIGHT'S SOLUTION. (Faugero) Coal tar solution 5%. Bot. 3, 8, 16 oz. See: U.S.P.

W-T TABLETS. (Warren-Teed) Phenobarbital 1/16 gr., belladonna ext. 1/24 gr., benzocaine 1/16 gr., Col. carb. 6 gr., Mag trisulfate 3½ gr., aluminum hydroxide 2½ gr., chlorophyll ext. 1%/Tab. Bot. 100s, 1000s. Also pow. (1 tsp. = 4 tabs.) Bot. 6 oz.

Use: Antacid & antispasmodic.

WYACORT. (Wyeth) Methylprednisolone 4 mg./Tab.

Use: Corticosteroid.

WYALIN. (Wyeth) Ox bile ext. 1 gr., cascara sagrada ext. 1 gr., nux vomica ext. 1/20 gr., pancreatic enzymes 2 gr./Tab. Bot. 50s, 500s.

Use: (1 l.i.d. & h.s.) Laxative.

WYAMINE, INHALER, N.N.D. 1962. (Wyeth) Mephentermine 250 mg., w/menthol & aromatics.

Use: Nasal congestion.

WYAMINE & PENICILLIN. (Wyeth) (Not listed) Contents of 1 Cap. in ½ oz. water makes Sol. containing 2000 I.U. penicillin & 0.5% mephentermine sulfate. Cap. 6s

Use: Spray; antibiotic & nasal decongestion.

WYAMINE SULFATE ELIXIR, N.N.D. 1962. (Wyeth) Mephentermine sulfate 25 mg./5 cc. in cherry wine base. Bot. 1 pt.

Use: To elevate mood in depressive states.

WYAMINE SULFATE INJECTION, N.N.D. 1962. (Wyeth) Mephentermine sulfate, 15 mg./cc. Vial 1 cc., 10 cc.

Use: I.V. or I.M.; in hypotensive states not associated with hemorrhage.

WYAMINE SULFATE, NASAL SOLUTION, N.N.D. 1962. (Wyeth) Wyamine sulfate 0.5% (isotonic sol.) containing 0.5% chlorobutanol. Bot. w/dropper 1 fl. oz., 1 pt

Use: Nasal decongestant.

WYAMINE SULFATE TABLETS, N.N.D. 1962. (Wyeth) Tab. 12.5 mg., Vial 100s. Tab. 25 mg., Bot. 24s, 100s.

Use: Treatment of mild mental depression.

WYAMINE-TYROTHRINICIN (Wyeth) Wyamine 0.5%, tyrothricin 0.02% Sol 1 oz, w/ jetomizer Nasol sol, Bot 1 oz

Use Antibiotic w/nasal decongestant

WYANOIDS (Wyeth) Ephedrine sulfate 0.1% w/belladonna alk, boric acid zinc oxide bismuth oxyiodide & subcarb bolsam peru & cocoa butter Supp, Box 12s Oint w/applicator 1 oz

Use: Hemorrhoids

WYANOIDS HC (Wyeth) Hydrocortisone acetate 10 mg, belladonna extract D.5%, ephedrine sulf 0.1%, zinc oxide, boric acid bismuth oxyiodide bismuth subcarb bonate/Supp Box 12s

Use Anorectal disorders

WYBIOTIC TROCHES (Wyeth) Bacitracin 300 u neomycin sulfate 5 mg polymyxin B sulfate 200 u in flavored base Can 48s

Use Oral anti infective

WYCHOL (Wyeth) Choline base 3 Gm, inositol 450 mg/15 cc Syr, Bot 1 pt, 1 gal Choline base 0.24 Gm, inositol 0.036 Gm/Cap Bot 100s 1000s

Use Restore normal fat metabolism

WYCHOL-B SYRUP (Wyeth) Choline base 3 Gm inositol 450 mg, Vit B 1 2.5 mg, B 2 0.5 mg, niacinamide 10 mg, Vit B 6 2.5 mg/15 cc Bot 1 pt

Use: Restore normal fat metabolism

WYCILLIN (Wyeth) Procaine penicillin G Aq inj, 300,000 u/dose Aq susp, 300,000 u/Tube Pkg 1s, 50s

WYCILLIN POSTIPIED, (Wyeth) Procaine penicillin G 300,000 u w/Pot penicillin G 100,000 u Vial 1, 5 10 doses

WYCILLIN 600 SUSPENSION, (Wyeth) Procaine penicillin G Aq inj 600,000 u/Tube Pkg 1s, 50s

WYDANE, LIQUID (Wyeth) Chlordane 24% chlordane like compounds 16%, pine oil 10% kerosene 38% Bot 2 oz

Use 2 tsp to 1 qt of watery, external parasiticide for dogs

WYDASE, NND 1962 (Wyeth) (Hyaluronic acid, lyophilized) Vial 150 TR u, Box 1s, 100s; 1500 TR u, Box 1s, 100s

Use Spreading factor

WYDASE IN SOLUTION, NND 1962 (Wyeth) Hyaluronic acid 150 TR u in sterile saline sol Vol 1 cc Store below 59° F, stable 2 yrs

Use: Hypodermoclysis

WY-KOFF (Lemmon) Phenylpropanolamine HCl 60 mg, methamphetamine HCl 6 mg sodium citrate 0.6 Gm alcohol 1%/oz Bot 3 R oz

Use Expectorant

* Generic name recognized by N.F., U.S.P. or A.M.A.

WYNESTRON, (Wyeth) Estrone Aq susp Vial (2 mg./cc) 10 cc

Use: I.M., relief of menopausal disturbances

XANCYLATE, (Pitman Moore) Phenobarbital ¼ gr, Cal theobromine Cal salicylate 5 gr, ethylenediamine dihydriodide ½ gr/Tab Bot 100s, 1000s

Use: Sedative w/diuretic.

XANIOMINE, (Pitman Moore) Theobromine 2½ gr, ethylenediamine dihydriodide ½ gr/Tab Bot 100s, 500s, 1000s

Use Ouretic & vasodilator

XANIOPHEN (Pitman Moore) Theobromine 2½ gr, ethylenediamine dihydriodide ½ gr, phenobarbital ¼ gr/Tab Bot 100s 1000s

Use: Diuretic & vasodilator

XANTHINE DERIVATIVES, NND 1960.

See: Caffeine

Theobromine

Theophylline

XANTHINYLX, (Cale Chem Co) Nux vomica ext 7 mg, caffeine theophylline comp 60 mg/Tab Bot 100s

Use: Stimulant to spinal cord reflexes

* **XANTHINOL HCl** 2-chloro-9-[3-[4-(3-hydroxypropyl)piperazinyl]propyl]thioxan (thene d HCl See Daxid (Roerig)

* **XANTHOCILLIN** Antibiotics obtained from the mycelium of *Penicillium notatum* (Xanthocillin X is 2,3-Diacetyloxy-1,4-d (p-hydroxyphenyl)buta 1,3 diene)

XANTHOTOXIN, NNR See: Methoxsolen NND 1959

Uses Antiseptic & local anesthetic.

* **XENTHORATE HCl** (Warner Lambert Res Inst) 2 diethylaminoethyl 2(4 biphenyl)lthiobutyrate HCl

XENTYSALATE Name approved by BPC 2 Diethylaminoethyl 3 phenylsaliolate 5s bacfen is the hydrochloride

XEROFORM OINTMENT, NNR 1951 (Merck) Tribromophenol bismuth 25%, opth oint base qx 100% Tube ½ oz. Use Eye antiseptic

XI-7, Bithional in Woodbury soap (Jergens)

XPECT (Brayton) Theophylline sod glycinate 60 mg phenylephrine HCl 5 mg, glyceryl guaicolate 50 mg, sod citrate 100 mg/5 cc Bot 8 oz

Use Treatment of cough

XPECTUSSIN (X & M Labs) Dextromethorphan HBr 15 mg chlorpheniramine maleate 2 mg phenylephrine HCl 5 mg pantoic acid 60 mg ascorbic acid 30 mg alcohol 5%/10 cc Bot pl

Use Cough

X-PREP. (Gray Lab.) Anthraquinoidal glycosides. Can $\frac{1}{4}$ oz.
 Use: Clean bowel for x-ray.

X-RAY CONTRAST MEDIA. See: Iodine Products, Diagnostic.

X-TRO. (Xitrium) Aminoxystropine tropate HCl 0.2 mg./Cap. Bot. 100s.
 Use: Anticholinergic.

X-TRO WITH PNEOBARBITAL. (Xitrium) Aminoxystropine tropate HCl, phenobarbital 15 mg./Cap. Bot. 100s.
 Use: Anticholinergic.

XYLOCAINE. (Astra) Lidocaine. Oint., Lidocaine base 2½ & 5%. Jar 35 Gm. Tube 35 Gm. Jelly, Lidocaine HCl 2%, Sod. carboxymethylcellulose, w/preservatives. Tube 30 Gm.

XYLOCAINE HYDROCHLORIDE, N.N.D. 1962, A.D.R. (Astra) Lidocaine HCl. Amp. (2%): 2 cc., 10s, 30s. Vial (0.3%): 20 & 30 cc., 5s, with or without epinephrine. Vial (1%): 20 & 30 cc., 5s, with or without epinephrine. Vial 100 cc.: without epinephrine. Vial (2%): 20 & 30 cc., 5s, w/epinephrine. Use: Local anesthetic.

W/Polymyxin B-Sulfate (Aerosporin), propylene glycol. See: Lidocaine. (Burroughs Wellcome)

W/N-pyrrolidinomethyl)tetracycline, ascorbic acid. See: Yelocycline Intramuscular w/ Xylocaine. (Squibb)

XYLOCAINE HYDROCHLORIDE 4% SOLUTION. (Astra) Etrabulbar injection and topical use. Ampules 5 cc. Box 10s. Topical use only. 50 cc. Screw top Bot.

XYLOCAINE HYDROCHLORIDE FOR SPINAL ANESTHESIA. (Astra) Xylocaine HCl 5%, glucose 7.5%. Specific Gravity 1.030-1.035. Ampules 2 cc. Box 10s, 100s.

XYLOCAINE OINT. 5% FLAVORED. (Astra) Lidocaine 5%/Tubes 3.5 Gm.
 Use: Dental procedures.

XYLOCAINE VISCOSUS. (Astra) Lidocaine HCl 0.02 Gm. w/CMC, preservatives and flavoring agents/cc. Bot. 100 cc., 450 cc. Use: Orally, 1 tabletspoonful for control of hiccup and reflex vomiting.

XYLOMETAZOLINE HCl, N.N.D. 1962. 2-(4-tert-butyl-2,6-dimethylbenzyl)-2-imidazoline HCl. See: Otrivin (Ciba)

YEAST ADENYLIC ACID. (No Mfr. currently lists.) An isomer of adenosine 5-monophosphate & has been found inactive. See: Adenosine 5-Monophosphate preps. for active compound.

YEAST CONCENTRATE. (Burroughs Wellcome) Vit. B-1 0.6 mg., B-2 1.2 mg., ni-

nic acid 6 mg., w/other Vit. B complex found in yeast/Tab. Bot. 100s, 500s, 1000s.
 Uses: Dietary supplement.

W/Methylcellulose. See: Cellulose Tablet (Dumas-Wilson)

YEAST, DRIED, N.F., N.N.R. 1952.

YEAST EXTRACT, N.N.R. 1952. (Kinney) Aq. ext. of brewer's yeast, thiamine HCl 0.375 mg., riboflavin 0.125 mg./5 cc. Bot. 4 fl. oz., 1 pt.
 Use: Dietary supplement.

YEAST EXTRACT, FORTIFIED LIQUID. (Kinney) Thiamine HCl 2 mg., riboflavin 3 mg., niacinamide 20 mg., panthoal 0.86 mg., pyridoxine HCl 0.5 mg./5 cc. Bot. 4 fl. oz., 1 pt.
 Use: Dietary supplement.

YEAST EXTRACT, FORTIFIED TABLET, N.N.R. 1952. (Kinney) Vit. B complex 5 gr., dried brewer's yeast w/thiamine HCl 2 mg., riboflavin 1.5 mg., niacinamide 10 mg., Cal. pantothenate 0.5 mg., pyridoxine HCl 0.25 mg./Tab. Bot. 100s, 1000s.
 Use: Dietary supplement.

YEAST TABLETS, DRIED, N.F. Dumas-Wilson—Tab. 7½ gr., Bot. 100s, 500s, 1000s. Upjohn—Tab. 7½ gr., Bot. 100s, 250s, 500s, 1000s.
 See: Brewer's Yeast, Tab.
 Use: Supplementary source of B complex vitamins.

YEAST, TORULA, N.F. See: Torula Yeast.

YELLOW ENZYME. See: Riboflavin.

YELLOW FEVER VACCINE, U.S.P.

YELLOW MERCURIC OXIDE, N.N.R. 1949. (Merck Sharp & Dohme) Ophth. soln. 1%, 2%, Tube ½ oz. Oint. topically, 2%, Tube ½ oz.

YERBASAL. (El Paso) Tab. Bot. 100s.

YODORIN, N.N.D. 1959. Dilodihydroxyquin.

YONIMBINE W/Thyroid and strychnine. See: Ando-Medicone, Tab. (Medicone)

YUYRAL. (Lederle) Vit. A 5000 u., D 500 u., B-12 1.0 mcg., thiamine HCl 3.0 mg., riboflavin 3.0 mg., niacinamide 20.0 mg., folic acid 0.2 mg., pyridoxine HCl 0.5 mg., Cal. pantothenate 1.0 mg., ascorbic acid 50 mg., Vit. E 5.0 I.U., Iron 15.0 mg., Iodine 0.15 mg., boron 0.1 mg., copper 1.0 mg., Boron 0.1 mg., purified Intrinsic factor conc. 0.5 mg., Mag. 1.0 mg., Mang. 1.0 mg., Pot. 5.0 mg., zinc 0.5 mg., Cal. 69.0 mg., Phos 53.8 mg., dibasic calcium phosphate 236.128 mg., molybdenum 0.2 mg./cap. Bot. 30s, 100s.

ZACTANE CITRATE, N.N.D. 1962. (Wyeth)

- Ethoheptazine citrate 75 mg /Tab Bot 100s
 Use: Analgesic
- ZACTIRIN.** (Wyeth) Ethoheptazine citrate in yellow layer 75 mg, acetyl salicylic acid in green layer 325 mg /Tab Bot 4Bs
 Use: Analgesic
- ZAMINE.** (Marlon) d Amphetamine sulf No 1: 10 mg, No 2: 15 mg /Delayed-action Cap Bot 30s, 500s
 Use: Treatment of obesity
- ZAMITAM No 1 AND ZAMITAM No. 2** (Marlon) Formerly known as Zamine
- ZAMITOL.** (Marlon) No 1: d Amphetamine sulf 10 mg, amobarbital 1 gr /Delayed action Cap No 2: d Amphetamine sulf 15 mg, amobarbital 1½ gr /Delayed action Cap Bot 30s, 500s
 Use: Treatment of obesity
- ZANCHOL.** NND 1962 (Searle) Floran tyrosine 250 mg /Tab Bot 100s
 Use: Hydrochlorotic and gastrointestinal disorders
- ZANTHIMAL.** (Haskell) Theobromine 5 gr, phenobarbital ½ gr /Tab Bot 50s, 500s, 1000s
 Use (111d) Diuretic & sedative
- ZANTRATE.** (Upjohn) Cyclopentenyl allyl barbituric acid ½ gr, ephedrine sulfate ½ gr, theophylline 2 gr /Tab Bot 100s, 500s
 Use: Sedative w/diuretic
- ZARONTIN** (Parke Davis) Ethosuximide α-ethyl α methylsuccinimide 0.25 Gm /Cap Bot 100s
 Use: Treatment of petit mal epilepsy
- ZASORB** (Stiefel) Cellulose plus hexa chlorophene Caps 2 oz
 Use: Moisture absorbent
- ZEMACIDE.** (Tilden) Calamine, zinc oxide glycerin & Col hydroxide w/1% phenol liq, Bot 1 pt, 1 gal
 Use: Antiseptic for skin infections
- ZEMALO.** (Barre) Sulfur, ZnO camphor, Ictium oxide Bot 6 oz
- ZEMARINE ELIXIR** (Zemmer) Butobarbital Sod 162 mg, hyoscyne HBr 0.0065 mg, hyoscyamine sulfate 0.1037, atropine sulfate 0.0194 /dr Bot Pts
 Use: Antispasmodic or sedative
- ZEM-HISTINE** (Zemmer) Methapyriene HCl Oint 5%, tube 1 oz Jar 1 lb Ophth oint 5%, Tube ½ oz Ophth Sol 5%, Bot ½ oz Tab 25 mg, Bot 100s, 1000s
 Use: Antihistaminic
- ZEM-HISTINONE SYRUP.** (Zemmer) Dihydrocadelone bitartrate 125 mg, ephedrine sulfate 5 mg, pyrimine maleate 10 mg, phenylpyramine HCl 5 mg /dr Bot pts
 Use: Bronchial irritations and upper respiratory conditions
- ZENTRON.** (Lilly) Ferrous sulfate 100 mg, thiamine hydrochloride 1 mg, riboflavin 1 mg, pyridoxine hydrochloride 0.5 mg, Vit B 12 crystalline 5 mcg, pantothenic acid 1 mg, nicotinamide 5 mg, ascorbic acid 35 mg, alcohol 2%/5 cc Bot 8 fl oz Box 12s
 Use: Liquid hematinic
- ZEPHIRAN CHLORIDE, NND 1962, ADR** (Winthrop Labs) Benzalkonium Chloride, USP Aq Sol 1:1000 Bot 8 oz, 1 gal Canc sol 17%, Bot 4 oz, 1 gal Tr, stain less 1:750 Bot 8 oz 1 gal tinted 1:150 Bot 8 oz 1 gal
 Use: Cationic detergent & germicide
- ZEPHIRAN CHLORIDE SPRAY.** (Winthrop Labs) Benzalkonium chloride tinted 1 ncture 1:750 (0.13%) Aerosol spray bot 1 fl oz, 6 fl oz
 Use: Skin disinfectants
- ZEPHIRAN TINT.** (Winthrop Labs) Vial 28 gr, Box 4s
 Use: To make United Zephiran tincture
- ZESTAMIN JUNIOR CAPTABS.** (Bryon) Vit A 5000 u, D 500 u, B-1 10 mg, B-2 2.5 mg, B 6 0.25 mg, B 12 5 mcg, folic acid 0.1 mg, niacinamide 10 mg, cal pantothenate 1 mg, Vit C 20 mg, E 2 mg, choline dihydrogen citrate 12.5 mg, Inositol 12.5 mg, dl methionine 12.5 mg, liver desiccated 25 mg, yeast 25 mg, dicalcium phos 50 mg, ferrous gluconate 15 mg, pot iodide 0.13 mg, mag sulf (dried) 2.5 mg, mang sulf 1.5 mg, cobalt sulf 0.5 mg, molybdenum trioxide 0.1 mg /Cap Bot 100s, 1000s
 Use: Supplement the pediatric diet
- ZESTAN.** (Standax) Acetaminophen 250 mg, phenylpropanolamine HCl 25 mg, ammonium chloride 200 mg /tab Bot 100s
 Use: Premenstrual tension and dysmenorrhea
- ZE-TAR-QUIN CREAM** (Dermik) Hydrocortisone alcohol 0.5%, diadaphydroxy quinoline 1%; zetar decolorized 2%; In a greaseless vegetable oil vehicle /Cream Tubes: 1 oz
 Use: Dermatologic
- ZILATONE.** (Drew) Bile salt comp 1½ gr, cascara ext ½ gr, nux vomica ext 1/16 gr, pancreatin ½ gr, phenolphthalein ½ gr, pepsin 3/20 gr, capsicum oleoresin 1/10 gr /Tab Bot 20s, 40s, 80s, 500s 1000s
 Use: Laxative
- ZINADON.** NND 1959 (Victor Pharmacal)

Tab., N.N.R. Isoniazid. Tab. 50 mg., 100 mg. Bot. 100s, 1000s.

Use: TB.

ZIN-AL-COL (Drug Products) Zinc hydroxide, aluminum subacetate, phenol 1%, menthol, glycerin, lime water, rose water. Bot. 1 pt., 1 gal.

Use: Antipruritic & antiseptic.

ZINAX, N.N.D. 1939. (Hynson, Wescott & Dunning) Partially hydrolyzed casein w/2% sod. lauryl sulfate. Gel 4 fl. oz. Tube $\frac{1}{4}$ oz. w/pads.

Use: Burn dressing.

ZINCASATE BURN DRESSING, N.N.D. 1939.

See: Zinox. (Hynson, Wescott & Dunning)

ZINC BACITRACIN W/Neomycin sulfate, polymyxin B, benzalkonium chloride. See: Biotres, Oint. (Central Pharmacal)

W/Neomycin HCl, tyrothricin, neomycin sulf., benzocaine. See: Pentasol, Trache. (Merck Sharp & Dahme)

ZINC BORATE W/Ephedrine. (Hille) Zinc borate 0.2%, ephedrine HCl 0.3%. Bot. 1 R. oz., 4 R. oz., 12 R. oz., 32 R. oz.

Use: Antiseptic.

ZINC CAPRYLATE, W/Sod. caprylate. See: Naprylate, Preps. (Strassenburgh)

W/Zinc propionate and tyrothricin. See: Gyna-Plex, Vog. Cr. (Rowell)

ZINCPRIN, (Alcon Labs.) Zinc sulfate $\frac{1}{4}$ %, phenylephrine HCl 0.12%, benzalkonium chloride 1:25,000 in buffered sol. Bot. 15 cc. & Drop-tainer 15 cc.

Use: Conjunctivitis & related infections.

ZINC GELATIN, U.S.P. Zinc gelatin boot.

ZINCHCHLORUNDESAL, N.N.R. 1955. * Zinchundesal. See: Salundek, New, Oint. (Wallace & Tiernan)

ZINC I SOL, (Ophthalmos Inc) Zinc sulf. $\frac{1}{4}$ % buffered sol. Bot. dropper 15 cc.

ZINC INSULIN, N.N.D. 1962. See: Insulin

Zinc, Prep. (Various Mfr.)

ZINCOFAX, (Burroughs Wellcome) Zinc oxide 15% in lanolin base. Oint., Tube 1 $\frac{1}{2}$ oz.

Use: Antiseptic & protective.

ZINC OXIDE, A.D.R. Flowers of zinc. (Merck Sharp & Dahme) Oint., Tube 1 oz. Jar 1 lb., Paste, Lassars plain zinc paste, Jar 1 lb.

See: Calamine Preps.

Use: Mild astringent & antiseptic in skin diseases.

W/Anapulgite activated, colloidal sulfur, resorcinol, Isopropanol. See: Sebasorb Lotion. (Summer Labs.)

W/Calamine, camphor, etc. See: Calomatum, Oint. (Toilby-Nason)

W/Cod liver oil, starch, talc, lanolin, alkyl-dimethylbenzyl ammonium chloride. See: Baby Care, Cream. (Invenax)

See: Baby Care, Cream. (Invenax)

* Generic name recognized by N.F.-U.S.P. or A.M.A.

W/Hexachlorophene, phenol, resorcinol, oil of tar. See: Retafen. (VanPelt & Brown)

W/Phenol, alum. See: Azinol, Oint. (Schief-felin)

ZINC OXIDE OINTMENT. See: Zincolax, Oint. (Burroughs Wellcome).

* **ZINC PEROXIDE**, U.S.P., N.N.R. 1955, A.O.R. (Mallinckrodt) Zinc superoxide. Bot. 25.3 Gm, 113.4 Gm., 454 Gm.

Use: Antiseptic, astringent, deodorant for wounds & skin diseases.

* **ZINC PHENOLSULFONATE**, N.F. W/Mag. trisulfate, col. carb., bis. subcarb., cerium oxalate. See: Bismosalite Comp., liq. (Blue Line)

W/Phenyl salicylate, bis. subcarb., col. carb., cerium oxalate. See: Bismosal, liq. (Blue Line)

W/Neomycin, koolin, pectin, salol. See: Neo-Kofetan. (Buffington)

ZINC PROPIONATE W/Zinc caprylate and tyrothricin. See: Gyna-Plex, Vog. Cr. (Rowell)

ZINC SULFANILATE. See: Nizix (Burroughs Wellcome)

ZINC SULFATE.

See: Bonoxin (Crookes-Barnes)

Eufapte Zinc Sulfate (Professional Pharm.)

Op-Thei-Zin Ophth. (Alcon)

Zinc I Sol (Ophthalmos Inc)

W/Methapyrilene HCl, phenylephrine HCl, boric acid, benzalkonium chloride. See: Optophed, Sol. (Lemmon)

W/Phenylephrine. See: Phenylzin (Iso-Sol)

W/Piperacaine HCl & phenylephrine HCl

Use: M-Z Solution (Crookes-Barnes)

W/Zinc chloride, pot. alum, boric acid. See: Astringent Comp., Pow. (Upjohn)

ZINC SULFIDE, LOTION. See: White Lotion, U.S.P.

ZINC SULFIDE COMPOUND LOTION.

(Upjohn) Zinc, sulfur as sulfide, polysulfides & thiosulfate, Sod. borate, boric acid, Sod. sulfate, Pot. sulfate, aluminum (hydroxide)

diethylene glycol. Bot. 4 oz.

Use: Antiseptic

ZINC SULFOCABBOLATE, W/Paregoric, pepsin, bismuth subsalicylate, salol & methyl salicylate. See: Parepaptal, liq. (Chicago Pharm.)

ZINCUNDECATE, N.N.R. 1955

See: Desenex (Wallace-Tiernan)

Undesol (Veltex)

* **ZINCUNDESAL**. See: Zinchloriendesal

* **ZINC UNDECYLENATE**, N.F., N.N.R. 1954. (Promo) Pow., Pkg. 25 lb.

W/Copper undecylenate & undecylenic acid

See: Undesilin, Oint. (Cutter)

W/Copper undecylenate, undecylenic ac²

- & phenylmercuric acetate See: Di Spar, Oint (Wintersmith)
- W/Hexachlorophene, undecylenic acid See: Undex Improved (VanPelt & Brown)
- W/Salicylanilide, 5 chloro-salicylanilide, 3,3' and 5,4' dichlorosalicylanilides & oedecylenic acid See: Salundek New, Oint (Wallace & Tiernan)
- W/Undecylenic acid
See: Deseex Paw (Wallace & Tiernan).
Epiderma Dex, Oint (Sulliff & Case)
Undex Paw (VanPelt & Brown)
- W/Undecylenic acid, hexachlorocyclohexene isomer & benzyl ammonium chloride See: Hexacid, Oint (Central Pharm)
- W/Undecylenic acid, menthol & benzocaine See: Deryl Cream (Paul Maney)
- ZINEFRIN** (Alcoa) Zinc sulfate 0.25% phenylephrine HCl 0.12% Dropper plastic bot 15 cc
Use: Astringent and decongestant
- ZIN-EMUL No. 1** (Texas Pharm Co) A non oily starch and zinc oxide lotion Bot 1 pt
- ZIN-EMUL No. 2** (Texas Pharm Co) A mineral oil lotion of starch, zinc oxide and zinc carbonate Bot 1 pt
- ZIRADRYL CREAM**, (Parke, Davis) Benadryl HCl 2% & zirconium oxide 4% in water miscible base Tube 1 oz
Use: Poison ivy
- ZIRADRYL LOTION** (Parke Davis) Benadryl HCl 1% zirconium oxide 2% Bot 6 oz Plastic Sq Bot 80 cc
Use: Poison ivy and related dermatitis
- ZIRAMEL** (Bruce) Zirconium carbonate 2%, hexachlorophene 1%, benzocaine 1%, thymol 0.7% menthol 0.7%, camphor 0.3%, isopropyl alcohol 8%, w/magnesium aluminum silicate and preservatives Cream, Plastic Bot 4 oz
Use: Topically, treatment of fungus infections
- ZIRCAL CREAM** (Schleffelin) Not listed Zirconium oxide 4% calamine 13% benzocaine 1%, menthol 0.2% merbak 0.1%, w/propylene glycol bentonite & water Cream Tube 1 oz
Use: Rhus dermatitis & aesthetic
- ZIRCONIUM CARBONATE OR OXIDE W/ Allantoin, trypsin, ethyl amleobenzoate, benzalkonium chloride Amerchol 9** See: Trypsin gel Creme (Bruce)
- W/Benadryl HCl See: Ziradryl, Cream & Lotion (Parke, Davis)
- W/Benzocaine See: Puragel (Bruce)
- W/Benzocaine menthol, camphor See: Rhuff cream (Lederle)
- W/Benzocaine menthol, camphor, calamine & Isopropyl alcohol
See: Rhulispay Aerosol (Lederle)
Rhuff Lotion (Lederle)
- W/Benzocaine, menthol, merbak, calamine & bentonite See: Zircal, Cream (Schleffelin)
- W/Bithioeol, tyrothricin, neomycin sulfate, phenyltoloxamine dihydrochloride zirconium oxide, diprodon HCl See: Allergetic, Lot (Bristol)
- W/Bristamine See: Bristamine Lotion W/Zircosam (Bristol)
- W/Calamine, benzocaine See: Rhuffoam (Lederle)
- W/Phenyltoloxamine dihydrochloride See: Zirnox, Lot (Bristol)
- W/Phenyltoloxamine dihydrochloride, calamine, camphor, phenyl, benzocaine See: Histonium, Sal (Columbus)
- W/Phenyltoloxamine dihydrochloride citrate oxyquinoline sulfate See: Ivy-Eze, liq (Lloyd, Dabney & Westerfield)
- W/Polythionates See: Dermasulf, Preps. (Smith Miller & Patch)
- W/Pyribenzamine See: Pyribenzamine w/ Zircosam, Cream (Ciba)
- W/Pyribenzamine hydrochloride See: Anliry, Lot (Ciba)
- W/Pyrimine maleate, benzocaine menthol & camphor See: Zirmolate, Cream (Jenkins Labs)
- W/Pyrimine maleate, p-Chloro-m-xyleneol & benzalkonium chloride See: Rhucen, Oint (Central Pharm)
- W/Sodium thiosulfate glycerin, isopropyl alcohol, benzocaine, and colloidal magnesium aluminum silicate See: Dermogel, Creme (Bruce)
- ZIRMOLATR** (Jenkins Labs) Zirconium oxide 3.5%, pyrimine maleate 1.5%, benzocaine 1%, menthol 0.7%, camphor 0.3% Cream Tube 1 oz
Use: Topically relief of itching and discomfort of ivy and oak poisoning insect bites & other minor skin irritations
- ZIRNOX LOTION** (Bristol) Phenyltoloxamine dihydrochloride citrate 1%, zirconium oxide 4% Bot 4 oz, 6 oz
Use: Topically; poison ivy dermatitis
- ZOLAMINE** (White Lab) N,N Dimethyl N' 2 thiazal N' p methoxybenzyl ethylene diamine HCl
Use: Aethiztamine and local anesthetic
- ZOLGER**, (Superior Pharm) Peetylenesterazol 100 mg nicotinic acid 50 mg, thiomine HCl 10 mg /Tab Bot 100s, 500s
Use: Respiratory and circulatory stimulation
- ZOLYSE** (Alcon) Lyophilized alpha chymotrypsin 750 u, balanced salt solution 10 cc./unit pack
Use: Cataract surgery

- ZONAZIDE.** (G. F. Harvey) Isoniazidic acid hydroxide. Vial (100 mg./ss.) 10 cc.
Use: I.M.; tuberculosis.
- ZONITE.** (Chemway Corp.) A stabilized solution of sodium hypochlorite. Approx. 1% NaClO.
Use: External antiseptic
- ZONIUM CHLORIDE.** (Lanett) Benzalkonium chloride 12.8%, Bot. 4 oz., pt., gal. Also as 1:1,000, Bot. 1 pt., 1 gal.
Use: Antiseptic.
- ZORANE.** (Lederle) Metaxalone.
- ZOXAZOLAMINE, N.F., N.N.D.** 1962. 2-Amino-5-chlorobenzoxazole. Withdrawn.
- ZYLASE TABS.** (Bryant) Amylolytic enzyme 7.5 mg., proteolytic enzyme 4 mg., cellulolytic enzyme 0.75 mg., calcium carbonate 300 mg., magnesium glycinate 50 mg./tab Bot. 30s (toiled), 100s.
Use: Relief of dyspepsia.
- ZYLATE.** (Upjohn) Benzyl benzoate 36%, Isopropyl alcohol 52%, inert ingredients 12%. Bot. 4 oz., 1 pt., 1 gal. Also Emulsion (30%) 1 pt., 1 gal.
Use: Treatment of scabies & pediculosis.
- ZYLAX.** (Glidden) Diacetylhydroxyphenylsolin (acaphenololatin) 5 mg., brewer's yeast dried 160 mg., sod. carboxymethylcellulose 300 mg./tab, Tins 12s.
Use: Orally, laxative.
- ZYLCAINE.** (Abbott) Procaine 75 mg., butesin 0.3 Gm., benzyl-alcohol in peanut oil/5 cc. Amp. 5 cc., 6s, 25s, 100s; 100 cc., 1s.
Use: Local anesthetic for hemorrhoids.
- ZYLOFURAMINE HCl.** (Winthrop Lab.) d-threo- α -benzyl-N-ethyltetrahydrofurfurylamine HCl.
Use: Antidepressant.
- ZYLOMETAZOLINE.** See: Otrivin, Sofn. (Ciba)
- ZYMABASIC.** (Upjohn) Vit. A 5000 u., D 1000 u., C 50 mg., B-6 1 mg./0.6 cc. Bot. 15 ss., 30 cc., 50 ss. w/saturated drapper
Use: Dietary supplement for infants
- ZYMACAP.** (Upjohn) Vit. A 12,500 u., D 1000 u., thiamine HCl 5 mg., riboflavin 5 mg., pyridoxine HCl 2 mg., Cal. pantothenate 10 mg., nicotinamide 30 mg., ascorbic acid 100 mg., folic acid 0.5 mg.
*Generic name recognized by N.F., U.S.F. or A.M.A.
- B-12 4 mg./Cap. Bot. 24s, 100s, 250s. Also avail. w/folic acid 2.5 mg./Cap.
Use: Vitamin deficiencies.
- ZYMAPROPS.** (Upjohn) Vit. A 5000 u., D 1000 u., B-1 1 mg., B-2 1 mg., B-6 1 mg., Cal. pantothenate 3 mg., nicotinamide 10 mg., ascorbic acid 60 mg./0.5 cc. Bot. w/dropper, 15 cc., 30 cc., 60 cc.
Use: Multivitamin supplement.
- ZYMAFOLIC.** (Upjohn) Vit. A 8500 u., D 850 u., folic acid 3.33 mg., ferrous sulfate 150 mg., liver cons. 330 mg., thiamine HCl 3.33 mg., riboflavin 3.33 mg., pyridoxine HCl 1.5 mg., Cal. pantothenate 10 mg., nicotinamide 30 mg., ascorbic acid 30 mg./Cap. Bot. 30s, 100s.
Use: Multiple vitamin therapy.
- ZYMALIXIR.** (Upjohn) Ferrous gluconate 130 mg., liver cons. 65 mg., Vit. B-1 1 mg., B-2 1 mg., nicotinamide 8 mg., Vit. B-6 0.5 mg., folic acid 1 mg., Vit. B-12 2 mcg./5 cc. Bot. 12 oz., 1 gal.
Use: Hematonic.
- ZYMABYRUP.** (Upjohn) Vit. A 5000 u., D 1000 u., B-1 1 mg., B-2 1 mg., B-6 0.5 mg., pantothenyl alcohol 3 mg., nicotinamide 10 mg., Vit. C 60 mg., folic acid 0.25 mg., Vit. B-12 3 mcg./5 cc. Syr., Bot. 3 oz.
Use: Dietary supplement.
- ZYMATINIC DROPS.** (Upjohn) liver conc. 65 mg., ferrous gluconate 1 mg., riboflavin 1 mg., thiamine HCl 1 mg., Vit. B-6 0.5 mg., nicotinamide 8 mg., folic acid 1 mg., Vit. B-12 1 mcg./cc. Bot. w/dropper, 30 cc., 60 cc.
Use: Anemias.
- ZYMELOSE.** (Glidden) Dried brewer's yeast 160 mg., Sod. carboxymethylcellulose 0.5 Gm./Tab. Bot. 84s, 200s. Granule, Bot. 70 Gm.
Use: (1/2 Tsp. or 2 Tab. w/water) Constipation.
- ZYMENOL.** (Glidden) Brewer's yeast 12%, mineral oil 50%, agar q.s. Bot. 8 oz., 14 oz.
Use: Bowel management.
- ZYOCEL.** (Keeb & Carnick) Methylcellulose 500 mg., thiamine HCl 2 mg./Tab. Bot. 50s, 100s, 500s, 1000s.
Use: Constipation.

PHARMACEUTICAL MANUFACTURERS

- A A D Drug Labs**
5602 Avenue N
Brooklyn 34 N Y
- A & S Pharmaceutical Co**
P O Box 1292
Birmingham Alabama
- Abbott Laboratories**
14th & Sheridan Road
North Chicago Ill
- Academy Mfg Drug Co p**
D v Sherman Drug &
Chem Co Inc
544 E 171st St
New York 57 N Y
- Adams Pharm Co**
838 Malcolm Avenue
Los Angeles 24 California
- Addison Drug Co**
403 N Addison St
Richmond 20 Virginia
- Adolph's Beauty Products Co Inc**
5509 Pearl St
Philadelphia 38 Pennsylvania
- Aeplest Corp**
420 Delaware Avenue
Dayton 3 Ohio
- Aetna Drug Co**
54 Madison Avenue
Jersey City 4 N J
- A G C Labs Corp**
P O Box 62
Middletown Connecticut
- Alan Insulated Mfg Co Inc**
1197 McCortner Highway
Newark 4 N J
- Albmarle Pharm Co**
130 E 35th St
New York 16 N Y
- Alcon Labs Inc**
P O Box 1358
Fort Worth 3 Texas
- Algo Corp**
3500 Lakeside Avenue
Richmond 28 Virginia
- Alkaloid Co**
141 Worthington St
Taunton Mass
- Allergon Pharmaceuticals**
1000 S Grand Ave
Santa Ana Calif
- Allied Biochemical Labs Inc**
925 Clay Street
San Francisco 11 California
- Allison Labs**
4543 Reisterstown Road
Baltimore 15 Maryland
- Almay Cosmetics**
28 Cooper Square
New York 3 N Y
- Alphaden Co Inc**
154 E Elm Street
Chicago 11 Illinois
- Alsa Pharmaceutical Co**
233 S Washington Avenue
Lansing 25 Michigan
- Amco Drug Products Co**
6124 Barton Rd P O Box 207
North Olmsted Ohio
- Amend Drug & Chemical Co Inc**
117 E 24th St
New York 10 N Y
- American Chemical Co**
30 20 Thomson Avenue
Long Island City 1 N Y
- American Cyanamid Co**
Surgical Products Div
20 Rockefeller Plaza
New York 20 N Y
- American Drug & Chem Co**
3723 N Racine Avenue
Chicago 13 Ill
- American Drug Co**
3713 New Hampshire Ave N W
Washington 10 D C
- American Drug Corp**
6060 Maple Ave
St Louis 12 Mo
- American Drug Industries Inc**
5510 S Perry Ave
Chicago 21 Ill
- American Drug Labs**
1294 Florida Street
Memphis 6 Tenn
- American Drug Products**
249 45 Lehigh Turnpike
Somerset 26 L I N Y
- American Drug & Chem Co**
213 Commerce Bldg
4th & Webster St
St Paul 1 Minn
- American Felsol Co**
200 1/2 North St
Lorain Ohio
- American Parental**
Div Benson Labs Inc
1450 Broadway
New York 18 N Y
- American Hospital Supply Corp**
Parenteral Products Div
Evanston Ill
- American Pharmaceutical Co**
120 Bruckner Blvd
New York 54 N Y
- Amerit Drug Co Inc**
112 E 19th St
New York 2 N Y
- Ames Co Inc**
819 McNaughton Ave
Elkhart Indiana
- Amfra Great Inc**
824 Rogers Avenue
Brooklyn 26 N Y
- Amnat Products Co**
1200 E Chicago Ave
Naperville Ill
- Anahist Co Inc**
21 Gray Oak Ave
Yonkers 2 N Y
- Anatex Co**
322 N Broadway
Santa Monica California
- Anglo French Drug Co (U S A) L**
75 Varick St
New York 13 N Y
- Approved Pharm Corp**
114 G Road St
Syracuse 2 N Y
- Arco Pharmaceuticals Inc**
111 05 Bockaway Blvd
South Ozone Park 20 N Y
- Arcum Pharm Corp**
1204 Wascott Ave N W
Washington 7 D C
- Artice Drug Co**
P O Box 451
Arlington 7 Va
- Ardan Pharmaceutical Co**
4504 S E Milwaukee Ave
Portland 2 Oregon
- Ar Ee Products Co**
1036 Van Buren St
Dept DT 60
Chicago 7 Ill
- Arlington Funk Labs**
Div of U S Vitamin &
Pharm Corp
250 E 43rd St
New York 17 N Y
- Arlington Labs Inc**
229 Cleveland Ave
Harrison N J
- Armour Pharm Co**
P O Box 311
Kankakee Ill
- Arner Stone Labs Inc**
225 E Prospect Ave
Mount Prospect Ill
- Arthritis Inc**
31 E Third St
Mil Varnon N Y
- Arwood**
Div Eiklan Inc
Boyle No 22
Seminole N J
- Arzel Chemical Co**
46 S Franklin St
Newark 10 N Y
- Ascher S F & Co Inc**
3100 E 59th St
Kansas City 30 Missouri
- Paul O Ashlan Pharmacist**
2724 Mahoning Ave
Yeaumertown 8 Ohio
- Associated Concentrates**
Div American Lection Co Inc
32 30 61st St
Woodside 77 L I N Y
- Associated Drug Co Inc**
P O Box 2712
Montgomery 3 Alabama
- Associated Products Co**
77 P St N E
Washington 2 D C

- Astra Pharm. Products, Inc.
7 Neponset St.
Worcester 6, Mass.
- Atlantic Sales Corp.
1 Merford St.
Rochester 9, N. Y.
- Allor Pharm. Labr., Inc.
13211 Canani Ave.
Detroit 2, Mich.
- Avenss Corp.
250 W. 57th St.
New York 19, N. Y.
- Ayert Labs.
Div. American Home Prod. Corp.
655 Third Ave.
New York 17, N. Y.
- Barnes-Hmd Labr., Inc.
895 Kiler Rd.
Sunnyvale, California
- Berrie Drug Co., The
4111 W. Belvedere Avenue
Baltimore 15, Md.
- Berrows Biochemical Prod. Corp.
203 Prospect St.
Inwood 96, L. I., N. Y.
- Berry Labs, Inc.
9100 Kercheval
Detroit 14, Michigan
- Berry Martin & Co.
4521 Pierre de Luan Blvd
Coral Gables 46, Fla.
- Baker Pharmaceutical Corp.
444 60th St.
West New York, N. J.
- Balle & Co.
4026 Olive St.
St. Louis 8, Mo.
- Baybank Drug Co., Inc.
84-40 101st St.
Richmond Hill 18, N. Y.
- Baxler, Dan, Inc.
101 S Grandview Avenue
Glendale 1, California
- Beach Pharmaceuticals
3401—85 MarDill Avenue
Innoc 11, Fla.
- Beard, Glynn A.
3064 S. Ash St.
Denver 22, Colorado
- Becton, Dickinson & Co.
Rutherford, N. J.
- Bell Pharmaceutical Co.
2503 Amherst Street
Lubbock, Texas
- Bell Corp., Inc.
270 Lafayette St.
New York 12, N. Y.
- Bestel Drug Corp.
1817 Mentor Avenue
Norwood 12, Ohio
- Bentex Pharm. Co.
3408 Caroline Street
Houston 4, Texas
- Benton Labs., Int.
34 South York Rd.
Hatboro, Pa.
- Berkeley Labs, Inc.
1148 Valley Road
Stirling, N. J.
- Bernhoff Labs, Inc.
P. O. Box 326
Bremerton, Washington
- Beeftich, Inc.
7006 N. Western Ave.
Chicago 45, Ill.
- Bilheber Knoll Corp.
Crane St.
Orange, N. J.
- Biochemical Research Labs, Inc.
1 East Walton Place
Chicago 11, Ill.
- Bio-Pharmaceuticals, Inc.
P. O. Box 1231
Warrenton, Oregon
- Bio-Product Research Labs
2526 N. Tyndall
P. O. Box 4547
Tucson, Arizona
- Bo-Rama Drug Co., Inc.
325 W. Baltimore St.
Baltimore 1, Maryland
- Biogenic Labs.
P. O. Box 194
East Paterson, N. J.
- Blockman, Stanley, Labs., Inc.
Wesley Street
South Hackensack, N. J.
- Blaine Co.
2700 Dixie Highway
South Fort Mitchell, Ky
- Blue Line Chemical Co., The
202 S. Broadway
St. Louis 2, Mo.
- Bonar Drugs, Inc.
4850 Madison St.
Chicago 46, Ill.
- Borhardt Co.
217 N. Westcott Ave.
Chicago 12, Ill.
- Borden Co., The Chemical Div.
350 Madison Avenue
New York 17, N. Y.
- Bowman Bros. Drug Co., The
965 Cleveland Ave., N. W.
Canton 2, Ohio
- Bayer Pharmaceutical Co.
5213 N. Hampshire
Ft. Worth 3, Texas
- Boyle & Co.
4855 E. Gage Ave.
Bull Garden, California
- Brayton Pharm. Co.
171 S. W. 38th St.
Cottonwood, Ariz.
- Brown, George A. & Co.
1450 Broadway
New York 18, New York
- Brewer & Co.
67 Union Street
Worcester, Mass.
- Brewer Pharmacal Co., Inc.
9140 West Chester Pike
Upper Darby, Pa.
- Britol Labs, Inc.
Div. Britol-Myers Co
630 Fifth Avenue
New York 20, New York
- British Drug Houses
(Canada) Ltd.
Toronto, Canada
- Brommel Pharmaceuticals
1235 Suller St.
San Francisco 9, California
- Brown Pharmaceutical Co., The
2500 W. 6th St
Los Angeles 57, California
- Bruce Pharmaceuticals
404 Park Avenue
Swarthmore, Pa
- Bryant Pharm. Corp.
70 S. MacQuarrie Parkway
Mt. Vernon, N. Y.
- Buflington's Int.
8 Sudbury St.
Worcester 8, Mass.
- Burnham E. Labs., Inc.
431 S. Dearborn St.
Chicago 5, Ill.
- Burrage Waller & Co.
(U.S.A.) Inc.
1 Scarsdale Road
Teterboro 7, N. Y.
- Burlin, Robert & Co.
1815 U St., N. W.
Washington 9, D. C.
- C. P. I. Lab.
554 E. 14th St.
New York 34, N. Y.
- C & M Pharmacal Inc
814 S. Seven Mile Rd.
Detroit 2, Mich.
- Caldwell & Bost Co., The
80 W. Third St.
Montfield, Ohio
- California Spray Chem. Corp.
Lector St. and Ortho Way
Richmond, California
- Calvert Chem. Corp.
234 N. E. 99th St.
P. O. Box 53 37
Miami Shores 33, Fla.
- Cambridge Co.
2561 N. Clark St.
Chicago 18, Ill.
- Campbell Pharmacal Co.
125 Berkley
Dearborn, Mich.
- Campbell Pharmaceutical, Inc.
121 E. 24th St.
New York 10, N. Y.
- C. R. Canfield & Co.
2739 Lyndale Avenue S.
Minneapolis 8, Minn.
- Canlight Corp.
Ginndale, Calif
- Carburelphal Co.
2917 Swiss Avenue
Dallas 4, Texas
- Carlton Corp., The
45 E. 17th St.

- Carnation Co
5045 Wilshire Blvd
Los Angeles 36, California
- Carrtens Labs, Inc
4936 Veterans Memorial Hwy
New Orleans, La
- Cavendish Pharmaceutical Corp
26 Cordier St
Irvington 11 N J
- Cenci H R Pharmacol Co
152 N Broadway
Fresno 14, California
- Central Pharm Co, The
120 128 E Third St
Seymour Ind
- Chal Yan Corp
45 Pine St
New York 5 N Y
- Chatham Pharmaceuticals, Inc
901 Broad St
Newark 2 N J
- Chemra Labs, Inc
7250 N E Fourth Avenue
Miami 38 Fla
- Chemo Pure Manufacturing Corp
150 Duramus Avenue
Newark 5 N J
- Chemway Corp
Household Prods Div
Fairfield Road
Wayne N J
- Chesebrough Ponds Inc
485 Lexington Ave
New York 17 N Y
- Chicago Pharmacol Co
5547 N Ravenswood Ave
Chicago 40, Ill
- Children's Hospital Res Found
Washington, D C
- Christine, Vincent & Co, Inc
401 Tray Ave
Brooklyn 13, N Y
- Church Chemical Co
75 B Werker St
Chicago 1, Ill
- Ciba Pharm Products Inc
Summit N J
- Clapp Otis & Sen, Inc
439 Bayliffe St
Boston 16, Mass
- Clerk B Clerk
Wenonah N J
- Clyde Labs, Inc.
1533 W Bone St
Oklahoma City 1, Okla
- Cole Chem Co
3727 Laclade Ave
St Louis 8 Mo
- Columbia Medical Co
34 East 19th St
New York 3, N Y
- Columbus Pharmacol Co, The
330 Oak St (Dir Phillips Boxes)
Columbus 18, Ohio
- Commercial Solvents Corp
260 Madison Ave
New York 16, N Y
- Consolidated Midland Corp
CMC Research Div
15 Parkway
Katonah, N Y
- Contra Company
Div Scientific Industries, Inc
P O Box 331
Wall St Station
New York 5, N Y
- Cook E K & Co, Inc
8218 Wisconsin Ave
Bethesda 14, Md
- Cooper Tinsley Laboratories Inc
229 Cleveland Avenue
Hornsea, N J
- Cooper Wm, & Nephews, Inc
1909 N Clifton Ave
Chicago 14 Ill
- Cost Labs Inc
6008 Wender Drive
Pl Wertz 13, Texas
- Crawley Pharmaceutical Inc
45 Southbridge St
Auburn, Mass
- Crane-Hall Corp
1832 14th St N W
Washington 9 D C
- Cromagol Labs, Inc
811 Fanam St
Houston 2, Texas
- Creamline Co
P O Box 1214
Atlanta 3, Ga
- Crimed Labs, Inc
205 Crescent Drive
Yakima Washington
- Crookas Barnes Labs, Inc
Fairfield Road
Wayne N J
- Crystal Pharm Co
222 S Elmwood Ave
Oak Park, Ill.
- Cross Prods, Inc
684 Broadway
New York 12, N Y
- Cutter Laboratories
4th & Parker Sts
Berkeley 10, California
- Cy Laboratories
22 Lewton St
New Rochelle, N Y
- Dallas Pharmacol Co
111 Mobil Bldg
Dallas 2, Texas
- Dela Pharm Corp
P O Box 599
Hempstead 1, N Y
- Dalton Edward Co
Div of Maud Johnson & Co
Evanville 12 Ind
- Darre Pharmacol Co, Inc
32 Broadway
New York & N Y
- Dartall
1226 South Flower Street
Los Angeles 15, California
- Darwin Labs, Inc.
4800 District Blvd
Los Angeles 38, California
- Davies, Ross & Company Ltd
22 Thayer Street
Boston 18 Mass
- Davis & Sly, Inc
2340 E Logan Street
Deratour, Ill
- Daywell Labs, Inc
46 Miller St
Fairfield, Conn
- Denab Labs, Inc
1420 E 18th Avenue
Denver 18, Colo
- Denver Chemical Mfg Co, Inc
35 Commerce Road
Stamford, Conn
- DePree Co, The
120 Central Avenue
Holland Mich
- Dermatologicals, Inc
28 Cord ar St
Irvington, N J
- Dermik Pharmaceutical Co, Inc.
150 Eileen Way
Spartan, I I, N Y
- Deseret Pharmaceutical Co, Inc
19 E Oakland Ave
Salt Lake City 15, Utah
- Derlin Chemical Co
812 Branch Ave
Providence 4, R I
- Diamond Labs
2538 S E 43rd St
Des Moines Iowa
- Distans Co
Hwy 100 at W 23rd St
Minneapolis 14, Minn
- Dix Chem Co
3954 W Pine Blvd
St Louis 8 Mo
- Ditaclo Labs, Inc
377 Geneva St
Buffalo 4, N Y
- Doak Pharmacol Co, Inc
99 Park Avenue
New York 18 N Y
- Ducre Distributing Co
P O Box 7131
Kansas City 13 Mo
- Duha Chem Corp, The
100 Varick St
New York 13, N Y
- Deicin Corp
295 Madison Avenue
New York 17, N Y
- Dole H Pineapple Co
215 Market St
San Francisco, California
- Dome Chemicals Inc
125 West End Avenue
New York 23, New York
- Domid Pharmaceuticals
442 Blackstone Avenue
Proser, California
- Domsco Inc
Box 541
Boles, Idaho
- Donley Evans & Co
4300 Oulde Avenue
St Louis 15, Mo

- Egan, Inc.
 179 Lexington Ave
 New York 17, N. Y.
- Egan Laboratories
 Dr. The Wander Co
 222 E 10th St
 Cicero 1, Mo.
- Drug Products Co., Inc.
 230 Broad St
 Newark 10, N. J.
- Drug Specialty, Inc.
 P. O. Box 220
 Winston-Salem 1, N. C.
- Drugmaster, Inc.
 1779 Brown Avenue
 St. Louis 11, Mo.
- Dumas Wilson & Co
 421 Fairview Avenue
 St. Louis 16, Mo.
- DuPont Pharmaceutical Co.
 1237 N. 9th St
 Philadelphia 22, Pa.
- Dunbar Labs
 Dr. Chas. W. Corp.
 Fairfield 64
 Warren, N. J.
- DuPont
 Wilmington, Delaware
- Durar Trade, Inc.
 646 Broadway
 New York 12, N. Y.
- Dur, S. P. & Co., Inc.
 3817 N. Third St.
 Philadelphia 20, Pa.
- Dwight, S. B. & Co.
 109 32th Street
 New York 17, N. Y.
- Eastern Research Lab., Inc.
 302 1/2 Central Ave.
 Baltimore 7, Md.
- Estra Chemical Co., Inc.
 625 W. 55th St.
 New York 19, N. Y.
- Estee Labs., Inc.
 17 East Avenue
 Norwich, N. Y.
- Egan-O-Lab. Co.
 1420 Union Ave., Belle 102
 Memphis 4, Tenn.
- Edson, S. M., Chem. Co.
 300 Park Avenue
 New York 22, N. Y.
- Edwards Pharmaceutical Co.
 1474 S. Travenant
 Memphis 14, Tenn.
- Elder, Paul R., Co.
 303 E. Mulberry St.
 Bryan, Ohio
- El Paso Chemical Co.
 512 S. Siesta Street
 El Paso, Texas
- Embur, Inc.
 4941 Tracy Street
 Kansas City 10, Mo.
- Emerson Labs.
 1407 Camp St.
 Dallas 27, Texas
- Enco Co.
 2910 Manchester St
 St. Louis 17, Mo.
- Ende Labs., Inc.
 84-23 101 St
 Fresh Pond M.C. 18, N. Y.
- Enderside Co.
 4429 Park Avenue
 Union City, N. J.
- Ensign Brand & Co.
 2043 S. Colorado Blvd.
 Pasadena, California
- Enfield Co., The
 18 Brown St
 Waterbury 2, Conn.
- Erie, Elb & Co., Inc.
 Box 32
 Cincinnati 41, Ohio
- Ethical Pharmacists of Cal
 P. O. Box 2658
 San Antonio 6, Texas
- Ethicon, Inc.
 Suite 22
 Summerville, N. J.
- Evans Co.
 2527 N. Clark St.
 Chicago 13, Ill.
- Everett Co.
 1128 E. Somerset St
 Philadelphia 34, Pa.
- Everette Pharmaceutical Lab.,
 Inc.
 1914 W. Richardson Ave
 Chicago 24, Ill.
- Fellows Medical Mfg. Co., Inc.
 1334 W. Lafayette Blvd
 Detroit 26, Michigan
- Ferrelle Labs., Inc.
 780 W. Eight Mile Rd.
 Ferrelle 20, Michigan
- Foster Co., Inc., The
 378 Fairfield Avenue
 Stamford, Conn.
- The Foster Company, Inc.
 Easton 10, N. J.
- Fidelity Pharmaceutical Co.
 331 Anthony
 Linden, N. J.
- Field Labs.
 475 S. Washington Ave.
 Bergenfield, N. J.
- Fine Products Co.
 4422 San Vicente Blvd
 Los Angeles 49, Calif.
- Finley, D. E. Co., Inc.
 7328 Dooling Avenue
 P. O. Box 627
 Condon Park, California
- First Texas Pharmaceuticals, Inc.
 1810 N. Lamar St.
 Dallas 2, Texas
- C. B. Flood Co., Inc.
 921 Commerce St.
 Lynchburg, Va.
- Flint, Eaton & Co.
 6301 Lincoln Ave
 Merion Grove, Ill.
- Forest Co.
 98 Crosby St
 New York 12, New York
- Foster Ditch Co.
 2704 S. Winchester Avenue
 Chicago 20, Illinois
- Fournier, E. & Co., Inc.
 P. O. Box 77, Collingswood Rd.
 McKeville, E. I., N. Y.
- Fore Pharm Co.
 8600 E. 33rd St
 Indianapolis, Mo.
- Fore Labs., Inc.
 331 W. Penn Ave
 Warrentonville, Pa.
- Frost Labs., Inc.
 430 Lexington St
 Auburndale 64, Mass.
- Fuller Pharmaceutical Co.
 3108 W. Lake St
 Minneapolis 18, Minn.
- Gardo Drug Co.
 377 W. Jackson St.
 Philadelphia 10, Pa.
- Galley Pharmaceuticals
 Div. of Galley Co., Inc.
 Box 430, Yankers, N. Y.
- General Vitamins Corp.
 1106 F St., N.W.
 Washington 4, D. C.
- Geber Products Co.
 445 Thos St
 Fremont, Mich.
- Geriatrics Pharm. Corp.
 43 Commonwealth Blvd
 Baltimore 10, E. I., N. Y.
- Giffels Labs
 508 Merchandise Mart
 Chicago 14, Ill.
- Givaudan Delaunoy, Inc.
 321 W. 44th St.
 New York 36, N. Y.
- Glenwood Labs., Inc.
 262 S. Washington Avenue
 Bergenfield, N. J.
- Gold Leaf Pharm. Co., Inc.
 26 Lawrence Street
 New Rochelle, N. Y.
- Goodrich Wright, Inc.
 3311 Bedford Drive
 Dallas 35, Texas
- Gordon Pharmaceuticals
 Div. Gordon Wheel Germ Co.
 441 Lexington Ave
 New York 17, N. Y.
- Gotham Lab., Corp.
 420 Lexington Avenue
 New York 17, N. Y. (New)
- Graft Corp., The
 P. O. Box 5103
 Abilene, Texas
- Grant Labs
 6020 Adeline Street
 Oakland 6, California
- Gray Pharmaceuticals
 18 E. 26th St
 New York 10, N. Y.

- Greco Pharmacol Inc
214 26 41st Avenue
Boys de 61, N Y
- Grave Lob Inc
8877 Ladue Rd
St Louis 24 Mo
- Guard on Chem Corp
38 15 30th Street
Long Island City 1, N Y
- Haack Labr, Inc
1415 S W Harbor Drive
Portland 1, Oregon
- Hoag Inc
1504 Main St
Richmond 19 Va
- Haller & Co, Inc
330 Lincoln Way East
Chambersburg 1, Pa
- Halperin A E, Co, Inc
75 Northampton St
Boston 18 Mass
- Hance Brer & White Co
12th and Hamilton Sts
Philadelphia 23, Pa
- Hanson Drug Prods
500 Church St
Fauquier Co, N Y
- Hart Drug Corp
25 N E 25th St
Miami 37, Fla
- Harvey (G F) Co, Inc The
11 E 24th St
New York 10 N Y
- Haskell Charles C & Co
Div Arner Stone Labs, Inc
9 S Horvie St
Richmond 20 Va
- Haug Drug Co
4905 N 31st St
Milwaukee 9, Wis
- Heikroff Medical Co, Inc
321 Talbot Ave
Boston 24, Mass
- Heinrich Bickensdt Corp
486 Emory Circle N E
Atlanta 7, Ga
- Herring Louis C & Co
25 E Underwood Ave
Orlando, Fla
- Herron Drug Co
P O Box 1162
Clearwater, Fla
- Heus, E W Co
1629 Worthington Avenue
St Louis 3, Mo.
- Heyden Chem Corp
342 Madison Ave
New York 17, N Y.
- Herzog Labs, Inc
438 Pratt St
Buffalo 4, N Y.
- Hille Labs, Inc
3477 N Clark St
Chicago 13, Ill
- Hirs Pharmacol Co, Inc
1221 Grey Ave
Utica New York
- Hobart Labs, Inc
900 N Franklin St
Chicago 10 Ill
- Hoffmann LaRoche, Inc
Roche Park
Nutley 10, N J
- Holland Santos Co, Inc
145 Hudson St
New York 13, N Y
- Hollister Sher Labs
2030 Wilshire Blvd
Los Angeles 57, California
Also—550 Industrial Park
Drive, Yeoman, Pa
- Homemakers Prods Div
Geo A Braas and Co
1450 Broadway
New York 18, N Y
- Horlicks Corp
2120 Northwester Ave
Eclise, Wisconsin
- Horton & Coevers
821 W Pico Blvd
Los Angeles 15, California
- Howe & Co
911 Western Avenue
Seattle 4, Washington
- Hoyt Pharmaceutical Corp
215 California Street
Newton 38 Mass
- Hemahrey Prods Co
1931 Third Ave
Detroit 30, Mich
- Huntington Labs, Inc
900 E Taton St
Huntington, Ind
- Huxley Pharm
2660 S Watson Rd
St Louis 24, Mo
- Hydrosol Co
717 Sycamore St
Cincinnati 2, Ohio
- Hylford Labs
4501 Colorado Blvd
Los Angeles 39, California
Also—160 Lockwood Ave
Yonkers, N Y
- Hynson, Warkent & Dunning, Inc
Charles & Chase Sts
Baltimore 1, Md
- Increte Products Corp
254 W 31st St
New York 1, N Y
- Ingram Pharm Co
340 Front St
San Francisco 11, California
- Inorganic Enforcements, Inc
15037 Shera Acres Drive
Cleveland 10, Ohio
- Intermedica Corp
118 Cherry Lane
Floral Park, L I, N Y
- International Pharmaceutical
Corp
415 Green St
Nantitown, Pa
- Invenex Pharmaceut cals
1163 Golden Gate Ave
San Francisco 15, California
- Irw n, Neisler & Co
P O Box 1110
Decatur, Ill
- Ives Cameron Co, Inc
685 Third Ave
New York 17, N Y
- Jackson Mitchell
Pharmaceutical Inc
10401 Virginia Avenue
Culver City, California
- Jamleson Pharmacol Co, Inc
439 Madison Avenue
New York 17, N Y.
- Jeffery Labs
13210 Oxnard St
Van Nuys, California
- Jenkins Labs, Inc
17 Wall St
Auburn, N Y
- Jergens Andrew Co
2535 Spring Grove Ave
Cincinnati 14 Ohio
- Jewell Pharmaceuticals, Inc
508 Franklin Avenue
Mt Vernon, N Y
- Johnson & Johnson
900 George St
New Brunswick, N J
- Jr League Blood Center
Milwaukee, Wis
- Kahlenberg Labs
41 N School Avenue
P O Box 1660
Sarasota, Fla
- Kasden Corp
50 Elm St
Huntington, N Y
- Keene Drug Co
202 N Delaware St
Indianapolis 4, Ind
- Kendall C B, Co
2039 Madison Avenue
Indianapolis 9, Ind
- Kent Labr
42 W 23rd St
New York 10, N Y
- Kenwood Labs, Inc
26 Court St
Brooklyn 1, N Y
- Kenyan Products
Box 284
Greenwich, Conn
- Key Corp
300 N E 59th St
Miami 37, Fla
- Kinney & Co, Inc
1013 14th St
Columbus Ind
- Kirk, C F, Labr
New York, N Y
Now—Kirk Labs, Inc
Warcenter 1, Mart
- Kirkman Pharmacol Co
1436 Elliott Ave, W
Seattle 99, Washington
- Klee Co, The
P O Box 5374
Phoenix 10, Ariz

- Bug, B. O. & Co.
 210 W. 8th St.
 Kansas City 4, Mo.
- Knoll Pharm. Co.
 377 Crane St.
 Orange, N. J.
- Kernal Inc.
 380 S. Sierra
 Reno, Nevada
- Kromers-Urban Co.
 141 W. Vase St.
 Milwaukee 1, Wis.
- Lactona Products Div
 236 E. Ninth Street
 St. Paul 1, Minn.
- Lafayette Pharm. Inc.
 522 N. Earl Ave.
 Lafayette, Ind.
- Leharde Laboratories, Inc.
 1707 E. North Ave.
 Milwaukee 1, Wis.
- Lennett Co., Inc., The
 Frankford Ave. & Allee St.
 Philadelphia 25, Pa.
- Lenteen Products
 (See Green) Labs., Inc.
- Lederle Labs.
 Div. of American Cyanamid Co.
 Pearl River, N. Y.
- Leeds Chemical Prods. Inc.
 831 Webster Ave
 Chicago 14, Ill.
- Leeming, Thomas & Co., Inc.
 185 E. 44th St.
 New York 17, N. Y.
- Lehn & Pink Peeds. Corp.
 445 Park Ave.
 New York 22, N. Y.
- Lemmon Pharmaceutical Co.
 Temple Ave
 Sellersville, Pa.
- Lemon Co., Victor
 206 N. Coast Blvd.
 Laguna Beach, California
- Lemus Pharmaceuticals
 P. O. Box 433
 Woodland Hills, California
- Lentharis Bling Men
 4401 W. North Ave.
 Chicago 39, Ill.
- Lewel Pharmaceutical Co.
 2735 N. Ashland Ave.
 Chicago 14, Ill.
- Leyden Labs.
 4802 N. 7th St
 Philadelphia 20, Pa.
- Liberty Labs
 Des Moines, Iowa
- Liens, E. C. Sales Co.
 P. O. Box 467
 Fillmore, California
- Lif O Gen Inc.
 1443 City Line Ave.
 Philadelphia 31, Pa.
- Lilly, Eli & Co
 Box 618
 Indianapolis 4, Ind.
- Lincoln Labs., Inc.
 Hickory Point Rd.
 Box 1139
 Decatur, Ill.
- Lipton, Theo J., Inc.
 Mohrhan, N. J.
- Litzell Labs.
 131 Belgrade Ave.
 Boston 32, Mass.
- I. E. Drug Co.
 4802 N. Seventh St.
 Philadelphia 20, Pa.
- Lloyd Bros., Inc.
 4527 Beading Rd.
 Cincinnati, Ohio
- Lloyd, Dabney & Westfield, Inc.
 3949 Brotherhood Bld.
 Cincinnati 9, Ohio
- Lubice DeBullio, Inc.
 1842 Broadway
 New York 23, N. Y.
- Lurie Corp., The
 3533 Easton Ave.
 St. Louis 12, Mo.
- Louisiana Pharmaceuticals
 11th Ave. at Ohio St.
 Evansville 4, Ind.
- Lowe Labs
 P. O. Box 1013
 Miami 34, Fla.
- Luke Pharm. Inc.
 Building 501 New Castle
 County Airport
 New Castle, Del
- Lynn Pharmaceuticals, Inc.
 433 S. LaBrea Avenue
 Los Angeles 24, California
- MacAllister Lab.
 9213 Wade Park Ave
 Cleveland 4, Ohio
- Mahan, Thomas J., Inc.
 75 Brayton St.
 Englewood, N. J.
- Mallard, Inc.
 3021 Wabash Ave
 Detroit 16, Mich.
- Mallinckrodt Chemical Works
 3400 N. Second St.
 St. Louis 7, Mo.
- Mallico Chem. Corp.
 100 Varick St.
 New York 13, N. Y.
- Maltbie Labs.
 Div. Wallens & Terman Inc.
 25 Main St
 Belleville 9, N. J.
- Mandelbeck, Paul J., Inc.
 6604 Newgard Ave.
 Chicago 28, Ill.
- Maney, Paul Labs., Inc.
 402 First St.
 S. E. Cedar Rapids, Iowa
- Mannheim Eye Salve Co., Inc.
 1063 Broadway Ed
 Louisville 4, Ky.
- Mann Fine Chemicals, Inc.
 736 Liberty St.
 New York 6, N. Y.
- Manolo Co
 4200 Laclede Ave.
 St. Louis 8, Mo.
- Marcello Cosmetics, Inc
 1741 North Western Ave
 Chicago 47, Illinois
- Marin Pharmaceuticals, Inc.
 P. O. Box 141
 N. Miami Beach, Fla
- Marlen Pharmacol Labs.
 Box 3 Sunnyside Station
 Long Island City 4, N. Y.
- Mason Pharm. Inc.
 5840 - 63rd St
 Sacramento, California
- Massena, S. E., Co., The
 527 Fifth St.
 Bristol, Tenn.
- Maury, H. E., Biological Co.
 4109 S. Western Avenue
 Los Angeles 47, California
- Maynard, Inc.
 Drows 2345
 Greensboro, N. C.
- McKesson & Robbins, Inc
 P. O. Box 548
 Bridgeport 1, Conn.
- McNeil Labs, Inc.
 Fort Washington, Pa.
- Mead Johnson & Co. Labs. Div. of
 Mead Johnson & Co.
 2404 W. Pennsylvania St
 Evansville 21, Ind
- Medco Co., Inc.
 3149 Calhoun St.
 P. O. Box 7083
 New Orleans 19, La
- Medtana Co
 225 Varick St.
 New York 14, N. Y.
- Medica Pharm. Corp.
 203 Big Circle
 Carroll, Ga.
- Menie Park Labs, Inc.
 P. O. Box 303
 Edison, N. J.
- Menhastam Co., The
 1340 Niagara St
 Buffalo 13, N. Y.
- Merrick Sharp & Dehne
 D. v. Merch & Co., Inc.
 West Point, Pa.
- Merrill, Wm. S., Co.
 Lakeland Station
 Cincinnati 15, Ohio
- Metabolis Products Corp
 37 Harley Street
 Cambridge, Mass
- Metro Med. Inc.
 2309 South Blvd.
 P. O. Box 2245
 Houston 2, Texas
- Metropolitan Labs, Inc.
 182 South St.
 Oyster Bay, N. Y.

- Mayer & Co
16261 Mack Avenue
Detroit 24 Mich
- Mayers Rasser Pharm Co
1515 N Washington
Dallas 4 Texas
- Mid Atlantic Pharmacol Co The
P O Box 117
Richmond 1 Va
- Miller & S Labs
Los Angeles California
New—Smith Miller & Patch Inc
902 Broadway
New York 10 N Y
- Miller Labs Inc
1627 Washington Ave
St Louis 3 Mo
- Miles Prod
Div Miles Labs Inc
1127 Myrtle St
Elkhart Ind
- Milwell Pharmacol Co
6 Seventh Ave
Brooklyn 17 N Y
- Minette Pharm Corp
P O Box 2696
Buffalo 26 N Y
- Misener Pharmaceuticals Inc
192d 36 E Meadowmere
Springfield 4 Mo
- Mason Pharmacol Co
1025 Victoria St
P O Box 1674
San Antonio 6 Texas
- Modern Products Inc
1428 N 24th St
Milwaukee 5 Wis
- Moffet Labs Inc
201 E First St
Dayton 2 Ohio
- Moore K k Labs Inc
326 Chandler St
Worcester 1 Mass
- Morton Pharm Inc
1425 N Highland St
Memphis 11 Tenn
- Mulford Collie d Labs
9 S 38th St
Philadelphia 4 Pa
- Mura Pharmacol Labs Inc
121 Liberty St
Quincy 69 Mass
- Myles Carter Labs Inc
1909 E McDowell Rd
Phoenix Arizona
- Nallick Chemical Industries Inc
9 Mayhaw St
Framington Center Mass
- National Drug Co
(D v Bichardson Marrell Inc)
4663 Stanton Ave
Philadelphia 44 Pa
- National Drug Labs Inc
700 W Division St
Chicago 10 Ill
- National Lacthin Inc
1552 W Howard St
Chicago 26 Ill
- National Research Inc
Springfield N J
- National Sanitary Laboratories
6652 N Western Ave
Chicago 45 Ill
- Nearly Laboratories
3908 Kilbourne Rd
Columbia S C
- Nelson Boyer Co
813 E Broadway
Glendale 3 Calif
- Nesco Corp
1001 N McCadden Pl
Los Angeles 38 Calif
- Neogra Laboratories
(See Warner Chilcott Labs)
- Neslie—Le Mer Co
902 Broadway
New York 10 N Y
- Neovane Products Co Inc
2103 Ferrysville Ave
Pittsburgh 14 Pa
- New England Pharmacol Co
14 Waterfield Rd
Hartford Conn
- New Medical Techniques Inc
142 Main St
Greenock Conn
- Newman Pharm Co Inc
316 W Main St
Louisville 2 Ky
- New York Pharmaceutical Co
Bedford Springs
Bedford Mass
- Nien Corp
1001 N McCadden Pl
Los Angeles 38 California
- Nordson Pharmaceutical
Laboratory
Div Cooper Tinsley Labora-
tories Inc
229 Cleveland Avenue
Harrison N J
- Noreen Inc
450 Ullcoate St
Denver 9 Colo
- Norgina Labs Inc
207 E 37th St
New York 16 N Y
- Norman Pharmacol Co
3101 Broadway
Oakland 7 California
- Normand Geo Pharmaceutical
Co Inc
314 10 S W 8th St
Miami 32 Fla
- Northfield Labs Inc
Box 51
Northfield Ill
- Northwest Chemical
Research Lab
3220 N Knapp Rd
Ottawa 4 Washington
- Norwich Pharm Co
17 Eaton Ave
Norwich N Y
- Novacol Chem Mfg Co Inc
2911 Atlantic Ave
Brooklyn 7 N Y
- Nasal Pharm Co
802 West 190th St
New York 40 N Y
- Nixsamo Chem Co
3100 Falls Cliff Rd
Baltimore 11 Md
- Noyes P J & Co
51 Main St
Lancaster N H
- N Y Quinine & Chem Works
Div S B Penick & Co
100 Church St
New York 8 N Y
- Obitrol Pharmaceuticals
Div Sener Pharm Corp
382 Schenck Ave
Brooklyn 7 N Y
- Ocycrystine Lab
41 08 149th Place
Flushing 55 N Y
- Ophthalmos Inc
D v of Deha Inc
100 Vail St
New York 13 N Y
- Organon Inc
375 Mt Pleasant Ave
West Orange N J
- Ortega Pharm Co
586 S Edgewood Ave
Jacksonville 3 Fla
- Ortho Pharm Corp
Highway 202
Raritan N J
- Oshold Hugs Inc
Box 23
Smithtown N Y
- Oveseas Pharm Co
2047 Broadway
New York 23 N Y
- Page Pharmacol Co Inc
251 Common St
Watertown 72 Mass
- Paindica Int
P O Box 3115
Columbia S C
- Pan American Pharm
1221 Cline St P O
New Orleans 25 La
- Pan Pharmaceuticals Inc.
285 Madison Ave
New York 17 N Y
- Panney/Palton Corp
269 S Dean St
Englewood N J
- Para Pharmaceutical Inc.
2440 Great St N E
Salem Oregon
- Patrick Davis & Co
Joseph Campau Ave
At the River
Detroit 32 Mich
- A J Parker Co
1238 Belmont Ave
Philadelphia 4 Pa
- Parlam Corp
266 S Dean St
Englewood N J
- Pascal Co Inc
331 Dexter Ave
Seattle 8 Wash

- Puchall Labs., Inc.
 4114 Raleigh Ave.
 Seattle 8, Wash.
- Patch, E. L. Co., The
 son-Smith, Miller & Patch, Inc.
 902 Broadway
 New York 10, N. Y.
- Paxton Enterprises
 308 5th St.
 Pecos, Texas
- Penick, S. P. & Co.
 100 Church St.
 New York 6, N. Y.
- Perantone Pharmacol Co., Inc.
 Florence, Mass.
- Perce & Cooney, Inc.
 224 S. Verdugo Rd.
 Glendale 5, California
- Peter, Strong & Co.
 413 Lexington Ave.
 New York 17, N. Y.
- Pfeiffer Labs., Inc.
 1219 Glen Rock Ave.
 Westgen, Ill.
- Pfizer Lab.
 Div. Chem. Pfizer & Co., Inc.
 235 E. 42nd St.
 New York 17, N. Y.
- Pharmacia Laboratories, Inc.
 301 Fifth Ave
 New York 17, New York
- Pharmacia, Inc.
 Gallopas Hill Road
 Kenilworth, N. J.
- Pharmacia, Inc.
 1310 W. 42nd St.
 Austin, Texas
- Pharmaceutical Labs
 1015 Grandview Ave.
 Glendale 1, California
- Philadelphia Ampebia Labs
 400 Green St.
 Philadelphia 23, Pa.
- Philips Research, Inc
 Pharmaceutical Div.
 330 Oak Street
 Columbus 14, Ohio
- Phillips Prods., Inc.
 1189 1/2 W. 4th St.
 Grand Island, Nebr.
- Physicians Prod. Co., Inc.
 2022 S. Syracuse St.
 Petersburg, Va.
- Physiological Chem. Co., Inc.
 3 Le Ferre Lane
 New Rochelle, N. Y.
- Pines Div
 Solvita Co.
 123 N. Columbia St.
 Ft. Wayne, Ind.
- Pilman-Moore Co.
 Div. of the Dow Chemical Co.
 1200 Madison Avenue
 Indianapolis 4, Ind
- Pivotal Co
 P. O. Box 3583
 Cleveland 18, Ohio
- Plassner, Paul, Co., The
 2028 13th Ave. No
 St. Petersburg 13, Fla.
- Plough, Inc.
 3022 Jackson Ave.
 Memphis 1, Tenn.
- Podiatree, Ltd
 1815 Flower St.
 Glendale, California
- Polter Drug & Chem. Corp.
 123 Medford St.
 Malden 48, Mass.
- Poythress, Wm. P. & Co., Inc.
 16 N. 22nd St.
 Richmond 17, Va.
- Premier Pharm. Corp.
 11 Niagara St., Rm. 309
 Buffalo 2, N. Y.
- Prama Labs, Inc.
 111 Laurel St.
 South Hackensack, N. J.
- Preston Drug Co.
 Box 44
 Oxford, Nebr.
- Prieler, R. E., Pharmaceuticals
 203 Chancellor Ave.
 Newark 12, N. J.
- Prichard Pharm. Prods., Inc.
 421 W. 66th Terrace
 P. I. Box 511
 Kansas City 43, Mo.
- Pro-Acal, Inc.
 2830 Seminary Ave.
 Oakland 5, California
- Progenaryl Co., Inc., The
 P. O. Box 2384
 Tulsa 3, Okla
- Professional Pharmacol Co
 300 W. Josephine St.
 P. O. Box 924
 San Antonio 12, Texas
- Purdue Brothers Co.
 135 Christopher St.
 New York 14, N. Y.
- Pyreox Corp.
 511 E. 72nd St.
 New York 21, N. Y.
- Radioactive Prods. Corp.
 49 Kisco Ave
 Mt. Kisco, N. Y.
- Roman Labs.
 27 N. Market St.
 Asheville 4, N. C.
- Rond Pharm. Co., Inc.
 333 Columbia St.
 Roseland, N. Y.
- Ross-Galen Div.
 Wallis Labs, Inc.
 Gallopas Hill Rd
 Kenilworth, N. J.
- Row Chemical Co., Inc.
 303 Fourth Ave.
 New York 10, N. Y.
- Royner Pharm. Co.
 Jasper and Willard Sts.
 Philadelphia 24, Pa.
- Sacral Labs, The
 433 Tabor Lane
 Santa Barbara, California
- Reed & Carrlich
 30 Barlight Ave.
 Kenilworth, N. J.
- Reid Labs, Inc.
 444 Gerard Ave., N. W.
 Atlanta 3, Ga
- Regue Mfg. Co., Inc.
 1193 Atlantic Ave.
 Brooklyn 14, N. Y.
- Research Supplies
 Five Station, 453 Western Ave.
 Albany 3, N. Y.
- Renell Drug Co.
 Rexall Square
 Beverly of La Cienega
 Los Angeles 54, California
- Rhinaclo Co., Inc., The
 3905 Cedar Springs Rd.
 Dallas 19, Texas
- Richard & Williams
 P. O. Box 64
 Sandston, Va.
- Richards Pharm. Co.
 P. O. Box 1129
 Santa Monica, California
- Richlyn Laboratories
 3725 Cantor Ave.
 Philadelphia 24, Pa.
- Riker Labs, Inc.
 19901 Northhoff St.
 Northridge, Calif.
- Robins, A. M. Co., Inc.
 1407 Cumming Drive
 Richmond 20, Va.
- Roche Labs. Div.
 Hoffman-La Roche, Inc.
 Roche Park
 Nutley 10, N. J.
- Rocky Mt. Pharmacol Co.
 1002 E. Indian School Rd.
 Phoenix, Arizona
- Roerich, J. R. & Co.
 Div. Chem. Pfizer & Co., Inc.
 235 E. 42nd St.
 New York 17, N. Y.
- Rorer, William H., Inc.
 500 Virginia Drive
 Ft. Washington, Pa
- Rosmas Mfg. Co.
 534 W. Adams St.
 Chicago 4, Ill.
- Roussel Corp.
 155 E. 44th St
 New York 17, N. Y.
- Rowell, E. H. Co., Inc.
 Elliott St.
 Bolivar, N. Y.
- Royce Pharmacol Co.
 10944 Ventura Blvd
 N. Hollywood, Calif.
- Ryan Labs.
 3434 S. E. Milwaukee
 Portland 2, Oregon
- Rytson Co.
 7 N. MacClellan Plwy.
 Mt. Vernon, New York
- Sahyun Labs.
 31 & Camille St
 Santa Barbara, Calif.

- Salem Pharmaceuticals
Naugatuck Conn
- Sandoz Pharmaceuticals
Div of Sandoz Chem Works, Inc.
Rowle 10 Hanover, N J
- Sanford & Son
6332 Warner Drive
Los Angeles 48 Calif
- Savage Labs Inc
P O Box 3103
Bellairs 101, Texas
- SchenLabs Pharmaceuticals Inc
See Riker Labs, Inc
- Scharing Corp
80 Orange St
Bloomfield N J
- Schleffelin & Co
28 Cooper Square
New York 3 N Y
- Schilling Labs, Inc
P O Box 503
Ojaiendole California
- Julius Schmid Inc
423 W 55th St
New York 19 N Y
- Schumann Jones Co, The
2184 B 9th St
Cleveland 15, Ohio
- Schwartz Research Labs Inc
230 Washington St
Mt Vernon, N Y
- Searle G D & Co
P O Box 5110
Chicago 80 Ill
- Sherman Labs
5031 Grandy Ave
Dearb 11 Mich
- Shield Labs
12850 Mansfield Ave
Detroit 27, Mich
- Sias Inc
710 Asher
Marshall, Ill
- Siler Med Corp
701 Welch Rd
Palo Alto, California
- Simsen Labs
5871 Jas Campau Ave
Detroit 11, Mich
- Sirell Laboratories Inc
2034 Broadway
Santo Monica California
- Smith D C, Co, The
233 S 3rd St
Grand Junction, Colo
- Smith Corral Dunham
Pharmaceutical Co
New Brunswick N J
New-Smith Miller & Patch, Inc
902 Broadway, New York 10, N Y
- Smith Dorsey
Div The Wander Co
233 S 10th St
Lincoln 1, Nebr
- Smith Kline & French Labs
1500 Spring Garden St
Philadelphia 1, Pa
- Smith Martin Co
Div Cooper, Tinsley Lab, Inc
229 Cleveland Avenue
Harrison N J
- Spanner G O, Inc
20 Church St
Mantoloking, N J
- Spartan Pharm Co
354 Mercat St
Jersey City 2, N J
- Spencer Labs, Inc
10 Pine St
Marristown N J
- Spirit & Co Inc
15 Brown St
Waterbury 2, Conn
- Squibb E B & Sons
Div Orla Matheson Chem Corp
745 Fifth Ave
New York 22, N Y
- Staff Research Labs, Inc
Div Leeds Pharmaceutical Corp
147 Frederick St
Garfield N J
- Standard Laboratories Inc
199 Tabor Rd
Morris Plains N J
- Strandex Laboratories
385 W Second Ave
Columbus Ohio
- Stanley Drug Products, Inc
934 N B 25th Ave
Portland 8 Oregon
- Starr Pharmacol Co
8822 Melrose Ave
Los Angeles 48 Calif
- Stoyner Corp
2531 Ninth St
Berkeley 10, Calif
- Stefel Labs
Oak Hill, New York
- Stiffco Labs
P O Box 991
Sarasota, Fla
- Stoddard, G S & Co
295 Lafayette St
New York 13, N. Y.
- Storch Pharmaceuticals, Inc
2328 Hampton Blvd
St Louis 10, Mo
- Strassenburgh Labs
P O Box 1710
Bethesda 3, N Y.
- Stuart Company, The
3360 E Penthill Blvd
Pasadena, California
- Stuart Chase Corp
7 Michigan Drive
Natick, Mass
- Success Chem Co, Inc
800 Hiersdale St
Brooklyn 7, N Y
- Summers Labs, Inc
Marris Rd & Wissahickon Creek
Part Washington, Pa
- Sun Lab, Inc
1036 S Orange Ave
Newark 6, N J
- Superior Pharamal Co, The
434 Maryland Ave
Dayton 4 Ohio
- Sulliff & Case Co, Inc
201 Spring St
Peoria 1, Ill
- Syntex Laboratories Inc
701 Walsh Blvd
Palo Alto Calif
- Tahlae Co
32 Union Square
New York 3, N Y
- Tahle Beck Labs, Inc
812 Hampton Ave
Greenville S C
- Taylor Mason Co
350 5th Ave
New York 1, N Y
- Tamm Inc
18 S Michigan Ave
Chicago 3, Ill
- Tannin Products
2039 Madison Ave
P O Box 543
Indianapolis 6 Ind
- Taylor Chem Co
7301 Melrose Ave
Hollywood 48, California
- Taylor Lab
Div Dumas Millner Corp
4359 Northview Dr
Jackson 12 Miss
- Tennessee Pharm Co, Inc
1655 Poplar Ave
Memphis, Tenn
- Tettager & Co Inc
1334 W Lafayette Blvd
Detroit 28 Mich
- Texas Pharmacol Co
307 E Josephine St
P O Box 1859
San Antonio 8 Texas
- Texaphyl Products
Bohling Texas
- Thompson M B
711 Fifth Ave
New York 22, N Y
- Tilden Co, The
Div Textran Pharmaceuticals, Inc
New Lebanon, N Y
- Tresh Labs, Inc
110 Chestnut St
Philadelphia 8, Pa
- Trialon Labs, Inc
218 20 98th Ave
Queens Village 9, L I, N Y
- Tyson Drug Co
3419 Broadway
Brooklyn 31, N Y.
- Tyso E & Co, Inc
924 Rogers Ave
Brooklyn 26 N Y
- Tweed Paulsen & Co, Inc.
140 E Duarte Rd
Monterey Calif
- Tracy Drug Prods, Inc.
2169 N Mission Ave
Altadena, California

- Travel Labs.
 6301 Lincoln Ave.
 Marlee Grove, Ill.
- Trilan Products
 2921 Santa Anita Ave.
 Altadena, California
- Tri-State Pharm. Co., Inc.
 131 S. N. Walker
 Oklahoma City 1, Okla.
- F. Trust Co., Inc.
 1628 Central Avenue
 St. Petersburg 12, Fla.
- Truxton, C. O. Inc.
 1146 Huddae Ave.
 Comdee 3, N. J.
- Tully Pharmaceutical Corp.
 48 Lincoln Park
 Newark 2, N. J.
- Tetas, S. J. Co., Inc.
 19180 Mi. Elliott Ave.
 Detroit 34, Mich.
- Tyler Labs.
 7207 Indianapolis Blvd.
 Hammond, Ind.
- Tyree, J. S., Chemist, Inc.
 743 15th St., N. E.
 Washington 2, D. C.
- Ulmer Pharmacal Co., The
 1400 Harmon Place
 Muncie 3, Ind.
- United Labs., Ltd.
 323 S. Fair Oaks Ave.
 Pasadena 1, California
- United Specialists
 P. O. Box 1277
 San Mateo, California
- University Laboratories, Inc.
 137 Alcazar Ave.
 Canal Quabes, Pa.
- Unjohn Co., The
 301 Henrietta St.
 Kalamazoo 99, Mich.
- U. S. Standard Product Co.
 225 E. Prospect Ave.
 Mount Prospect, Ill.
- U. S. Vitamin B Pharm. Corp.
 800 Second Ave.
 New York 17, N. Y.
- V. C. A. Labr.
 Div. Vitamins Corp. of America
 37 Emalee St.
 Newark 12, N. J.
- Vaia Chemical Co. Inc., The
 1201 Liberty St.
 Allentown, Pa.
- Valentina Co., P. O. Box 7306
 3414 W. Larch St.
 Richmond 21, Va.
- Vanguard Pharmaceutical Corp.
 410 Pamplin Ave.
 Cedar Grove, N. J.
- Vee Futtes Pharm. Co.
 3208 N. Clinton Ave.
 Chicago 13, Ill.
- VeePitt & Brown, Inc.
 1322 E. Main St.
 Richmond 4, Va.
- Veltex, Inc.
 1811 First Ave., N.
 Birmingham 3, Ala.
- Veritas Products Co., Inc.
 30 Conder St.
 Irvington 19, N. J.
- Vertal, Inc.
 4943 Manchester Ave.
 St. Louis 10, Mo.
- Viemary Co., The
 P. O. Box 84
 Orange, N. J.
- Via Bio Corp.
 224 W. Livingston
 Meadville, Ill.
- Vitamix Pharm., Inc.
 5051 Lancaster Ave.
 Philadelphia 31, Pa.
- Vriarlon Co., Inc., The
 625 W. 55th St.
 N. Y. 19, N. Y.
- Vagorell Prods. Co.
 1212 W. Washington Blvd.
 Los Angeles 7, California
- WIS Pharmaceuticals, Div.
 Wallace & Tiernon, Inc.
 331 Clay Road
 Rochester 23, N. Y.
- Walden Co., The
 715 Dexter Ave.
 Seattle 9, Washington
- Walker Labs., Inc.
 Bradford Rd.
 Mt. Vernon, N. Y.
- Wallace Labs.
 Half Acre Road
 Cranbury, N. J.
- Wallace, Geo. J., Inc.
 153 Waverly Pl.
 New York 14, N. Y.
- Wampala, Henry F. & Co., Div.
 The Denver Chem. Mfg. Co.
 25 Commerce Road
 Stamford, Conn.
- Warner-Chilcott Lab.
 Div. Warner Lambert Pharm. Co.
 201 Talar Rd.
 Morris Plains, N. J.
- Warren Pharmacal Corp.
 8317 Manchester Rd.
 St. Louis 17, Mo.
- Warren-Toed Prods. Co., The
 582 W. Goodale St.
 Columbus 15, Ohio
- Webrier, William A. Co.
 224 Earl St.
 Memphis 3, Tenn.
- Wende, Alan, Co., Inc.
 84 Willis Ave.
 Mineola, L. I., New York
- Werl-Ward, Inc.
 745 Eagle Ave.
 New York 58, N. Y.
- Werlwood Pharmaceuticals
 468 Dewitt St.
 Buffalo 13, N. Y.
- White Laboratories, Inc.
 Kenilworth, N. J.
- Whitehall Labr.
 Div. American Home
 Products Corp.
 885 Third Ave.
 New York 17, N. Y.
- Whittier Labs.
 2101 Dempster St.
 Evanston, Ill.
- Wilfred Pharm. Co., Inc.
 P. O. Box 3304
 Mt. Carmel, Conn.
- Wilren Labr.
 4221 South Western Blvd.
 Chicago 9, Illinois
- Winarsmith Chem. Co.
 649 W. Hill St.
 Louisville 8, Ky.
- Winthrop Labs.
 1450 Broadway
 New York 18, N. Y.
- Wirth Labs.
 1030 S. Main
 Rockford, Ill.
- Wally Pharmed Co., Inc.
 P. O. Box 1252
 Charlotte 1, N. C.
- Woods, C. G. Chemical Co., Inc.
 711 N. Isabel St.
 Glendale 6, California
- Wyeth Labr.
 Div. American Home Prodr. Corp.
 P. O. Box 8298
 Philadelphia 1, Pa.
- Wynll Pharmaceuticals, Int.
 91 Main St.
 Medford, N. J.
- Wyns Pharmacal Corp.
 5051 Lancaster Ave.
 Philadelphia 31, Pa.
- F. E. Yenna & Co.
 8057 Stony Island Ave.
 Chicago 17, Ill.
- Younger Babber Corp.
 145 Hudson St.
 New York 13, N. Y.
- Zammer Co., Inc., The
 221 Melton Rd.
 Oakmont, Pa.

- Salem Pharmaceuticals
Naugatuch Conn
- Sandoz Pharmaceuticals
Div of Sandoz Chem Works, Inc.
Boule 10, Hanover, N J
- Sanford & Son
6332 Warner Drive
Los Angeles 48 Calif
- Savage Labs Inc
P O Box 3103
Belleire 101, Texas
- SchenLabs Pharmaceuticals, Inc
See Siker Labs Inc
- Schering Corp
60 Orange St
Bloomfield N J
- Schleffelin & Co
28 Cooper Square
New York 3 N Y
- Schilling Labs, Inc
P O Box 303
Glendale California
- Julius Schmid Inc
423 W 55th St
New York 19, N Y
- Schuemone Jones Co, The
2134 S 9th St
Cleveland 13 Ohio
- Schwartz Research Labs, Inc
230 Washington St
HI Vernon, N Y
- Searle G D & Co
P O Box 5110
Chicago 80 Ill
- Sherman Labs
5031 Grandy Ave
Detroit 11 Mich
- Shield Labs
12850 Montefield Ave
Detroit 27 Mich
- Sig Inc
710 Archer
Marshall, Ill
- Siler Med Corp
701 Welch Rd
Falo Alto, California
- Simpson Labs
5871 Jose Campus Ave
Detroit 11, Mich
- Siral Laboratories, Inc
2034 Broadway
Santa Monica, California
- Smith D C, Co The
233 S 5th St
Grand Junction Colo
- Smith Corral Dunham
Pharmaceutical Co
New Brunswick N J
Now—Smith Miller & Petch Inc
902 Broadway New York 10 N Y
- Smith Daney
Div The Wander Co
233 S 10th St
Lincoln 1, Nebr
- Smith Kline & French Labs
1500 Spring Garden St
Philadelphia 1, Pa
- Smith Martin H, Co
Div Cooper, Trusley Lab, Inc
229 Cleveland Avenue
Morriston N J
- Spanner, G O, Inc
20 Church St
Mentclair, N J
- Seaton Pharm Co
354 Mercer St
Jersey City 2, N J
- Spencer Labs, Inc
10 Pine St
Morristown, N J
- Spirit & Co Inc
15 Brown St
Waterbury 2, Conn
- Squibb S R & Sons
Div Otto Mathieson Chem Corp
745 Fifth Ave
New York 22 N Y
- Staff Research Labs, Inc
Div Leeds Pharmaceutical Corp
147 Frederick St
Garfield, N J
- Standard Laboratories Inc
199 Tabor Rd
Morris Plains N J
- Standax Laboratories
585 W Second Ave
Columbus Ohio
- Stanley Drug Products Inc
934 N E 25th Ave
Portland 8 Oregon
- Starr Pharmacol Co
6822 Malrose Ave
Los Angeles 46 Calif
- Stoyner Corp
2531 North St
Berkeley 10, Calif
- Stefel Labs
Oak Hill, New York
- Stillec Labs
P O Box 991
Sarasota Fla
- Stoddard, O S & Co
293 Lafayette St
New York 12, N Y
- Storch Pharmaceuticals, Inc
2326 Huxton Blvd
St Louis 10, Mo
- Stresenburgh, Labs
P O Box 1710
Rochester 3 N Y
- Stearns Company The
3360 E Foothill Blvd
Pasadena California
- Stuart Chase Corp
7 Michigan Drive
North Mass
- Success Chem Co, Inc
800 Maryland St
Brooklyn 7, N Y
- Sommers Labs Inc
Mart 2 Rd & Wissahickon Creek
Fort Washington, Pa
- See Lab Inc
1036 S Orange Ave
Newark 6 N J
- Superior Pharmacol Co, The
434 Maryland Ave
Dayton 4, Ohio
- Sutcliffe & Cose Co, Inc
201 Spring St
Peoria 1, Ill
- Syntex Laboratories, Inc
701 Welch Road
Falo Alto, Calif
- Tablax Co
32 Union Square
New York 3, N Y,
- Table Rock Labs, Inc
812 Hamden Ave
Greenville, S C
- Terby Masen Co
350 5th Ave
New York 1, N Y.
- Tema Inc
12 S Michigan Ave
Chicago 3, Ill
- Texnia Products
2039 Madison Ave
P O Box 543
Indianapolis 6, Ind
- Taylor Chem Co
7301 Melrose Ave
Hollywood 46, California
- Taylor Lab
Div Dumas Milner Corp
4359 Northview Dr
Jackson 12 Miss
- Tennessee Pharm Co, Inc
1655 Penlar Ave
Memphis Tenn
- Testager & Co, Inc
1354 W Lafayette Blvd
Detroit 28 Mich
- Texas Pharmacol Co
307 E Josephine St
P O Box 1659
San Antonio 6, Texas
- Texaphyl Products
Sahling, Texas
- Thompson M S
711 Fifth Ave
New York 22, N Y.
- Tilden Co, The
Div Textren Pharmaceuticals, Inc
New Lebanon, N Y
- Torch Labs, Inc
110 Chestnut St
Philadelphia 6, Pa
- Torlon Labs, Inc
218 20 98th Ave
Queens Village 9, L I, N Y
- Toson Drug Co
3419 Broadway
Brooklyn 37, N Y
- Torre E & Co, Inc
924 Bowers Ave
Brooklyn 26, N Y
- Towne Paulsen & Co, Inc
140 E Duarte Rd
Monrovia Calif
- Tracy Drug Prods, Inc
2169 N Maranae Ave
Altadena California

- Trovoni Labs
4301 Lincoln Ave.
Marion Grove, Ill.
- Trifolt Products
2921 Santa Anita Ave.
Athens, California
- Tri State Pharm. Co., Inc.
1314 N. Walker
Oklahoma City 1, Okla.
- F. Trout Co., Inc.
1428 Central Avenue
St. Petersburg 12, Fla.
- Troster, C. O. Inc.
1146 Hodson Ave.
Camden 3, N. J.
- Tully Pharmaceuticals Corp.
49 Lincoln Park
Newark 2, N. J.
- Tuma, S. J. Co., Inc.
19180 Mt. Billitt Ave.
Detroit 34, Mich.
- Tyler Labs
7207 Indianapolis Blvd.
Hammond, Ind.
- Tyree, J. S., Chemist, Inc.
743 15th St., N. B.
Washington 2, D. C.
- Ulmer Pharmacol. Co., The
1400 Holmes Place
Minneapolis 2, Minn.
- United Labs., Ltd.
333 S. Fair Oaks Ave.
Pasadena 1, California
- United Specifier
P. O. Box 1277
San Mateo, California
- University Laboratories, Inc.
157 Alcegar Ave.
Coral Gables, Fla.
- Uolehn Co., The
301 Menzies St.
Kalamazoo 99, Mich.
- U. S. Standard Products Co.
225 E. Franklin Ave.
Mount Prospect, Ill.
- U. S. Vitamin & Pharm. Corp.
800 Seared Ave.
New York 17, N. Y.
- V. C. A. Labs
Div. Vitamin Corp. of America
37 Empire St.
Newark 12, N. J.
- Vale Chemical Co. Inc., The
1201 Liberty St.
Allentown, Pa.
- Valentia Co., P. O. Box 7300
3414 W. 14th St.
Richmond 21, Va.
- Vanguard Pharmaceuticals Corp.
410 Pamolan Ave.
Cedar Grove, N. J.
- Vee Parke Pharm. Co.
3208 N. Clifton Ave.
Chicago 12, Ill.
- Vee-Pelt & Brown, Inc.
1323 B. Main St.
Richmond 4, Va.
- Velton, Inc.
1811 First Ave., N.
Birmingham 3, Ala.
- Velton Products, Co., Inc.
30 Cardier St.
Irvington 11, N. J.
- Verbal, Inc.
4943 Manchester Ave.
St. Louis 10, Mo.
- Viamary Co., The
P. O. Box 84
Orange, N. J.
- Via Bio Corp.
274 W. Irvington
Montreal, Ill.
- Vitamin Pharm., Inc.
5051 Lancaster Ave.
Philadelphia 31, Pa.
- Vitarina Co., Inc., The
425 W. 35th St.
N. Y. 19, N. Y.
- Vasorell Prods. Co.
1212 W. Washington Blvd.
Los Angeles 7, California
- WTS Pharmaceuticals, Div.
Wallace & Tiernan, Inc.
331 Clay Road
Bechtelar 23, N. Y.
- Walden Co., The
713 Dexter Ave.
Seattle 9, Washington
- Walker Labs., Inc.
Bardard Rd
Mt. Vernon, N. Y.
- Wallace Labs.
Hull Acre Road
Cranbury, N. J.
- Wallon, Geo. J., Inc.
153 Waverly Pl.
New York 14, N. Y.
- Wampala, Henry K. & Co., Div.
The Denver Chem. Mfg. Co.
35 Commerce Road
Stamford, Conn.
- Warner Chilcott Lab.
Div. Warner Lambert Pharm. Co.
201 Tubor Rd.
Morris Plains, N. J.
- Warren Pharmacol Corp
8517 Manchester Rd
St. Louis 37, Mo
- Warren-Yeard Prods. Co., The
583 W. Goodale St.
Columbus 15, Ohio
- Webster, William A. Co.
224 East St.
Memphis 3, Tenn.
- Wende, Alon, Co., Inc.
96 Willis Ave.
Missale, I. I., New York
- West Ward, Inc.
745 Eagle Ave.
New York 56, N. Y.
- Westwood Pharmaceuticals
468 Dawitt St.
Buffalo 17, N. Y.
- White Laboratories, Inc.
Kenilworth, N. J.
- Whitehall Labs.
Div. American Home
Products Corp
485 Third Ave
New York 17, N. Y.
- Whitler Labs.
2101 Dampier St.
Evanston, Ill.
- Wilfrid Pharm. Co., Inc.
P. O. Box 5204
Mt. Carmel, Conn.
- Wilson Labs.
4221 South Western Blvd.
Chicago 9, Illinois
- Wintermilk Chem. Co.
449 W. Hill St
Louisville 8, Ky.
- Winthrop Lab.
1450 Broadway
New York 18, N. Y.
- Wirth Labs.
1030 S. Main
Rockford, Ill.
- Wolfe Pharmacol Co., Inc.
P. O. Box 1252
Charlotte 1, N. C.
- Woods, C. G. Chemical Co., Inc.
711 N. Isabel St.
Glendale 4, California
- Wyeth Labs.
Div. American Home Prods. Corp.
P. O. Box 8299
Philadelphia 1, Pa
- Wynlit Pharmaceuticals, Inc.
91 Main St.
Madison, N. J.
- Wynn Pharmacol Corp.
3051 Lancaster Ave.
Philadelphia 31, Pa.
- P. E. Young & Co.
8057 Stany Island Ave.
Chicago 17, Ill.
- Youngs Rubber Corp
145 Hudson St.
New York 13, N. Y.
- Zimmer Co., Inc., The
231 Hulton Bc.
Oakman, Pa.