

Nutritional value of orange fleshed sweet potatoes pdf

Nutritional value of orange-fleshed sweet potatoes pdf. Nutritional value of orange-fleshed sweet potatoes.

A wide variety of the roots and tubers plays a major role in human diet, animal feed, and industrial raw materials. Sweet potatoes (SPs) play an immense role in human diet and considered as second staple food in developed and underdeveloped countries.

Moreover, SP production and management need low inputs compared to the other staple crops. The color of SP flesh varied from white, yellow, purple, and orange. Scientific studies reported the diversity in SP flesh color and connection with nutritional and sensory acceptability. Among all, orange-fleshed sweet potato (OFSP) has been attracting food technologists and nutritionists due to its high content of carotenoids and pleasant sensory characteristics with color. Researchers reported the encouraging health effects of OFSP intervention into the staple food currently practicing in countries such as Uganda, Mozambique, Kenya, and Nigeria. Scientific reviews on the OFSP nutritional composition and role in vitamin A management (VAM) are hardly available in the published literature. So, this review is conducted to address the detailed nutritional composition (proximate, mineral, carotenoids, vitamins, phenolic acids, and antioxidant properties), role in vitamin A deficiency (VAD) management, and different food products that can be made from OFSP.

Keywords: antioxidants; orange-fleshed sweet potato; polyphenols; sweet potato; β-carotene. Introduction Bread is a baked product of flour or meal of cereals, especially wheat which includes ordinary, unleavened and leavened types. Bread production dates back to over 12,000 years ago (around 10000 BC) with the Egyptians as the pioneers, and was probably a deliberate experimentation with water and grain flour.2 Bread baking is a widely done practice globally with a quite diverse recipe. Baking of leavened bread must have been developed accidentally through the exposure of crushed grain to yeast cultures. It is estimated that around 1,000 BC, the Egyptians isolated yeast and used it in bread baking, helping them to produce bread to the tune of thirty varieties. 3 The spontaneous fermentation of bread was replaced with controlled process of fermentation in late 19th Century yielding increased fermentation of bread was replaced with controlled process of fermentation in late 19th Century yielding increased fermentation in late 19th Century yielding increased fermentation in late 19th Century yielding increased fermentation of bread was replaced with controlled process of fermentation in late 19th Century yielding increased fermentation in different kinds of domesticated bread production methods.5 Bread production has evolved over the ages in terms of ingredients used. Latest developments in the bread industry that have enabled the use of various composite flours including purees to produce bread with improved sensory acceptability and physico-chemical quality.6,7 Latest advances in bread baking have enabled the incorpotation of OFSP puree into bread to improve the vitamin A content of these breads.8 This had the aim of extending the value-chain of OFSP puree into bread to improve the vitamin A content of these breads.8 This had the aim of extending the value-chain of OFSP puree into bread to improve the vitamin A content of these breads.8 This had the aim of extending the value-chain of OFSP puree into bread to improve the vitamin A content of these breads.8 This had the aim of extending the value-chain of OFSP puree into bread to improve the vitamin A content of these breads.8 This had the aim of extending the value-chain of OFSP puree into bread to improve the vitamin A content of these breads.8 This had the aim of extending the value-chain of OFSP puree into bread to improve the vitamin A content of these breads.8 This had the aim of extending the value-chain of OFSP puree into bread to improve the vitamin A content of these breads.8 This had the aim of extending the value-chain of OFSP puree into bread to improve the vitamin A content of these breads.8 This had the aim of extending the value-chain of OFSP puree into bread to improve the vitamin A content of th OFSP puree bread is in practice across many countries in sub-Saharan Africa, including Kenya. However, the utilization of this β-carotene rich bread is still low as a result of limited production. Technological and marketing challenges have also occasioned the low utilization of this puree in bread baking. carotene rich OFSP-puree bread, its nutritional benefits as well as the prospects and opportunities to improve its uptake and utilization. Orange-Fleshed Sweetpotatoes (OFSP) OFSP variety of sweetpotato was developed through conventional breeding in 1995.9 The utilization of OFSP roots in the alleviation of Vitamin A deficiency (VAD) in countries like Guatemala was recorded in the early as in the mid-1990s.10 Sweetpotatoes have been promoted for their calorie-rich value, but the OFSP has been greatly recommended for the reduction of VAD in developing countries, 11 The chemical composition as well as the nutritional composition of the OFSP roots is largely influenced by the genotype and growing conditions, 12, 13 thus the variety grown must be suited to the intended purpose. Originally in SSA, great reliance was on white-fleshed or yellow-fleshed that had established markets; however, the beta-carotene content of both the white and yellow-fleshed sweetpotato roots is low as compared to the OFSP.14 Over forty cultivars of OFSP roots have been introduced in Africa.15 Some of the sweet potato varieties that have been tried and grown in Kenya include KENSPOT-3, KENSPOT-5, Kabode (NASPOT 10-0), Vitaa, Simama, Pumpkin, Japanese, Kakamega 4, Local check, NASPOT 9-0, Vindolotamu and Vitamu A.16 The varieties that have been tried and grown in Kenya include KENSPOT-5, Kabode (NASPOT 10-0), Vitaa, Simama, Pumpkin, Japanese, Kakamega 4, Local check, NASPOT 9-0, Vindolotamu and Vitamu A.16 The varieties that have been tried and grown in Kenya include KENSPOT-5, Kabode (NASPOT 10-0), Vitaa, Simama, Pumpkin, Japanese, Kakamega 4, Local check, NASPOT 9-0, Vindolotamu and Vitamu A.16 The varieties that have been tried and grown in Kenya include KENSPOT-5, Kabode (NASPOT 10-0), Vitaa, Simama, Pumpkin, Japanese, Kakamega 4, Local check, NASPOT 9-0, Vindolotamu and Vitamu A.16 The varieties that have been exploited in purce processing by the sole processor in Homabay County of Kenya, are Vitaa and Kabode.17 The β-carotene colour on the flesh, with lighter orange flesh coloured varieties having lower β-carotene levels.18,19 The nutritional composition of orange fleshed sweetpotato is as summarized in Table 1. OFSP roots can be exploited for their calorie-rich property to alleviate protein-energy malnutrition (PEM), but has majorly been recommended for fighting VAD.20 OFSP roots are rich in provitamin A carotenoids than other sweetpotato varieties including cream and white fleshed varieties. β-carotene in its form of 13-cis, all trans and 9-cis comprise 10-93% of total carotenoid in OFSP roots.9 Studies have reported a beta-carotene content ranging between 0.03 and 13.63 mg/100g fresh weight for different varieties of OFSP roots.13,21-23 A study by Islam 24 that compared the beta-carotene content of different sweetpotato varieties reported a higher beta-carotene content of 19.31-61.39μg/g in OFSP roots as compared to those of the white, yellow and cream fleshed sweetpoatoes which were 1.02, 3.28-5.64 and 3.7839μg/g respectively. Even upon cooking, the OFSP roots still provide high levels of beta-carotene.22 Mitra19 reported to those of the white, yellow and cream fleshed sweetpoatoes which were 1.02, 3.28-5.64 and 3.7839μg/g respectively. varieties of sweetpotatoes grown in India as 76.56-87.76%; providing beta-carotene content of 4060-9740µg/100g fresh weight. However, it has been reported that with increasing beta-carotene content in OFSP the dry matter content decreases.25 In another study, it was established that the acceptability of OFSP roots is not only positively influenced by appealing colour and taste, but also a higher dry matter content.26 Additionally, Bengtsson et al., 27 reported an increase of up to 46% in the in vitro bio accessibility of β-carotene for OFSP roots prepared with fat as compared to just 41% for OFSP roots without fat. This gives credence to the use of OFSP roots with lower dry matter but a higher beta-carotene content as a functional ingredient in bread baking. Table 1: Nutritional Composition of Sweet Potato Roots (per 100g Fresh Weight) Nutrient Orange-Fleshed White-Fleshed Vellow-fleshed Dry matter (g) 25.09-38.56 27.79-31.64 35.46-40.3 31.26-41.12 Protein (g) 2.03-4.19 3.35-3.40 3.61-4.27 3.20-4.09 Fat (g) 1.74-2.22 0.67-8.88 1.29-1.45 0.54-1.81 Carbohydrates (g) 14.46-22.86 15.71-20.19 27.39-33.22 22.36-38.92 Ash (g) 3.03-4.70 3.53-3.70 1.52-3.00 2.35-2.51 2.00-2.11 1.81-2.56 Energy value (kCal) 90.24-153.63 100.39-156.18 137.06-161.49 115.68-176.96 Starch (g) 7.840-16.934 7.056-8.31 14.465-26.64 17.287-32.673 Reducing sugars (mg) 335-1086 122-270 114-325.8 38-1211 Total sugars (mg) 610-24268 700-2742 163-3370 2423-68443 Adapted from Sanoussi et al., 28 Omodamiro et al., 29 and Nabubuya et al., 29 Omodamiro et al., 29 Omodami mashed (commonly known as puree) forms.31-33 OFSP flour has been used locally at domestic level and in industrial production of bakery products.

## Challenges to sweetpotato production



Lack of access to "clean" planting material.
Lack of improved varieties that meet consumer preferences.
Sweetpotato pests & diseases

 Insufficient knowledge and poor agronomic practices.
 Poor development of products and potential markets.
 Limited access to the marshlandplant in marginalized land



Some of the bakery products in which OFSP flour is incorporated as an ingredient are cakes, bread, muffins and buns.34 Mashed OFSP has also been used for flour substitution in Golden bread.35 OFSP has been used to produce products such as chips, crisps, flour, puree, juice, bread and other bakery products.37,38 In Asian countries, sweetpotato pickles and cubes are produced commercially and are known for their β-carotene rich property.39 Some of the OFSP roots such as Beta 1 and Beta 2, main varieties grown in Indonesia, are high in moisture thus are not consumed directly as roots but in derivative products40 In such cases, OFSP roots are processed and serve as functional ingredients. Production and use of OFSP puree as functional ingredients in food processing has been done for over three decades in the United States (US).33 OFSP roots have also been exploited in stock feeds as root meal especially in the raising of pigs and in other food processing such as starch extraction is limited as the dry matter content of these roots are quite low.42 OFSP puree is incorporated into bread as a functional ingredient to increase the provitamin A composition of this bread. The substitution of wheat flour with OFSP puree in bread baking has been done at varied levels in different studies.43,44 The use of OFSP to substitute wheat flour in bread is aimed at increasing the uptake and utilization of the OFSP roots. rural economies of households involved in the production of OFSP roots. Orange-Fleshed Sweet Potato Puree Based Bread OFSP roots have been exploited as functional ingredients in bread both as flours and puree.29,45 Since 1990s, OFSP had been incorporated into bread production in Peru in grated form as little sun was available for sun-drying to produce flour.46 Incorporation of OFSP in wheat products also has economic advantages as it was the case of cassava that helped reduce reliance on imported expensive wheat flour. The incorporation of either OFSP flour or OFSP flour. in bread baking presents an economic and nutritional advantage : the flour has a lower conversion rate of 4.5-5kg to produce a kilogram of flour compared to 1.3-1.6 kg of fresh root required to produce a kilogram of glour compared to 1.3-1.6 kg of fresh root required to produce a kilogram of glour compared to 1.3-1.6 kg of fresh root required to produce a kilogram of glour compared to 1.3-1.6 kg of fresh root required to produce a kilogram of glour compared to 1.3-1.6 kg of fresh root required to produce a kilogram of glour compared to 1.3-1.6 kg of fresh root required to produce a kilogram of glour compared to 1.3-1.6 kg of fresh root required to produce a kilogram of glour compared to 1.3-1.6 kg of fresh root required to produce a kilogram of glour compared to 1.3-1.6 kg of fresh root required to 1.3-1.6 kg of fresh root required to produce a kilogram of glour compared to 1.3-1.6 kg of fresh root required to 1.3-1.6 kg o reported a flour yield as low as 21.5% from some roots of OFSP varieties. This necessitates alternative economical ways such as puree or paste for bakery products. 18,40,50 Production of OFSP varieties. Figure 1. The puree is usually stored frozen for later use, thus preserving most of its chemical and physical properties. The use of OFSP puree subjected to cold-chain storage presents a challenge in terms of additional costs as the storage demands electric power and additional equipment if the long term production is to be sustained, thus limiting the use of this puree by small scale bakeries that cannot afford the cold chain storage. Moreover, it has limited the quantity of puree that can be stored due to its resource intensive nature, thus resulting into inconsistent supply of the puree occasioned with massive spoilage of roots during glut and scarce supply of roots during shortage. This has occasioned limited production of this bread. Current advances in research have enabled the production of preservative-treated shelf-life of 4 months at 230C .17 Potassium sorbate, sodium benzoate and citric acid each singly has antimicrobial effect.51,52 Synergism has been shown to increase the effectiveness of food preservatives.53 A microbial challenge study by Musyoka et al.,8 found that the combination of potassium sorbate, sodium benzoate and citric acid result in a decline of 4 log cycles for Escherichia coli and Staphylococcus aureus. However, only the cold chain stored puree has been evaluated in bread baking. Moreover, OFSP puree bread production in Kenya is only limited. Marketability of bread made from shelf-storable OFSP puree also needs to be evaluated. Production of Orange-Fleshed Sweet potato Puree Bread In developed countries and the US, nutrition awareness has served to spur demand for OFSP thus increasing production of novel foods.55 Utilization of OFSP as a fortifying agent in various foods in these countries has been a practice over time. OFSP puree has been used as an ingredient in commercial bread production in Kenya as a substitution.56 In Ghana, the bread is on the roll out for industrial production under the trade name "vitabread".31 In Rwanda, wheat flour substitution. with OFSP puree in bread baking has been done in the proportions ranging from 20 to 45%.56 Studies have documented that 30% OFSP puree incorporation into bread using OFSP puree. The use of OFSP puree as a substitute for flour has provided both economic and nutritional advantages.31 The incorporation of OFSP puree in bread targeted the high end market who could not easily access the OFSP roots and benefit from its β-carotene rich property.34 The optimal product formulation is as shown in Table 2. Decline in the gluten impacts on the physical attributes of the bread especially the loaf volume of the bread. The vital gluten and improver are added to the formulation to undo the possible effects of the wheat flour makes it virtually impossible to produce a quality bread. The baking of the OFSP puree bread follows the schematic representation showed in Figure 2. Table 2: Optimal Formulation for OFSP Puree Bread follows the schematic representation showed in Figure 2. Table 2: Optimal Formulation for OFSP Puree Bread follows the schematic representation showed in Figure 2. Table 2: Optimal Formulation for OFSP Puree Bread follows the schematic representation showed in Figure 2. Table 2: Optimal Formulation for OFSP Puree Bread follows the schematic representation showed in Figure 2. Table 2: Optimal Formulation for OFSP Puree Bread follows the schematic representation showed in Figure 2. Table 2: Optimal Formulation for OFSP Puree Bread follows the schematic representation showed in Figure 2. Table 2: Optimal Formulation for OFSP Puree Bread follows the schematic representation showed in Figure 2. Table 2: Optimal Formulation for OFSP Puree Bread follows the schematic representation showed in Figure 2. Table 2: Optimal Formulation for OFSP Puree Bread follows the schematic representation showed in Figure 2. Table 2: Optimal Formulation for OFSP Puree Bread follows the schematic representation showed in Figure 2. Table 2: Optimal Formulation for OFSP Puree Bread follows the schematic representation showed in Figure 2. Table 2: Optimal Formulation for OFSP Puree Bread follows the schematic representation showed in Figure 2. Table 2: Optimal Formulation for OFSP Puree Bread follows the schematic representation showed in Figure 2. Table 2: Optimal Formulation for OFSP Puree Bread follows the schematic representation showed in Figure 2. Table 2: Optimal Formulation for OFSP Puree Bread follows the schematic representation showed in Figure 2. Table 2: Optimal Formulation for OFSP Puree Bread follows the schematic representation showed in Figure 2. Table 2: Optimal Formulation for OFSP Puree Bread follows the schematic representation showed in Figure 2. Table 2: Optimal Formulation for OFSP Puree Bread follows the schematic representation showed in flour weight) 1% Shortening fat (based composite flour weight) 0.5% Dough Improvers (based composite flour weight) 0.5% Dough Improvers (based composite flour weight) 0.5% Dough Improvers (based composite flour weight) 18% Adapted from Wanjuu58 Another great challenge that has greatly hindered the production of OFSP pure bread is that the production OFSP roots are seasonal. Such a challenge can be overcome by the utilization of preservative treated (shelf-storable) OFSP puree.17 Bread made from shelf-stability. The adoption for use of this puree in bread production has advantages including high economic returns as there would be reduced spoilage of roots during glut, consistent supply of puree to bakeries and reduced bread production costs as the shelf-storable puree is also a better alternative for bakers who lack puree processing plants or cannot access fresh puree.17 Figure 2: Production of OFSP puree bread Click here to View figure Shelf-Stability of OFSP Puree Bread The shelf-life.59 The shelf-life of bread depends on processing, packaging, formulation and storage conditions.60 Preservation of bread seeks to control rancidity, moisture migration, development of off-flavours, crystallization, grittiness and structural weakness in bread.61 Preservation techniques including physical methods such as infrared, ultraviolet light, ultra high pressure (UHP) and microwave heating; chemical preservatives such as sodium acetate, acetic acid, potassium acetate among others; and bio-preservation as in the case of sour bread.62 Advances in research are exploiting the use of nanotechnology and packaging to improve shelf-life of bread.62 Little of this has been tested on the OFSP purce bread, but presents an opportunity of extending the keeping quality of this bread



Stability of OFSP puree bread in storage is defined by moisture content, microbial growths and staling.63 The microbiological and physico-chemical changes in bread result into firming of bread in a process known as bread staling.62 Wanjuu58 reported the shelf-life of OFSP puree bread with no added chemical preservatives to be six days as compared to that of white bread that was four days. This was attributed to the lower water activity of OFSP puree bread as the water binding capacity of the nutritional quality of the bread under refrigerated storage, however, the bread stale faster under such conditions. Spoilage and Safety of Orange-Fleshed Sweet Potato Puree Bread Baking ensures that the bacteria and mould spores which were in the flour are destroyed.58 However, some of the bacteria apores are known to survive the heat treatment and under favourable conditions grow to cause roping of the bread.64 Bacillus sp. are known to form spores which can withstand the baking process.6,65 OFSP puree based bread is subject to microbial contaminated ingredients such as flour or OFSP puree would easily cause microbial contamination of the bread.66 It is, therefore, important to ensure that the raw materials are free from mycotoxin or fungal contamination in bread. Post-thermal processing contamination is also possible.67 Proper hygiene in food handling of food processing equipment ensures limitation of microbial contamination of the product.68 This aims at improving the keeping quality of the bread. OFSP puree based bread also undergoes spoilage in bread.69 Bread staling involves modifications of the product matrix, both the macroscopic and the molecular structures. Staling affects the firmness of the crumb, deterioration of the product flavour and softening of the crust in the macroscopic structure, staling causes retro gradation and crystallinity of starch, water redistribution within the molecules of bread, decreases the amount of soluble amy lose and changes the gluten (wheat protein) network and the interactions between gluten protein and starch granules in wheat bread.63 The staling of bread is as a result of the slow change of the starch at temperatures below 550C from a moistened amorphous form to a hard less moistened crystalline form. This change causes the rapid hardening and shrinking of starch granules away from the gluten skeleton resulting into crumbliness.71 It is important to ensure proper preservation to avoid bread staling as it results into deterioration of QFSP Puree Bread Promotion of OFSP roots as a functional food in India proved a succesful strategy towards eliminating VAD while cost-effectiness is met.73 Extending such efforts to other resource poor countries has the potential of improving nutrition status of various communities. Bread made by substituting 38% of the wheat with OFSP puree was found to be high in vitamin A (135 RAE/110 g) and had a deep-yellow colour which was highly acceptable among consumers in Mozambique.37 Other studies done in Rwanda showed that there is great consumer preference of bread made of composite mixture of 30% OFSP-puree and 70% wheat-flour.74 A study by Bonsi et al.,75 found that 30% incorporation of OFSP puree into bread obtained the highest overall acceptability but texture had the lowest acceptability as compared to the white bread and other OFSP puree breads. The study also established that increasing the level of OFSP incorporated into bread resulted into firmer breads. Incorporated into bread energy and nutrients such as vitamins (pro-vitamin A) and minerals (Ca. P. Fe. Zn and K) and also add natural sweetness, color. flavor and dietary fiber.19,76 The "vitabread" has a trans β-carotene content of 1.333 mg/100 g meeting about 21% of daily requirement (1,300 μg RAE/day) of nursing mothers.44 Studies in the US have proven OFSP roots as worthy fortifying agents to help ameliorate nutritional status 77: in SSA, the bread offers that opportunity. OFSP puree also gives the bread a golden yellow colour, thus resulting in high consumer acceptability. 34 OFSP puree bread also had desirable qualities in terms of shelf-stability, moisture content, water activity, texture and microbial growths. 58 This has encouraged its wider use as compared to OFSP flour in production of bread and other bakery products. Marketability of OFSP-Puree Based Bread Commercialization of a product should be demand driven rather than a push by a "champion". Other factors can come into play in commercialization of a product should be demand driven rather than a push by a "champion". through selected retail chains.17 OFSP puree bread has been in commercial production in Japan since the 1990s where it happens to be a renowned functional food consumers were willing to pay for such products, 45 thus a proof of likelihood of success in adopting shelf-storable OFSP puree based bread by the consumers in Kenya revealed that the willingness to pay for OFSP puree based bread by the consumers in Kenya revealed that the willingness to pay for OFSP puree based bread by the consumers in Kenya revealed that the willingness to pay for OFSP puree based bread by the consumers in Kenya revealed that the willingness to pay for OFSP bread developed from shelf storable puree, with evaluation of its nutritional content, can have the market demand created through awareness creation.79 With the success of marketability of shelf-storable OFSP puree based bread, the economic gains can be realized across the OFSP puree in bread baking has been an avenue for food fortification and product diversification. OFSP puree based bread can enhance vitamin A intake among the population especially the consumption of this nutritious bread is still being witnessed due to limitations resulting from the use of cold chain stored OFSP puree. The challenge posed by the use of fresh OFSP puree in the production of bread can be overcome by the use of shelf-storable preservative treated OFSP puree. However, the marketability, physico-chemical properties, shelf-stability and sensory attributes of the bread produced from the shelf-storable puree require systematic evaluation in order to establish a substantive argument for its use. With the high consumption of bread among populations, OFSP puree based bread can be part of efforts towards improving the vitamin A status of the population. Acknowledgements The research was not funded by any organization or person but was done on scholarly basis.

References FAO. Definition and classification of commoditie: Cereals and cereal products. Published 1994. Accessed April 23, 2017. Khater E. S. G. Heat and Mass Balance for Baking Process. J Bioprocess Biotech. 2014;4(7). doi:10.4172/2155-9821.1000190. CrossRef Salem S. H., Naguib M. M., El-Sheikh H. .H, Heikal Y. A. Determination of the Microbial Profile of Compressed Baker's Yeast and Its Symbiotic Interractions. Int J Adv Res. 2015;3(2):303-312.



## Aslankoohi E., Herrera-Malaver B., Rezaei M. N., Steensels J., Courtin C.



M., Verstrepen K. J. Non-conventional yeast strains increase the aroma complexity of bread. PLoS One. 2016;11(10):1-18. doi:10.1371/journal.pone.0165126. CrossRef Gori K., Cantor M. D., Jakobsen M., Jespersen L. Production of bread, cheese and meat. Mycota Ind Appl. 2010:pp 3-27. doi:10.1007/978-3-642-11458-8\_1. CrossRef Ijah U. J. J., Auta H. S., Aduloju M. O., Aransiola S. A. Microbiological, nutritional, and sensory quality of bread produced from wheat and potato flour blends. Int J Food Sci. 2014;2014:1-6. doi:10.1155/2014/671701. CrossRef Nwosu U. L., Elochukwu C. U., Onwurah C.



O. Physical characteristics and sensory quality of bread produced from wheat/African oil bean flour blends. African J Food Sci. 2014;8(6):351-355.

doi:10.5897/AJFS2013.1079. CrossRef Musyoka J. N., Abong G. O., Mbogo D. M. et al., Effects of acidification and preservatives on microbial growth during storage of orange fleshed sweetpotato puree. Int J Food Sci Vol.

2018;2018:1-11. doi:10.1155/2018/8410747. CrossRef van Jaarsveld P. J., Marais D. W., Harmse E., Nestel P., Rodriguez-Amaya D. B. Retention of β-carotene in boiled, mashed orange-fleshed sweet potato. J Food Compos Anal. 2006;19(4):321-329. doi:10.1016/j.jfca.2004.10.007. CrossRef Talsma E. F., Melse-Boonstra A., Brouwer I. D. Acceptance and adoption of biofortified crops in low- and middle-income countries: A systematic review. Nutr Rev. 2017;75(10):798-829. doi:10.1093/nutrit/nux037. CrossRef Mahmoud A.

H., Anany A. M El. Nutritional and sensory evaluation of a complementary food formulated from rice, faba beans, sweet potato flour, and peanut oil. Food Nutr Bull. 2014;35(4):403-413.

doi:10.1177/156482651403500402. CrossRef Vizzotto M., Pereira E dos S., Vinholes J. R. et al., Physicochemical and antioxidant capacity analysis of colored sweet potato genotypes: in natura and thermally processed. Cienc Rural St Maria. 2017;47(4):1-8. doi:10.1590/0103-8478cr20151385. CrossRef Kathabwalika D. M., Chilembwe E. H. C, Mwale V. M. Evaluation of dry matter, starch and beta-carotene content in orange-fleshed sweet potato (Ipomoea batatas L.) genotypes tested in three agro-ecological zones of Malawi. African J Food Sci. 2016;10(11):320-326. doi:10.5897/AJFS2016.1471. CrossRef Bouis H., Low J., McEwan M., Tanumihardjo S. Biofortification: Evidence and lessons learned linking agriculture and nutrition. FAO WHO. 2013:1-23. Low J. W., Mwanga R. O.

M., Andrade M., Carey E., Ball A. M. Tackling vitamin A deficiency with biofortified sweetpotato in sub-Saharan Africa. Glob Food Sec. 2017;14:23-30.

doi:10.1016/j.gfs.2017.01.004. CrossRef Abong' G. O., Ndanyi V. C. M., Kaaya A. et al., A review of production, post-harvest handling and marketing of sweetpotatoes in Kenya and Uganda. Curr Res Nutr Food Sci. 2016;4(3):162-181. doi:10.12944/CRNFSJ.4.3.03. CrossRef Bocher T., Low J. W., Muoki P., Magnaghi A., Muzhingi T. From lab to life : Making storable orange-fleshed sweetpotato purée a commercial reality. Open Agric. 2017;2:148-154. doi:10.1515/opag-2017-0014. CrossRef Alam M., Rana Z., Islam S.

Comparison of the proximate composition, total carotenoids and total polyphenol content of nine orange-fleshed sweetpotato varieties grown in Bangladesh. Foods. 2016;5(3):64. doi:10.3390/foods5030064. CrossRef Mitra S. Nutrition status of orange-fleshed sweetpotatoes in alleviating vitamin A malnutrition through a food-based approach. J Nutr Food Sci. 2012;2(8):8-10. doi:10.4172/2155-9600.1000160.

CrossRef Honi B., Mukisa I. M., Mongi R.

J. Proximate composition, provitamin A retention, and shelf life of extruded orange-fleshed sweet potato and bambara groundnut-based snacks. J Food Process Preserv. 2017;42(1):1-8. doi:10.1111/jfpp.13415. CrossRef Mbusa H., Ngugi K., Olubayo F., Kivuva B., Muthomi J., Nzuve F. Agronomic performance of Kenyan orange fleshed sweet potato varieties. J Plant Stud. 2018;7(2):11. doi:10.5539/jps.v7n2p11. CrossRef Vimala B., Nambisan B., Hariprakash B. Retention of carotenoids in orange-fleshed sweet potato during processing. J Food Sci Technol. 2011;48(4):520-524.

doi:10.1007/s13197-011-0323-2. CrossRef Oloo B. O., B. M. J, Rose O. B. Effects of lactic acid fermentation on the retention of beta-carotene content in orange fleshed sweet potatoes. Int J Food Stud. 2014;3:13-33. doi:10.7455/ijfs/3.1.2014.a2. CrossRef Islam S. N, Nusrat T., Begum P., Ahsan M. Carotenoids and β-carotene in orange fleshed sweet potatoes. Int J Food Stud. 2014;3:13-33. doi:10.7455/ijfs/3.1.2014.a2. CrossRef Islam S. N, Nusrat T., Begum P., Ahsan M. Carotenoids and β-carotene in orange fleshed sweet potato: A possible solution to Vitamin A deficiency. Food Chem. 2016;199:628-631. doi:10.1016/j.foodchem.2015.12.057. CrossRef Tomlins K., Owori C., Bechoff A., Menya G., Westby A. Relationship among the carotenoid content, dry matter content and sensory attributes of sweet potato. Food Chem. 2012;131(1):14-21. doi:10.1016/j.foodchem.2011.07.072.

CrossRef Dibi K. E. B., Essis B. S., N'Zué B. et al. Participatory selection of orange-fleshed sweetpotato varieties in north and north-east Côte d'Ivoire. Open Agric. 2017;2(1):83-90. doi:10.1515/opag-2017-0009. CrossRef Bengtsson A., Alminger M. L., Svanberg U. In vitro bioaccessibility of β-carotene from heat-processed orange-fleshed sweet potato. J Agric Food Chem. 2009;57(20):9693-9898. doi:10.1021/jf901692r. CrossRef Sanoussi a F., Dansi A., Ahissou H. et al., Possibilities of sweet potato [Ipomoea batatas (L.) Lam] value chain upgrading as revealed by physico-chemical composition of ten elites landraces of Benin. AfriJ of Biotechnol. 2016;15(13):481-489. doi:10.5897/AJB2015.15107. CrossRef Omodamiro R. M., Afuape S.

O., Njoku C. J., Nwankwo I. I. M., Echendu T. N. C. Acceptability and proximate composition of some sweet potato genotypes : Implication of breeding for food security and industrial quality. Int J Biotechnol Food Sci. 2013;1(5):97-101. Nabubuya A., Namutebi A., Byaruhanga Y., Narvhus J., Wicklund T. Potential Use of Selected Sweetpotato (Ipomea batatas Lam) Varieties as Defined by Chemical and Flour Pasting Characteristics. Food Nutr Sci. 2012;889-896. doi:10.4236/fns.2012.37118. CrossRef Abidin P. E., Dery E., Amagloh F. K., Asare K., Amoaful E. F., Care E. E. Training of Trainers ' Module for Orange-Fleshed Sweetpotato (OFSP) Utilization and Processing. Tamale (Ghana): Nutrition Department of the Ghana Health Service; 2015. doi:10.4160/9789290604624. CrossRef Stathers T., Carey E., Mwanga R. et al., Everything You Ever Wanted to Know about Sweetpotato: Reaching Agents of Change Tot Training Manual.

Vol 4. Nairobi, Kenya: International Potato Centre. 2013. doi:10.4160/9789290604273.v4.

CrossRef Truong V-D., Avula R. Y. Sweetpotato purees and powders for functional food ingredients.

In: Ray RC, Tomlins KI, eds. Sweetpotato: Post-Harvest Aspects in Food. New York, USA: Nova Science Publishers, Inc. 2010:117-161. Sindi K., Kirimi L., Low J. Can biofortified orange-fleshed sweetpotato make commercially viable products and help in combatting vitamin A deficiency? In: 4th International Conference of the African Association of Agricultural Economists. Hammamet. 2013:1-17.

Jenkins M., Shanks C.

B., Houghtaling B.

Orange-fleshed sweet potato: Successes and remaining challenges of the introduction of a nutritionally superior staple crop in Mozambique. Food Nutr Bull. 2015;36(3):327-353. doi:10.1177/0379572115597397. CrossRef van Vugt D., Franke A. C. Exploring the yield gap of orange-fleshed sweet potato varieties on smallholder farmers' fields in Malawi. F Crop Res.

2018;221(36):245-256. doi:10.1016/j.fcr.2017.11.028. CrossRef Andrade M., Barker I., Cole D. et al., Unleashing the Potential of Diversity. (Barker C, ed.). Lima, Peru: International Potato Center (CIP); 2016. doi:10.1016/S0965-2590(06)70509-6. CrossRef Dooshima I. B. Quality evaluation of composite bread produced from wheat, defatted soy and banana flours.

Int J Nutr Food Sci. 2014;3(5):471. doi:10.11648/j.ijnfs.20140305.26. CrossRef Padmaja G., Jaffer T. S., Moothandassery S. S. Food uses and nutritional benefits of sweetpotato. Fruit, Veg Cereal Sci Biotechnol 6. 2012;(1):115-123. SI1)/FVCSB\_6(SI1)115-1230.pdf. Ginting E., Yulifianti R. Characteristics of noodle prepared from orange-fleshed sweet potato and domestic wheat flour. Procedia Food Sci. 2015;3:289-302. doi:10.1016/j.profoo.2015.01.032. CrossRef Chang H-SC. Future prospects for sweetpotato processing in P apua New Guinea. Australas Agribus Perspect Pap 99, ISSN 1442-6951. 2014;(December):1-17.

Nair A. G. H., Vidya P., Ambu V., Sreekumar J., Mohan C. Evaluation of orange fleshed sweet potato genotypes for storage root yield and dry matter content. Int J Appl Pure Sci Agric. 2017;3(9):82-85.

doi:10.22623/IJAPSA.2017.3103.LFVRQ. CrossRef Kariuki S., Onsare R., Mwituria J. et al., FAO/WHO project report improving food safety in meat value chains in Kenya. Food Prot Trends. 2013;33(3):172-179. Awuni V., Alhassan M. W., Amagloh F. K.

Orange-fleshed sweet potato (Ipomoea batatas) composite bread as a significant source of dietary vitamin A. Food Sci Nutr. 2017;1-6. doi:10.1002/fsn3.543. CrossRef Low J. W., Mwanga R. O. M., Andrade M., Carey E., Ball A. Tackling vitamin A de fi ciency with biofortified sweetpotato in sub-Saharan Africa. Glob Food Sec. 2017.

doi:10.1016/j.gfs.2017.01.004. CrossRef Sheikha A. F El., Ray R. C. Potential impacts of bio-processing of sweet potato: Review. Crit Rev Food Sci Nutr.

2015;(May 2015):37-41. doi:10.1080/10408398.2014.960909.

CrossRef Kamal M. S., Islam M.

N., Aziz M. G. Effect of variety and effect of sweet potato flour of two lo cal varieties on quality of breads. J Bangladesh Agril Univ. 2013;11(2):301-306.

doi:10.1007/s13197-010-0217-8. CrossRef Bonsi E., Chibuzo E., Robert Z. The preliminary study of the acceptability of Ghana bread made with orange sweet potato puree.

J Hum Nutr Food Sci. 2014;2(4):1-5. doi:10.13140/RG.2.1.2776.5523. Low J., Jaarsveld P. The potential contribution of bread buns fortified with β - Carotene - rich sweetpotato in Central Mozambique. 2008. doi:10.1177/156482650802900203. CrossRef Mwanga R. O. M., Ssemakula G. Orange-fleshed sweetpotatoes for food, health and wealth in Uganda. Int J Agric Sustain.

2011;9(1):42-49. doi:10.3763/ijas.2010.0546. CrossRef Biswas G., Das S., Nipa M. N., Patwary R. H. A comparative study for the determination of of commonly used antimicrobial against specific bacterial strains in tomato (Solanum lycopersicum L.) juice. J Glob Biosci. 2015;4(8):3094-3103.

Su L-C., Xie Z., Zhang Y., Nguyen K. T., Yang J. Study on the antimicrobial properties of citrate-based biodegradable polymers. Front Bioeng Biotechnol. 2014;23. doi:10.3389/fbioe.2014.00023.

CrossRef Baljeet S.

Y., Simmy G., Ritika Y., Roshanlal Y.

Antimicrobial activity of individual and combined extracts of selected spices against some pathogenic and food spoilage microorganisms. Int Food Res J. 2015;22(6):2594-2600.

Tedesco I., Stathers T. Sweetpotato Value Chains in Kenya : A Business Opportunity for Puree Processing and the Potential Role for Commercial Fresh Root Storage. Chatham: UK. 2015. Gruneberg W. J., Ma D., Mwanga R. O. M. et al., Advances in sweetpotato breeding from 1993 to 2012. In: Low J, Nyongesa M, Quinn S, Parker M, eds. Potato and Sweetpotato in Africa. Transforming the Value Chains for Food and Nutrition Security. Oxfordshire (UK). 2012:1-77. doi: . CrossRef Bukania C. Marketing , processing and utilization community of practice rebranding OFSP for health and wealth proceedings of cop meeting held in Kunduchi beach hotel , Dar es Salaam. In: Amagloh F, Mzamwita M, eds. Proceedings of CoP Meeting Held in Kunduchi Beach Hotel, Dar Es Salaam 14-16 March 2016 Compiled. Dar es Salaam. 2016. Nwanekezi E. C. Composite flours for baked products and possible challenges – A review.

Niger Food J. 2013;31(2):8-17.

doi:10.1016/S0189-7241(15)30071-0. CrossRef Wanjuu C., Abong G., Mbogo D., Heck S., Low J., Muzhingi T. The physiochemical properties and shelf-life of orange-fleshed sweet potato puree composite bread. Food Sci Nutr. 2018;1555-1563. doi:10.1002/fsn3.710. CrossRef Smith J. P., Daifas D. P., El-Khoury W., Koukoutsis J., El-Khoury A. Shelf life of safety concerns of bakery products – A review. Crit Rev Food Sci Nutr. 2004;44(1):19-55. doi:10.1080/10408690490263774. CrossRef Galic K., Curic D., Gabri D. Shelf life of packaged bakery goods — A review. Crit Rev Food Sci Nutr. 2009;49(5):405-426. doi:10.1080/10408390802067878. CrossRef Bhise S., Kaur A. Baking quality, sensory properties and shelf life of bread with polyols. J Food Sci Technol. 2014;51(9):2054-2061. doi:10.1007/s13197-014-1256-3. CrossRef Melini V. Strategies to extend bread and GF bread shelf-Life: From sourdough to antimicrobial active packaging and nanotechnology. Fermentation. 2018;4(1):9. doi:10.3390/fermentation4010009. CrossRef Kwaśniewska I., Rosicka-kaczmarek J., Krala L. Factors influencing quality and shelf-life of baking products. J Process Energy Agric 18.

2014;18(1):1-7. Saranraj P., Geetha M. Microbial spoilage of bakery products and its control by preservatives. Int J Pharm Biol Arch 2012. 2012;3(1):38-48. Rumeus I., Turtoi M.

Influence of sourdough use on rope spoilage of wheat bread. J Agroaliment Process Technol. 2013;19(1):94-98.

Madani H El., Taouda H., Aarab L. Evaluation of contamination of wheat and bread by fungi and mycotoxins in Fez region of Morocco. Eur J Adv Res Biol Life Sci. 2016;4(June):43-52. doi:10.13140/RG.2.1.1287.1288. Chavan R. S., Chavan S. R. Sourdough technology — A traditional way for wholesome foods : A Review. Compr Rev Food Sci Food Saf. 2011;10:170-183. doi:10.1111/j.1541-4337.2011.00148.x. CrossRef Al-Bahry S. N., Mahmoud I.

Y., Al-Musharafi S. K., Sivakumar N. Staphylococcus aureus contamination during food preparation, processing and handling. Int J Chem Eng Appl. 2014;5(5):388-392. doi:10.7763/IJCEA.2014.V5.415. CrossRef Hiroaki Y., Daijyu Y., Daiki M. et al., The staling and texture of bread made using the Yudane dough method. Food Sci Technol Res. 2014;20(5):1071-1078. doi:10.3136/fstr.20.1071. CrossRef Ho L. H., Abdul Aziz N. A., Bhat R., Azahari B. Storage studies of bread prepared by incorporation of the banana pseudo-stem flour and the composite breads containing hydrocolloids. CYTA - J Food. 2014;12(2):141-149. doi:10.1080/19476337.2013.806597. CrossRef López E. P., Pérez G. T., de Erramouspe P. L. J., Cuevas C. M. Effect of brea gum on the characteristics of wheat bread at different storage times. Food Sci Technol. 2013;33(4):745-752. CrossRef Lee M. L. W. Wheat quality and its effect on bread staling. J Agric Life Sci.

2012;46(1):153-161. Attaluri S., Ilangantileke S. Evaluation and promotion of orange-fleshed sweetpotato to alleviate Vitamin A deficiency in Orissa and Eastern Uttar Pradesh of India. In: Proceedings of the 13th ISTRC Symposium.

Arusha, Tanzania. 2007:732-736. Sindi K., Ndirigwe J., Mukantwali C., Low J., Kirimi L. What is the consumers' perception of bakery products made with vitamin A rich sweetpotato and wheat? In: 16th Triennial Symposium International Society for Tropical Root Crops Federal University of Agriculture. Abeokuta, Nigeria. 2012:2012. Bonsi E. A., Zabawa R., Mortley D. et al., Nutrient composition and consumer acceptability of bread made with orange sweet potato puree. Acta Hortic. 2016:7-14. doi:10.17660/ActaHortic.2016.1128.2. CrossRef Laelago T., Haile A., Fekadu T. Production and quality evaluation of cookies enriched with β-carotene by blending orange-fleshed sweetpotato and wheat flours for alleviation of nutritional insecurity. Int J Food Sci Nutr Eng. 2015;5(5):209-217. doi:10.5923/j.food.20150505.05. Burri B.

J. Evaluating sweetpotato as an intervention food to prevent vitamin A deficiency. Compr Rev Food Sci Food FaciFiFood Sci Food Sci Food Sc