



This work is licensed under a Creative Commons Attribution 3.0 License.

Research article

urn.lsid:zoobank.org:pub:58777EEB-7C5D-4887-8BE2-CEE4A4115CF5

Unexpected discovery of six new species of *Aphyosemion* (Cyprinodontiformes, Aplocheilidae) in the Wonga-Wongué Presidential Reserve in Gabon

Jean-François AGNÈSE^{1,*}, Laurent CHIRIO², Olivier LEGROS³,
Richard OSLISLY⁴ & Hervé Mvé BHÉ⁵

¹ISEM, CNRS, Univ. Montpellier, IRD, EPHE, Montpellier, France.

²14, rue des Roses, 06130 Grasse, France.

³Floralaan 51, 1501 Buizingen, Belgium.

⁴Patrimoines Locaux et Gouvernance UMR 208, IRD, MNHN, Paris, France.

⁴Agence Nationale des Parcs Nationaux (ANPN), BP 20379, Libreville, Gabon.

⁵Institut de Recherches Agronomiques et Forestières (IRAF), Centre National de la Recherche Scientifique et Technologique (CENAREST), BP 2246, Libreville, Gabon.

* Corresponding author: jean-francois.agnese@ird.fr

²Email: lichirio@hotmail.com

³Email: olivier.legros@belgacom.net

⁴Email: richard.oslisly@ird.fr

⁵Email: woleuntem@hotmail.com

¹urn.lsid:zoobank.org:author:8AEF0368-6444-4E50-8C72-23CCC65E69A5

²urn.lsid:zoobank.org:author:33A241AD-4C70-4A0A-8E30-B137E4FC771F

³urn.lsid:zoobank.org:author:F13EB21F-A391-451B-8139-9AA0F90AEF52

⁴urn.lsid:zoobank.org:author:1DC16C89-E5DB-43F2-874D-AAB2DC8BAF50

⁵urn.lsid:zoobank.org:author:2F0F18BD-C1AF-401B-94FE-0CB0AD1A8DE5

Abstract. During a survey of the fishes in the region of the Wonga-Wongué Presidential Reserve, 14 new populations of the subgenus *Chromaphyosemion* Myers, 1924 were found. These observations extend the previously known distribution range of the subgenus 120 kilometres southward. None of these populations could be related to any described species. Based on the colouration of the males and females, together with a genetic marker (mitochondrial DNA cytochrome b sequences), the populations studied are grouped into six new species which are described in this article, all close to *Aphyosemion alpha* Huber, 1998 with which they share the presence of a black alpha-shaped mark on the pre- and post-opercular region. The group composed of *A. alpha* and the six new species is referred to here as the *A. alpha* species group. All the new species, *A. aurantiacum* Chirio, Legros & Agnès sp. nov., *A. barakoniense* Chirio, Legros & Agnès sp. nov., *A. flammulatum* Chirio, Legros & Agnès sp. nov., *A. flavocyaneum* Chirio, Legros & Agnès sp. nov., *A. pusillum* Chirio, Legros & Agnès sp. nov. and *A. rubrogaster* Chirio, Legros & Agnès sp. nov., are further unambiguously diagnosed by unique combinations of colour patterns, making it possible to generate an identification key for the *A. alpha* species group. It is likely that the coastal dunes of Wonga-Wongué that form a sandy relief, could have

led to the fragmentation and then isolation of the hydrographical networks that flow into the Atlantic Ocean, making possible a significant number of allopatric speciations.

Keywords. Killifish, mtDNA, Nothobranchiidae, taxonomy.

Agnèse J.-F., Chirio L., Legros O., Oslisly R. & Mvé Bhé H. 2018. Unexpected discovery of six new species of *Aphyosemion* (Cyprinodontiformes, Aplocheilidae) in the Wonga-Wongué Presidential Reserve in Gabon. *European Journal of Taxonomy* 471: 1–28. <https://doi.org/10.5852/ejt.2018.471>

Introduction

Fishes of the genus *Aphyosemion* Myers, 1924 inhabit small rivers and freshwater streams in tropical and equatorial Africa, from Togo to Angola (Huber 2006). The genus is composed of more than 90 species. Since there is a considerable overlapping in meristic characters among species of *Aphyosemion* (Huber 2000; Scheel 1968, 1990; Wildekamp 1993), diagnoses generally rely on the extraordinary colour patterns of males. Scheel (1968, 1990) combined information from colour patterns, karyotypes, hybridization experiments and distribution to discern sets of related species, which he called species groups. Scheel (1990) recognized nine species groups, named after their emblematic species, *A. batesii* (Boulenger, 1911), *A. bivittatum* (Lönnerberg, 1895), *A. calliurum* (Boulenger, 1911), *A. cameronense* (Boulenger, 1903), *A. coeleste* Huber & Radda, 1977, *A. elegans* (Boulenger, 1899), *A. exiguum* (Boulenger, 1911), *A. georgiae* Lambert & Géry, 1968 and *A. striatum* (Boulenger, 1911), but admitted that certain species were difficult to place definitively into any one of these groups. Species from the *A. bivittatum* group (Scheel 1968) can easily be distinguished from all the other species of *Aphyosemion* through several characters, including the presence of two dark lateral bands in both sexes (vs one or none) and the ability of males to change their colouration rapidly depending on stress or their hierarchical status (vs less conspicuous changes in other *Aphyosemion*). The *A. bivittatum* group is now placed in *Chromaphyosemion* Radda, 1971 which is considered as a subgenus of *Aphyosemion* (Agnèse *et al.* 2006; Collier 2006) or a full genus (Legros *et al.* 2005; Sonnenberg 2000, 2007a, 2007b; Völker 2006). Agnèse *et al.* (2006) and Collier (2006) demonstrated that *Chromaphyosemion* is a monophyletic group, but still considered it as a subgenus of *Aphyosemion* until a complete taxonomic revision of the genus is done. The subgenus *Chromaphyosemion* is distributed from Togo to Gabon (Huber 2006), north of the equator, but most of its diversity (12 of 19 valid species) is found in Cameroon.

Until the present study, a population of *A. alpha* from Gabon, in the Owendo area south of Libreville (0.30750 N and 9.50972 E), was the southernmost population known. During our survey of fishes from south of Libreville, with emphasis on the region of the Wonga-Wongué Presidential Reserve, we found new populations of *Chromaphyosemion* all the way from Libreville to Owendo, and then south to the equator, from 0.04404 S to 0.70581 S.

Obviously, some of them are not conspecific with any described species. Within *Chromaphyosemion* it is not always easy to distinguish species, because there are no meristic differences between them. Species are only distinguished by male colouration and sometimes female colour patterns as well. Recently, the use of genetic markers (mitochondrial DNA sequences) has been very useful for solving many of the questions about the taxonomic status of certain populations (Agnèse *et al.* 2006, 2013; Sonnenberg 2007a, 2007b). Morphological, phenotypical and genetic analyses were performed in order to evaluate the taxonomic status of these newly-discovered populations in the coastal plains of Gabon.

Material and methods

Fish samples

Samples were collected in Gabon (fishing and export permit numbers AE0004/14/MESRS/CENAREST/CG/CST/CSAR and AE0006/16/MESRS/ CENAREST/CG/CST/CSAR) using dipnets, between April




Fig. 1. Collecting sites of the different populations. Black circles: *Aphyosemion alpha* Huber, 1998; Black triangles: *A. flavocyaneum* Chirio, Legros & Agnèse sp. nov.; red triangle: *A. flammulatum* Chirio, Legros & Agnèse sp. nov.; yellow triangles: *A. barakoniense* Chirio, Legros & Agnèse sp. nov.; red circles: *A. pusillum* Chirio, Legros & Agnèse sp. nov.; yellow circles: *A. rubrogaster* Chirio, Legros & Agnèse sp. nov.; red squares: *A. aurantiacum* Chirio, Legros & Agnèse sp. nov.

Table 1. Species identification; population number (N°) on the map (Fig. 1) and on the phylogenetic network (Fig. 2); voucher specimen references registered at the Royal Museum for Central Africa (MRAC), Tervuren (Belgium); coordinates.

Species name	N°	Voucher MRAC specimen 2016-019-P	Coordinates	
<i>A. flavocyaneum</i>	6	1–10	0.42874 S	9.54502 E
	9		0.43460 S	9.38216 E
	5		0.38381 S	9.39902 E
	3		0.32762 S	9.39489 E
<i>A. flammulatum</i>	4	11–18	0.39594 S	9.30772 E
<i>A. barakoniense</i>	7	19–36	0.47664 S	9.26483 E
	8	37–56	0.45815 S	9.33465 E
<i>A. pusillum</i>	10	57–59	0.55590 S	9.21322 E
	13	60–63	0.61282 S	9.15650 E
<i>A. aurantiacum</i>	12	64–73	0.58186 S	9.46718 E
	11	74–91	0.56336 S	9.33732 E
	14	92	0.66401 S	9.42652 E
<i>A. rubrogaster</i>	16	93–108	0.65524 S	9.57355 E
	15	109–121	0.70581 S	9.47592 E
<i>A. alpha</i>	1		0.56932 N	9.33858 E
	2		0.04404 S	9.34590 E

2014 and March 2016, from 16 natural populations, of which 14 (3 to 16) were from the Wonga-Wongué Presidential Reserve. The fish were anaesthetized with phenoxyethanol and then preserved in 90% alcohol. Collection localities and taxonomic identification of the samples are shown in Fig. 1 and Table 1.

Morphometric and phenotypic studies

Descriptions are based on wild-caught specimens registered in the collection of the Royal Museum of Central Africa (MRAC) in Tervuren, Belgium (Table 1). Phenotypic studies, diagnoses, descriptions and the identification key were made using Xper² (Ung *et al.* 2010). Morphometric data were obtained as described in Huber (2000) using a stereo microscope. Measurements were made with a digital calliper, corrected to the nearest 0.1 mm. All ratios are expressed as percentages of standard length (SL). Colouration data and live specimen photographs were obtained from wild fish after a few weeks to a few months in captivity. No colour variation was observed between individuals in the same population as is usually the case with this group of species.

Genetic study

Total genomic DNA was extracted from fin tissues (voucher specimens were deposited at the ISE-M, Montpellier, reference numbers are listed in Table 1) using the protocol described in Sambrook *et al.* (1989). A fragment of the mitochondrial gene cytochrome *b* was sequenced and used to both reconstruct a mitochondrial gene tree and evaluate genetic divergence. PCR was performed with two specific primers (ctb-F1 5'AACCACCGTTGTTATTCAAC3' forward and ctb-R1 5'CTCCCAAAGCCAGAATTCTAAA3' reverse). The amplification protocol consisted of 3 min at 93°C for initial denaturation, followed by 30 cycles of 30 sec at 93°C, 30 sec at 53°C for annealing, 1 min 30 sec at 72°C for extension, and a final 5 min extension step at 72°C.

AGNÈSE J.-F. *et al.*, Six new species of *Aphyosemion* (Cyprinodontiformes, Aplocheilidae)**Table 2.** GenBank accession numbers, species identification, voucher specimen references registered at the Institut des Sciences de l'Évolution de Montpellier (ISEM).

GenBank numbers	Species name	Voucher reference
MG779236	<i>A. flavocyaneum</i>	ISEM-LCH-108
MG779237		ISEM-LCH-138
MG779238		ISEM-LCH-171
MG779239		ISEM-LCH-107
MG779240		ISEM-LCH-141
MG779241		ISEM-LCH-139
MG779242		ISEM-LCH-168
MG779243		ISEM-LCH-167
MG779244	<i>A. flammulatum</i>	ISEM-LCH-162
MG779245	<i>A. barakoniense</i>	ISEM-LCH-354
MG779246		ISEM-LCH-352
MG779247	<i>A. pusillum</i>	ISEM-LCH-356
MG779248		ISEM-LCH-351
MG779249	<i>A. aurantiacum</i>	ISEM-LCH-144
MG779250		ISEM-LCH-355
MG779251		ISEM-LCH-353
MG779252	<i>A. rubrogaster</i>	ISEM-LCH-143
MG779253		ISEM-LCH-350
MG779254	<i>A. alpha</i>	ISEM-LCH-169
MG779255		ISEM-LCH-145

Phylogenetic analyses

All sequences were edited and aligned using Seqscape version 2.5 and subsequently inspected and corrected manually (Table 2). The sequences of all the valid species, obtained in a previous work (Agnèse *et al.* 2013) from the subgenus *Chromaphyosemion*, were included in the analyses: *A. alpha* Huber, 1998 (GenBank reference KC893920), *A. bitaeniatum* (Ahl, 1924) (KC893922), *A. bivittatum* (Lönnerberg, 1895) (KC893813), *A. ecucuense* (Sonnenberg, 2007) (KC893923), *A. erythron* (Sonnenberg, 2007) (KC893926), *A. kouamense* Legros, 1999 (KC893921), *A. koungeense* (Sonnenberg, 2007) (KC893896), *A. loennbergii* (Boulenger, 1903) (KC893884), *A. lugens* Amiet, 1991 (KC893894), *A. malumbresi* Legros & Zentz, 2007 (KC893927), *A. melanogaster* (Legros, Zentz & Agnèse, 2005) (KC893892), *A. melinoeides* (Sonnenberg, 2007) (KC893892), *A. omega* (Sonnenberg, 2007) (KC893882), *A. pamaense* Agnèse, Legros, Cazaux & Estivals, 2013 (KC893888), *A. poliaki* Amiet, 1991 (KC893913), *A. punctulatum* (Legros, Zentz & Agnèse, 2005) (KC893907), *A. riggenbachi* (Ahl, 1924) (KC893889), *A. splendopleure* (Brüning, 1929) (KC893904) and *A. volcanum* Radda & Wildekamp, 1977 (KC893879).

Aphyosemion franzwernerii Scheel, 1971 (KC893930), *A. ahli* Myers, 1933 (KC893931), *A. exiguum* (Boulenger, 1911) (KC893928), *A. elberti* (Ahl, 1924) (KC893929), *A. amoenum* Radda & Purzl, 1976 (KC893932) and *A. cameronense* (Boulenger, 1903) (KC893993) are species closely-related to *Chromaphyosemion* (Agnèse *et al.* 2006; Collier 2006) and were used as an out-group in the analysis. Tree search analyses were performed with (1) Maximum Likelihood (ML) (Stamakis 2014, RAxML, ver. 8), (2) Distance (Kimura 1980) and Neighbour Joining methods (Saitou & Nei 1987) and (3) Minimum Evolution approaches (Nei *et al.* 1998; Takahashi & Nei 2000). Supports for inferred clades were obtained through the nonparametric bootstrap (Felsenstein 1985), 2000 replicates for the three methods.




Fig. 2. Consensus tree based on Maximum Likelihood, Distance and Minimum Evolution. Numbers above the branches are percentages of bootstrap values based on 2000 replicates for each method. Numbers correspond to sample location (Fig. 1 and Table 1). Scale refers to Kimura 2 distance.

An evolutionary model for nucleotide substitution was chosen with MEGA 5 (Tamura *et al.* 2011) using the Bayesian information criterion (BIC) (Schwarz 1978). The optimal model was TN93+G+I (G=1.40, I=0.51). This model (Tamura & Nei 1993) accounted for the difference between transitions and transversions, and differentiated the two kinds of transition. The G parameter indicates that the non-uniformity of the evolutionary rates in the sites was modelled using a discrete Gamma distribution, and the I parameter indicates that there is a proportion of invariant sites. This model was used for the subsequent ME analysis. For RAxML, data were partitioned into three matrices corresponding to the three different codon position allowing the program to choose an independent evolutionary model for each partition. The different trees were rooted using *Aphyosemion franzwernerii*, *A. ahli*, *A. exiguum*, *A. elberti*, *A. amoenum* and *A. cameronense*.

To determine the delimitation of species, we used the Bayesian program bPTP (Zhang *et al.* 2013) on the web-server (<http://species.h-its.org/>) to estimate a species tree using our maximum likelihood tree, without the root species (i.e., only using the sequences of the *Chromaphyosemion* subgenus).

Results

Molecular analysis

A 1028 base-pair alignment for the Cytochrome *b* region from the mitochondrial genome was obtained after trimming the ends of each sequence. A total of 45 different sequences were obtained. A total of 456 variable sites were identified, from which 413 were parsimony informative (i.e., shared by at least two different sequences). The three different methods used, Maximum Likelihood, Distance and Minimum Evolution, gave congruent results, summarised in Fig. 2. Phylogenetic relationships within previously described species of *Chromaphyosemion* were congruent with the previous studies (Agnès *et al.* 2006; Collier 2006).

Aphyosemion alpha and all the newly-discovered populations were grouped in a monophyletic assemblage that appeared to be the sister group of all the other species of *Chromaphyosemion*.

This clade, which clustered all the populations or species from Gabon except for *A. kouamense*, was strongly supported by high bootstrap values (99/100/100 for ML, NJ and ME respectively).

Within this clade, the populations were clearly separated into seven groups. Two of these groups (A and B) were highly supported (bootstraps from 80 to 100) and composed of populations 15 and 16 on the one hand, and 10 and 13 on the other. Both these groups clustered together and might have represented the sister group of all the other populations, but this position was not supported by high bootstrap values (89/./67). Group C (populations 1, 2 and *A. alpha* (populations from Cap Esterias, LEC 96-26, GenBank reference KC893920)), group D (7, 8), group E (4), group F (11, 12, 14) and group G (3, 5, 6, 9) were all supported by very high bootstrap values (> 93). Groups D and E both formed a monophyletic assemblage (100/100/100) and, together with group F, another clade that was highly supported (98/93/98).

All the other species of *Chromaphyosemion*, except for *A. alpha*, were grouped into a monophyletic cluster that was strongly supported by high bootstrap values (100/92/100). The topology of the phylogenetic tree was similar to what has been observed previously (Agnès *et al.* 2013). *Aphyosemion lugens* and *A. melinoeides* occupied a basal position as the sister group of all the other species. The species from Equatorial Guinea: *A. erythron*, *A. ecucuense*, and *A. malumbresi*, plus *A. melanogaster* from Cameroon, appeared to form a monophyletic group (75/86/100).

Genetic distances (Kimura 2) observed between the haplotypes were used to create the histogram in Fig. 3. This Figure represents the distribution of distances between every pair of previously-described species, and between pairs of populations from the different groups of newly-discovered populations A

to G (one population per group, here 4, 8, 9, 13, 14, 16 and *A. alpha* (populations from Cap Esterias)). Genetic distances among the 18 previously-described species were distributed in two (yellow) groups. On the one hand, the distances ranged from 0.013 to 0.302, and on the other, the distances ranged from 0.412 to 0.620. The first group corresponds to the distances between all the pairs of species, except those including *A. alpha*, and the second group corresponds to the sequences between *A. alpha* and all the other 18 species of *Chromaphyosemion*. Genetic distances (red) in the batch of newly-discovered populations, including *A. alpha*, ranged from 0.016 and 0.253.




Fig. 3. Genetic distances (Kimura 2) between all pairs of species or populations. Only one population was used for group A to G, respectively population 4, 8, 9, 13, 14, 16 and *Aphyosemion alpha* Huber, 1998.

Even though there is no direct relationship between genetic distance and taxonomic distance, it can be seen that genetic divergences in the different groups of newly-discovered populations (A to G) appeared to be comparable with what has been already observed at the species level in *Chromaphyosemion*. For example, well-recognized species such as *A. splendopleure* and *A. omega*, or *A. splendopleure* and *A. koungueense*, or *A. koungueense* and *A. omega* exhibited low genetic divergences (0.013, 0.017, and 0.020 respectively) that were comparable to the distances observed in the present study between groups D and E (0.016).

Taking these observations into account, it is likely that all seven taxa (groups) of the southern Gabon species may represent seven different species: *A. alpha* and six others new to science. In order to verify this hypothesis, we have tried to confirm the number and the limits of the different new species present in southern Gabon using bPTP (Zhang *et al.* 2013). The most supported partition found by simple heuristic search, identified all previously described species except *A. omega*, *A. koungueense* and *A. splendopleure* which clustered together. For populations from southern Gabon, bPTP recognized seven species corresponding to the seven groups previously observed: group A (populations 15 and 16), group B (10, 13), group C (*A. alpha* from Cap Esterias together with populations 1 and 2), group D (7 and 8), group E (4), group F (11, 12 and 14) and group G (3, 5, 6 and 9).

Consequently, six species that are new to science are described below.

It should be noted that the authors of the new taxa are different from the authors of this paper (reference to Article 50.1 and Recommendation 50A of the International Code of Zoological Nomenclature; ICZN 1999).

Species descriptions

Order Cyprinodontiformes Berg, 1940
Suborder Aplocheiloidei Bleeker, 1859
Family Nothobranchiidae Garman 1895
Subfamily Nothobranchiinae Garman, 1895
Tribe Nothobranchiini Garman, 1895
Subtribe Aphyosemiina Huber, 2000
Genus *Aphyosemion* Myers, 1924

Aphyosemion flavocyaneum Chirio, Legros & Agnès sp. nov.
urn:lsid:zoobank.org:act:1C559D3D-38A3-4322-A7CF-BEBC547D4D8D

Fig. 4A–D, Table 3

Etymology

The specific epithet refers to the main colours of this species (blue and yellow).

Material examined

Holotype

GABON: adult ♂, 32.3 mm SL (43.2 mm TL), Lake Ndaminzé 0.42874 S, 9.54502 E, 115 m a.s.l., field reference code CHRSP1-Lac Ndaminzé, 13 Apr. 2014, Laurent Chirio leg. (MRAC 2016-019-P-1).

Paratypes

GABON: 3 ♂♂, 6 ♀♀, 21.3–27.5 mm SL, same collection data as for holotype (MRAC 2016-019-P-2–10).

Differential diagnosis

Distinguished from other species of *Chromaphyosemion* by its blue anal fin with no submarginal red band (vs species with a submarginal red band present), except for *A. poliaki*, and very rare specimens of *A. melanogaster* and *A. malumbresi*. Differs from the latter species through a combination of the following features: flanks and venter yellow-orange with no red punctuation (vs flanks bronze, venter brown-orange, regular red dots; flanks yellow-green, venter yellow-green with a black zone, irregular lines of red dots; flanks bluish-white, venter bluish with a black zone, regular red dots), blue iridescent scales on the ventral region close to the caudal peduncle (vs red scales or black scales), black alpha-shaped mark on the pre- and post-opercular region (vs no alpha-shaped mark, only a few red macules), anal fin blue with an orange portion on the basal region with no punctuation except black dots on basal and/or posterobasal portions (vs anal fin blue-green with many red dots; yellow-green with red dots; bluish-white with red dots).

Colouration of live males (Fig. 4A)

FLANKS AND VENTER. Yellow-orange with no red punctuation, two greyish lateral stripes on ventral region close to caudal peduncle with some blue iridescent scales. Two rows of paradorsal golden scales in dorsal region from operculum to caudal peduncle with few black macules.

HEAD. Yellow-orange. Premaxilla and mandible yellow-orange with black lower lips, orange infraorbital region with black macule, orange post-orbital region with two black macules, yellow-orange pre-opercle with two black macules, yellow-orange opercle with two black macules; black macules on pre- and post-opercular region reveal an alpha-shaped mark.

FINS. The dorsal fin is blue, but more orange near the insertion of the fin, and rows of black dots between rays. Anal fin blue with orange portion on basal region of fin, with no punctuation and no submarginal or marginal bands. Certain individuals may have some small black dots on basal region. Caudal fin blue with many carmine red dots and some carmine red streaks in upper and lower lobes; submarginal and marginal red bands absent. Acumens at apex of unpaired fins orange or white for dorsal fin, very short blue acumen for anal fin, and orange or white for caudal fin. Colour pattern of pelvic fins identical to anal fin with blue background and no submarginal band or marginal band. Pectoral fins translucent.

Colouration of live females (Fig. 4B)

FLANKS AND VENTER. Pale brown, four lines of red dots from opercle to caudal peduncle, two greyish lateral stripes, venter yellow-green, white on basal portion. Two rows of paradorsal golden scales in dorsal region, from operculum to caudal peduncle with red dots.

HEAD. Premaxilla brown, mandible pale orange with black lower lips, brown supraorbital region, white infraorbital region with red macule, grey postorbital region with two macules, white pre-opercle, yellow-orange opercle with one red dot.

FINS. Dorsal fin yellow-green, rows of red dots between rays. Anal fin yellow-green with rows of red dots between rays. Caudal fin yellow-green, blue-green on distal portion, with rows of red dots between rays. Pelvic fins translucent with yellow-green reflection, red dots, blue reflection on edges. Pectoral fins are translucent.

Colouration of ethanol-preserved males and females (Fig. 4C–D)

FLANKS AND VENTER. Flanks of males grey-yellow with three short lines of red dots, venter grey-yellow with no red punctuation. Two rows of paradorsal red scales in brown dorsal region from operculum to caudal peduncle. Flanks of females grey-yellow, three lines of red dots from the opercle to the caudal peduncle. Two rows of paradorsal red scales in the brown dorsal region, from the operculum to the caudal peduncle.




Fig. 4. A–D. *Aphyosemion cyanoflavum* Chirio, Legros & Agnèse sp. nov. **A.** Adult, ♂, from locality 9, not preserved. Photo by O. Buisson. **B.** Adult, ♀, from locality 9, not preserved. Photo by O. Buisson. **C.** Holotype, adult, ♂, from Lake Ndaminzé (MRAC 2016-019-P-1). **D.** Paratype, adult, ♀, from Lake Ndaminzé (MRAC 2016-019-P-2-10). E–H. *A. flammulatum* Chirio, Legros & Agnèse sp. nov. **E.** Adult, ♂, from locality 4, not preserved. **F.** Adult, ♀, from locality 4, not preserved. **G.** Holotype, adult, ♂, from the lower Aloumbé River (MRAC 2016-019-P-11). **H.** Paratype, adult, ♀, from the lower Aloumbé River (MRAC 2016-019-P-12-18).

Table 3. Meristic and morphometric data of *Aphyosemion flavocyaneum* Chirio, Legros & Agnès sp. nov. Abbreviations: A = anal-fin rays; D = Dorsal-fin rays; D/A = displacement of the dorsal in relation to the anal fin expressed as the number of anal fin rays; dcp = caudal peduncle height; Eye = eye diameter; Hd = head length; HT = holotype values; Ht = height at the anal; I.O. = interorbital space; L.L. = lateral line scales, between brackets scales on the caudal fin; Max = maximum observed value; Mean = mean of observed value; Min = minimum observed value; N = number of specimens studied; P.A. = preanal distance; P.D. = predorsal distance; pDor = predorsal scales; P.V. = preventral distance; SD = standard deviation; S.L. = standard length in mm; all other measurements in percentage from S.L.; T.L. = total length; TRAV = transversal scales.

Character	N	HT	Min	Max	Mean	SD
D	10	12	10	12	10.80	0.63
A	10	13	11	14	12.60	1.07
D/A	10	-1	-1	1	0	0.82
L.L.	10	24 (+3)	22 (+1)	26 (+3)	24.2 (+2)	1.55 (+0.67)
pDor	10	12	11	13	11.8	0.63
TRAV	10	8	6	8	7.1	0.57
S.L.	10	32.3	21.3	32.3	24.59	3.45
T.L. (%)	10	134	126	136	130.87	4.19
P.D. (%)	10	62	56	68	60.07	3.32
P.A. (%)	10	65	59	66	61.65	2.72
P.V. (%)	10	51	47	53	50.05	1.99
Ht (%)	10	22	17	22	18.84	1.33
dcp (%)	10	12	11	14	12.23	0.97
Hd (%)	10	31	30	36	32.82	2.32
I.O. (%)	10	15	13	17	14.93	1.31
Eye (%)	10	7	7	9	7.53	0.48

HEAD. Premaxilla and mandible of male grey with black lower lips, infraorbital region has red macule, opercle with one red macule; very dark macules in post-opercular region. Head of female with a black lower lips, infraorbital region with a red macule. Head of female with black lower lips, infraorbital region with red macule.

FINS. Male dorsal fin grey-blue with rows of red dots between rays. Anal fin grey-blue with some small red dots in posterobasal region. Caudal fin grey-blue with many red dots and some carmine red streaks in upper and lower lobes. Female dorsal fin greyish, with rows of red dots between rays. Anal fin greyish, with rows of red dots between rays. Caudal fin greyish, with rows of red dots between rays.

Distribution and habitat

This species has been found at seven localities (for three of them, specimens have not been studied or deposited in a museum) in the Awagné River Basin (Table 1, Fig. 1), has never been found elsewhere and seems to be endemic to this hydrographic basin. It lives in small, secondary rivers and small, forest streams, often with a sandy bottom, where it can be found along the riverbanks. It hides quickly in water plants or dead leaves. In one locality, the bottom is muddy, the water quite stagnant and some fish have even been observed in small water holes. The uppermost locality is Lake Ndaminzé, which is about 1 km long: the species was found only under the shadow of large trees, hidden in the roots near the banks. This species was found in syntopy with an undescribed species of *Aphyosemion* and an undescribed species of *Epiplatys*. At one single locality in the Mbomba River, a tributary of the Awagné River, it was found in syntopy with *Poropanchax stigmatopygus* (Wildekamp & Malumbres, 2004).

Aphyosemion flammulatum Chirio, Legros & Agnès sp. nov.
urn:lsid:zoobank.org:act:EEF24797-1B05-4D20-BCCB-A3336028C798

Fig. 4E–H, Table 4

Etymology

The specific epithet refers to the colour of the tip of the dorsal fin in this species (bright orange).

Material examined

Holotype

GABON: adult ♂, 36.2 mm SL (42 mm TL), lower Aloumbé River, 0.39594 S, 9.30772 E, 30 m a.s.l., field reference code CHRSP2-Aloumbé, 11 Apr. 2014, Laurent Chirio leg. (MRAC 2016-019-P-11).

Paratypes

GABON: 3 ♂♂, 4 ♀♀, 29.4–42.4 mm SL, same collection data as for holotype (MRAC 2016-019-P-12–18).

Differential diagnosis

Differs from all other species in the *A. alpha* group through combination of following features: head intense orange, flanks purplish-blue, ventral region bluish (vs never same colour pattern), anal fin purplish-blue, with no punctuation except for some black macules in basal region and irregular submarginal red band (vs anal fin orange with or without punctuation, anal fin blue-orange with no punctuation; regular or no submarginal red band). Differs from all other species of *Chromaphyosemion* with black alpha-shaped mark in pre- and post-opercular region (vs no alpha-shaped mark, only a few red macules).

Colouration of live males (Fig. 4E)

FLANKS AND VENTER. Flanks purplish-blue with two very dark and often visible lateral stripes, three lines of very dark red dots from operculum to caudal peduncle. Ventral region bluish-white with some blue iridescent scales between end of anal fin and caudal peduncle. Dorsal region brown, two rows of paradorsal yellow-orange scales in dorsal region, from operculum to caudal peduncle, with two lines of carmine red dots.

HEAD. Very intense orange: premaxilla brown, mandible orange with black lower lips, brown supraorbital region, orange infraorbital region with black macule, orange pre-opercle, orange operculum with three black macules; black macules on pre- and post-opercular region reveal alpha-shaped mark.

FINS. Dorsal fin blue, orange distal portion near apex, rows of black dots between rays; very long orange acumen on apex. Anal fin purplish-blue, with no punctuation except for few black macules in basal region of fin, red, irregular submarginal band, blue marginal band, blue acumen. Caudal fin blue with orange portion in peduncle region, few red dots in basal region, very long red streaks in distal region and in upper and lower lobes, red, irregular submarginal band, blue marginal band, and acumen of very intense orange. Pelvic fins blue-orange, submarginal red band and blue marginal band. Pectoral fins with orange reflections.

Colouration of live females (Fig. 4F)

FLANKS AND VENTER. Pale beige, with four lines of red and orange dots from opercle to caudal peduncle, and two very dark lateral stripes, venter yellow, white on basal portion. Two rows of paradorsal golden scales in dorsal region from operculum to start of anal fin with red dots, copper scales from start of anal fin to caudal peduncle, with red dots.

Table 4. Meristic and morphometric data of *Aphyosemion flammulatum* Chirio, Legros & Agnès sp. nov. See Table 3 for the abbreviations used.

Character	N	HT	Min	Max	Mean	SD
D	8	14	11	14	11.75	1.04
A	8	13	13	14	13.13	0.35
D/A	8	2	1	3	1.88	0.64
L.L.	8	25 (+2)	24 (+1)	28 (+2)	25.75 (+1.75)	1.28 (+0.46)
pDor	8	13	12	13	12.88	0.35
TRAV	8	7	7	7	7	0
S.L.	8	36.2	25.7	36.2	30.35	3.97
T.L. (%)	8	116	114	125	119.37	3.59
P.D. (%)	8	53	52	59	55.14	2.14
P.A (%)	8	53	51	64	56.33	3.84
P.V.(%)	8	41	41	50	45.4	2.56
Ht (%)	8	20	16	26	19.31	3.2
dcp (%)	8	12	8	18	11.68	2.65
Hd (%)	8	30	10	30	25.59	6.22
I.O. (%)	8	13	12	15	13.58	0.75
Eye (%)	8	7	5	7	5.84	0.38

HEAD. Premaxilla brown, mandible yellow with black lower lips, brown supraorbital region, white infraorbital region with black macule, orange postorbital region with two red macules, orange preopercle, and orange opercle with five red dots.

FINS. Dorsal fin orange, rows of carmine red dots between rays. Anal fin orange in median portion, some red dots between the rays in posteromedian portion. Caudal fin orange near caudal peduncle, yellow-orange with red dots in median portion, and translucent in distal portion. Pelvic fins translucent with orange reflection, blue reflection on edges. Pectoral fins translucent with orange reflection.

Colouration of ethanol-preserved males and females (Fig. 4G–H)

FLANKS AND VENTER. Flanks of males yellowish with no red dots, upper grey visible upper lateral stripe, venter yellowish. Two rows of paradorsal black scales in brown dorsal region from operculum to caudal peduncle. Flanks of females yellowish, four lines of red dots from opercle to caudal peduncle. Two rows of paradorsal black scales in yellowish dorsal region from operculum to the caudal peduncle.

HEAD. Head of male with black lower lips, infraorbital region with black macule, dark macules in postopercular region. Head of female with black lower lips, infraorbital region with red macule, and postopercular region with dark macules.

FINS. Male dorsal fin grey with rows of red dots between rays. Anal fin grey with no dots. Caudal fin grey with red dots in median portion and some carmine red streaks in upper and lower lobes. Female dorsal fin blue-orange with red dots. Anal fin blue-orange, caudal fin blue-orange in upper and lower regions, some red dots near peduncle region.

Distribution and habitat

This species has been found at only three localities (for two of them, specimens have not been studied or deposited in a museum) in the Aloumbé Basin or from a small, unnamed river just north of the Aloumbé

AGNÈSE J.-F. *et al.*, Six new species of *Aphyosemion* (Cyprinodontiformes, Aplocheilidae)

River (Table 1, Fig. 1) and never elsewhere. It also seems to be endemic to this very small, littoral hydrographic basin. It lives in very small forest streams (less than 1 metre wide), with sandy or rocky bottoms. It hides among roots or dead leaves. This species was found in syntopy with an undescribed species of *Aphyosemion*.

Aphyosemion barakoniense Chirio, Legros & Agnès sp. nov.

[urn:lsid:zoobank.org:act:93CAFA79-3B4F-4F39-8430-D9A0DB9A58C5](https://zoobank.org/act:93CAFA79-3B4F-4F39-8430-D9A0DB9A58C5)

Fig. 5A–D, Table 5

Etymology

The specific epithet refers to the river in which the species was found.

Material examined

Holotype

GABON: adult ♂, 39.9 mm SL (48.9 mm TL), lower Barakonié River, 0.47664 S, 9.26483 E, 5 m a.s.l., field reference code CHRSP3-Basse Barakonié, 2 Mar. 2016, Laurent Chirio leg. (MRAC 2016-019-P-19).

Paratypes

GABON: 7 ♂♂, 10 ♀♀, 21.5–46.6 mm SL, same data as for holotype (MRAC 2016-019-P-20-36); 12 ♂♂, 8 ♀♀, 20.7–37.5 mm SL, upper Barakonié River, 0.45815 S, 9.33465 E, 55 m a.s.l., field reference code CHRSP3-Haute Barakonié, 3 Mar. 2016, Laurent Chirio leg. (MRAC 2016-019-P-37-56).

Differential diagnosis

Differs from all other species in *A. alpha* group through combination of following features: head orange, flanks grey-mauve, ventral region orange (vs never same colour pattern), anal fin orange with some red dots (vs no punctuation or only black dots on posterobasal portion for *A. flavocyaneum* Chirio, Legros & Agnès sp. nov. and *A. flammulatum* Chirio, Legros & Agnès sp. nov.). Distinguished from all other species of *Chromaphyosemion* by alpha-shaped mark in pre- and post-opercular region (vs no alpha-shaped mark, only some red macules).

Colouration of live males (Fig. 5A)

FLANKS AND VENTER. Flanks grey-mauve with two dark lateral stripes, three lines of very dark red dots from opercle to caudal peduncle. Ventral region orange with some blue iridescent scales between end of anal fin and caudal peduncle. Dorsal region brown, two rows of paradorsal copper scales in dorsal region from operculum to caudal peduncle with two lines of carmine red dots.

HEAD. Orange, premaxilla brown, mandible orange with black lower lips, with brown supraorbital region, orange infraorbital region with black macule, postorbital region orange with two black macules, orange pre-opercle with two black macules, orange opercle with three black macules; black macules on pre- and post-opercular regions reveal alpha-shaped drawings.

FINS. Dorsal fin blue on basal portion, orange on upper portion, yellowish on edge of fin, with rows of carmine red dots between rays, orange acumen on apex. Anal fin mauve on basal portion, orange on median portion, yellowish portion near the submarginal band, with some carmine red macules, regular, red submarginal band, blue marginal band, small orange acumen. Caudal fin light mauve, with tips of upper and lower lobes orange, red dots on median region, short red streaks in distal region and in upper and lower lobes, irregular red submarginal band, blue marginal band, orange acumens. Pelvic fins orange with no punctuation, submarginal red band, blue marginal band. Pectoral fins with orange reflections.

Colouration of live females (Fig. 5B)

FLANKS AND VENTER. Flanks beige-pink, four lines of red dots from opercle to caudal peduncle, two greyish lateral stripes, venter yellow, white on basal portion. Two rows of paradorsal copper scales in dorsal region from operculum to caudal peduncle with red dots.

HEAD. Premaxilla brown, mandible yellow; orange zone on side portion with black lower lips, brown supraorbital region, white infraorbital region with black macule, orange postorbital region with two black macules, orange pre-opercle, and orange opercle with two black dots.




Fig. 5. A–D. *Aphyosemion barakoniense* Chirio, Legros & Agnèse sp. nov. **A.** Adult, ♂, from locality 7, not preserved. **B.** Adult, ♀ from locality 7, not preserved. **C.** Holotype, adult, ♂, from the lower Barakonié River (MRAC 2016-019-P-19). **D.** Paratype, adult, ♀, from the lower Barakonié River (MRAC 2016-019-P-20-36). E–F. *A. pusillum* Chirio, Legros & Agnèse sp. nov. **E.** Adult, ♂, from locality 10, not preserved. **F.** Holotype, adult, ♂, from the Okoyo River (MRAC 2016-019-P-57).

AGNÈSE J.-F. *et al.*, Six new species of *Aphyosemion* (Cyprinodontiformes, Aplocheilidae)**Table 5.** Meristic and morphometric data of *Aphyosemion barakoniense* Chirio, Legros & Agnès sp. nov. See Table 3 for the abbreviations used.

Character	N	HT	Min	Max	Mean	SD
D	38	11	10	14	11.42	0.83
A	38	13	11	15	12.71	0.96
D/A	38	3	0	3	1.37	1.1
L.L.	38	26 (+2)	23 (+1)	26 (+4)	24.84 (+2.03)	0.95 (+0.49)
pDor	38	13	11	14	12.58	0.68
TRAV	38	8	7	8	7.18	0.39
S.L.	38	39.9	20.7	39.9	28.34	4.91
T.L. (%)	38	123	111	126	119.18	3.48
P.D. (%)	38	58	52	61	56.91	2.17
P.A. (%)	38	57	49	60	55.04	2.8
P.V. (%)	38	47	41	55	46.12	2.9
Ht (%)	38	18	14	21	17.4	1.57
dcp (%)	38	12	9	14	11.03	1.06
Hd (%)	38	30	23	33	29.58	2.26
I.O. (%)	38	12	10	15	12.15	0.97
Eye (%)	38	5	5	11	6.43	1.14

FINS. Dorsal fin intense orange, with rows of carmine red dots between rays. Anal fin orange on the median portion with no dots. Caudal fin intense orange near caudal peduncle, red dots between rays. Pelvic fins with orange reflection, blue reflection on edges. Pectoral fins translucent.

Colouration of ethanol-preserved males and females (Fig. 5C–D)

FLANKS AND VENTER. Flanks of males dark grey with three lines of red dots, venter grey-yellow. Two rows of paradorsal red scales in brown dorsal region from operculum to caudal peduncle. Flanks of females beige with two dark lateral stripes, six lines of red dots from opercle to caudal peduncle, venter grey-yellow. Two rows of paradorsal red scales in grey dorsal region, from operculum to caudal peduncle.

HEAD. Premaxilla and mandible of male grey, with black lower lips, orange infraorbital region with black macule, orange pre-opercular region, orange opercle with one red macule; dark macules in post-opercular region. Head of female with black lower lips, infraorbital region with red macule.

FINS. Dorsal fin of male orange with rows of red dots between rays. Anal fin orange with some small red dots, red sumarginal band. Caudal fin orange with many red dots on basal portion, red streaks on edge and in upper and lower lobes. Dorsal fin of female orange, with rows of red dots between rays. Anal fin orange, with red streaks between rays. Caudal fin orange, with rows of red dots between rays on median portion, red streaks on edge.

Distribution and habitat

This species has been found at only two localities in the Barakonié Basin (Table 1, Fig. 1) and never elsewhere. It seems to be endemic to this small, coastal hydrographic basin. In the upper Barakonié, where the river is less than 1 m wide, the fish live among roots or dead leaves in the middle of the river. In the lower Barakonié, where the river can be 2 to 3 m wide, they are not found in the main course, but only in small water holes of stagnant water close to the river, hidden among dead leaves and mud, and there, they can be very abundant. This species was found in syntopy with an undescribed species of *Aphyosemion*.

Aphyosemion pusillum Chirio, Legros & Agnès sp. nov.
[urn:lsid:zoobank.org:act:529ADFCC-1E00-4715-A157-1281E749DE48](https://doi.org/10.21203/europeanjournaloftaxonomy.eu/471/157)

Fig. 5E–F, Table 6

Etymology

The specific epithet refers to the small size of this species.

Material examined

Holotype

GABON: adult ♂, 28.5 mm SL (33.8 mm TL), bridge on Okoyo River, 0.55590 S, 9.21322 E, 10 m a.s.l., field reference code CHRSP4-Okoyo, 1 Mar. 2016, Laurent Chirio leg. (MRAC 2016-019-P-57).

Paratypes

GABON: 2 ♂♂, 24.5–24.8 mm SL, same collection data as for holotype (MRAC 2016-019-P-58-59); 4 ♂♂, 20.6–23 mm SL, bridge on Pembé River, 0.61282 S, 9.15650 E, 5 m a.s.l., field reference code CHRSP4-Pembé, 1 Mar. 2016, Laurent Chirio leg. (MRAC 2016-019-P-60-63).

Differential diagnosis

Distinguished from all other species in *A. alpha* group by two delimited orange zones on body: orange portion on opercle and orange portion above pelvic fins (vs no orange portion) with beige ventral region (vs yellow-orange, white bluish or orange), anal fin orange, blue near basal portion and near submarginal red band, no punctuation (vs never a similar colour pattern). Distinguished from all other species of *Chromaphyosemion* by alpha-shaped mark on pre- and postopercular region, especially visible on preserved specimens (vs no alpha-shaped mark, only some red macules).

Colouration of live males (Fig. 5E)

FLANKS AND VENTER. Flanks beige with two dark lateral stripes, two lines of small red dots from opercle to caudal peduncle. Ventral region beige with some blue iridescent scales between end of anal fin and caudal peduncle, orange portion above pelvic fins. Dorsal region light brown, two rows of paradorsal copper scales from operculum to caudal peduncle with two lines of red dots.

HEAD. Beige with premaxilla brown, mandible yellow orange with black lower lips, brown supraorbital region, beige infraorbital region with black macule, postorbital region beige with one black macule, beige preopercle with one black macule, opercle yellow orange in background with orange zone with four black macules; black macules on pre- and postopercular region reveal incomplete alpha-shaped mark.

FINS. Dorsal fin orange, yellowish on edge of fin, blue on posterobasal portion of fin, rows of carmine red dots between rays, red streaks on edge of fin, orange acumen on apex. Anal fin orange, blue near basal portion of fin and near submarginal band, regular red submarginal band, yellow green marginal band, small orange acumen. No punctuation. Caudal fin blue with beginning of upper and lower lobes light orange, red dots on upper part of fin, long red streaks on upper and median regions, red regular submarginal band, blue marginal band, orange acumens. Pelvic fins orange without punctuation, blue near submarginal band, submarginal red band and yellow green marginal band. Pectoral fins translucent.

Colouration of live females

Unknown

AGNÈSE J.-F. *et al.*, Six new species of *Aphyosemion* (Cyprinodontiformes, Aplocheilidae)**Table 6.** Meristic and morphometric data of *Aphyosemion pusillum* Chirio, Legros & Agnès sp. nov. See Table 3 for the abbreviations used.

Character	N	HT	Min	Max	Mean	SD
D	7	10	9	12	10.43	0.98
A	7	11	11	14	12.29	1.11
D/A	7	0	0	2	1.43	0.98
L.L.	7	27 (+2)	24 (+2)	27 (+3)	26.14 (+2.29)	1.21 (+0.49)
pDor	7	13	13	15	13.57	0.79
TRAV	7	8	7	8	7.43	0.53
S.L.	7	28.5	20.6	28.5	23.59	2.68
T.L. (%)	7	119	114	122	117.72	2.6
P.D. (%)	7	56	56	61	58.08	2.52
P.A. (%)	7	65	57	65	60.48	2.99
P.V. (%)	7	52	43	52	47.3	3.19
Ht (%)	7	21	13	21	16.63	2.44
dcp (%)	7	11	7	13	10.29	1.84
Hd (%)	7	27	27	36	30.27	3.15
I.O. (%)	7	14	12	15	13.91	1.16
Eye (%)	7	6	6	9	7.53	0.99

Colouration of ethanol preserved males (Fig. 5F)

FLANKS AND VENTER. Grey with three incomplete lines of red dots, venter beige with orange portion, white lower part, two grey lateral stripes. Two rows of paradorsal red scales in grey dorsal region from operculum to caudal peduncle.

HEAD. White mandible with black lower lips, white infraorbital region with black macule, white preopercular region, orange opercle; one dark macule in postopercular region.

FINS. Dorsal fin grey with rows of grey dots between rays. Anal fin with grey area on basal portion. Caudal fin grey with some red streaks on upper lobe, red dots in median portion.

Colouration of preserved females

Unknown.

Distribution and habitat

This small species has been found at only two localities in the Okoyo and Pembé Basins (Table 1, Fig. 1) and never elsewhere. The species seems to be endemic to these two small, coastal hydrographic basins, situated just north of the Wézé drainage. In the Okoyo Basin, *A. pusillum* Chirio, Legros & Agnès sp. nov. was found out of the main course of the river, in small, muddy streams less than 1 m wide. In the Pembé Basin, it was found directly in the course of this small river, hidden among dead leaves and water plants, in quiet places. This species was found in syntopy with an undescribed species of *Aphyosemion*, two undescribed species of *Epiplatys* and an undescribed species of *Plataplochilus*.

Aphyosemion aurantiacum Chirio, Legros & Agnès sp. nov.
urn:lsid:zoobank.org:act:93840539-567F-4817-B774-3568D388E094

Fig. 6A–D, Table 7

Etymology

The specific epithet refers to the main colour of this species (dark orange).

Material examined

Holotype

GABON: adult ♂, 33 mm SL (39.7 mm TL), Wézé spring, 0.58186 S, 9.46718 E, 89 m a.s.l., field reference code CHRSP5-sources Wézé, 3 Jul. 2014, Laurent Chirio leg. (MRAC 2016-019-P-64).

Paratypes

GABON: 4 ♂♂, 5 ♀♀, 23.4–29.9 mm SL, same collection data as for holotype (MRAC 2016-019-P-65-73); 18 ♂♂, 22.2–32.1 mm SL, northern tributary of Wézé River, 0.56336 S, 9.33732 E, 45 m a.s.l., field reference code CHRSP5-Wézé2, 1 Mar. 2016, Laurent Chirio leg. (MRAC 2016-019-P-74-91); 1 ♀, 23 mm SL, southern tributary of Wézé River, 0.66401 S, 9.42652 E, 87 m a.s.l., field reference code CHRSP5-Wézé3, 5 Mar. 2016, Laurent Chirio leg. (MRAC 2016-019-P-92).

Differential diagnosis

Differs from all other species in *A. alpha* group through combination of following features: head orange, flanks beige, ventral region orange (vs never the same colour pattern), orange anal fin with no punctuation (vs blue-orange, purplish-blue or orange with punctuation). Distinguished from all other species of *Chromaphyosemion* by an alpha-shaped drawing in pre- and post-opercular region (vs no alpha-shaped drawing, only some red macules).

Colouration of live males (Fig. 6A)

FLANKS AND VENTER. Flanks beige with two grey, rarely visible lateral stripes, four lines of red dots from opercle to caudal peduncle. Ventral region orange and region between end of anal fin and caudal peduncle intense orange, two rows of blue iridescent scales between end of anal fin and caudal peduncle. Dorsal region brown, with two rows of paradorsal copper scales from operculum to caudal peduncle with two lines of carmine red dots.

HEAD. Orange with premaxilla brown, mandible orange with black lower lips, brown supraorbital region, orange infraorbital region with black macule, postorbital region orange with two black macules, orange pre-opercle with one black macule, orange opercle with two black macules; black macules on pre- and post-opercular region reveal an alpha-shaped drawing.

FINS. Dorsal fin blue, large orange zone on rays, rows of carmine red dots between rays, and carmine red streaks on distal and posterobasal portions of fin. Orange acumen on apex. Anal fin orange, blue zone near submarginal band, regular, red submarginal band, blue marginal band, orange acumen. No punctuation. Caudal fin blue with orange caudal peduncle, 5–6 red dots in median region, very long red streaks in median region and in upper and lower lobes, regular, red submarginal band, blue marginal band, and orange acumens. Pelvic fins orange with no punctuation, submarginal red band and blue marginal band. Pectoral fins with orange reflections.

Colouration of live females (Fig. 6B)

FLANKS AND VENTER. Flanks pale yellow, with four lines of red dots from opercle to caudal peduncle, two greyish lateral stripes, yellow venter, white on basal portion. Two rows of paradorsal golden scales

AGNÈSE J.-F. *et al.*, Six new species of *Aphyosemion* (Cyprinodontiformes, Aplocheilidae)

in dorsal region from operculum to start of anal fin with red dots, copper scales from start of anal fin to caudal peduncle with red dots.

HEAD. Premaxilla brown, mandible pale yellow with black lower lips, brown supraorbital region, white infraorbital region with black macule, white postorbital region with two red macules, yellow pre-opercle with red dot, yellow opercle with two red dots.

FINS. Dorsal fin yellow-green, with rows of carmine red dots between rays, red streaks on distal portion. Anal fin blue on basal portion with row of red dots between rays, some red streaks between rays on median portion, pale orange on distal portion. Caudal fin yellow-green near caudal peduncle, with some red dots on basal portion, red streaks between rays from median to distal portion. Pelvic fins translucent with blue reflection on the edge. Pectoral fins translucent.

Colouration of ethanol-preserved males and females (Fig. 6C–D)

FLANKS AND VENTER. Flanks of males grey-yellow with no red dots, venter grey-yellow. Two rows of paradorsal black scales in grey-yellow dorsal region from operculum to caudal peduncle, visible greyish upper lateral stripe. Flanks of females beige with two grey lateral stripes and two lines of two to three red dots behind opercle. Venter beige, with two grey lateral stripes. Two rows of paradorsal red scales in grey dorsal region from operculum to caudal peduncle.

HEAD. Head of male grey-yellow mandible with black lower lips, grey-yellow infraorbital region with small black macule, grey-yellow pre-opercular region, opercle with two black macules; dark macules in post-opercular region. Head of female with black lower lips, infraorbital region with black macule, dark opercle.

FINS. Dorsal fin of male orange with two to three black dots on basal portion, streaks between rays on distal portion. Anal fin orange with dark rays. Caudal fin orange with many red streaks on distal portion, red submarginal band. Dorsal fin of female light orange, rows of red dots between rays. Anal fin greyish with no red dots. Caudal fin light orange, with some red dots on upper portion, red streaks on edge.

Distribution and habitat

This species has been found at three localities in the Wézé Basin (Table 1, Fig. 1) and never elsewhere. The species seems to be endemic to this hydrographic basin. It lives only in very small forest streams and water holes, often with a bottom consisting of roots and dead leaves, where it can be very abundant. In the southern-most locality, it was found hidden under dead leaves along the river banks in a 2 metre wide river with a sandy bottom. This species was found in syntopy with three undescribed species of *Aphyosemion*, *Epiplatys* and *Plataplochilus*.

Aphyosemion rubrogaster Chirio, Legros & Agnès sp. nov.
urn:lsid:zoobank.org:act:1778A417-0D05-4C16-BE75-00B6DE36398C

Fig. 6E–H, Table 8

Etymology

The specific epithet refers to the colour of the ventral region of this species (red).

Material examined

Holotype

GABON: adult ♂, 25.2 mm SL (31.3 mm TL), bridge on Niengé River, 0.65524 S, 9.57355 E, 61 m a.s.l., field reference code CHRSP6-Niengué, 4 Jul. 2014, Laurent Chirio leg. (MRAC 2016-019-P-93).




Fig. 6. A–D. *Aphyosemion aurantiacum* Chirio, Legros & Agnèse sp. nov. **A.** Adult, ♂, from locality 12, not preserved. Photo O. Buisson. **B.** Adult, ♀, from locality 12, not preserved. Photo O. Buisson. **C.** Holotype, adult, ♂, from Wézé Spring (MRAC 2016-019-P-64). **D.** Paratype, adult, ♀, from Wézé Spring (MRAC 2016-019-P-65-73). **E–H.** *A. rubrogaster* Chirio, Legros & Agnèse sp. nov. **E.** Adult, ♂, from locality 16, not preserved. Photo O. Buisson. **F.** Adult, ♀, from locality 16, not preserved. Photo O. Buisson. **G.** Holotype, adult, ♂, from the Niengé River (MRAC 2016-019-P-93). **H.** Paratype, adult, ♀, from the Niengé River (MRAC 2016-019-P-94-108).

AGNÈSE J.-F. *et al.*, Six new species of *Aphyosemion* (Cyprinodontiformes, Aplocheilidae)**Table 7.** Meristic and morphometric data of *Aphyosemion aurantiacum* Chirio, Legros & Agnès sp. nov. See Table 3 for the abbreviations used.

Character	N	HT	Min	Max	Mean	SD
D	29	12	10	13	11.21	0.86
A	29	12	12	15	13.21	0.98
D/A	29	1	-1	2	0.86	0.88
L.L.	29	26 (+2)	24 (+1)	27 (+3)	25.52 (+1.86)	0.83 (+0.52)
pDor	29	12	11	13	12.38	0.56
TRAV	29	8	7	8	7.1	0.31
S.L.	29	33	22.2	33	26.85	2.84
T.L. (%)	29	120	112	135	122.92	5.22
P.D. (%)	29	58	47	67	56.52	4.21
P.A. (%)	29	55	47	65	56.78	3.22
P.V.(%)	29	45	39	51	46.31	2.69
Ht (%)	29	21	16	21	18.15	1.46
dcp (%)	29	13	9	14	11.57	1.2
Hd (%)	29	30	22	33	29.33	2.29
I.O. (%)	29	16	12	19	14.3	1.68
Eye (%)	29	7	6	9	6.78	0.81

Paratypes

GABON: 12 ♂♂, 3 ♀♀, 16.1–23.5 mm SL, same collection data as for holotype (MRAC 2016-019-P-94-108); 13 ♂♂, 18.9–26.6 mm SL, upper Alowé River, 0.70581 S, 9.47592 E, 48 m a.s.l., field reference code CHRSP6-Alowé, 4 Mar. 2016, Laurent Chirio leg. (MRAC 2016-019-P-109-121).

Differential diagnosis

Differs from all other species in *A. alpha* group through combination of following features: head with very intense reddish colour, flanks bluish-orange, ventral region orange (vs never same colour pattern). Distinguished from all other species of *Chromaphyosemion* by an alpha-shaped mark on pre- and post-opercular region (vs no alpha-shaped mark, only some red macules).

Colouration of live males (Fig. 6E)

FLANKS AND VENTER. Flanks orange and bluish with two often visible dark lateral stripes, four lines of red dots from opercle to caudal peduncle. Ventral region orange, region between end of the anal fin and caudal peduncle is intense orange. Dorsal region brown and bluish, with two rows of paradorsal copper scales from the operculum to caudal peduncle with two lines of carmine red dots.

HEAD. Very intense red, premaxilla reddish, mandible reddish with black lower lips, reddish supraorbital region, reddish infraorbital region with black macule, postorbital region bluish-grey with black macule, orange pre-opercle with two black macules, orange opercle with one black macule; black macules on pre- and post-opercular regions, revealing alpha-shaped mark.

FINS. Dorsal fin orange, blue on posterobasal portion of fin, with rows of carmine red dots between rays, carmine red streaks on distal and posterobasal portions of fin. Orange acumen on apex. Anal fin orange, with blue zone near submarginal band, regular, red submarginal band, blue marginal band and

orange acumen. No punctuation. Caudal fin blue, orange on upper and lower lobes, with some red dots on median region, very long red streaks on median region and in upper and lower lobes, irregular, red submarginal band, blue marginal band and orange acumens. Pelvic fins orange with no punctuation, blue zone near submarginal band, submarginal red band, blue marginal band. Pectoral fins with orange reflections.

Colouration of live females (Fig. 6F)

FLANKS AND VENTER. Flanks pale beige, four lines of red dots from opercle to caudal peduncle, two dark lateral stripes, venter pale beige, white on basal portion. Two rows of paradorsal brown scales in dorsal region from operculum to caudal peduncle with red dots.

HEAD. Premaxilla brown, mandible yellow-orange with black lower lips, with brown supraorbital region, white infraorbital region with black macule, white postorbital region with one black macule, pale blue pre-opercle, orange opercle with two macules.

FINS. Dorsal fin yellow-green, orange on distal portion, with rows of carmine red dots between rays. Anal fin translucent, orange zone on distal portion, blue on edge. Caudal fin translucent, orange on ventral portion of fin, red dots on dorsal portion of the fin near caudal peduncle. Pelvic fins translucent with orange reflection, blue reflection on edges. Pectoral fins translucent.

Colouration of ethanol-preserved males and females (Fig. 6G–H)

FLANKS AND VENTER. Flanks of males grey with four lines of red dots, venter whitish. Two rows of paradorsal red scales in grey dorsal region, the operculum to caudal peduncle, with two visible greyish lateral stripes. Flanks of females beige with two grey lateral stripes, four lines of red dots from opercle to caudal peduncle, incomplete near the peduncle; venter yellowish, with two grey lateral stripes. Two rows of paradorsal red scales in brown dorsal region, from operculum to caudal peduncle.

HEAD. Male with whitish mandible with black lower lips (Alowé specimens: orange below mandible), infraorbital region with one red macule, orange pre-opercular region and opercle region, and opercle with two small black macules. Head of female with black lower lips, infraorbital region with red macule, orange opercle.

FINS. Dorsal fin of male light orange with red dots and some red streaks on edge. Anal fin orange with red submarginal band. Caudal fin orange on median portion, with red streaks on upper portion, and red submarginal band. Dorsal fin of female whitish, rows of red dots between rays. Anal fin whitish, with no red dots. Caudal fin whitish, with some red dots on upper portion, red streaks on edge.

Distribution and habitat

This species has been found at two localities, one in the Niengé River (Table 1, Fig. 1) flowing towards Lake Gomé, itself flowing into the lower Ogooué River, and one in the Alowé River, flowing towards Lake Alombié, also flowing into the lower Ogooué River. The species has never been found elsewhere and seems to be endemic to this part of the lower Ogooué hydrographic basin. The habitat of this species differs quite significantly from that of the five other species: all specimens were found in 3 to 5 metre wide rivers with fast running water, and sandy bottoms with no aquatic vegetation. They were hidden in tree roots along the riverbanks, or under dead leaves at the mouth of small secondary brooks, but they were not found in the small brooks themselves. This species was found in syntopy with an undescribed species of *Aphyosemion* and an undescribed species of *Plataplochilus*.

AGNÈSE J.-F. *et al.*, Six new species of *Aphyosemion* (Cyprinodontiformes, Aplocheilidae)**Table 8.** Meristic and morphometric data of *Aphyosemion rubrogaster* Chirio, Legros & Agnès sp. nov. See Table 3 for the abbreviations used.

Character	N	HT	Min	Max	Mean	SD
D	29	12	11	14	12,07	0.7
A	29	13	12	15	13.69	0.89
D/A	29	1	0	3	1.72	0.88
L.L.	29	27+1	23 (+1)	27 (+3)	25.1 (+1.83)	1.08 (+0.54)
pDor	29	13	11	13	12.07	0.65
TRAV	29	8	6	8	7	0.38
S.L.	29	25.2	16.1	26.6	21.13	2.65
T.L. (%)	29	124	111	130	120.68	4.47
P.D. (%)	29	57	47	65	56.87	3.78
P.A. (%)	29	53	22	62	54.03	6.99
P.V. (%)	29	44	39	52	45.75	2.98
Ht (%)	29	12	12	42	17.64	4.92
dcp (%)	29	17	9	17	11.31	1.58
Hd (%)	29	27	15	34	28.62	3.49
I.O. (%)	29	12	11	31	13.24	3.64
Eye (%)	29	7	6	11	7.76	1.04

Identification key of the *Aphyosemion* alpha group species

1. Anal fin without punctuation 2
 - Anal fin with punctuation of black or red dots 3
2. Flanks orange and bluish, ventral region orange, very intense orange portion between the end of the anal fin and the caudal peduncle, head highly red-orange
 - *A. rubrogaster* Chirio, Legros & Agnès sp. nov.
 - Flanks beige, ventral region orange or beige, blue iridescent scales between the end of the anal fin and the caudal peduncle 4
 - Flanks blue or purplish blue, ventral region orange or bluish white, blue iridescent scales between the end of the anal fin and the caudal peduncle 5
3. Anal fin with some red carmine macules, mauve on basal portion, orange on median portion, a yellowish portion near the submarginal band, red and regular submarginal band, blue marginal band, orange little acumen *A. barakoniense* Chirio, Legros & Agnès sp. nov.
 - Anal fin blue with orange portion on basal region, without punctuation except black dots on basal and/or postero-basal portions, without submarginal and marginal bands. Some individuals may have some small black dots instead of the submarginal band
 - *A. flavocyaneum* Chirio, Legros & Agnès sp. nov.
4. Two orange zones: orange portion on the opercle, venter beige with an orange portion above the pelvic fins, anal fin orange, blue near the basal portion and near the red submarginal band, dorsal fin orange, caudal fin blue with begin of upper and lower lobes light orange *A. pusillum* Chirio, Legros & Agnès sp. nov.
 - No delimited orange zone, opercle orange, ventral region entirely orange, anal fin orange with a blue zone upper the red submarginal band, dorsal fin blue with large orange zone on rays, caudal fin blue with an orange caudal peduncle *A. aurantiacum* Chirio, Legros & Agnès sp. nov.

5. Dorsal fin brown-orange, greenish on basal portion, red dots between rays, little orange apex, caudal fin light blue, some red dots on basal region, very long red streaks on distal region, red and irregular submarginal band, blue marginal band, little acumens light orange, flanks blue, ventral region orange *A. alpha* Huber, 1998
- Dorsal fin blue, orange distal portion near the apex, black dots between rays, very long orange acumen on the apex, caudal fin blue with orange portion on peduncle region, some red dots on basal region, very long red streaks on distal region, red and irregular submarginal band, blue marginal band, acumen orange very intense, flanks purplish blue, ventral region bluish white *A. flammulatum* Chirio, Legros & Agnès sp. nov.

Discussion

This high diversity present in the Wonga-Wongué Presidential Reserve is based on a single monophyletic group of species of the subgenus *Chromaphyosemion*. There are at least six different species in this small area (4300 km², approximately 80 km north to south and west to east). This diversity may be underestimated, however, because three drainages (Liamé, Sangatanga and Ngélié) have not yet been investigated. The simplest hypothesis to explain this pattern is to consider that all these species originated from vicariance events. The Wonga-Wongué Presidential Reserve is composed of coastal hills, rising to 284 m. These hills are ancient coastal dunes covered with a savannah and forest mosaic. Today, these hills separate rivers that were probably in contact with each other when the sea level was much lower. During the Pleistocene glaciation periods, the sea level dropped by more than 100 metres (Van Handel 1989). These level drops moved the coastline westward and allowed separate rivers to join and form new rivers or deltas. This has been demonstrated for example for the Seine and the Somme Rivers in France that joined in the English Channel during the cold periods (Antoine *et al.* 2000). The alternating cold and warm periods, corresponding to contact and separation periods for the rivers, may have generated multiple speciation events through accumulation of slight phenotypic differences (during isolation times) and reinforcement (during secondary contact) because of the ability of the female *Chromaphyosemion* to distinguish and preferentially mate with males of its own species and population (Kullmann & Klemme 2007).

Acknowledgements

The authors would like to thank Nicolas Hubert for his help with the genetic analysis. This work was financed by the Institut des Sciences de l'Evolution de Montpellier (genetic study) (ISEM), the Institut de Recherche pour le Développement (IRD) and the Agence Nationale des Parcs Nationaux (ANPN). The latter allowed us to collect specimens from the Presidential Reserve of Wonga-Wongué. We also thank Flore Koumba Pambo from CENAREST for her help in obtaining the research permits. DNA sequences were produced through molecular genetic analysis in the technical facilities of the Centre Méditerranéen de l'Environnement et de la Biodiversité (CeMEB). We are grateful to Thibault Cavelier de Cuverville for his valuable help in collecting the fish and to Kate Abernethy for the corrections made to the last proofs of this text.

References

- Agnès J.-F., Zentz F., Legros O. & Sellos D. 2006. Phylogenetic relationships and phylogeography of the killifish species of the subgenus *Chromaphyosemion* (Radda, 1971) in West Africa, inferred from mitochondrial DNA sequences. *Molecular Phylogenetics and Evolution* 40: 332–346. <https://doi.org/10.1016/j.ympev.2006.03.018>
- Agnès J.-F., Legros O., Cazaux B. & Estivals G. 2013. *Aphyosemion pamaense*, a new killifish species (Cyprinodontiformes: Nothobranchiidae) from Cameroon. *Zootaxa* 3670 (4): 516–530. <https://doi.org/10.11646/zootaxa.3670.4.6>

- Antoine P., Lautridou J.-P. & Laurent M. 2000. Long-term fluvial archives in NW France: response of the Seine and Somme Rivers to tectonic movements, climatic variations and sea-level changes. *Geomorphology* 33: 183–207. [https://doi.org/10.1016/S0169-555X\(99\)00122-1](https://doi.org/10.1016/S0169-555X(99)00122-1)
- Collier G. 2006. The genus *Aphyosemion*: taxonomic history and molecular phylogeny. *Journal of the American Killifish Association* 39 (5–6) : 147–168.
- Felsenstein J. 1985. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* 39: 783–791. <https://doi.org/10.2307/2408678>
- Huber J.H. 2000. Killi-Data 2000. *Updated Checklist of taxonomic Names, collecting Localities and bibliographic References of oviparous Cyprinodont Fishes (Cyprinodontiformes); in French, English, German and Spanish*. Société française d’Ichtyologie, Paris.
- Huber J.H. 2006. *Killi-Data Catalogue*. Paris, France.
- ICZN. 1999. *International Code of Zoological Nomenclature*. Fourth Edition. The International Trust for Zoological Nomenclature, London.
- Kimura M. 1980. A simple model for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *Journal of Molecular Evolution* 16: 111–120. <https://doi.org/10.1007/bf01731581>
- Kullmann H. & Klemme B. 2007. Female mating preference for own males on species and population level in *Chromaphyosemion* killifishes (Cyprinodontiformes, Nothobranchiidae). *Zoology* 110: 377–386. <https://doi.org/10.1016/j.zool.2007.08.001>
- Legros O., Zentz F. & Agnès J.-F. 2005. Description de deux nouveaux *Chromaphyosemion* (Teleostei: Aplocheilidae) du sud de la Plaine littorale camerounaise. *Killi-Contact* 33: 3–30.
- Nei M., Kumar S. & Takahashi K. 1998. The optimization principle in phylogenetic analysis tends to give incorrect topologies when the number of nucleotides or amino acids used is small. *Proceedings of National Academy of Sciences* 95: 12390–12397. <https://doi.org/10.1073/pnas.95.21.12390>
- Saitou N. & Nei M. 1987. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Molecular Biology and Evolution* 4: 406–425. <https://doi.org/10.1093/oxfordjournals.molbev.a040454>
- Sambrook J., Fritsch E.F. & Maniatis T. 1989. *Molecular Cloning: a Laboratory Manual*. Cold Spring Harbor Laboratory Press, New York.
- Scheel J.J. 1968. *Rivulins of the Old World*. TFH publications, New Jersey.
- Scheel J.J. 1990. *Atlas of Killifishes of the Old World*. TFH publications, New Jersey.
- Schwarz G. 1978. Estimating the dimension of a model. *Annals of Statistics* 6 (2): 461–464. <https://doi.org/10.1214/aos/1176344136>
- Sonnenberg R. 2000. The distribution of *Chromaphyosemion* Radda, 1971 (Teleostei: Cyprinodontiformes) on the Coastal Plains of West and Central Africa. In: Rheinwald G. (ed.) *Isolated Vertebrate Communities in the Tropics*. Proceedings of the 4th International Symposium, Bonn. *Bonner zoologische Monographien* 46: 79–94.
- Sonnenberg R. 2007a. Two new species of *Chromaphyosemion* (Cyprinodontiformes: Nothobranchiidae) from the coastal plain of Equatorial Guinea. *Ichthyological Exploration of Freshwaters* 18: 359–373.
- Sonnenberg R. 2007b. Description of three new species of the genus *Chromaphyosemion* Radda, 1971 (Cyprinodontiformes: Nothobranchiidae) from the coastal plains of Cameroon with a preliminary review of the *Chromaphyosemion splendopleure* complex. *Zootaxa* 1591: 1–38.

- Stamatakis A. 2014. RAxML Version 8: A tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 30: 1312–1313. <https://doi.org/10.1093/bioinformatics/btu033>
- Takahashi K. & Nei M. 2000. Efficiencies of fast algorithms of phylogenetic inference under the criteria of maximum parsimony, minimum evolution, and maximum likelihood when a large number of sequences are used. *Molecular Biology and Evolution* 17: 1251–1258. <https://doi.org/10.1093/oxfordjournals.molbev.a026408>
- Tamura K. & Nei M. 1993. Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. *Molecular Biology and Evolution* 10 (3): 512–526. <https://doi.org/10.1093/oxfordjournals.molbev.a040023>
- Tamura K., Peterson D., Peterson N., Stecher G., Nei M. & Kumar S. 2011. MEGA5: Molecular Evolutionary Genetics Analysis using Maximum Likelihood, Evolutionary Distance, and Maximum Parsimony Methods. *Molecular Biology and Evolution* 28: 2731–2739. <https://doi.org/10.1093/molbev/msr121>
- Ung V., Dubus G., Zaragüeta Bagils R. & Vignes-Lebbe R. 2010. Xper2: introducing e-taxonomy. *Bioinformatics* 26 (5): 703–704. <https://doi.org/10.1093/bioinformatics/btp715>
- Van Handel T.H. 1989. Late Quaternary sea-level changes and archaeology. *Antiquity* 63: 733–745.
- Völker M.E. 2006. *Karyotype Differentiation in Chromaphyosemion Killifishes (Cyprinodontiformes, Nothobranchiidae): Patterns, Mechanisms and Evolutionary Implications*. PhD Thesis, University of Bonn, Germany.
- Wildekamp R.H. 1993. *A World of Killies. Atlas of the Oviparous Cyprinodontiform Fishes of the World*. Volume 1. The American Killifish Association, Mishawaka, Indiana.
- Zhang J., Kapli P., Pavlidis P. & Stamatakis A. 2013. A general species delimitation method with applications to phylogenetic placements. *Bioinformatics* 29: 2869–2876. <https://doi.org/10.1093/bioinformatics/btt499>

Manuscript received: 18 January 2018

Manuscript accepted: 18 June 2018

Published on: 30 October 2018

Topic editor: Rudy Jocqué

Desk editor: Kristiaan Hoedemakers

Printed versions of all papers are also deposited in the libraries of the institutes that are members of the *EJT* consortium: Muséum national d'Histoire naturelle, Paris, France; Botanic Garden Meise, Belgium; Royal Museum for Central Africa, Tervuren, Belgium; Natural History Museum, London, United Kingdom; Royal Belgian Institute of Natural Sciences, Brussels, Belgium; Natural History Museum of Denmark, Copenhagen, Denmark; Naturalis Biodiversity Center, Leiden, the Netherlands; Museo Nacional de Ciencias Naturales-CSIC, Madrid, Spain; Real Jardín Botánico de Madrid CSIC, Spain; Zoological Research Museum Alexander Koenig, Bonn, Germany.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [European Journal of Taxonomy](#)

Jahr/Year: 2018

Band/Volume: [0471](#)

Autor(en)/Author(s): Agnese Jean-Francois, Chirio Laurent, Legros Olivier, Oslisly Richard, Bhe Herve Mve

Artikel/Article: [Unexpected discovery of six new species of Aphyosemion \(Cyprinodontiformes, Aplocheilidae\) in the Wonga-Wongué Presidential Reserve in Gabon 1-28](#)