

Contributions to Indian Myriangiaeae II

By J. S. Tendulkar,

Maharashtra Association for the Cultivation of Science, Poona—4 (India)

With three Figures

This is the 2nd contribution to this aspect of studies and presents three new species of *Myriangium* collected by the writer at Tungareshwar near Bassein, Maharashtra State, during the months of September—October 1969, parasitizing *Tamarindus indica* L., *Abutilon indicum* Sweet, and *Zyziphus rotundifolia* Lamk. since no species of *Myriangium* had been previously reported on the three host families, a detailed comparative study was undertaken with the type species in respect of habit, nature of stroma, morphological characters, host relationships and nature of parasitism.

A review of the literature bearing on the taxonomy of the genus *Myriangium* has already been presented by the writer (1969) in his 1st contribution of this series and needs no repetition.

The three species described below were compared with the type *M. duriae* and other Indian species and found to be significantly distinct in respect of habit, nature of stroma, arrangement of locules within the stroma and dimensions besides being collected on hitherto unreported hosts. These species are essentially phytopathogenic unlike the type species which is entomogenous in its parasitism. The species accordingly are accommodated as new taxa.

1. *Myriangium tamarindi* Tendulkar sp. nov. (Fig. 1).

Stromata black, rigid with distinctly raised margins having wavy base, generally uniloculate, hypostroma present. Locules allantoid, subiculum lacking, measuring 352—368 μ in length and 144—160 μ in height. Ascii globose to obovate, in uniascal cavities, arranged regularly in 2 to 3 tiers, 8-spored, measuring 28—44 μ \times 26—32 μ . Ascospores muriform with one vertical and 5 to 6 transverse septa, constricted at septa, light yellow, measuring 24—28 μ \times 7.6—8 μ .

Stromata dispersa, superficialia, ambitu orbicularia, tuberculiformia vel pulvinata, superne plus minusve convexa ex hypostromate circumcirca explanato et extenso, pseudoparenchymatico emersa, contextu pseudoparenchymatico, 352—368 μ diam., 144—160 μ crassa, plurilocularia, loculi irregulariter di- vel tritichi, monoascigeri; ascii subglobosi vel late ovoidei, sessiles, crassiuscule tunicati, 8-spori, 28—44 \times

26—32 μ ; sporeæ congregatae vel indistincte tristichæ, oblongæ, utrinque vix vel parum attenuatae, rectæ vel leniter curvulae, longitudinaliter septo unico, transverse septis 5—6 praeditæ, in medio plus minusve, ceterum non vel lenissime constrictæ, pallide luteolæ, 24—28 \times 7.6—8 μ .




Fig. 1. *Myriangium tamarindi* — a. Habit showing unilocate stroma (Schematic) — b. Stroma with ascii in uniascal cavities arranged in 2 to 3 tiers. — c. Ascus. — d. Ascospores.

Fig. 2. *Myriangium kamattii* — a. Habit showing multiloculate stroma (Schematic) — b. Stroma with allantoid locule and ascii in uniascal cavities packed in 2 to 3 tiers. — c. Ascus. — d. Ascospores.

Fig. 3. *Myriangium zyziphi* — a. Habit showing biloculate stroma (Schematic). — b. Stroma showing ascii, in uniascal cavities, scattered in the locule. — c. Ascus. — d. Ascospores.

Parasitic on the bark of *Tamarindus indica* L. collected by J. S. Tendulkar at Tungareshwar, Bassein, Maharashtra, India on 11. 9. 1969, M. A. C. S. Type No. 819.

2. *Myriangium kamati* Tendulkar sp. nov. (Fig. 2).

Stromata black, rigid with distinctly raised margin and wavy bottom, multiloculate, generally 3, hypostroma present. Locules allantoid with basal subiculum, measuring $416-624 \mu \times 272-320 \mu$. Ascii globose to obovate, in uniascal cavities, arranged regularly in 2 to 3 tiers, 8-spored, measuring $40-44 \mu \times 32-36 \mu$. Ascospores muri-form with 1 vertical and 4 to 5 transverse septa, constricted at septa, yellowish in colour, measuring $26-30 \mu \times 8 \mu$.

Stromata dispersa, superficialia, ambitu plus minusve orbicularia, tuberculiformia vel pulvinata, superne plus minusve convexa, contextu pseudoparenchymatico, ex hypostromate circumcirca explanato et plus minusve extenso pseudoparenchymatico emersa, $416-624 \times 272-320 \mu$, plurilocularia; loculi subregulariter di- vel tristichi, monoascigeri; asci subglobosi vel late ovoidei, sessiles, crassiuscule tunicati, 8-spori, $40-44 \times 32-36 \mu$; sporae congregatae, oblongae, utrinque obtusae, vix vel parum et abruptiuscule attenuatae, rectae vel leniter curvulae, longitudinaliter septo unco, transverse septis 5-6 praeditae, in medio plus minusve, ceterum non vel lenissime constrictae, luteolae, $26-30 \times 8 \mu$.

Parasitic on the bark of *Abutilon indicum* Sweet, collected by J. S. Tendulkar at Sateri, Bassein, Maharashtra, India, on 13. 10. 1969. M. A. C. S. Type No. 820.

The species is described after Prof. M. N. Kamat, Head of the Mycology Department, M. A. C. S., Poona 4, India, in recognition of his outstanding contributions to the Ascomycetes.

3. *Myriangium zyziphi* Tendulkar sp. nov. (Fig. 3).

Stromata black, rigid with wavy margins, biloculate, hypostroma present. Locules linearly disc-shaped, subiculum lacking, convex at the upper surface and concave towards the lower part, measuring $880-1056 \mu \times 160-240 \mu$. Ascii in uniascal cavities, scattered, globose to obovate, 8-spored, measuring $36-48 \mu \times 32-40 \mu$. Ascospores muri-form with 1 vertical and 4 transverse septa, constricted at septa, yellowish in colour, measuring $22-32 \mu \times 8 \mu$.

Stromata dispersa, superficialia, lineari-tuberculiformia vel elongato-pulvinata, superne convexa, inferne concava, contextu pseudoparenchymatico, ex hypostromate circumcirca explanato et plus minusve extenso, pseudoparenchymatico emersa, $880-1056 \times 160-240 \mu$, plurilocularia, loculi subregulariter di- vel tristichi, monoascigeri; asci plus minusve globosi vel late ovoidei, sessiles, crassiuscule tunicati, 8-spori, $36-48 \times 32-40 \mu$; sporae congregatae, oblongae, utrin-

que abruptiuscule attenuatae, tunc oblongo-fusoideae, rectae, raro inaequilaterae vel curvulae, longitudinaliter septo unico, transverse septis 4—5 praeditae, in medio plus minusve, ceterum non vel lenissime constrictae, luteolae, 22—32 × 8 μ.

Parasitic on the bark of *Zyziphus rotundifolia* Lamk. collected by J. S. Tendulkar at Sateri, Bassein, Maharashtra, India, on 13. 10. 1969. M. A. C. S. Type No. 821.

The genus is essentially known to parasitize scale insects. However, it is interesting to note that out of the nine species of *Myriangium* reported from India only one viz. *M. cinchonae* has been reported to be associated with scale insects.

This contribution brings the total number of species of *Myriangium* in India to 9 all of which are essentially parasitic on the bark of several plants of economic importance. The infection, however, appeared to be of sporadic nature with little or no damage to the hosts.

The type materials of the three new species are being deposited at Herb. Orientalis, New Delhi (India) and C. M. I., Kew, England.

Acknowledgement

The writer is deeply indebted to Prof. M. N. Kamat for his valuable guidance, to Dr. G. B. Deodikar, the Director, M. A. C. S. for laboratory and library facilities, to the Ministry of Education, Govt. of India for financial assistance and to Dr. F. Petrank for his valuable assistance in Latin rendering of the new species and to Shri D. N. Na gpure, the Artist of this Institute, for the preparation of figures.

References

1. Arx von, J. A., 1963: Die Gattungen der Myriangiales Persoonia, 2: 421—475.
2. Chona, B. L., R. L. Munjal, 1950: Notes on Indian Miscellaneous fungi. Ind. Phytopath 3: 105—116.
3. Miller, J. H., 1938: Studies in the development of the two *Myriangium* species and the systematic position of the order Myriangiales. Mycologia 30: 158—181.
4. Seshadri, V. S., 1967: Studies in Indian Ascomycetes. Ph. D. Thesis — University of Poona.
5. Tendulkar, J. S., 1969: Three new species of *Myriangium* from India. Sydowia (in Press).

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Sydowia](#)

Jahr/Year: 1970/1971

Band/Volume: [24](#)

Autor(en)/Author(s): Tendulkar J. S.

Artikel/Article: [Contributions to Indian Myriangiabeae II. 290-293](#)